

POLICY ITERATION FOR EXPLORATORY HAMILTON–JACOBI–BELLMAN EQUATIONS

HUNG VINH TRAN, ZHENHUA WANG, YUMING PAUL ZHANG

ABSTRACT. We study the policy iteration algorithm (PIA) for entropy-regularized stochastic control problems on an infinite time horizon with a large discount rate, focusing on two main scenarios. First, we analyze PIA with bounded coefficients where the controls applied to the diffusion term satisfy a smallness condition. We demonstrate the convergence of PIA based on a uniform $C^{2,\alpha}$ estimate for the value sequence generated by PIA, and provide a quantitative convergence analysis for this scenario. Second, we investigate PIA with unbounded coefficients but no control over the diffusion term. In this scenario, we first provide the well-posedness of the exploratory Hamilton–Jacobi–Bellman equation with linear growth coefficients and polynomial growth reward function. By such a well-posedness result we achieve PIA’s convergence by establishing a quantitative locally uniform $C^{1,\alpha}$ estimates for the generated value sequence.

Key words: Hamilton–Jacobi–Bellman equations, policy iteration algorithm, stochastic control, reinforcement learning, entropy regularization, convergence rate.

AMS subject classifications: 35F21, 60J60, 68W40, 93E20.

1. INTRODUCTION

Policy improvement involves updating the current strategy to enhance performance. This iterative process, known as a policy improvement algorithm (PIA, also called policy iteration algorithm), aims at converging towards an optimal policy through successive refinements. Rooted in Dynamic Programming, PIA plays an important role in Markov decision processes problems (MDPs) and reinforcement learning (RL), which could be dated back to Bellman [2] and Howard [15]. The convergence rate of PIA for an infinite time horizon was investigated in Puterman, Brumelle [25]. For discrete-time MDPs, PIA has been well explored and its convergence has been established under suitable conditions on the model parameters, see e.g., Puterman [24], Sutton, Barto [26], and Bertsekas [3], among many others.

In the framework of controlled ODEs or deterministic optimal control, the convergence of PIA has been studied in linear quadratic settings, see Abu-Khalaf, Lewis [1], Vrabie, Pastravanu, Abu-Khalaf, Lewis [30], and the references therein. Lee and Sutton [20] proved the convergence under certain regularity and fixed point assumptions. To overcome the ill-posedness of PIA for general controlled ODEs, Tang, Tran, and Zhang [27] proposed a semi-discrete scheme for the PIA and showed its general exponential convergence rate. Lee and Kim [21] incorporated a deep operator network with the scheme in [27] to numerically solve the PIA and the optimal control problems.

For stochastic control problems (without entropy-regularization), Krylov [19] and Puterman [23] showed that the optimal value is recovered under the PIA for a specific control problem with a compact space-time domain. Jacka and Mijatović [17] outlined a list of assumptions leading to PIA’s convergence towards optimal control, offering illustrative examples. Afterward, Kerimkulov, Šiška, and Szpruch [18] established the convergence and studied the convergence rate by assuming a certain regularity of the optimal value function and the control does not appear in the diffusion

The work of H. Tran is partially supported by NSF CAREER grant DMS-1843320 and a Vilas Faculty Early-Career Investigator Award. The work of Y. P. Zhang is partially supported by a start-up grant at Auburn University.

term. The convergence of PIA was further studied in some problems in mean-field games, see Cacace, Camilli, Goffi [4], and Camilli, Tang [5].

In the RL literature, it is now well-known that the entropy regularization (also termed the “softmax” criterion) encourages exploration of the unknown environment through measure-valued control strategies (i.e., relaxed controls). This approach prevents early settlement to suboptimal strategies, a problem known as the curse of optimality, for a detailed explanation, see Zhou [34]. With entropy regularization, Wang, Zariphopoulou, and Zhou [31] opened the door to incorporating continuous-time stochastic control problems into the exploratory framework of RL. The following so-called *exploratory* Hamilton-Jacobi-Bellman (HJB) equation thereby was introduced and first studied in a linear-quadratic setting in [31].

$$\rho v - \sup_{\pi \in \mathcal{P}(U)} \left\{ \int_U \left[b(x, u) \cdot Dv + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u) D^2 v(x)) + r(x, u) - \lambda \ln(\pi(u)) \right] \pi(u) du \right\} = 0. \quad (1.1)$$

Here, $\mathcal{P}(U)$ represents all probability densities on the action space U . Moreover, as the weight λ tends to zero, the optima and optimal relaxed strategy for an exploratory stochastic control problem converge to the optima and optimal feedback control for the corresponding standard stochastic control problem, as proved by Tang, Zhang, and Zhou [28]. Such entropy-regularization formulation has been further extended to various settings, such as mean-variance problems (Wang, Zhou [32]), stopping problems (Dong [9]), and mean-field games/mean-field controls (Guo, Xu, Zariphopoulou [14], Firoozi, Jaimungal [12], Frikha, Germain, Laurière, Pham, Song [13], Wei, Yu [33]).

The PIA for the HJB equation (1.1) can be written as follows in a PDE framework.

Algorithm Policy Iteration Algorithm for (1.1)

Initialization: Take suitable $v^0 \in \mathcal{C}^2$.

Iteration: For $n \in \mathbb{N}$, **do**

$$\text{Set } \pi^n(x, u) := \frac{\exp\left(\frac{1}{\lambda} b(x, u) \cdot Dv^{n-1}(x) + 2^{-1} \text{tr}(\sigma \sigma^T(x, u) D^2 v^{n-1}(x)) + r(x, u)\right)}{\int_U \exp\left(\frac{1}{\lambda} b(x, u) \cdot Dv^{n-1}(x) + 2^{-1} \text{tr}(\sigma \sigma^T(x, u) D^2 v^{n-1}(x)) + r(x, u)\right) du}; \quad (1.2)$$

Solve $v^n(x)$ for

$$\rho v^n - \int_U \left[b(x, u) \cdot Dv^n + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u) D^2 v^n(x)) + r(x, u) - \lambda \ln(\pi^n(u)) \right] \pi^n(u) du = 0. \quad (1.3)$$

For entropy-regularized PIA, [32] demonstrated its policy improvement property for linear-quadratic mean-variance problems and shows that optimal solutions can be achieved within two iterations due to the Gaussian densities of the optimal feedback control in this setting. [9] introduced the PIA for entropy-regularized stopping problems under a finite horizon and proved its convergence, where the underlying process is a geometric Brownian motion. Notably, the policy improvement property may fail when seeking Nash equilibrium strategies in a time-inconsistent stochastic control problem, as discussed in Dai, Dong, and Jia [8]. More recently, Huang, Wang, and Zhou [16] proved a qualitative convergence result with bounded coefficients when the diffusion term is not controlled (i.e., when σ is independent of u).

Despite these developments, the general convergence of Algorithm 1 remains largely unexplored, especially when the coefficients b, σ, r are unbounded and controls appear in the diffusion term. The first question concerns the well-posedness of PIA: whether the updated control π^n in (1.2) is well-defined and whether the elliptic equation (1.3) admits a unique solution. Once this is resolved, the next goal is to investigate the convergence of the generated sequence $\{v^n\}_n$ to the optimal solution of the entropy-regularized stochastic control, which should solve the HJB equation (1.1). As indicated in (1.2), entropy regularization makes the policy sequence $\{\pi^n\}_n$ less singular, but it

introduces second-order derivatives of the value sequences into the entropy of the policy sequence. This, in turn, makes achieving the compactness of the policy sequence more challenging.

In this paper, we investigate PIA with sufficiently large discounting rates in two scenarios. Firstly, we study PIA with bounded coefficients, allowing the control to influence the diffusion terms. This generalization is significant and challenging, as most literature on PIA considers problems with controls in the source or drift terms, resulting in linear or semilinear partial differential equations. Specifically, in their iterations, the updated policies are independent of the second-order derivatives of the solutions. In contrast, in our setting, the limiting equation is fully nonlinear, and the iterated controls depend also on the second derivatives.

So far, we have proven the convergence of PIA under the condition that the control has a small effect on the diffusion. The key step is to obtain a uniform $\mathcal{C}^{2,\alpha}$ estimate for the solutions in the iteration (see Theorem 2.1). This result can be viewed as a version of the Evans-Krylov theorem for PIA. The smallness assumption is essential in the proof of the uniform regularity result, as there is no obvious regularizing effect from the iteration process. It is unclear whether PIA converges without the smallness assumption and regardless of uniform regularity. Furthermore, we provide quantitative results on the convergence in Theorem 3.2 and Remark 3.1. In particular, Remark 3.1 gives an exponential convergence rate for the result in [16] when the diffusion term is not controlled.

The second objective is to investigate PIA with unbounded coefficients in \mathbb{R}^d , where the controls only appear in the lower-order terms. In this scenario, we allow the coefficients b and σ to exhibit linear growth in x , and the reward function r to have polynomial growth, encompassing linear quadratic settings. It is important to note that under such settings, the well-posedness of solutions to elliptic equations or PIA is not guaranteed without additional conditions. Specifically, ρ needs to be large, and our results are optimal in terms of the growth rate of r (see Proposition 4.1 and Remark 4.1). To show the convergence of PIA, we obtain quantitative locally uniform $\mathcal{C}^{1,\alpha}$ estimates for the solutions. Here, the analysis is localized, and the estimates must be done carefully with explicit dependence on the size of the coefficients.

After our paper was finished, we learned that Ma, Wang, and Zhang [22] obtained some very interesting related results independently by using a different approach when the diffusion term is not controlled. Specifically, the authors employed Feynman-Kac type probabilistic representation formulas for solutions of the iterative PDEs and their derivatives. By this method, they first provided a simple proof of the convergence result in [16], and then achieved an exponential convergence rate in the infinite horizon model with a large discount factor (which is essentially the same as Remark 3.1) and in the finite horizon model.

1.1. Model formulation. Denote by $|\cdot|$ the Euclidean norm. For a fixed dimension $d \in \mathbb{N}$, we denote by $B_R(x)$ the open ball centered at $x \in \mathbb{R}^d$ with radius R , and we simply write B_R for $B_R(0)$. We further use \mathcal{S}^d to denote the set of all symmetric, non-negative definite matrices of size d . We use \mathbb{I}_d to denote the identity matrix of size d .

For a multi-index $a = (a_1, \dots, a_m)$, denote $|a|_1 = \sum_{i=1}^m a_i$. Given a non-empty connected open set $V \subseteq \mathbb{R}^d$, for $k \in \mathbb{N} \cup \{0\}$ and $\alpha \in (0, 1)$, we define

$$[w]_{\mathcal{C}^{k,\alpha}(V)} := \sup_{x,y \in V, |x-y| \leq 1, |a|_1 = k} \frac{|D^a w(x) - D^a w(y)|}{|x-y|^\alpha}, \quad \|w\|_{L^\infty(V)} := \sup_{x \in V} |w(x)|,$$

$$\llbracket w \rrbracket_{\mathcal{C}^k(V)} := \sup_{|a|_1 = k} \|D^a w\|_{L^\infty(V)}, \quad \|w\|_{\mathcal{C}^{k,\alpha}(V)} := \sum_{0 \leq j \leq k} (\llbracket w \rrbracket_{\mathcal{C}^j(V)} + [w]_{\mathcal{C}^{j,\alpha}(V)}),$$

and

$$\|w\|_{\mathcal{C}^k(V)} := \sum_{0 \leq j \leq k} \llbracket w \rrbracket_{\mathcal{C}^j(V)},$$

and denote by $\mathcal{C}^{k,\alpha}(V)$ the set of all w such that $\|w\|_{\mathcal{C}^{k,\alpha}(V)} < \infty$. For any vector-valued or matrix-valued function $w = (w_{ij}(x))$, we denote

$$|w(x)| := \max_{ij} |w_{ij}(x)| \quad \text{and} \quad \|w\|_{L^\infty(V)} := \sup_{x \in V} |w(x)|.$$

When V equals to the whole space \mathbb{R}^d , we simply write

$$\begin{aligned} [w]_{k,\alpha} &= [w]_{\mathcal{C}^{k,\alpha}(\mathbb{R}^d)}, \quad \|w\|_\infty = \|w\|_{L^\infty(\mathbb{R}^d)}, \quad \llbracket w \rrbracket_k = \llbracket w \rrbracket_{\mathcal{C}^k(\mathbb{R}^d)}, \\ \|w\|_{k,\alpha} &= \|w\|_{\mathcal{C}^{k,\alpha}(\mathbb{R}^d)}, \quad \|w\|_k = \|w\|_{\mathcal{C}^k(\mathbb{R}^d)}, \end{aligned}$$

For any $q \geq 1$, we will also use the Sobolev norm $\|\cdot\|_{W^{k,q}(V)}$.

Fix positive integers d, m , and l . Consider an m -dimensional Brownian motion and $\mathcal{F} = (\mathcal{F}_t)_{t \geq 0}$ as the natural filtration generated by $(W_t)_{t \geq 0}$, i.e., $\mathcal{F}_t = \sigma(W_s, 0 \leq s \leq t)$ for all $t \geq 0$. Let U be a bounded domain in \mathbb{R}^l with its Lebesgue measure $|U| \in (0, \infty)$. Without loss of generality, we simply assume $|U| = 1$ throughout the paper. Denote by $\mathcal{P}(U)$ the set of all probability density functions on U . Given a relaxed control $\pi = (\pi_t)_{t \geq 0}$, which is \mathcal{F} -adapted and $\pi_t \in \mathcal{P}(U)$ for all $t \geq 0$, we consider the controlled process

$$dX_t^\pi = \left(\int_U b(X_t^\pi, u) \pi_t(u) du \right) dt + \sqrt{\left(\int_U \sigma(X_t^\pi, u) \sigma(X_t^\pi, u)^T \pi_t(u) du \right)} dW_t,$$

with $X_0 = x \in \mathbb{R}^d$ and functions $b(x, u) = (b_1(x, u), \dots, b_d(x, u)) : \mathbb{R}^d \times U \rightarrow \mathbb{R}^d$, $\sigma(x) = (\sigma_{ij}(x))_{i,j} : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$. The value function under π is given by

$$V^\pi(x) := \mathbb{E}_x \left[\int_0^\infty e^{-\rho t} \left(\int_U r(X_t^\pi, u) \pi_t(u) du - \lambda \int_U \ln(\pi_t(u)) \pi_t(u) du \right) dt \right].$$

For any $x, p \in \mathbb{R}^d$, $X \in \mathcal{S}^d$ and $\pi \in \mathcal{P}(U)$, define

$$F_\pi(x, p, X) := \int_U \left[r(x, u) + b(x, u) \cdot p + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u) X) - \lambda \ln(\pi(u)) \right] \pi(u) du. \quad (1.4)$$

and

$$F(x, p, X) := \sup_{\pi \in \mathcal{P}(U)} F_\pi(x, p, X), \quad (1.5)$$

By Dynamic Programming argument, $v^*(x) := \sup_\pi V^\pi(x)$ is a viscosity solution to the HJB equation (1.1), which is now rewritten as

$$\rho v - F(x, Dv, D^2v) = 0 \quad (1.6)$$

Given $x, p \in \mathbb{R}^d$, $X \in \mathcal{S}^d$, set

$$\Gamma(x, p, X)(u) := \frac{\exp\left(\frac{1}{\lambda}(r(x, u) + b(x, u) \cdot p + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u) X))\right)}{\int_U \exp\left(\frac{1}{\lambda}(r(x, u') + b(x, u') \cdot p + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u') X))\right) du'}. \quad (1.7)$$

We claim that $\pi(u) := \Gamma(x, p, X)(u)$ is the maximizer of (1.5) (see the variational principle in [10, Proposition 1.4.2]). Indeed, let $\phi(u)$ be such that $\int_U \phi(u) du = 0$ and $\pi + \varepsilon \phi \geq 0$ for all $\varepsilon \in (-1, 1)$. Then $\varepsilon = 0$ is a critical point of $G(\pi + \varepsilon \phi) := F_{\pi + \varepsilon \phi}(x, p, X)$, and the first variation of G satisfies

$$\int_U \left[r(x, u) + b(x, u) \cdot p + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u) X) - \lambda \ln(\pi(u)) \right] \phi(u) du = 0.$$

This holds for all such ϕ , so $r(x, u) + b(x, u) \cdot p + \frac{1}{2} \text{tr}(\sigma \sigma^T(x, u) X) - \lambda \ln(\pi(u))$ is constant in u . Using that $\int_U \pi(u) du = 1$ yields the formula (1.7).

In particular, we have

$$F_{\pi^*}(x, Dv^*, D^2v^*) = F(x, Dv^*, D^2v^*) \quad \text{with} \quad \pi^*(x, u) := \Gamma(x, Dv^*(x), D^2v^*(x))(u). \quad (1.8)$$

Recall the PIA, and π^n defined in (1.2) can be rewritten as

$$\begin{aligned}\pi^n(x, u) &= \Gamma(x, Dv^{n-1}(x), D^2v^{n-1}(x))(u) \\ &= \arg \max_{\pi(\cdot) \in \mathcal{P}(U)} F_\pi(x, Dv^{n-1}(x), D^2v^{n-1}(x)),\end{aligned}\tag{1.9}$$

and (1.3) becomes

$$\rho v^n - F_{\pi^n}(x, Dv^n, D^2v^n) = 0.\tag{1.10}$$

We will require ρ to be large in both the bounded and unbounded settings, which is essential for better regularity of the solutions. This is known in the literature, and we refer readers to [29, Theorem 2.8], which discussed the first-order case.

1.2. Organization of the paper. The paper is organized as follows. Sections 2 and 3 are dedicated to the first objective for bounded coefficients where the controls applied to the diffusion term satisfy a smallness condition. In Section 2, Theorem 2.1 demonstrates the locally uniform $\mathcal{C}^{2,\alpha}$ bound for the sequence $\{v^n\}_n$ generated by PIA. Section 3 establishes the convergence of PIA in Theorem 3.1 and provides a quantitative result in Theorem 3.2 for this scenario. Section 4 studies the well-posedness of the nonlinear elliptic equation (1.6) for the case of unbounded coefficients. Section 5 then addresses the well-posedness of PIA and its convergence for the second scenario with unbounded coefficients where the diffusion term is independent of the control.

2. UNIFORM $\mathcal{C}^{2,\alpha}$ ESTIMATES FOR BOUNDED EQUATIONS

We first estimate several norms of solutions with explicit dependence on $\rho \geq 1$.

Lemma 2.1. *Let $\rho \geq 1$, and let v be a solution to*

$$\rho v(x) - \tilde{r}(x) - \tilde{b}(x) \cdot Dv(x) - 2^{-1} \operatorname{tr}(\tilde{\Sigma}(x)D^2v(x)) = 0.$$

Assume that $\tilde{\Sigma} \geq \mathbb{I}_d/C_0$ for some $C_0 > 0$. Then there exists an increasing function $\eta : [1, \infty) \rightarrow [1, \infty)$ independent of ρ such that if

$$\|\tilde{r}(\cdot)\|_{0,\alpha}, \|\tilde{b}(\cdot)\|_{0,\alpha}, \|\tilde{\Sigma}(\cdot)\|_{0,\alpha} \leq A \quad \text{for some } A \geq 1 \text{ and } \alpha \in (0, 1),\tag{2.1}$$

we have

$$\rho \|v\|_\infty, \rho^{1-\frac{\alpha}{2}} [v]_{0,\alpha}, \rho^{\frac{1}{2}-\frac{\alpha}{2}} [v]_{1,\alpha}, \rho^{-\frac{\alpha}{2}} [v]_{2,\alpha}, \rho^{\frac{1}{2}} [v]_1, [v]_2 \leq \eta(A).\tag{2.2}$$

Proof. By comparing v with $\pm \|\tilde{r}\|_\infty / \rho$, we obtain $\|v\|_\infty \leq \|\tilde{r}\|_\infty / \rho$.

To prove (2.2), we apply a scaling argument. Note that $w(x) := v(x/\sqrt{\rho})$ solves

$$w - \frac{\tilde{r}(x/\sqrt{\rho})}{\rho} - \frac{\tilde{b}(x/\sqrt{\rho})}{\sqrt{\rho}} \cdot Dw - \frac{1}{2} \operatorname{tr}(\tilde{\Sigma}(x/\sqrt{\rho})D^2w) = 0.$$

It is direct to see that the α -Hölder norms of

$$\frac{\tilde{r}(x/\sqrt{\rho})}{\rho}, \quad \frac{\tilde{b}(x/\sqrt{\rho})}{\sqrt{\rho}}, \quad \tilde{\Sigma}(x/\sqrt{\rho})$$

are non-increasing as $\rho \geq 1$ increases, and also that $\tilde{\Sigma}(x/\sqrt{\rho}) \geq \mathbb{I}_d/C_0$ is preserved. Thus, the famous interior Schauder estimates state

$$\|w\|_{2,\alpha} \leq C_A (\|w\|_\infty + \rho^{-1} \|\tilde{r}(\cdot/\sqrt{\rho})\|_{0,\alpha}) \leq C_A / \rho$$

where C_A only depends on A and the dimension. This implies that

$$[v]_{0,\alpha} \leq C_A \rho^{-1+\frac{\alpha}{2}}, \quad [v]_{1,\alpha} \leq C_A \rho^{-\frac{1}{2}+\frac{\alpha}{2}}, \quad [v]_{2,\alpha} \leq C_A \rho^{\frac{\alpha}{2}}, \quad [v]_1 \leq C_A \rho^{-\frac{1}{2}}, \quad [v]_2 \leq C_A.$$

□

Remark 2.1. We note that the bound (2.2) is essentially optimal thanks to the scaling approach. In particular, the bound $[v]_{2,\alpha} \leq C_A \rho^{\frac{\alpha}{2}}$ cannot be improved in general. Here is another way to see it. Indeed, (2.2) gives $\rho \|v\|_{0,\alpha} \leq C_A \rho^{\frac{\alpha}{2}}$. Hence, by applying the interior Schauder estimates directly to v , we get

$$\|v\|_{2,\alpha} \leq C_A (\rho \|v\|_{0,\alpha} + \|\tilde{r}\|_{0,\alpha}) \leq C_A \rho^{\frac{\alpha}{2}}.$$

Recall $\Gamma(x, p, X)(u)$ from (1.7), and define

$$\pi(x, u) := \Gamma(x, Dv, D^2v)(u).$$

Denoting $\Sigma = \sigma\sigma^T$, we assume that there exists $C_0 \geq 1$ such that uniformly for all $u \in U$,

$$\Sigma(\cdot, u) \geq \mathbb{I}_d/C_0, \quad \|r(\cdot, u)\|_{0,\alpha}, \|b(\cdot, u)\|_{0,\alpha}, \|\Sigma(\cdot, u)\|_{0,\alpha} \leq C_0. \quad (2.3)$$

We have the following estimates for π and the entropy.

Lemma 2.2. Let v satisfy (2.2) with A_1 in place of $\eta(A)$. Assume (2.3), $|U| = 1$, $\rho \geq 1$, and that there exist $\Sigma_0(x)$ and $\varepsilon_0, \varepsilon_1 \in (0, 1)$ such that for $M(x, u) := \Sigma(x, u) - \Sigma_0(x)$,

$$\sup_{u \in U} \|M(\cdot, u)\|_{\infty} \leq \varepsilon_0, \quad \sup_{u \in U} [M(\cdot, u)]_{0,\alpha} \leq \varepsilon_1. \quad (2.4)$$

Then there exists C depending only on λ, C_0 such that $\pi(x, u) = \Gamma(x, Dv, D^2v)(u)$ satisfies

$$\begin{aligned} \|\pi(\cdot, u)\|_{\infty} &\leq \exp \left[C(1 + A_1(\rho^{-\frac{1}{2}} + \varepsilon_0)) \right], \\ [\pi(\cdot, u)]_{0,\alpha} &\leq \left(1 + A_1(\rho^{\frac{\alpha}{2}-\frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}}\varepsilon_0) \right) \exp \left[C(1 + A_1(\rho^{-\frac{1}{2}} + \varepsilon_0)) \right], \\ \left\| \lambda \int_U \pi(\cdot, u) \ln \pi(\cdot, u) du \right\|_{\infty} &\leq C \left(1 + A_1(\rho^{-\frac{1}{2}} + \varepsilon_0) \right), \\ \left[\lambda \int_U \pi(\cdot, u) \ln \pi(\cdot, u) du \right]_{0,\alpha} &\leq \exp \left[C \left(1 + A_1(\rho^{\frac{\alpha}{2}-\frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}}\varepsilon_0) \right) \right]. \end{aligned}$$

Proof. Let us write

$$f(x, u) = \frac{1}{\lambda} \left(r(x, u) + b(x, u) \cdot Dv + \frac{1}{2} \text{tr}(\Sigma(x, u)D^2v) \right),$$

and

$$g(x, u) = \frac{1}{\lambda} \left(r(x, u) + b(x, u) \cdot Dv + \frac{1}{2} \text{tr}((\Sigma(x, u) - \Sigma_0(x))D^2v) \right).$$

It is easy to see that

$$\pi(x, u) = \frac{\exp(f(x, u))}{\int_U \exp(f(x, u')) du'} = \frac{\exp(g(x, u))}{\int_U \exp(g(x, u')) du'}. \quad (2.5)$$

By the assumptions of (2.3), (2.4) and that (2.2) holds with A_1 in place of $\eta(A)$, it follows that for some $C = C(C_0, \lambda)$,

$$|g(x, u)| \leq C(1 + \rho^{-\frac{1}{2}}A_1 + \varepsilon_0A_1). \quad (2.6)$$

Therefore, also using $|U| = 1$, we get

$$\exp \left[-C(1 + \rho^{-\frac{1}{2}}A_1 + \varepsilon_0A_1) \right] \leq \pi(x, u) \leq \exp \left[C(1 + \rho^{-\frac{1}{2}}A_1 + \varepsilon_0A_1) \right] \quad (2.7)$$

Next, note that

$$\begin{aligned} |\pi(x, u) - \pi(y, u)| &= \left| \frac{\exp(g(x, u)) \int_U \exp(g(y, u')) du' - \exp(g(y, u)) \int_U \exp(g(x, u')) du'}{\int_U \exp(g(x, u')) du' \int_U \exp(g(y, u')) du'} \right| \\ &\leq \left| \frac{\int_U \exp(g(y, u')) - \exp(g(x, u')) du'}{\int_U \exp(g(y, u')) du'} \right| \pi(x, u) + \left| \frac{\exp(g(x, u)) - \exp(g(y, u))}{\int_U \exp(g(y, u')) du'} \right|. \end{aligned}$$

Because (2.2), (2.3) and (2.4) yield

$$\begin{aligned} [g(\cdot, u)]_{0,\alpha} &\leq \lambda^{-1} ([r]_{0,\alpha} + [b \cdot Dv]_{0,\alpha} + [\Sigma - \Sigma_0]_{0,\alpha} \|v\|_2 + \|\Sigma - \Sigma_0\|_\infty \|v\|_{2,\alpha}) \\ &\leq C \left(1 + A_1 \rho^{\frac{\alpha}{2} - \frac{1}{2}} + A_1 \varepsilon_1 + A_1 \rho^{\frac{\alpha}{2}} \varepsilon_0 \right), \end{aligned} \quad (2.8)$$

also using (2.6) and (2.7), we obtain

$$[\pi(\cdot, u)]_{0,\alpha} \leq \left(1 + A_1 (\rho^{\frac{\alpha}{2} - \frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}} \varepsilon_0) \right) \exp \left[C (1 + A_1 (\rho^{-\frac{1}{2}} + \varepsilon_0)) \right].$$

By (2.7) again and that π is a probability distribution, we get

$$\int_U |\ln \pi(x, u)| \pi(x, u) du \leq \|\ln \pi(\cdot, \cdot)\|_\infty \leq C \left(1 + A_1 (\rho^{-\frac{1}{2}} + \varepsilon_0) \right).$$

For the last claim, note that

$$\begin{aligned} &\left| \int_U \pi(x, u) \ln \pi(x, u) du - \int_U \pi(y, u) \ln \pi(y, u) du \right| \\ &\leq \sup_{u \in U} |\ln \pi(x, u) - \ln \pi(y, u)| + \sup_{u \in U} \left| \frac{\pi(y, u)}{\pi(x, u)} - 1 \right| |\ln \pi(y, u)|. \end{aligned}$$

So let us estimate the right-hand side two terms below. Because of (2.5) and (2.8), we get

$$\begin{aligned} |\ln \pi(x, u) - \ln \pi(y, u)| &= \left| g(x, u) - g(y, u) - \ln \int_U e^{g(x, u') - g(y, u')} du' \right| \\ &\leq C \left(1 + A_1 (\rho^{\frac{\alpha}{2} - \frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}} \varepsilon_0) \right) |x - y|^\alpha. \end{aligned}$$

To estimate $\left| \frac{\pi(y, u)}{\pi(x, u)} - 1 \right|$, let us assume without loss of generality that $\pi(y, u) \geq \pi(x, u)$. Then since $|g(x, u) - g(y, u)| \leq [g(\cdot, u)]_{0,\alpha} |x - y|^\alpha$ and $|U| = 1$, we find

$$\begin{aligned} \left| \frac{\pi(y, u)}{\pi(x, u)} - 1 \right| &= \frac{\exp[g(y, u)] \int_U \exp[g(x, u')] du'}{\int_U \exp[g(y, u')] du' \exp[g(x, u)]} - 1 \\ &\leq \frac{\exp[g(y, u)] \int_U \exp[g(y, u') + [g(\cdot, u)]_{0,\alpha} |x - y|^\alpha] du'}{\int_U \exp[g(y, u')] du' \exp[g(x, u)]} - 1 \\ &= \exp[g(y, u) - g(x, u)] \exp \left[\sup_{u \in U} [g(\cdot, u)]_{0,\alpha} |x - y|^\alpha \right] - 1 \\ &\leq \exp \left[C \left(1 + A_1 (\rho^{\frac{\alpha}{2} - \frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}} \varepsilon_0) \right) \right] |x - y|^\alpha, \end{aligned}$$

for all $|x - y| \leq 1$. Finally, combining these with (2.7), we obtain

$$\left| \int_U (\pi(x, u) \ln \pi(x, u) - \pi(y, u) \ln \pi(y, u)) du \right| \leq \exp \left[C \left(1 + A_1 (\rho^{\frac{\alpha}{2} - \frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}} \varepsilon_0) \right) \right] |x - y|^\alpha. \quad \square$$

Recall that, by (1.10), v^n satisfies the linear parabolic equation:

$$\rho v^n(x) - r^n(x) - b^n(x) \cdot Dv^n(x) - 2^{-1} \operatorname{tr}(\Sigma^n(x) D^2 v^n(x)) + \lambda \mathcal{H}^n(x) = 0, \quad (2.9)$$

where π^n is given in (1.9), and

$$\begin{aligned} r^n(x) &:= \int_U r(x, u) \pi^n(x, u) du, & b^n(x) &:= \int_U b(x, u) \pi^n(x, u) du, \\ \Sigma^n(x) &:= \int_U \sigma \sigma^T(x, u) \pi^n(x, u) du, & \mathcal{H}^n(x) &:= \int_U \ln(\pi^n(x, u)) \pi^n(x, u) du. \end{aligned} \quad (2.10)$$

We can prove the following result.

Theorem 2.1. *Let $\alpha \in (0, 1)$, and assume (2.3), (2.4) and $|U| = 1$. Then there exists $A_1 \geq 1$ depending only on $\|v^0\|_{2,\alpha}$, C_0 , λ and η from Lemma 2.1 such that if*

$$\rho \geq A_1^{2/(1-\alpha)}, \quad \varepsilon_0 A_1 \rho^{\frac{\alpha}{2}} \leq 1 \quad \text{and} \quad \varepsilon_1 A_1 \leq 1, \quad (2.11)$$

we have for v^n from PIA,

$$\|v^n\|_2 \leq A_1 \quad \text{and} \quad \rho^{-\frac{\alpha}{2}} [v^n]_{2,\alpha} \leq A_1 \quad \text{for all } n \geq 1.$$

We note that both ε_0 and ε_1 are quantifiable from the proof using Schauder estimates, and ε_1 can be chosen independently of ρ . In other words, $M(x, u) = \Sigma(x, u) - \Sigma_0(x)$ needs to be small for large ρ , although it is allowed to have certain oscillations independent of ρ .

Proof. Let $A \geq 1$ to be determined depending only on λ and (2.3) and let $A_1 = \eta(A) > 0$. Since $v^0 \in \mathcal{C}^{2,\alpha}(\mathbb{R}^d)$, then we can have that the Hölder norms of $r^1, b^1, \Sigma^1, \mathcal{H}^1$ are bounded by A , i.e. (2.1) holds with $r^1 - \lambda \mathcal{H}^1, b^1, \Sigma^1$ in place of $\tilde{r}, \tilde{b}, \tilde{\Sigma}$. Thus Lemma 2.1 implies that

$$\rho \|v^1\|_\infty, \rho^{1-\frac{\alpha}{2}} [v^1]_{0,\alpha}, \rho^{\frac{1}{2}-\frac{\alpha}{2}} [v^1]_{1,\alpha}, \rho^{-\frac{\alpha}{2}} [v^1]_{2,\alpha}, \rho^{\frac{1}{2}} [v^1]_1, [v^1]_2 \leq A_1.$$

Let us assume for induction that for some $n \geq 2$, the following holds

$$\rho \|v^{n-1}\|_\infty, \rho^{1-\frac{\alpha}{2}} [v^{n-1}]_{0,\alpha}, \rho^{\frac{1}{2}-\frac{\alpha}{2}} [v^{n-1}]_{1,\alpha}, \rho^{-\frac{\alpha}{2}} [v^{n-1}]_{2,\alpha}, \rho^{\frac{1}{2}} [v^{n-1}]_1, [v^{n-1}]_2 \leq A_1.$$

We now prove that the above inequalities all hold with v^n in place of v^{n-1} .

It follows from Lemma 2.2 and $|U| = 1$ that

$$\begin{aligned} \|r^n(\cdot)\|_{0,\alpha} &\leq \|r(\cdot)\|_{0,\alpha} + \|r(\cdot)\|_\infty \sup_{u \in U} [\pi^n(\cdot, u)]_{0,\alpha} \\ &\leq \left(1 + A_1(\rho^{\frac{\alpha}{2}-\frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}} \varepsilon_0)\right) \exp \left[C(1 + A_1(\rho^{-\frac{1}{2}} + \varepsilon_0)) \right], \end{aligned}$$

where C only depending on λ and C_0 in (2.3). Since $A_1 = \eta(A)$, (2.11) and the above yield

$$\|r^n(\cdot)\|_{0,\alpha} \leq 4 \exp(3C) \leq A/2$$

if A is sufficiently large compared with C . For the entropy, Lemma 2.2 and (2.11) yield

$$\left\| \lambda \int_U \pi^n(\cdot, u) \ln \pi^n(\cdot, u) du \right\|_{0,\alpha} \leq \exp \left[C \left(1 + A_1(\rho^{\frac{\alpha}{2}-\frac{1}{2}} + \varepsilon_1 + \rho^{\frac{\alpha}{2}} \varepsilon_0)\right) \right] \leq \exp(4C) \leq A/2$$

if A is sufficiently large depending only on C . Thus, $\|r^n - \lambda \mathcal{H}^n\|_{0,\alpha} \leq A$. Similarly, can we get

$$\|b^n(\cdot)\|_{0,\alpha}, \quad \|\Sigma^n(\cdot)\|_{0,\alpha} \leq A.$$

Finally, we apply Lemma 2.1 to conclude that

$$\rho \|v^n\|_\infty, \rho^{1-\frac{\alpha}{2}} [v^n]_{0,\alpha}, \rho^{\frac{1}{2}-\frac{\alpha}{2}} [v^n]_{1,\alpha}, \rho^{-\frac{\alpha}{2}} [v^n]_{2,\alpha}, \rho^{\frac{1}{2}} [v^n]_1, [v^n]_2 \leq \eta(A) = A_1.$$

Since the constants C, A, A_1 are independent of n , the proof is finished by induction. \square

3. CONVERGENCE FOR PIA WITH BOUNDED COEFFICIENTS

3.1. Convergence for uniform $\mathcal{C}^{2,\alpha}$ solutions. Theorem 2.1 has proved that, under certain conditions, v^n from PIA are uniformly $\mathcal{C}^{2,\alpha}$. The following theorem shows that such solutions converge to the solution v^* of (1.6).

Theorem 3.1. *Assume (2.3). Let v^* solve (1.6) and let v^n be the solution to (1.10) for $n \in \mathbb{N}$. If v^n are uniformly bounded in $\mathcal{C}^{2,\alpha}(\mathbb{R}^d)$ for all $n \in \mathbb{N}$, then $\lim_{n \rightarrow \infty} v^n = v^*$ as $n \rightarrow \infty$ locally uniformly in $\mathcal{C}^{2,\alpha}(\mathbb{R}^d)$.*

Proof. Since π^{n+1} is a maximizer for $F_\pi(x, Dv^n, D^2v^n)$ for each $n \geq 0$, it follows from (1.5) and the equation (1.10) that

$$\rho v^n - F(x, Dv^n, D^2v^n) = \rho v^n - F_{\pi^{n+1}}(x, Dv^n, D^2v^n) \leq 0.$$

Thus, the comparison principle yields

$$v^n \leq v^{n+1} \leq v^*.$$

We can then take $\bar{v} := \lim_{n \rightarrow \infty} v^n$. Since v^n are uniformly bounded in $\mathcal{C}^{2,\alpha}(\mathbb{R}^d)$ for all n , we get that $v^n \rightarrow \bar{v}$ as $n \rightarrow \infty$ locally uniformly in $\mathcal{C}^{2,\alpha}(\mathbb{R}^d)$, and $\bar{v} \in \mathcal{C}^{2,\alpha}(\mathbb{R}^d)$.

Note that, for each $u \in U$,

$$\pi^n(x, u) = \Gamma(x, Dv^n, D^2v^n)(u) \rightarrow \Gamma(x, D\bar{v}, D^2\bar{v})(u) \quad \text{locally uniformly.}$$

By the equations of v^n and the stability of viscosity solutions (under locally uniform convergence), we get that \bar{v} is a viscosity to

$$\begin{aligned} \rho v - \int_U [r(x, u) + b(x, u) \cdot Dv + 2^{-1} \operatorname{tr}(\sigma \sigma^T(x) D^2v) \\ - \lambda \ln(\Gamma(x, Dv, D^2v)(u))] \Gamma(x, Dv, D^2v)(u) du = 0. \end{aligned}$$

In view of the definition of Γ in (1.7), this shows that \bar{v} is a solution to (1.6). By the comparison principle again, we have $\bar{v} = v^*$, which finishes the proof. \square

3.2. Quantitative convergence results. In this subsection, we estimate the convergence rate of v^n to v^* , allowing some errors. Let $\pi^*(x, u) = \Gamma(x, Dv^*, D^2v^*)(u)$ be the unique optimal feedback control, and let

$$\begin{aligned} r^*(x) &= \int_U r(x, u) \pi^*(x, u) du, & b^*(x) &= \int_U b(x, u) \pi^*(x, u) du, \\ \Sigma^*(x) &= \int_U \sigma \sigma^T(x, u) \pi^*(x, u) du, & \mathcal{H}^*(x) &= \int_U \ln(\pi^*(x, u)) \pi^*(x, u) du, \end{aligned} \quad (3.1)$$

and then (1.6) is reduced to

$$\rho v^*(x) - r^*(x) - b^*(x) \cdot Dv^*(x) - 2^{-1} \operatorname{tr}(\Sigma^*(x) D^2v^*(x)) + \lambda \mathcal{H}^*(x) = 0. \quad (3.2)$$

Theorem 3.2. *Under the assumptions of Theorem 2.1 and also assuming that $\Sigma_0(\cdot)$ is Lipschitz continuous, then there exist $\rho_0, C > 0$ such that if $\rho \geq \rho_0$, we have for all $n \geq 1$ and $x_0 \in \mathbb{R}^n$,*

$$\rho \int_{B_1(x_0)} |v^n - v^*|^2 dx + \int_{B_1(x_0)} |D(v^n - v^*)|^2 dx \leq C(2^{-n} + \varepsilon_0^2/\rho).$$

As a consequence of this, we have for all $n \geq 1$,

$$\|D(v^n - v^*)\|_\infty \leq C(2^{-n} + \varepsilon_0^2/\rho)^{\frac{1}{d+2}}$$

and

$$\|v^n - v^*\|_\infty \leq C \rho^{-\frac{1}{d+2}} (2^{-n} + \varepsilon_0^2/\rho)^{\frac{1}{d+2} + \frac{d}{(d+2)^2}}.$$

Remark 3.1. *If σ is independent of u , then $\varepsilon_0 = \varepsilon_1 = 0$, so we obtain the exponential convergence rate for PIA as a corollary of the result. Also, we remark that the smallness of ε_1 is only used to guarantee that $v^n \in \mathcal{C}^{2,\alpha}(\mathbb{R}^d)$ uniformly for all $n \geq 1$.*

Proof. Without loss of generality, let us assume that $x_0 = 0$. The proof will be divided into three steps.

Step 1. Recall that, under the assumptions, v^n and v^* are uniformly bounded in $\mathcal{C}^{2,\alpha}(\mathbb{R}^d)$ for all $n \geq 1$. Denote $M(x, u) = \Sigma(x, u) - \Sigma_0(x)$ and then we have

$$|M(x, u)| \leq \varepsilon_0 \quad \text{for all } x, u.$$

As done before, if we write

$$\begin{aligned} g^n(x, u) &:= \frac{1}{\lambda} \left(r(x, u) + b(x, u) \cdot Dv^n + \frac{1}{2} \operatorname{tr}(M(x, u)D^2v^n) \right), \\ g^*(x, u) &:= \frac{1}{\lambda} \left(r(x, u) + b(x, u) \cdot Dv^* + \frac{1}{2} \operatorname{tr}(M(x, u)D^2v^*) \right), \end{aligned}$$

then

$$\pi^n(x, u) = \frac{\exp(g^{n-1}(x, u))}{\int_U \exp(g^{n-1}(x, u')) du'}, \quad \pi^*(x, u) = \frac{\exp(g^*(x, u))}{\int_U \exp(g^*(x, u')) du'}.$$

In view of (2.9) and (3.2), and setting $\bar{v}^n := v^* - v^n$, we obtain

$$\rho \bar{v}^n(x) - \frac{1}{2} \operatorname{tr}(\Sigma_0(x)D^2\bar{v}^n) = \lambda \int_U (g^*(x, u)\pi^*(x, u) - g^n(x, u)\pi^n(x, u)) du - \lambda(\mathcal{H}^*(x) - \mathcal{H}^n(x)). \quad (3.3)$$

It follows from the proof of (2.6) and the assumption of (2.11) that $|g^n|, |g^*| \leq C$ uniformly for all $n \geq 1$. By (1.9) and the uniform regularity of v^n, v^* , we get

$$|\pi^n(x, u) - \pi^*(x, u)| \leq C (|D\bar{v}^{n-1}| + \varepsilon_0 |D^2\bar{v}^{n-1}|).$$

Let us remark that ε_1 is not needed here. For simplicity of notation, we denote

$$\mathcal{E}_{n-1} := \mathcal{E}_{n-1}(x) = C (|D\bar{v}^{n-1}| + \varepsilon_0 |D^2\bar{v}^{n-1}|)(x)$$

with possibly different constant C in \mathcal{E}_n from one line to another. Recall (2.10) and (1.9). By the assumptions, it follows that

$$|b^* - b^n|, |r^* - r^n|, |M^* - M^n| \leq \mathcal{E}_{n-1}.$$

Thus, there is C and \mathcal{E}_{n-1} such that for all $u \in U$, $|\pi^*(x, u) - \pi^n(x, u)| \leq \mathcal{E}_{n-1}$. Then

$$\begin{aligned} &|g^*(x, u)\pi^*(x, u) - g^n(x, u)\pi^n(x, u)| \\ &\leq |g^*(x, u)| |\pi^*(x, u) - \pi^n(x, u)| + |\pi^n(x, u)| |g^*(x, u) - g^n(x, u)| \leq \mathcal{E}_{n-1} + \mathcal{E}_n. \end{aligned}$$

To estimate $|\mathcal{H}^n - \mathcal{H}^*|$, note that

$$|\ln \pi^n(x, u) - \ln \pi^*(x, u)| = \left| g^{n-1}(x, u) - g^*(x, u) - \ln \int_U e^{g^{n-1}(x, u') - g^*(x, u')} du' \right| \leq \mathcal{E}_{n-1},$$

and, since π^n is strictly positive by the uniform regularity of v^n ,

$$\left| \frac{\pi^*(x, u)}{\pi^n(x, u)} - 1 \right| \leq C |\pi^*(x, u) - \pi^n(x, u)| \leq \mathcal{E}_{n-1}.$$

Thus, we obtain

$$\begin{aligned} |\mathcal{H}^n(x) - \mathcal{H}^*(x)| &\leq \left| \int_U \pi^n(x, u) \ln \pi^n(x, u) du - \int_U \pi^*(x, u) \ln \pi^*(x, u) du \right| \\ &\leq \sup_{u \in U} |\ln \pi^n(x, u) - \ln \pi^*(x, u)| + \sup_{u \in U} \left| \frac{\pi^*(x, u)}{\pi^n(x, u)} - 1 \right| |\ln \pi^*(x, u)| \leq \mathcal{E}_{n-1}. \end{aligned}$$

Putting these into (3.3) yields

$$\rho \bar{v}^n(x) - 2^{-1} \operatorname{tr}(\Sigma_0(x)D^2\bar{v}^n) \leq \mathcal{E}_{n-1} + \mathcal{E}_n \quad (3.4)$$

for some C depending only on the assumptions.

Step 2. Next, let $\phi \in [0, 1]$ be a smooth function on \mathbb{R}^n such that for some $C > 0$,

$$\phi(\cdot) \equiv 1 \text{ on } B_1, \quad |D\phi(x)| \leq C\phi(x) \text{ for all } x \in \mathbb{R}^d, \quad \int_{\mathbb{R}^d} \phi(x) dx < \infty.$$

Such ϕ exists as one can take a smooth version of $\min\{1, e^{2-|x|}\}$.

From (3.4), multiply $\bar{v}^n \phi$ on both sides of (3.3) and integrate over \mathbb{R}^n to get

$$\rho \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx - \frac{1}{2} \int_{\mathbb{R}^d} \operatorname{tr}(\Sigma_0 D^2 \bar{v}^n) \bar{v}^n \phi \, dx \leq \int_{\mathbb{R}^d} (\mathcal{E}_{n-1} + \mathcal{E}_n) \bar{v}^n \phi \, dx. \quad (3.5)$$

Since $\Sigma_0 = \Sigma_0(x)$ is uniformly Lipschitz continuous and uniformly elliptic, direct computation yields

$$\begin{aligned} - \int_{\mathbb{R}^d} \operatorname{tr}(\Sigma_0 D^2 \bar{v}^n) \bar{v}^n \phi \, dx &= \sum_{1 \leq i, j \leq d} \int_{\mathbb{R}^d} \partial_{x_j} (\Sigma_0 \bar{v}^n \phi)_{ij} \partial_{x_i} \bar{v}^n \, dx \\ &= \sum_{1 \leq i, j \leq d} \int_{\mathbb{R}^d} \partial_{x_j} ((\Sigma_0)_{ij} \phi) \bar{v}^n \partial_{x_i} \bar{v}^n \, dx + \sum_{1 \leq i, j \leq d} \int_{\mathbb{R}^d} \partial_{x_j} \bar{v}^n (\Sigma_0)_{ij} \phi \partial_{x_i} \bar{v}^n \, dx \\ &\geq c \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx - C \int_{\mathbb{R}^d} |\bar{v}^n| |D \bar{v}^n| \phi \, dx \geq c \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx - C \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx. \end{aligned}$$

In the first inequality, we used that $|D\phi| \leq C\phi$ and uniform ellipticity; and we applied Young's inequality in the second inequality. The positive constants c, C depend only on ϕ and the C^1 norm of Σ_0 . With this, (3.5) implies that for some positive constants $C_1, c_1, C_2 > 0$,

$$\rho \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx + c_1 \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx \leq C_2 \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx + C_1 \int_{\mathbb{R}^d} (\mathcal{E}_{n-1} + \mathcal{E}_n) |\bar{v}^n| \phi \, dx.$$

By Young's inequality and that $|D^2 \bar{v}^{n-1}|, |D^2 \bar{v}^n| \leq C$, we obtain

$$\begin{aligned} (\rho - C_2) \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx + c_1 \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx \\ \leq C_1 \int_{\mathbb{R}^d} (|D \bar{v}^{n-1}| + |D \bar{v}^n|) |\bar{v}^n| \phi \, dx + CC_1 \varepsilon_0 \int_{\mathbb{R}^d} |\bar{v}^n| \phi \, dx \\ \leq \frac{c_1}{2} \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx + \frac{c_1}{4} \int_{\mathbb{R}^d} |D \bar{v}^{n-1}|^2 \phi \, dx + \frac{3C_1^2}{2c_1} \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx + C'_1 \varepsilon_0 \left(\int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx \right)^{1/2} \end{aligned}$$

with $C'_1 = CC_1$. Let us assume $\rho \geq 2C_2 + 3C_1^2/c_1$, and so

$$\rho \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx + c_1 \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx \leq \frac{c_1}{2} \int_{\mathbb{R}^d} |D \bar{v}^{n-1}|^2 \phi \, dx + 2C'_1 \varepsilon_0 \left(\int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx \right)^{1/2}. \quad (3.6)$$

If $\int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx \geq (\frac{2C'_1 \varepsilon_0}{\rho})^2$, (3.6) is reduced to

$$\int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx \leq \frac{1}{2} \int_{\mathbb{R}^d} |D \bar{v}^{n-1}|^2 \phi \, dx.$$

Otherwise, $\int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi \, dx \leq (\frac{2C'_1 \varepsilon_0}{\rho})^2$, so for both cases we have

$$\int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx \leq \frac{1}{2} \int_{\mathbb{R}^d} |D \bar{v}^{n-1}|^2 \phi \, dx + \frac{(2C'_1 \varepsilon_0)^2}{\rho c_1}.$$

It follows that

$$\begin{aligned} \int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx &\leq \frac{1}{2} \int_{\mathbb{R}^d} |D \bar{v}^{n-1}|^2 \phi \, dx + \frac{(2C'_1 \varepsilon_0)^2}{\rho c_1} \leq \frac{1}{4} \int_{\mathbb{R}^d} |D \bar{v}^{n-2}|^2 \phi \, dx + (1 + \frac{1}{2}) \frac{(2C'_1 \varepsilon_0)^2}{\rho c_1} \\ &\leq \dots \leq 2^{-n} \int_{\mathbb{R}^d} |D \bar{v}^0|^2 \phi \, dx + \frac{8(C'_1 \varepsilon_0)^2}{\rho c_1}. \end{aligned}$$

Since $\|v^0\|_2 < \infty$ and $\int_{\mathbb{R}^d} |\phi(x)| \, dx < \infty$, we proved that there exists $C_3 > 0$ such that for all $n \geq 1$,

$$\int_{\mathbb{R}^d} |D \bar{v}^n|^2 \phi \, dx \leq C_3 (2^{-n} + \varepsilon_0^2 / \rho). \quad (3.7)$$

Step 3. Finally, it follows from (3.6) that

$$\rho \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi dx \leq C_3 c_1 (2^{-n} + \varepsilon_0^2/\rho) + 2C'_1 \varepsilon_0 \left(\int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi dx \right)^{1/2}.$$

This shows that

$$\rho \int_{\mathbb{R}^d} |\bar{v}^n|^2 \phi dx \leq \max \{ 2C_3 c_1 (2^{-n} + \varepsilon_0^2/\rho), (4C'_1 \varepsilon_0)^2/\rho \},$$

which, combining with (3.7) and the fact that $\phi = 1$ in B_1 , finishes the proof of the first claim.

Because $\sup_n \|v^n - v^*\|_{2,\alpha} < \infty$, if $|D(v^n - v^*)|(0) = \delta > 0$ for some $\delta > 0$, then $|D(v^n - v^*)| \geq \frac{\delta}{2}$ in $B_{c\delta}$ for some $c > 0$. Therefore $\|D(v^n - v^*)\|_{L^2(B_1)}^2 \leq C(2^{-n} + \varepsilon_0^2/\rho)$ from the theorem yields

$$|D(v^n - v^*)|(0) \leq C(2^{-n} + \varepsilon_0^2/\rho)^{\frac{1}{d+2}} =: \varepsilon_2.$$

With this, if $|v^n - v^*|(0) = \delta' > 0$ for some $\delta' > 0$, then $|v^n - v^*| \geq \frac{\delta'}{2}$ in $B_{c\delta'/\varepsilon_2} \cap B_1$ for some $c > 0$. We obtain the following pointwise estimate:

$$|v^n(0) - v^*(0)| \leq C\rho^{-\frac{1}{d+2}} (2^{-n} + \varepsilon_0^2/\rho)^{\frac{1}{d+2} + \frac{d}{(d+2)^2}}.$$

After shifting the solutions, these finish the proof. □

4. UNBOUNDED NON-DEGENERATE ELLIPTIC EQUATIONS

In this section, we study non-degenerate elliptic equations, in which the coefficients might be unbounded in \mathbb{R}^d .

Recall that, by (1.8), $v^*(x) = \sup_{\pi \in \mathcal{P}(U)} V^\pi(x)$ is a viscosity solution to

$$\rho v - F(x, Dv, D^2v) = 0 = \rho v - F_{\pi^*}(x, Dv, D^2v) \quad \text{with } \pi^* = \Gamma(x, Dv^*, D^2v^*).$$

Plugging in the definition of Γ in (1.7) into the definition of F_π in (1.4) yields the equation

$$\rho v - \lambda \ln \int_U \exp \left[\frac{1}{\lambda} \left(r(x, u) + b(x, u) \cdot Dv + \frac{1}{2} \text{tr}(\Sigma(x, u) D^2v) \right) \right] du = 0. \quad (4.1)$$

In this section, we study (4.1) and only assume $\Sigma \geq 0$. If r, b and Σ are independent of u , the equation becomes a linear equation, which includes (1.10).

Recall $|U| = 1$ and $\Sigma(x, u) = \sigma(x, u)\sigma(x, u)^T$. We make the following assumptions:

$$r(x, u), b(x, u), \sigma(x, u) \text{ are locally uniformly Lipschitz continuous in } x, \quad (4.2)$$

and there exist $N > 0$ and $A_0, A_1 \geq 1$ such that for all $(x, u) \in \mathbb{R}^d \times U$,

$$|r(x, u)| \leq A_1(1 + |x|^N), \quad |b(x, u)| \leq A_2(1 + |x|), \quad |\sigma(x, u)| \leq A_3(1 + |x|), \quad (4.3)$$

where, for $X = (x_{ij})$ a vector or a matrix, $|X| := \max_{i,j} |x_{ij}|$.

4.1. Existence and uniqueness. We start with a comparison principle in bounded domains. Throughout this section, we allow degenerate diffusion, that is, we only require $\Sigma \geq 0$.

Lemma 4.1. *Let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set, and assume (4.2). Let μ (resp. v) be a bounded subsolution (resp. supersolution) to*

$$\rho\mu - F(x, D\mu, D^2\mu) = 0 \quad \text{in } \Omega, \quad (4.4)$$

such that

$$\sup_{x \in \partial\Omega} (\mu(x) - v(x)) \leq 0.$$

Then $\mu \leq v$ in Ω .

Proof. In view of [7, Theorem 3.3], it suffices to check that the operator $\rho\mu - F(x, p, X)$ is proper and satisfies conditions (3.13) and (3.14) in [7]. By the definition of the operator, we only need to verify (3.14): there is a function $\omega : [0, \infty) \rightarrow [0, \infty)$ such that $\omega(0+) = 0$ and

$$F(x, \alpha(x - y), X_\alpha) - F(y, \alpha(x - y), Y_\alpha) \leq \omega(\alpha|x - y|^2 + |x - y|) \quad (4.5)$$

whenever $x, y \in \Omega$, $r \in \mathbb{R}$, and $X_\alpha, Y_\alpha \in \mathcal{S}^d$ are such that

$$-3\alpha \begin{pmatrix} \mathbb{I}_d & 0 \\ 0 & \mathbb{I}_d \end{pmatrix} \leq \begin{pmatrix} X_\alpha & 0 \\ 0 & -Y_\alpha \end{pmatrix} \leq 3\alpha \begin{pmatrix} \mathbb{I}_d & -\mathbb{I}_d \\ -\mathbb{I}_d & \mathbb{I}_d \end{pmatrix}. \quad (4.6)$$

Indeed from [7, Example 3.6], (4.6) and the uniform Lipschitz continuity of $\sigma(\cdot, u)$ yield for any $u \in U$ and $x, y \in \Omega$,

$$\text{tr}(\sigma(x, u)\sigma^T(x, u)X_\alpha) - \text{tr}(\sigma(y, u)\sigma^T(y, u)Y_\alpha) \leq C\alpha|x - y|^2$$

where $C > 0$ only depends on (4.2), A_1 and R , where R is such that $\Omega \subseteq B_R$. Moreover, using (4.2) yields, for

$$f_u(x, p, X) := r(x, u) + b(x, u) \cdot p + \frac{1}{2} \text{tr}(\Sigma(x, u)X), \quad (4.7)$$

that

$$f_u(x, p, X_\alpha) - f_u(y, p, Y_\alpha) \leq C(|x - y| + |x - y||p| + \alpha|x - y|^2)$$

for some $C > 0$, and so

$$\begin{aligned} & \lambda \ln \int_U \exp \left[\frac{1}{\lambda} f_u(x, \alpha(x - y), X_\alpha) \right] du - \lambda \ln \int_U \exp \left[\frac{1}{\lambda} f_u(y, \alpha(x - y), Y_\alpha) \right] du \\ & \leq \lambda \ln \int_U \exp \left(\frac{1}{\lambda} C(|x - y| + \alpha|x - y|^2) \right) du = C(|x - y| + \alpha|x - y|^2). \end{aligned}$$

This and the equation (4.1) show (4.5) with $\omega(z) := Cz$. \square

Lemma 4.2 (Comparison principle in \mathbb{R}^d). *Assume (4.2), (4.3) and that*

$$\rho \geq 4(N + 1)(A_2 + NA_3). \quad (4.8)$$

Let μ and v be, respectively, a subsolution and a supersolution to (4.1) in \mathbb{R}^d such that

$$\limsup_{|x| \rightarrow \infty} \frac{\mu(x) - v(x)}{|x|^{N+1}} \leq 0. \quad (4.9)$$

Then $\mu \leq v$ in \mathbb{R}^d .

Proof. For any $\varepsilon > 0$, define

$$\mu_\varepsilon(x) := \mu(x) - \varepsilon(1 + |x|^{N+1}).$$

We claim that μ_ε is a (viscosity) subsolution to (4.1) in \mathbb{R}^d . Indeed, if $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ is such that $\mu_\varepsilon - \varphi$ has a local maximum at $x_0 \in \mathbb{R}^d$. Then $\mu - \varphi_\varepsilon$ with $\varphi_\varepsilon := \varphi + \varepsilon(1 + |x|^{N+1})$ has a local maximum at x_0 . Using the notation from (4.7) and the assumptions (4.2)–(4.3) implies that

$$\begin{aligned} & f_u(x_0, D\varphi_\varepsilon(x_0), D^2\varphi_\varepsilon(x_0)) - f_u(x_0, D\varphi(x_0), D^2\varphi(x_0)) \\ & \leq \varepsilon(N + 1)|b(x_0, u)||x_0|^N + \varepsilon(N + 1)N|\Sigma(x_0, u)||x_0|^{N-1} \\ & \leq \varepsilon(N + 1)A_2(1 + |x_0|)|x_0|^N + \varepsilon(N + 1)NA_3(1 + |x_0|)^2|x_0|^{N-1}. \end{aligned}$$

Recall that $F(x, p, X) = \lambda \ln \int_U \exp[\frac{1}{\lambda} f_u(x, p, X)] du$. Since μ is a subsolution to (4.1),

$$\rho\mu(x_0) - F(x_0, D\varphi_\varepsilon(x_0), D^2\varphi_\varepsilon(x_0)) \leq 0.$$

We obtain at $x = x_0$,

$$\begin{aligned} \rho\mu_\varepsilon - F(x_0, D\varphi, D^2\varphi) &\leq \rho(\mu - \varepsilon(1 + |x_0|^{N+1})) - F(x_0, D\varphi_\varepsilon, D^2\varphi_\varepsilon) \\ &\quad + \varepsilon(N+1)A_2(1 + |x_0|)|x_0|^N + \varepsilon(N+1)NA_3(1 + |x_0|)^2|x_0|^{N-1} \\ &\leq -\varepsilon\rho(1 + |x_0|^{N+1}) + \varepsilon(N+1)(A_2 + NA_3)(1 + |x_0|)^2|x_0|^{N-1}. \end{aligned}$$

Hence by (4.8), we get from the above that

$$\rho\mu_\varepsilon - F(x_0, D\varphi, D^2\varphi) \leq 0.$$

Therefore, for all $\varepsilon > 0$, μ_ε is a subsolution to (4.1).

Now by (4.9), there exists $R_\varepsilon > 0$ such that $\lim_{\varepsilon \rightarrow 0} R_\varepsilon = \infty$ and $\mu_\varepsilon(x) \leq v(x)$ for all $|x| \geq R_\varepsilon$. Therefore applying Lemma 4.1 to v, μ_ε with $\Omega = B_{R_\varepsilon}$ yields

$$\mu_\varepsilon(x) \leq v(x) \quad \text{for all } x \in B_{R_\varepsilon}.$$

Taking $\varepsilon \rightarrow 0$ leads to $\mu \leq v$ in \mathbb{R}^d . \square

Proposition 4.1. *Under the assumption of Lemma 4.2, there exists a unique (viscosity) solution v to (4.1) such that for all $x \in \mathbb{R}^d$,*

$$|v(x)| \leq 2A_1\rho^{-1}(1 + |x|^2)^{N/2}. \quad (4.10)$$

Proof. To prove the existence and uniqueness of solutions, in view of the comparison principle, it suffices to produce a supersolution and a subsolution with a polynomial growth at infinity and invoke Perron's method.

By (4.1) and (4.3), for any $(x, p, X) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathcal{S}^d$, we have

$$F(x, p, X) \leq A_1(1 + |x|^2)^{N/2} + A_2(1 + |x|^2)^{1/2}|p| + A_3(1 + |x|^2)|X|. \quad (4.11)$$

Set $\phi(x) := (1 + |x|^2)^{N/2}$, and then define $\bar{\mu}(x) := A_0\phi(x)$ with $A_0 := 2A_1/\rho$. For simplicity, below we drop (x) from the notations of $\bar{\mu}(x)$, $\phi(x)$. We have from (4.11) and direct computations that

$$\begin{aligned} \rho\bar{\mu} - F(x, D\bar{\mu}, D^2\bar{\mu}) &\geq \rho A_0(1 + |x|^2)^{N/2} - A_1(1 + |x|^2)^{N/2} - A_0N(A_2 + A_3(N-1))(1 + |x|^2)^{N/2} \\ &\geq (\rho A_0/2 - A_1)(1 + |x|^2)^{N/2} \geq 0, \end{aligned}$$

thanks to (4.8). Thus, $\bar{\mu}$ is a supersolution.

Similarly, it can be shown that $\underline{\mu} := -\bar{\mu}$ is a subsolution. It is clear that

$$\lim_{|x| \rightarrow \infty} \frac{|\bar{\mu}(x)| + |\underline{\mu}(x)|}{|x|^{N+1}} = 0.$$

Thus by Perron's method and Lemma 4.2, we obtain the unique solution v to (4.1) and it satisfies $\underline{\mu} \leq v \leq \bar{\mu}$, which yields (4.10). \square

Remark 4.1. *The following comments are in order.*

1. *If we do not assume ρ to be sufficiently large, then the uniqueness of solutions might fail. For example (when $d = 1$) both $v \equiv 0$ and $v = x$ are solutions to $v - xv_x = 0$, and both $v \equiv 0$ and $v = x^2 + 1$ are solutions to $v - \frac{1}{2}(1 + x^2)v_{xx} = 0$.*
2. *For any fixed large $\rho = N(N-1)$ with $N \geq 2$ an integer, the uniqueness of solutions still fails in general. To see this, we first construct the following $N+1$ numbers: $a_N = 1$, $a_{N-1} = \frac{N}{2(N-1)}$ and define iteratively for $k = N-2, \dots, 0$,*

$$a_k = \frac{(k+1)a_{k+1} + (k+2)(k+1)a_{k+2}}{N(N-1) - k(k-1)}.$$

Then one can check directly that $v = \sum_{k=0}^N a_k x^k$ and $v \equiv 0$ are both solutions to $\rho v - v_x - \frac{1}{2}(1 + x^2)v_{xx} = 0$. This does not contradict Proposition 4.1 because our result claims

the unique solution among functions that grow as fast as a polynomial of power $c\sqrt{\rho}$ for some possibly small $c > 0$ (depending on the assumption (4.3)). However, in the example, $\sqrt{\rho} = \sqrt{N(N-1)}$.

Also, we cannot allow exponential growth of solutions (otherwise uniqueness fails). For instance, both 0 and e^x are solutions to $v - v_{xx} = 0$. These examples show the optimality of our assumptions on ρ in terms of N in Lemma 4.2 and Proposition 4.1.

3. If $b(\cdot, u)$ and $\sigma(\cdot, u)$ are only allowed to have a sublinear growth in x , that is

$$\lim_{R \rightarrow \infty} \sup_{(x,u) \in B_R \times U} \frac{|b(x, u)| + |\sigma(x, u)|}{R} = 0,$$

then the existence and uniqueness of solutions, and the comparison principle hold the same without having to assume ρ to be sufficiently large. The proof is similar to the one presented in the paper. We also refer readers to [28, Theorem 8].

5. CONVERGENCE OF PIA WITH UNBOUNDED COEFFICIENTS

This section concerns the case when σ is independent of the control, and the coefficients can be unbounded.

Let us start with the following interior $W^{2,p}$ estimate. The classical result can be found, for example, in [6, Chapter 3, Theorem 4.2]). However, it is not sufficient for us as we need to carefully track the dependence of the constants on the size of the coefficients of the equation.

Let \tilde{v} be a solution to

$$\rho \tilde{v}(x) - \tilde{r}(x) - \tilde{b}(x) \cdot D\tilde{v}(x) - 2^{-1} \operatorname{tr} \left(\tilde{\Sigma}(x) D^2 \tilde{v}(x) \right) = 0. \quad (5.1)$$

Lemma 5.1. *Assume that $\rho \geq 1$, and for some $\tilde{C}_0 > 0$, $\tilde{\Sigma} = \tilde{\sigma} \tilde{\sigma}^T \geq \mathbb{I}_d / \tilde{C}_0$ and $\tilde{\sigma}$ is Lipschitz continuous with constant \tilde{C}_0 . Consider the domain $B_2(x_0)$ for an arbitrary $x_0 \in \mathbb{R}^d$. Suppose there exist $\tilde{A}_1, \tilde{A}_2, \tilde{A}_3 \geq 1$ such that*

$$\|\tilde{r}(\cdot)\|_{L^\infty(B_2(x_0))} \leq \tilde{A}_1, \quad \|\tilde{b}(\cdot)\|_{L^\infty(B_2(x_0))} \leq \tilde{A}_2, \quad \|\tilde{\sigma}(\cdot)\|_{L^\infty(B_2(x_0))} \leq \tilde{A}_3. \quad (5.2)$$

Then for any $p > d$, there exists $C = C(\tilde{C}_0, p, d) > 0$ but independent of $\tilde{A}_1, \tilde{A}_2, \tilde{A}_3, \rho$ and x_0 such that the solution \tilde{v} to (5.1) satisfies

$$\|D^2 \tilde{v}\|_{W^{1,p}(B_1(x_0))} \leq C \left[\tilde{A}_1 + (\tilde{A}_2^2 + \tilde{A}_3^{4+\frac{2d}{p}} + \rho) \|\tilde{v}\|_{L^\infty(B_2(x_0))} \right]. \quad (5.3)$$

Proof. Without loss of generality, we only prove the result for $x_0 = 0$. Below, we divide the proof into three steps.

Step 1. We first derive a Sobolev inequality over B_R with explicit dependence of constants on $R \in (0, \frac{1}{2})$. Since $p > d$, we can apply Gagliardo-Nirenberg interpolation inequality to $\mu(x) := \tilde{v}(Rx)$ in B_1 to get

$$\|D\mu\|_{L^p(B_1)} \leq C \|D^2 \mu\|_{L^p(B_1)}^\theta \|\mu\|_{L^\infty(B_1)}^{1-\theta} + C \|\mu\|_{L^\infty(B_1)},$$

where the parameters satisfy

$$\frac{1}{p} = \frac{1}{d} + \theta \left(\frac{1}{p} - \frac{2}{d} \right),$$

and C only depends p and d . By Young's inequality, we have for any $\varepsilon \in (0, 1]$ that

$$\|D\mu\|_{L^p(B_1)} \leq \varepsilon \|D^2 \mu\|_{L^p(B_1)} + C \varepsilon^{-\frac{\theta}{1-\theta}} \|\mu\|_{L^\infty(B_1)},$$

which implies, for C only depending on p, d and independent of R, ε that

$$\|D\tilde{v}\|_{L^p(B_R)} \leq \varepsilon R \|D^2 \tilde{v}\|_{L^p(B_R)} + C \varepsilon^{-1+\frac{d}{p}} R^{-1+\frac{d}{p}} \|\tilde{v}\|_{L^\infty(B_R)}. \quad (5.4)$$

Step 2. In this step, we derive a uniform Sobolev estimate for a transformed function $\tilde{w}(x)$ defined below on small balls.

First, since $\tilde{\Sigma} = \tilde{\sigma}\tilde{\sigma}^T$ and (5.2), there exists $c \in (0, 1)$ depending on d such that

$$\left\{ \tilde{\Sigma}(x_1)^{1/2}x + x_1 \mid x \in B_{c/\tilde{A}_3} \right\} \subset B_2 \quad \forall x_1 \in B_1. \quad (5.5)$$

Set $\tilde{R} := c'(\tilde{A}_2 + \tilde{A}_3^2)^{-1}$ for some $c' \in (0, c)$ to be determined in the proof. Then, by the assumption that $\tilde{\sigma}$ is Lipschitz continuous, there exists C' depending only on \tilde{C}_0 such that for any $0 < R < \tilde{R}$,

$$\left| \tilde{\Sigma} \left(\tilde{\Sigma}(x_1)^{1/2}x + x_1 \right) - \tilde{\Sigma}(x_1) \right| \leq C' \tilde{A}_3^2 |x| \leq C' \tilde{A}_3^2 R. \quad (5.6)$$

From now on, fix $x_1 \in B_1$ and define $\tilde{w}(x) := v(\tilde{\Sigma}(x_1)^{1/2}x + x_1)$. (5.5) yields that \tilde{w} is well-defined in a neighbourhood of $B_{\tilde{R}}$. A direct calculation shows that \tilde{w} satisfies $-\frac{1}{2}\Delta\tilde{w}(x) = g(x)$ with

$$\begin{aligned} g(x) := & \tilde{r} \left(\tilde{\Sigma}(x_1)^{1/2}x + x_1 \right) + \tilde{b} \left(\tilde{\Sigma}(x_1)^{1/2}x + x_1 \right) \cdot \tilde{\Sigma}(x_1)^{-1/2} D\tilde{w}(x) \\ & + 2^{-1} \operatorname{tr} \left(\left(\tilde{\Sigma}(\tilde{\Sigma}(x_1)^{1/2}x + x_1)\tilde{\Sigma}(x_1)^{-1} - \mathbb{I}_d \right) D^2\tilde{w}(x) \right) - \rho\tilde{w}(x). \end{aligned} \quad (5.7)$$

Next, for any $R \in (0, \tilde{R}]$ and $\eta \in (0, R)$, take $\zeta \in C_0^\infty(B_R)$ a cutoff function such that

$$\zeta \leq 1 \text{ in } B_R, \quad \zeta = 1 \text{ in } B_\eta \quad \text{and} \quad |D^k\zeta| \leq C(R - \eta)^{-k} \text{ with } k = 1, 2.$$

Here the constant C can be taken independent of R and η . Then $\hat{w}(x) := \tilde{w}(x)\zeta(x)$ satisfies

$$2^{-1}\Delta\hat{w}(x) = g(x)\zeta(x) - D\tilde{w}(x) \cdot D\zeta(x) - 2^{-1}\tilde{w}(x)\Delta\zeta(x) =: \hat{g}(x).$$

By classical results for elliptic equations (e.g., [6, Chapter 3, Theorem 3.6]), we have that

$$\|D^2\hat{w}\|_{L^p(B_R)} \leq C\|\hat{g}\|_{L^p(B_R)},$$

where C depends only on p, d . Then by the definition of ζ , the above estimate implies that

$$\|D^2\tilde{w}\|_{L^p(B_\eta)} \leq C\|g\|_{L^p(B_R)} + C(R - \eta)^{-1}\|D\tilde{w}\|_{L^p(B_R)} + C(R - \eta)^{-2}\|\tilde{w}\|_{L^\infty(B_R)}, \quad (5.8)$$

with C independent of $0 < \eta < R \leq \tilde{R}$. Then (5.6)–(5.8) together yield that

$$\begin{aligned} \|D^2\tilde{w}\|_{L^p(B_\eta)} & \leq C \left(\|\tilde{r}\|_{L^\infty(B_1)}|B_R|^{1/p} + \|\tilde{b}\|_{L^\infty(B_1)}\|D\tilde{w}\|_{L^p(B_R)} + \tilde{A}_3^2 R \|D^2\tilde{w}\|_{L^p(B_R)} + \rho\|\tilde{w}\|_{L^p(B_R)} \right) \\ & \quad + C(R - \eta)^{-1}\|D\tilde{w}\|_{L^p(B_R)} + C(R - \eta)^{-2}\|\tilde{w}\|_{L^\infty(B_R)} \\ & \leq C \left(\tilde{A}_1 R^{d/p} + \tilde{A}_2 \|D\tilde{w}\|_{L^p(B_R)} + \tilde{A}_3^2 R \|D^2\tilde{w}\|_{L^p(B_R)} + \rho R^{d/p} \|\tilde{w}\|_{L^\infty(B_R)} \right) \\ & \quad + C(R - \eta)^{-1}\|D\tilde{w}\|_{L^p(B_R)} + C(R - \eta)^{-2}\|\tilde{w}\|_{L^\infty(B_R)}. \end{aligned} \quad (5.9)$$

where C only depends on \tilde{C}_0, p and d . In particular, it is independent of $0 < \eta < R \leq \tilde{R}$ and $x_1 \in B_1$.

Now, it follows from (5.4) that there exists C independent of $R \in (0, \tilde{R})$ and $\varepsilon \in (0, 1]$ such that

$$\|D\tilde{w}\|_{L^p(B_R)} \leq \varepsilon R \|D^2\tilde{w}\|_{L^p(B_R)} + C\varepsilon^{-1+\frac{d}{p}}R^{-1+\frac{d}{p}}\|\tilde{w}\|_{L^\infty(B_R)}.$$

Since $R \leq \tilde{R} \leq (\tilde{A}_2 + \tilde{A}_3^2)^{-1}$, applying the above estimate with $\varepsilon = 1$ in the last but one line of (5.9) and with $\varepsilon = (\tilde{A}_2 + \tilde{A}_3^2)(R - \eta) \leq 1$ in the last line of (5.9) yields

$$\begin{aligned} \|D^2\tilde{w}\|_{L^p(B_\eta)} & \leq C_1(\tilde{A}_2 + \tilde{A}_3^2)R \|D^2\tilde{w}\|_{L^p(B_R)} + C_1 \left(\tilde{A}_1 R^{\frac{d}{p}} + (\tilde{A}_2 R^{-1+\frac{d}{p}} + \rho R^{\frac{d}{p}}) \|\tilde{w}\|_{L^\infty(B_R)} \right) \\ & \quad + C_1(1 + ((\tilde{A}_2 + \tilde{A}_3^2)R)^{-1+\frac{d}{p}})(R - \eta)^{-2}\|\tilde{w}\|_{L^\infty(B_R)}, \end{aligned}$$

where $C_1 \geq 1$ depends only on \tilde{C}_0, p and d .

Let us now take $c' = \min\{c, 1/(2C_1)\}$, $\tilde{R} = c'(\tilde{A}_2 + \tilde{A}_3^2)^{-1}$ and $R_0 := \tilde{R}/2$. Thus, for all $R_0 \leq \eta < R \leq \tilde{R}$, the above inequality yields for some $C'_1 > 0$ depending only on C_1, c', d, p ,

$$\begin{aligned} \|D^2\tilde{w}\|_{L^p(B_\eta)} &\leq \frac{1}{2}\|D^2\tilde{w}\|_{L^p(B_R)} + C_1 \left(\tilde{A}_1\tilde{R}^{\frac{d}{p}} + (\tilde{A}_2\tilde{R}^{-1+\frac{d}{p}} + \rho\tilde{R}^{\frac{d}{p}})\|\tilde{w}\|_{L^\infty(B_{\tilde{R}})} \right) \\ &\quad + C'_1(R - \eta)^{-2}\|\tilde{w}\|_{L^\infty(B_{\tilde{R}})}. \end{aligned}$$

We use [6, Chapter 2, Lemma 4.1] to get for all $R_0 \leq \eta < R \leq \tilde{R}$ that

$$\|D^2\tilde{w}\|_{L^p(B_\eta)} \leq C \left(\tilde{A}_1R_0^{\frac{d}{p}} + (\tilde{A}_2R_0^{-1+\frac{d}{p}} + \rho R_0^{\frac{d}{p}})\|\tilde{w}\|_{L^\infty(B_{2R_0})} \right) + C(R - \eta)^{-2}\|\tilde{w}\|_{L^\infty(B_{2R_0})},$$

which yields

$$\|D^2\tilde{w}\|_{L^p(B_{R_0})} \leq C \left(\tilde{A}_1R_0^{\frac{d}{p}} + (\tilde{A}_2R_0^{-1+\frac{d}{p}} + \rho R_0^{\frac{d}{p}} + R_0^{-2})\|\tilde{w}\|_{L^\infty(B_{2R_0})} \right) \quad (5.10)$$

where C only depends on \tilde{C}_0, p, d . For readers' convenience, we copy [6, Chapter 2, Lemma 4.1] after the proof.

Step 3. Let us turn back to v . Notice that there exists $\tilde{C} \geq 1$ only depending on C_0 such that $\tilde{\Sigma}(x_1)^{-1/2}B_R := \{\tilde{\Sigma}(x_1)^{-1/2}x \mid x \in B_R\} \subset B_{\tilde{C}R}$ for all $x_1 \in B_1$. Set $r_0 = R_0/\tilde{C}$, and then (5.10) and $\tilde{w}(x) = v(\tilde{\Sigma}(x_1)^{1/2}x + x_1)$ together give that

$$\begin{aligned} \|D^2v\|_{L^p(B_{r_0}(x_1))} &= \left(\int_{\tilde{\Sigma}(x_1)^{-1/2}B_{r_0}} |D^2\tilde{w}(z)|^p \tilde{\Sigma}(x_1)^{-1/2} dz \right)^{1/p} \leq C\|D^2\tilde{w}\|_{L^p(B_{R_0})} \\ &\leq C \left(\tilde{A}_1r_0^{\frac{d}{p}} + (\tilde{A}_2r_0^{-1+\frac{d}{p}} + \rho r_0^{\frac{d}{p}} + r_0^{-2})\|v\|_{L^\infty(B_2)} \right), \end{aligned} \quad (5.11)$$

where we also used (5.5) in the last inequality, and the constant C only depends on \tilde{C}_0, p, d .

Next, we can take $N := (\lfloor \sqrt{d}/r_0 \rfloor + 1)^d$ balls centered at $\{y_1, y_2, \dots, y_N\} \subseteq B_1$ such that $B_1 \subseteq \bigcup_{i=1}^N B_{r_0}(y_i)$. By applying (5.11) to $x_1 = y_1, \dots, y_N$, we have that

$$\begin{aligned} \|D^2v\|_{L^p(B_1)} &\leq \left(\sum_{i=1}^N \|D^2v\|_{L^p(B_{r_0}(y_i))}^p \right)^{1/p} \leq CN^{1/p} \left[\tilde{A}_1r_0^{\frac{d}{p}} + (\tilde{A}_2r_0^{-1+\frac{d}{p}} + \rho r_0^{\frac{d}{p}} + r_0^{-2})\|v\|_{L^\infty(B_2)} \right] \\ &\leq C \left[\tilde{A}_1 + (\tilde{A}_2r_0^{-1} + \rho + r_0^{-2-\frac{d}{p}})\|v\|_{L^\infty(B_2)} \right]. \end{aligned}$$

Recall that $r_0 = c'/\tilde{C}(\tilde{A}_2 + \tilde{A}_3^2)^{-1}$. By (5.4) again, we obtain

$$\|D^2v\|_{W^{1,p}(B_1)} \leq C \left[\tilde{A}_1 + (\tilde{A}_2^2 + \tilde{A}_2\tilde{A}_3^2 + \tilde{A}_3^{4+\frac{2d}{p}} + \rho)\|v\|_{L^\infty(B_2)} \right]$$

where C is independent of $\tilde{A}_1, \tilde{A}_2, \tilde{A}_3$ and ρ . This finishes the proof by Young's inequality. \square

Let us state Lemma 4.1 from [6, Chapter 2] that was used in the proof of Lemma 5.1.

Lemma 5.2. *Let $\varphi(r)$ be a bounded nonnegative function defined on the interval $[R_0, R_1]$, where $R_1 > R_0 \geq 0$. Suppose that for any $R_0 \leq \eta < R \leq R_1$, φ satisfies*

$$\varphi(\eta) \leq \theta\varphi(R) + \frac{A}{(R - \eta)^\alpha} + B$$

where θ, A, B and α are nonnegative constants, $\theta < 1$. Then

$$\varphi(\eta) \leq C \left[\frac{A}{(R - \eta)^\alpha} + B \right], \quad \text{for all } R_0 \leq \eta < R \leq R_1,$$

where C depends only on α, θ .

Now we study PIA. We make the following assumptions:

(H1) Assume $\Sigma(\cdot, u) \geq \mathbb{I}_d/C_0$, (4.2), (4.3), and (4.8). Suppose that v^0 is locally uniformly $\mathcal{C}^{1,\alpha}$ for some $\alpha \in (0, 1)$ and

$$\sup_{R \geq 1} R^{-N} \|v^0\|_{\mathcal{C}^{1,\alpha}(B_R)} < \infty.$$

(H2) $U = [0, 1]^L$ for some $L \geq 1$.

(H3) The maps $u \rightarrow r(x, u)$ and $u \rightarrow b(x, u)$ are uniformly Lipschitz continuous. And $\sigma = \sigma(x)$ is independent of u .

We comment that (H2) can be generalized to a cone test condition [16, Assumption 4.2].

Since σ is independent of u , we have a simpler formula for π^n . Indeed, for $n \geq 1$, define

$$\pi^n(x)(u) := \Gamma(x, Dv^{n-1}(x))(u) = \frac{\exp\left(\frac{1}{\lambda}(r(x, u) + b(x, u) \cdot Dv^{n-1}(x))\right)}{\int_U \exp\left(\frac{1}{\lambda}(r(x, u') + b(x, u') \cdot Dv^{n-1}(x))\right) du'}, \quad (5.12)$$

and r^n, b^n, \mathcal{H}^n the same as in (2.10) with the above π^n . We then look for v^n from the equation:

$$\rho v^n(x) - r^n(x) - b^n(x) \cdot Dv^n(x) - 2^{-1} \operatorname{tr}(\Sigma(x) D^2 v^n(x)) + \lambda \mathcal{H}^n(x) = 0. \quad (5.13)$$

Below we show the well-posedness of (5.13) and thus the PIA, and we use the uniform ellipticity of the equation to obtain some uniform estimates on v^n for all $n \geq 0$.

Lemma 5.3. *Under the assumptions of (H1)-(H3), there exists $C > 0$ independent of ρ such that for all $n \geq 1$ and any $R > 0$,*

$$\rho \|v^n\|_{L^\infty(B_R)} \leq C(1 + R^N) \quad \text{and} \quad \|Dv^n\|_{\mathcal{C}^\alpha(B_R)} \leq C(1 + |R|^{N+5}).$$

Proof. Let $M \geq 2$ be a large constant, so that it satisfies for some $\tilde{C} \geq 1$ to be determined (independent of n, ρ),

$$\sup_{R \geq 1} R^{-N} \|v^0\|_{\mathcal{C}^{1,\alpha}(B_R)} \leq M \quad \text{and} \quad M \geq \tilde{C}(1 + \ln M).$$

Assume for induction that for some $n \geq 1$, v^{n-1} exists, and for all $x_0 \in \mathbb{R}^d$ we have

$$\|Dv^{n-1}\|_{\mathcal{C}^\alpha(B_1(x_0))} \leq M(1 + |x_0|^{N+5}). \quad (5.14)$$

Recall (2.10) and (5.12). We get from (4.3) that

$$|r^n(x)| \leq \sup_{u \in U} |r(x, u)| \leq A_1(1 + |x|^N) \quad \text{and} \quad |b^n(x)| \leq \sup_{u \in U} |b(x, u)| \leq A_2(1 + |x|). \quad (5.15)$$

Since $U = [0, 1]^L$, the cone test condition in [16, Corollary 4.2] is satisfied. Thus, the corollary yields that there exists C depending on L, λ, N and other constants in the assumptions such that

$$\begin{aligned} |\lambda \mathcal{H}^n(x)| &= \left| \lambda \int_U \ln(\Gamma(x, Dv^{n-1}(x))) \Gamma(x, Dv^{n-1}(x)) du \right| \\ &\leq C(1 + \ln(1 + |Dv^{n-1}(x)|)) \leq C(\ln M + \ln(1 + |x|)) \end{aligned}$$

where, in the second inequality, we applied the induction hypothesis (5.14). Thus,

$$|r^n(x) - \lambda \mathcal{H}^n(x)| \leq (A_1 + 2C \ln M)(1 + |x|^N). \quad (5.16)$$

Then it follows from (4.8) and Proposition 4.1 (with $r^n(x) - \lambda \mathcal{H}^n(x), b^n(x)$ in place of $r(x, u), b(x, u)$) that there exists a unique solution v^n to (5.13) such that

$$\rho |v^n(x)| \leq C_1(A_1 + \ln M) (1 + |x|^N) \quad \forall x \in \mathbb{R}^d, \quad (5.17)$$

where C_1 depends only on L, λ, N and the assumptions. By applying Lemma 5.1 to v^n with an arbitrary $p > d$, we have for any point $x_0 \in \mathbb{R}^d$ that,

$$\|v^n\|_{W^{2,p}(B_2(x_0))} \leq C \left[\|r^n - \lambda \mathcal{H}^n\|_{L^\infty(B_4(x_0))} + \left(\rho + \|b^n\|_{L^\infty(B_4(x_0))}^2 + \|\sigma\|_{L^\infty(B_4(x_0))}^{4+\frac{2d}{p}} \right) \|v^n\|_{L^\infty(B_4(x_0))} \right],$$

Recall $\rho \geq 4(N+1)(A_2 + NA_3)$ from (4.8). We apply (4.3), (5.15), (5.16) and (5.17) to get

$$\|v^n\|_{W^{2,p}(B_1(x_0))} \leq C(A_1 + \ln M) \left(A_2 + A_3^{3+\frac{2d}{p}} \right) \left(1 + |x_0|^{N+4+\frac{2d}{p}} \right) \quad (5.18)$$

with C depending only on C_0, p, d (independent of $\rho \geq 1$).

Now, taking $p = \max\{\frac{d}{1-\alpha}, 2d\}$ and using (5.18), we apply Sobolev embedding (e.g., [11, Section 5.6.2, Theorem 5]) to get

$$\|v^n\|_{C^{1,\alpha}(B_1(x_0))} \leq C \|v^n\|_{W^{2,p}(B_2(x_0))} \leq C(A_1 + \ln M) (A_2 + A_3^4) (1 + |x_0|^{N+5}), \quad (5.19)$$

where C is independent of x_0, ρ and $n \geq 1$. Hence if M is sufficiently large such that

$$C(A_1 + \ln M) (A_2 + A_3^4) \leq M, \quad (5.20)$$

then

$$\|Dv^n\|_{C^\alpha(B_1(x_0))} \leq M(1 + |x_0|^{N+5}).$$

By induction, we finished the proof of (5.14).

Finally, (5.17) finishes the proof of the lemma. \square

Corollary 5.1. *Under the assumptions of Lemma 5.3, then we have*

$$v^{n-1} \leq v^n, \quad \forall n \geq 1.$$

Proof. By Proposition 4.1 and Lemma 5.3

$$\rho \|v^n\|_{L^\infty(B_R)} \leq C(1 + R^N) \quad \forall R > 0,$$

for some constant C independent of n and ρ , and v^n is a subsolution for (4.1). Notice that v^{n-1} being a subsolution to (5.13) for any $n \geq 1$. Thus, with $r^n(x) - \lambda \mathcal{H}^n(x), b^n(x)$ in place of $r(x, u), b(x, u)$ in (4.4), we apply Lemma 4.2 to (5.13) to get that $v^n \geq v^{n-1}$ for any $n \geq 1$. \square

The next goal is to obtain the convergence of v^n as $n \rightarrow \infty$.

Theorem 5.1. *Under the assumptions (H1)–(H3), let v^* solve (3.2) with σ independent of u , and let v^n from PIA. Then $v^n \rightarrow v$ as $n \rightarrow \infty$ locally uniformly in $C^{1,\alpha}$ over \mathbb{R}^d .*

Proof. By the uniform local bound of $\{v^n\}_n$ in Lemma 4.2, Corollary 5.1 and the Monotone Convergence Theorem, v^n converges locally uniformly to some function, denoted as \bar{v} .

By Lemma 5.3, v^n and \bar{v} are locally uniformly bounded in $C^{1,\alpha}$. Therefore, we actually have that $v^n \rightarrow \bar{v}$ locally uniformly in $C^{1,\alpha}$. This implies that $\pi^n(x, u) \rightarrow \Gamma(x, D\bar{v})(u)$ locally uniformly as $n \rightarrow \infty$. By the definition of r^n, b^n and the stability of viscosity solutions (under locally uniform convergence), we get that \bar{v} is a viscosity to

$$\rho v - \int_U \left[r(x, u) + b(x, u) \cdot Dv + \frac{1}{2} \text{tr}(\sigma \sigma^T(x) D^2 v) - \lambda \ln(\Gamma(x, Dv)(u)) \right] \Gamma(x, Dv)(u) du = 0.$$

The definition of Γ then yields that \bar{v} is a viscosity solution to (3.2). Thus, by the comparison principle, $\bar{v} = v^*$, which finishes the proof. \square

REFERENCES

- [1] M. Abu-Khalaf and F. L. Lewis. Nearly optimal control laws for nonlinear systems with saturating actuators using a neural network hjb approach. *Automatica*, 41(5):779–791, 2005.
- [2] R. Bellman. Dynamic programming. *Princeton University Press*, 1957.
- [3] D. P. Bertsekas. Value and policy iterations in optimal control and adaptive dynamic programming. *IEEE transactions on neural networks and learning systems*, 28(3):500–509, 2015.
- [4] S. Cacace, F. Camilli, and A. Goffi. A policy iteration method for mean field games. *ESAIM Control Optim. Calc. Var.*, 27:Paper No. 85, 19, 2021.
- [5] F. Camilli and Q. Tang. Rates of convergence for the policy iteration method for mean field games systems. *J. Math. Anal. Appl.*, 512(1):Paper No. 126138, 18, 2022.
- [6] Y.-Z. Chen and L.-C. Wu. *Second order elliptic equations and elliptic systems*, volume 174. American Mathematical Soc., 1998.
- [7] M. G. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [8] M. Dai, Y. Dong, and Y. Jia. Learning equilibrium mean-variance strategy. *Mathematical Finance*, 33(4):1166–1212, 2023.
- [9] Y. Dong. Randomized optimal stopping problem in continuous time and reinforcement learning algorithm. *arXiv preprint arXiv:2208.02409*, 2022.
- [10] P. Dupuis and R. S. Ellis. *A Weak Convergence Approach to the Theory of Large Deviations*, volume 313. John Wiley & Sons, 1997.
- [11] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1998.
- [12] D. Firoozi and S. Jaimungal. Exploratory LQG mean field games with entropy regularization. *Automatica J. IFAC*, 139:Paper No. 110177, 12, 2022.
- [13] N. Frikha, M. Germain, M. Laurière, H. Pham, and X. Song. Actor-critic learning for mean-field control in continuous time. *arXiv preprint arXiv:2303.06993*, 2023.
- [14] X. Guo, R. Xu, and T. Zariphopoulou. Entropy regularization for mean field games with learning. *Mathematics of Operations research*, 47(4):3239–3260, 2022.
- [15] R. A. Howard. Dynamic programming and markov processes. 1960.
- [16] Y.-J. Huang, Z. Wang, and Z. Zhou. Convergence of policy improvement for entropy-regularized stochastic control problems. *arXiv preprint arXiv:2209.07059*, 2022.
- [17] S. D. Jacka and A. Mijatović. On the policy improvement algorithm in continuous time. *Stochastics*, 89(1):348–359, 2017.
- [18] B. Kerimkulov, D. Šiška, and L. Szpruch. Exponential convergence and stability of Howard’s policy improvement algorithm for controlled diffusions. *SIAM J. Control Optim.*, 58(3):1314–1340, 2020.
- [19] N. V. Krylov. *Controlled diffusion processes*, volume 14. Springer Science & Business Media, 2008.
- [20] J. Lee and R. S. Sutton. Policy iterations for reinforcement learning problems in continuous time and space—fundamental theory and methods. *Automatica*, 126:109421, 2021.
- [21] J. Y. Lee, Y. Kim. Hamilton-Jacobi Based Policy-Iteration via Deep Operator Learning. *arXiv:2406.10920 [math.OC]*, preprint.
- [22] J. Ma, G. Wang, and J. Zhang. On Convergence and Rate of Convergence of Policy Improvement Algorithms. *arXiv:2406.10959 [math.OC]*, preprint.
- [23] M. L. Puterman. On the convergence of policy iteration for controlled diffusions. *J. Optim. Theory Appl.*, 33(1):137–144, 1981.
- [24] M. L. Puterman. *Markov decision processes: discrete stochastic dynamic programming*. John Wiley & Sons, 2014.
- [25] M. L. Puterman and S. L. Brumelle. On the convergence of policy iteration in stationary dynamic programming. *Mathematics of Operations Research*, 4(1):60–69, 1979.
- [26] R. S. Sutton and A. G. Barto. *Reinforcement learning: An introduction*. MIT press, 2018.
- [27] W. Tang, H. V. Tran, and Y. P. Zhang. Policy iteration for the deterministic control problems—a viscosity approach. *arXiv preprint arXiv:2301.00419*, 2023.
- [28] W. Tang, Y. P. Zhang, and X. Y. Zhou. Exploratory hjb equations and their convergence. *SIAM Journal on Control and Optimization*, 60(6):3191–3216, 2022.
- [29] H. V. Tran. *Hamilton-Jacobi equations—theory and applications*, volume 213 of *Graduate Studies in Mathematics*. American Mathematical Society, 2021.
- [30] D. Vrabie, O. Pastravanu, M. Abu-Khalaf, and F. L. Lewis. Adaptive optimal control for continuous-time linear systems based on policy iteration. *Automatica*, 45(2):477–484, 2009.
- [31] H. Wang, T. Zariphopoulou, and X. Y. Zhou. Reinforcement learning in continuous time and space: A stochastic control approach. *J. Mach. Learn. Res.*, 21:198:1–198:34, 2020.

- [32] H. Wang and X. Y. Zhou. Continuous-time mean–variance portfolio selection: A reinforcement learning framework. *Mathematical Finance*, 30(4):1273–1308, 2020.
- [33] X. Wei and X. Yu. Continuous-time q-learning for mean-field control problems. *arXiv preprint arXiv:2306.16208*, 2023.
- [34] X. Y. Zhou. Curse of optimality, and how we break it. *Available at SSRN 3845462*, 2021.

(H. V. Tran) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN MADISON, VAN VLECK HALL, 480 LINCOLN DRIVE, MADISON, WI 53706
Email address: `hung@math.wisc.edu`

(Z. Wang) DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, 396 CARVER HALL, 411 MORRILL ROAD, AMES, IA 50011
Email address: `zhenhuaw@iastate.edu`

(Y. P. Zhang) DEPARTMENT OF MATHEMATICS AND STATISTICS, AUBURN UNIVERSITY, PARKER HALL, 221 ROOSEVELT CONCOURSE, AUBURN, AL 36849
Email address: `yzhangpaul@auburn.edu`