

# Green's functions in the presence of a bubble wall

---

**Takahiro Kubota** <sup>a</sup>

<sup>a</sup>*CELAS, Osaka University, Toyonaka, Osaka, 560-0043, Japan*

*E-mail:* [takahirokubota859@hotmail.com](mailto:takahirokubota859@hotmail.com)

ABSTRACT: Field theoretical tools are developed so that one can analyze quantum phenomena such as transition radiation that must have occurred during the Higgs condensate bubble expansion through plasma in the early universe. Integral representations of Bosonic and Fermionic propagators are presented for the case that particle masses are varied continuously during the passage through the bubble wall interface between symmetry-restored and symmetry-broken regions. The construction of propagators is based on the so-called eigenfunction expansion method associated with self-adjoint differential operators, developed by Weyl, Stone, Titchmarsh, Kodaira and several others. A novel method of field quantization in the presence of the bubble wall is proposed by using the spectral functions introduced in constructing the two-point Green's functions.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The scalar particle's two-point Green's function</b>	<b>4</b>
2.1	The method à la Weyl, Stone, Titchmarsh and Kodaira	4
2.2	The constant mass case (no bubble wall)	7
2.3	A solvable model of the bubble wall	8
<b>3</b>	<b>The Dirac Fermion's two-point Green's function</b>	<b>14</b>
3.1	Preliminaries	14
3.2	The canonical form	14
3.3	The transverse momentum dependence of the energy $E$	17
3.4	The general expansion method and completeness	18
3.5	Construction of the Propagator	19
3.6	The constant mass case (no bubble wall)	20
<b>4</b>	<b>The vector boson's two-point Green's function</b>	<b>23</b>
4.1	Mixing between gauge boson and unphysical scalar fields	23
4.2	The general expansion method for two-component fields	25
<b>5</b>	<b>Field quantization in the presence of the bubble wall</b>	<b>27</b>
5.1	Real scalar field	28
5.2	The Dirac field	30
<b>6</b>	<b>Concluding Remarks</b>	<b>31</b>

---

## 1 Introduction

The discoveries of the gravitational waves by LIGO and Virgo collaboration [1] - [4] are apparently a strong driving force, urging us not only to look for further astrophysical sources but also to gain a direct access to the physics of the early universe by observing gravitational radiation [5] - [8]. The standard model phase transition is smooth cross-over [9] - [17], but extension of the electroweak model can lead to first-order phase transitions [18] - [20] that could have produced observable gravitational waves in accessible frequency ranges for future experiments. The LISA space-based interferometer [21] - [24], which is planned to have sensitivity band peaks at lower frequencies than ground based detectors, is expected to probe first-order phase transitions of the extended electroweak models in the early universe. Other planned space-based gravitational wave observation projects include DECIGO [25], Taiji [26] and TianQin [27].

The strength and spectral shape of the gravitational wave in the phase transitions depend on several parameters, which include (i) the characteristic time duration of the phase transition, (ii) the ratio of the vacuum energy density released in the transition to the radiation bath, (iii) the temperature and the Hubble parameter at the time of the phase transition, (iv) the fraction of vacuum energy converted into bulk motion of the fluid and into the gradient energy of the Higgs field and (v) the bubble wall velocity in the plasma rest frame [21] - [24].

Among these parameters the bubble wall velocity is the least clearly understood quantity. Unless the microscopic mechanism of friction exerted by the fluid is fully understood, the wall velocity is basically left as an ambiguous parameter in the hydrodynamical treatment of the plasma. The wall velocity is also an important quantity for the generation of the matter-antimatter asymmetry [28] - [33]. Bödeker and Moore [34] have once investigated the propagation speed of the interface between symmetry-restored and symmetry-broken regions. Particles crossing the bubble wall lose some part of their momentum in the direction perpendicular to the wall, thereby impeding the expansion of the bubble wall. The question was whether the bubble wall would become relativistic or even ultra-relativistic and therefore run away. They extended the standard electroweak theory with extra scalar fields and looked for a simple criterion to determine whether or not the runaway of the bubble wall could occur. Espinosa et al [35] also made a comprehensive study of the bubble wall velocity, classifying the bubble expansion regimes into deflagrations, detonations, hybrids and runaway.

Several years later, Bödeker and Moore revisited the bubble wall runaway problem [36], by considering transition radiation effects [37, 38], which had been neglected in their previous work [34]. When particles hit the bubble wall, they may well radiate particles with phase-dependent mass, such as  $W$ - and  $Z$ -bosons, and the emission induces important effects on the bubble wall. It was argued that the force is dominated by soft particle radiation and that the friction force could be significantly altered due to logarithmically-enhanced radiative corrections. The appearance of the large logarithms is traced back to the soft momentum region in the phase space and is therefore of infrared character. The analyses in [36] showed that without considering the transition radiation the friction on the wall would be underestimated.

The transition radiation considered in [36] is due to the process of a particle decaying into two, one of which is a soft vector boson. If such a process is important, then a question naturally arises whether the processes of a particle decaying into several soft particles could be equally important. Fixed order calculations will quite probably break down for very large wall velocity. Höche et al [39] therefore attempted to perform the all-order summation of the large logarithmic terms due to the multiple emission of soft gauge bosons. In the course of the calculation they also paid attention to double logarithms of the Sudakov type [40], whose all-order resummation and exponentiation had been extensively studied in QED [41, 42] and QCD [43] - [46]. After all they concluded that the logarithmic enhancement implies significantly slower terminal velocities than previously thought, to find a peculiar scaling law of the pressure. (See also [47] for similar scaling law.)

Their formulation in [39] reminds us of the parton shower algorithm [48, 49] in hadron

collisions formulated as an evolution in momentum transfer from high energy scale to the lower. In the QCD-based resummation technique, the experimental resolution scale plays the role of regulator of infrared singularities. In the electroweak radiative corrections, the dominant contributions are also expected to come from double logarithms of the Sudakov type as well as single logarithms involving  $\sqrt{s}/M_Z$ , when the energy  $\sqrt{s}$  of the system is much larger than the electroweak energy scale, say, the  $Z$ -boson mass  $M_Z$  ( i.e.,  $\sqrt{s} \gg M_Z$ ). The origin of these logarithms is in close correspondence with the soft and collinear singularities known well in massless gauge theories. In contrast to the massless gauge theories such as QED and QCD where cancellation occurs among virtual corrections and real gauge boson radiation, the masses of  $Z$ - and  $W$ -bosons in the electroweak theory provide the physical cutoff and the large logarithms originating from virtual corrections are of direct physical significance [50]- [57].

The processes that must have occurred during the bubble expansion in the early universe are somewhere between massless and massive gauge boson theories, and we have to take care of the fact that the masses of the standard model particles, irrespectively of real or virtual ones, may change near the bubble wall. In Ref. [39] real emissions of gauge bosons and virtual corrections were combined together to see infrared divergence cancellations, while neglecting the change of masses of vector bosons at the domain wall. In order to see the varying mass effects, it might be tempting to use the so-called vacuum expectation value insertion approximation, but we must exert great care in using such an approximation when infrared singularities are involved.

In the last several year a lot of efforts have been made [58] - [73] in order to understand in a better way the dynamics occurring between plasma particles and bubble wall. We must say, however, that we are not yet fully equipped with theoretical tools to perform quantum field theoretical analyses such as loop calculations. One of the sources of difficulties is that we do not have at hand a suitable way of constructing field propagators in which the bubble wall background effects are fully taken into account. This unsatisfactory situation is closely connected with the lack of suitable field quantization methods for the case in which particles masses change between symmetry-restored and symmetry-broken regions.

The purpose of the present paper is to improve the present status quo by constructing explicitly the two-point Green's functions, i.e., propagators of scalar, spinor and vector fields for the case that particle masses are position-dependent. Throughout the present paper we will work in the Lorentz frame in which the bubble wall is at rest. We will solve the partial differential equations for the propagators with position-dependent mass by making use of the so-called eigenfunction expansion method associated with self-adjoint differential operators developed several decades ago in mathematical literatures [74] - [78]. Our method is general enough and is applicable irrespectively of the details of the shape of the bubble wall. Our two-point Green's functions are expressed in the form of integral representations over the spectrum described by spectral functions. On the basis of the spectral functions, the field quantization method is also reexamined for the case that masses are position-dependent. We will introduce operators that are analogous to the ordinary creation and annihilation operators of particles and will give the definition of the vacuum using these operators. It is argued that the propagators to be derived in this work are the expectation

value of time-ordered product of fields with respect to this vacuum.

The first-principle method of field quantization in the presence of a bubble wall was recently discussed in detail by Azatov et al [73]. We agree completely with their view on the necessity of the first-principle field-quantization method, while the approach taken in the present paper differs from theirs in that emphasis is placed on the roles played by the spectral functions to be discussed below.

## 2 The scalar particle's two-point Green's function

### 2.1 The method à la Weyl, Stone, Titchmarsh and Kodaira

The method of constructing the propagators in the presence of the bubble wall is most easily explained for the case of the spin zero particle obeying the Klein-Gordon equation. We suppose that the bubble wall is so large in comparison with our microscopic length scale and that the front of the wall can be approximated by a nearly planar wall. For the sake of simplicity we assume that the wall is spreading infinitely along the  $x - y$ -plane. This means that the expectation value of the Higgs field as well as particle masses depend on the  $z$ -coordinate only. Denoting the mass by  $M(z)$ , we identify  $M(+\infty)$  and  $M(-\infty)$  with the mass in the symmetry-broken and unbroken regions, respectively. For spineless particles satisfying the Klein-Gordon equation, we would like to look for the Green's function that satisfies

$$\begin{aligned} \{\square + M(z)^2\} \mathcal{G}(t - t', x - x', y - y', z, z'; M(*)) \\ = -i \delta(t - t') \delta(x - x') \delta(y - y') \delta(z - z'), \end{aligned} \quad (2.1)$$

$$\begin{aligned} \{\square' + M(z')^2\} \mathcal{G}(t - t', x - x', y - y', z, z'; M(*)) \\ = -i \delta(t - t') \delta(x - x') \delta(y - y') \delta(z - z'), \end{aligned} \quad (2.2)$$

where  $\square$  and  $\square'$  are the usual d'Alembertian operators

$$\square = \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2}, \quad \square' = \frac{\partial^2}{\partial t'^2} - \frac{\partial^2}{\partial x'^2} - \frac{\partial^2}{\partial y'^2} - \frac{\partial^2}{\partial z'^2}. \quad (2.3)$$

Our notation  $M(*)$  as opposed to  $M(z)$  or  $M(z')$  in (2.1) and (2.2) is meant that the Green's function depends only indirectly on the functional form of the  $z$ -dependent mass. Note that the translational invariances in the  $t$ -,  $x$ - and  $y$ - directions are just as usual and one can express the solution of (2.1) and (2.2) as a superposition of the plane waves in the  $x$ - and  $y$ -directions. The translational invariance in the  $z$ -direction, however, is not respected due to the presence of the wall and the mass term  $M(z)^2$ , and therefore (2.2) does not follow from (2.1) immediately.

We now argue that a certain mathematical framework developed in Refs. [74] - [78] is quite useful to give a closed form of the solutions to (2.1) and (2.2). It is a generalized expansion method associated with second-order differential equations and is applicable respectively of the detailed form of the function  $M(z)$ . We will look for the solution to

(2.1) and (2.2) in the following integral representation

$$\begin{aligned} & \mathcal{G}(t-t', x-x', y-y', z, z'; M(*)) \\ &= i \int \frac{dE d^2\vec{p}_\perp}{(2\pi)^3} e^{-iE(t-t')} e^{i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)} \int \sum_{k,l=1,2} \frac{\phi_k(z; \lambda) d\rho_{kl}(\lambda) \phi_l(z'; \lambda)}{E^2 - \vec{p}_\perp^2 - \lambda + i\varepsilon}, \end{aligned} \quad (2.4)$$

where  $\phi_k(z; \lambda)$  is the solution to the following differential equation in the infinite region  $-\infty < z < +\infty$

$$\left\{ -\frac{d^2}{dz^2} + M(z)^2 \right\} \phi_k(z; \lambda) = \lambda \phi_k(z; \lambda), \quad (k = 1, 2) \quad (2.5)$$

and  $\lambda$  is a parameter of the theory. We have introduced in (2.4) the two-dimensional position and momentum vectors,  $\vec{x}_\perp = (x, y, 0)$ ,  $\vec{x}'_\perp = (x', y', 0)$  and  $\vec{p}_\perp = (p_x, p_y, 0)$  together with the two-dimensional integration volume  $d^2\vec{p}_\perp = dp_x dp_y$ .

We notice immediately that Eq. (2.5) is the one-dimensional Schrödinger-type equation with “potential term”  $M(z)^2$  and the “energy eigenvalue”  $\lambda$ . Since this is the second-order differential equation, we have two independent solutions, which we denote by  $\phi_1(z; \lambda)$  and  $\phi_2(z; \lambda)$ , respectively, and will be called fundamental solutions. The normalization of these solutions is fixed by the initial conditions at  $z = 0$ , i.e.,

$$\phi_1(0; \lambda) = 1, \quad \phi_1'(0; \lambda) = 0, \quad (2.6)$$

$$\phi_2(0; \lambda) = 0, \quad \phi_2'(0; \lambda) = 1. \quad (2.7)$$

The derivative with respect to  $z$  is denoted by prime ( $\prime$ ) in (2.6) and (2.7). The spectrum of  $\lambda$  is described by  $2 \times 2$  spectral function matrix  $\rho_{kl}(\lambda)$  and Eq. (2.4) is the Stieltjes integral with respect to  $\rho_{kl}(\lambda)$ . This spectral function will play the most important roles throughout the present work. Since we are considering Eq. (2.5) in the infinite region  $-\infty < z < +\infty$ , our spectral problem is more general and is more singular in a sense than the Sturm-Liouville boundary value problem, in which we usually deal with finite regions. As we will see later, (2.4) should go over to the conventional Feynman’s propagator if the mass is  $z$ -independent, and we have added  $+i\varepsilon$  by hand in the denominator of the integrand of (2.4). This  $+i\varepsilon$  prescription indicates that the Green’s function (2.4) corresponds to the vacuum expectation value of the time-ordered product of relevant scalar fields, but the justification of the prescription will be given only after we specify what we mean by the vacuum in the presence of the bubble wall.

Let us apply the differential operator  $\{\square + M(z)^2\}$  on both sides of (2.4) and we then obtain

$$\begin{aligned} & \{\square + M(z)^2\} \mathcal{G}(t-t', x-x', y-y', z, z'; M(*)) \\ &= i \int \frac{dE d^2\vec{p}_\perp}{(2\pi)^3} e^{-iE(t-t')} e^{i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)} \\ & \quad \times \int \sum_{k,l=1,2} \left\{ -E^2 + \vec{p}_\perp^2 - \frac{\partial^2}{\partial z^2} + M(z)^2 \right\} \frac{\phi_k(z; \lambda) d\rho_{kl}(\lambda) \phi_l(z'; \lambda)}{E^2 - \vec{p}_\perp^2 - \lambda + i\varepsilon}. \end{aligned} \quad (2.8)$$

Using the differential equation (2.5) in (2.8) we immediately arrive at

$$\begin{aligned} & \{\square + M(z)^2\} \mathcal{G}(t - t', x - x', y - y', z, z'; M(*)) \\ &= -i \delta(t - t') \delta(x - x') \delta(y - y') \int \sum_{k,l=1,2} \phi_k(z; \lambda) d\rho_{kl}(\lambda) \phi_l(z'; \lambda). \end{aligned} \quad (2.9)$$

By virtue of the work of Weyl [74] and of Stone [75] it has been known that there exists the matrix  $\rho_{kl}(\lambda)$  with such a nice property that any real-valued *square integrable* function  $f(z)$  can be expressed as a superposition of the fundamental solutions  $\phi_k(z; \lambda)$  ( $k = 1, 2$ ), i.e.,

$$f(z) = \int \sum_{k,l=1,2} \phi_k(z; \lambda) d\rho_{kl}(\lambda) \int_{-\infty}^{\infty} dz' \phi_l(z'; \lambda) f(z'). \quad (2.10)$$

In other words we are able to write down the formula

$$\int \sum_{k,l=1,2} \phi_k(z; \lambda) d\rho_{kl}(\lambda) \phi_l(z'; \lambda) = \delta(z - z'), \quad (2.11)$$

in the sense of (2.10). Eq. (2.11) being substituted into (2.9), we are able to go back to the defining equation (2.1) of the Green's function. It is obvious that (2.4) also satisfies (2.2).

While it had been shown in [74] that  $\rho_{kl}(\lambda_2) - \rho_{kl}(\lambda_1)$  is a positive semi-definite matrix for  $\lambda_2 > \lambda_1$ , Titchmarsh [76] and Kodaira [77] went one step further to derive the relations

$$\rho_{11}(\lambda_2) - \rho_{11}(\lambda_1) = \lim_{\varepsilon \rightarrow +0} \frac{1}{\pi} \int_{\lambda_1}^{\lambda_2} \Im \left( \frac{1}{m_1(\lambda + i\varepsilon) - m_2(\lambda + i\varepsilon)} \right) d\lambda, \quad (2.12)$$

$$\begin{aligned} \rho_{12}(\lambda_2) - \rho_{12}(\lambda_1) &= \rho_{21}(\lambda_2) - \rho_{21}(\lambda_1) \\ &= \lim_{\varepsilon \rightarrow +0} \frac{1}{2\pi} \int_{\lambda_1}^{\lambda_2} \Im \left( \frac{m_1(\lambda + i\varepsilon) + m_2(\lambda + i\varepsilon)}{m_1(\lambda + i\varepsilon) - m_2(\lambda + i\varepsilon)} \right) d\lambda, \end{aligned} \quad (2.13)$$

$$\rho_{22}(\lambda_2) - \rho_{22}(\lambda_1) = \lim_{\varepsilon \rightarrow +0} \frac{1}{\pi} \int_{\lambda_1}^{\lambda_2} \Im \left( \frac{m_1(\lambda + i\varepsilon) m_2(\lambda + i\varepsilon)}{m_1(\lambda + i\varepsilon) - m_2(\lambda + i\varepsilon)} \right) d\lambda, \quad (2.14)$$

and thereby paved the way for determining the spectral function  $\rho_{kl}(\lambda)$ . Here the symbol  $\Im$  denotes the imaginary part and the two quantities  $m_1(\lambda + i\varepsilon)$  and  $m_2(\lambda + i\varepsilon)$  are called characteristic functions in [77]. The residue theorem in the complex function theory is made use of extensively in [76], while the spectral theorem for the self-adjoint differential operator plays the key role in [77]. See Eqs.(3.1.5) - (3.1.8) in Ref.[76] and Eq.(1.15) in Ref.[77].

Weyl classified the properties of the boundary points ( $z = +\infty$  and  $z = -\infty$  in the present case) into two types, "limit circle type" and "limit point type" [74]. If both of  $z = +\infty$  and  $z = -\infty$  are the limit point type, then the characteristic functions are given implicitly in [76] and explicitly in [77] in the following form

$$m_1(\lambda + i\varepsilon) = - \lim_{a \rightarrow -\infty} \frac{\phi_1(a; \lambda + i\varepsilon)}{\phi_2(a; \lambda + i\varepsilon)}, \quad m_2(\lambda + i\varepsilon) = - \lim_{b \rightarrow +\infty} \frac{\phi_1(b; \lambda + i\varepsilon)}{\phi_2(b; \lambda + i\varepsilon)}. \quad (2.15)$$

See also Refs. [79, 80] for an alternative and simpler derivation of (2.15). Procedures of computing the characteristic functions for the case of the limit circle type have also been worked out, although a little involved. For more details, see [77].

If  $M(z)^2$  is such that the Schrödinger type equation (2.5) admits bound state solutions, the spectral functions increase stepwise as  $\lambda$  goes up through the corresponding binding energy level. If there is no bound states, on the other hand, the spectral functions are continuous and differentiable. Then the Stieltjes integral in (2.4) turns out to be the ordinary Riemann integral over  $\lambda$  with the following spectral weight,

$$d\rho_{11}(\lambda) = \lim_{\varepsilon \rightarrow +0} \frac{1}{\pi} \Im \left( \frac{1}{m_1(\lambda + i\varepsilon) - m_2(\lambda + i\varepsilon)} \right) d\lambda, \quad (2.16)$$

$$\begin{aligned} d\rho_{12}(\lambda) &= d\rho_{21}(\lambda) \\ &= \lim_{\varepsilon \rightarrow +0} \frac{1}{2\pi} \Im \left( \frac{m_1(\lambda + i\varepsilon) + m_2(\lambda + i\varepsilon)}{m_1(\lambda + i\varepsilon) - m_2(\lambda + i\varepsilon)} \right) d\lambda, \end{aligned} \quad (2.17)$$

$$d\rho_{22}(\lambda) = \lim_{\varepsilon \rightarrow +0} \frac{1}{\pi} \Im \left( \frac{m_1(\lambda + i\varepsilon) m_2(\lambda + i\varepsilon)}{m_1(\lambda + i\varepsilon) - m_2(\lambda + i\varepsilon)} \right) d\lambda. \quad (2.18)$$

The formulae (2.15) - (2.18) tell us that we can evaluate the spectral density  $d\rho_{kl}(\lambda)/d\lambda$ , once we know the asymptotic behavior at infinity of the fundamental solutions defined by (2.6) and (2.7). This is the key observation in the spectral theory.

## 2.2 The constant mass case (no bubble wall)

As the simplest exercise, let us consider the well-known case, i.e.,  $M(z)^2 = M^2$  (constant), for which there is no bubble wall. The fundamental solutions to (2.5) satisfying (2.6) and (2.7) for  $\lambda > M^2$  are simply given by

$$\phi_1(z; \lambda) = \cos \left( \sqrt{\lambda - M^2} z \right), \quad \phi_2(z; \lambda) = \frac{1}{\sqrt{\lambda - M^2}} \sin \left( \sqrt{\lambda - M^2} z \right), \quad (2.19)$$

and those for  $\lambda < M^2$  are obtained by analytic continuation of (2.19) with respect to  $\lambda$  or more explicitly are given by

$$\phi_1(z; \lambda) = \cosh \left( \sqrt{M^2 - \lambda} z \right), \quad \phi_2(z; \lambda) = \frac{1}{\sqrt{M^2 - \lambda}} \sinh \left( \sqrt{M^2 - \lambda} z \right), \quad (2.20)$$

The characteristic functions are computed by using (2.15) and we get

$$m_1(\lambda) = -i\sqrt{\lambda - M^2}, \quad m_2(\lambda) = i\sqrt{\lambda - M^2}, \quad \text{for } \lambda > M^2, \quad (2.21)$$

$$m_1(\lambda) = \sqrt{M^2 - \lambda}, \quad m_2(\lambda) = -\sqrt{M^2 - \lambda}, \quad \text{for } \lambda < M^2. \quad (2.22)$$

Therefore the integral in (2.4) should be done by setting

$$d\rho_{11}(\lambda) = \begin{cases} \frac{1}{2\pi} \frac{1}{\sqrt{\lambda - M^2}} d\lambda, & \text{for } \lambda > M^2, \\ 0 & \text{for } \lambda < M^2, \end{cases} \quad (2.23)$$

$$d\rho_{12}(\lambda) = d\rho_{21}(\lambda) = 0, \quad (2.24)$$

$$d\rho_{22}(\lambda) = \begin{cases} \frac{1}{2\pi} \sqrt{\lambda - M^2} d\lambda, & \text{for } \lambda > M^2, \\ 0 & \text{for } \lambda < M^2. \end{cases} \quad (2.25)$$



Note that the integration region in (2.4) is automatically restricted to  $\lambda > M^2$ , by virtue of the above formulae.

Combining the solutions (2.19) with (2.23), (2.24) and (2.25) we get

$$\begin{aligned} \int \sum_{k,l=1,2} \phi_k(z; \lambda) d\rho_{kl}(\lambda) \phi_l(z'; \lambda) &= \frac{1}{2\pi} \int_{M^2}^{\infty} \cos \left\{ \sqrt{\lambda - M^2} (z - z') \right\} \frac{d\lambda}{\sqrt{\lambda - M^2}} \\ &= \frac{1}{2\pi} \int_0^{\infty} \left\{ e^{ip_z(z-z')} + e^{-ip_z(z-z')} \right\} dp_z \\ &= \delta(z - z'), \end{aligned} \quad (2.26)$$

where we set  $p_z = \sqrt{\lambda - M^2}$ . We can easily confirm in this example that our Green's function (2.4) turns out to be the familiar Feynman's scalar field propagator, i.e.,

$$\Delta_F(x - x'; M) = \int \frac{d^4 p}{(2\pi)^4} e^{-ip \cdot (x - x')} \frac{i}{p^2 - M^2 + i\varepsilon}. \quad (2.27)$$

The method of the general expansion method explained in Sec. 2.1 is thus reduced for the constant mass case to the familiar Fourier expansion. If the mass is position-dependent, however, the Fourier expansion is no more useful and the general expansion method in Sec. 2.1 is an appropriate tool for diagrammatical analyses.

### 2.3 A solvable model of the bubble wall

The second example which is analytically solvable is given by the following mass function

$$M(z)^2 = \mu^2(1 - s) + M_0^2 s + u s(1 - s), \quad (2.28)$$

where we have introduced a new variable

$$s = \frac{1}{2} \left\{ 1 + \tanh \left( \frac{z}{2l} \right) \right\} \quad (2.29)$$


in place of  $z$ , thereby mapping the whole  $z$ -space ( $-\infty < z < +\infty$ ) into the finite region  $0 < s < 1$ . This model has once been called attention to in [81] and was studied thoroughly in Refs. [82, 83] in connection with a scattering problem in the presence of a domain wall. The profile of the bubble wall is described by four parameters,  $\mu^2$ ,  $M_0^2$ ,  $u$ , and  $l$ . The gross shape of the bubble wall is illustrated in Figure 1 for  $u = 0$ . Note that for small values of  $u$ ,  $M(z)^2$  is monotonically increasing and

$$\lim_{z \rightarrow +\infty} M(z)^2 = M_0^2, \quad \lim_{z \rightarrow -\infty} M(z)^2 = \mu^2. \quad (2.30)$$

The size of the bubble wall interface is determined by the length scale  $l$ . The parameter  $M_0$  is the mass observed in the symmetry-broken phase and  $\mu$  is expected to play the role of infrared regularization and should be taken to be zero eventually.

Let us seek for the solutions to (2.5) by setting

$$\phi_k(z; \lambda) = s^{\kappa_0} (1 - s)^{\kappa_1} \psi_k(s; \lambda), \quad (k = 1, 2). \quad (2.31)$$



**Figure 1.** The gross shape of the bubble wall given by (2.28) with  $u = 0$ .

The two exponents  $\kappa_0$  and  $\kappa_1$  are determined by

$$\kappa_0^2 = (\mu^2 - \lambda) l^2, \quad \kappa_1^2 = (M_0^2 - \lambda) l^2, \quad (2.32)$$

and the choice of Riemann sheets when taking the square root will be discussed later. Eq. (2.5) is now transformed into the following form

$$\begin{aligned} 0 &= \left\{ -\frac{d^2}{dz^2} + M(z)^2 - \lambda \right\} \phi_k(z; \lambda) \\ &= -\frac{1}{l^2} s^{\kappa_0+1} (1-s)^{\kappa_1+1} \left[ s(1-s) \frac{d^2}{ds^2} + \{\gamma - (\alpha + \beta + 1)s\} \frac{d}{ds} - \alpha\beta \right] \psi_k(s; \lambda), \end{aligned} \quad (2.33)$$

where we have defined  $\alpha$ ,  $\beta$  and  $\gamma$  as

$$\alpha = \kappa_0 + \kappa_1 + \frac{1}{2} + \delta, \quad (2.34)$$

$$\beta = \kappa_0 + \kappa_1 + \frac{1}{2} - \delta, \quad (2.35)$$

$$\gamma = 1 + 2\kappa_0, \quad (2.36)$$

$$\delta = \sqrt{\frac{1}{4} - ul^2}. \quad (2.37)$$

Note that  $\kappa_0$  and  $\kappa_1$  together with  $\alpha$ ,  $\beta$  and  $\gamma$  all depend on  $\lambda$ .

Eq. (2.33) is the Gauss's hypergeometric equation and there are two independent solutions around  $s = 0$ , one of which is given by

$$F(\alpha, \beta, \gamma; s) = 1 + \frac{\alpha\beta}{\gamma} s + \frac{\alpha(\alpha+1)\beta(\beta+1)}{\gamma(\gamma+1)} \frac{s^2}{2!} + \dots \quad (2.38)$$

There are several ways of expressing the second solution and here we take the following one,

$$s^{1-\gamma} F(1 + \alpha - \gamma, 1 + \beta - \gamma, 2 - \gamma; s). \quad (2.39)$$

Thus we express  $\psi_k(s; \lambda)$  as a linear combination of (2.38) and (2.39), namely,

$$\begin{aligned} \psi_k(s; \lambda) = & c_{k1}(\lambda) F(\alpha, \beta, \gamma; s) \\ & + c_{k2}(\lambda) s^{1-\gamma} F(1 + \alpha - \gamma, 1 + \beta - \gamma, 2 - \gamma; s), \quad (k = 1, 2). \end{aligned} \quad (2.40)$$

Since we are interested in the characteristic functions (2.15), we ought to examine the  $z \rightarrow -\infty$  and  $z \rightarrow +\infty$  behavior of (2.31) or equivalently  $s \rightarrow 0$  and  $s \rightarrow 1$  behavior of (2.40). The first term of (2.40) in the  $s \rightarrow 0$  limit is simply determined by the formula  $F(\alpha, \beta, \gamma; 0) = 1$ , while the behavior of the second term depends on the factor  $s^{1-\gamma}$ . Here we require the following inequality for the real part ( $\Re$ ) of the exponent

$$\Re(1 - \gamma) = -2\Re(\kappa_0) > 0, \quad (2.41)$$

under which the first term in (2.40) is more dominant than the second as  $s \rightarrow 0$ . This indicates that we should choose  $\kappa_0$  as

$$\kappa_0 = \begin{cases} -\sqrt{\mu^2 - \lambda} l & \text{for } \mu^2 > \lambda, \\ i\sqrt{\lambda + i\varepsilon - \mu^2} l & \text{for } \lambda > \mu^2, \end{cases} \quad (2.42)$$

where we always take the  $\varepsilon \rightarrow +0$  limit.

Next let us turn to the  $s \rightarrow 1$  limit of (2.40). We know that  $F(\alpha, \beta, \gamma; 1)$  is finite and is given by the Gauss's formula

$$F(\alpha, \beta, \gamma; 1) = \frac{\Gamma(\gamma)\Gamma(\gamma - \alpha - \beta)}{\Gamma(\gamma - \alpha)\Gamma(\gamma - \beta)}, \quad (2.43)$$

if  $\Re\gamma > 0$  and the following inequality is satisfied

$$\Re(\gamma - \alpha - \beta) = -2\Re(\kappa_1) > 0. \quad (2.44)$$

Note that  $F(1 + \alpha - \gamma, 1 + \beta - \gamma, 2 - \gamma; s)$  in the second term of (2.40) is also finite as  $s \rightarrow 1$  if  $\Re(2 - \gamma) > 0$  and  $\Re(\gamma - \alpha - \beta) > 0$ . Thus in order to meet the inequality (2.44),  $\kappa_1$  has to be

$$\kappa_1 = \begin{cases} -\sqrt{M_0^2 - \lambda} l & \text{for } M_0^2 > \lambda, \\ i\sqrt{\lambda + i\varepsilon - M_0^2} l & \text{for } \lambda > M_0^2. \end{cases} \quad (2.45)$$

With these choices of the Riemann sheets, the characteristic functions (2.15) turn out to be

$$\begin{aligned} m_1(\lambda) &= -\frac{c_{11}(\lambda)}{c_{21}(\lambda)}, \\ m_2(\lambda) &= -\frac{c_{11}(\lambda) F(\alpha, \beta, \gamma; 1) + c_{12}(\lambda) F(1 + \alpha - \gamma, 1 + \beta - \gamma, 2 - \gamma; 1)}{c_{21}(\lambda) F(\alpha, \beta, \gamma; 1) + c_{22}(\lambda) F(1 + \alpha - \gamma, 1 + \beta - \gamma, 2 - \gamma; 1)}. \end{aligned} \quad (2.46)$$

The coefficients  $c_{kl}(\lambda)$  in (2.40) are determined by the conditions (2.6) and (2.7) at  $z = 0$  (or equivalently at  $s = 1/2$ ), and we list up the results:

$$c_{11}(\lambda) = \frac{2^{\kappa_0 + \kappa_1 + \gamma - 1}}{W} \left[ -2(\kappa_0 + \kappa_1)F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; 1/2) + F'(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; 1/2) \right], \quad (2.47)$$

$$c_{12}(\lambda) = -\frac{2^{\kappa_0 + \kappa_1}}{W} \left[ 2(\kappa_0 - \kappa_1)F(\alpha, \beta, \gamma; 1/2) + F'(\alpha, \beta, \gamma; 1/2) \right], \quad (2.48)$$

$$c_{21}(\lambda) = -\frac{2^{\kappa_0 + \kappa_1 + \gamma + 1}}{W} F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; 1/2) l, \quad (2.49)$$

$$c_{22}(\lambda) = \frac{2^{\kappa_0 + \kappa_1 + 2}}{W} F(\alpha, \beta, \gamma; 1/2) l. \quad (2.50)$$

Here  $W$  in the denominator is given by

$$W = 2^{\gamma - 1} \left[ F(\alpha, \beta, \gamma; 1/2)F'(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; 1/2) - F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; 1/2)F'(\alpha, \beta, \gamma; 1/2) + 2(1 - \gamma)F(\alpha, \beta, \gamma; 1/2)F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; 1/2) \right]. \quad (2.51)$$


With these coefficients, we illustrate the behavior of  $\phi_1(z; \lambda)$  and  $\phi_2(z; \lambda)$  in Figure 2 for sample values of  $M_0$ ,  $\mu$ ,  $\lambda$  and  $u$  in the mass unit  $1/l$ . Figure 2 exhibits oscillatory behaviors like sinusoidal waves. The hypergeometric function shows similar behavior in the finite  $z$  region, but at infinity ( $z \rightarrow \pm\infty$ , i.e.,  $s \approx 0$  and  $s \approx 1$ ), the prefactor  $s^{\kappa_0}(1-s)^{\kappa_1}$  in Eq. (2.31) and  $s^{1-\gamma} = s^{-2\kappa_0}$  in the second term in (2.40) are also responsible for such behaviors. Note that both  $\kappa_0$  and  $\kappa_1$  are pure imaginary for our choice of the parameters in Figure 2. For large  $|z|$ , we have

$$\begin{aligned} (1-s) &\sim e^{-z/l}, & (1-s)^{\kappa_1} &\sim e^{-\kappa_1 z/l} & \text{for } z &\rightarrow +\infty, \\ s &\sim e^{z/l}, & s^{\kappa_0} &\sim e^{\kappa_0 z/l} & \text{for } z &\rightarrow -\infty, \end{aligned} \quad (2.52)$$

and therefore the large negative  $z$  behavior of  $\phi_k(z; \lambda)$  is given by a linear combination of  $\cos(|\kappa_0|z/l)$  and  $\sin(|\kappa_0|z/l)$ . Likewise the large positive  $z$  behavior is a combination of  $\cos(|\kappa_1|z/l)$  and  $\sin(|\kappa_1|z/l)$ .

The region sandwiched by the two dashed lines in Figure 2 ( $-1 < z/l < +1$ ) is the bubble wall interface and we can see obviously that the wave length in the  $z \rightarrow +\infty$  region is longer than in the  $z \rightarrow -\infty$  region. This means that the  $z$ -component of momentum carried by the propagating wave is not conserved and some amount of  $z$ -momentum is transferred to the bubble wall when moving toward the  $z \rightarrow +\infty$  region through the interface.


With the explicit formulae (2.46), we can compute the  $2 \times 2$  spectral function matrix  $\rho_{kl}(\lambda)$  by using the formulae (2.12), (2.13) and (2.14). Without entering detailed calculations, however, one thing can be seen clearly. We notice that, for  $M_0^2 > \mu^2 > \lambda$ , both  $\kappa_0$  and  $\kappa_1$  are real numbers and that  $\alpha$ ,  $\beta$  and  $\gamma$  are all real numbers accordingly. This tells us



**Figure 2.** The typical behavior of the fundamental solutions  $\phi_1(z; \lambda)$  (red curve) and  $\phi_2(z; \lambda)/l$  (blue curve) in the model (2.28) for  $\lambda > M_0^2 > \mu^2$ . Note that both  $\phi_1(z; \lambda)$  and  $\phi_2(z; \lambda)/l$  are dimensionless. We take  $1/l$  as the unit of the mass scale, and other parameters are chosen as follows:  $M_0 = 10/l$ ,  $\mu = 10^{-4}/l$ ,  $\lambda = 110/l^2$  and  $u=0$ . The region  $-1 < z/l < +1$  sandwiched by the two dashed lines is the bubble wall interface.

that there does not exist any source giving rise to imaginary part on the right hand side of (2.12), (2.13) and (2.14). Therefore we can safely say that the integration  $d\rho_{kl}(\lambda)$  in (2.4) is restricted in the region  $\lambda > \mu^2$ .

The characteristic functions in the regions  $M_0^2 > \lambda > \mu^2$  and  $\lambda > M_0^2$  can be studied only numerically and we illustrate the behavior of  $d\rho_{11}(\lambda)/d\lambda$  in Figure 3. Looking at the behavior shown in Figure 3 (a) for the region  $M_0^2 > \lambda > \mu^2$ , we immediately notice the existence of several ups and downs in sharp contrast with the smooth behavior in the region  $\lambda > M_0^2$  in Figure 3 (b). The contrast of similar kind can be seen in  $d\rho_{12}(\lambda)/d\lambda = d\rho_{21}(\lambda)/d\lambda$  and  $d\rho_{22}(\lambda)/d\lambda$ . Although origins of this contrast have to be scrutinized further, this could be a warning message that we have to be careful of the soft (i.e., small  $\lambda$ ) region integration in the loop diagrams. Large single and double logarithmic effects are closely connected with integration over soft and/or collinear regions in phase space, and such a behavior as in Figure 3 (a) could have important influences in analogous calculations in the presence of a bubble wall.



**Figure 3.** The spectral function  $\frac{1}{l} \frac{d\rho_{11}(\lambda)}{d\lambda}$  for the region (a)  $0.05 < \lambda/M_0^2 < 0.99$  and (b)  $1.05 < \lambda/M_0^2 < 7.0$ . By taking  $1/l$  as the mass unit, other parameters are chosen as  $M_0 = 10/l$ ,  $\mu = 10^{-4}/l$  and  $u = 0$ . All of the numbers on the vertical and horizontal axes are dimensionless.

### 3 The Dirac Fermion's two-point Green's function

#### 3.1 Preliminaries

Let us now move to the construction of the Fermion's propagator, whose mass is assumed to be  $z$ -dependent, i.e.,  $m(z)$ . We are going to solve the following equation

$$\left\{ i\gamma^\mu \vec{\partial}_\mu - m(z) \right\} \mathcal{F}(t-t', x-x', y-y', z, z'; m(*)) = i\delta(t-t')\delta(x-x')\delta(y-y')\delta(z-z'), \quad (3.1)$$

$$\mathcal{F}(t-t', x-x', y-y', z, z'; m(*)) \left\{ i\gamma^\mu \overleftarrow{\partial}'_\mu + m(z') \right\} = -i\delta(t-t')\delta(x-x')\delta(y-y')\delta(z-z'). \quad (3.2)$$

The derivatives in (3.1) and (3.2) are given as usual by

$$\partial_\mu = \left( \frac{\partial}{\partial t}, \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right), \quad \partial'_\mu = \left( \frac{\partial}{\partial t'}, \frac{\partial}{\partial x'}, \frac{\partial}{\partial y'}, \frac{\partial}{\partial z'} \right) \quad (3.3)$$

and are acting on the right ( $\vec{\partial}_\mu$ ) and left ( $\overleftarrow{\partial}_\mu$ ) quantities, respectively. As in the Bosonic case, our notation  $m(*)$  instead of  $m(z)$  or  $m(z')$  in (3.1) and (3.2) is to remind us that the two-point function  $\mathcal{F}$  depends on the mass function only implicitly. Note that  $\mathcal{F}$  carries two spinor indices that are, however, suppressed in (3.1) and (3.2). Our propagator is necessarily a generalization of the well-known one

$$S_F(x-x') = i \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot (x-x')} \frac{\gamma^\mu p_\mu + m}{p^2 - m^2 + i\varepsilon}, \quad (3.4)$$

and is supposed to replace (3.4) in the presence of the electroweak bubble wall.

#### 3.2 The canonical form

The ways of handling the dynamics along the  $z$ -axis will differ from those in the  $x$ - and  $y$ -axes and the most convenient choice of the gamma-matrices that we found is rather unconventional, i.e.,

$$\begin{aligned} \gamma^0 &= \begin{pmatrix} \mathbf{0} & \mathbf{1} \\ \mathbf{1} & \mathbf{0} \end{pmatrix}, & \gamma^1 &= \begin{pmatrix} \mathbf{0} & \sigma^2 \\ -\sigma^2 & \mathbf{0} \end{pmatrix}, \\ \gamma^2 &= \begin{pmatrix} \mathbf{0} & -\sigma^1 \\ \sigma^1 & \mathbf{0} \end{pmatrix}, & \gamma^3 &= -i \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & -\mathbf{1} \end{pmatrix}. \end{aligned} \quad (3.5)$$

Here  $\mathbf{1}$  and  $\mathbf{0}$  are the  $2 \times 2$  unit matrix and zero-matrix, respectively. The Pauli matrices are denoted by  $\sigma^i$  ( $i = 1, 2, 3$ ). This set of gamma-matrices is obtained by applying a unitary transformation to the Dirac's representation and is characteristic in that the third Pauli matrix  $\sigma^3$  does not appear in (3.5).

In constructing the Bosonic Green's function in Sec.2, we learned that the spectral problem of the second-order differential operator and in particular the determination of  $d\rho_{kl}(\lambda)/d\lambda$  was our central concern. We have to think about similar ways of formulation

that is suitable for the first-order Dirac operator. Fortunately, spectral problems for a first-order differential equation for a pair of functions have been studied in mathematical literatures for the case of half finite interval, i.e.,  $z \in (0, +\infty)$  in Refs.[84–86] and for infinite open interval, i.e.,  $z \in (-\infty, +\infty)$  in Ref. [86]. Self-contained monographs on this subject are also available [87, 88]. Bearing these mathematical knowledge in our mind, let us disentangle the Dirac equation

$$\{i\gamma \cdot \partial - m(z)\} \psi(z) e^{-iEt+i\vec{p}_\perp \cdot \vec{x}_\perp} = \{\gamma \cdot p - m(z)\} \psi(z) e^{-iEt+i\vec{p}_\perp \cdot \vec{x}_\perp} = 0 ,$$

$$p^\mu = (E, p_x, p_y, -i\frac{\partial}{\partial z}) , \quad (3.6)$$

in a simpler form. Here  $E$  and  $\vec{p}_\perp = (p_x, p_y, 0)$  are just numbers, but the  $z$ -component of the momentum operator gives rise to a non-trivial differential equation for  $\psi(z)$  due to the  $z$ -dependent mass  $m(z)$ . We now put (3.6) into a form that is suitable to the analyses in Ref. [86],

By decomposing  $\psi(z)$  into upper and lower two component spinors,  $\xi_1$  and  $\xi_2$ ,

$$\psi(z) = \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} , \quad (3.7)$$

the Dirac equation (3.6) now becomes

$$(E + \sigma^2 p_x - \sigma^1 p_y) \xi_1 + \left\{ -\frac{d}{dz} - m(z) \right\} \xi_2 = 0 , \quad (3.8)$$

$$\left\{ \frac{d}{dz} - m(z) \right\} \xi_1 + (E - \sigma^2 p_x + \sigma^1 p_y) \xi_2 = 0 . \quad (3.9)$$

Thanks to the use of the gamma-matrix representation as in (3.5), we find that the Pauli matrix appears in (3.8) and (3.9) only in the combination of  $(\sigma^2 p_x - \sigma^1 p_y)$ . This suggest that (3.8) and (3.9) will become much simpler if we use the eigenstates of  $(\sigma^2 p_x - \sigma^1 p_y)$  as the basis of our calculation. The eigenstates of  $(\sigma^2 p_x - \sigma^1 p_y)$  are given by

$$\eta^{(\pm)} = \frac{1}{\sqrt{2}|\vec{p}_\perp|} \begin{pmatrix} \mp(i p_x + p_y) \\ |\vec{p}_\perp| \end{pmatrix} , \quad |\vec{p}_\perp| = \sqrt{p_x^2 + p_y^2} , \quad (3.10)$$

which are shown to satisfy

$$\frac{1}{|\vec{p}_\perp|} (\sigma^2 p_x - \sigma^1 p_y) \eta^{(\pm)} = \pm \eta^{(\pm)} \quad (3.11)$$

together with

$$\eta^{(+)\dagger} \eta^{(+)} = \eta^{(-)\dagger} \eta^{(-)} = 1, \quad \eta^{(+)\dagger} \eta^{(-)} = \eta^{(-)\dagger} \eta^{(+)} = 0 , \quad (3.12)$$

$$\eta^{(+)} \eta^{(+)\dagger} + \eta^{(-)} \eta^{(-)\dagger} = \mathbf{1} , \quad (3.13)$$

$$\eta^{(+)} \eta^{(+)\dagger} - \eta^{(-)} \eta^{(-)\dagger} = \frac{1}{|\vec{p}_\perp|} (\sigma^2 p_x - \sigma^1 p_y) . \quad (3.14)$$



We now eliminate the matrix  $(\sigma^2 p_x - \sigma^1 p_y)$  from (3.8) and (3.9) by setting  $\xi_1$  and  $\xi_2$  in the following way,

$$\xi_1 = f^{(\pm)}(z; E) \eta^{(\pm)}, \quad \xi_2 = g^{(\pm)}(z; E) \eta^{(\pm)}. \quad (3.15)$$

Here our notations,  $f^{(\pm)}(z; E)$  and  $g^{(\pm)}(z; E)$ , are to keep in our mind that these two functions constitute the solution belonging to the energy  $E$ . They are determined by the coupled differential equation

$$(E \pm |\vec{p}_\perp|) f^{(\pm)}(z; E) + \left\{ -\frac{d}{dz} - m(z) \right\} g^{(\pm)}(z; E) = 0, \quad (3.16)$$

$$\left\{ \frac{d}{dz} - m(z) \right\} f^{(\pm)}(z; E) + (E \mp |\vec{p}_\perp|) g^{(\pm)}(z; E) = 0. \quad (3.17)$$

This may be rewritten in the matrix form

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \frac{d}{dz} \begin{pmatrix} f^{(\pm)}(z; E) \\ g^{(\pm)}(z; E) \end{pmatrix} + \begin{pmatrix} \mp |\vec{p}_\perp| & m(z) \\ m(z) & \pm |\vec{p}_\perp| \end{pmatrix} \begin{pmatrix} f^{(\pm)}(z; E) \\ g^{(\pm)}(z; E) \end{pmatrix} = E \begin{pmatrix} f^{(\pm)}(z; E) \\ g^{(\pm)}(z; E) \end{pmatrix}. \quad (3.18)$$

Now we refine Eq. (3.18) further by applying an orthogonal transformation

$$\begin{pmatrix} f^{(\pm)}(z; E) \\ g^{(\pm)}(z; E) \end{pmatrix} = \begin{pmatrix} \cos \theta^{(\pm)}(z) & -\sin \theta^{(\pm)}(z) \\ \sin \theta^{(\pm)}(z) & \cos \theta^{(\pm)}(z) \end{pmatrix} \begin{pmatrix} F^{(\pm)}(z; E) \\ G^{(\pm)}(z; E) \end{pmatrix}. \quad (3.19)$$

Here we set

$$\tan 2\theta^{(\pm)}(z) = \mp \frac{m(z)}{|\vec{p}_\perp|} \quad (3.20)$$

to find Eq. (3.18) transformed into

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \frac{d}{dz} \begin{pmatrix} F^{(\pm)}(z; E) \\ G^{(\pm)}(z; E) \end{pmatrix} + \begin{pmatrix} \mp Q(z) & 0 \\ 0 & \pm Q(z) \end{pmatrix} \begin{pmatrix} F^{(\pm)}(z; E) \\ G^{(\pm)}(z; E) \end{pmatrix} = E \begin{pmatrix} F^{(\pm)}(z; E) \\ G^{(\pm)}(z; E) \end{pmatrix}, \quad (3.21)$$

with

$$Q(z) = \sqrt{m(z)^2 + |\vec{p}_\perp|^2} + \frac{m'(z)|\vec{p}_\perp|}{2\{m(z)^2 + |\vec{p}_\perp|^2\}}. \quad (3.22)$$

This simplified form of the Dirac equation is called ‘‘canonical’’ in Ref. [88], where the spectra of the energy  $E$  were fully analyzed for general ‘‘potential’’  $Q(z)$ . This canonical

form (3.21) is useful to look into the symmetry property of the  $E$  spectra. In fact we can rewrite (3.21) as

$$\begin{aligned} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \frac{d}{dz} \begin{pmatrix} G^{(\pm)}(z; E) \\ F^{(\pm)}(z; E) \end{pmatrix} + \begin{pmatrix} \mp Q(z) & 0 \\ 0 & \pm Q(z) \end{pmatrix} \begin{pmatrix} G^{(\pm)}(z; E) \\ F^{(\pm)}(z; E) \end{pmatrix} \\ = -E \begin{pmatrix} G^{(\pm)}(z; \lambda) \\ F^{(\pm)}(z; \lambda) \end{pmatrix}, \end{aligned} \quad (3.23)$$

and we note the sign change,  $E \rightarrow -E$ , on the right hand side of (3.23). In other words, the coupled equation (3.23) for a pair of functions  $F^{(\pm)}(z; \lambda)$  and  $G^{(\pm)}(z; \lambda)$  is form-invariant under the following replacement

$$E \rightarrow -E, \quad \begin{pmatrix} F^{(\pm)}(z; E) \\ G^{(\pm)}(z; E) \end{pmatrix} \rightarrow \begin{pmatrix} G^{(\pm)}(z; E) \\ F^{(\pm)}(z; E) \end{pmatrix}. \quad (3.24)$$

This is the symmetry between positive and negative energy spectra in the usual sense.

### 3.3 The transverse momentum dependence of the energy $E$

The energy  $E$  that appeared in (3.21) is inter-connected with the magnitude of the transverse momentum  $|\vec{p}_\perp|$ . This connection can be made more explicit in the following way. Let us now deduce second order differential equations for  $f^{(\pm)}(z; E)$  and  $g^{(\pm)}(z; E)$  from (3.16) and (3.17), which turn out to be the following one-dimensional Schrödinger type equations,

$$\left\{ -\frac{d^2}{dz^2} + U_f(z) \right\} f^{(\pm)}(z; E) = \nu_f f^{(\pm)}(z; E), \quad (3.25)$$

$$\left\{ -\frac{d^2}{dz^2} + U_g(z) \right\} g^{(\pm)}(z; E) = \nu_g g^{(\pm)}(z; E). \quad (3.26)$$

Here the ‘‘potential’’ terms are given by

$$U_f(z) = m(z)^2 + \frac{dm(z)}{dz}, \quad U_g(z) = m(z)^2 - \frac{dm(z)}{dz}, \quad (3.27)$$

respectively and the ‘‘energy’’  $\nu_f$  and  $\nu_g$ , of these potential problems are connected with  $E$  by a common relation

$$\nu_f = E^2 - |\vec{p}_\perp|^2, \quad \nu_g = E^2 - |\vec{p}_\perp|^2. \quad (3.28)$$

In general the allowed region of the spectra of  $\{\nu_f\}$  and  $\{\nu_g\}$  do not necessarily coincide with each other since the potentials are different. Only when they share common region of the spectra i.e.,  $\nu_f = \nu_g = \nu$ , we are able to obtain the energy  $E$  of the coupled equations (3.16) and (3.17) by the formula

$$E = \pm \sqrt{\nu + |\vec{p}_\perp|^2}. \quad (3.29)$$

Here it should be stressed that the spectra of  $\{\nu_f\}$  and  $\{\nu_g\}$  are independent of  $|\vec{p}_\perp|$  since the potentials,  $U_f(z)$  and  $U_g(z)$ , do not contain  $|\vec{p}_\perp|$ . The energy  $E$  in (3.16), (3.17) and (3.21) depend on  $|\vec{p}_\perp|$  only through the restricted form of (3.29) irrespectively of the details of the form of  $m(z)$ . This observation could be useful when we look at infrared singularities associated with loop diagrams, since singularities always come from limited corners in the phase space, i.e.,  $|\vec{p}_\perp|^2 \approx 0$

### 3.4 The general expansion method and completeness

Let us now discuss an expansion of a pair of arbitrary real-valued functions  $(A(z) B(z))$  in terms of the set of solutions  $(F^{(\pm)}(z; E) G^{(\pm)}(z; E))$ . We assume that  $A(z)$  and  $B(z)$  are both *square-integrable* in  $(-\infty, +\infty)$ . We introduce two types of solutions to Eq. (3.21)  $(F_1^{(\pm)}(z; E) G_1^{(\pm)}(z; E))$  and  $(F_2^{(\pm)}(z; E) G_2^{(\pm)}(z; E))$ , satisfying the conditions at  $z = 0$ ,

$$F_1^{(\pm)}(0; E) = 1, \quad G_1^{(\pm)}(0; E) = 0, \quad (3.30)$$

and

$$F_2^{(\pm)}(0; E) = 0, \quad G_2^{(\pm)}(0; E) = 1, \quad (3.31)$$

analogously to Eqs. (2.6) and (2.7) in the Bosonic case. It has been argued in [86] that a pair of real-valued functions  $(A(z) B(z))$  can be expanded as a superposition of (3.30) and (3.31) in the following form

$$\begin{aligned} \begin{pmatrix} A(z) \\ B(z) \end{pmatrix} &= \sum_{k,l=1,2} \int \begin{pmatrix} F_k^{(\pm)}(z; \Lambda) \\ G_k^{(\pm)}(z; \Lambda) \end{pmatrix} d\rho_{kl}^{(\pm)}(\Lambda) \\ &\quad \times \int_{-\infty}^{+\infty} dz' \left( F_l^{(\pm)}(z'; \Lambda) G_l^{(\pm)}(z'; \Lambda) \right) \begin{pmatrix} A(z') \\ B(z') \end{pmatrix}. \end{aligned} \quad (3.32)$$

Here  $d\rho_{kl}^{(\pm)}(\Lambda)$  describes the distribution of  $\Lambda$  in connection with the differential equation (3.21). We use the letter  $\Lambda$  instead of  $E$  for the integration variable in (3.32) in order to avoid possible confusion that could perhaps arise later in (3.48). According to Ref. [86], the spectral functions are given by

$$d\rho_{11}^{(\pm)}(\Lambda) = \lim_{\varepsilon \rightarrow +0} \frac{1}{\pi} \Im \left( \frac{1}{m_2^{(\pm)}(\Lambda + i\varepsilon) - m_1^{(\pm)}(\Lambda + i\varepsilon)} \right) d\Lambda, \quad (3.33)$$

$$\begin{aligned} d\rho_{12}^{(\pm)}(\Lambda) &= d\rho_{21}^{(\pm)}(\Lambda) \\ &= \lim_{\varepsilon \rightarrow +0} \frac{1}{2\pi} \Im \left( \frac{m_2^{(\pm)}(\Lambda + i\varepsilon) + m_1^{(\pm)}(\Lambda + i\varepsilon)}{m_2^{(\pm)}(\Lambda + i\varepsilon) - m_1^{(\pm)}(\Lambda + i\varepsilon)} \right) d\Lambda, \end{aligned} \quad (3.34)$$

$$d\rho_{22}^{(\pm)}(\Lambda) = \lim_{\varepsilon \rightarrow +0} \frac{1}{\pi} \Im \left( \frac{m_1^{(\pm)}(\Lambda + i\varepsilon) m_2^{(\pm)}(\Lambda + i\varepsilon)}{m_2^{(\pm)}(\Lambda + i\varepsilon) - m_1^{(\pm)}(\Lambda + i\varepsilon)} \right) d\Lambda, \quad (3.35)$$

for  $\Lambda > 0$ , in case of continuous spectra. (See Eq. (6.2) in Ref. [86].) The characteristic functions  $m_1^{(\pm)}(\Lambda + i\varepsilon)$  and  $m_2^{(\pm)}(\Lambda + i\varepsilon)$  are given by

$$\begin{aligned} m_1^{(\pm)}(\Lambda + i\varepsilon) &= - \lim_{a \rightarrow -\infty} \frac{F_1^{(\pm)}(a; \Lambda + i\varepsilon)}{F_2^{(\pm)}(a; \Lambda + i\varepsilon)}, \\ m_2^{(\pm)}(\Lambda + i\varepsilon) &= - \lim_{b \rightarrow +\infty} \frac{F_1^{(\pm)}(b; \Lambda + i\varepsilon)}{F_2^{(\pm)}(b; \Lambda + i\varepsilon)}, \end{aligned} \quad (3.36)$$

in a similar way as in the Bosonic case. For  $\Lambda < 0$ , the spectral functions are determined by the symmetry property between positive and negative energy solutions mentioned before.

The expansion (3.32) may be equivalently expressed by concise formulas

$$\sum_{k,l=1,2} \int F_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) F_l^{(\pm)}(z'; \Lambda) = \delta(z - z'), \quad (3.37)$$

$$\sum_{k,l=1,2} \int F_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) G_l^{(\pm)}(z'; \Lambda) = 0, \quad (3.38)$$

$$\sum_{k,l=1,2} \int G_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) F_l^{(\pm)}(z'; \Lambda) = 0, \quad (3.39)$$

$$\sum_{k,l=1,2} \int G_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) G_l^{(\pm)}(z'; \Lambda) = \delta(z - z'). \quad (3.40)$$

These formulas are further rephrased, through the orthogonal rotation like (3.19),

$$\begin{pmatrix} f_k^{(\pm)}(z; \Lambda) \\ g_k^{(\pm)}(z; \Lambda) \end{pmatrix} = \begin{pmatrix} \cos \theta^{(\pm)}(z) & -\sin \theta^{(\pm)}(z) \\ \sin \theta^{(\pm)}(z) & \cos \theta^{(\pm)}(z) \end{pmatrix} \begin{pmatrix} F_k^{(\pm)}(z; \Lambda) \\ G_k^{(\pm)}(z; \Lambda) \end{pmatrix}, \quad (k = 1, 2), \quad (3.41)$$

by the formulas

$$\sum_{k,l=1,2} \int f_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) f_l^{(\pm)}(z'; \Lambda) = \delta(z - z'), \quad (3.42)$$

$$\sum_{k,l=1,2} \int f_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) g_l^{(\pm)}(z'; \Lambda) = 0, \quad (3.43)$$

$$\sum_{k,l=1,2} \int g_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) f_l^{(\pm)}(z'; \Lambda) = 0, \quad (3.44)$$

$$\sum_{k,l=1,2} \int g_k^{(\pm)}(z; \Lambda) d\rho_{kl}^{(\pm)}(\Lambda) g_l^{(\pm)}(z'; \Lambda) = \delta(z - z'). \quad (3.45)$$

### 3.5 Construction of the Propagator

We are now in a position to construct the Fermion's two-point Green's function. Let us write down the solutions to the Dirac equation (3.6) as

$$\psi_k^{(\pm)}(z; \Lambda) \equiv \begin{pmatrix} f_k^{(\pm)}(z; \Lambda) \eta^{(\pm)} \\ g_k^{(\pm)}(z; \Lambda) \eta^{(\pm)} \end{pmatrix} \quad (k = 1, 2) \quad (3.46)$$

and pay attention to the following formula

$$\begin{aligned} & \sum_{k,l=1,2} \sum_{s=+,-} \int \psi_k^{(s)}(z; \Lambda) d\rho_{kl}^{(s)}(\Lambda) \psi_l^{(s)}(z'; \Lambda)^\dagger \\ &= \begin{pmatrix} \eta^{(+)} \eta^{(+)\dagger} + \eta^{(-)} \eta^{(-)\dagger} & 0 \\ 0 & \eta^{(+)} \eta^{(+)\dagger} + \eta^{(-)} \eta^{(-)\dagger} \end{pmatrix} \delta(z - z') \\ &= \begin{pmatrix} \mathbf{1} & 0 \\ 0 & \mathbf{1} \end{pmatrix} \delta(z - z'). \end{aligned} \quad (3.47)$$

This formula can be derived by using (3.42) - (3.45) together with (3.13). Noting that  $\psi_k^{(\pm)}(z; \Lambda)$  satisfies the differential equation

$$\gamma^0 \{i\gamma \cdot \partial - m(z)\} \psi_k^{(\pm)}(z; \Lambda) e^{-iEt + i\vec{p}_\perp \cdot \vec{x}_\perp} = (E - \Lambda) \psi_k^{(\pm)}(z; \Lambda) e^{-iEt + i\vec{p}_\perp \cdot \vec{x}_\perp}, \quad (3.48)$$

we also find the formula

$$\begin{aligned} \gamma^0 \{i\gamma \cdot \partial - m(z)\} \sum_{k,l=1,2} \sum_{s=+,-} \int \psi_k^{(s)}(z; \Lambda) \frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda} \psi_l^{(s)}(z'; \Lambda)^\dagger e^{-iE(t-t') + i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)} \\ = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix} \delta(z - z') e^{-iE(t-t') + i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)}. \end{aligned} \quad (3.49)$$

By multiplying  $\gamma^0$  from the left and another  $\gamma^0$  from the right, we get

$$\begin{aligned} \{i\gamma \cdot \partial - m(z)\} \int \frac{dE d^2\vec{p}_\perp}{(2\pi)^3} e^{-iE(t-t') + i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)} \\ \times \sum_{k,l=1,2} \sum_{s=+,-} \int \psi_k^{(s)}(z; \Lambda) \frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda} \psi_l^{(s)}(z'; \Lambda)^\dagger \gamma^0 \\ = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix} \delta(z - z') \delta(t - t') \delta^2(\vec{x}_\perp - \vec{x}'_\perp). \end{aligned} \quad (3.50)$$

We have thus arrived finally at the integral representation for the Fermion's propagator

$$\begin{aligned} \mathcal{F}(t - t', x - x', y - y', z, z'; m(*)) \\ = i \int \frac{dE d^2\vec{p}_\perp}{(2\pi)^3} e^{-iE(t-t')} e^{i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)} \sum_{k,l=1,2} \sum_{s=+,-} \int \psi_k^{(s)}(z; \Lambda) \frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda} \psi_l^{(s)}(z'; \Lambda)^\dagger \gamma^0. \end{aligned} \quad (3.51)$$

In this expression we have not yet specified the  $+i\varepsilon$  prescription in the energy denominator. In order for the above Green's function to be the time-ordered vacuum expectation value of the Fermion's bilinear operators we must go into the detail of the definition of the vacuum in the presence of the bubble wall. We will come to this point later in Sect. 5, and for now suffice it to say that the singularity at  $E = \Lambda$  in (3.51) should be avoided by setting

$$\frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda} \rightarrow \frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda + i\varepsilon} \quad \text{for } \Lambda > 0, \quad (3.52)$$

and

$$\frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda} \rightarrow \frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda - i\varepsilon} \quad \text{for } \Lambda < 0. \quad (3.53)$$

### 3.6 The constant mass case (no bubble wall)

Let us put our formula (3.51) into an explicit one for the case that the mass is just a constant, i.e.,  $m(z) = m$ . In this case  $Q(z)$  in (3.22) is just a constant  $\sqrt{m^2 + |\vec{p}_\perp|^2}$  and

it is a simple matter to get the solutions to (3.21). For  $\Lambda^2 > (m^2 + |\vec{p}_\perp|^2)$ , the solutions become

$$F_1^{(\pm)}(z; \Lambda) = \cos(qz), \quad G_1^{(\pm)}(z; \Lambda) = \frac{\Lambda \pm \sqrt{m^2 + |\vec{p}_\perp|^2}}{q} \sin(qz), \quad (3.54)$$

$$F_2^{(\pm)}(z; \Lambda) = -\frac{\Lambda \mp \sqrt{m^2 + |\vec{p}_\perp|^2}}{q} \sin(qz), \quad G_2^{(\pm)}(z; \Lambda) = \cos(qz), \quad (3.55)$$

where we defined  $q$  as follows

$$q = \sqrt{\Lambda^2 - (m^2 + |\vec{p}_\perp|^2)}. \quad (3.56)$$

For  $\Lambda^2 < (m^2 + |\vec{p}_\perp|^2)$ , on the other hand, the solutions are

$$F_1^{(\pm)}(z; \Lambda) = \cosh(\tilde{q}z), \quad G_1^{(\pm)}(z; \Lambda) = \frac{\Lambda \pm \sqrt{m^2 + |\vec{p}_\perp|^2}}{\tilde{q}} \sinh(\tilde{q}z), \quad (3.57)$$

$$F_2^{(\pm)}(z; \Lambda) = -\frac{\Lambda \mp \sqrt{m^2 + |\vec{p}_\perp|^2}}{\tilde{q}} \sinh(\tilde{q}z), \quad G_2^{(\pm)}(z; \Lambda) = \cosh(\tilde{q}z). \quad (3.58)$$

Here  $\tilde{q}$  is defined by

$$\tilde{q} = \sqrt{(m^2 + |\vec{p}_\perp|^2) - \Lambda^2}. \quad (3.59)$$

The spectral functions (3.33) - (3.35), are also explicitly computed by using the formula (3.36) as follows,

$$d\rho_{11}^{(\pm)}(\Lambda) = \begin{cases} \frac{1}{2\pi} \frac{\Lambda \mp \sqrt{m^2 + |\vec{p}_\perp|^2}}{q} d\Lambda & \text{for } \Lambda > \sqrt{m^2 + |\vec{p}_\perp|^2}, \\ 0 & \text{for } \Lambda^2 < (m^2 + |\vec{p}_\perp|^2), \\ \frac{1}{2\pi} \frac{-\Lambda \pm \sqrt{m^2 + |\vec{p}_\perp|^2}}{q} d\Lambda & \text{for } \Lambda < -\sqrt{m^2 + |\vec{p}_\perp|^2}, \end{cases} \quad (3.60)$$

$$d\rho_{12}^{(\pm)}(\Lambda) = d\rho_{21}^{(\pm)}(\Lambda) = 0, \quad (3.61)$$

$$d\rho_{22}^{(\pm)}(\Lambda) = \begin{cases} \frac{1}{2\pi} \frac{\Lambda \pm \sqrt{m^2 + |\vec{p}_\perp|^2}}{q} d\Lambda & \text{for } \Lambda > \sqrt{m^2 + |\vec{p}_\perp|^2}, \\ 0 & \text{for } \Lambda^2 < (m^2 + |\vec{p}_\perp|^2), \\ \frac{1}{2\pi} \frac{-\Lambda \mp \sqrt{m^2 + |\vec{p}_\perp|^2}}{q} d\Lambda & \text{for } \Lambda < -\sqrt{m^2 + |\vec{p}_\perp|^2}. \end{cases} \quad (3.62)$$

Here we notice a simple relation  $d\rho_{11}^{(\pm)}(-\Lambda)/d(-\Lambda) = d\rho_{22}^{(\pm)}(\Lambda)/d\Lambda$  which is due to the positive- and negative energy correspondence. With these spectral functions we can confirm the completeness relations (3.37) - (3.40) explicitly.

In order to put the formula (3.51) into a more familiar form, we have to use the solutions (3.54) and (3.55) together with the spectral functions to get the integrations

$$\begin{aligned} & \sum_{k,l=1,2} \int F_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} F_l^{(\pm)}(z'; \Lambda) \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{E \mp \sqrt{m^2 + |\vec{p}_\perp|^2}}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}, \end{aligned} \quad (3.63)$$

$$\begin{aligned}
& \sum_{k,l=1,2} \int F_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} G_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{iq}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}, \quad (3.64)
\end{aligned}$$

$$\begin{aligned}
& \sum_{k,l=1,2} \int G_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} F_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{-iq}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}, \quad (3.65)
\end{aligned}$$

$$\begin{aligned}
& \sum_{k,l=1,2} \int G_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} G_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{E \pm \sqrt{m^2 + |\vec{p}_\perp|^2}}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}. \quad (3.66)
\end{aligned}$$

According to the original definition (3.56),  $q$  is positive, but the integration region of  $q$  in the above formulas is extended to  $(-\infty, +\infty)$  by using symmetry arguments. These formulas may be transformed via orthogonal transformation (3.41) into the formulas involving  $f_k^{(\pm)}(z; \Lambda)$  and  $g_k^{(\pm)}(z; \Lambda)$ ,

$$\begin{aligned}
& \sum_{k,l=1,2} \int f_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} f_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{E \mp |\vec{p}_\perp|}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}, \quad (3.67)
\end{aligned}$$

$$\begin{aligned}
& \sum_{k,l=1,2} \int f_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} g_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{m + iq}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}, \quad (3.68)
\end{aligned}$$

$$\begin{aligned}
& \sum_{k,l=1,2} \int g_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} f_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{m - iq}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}, \quad (3.69)
\end{aligned}$$

$$\begin{aligned}
& \sum_{k,l=1,2} \int g_k^{(\pm)}(z; \Lambda) \frac{d\rho_{kl}^{(\pm)}(\Lambda)}{E - \Lambda} g_l^{(\pm)}(z'; \Lambda) \\
&= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{E \pm |\vec{p}_\perp|}{E^2 - (m^2 + |\vec{p}_\perp|^2 + q^2) + i\varepsilon}. \quad (3.70)
\end{aligned}$$

Combining all these formulas, we end up with

$$\begin{aligned} & \sum_{k,l=1,2} \sum_{s=+,-} \int \psi_k^{(s)}(z; \lambda) \frac{d\rho_{kl}^{(s)}(\Lambda)}{E - \Lambda} \psi_l^{(s)}(z'; \lambda)^\dagger \gamma^0 \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dq e^{iq(z-z')} \frac{E\gamma^0 - \gamma^1 p_x - \gamma^2 p_y - q\gamma^3 + m}{E^2 - |\vec{p}_\perp|^2 - q^2 - m^2 + i\varepsilon}. \end{aligned} \quad (3.71)$$

Here use has been made of the identities (3.13) and (3.14). Note, in particular, that the momenta  $p_x$  and  $p_y$  in the numerator of (3.71) show up through the use of the formula (3.14). Putting the above relation into (3.51), we finally arrive at the familiar propagator (3.4) with  $p^\mu = (E, p_x, p_y, q)$ . We have thus successfully reproduced the well-known formula with the aid of less-known method using the spectral function  $\rho_{kl}(\Lambda)$ . In the presence of the bubble wall, it seems absolutely necessary to make a full use of the spectral function to construct Green's functions.

## 4 The vector boson's two-point Green's function

### 4.1 Mixing between gauge boson and unphysical scalar fields

Now let us turn to the Green's functions of the gauge bosons, in which we encounter with some peculiar phenomena in contrast to the scalar case. The polarization vectors of the vector bosons are usually defined referring to the four-momentum, whose  $z$ -component in the present case, however, is changing during the passage of the bubble wall. The separation into transverse and longitudinal polarization states becomes therefore more involved than usual. Moreover the unphysical scalar fields are also mixed up with the vector bosons through the variation of the Higgs vacuum expectation value. Farrar and McIntosh [83] once discussed the change of the polarization states that vector bosons meet with in the course of passing through the bubble wall. Azatov et al also discussed in their work [73] the alteration of the polarization vector by choosing the unitary gauge. We will go along similar lines as theirs by making use of the 't Hooft-Feynman gauge which is more easy to deal with to construct our propagators.

Let us consider specifically the standard  $SU(2)_L \times U(1)_Y$  electroweak theory in order to see the mixing between the unphysical scalar and gauge boson fields. The Higgs Lagrangian is given as usual by

$$\mathcal{L}_{\text{scalar}} = (\nabla_\mu \Phi)^\dagger (\nabla^\mu \Phi) - V(\Phi), \quad V(\Phi) = \lambda(\Phi^\dagger \Phi)^2 + \mu_0^2 \Phi^\dagger \Phi, \quad (4.1)$$

where  $\nabla_\mu$  is the gauge covariant derivative and the Higgs doublet field  $\Phi$  is parametrized as

$$\Phi = \begin{pmatrix} w^\dagger \\ \frac{1}{\sqrt{2}}(v + H + i\chi) \end{pmatrix}. \quad (4.2)$$

The physical Higgs field is denoted by  $H$  and  $w$ ,  $w^\dagger$  and  $\chi$  are the unphysical scalar fields. In contrast with the usual case, the vacuum expectation value  $v$  is not a constant but is



supposed to be position dependent, i.e.,  $\partial_\mu v \neq 0$ . Putting (4.2) into (4.1) and keeping terms up to the quadratic ones, we get

$$\begin{aligned}
\mathcal{L}_{\text{scalar}} = & \partial_\mu w^\dagger \partial^\mu w + \frac{1}{2} \partial_\mu (v + H - i\chi) \partial^\mu (v + H + i\chi) \\
& - M_Z Z^\mu \left( \partial_\mu \chi - \chi \frac{\partial_\mu v}{v} \right) \\
& + i M_W W^{+\mu} \left( \partial_\mu w - w \frac{\partial_\mu v}{v} \right) - i M_W W^{-\mu} \left( \partial_\mu w^\dagger - w^\dagger \frac{\partial_\mu v}{v} \right) \\
& + M_W^2 W_\mu^+ W^{-\mu} + \frac{1}{2} M_Z^2 Z_\mu Z^\mu \\
& + (\lambda v^2 + \mu_0^2) \left( w^\dagger w + \frac{1}{2} \chi^2 \right) + \left( \frac{3}{2} \lambda v^2 + \frac{1}{2} \mu_0^2 \right) H^2 + (\lambda v^3 + v \mu_0^2) H \\
& + \dots, \tag{4.3}
\end{aligned}$$

where ellipses denote cubic and quartic terms with respect to fields. Note that the  $W$  and  $Z$  boson masses

$$M_W^2 = \frac{1}{4} g^2 v^2, \quad M_Z^2 = \frac{1}{4} (g^2 + g'^2) v^2 \tag{4.4}$$

are also position dependent. Looking at (4.3), we immediately notice that there are extra terms

$$\left( M_Z Z^\mu \chi - i M_W W^{+\mu} w + i M_W W^{-\mu} w^\dagger \right) \frac{\partial_\mu v}{v} \tag{4.5}$$

for  $\partial_\mu v \neq 0$  in comparison with the conventional case. This describes the mixing between gauge bosons and associated unphysical scalar fields.

In the following we would like to employ the 't Hooft-Feynman ( $\xi = 1$  in  $R_\xi$ ) gauge and add the following gauge fixing (GF) Lagrangian

$$\mathcal{L}_{\text{GF}} = -(\partial \cdot W^+ + i M_W w^\dagger)(\partial \cdot W^- - i M_W w) - \frac{1}{2} (\partial \cdot Z + M_Z \chi)^2 - \frac{1}{2} (\partial \cdot A)^2. \tag{4.6}$$

(The gauge fixing term for the electromagnetic field  $A_\mu$  is included in (4.6) for completeness, but will be irrelevant to the subsequent analyses.) The  $R_\xi$  gauge is usually arranged to eliminate the mixing terms between gauge boson and associated unphysical scalar fields. In the present case, however, the mixing does not disappear if  $\partial_\mu v \neq 0$ . In fact, in the partial integration in (4.6) we use the relations

$$\partial_\mu M_W = M_W \frac{\partial_\mu v}{v}, \quad \partial_\mu M_Z = M_Z \frac{\partial_\mu v}{v}, \tag{4.7}$$

and we notice that the mixing terms in (4.5) are not cancelled but are doubled. The free equations of motion of the  $W$ - and  $Z$ -bosons and associated unphysical scalars turn out to

be

$$(\square + M_Z^2) Z_\mu + 2M_Z \frac{\partial_\mu v}{v} \chi = 0, \quad (4.8)$$

$$(\square + M_W^2) W_\mu^- - 2iM_W \frac{\partial_\mu v}{v} w = 0, \quad (\square + M_W^2) W_\mu^+ + 2iM_W \frac{\partial_\mu v}{v} w^\dagger = 0, \quad (4.9)$$

$$(\square + M_1^2) \chi - 2M_Z \frac{\partial_\mu v}{v} Z^\mu = 0, \quad (4.10)$$

$$(\square + M_2^2) w - 2iM_W \frac{\partial_\mu v}{v} W^{-\mu} = 0, \quad (\square + M_2^2) w^\dagger + 2iM_W \frac{\partial_\mu v}{v} W^{+\mu} = 0, \quad (4.11)$$

and show mixing apparently, where use has been made of the notations

$$M_1^2 = M_Z^2 + (\lambda v^2 + \mu_0^2), \quad M_2^2 = M_W^2 + (\lambda v^2 + \mu_0^2). \quad (4.12)$$

Note that the term  $\lambda v^2 + \mu_0^2$  in (4.12) goes to zero in the symmetry-broken region, but we have to keep it in general.

The mixing is, however, rather limited, if  $v$  depends on the  $z$ -coordinate only, i.e.,  $v = v(z)$ . In fact the components of vector fields with  $\mu = 0, 1, 2$  are free from mixing

$$(\square + M_Z^2) Z^i = 0, \quad (\square + M_W^2) W^{\pm i} = 0, \quad (i = 0, 1, 2), \quad (4.13)$$

and the two-point Green's functions  $\langle\langle T(Z^i Z^j) \rangle\rangle$  and  $\langle\langle T(W^{\pm i} W^{\mp j}) \rangle\rangle$  ( $i, j = 0, 1, 2$ ) are easily worked out in the same way as in the scalar field case described in Sect. 2. The mixing occurs only in the third components,  $Z^3$  and  $W^{\pm 3}$ , with their respective unphysical scalar fields. The equations of motion for the pair  $(Z^3 \ \chi)$  become

$$(\square + M_Z^2) Z^3 - 2M_Z \frac{v'(z)}{v(z)} z = 0, \quad (\square + M_1^2) \chi - 2M_Z \frac{v'(z)}{v(z)} Z^3 = 0, \quad (4.14)$$

and similar equations can be written down for the pairs,  $(W^{-3} \ w)$  and  $(W^{+3} \ w^\dagger)$  as follows

$$(\square + M_W^2) W^{-3} + 2iM_W \frac{v'(z)}{v(z)} w = 0, \quad (\square + M_2^2) w - 2iM_W \frac{v'(z)}{v(z)} W^{-3} = 0, \quad (4.15)$$

$$(\square + M_W^2) W^{+3} - 2iM_W \frac{v'(z)}{v(z)} w^\dagger = 0, \quad (\square + M_2^2) w^\dagger + 2iM_W \frac{v'(z)}{v(z)} W^{+3} = 0. \quad (4.16)$$

## 4.2 The general expansion method for two-component fields

Since the third component of the gauge boson is mixed with the associated unphysical scalar boson, the two-point function that we are seeking for becomes necessarily a  $2 \times 2$  matrix and let us explain what the matrix looks like for the pair  $(Z^3 \ \chi)$ . (The case of the  $W$  boson and  $w$  can be handled exactly in the same way.)

Looking at Eq. (4.14), we notice that the  $2 \times 2$  Green's function matrix has to satisfy the following equation,

$$\begin{aligned} & \begin{pmatrix} \square + M_Z^2 & -2M_Z \frac{v'(z)}{v(z)} \\ -2M_Z \frac{v'(z)}{v(z)} & \square + M_1^2 \end{pmatrix} \begin{pmatrix} \mathcal{G}_{11} & \mathcal{G}_{12} \\ \mathcal{G}_{21} & \mathcal{G}_{22} \end{pmatrix} \\ & = -i \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \delta(t - t') \delta(x - x') \delta(y - y') \delta(z - z'). \end{aligned} \quad (4.17)$$

The translation invariance in time and the transverse  $\vec{x}_\perp$ -directions is respected and each component of the matrix is a function of the following variables,

$$\mathcal{G}_{\alpha\beta} = \mathcal{G}_{\alpha\beta}(t - t', x - x', y - y', z, z'; v(*)), \quad (\alpha, \beta = 1, 2), \quad (4.18)$$

and is expressed as a superposition of the plane wave

$$e^{-iE(t-t')} e^{i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)}. \quad (4.19)$$

The d'Alembertian is therefore going to be replaced as

$$\square \rightarrow -E^2 + \vec{p}_\perp^2 - \frac{\partial^2}{\partial z^2} \quad (4.20)$$

and we are naturally led to consider the differential equation of the following type

$$\left\{ -\frac{d^2}{dz^2} + V(z) \right\} \phi(z; \lambda) = \lambda \phi(z; \lambda). \quad (4.21)$$

This is again the Schrödinger-type equation with the “potential”  $V(z)$ , which is a  $2 \times 2$  matrix defined by

$$V(z) = \begin{pmatrix} M_Z^2 & -2M_Z \frac{v'(z)}{v(z)} \\ -2M_Z \frac{v'(z)}{v(z)} & M_1^2 \end{pmatrix}, \quad (4.22)$$

The “energy” is denoted by  $\lambda$  in (4.21), and  $\phi(z; \lambda)$  is the corresponding wave function with two components

$$\phi(z; \lambda) = \begin{pmatrix} \phi_1(z; \lambda) \\ \phi_2(z; \lambda) \end{pmatrix}. \quad (4.23)$$

Note that the matrix (4.22) is real-symmetric, and the operator on the left hand side of (4.21) is self-adjoint in the space of functions with favorable behaviors at  $z \rightarrow \pm\infty$ . The

differential equation (4.21) can be rewritten equivalently in the following form,

$$\begin{pmatrix} \square + M_Z^2 & -2M_Z \frac{v'(z)}{v(z)} \\ -2M_Z \frac{v'(z)}{v(z)} & \square + M_1^2 \end{pmatrix} \phi(z; \lambda) e^{-iEt} e^{i\vec{p}_\perp \cdot \vec{x}_\perp} \\ = (-E^2 + \vec{p}_\perp^2 + \lambda) \phi(z; \lambda) e^{-iEt} e^{i\vec{p}_\perp \cdot \vec{x}_\perp}. \quad (4.24)$$

At this point let us recall that Kodaira, in his follow-up paper [78], extended the work done in [74], [75], [76] and [77] on the general expansion method associated with the second-order differential operator to the case of any even-order self-adjoint differential operator. As remarked almost at the end of Ref. [78], the expansion method developed there can be carried to the case of coupled differential equation of lower-order for multi-component fields and our present problem belongs to such a category of the two-component case.

We can proceed similarly as in Sect. 2 and Sect. 3 by preparing fundamental solutions to (4.21), which we denote  $\phi^{(k)}(z; \lambda)$  ( $k = 1, \dots, 4$ ). They are defined by the initial conditions at  $z = 0$ . All the solutions to the Schrödinger equation (4.21) are expressed as a linear combination of  $\phi^{(k)}(z; \lambda)$ . It has been shown that there exists the spectral function,  $\rho_{kl}(\lambda)$  with such a property that any pair of square-integrable functions  $u_\alpha(z)$  ( $\alpha = 1, 2$ ), can be expressed as a superposition of the fundamental solutions

$$u_\alpha(z) = \int \sum_{k,l} \phi_\alpha^{(k)}(z; \lambda) d\rho_{kl}(\lambda) \int_{-\infty}^{+\infty} dz' \sum_{\beta=1,2} \phi_\beta^{(l)}(z'; \lambda) u_\beta(z'). \quad (4.25)$$

This indicates that we may rewrite this formula in a simple form as

$$\int \sum_{k,l} \phi_\alpha^{(k)}(z; \lambda) d\rho_{kl}(\lambda) \phi_\beta^{(l)}(z'; \lambda) = \delta_{\alpha\beta} \delta(z - z'). \quad (4.26)$$

With the help of (4.26), it is almost straightforward to write down the Green's function in the following integral form

$$\mathcal{G}_{\alpha\beta}(t - t', x - x', y - y', z, z'; v(*)) \quad (4.27)$$

$$= i \int \frac{dE d^2\vec{p}_\perp}{(2\pi)^3} e^{-iE(t-t')} e^{i\vec{p}_\perp \cdot (\vec{x}_\perp - \vec{x}'_\perp)} \int \sum_{k,l} \frac{\phi_\alpha^{(k)}(z; \lambda) d\rho_{kl}(\lambda) \phi_\beta^{(l)}(z'; \lambda)}{E^2 - \vec{p}_\perp^2 - \lambda + i\varepsilon}. \quad (4.28)$$

We have added  $+i\varepsilon$  in the denominator in (4.28) just by hand so that this should be the vacuum expectation value of the time-ordered product of fields in the sense to be explained next.

## 5 Field quantization in the presence of the bubble wall

In constructing the two-point functions, we made use of various propagation modes of scalar and vector fields labelled by  $\lambda$ , and of spinor fields labelled by  $\Lambda$ . We are now well-prepared

to discuss how to quantize these fields by taking the existence of the bubble wall into account. For the sake of simplicity we consider the case in which the spectral functions admit only continuous spectra. We will discuss below the quantization of scalar and spinor fields, while the vector fields will be dealt with in our separate publication since the physical state conditions are complicated and we have to prepare other tools to explain them.

### 5.1 Real scalar field

Let us begin with the scalar field satisfying the free equations of motion

$$\{\square + M(z)^2\} \varphi(t, \vec{x}_\perp, z) = 0, \quad (5.1)$$

in the presence of the bubble wall. Our task now is to realize the canonical commutation relations

$$[\dot{\varphi}(t, \vec{x}_\perp, z), \varphi(t, \vec{x}'_\perp, z')] = -i \delta^2(\vec{x}_\perp - \vec{x}'_\perp) \delta(z - z'), \quad (5.2)$$

$$[\varphi(t, \vec{x}_\perp, z), \varphi(t, \vec{x}'_\perp, z')] = [\dot{\varphi}(t, \vec{x}_\perp, z), \dot{\varphi}(t, \vec{x}'_\perp, z')] = 0, \quad (5.3)$$

by introducing commutation relations among some operators which are analogous to the conventional creation and annihilation operators. Note that the time derivative is denoted by dot in (5.2) and (5.3).

The solutions to the equations of motion (5.1) are given in the form of a superposition of the fundamental solutions  $\phi_k(z; \lambda)$  ( $k = 1, 2$ ) in (2.5) combined with the plane waves in the  $x$ - and  $y$ - directions. Therefore the scalar field is expanded generally as

$$\varphi(t, \vec{x}_\perp, z) = \int d\lambda \sum_{k=1,2} \int \frac{d^2 \vec{p}_\perp}{\sqrt{(2\pi)^2 2E}} \left\{ \alpha_k(\vec{p}_\perp, \lambda) e^{i\vec{p}_\perp \cdot \vec{x}_\perp - iEt} \phi_k(z; \lambda) + \alpha_k^\dagger(\vec{p}_\perp, \lambda) e^{-i\vec{p}_\perp \cdot \vec{x}_\perp + iEt} \phi_k(z; \lambda) \right\}, \quad (5.4)$$

where  $E$  is defined by

$$E = \sqrt{\vec{p}_\perp^2 + \lambda}. \quad (5.5)$$

The coefficients,  $\alpha_k(\vec{p}_\perp, \lambda)$  and  $\alpha_k^\dagger(\vec{p}_\perp, \lambda)$  ( $k = 1, 2$ ) in (5.4), are regarded as operators, which we postulate to satisfy the following commutation relations,

$$\left[ \alpha_k(\vec{p}_\perp, \lambda), \alpha_l^\dagger(\vec{p}'_\perp, \lambda') \right] = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(\lambda - \lambda') \frac{d\rho_{kl}(\lambda)}{d\lambda}, \quad (5.6)$$

$$\left[ \alpha_k(\vec{p}_\perp, \lambda), \alpha_l(\vec{p}'_\perp, \lambda') \right] = \left[ \alpha_k^\dagger(\vec{p}_\perp, \lambda), \alpha_l^\dagger(\vec{p}'_\perp, \lambda') \right] = 0. \quad (5.7)$$

Note that the spectral density matrix  $d\rho_{kl}(\lambda)/d\lambda$  in (5.6) is symmetric and positive semi-definite. Employing (5.6) and (5.7) we immediately get

$$[\dot{\varphi}(t, \vec{x}_\perp, z), \varphi(t, \vec{x}'_\perp, z')] = -i \delta^2(\vec{x}_\perp - \vec{x}'_\perp) \sum_{k,l=1,2} \int d\lambda \phi_k(z, \lambda) \frac{d\rho_{kl}(\lambda)}{d\lambda} \phi_l(z'; \lambda), \quad (5.8)$$

and are led to the canonical commutation relation (5.2) on using the completeness (2.11). The relation (5.3) can also be derived straightforwardly by using (5.6) and (5.7).

In case that there does not exist a bubble wall, i.e.,  $M(z) = M$  (constant), the operators  $\alpha_k^\dagger(\vec{p}_\perp, \lambda)$  and  $\alpha_k(\vec{p}_\perp, \lambda)$  have to reduce to the conventional creation and annihilation operators. In fact putting the solutions (2.19) together with the spectral functions (2.23), (2.24) and (2.25), we are able to rewrite (5.4) in the following form

$$\begin{aligned} \varphi(t, \vec{x}_\perp, z) = & \int_{M^2}^{\infty} \frac{d\lambda}{2\sqrt{\lambda - M^2}} \int \frac{d^2\vec{p}_\perp}{\sqrt{(2\pi)^3 2E}} \left\{ \right. \\ & a_-(\vec{p}_\perp, \lambda) e^{i\vec{p}_\perp \cdot \vec{x}_\perp + i\sqrt{\lambda - M^2} z} e^{-iEt} + a_+(\vec{p}_\perp, \lambda) e^{i\vec{p}_\perp \cdot \vec{x}_\perp - i\sqrt{\lambda - M^2} z} e^{-iEt} \\ & \left. + a_-^\dagger(\vec{p}_\perp, \lambda) e^{-i\vec{p}_\perp \cdot \vec{x}_\perp - i\sqrt{\lambda - M^2} z} e^{iEt} + a_+^\dagger(\vec{p}_\perp, \lambda) e^{-i\vec{p}_\perp \cdot \vec{x}_\perp + i\sqrt{\lambda - M^2} z} e^{iEt} \right\}, \end{aligned} \quad (5.9)$$

where we have introduced two types of linear combinations of  $\alpha_1(\vec{p}_\perp, \lambda)$  and  $\alpha_2(\vec{p}_\perp, \lambda)$ , i.e.,

$$a_\pm(\vec{p}_\perp, \lambda) \equiv \sqrt{2\pi} \left\{ \sqrt{\lambda - M^2} \alpha_1(\vec{p}_\perp, \lambda) \pm i\alpha_2(\vec{p}_\perp, \lambda) \right\}. \quad (5.10)$$

The commutation relations among these combinations turn out to be

$$\left[ a_+(\vec{p}_\perp, \lambda), a_+^\dagger(\vec{p}'_\perp, \lambda') \right] = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(\sqrt{\lambda - M^2} - \sqrt{\lambda' - M^2}), \quad (5.11)$$

$$\left[ a_-(\vec{p}_\perp, \lambda), a_-^\dagger(\vec{p}'_\perp, \lambda') \right] = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(\sqrt{\lambda - M^2} - \sqrt{\lambda' - M^2}), \quad (5.12)$$

and all of the other types of commutators vanish. Once we identify  $\sqrt{\lambda - M^2}$  as the  $z$ -component of the momentum  $\vec{p}$ , then it is easy to see that  $a_-(\vec{p}_\perp, \lambda)$  and  $a_+(\vec{p}_\perp, \lambda)$  are the usual annihilation operator of a particle with momentum  $\vec{p} = (\vec{p}_\perp, \sqrt{\lambda - M^2})$  and  $\vec{p} = (\vec{p}_\perp, -\sqrt{\lambda - M^2})$ , respectively.

In the presence of the bubble wall, the interpretation of operators  $\alpha_k^\dagger(\vec{p}_\perp, \lambda)$  and  $\alpha_k(\vec{p}_\perp, \lambda)$  is subtle and is not so simple. The notion of a particle itself is obscured, since the mass is position-dependent. We may, however, call  $\alpha_k^\dagger(\vec{p}_\perp, \lambda)$  and  $\alpha_k(\vec{p}_\perp, \lambda)$  as creation and annihilation operators of some ‘‘quanta’’ labelled by  $k$ ,  $\vec{p}_\perp$  and  $\lambda$ . Suppose that we define the vacuum  $|0\rangle$  by

$$\alpha_k(\vec{p}_\perp, \lambda)|0\rangle = 0, \quad \text{for } k = 1, 2, \quad \forall \vec{p}_\perp, \quad \forall \lambda. \quad (5.13)$$

In the symmetry-broken region, the state  $\alpha_k^\dagger(\vec{p}_\perp, \lambda)|0\rangle$  behaves like a one-particle state with a definite mass, say,  $M$ , and looks like a superposed state with four-momentum of  $(E, \vec{p}_\perp, +\sqrt{\lambda - M^2})$  and  $(E, \vec{p}_\perp, -\sqrt{\lambda - M^2})$ . We have to keep in our mind, however, that the normalization of this state differs from the usual one due to the  $\lambda$ -dependent quantity  $d\rho_{kl}(\lambda)/d\lambda$  in (5.6) and that this should be taken into account in quantitative calculations. We should also notice that the two-point function (2.4) is the vacuum expectation value of the time-ordered product of the scalar fields where the vacuum is defined by (5.13).

## 5.2 The Dirac field

Let us now turn our attention to the Dirac field, whose general solution to the free equations of motion (3.6) is given by a superposition of (3.46). We express the solution by separating the part belonging to the positive ( $E > 0$ ) and negative ( $E < 0$ ) energies in the following way, i.e.,

$$\begin{aligned} \psi(t, \vec{x}_\perp, z) = & \sum_{k=1,2} \sum_{s=+,-} \left\{ \int_{E>0} dE \int \frac{d^2\vec{p}_\perp}{2\pi} \beta_k^{(s)}(\vec{p}_\perp, E) \psi_k^{(s)}(z; E) e^{i\vec{p}_\perp \cdot \vec{x}_\perp - iEt} \right. \\ & \left. + \int_{E<0} dE \int \frac{d^2\vec{p}_\perp}{2\pi} \gamma_k^{(s)\dagger}(\vec{p}_\perp, E) \psi_k^{(s)}(z; E) e^{i\vec{p}_\perp \cdot \vec{x}_\perp - iEt} \right\}. \end{aligned} \quad (5.14)$$

The coefficients  $\beta_k^{(s)}(\vec{p}_\perp, E)$  and  $\gamma_k^{(s)\dagger}(\vec{p}_\perp, E)$  are operators whose anti-commutation relations are arranged so that the basic relations

$$\{\psi(t, \vec{x}_\perp, z), \psi^\dagger(t, \vec{x}'_\perp, z')\} = \delta^2(\vec{x}_\perp - \vec{x}'_\perp) \delta(z - z'), \quad (5.15)$$

$$\{\psi(t, \vec{x}_\perp, z), \psi(t, \vec{x}'_\perp, z')\} = 0, \quad (5.16)$$

$$\{\psi^\dagger(t, \vec{x}_\perp, z), \psi^\dagger(t, \vec{x}'_\perp, z')\} = 0, \quad (5.17)$$

are realized. Here we would like to point out that once we postulate the following anti-commutation relations

$$\{\beta_k^{(r)}(\vec{p}_\perp, E), \beta_l^{(s)\dagger}(\vec{p}'_\perp, E')\} = \delta^{rs} \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(E - E') \frac{d\rho_{kl}^{(s)}(E)}{dE}, \quad (5.18)$$

$$\{\gamma_k^{(r)}(\vec{p}_\perp, E), \gamma_l^{(s)\dagger}(\vec{p}'_\perp, E')\} = \delta^{rs} \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(E - E') \frac{d\rho_{kl}^{(s)}(E)}{dE}, \quad (5.19)$$

and vanishing of all other types of anti-commutators, then (5.15), (5.16) and (5.17) are all satisfied.

As in the scalar case we may call  $\beta_k^{(r)\dagger}(\vec{p}_\perp, E)$  and  $\beta_k^{(r)}(\vec{p}_\perp, E)$  as creation and annihilation operators of Fermionic “quanta” labelled by  $k, r, \vec{p}_\perp$  and  $E$ . Likewise we may call  $\gamma_k^{(r)\dagger}(\vec{p}_\perp, E)$  and  $\gamma_k^{(r)}(\vec{p}_\perp, E)$  as those of anti-Fermionic “quanta” labelled by  $k, r, \vec{p}_\perp$  and  $E$ . We define the vacuum by the conditions

$$\beta_k^{(r)}(\vec{p}_\perp, E)|0\rangle = 0, \quad \gamma_k^{(r)}(\vec{p}_\perp, E)|0\rangle = 0, \quad \text{for } k = 1, 2, \quad r = \pm, \quad \forall \vec{p}_\perp, \quad \forall E. \quad (5.20)$$

Then it turns out that the Fermion’s propagator (3.51) supplemented by the  $+i\varepsilon$  prescriptions (3.52) and (3.53) is an expectation value of the time-ordered product of Fermion’s bilinear operators with respect to this vacuum. In the symmetry-broken region, the states  $\beta_k^{(r)\dagger}(\vec{p}_\perp, E)|0\rangle$  and  $\gamma_k^{(r)\dagger}(-\vec{p}_\perp, E)|0\rangle$  behave like a one-particle Fermion and anti-Fermion states, respectively, with a definite mass, say,  $m$  and are a superposition of the four-momentum states of  $(|E|, \vec{p}_\perp, +q)$  and  $(|E|, \vec{p}_\perp, -q)$ , as a linear combination of  $k = 1$  and  $k = 2$  states. Here  $q$  is given by

$$q = \sqrt{E^2 - (m^2 + |\vec{p}_\perp|^2)}. \quad (5.21)$$

In the absence of the bubble wall, we have  $m(z) = m$  (constant) and  $\beta_k^{(s)}(\vec{p}_\perp, E)$  and  $\gamma_k^{(s)}(\vec{p}_\perp, E)$  may be shown to go over to the usual annihilation operators. In fact

$$b^{(\pm)}(\vec{p}_\perp, q) = \kappa^{(\pm)} \beta_1^{(\pm)}(\vec{p}_\perp, E) + i\kappa^{(\mp)} \beta_2^{(\pm)}(\vec{p}_\perp, E), \quad (5.22)$$

$$b^{(\pm)}(\vec{p}_\perp, -q) = \kappa^{(\pm)} \beta_1^{(\pm)}(\vec{p}_\perp, E) - i\kappa^{(\mp)} \beta_2^{(\pm)}(\vec{p}_\perp, E), \quad (5.23)$$

are the annihilation operators of the Fermion with the momentum  $\vec{p} = (\vec{p}_\perp, q)$  and  $\vec{p} = (\vec{p}_\perp, -q)$ , respectively, where we defined

$$\kappa^{(\pm)} = \sqrt{2\pi} \sqrt{\frac{|E| \pm \sqrt{m^2 + |\vec{p}_\perp|^2}}{2|E|}}. \quad (5.24)$$

We are able to confirm the usual anti-commutators

$$\left\{ b^{(\pm)}(\vec{p}_\perp, q), b^{(\pm)\dagger}(\vec{p}'_\perp, q') \right\} = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(q - q'), \quad (5.25)$$

$$\left\{ b^{(\pm)}(\vec{p}_\perp, -q), b^{(\pm)\dagger}(\vec{p}'_\perp, -q') \right\} = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(q - q'), \quad (5.26)$$

and all other types anti-commutators vanish.

Similarly, the creation operators of the anti-Fermion with momentum  $\vec{p} = (\vec{p}_\perp, q)$  and  $\vec{p} = (\vec{p}_\perp, -q)$  are expressed, respectively, by

$$c^{(\pm)\dagger}(-\vec{p}_\perp, -q) = \kappa^{(\pm)} \gamma_1^{(\mp)\dagger}(\vec{p}_\perp, E) - i\kappa^{(\mp)} \gamma_2^{(\mp)\dagger}(\vec{p}_\perp, E), \quad (5.27)$$

$$c^{(\pm)\dagger}(-\vec{p}_\perp, q) = \kappa^{(\pm)} \gamma_1^{(\mp)\dagger}(\vec{p}_\perp, E) + i\kappa^{(\mp)} \gamma_2^{(\mp)\dagger}(\vec{p}_\perp, E), \quad (5.28)$$

and are shown to satisfy the anti-commutators

$$\left\{ c^{(\pm)}(-\vec{p}_\perp, -q), c^{(\pm)\dagger}(-\vec{p}'_\perp, -q') \right\} = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(q - q'), \quad (5.29)$$

$$\left\{ c^{(\pm)}(-\vec{p}_\perp, q), c^{(\pm)\dagger}(-\vec{p}'_\perp, q') \right\} = \delta^2(\vec{p}_\perp - \vec{p}'_\perp) \delta(q - q'), \quad (5.30)$$

together with vanishing of all other types of anti-commutators.

## 6 Concluding Remarks

In the presence of the Higgs condensate bubble, we have successfully constructed the two-point Green's functions of scalar, spinor and vector fields in the form of integral representations. The key role of the construction is played by the spectral functions, which we are in principle able to compute provided that the large  $|z|$  behavior of fundamental solutions are known. The spectral functions also tell us about the range of the spectra and therefore we can identify the infrared region simply by looking at the integral representation. With the help of these propagators, we are able to check the interplay between virtual and real soft emitted gauge bosons, and natures of cancellation of infrared singularities.

The Sudakov-type double logarithms and variations thereof may also be investigated by using our Green's functions. As we see in (3.51), the denominators of the Fermion's propagator look like those in the old-fashioned perturbation. Of course the denominators



of the Bosonic propagator (2.4) may also be divided into two to cast them into the form of old-fashioned perturbation. It has been known that, in the very high energy limit, only a few among many terms in the old-fashioned perturbation survive the limit and perturbative calculation rules become very much simplified [89]. It has also been known that the derivation of the Sudakov-type double logarithms and their all-order resummation are possible by picking up only limited number of terms in the old-fashioned perturbation theory [90]. The propagators that we derived in the present paper may be adapted to such analyses.

For more dedicated analyses on the dynamics near the bubble wall interface, finite temperature Green's functions must be worked out, which, however, we will report in separate publications. In constructing finite temperature Green's functions, we have to take care of the operator ordering. The commutation and anti-commutation relations that we have postulated in Sect. 5 may provide us with a good testing-ground of their usefulness.

## References

- [1] B.P. Abbott et al., “Observation of Gravitational Waves from a Binary Black Hole Merger” Phys. Rev. Letters 116, 061102 (2016), arXiv:1602.03837 [gr-qc].  
DOI: <https://doi.org/10.1103/PhysRevLett.116.061102>
- [2] B.P. Abbott et al., “GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence” Phys. Rev. Letters 116, 241103 (2016), arXiv:1606.04855 [gr-qc].  
DOI: <https://doi.org/10.1103/PhysRevLett.116.241103>
- [3] B.P. Abbott et al., “GW170104: Observation of a 50-Solar-Mass Binary Black Hole Coalescence at Redshift 0.2” Phys. Rev. Letters 118, 221101 (2017), arXiv:1706.01812 [gr-qc].  
DOI: <https://doi.org/10.1103/PhysRevLett.118.221101>
- [4] B.P. Abbott et al., “GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral” Phys. Rev. Letters 119, 161101 (2017), arXiv:1710.05832 [gr-qc].  
DOI: <https://doi.org/10.1103/PhysRevLett.119.161101>
- [5] E. Witten, “Cosmic separation of phases” Phys. Rev. D 30 (1984) 272 - 285.  
DOI: <https://doi.org/10.1103/PhysRevD.30.272>
- [6] C. J. Hogan, “Gravitational radiation from cosmological phase transitions”, Mon. Not. Roy. Astron. Soc. 218 (1986) 629 - 636.  
DOI: <https://doi.org/10.1093/mnras/218.4.629>
- [7] A. Kosowsky, M.S. Turner and R. Watkins, “Gravitational waves from first order cosmological phase transitions” Phys. Rev. Letters 69 (1992) 2026.  
DOI: <https://doi.org/10.1103/PhysRevLett.69.2026>
- [8] M. Kamionkowski, A. Kosowsky, and M.S. Turner, “Gravitational radiation from first order phase transitions” Phys. Rev. D 49 (1994) 2837 - 2851.  
DOI: <https://doi.org/10.1103/PhysRevD.49.2837>  
astro-ph/9310044
- [9] K. Kajantie, M. Laine, K. Rummukainen and M.E. Shaposhnikov, “Is There a Hot Electroweak Phase Transition at  $m_H \geq m_W$  ?” Phys. Rev. Letters 77 (1996) 2887.  
DOI: <https://doi.org/10.1103/PhysRevLett.77.2887>

- [10] F. Karsch, T. Neuhaus, A. Patkos and J. Rank, “Critical Higgs Mass and Temperature Dependence of Gauge Boson Masses in the  $SU(2)$  Gauge-Higgs Model” Nucl. Phys. B Proc. Suppl. 53 (1997) 623 - 625.  
DOI: [https://doi.org/10.1016/S0920-5632\(96\)00736-0](https://doi.org/10.1016/S0920-5632(96)00736-0)
- [11] M. Gurtler, E.-M. Ilgenfritz and A. Schiller, “Where the electroweak phase transition ends” Phys. Rev. D 56 (1997) 3888.  
DOI: <https://doi.org/10.1103/PhysRevD.56.3888>
- [12] K. Rummukainen, M. Tsy-pin, K. Kajantie, M. Laine and M.E. Shaposhnikov, “The Universality Class of the Electroweak Theory”, Nucl. Phys. B 532 (1998) 283 - 314.  
DOI: [https://doi.org/10.1016/S0550-3213\(98\)00494-5](https://doi.org/10.1016/S0550-3213(98)00494-5)
- [13] F. Csikor, Z. Fodor and J. Heitger, “The Strength of the Electroweak Phase Transition at  $m(H) \approx 80$  GeV” Phys. Letters B441 (1999) 354 - 362.  
DOI: [https://doi.org/10.1016/S0370-2693\(98\)01127-7](https://doi.org/10.1016/S0370-2693(98)01127-7)
- [14] Y. Aoki, F. Csikor, Z. Fodor and A. Ukawa, “The end point of the first-order phase transition of the  $SU(2)$  gauge-Higgs model on a 4-dimensional isotropic lattice”, Phys. Rev. D 60 (1999) 0130001.  
DOI: <https://doi.org/10.1103/PhysRevD.60.013001>
- [15] M. Laine and K. Rummukainen, Nucl. Phys. B Proc. Suppl. 73 1999 180 - 185, “What’s new with the electroweak phase transition?” [hep-lat/9809045].  
DOI: [https://doi.org/10.1016/S0920-5632\(99\)85017-8](https://doi.org/10.1016/S0920-5632(99)85017-8)
- [16] M. D’Onofrio and K. Rummukainen, “Standard model cross-over on the lattice” Phys. Rev D 93 (2016) 025003, arXiv:1508.07161 1  
DOI: <https://doi.org/10.1103/PhysRevD93.025003>
- [17] O. Gould, S. Güyer and K. Rummukainen, “First-order electroweak phase transitions: A nonperturbative update”, Phys. Rev. D 106(2022) 114507, arXiv:2205.07238 [hep-lat].  
DOI: [10.1103/PhysRevD.106.114507](https://doi.org/10.1103/PhysRevD.106.114507)
- [18] G.W. Anderson and L.J. Hall, “The electroweak phase transition and baryogenesis” Phys. Rev. D 45 (1992) 2685.  
DOI: <https://doi.org/10.1103/PhysRevD.45.2685>
- [19] M. Quirós, “Finite temperature field theory and phase transitions” ICTP summer school in high-energy physics and cosmology (1999) p. 187  
hep-ph/9901312
- [20] C. Grojean, G. Servant and J.D. Wells, “First-order electroweak phase transition in the standard model with a low cutoff” Phys. Rev. D 71 (2005) 036001, hep-th/0407019.  
DOI: <https://doi.org/10.1103/PhysRevD.71.036001>
- [21] C. Caprini et al, “Science with the space-based interferometer eLISA. II: gravitational waves from cosmological phase transitions” JCAP 04 (2016) 001, arXiv: 1512.06239 [astro-ph.CO]  
DOI: [10.1088/1475-7516/2016/04/001](https://doi.org/10.1088/1475-7516/2016/04/001)
- [22] C. Caprini et al, “Detecting gravitational waves from cosmological phase transitions with LISA: an update”, JCAP 03 (2020) 024, arXiv: 1910.13125 [astro-ph.CO]  
DOI: [10.1088/1475-7516/2020/03/024](https://doi.org/10.1088/1475-7516/2020/03/024)
- [23] M. Hindmarsh, M. Luben, J. Lumma and M. Pauly, “Phase transitions in the early universe” SciPost Physics Lecture Notes, 24 (2021) 1-61, arXiv:2008.09136 [astro-ph.CO]

doi:10.21468/SciPostPhysLecNotes.24

- [24] LISA Cosmology Working Group, P. Auclair et al., “Cosmology with the Laser Interferometer Space Antenna”, *Living Rev. Rel.* 26 (2023) 1 - 254, arXiv: 2204.05434 [astro-ph.CO].  
<https://doi.org/10.1007/s41114-023-00045-2>
- [25] S. Kawamura et al, “Space gravitational-wave antennas DECIGO and B-DECIGO” , *Int. J. Mod. Phys. D* 28 (2019) 12, 1845001.  
doi:101142/S0218271818450013
- [26] W.-H. Ruan, Z.-K. Guo, R.-G.Cai and Y.-Z. Zhang, “Taiji program: Gravitational-wave sources”, *Int. J. Mod. Phys. A* 35 (2020) 17, 2050075, arXiv:1807.09495 [gr-qc].  
doi:101142/S0217751X2050075X
- [27] J. Luo et al., “TianQin: a space-borne gravitational wave detector”, *Class. Quantum Grav.* 33 (2016) 3, 035010, arXiv:1512.02076 [astro-ph.IM].  
doi: 101088/0264-9381/33/3/035010
- [28] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, “On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe” *Phys. Lett B* 155 (1985) 36 - 42.  
[https://doi.org/10.1016/0370-2693\(85\)91028-7](https://doi.org/10.1016/0370-2693(85)91028-7)
- [29] M.E. Shaposhnikov, “Baryon Asymmetry of the Universe in Standard Electroweak Theory” *Nucl. Phys. B* 287 (1987) 757- 775.  
[https://doi.org/10.1016/0550-3213\(87\)90127-1](https://doi.org/10.1016/0550-3213(87)90127-1)
- [30] A.G. Cohen, D.B. Kaplan and A.E. Nelson, “Progress in Electroweak Baryogenesis”, *Annu. Rev. Nucl. Part. Sci.*, 43 (1993) 27 - 70, hep-ph/9302210 [hep-ph] .  
<https://doi.org/10.1146/annurev.ns.43.120193.000331>
- [31] T. Konstandin, “Quantum Transport and Electroweak Baryogenesis” *Phys. Usp.* 56 (2013) 747, arXiv: 1302.6713 [hep-ph].  
<https://doi.org/10.3367/ufne.0183.201308a.0785>
- [32] D.E. Morrissey and M.J. Ramsey-Musolf, “Electroweak Baryogenesis” *New J. Phys.* 14 (2012) 125003, arXiv: 1206.2942 [hep-ph]  
DOI:10.1088/1367-2630/14/12/125003
- [33] B. Garbrecht, “Why is there more matter than antimatter ?” , *Progr. Part. Nucl. Phys.* 110 (2020) 103727 1-46, arXiv: 1812.02651 [hep-ph]  
<https://doi.org/10.1016/j.ppnp.2019.103727>
- [34] D. Bödeker and G.D. Moore, “Can electroweak bubble walls run away?”, *JCAP* 05 (2009) 09, arXiv: 0903.4099 [hep-ph].  
DOI: 10.1088/1475-7516/2009/05/009
- [35] J.R. Espinoza, T. Konstandin, J.M. No and G. Servant, “Energy budget of cosmological first-order phase transitions” *JCAP* 06 (2010) 028, arXiv: 1004.4187 [hep-ph].  
<https://doi.org/10.1088/1475-7516/2010/06/028>
- [36] D. Bödeker and G.D. Moore, “ Electroweak bubble wall speed limit”, *JCAP* 05 (2017) 025, arXiv: 1703.08215 [hep-ph]  
DOI: 10.1088/1475-7516/2017/05/025
- [37] V.L. Ginzburg and I.M. Frank, “Radiation of a uniformly moving electron due to its transition from one medium into another” *J. Phys. (USSR)* 9 (1945) 353 - 362; *Zh. Eksp. Teor. Fiz.* 16 (1946) 15-28.

- [38] J.D. Jackson, “Classical Electrodynamics” (Wiley, 1998)  
ISBN-10:9780471309321
- [39] S. Höche, J. Kozaczuk, A.J. Long, J. Turner and Y. Wang, “Towards an all-orders calculation of the electroweak bubble wall velocity”, JCAP 03 (2021) 009, arXiv: 2007.10343 [hep-ph].  
DOI: 10.1088/1475-7516/2021/03/009
- [40] V.V. Sudakov, “Vertex Parts at Very High Energies in Quantum Electrodynamics” Soviet Physics JETP 3 (1956) 65 - 71.
- [41] A.H. Mueller, “Asymptotic behavior of the Sudakov form factor” Phys. Rev. D20 (1979) 2037 - 2046.  
DOI: <https://doi.org/10.1103/PhysRevD.20.2037>
- [42] J.C. Collins, “Algorithm to compute corrections to the Sudakov form factor” Phys. Rev. D22 (1980) 1478 - 1489.  
DOI: <https://doi.org/10.1103/PhysRevD.22.1478>
- [43] A. Sen, “Asymptotic behavior of the Sudakov form factor in quantum chromodynamics” Phys. Rev. D24 (1981) 3281 - 3304.  
DOI: <https://doi.org/10.1103/PhysRevD.24.3281>
- [44] A. Sen, “Asymptotic behavior of the fixed-angle on-shell scattering amplitudes in non-Abelian gauge theories” Phys. Rev. D28 (1983) 860 - 875.  
DOI: <https://doi.org/10.1103/PhysRevD.28.860>
- [45] J.C. Collins, “Sudakov Form Factor” in “Perturbative QCD” (ed. by A.H. Mueller) Advanced Series on Directions in High Energy Physics 5, 1 (1988, ISBN: 978-981-4503-26-6)
- [46] H. Contopanagos, E. Laenen, and G.F. Sterman, “Sudakov factorization and resummation” Nucl. Phys. B 484 (1997) 303 - 330, hep-ph/9604313  
[https://doi.org/10.1016/50550-3213\(96\)00567-6](https://doi.org/10.1016/50550-3213(96)00567-6)
- [47] M.B. Mancha, T. Prokopec and B. Świążewska, “Field-theoretic derivation of bubble-wall force” JHEP 01 (2021) 070, arXiv:2005.10875 [hep-th]  
DOI: [https://doi.org/10.1007/JHEP01\(2021\)070](https://doi.org/10.1007/JHEP01(2021)070)
- [48] A. Buckley et al., “General-purpose event generators for LHC physics”, Physics Reports 504 (2011) 145-233, arXiv:1101.2599 [hep-ph].  
doi: 10.1016/j.physrep.2011.03.005
- [49] B.R. Webber, “Monte Carlo Simulation of Hard Hadronic Processes” Ann. Rev. Nucl. Part. Sci. 36 (1986) 253 - 286, arXiv: hep-ph/0312336  
<https://doi.org/10.1146/annurev.ns.36.120186.001345>
- [50] A. Denner and S. Pozzorini, “One-loop leading logarithms in electroweak radiative corrections I. Results” Eur. Phys. J. C 18 (2001) 461- 480, arXiv:hep-ph/0010201  
<https://doi.org/10.1007/s100520100551>
- [51] A. Denner and S. Pozzorini, “One-loop leading logarithms in electroweak radiative corrections II. Factorization of collinear singularities” Eur. Phys. J. C 21 (2001) 63-79, arXiv:hep-ph/0104127  
<https://doi.org/10.1007/s100520100721>
- [52] A. Denner and S. Pozzorini, “An algorithm for the high-energy expansion of multi-loop diagrams to next-to-leading logarithmic accuracy” Nucl. Phys. B717 (2005) 48-85, arXiv:hep-ph/0408068

<https://doi.org/10.1016/j.nuclphysb.2005.03.036>

- [53] M. Roth and A. Denner, “High-Energy Approximation of One-Loop Feynman Integrals” Nucl. Phys. B 479 (1996) 495 - 514, hep-ph/9605420.  
[https://doi.org/10.1016/0550-3213\(96\)00435-X](https://doi.org/10.1016/0550-3213(96)00435-X)
- [54] V.S. Fadin, L.N. Lipatov, A.D. Martin and M. Melles, “ Resummation of double logarithms in electroweak high energy processes”, Phys. Rev. D 61 (2000) 094002 1-13,  
arXiv:hep-ph/9910338  
DOI: <https://doi.org/10.1103/PhysRevD.61.094002>
- [55] M. Melles, “Subleading Sudakov logarithms in electroweak high energy processes to all orders” Phys. Rev. D 63 (2001) 034003, arXiv:hep-ph/0004056  
DOI: <https://doi.org/10.1103/PhysRevD.63.034003>
- [56] D. Pagani and M. Zaro, “One-loop electroweak Sudakov logarithms: a revisit and automation” JHEP 02 (2022) 161, arXiv: 2110.03714 [hep-ph]  
[https://doi.org/10.1007/JHEP02\(2022\)161](https://doi.org/10.1007/JHEP02(2022)161)
- [57] J.M. Lindert and L. Mai, “Logarithmic EW corrections at one-loop”  
arXiv: 2312.07927 [hep-ph]
- [58] B. Laurent and J.M. Cline, “Fluid equations for fast-moving electroweak walls” Phys. Rev. D 102 (2020) 063516, arXiv:2007.10935 [hep-ph]  
DOI: <https://doi.org/10.1103/PhysRevD.102.063516>
- [59] B. Laurent and J.M. Cline, “First principles determination of bubble wall velocity” Phys. Rev. D 106 (2022) 023501, arXiv:2204.13120 [hep-ph].  
DOI: <https://doi.org/10.1103/PhysRevD.106.023501>
- [60] S. De Curtis, L.Delle Rose, A. Guiggiani, A.G. Muyor and G. Panico, “Bubble wall dynamics at the electroweak phase transition” JHEP 03 (2022) 163, arXiv:2201.08220 [hep-ph]  
[https://doi.org/10.1007/JHEP03\(2022\)163](https://doi.org/10.1007/JHEP03(2022)163)
- [61] S. De Curtis, L.Delle Rose, A. Guiggiani, A.G. Muyor and G. Panico, “Collision Integrals for Cosmological Phase Transitions” JHEP 05 (2023) 194, arXiv:2303.05846 [hep-ph]  
[https://doi.org/10.1007/JHEP05\(2023\)194](https://doi.org/10.1007/JHEP05(2023)194)
- [62] S. De Curtis, L.Delle Rose, A. Guiggiani, A.G. Muyor and G. Panico, “Non-linearities in cosmological bubble wall dynamics” arXiv:2401.13522 [hep-ph]
- [63] S. Balaji, M. Spannowsky and C. Tamarit, “Cosmological bubble friction in local equilibrium” JCAP 03 (2021) 051, arXiv:2010.08013 [hep-ph].  
DOI: 10.1088/1475-7516/2021/03/051
- [64] W.-Y.Ai, B. Garbrecht and C. Tamarit, “Bubble wall velocities in local equilibrium” JCAP 03 (2022) 015, arXiv: 2109.13710 [hep-ph]  
DOI: 10.1088/1475-7516/2022/03/015
- [65] W.-Y.Ai, B. Laurent and J. van de Vis, “Model-independent bubble wall velocities in local thermal equilibrium”, JCAP 07 (2023) 002, arXiv: 2303.10171 [astro-ph.CO].  
DOI: 10.1088/1475-7516/2023/07/002
- [66] A. Azatov and M. Vanvlasselaer, “Phase transitions in perturbative walking dynamics” JHEP 09 (2020) 085, arXiv:2003.10265 [hep-ph]  
[https://doi.org/10.1007/JHEP09\(2020\)085](https://doi.org/10.1007/JHEP09(2020)085)

- [67] Y. Gouttenoire, R. Jinno and F. Sala, “Friction pressure on relativistic bubble walls”, JHEP 05 (2022) 004, arXiv:2112.07686 [hep-ph].  
[https://doi.org/10.1007/JHEP05\(2022\)004](https://doi.org/10.1007/JHEP05(2022)004)
- [68] I.G. Garcia, G. Koszegi and R. Petrossian-Byrne, “Reflections on Bubble Walls” JHEP 09 (2023) 013, arXiv: 2212.10572 [hep-ph]  
[https://doi.org/10.1007/JHEP09\(2023\)013](https://doi.org/10.1007/JHEP09(2023)013)
- [69] A. Azatov and M. Vanvlasselaer, “Bubble wall velocity: heavy physics effects” JCAP 01 (2021) 058, arXiv:2010.02590 [hep-ph]  
 DOI: 10.1088/1475-7516/2021/01/058
- [70] G. C. Dorsch, S.J. Huber and T. Konstandin, “On the wall velocity dependence of electroweak baryogenesis” JCAP 08 (2021) 020, arXiv:2106.06547 [hep-ph].  
 DOI: <https://doi.org/10.1088/1475-7516/2021/08/020>
- [71] G. C. Dorsch, S.J. Huber and T. Konstandin, “A sonic boom in bubble wall friction” JCAP 04 (2022) 010, arXiv:2112.12548 [hep-ph]  
 DOI: 10.1088/1475-7516/2022/04/010
- [72] G. C. Dorsch and D.A. Pinto, “Bubble wall velocities with an extended fluid Ansatz” arXiv:2312.02354 [hep-ph]
- [73] A. Azatov, G. Barni, R. Petrossian-Byrne and M. Vanvlasselaer, “Quantisation Across Bubble Walls and Friction” arXiv: 2310.06972 [hep-ph]
- [74] H. Weyl, “Über gewöhnliche Differentialgleichungen mit Singularitäten und die zugehörigen Entwicklungen willkürlicher Funktionen”, *Mathematische Annalen* **68** (1910) 220 - 269.  
<https://doi.org/10.1007/BF01474161>.
- [75] M.H. Stone, “Linear Transformation in Hilbert Space and their Applications to Analysis”, American Mathematical Society Colloquium Publication, vol. 15 (New York, 1932)  
 ISBN: 9780821810156[0821810154]
- [76] E.C. Titchmarsh, “Eigenfunction expansions associated with second-order differential equation” (Oxford, Clarendon Press, 1946)
- [77] K. Kodaira, “The Eigenvalue Problem for Ordinary Differential Equation of the Second Order and Heisenberg’s Theory of S-Matrices” *American Journal of Mathematics*, **71** (1949) 921 - 945.  
<https://www.jstor.org/stable/2372377>
- [78] K. Kodaira, “On ordinary differential equations of any even order and corresponding eigenfunction expansion” *American Journal of Mathematics*, **72** (1950) 502 - 544.  
<https://www.jstor.org/stable/2372051>
- [79] K. Yosida, “On Titchmarsh-Kodaira’s formula concerning Weyl-Stone’s eigenfunction expansion” *Nagoya Mathematical Journal* 1 (1950) 49-58.  
<https://doi.org/10.1017/S0027763000022820>
- [80] K. Yosida, “Lectures on Differential and Integral Equations” (Dover Publication, New York, 1991)  
 ISBN-13: 978-0486666792
- [81] A. M. Polyakov, “Particle spectrum in quantum field theory” *JETP Lett.* 20 (1974) 194 - 195.

- [82] A. Ayala, J.J-Marlan, L. McLerran and A.P. Vischer, “Scattering in the presence of electroweak phase transition bubble wall” *Phys. Rev. D* 49 (1994) 5559 - 5570.  
DOI: <https://doi.org/10.1103/PhysRevD.49.5559>
- [83] G.R. Farrar and J.W. McIntosh, “Scattering from a domain wall in a spontaneously broken gauge theory” *Phys. Rev. D* 51 (1995) 5889 - 5904, arXiv:hep-ph/9412270.  
DOI: <https://doi.org/10.1103/PhysRevD.51.5889>
- [84] S.D. Conte and W.C. Sangren, “An Expansion Theorem for a Pair of Singular First Order Equations” *Canadian Journal of Mathematics*, 6 (1954) 554 - 560.  
<https://doi.org/10.4153/CJM-1954-060-0>
- [85] B.W. Roos and W.C. Sangren, “Spectra for a Pair of Singular First Order Differential Equations” *Proceedings of the American Mathematical Society*, 12 (1961) 468 - 476.  
<https://www.jstor.org/stable/2034220>
- [86] E.C. Titchmarsh, “ Some Eigenfunction Expansion Formulae” *Proceedings of the London Mathematical Society* (3) 11 (1961) 159-168.  
<https://doi.org/10.1112/plms/s3-11.1.159>
- [87] B.M. Levitan and I.S. Sargsjan, “Introduction to Spectral Theory: Selfadjoint Ordinary Differential Operators” (Translations of Mathematical Monographs, Vol. 39, American Mathematical Society, 1975)  
ISBN O-8218-1589-X
- [88] B.M. Levitan and I.S. Sargsjan, “Sturm-Liouville and Dirac Operators” ( Kluwer Academic Publishers, 1991) .  
ISBN 978-94-010-5667-0
- [89] S. Weinberg, “Dynamics at Infinite Momentum”, *Phys. Rev.* 150 (1966) 1313 - 1318.  
DOI: <https://doi.org/10.1103/PhysRev.150.1313>
- [90] R. Jackiw, “Dynamics at High Momentum and the Vertex Function of Spinor Electrodynamics”, *Annals of Physics* 48 (1968) 292 - 321.  
[https://doi.org/10.1016/0003-4916\(68\)90087-0](https://doi.org/10.1016/0003-4916(68)90087-0)