

# Strangeness Enhancement in Proton-Proton Collisions

Ahmed. M. Hamed\*and Fatma. H. Sawy†  
American University in Cairo, Cairo, Egypt

June 4, 2024

## Abstract

The analysis for the strange and multi-strange hadrons production is presented for proton-proton (pp) collisions at  $\sqrt{s_{NN}} = 200$  GeV as reported by the PYTHIA 8 event generator simulations. The strangeness enhancement is studied in details at  $\sqrt{s_{NN}} = 200$  GeV in proton-proton (pp). We preform the transverse momentum distributions for high and low multiplicity events of the strange and multi-strange hadrons. Here we present the observation of strangeness enhancement in pp collisions for  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$ ,  $\Omega^-(sss)$  production. At the RHIC energy, the strangeness content yield increases with the event multiplicity.

## Introduction

The strangeness enhancement as a signature of QGP has been discussed for nearly 40 years ago[1] [2]. In 1982, Rafelski and Müller proposed that QGP formation should lead to a higher abundance of strangeness per participating nucleon than what is seen in pp interactions [3] [4]. In 1995, the series of International Conference on Strangeness in Quark Matter first organized in Tucson, Arizona [5]. Strangeness enhancement was indeed observed for the first time by comparing central nucleus-nucleus collisions with proton-ion and pp reactions at the SPS [3]. In 2005, the Brookhaven National Laboratory (BNL) announced that forming the perfect liquid phase in nucleus-nucleus collisions [6]. The researchers discovered spatial correlations reminiscent of the high multiplicity events attributed to QGP formation in nucleus-nucleus collisions [6] [7]. In 2010, the CMS collaboration spotted something unexpected in pp collisions, the experiment reported flow-like features of particle production in high multiplicity events in pp collisions [8]. In 2017, the ALICE collaboration reports the first observation of strangeness enhancement in pp collisions, a signature of QGP [1]. At the LHC energies, the high multiplicity events in pp collisions show features which were considered as signatures of QGP formation [3]. Since the QGP is formed in pp collisions, physicists could probe the properties of QGP in a far simpler system than the one produced in nucleus-nucleus collisions.

Here we present the transverse momentum distributions for strange hadrons,  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$ ,  $\Omega^-(sss)$ , for PYTHIA 8 simulated pp collisions at the center of mass energy  $\sqrt{s_{NN}} = 200$  GeV. The normalized transverse momentum distributions to total number of events are provided for all of the six considered particles. The simulated data are preformed for low and high multiplicity events, from event multiplicity less than 70 to higher than 200, to compare between them. The ratio between the production for low and high multiplicity events for  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$ ,  $\Omega^-(sss)$  are examined in detail. In addition, we study the relation between the strange content yield and the event multiplicity. The interest in these studies lies deep understanding of matter phase transitions and dynamics of strangeness creation. We show the production mechanism/s of strange and multi-strange particles. Our main objectives are introducing full complete study about QGP formation signature, the strangeness enhancement. The primary physics task of STAR experiment is to study the formation and characteristics of the quark gluon plasma. Details of the STAR experiment are in [9] [10]. Since PYTHIA 8 event generator simulations cannot be performed for nucleus-nucleus collisions, we consider the high multiplicity events in pp

---


\*ahmed.mhamed@aucegypt.edu

†fatma.helal@aucegypt.edu

collisions as heavy ion events. The PYTHIA 8 generator is an ideal system where the final state interactions are not included i.e the hadrons stream freely and do not talk to each other.

## The Strangeness Enhancement

The strangeness enhancement in high energy nuclear interactions provides one of the tools to investigate properties of QGP [2]. An increase of the strange quark density in the medium result in an increase of the strange hadron production. The mass of strange quark,  $95 \text{ MeV}/c^2$ , is equivalent to the temperature or energy at which hadrons dissolve into quarks. This means that the abundance of strange quarks is sensitive to the conditions of the deconfined matter phase. If the strange quark production is large, it can be assumed that deconfinement conditions were reached. As the lifetime of QGP is ultra short, an order of  $10^{-23}$  of a second, the signatures of its creation are indirect in nature. The quark flavor composition of the plasma varies during its ultra short lifetime as new flavors of quarks such as strangeness are cooked up inside. Since the initial colliding protons have no strangeness content, all new fresh strange quarks must be formed from the new plasma medium produced. The plasma evolves in time to form a liquid-like medium of quarks and gluons. The new strange quarks are created from the medium provided that there is enough time and that the temperature is high enough. T. Biro and J. Zimanyi demonstrated that strange quarks could not be produced fast enough alone by quark anti-quark pair production. A new mechanism get alone in QGP was proposed. Rafelski and Müller showed that the equilibrium of the strangeness yield in QGP is only possible due to gluon fusion. The gluon fusion process occurs almost ten times faster than quark - antiquark pair production. The strangeness enhancement is combined with gluon existence in QGP, in which the gluon dissolves to a pair of strange quarks rapidly as shown in Figure 1. The up and down quarks are easily produced as quark - antiquark pairs in the hot fireball because they have small masses. The Wroblewski ratio of newly produced  $s\bar{s}$  pairs with the normalized light quark pairs  $\bar{u}u$  and  $\bar{d}d$  is considered a measure of efficiency of strangeness production. Since  $s\bar{s}$  pairs are generated together from the same medium by the same way, the both strange matter and antimatter must be originated from a common source by the same creation mechanism [11]. Shortly, the strange (anti)baryon production processes in the two successive steps. The strange quarks are produced inside the fireball and later on, in an independent process, they hadronized from several (anti)strange quarks to (anti)baryons. Since the production of strangeness occurs predominantly in thermal gluon fusion, as discussed before, the overabundant presence of strangeness is theoretical linked to presence of thermal gluons, and thus to QGP [12].



**Figure 1:** Feynman diagrams for production of strange quarks in the QGP.

At sufficient high temperature and energy density, the ordinary nuclear phase goes to plasma phase [1]. After an initial pre-equilibrium stage, the colored strongly interacting medium produced in the collisions of the two interacting protons quickly achieves local thermodynamic equilibrium and, hence, reaches a plasma state [13]. The deconfined interacting QGP phase is the equilibrium state of matter at high temperature ( $T$ ) and/or particle density ( $\epsilon$ ) [11]. An early study explored the phase transition from hadronic matter to QGP and postulated the idea of chemical and thermal equilibrium [14]. The gluons and light quarks reach chemical equilibrium during the beginning stage of the plasma state. The heavier quark flavors such as strange typically approach chemical equilibrium in a thermodynamic evolution process. To obtain the measurable abundance of strange quarks, the lifetime of the plasma should reach after  $2 \times 10^{-23}$ s. At  $T = T_{ch} = 160 - 170 \text{ MeV}$ , the chemical freeze-out (chemical equilibrium) is assumed to take place immediately after hadronization of the QGP, the strangeness enhancement chemically equilibrated in plasma and the strange abundance saturation is clearly visible as well [15] [16]. The chemical freeze-out temperature  $T_{ch}$  and the baryon chemical potential  $\mu_B$ , describes the net baryon content of the system, mainly govern the statistical models. As the initial quark chemical potential increases, the production rate of gluons increases and the gluon equilibration rate necessarily decreases, thus leading to the little energy consumption of the system, i.e., slow cooling

of the medium [17]. The increase of the quark phase lifetime with increasing initial quark chemical potential obviously increases strangeness production <sup>1</sup> [17]. The relative chemical equilibrium happens when the strangeness is originated from the quarks and gluons exchanging where the dominant mechanism of strangeness production involves gluons only present at pure QGP phase. The absolute chemical equilibrium happens when the strangeness is originated from pair production between constituents of the plasma. At  $T \approx 120 - 160$  MeV, the strangeness abundance shows a pronounced threshold behavior. With decreasing the temperature of the medium, the phase transition between plasma phase to hadronic phase occurs at a certain critical temperature ( $T_c$ ). As time evolves and the plasma medium rapidly expands and cools, remaining close to local equilibrium, the hadron gas phase is produced where the degrees of freedom are the hadronic ones, as quarks and gluons are confined [18] [13]. During the expansion phase of the hadron gas, collective hydrodynamic flow develops from the initially generated pressure gradients in the strongly interacting system. This results in a characteristic dependence of the shape of the transverse momentum ( $p_T$ ) distribution on the particle mass, which can be described with a common kinetic freeze out temperature parameter  $T_{kin}$  and a collective average expansion velocity [18]. After a short kinetic stage of interacting hadron gas, the strongly stable particles stop colliding (kinetic freeze-out) and freely travel toward the detectors. The hadron gas is born into equilibrium and the hadrons escape from the reaction medium immediately after kinetic freeze-out. [13].

From thermodynamics point of view, the QGP is formed where the degrees of freedom are at partonic level (free quarks and gluons). The initial hot and dense partonic matter can be described in terms of hydrodynamic and statistical models [18]. The signature of the formation of QGP lies exactly on the overabundance of strangeness production in the medium. The energy needed to produce strange hadrons in a thermally equilibrated hadron gas is significantly higher than in the case of a QGP therefore the production of strange hadrons in hadron gas would be much more difficult [19]. The strangeness in the plasma phase is more abundant than in the hadronic gas phase, even if the gas is saturated. Originally, this enhancement is due to the high production rate of  $gg \rightarrow s\bar{s}$  in a QGP, a process absent in the hadronic state. The subsequent hadronization of these (anti)strange quarks results in a significant increase in strange hadron production, thus signaling a QGP was formed [9]. At certain lower temperature, the fast expansion of the fireball will lead to super cooling which may help to understand the sudden nature of the fireball breakup that effectively described by relativistic hydrodynamic equations [13]. Features of strange baryon and (anti)baryon spectra demonstrate sudden breakup of the fireball into hadrons, absence of re-equilibration. The gluons attract each other which gives rise to the confinement of quarks inside hadrons. When QGP hadronized, the high availability of strange quarks and anti-quarks helps to produce matter and antimatter containing multiple strange quarks, which is otherwise rarely made. The hadronization of the QGP phase is direct, fast and sudden [11]. Many strange hadrons decay within several centimeters from the point of production [12]. Strange hadrons are naturally radioactive and decay by weak interactions into lighter hadrons on a time scale that is extremely long compared with the nuclear collision times. This makes it relatively easy to detect strange particles through the tracks left by their decay products <sup>2</sup>.


## The Transverse Momentum and QGP

We work to study strangeness enhancement from PYTHIA 8 event generator data sets at RHIC energy for pp collisions. In order to be more specific about the nature of mechanism/s of strangeness creation, we study the transverse momentum spectrum at low multiplicity and high multiplicity events for mesons  $K_s^+(u\bar{s})$  and  $K_s^-(s\bar{u})$  and for baryons  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$  and  $\Omega^-(sss)$ . The evidence for strangeness enhancement is obtained from the study of strange hadron transverse momentum spectra. The measurements of transverse momentum distributions of  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$  and  $\Omega^-(sss)$  are an important corner-stones of the claim that QGP has been formed in pp collisions at  $\sqrt{s_{NN}} = 200$  GeV. The Figure 2 and Figure 3 prove that  $K_s^+(u\bar{s})$  and  $K_s^-(s\bar{u})$  can


<sup>1</sup>From RHIC to LHC energies, the abundances of strange particles relative to pions in nucleus-nucleus collisions do not show a significant dependence on either the initial volume (collision centrality) or the initial energy density (collision energy)[1].

<sup>2</sup>Strangeness enhancement has been experimentally observed for CERN SPS, RHIC and LHC energies in nucleus-nucleus collisions [20]

be produced in events have multiplicity less than 70 pronouncing that the strangeness is originated.



**Figure 2:** The transverse momentum distribution of PYTHIA 8 simulations for  $k^+(u\bar{s})$  particles at  $\sqrt{s_{NN}} = 200$  GeV for low multiplicity events, less than 70, in the left pannel and for low multiplicity events normalized to total number of simulated events in the right pannel.




**Figure 3:** The transverse momentum distribution of PYTHIA8 simulations for  $k^-(s\bar{u})$  particles at  $\sqrt{s_{NN}} = 200$  GeV for low multiplicity events, less than 90, in the left pannel and for low multiplicity events normalized to total number of simulated events in the right pannel.


The strangeness enhancement is observed in events having multiplicity less than 120 emphasized through the  $\Lambda^0(uds)$  transverse momentum distribution, Figure 4. This observation is supported when the distributions Figure 5 and Figure 6 are originated proving the enhancement in  $\Xi^0(uss)$  and  $\Xi^-(dss)$  production for events having multiplicity less than 150. The Figure 7 establishes the overabundant of strangeness through the higher production of  $\Omega^-(sss)$  for events having multiplicity less than 200. Since an observation of enhanced strange and multi-strange particles is called strangeness enhancement, we confirm that for events, have multiplicity less than 200, the strangeness is produced as it is proved in Figure 2, Figure 3, Figure 4, Figure 5, Figure 6 and Figure 7. As the event multiplicity increases, the production of strange and multi-strange particle increases for these events. Additionally, we find out that for these events, as the strangeness content increases, the  $dn/dp_T$  is suppressed at high  $p_T$ . As already known, at high  $p_T$ , the strangeness is produced in perturbative processes by flavor creation ( $q\bar{q} \rightarrow s\bar{s}$ ) or ( $gg \rightarrow s\bar{s}$ ), flavor excitation<sup>3</sup> ( $qs \rightarrow qs$ ) and gluon splittings ( $g \rightarrow s\bar{s}$ ) as shown in Figure 1. From intermediate to high  $p_T$ , the produced hadrons are suppressed and this effect is attributed to the energy loss of partons as they move through the hot and dense medium produced [9]. The details of the energy loss mechanism guide us to understand dynamics of strangeness creation. At critical sharp temperature, the de-confined phase of the interacting protons is reached producing

<sup>3</sup>In the flavor excitation process, a heavy sea quark belonging to an incoming hadron scatters off a quark or a gluon from the other hadron[21].


medium of interacting partons. The fireball formed from the colliding constituents originates the new quarks heavier than interacting partons through soft and hard collisions. Since  $s$  quark is heavier than the lightest quarks,  $u$  and  $d$ , the fireball starts to produce it. As the temperature of the system is higher than the mass of a strange quark, the strange quarks and anti-quarks can be abundantly produced through several processes leading to strangeness enhancement manifesting equilibrated QGP formation. So that we had to study the low multiplicity events for strange  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$  and multi-strange  $\Xi^0(uss)$ ,  $\Xi^-(dss)$ ,  $\Omega^-(sss)$  and they are reported in Figure 2, Figure 3, Figure 4, Figure 5, Figure 6 and Figure 7. From this deep study, equilibrated QGP formation could be occurred in low multiplicity events, less than 200, in pp collisions. At low  $p_T$ , non-perturbative processes dominate the production of strange hadrons through soft creation mechanism/s that is explained only by phenomenology models [22]. As it notices from the  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$  and  $\Omega^-(sss)$  distributions, Figure 3, Figure 2 Figure 4, Figure 5, Figure 6 and Figure 7, that a high strange particle production is presented at low momentum. This observation is consist with the PYTHIA 8 event generator nature i.e the generator produces non interacting final state particles by soft creation mechanism [22].




**Figure 4:** The transverse momentum distribution of PYTHIA8 simulations for  $\Lambda^0(uds)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for low multiplicity events, less than 120, in left pannel and or low multiplicity events normalized to total number of simulated events in the right pannel.



**Figure 5:** The transverse momentum distribution of PYTHIA8 simulations for  $\Xi^-(dss)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for low multiplicity events, less than 150, in the left pannel and for low multiplicity events normalized to total number of simulated events in the right pannel.




**Figure 6:** The transverse momentum distribution of PYTHIA8 simulations for  $\Xi^0(uss)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for low multiplicity events, less than 150, in the left pannel and for low multiplicity events normalized to total number of simulated events in the right pannel.




**Figure 7:** The transverse momentum distribution of PYTHIA8 simulations for  $\Omega^-(sss)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for low multiplicity events, less than 200, in the left pannel and for low multiplicity events normalized to total number of simulated events in the right pannel.

The suppression of high  $p_T$  particles identified the collective medium as a thermal medium of QGP [8]. To understand this behavior, we studied PYTHIA 8 simulations for high multiplicity events in pp collisions at STAR energy and draw the results for  $k^+(s\bar{u})$  in Figure 8,  $k^-(u\bar{s})$  in Figure 9, for  $\Lambda^0(uds)$  in Figure 10, for  $\Xi^-(dss)$  in Figure 11, for  $\Xi^0(uss)$  in Figure 12 and  $\Omega^-(sss)$  in Figure 13. All these distributions confirm the suppression behavior for the  $K_s^+(u\bar{s})$ ,  $K_s^-(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^0(uss)$ ,  $\Xi^-(dss)$ ,  $\Omega^-(sss)$  particles which manifests itself as QGP phase formation. Where the plasma present steals the high transverse momentum from the interacting fireball. The azimuthal correlations between produced particles through hard collisions are typically attributed to the formation of a strongly interacting quark gluon medium [1] [23]. For mesons  $K_s^+(u\bar{s})$  and  $K_s^-(s\bar{u})$ , Figure 8 and Figure 9 introduce high multiplicity events, higher than 70, normalized to total number of simulated events distributions.




**Figure 8:** The transverse momentum distribution of PYTHIA8 simulations for  $k^+(u\bar{s})$  particles at  $\sqrt{s_{NN}} = 200$  GeV for high multiplicity events, higher than 50, in the left pannel and for high multiplicity events normalized to total number of simulated events in the right pannel.



**Figure 9:** The transverse momentum distribution of PYTHIA8 simulations for  $k^-(s\bar{u})$  particles at  $\sqrt{s_{NN}} = 200$  GeV for high multiplicity events, higher than 50, in the left pannel and for high multiplicity events normalized to total number of simulated events in the right pannel.


For strange  $\Lambda^0(uds)$ , we study the high multiplicity events, higher than 120, normalized to total number of simulated events transverse momentum distributions and draw them in Figure 10.




**Figure 10:** The transverse transverse momentum distribution of PYTHIA8 simulations for  $\Lambda^0(uds)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for high multiplicity events, higher than 120, in left pannel and or high multiplicity events normalized to total number of simulated events in right pannel.

For events having multiplicity higher than 150 normalized to total number of simulated events,

we introduce transverse momentum distributions of  $\Xi^-(dss)$  and  $\Xi^0(uss)$  confirming overabundant of strangeness. Our results are consistent with the enhanced production of multi-strange hadrons in high-multiplicity events in pp collisions that is confirmed in [1].




**Figure 11:** The transverse momentum distribution of PYTHIA8 simulations for  $\Xi^-(dss)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for high multiplicity events, higher than 150, in the left pannel and for high multiplicity events normalized to total number of simulated events in the right pannel.



**Figure 12:** The transverse momentum distribution of PYTHIA8 simulations for  $\Xi^0(uss)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for high multiplicity events, higher than 150, in the left pannel and for high multiplicity events normalized to total number of simulated events in the right pannel.

An even stronger signature of strangeness enhancement is the highly enhanced production of multi-strange baryons. The production of triple strange  $\Omega^-(sss)$  is the strongest signature to pronounce QGP formation. For high multiplicity events, higher than 200, normalized to total number of simulated events distribution of  $\Omega^-(sss)$ , we draw Figure 13.






**Figure 13:** The transverse momentum distribution of PYTHIA8 simulations for  $\Omega^- (sss)$  particles at  $\sqrt{s_{NN}} = 200$  GeV for high multiplicity events, higher than 200, in left pannel and for high multiplicity events, higher than 200, normalized to total number of simulated events in right pannel.

## Analysis and Results


The PYTHIA 8 (version 8.309) simulation measurements are performed for  $10^5$  pp generated events collected at  $\sqrt{s_{NN}} = 200$  GeV. The particle decay channels in PYTHIA 8 was set to "on" and the particle of interest was set to "off". the events that have total charges equal to zero are removed from the data sets. The results presented here are for the considered particles at events having low multiplicity less than 200 and events having high multiplicity higher than 200. For meson  $k^+(s\bar{u})$  and  $k^-(s\bar{u})$ , the events at low multiplicity, less than 200, show lower production than the events at high multiplicity, higher than 200. The figure Figure 14 finds out the ratio between the  $k^-(s\bar{u})$  and  $k^+(s\bar{u})$  production of events having multiplicity lower than 200 to their production at events having multiplicity higher than 200 is around 1.0. According to [22], the threshold energy creation of kaons is 13.5 GeV and the nature of the strangeness production comes from the color field nature which loses its energy bit by bit as excitations (kinks) of soft gluons rather than by hard single gluon radiation. This means that the soft gluons massively created hadronizing  $k^-(s\bar{u})$  and  $k^+(s\bar{u})$  in low and high multiplicity events; this behavior explains the higher production of  $k^-(s\bar{u})$  and  $k^+(s\bar{u})$  even the pionasatain dominates. Experimentally, the average multiplicity of keon production is 1.47 and pion production is 17.9 for full phase space at  $\sqrt{s_{NN}} = 200\text{GeV}$  [22]. The Wroblewski ratio between kaons to pions is about 12.17.



**Figure 14:** The production ratio for  $k^-(s\bar{u})$ ,  $k^+(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^-(dss)$ ,  $\Xi^0(uss)$  and  $\Omega^-(sss)$ .

For  $\Lambda^0(uds)$  particle, the production from high multiplicity events is higher than the production

from low multiplicity events where the ratio around 1.062. This ratio presents that the strange production mechanism of  $\Lambda^0(uds)$  is the same as  $k^-(s\bar{u})$ . Experimentally, the average multiplicity of lambda is 0.23 for full phase space at RHIC energy [22]. The ratio between high multiplicity events to low multiplicity events is higher than 1, for  $\Xi^-(dss)$  is 1.072 and for  $\Xi^0(uss)$  is 1.097. The Figure 14 confirms that the high production in strangeness for  $\Xi^-(dss)$  and  $\Xi^0(uss)$  for high multiplicity events is more than low multiplicity events. The multi-strange  $\Omega^-(sss)$  shows enhancement of strange quark that is approved in Figure 14. The high multiplicity events to low multiplicity ration is 1.02 that strengthen the result.



**Figure 15:** The relation between the event multiplicity and the strangeness content.

We find out that in pp collisions, the strange and multi-strange particle yield increases with the event multiplicity as proved in Figure 15. The observations we confirmed agreed with The ALICE researchers found that in proton–lead ion collisions, the relative yield of particles that contain strange quarks increases with event multiplicity [7].

## Conclusion

At the RHIC energy, the observed strange yield enhancements increase with the strangeness contained in hadrons. There is a linear relation between the event multiplicity and the strangeness content of the produced particles as Figure 15 demonstrated. The suppression of the  $p_T$  spectrum at high  $p_T$  is observed for  $k^-(s\bar{u})$ ,  $k^+(s\bar{u})$ ,  $\Lambda^0(uds)$ ,  $\Xi^-(dss)$ ,  $\Xi^0(uss)$  and  $\Omega^-(sss)$  for high and low multiplicity events. For  $k^-(s\bar{u})$  and  $k^+(s\bar{u})$  meson, the low multiplicity events give less strange production than the high multiplicity events. For  $\Lambda^0(uds)$ ,  $\Xi^-(dss)$ ,  $\Xi^0(uss)$  and  $\Omega^-(sss)$  baryons, the high multiplicity events enhance the production of multi-strange quarks than low multiplicity events. In high multiplicity events in pp collisions, the strangeness production reaches values similar to those observed in nucleus-nucleus collisions, where a QGP is formed [1]. We conclude that pp collisions at RHIC energy produce QGP at low multiplicity events, less than 70 and high multiplicity events, higher than 200.

## References

- [1] “Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions”. In: *Nature Physics* 13.6 (2017), pp. 535–539.
- [2] Sarita Sahoo et al. “Enhanced Production of Multi-strange Hadrons in Proton-Proton Collisions”. In: *Springer Proc. Phys.* 261 (2021). Ed. by Prafulla Kumar Behera et al., pp. 1043–1046. DOI: 10.1007/978-981-33-4408-2\_152.

- [3] Livio Bianchi, ALICE collaboration, et al. “Strangeness production as a function of charged particle multiplicity in proton–proton collisions”. In: *Nuclear Physics A* 956 (2016), pp. 777–780.
- [4] Johann Rafelski and Berndt Müller. “Strangeness production in the quark gluon plasma”. In: *Physical Review Letters* 48.16 (1982), p. 1066.
- [5] Johann Rafelski. “Strangeness in Hadronic Matter”. In: (1995).
- [6] Barbara Jacak and Peter Steinberg. “Creating the perfect liquid in heavy-ion collisions”. In: *Physics today* 63.BNL-93753-2010-JA (2010).
- [7] Sung Chang. “When proton–proton collisions turn strange”. In: (2017).
- [8] Samrangy Sadhu and Premomoy Ghosh. “Anomalous features of particle production in high-multiplicity events of p p collisions at the LHC energies”. In: *Physical Review D* 99.3 (2019), p. 034020.
- [9] BI Abelev et al. “Enhanced strange baryon production in Au–Au collisions compared to p–p at  $\sqrt{s_{NN}} = 200$  GeV”. In: *Physical Review C* 77.4 (2008), p. 044908.
- [10] KH Ackermann et al. “The STAR time projection chamber”. In: *Nuclear Physics A* 661.1–4 (1999), pp. 681–685.
- [11] Johann Rafelski and Jean Letessier. “Strangeness and the discovery of quark gluon plasma”. In: *Journal of Physics: Conference Series*. Vol. 50. 1. IOP Publishing. 2006, p. 176.
- [12] Johann Rafelski and Jean Letessier. “Strangeness and quark gluon plasma”. In: *Journal of Physics G: Nuclear and Particle Physics* 30.1 (2003), S1.
- [13] Francesco Becattini and Michael A Lisa. “Polarization and vorticity in the quark gluon plasma”. In: *Annual Review of Nuclear and Particle Science* 70 (2020), pp. 395–423.
- [14] Mahmoud Hanafy, Omnia SA Qandil, and Asmaa G Shalaby. “Strangeness Enhancement at LHC Energies Using the Thermal Model and EPOS LHC Event Generator”. In: *Advances in High Energy Physics* 2021 (2021), pp. 1–13.
- [15] Johann Rafelski. *Strangeness in quark gluon plasma*. Tech. rep. 1982.
- [16] Pasi Huovinen. “Chemical freeze-out temperature in the hydrodynamical description of Au+ Au collisions at  $\sqrt{s_{NN}} = 200$  GeV”. In: *The European Physical Journal A* 37.1 (2008), pp. 121–128.
- [17] ZJ He et al. “Strangeness in a chemically equilibrating quark gluon plasma”. In: *Physical Review C* 69.3 (2004), p. 034906.
- [18] Betty Bezverkhny Abelev et al. “Multiplicity Dependence of Pion, Kaon, Proton and Lambda Production in p–Pb Collisions at  $\sqrt{s_{NN}} = 5.02$  TeV”. In: *Phys. Lett. B* 728.arXiv: 1307.6796 (2014), pp. 25–38.
- [19] Dott Francesco Noferini and Francesca Ercolessi. “Strangeness enhancement with multiplicity and effective energy in pp collisions at  $\sqrt{s} = 13$  TeV with the ALICE detector at the LHC”. In: ().
- [20] Raghunath Sahoo. “Possible formation of QGP-droplets in proton-proton collisions at the CERN Large Hadron Collider”. In: *arXiv preprint arXiv:1908.10566* (2019).
- [21] Marius Ole Johansen. “Heavy flavor azimuthal correlations in PYTHIA generated proton-proton collisions at LHC energies”. MA thesis. The University of Bergen, 2014.
- [22] M. T. Ghoneim, F. H. Sawy, and H. T. Hussein. “Particle production in proton-proton collisions”. In: *International Journal of Physical Sciences* (2015).
- [23] Krzysztof W Wozniak, ATLAS Collaboration, et al. “Study of the long-range azimuthal correlations in pp and p Pb collisions with the ATLAS detector at the LHC”. In: *Journal of Physics: Conference Series*. Vol. 779. 1. IOP Publishing. 2017, p. 012057.