A Worldsheet Derivation of the Classical Off-shell Boundary Action for the Dilaton in Half-Space

Amr Ahmadain,^a Rifath Khan^b

^a Department of Physics, Swansea University, Swansea, SA2 8PP, UK ^b Stanford Institute for Theoretical Physics, 382 Via Pueblo, Stanford, CA 94305

E-mail: amrahmadain@gmail.com, rifathkhantheo@gmail.com

ABSTRACT: We use the method of images to present a worldsheet derivation of the sphere partition function for the dilaton in half-space to leading order in α' with Neumann boundary conditions. We use Tseytlin's sphere prescription to obtain the total (bulk and boundary) off-shell classical bosonic string action for the dilaton in half-space and show that it satisfies the requirement for a well-defined variational principle.

Contents

1	Introduction	1
	1.1 Related Work	2
	1.2 Strategy and summary of results	4
2	The sphere partition function in $\mathbb{R}_+ \times \mathbb{R}^{D-1}$	7
	2.1 The method of images	8
3	Setup of calculation	9
4	The classical boundary dilaton action in half-space	11
	4.1 The classical on-shell action	15
5	Discussion	16
	5.1 The probabilistic interpretations of boundary conditions	18
	5.2 Reflected Brownian motion, local boundary time and running maximum	20
	5.3 The probabilistic nature of the $\left\langle \left Y^D + \sqrt{\alpha'} \eta^D(z) \right \right\rangle$	21
	5.4 Outlook and Future Directions	23
A	The computation of $\left\langle \left Y^D + \sqrt{lpha'} \eta^D(z) \right \right\rangle$	24

1 Introduction

The sphere partition function is a mysterious yet important object in string theory. Generally speaking, a worldsheet path integral calculation of the sphere partition function should produce the classical (tree-level) *bulk* gravitational effective action, I_{bulk} , to all orders in α' , in addition to all boundary terms, I_{bdy} , which are required to have a sound variational principle [1]

$$I_{\rm sphere} = I_{\rm bulk} + I_{\rm bdy}.\tag{1.1}$$

 I_{bulk} has been successfully derived from the worldsheet to leading order in α' [2–4] using Tseytlin's off-shell sphere prescription [5] and it was found to vanish on solutions of the equations of motion, i.e. on-shell. To date, as far as the authors know, there is no consistent way to derive or extract the boundary action I_{bdy} from the worldsheet and Tseytlin's off-shell prescriptions are not successful either in doing that.

This paper takes a very first step in that direction. We use Tseytlin's off-shell sphere prescription [4–6] and the method of images to evaluate the *off-shell* classical bosonic string action for the dilaton in half-space. We impose Neumann boundary conditions on the dilaton at the codimension-1 boundary (wall) and obtain the dilaton boundary term. We then show that the total (bulk plus boundary) off-shell action we derive satisfies the requirement for a well-defined variational principle.

Concretely, we take our target space \mathcal{M} to be $\mathbb{R}_+ \times \mathbb{R}^{D-1}$. The total off-shell classical string action that we derive from the worldsheet is the sum of the bulk (kinetic) and boundary action for the dilaton

$$I_{\rm sphere} = -\tilde{Z}_{\rm nz} \alpha' \int_{\mathcal{M}} d^D Y e^{-2\Phi} \partial^2 \Phi + \alpha' \tilde{Z}_{\rm nz} \int_{\partial \mathcal{M}} d^{D-1} Y \ e^{-2\Phi} \partial_n \Phi \,, \tag{1.2}$$

where $\partial_n \Phi$ is the normal derivative of Φ to the boundary. I_{bulk} does not have a well-posed variational principle since it is second order in derivatives. I_{bdy} was added by hand [1] such that I_{sphere} has a well-defined variational principle in target spaces with boundaries. In this paper, we however provide an exact first-principles worldsheet derivation of I_{bdy} . Equivalently, integrating I_{bulk} by parts gives the standard dilaton kinetic term $2e^{-2\Phi}(\partial\Phi)^2$ in addition to $-I_{\text{bdy}}$. To be precise, by off-shell, we mean the nonconstant dilaton $\Phi(Y)$ is *not* constrained to satisfy any equation of motion, bulk or boundary, during any step of our derivation. It is only in the last step that we impose the equations of motion to get the classical on-shell action.

It is also important to define what we mean by a Neumann boundary condition and more importantly, when, i.e. at which step of the derivation we impose it on $\Phi(Y)$. Physically, a Neumann boundary condition means having a constant flux at the boundary, i.e. $\partial_n \Phi = \text{constant}$. In this paper, we impose the condition $\partial_n \Phi = 0$ only after as a boundary equation of motion but not while varying the total action.

1.1 Related Work

Below, we give a quick overview of Tseytlin's off-shell prescription and summarize efforts to derive both terms in I_{sphere} from the worldsheet.

It is a standard result in string theory that the classical bulk action I_{bulk} (tree-level cosmological constant), the one-point function (tree-level tadpole) and two-point function¹ all vanish on-shell (on a string background defined by a CFT) in *compact* target spacetime [8]. From the worldsheet perspective, the reason for this is simple. It is because after fixing the Diff and Weyl gauge symmetries of the genus-0 (sphere) worldsheet theory, one still has to divide by the infinite volume of the noncompact $SL(2,\mathbb{C})$. As a result, the on-shell sphere partition function with zero number of insertions vanishes

$$-I_{\text{bulk}} = Z_{\text{CFT}} = \frac{1}{\text{SL}(2,\mathbb{C})} Z_{\text{ghost}} Z_{\text{matter}}[\hat{g}] = \frac{K_0}{\infty}, \qquad (1.3)$$

where K_0 is the genus-0 partition function without the CKG factor and \hat{g} is a gauge-fixed metric.

Within his first-quantized off-shell nonlinear sigma model (NSLM) formalism [9–11], Tseytlin made two proposals [5, 12, 13] to obtain the *bulk* classical off-shell action from the worldsheet and address the noncompactess of the SL(2, \mathbb{C}) group in the denominator of the sphere partition function. The regularized volume of SL(2, \mathbb{C}) has a logarithmically divergent piece [5, 6, 8, 14]. At the heart of both proposals, Tseytlin integrates over all *n* vertex operator insertions and produces a logarithmically divergent term to cancel the log divergence of SL(2, \mathbb{C}) in (1.3). In the integrated vertex operator formalism, this directly leads to different types of divergences, logarithmic and power-law, in the *n*point correlation function $K_{0,n}$ as n, n-1 or fewer operators approach each other in the integration region respectively. Regularizing these divergences require introducing a UV length cutoff ϵ on the worldsheet. To obtain the off-shell sphere partition function (the classical action), Tseytlin takes the

¹Except for a string that comes in and back out without no interactions [7]. In this case, there is an additional delta function in the numerator of the connected S-matrix that causes the two-function to diverge.

derivative of the log of the UV cutoff with respect to K_0 . This is Tseytlin's first prescription²

$$-I_{\text{bulk}} = Z_{\text{QFT}} = \frac{\partial}{\partial \log \epsilon} K_0.$$
(1.4)

In the NLSM, it is possible to find a renormalization group scheme where, to an arbitrary order in α' , the QFT sphere partition function is defined to be

$$K_0 = V_{\text{gen}} \coloneqq \int_{\mathcal{M}} d^D Y e^{-2\Phi}$$
(1.5)

where V_{gen} is the generalized volume factor [1, 6, 16].

In his first-quantized off-shell string theory formalism, Tseytlin breaks the sacred local conformal invariance of only the matter part worldsheet NLSM.³ As a result, all calculable quantities, e.g. correlation functions, in the now worldsheet QFT will depend on the choice of the Weyl frame ω when the worldsheet metric is expressed as $g_{ab} = e^{2\omega}\gamma_{ab}$. In QFT, breaking a local gauge symmetry is catastrophic since it immediately renders the underlying theory inconsistent. But it was shown in [6] that infinitesimal changes in ω on the worldsheet are equivalent to field redefinitions in target space, which corresponds to RG flow of the QFT. The conceptual foundations of Tseytlin's off-shell formalism and sphere prescriptions proposed in [5, 12, 13] have been studied, analyzed and generalized in [6].

This is the story in compact spacetime. However, in noncompact target spacetime (those with boundaries), there are subtleties [17]. To begin with, it is not yet known how to use Tseytlin's off-shell sphere prescription (1.3) to obtain I_{bdy} directly from the worldsheet [1]. As far as the authors know, there is no consistent way yet to derive or extract the boundary action I_{bdy} from the worldsheet.

As we have discussed above, from the worldsheet perspective, the classical closed string off-shell action vanishes on-shell due to the infinite volume of $SL(2, \mathbb{C})$. However, from the spacetime point of view, there is another reason why the classical string action is zero on-shell modulo a boundary term. It is because the nonconstant dilaton in the classical action transforms in spacetime in such a way that only changes the overall normalization of the action. This property is also true for the closed string field theory action. [18]⁴ and to at any order in α' in perturbative string theory [20].

In noncompact target spaces, the action is only stationary with respect to normalizable deformations of the CFT defining the on-shell string background but not for nonnormalizable deformations that survive at infinity. As pointed out in [21], a constant shift of the dilaton is not a variation of this form and thus, can receive nonzero boundary contributions from nonnormalizable modes in a noncompact CFT [17]. In the AdS₃, this also leads to a non-zero one-point function [22].

A peculiarity of the classical string action is that the generalized volume factor $e^{-2\Phi}$ is the only term that depends on the nonconstant dilaton $\Phi(Y)$ and does not include derivatives

$$I_{\text{bulk}} = \frac{1}{16\pi G_N} \left[\int_{\mathcal{M}} d^D Y \sqrt{G} e^{-2\Phi} \mathcal{L}(\partial_\mu \Phi, \phi_i) \right], \qquad (1.6)$$

$$-I_0^{\mathbf{T2}} = Z_0^{\mathbf{T2}} = \left(\frac{\partial}{\partial \log \epsilon} + \frac{1}{2} \frac{\partial^2}{(\partial \log \epsilon)^2}\right) K_0.$$
 (**T2**)

²The prescription (1.4) does not give the correct string vacuum for tachyons; it does not remove the tachyon tadpole. To deal with this problem, Tseytlin proposed his second prescription [12, 15]

³Not all of $SL(2,\mathbb{C})$ is broken though. Only local Weyl invariance is broken whereas translation and special conformal symmetries are left intact. The *bc* ghost sector of the off-shell sphere partition function is never deformed away from the CFT in Tseytlin's off-shell formalism.

⁴Deriving the on-shell action from the classical string field theory action is still a challenge (see section 10 in [19]).

where \mathcal{L} depends only on $\partial_{\mu}\Phi$ and some other fields ϕ_i (but not on Φ itself). This feature of the classical action leads to an interesting consequence: the dilaton equation of motion implies the Lagrangian is a total derivative, and hence the on-shell action is a boundary term

$$-2e^{-2\Phi}\mathcal{L} = \partial_{\mu} \left(e^{-2\Phi} \frac{\delta \mathcal{L}}{\delta \partial_{\mu} \Phi} \right).$$
(1.7)

For an asymptotically flat boundary, where gravity is weakly coupled, the classical string action to first order in α' , including the Gibbons-Hawking-York boundary term [23, 24] is given by

$$I_{\text{sphere}} = \frac{1}{16\pi G_N} \left[\int_{\mathcal{M}} d^D Y \sqrt{G} e^{-2\Phi} \left(-R - 4(\nabla \Phi)^2 + \cdots \right) - 2 \int_{\partial \mathcal{M}} d^{D-1} Y \sqrt{h} e^{-2\Phi} K \right], \qquad (1.8)$$

in which case $I_{\rm bdy}$ becomes

$$-I_{\rm bdy} = \frac{1}{8\pi G_N} \int_{\partial \mathcal{M}} d^{D-1} Y \,\partial_n \left(e^{-2\Phi} \sqrt{h} \right). \tag{1.9}$$

In this paper, we give an off-shell worldsheet derivation of I_{bdy} in half-space (in flat spacetime, so without the GHY boundary term.) and impose Neumann boundary conditions on $\Phi(Y)$ at the codimension-1 wall.

The sphere and disk partition functions have been recently calculated for different backgrounds using methods different from those we use in this paper. The string partition function on the disk (for the open string) has been calculated in [25] where it was shown to give a finite nonzero value, contrary to the naive expectation that it vanishes due to the division by the infinite volume of the PSL(2, \mathbb{R}) group [8]. The sphere partition function has been calculated in minimal models [26] and shown to have unambiguous finite values and in two-dimensional quantum gravity [27]. More recently, the sphere partition function for global AdS₃ with pure NS flux has been studied [28] and found to match the conformal anomaly of the boundary holographic CFT as it should only in the supersymmetric string. Because of the tachyon in bosonic string theory, a mismatch was found.

1.2 Strategy and summary of results

In this paper, we do not derive the on-shell classical boundary action from the worldsheet for general noncompact target spaces, e.g. cigar or $AdS_3 \times S^3 \times X$ with pure NS-NS flux, but we take an initial step in this direction. We use Tseytlin's sphere prescription to derive, from the worldsheet, the *off-shell* classical boundary action for the dilaton in half-space (with a flat target spacetime metric) with Neumann boundary conditions.

We take our target space \mathcal{M} be the half-space $\mathbb{R}_+ \times \mathbb{R}^{D-1}$ with coordinates $X^{\mu} = (X^i, X^D)$ where $X^i \in (1, \dots D-1)$ are coordinates along a codimension-1 wall at $X_D = 0$. Strings in half-space is a \mathbb{Z}_2 orbifold of flat spacetime since $\mathbb{R}_+ = \mathbb{R}/\mathbb{Z}_2$ [29]. In this half-space, the spherical worldsheet is *restricted* to the right of the hard wall (boundary) with coordinates $X^D \ge 0$ that splits the entire spacetime into two halves.

Our goal is to compute sphere partition function in half-space Z_{HS} . We use the method of images to evaluate it in a *reflected* (doubled) space with a reflected dilaton $\tilde{\Phi}$ and impose Neumann boundary condition on $\tilde{\Phi}$ on the codimension-1 wall (the orbifold fixed point) at $X_D = 0$ such that Z_{HS} can be expressed as $\frac{1}{2} Z_{\rm RS}$

$$Z_{\rm HS} = \int_{X^D \ge 0} [DX] e^{-I_{\rm NLSM}[X, \Phi, \omega, \gamma]}$$

$$= \frac{1}{2} Z_{\rm RS} = \frac{1}{2} \int [DX] e^{-I_{\rm NLSM}[X, \tilde{\Phi}, \omega, \gamma]}$$
(1.10)

where the dilaton $\tilde{\Phi}$ in reflected space is an infinitesimal deformation about the constant mode Φ_0

$$\tilde{\Phi}(X^{\mu}) = \Phi_0 + \phi(X^i, X^D) \Theta(X^D) + \phi(X^i, -X^D) \Theta(-X^D).$$
(1.11)

As in [2, 4], we use heat kernel regularization to regulate divergences in Z_{HS} and [DX] by a inserting a factor of $e^{\epsilon \Delta}$ into divergent expressions, with $\Delta = -\nabla^2$.

We decompose the string into a zero mode Y^{μ} and a nonzero mode $\eta^{\mu}(z)$. Because of the kink in $\Phi(Y^D)$ at the wall, we find that $\langle |\eta^D(z)| \rangle$ does not vanish as it always does in the NLSM expansion of deformations around *compact* CFT with no target space boundary. We show that the near-wall (in terms of $\sqrt{\alpha'}$) part of $Z_{\rm HS}$ is given by integral of the one-point function $\langle |Y^D + \sqrt{\alpha'}\eta^D(z)| \rangle$ over the zero mode of the string Y^D normal to the boundary in target spacetime

$$Z_{\text{near}} := \frac{1}{2} Z_{\text{nz}} \int_{\frac{Y^D}{\sqrt{\alpha'}} \sim 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left(\phi \left(Y^i, 0 \right) + O(\alpha') \right) \right) + Z_{\text{wall}}, \tag{1.12}$$

where Z_{nz} is the nonzero mode piece and Z_{wall} is (after taking some appropriate limits) is given by

$$Z_{\text{wall}} = \frac{1}{2} \times 2Z_{\text{nz}} \int_{\frac{Y^D}{\sqrt{\alpha'}} \sim 1} d^D Y \partial_D \phi \left(Y^i, 0 \right) \left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle$$
(1.13)

$$= -\frac{1}{2} \times 2\tilde{Z}_{\mathrm{nz}} \,\alpha' \int d^{D-1}Y \, e^{-2\Phi} \partial_D \Phi\left(Y^i, 0\right) \left(\frac{\alpha' Y_c^2}{4} + \frac{-2\ln\epsilon}{2}\right). \tag{1.14}$$

We then use Tseytlin's prescription (1.4) to obtain the classical boundary action for the dilaton in half-space

$$I_{\text{wall}} = -\frac{\partial}{\partial \log \epsilon} Z_{\text{wall}} = -\alpha' \tilde{Z}_{\text{nz}} \int d^{D-1} Y \ e^{-2\Phi} \partial_D \Phi \left(Y^i, 0 \right) = +\alpha' \tilde{Z}_{\text{nz}} \int d^{D-1} Y \ e^{-2\Phi} \partial_n \Phi \left(Y^i, 0 \right),$$
(1.15)

where ∂_n is the normal derivative in the direction of the outward-pointing normal n^{μ} i.e. $\partial_n = -\partial_D$.

Next, we again use Tseytlin's prescription to obtain the *bulk* dilaton action in half-space from Z_{far} , the sphere partition function in the target spacetime region far from the wall

$$I_{\text{bulk}} := I_{\text{far}} = -\frac{\partial}{\partial \log \epsilon} Z_{\text{far}} = -\tilde{Z}_{\text{nz}} \alpha' \int_{\frac{Y^D}{\sqrt{\alpha'}} \gg 1} d^D Y e^{-2\Phi} \partial^2 \Phi$$
(1.16)

In spaces with boundaries, such as half-space, I_{bulk} does not have a well-defined variational prin-

ciple, since varying it yields

$$\delta I_{\text{bulk}} = 2\alpha' \tilde{Z}_{\text{nz}} \int_{\frac{Y^D}{\sqrt{\alpha'}} \gg 1} d^D Y \left[e^{-2\Phi} \partial^2 \Phi + \partial_\mu \left(e^{-2\Phi} \partial^\mu \Phi \right) \right] \delta \Phi$$
$$- \alpha' \tilde{Z}_{\text{nz}} \int_{\partial \mathcal{M}} d^{D-1} Y e^{-2\Phi} \,\delta(\partial_n \Phi) - 2\alpha' \tilde{Z}_{\text{nz}} \int_{\partial \mathcal{M}} d^{D-1} Y e^{-2\Phi} \partial_n \Phi \,\delta \Phi. \tag{1.17}$$

The first boundary term in I_{bulk} spoils the variational principle. It is the GHY-like term of the free two-derivative action which must be accounted for, i.e. by adding an *additional* boundary term whose variation exactly cancels with it to have a well-posed variational principle. This additional boundary term is I_{wall} .

Hence, the total classical string action in half-space, that we derive from the worldsheet in this paper, is the sum of the bulk and boundary parts for the dilaton

$$I_{\text{sphere}} := I_{\text{HS}} = I_{\text{bulk}} + I_{\text{wall}}$$
$$= -\tilde{Z}_{nz} \alpha' \int_{\frac{YD}{\sqrt{\alpha'}} \gg 1} d^D Y e^{-2\Phi} \partial^2 \Phi + \alpha' \tilde{Z}_{nz} \int d^{D-1} Y \ e^{-2\Phi} \partial_n \Phi \left(Y^i, 0 \right). \tag{1.18}$$

Varying I_{sphere} gives

$$\delta I_{\text{sphere}} = \delta I_{\text{bulk}} + \delta I_{\text{wall}}$$
$$= 2\alpha' \tilde{Z}_{\text{nz}} \int_{\frac{Y^D}{\sqrt{\alpha'}} \gg 1} d^D Y \left[e^{-2\Phi} \partial^2 \Phi + \partial_\mu \left(e^{-2\Phi} \partial^\mu \Phi \right) \right] \delta \Phi$$
(1.19)

$$-\alpha'\tilde{Z}_{nz}\int_{\partial\mathcal{M}}d^{D-1}Ye^{-2\Phi}\,\delta(\partial_n\Phi) + \alpha'\tilde{Z}_{nz}\int_{\partial\mathcal{M}}d^{D-1}Ye^{-2\Phi}\,\delta(\partial_n\Phi) \tag{1.20}$$

$$-4\alpha'\tilde{Z}_{nz}\int_{\partial\mathcal{M}}e^{-2\Phi}\partial_n\Phi\,\delta\Phi\tag{1.21}$$

$$= 2\alpha' \tilde{Z}_{\rm nz} \int_{\frac{YD}{\sqrt{\alpha'}} \gg 1} d^D Y \left[e^{-2\Phi} \partial^2 \Phi + \partial_\mu \left(e^{-2\Phi} \partial^\mu \Phi \right) \right] \delta \Phi - 4\alpha' \tilde{Z}_{\rm nz} \int_{\partial \mathcal{M}} e^{-2\Phi} \partial_n \Phi \,\delta \Phi. \quad (1.22)$$

In the second line, we observe that the variation δI_{wall} exactly cancels its counterpart in δI_{bulk} i.e. $e^{-2\Phi} \,\delta(\partial_n \Phi)$ such that we are left in the end with only the bulk and boundary dilaton equations of motion. Therefore, I_{sphere} has a well-defined variational principle only with the addition of I_{wall} as a boundary term.

We then impose the *bulk* dilaton equation of motion in I_{sphere} to obtain the classical on-shell (in the bulk) action for the dilaton in half-space

$$I_{\rm sphere} := I_{\rm HS} = I_{\rm bulk} + I_{\rm wall} \tag{1.23}$$

$$= 2 \alpha' \tilde{Z}_{nz} \int d^{D-1} Y e^{-2\Phi} \partial_n \Phi = 2I_{\text{wall}} .$$
 (1.24)

In this paper, we were inspired by the work in [30–32] which employ the apparatus of the method of images within the worldline formalism to calculate particle path integral in $\mathbb{R}_+ \times \mathbb{R}^{D-1}$ using heat kernel methods.

Plan of paper: In section 2, we present the conceptual foundations of calculating $Z_{\rm HS}$ and $I_{\rm HS}$ using the method of images with Neumann boundary conditions imposed on the dilaton at the wall.

We will comment in this section on how to still use the method of images to impose Dirichlet boundary conditions on target space fields. We will have more to say about this in the discussion section.

In section 3, we setup the calculation that we do in section 4. In appendix A, we present the details of computing the one-point function $\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle$. We first expand $\eta^D(z)$ and the Laplacian in real spherical harmonics and then constrain the path integral by inserting a delta function to compute $\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle$.

In section 4, we finally put things together to arrive at the central result of this work. We write down the expression for I_{sphere} , the total off-shell classical action in half-space and demonstrate that it has a well-posed variational principle.

Finally, in section 5, we highlight the relationship between $\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z) \right| \right\rangle$ and the important concept of boundary local time $\left\langle L_{t}^{0} \right\rangle$ in *reflected* Brownian motion. Specifically, we show that $\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z) \right| \right\rangle$ and $\left\langle L_{t}^{0} \right\rangle$ are equal in distributional law with both quantities having the halfnormal probability distribution. In this probabilistic sense, we argue that the one-point function $\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z_{0}) \right| \right\rangle$ can be interpreted as a measure of the average distance (length) of the string near (a few string-length away from) the wall taking all reflection physics into account.

We end with an outlook on future directions and open questions.

2 The sphere partition function in $\mathbb{R}_+ \times \mathbb{R}^{D-1}$

Let our target space \mathcal{M} be the half-space $\mathbb{R}_+ \times \mathbb{R}^{D-1}$ with coordinates X^{μ} . Half-space is a \mathbb{Z}_2 orbifold of flat spacetime since $\mathbb{R}_+ = \mathbb{R}/\mathbb{Z}_2$ [29]. A co-dimension one wall sits at $X^D = 0$ and splits \mathcal{M} into two halves. In this half-space, the spherical worldsheet will be *restricted* to the right of the wall with coordinates $X^D \ge 0$.

Consider the worldsheet NLSM path integral in $\mathbb{R}_+ \times \mathbb{R}^{D-1}$ given by the off-shell sphere partition function

$$Z_{\rm HS}[G,\Phi,\omega] = \int_{X^D \ge 0} \frac{[D\gamma][DX]}{\text{Diff}} e^{-I_{\rm NLSM}[X,G,\Phi,\omega,\gamma]},\tag{2.1}$$

where γ is the conformal worldsheet metric, ω is the worldsheet conformal factor, and X are the embedding fields on the worldsheet. The path integral with the restriction $X^D \ge 0$ ensures that the worldsheet lives only on one (the right) side of the wall. For a sphere, the conformal metric is the metric on a round sphere.

In this work, we take the target spacetime metric to be flat, and the dilaton field is expanded about a constant mode⁵

$$G_{\mu\nu}(X) = \delta_{\mu\nu}, \quad \Phi(X) = \Phi_0 + \phi(X), \quad \text{and} \quad \phi(X) \ll 1$$
 (2.2)

Here, the function $\Phi(X)$ is defined only for $X^D \ge 0$, on the right side of the wall.

If we could find a way evaluate $Z_{\rm HS}$ with the restriction $X^D \ge 0$ i.e. restricting the worldsheet to the right side of the wall, then we would obtain the classical boundary action for the dilaton. Evaluating the worldsheet NLSM path integral with this restriction to half-space is highly non-trivial. To do this, we combine the methods of off-shell string theory studied in [4, 6] with the method of images to evaluate the NLSM worldsheet string path integral in $\mathbb{R}_+ \times \mathbb{R}^{D-1}$.

⁵The mass dimensions of the metric and dilaton are [G] = 0, $[\Phi] = 0$, and $[\alpha'] = -2$.

Precisely, we use the method of images to evaluate the sphere partition function in a *reflected* (doubled) space with a reflected dilaton $\tilde{\Phi}$ defined in (1.11) and impose Neumann boundary condition on $\tilde{\Phi}$ at $X_D = 0$ such that $Z_{\rm HS}$ can be expressed as $\frac{1}{2}Z_{\rm RS}$

$$Z_{\rm HS} = \int_{X^D \ge 0} [DX] e^{-I_{\rm NLSM}[X, \Phi, \omega, \gamma]}$$

$$= \frac{1}{2} Z_{\rm RS} = \frac{1}{2} \int [DX] e^{-I_{\rm NLSM}[X, \tilde{\Phi}, \omega, \gamma]}$$
(2.3)

We explain next how to do this using the method of images.

2.1 The method of images

The quantity we want calculate is

$$Z_{\rm HS} = \int_{X^D \ge 0} [DX] e^{-I_{\rm NLSM}[X, \Phi, \omega, \gamma]}.$$
(2.4)

This is the path integral of embeddings of the spherical worldsheet in half-space i.e. on the right side of the wall $(X^D \ge 0)$. Let \mathcal{C} be the set of all embeddings of the worldsheet on the right side of the wall. We aim to classify this set first by the number of bounces of the worldsheet off the wall. An example of an element of this set is an embedding for which no points on the worldsheet touch the wall (i.e. $X^D > 0$ everywhere on the worldsheet). Denote this class of embeddings as \mathcal{C}_0 (see fig. 1). There is also a class of embeddings in which the worldsheet bounces off the wall *once*, by which we mean that the worldsheet intersects the wall in one circle at the wall, but the worldsheet doesn't pass through it. Denote this class as \mathcal{C}_1 . The worldsheet can also bounce twice off the wall, i.e. the worldsheet intersects the wall in two circles. Denote the class of all embeddings based on the number of bounces off the wall by \mathcal{C}_n for n bounces.

However, this is not all of the embeddings one can consider. For example, one can also consider an embedding which intersects the wall in only one point, or in a disc. However, all such embeddings would be a measure zero set in the path integral and so can be thrown away. So all the embeddings that give a nonzero contribution can be classified by the number of bounces off the wall.

It is important to note that Whenever the worldsheet bounces from the wall, there is a contribution to the path integral. This contribution depends on the boundary conditions. For Dirichlet boundary conditions, it comes with a (-) sign while for Neumann boundary conditions, it comes with a (+) sign [30-32]. We will be choosing Neumann boundary condition in this work.

Consider the embedding that bounces off the wall once. The worldsheet intersects the wall in a circle. The action being an integral on the worldsheet can be split into two pieces, each for a piece emanating from the circle. Now, if we reflect the spacetime and a single piece, the action remains the same. So we get a embedding in the reflected spacetime that crosses the wall once. But, there are exactly two embeddings in the reflected target space that correspond to a single embedding in half space. The same argument holds for any number of bounces (including zero bounces). Therefore, we can express the path integral of embeddings in half-space as one half the path integral in the reflected (doubled) space. We will explore in a bit more detail the relationship between half-space and reflected space in light of reflected Brownian motion and stochastic dynamics in section 5.



Figure 1: Method of images in target spacetime with Neumann boundary conditions. The partition function in half-space $Z_{\rm HS}$ is one-half its counterpart in reflected (doubled) space $Z_{\rm RS}$.

$$Z_{\rm HS} = \int_{X^D \ge 0} [DX] e^{-I_{\rm NLSM}[X, \Phi, \omega, \gamma]}$$

$$= \frac{1}{2} Z_{\rm RS} = \int \frac{1}{2} [DX] e^{-I_{\rm NLSM}[X, \tilde{\Phi}, \omega, \gamma]}$$
(2.5)

Next, we explain the set-up of the calculation of $Z_{\rm RS}$.

3 Setup of calculation

In terms of the NLSM worldsheet action, both the metric and dilaton are reflected *about* the wall to the other side. The spacetime metric is flat *everywhere*, but the dilaton in this reflected space $\tilde{\Phi}(X^{\mu})$ is defined for $X^D \in \mathbb{R}$. More concretely, $\tilde{\Phi}(X^{\mu})$ is the sum of two terms, each takes values on one side of the wall

$$\tilde{\Phi}(X^{\mu}) = \Phi_0 + \tilde{\phi}(X^{\mu}) = \Phi(X^i, X^D)\Theta(X^D) + \Phi(X^i, -X^D)\Theta(-X^D)$$

$$= \Phi_0 + \phi(X^i, X^D)\Theta(X^D) + \phi(X^i, -X^D)\Theta(-X^D).$$
(3.1)

To carry out the NLSM expansion, we split the string X^{μ} into a zero mode Y^{μ} and a non-zero mode $\eta^{\mu}(z)$

$$X^{\mu}(z) = Y^{\mu} + \sqrt{\alpha'} \eta^{\mu}(z).$$
 (3.2)

The deformed NLSM action is given by

$$I_{\rm NLSM} = \frac{1}{4\pi\alpha'} \int d^2 z \sqrt{g} \left(g^{ab} \partial_a X^{\mu} \partial_b X^{\nu} \delta_{\mu\nu} + \alpha' R^{(2)} \tilde{\Phi}(X) \right)$$
(3.3)

$$=\frac{1}{4\pi\alpha'}\int d^2z\sqrt{g}\left(\alpha'g^{ab}\partial_a\eta^\mu\partial_b\eta^\nu\delta_{\mu\nu} + \alpha'R^{(2)}\tilde{\Phi}\left(Y+\sqrt{\alpha'}\eta\right)\right)$$
(3.4)

$$=\frac{1}{4\pi\alpha'}\int d^2z\sqrt{g}\left(\alpha'g^{ab}\partial_a\eta^{\mu}\partial_b\eta^{\nu}\delta_{\mu\nu}+\alpha'R^{(2)}\Phi_0\right)+\frac{1}{4\pi\alpha'}\int d^2z\sqrt{g}\left(\alpha'R^{(2)}\tilde{\phi}(Y+\sqrt{\alpha'}\eta)\right)$$
(3.5)

$$=\frac{1}{4\pi\alpha'}\int d^2z\sqrt{g}\left(\alpha'g^{ab}\partial_a\eta^{\mu}\partial_b\eta^{\nu}\delta_{\mu\nu}\right)+2\Phi_0+\frac{1}{4\pi\alpha'}\int d^2z\sqrt{g}\left(\alpha'R^{(2)}\tilde{\phi}(Y+\sqrt{\alpha'}\eta)\right)$$
(3.6)

$$= I_{\rm P}[\eta, g] + V[\eta, g, \phi], \tag{3.7}$$

where

$$V[\eta, g, \tilde{\phi}] = \frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \tilde{\phi}(Y + \sqrt{\alpha'} \eta(z)).$$
(3.8)

The path integral measure is given by

$$[DX] = d^D Y [D\sqrt{\alpha'}\eta] \,. \tag{3.9}$$

and the matter partition function in half-space becomes

$$Z_{\rm HS} = \frac{1}{2} \int [DX] e^{-I_{\rm NLSM}[X,\tilde{\Phi},\omega,\gamma]},\tag{3.10}$$

$$=\frac{1}{2}\int d^{D}Y\int [D\sqrt{\alpha'}\eta]e^{-I_{\mathrm{p}}[\eta,g]}e^{-V[\eta,g,\tilde{\phi}]}$$
(3.11)

$$= \frac{1}{2} \int d^{D}Y \int [D\sqrt{\alpha'}\eta] e^{-I_{\rm P}[\eta,g]} \left(1 - V + O(V^{2})\right)$$
(3.12)

$$=\frac{1}{2}Z_{\rm nz}\int d^D Y\left(1-\langle V\rangle+\langle O(V^2)\rangle\right),\tag{3.13}$$

where $Z_{nz} := \int \left[D\sqrt{\alpha'}\eta \right] e^{-I_p[\eta,g]}$ is the path integral for the non-zero modes and $\langle V \rangle$ is defined by

$$\langle V \rangle = \frac{1}{Z_{\rm nz}} \int [D\sqrt{\alpha'}\eta] e^{-I_{\rm p}[\eta,g]} V \,. \tag{3.14}$$

Since $\phi \ll 1$, the higher order terms in V are suppressed, so we can drop the $O(V^2)$ terms.

Now we compute $\langle V \rangle$.

$$\langle V \rangle = \frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \int [D\sqrt{\alpha'}\eta] e^{-I_{\rm p}[\eta,g]} \tilde{\phi}(Y^{\mu} + \sqrt{\alpha'}\eta^{\mu}(z))$$
(3.15)

$$= \frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left\langle \tilde{\phi}(Y^{\mu} + \sqrt{\alpha'} \eta^{\mu}(z)) \right\rangle.$$
(3.16)

If Y^{μ} is far away from the wall in string units, i.e. $\frac{|Y^{D}|}{\sqrt{\alpha'}} \gg 1$, then one can Taylor expand $\tilde{\phi}(X^{\mu})$ about the point Y^{μ}

$$\left\langle \tilde{\phi} \left(Y^{\mu} + \sqrt{\alpha'} \eta^{\mu}(z) \right) \right\rangle = \tilde{\phi} \left(Y^{\mu} \right) \left\langle \mathbf{1} \right\rangle + \sqrt{\alpha'} \partial_{\mu} \tilde{\phi} \left(Y^{\mu} \right) \left\langle \eta^{\mu}(z) \right\rangle + \frac{\alpha'}{2} \partial_{\mu} \partial_{\nu} \tilde{\phi} \left(Y^{\mu} \right) \left\langle \eta^{\mu}(z) \eta^{\nu}(z) \right\rangle + O(\alpha'^{3/2}).$$

$$(3.17)$$

Here, $\langle \mathbf{1} \rangle = 1$ and the one-point function vanishes (because it an odd integral). Then

$$\left\langle \tilde{\phi} \left(Y^{\mu} + \sqrt{\alpha'} \eta^{\mu}(z) \right) \right\rangle = \tilde{\phi} \left(Y^{\mu} \right) + \frac{\alpha'}{2} \partial_{\mu} \partial_{\nu} \tilde{\phi} \left(Y^{\mu} \right) \left\langle \eta^{\mu}(z) \eta^{\nu}(z) \right\rangle + O(\alpha'^{3/2}).$$
(3.18)

It may be worth pointing out that this far-from-the-wall region where the worldsheet path integral is *not* restricted is the case considered in [4], and this is the reason why there is no notion of a one-point to begin with.

However, if Y^{μ} is close to the wall, i.e $\frac{|Y^{D}|}{\sqrt{\alpha'}} \sim 1$, then the kink in $\tilde{\phi}(X^{i}, X^{D})$ at the wall does matter. In this near wall region, the dilaton will approximately be a linear function of the absolute value of X_{D} on either side of the wall

$$\tilde{\phi}\left(X^{i}, X^{D}\right) \approx \phi\left(X^{i}, 0\right) + \partial_{D}\phi\left(X^{i}, 0\right) \left|X^{D}\right|, \qquad (3.19)$$

where $\partial_D \phi(X^i, 0)$ is the right limit of the first derivative (as strictly speaking, the derivative doesn't exist on the wall due to the kink):

$$\partial_D \phi \left(X^i, 0 \right) := \lim_{X^D \to 0^+} \partial_D \phi \left(X^i, X^D \right).$$
(3.20)

Now we can Taylor expand along the tangential direction to the wall about the point Y^i

$$\tilde{\phi}\left(Y^{i} + \sqrt{\alpha'}\eta^{i}(z), Y^{D} + \sqrt{\alpha'}\eta^{D}(z)\right) \approx \phi\left(Y^{i} + \sqrt{\alpha'}\eta^{i}(z), 0\right) + \partial_{D}\phi\left(Y^{i} + \sqrt{\alpha'}\eta^{i}(z), 0\right) \left|Y^{D} + \sqrt{\alpha'}\eta^{D}(z)\right|$$
(3.21)

$$=\phi\left(Y^{i},0\right)+\sqrt{\alpha'}\partial_{i}\phi\left(Y^{i},0\right)\eta^{i}(z)$$
(3.22)

$$+ \partial_D \phi \left(Y^i, 0 \right) \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right|$$
(3.23)

$$+ \sqrt{\alpha'} \partial_i \partial_D \phi \left(Y^i, 0 \right) \eta^i(z) \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| + O(\alpha').$$
(3.24)

The expectation value of all terms odd in $\eta^i(z)$ vanishes in a free theory. Hence,

$$\left\langle \tilde{\phi} \left(Y^{\mu} + \sqrt{\alpha'} \eta^{\mu}(z) \right) \right\rangle = \phi \left(Y^{i}, 0 \right) + \partial_{D} \phi \left(Y^{i}, 0 \right) \left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z) \right| \right\rangle + O(\alpha').$$
(3.25)

Thus, it is the near-wall region that leads to a non-zero one-point function in the limit $X^D \to 0^+$ because the wiggling part of the string $\eta^D(z)$ is restricted to half-space by the presence of the wall.

An explicit calculation of the exact one-point function $\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle$ is presented in appendix A.

4 The classical boundary dilaton action in half-space

In this section, we put things together to compute the total classical off-shell action in half-space, $I_{\rm HS}$, and show it obeys a well-defined variational principle for Neumann boundary conditions. We first

use the result in appendix A for the one-point $\langle |Y^D + \sqrt{\alpha'}\eta^D(z_0)| \rangle$ to write down an expression for the boundary dilaton action from the near-wall spacetime region. We then compute the bulk dilaton action I_{bulk} , i.e. the dilaton kinetic term, from the far-wall region. This bulk action comes with its own boundary contribution [4]. To obtain the total action, we add all bulk and boundary contributions and apply Tsyetlin's prescriptions. We then find that the two boundary terms cancel out (as they should) and we are left with an off-shell action for the dilaton in half-space that has a well-defined variational principle for a Neumann boundary condition.

In appendix A, we show that

$$\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z_{0}) \right| \right\rangle = \sqrt{\alpha'} \left(\frac{\sqrt{\Omega}}{\sqrt{\pi}} e^{-\frac{(Y^{D})^{2}}{\alpha'\Omega}} + \frac{Y^{D}}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^{D}}{\sqrt{\alpha'\Omega}}\right) \right)$$
(4.1)

$$= \sqrt{\alpha'} \left(\left\langle \left| \eta^D(z_0) \right| \right\rangle e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{Y^D}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right) \right).$$
(4.2)

The sphere partition function in half-space (3.10) splits into two pieces, each in a different spacetime region: near the wall and far from the wall

$$Z_{\rm HS} = \frac{1}{2} Z_{\rm nz} V_{Y^D} + Z_{\rm near} + Z_{\rm far} + O(V^2) \,, \tag{4.3}$$

where $V_{Y^D} = \int d^D Y$ and

$$Z_{\text{near}} := \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \sim 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left\langle \tilde{\phi}(Y^\mu + \sqrt{\alpha'} \eta^\mu(z)) \right\rangle \right).$$
(4.4)

$$Z_{\text{far}} := \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \gg 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left\langle \tilde{\phi}(Y^\mu + \sqrt{\alpha'} \eta^\mu(z)) \right\rangle \right).$$
(4.5)

Far from the wall, Z_{far} gives the bulk contribution to the action while, due to the presence of the wall, Z_{near} gives an additional contribution which is given by

$$Z_{\text{near}} := \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \sim 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left(\phi \left(Y^i, 0 \right) + \partial_D \phi \left(Y^i, 0 \right) \left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle + O(\alpha') \right) \right)$$

$$(4.6)$$

$$= \frac{1}{2} Z_{\rm nz} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \sim 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left(\phi \left(Y^i, 0 \right) + O(\alpha') \right) \right) + Z_{\rm wall} \,, \tag{4.7}$$

where Z_{wall} is

$$Z_{\text{wall}} = \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \sim 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \partial_D \phi \left(Y^i, 0 \right) \left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle \right)$$
(4.8)

$$= \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \sim 1} d^D Y(-2) \partial_D \phi\left(Y^i, 0\right) \left\langle \left|Y^D + \sqrt{\alpha'} \eta^D(z)\right| \right\rangle$$

$$\tag{4.9}$$

$$= -\frac{1}{2} \times 2Z_{\rm nz} \int d^{D-1}Y \,\partial_D \phi\left(Y^i, 0\right) \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \sim 1} dY^D \sqrt{\alpha'} \left(\frac{\sqrt{\Omega}}{\sqrt{\pi}} e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{Y^D}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right)\right). \tag{4.10}$$

The integrand in (4.10) is valid *only* near the wall where $Y^D \sim \sqrt{\alpha'}$ and thus we integrate it from $-\frac{Y_c\sqrt{\alpha'}}{2}$ to $\frac{Y_c\sqrt{\alpha'}}{2}$ where the cutoff Y_c is a dimensionless constant.

$$Z_{\text{wall}} = -\frac{1}{2} \times 2Z_{\text{nz}} \int d^{D-1}Y \ \partial_D \phi \left(Y^i, 0\right) \int_{-\frac{Y_c \sqrt{\alpha'}}{2}}^{\frac{Y_c \sqrt{\alpha'}}{2}} dY^D \sqrt{\alpha'} \left(\frac{\sqrt{\Omega}}{\sqrt{\pi}} e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{Y^D}{\sqrt{\alpha'}} \text{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right)\right)$$
(4.11)

$$= -\frac{1}{2} \times 2Z_{\rm nz} \int d^{D-1}Y \,\partial_D \phi\left(Y^i, 0\right) 2 \int_0^{\frac{Y_c \sqrt{\alpha'}}{2}} dY^D \sqrt{\alpha'} \left(\frac{\sqrt{\Omega}}{\sqrt{\pi}} e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{Y^D}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right)\right)$$
(4.12)

$$= -\frac{1}{2} \times 2Z_{\rm nz} \int d^{D-1}Y \ \partial_D \phi \left(Y^i, 0\right) \frac{\alpha'}{4} \left(\frac{2e^{-\frac{Y_c^2}{4\Omega}}Y_c\sqrt{\Omega}}{\sqrt{\pi}} + (Y_c^2 + 2\Omega) \mathrm{erf}\left(\frac{Y_c}{2\sqrt{\Omega}}\right)\right). \tag{4.13}$$

Expanding near $Y_c \to \infty$ and keeping only the leading and subleading terms, the first term vanishes and erf $\left(\frac{Y_c}{2\sqrt{\Omega}}\right) \to 1$ and thus we obtain

$$Z_{\text{wall}} = -\frac{1}{2} \times 2Z_{\text{nz}} \int d^{D-1}Y \ \partial_D \phi \left(Y^i, 0\right) \left(\frac{\alpha' Y_c^2}{4} + \frac{\alpha' \Omega}{2}\right). \tag{4.14}$$

There is a factor of $e^{-2\Phi_0}$ sitting inside Z_{nz} . Extracting it gives $Z_{nz} = \tilde{Z}_{nz}e^{-2\Phi_0}$, where \tilde{Z}_{nz} does not depend on the dilaton

$$Z_{\text{wall}} = -\frac{1}{2} \times 2\tilde{Z}_{\text{nz}} \int d^{D-1}Y \ e^{-2\Phi_0} \partial_D \phi \left(Y^i, 0\right) \left(\frac{\alpha' Y_c^2}{4} + \frac{\alpha' \Omega}{2}\right).$$
(4.15)

We can express the derivative of ϕ as the derivative of Φ (as Φ_0 is just a constant) and also replace $e^{-2\Phi_0}$ with $e^{-2\Phi}$ (as the difference will just be higher order) to finally get

$$Z_{\text{wall}} = -\frac{1}{2} \times 2\tilde{Z}_{\text{nz}} \int d^{D-1}Y \ e^{-2\Phi} \partial_D \Phi \left(Y^i, 0\right) \left(\frac{\alpha' Y_c^2}{4} + \frac{\alpha' \Omega}{2}\right). \tag{4.16}$$

Using that $\Omega = -2 \ln \epsilon$, Z_{wall} takes the final form

$$Z_{\text{wall}} = -\frac{1}{2} \times 2\tilde{Z}_{\text{nz}} \,\alpha' \int d^{D-1}Y \, e^{-2\Phi} \partial_D \Phi \left(Y^i, 0\right) \left(\frac{Y_c^2}{4} + \frac{-2\ln\epsilon}{2}\right). \tag{4.17}$$

The above form should be familiar. It has a structure similar to that of the bulk sphere partition function: there is an overall log divergence multiplying the boundary dilaton sphere effective action.⁶

As in the bulk off-shell action, we apply Tseytlin's prescription $\frac{\partial}{\partial \log \epsilon}$ to eliminate overall the log divergence and the divergent Y_c^2 term to obtain the classical off-shell boundary action of the dilaton

⁶Although we emphasize that in this work, we don't have an independent NLSM derivation of $\partial_D \Phi(Y^i, 0)$ in terms of the dilaton beta function.

in half-space

$$I_{\text{wall}} = -\frac{\partial}{\partial \log \epsilon} Z_{\text{wall}} = -\alpha' \tilde{Z}_{\text{nz}} \int d^{D-1} Y \ e^{-2\Phi} \partial_D \Phi \left(Y^i, 0 \right) = +\alpha' \tilde{Z}_{\text{nz}} \int d^{D-1} Y \ e^{-2\Phi} \partial_n \Phi \left(Y^i, 0 \right),$$

$$(4.18)$$

where ∂_n is the normal derivative in the direction of the outward-pointing normal n^{μ} i.e. $\partial_n = -\partial_D$.

Now, we compute Z_{far} to obtain the bulk contribution to Z_{HS}

$$Z_{\text{far}} = \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \gg 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left\langle \tilde{\phi}(Y^\mu + \sqrt{\alpha'} \eta^\mu(z)) \right\rangle \right)$$

$$= \frac{1}{2} Z_{\text{nz}} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \gg 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \left(\tilde{\phi}(Y^\mu) + \frac{\alpha'}{2} \partial_\mu \partial_\nu \tilde{\phi}(Y^\mu) \left\langle \eta^\mu(z) \eta^\nu(z) \right\rangle + O(\alpha'^{3/2}) \right) \right).$$
(4.19)
$$(4.20)$$

Keeping only the relevant term (with a log ϵ^{-1} coefficient) in Z_{far} is

$$Z_{\text{far}} = \frac{1}{2} Z_{\text{nz}} \int_{\frac{|YD|}{\sqrt{\alpha'}} \gg 1} d^D Y \left(-\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \frac{\alpha'}{2} \partial_\mu \partial_\nu \tilde{\phi} \left(Y^\mu \right) \delta^{\mu\nu} \left(-\log \epsilon + O(\epsilon^2) \right) \right).$$
(4.21)

Applying Tseytlin's prescription to Z_{far} gives I_{bulk}

$$I_{\text{bulk}} := I_{\text{far}} = -\frac{\partial}{\partial \log \epsilon} Z_{\text{far}}$$
(4.22)

$$= -\frac{1}{2} Z_{\mathrm{nz}} \frac{\alpha'}{2} \int_{\frac{|Y^D|}{\sqrt{\alpha'}} \gg 1} d^D Y \left(\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \partial^2 \tilde{\phi} \left(Y^{\mu} \right) \right)$$
(4.23)

$$= -\frac{2}{2} Z_{\rm nz} \frac{\alpha'}{2} \int_{\frac{Y^D}{\sqrt{\alpha'}} \gg 1} d^D Y \left(\frac{1}{4\pi} \int d^2 z \sqrt{g} R^{(2)} \partial^2 \tilde{\phi} \left(Y^{\mu} \right) \right)$$
(4.24)

$$= -\frac{2}{2} Z_{\rm nz} \frac{\alpha'}{2} \int_{\frac{YD}{\sqrt{\alpha'}} \gg 1} d^D Y \left(\frac{8\pi}{4\pi} \partial^2 \tilde{\phi} \left(Y^{\mu} \right) \right)$$
(4.25)

$$= -\tilde{Z}_{nz}\alpha' \int_{\frac{YD}{\sqrt{\alpha'}} \gg 1} d^D Y e^{-2\Phi} \partial^2 \Phi$$
(4.26)

This free action for the dilaton (4.26) does *not* have a well-defined variational principle since upon variation, it gives

$$\delta I_{\text{bulk}} = 2\alpha' \tilde{Z}_{\text{nz}} \int_{\frac{YD}{\sqrt{\alpha'}} \gg 1} d^D Y \left[e^{-2\Phi} \partial^2 \Phi + \partial_\mu \left(e^{-2\Phi} \partial^\mu \Phi \right) \right] \delta \Phi$$
$$- \alpha' \tilde{Z}_{\text{nz}} \int_{\partial \mathcal{M}} d^{D-1} Y e^{-2\Phi} \,\delta(\partial_n \Phi) - 2\alpha' \tilde{Z}_{\text{nz}} \int_{\partial \mathcal{M}} d^{D-1} Y e^{-2\Phi} \partial_n \Phi \,\delta \Phi. \tag{4.27}$$

The first term in (4.27) is the bulk dilaton equation of motion while the second and third are boundary terms. The second boundary term in (4.27) is where we impose boundary conditions on $\Phi(Y^i, 0)$. It vanishes for Dirichlet boundary conditions ($\delta \Phi = 0$) while imposing Neumann boundary conditions ($\partial_n \Phi = 0$) guarantees a consistent boundary equation of motion. However, this is true *only if* the dilaton $\Phi(Y^i, 0)$ satisfies the boundary conditions on the wall. In our case, it does *not* simply because $\Phi(Y^i, 0)$ is off-shell. The first boundary term is what spoils the variational principle. It is the GHY-like term of the free two-derivative action (4.26) which must be accounted for, i.e. by adding an *additional* boundary term whose variation exactly cancels with it to have a well-posed variational principle. This additional boundary term is I_{wall} in (4.18). Let's see how this works.

The total classical off-shell string action for the dilaton in half-space is the sum of bulk and boundary contributions (4.18)

$$I_{\text{sphere}} := I_{\text{HS}} = I_{\text{bulk}} + I_{\text{wall}}$$

$$= -\tilde{Z}_{\text{nz}} \alpha' \int_{\frac{Y^{D}}{\sqrt{\alpha'}} \gg 1} d^{D} Y e^{-2\Phi} \partial^{2} \Phi + \alpha' \tilde{Z}_{\text{nz}} \int d^{D-1} Y \ e^{-2\Phi} \partial_{n} \Phi \left(Y^{i}, 0\right) \qquad (4.28)$$

$$= -2\tilde{Z}_{\text{nz}} \alpha' \int_{\frac{Y^{D}}{\sqrt{\alpha'}} \gg 1} d^{D} Y e^{-2\Phi} \partial_{\mu} \Phi \partial^{\mu} \Phi, \qquad (4.29)$$

where integration by parts is used to get the last line.

Vorwing I gives

Varying I_{sphere} gives

$$\delta I_{\text{sphere}} = \delta I_{\text{bulk}} + \delta I_{\text{wall}}$$
$$= 2\alpha' \tilde{Z}_{\text{nz}} \int_{\frac{Y^D}{\sqrt{\alpha'}} \gg 1} d^D Y \left[e^{-2\Phi} \partial^2 \Phi + \partial_\mu \left(e^{-2\Phi} \partial^\mu \Phi \right) \right] \delta \Phi$$
(4.30)

$$-\alpha'\tilde{Z}_{nz}\int_{\partial\mathcal{M}} d^{D-1}Y e^{-2\Phi}\,\delta(\partial_n\Phi) + \alpha'\tilde{Z}_{nz}\int_{\partial\mathcal{M}} d^{D-1}Y e^{-2\Phi}\,\delta(\partial_n\Phi) \tag{4.31}$$

$$-4\alpha'\tilde{Z}_{nz}\int_{\partial\mathcal{M}}d^{D-1}Ye^{-2\Phi}\partial_n\Phi\,\delta\Phi\tag{4.32}$$

$$= 2\alpha' \tilde{Z}_{nz} \int_{\frac{YD}{\sqrt{\alpha'}} \gg 1} d^D Y \left[e^{-2\Phi} \partial^2 \Phi + \partial_\mu \left(e^{-2\Phi} \partial^\mu \Phi \right) \right] \delta \Phi - 4\alpha' \tilde{Z}_{nz} \int_{\partial \mathcal{M}} d^{D-1} Y e^{-2\Phi} \partial_n \Phi \, \delta \Phi.$$

$$\tag{4.33}$$

In the second line, we notice that the variation δI_{wall} exactly cancels its counterpart in δI_{bulk} i.e. $e^{-2\Phi} \,\delta(\partial_n \Phi)$ such that we are left in the end with only the bulk and boundary dilaton equations of motion. Therefore, I_{sphere} has a well-defined variational principle only with the addition of I_{wall} as a boundary term.

This is the main result of this paper.

4.1 The classical on-shell action

Let us now impose the dilaton *bulk* equation of motion in I_{bulk} to obtain the on-shell action. To do that, we first integrate by parts (4.26) *before* varying to obtain

$$I_{\text{bulk}} = -\tilde{Z}_{\text{nz}} \alpha' \int_{\frac{Y^D}{\sqrt{\alpha'}} \gg 1} d^D Y e^{-2\Phi} \partial^2 \Phi$$
(4.34)

$$= -\tilde{Z}_{nz}\alpha'\int_{\frac{Y^D}{\sqrt{\alpha'}}\gg 1} d^D Y \,\partial_\mu \left(e^{-2\Phi}\partial^\mu\Phi\right) - 2\tilde{Z}_{nz}\alpha'\int_{\frac{Y^D}{\sqrt{\alpha'}}\gg 1} d^D Y e^{-2\Phi}(\partial\Phi)^2 \tag{4.35}$$

$$= -\tilde{Z}_{nz}\alpha'\int d^{D-1}Y e^{-2\Phi}\partial_n\Phi - 2\tilde{Z}_{nz}\alpha'\int_{\frac{YD}{\sqrt{\alpha'}}\gg 1} d^DY e^{-2\Phi}(\partial\Phi)^2.$$
(4.36)

Imposing the dilaton equation of motion in I_{bulk}

$$e^{-2\Phi}(\partial\Phi)^2 = -\partial_\mu \left(e^{-2\Phi}\partial^\mu\Phi\right) \tag{4.37}$$

leaves us with the following bulk action

$$I_{\text{bulk}} = +\tilde{Z}_{\text{nz}}\alpha' \int d^{D-1}Y e^{-2\Phi} \partial_n \Phi$$
(4.38)

The classical on-shell action for the dilaton in half-space is therefore the sum of the two boundary contributions (4.38) and (4.18)

$$I_{\rm sphere} := I_{\rm HS} = I_{\rm bulk} + I_{\rm wall} \tag{4.39}$$

$$= 2 \alpha' \tilde{Z}_{\rm nz} \int d^{D-1} Y e^{-2\Phi} \partial_n \Phi = 2I_{\rm wall} \,. \tag{4.40}$$

Therefore, we observe that the two boundary terms in I_{sphere} add up to give the on-shell classical action.

Note that if we impose the *boundary* equation for the motion for the dilaton, i.e. the Neumann boundary condition, then $I_{\text{wall}} = 0$ and hence $I_{\text{sphere}} = 0$. This is of course another way to say our worldsheet derivation of I_{wall} from the outset is off-shell.

5 Discussion

In this section, we begin by making the observation that the N-th point correlation function $\left\langle \left| \eta^{D}(z_{0}) \right|^{N} \right\rangle$ in (A.56)

$$\left\langle \left| \eta^{D}(z_{0}) \right|^{N} \right\rangle = \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{1+N}{2}\right) \Omega^{N/2},$$
(5.1)

equals the N-th moment of the half-normal distribution for the absolute value of a random variable |X|. We will comment on the relationship between the two quantities as well as the relationship between the string's zero and nonzero mode to reflected Brownian motion and one-dimensional random walks in spaces with a totally reflecting barrier. We begin first by defining the half-normal distribution and its main properties.

Let X_t be a normally distributed random variable with a standard deviation $t \in [0, \infty)$. The half-normal distribution is a one-parameter scale family of distributions with scale (time) parameter twhich is closed under scale transformations in the sense that for $t \in [0, \infty)$, $Y_t = t|X_t|$. If t = 1, then Y_t has the standard half-normal distribution.⁷

The probability density function (PDF) f(y;t) of Y with scale parameter σ is given by

$$f(y;t) = \frac{1}{t}\sqrt{\frac{2}{\pi}}\exp\left(-\frac{y^2}{2t^2}\right), \quad y \in [0,\infty),$$
 (5.2)

⁷The half-normal distribution is a special case of the folded normal distribution with $\mu = 0$: = $\frac{1}{t\sqrt{2\pi}} \left\{ \exp\left[-\frac{1}{2}\left(\frac{x+\mu}{t}\right)^2\right] + \exp\left[-\frac{1}{2}\left(\frac{x-\mu}{t}\right)^2\right] \right\}.$

while the cumulative distribution function (CDF) F(y;t) is

$$F(y;t) = \int_0^y \frac{1}{t} \sqrt{\frac{2}{\pi}} \exp\left(-\frac{x^2}{2t^2}\right) \, dx \,.$$
 (5.3)

Changing variables to $w = x/(\sqrt{2}t)$, the CDF can be written as

$$\mathbb{P}\left(Y_t \le y\right) := F_Y(y;t) = \operatorname{erf}\left(\frac{y}{\sqrt{2t}}\right).$$
(5.4)

For any (even or odd) $N \ge 0$, the absolute central moments are given by [33]⁸

$$\left\langle Y_t^N \right\rangle = \left\langle |X|_t^N \right\rangle = t^N \cdot \frac{2^{N/2} \Gamma\left(\frac{N+1}{2}\right)}{\sqrt{\pi}},\tag{5.5}$$

with variance $\operatorname{var}(Y_t) = t^2 \left(1 - \frac{2}{\pi}\right)$. For N = 1, $\langle |X| \rangle = \frac{t\sqrt{2}}{\sqrt{\pi}}$ is the mean of the half-normal distribution. Comparing (5.1) with (5.5), and using $\Omega = 2 \log \epsilon^{-1}$, we observe that they are identical

$$\left\langle \left| \eta^{D}(z_{0}) \right|^{N} \right\rangle = \left\langle \left| X \right|_{t}^{N} \right\rangle,$$
(5.6)

once we identify the scale parameter of the distribution t with the square root of the log of the worldsheet UV cutoff ϵ

$$t\sqrt{2} = \sqrt{\Omega} = \sqrt{-\log\epsilon}\sqrt{2}$$
, and hence, $t = \sqrt{-\log\epsilon}$. (5.7)

In particular, the one-point function of the string nonzero mode $|\eta^D|$ is trivially equal to the first moment (N = 1) of $|X_t|$

$$\left\langle \left| \eta^D(z_0) \right| \right\rangle = \left\langle |X_t| \right\rangle = \frac{t\sqrt{2}}{\sqrt{\pi}}.$$
 (5.8)

This observation gives a probabilistic interpretation of I_{HS} and connects it to the vast and interesting literature on the theory of reflecting Brownian motion [34, 35]. Specifically, the half-normal distribution naturally appears in the theory of reflected Brownian motion (RBM) of stochastic (Wiener) processes of $\langle |X_t| \rangle$ in spaces with reflecting boundaries. It arises as the probability distribution for the *local boundary time* [36–40] and running maximum [34, 35, 41, 42]. See also [43]. In [44, 45], it is also shown that the half-normal distribution emerges as the limit law for the *local time* at x = 0 in directed lattice path on \mathbb{Z} with no drift. The local time at zero in this type of random walk is the sum the number of returns to zero and number of x-axis crossings.

After giving a short introduction to the probabilistic interpretations of boundary conditions in section 5.1, we give in section 5.2 a lightening overview of reflected Brownian motion (RBM), the Skorokhod construction [46, 47], Tanakas's formula [48] and the important notion of local boundary time at a point $x L_t^x$ [36, 37].

We will then argue in section 5.3 that $|\sqrt{\alpha'}\eta^D(z)|$ has the same probability distribution of the boundary local time and that

$$\left\langle \left| \sqrt{\alpha'} \eta^D(z) \right| \right\rangle = \left\langle L_t^x \right\rangle,$$
(5.9)

⁸We assume the expectation value (first central moment) of X is zero. Also, we are using $\langle X^N \rangle$ to denote the moments of X rather than $\mathbb{E}[X^N]$.

thereby giving a purely probabilistic (random walk) description of the string dynamics in target spacetime.

We would like to comment on the fact that the value of the square root of the worldsheet length (UV) cutoff ϵ^{-1} acts in target space as a dial that controls the *overall* scale t of the half-normal distribution. Each value of ϵ^{-1} describes a worldsheet theory at a particular renormalization group (RG) scale. Fig. 2 shows the PDF and CDF of the half-normal probability distribution with different scale parameter $t = \sqrt{-\log \epsilon}$. The plots can be interpreted as the RG flow from the UV with larger value of ϵ^{-1} (or smaller value of $\log \epsilon^{-1}$) to the IR with a small value of $\epsilon^{-1} \rightarrow 0$. The RG flow renormalizes the length of the Euclidean Schwinger propagator or equivalently the Schwinger time, as the worldsheet theory flows from the UV to the IR. The $\log \epsilon^{-1} \rightarrow 0$ limit corresponds to the S-matrix regime whereas, when $\log \epsilon^{-1} \rightarrow 0$ when ($\epsilon \sim 1$), the worldsheet theory is in the extreme UV limit where the worldsheet theory is nonlocal [6]. The latter is the limit where the string nonzero mode $\langle |\eta(z)| \rangle$ vanishes and thus, $\eta^D(z)$ spends all its time (with probability 1) at the boundary $X_D = 0$.

To establish a more visible connection between the value of $\log \epsilon^{-1}$ and RG time, we note that the solution to one-dimensional heat diffusion equation

$$\frac{\partial k}{\partial t} = \frac{\partial^2 k}{\partial x^2},$$
(5.10)

with a step function as the initial condition

$$k(x,0) = \begin{cases} 0 & \text{if } x < 0\\ 1 & \text{if } x > 0 \end{cases}$$
(5.11)

is given by

$$k(x,t) = \frac{1}{2} \left[1 - \operatorname{erf}\left(\frac{x}{\sqrt{4t}}\right) \right], \qquad (5.12)$$

for $x \in \mathbb{R}$ and t > 0. This solution satisfies the heat equation and approaches the initial condition as $t \to 0^+$. It shows how the initial discontinuity (kink) at x = 0 diffuses (renormalized) over time, gradually smoothing out the step transition from k = 0 to k = 1 as heat spreads out into the bulk. Note that *except* right at the boundary at x = 0, the motion k(x, t) is standard Brownian. The discontinuity at the x = 0 breaks the the *spatial homogeneity* assumption of standard Brownian motion. In other words, scale invariance is broken at the target space boundary. This is a fundamental principle of the reflected Brownian motion. In a *distributional* sense, function $k(x, t = t_0)$ corresponds to the dilaton $\tilde{\Phi}(Y^i, 0)$.

5.1 The probabilistic interpretations of boundary conditions

It is well known that with a Neumann boundary condition at x = a, $\frac{\partial k}{\partial x}\Big|_{x=a} = 0$, the solution to the one-dimensional heat kernel equation is the *sum* of k(x,t) + k(x-a,t) since reflecting about the line x = a, we simply have

$$\frac{\partial k}{\partial x}\Big|_{x=a} = k(x,t)\Big|_{x=a} + k(x-a,t)\Big|_{x=a}$$
(5.13)

$$= k(x,t)\Big|_{x=a} + k(a-x,t)\Big|_{x=a}$$
(5.14)

$$= k(x,t)\Big|_{x=a} - k(x,t)\Big|_{x=a} = 0, \qquad (5.15)$$



Figure 2: The half-normal probability density f(y;t) and cumulative distribution F(y;t) functions for different scale parameters $t = \sqrt{\log \epsilon^{-1}}$. The effect of changing the value of the UV position cutoff ϵ is to change the overall size of the distribution. A larger value of t (or equivalently, a smaller value of ϵ) brings the probability density on the left closer to a delta function and distribution on the right closer to a step function $\theta(Y)$. In terms of the local boundary time (5.19), a larger ϵ is a less diffusive regime that describes a *typical* string (in the distribution) $\eta^D(z_0)$ which is more *bound* to the wall at $X^D = 0$, i.e. spends most of its time in the near the wall within a string length $\sqrt{\alpha'}$. In this region, the string hits the wall several times before getting diffused away. Contrast this with other distributions with smaller values of ϵ where the string is less localized to the boundary in a more diffusive medium. Figures adapted from https://www.mathworks.com/help/stats/half-normal-distribution.html.

where $k(x,t) = \frac{1}{2\sqrt{t}} \exp\left(-x^2/4t\right)$ is the solution with no boundaries and a delta function $\delta(x-a)$ as an initial condition. With Dirichlet boundary condition on the other hand, it is the difference instead k(x,t) - k(x-a,t) that satisfies k(a,t) = 0, as can be easily checked.

The probabilistic interpretations of boundary conditions e.g. Neumann [40] and Dirichlet boundary conditions provide a fascinating link between stochastic processes, particularly Brownian motion, and partial differential equations. These interpretations can offer intuitive insights into how boundary conditions affect the behavior of random walks or diffusions, which parallel solutions to PDEs under similar conditions.

Neumann boundary conditions specify the derivative of a function along the boundary of its domain D, typically the normal derivative. In probabilistic terms, Neumann boundary conditions are totally reflecting barriers, in the sense that particle paths of a Brownian motion or another diffusion process (or in our case string worldsheets) that reach the wall are reflected *instantaneously* back into D with probability 1.

Dirichlet boundary conditions, on the other hand, specify the value of a function at the boundary of its domain. In the context of stochastic processes, Dirichlet boundary conditions correspond to the behavior of absorbing barriers where the paths of Brownian motion are totally absorbed and the random process completely stops upon reaching the boundary ∂D [49].

Reflected Brownian motion is the formal mathematical framework used to describe this probabilistic interpretation of the stochastic string dynamics in target spacetime. This is what we review next.

5.2 Reflected Brownian motion, local boundary time and running maximum

In this section, we give the bare minimum review of reflected Brownian motion and the associated notion of boundary local time. We will state known results, propositions and theorems without proof and omit any discussion of mathematical subtleties. However, plenty of references will be given for the readers interested in the details. ⁹

Let $\mathbf{X} = \{X_t : t \in [0, \infty)\}$ be a standard Brownian motion (Weiner) process and let $S_t = \sup_{s \leq t} \mathbf{X}_s$ of \mathbf{X} be the maximum value of \mathbf{X} on the interval [0, t]. Then, the reflection principle states that if X_t is a Brownian motion and T is the time at which X_t first hits level $a > 0^{10}$, then the reflected process defined by

$$X'(t) = \begin{cases} X_t & \text{if } t < T, \\ 2X_T - X_t & \text{if } t \ge T, \end{cases}$$
(5.16)

is also a Brownian motion. This principle is used to derive properties of Brownian motion, including the distribution of S_t over certain intervals. Specifically, it refers to a lemma that states

$$\mathbb{P}\left(S_t \ge a\right) = 2\mathbb{P}(W(t) \ge a) \tag{5.17}$$

The boundary local time [36, 37, 51] is the quantity that encodes all the boundary physics. It is the term you add to a (stochastic) differential equation or a classical action to impose a specific boundary condition in spaces with boundaries. It represents the amount of time that a stochastic process, such as Brownian motion, spends at a boundary of its domain. In more technical terms, it quantifies the *intensity* or of the path's contact with the boundary.

In path integrals, the exponential of the boundary local time gives the total probability of particle paths reflecting from the wall [52]. In reflected Brownian motion, for example, it ensures that worldlines of particles (and of course string worldsheets) in a diffusive medium do not cross a reflecting boundary, i.e. reflected off the boundary. Let us make this notion a bit more precise and see how it relates to $\langle |\eta^D(z)| \rangle$.

The boundary local time measures the amount of time a Brownian motion X spends at level x. Informally, it is defined as

$$L_t^x = \int_0^t \delta_x \left(X_s \right) ds.$$
(5.18)

A more formal definition of L_t is given in terms of the limit of the occupation time T_t of the Brownian motion \mathbf{X}_t in a region ε near the boundary $\partial \mathcal{M}_{\varepsilon} = \{X_t \in \mathcal{M} : |x - \partial \mathcal{M}| < \varepsilon\}$

$$L_{t} = \lim_{\varepsilon \to 0} \frac{1}{2\varepsilon} \int_{0}^{t} ds \, \mathbb{1}_{\partial \mathcal{M}_{\varepsilon}} \left(X_{s} \right), \tag{5.19}$$

where $\mathbb{1}_{\partial \mathcal{M}_{\varepsilon}}(X_s)$ is the indicator function of the boundary (i.e., $\mathbb{1}_{\partial \mathcal{M}_{\varepsilon}}(X_t) = 1$ if $X_t \in \partial \mathcal{M}_{\varepsilon}$, and 0 otherwise). With this definition, the local boundary time L_t measures the amount of time X_t spends in a region ε near the boundary $\partial \mathcal{M}$, and thus increases only when $X_s \in \partial \mathcal{M}_{\varepsilon}$.

A reflected Brownian motion Z is a Brownian process X regulated by another process F such that Z remains always positive. More precisely, if the process $X : \mathbb{R} \to \mathbb{R}$, the Skorokhod construction [46, 47] guarantees that there always exists a pair of functions Z, F such that

$$Z_t = X_t + F_t(X) \ge 0$$
, for all $t \ge 0$. (5.20)

 $^{^{9}}$ We find [50] an accessible pedagogical introduction to the subject of reflected Brownian motion.

¹⁰The first time at which X hits level a is $T = \inf \{t \ge 0 : X_t \ge a\}$, which is a stopping time.

The process F_t is nondecreasing with $F_0 = 0$ and increases only when Z_t reaches the boundary, i.e. when $Z_t = 0$. Intuitively, F_t pushes the process X_t away from the boundary such that Z_t is always positive at any point in time.

All three processes Z_t , X_t and F_t appear in *Skorokhod* stochastic differential equation [46] that describes motion in spaces (R^D) with reflecting obstacles [38]

$$dZ_t = dX_t + \boldsymbol{n} \left(X_t \right) \mathbb{1}_{\partial \mathcal{M}} \left(X_t \right) dL_t, \quad \text{with } Z_0 = z_0 \,, \tag{5.21}$$

where $\boldsymbol{n}(X_t)$ is the normal unit vector perpendicular to the boundary at X_t .

5.3 The probabilistic nature of the $\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle$

Let our bounded region in \mathbb{R}^D be the half-space $\mathbb{R}_+ \times \mathbb{R}^{D-1}$. In this case, Tanaka's formula solves the Skorokhod equation for the reflected process $Z_t = |X_t|$ and $F_t = L_t^0$

$$|X_t| = |Y^D| + \int_0^t \operatorname{sgn}(X_s) \, dX_s + L_t^0,$$
(5.22)

where sgn (X_s) is the sign function of X_s and L_t^0 is the local time of X_t at 0 up to time t. Skipping details of the proof [51], then it can proved using the reflection principle the following three processes have the same probability distribution

$$L_t^0(X) \stackrel{d}{=} S_t \stackrel{d}{=} |X_t|, \qquad (5.23)$$

where in the notation of [50], $\stackrel{d}{=}$ denotes equality in distribution.

Let the motion start at $Y^D = 0$, right at the wall. Taking the spatial expectation value (the first moment) of (5.23) in half-space, we then get at a given fixed time t

$$\langle |X_t|\rangle = \langle S_t\rangle = \langle L_t^0(X)\rangle.$$
(5.24)

The running maximum $S_t = \sup_{s \le t} \mathbf{X}_s$ of \mathbf{X} is known to have the half-normal distribution in half-space [43, 53]. Therefore, we directly conclude that the expectation value of the string nonzero $\langle |\eta^D(z_0)| \rangle$ (starting at $Y^D = 0$) is the same as that of the boundary local time $L_t^0(X)$ in half-space

$$\left\langle \left| \eta^D(z_0) \right| \right\rangle = \left\langle L_t^0(X) \right\rangle = \frac{\sqrt{\Omega}}{\sqrt{\pi}} = \frac{\sqrt{-2\log\epsilon}}{\pi} \,.$$
 (5.25)

This equivalence in distribution law extends to the N-th moment [53]

$$\left\langle \left| \eta^{D}(z_{0}) \right|^{N} \right\rangle = \left\langle L_{t}^{0}(X)^{N} \right\rangle.$$
 (5.26)

In the reflected (doubled) space, we must divide $\langle L_t^0(X) \rangle$ by a factor of 2

$$\left\langle \left| \eta^{D}(z_{0}) \right|^{N} \right\rangle = \frac{1}{2} \left\langle L_{t}^{0}(X)^{N} \right\rangle.$$
 (5.27)

Now suppose our Brownian motion starts on the right side of the boundary at a particular point

 $Y^D > 0$. Taking expectation value of (5.22) and using linearity gives¹¹

$$\langle |X_t|\rangle = \left\langle \left|Y^D\right|\right\rangle + \left\langle \int_0^t \operatorname{sgn}\left(X_s\right) dX_s \right\rangle + \left\langle L_t^0(X)\right\rangle$$
(5.28)

$$=Y^{D}+\left\langle L_{t}^{0}(X)\right\rangle . \tag{5.29}$$

We now comment on the properties of the one-point function we computed in appendix A

$$\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z_{0}) \right| \right\rangle = \sqrt{\alpha'} \left(\frac{Y^{D}}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^{D}}{\sqrt{\alpha'\Omega}} \right) + \left\langle \left| \eta^{D}(z_{0}) \right| \right\rangle e^{-\frac{(Y^{D})^{2}}{\alpha'\Omega}} \right).$$
(5.30)

The first term in (5.30) is the half-normal probability density of finding the string at a point $Y^D > 0$ times the average length or scale of the string nonzero mode. The second term is the probability to find $\eta^D(z_0)$ within a small region $(0, Y^D]$ in half-space or $[-Y^D, Y^D]$ in the reflected space times the length of that region

$$Y^{D} \mathbb{P}\left(-Y^{D} \leq \eta^{D}(z_{0}) \leq Y^{D}\right) = 2Y^{D} \mathbb{P}\left(\eta^{D}(z_{0}) \leq Y^{D}\right).$$

$$(5.31)$$

Using the fact that $\langle L_t^0 \rangle = \int_0^t f(s, Y^D) ds = f(Y^D; t)$ for any probability density $f(Y^D; t)$, we see that the second term in (5.30) is nothing but $\langle L_t^0(X) \rangle$ taking into account the (uncertainty) that the motion started at $Y^D > 0$.¹² The first term in (5.30) becomes Y^D in (5.28) in the limit that $Y^D \to \infty$

$$\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z_{0}) \right| \right\rangle = \begin{cases} \sqrt{\alpha'} \left\langle \left| \eta^{D}(z_{0}) \right| \right\rangle & \text{if } Y^{D} = 0, \\ Y^{D} & \text{as } Y^{D} \to \infty. \end{cases}$$
(5.32)

To recap, we have argued that the string nonzero mode in spacetime has the dynamics of a standard Brownian motion (with zero mean) away from the wall whereas near the boundary, $|\eta(z)|$ it is equivalent to the boundary local time $L_t^0(X)$ in distribution law. This provides a justification for splitting $Z_{\rm HS}$ in section 4 into $Z_{\rm near}$ and $Z_{\rm far}$. In this distributional sense, $||Y^D + \sqrt{\alpha'}\eta^D(z_0)|$ measures the time (or length of) the string spends near (a few string-length away from) the boundary sitting at $X_D = 0$. The one-point function $\langle |Y^D + \sqrt{\alpha'}\eta^D(z_0)| \rangle$ measures the average distance (length) of the string near the wall taking into account all number of bounces from it. Therefore, generally speaking, in spaces with reflecting barriers, for all types of reflecting boundary conditions¹³, we expect all boundary physics to be encoded in the probabilistic (distributional) spacetime dynamics of the string nonzero mode.

Note that
$$\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z_0) \right| \right\rangle$$
 can be expressed as the indefinite integral of $\operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right)$

$$\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z_{0}) \right| \right\rangle = \sqrt{\Omega} \frac{\sqrt{\alpha'}}{\sqrt{\pi}} e^{-\frac{(Y^{D})^{2}}{\alpha'\Omega}} + Y^{D} \operatorname{erf}\left(\frac{Y^{D}}{\sqrt{\alpha'\Omega}}\right)$$
(5.33)

$$= \int dY^D \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right) + \operatorname{constant} .$$
 (5.34)

 $^{^{11}\}Big<\int_{0}^{t}\mathrm{sgn}\left(X_{s}\right)dX_{s}\Big>=0$ for an Ito integral with respect to Brownian motion.

¹²There is a slight difference however. The half-normal probability density defined in (5.2) divides by $\frac{1}{\sqrt{\Omega}}$ whereas the second term in (5.30) multiplies by $\sqrt{\Omega}$.

 $^{^{13}}$ For example, a sticky motion is a boundary condition where the string can stick to the boundary for some time before getting reflected.



Figure 3: A directed random walk with 3 returns (R) to and 5 crossings (C) of the line x = 0. The boundary local time = 8.

Taking the two derivatives of both sides of (5.33) with respect to Y^D gives the following equation of motion for the one-point function

$$\Box \left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z_0) \right| \right\rangle = \frac{2}{\sqrt{\alpha' \pi \Omega}} e^{-\frac{(Y^D)^2}{\alpha' \Omega}}, \qquad (5.35)$$

where $\Box \coloneqq \partial_{Y^D}^2$. Therefore, we see that $\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z_0) \right| \right\rangle$ is the general solution of an equation with a Gaussian source, which is a smeared out (regularized) delta function potential. This is not unexpected. After all, the dilaton equation of motion near the boundary for $\Phi = |Y|$ is $\Box |Y| = 2\delta(Y)$.

We end this discussion by illustrating the random walk behavior of the string in target spacetime and what boundary local time means in this case. As mentioned in the beginning of this section, the half-normal distribution emerges as the *limit* law for the *local time* at x = 0 in directed lattice paths (random walks) on \mathbb{Z} with no drift in the limit of large number of steps. Specifically,

local time at zero = the number of returns to zero + number of x-axis crossings . (5.36)

The discrete nature of the boundary local time in directed random walks reveals the explicit random walk behavior of the string in target spacetime. In particular, the expectation value of the string nonzero mode can therefore, be understood as the average number of returns to the wall at zero. The larger this number is, the larger the average length of the string's random walk at a given RG time ϵ^{-1} .

5.4 Outlook and Future Directions

A generalization of the current work to S^1/\mathbb{Z}_2 and curved target spaces with boundaries is an interesting direction. The generalization to codimension-2 boundaries, especially conical geometries are all interesting future directions. A worldsheet understanding of how to impose Dirichlet boundary conditions on propagating fields in target spacetime is required to derive the GHY boundary term in the on-shell classical action. This, in turn, is important for deriving the classical black hole entropy [54].

A probabilistic understanding of the boundary local time in conical geometries may give us some clues that might help us unravel the nature of the boundary conditions required for a worldsheet derivation of the classical black hole entropy $\frac{A}{4G_N}$ and the Ryu-Takayanagi formula [55] as presented in [56]. In particular, figuring out the boundary local time in conical geometries near the tip of the cone may guide us to understand the boundary condition in the NLSM limit of the winding condensate [57, 58]. The authors of this article are now actively working on this direction.

In a forthcoming work [59], it will be shown that the metric total derivative terms for the metric obtained in [4] gives the GHY boundary term [23] (more precisely the Einstein boundary term [60]) in the classical action but with *one-half* the coefficient required to have a well-defined variational principle, if we assume Dirichlet boundary conditions. A proper worldsheet path integral calculation similar to the one in this work, but for Dirichlet boundary conditions, is therefore required to obtain the correct factor of "2" TrK and obtain a total action (EH plus GHY) with a well-defined variational principle.

Another very interesting direction involves understanding the behavior of strings at a *finite non-asymptotic* Dirichlet boundary on a bulk Cauchy slice in AdS [61–65] for $T\bar{T}$ -deformed CFTs. The related question of placing a finite timelike boundary in an AdS black hole and in a dS static patch requires an understanding of the behavior of strings near totally absorbing walls with a Dirichlet boundary condition [66, 67].¹⁴ It would be very interesting if the work in this paper can be generalized to address this question.

Acknowledgements

We are grateful to Aron Wall, Eva Silverstein, Daniel Jeffries, Alex Frenkel, Raghu Mahajan, Prahar Mitra, Arkadi Tseytlin, Edward Witten, Jose Sa, Xi Yin, Gabriel Wong, Yiming Chen, Douglas Stanford, Shoaib Akhtar, Ronak Soni, David Kolchmeyer, and Juan Maldacena for insightful discussions and comments. AA thanks Minjae Cho, Harold Erbin, Lorenz Eberhardt, and Daniel Thompson for interesting discussions. We are especially grateful to Aron Wall for very insightful discussions during the course of this project, reviewing parts of the manuscript and providing valuable feedback and comments. We are also grateful to Eva Silverstein for reviewing parts of the manuscript and giving valuable comments. AA is grateful to Edward Witten and Juan Maldacena for the opportunity to give a talk about this work while it was still ongoing. AA is supported by The Royal Society and by STFC Consolidated Grant No. ST/X000648/1.

Open Access Statement - For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Data access statement: no new data were generated for this work.

Appendix A The computation of $\left< \left| Y^D + \sqrt{lpha'} \eta^D(z) \right| \right>$

In this section we show how to compute the one-point function $\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z) \right| \right\rangle$. Let us in fact be more general and compute $\left\langle f\left(\eta^D(z)\right) \right\rangle$ for an arbitrary function $f\left(\eta^D(z)\right)$.

The first step is to expand $\eta^D(z)$ in spherical harmonics

$$\eta^{D}(z) = \sum_{l=1}^{\infty} \sum_{m=-l}^{m=l} x_{lm} Y_{lm}(z), \tag{A.1}$$

 $^{^{14}\}mathrm{We}$ thank Eva Silverstein and Aron Wall for a discussion of this point.

where $Y_{lm}(z)$ are the real spherical hamonics

$$Y_{\ell m} = \begin{cases} (-1)^m \sqrt{2} \sqrt{\frac{2\ell+1}{4\pi} \frac{(\ell-|m|)!}{(\ell+|m|)!}} P_{\ell}^{|m|}(\cos\theta) \sin(|m|\varphi) & \text{if } m < 0\\ \sqrt{\frac{2\ell+1}{4\pi}} P_{\ell}^m(\cos\theta) & \text{if } m = 0\\ (-1)^m \sqrt{2} \sqrt{\frac{2\ell+1}{4\pi} \frac{(\ell-m)!}{(\ell+m)!}} P_{\ell}^m(\cos\theta) \cos(m\varphi) & \text{if } m > 0 \,. \end{cases}$$
(A.2)

They satisfy the Laplace equation in spherical coordinates

$$\nabla^2 Y_{lm}(z) = -l(l+1)Y_{lm}(z), \qquad (A.3)$$

and are orthogonal

$$\int_0^{2\pi} d\phi \int_0^{\pi} d\theta \,\sin\theta \,Y_{lm}(\theta,\phi)Y_{l'm'}(\theta,\phi) = \delta_{ll'}\delta_{mm'}\,,\tag{A.4}$$

where x_{lm} are the expansion coefficients, and the sum runs from l = 1 to ∞ , not including the zero mode. Now we have

$$\left\langle f\left(\eta^{D}(z)\right)\right\rangle = \frac{1}{Z_{\rm nz}} \int [D\sqrt{\alpha'}\eta] e^{-I_{\rm p}[\eta,g]} f\left(\eta^{D}(z)\right) \tag{A.5}$$

$$= \frac{1}{Z_{\rm nz}^{(D)}} \int [D\sqrt{\alpha'}\eta^D] e^{-I_{\rm p}^{(D)}[\eta^D,g]} f\left(\eta^D(z)\right), \tag{A.6}$$

where in the second line, we used the fact that the Polyakov action factorises into a sum of two terms: a term containing η^D and another term containing η^i . Now we multiply by 1 and express 1 as a Dirac delta function

$$\left\langle f\left(\eta^{D}(z)\right)\right\rangle = \frac{1}{Z_{\rm nz}^{(D)}} \int [D\sqrt{\alpha'}\eta^{D}] e^{-I_{\rm p}^{(D)}[\eta^{D},g]} f\left(\eta^{D}(z)\right) 1 \tag{A.7}$$

$$=\frac{1}{Z_{\rm nz}^{(D)}}\int [D\sqrt{\alpha'}\eta^D]e^{-I_{\rm p}^{(D)}[\eta^D,g]}f\left(\eta^D(z)\right)\int_{-\infty}^{\infty}d\xi\,\,\delta\left(\xi-\eta^D(z)\right)\tag{A.8}$$

$$=\frac{1}{Z_{\rm nz}^{(D)}}\int_{-\infty}^{\infty}d\xi\int [D\sqrt{\alpha'}\eta^D]e^{-I_{\rm p}^{(D)}[\eta^D,g]}f\left(\sum_{l=1}^{\infty}\sum_{m=-l}^{m=l}x_{lm}Y_{lm}(z_0)\right)\ \delta\left(\xi-\sum_{l=1}^{\infty}\sum_{m=-l}^{m=l}x_{lm}Y_{lm}(z)\right),\tag{A.9}$$

where the measure in terms of the expansion coefficients is now

$$[D\sqrt{\alpha'}\eta^D] = \prod_{l=1}^{\infty} \prod_{m=-l}^{m=l} \sqrt{\alpha'} dx_{lm} \,. \tag{A.10}$$

The Polyakov action of the $\eta^D(z_0)$ is given by (using the round sphere worldsheet metric)

$$I_{\rm p}^{(D)}[\eta^D, g] = \frac{1}{4\pi\alpha'} \int d^2 z \sqrt{g} \left(\alpha' g^{ab} \partial_a \eta^D \partial_b \eta^D \right) + 2\Phi_0 \tag{A.11}$$

$$= -\frac{1}{4\pi\alpha'} \int d^2 z \sqrt{g} \left(\alpha' \eta^D \nabla^2 \eta^D \right) + 2\Phi_0 \,. \tag{A.12}$$

The heat kernel-regularized $I^{(D)}_{\rm p}[\eta^D,g]$ is given by

$$I_{\rm p}^{(D)}[\eta^D, g] = -\frac{1}{4\pi} \int d^2 z \sqrt{g} \left(\eta^D e^{-\epsilon \nabla^2} \nabla^2 \eta^D \right) + 2\Phi_0 \tag{A.13}$$

$$= -\frac{1}{4\pi} \sum_{l=1}^{\infty} \sum_{m=-l}^{m=l} (-l(l+1))e^{\epsilon l(l+1)}x_{lm}^2 + 2\Phi_0.$$
(A.14)

Thus, $\left\langle f\left(\eta^{D}(z)\right) \right\rangle$ is now given by

$$\left\langle f\left(\eta^{D}(z)\right)\right\rangle = \frac{e^{-2\Phi_{0}}}{Z_{nz}^{(D)}} \int_{-\infty}^{\infty} d\xi \int \left(\prod_{l=1}^{\infty} \prod_{m=-l}^{m=l} \sqrt{\alpha'} dx_{lm}\right) e^{-\frac{1}{4\pi} \sum_{l=1}^{\infty} \sum_{m=-l}^{m=l} l(l+1)e^{\epsilon l(l+1)} x_{lm}^{2}}$$
(A.15)

$$f\left(\sum_{l=1}^{\infty}\sum_{m=-l}^{m=l} x_{lm} Y_{lm}(z_0)\right) \ \delta\left(\xi - \sum_{l=1}^{\infty}\sum_{m=-l}^{m=l} x_{lm} Y_{lm}(z_0)\right).$$
(A.16)

Without loss of generality, we can take z to be the north pole.¹⁵ This is because $\langle f(\eta^D(z)) \rangle$ is independent of z due to translation symmetry of the scalar field $\eta^D(z)$ on the worldsheet, which is *not* broken even off-shell. Thus, we can just evaluate it at $z = z_0 = 0$

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{e^{-2\Phi_{0}}}{Z_{\mathrm{nz}}^{(D)}} \int_{-\infty}^{\infty} d\xi \int \left(\prod_{l=1}^{\infty} \prod_{m=-l}^{m=l} \sqrt{\alpha'} dx_{lm}\right) e^{-\frac{1}{4\pi} \sum_{l=1}^{\infty} \sum_{m=-l}^{m=l} l(l+1)e^{\epsilon l(l+1)} x_{lm}^{2}}$$
(A.17)

$$f\left(\sum_{l=1}^{\infty}\sum_{m=-l}^{m=l}x_{lm}Y_{lm}(0)\right) \ \delta\left(\xi - \sum_{l=1}^{\infty}\sum_{m=-l}^{m=l}x_{lm}Y_{lm}(0)\right).$$
(A.18)

At the north pole $(z_0 = 0)$, the spherical hamonics vanish unless m = 0, i.e. $Y_{lm}(0) = 0$ when $m \neq 0$. So we get

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{e^{-2\Phi_{0}}}{Z_{\mathrm{nz}}^{(D)}} \int_{-\infty}^{\infty} d\xi \int \left(\prod_{l=1}^{\infty} \prod_{m=-l}^{m=l} \sqrt{\alpha'} dx_{lm}\right) e^{-\frac{1}{4\pi} \sum_{l=1}^{\infty} \sum_{m=-l}^{m=l} l(l+1)e^{\epsilon l(l+1)} x_{lm}^{2}}$$
(A.19)

$$f\left(\sum_{l=1}^{\infty} x_{l0} Y_{l0}(0)\right) \ \delta\left(\xi - \sum_{l=1}^{\infty} x_{l0} Y_{l0}(0)\right).$$
(A.20)

The $m \neq 0$ terms of the above integral factorises out and cancels the corresponding part in $Z_{nz}^{(D)}$ and we obtain

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{e^{-2\Phi_{0}}}{Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \int \left(\prod_{l=1}^{\infty} \sqrt{\alpha'} dx_{l0}\right) e^{-\frac{1}{4\pi} \sum_{l=1}^{\infty} l(l+1)e^{\epsilon l(l+1)}x_{l0}^{2}}$$
(A.21)

$$f\left(\sum_{l=1}^{\infty} x_{l0} Y_{l0}(0)\right) \ \delta\left(\xi - \sum_{l=1}^{\infty} x_{l0} Y_{l0}(0)\right).$$
(A.22)

Also, we have $Y_{l0}(0) = \sqrt{\frac{2l+1}{4\pi}} =: \omega_l$. So we have

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{e^{-2\Phi_{0}}}{Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \int \left(\prod_{l=1}^{\infty} \sqrt{\alpha'} dx_{l}\right) e^{-\frac{1}{4\pi} \sum_{l=1}^{\infty} l(l+1)e^{\epsilon l(l+1)}x_{l}^{2}}$$
(A.23)

$$f\left(\sum_{l=1}^{\infty}\omega_l x_l\right) \ \delta\left(\xi - \sum_{l=1}^{\infty}\omega_l x_l\right),\tag{A.24}$$

where we have dropped the subscript "0" from x_{l0} and Y_{l0} . We can absorb ω_l into x_l by a change of integration variable to get

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{e^{-2\Phi_{0}}}{Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \int \left(\prod_{l=1}^{\infty} \frac{\sqrt{\alpha'}}{\omega_{l}} dx_{l}\right) e^{-\frac{1}{4\pi} \sum_{l=1}^{\infty} \frac{l(l+1)}{\omega_{l}^{2}} e^{\epsilon l(l+1)} x_{l}^{2}}$$
(A.25)

$$f\left(\sum_{l=1}^{\infty} x_l\right) \ \delta\left(\xi - \sum_{l=1}^{\infty} x_l\right). \tag{A.26}$$

 15 We could have proceeded without picking any point on the sphere at the expense of complicating the calculation bit.

Now we shall pick a mode to integrate over, say x_1 . Integrating over x_1 , we obtain

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{\sqrt{\alpha'}e^{-2\Phi_{0}}}{\omega_{1}Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \ f\left(\xi\right) \int \left(\prod_{l=2}^{\infty} \frac{\sqrt{\alpha'}}{\omega_{l}} dx_{l}\right)$$
(A.27)

$$\exp\left\{-\frac{1}{4\pi}\sum_{l=2}^{\infty}\frac{l(l+1)}{\omega_l^2}e^{\epsilon l(l+1)}x_l^2 - \frac{2}{4\pi\omega_1^2}e^{2\epsilon}\left(\xi - \sum_{l=2}^{\infty}x_l\right)^2\right\}$$
(A.28)

$$= \frac{\sqrt{\alpha'}e^{-2\Phi_0}}{\omega_1 Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \ f(\xi) \ e^{-\frac{2}{4\pi\omega_1^2}e^{2\epsilon}\xi^2} \int \left(\prod_{l=2}^{\infty} \frac{\sqrt{\alpha'}}{\omega_l} dx_l\right)$$
(A.29)

$$\exp\left\{-\frac{1}{4\pi}\sum_{l=2}^{\infty}\frac{l(l+1)}{\omega_l^2}e^{\epsilon l(l+1)}x_l^2 - \frac{2}{4\pi\omega_1^2}e^{2\epsilon}\left(\sum_{l=2}^{\infty}x_l\right)^2 - \frac{4}{4\pi\omega_1^2}e^{2\epsilon}\xi\sum_{l=2}^{\infty}x_l\right\}$$
(A.30)

$$=\frac{\sqrt{\alpha'}e^{-2\Phi_0}}{\omega_1 Z_{\mathrm{nz}(m\neq0)}^{(D)}}\int_{-\infty}^{\infty} d\xi \ f(\xi) \ e^{-\frac{2}{4\pi\omega_1^2}e^{2\epsilon}\xi^2} \int \left(\prod_{l=2}^{\infty}\frac{\sqrt{\alpha'}}{\omega_l}dx_l\right) e^{\Gamma},\tag{A.31}$$

where

$$\Gamma := -\frac{1}{4\pi} \sum_{l=2}^{\infty} \frac{l(l+1)}{\omega_l^2} e^{\epsilon l(l+1)} x_l^2 - \frac{2}{4\pi\omega_1^2} e^{2\epsilon} \left(\sum_{l=2}^{\infty} x_l\right)^2 - \frac{4}{4\pi\omega_1^2} e^{2\epsilon} \xi \sum_{l=2}^{\infty} x_l \tag{A.32}$$

$$= -\frac{1}{4\pi} \sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{n(n+1)}{\omega_n^2} e^{\epsilon n(n+1)} x_n \delta_{nm} x_m - \frac{2}{4\pi\omega_1^2} e^{2\epsilon} \sum_{n=2}^{\infty} \sum_{m=2}^{\infty} x_n x_m - \frac{4}{4\pi\omega_1^2} e^{2\epsilon} \xi \sum_{l=2}^{\infty} x_l$$
(A.33)

$$= -\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{1}{2\pi\omega_1^2} e^{2\epsilon} \left(\frac{\omega_1^2}{\omega_n^2} \frac{n(n+1)}{2} e^{\epsilon n(n+1)-2\epsilon} \delta_{nm} + 1 \right) x_n x_m - \frac{4}{4\pi\omega_1^2} e^{2\epsilon} \xi \sum_{l=2}^{\infty} x_l$$
(A.34)

$$= -M_{nm}\tilde{x}_n\tilde{x}_m - \frac{1}{\omega_1}\sqrt{\frac{2}{\pi}}e^{\epsilon}\xi\sum_{l=2}^{\infty}\tilde{x}_l,\tag{A.35}$$

and

$$M_{nm} := \left(\tilde{\lambda}_n \delta_{nm} + 1\right) \tag{A.36}$$

$$\tilde{\lambda}_n := \frac{\omega_1^2}{\omega_n^2} \frac{n(n+1)}{2} e^{\epsilon n(n+1)-2\epsilon}$$
(A.37)

$$\tilde{x}_n := \frac{1}{\sqrt{2\pi\omega_1}} e^{\epsilon} x_n \tag{A.38}$$

So we get

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \left(\prod_{l=2}^{\infty} \frac{\omega_{1}}{\omega_{l}}\right) \frac{\sqrt{\alpha'}e^{-2\Phi_{0}}}{\omega_{1}Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \ f\left(\xi\right) e^{-\frac{1}{2\pi\omega_{1}^{2}}e^{2\epsilon}\xi^{2}} \int \left(\prod_{l=2}^{\infty} \sqrt{2\pi\alpha'}e^{-\epsilon}d\tilde{x}_{l}\right) e^{\Gamma},\tag{A.39}$$

Now let us diagonalise the real symmetric matrix M by an orthogonal transformation $\tilde{x}_n = \Lambda_{nm} y_m$ to get a diagonal matrix $D = \Lambda^T M \Lambda$

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \left(\prod_{l=2}^{\infty} \frac{\omega_{1}}{\omega_{l}}\right) \frac{\sqrt{\alpha'}e^{-2\Phi_{0}}}{\omega_{1}Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \ f\left(\xi\right) e^{-\frac{1}{2\pi\omega_{1}^{2}}e^{2\epsilon}\xi^{2}} \frac{1}{\mathrm{Det}(\Lambda)} \int \left(\prod_{n=2}^{\infty} \sqrt{2\pi\alpha'}e^{-\epsilon}dy_{n}\right)$$
(A.40)

$$\exp\left\{-\sum_{n=2}^{\infty}D_ny_n^2-\xi\sum_{n=2}^{\infty}k_ny_n\right\},\tag{A.41}$$

where $k_n := \frac{1}{\omega_1} \sum_{m=2}^{\infty} \sqrt{\frac{2}{\pi}} e^{\epsilon} \Lambda_{mn}$ and a factor of $\text{Det}(\Lambda)$ comes from the Jacobian. Now one can complete the squares

and shift the mean to get (one can check that all $D_n > 0$)

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \left(\prod_{l=2}^{\infty} \frac{\omega_{1}}{\omega_{l}}\right) \frac{\sqrt{\alpha'}e^{-2\Phi_{0}}}{\omega_{1}Z_{\mathrm{nz}(m\neq0)}^{(D)}} \int_{-\infty}^{\infty} d\xi \ f\left(\xi\right) e^{-\frac{1}{2\pi\omega_{1}^{2}}e^{2\epsilon}\xi^{2}} \frac{1}{\mathrm{Det}(\Lambda)} \exp\left\{\sum_{n=2}^{\infty} \frac{k_{n}^{2}}{4D_{n}}\xi^{2}\right\}$$
(A.42)

$$\prod_{n=2}^{\infty} \int \sqrt{2\pi\alpha'} e^{-\epsilon} dy_n \exp\left\{-D_n \left(y_n + \xi \frac{k_n}{2D_n}\right)^2\right\}$$
(A.43)

$$= \left(\prod_{l=2}^{\infty} \frac{\omega_1}{\omega_l}\right) \frac{\sqrt{\alpha'} e^{-2\Phi_0}}{\omega_1 Z_{\mathrm{nz}(m\neq0)}^{(D)}} \frac{1}{\mathrm{Det}(\Lambda)} \frac{1}{\sqrt{\mathrm{Det}(M)}} \left(\prod_{n=2}^{\infty} \sqrt{2\alpha'} \pi e^{-\epsilon}\right)$$
(A.44)

$$\int_{-\infty}^{\infty} d\xi \ f(\xi) \ e^{-\frac{1}{2\pi\omega_1^2}e^{2\epsilon}\xi^2} \exp\left\{\sum_{n=2}^{\infty} \frac{k_n^2}{4D_n}\xi^2\right\}.$$
(A.45)

All of the annoying factors in (A.44) gets absorbed into $Z_{\mathrm{nz}(m\neq 0)}^{(D)}$ and we finally get

$$\left\langle f\left(\eta^{D}(z_{0})\right)\right\rangle = \frac{\int_{-\infty}^{\infty} d\xi \ f\left(\xi\right) e^{-\xi^{2}/\Omega}}{\int_{-\infty}^{\infty} d\xi \ e^{-\xi^{2}/\Omega}},\tag{A.46}$$

where

$$\Omega^{-1} := \frac{e^{2\epsilon}}{2\pi\omega_1^2} \left(1 - \mathbb{M}\right) \,, \tag{A.47}$$

$$\mathbb{M} := \omega_1^2 \frac{\pi}{2} e^{-2\epsilon} \sum_{n=2}^{\infty} \frac{k_n^2}{D_n} \,. \tag{A.48}$$

Now let us calculate $\mathbb M.$

$$\mathbb{M} = \omega_1^2 \frac{\pi}{2} e^{-2\epsilon} \sum_{n=2}^{\infty} \frac{k_n^2}{D_n} = \frac{\pi}{2} e^{-2\epsilon} \sum_{n=2}^{\infty} \sum_{\bar{m}=2}^{\infty} \sum_{\bar{m}=2}^{\infty} \sqrt{\frac{2}{\pi}} e^{\epsilon} \Lambda_{\bar{m}n} \frac{1}{D_n} \sqrt{\frac{2}{\pi}} e^{\epsilon} \Lambda_{\bar{m}n}$$
(A.49)

$$=\sum_{n=2}^{\infty}\sum_{m=2}^{\infty}[\Lambda D^{-1}\Lambda^{T}]_{nm} = \sum_{n=2}^{\infty}\sum_{m=2}^{\infty}[M^{-1}]_{nm},$$
(A.50)

which follows from Λ being an orthogonal matrix. So M is the sum of all the entries of the inverse of the matrix M. One can show that

$$\mathbb{M} = \frac{1}{1 + \frac{1}{\sum_{n=2}^{\infty} \tilde{\lambda}_n^{-1}}} \ . \tag{A.51}$$

One can also show that

$$\Omega = \left(\frac{e^{2\epsilon}}{2\pi\omega_1^2} \left(1 - \mathbb{M}\right)\right)^{-1} = \frac{2\pi\omega_1^2}{e^{2\epsilon}} \left(1 + \sum_{n=2}^{\infty} \tilde{\lambda}_n^{-1}\right) = \frac{3}{2}e^{-2\epsilon} + \sum_{n=2}^{\infty} \frac{2n+1}{n(n+1)}e^{-\epsilon n(n+1)}$$
(A.52)

$$=\sum_{n=1}^{\infty} \frac{2n+1}{n(n+1)} e^{-\epsilon n(n+1)} \,. \tag{A.53}$$

If we specialize to $f(\eta^D(z_0)) = |\eta^D(z_0)|^N$, then the general formula in equation (A.46) reduces to

$$\left\langle \left| \eta^{D}(z_{0}) \right|^{N} \right\rangle = \frac{\int_{-\infty}^{\infty} d\xi \, |\xi|^{N} e^{-\xi^{2}/\Omega}}{\int_{-\infty}^{\infty} d\xi \, e^{-\xi^{2}/\Omega}} = \frac{\int_{0}^{\infty} d\xi \, \xi^{N} e^{-\xi^{2}/\Omega}}{\int_{0}^{\infty} d\xi \, e^{-\xi^{2}/\Omega}} \tag{A.54}$$

$$=\frac{1}{\sqrt{\pi}}\Gamma\left(\frac{1+N}{2}\right)\Omega^{N/2}.$$
(A.55)

We thus get

$$\left\langle \left| \eta^D(z_0) \right|^N \right\rangle = \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{1+N}{2}\right) \left(\sum_{n=1}^\infty \frac{2n+1}{n(n+1)} e^{-\epsilon n(n+1)} \right)^{N/2} . \tag{A.56}$$

The N = 2 case is a standard result, so we shall compare our result with it. We know that [3]

$$\begin{split} \langle \eta^{\mu}(z)\eta^{\nu}(z) \rangle &= 2\pi \alpha' \delta^{\mu\nu} \left(\frac{1}{4\pi} \sum_{n=1}^{\infty} \frac{2n+1}{n(n+1)} e^{-\epsilon n(n+1)} \right) \\ &= 2\pi \alpha' \delta^{\mu\nu} \left(-\frac{1}{2\pi} \alpha' \log \epsilon + O(\epsilon^2) + \text{finite term} \right) \,. \end{split}$$

Thus, we find from (A.52) that $\Omega = 2 \log e^{-1}$.

Now we focus on the original term we set out to compute: $\langle |Y^D + \sqrt{\alpha'}\eta^D(z_0)| \rangle$. For this, the general formula in equation (A.46) reduces to

$$\left\langle \left| Y^D + \sqrt{\alpha'} \eta^D(z_0) \right| \right\rangle = \frac{\int_{-\infty}^{\infty} d\xi \left| Y^D + \sqrt{\alpha'} \xi \right| e^{-\xi^2 / \Omega}}{\int_{-\infty}^{\infty} d\xi \left| e^{-\xi^2 / \Omega} \right|} = \frac{1}{\sqrt{\pi\Omega}} \int_{-\infty}^{\infty} d\xi \left| Y^D + \sqrt{\alpha'} \xi \right| e^{-\xi^2 / \Omega}$$
(A.57)

$$=\frac{1}{\sqrt{\pi\Omega}}\int_{-\frac{YD}{\sqrt{\alpha'}}}^{\infty}d\xi \left(Y^D + \sqrt{\alpha'}\xi\right)e^{-\xi^2/\Omega} - \frac{1}{\sqrt{\pi\Omega}}\int_{-\infty}^{-\frac{YD}{\sqrt{\alpha'}}}d\xi \left(Y^D + \sqrt{\alpha'}\xi\right)e^{-\xi^2/\Omega} \quad (A.58)$$

$$=\frac{Y^D}{\sqrt{\pi\Omega}}\int_{-\frac{Y^D}{\sqrt{\alpha'}}}^{\infty} d\xi \ e^{-\xi^2/\Omega} - \frac{Y^D}{\sqrt{\pi\Omega}}\int_{-\infty}^{-\frac{Y^D}{\sqrt{\alpha'}}} d\xi \ e^{-\xi^2/\Omega}$$
(A.59)

$$+\frac{\sqrt{\alpha'}}{\sqrt{\pi\Omega}}\int_{-\frac{YD}{\sqrt{\alpha'}}}^{\infty}d\xi\;\xi e^{-\xi^2/\Omega} - \frac{\sqrt{\alpha'}}{\sqrt{\pi\Omega}}\int_{-\infty}^{-\frac{YD}{\sqrt{\alpha'}}}d\xi\;\xi e^{-\xi^2/\Omega} \tag{A.60}$$

$$=\frac{Y^{D}}{\sqrt{\pi\Omega}}\frac{\sqrt{\pi\Omega}}{2}\left(1+\operatorname{erf}\left(\frac{Y^{D}}{\sqrt{\alpha'\Omega}}\right)\right)-\frac{Y^{D}}{\sqrt{\pi\Omega}}\frac{\sqrt{\pi\Omega}}{2}\left(\operatorname{erfc}\left(\frac{Y^{D}}{\sqrt{\alpha'\Omega}}\right)\right) \tag{A.61}$$

$$+\frac{\sqrt{\alpha'}}{\sqrt{\pi\Omega}}\frac{\Omega}{2}e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{\sqrt{\alpha'}}{\sqrt{\pi\Omega}}\frac{\Omega}{2}e^{-\frac{(Y^D)^2}{\alpha'\Omega}}$$
(A.62)

$$=\sqrt{\Omega}\frac{\sqrt{\alpha'}}{\sqrt{\pi}}e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{Y^D}{2}\left(1 + \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right) - \operatorname{erfc}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right)\right)$$
(A.63)

$$=\sqrt{\Omega}\frac{\sqrt{\alpha'}}{\sqrt{\pi}}e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + Y^D \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right).$$
(A.64)

So finally, we obtain our desired expression

$$\left\langle \left| Y^{D} + \sqrt{\alpha'} \eta^{D}(z_{0}) \right| \right\rangle = \sqrt{\alpha'} \left(\frac{\sqrt{\Omega}}{\sqrt{\pi}} e^{-\frac{(Y^{D})^{2}}{\alpha'\Omega}} + \frac{Y^{D}}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^{D}}{\sqrt{\alpha'\Omega}}\right) \right)$$
(A.65)

$$= \sqrt{\alpha'} \left(\left\langle \left| \eta^D(z_0) \right| \right\rangle e^{-\frac{(Y^D)^2}{\alpha'\Omega}} + \frac{Y^D}{\sqrt{\alpha'}} \operatorname{erf}\left(\frac{Y^D}{\sqrt{\alpha'\Omega}}\right) \right).$$
(A.66)

where we used (A.54) in the last line that $\langle |\eta^D(z_0)| \rangle = \frac{\sqrt{\Omega}}{\sqrt{\pi}}$.

References

- V.A. Kazakov and A.A. Tseytlin, On free energy of 2-D black hole in bosonic string theory, JHEP 06 (2001) 021 [hep-th/0104138].
- [2] A.A. Tseytlin, Partition Function of String σ Model on a Compact Two Space, Phys. Lett. B 223 (1989) 165.
- [3] O.D. Andreev, R.R. Metsaev and A.A. Tseytlin, Covariant calculation of the statistical sum of the two-dimensional sigma model on compact two surfaces., Sov. J. Nucl. Phys. 51 (1990) 359 [2301.02867].
- [4] A. Ahmadain and A.C. Wall, Off-Shell Strings II: Black Hole Entropy, 2211.16448.
- [5] A.A. Tseytlin, Mobius Infinity Subtraction and Effective Action in σ Model Approach to Closed String Theory, Phys. Lett. B 208 (1988) 221.

- [6] A. Ahmadain and A.C. Wall, Off-Shell Strings I: S-matrix and Action, 2211.08607.
- [7] H. Erbin, String Field Theory: A Modern Introduction, vol. 980 of Lecture Notes in Physics (3, 2021), 10.1007/978-3-030-65321-7.
- [8] J. Liu and J. Polchinski, Renormalization of the Mobius Volume, Phys. Lett. B 203 (1988) 39.
- [9] E.S. Fradkin and A.A. Tseytlin, Quantum String Theory Effective Action, Nucl. Phys. B 261 (1985) 1.
 [Erratum: Nucl.Phys.B 269, 745–745 (1986)].
- [10] E.S. Fradkin and A.A. Tseytlin, Effective Field Theory from Quantized Strings, Phys. Lett. B 158 (1985) 316.
- [11] E.S. Fradkin and A.A. Tseytlin, Effective Action Approach to Superstring Theory, Phys. Lett. B 160 (1985) 69.
- [12] A.A. Tseytlin, Tachyon effective actions in string theory, Theor. Math. Phys. 128 (2001) 1293.
- [13] A. Adams, J. Polchinski and E. Silverstein, Don't panic! Closed string tachyons in ALE space-times, JHEP 10 (2001) 029 [hep-th/0108075].
- [14] L. Eberhardt and S. Pal, Holographic Weyl anomaly in string theory, SciPost Phys. 16 (2024) 027 [2307.03000].
- [15] A.A. Tseytlin, Sigma model approach to string theory effective actions with tachyons, J. Math. Phys. 42 (2001) 2854 [hep-th/0011033].
- [16] A.A. Tseytlin, On sigma model RG flow, 'central charge' action and Perelman's entropy, Phys. Rev. D 75 (2007) 064024 [hep-th/0612296].
- [17] P. Kraus, A. Ryzhov and M. Shigemori, Strings in noncompact space-times: Boundary terms and conserved charges, Phys. Rev. D 66 (2002) 106001 [hep-th/0206080].
- [18] T. Erler, The closed string field theory action vanishes, JHEP 10 (2022) 055 [2204.12863].
- [19] A. Sen and B. Zwiebach, String Field Theory: A Review, 2405.19421.
- [20] Y. Chen, J. Maldacena and E. Witten, On the black hole/string transition, Journal of High Energy Physics 2023 (2023) 1.
- [21] T. Erler, The closed string field theory action vanishes, 2204.12863.
- [22] J. Troost, The ads3 central charge in string theory, Physics Letters B 705 (2011) 260.
- [23] J.W. York, Jr., Role of conformal three geometry in the dynamics of gravitation, Phys. Rev. Lett. 28 (1972) 1082.
- [24] G.W. Gibbons and S.W. Hawking, Action Integrals and Partition Functions in Quantum Gravity, Phys. Rev. D 15 (1977) 2752.
- [25] L. Eberhardt and S. Pal, The disk partition function in string theory, JHEP 08 (2021) 026 [2105.08726].
- [26] R. Mahajan, D. Stanford and C. Yan, Sphere and disk partition functions in Liouville and in matrix integrals, JHEP 07 (2022) 132 [2107.01172].
- [27] B. Mühlmann, The two-sphere partition function in two-dimensional quantum gravity at fixed area, JHEP 09 (2021) 189 [2106.04532].
- [28] L. Eberhardt and S. Pal, Holographic weyl anomaly in string theory, arXiv preprint arXiv:2307.03000 (2023).
- [29] T.G. Mertens, Hagedorn String Thermodynamics in Curved Spacetimes and near Black Hole Horizons, Ph.D. thesis, Gent U., 2015. 1506.07798.

- [30] F. Bastianelli, O. Corradini and P.A.G. Pisani, Worldline approach to quantum field theories on flat manifolds with boundaries, JHEP 02 (2007) 059 [hep-th/0612236].
- [31] F. Bastianelli, O. Corradini, P.A.G. Pisani and C. Schubert, Scalar heat kernel with boundary in the worldline formalism, JHEP 10 (2008) 095 [0809.0652].
- [32] F. Bastianelli, O. Corradini, P.A.G. Pisani and C. Schubert, Worldline Approach to QFT on Manifolds with Boundary, in 9th Conference on Quantum Field Theory under the Influence of External Conditions (QFEXT 09): Devoted to the Centenary of H. B. G. Casimir, pp. 415–420, 2010, DOI [0912.4120].
- [33] E.W. Weisstein, "Half-normal distribution." from mathworld-a wolfram web resource." https://mathworld.wolfram.com/Half-NormalDistribution.html.
- [34] P. Mörters and Y. Peres, Brownian motion, vol. 30, Cambridge University Press (2010).
- [35] I. Karatzas and S. Shreve, Brownian motion and stochastic calculus, vol. 113, springer (2014).
- [36] P. Lévy, Processus stochastiques et mouvement brownien (paris: Gauthier-villars) google scholar lévy p 1951, in Proc. 2nd Berkeley Symp. on Mathematical Statistics and Probability, 1965.
- [37] K. Ito and H. McKean, Diffusion processes and their sample paths, Die Grundlehren der Mathematischen Wissenschaften in Einzeldarstellungen 125 (1965).
- [38] P.-L. Lions and A.-S. Sznitman, Stochastic differential equations with reflecting boundary conditions, Communications on pure and applied Mathematics 37 (1984) 511.
- [39] Y. Saisho, Stochastic differential equations for multi-dimensional domain with reflecting boundary, Probability Theory and Related Fields 74 (1987) 455.
- [40] P. Hsu, Probabilistic approach to the neumann problem, Communications on pure and applied mathematics 38 (1985) 445.
- [41] A.N. Borodin and P. Salminen, Handbook of Brownian motion-facts and formulae, Springer Science & Business Media (2015).
- [42] D.S. Grebenkov, Probability distribution of the boundary local time of reflected brownian motion in euclidean domains, Physical Review E 100 (2019) 062110.
- [43] K. Siegrist, "Probability, mathematical statistics, stochastic processes." https://www.randomservices.org/random/special/FoldedNormal.html/.
- [44] C. Banderier and M. Wallner, Local time for lattice paths and the associated limit laws, arXiv preprint arXiv:1805.09065 (2018).
- [45] M. Wallner, A half-normal distribution scheme for generating functions, European Journal of Combinatorics 87 (2020) 103138.
- [46] A.V. Skorokhod, Stochastic equations for diffusion processes in a bounded region, Theory of Probability & Its Applications 6 (1961) 264.
- [47] A.V. Skorokhod, Stochastic equations for diffusion processes in a bounded region. ii, Theory of Probability & Its Applications 7 (1962) 3.
- [48] H. Tanaka, Stochastic differential equations with reflecting, Stochastic Processes: Selected Papers of Hiroshi Tanaka 9 (1979) 157.
- [49] K.L. Chung, Z. Zhao, K.L. Chung and Z. Zhao, Killed brownian motion, From Brownian Motion to Schrödinger's Equation (1995) 31.
- [50] T. Björk, The pedestrian's guide to local time, 1512.08912.
- [51] H.P. McKean, *Stochastic integrals*, vol. 353, American Mathematical Soc. (1969).

- [52] T.E. Clark, R. Menikoff and D.H. Sharp, Quantum mechanics on the half-line using path integrals, Phys. Rev. D 22 (1980) 3012.
- [53] L. Takács, On the local time of the brownian motion, The Annals of Applied Probability (1995) 741.
- [54] G.W. Gibbons and S.W. Hawking, Action integrals and partition functions in quantum gravity, Phys. Rev. D 15 (1977) 2752.
- [55] S. Ryu and T. Takayanagi, Holographic derivation of entanglement entropy from the anti-de sitter space/conformal field theory correspondence, Phys. Rev. Lett. 96 (2006) 181602.
- [56] A. Lewkowycz and J. Maldacena, Generalized gravitational entropy, JHEP 08 (2013) 090 [1304.4926].
- [57] D.L. Jafferis and E. Schneider, Stringy ER=EPR, 2104.07233.
- [58] I. Halder and D.L. Jafferis, Thermal Bekenstein-Hawking entropy from the worldsheet, 2310.02313.
- [59] A. Ahmadain, V. Shyam and Y. Zihan, One-half the Einstein boundary term from the string worldsheet, forthcoming, .
- [60] A. Einstein, Hamilton's Principle and the General Theory of Relativity, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) 1916 (1916) 1111.
- [61] A.B. Zamolodchikov, Expectation value of composite field T anti-T in two-dimensional quantum field theory, hep-th/0401146.
- [62] F.A. Smirnov and A.B. Zamolodchikov, On space of integrable quantum field theories, Nucl. Phys. B 915 (2017) 363 [1608.05499].
- [63] A. Cavaglià, S. Negro, I.M. Szécsényi and R. Tateo, TT-deformed 2D Quantum Field Theories, JHEP 10 (2016) 112 [1608.05534].
- [64] L. McGough, M. Mezei and H. Verlinde, Moving the CFT into the bulk with $T\overline{T}$, JHEP **04** (2018) 010 [1611.03470].
- [65] G. Araujo-Regado, R. Khan and A.C. Wall, Cauchy slice holography: a new AdS/CFT dictionary, JHEP 03 (2023) 026 [2204.00591].
- [66] E. Silverstein, Black hole to cosmic horizon microstates in string/m theory: timelike boundaries and internal averaging, Journal of High Energy Physics **2023** (2023) 1 [2212.00588].
- [67] G. Batra, G.B. De Luca, E. Silverstein, G. Torroba and S. Yang, Bulk-local dS₃ holography: the Matter with $T\bar{T} + \Lambda_2$, 2403.01040.