W¹,¹ STABILITY FOR THE LSI

EMANUEL INDREI

Abstract. The logarithmic Sobolev inequality is fundamental in mathematical physics. Associated stability estimates are equivalent to uncertainty principles. Via a second moment bound, $W^{1,1}$ estimates are obtained in one dimension and similar W_1 quantitative estimates are investigated.

1. INTRODUCTION

The classical logarithmic Sobolev inequality LSI attributed to L. Gross states

(1.1)
$$
\delta(f) := \frac{1}{2}\mathbf{I}(f) - \mathbf{H}(f) = \frac{1}{2}\int \frac{|\nabla f|^2}{f} d\gamma - \int f \log f d\gamma \ge 0,
$$

where $d\gamma = (2\pi)^{-\frac{n}{2}}e^{-\frac{|x|^2}{2}}dx$, f is normalized, $\sqrt{f} \in W^{1,2}(\mathbb{R}^n, d\gamma)$, & δ is the LSI deficit [\[3,](#page-9-0) [6\]](#page-9-1). Carlen [\[2\]](#page-9-2) characterized the equality cases: equality holds in [\(1.1\)](#page-0-0) if and only if $f_b(x) = e^{b \cdot x - \frac{b^2}{2}}$, $b \in \mathbb{R}^n$. Note that if f is normalized and centered, equality is valid if and only if $f = 1$. Therefore, a natural problem is to identify a metric $d, a > 0$, and $\alpha > 0$ such that

(1.2)
$$
\delta(f) \ge ad^{\alpha}(f, 1),
$$

with f normalized and centered. In a recent paper, a variant of (1.2) with $\alpha = 2$, a constant $a > 0$, and $d(f, h) = ||f - h||_{L^1(\mathbb{R}^n, d\gamma)}$ was established [\[4\]](#page-9-3). In [\[7\]](#page-9-4), $\alpha = 2$ was proven to be sharp. Thus, the problem of identifying the strongest norm was a natural consequence which was addressed in [\[7\]](#page-9-4). Set

$$
f = |u|^2 \left(\frac{x}{\sqrt{2\pi}}\right).
$$

Then

$$
\int f e^{-\frac{|x|^2}{2}} (2\pi)^{-\frac{n}{2}} dx = \int |u|^2 e^{-\pi |x|^2} dx.
$$

$$
\frac{1}{2} \int \frac{|\nabla f|^2}{f} d\gamma - \int f \ln f d\gamma = \frac{1}{\pi} \int |\nabla u|^2 e^{-\pi |x|^2} dx - \int |u|^2 \ln |u|^2 e^{-\pi |x|^2} dx.
$$

Therefore [\(1.2\)](#page-0-1) has an equivalent version

$$
\pi \delta_*(u) := \int |\nabla u|^2 e^{-\pi |x|^2} dx - \pi \int |u|^2 \ln |u|^2 e^{-\pi |x|^2} dx \ge \kappa \int |u-1|^2 e^{-\pi |x|^2} dx
$$

in the space of non-negative functions which satisfy

$$
||u||_{L^2(\mathbb{R}^n, e^{-\pi |x|^2} dx)} = 1
$$

$$
\int x|u|^2 e^{-\pi |x|^2} dx = 0.
$$

Suppose $\delta_*(u_k) \to 0$ as $k \to \infty$, then [\[7\]](#page-9-4) yields

 $|u_k| \to 1$

in $H^1(e^{-\pi|x|^2}dx)$ if and only if

$$
\int |x|^2 |u_k(x)|^2 e^{-\pi |x|^2} dx \to \int |x|^2 e^{-\pi |x|^2} dx.
$$

Thus, some additional assumptions are necessary to prove

(1.3)
$$
\delta_*(u) \ge ad^\alpha(u,1)
$$

with $d(u, g) = ||u - g||_{H^1(e^{-\pi |x|^2} dx)}$. Supposing the associated measures satisfy a Poincaré inequality via $\lambda > 0$, the stability with the sharp $\alpha = 2$ and also best constant $a =$ $a(\lambda) > 0$ was proven in [\[5\]](#page-9-5). It has already been underscored in [\[1\]](#page-9-6) that a uniform upper bound on the second moment is necessary but not sufficient to prove [\(1.3\)](#page-1-0) with the $H^1 = W^{1,2}$ norm if $\delta_*(u) \to 0$. A sufficient condition was shown with some type of uniform exponential moment condition. Observe a fourth moment bound also yields [\(1.3\)](#page-1-0) [\[7\]](#page-9-4). The problem of understanding the extra information that is needed with assuming a second moment bound was discussed on [\[1,](#page-9-6) p. 6]. The main result in my paper is that if $n = 1$, while the second moment control does not imply stability in $W^{1,2}$, it indeed implies stability in $W^{1,1}$:

Theorem 1.1. *(1)* If *f is normalized and centered in* $L^1(\mathbb{R}, d\gamma)$ &

$$
m_2(fd\gamma) := \int_{\mathbb{R}} |x|^2 f d\gamma \le \alpha < \infty,
$$

then there exists $a_{\alpha} > 0$ *so that*

$$
||f-1||_{W^{1,1}(\mathbb{R},d\gamma)} \leq a_{\alpha} \left(\delta^{\frac{1}{4}}(f) + \delta^{\frac{3}{4}}(f) \right).
$$

(2) If *u is normalized and centered in* $L^2(\mathbb{R}, e^{-\pi|x|^2}dx)$ &

$$
\int |x|^2 |u(x)|^2 e^{-\pi |x|^2} dx \le \alpha < \infty,
$$

then there exists $\overline{a}_{\alpha} > 0$ *so that*

$$
||u-1||_{W^{1,1}(\mathbb{R},e^{-\pi|x|^2}dx)} \leq \overline{a}_{\alpha}(\delta_*(u)^{\frac{1}{4}} + \delta_*(u)).
$$

Therefore a natural curiosity is the higher dimensional analog but that remains an open problem. However the following can be proven:

Corollary 1.2. (1) If $f(x_1, x_2, \ldots, x_n) = \prod_{k=1}^n f_k(x_k)$, f_k is normalized and centered in $L^1(\mathbb{R},d\gamma)$ &

$$
m_2(f_k) \leq \alpha,
$$

then

$$
||f - 1||_{W^{1,1}(\mathbb{R}^n, d\gamma)} \le a_\alpha n^{3/4} \left(\delta^{\frac{1}{4}}(f) + \delta^{\frac{3}{4}}(f)\right),
$$

with a_{α} *as in Theorem 1.1.*

(2) If
$$
u(x_1, x_2,..., x_n) = \prod_{k=1}^n u_k(x_k)
$$
, u_k is normalized and centered in $L^2(\mathbb{R}, e^{-\pi |x|^2} dx)$

$$
\int |x_k|^2 |u_k(x_k)|^2 e^{-\pi |x_k|^2} dx_k \leq \alpha < \infty,
$$

then

$$
||u-1||_{W^{1,1}(\mathbb{R}^n,e^{-\pi|x|^2}dx)} \leq n\overline{a}_{\alpha}(\delta_*^{\frac{1}{4}}(u)+\delta_*(u)),
$$

with \overline{a}_{α} *as in Theorem 1.1.*

 $\int_{\mathbb{R}^n} |x|^p d\mu < \infty$, $m_p(d\nu) = \int_{\mathbb{R}^n} |x|^p d\nu < \infty$ is The *Wasserstein distance* between two probability measures μ , ν with $p \geq 1$, $m_p(d\mu)$

$$
W_p(d\mu, d\nu) = \inf_{\pi} \left(\iint_{\mathbb{R}^n \times \mathbb{R}^n} |x - y|^p d\pi(x, y) \right)^{\frac{1}{p}}
$$

where the infimum is taken over all probability measures π on $\mathbb{R}^n \times \mathbb{R}^n$ with marginals μ and ν . In particular, W_1 is called the Kantorovich–Rubinstein distance. The stability for W_1 has already appeared in [\[8\]](#page-9-7): let $fd\gamma$ be a centered probability measure, $m_2(fd\gamma) = \int_{\mathbb{R}^n} |x|^2fd\gamma \leq M < \infty$. There exists a constant $C = C(n, M) > 0$ such that

 $\delta(f) \ge C \min\{W_1(fd\gamma,d\gamma),W_1^4(fd\gamma,d\gamma)\}.$

 W_1 -stability is not true if one merely has finite second moments, therefore $C = C(n, M)$ cannot be taken independent of M [\[9,](#page-9-8) Theorem 1.2]. Examples in [\[9\]](#page-9-8) allude to

 $\delta(f) \geq CW_1^2(fd\gamma,d\gamma),$

with $f \in \{f \geq 0 : m_2(f d\gamma) \leq M, ||f||_{L^1(\mathbb{R}^n,d\gamma)} = 1, \int_{\mathbb{R}^n} x f d\gamma = 0\}$. Supposing a type of uniform exponential moment condition, this holds:

Theorem 1.3. If f is normalized and centered in $L^1(\mathbb{R}^n, d\gamma)$, $\alpha > 0$, $\epsilon > 0$, &

$$
\int \int f(x)f(y)e^{\epsilon|x-y|^2}d\gamma(x)d\gamma(y) \le \alpha,
$$

then there exists $a_{\alpha,\epsilon} > 0$ *so that*

$$
W_1(fd\gamma,d\gamma) \le a_{\alpha,\epsilon} \delta^{\frac{1}{2}}(u).
$$

Also, supposing ϵ < .25 *combined with*

$$
\alpha > \frac{1}{(2\pi)^n} \left(\frac{\pi}{-2\epsilon + .5}\right)^n,
$$

the best possible $a_{\alpha,\epsilon}$ *necessarily has a lower bound:*

$$
a_{\alpha,\epsilon} \ge \frac{|nm_1(\gamma) - m_3(\gamma)|}{\sqrt{n}}.
$$

When $n = 1$ *,*

$$
a_{\alpha,\epsilon}\geq \sqrt{\frac{2}{\pi}}.
$$

In addition, the exponent on the deficit is sharp.

2. The Proofs

Proof of Theorem 1.1. The first step is to consider a probability measure $fd\gamma$ & T = $\nabla \Phi$ the Brenier map which pushes forward $fd\gamma$ to $d\gamma$. The proof of [\(1.1\)](#page-0-0) via optimal transport [\[3\]](#page-9-0) implies

$$
2\delta(f) \ge \int |T(x) - x + \nabla \ln f|^2 f d\gamma :
$$

define $\mu := \Phi - \frac{1}{2}$ $\frac{1}{2}|x|^2$ so that

$$
\det(I + D^2 \mu(x))e^{-|x + \nabla \mu(x)|^2/2} = f(x)e^{-|x|^2/2}.
$$

Therefore taking the ln and integrating,

$$
\int f \ln f d\gamma \le \int f \left[\Delta \mu - x \cdot \nabla \mu \right] d\gamma - \frac{1}{2} \int |\nabla \mu|^2 f d\gamma
$$

=
$$
- \int \nabla \mu \cdot \nabla f d\gamma - \frac{1}{2} \int |\nabla \mu|^2 f d\gamma
$$

=
$$
- \frac{1}{2} \int \left| \nabla \mu + \frac{\nabla f}{f} \right|^2 f d\gamma + \frac{1}{2} \int \frac{|\nabla f|^2}{f} d\gamma.
$$

Note this readily implies

$$
\frac{1}{2}\int \left|T - x + \nabla \ln f\right|^2 f d\gamma \le \frac{1}{2}\int \frac{|\nabla f|^2}{f} d\gamma - \int f \ln f d\gamma,
$$

which thanks to Jensen's inequality yields

(2.1)
$$
\delta(f) \ge \frac{1}{2} \int |T(x) - x + \nabla \ln f|^2 f d\gamma \ge \frac{1}{2} \Big(\int |T(x) - x + \nabla \ln f| f d\gamma \Big)^2,
$$

$$
\int |\nabla \ln f| f d\gamma - \int |T(x) - x| f d\gamma \le \int |T(x) - x + \nabla \ln f| f d\gamma \le \sqrt{2\delta(f)}.
$$

Hence since $n = 1$

Hence since $n = 1$,

$$
\int |\nabla \ln f| f d\gamma \le \int |T(x) - x| f d\gamma + \sqrt{2\delta(f)} = W_1(f d\gamma, d\gamma) + \sqrt{2\delta(f)}.
$$

Furthermore, the end of the proof of $[8,$ Proposition C.1] in addition to $[5,$ Corollary 6] (in the one-dimensional argument, positivity & the local boundedness can be removed) imply

(2.2)
$$
\frac{1}{2} \int \frac{|\nabla f|^2}{f} d\gamma - \int f \ln f d\gamma \ge \frac{1}{4n} \Big(2 \int f \ln f d\gamma + (m_2(\gamma) - m_2(f d\gamma)) \Big)^2,
$$

(2.3)
$$
W_1(fd\gamma,d\gamma) \le a \max\{(H(f)\delta(f))^{\frac{1}{4}}, (H(f)\delta(f))^{\frac{1}{2}}\}.
$$

Since

$$
m_2(fd\gamma) \le \alpha,
$$

\n
$$
2H = 2 \int f \ln f d\gamma
$$

\n
$$
\le 2\delta^{\frac{1}{2}}(f) + \alpha + m_2(d\gamma).
$$

Note

$$
\max\{(H(f)\delta(f))^{\frac{1}{4}}, (H(f)\delta(f))^{\frac{1}{2}}\}
$$

\n
$$
\leq t_a \max\{(\delta^{\frac{1}{2}}(f) + r_a)\delta(f))^{\frac{1}{4}}, (\delta^{\frac{1}{2}}(f) + r_a)\delta(f))^{\frac{1}{2}}\}
$$

\n
$$
= t_a \max\{(\delta^{\frac{3}{2}}(f) + r_a\delta(f))^{\frac{1}{4}}, (\delta^{\frac{3}{2}}(f) + r_a\delta(f))^{\frac{1}{2}}\};
$$

assuming

$$
l_a \leq \delta(f) \leq L_a,
$$

$$
W_1(fd\gamma,d\gamma) \le a \max\{(H(f)\delta(f))^{\frac{1}{4}}, (H(f)\delta(f))^{\frac{1}{2}}\} \le c_a = \frac{c_a}{l_a}l_a \le \frac{c_a}{l_a}\delta(f).
$$

Supposing

$$
\delta(f) \le l_a
$$

and $l_a<<1,$

$$
\max\{(\delta^{\frac{3}{2}}(f) + r_a \delta(f))^{\frac{1}{4}}, (\delta^{\frac{3}{2}}(f) + r_a \delta(f))^{\frac{1}{2}}\} \leq x_a \delta(f)^{\frac{1}{4}}.
$$

Supposing

$$
\delta(f) \ge l_a
$$

 $l_a \gg 1$ (observe this case is easy since if $\delta(f) \geq l_a$, subject to the second moment assumption, $W_1(f d\gamma, d\gamma) \leq t_a$,

$$
\max\{(\delta^{\frac{3}{2}}(f) + r_a \delta(f))^{\frac{1}{4}}, (\delta^{\frac{3}{2}}(f) + r_a \delta(f))^{\frac{1}{2}}\} \le j_a \delta^{\frac{3}{4}}(f).
$$

Therefore

$$
\int |\nabla f| d\gamma = \int |\nabla \ln f| f d\gamma
$$

\n
$$
\leq \int |T(x) - x| f d\gamma + \sqrt{2\delta(f)}
$$

\n
$$
= W_1 (f d\gamma, d\gamma) + \sqrt{2\delta(f)}
$$

\n
$$
\leq \max\{x_a, j_a\} (\delta^{\frac{1}{4}}(f) + \delta^{\frac{3}{4}}(f)) + \sqrt{2\delta(f)}.
$$

Thus set

$$
f = |u|^2 \left(\frac{x}{\sqrt{2\pi}}\right).
$$

Then

$$
\int f e^{-\frac{|x|^2}{2}} (2\pi)^{-\frac{n}{2}} dx = \int |u|^2 e^{-\pi |x|^2} dx = 1,
$$

$$
\int |\nabla f| d\gamma = \sqrt{\frac{2}{\pi}} \int |\nabla u| |u| e^{-\pi |x|^2} dx.
$$

Hence utilizing [\[4\]](#page-9-3)

$$
\int |\nabla u| e^{-\pi |x|^2} dx
$$
\n
$$
= \int |\nabla u| |u| e^{-\pi |x|^2} dx + \int |\nabla u| (1 - |u|) e^{-\pi |x|^2} dx
$$
\n
$$
\leq \sqrt{\frac{\pi}{2}} \int |\nabla f| d\gamma + \int |\nabla u| |1 - u| e^{-\pi |x|^2} dx
$$
\n
$$
\leq \sqrt{\frac{\pi}{2}} \int |\nabla f| d\gamma + (\int |\nabla u|^2 e^{-\pi |x|^2} dx)^{1/2} (\int |1 - u|^2 e^{-\pi |x|^2} dx)^{1/2}
$$
\n
$$
\leq m_a(\delta^{\frac{1}{4}}(f) + \delta^{\frac{3}{4}}(f)) + (\int |\nabla u|^2 e^{-\pi |x|^2} dx)^{1/2} \sqrt{\kappa} \Big[\frac{1}{\pi} \int |\nabla u|^2 e^{-\pi |x|^2} dx - \int |u|^2 \ln |u|^2 e^{-\pi |x|^2} dx \Big]^{1/2}.
$$
\nNote

Note

$$
\frac{1}{2} \int \frac{|\nabla f|^2}{f} d\gamma - \int f \ln f d\gamma = \frac{1}{\pi} \int |\nabla u|^2 e^{-\pi |x|^2} dx - \int |u|^2 \ln |u|^2 e^{-\pi |x|^2} dx
$$

$$
\int |x|^2 |u(x)|^2 e^{-\pi |x|^2} dx = \frac{1}{2\pi} \int |x|^2 f d\gamma \le \frac{1}{2\pi} \alpha,
$$

which combined with [\[8,](#page-9-7) Theorem 1.17] implies

$$
\frac{1}{\pi} \int |\nabla u(x)|^2 e^{-\pi |x|^2} dx
$$
\n
$$
\leq |\pi \int |x|^2 e^{-\pi |x|^2} dx - \pi \int |x|^2 |u(x)|^2 e^{-\pi |x|^2} dx + \sqrt{2n} \Big(\frac{1}{\pi} \int |\nabla u|^2 e^{-\pi |x|^2} dx - \int |u|^2 \ln |u|^2 e^{-\pi |x|^2} dx \Big)^{\frac{1}{2}}
$$
\n
$$
+ \Big(\frac{1}{\pi} \int |\nabla u|^2 e^{-\pi |x|^2} dx - \int |u|^2 \ln |u|^2 e^{-\pi |x|^2} dx \Big)
$$
\n
$$
\leq q_\alpha (1 + \delta_*^{\frac{1}{2}}(u) + \delta_*(u)).
$$

Therefore utilizing the above estimates,

$$
\int |\nabla u| e^{-\pi |x|^2} dx
$$

\n
$$
\leq \overline{m}_a(\delta_*^{\frac{1}{4}}(u) + \delta_*^{\frac{3}{4}}(u)) + (\pi q_\alpha (1 + \delta_*^{\frac{1}{2}}(u) + \delta_*(u)))^{1/2} \sqrt{\overline{\kappa}} \Big[\delta_*(u)\Big]^{1/2}
$$

\n
$$
\leq a_\alpha(\delta_*^{\frac{1}{4}}(u) + \delta_*(u)).
$$

Proof of Corollary 1.2. Observe that

$$
||f_k - 1||_{W^{1,1}(\mathbb{R}, d\gamma)} \le a_\alpha \left(\delta^{\frac{1}{4}}(f_k) + \delta^{\frac{3}{4}}(f_k)\right),
$$

$$
\partial_{x_k} f = f'_k(x_k) \Pi_{i \neq k} f_i(x_i),
$$

$$
\delta(f) = \sum_k \delta(f_k),
$$

yield

$$
\int |\nabla f| d\gamma = \int \sqrt{\sum_{k} (\partial_{x_k} f)^2} d\gamma
$$
\n
$$
\leq \int \sum_{k} |\partial_{x_k} f| d\gamma
$$
\n
$$
= \sum_{k} \int_{\mathbb{R}^{n-1}} \left(\int_{\mathbb{R}} |f'_k(x_k)| \frac{e^{-\frac{x_k^2}{2}}}{\sqrt{2\pi}} dx_k \right) \prod_{i \neq k} f_i(x_i) \frac{e^{-\sum_{i \neq k} \frac{x_i^2}{2}}}{(\sqrt{2\pi})^{n-1}} dx_1 dx_2 \dots dx_{k-1} dx_{k+1} \dots dx_n
$$
\n
$$
= \sum_{k} \int_{\mathbb{R}} |f'_k(x_k)| \frac{e^{-\frac{x_k^2}{2}}}{\sqrt{2\pi}} dx_k
$$
\n
$$
\leq \sum_{k} a_\alpha \left(\delta^{\frac{1}{4}}(f_k) + \delta^{\frac{3}{4}}(f_k) \right)
$$
\n
$$
\leq a_\alpha n^{3/4} (\delta^{\frac{1}{4}}(f) + \delta^{\frac{3}{4}}(f)).
$$

The proof of (2) is similar. \square

Proof of Theorem [1.3.](#page-2-0) Observe thanks to [\(2.1\)](#page-3-0),

$$
\delta^{\frac{1}{2}}(f) + \frac{1}{\sqrt{2}} \int |\nabla f| d\gamma \ge \frac{1}{\sqrt{2}} \int |T(x) - x| f d\gamma \ge \frac{1}{\sqrt{2}} W_1(f d\gamma, d\gamma);
$$

lar let

in particular, let

$$
f=|u|^2.
$$

Via [\[1,](#page-9-6) Theorem 1]

$$
\int |\nabla f| d\gamma = 2 \int |u| |\nabla u| d\gamma
$$

\n
$$
\leq 2 \sqrt{\int |\nabla u|^2 d\gamma}
$$

\n
$$
\leq 2(2/\eta(\alpha, \epsilon))^{\frac{1}{2}} \delta^{\frac{1}{2}}(f).
$$

This yields the conclusion via

$$
a_{\alpha,\epsilon} = 2(2/\eta(\alpha,\epsilon))^{1/2} + \sqrt{2}.
$$

 \Box

Set

$$
f_a(x) := (2a+1)^{\frac{n}{2}} e^{-a|x|^2}.
$$

Several calculations yield

 $\overline{\mathsf{I}}$

$$
\delta(f_a) = na - \frac{n}{2}\ln(2a+1).
$$

Assuming $\Gamma_1 = f d\gamma$ is the first marginal of Γ , $\Gamma_2 = d\gamma$ is the second marginal of Γ ,

$$
W_1(fd\gamma, d\gamma) = \inf \int \int |x - y| d\Gamma \ge |\int |x| f d\gamma - \int |y| d\gamma|,
$$

$$
|\int |x| f_a d\gamma - \int |y| d\gamma| = |\int |x| ((2a + 1)^{\frac{n}{2}} e^{-a|x|^2} - 1) d\gamma|
$$

$$
= |\int |x| (a(n - |x|^2) + o(a)) d\gamma|,
$$

$$
\frac{\int |x|f_a d\gamma - \int |y|d\gamma|}{a} = |\int |x|(n - |x|^2 + \frac{o(a)}{a})d\gamma|,
$$

$$
\frac{\delta^{\frac{1}{2}}(f_a)}{W_1(f_a d\gamma, d\gamma)} \le \frac{\delta^{\frac{1}{2}}(f_a)}{|\int |x|f_a d\gamma - \int |y|d\gamma|}
$$

$$
= \frac{\delta^{\frac{1}{2}}(f_a)}{|\int |x|(a(n - |x|^2) + o(a))d\gamma|}
$$

$$
= \frac{(\frac{\delta(f_a)}{a^2})^{\frac{1}{2}}}{|\int |x|(n - |x|^2 + \frac{o(a)}{a})d\gamma|}
$$

$$
= \frac{(\frac{na - \frac{n}{2}\ln(2a+1)}{a^2})^{\frac{1}{2}}}{|\int |x|(n - |x|^2 + \frac{o(a)}{a})d\gamma|},
$$

$$
= \frac{na - \frac{n}{2}\ln(2a + 1)}{a^2}\frac{1}{a^2} \rightarrow \sqrt{n},
$$

with $a \to 0^+$, &

$$
\left| \int |x|(n-|x|^2 + \frac{o(a)}{a})d\gamma \right| \to |nm_1(d\gamma) - m_3(d\gamma)|.
$$

Observe now that if $\epsilon < .25$ &

$$
\alpha > \frac{1}{(2\pi)^n} \left(\frac{\pi}{-2\epsilon + .5}\right)^n,
$$

one has assuming $a > 0$ is small,

$$
W_1(f_a d\gamma, d\gamma) \le a_{\alpha,\epsilon} \delta^{\frac{1}{2}}(f_a).
$$

The argument is: observe that

$$
e^{\epsilon |x-y|^2} \le e^{2\epsilon (|x|^2 + |y|^2)},
$$

which yields

$$
\int \int f_a(x)f_a(y)e^{\epsilon|x-y|^2}d\gamma(x)d\gamma(y) \leq \left(\frac{2a+1}{2\pi}\right)^n\left(\frac{\pi}{a-2\epsilon+1.5}\right)^n;
$$

supposing

$$
\alpha > \frac{1}{(2\pi)^n} \left(\frac{\pi}{-2\epsilon + .5}\right)^n,
$$

then when $a > 0$ is small,

$$
\left(\frac{2a+1}{2\pi}\right)^n \left(\frac{\pi}{a-2\epsilon+.5}\right)^n < \alpha
$$

and that implies

$$
W_1(f_a d\gamma, d\gamma) \le a_{\alpha,\epsilon} \delta^{\frac{1}{2}}(f_a).
$$

When

$$
W_1(f d\gamma, d\gamma) \le a_{\alpha,\epsilon} \delta^{\frac{1}{2}}(f),
$$

and $a_{\alpha,\epsilon}$ is the sharp constant,

$$
\frac{1}{a_{\alpha,\epsilon}} \le \frac{\delta^{\frac{1}{2}}(f_a)}{W_1(f_a d\gamma, d\gamma)}
$$
\n
$$
\le \frac{\left(\frac{na - \frac{n}{2}\ln(2a+1)}{a^2}\right)^{\frac{1}{2}}}{\left|\int |x|(n - |x|^2 + \frac{o(a)}{a})d\gamma\right|} \to \frac{\sqrt{n}}{|nm_1(d\gamma) - m_3(d\gamma)|}
$$

.

When $n = 1$,

$$
\frac{\sqrt{n}}{|nm_1(\gamma) - m_3(\gamma)|} = \frac{\sqrt{2\pi}}{2}.
$$

In order to finish the argument, assume via contradiction that there is the estimate

$$
W_1(fd\gamma,d\gamma) \le a_{\alpha,\epsilon} \mu(\delta^{\frac{1}{2}}(f)),
$$

1

 $\mu(a) = o(a)$. Therefore

$$
W_1(f_a d\gamma, d\gamma) \le a_{\alpha,\epsilon} \mu(\delta^{\frac{1}{2}}(f_a))
$$

= $a_{\alpha,\epsilon} \frac{\mu(\delta^{\frac{1}{2}}(f_a))}{\delta^{\frac{1}{2}}(f_a)} \delta^{\frac{1}{2}}(f_a),$

and

$$
\limsup_{a \to 0^+} \frac{\delta^{\frac{1}{2}}(f_a)}{W_1(f_a d\gamma, d\gamma)} \le \frac{\sqrt{2\pi}}{2},
$$

easily imply

$$
\frac{1}{a_{\alpha,\epsilon}} \leq \limsup_{a \to 0^+} \frac{\mu(\delta^{\frac{1}{2}}(f_a))}{\delta^{\frac{1}{2}}(f_a)} \frac{\delta^{\frac{1}{2}}(f_a)}{W_1(f_a d\gamma, d\gamma)}
$$

= 0.

In particular, this yields the contradiction.

REFERENCES

- [1] Giovanni Brigati, Jean Dolbeault, and Nikita Simonov, Stability for the logarithmic Sobolev inequality, arXiv:2303.12926v2 (2024).
- [2] Eric A. Carlen, Superadditivity of Fisher's information and logarithmic Sobolev inequalities, J. Funct. Anal. 101 (1991), no. 1, 194–211. MR 1132315
- [3] Dario Cordero-Erausquin, Some applications of mass transport to Gaussian-type inequalities, Arch. Ration. Mech. Anal. 161 (2002), no. 3, 257–269. MR 1894593
- [4] Jean Dolbeault, Maria J. Esteban, Alessio Figalli, Rupert L. Frank, and Michael Loss, Sharp stability for sobolev and log-sobolev inequalities, with optimal dimensional dependence, arXiv:2209.08651v4 (2023).
- [5] Max Fathi, Emanuel Indrei, and Michel Ledoux, Quantitative logarithmic Sobolev inequalities and stability estimates, Discrete Contin. Dyn. Syst. 36 (2016), no. 12, 6835–6853. MR 3567822
- [6] L. Gross, Logarithmic Sobolev Inequalities, Amer. J. Math 97 (1975), 1061–1083.
- [7] Emanuel Indrei, *Sharp Stability for LSI*, Mathematics **11 (12), 2670** (2023).
- [8] Emanuel Indrei and Daesung Kim, *Deficit estimates for the logarithmic sobolev inequality*, Differential and Integral Equations 34 (2021), no. 7-8, 437–466.
- [9] Daesung Kim, Instability results for the logarithmic Sobolev inequality and its application to related inequalities, Discrete Contin. Dyn. Syst. 42 (2022), no. 9, 4297–4320. MR 4455233

Department of Mathematics and Statistics, Sam Houston State University, Huntsville, TX, USA.