Temperature and pressure jump coefficients at a liquid-vapor interface

Denize Kalempa^a, Irina Graur^b

^aDepartamento de Ciências Básicas e Ambientais, Escola de Engenharia de Lorena, Universidade de São Paulo, 12602-810, Lorena, Brazil ^bAix-Marseille Université, CNRS, IUSTI UMR Marseille 7343, 13453, France

Abstract

The temperature and pressure jump coefficients at a liquid-vapor interface are calculated from the solution of the Shakhov kinetic model for the linearized Boltzmann equation. Complete and partial evaporation/condensation at the vapor-liquid interface are assumed as the boundary condition. The discrete velocity method is used to solve the problem numerically. The jump coefficients are tabulated as functions of the evaporation/condensation coefficient. The profiles of the vapor temperature and pressure deviations from that values at the interface corresponding to the liquid temperature and saturation pressure are plotted and the solutions obtained from kinetic theory and continuum approach are shown to underline the effect of the jumps at the interface. The obtained results have been compared to those given by other authors, who applied the linearized Boltzmann equation as well as the model proposed by Bhatnagar, Gross and Krook to it, and it was found that the pressure and temperature jump coefficients are relatively insensitive to the collision laws.

1. Introduction

In kinetic theory of gases, the problem of evaporation and condensation of a vapor on its condensed phase has been studied by many authors over the years, e.g. Refs.[1, 2, 3, 4, 5, 6].

As is known, from the kinetic viewpoint, during evaporation and condensation processes the vapor in the thin layer near the interface, called as Knudsen layer and whose thickness has the magnitude of a few molecular mean free paths, is in a non-equilibrium state. In the Knudsen layer, the continuum models based on the assumption of continuity of thermodynamic variables at the interface fail to predict the vapor behavior. Thus, to simulate the vapor behavior properly the methods of kinetic theory of gases based on either the solution of the Boltzmann equation and its related models, such as those proposed by Bhatnagar, Gross and Krook [7] and Shakhov [8], referred to as BGK-model and S-model, or the Direct Simulation Monte Carlo method [9] must be applied. Since the methods of kinetic theory of gases require a great computational effort, for practical purposes it is still worth taking into account the gas rarefaction effects by modest computational effort and it can be done by using the special boundary conditions in the frame of the continuum approach, i.e. with the Navier-Stokes-Fourier equations. Thus, similar to the temperature jump on a gas-solid interface, e.g. Refs. [10, 11, 12], in case of a slightest gas rarefaction the continuum equations with proper boundary conditions which take into account the jumps of thermodynamic variables at the interface can be used to describe the gas behavior outside the Knudsen layer.

In the pioneering works by Pao [1, 2] and Cipolla *et al.* [4] the authors, by analogy with the temperature jump on a solid surface, assumed that vapor parameters could be different from that of the condensed phase and proposed the so called pressure and temperature jump boundary conditions. From the macroscopic point of view, the implementation of the jump conditions on the continuum type Navier-Stokes-Fourier equations allows to simulate correctly the vapor behavior outside the Knudsen layer. In the aforementioned papers, the collision term of the linearized Boltzmann equation was simulated via the BGK-model. However, while Pao used the integral moment method to solve the linearized kinetic equation, which consists on obtaining a set of integral equations for the moments of the distribution function, Cipolla et al. used the variational method which implies the use of trial functions for the moments of the distribution function. The temperature and pressure jump coefficients arising from the conductive heat flux and mass flux were calculated by the authors and the comparison between their results obtained from different numerical techniques show a maximum difference of about 1%for the temperature jump due to the conductive heat flux.

The pressure and temperature jump coefficients of a gas on its plane condensed phase were also obtained by Sone and Onishi [5] and Siewert and Thomas [3] from the BGK model for the linearized Boltzmann equation. However, the authors considered only the mass flux as the driving force which causes the jumps at the interface. Later, the same problem was solved by Sone *et al.* [13] with basis on the linearized Boltzmann equation. The comparison between the results obtained from the BGK kinetic model [3, 5] and Boltzmann equation [13] shows a difference of about 2% for the temperature jump at the interface, while the difference for the density jump is negligible.

The aim of the preset paper is to calculate the pressure and temperature jumps at the interface, caused by both heat and mass flux in the Knudsen layer, from the S-model [8] for the linearized Boltzmann equation. The S-model is considered the most suitable to deal with problems concerning both heat and mass transfer because it maintains the original properties of the Boltzmann equation and provides the correct Prandtl number. Complete and partial evaporation/condensation at the interface is assumed as the boundary condition. The discrete velocity method, which has been widely used to solve rarefied gas problems, is employed to solve the problem numerically. Explicit expressions of the pressure and temperature jumps coefficients related to the evaporation and heat fluxes through the interface based on the Onsager reciprocity relations are given and used as an additional criterium of the convergence of the numerical scheme. The implementation of the jump conditions in the continuum models is discussed.

2. Statement of the problem

Let us consider the heat and mass transfer problem in the two-phase system. A gas in the semi-infinite space $x' \geq 0$ is bounded by its plane condensed phase at x' = 0. The liquid layer has a uniform temperature T_0 and saturation pressure p_0 . The number density of saturated vapor at T_0 is denoted by n_0 . Far from the plane condensed phase, i.e. outside the Knudsen layer, the gas is in equilibrium with number density n_{∞} , temperature T_{∞} , pressure $p_{\infty} = n_{\infty}kT_{\infty}$, and flow velocity $(U_{\infty}, 0, 0)$. The slow phase change is due to small density and temperature gradients in the vapor, in the x'direction, which are given as

$$X_n = \left| \frac{\ell_0}{n_0} \frac{dn}{dx'} \right| \ll 1, \quad X_T = \left| \frac{\ell_0}{T_0} \frac{dT}{dx'} \right| \ll 1, \tag{1}$$

so the density, temperature and pressure vary on x'-coordinate. The length ℓ_0 is equivalent to the molecular mean free path of vapor defined as

$$\ell_0 = \frac{\mu_0 v_0}{p_0}, \quad v_0 = \sqrt{\frac{2kT_0}{m}},\tag{2}$$

where μ_0 is the vapor viscosity and v_0 is the most probable speed of vapor molecules at the temperature T_0 .

Let us assume that far from the plane condensed phase the temperature and density vary linearly on the x'-coordinate, while the pressure is constant. Thus, the asymptotic behavior of the gas temperature and density read

$$T(x') \to T_w \left(1 + \frac{x'}{\ell_0} X_T \right) \quad \text{at} \quad \frac{x'}{\ell_0} \to \infty,$$
(3)

$$n(x') \to n_w \left(1 - \frac{x'}{\ell_0} X_T\right) \quad \text{at} \quad \frac{x'}{\ell_0} \to \infty.$$
 (4)

Note that, since the pressure is constant far from the surface, $X_n = -X_T$ when $x' \to \infty$. T_w and n_w are the temperature and density of the vapor at the plane condensed phase and their values are different from the temperature T_0 and density n_0 at the plane condensed phase. In fact, there is a jump of both quantities at the interface so that

$$T_w = T_0(1 + \epsilon_T), \quad n_w = n_0(1 + \epsilon_n), \tag{5}$$

where ϵ_T and ϵ_n denote the macroscopic temperature and density jumps at the interface.

When no jump condition is assumed at the interface, the asymptotic behavior of temperature and density are denoted by

$$T_{NJ}(x') \to T_0\left(1 + \frac{x'}{\ell_0}X_T\right) \quad \text{at} \quad \frac{x'}{\ell_0} \to \infty,$$
(6)

$$n_{NJ}(x') \to n_0 \left(1 - \frac{x'}{\ell_0} X_T\right) \quad \text{at} \quad \frac{x'}{\ell_0} \to \infty,$$
(7)

where the subscript NJ means no jump.

Thus, from (3), (4), (6) and (7), the macroscopic jumps can be calculated as

$$\epsilon_T = \lim_{x' \to \infty} \left[\frac{T(x') - T_{NJ}(x')}{T_0} \right], \quad \epsilon_n = \lim_{x' \to \infty} \left[\frac{n(x') - n_{NJ}(x')}{n_0} \right].$$
(8)

It is worth noting that although the jumps are used in the boundary condition, i.e. at x' = 0, they are calculated outside the Knudsen layer. This is valid because the thickness of the Knudsen layer has the order of few molecular mean free paths. Moreover, it is very thin compared to the characteristic size of the region occupied by the vapor.

In order to calculate the jumps given in (8), the temperature T(x') and the density n(x') must be calculated from methods of kinetic theory of gases.

The vapor pressure at the interface also experiences a jump so that

$$p_w = p_0(1 + \epsilon_p),\tag{9}$$

where ϵ_p is the pressure jump. Therefore, from the ideal gas law, $p_w = n_w k T_w$, the pressure, temperature and number density jumps are related as

$$\epsilon_p = \epsilon_n + \epsilon_T. \tag{10}$$

Here, we are going to present the values for ϵ_p and ϵ_T because, in practice, these jumps can be measured.

3. Kinetic equation

For further derivation we will use the Boltzmann kinetic equation with the Shakhov model [8] for the collisional term. For convenience, let us introduce the dimensionless x-coordinate and molecular velocity as

$$x = \frac{x'}{\ell_0}, \quad \mathbf{c} = \frac{\mathbf{v}}{v_0},\tag{11}$$

where ℓ_0 and v_0 are defined in (2).

Since we are going to calculate the pressure, temperature and density jumps with basis on macroscopic quantities outside the Knudsen layer, two driving forces are introduced as

$$X_T = \frac{1}{T_0} \left(\frac{dT}{dx}\right)_{\infty}, \quad X_u = \frac{U_{\infty}}{v_0}, \tag{12}$$

which are assumed as being very small, i.e.

$$|X_T| \ll 1, \quad |X_u| \ll 1. \tag{13}$$

Note that, outside the Knudsen layer the pressure is constant and, as a consequence, the thermodynamic forces given in (1) are related as $X_T = -X_n$. The assumption of smallness of the driving forces allows us to solve the problem on the basis of a linearized approach. Thus, the distribution function of molecular velocities can be represented as

$$f(x, \mathbf{c}) = f_0^M [1 + h^{(T)}(x, \mathbf{c})X_T + h^{(u)}(x, \mathbf{c})X_u],$$
(14)

where

$$f_0^M = \frac{n_0}{(\sqrt{\pi}v_0)^3} e^{-c^2} \tag{15}$$

is the Maxwellian function with the plane condensed phase characteristics, and $h^{(i)}(x, \mathbf{c})$ (i=T, u) is the perturbation function due to the corresponding thermodynamic force.

Thus, in the absence of external forces, after the substitution of the representation (14) into the Shakhov model kinetic equation [8], two independent kinetic equations are obtained for the problem in question as

$$c_x \frac{\partial h^{(i)}}{\partial x} = \hat{L}h^{(i)}, \quad i = T, u,$$
(16)

where the collisional operator reads

$$\hat{L}h^{(i)} = \nu^{(i)} + \left(c^2 - \frac{3}{2}\right)\tau^{(i)} + 2c_x u_x^{(i)} + \frac{4}{15}c_x\left(c^2 - \frac{5}{2}\right)q_x^{(i)} - h^{(i)}.$$
 (17)

The dimensionless quantities $\nu^{(i)}$, $\tau^{(i)}$, $u_x^{(i)}$ and $q_x^{(i)}$ appearing in (17) correspond to the density and temperature deviations from the values n_0 and T_0 at the plane condensed phase, bulk velocity and heat flux in the x-direction, respectively, caused by the corresponding driving force. According to Refs. [14, 15], in kinetic theory of gases the macroscopic quantities which characterize the gas flow are calculated in terms of the distribution function of molecular velocities. In our notation, the representation (14) allows to write the density and temperature deviations as

$$\nu(x) = \frac{n(x) - n_0}{n_0} = \nu^{(T)}(x)X_T + \nu^{(u)}(x)X_u,$$
(18)

$$\tau(x) = \frac{T(x) - T_0}{T_0} = \tau^{(T)}(x)X_T + \tau^{(u)}(x)X_u,$$
(19)

where

$$\nu^{(i)}(x) = \frac{1}{\pi^{3/2}} \int h^{(i)}(x, \mathbf{c}) e^{-c^2} \,\mathrm{d}\mathbf{c},\tag{20}$$

$$\tau^{(i)}(x) = \frac{2}{3\pi^{3/2}} \int h^{(i)}(x, \mathbf{c}) \left(c^2 - \frac{3}{2}\right) e^{-c^2} d\mathbf{c}.$$
 (21)

Moreover, the bulk velocity and heat flux read

$$u(x) = \frac{U_x(x)}{v_0} = u_x^{(T)}(x)X_T + u_x^{(u)}(x)X_u,$$
(22)

$$q(x) = \frac{Q_x(x)}{v_0 p_0} = q_x^{(T)}(x) X_T + q_x^{(u)}(x) X_u,$$
(23)

where

$$u_x^{(i)}(x) = \frac{1}{\pi^{3/2}} \int h^{(i)}(x, \mathbf{c}) c_x \mathrm{e}^{-c^2} \,\mathrm{d}\mathbf{c},\tag{24}$$

$$q_x^{(i)}(x) = \frac{1}{\pi^{3/2}} \int h^{(i)}(x, \mathbf{c}) \left(c^2 - \frac{5}{2}\right) c_x \mathrm{e}^{-c^2} \,\mathrm{d}\mathbf{c}.$$
 (25)

Therefore, from (8) one can easily see that the temperature and density jumps can be calculated in terms of the density and temperature deviations defined in (20) and (21) as

$$\epsilon_T = \lim_{x \to \infty} \left[(\tau^{(T)} - x) X_T + \tau^{(u)} X_u \right], \qquad (26)$$

$$\epsilon_n = \lim_{x \to \infty} \left[(\nu^{(T)} + x) X_T + \nu^{(u)} X_u \right].$$
(27)

4. Boundary condition

It is assumed that a fraction σ of incident molecules condensates at the plane condensed phase and then evaporates from it with the Maxwellian distribution function f_0^M given in (15). Meanwhile, the fraction $1-\sigma$ is reflected from the condensed phase under the assumption of diffuse scattering. Therefore, the disbribution function of emmited molecules from the condensed phase is written as

$$f(0,\mathbf{c}) = \sigma f_0^M + (1-\sigma)f_r \quad \text{at} \quad c_x > 0,$$
(28)

where

$$f_r = \frac{n_r}{n_0} f_0^M \tag{29}$$

is the distribution function of the molecules diffusely reflected from the condensed phase. The constant n_r is the number density calculated from the impermeability condition for the molecules diffusely reflected from the plane condensed phase. Since the representation (14) is used to linearize the Shakhov kinetic equation, the boundary condition (28) to solve the equations given in (16) reads

$$h^{(i)}(0, \mathbf{c}) = -(1 - \sigma)\xi^{(i)}$$
 at $c_x > 0,$ (30)

where

$$\xi^{(i)} = 1 - \frac{n_r^{(i)}}{n_0} = \frac{2}{\pi} \int_{c_x < 0} c_x h^{(i)}(0, \mathbf{c}) e^{-c^2} \,\mathrm{d}\mathbf{c}.$$
(31)

It is worth underling that in the case of slow evaporation and condensation these processes are symmetric, i.e. the results can be obtained just by changing the sign of the evaporation (condensation) velocity.

5. Condition far from the condensed phase $(x \to \infty)$

Far from the plane condensed phase, the solution of the problem is obtained from the Chapman-Enskog approach to the kinetic equation. Thus, the distribution function of molecular velocities can be written as

$$f = f_{\infty}^{M} (1 + h_{CE}^{(T)} X_{T} + h_{CE}^{(u)} X_{u})$$
(32)

where the Maxwellian function corresponding to the equilibrium reads

$$f_{\infty}^{M} = n_{\infty} \left(\frac{m}{2\pi k T_{\infty}}\right)^{3/2} \exp\left[-\frac{m}{2k T_{\infty}} (\mathbf{v} - \mathbf{U}_{\infty})^{2}\right],\tag{33}$$

with T_{∞} and n_{∞} given in (3) and (4), while $\mathbf{U}_{\infty} = (U_{\infty}, 0, 0)$. This function is related to the distribution function f_0^M , defined in (15), as follows

$$f_{\infty}^{M} = f_{0}^{M} \left[1 + \epsilon_{n} + \left(c^{2} - \frac{3}{2} \right) \epsilon_{T} + x \left(c^{2} - \frac{5}{2} \right) X_{T} + 2c_{x} X_{u} \right].$$
(34)

The use of the representation (32) into the Shakhov kinetic equation allows to find the Chapman-Enskog solution as

$$h_{CE}^{(T)} = -\frac{3}{2}c_x \left(c^2 - \frac{5}{2}\right), \quad h_{CE}^{(u)} = 0.$$
(35)

Note that, from (26) and (27), the temperature and density jumps can be written as

$$\epsilon_T = \epsilon_T^{(T)} X_T + \epsilon_T^{(u)} X_u, \tag{36}$$

$$\epsilon_n = \epsilon_n^{(T)} X_T + \epsilon_n^{(u)} X_u, \tag{37}$$

where the jump coefficients are defined as

$$\epsilon_T^{(T)} = \lim_{x \to \infty} (\tau^{(T)} - x), \quad \epsilon_T^{(u)} = \lim_{x \to \infty} \tau^{(u)}, \tag{38}$$

$$\epsilon_n^{(T)} = \lim_{x \to \infty} (\nu^{(T)} + x), \quad \epsilon_n^{(u)} = \lim_{x \to \infty} \nu^{(u)}.$$
(39)

Thus, after substituting the representation (36) and (37) into (34), the following asymptotic behaviors of the perturbation functions are obtained from (32) as

$$h^{(T)}(x,\mathbf{c}) = \epsilon_n^{(T)} + \left(c^2 - \frac{3}{2}\right)\epsilon_T^{(T)} + x\left(c^2 - \frac{5}{2}\right) + h_{CE}^{(T)} \quad \text{at} \quad x \to \infty, \ (40)$$

$$h^{(u)}(x,\mathbf{c}) = \epsilon_n^{(u)} + \left(c^2 - \frac{3}{2}\right)\epsilon_T^{(u)} + 2c_x + h_{CE}^{(u)} \quad \text{at} \quad x \to \infty.$$
(41)

where $h_{CE}^{(T)}$ and $h_{CE}^{(u)}$ are given in (35). Note that, from the relation (10),

$$\epsilon_n^{(i)} = \epsilon_p^{(i)} - \epsilon_T^{(i)}, \quad i = u, T.$$
(42)

6. Numerical solution

The linearized kinetic equations given in (16) subject to the corresponding conditions given in (30), (40) and (41), are solved numerically by using the discrete velocity method whose details can be found in the literature, see e.g. Ref. [16]. Firstly, to reduce the number of variables in the molecular velocity space and, consequently, to reduce the computational effort to solve the problem, new functions are introduced as

$$\phi^{(i)}(x, c_x) = \frac{1}{\pi} \int \int h^{(i)}(x, \mathbf{c}) e^{-c_y^2 - c_z^2} \, \mathrm{d}c_y \mathrm{d}c_z, \tag{43}$$

$$\psi^{(i)}(x,c_x) = \frac{1}{\pi} \int \int h^{(i)}(x,\mathbf{c}) e^{-c_y^2 - c_z^2} (c_y^2 + c_z^2 - 1) \, \mathrm{d}c_y \mathrm{d}c_z.$$
(44)

The substitution of these new functions into (16) leads to the following system of kinetic equations for each thermodynamic force

$$c_x \frac{\partial \phi^{(i)}}{\partial x} = \hat{L} \phi^{(i)},\tag{45}$$

$$c_x \frac{\partial \psi^{(i)}}{\partial x} = \hat{L} \psi^{(i)},\tag{46}$$

where

$$\hat{L}\phi^{(i)} = \nu^{(i)} + \left(c_x^2 - \frac{1}{2}\right)\tau^{(i)} + 2c_x u_x^{(i)} + \frac{4}{15}c_x\left(c_x^2 - \frac{3}{2}\right)q_x^{(i)} - \phi^{(i)}, \quad (47)$$

$$\hat{L}\psi^{(i)} = \tau^{(i)} + \frac{4}{15}c_x q_x^{(i)} - \psi^{(i)}.$$
(48)

The moments of the perturbation function given in (20)-(25) are written in terms of the new functions as

$$\nu^{(i)}(x, c_x) = \frac{1}{\sqrt{\pi}} \int \phi^{(i)} e^{-c_x^2} \, \mathrm{d}c_x, \tag{49}$$

$$\tau^{(i)}(x,c_x) = \frac{2}{3\sqrt{\pi}} \int \left[\left(c_x^2 - \frac{1}{2} \right) \phi^{(i)} + \psi^{(i)} \right] e^{-c_x^2} dc_x,$$
(50)

$$u_x^{(i)}(x, c_x) = \frac{1}{\sqrt{\pi}} \int \phi^{(i)} c_x \mathrm{e}^{-c_x^2} \,\mathrm{d}c_x,\tag{51}$$

$$q_x^{(i)}(x, c_x) = \frac{1}{\sqrt{\pi}} \int \left[\left(c_x^2 - \frac{3}{2} \right) \phi^{(i)} + \psi^{(i)} \right] c_x e^{-c_x^2} dc_x.$$
(52)

Moreover, from (30), the boundary conditions at x=0 are written in terms of the new functions as

$$\phi^{(i)}(0, c_x) = -(1 - \sigma)\xi^{(i)}, \quad c_x > 0$$
(53)

$$\psi^{(i)}(0, c_x) = 0, \quad c_x > 0, \tag{54}$$

where

$$\xi^{(i)} = 2 \int_{c_x < 0} \phi^{(i)}(0, c_x) c_x e^{-c_x^2} dc_x.$$
(55)

Note that, the boundary condition depends on $\xi^{(i)}$ only when $\sigma \neq 1$, which corresponds to the partial, or non-complete, condensation-evaporation at the interface.

Far from the boundary, i.e. in the limit $x \to \infty$, the conditions (40) and (41) are written in terms of the functions as

$$\phi^{(T)}(x,c_x) = \epsilon_n^{(T)} + \left(c_x^2 - \frac{1}{2}\right)\epsilon_T^{(T)} + x\left(c_x^2 - \frac{3}{2}\right) - \frac{3}{2}c_x\left(c_x^2 - \frac{3}{2}\right), \quad (56)$$

$$\psi^{(T)}(x, c_x) = \epsilon_T^{(T)} + x - \frac{3}{2}c_x, \tag{57}$$

$$\phi^{(u)}(x, c_x) = \epsilon_n^{(u)} + \left(c_x^2 - \frac{1}{2}\right)\epsilon_T^{(u)} + 2c_x,$$
(58)

$$\psi^{(u)}(x,c_x) = \epsilon_T^{(u)}.$$
(59)

Thus, for each thermodynamic force, the system of kinetic equations given by (45) and (46) with the corresponding boundary condition and asymptotic behavior was solved numerically via the discrete velocity method with an numerical error of 0.1% for the temperature and density jump coefficients at the vapor-liquid interface. The Gaussian-Hermite quadrature was employed to discretize the x-component of the molecular velocity space and calculate the macroscopic characteristics of the gas flow corresponding the the density and temperature deviations, bulk velocity and heat flux. Details concerning the numerical technique to found the nodes and weights can be found in Ref. [17]. A finite difference scheme was used to calculate the derivatives in the kinetic equations. The accuracy was estimated by varying the grid parameters N_x and N_c , corresponding to the number of nodes in the xcoordinate and molecular velocity component c_x , as well and the maximum distance x_{max} from the interface. The value of N_c was fixed at 12, while the value of N_r varied according to x_{max} so that the increment $\Delta x \sim 10^{-3}$.

7. Thermodynamic analysis

As it is known from the thermodynamic of irreversible processes, e.g. Ref.[18], in a weakly disturbed system, the thermodynamic fluxes are linearly related to thermodynamic forces. In the present problem, the linear relations read

$$J'_M = \Lambda'_{11}X_p + \Lambda'_{12}X_T, \tag{60}$$

$$J_T' = \Lambda_{21}' X_p + \Lambda_{22}' X_T, \tag{61}$$

where the thermodynamic fluxes conjugated to the pressure X_p and temperature X_T gradients are introduced as

$$J'_{M} = n_{0}U_{x} = n_{0}v_{0}u, \quad J'_{T} = \frac{Q_{x}}{kT_{0}} = n_{0}v_{0}q.$$
(62)

The dimensionless quantities u and q are given in (22) and (23), and outside of the Knudsen layer both quantities are constant. Λ'_{ij} (i, j=1, 2) are referred to as kinetic coefficients which satisfy the Onsager reciprocity relation $\Lambda'_{12} = \Lambda'_{21}$.

Moreover, inverted linear relations can be obtained from (60) and (61), so that the thermodynamic forces can be written in terms of the thermodynamic fluxes as

$$X_p = a'_{11}J'_M + a'_{12}J'_T, (63)$$

$$X_T = a'_{21}J'_M + a'_{22}J'_T, (64)$$

where the new kinetic coefficients read

$$a'_{11} = \frac{\Lambda'_{22}}{\mathcal{D}}, \quad a'_{12} = a'_{21} = -\frac{\Lambda'_{12}}{\mathcal{D}}, \quad a'_{22} = \frac{\Lambda'_{11}}{\mathcal{D}},$$
 (65)

and

$$\mathcal{D} = \Lambda_{11}' \Lambda_{22}' - \Lambda_{12}' \Lambda_{21}'. \tag{66}$$

In our approach, the jump coefficients are calculated in terms of macroscopic quantities outside the Knudsen layer. However, in the macroscopic point of view, we are extrapolating the continuum profiles of thermodynamic variables through the Knudsen layer to the interface. Then, the pressure and temperature jump coefficients can also be written as

$$\epsilon_p = \frac{p_w - p_0}{p_0}, \quad \epsilon_T = \frac{T_w - T_0}{T_0}.$$
 (67)

Thus, after some algebraic manipulation, the representations (36) and (37) allow to obtain relations similar to those given in (63) and (64) as

$$\frac{p_w - p_0}{p_0} = \epsilon_p^{(u)} \frac{J'_M}{n_0 v_0} - \frac{8}{15} \epsilon_p^{(T)} \frac{J'_T}{n_0 v_0},\tag{68}$$

$$\frac{T_w - T_0}{T_0} = \epsilon_T^{(u)} \frac{J'_M}{n_0 v_0} - \frac{8}{15} \epsilon_T^{(T)} \frac{J'_T}{n_0 v_0},\tag{69}$$

where the Fourier law was used to write the conductive heat flux as $q=-15X_T/8$. These relations allow us to verify the fullfillment of the reciprocity relation

$$\epsilon_T^{(u)} = -\frac{8}{15} \epsilon_p^{(T)}.$$
(70)

The fullfillment of the reciprocity relation (70) is verified numerically as an additional criterium of the convergence of the numerical scheme.

8. Results

Table 1 presents the pressure and temperature jump coefficients obtained in the present work under the assumption of complete evaporation and condensation at the interface. The results on the jumps provided by Pao [1] and Cipolla *et al.* [4] due to both heat and mass transfer are given in Table 1 for comparison. In both papers the BGK kinetic model was used to solve the problem, but the numerical techniques are different. Pao used the integral moment method, which consists on obtaining a set of integral equations for the moments of the distribution function, while Cipolla *et al.* used the variational method which implies the use of trial functions for the moments of the distribution function. The results for the jumps due to only the mass flux obtained by Sone and Onishi [5] from the BGK kinetic model and those obtained by Sone *et al.* [13] from the Boltzmann equation with the original collision term are also given in Table 1. It is worth noting that, in Ref. [13] the rigid-spheres intermolecular model was used to solve the Boltzmann equation. Loyalka [6] analyzed the influence of intermolecular force laws on the values of the jump coefficients and he concluded that these coefficients are relatively insensitive to them.

According to the tabulated results, there is a good agreement between the present results and those given in Ref. [4]. The maximum difference between the results is approximately equal to 1.2% for the coefficient $\epsilon_T^{(T)}$. However, it is interesting to note that the coefficient $\epsilon_T^{(T)}$ which arises due to the conductive heat flux corresponds to the temperature jump coefficient calculated in case of diffuse scattering of gas molecules from a solid surface. According to the literature review on slip and temperature jumps [19], for practical applications the value of 1.95 is recommended for this jump coefficient. Concerning the comparison with the results given in Refs. [1, 5], the maximum difference is negligible. Thus, Table 1 shows that even if the collision terms of the BGK and S models are different both kinetic models provide the same results for the jump coefficients. The small difference in the comparison with the results given by Cipolla *et al.* is due to the use of trial functions in the variational method which introduces numerical error.

The comparison with the results obtained from the Boltzmann equation [13] shows a difference of about 0.5% for the pressure jump and 2% for the temperature jump due to the mass flux in the Knudsen layer. Thus, since to solve the Boltzmann equation is still a difficult task which demands a great computational effort, the use of kinetic models plays an important role in the solution of practical problems.

It is worth noting that the Shakhov model is the most suitable for solving problems concerning both heat and mass transfer. The collision term depends on the collision frequency f. In the S-model the collision frequency is given by the ratio between the pressure and viscosity of the gas, i.e. $f = p/\mu$. When the BGK model is used with the same collision frequency, the results for $\epsilon_p^{(u)}$ and $\epsilon_T^{(u)}$ are equal to those obtained from the S-model. Nonetheless, the results for $\epsilon_p^{(T)}$ and $\epsilon_T^{(T)}$ must be multiplied by 3/2 due to the fact that the BGK model provides the correct Prandt number when the collision frequency is chosen as $f = 2p/(3\mu)$.

From the provided analysis we may conclude that the values of jump coefficients are less sensitive not only to the intermolecular force laws as it was found in Ref. [6], but also to the collisional laws.

The results for partial or non-complete evaporation-condensation at the interface are provided in Table 2. According to these results, only the coefficients $\epsilon_p^{(T)}$ and $\epsilon_p^{(u)}$ are impacted by the non-complete evaporation-condensation condition. The smaller the evaporation/condensation coefficient σ the larger the magnitude of these jump coefficients. The maximum deviation of $\epsilon_p^{(T)}$ from the corresponding value for complete evaporation/condensation is less than 0.5% so that the influence of σ in such jump coefficient is negligible. Meanwhile, the deviation of $\epsilon_p^{(u)}$ from that value for complete evaporation/condensation at the interface is larger than 100% when $\sigma \leq 0.6$. Therefore, one can say that only the jump coefficient $\epsilon_p^{(u)}$ is indeed impacted by the evaporation/condensation coefficient σ . A similar conclusion is given by Cipolla *et al.* [4] and their results for $\epsilon_p^{(u)}$ are shown in Table 2 for comparison.

The significant dependence of $\epsilon_p^{(u)}$ on the evaporation/condensation co-

efficient σ is due to the dependence of the density deviation $\nu^{(u)}$ on such coefficient. Figure 1 shows the profile of $\nu^{(u)}$ as function of the *x*-coordinate for $\sigma=0.1, 0.4, 0.8$ and 1.

In the following it is shown how the derived jump conditions modify the temperature and pressure profiles obtained from the continuum equations.

In the case of continuum solution, no temperature jump is assumed on the interface and the vapor temperature at the interface, T_w , is equal to the temperature of the interface, i.e. $T_w = T_0$. In this case, the temperature deviation given in (19) is obtained from (6) as

$$\tau(x) = xX_T,\tag{71}$$

so that

$$\tau^{(T)}(x) = x$$
 and $\tau^{(u)} = 0.$ (72)

Figure 2 shows the temperature deviations given in (72) by green lines.

When the temperature jump condition is used, the temperature deviation obtained in the frame of continuum approach with the vapor temperature at the interface, T_w , given in (5) reads

$$\tau(x) = (\epsilon_T^{(T)} + x)X_T + \epsilon_T^{(u)}X_u, \tag{73}$$

where

$$\tau^{(T)}(x) = \epsilon_T^{(T)} + x \quad \text{and} \quad \tau^{(u)}(x) = \epsilon_T^{(u)}.$$
(74)

These temperature deviations are also shown in Figure 2 by red lines.

The temperature deviations calculated numerically from the solution of the Shakhov kinetic equation are also shown in Figure 2 by blue lines. As one can see from Figure 2, outside the Knudsen layer the use of the temperature jump boundary condition leads to the same solution obtained from kinetic theory. Inside the Knudsen layer the profile obtained from kinetic theory is different from that obtained in the frame of continuum approach, but when rarefaction is small such an effect is neglected because the thickness of the Knudsen layer.

Similarly, the pressure deviation can also be plotted for comparison. Let us write the pressure deviation as

$$\eta(x) = \frac{p(x) - p_0}{p_0} = \eta^{(T)} X_T + \eta^{(u)} X_u.$$
(75)

When no jump condition is assumed so that $p_w = p_0$, the profiles given in (6) and (7) allow us to obtain the pressure deviations as

$$\eta^{(T)}(x) = 0, \qquad \eta^{(u)}(x) = 0.$$
(76)

Otherwise, when the pressure jump condition is assumed, the vapor pressure at the interface, p_w , is given in (9). Thus, the pressure deviations obtained from (3) and (4) read

$$\eta^{(T)}(x) = \epsilon_p^{(T)}, \qquad \eta^{(u)}(x) = \epsilon_p^{(u)}.$$
(77)

Figure 3 shows the pressure deviations profiles given in (76) and (77) by green and red lines, respectively. The pressure deviations obtained numerically from the Shakhov kinetic equation are also plotted for comparison.

9. Application to evaporation problem

The derived expressions (68) and (69) allow to relate the evaporation and heat fluxes through an interface to the pressure and temperature jumps. We can write them in another form as

$$\frac{p_{\rm v} - p_{\rm sat}(T_{\rm L})}{p_{\rm sat}(T_{\rm L})} = \epsilon_p^{(u)} \frac{J}{\rho_{\rm sat}(T_{\rm L})\sqrt{2\mathcal{R}T_{\rm L}}} - \frac{8}{15} \epsilon_p^{(T)} \frac{q_{\rm v}}{p_{\rm sat}(T_{\rm L})\sqrt{2\mathcal{R}T_{\rm L}}},\tag{78}$$

$$\frac{T_{\rm v} - T_{\rm L}}{T_{\rm L}} = \epsilon_T^{(u)} \frac{J}{\rho_{\rm sat}(T_{\rm L})\sqrt{2\mathcal{R}T_{\rm L}}} + -\frac{8}{15}\epsilon_T^{(T)} \frac{q_{\rm v}}{p_{\rm sat}(T_{\rm L})\sqrt{2\mathcal{R}T_{\rm L}}}.$$
(79)

Here $T_{\rm L}$ is the liquid-vapor interface temperature from the liquid side, $p_{\rm sat}(T_{\rm L})$ is the saturation pressure at this temperature, $\rho_{\rm sat}(T_{\rm L})$ is the saturation density, $p_{\rm v}$ is the vapor pressure, $T_{\rm v}$ is the liquid-vapor interface temperature from the vapor side, \mathcal{R} is the gas-specific constant. The specific quantities such as the saturation pressure and density for a given liquid temperature can be found in handbooks for a given liquid nature. As during the evaporation or condensation a vapor near a liquid interface is in its non-equilibrium state these conditions must be applied when simulating the evaporation and condensation using the continuum approach.

One example of application of these conditions to the FC-72 liquid evaporation can be found in Ref. [20]. The results obtained by using the jump conditions (78)-(79) are compared to that obtained from the widely used Schrage expression.

10. Conclusion

The temperature and pressure jump coefficients at the liquid vapor interface are calculated from the numerical solution of the Shakhov model kinetic equation. The cases of complete and partial evaporation and condensation processes are considered. The numerical values of the obtained coefficients are very close to those available in the literature, obtained by using another approaches such as the variational method and the integral moment method to solve the BGK kinetic model. The comparison with results obtained from the Boltzmann equation shows a good agreement for the jump coefficients due the mass flux in the Knudsen layer. Therefore, one may conclude that the jump coefficients are less sensitive not only to the intermolecular force laws as it was found previously, but also to the collisional laws. For practical purposes, the Shakhov model is the most suitable for solving problems concerning both heat and mass transfer. Moreover, its solution via the discrete velocity method is more attractive because it provides reliable results with less computational effort compared to that required when other numerical approaches are used.

Acknowledgments

The authors acknowledge the Fundação de Pesquisa do Estado de São Paulo (FAPESP, Brazil), grant 2022/10476-1, for the support of the research.

References

- [1] Y-P Pao. Temperature and density jumps in the kinetic theory of gases and vapors. Phys. Fluids, 14(7):1340–1346, 1971.
- [2] Y-P Pao. Application of kinetic theory to the problem of evaporation and condensation. Phys. Fluids, 14(2):306–312, 1971.
- [3] C E Siewert and J R Thomas. Half-space problems in the kinetic theory of gases. Phys. Fluids, 16:1557–1559, 1973.
- [4] J W Cipolla Jr, H Lang, and S K Loyalka. Kinetic theory of condensation and evaporation. II. J. Chem. Phys., 61:69–77, 1974.
- Y Sone and Y Onishi. Kinetic theory of evaporation and condensation. J. Phys. Soc. Jpn., 35:1773–1776, 1973.
- [6] S. K. Loyalka. Kinetic theory of planar condensation and evaporation. Transport theory and statistical physics, 20(2-3):237–249, 1991.
- [7] P L Bhatnagar, E P Gross, and M A Krook. A model for collision processes in gases. Phys. Rev., 94:511–525, 1954.
- [8] E M Shakhov. Generalization of the Krook kinetic relaxation equation. Fluid Dynamics, 3(5):95–96, 1968.

- [9] G A Bird. <u>Molecular Gas Dynamics and the Direct Simulation of Gas</u> <u>Flows</u>. Oxford University Press, Oxford, 1994.
- [10] S K Loyalka. Temperature jump and thermal creep slip: Rigid sphere gas. Phys. Fluids A, 1:403–408, 1989.
- [11] Y Sone, T Ohwada, and K Aoki. Temperature jump and Knudsen layer in a rarefied gas over a plane wall: Numerical analysis of the linearized Boltzmann equation for hard-sphere molecules. <u>Phys. Fluids A</u>, 1(2):363–370, 1989.
- [12] C E Siewert. The linearized Boltzmann equation: A concise and accurate solution of the temperature-jump problem. <u>J. Quant. Spec. Rad. Tran</u>, 77:417–432, 2003.
- [13] Y Sone, T Ohwada, and K Aoki. Evaporation and condensation on a plane condensed phase: numerical analysis of the linearized Boltzmann equation for hard-sphere molecules. Phys. Fluids A, 1:1398–1405, 1989.
- [14] J H Ferziger and H G Kaper. <u>Mathematical Theory of Transport</u> <u>Processes in Gases</u>. North-Holland Publishing Company, Amsterdam, 1972.
- [15] S Chapman and T G Cowling. <u>The Mathematical Theory of</u> Non-Uniform Gases. University Press, Cambridge, 3 edition, 1970.
- [16] F Sharipov. <u>Rarefied Gas Dynamics. Fundamentals for Research and</u> Practice. Wiley-VCH, Berlin, 2016.
- [17] V I Krylov. <u>Approximate Calculation of Integrals</u>. Dover Publication Inc., Mineola, 2005.
- [18] S R De Groot and P Mazur. <u>Non-Equilibrium Thermodynamics</u>. Dover Publications, Inc., New York, 1984.
- [19] F Sharipov. Data on the velocity slip and temperature jump on a gassolid interface. J. Phys. Chem. Ref. Data, 40(2):023101, 2011.
- [20] I. Graur, A. Rednikov, and L. Ronshin, F.and Tadrist. Numerical modeling and simulation of near-contact-line evaporation kinetics. In <u>Proceedings of 9th European Thermal Sciences Conference (Eurotherm</u> 2024), Slovenia, 2024.

1					
Jump	Present	Cipolla <i>et al.</i>	Pao	Sone/Onishi	Sone <i>et al.</i>
Coefficient	Work	$\operatorname{Ref.}[4]$	$\operatorname{Ref.}[1]$	Ref. $[5]$	Ref. [13]
$\epsilon_T^{(T)}$	1.9540	1.9309	1.9540		
$\epsilon_P^{(T)}$	0.8376	0.8385	0.8376		
$\epsilon_T^{(u)}$	-0.4467	-0.4472	-0.4468	-0.4467	-0.4557
$\epsilon_{P}^{(u)}$	-2.1320	-2.1254	-2.1322	-2.1320	-2.1413

Table 1: Temperature and pressure jump coefficients in the case of complete condensation and evaporation at the interface (σ =1).

Table 2: Temperature and pressure jump coefficients for complete ($\sigma = 1$)and non-complete ($\sigma \neq 1$) condensation and evaporation at the interface.

σ	$\epsilon_T^{(T)}$	$\epsilon_P^{(T)}$	$\epsilon_T^{(u)}$	$\epsilon_P^{(u)}$	$\epsilon_P^{(u)}$ [4]
0.1	1.9540	0.8404	-0.4467	-33.9467	-34.0296
0.2	1.9540	0.8389	-0.4468	-16.3102	-16.3050
0.4	1.9540	0.8381	-0.4468	-7.4489	-7.4428
0.6	1.9540	0.8379	-0.4468	-4.4951	-4.4887
0.8	1.9540	0.8377	-0.4468	-3.0182	-3.0116
1.0	1.9540	0.8376	-0.4467	-2.1320	-2.1254



Figure 1: Density deviation due to X_u .



Figure 2: Temperature deviation due to X_T and X_u .



Figure 3: Pressure deviation due to X_T and X_u .