

INFINITELY BADLY APPROXIMABLE AFFINE FORMS

TAEHYEONG KIM

ABSTRACT. A pair (A, \mathbf{b}) of a real $m \times n$ matrix A and $\mathbf{b} \in \mathbb{R}^m$ is said to be *infinitely badly approximable* if

$$\liminf_{\mathbf{q} \in \mathbb{Z}^n, \|\mathbf{q}\| \rightarrow \infty} \|\mathbf{q}\|^{\frac{n}{m}} \|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} = \infty,$$

where $\|\cdot\|_{\mathbb{Z}}$ denotes the distance from the nearest integer vector. In this article, we introduce a novel concept of singularity for (A, \mathbf{b}) and characterize the infinitely badly approximable property by this singular property. As an application, we compute the Hausdorff dimension of the infinitely badly approximable set. We also discuss dynamical interpretations on the space of grids in \mathbb{R}^{m+n} .

1. INTRODUCTION

1.1. Background and motivation. Let m, n be positive integers, and denote by $M_{m,n}(\mathbb{R})$ the set of $m \times n$ real matrices. We start by introducing the following classical definition. For any $\epsilon > 0$, we say that $A \in M_{m,n}(\mathbb{R})$ is *ϵ -badly approximable* if

$$\liminf_{\mathbf{q} \in \mathbb{Z}^n, \|\mathbf{q}\| \rightarrow \infty} \|\mathbf{q}\|^{\frac{n}{m}} \|A\mathbf{q}\|_{\mathbb{Z}} \geq \epsilon.$$

Here and hereafter, $\|\mathbf{x}\| = \max_{1 \leq i \leq k} |x_i|$ and $\|\mathbf{x}\|_{\mathbb{Z}} = \min_{\mathbf{n} \in \mathbb{Z}^k} \|\mathbf{x} - \mathbf{n}\|$ for $\mathbf{x} \in \mathbb{R}^k$. We say that $A \in M_{m,n}(\mathbb{R})$ is *badly approximable* if it is ϵ -badly approximable for some $\epsilon > 0$. It is well-known that the set of badly approximable matrices is of zero Lebesgue measure [Khi26], but has full Hausdorff dimension [Sch69]. See [BK15, Sim18] for the set of ϵ -badly approximable matrices.

Dirichlet's theorem (1842) says that for any $A \in M_{m,n}(\mathbb{R})$ there exist infinitely many $\mathbf{q} \in \mathbb{Z}^n$ such that $\|A\mathbf{q}\|_{\mathbb{Z}} \leq \|\mathbf{q}\|^{-\frac{n}{m}}$, hence there is no ϵ -badly approximable matrix for any $\epsilon > 1$. However, if we consider inhomogeneous Diophantine approximation, the situation becomes completely different, and investigating this situation is the goal of the present paper.

More precisely, given $\epsilon > 0$, we say that a pair $(A, \mathbf{b}) \in M_{m,n}(\mathbb{R}) \times \mathbb{R}^m$ is *ϵ -badly approximable* if

$$\liminf_{\mathbf{q} \in \mathbb{Z}^n, \|\mathbf{q}\| \rightarrow \infty} \|\mathbf{q}\|^{\frac{n}{m}} \|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} \geq \epsilon.$$

2020 *Mathematics Subject Classification*: Primary 11J20 ; Secondary 37A17, 28A78
The author is supported by the ERC grant HomDyn, ID 833423.

Denote by $\mathbf{Bad}(\epsilon)$ the set of ϵ -badly approximable pairs in $M_{m,n}(\mathbb{R}) \times \mathbb{R}^m$. For $A \in M_{m,n}(\mathbb{R})$ and $\mathbf{b} \in \mathbb{R}^m$, consider the slices of $\mathbf{Bad}(\epsilon)$ as follows:

$$\begin{aligned}\mathbf{Bad}_A(\epsilon) &= \{\mathbf{b} \in \mathbb{R}^m : (A, \mathbf{b}) \in \mathbf{Bad}(\epsilon)\}, \\ \mathbf{Bad}^{\mathbf{b}}(\epsilon) &= \{A \in M_{m,n}(\mathbb{R}) : (A, \mathbf{b}) \in \mathbf{Bad}(\epsilon)\}.\end{aligned}$$

A pair $(A, \mathbf{b}) \in M_{m,n}(\mathbb{R}) \times \mathbb{R}^m$ is said to be *infinitely badly approximable* if the pair (A, \mathbf{b}) is ϵ -badly approximable for all $\epsilon > 0$, or equivalently,

$$\liminf_{\mathbf{q} \in \mathbb{Z}^n, \|\mathbf{q}\| \rightarrow \infty} \|\mathbf{q}\|^{\frac{n}{m}} \|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} = \infty.$$

Denote

$$\mathbf{Bad}(\infty) = \bigcap_{\epsilon > 0} \mathbf{Bad}(\epsilon), \quad \mathbf{Bad}_A(\infty) = \bigcap_{\epsilon > 0} \mathbf{Bad}_A(\epsilon), \quad \mathbf{Bad}^{\mathbf{b}}(\infty) = \bigcap_{\epsilon > 0} \mathbf{Bad}^{\mathbf{b}}(\epsilon).$$

We first remark that if $\mathbf{b} \in \mathbb{Z}^m$, we return to the homogeneous Diophantine approximation, so we will usually only consider the case $\mathbf{b} \in \mathbb{R}^m \setminus \mathbb{Z}^m$. Let us discuss the following motivating observation: For any $A \in M_{m,n}(\mathbb{R})$, Kronecker's theorem (see e.g. [Cas57, Chapter III, Theorem IV]) asserts that the sequence $\{A\mathbf{q}\}_{\mathbf{q} \in \mathbb{Z}^n}$ modulo 1 is dense in $\mathbb{R}^m / \mathbb{Z}^m$ if and only if the subgroup ${}^tAZ^m + \mathbb{Z}^n$ of \mathbb{R}^n has maximal rank $m+n$ over \mathbb{Z} , where tA denotes the transpose of the matrix A . It follows that if tA is rational in the sense that ${}^tA\mathbf{y} \in \mathbb{Z}^n$ for some nonzero $\mathbf{y} \in \mathbb{Z}^m$, then there is an open subset U of $\mathbb{R}^m / \mathbb{Z}^m$ not including the sequence $\{A\mathbf{q}\}_{\mathbf{q} \in \mathbb{Z}^n}$ modulo 1, hence for any $\mathbf{b} \in U$ the pair (A, \mathbf{b}) is infinitely badly approximable.

Through this observation, one may expect

$$(1.1) \quad \begin{aligned} &\text{a certain connection between "singularity" of } {}^tA \\ &\text{and infinitely badly approximability of } (A, \mathbf{b}). \end{aligned}$$

Let us recall the classical definition of singularity. We say that $A \in M_{m,n}(\mathbb{R})$ is *singular* if for any $\epsilon > 0$ for all sufficiently large X the inequalities

$$(1.2) \quad \|A\mathbf{q}\|_{\mathbb{Z}} < \epsilon X^{-\frac{n}{m}} \quad \text{and} \quad 0 < \|\mathbf{q}\| < X$$

have an integer solution $\mathbf{q} \in \mathbb{Z}^n$.

One possible answer to (1.1) comes from [BL05] although it is not written in that paper. We denote by $w(A, \mathbf{b})$ the supremum of the real numbers w for which, for arbitrarily large X , the inequalities

$$\|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} < X^{-w} \quad \text{and} \quad \|\mathbf{q}\| < X$$

have an integer solution $\mathbf{q} \in \mathbb{Z}^n$. We also denote by $\widehat{w}(A)$ the supremum of the real numbers w for which, for all sufficiently large X , the inequalities

$$\|A\mathbf{q}\|_{\mathbb{Z}} < X^{-w} \quad \text{and} \quad 0 < \|\mathbf{q}\| < X$$

have an integer solution $\mathbf{q} \in \mathbb{Z}^n$. It follows from [BL05, Theorem] that for almost all $\mathbf{b} \in \mathbb{R}^m$,

$$w(A, \mathbf{b}) = \frac{1}{\widehat{w}({}^tA)}.$$

If $\widehat{w}({}^tA) > \frac{m}{n}$, the set of $\mathbf{b} \in \mathbb{R}^m$ such that $w(A, \mathbf{b}) < \frac{n}{m}$ is of full Lebesgue measure, hence $\mathbf{Bad}_A(\infty)$ is of full Lebesgue measure. Note that $\widehat{w}({}^tA) > \frac{m}{n}$ implies that tA is singular, or equivalently, A is singular.

The author [Kim24] strengthened this observation under a certain weaker assumption. We remark that even if A is singular, the set $\mathbf{Bad}_A(\infty)$ may have zero Lebesgue measure (see [MRS, Theorem 8.4]).

In terms of Hausdorff dimension, it was shown in [ET11, Theorem 1.5] that if $A \in M_{m,n}(\mathbb{R})$ is singular, then the set $\mathbf{Bad}_A(\infty)$ has full Hausdorff dimension (in fact, it is winning on some fractals).

To the authors' knowledge, these are the only known results in this direction. In this paper, we will give a complete answer to (1.1), compute Hausdorff dimension for infinitely badly approximable affine forms, and discuss dynamical interpretations on the space of grids in \mathbb{R}^{m+n} .

1.2. Main results. One of the main results of this paper is to answer to (1.1) completely with the following new definition of singularity: For fixed $\mathbf{b} \in \mathbb{R}^m$, we say that a matrix ${}^tA \in M_{n,m}(\mathbb{R})$ is *singular for \mathbf{b}* if for any $\epsilon > 0$ for all sufficiently large X the inequalities

$$(1.3) \quad \|{}^tA\mathbf{y}\|_{\mathbb{Z}} < \epsilon |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} X^{-\frac{m}{n}} \quad \text{and} \quad \|\mathbf{y}\| < |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} X$$

have an integer solution $\mathbf{y} \in \mathbb{Z}^m$, where \cdot stands for the usual dot product and $|\cdot|_{\mathbb{Z}}$ denotes the distance from the nearest integer. Denote by $\mathbf{Sing}(\mathbf{b})$ the set of $A \in M_{m,n}(\mathbb{R})$ such that tA is singular for \mathbf{b} .

The following theorem can be seen as the transference principle between infinitely badly approximable affine forms and singular linear forms for translation vectors, which answers to (1.1).

Theorem 1.1. *For any $(A, \mathbf{b}) \in M_{m,n}(\mathbb{R}) \times \mathbb{R}^m$, the pair (A, \mathbf{b}) is infinitely badly approximable if and only if tA is singular for \mathbf{b} . In particular, for any $\mathbf{b} \in \mathbb{R}^m$,*

$$\mathbf{Bad}^{\mathbf{b}}(\infty) = \mathbf{Sing}(\mathbf{b}) \quad \text{and} \quad \mathbf{Bad}(\infty) = \bigcup_{\mathbf{b} \in \mathbb{R}^m} \mathbf{Sing}(\mathbf{b}) \times \{\mathbf{b}\}.$$

Remark 1.2. Since $|\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} \leq 1/2$ for any $\mathbf{b} \in \mathbb{R}^m$ and $\mathbf{y} \in \mathbb{Z}^m$, if tA is singular for \mathbf{b} , then tA is singular, hence A is singular. Therefore, for any $\mathbf{b} \in \mathbb{R}^m$, the set $\mathbf{Bad}^{\mathbf{b}}(\infty)$ is contained in the set of singular matrices.

It is well-known that the singular set is just rationals in the one dimensional case, but this set is nontrivial in higher dimensional cases. Hence we consider $(m, n) = (1, 1)$ and $(m, n) \neq (1, 1)$, separately. The one dimensional case is simple.

Corollary 1.3. *When $(m, n) = (1, 1)$, for any $b \in \mathbb{R} \setminus \mathbb{Z}$,*

$$\begin{cases} \mathbf{Bad}^b(\infty) = \mathbb{Q} & \text{if } b \notin \mathbb{Q}, \\ \mathbf{Bad}^b(\infty) = \{r/s \in \mathbb{Q} : \gcd(r, s) = 1, q \nmid s\} & \text{if } b = p/q \text{ with } \gcd(p, q) = 1. \end{cases}$$

Proof. Using Theorem 1.1, it is enough to show that $\mathbf{Sing}(b)$ is equal to the right hand side, and it can be easily checked. \square

Now assume $(m, n) \neq (1, 1)$. If $\mathbf{b} \in \mathbb{R}^m \setminus \mathbb{Z}^m$, then there exists nonzero $\mathbf{y} \in \mathbb{Z}^m$ such that $|\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} > 0$. Since the set of A such that ${}^t A \mathbf{y} \in \mathbb{Z}^n$ is contained in $\mathbf{Sing}(\mathbf{b})$, it follows from Theorem 1.1 that

$$\dim_H \mathbf{Bad}^{\mathbf{b}}(\infty) \geq n(m-1) \quad \text{and} \quad \dim_H \mathbf{Bad}(\infty) \geq n(m-1) + m.$$

Here and hereafter, \dim_H refers to the Hausdorff dimension. On the other hand, using Hausdorff dimension of singular sets [Che11, CC16, KKLM17, DFSU24], it follows from Remark 1.2 that

$$\dim_H \mathbf{Bad}^{\mathbf{b}}(\infty) \leq mn - \frac{mn}{m+n} \quad \text{and} \quad \dim_H \mathbf{Bad}(\infty) \leq mn + m - \frac{mn}{m+n}.$$

It is natural to ask the exact Hausdorff dimension for $\mathbf{Bad}^{\mathbf{b}}(\infty)$ and $\mathbf{Bad}(\infty)$ and our main application of Theorem 1.1 answers this question in the case that $m = 1$ and $n \geq 2$.

Theorem 1.4. *Assume that $m = 1$ and $n \geq 2$. For any $b \in \mathbb{R} \setminus \mathbb{Z}$,*

$$\dim_H \mathbf{Bad}^b(\infty) = \frac{n^2}{n+1} \quad \text{and} \quad \dim_H \mathbf{Bad}(\infty) = \frac{n^2}{n+1} + 1.$$

Moreover, if $b \in \mathbb{Q} \setminus \mathbb{Z}$, there is a constant $C > 0$ such that for any $t > 1$ and all small enough $\epsilon > 0$,

$$\frac{n^2}{n+1} + \epsilon^t \leq \dim_H \mathbf{Bad}^b(1/\epsilon) \leq \frac{n^2}{n+1} + C\epsilon^{\frac{1}{2}}.$$

It is worth noting that Theorem 1.4 is somewhat similar to [CC16, Theorems 1.1 and 1.3]. Indeed, our method relies on their method. This will be discussed at the end of the introduction.

In the view of Theorem 1.4, we may expect

Conjecture 1. *When $(m, n) \neq (1, 1)$, for any $\mathbf{b} \in \mathbb{R}^m \setminus \mathbb{Z}^m$,*

$$\dim_H \mathbf{Bad}^{\mathbf{b}}(\infty) = mn - \frac{mn}{m+n} \quad \text{and} \quad \dim_H \mathbf{Bad}(\infty) = mn + m - \frac{mn}{m+n}.$$

Now we focus on the set $\mathbf{Bad}_A(\infty)$. Following [DFSU24], we say that $A \in M_{m,n}(\mathbb{R})$ is *very singular* if $\widehat{w}(A) > \frac{n}{m}$. It can be easily checked that A is very singular if and only if ${}^t A$ is very singular (see Proposition 5.1). Following the discussion in Subsection 1.1, [BL05, Theorem] implies that if A is very singular, then $\mathbf{Bad}_A(\infty)$ is of full Lebesgue measure. The following theorem says that the complement of $\mathbf{Bad}_A(\infty)$ should be small with respect to $\widehat{w}({}^t A)$ in terms of Hausdorff dimension.

Theorem 1.5. *For any $A \in M_{m,n}(\mathbb{R})$,*

$$\dim_H(\mathbb{R}^m \setminus \mathbf{Bad}_A(\infty)) \leq m - \frac{\widehat{w}({}^t A) - \frac{m}{n}}{\widehat{w}({}^t A) + 1}.$$

In particular, if A is very singular, or equivalently, ${}^t A$ is very singular, then the complement of $\mathbf{Bad}_A(\infty)$ cannot have full Hausdorff dimension.

Remark 1.6.

- (1) In the view of [ET11, Kim24] and Theorem 1.5, the more singular A is, the larger $\mathbf{Bad}_A(\infty)$ becomes.
- (2) It seems interesting to give a lower bound for the Hausdorff dimension of the complement of $\mathbf{Bad}_A(\infty)$ or calculate its dimension precisely.
- (3) It was proved in [BKLR21, KKL] that A is *singular on average* if and only if there exists $\epsilon > 0$ such that $\mathbf{Bad}_A(\epsilon)$ has full Hausdorff dimension. Thus it seems interesting to reveal a certain equivalence between a Diophantine property of A and a metrical property of $\mathbf{Bad}_A(\infty)$.

1.3. Dynamical discussion. In this subsection, we will discuss the dynamical interpretation of infinitely badly approximability following classical relations between homogeneous dynamics and Diophantine approximation.

Consider the homogeneous space $\mathcal{X}_0 = \mathrm{SL}_{m+n}(\mathbb{R})/\mathrm{SL}_{m+n}(\mathbb{Z})$, which can be identified with the space of unimodular lattices in \mathbb{R}^{m+n} . For any $t \in \mathbb{R}$ and $A \in M_{m,n}(\mathbb{R})$, let us denote

$$a_t = \begin{pmatrix} e^{t/m} I_m & \\ & e^{-t/n} I_n \end{pmatrix} \quad \text{and} \quad u_A = \begin{pmatrix} I_m & A \\ & I_n \end{pmatrix}.$$

Define the function $\Delta_0 : \mathcal{X}_0 \rightarrow \mathbb{R}_{>0}$ by $\Delta_0(x) = \min_{v \in x \setminus \{0\}} \|v\|$. Dani's correspondence [Dan85] says that

$$A \text{ is singular} \iff \Delta_0(a_t u_A \mathbb{Z}^{m+n}) \rightarrow 0 \text{ as } t \rightarrow \infty.$$

Let $\mathrm{ASL}_{m+n}(\mathbb{R}) = \mathrm{SL}_{m+n}(\mathbb{R}) \ltimes \mathbb{R}^{m+n}$ and $\mathrm{ASL}_{m+n}(\mathbb{Z}) = \mathrm{SL}_{m+n}(\mathbb{Z}) \ltimes \mathbb{Z}^{m+n}$. The homogeneous space $\mathcal{X} = \mathrm{ASL}_{m+n}(\mathbb{R})/\mathrm{ASL}_{m+n}(\mathbb{Z})$ can be identified with the space of unimodular grids in \mathbb{R}^{m+n} , i.e. affine shifts of unimodular lattices in \mathbb{R}^{m+n} . Define the function $\Delta : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$ by $\Delta(x) = \min_{v \in x} \|v\|$. For $A \in M_{m,n}(\mathbb{R})$ and $\mathbf{b} \in \mathbb{R}^m$, consider the grid

$$\Lambda_{A,\mathbf{b}} = u_A \mathbb{Z}^{m+n} + \begin{pmatrix} \mathbf{b} \\ \mathbf{0} \end{pmatrix} \in \mathcal{X}.$$

Note that the diagonal matrix a_t acts naturally on \mathcal{X} .

Following [Kle99, Subsection 1.3], we say that $(A, \mathbf{b}) \in M_{m,n}(\mathbb{R}) \times \mathbb{R}^m$ is *rational* if $\|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} = 0$ for some $\mathbf{q} \in \mathbb{Z}^n$, and *irrational* otherwise. Using the proof of [Kle99, Theorem 4.4], we have the following characterization:

$$(A, \mathbf{b}) \in \mathbf{Bad}(\infty) \text{ and irrational} \iff \Delta(a_t \Lambda_{A,\mathbf{b}}) \rightarrow \infty \text{ as } t \rightarrow \infty.$$

As stated in [Kle99, Subsection 1.3], if (A, \mathbf{b}) is rational, then the badly approximability of (A, \mathbf{b}) is the same as that of A . Since there is no infinitely badly approximable linear form, we indeed have

$$(1.4) \quad (A, \mathbf{b}) \in \mathbf{Bad}(\infty) \iff \Delta(a_t \Lambda_{A,\mathbf{b}}) \rightarrow \infty \text{ as } t \rightarrow \infty.$$

Consider the natural projection $\pi : \mathcal{X} \rightarrow \mathcal{X}_0$ sending a grid $x = x_0 + v$ to the lattice x_0 . Then \mathcal{X} can be seen as the torus bundle of the base space

\mathcal{X}_0 . Since $\Delta^{-1}(0) = \mathcal{X}_0$, the dynamical property that $\Delta(a_t x) \rightarrow \infty$ as $t \rightarrow \infty$ means that the orbit $a_t x$ moves away from the base space \mathcal{X}_0 as $t \rightarrow \infty$.

The dynamical interpretation of Remark 1.2, which is weaker than Theorem 1.1, is the following (see Proposition 5.1):

$$\Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty \implies \Delta_0(a_t \pi(x)) \rightarrow 0 \text{ as } t \rightarrow \infty.$$

Using this implication, it follows from [KKLM17, Theorem 1.1] that

$$(1.5) \quad \dim_H \{x \in \mathcal{X} : \Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty\} \leq \dim_H \mathcal{X} - \frac{mn}{m+n}.$$

As an application of Theorem 1.4 and this upper bound, we have

Theorem 1.7. *When $m = 1$ and $n \geq 2$,*

$$\dim_H \{x \in \mathcal{X} : \Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty\} = \dim_H \mathcal{X} - \frac{n}{n+1}.$$

Similar to Conjecture 1, we may expect

Conjecture 2. *When $(m, n) \neq (1, 1)$,*

$$\dim_H \{x \in \mathcal{X} : \Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty\} = \dim_H \mathcal{X} - \frac{mn}{m+n}.$$

1.4. Discussion of the proof of Theorem 1.4. In this paper, although Theorem 1.1 is philosophically significant, the most technically important part is Theorem 1.4. Thus, we will discuss the proof of Theorem 1.4.

Thanks to Theorem 1.1, it is enough to estimate the Hausdorff dimension of **Sing(b)**. For this, we will modify the method of [CC16] as follows. Comparing the new singular definition (1.3) with the classical one (1.2), we observe a difference of $|\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}}$ -term. Thus, if we consider the set of A in $M_{m,n}(\mathbb{R})$ such that for any $\epsilon > 0$ for all sufficiently large X the inequalities

$$(1.6) \quad \|{}^t A \mathbf{y}\|_{\mathbb{Z}} < \epsilon \delta X^{-\frac{m}{n}}, \quad \|\mathbf{y}\| < \delta X, \quad \text{and} \quad |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} > \delta$$

have an integer solution $\mathbf{y} \in \mathbb{Z}^m$, then this set is contained in **Sing(b)**. We can also consider this set as the set of singular matrices with solutions restricted on the set $\{\mathbf{y} \in \mathbb{Z}^m : |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} > \delta\}$. Therefore, we will modify the fractal structure developed in [CC16] so that the modified structure is contained in the singular set with restricted solutions, and show that this structure has the same dimension as the original structure.

Now let us discuss why we need the condition $b \in \mathbb{Q} \setminus \mathbb{Z}$ in the second argument of Theorem 1.4. In order to bound the Hausdorff dimension of **Bad** ^{b} ($1/\epsilon$), we need to bound the Hausdorff dimension of the set of ϵ -Dirichlet improvable vectors for b , which will be defined in Section 2. For this, δ in (1.6) should be fixed and it is the main difference from the case **Sing(b)**. If $b = c/d$ with $\text{gcd}(c, d) = 1$ and $d \geq 2$, then $|bk|_{\mathbb{Z}} < 1/d$ if and only if $|bk|_{\mathbb{Z}} = 0$ for any $k \in \mathbb{Z}$. This property eventually enable us to bound the Hausdorff dimension of **Bad** ^{b} ($1/\epsilon$) even if δ in (1.6) is fixed.

A similar idea to the above was used significantly in [KK22]. Of course, the dimension of the singular matrices was calculated in [DFSU24], but the

fractal structure remains unclear since they used the variational principle in the parametric geometry of numbers. On the other hand, the fractal structure was clearly given in [CC16] so that we can modify it. If one can reveal the fractal structure in [DFSU24], it seems plausible to modify the structure as above to obtain Conjectures 1 and 2.

Structure of the paper. In Section 2, we recall the transference principle following [Cas57] and prove Theorem 1.1. Section 3 is devoted to the modification of fractal structures of [CC16] in order to prove Theorem 1.4. Section 4 is devoted to the proof of Theorem 1.5. Section 5 is devoted to dynamical discussion including the proof of Theorem 1.7.

Convention. In what follows, the notation $A \ll B$ means that there exists a constant C (the *implied constant*) such that $A \leq CB$. The notation $A \asymp B$ means $A \ll B \ll A$.

Acknowledgments. I would like to thank Elon Lindenstrauss for his valuable comments and suggestion to consider Theorem 1.5. I further thank Yitwah Cheung for generously sharing the idea of [CC16].

2. TRANSFERENCE PRINCIPLE

In this section, we will prove a certain stronger transference argument than Theorem 1.1. For this, we introduce the following definition: For given $\epsilon > 0$ and $\mathbf{b} \in \mathbb{R}^m$, we say that ${}^tA \in M_{n,m}(\mathbb{R})$ is ϵ -Dirichlet improvable for \mathbf{b} if for all sufficiently large X the inequalities

$$\|{}^tA\mathbf{y}\|_{\mathbb{Z}} < \epsilon |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} X^{-\frac{m}{n}} \quad \text{and} \quad \|\mathbf{y}\| < |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} X$$

have an integer solution $\mathbf{y} \in \mathbb{Z}^m$. Denote by $\mathbf{DI}_{\epsilon}(\mathbf{b})$ the set of $A \in M_{m,n}(\mathbb{R})$ such that tA is ϵ -Dirichlet improvable for \mathbf{b} . Note that $\mathbf{Sing}(\mathbf{b}) = \bigcap_{\epsilon > 0} \mathbf{DI}_{\epsilon}(\mathbf{b})$.

The following is the main theorem of this section.

Theorem 2.1. *There are constants $c_1, c_2 > 0$ such that for any $\epsilon > 0$ and any $(A, \mathbf{b}) \in M_{m,n}(\mathbb{R}) \times \mathbb{R}^m$ the following holds: If tA is $c_1\epsilon^{m/n}$ -Dirichlet improvable for \mathbf{b} , then (A, \mathbf{b}) is $1/\epsilon$ -badly approximable. On the other hand, if (A, \mathbf{b}) is $1/\epsilon$ -badly approximable, then tA is $c_2\epsilon^{m/n}$ -Dirichlet improvable for \mathbf{b} . In particular, for any $\mathbf{b} \in \mathbb{R}^m$*

$$\mathbf{DI}_{c_1\epsilon^{\frac{m}{n}}}(\mathbf{b}) \subset \mathbf{Bad}^{\mathbf{b}}(1/\epsilon) \subset \mathbf{DI}_{c_2\epsilon^{\frac{m}{n}}}(\mathbf{b}).$$

Proof of Theorem 1.1. Taking intersection over all $\epsilon > 0$, Theorem 2.1 implies Theorem 1.1. \square

The following transference principle is the main ingredient of the proof of Theorem 2.1.

Theorem 2.2. [Cas57, Theorem XVII in Chapter V] *Let $A \in M_{m,n}(\mathbb{R})$, $\mathbf{b} \in \mathbb{R}^m$, $C > 0$, $X > 1$ be given.*

(1) A necessary condition that

$$(2.1) \quad \|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} \leq C \quad \text{and} \quad \|\mathbf{q}\| \leq X,$$

for some $\mathbf{q} \in \mathbb{Z}^n$ is that

$$(2.2) \quad |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} \leq \gamma \max(X \|{}^t A \mathbf{y}\|_{\mathbb{Z}}, C \|\mathbf{y}\|),$$

hold for all $\mathbf{y} \in \mathbb{Z}^m$ with $\gamma = m + n$.

(2) A sufficient condition that (2.1) has a solution $\mathbf{q} \in \mathbb{Z}^n$ is that (2.2) hold for all $\mathbf{y} \in \mathbb{Z}^m$ with $\gamma = 2^{m-1}((m+n)!)^{-2}$.

Proof of Theorem 2.1. Observe that (A, \mathbf{b}) is $1/\epsilon$ -badly approximable if and only if for all large enough $T > 1$, there is no solution $\mathbf{q} \in \mathbb{Z}^n$ such that

$$\|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} \leq \epsilon^{-1} T^{-\frac{n}{m}} \quad \text{and} \quad \|\mathbf{q}\| \leq T.$$

By Theorem 2.2 (2), this implies that for all large enough $T > 1$, there exists $\mathbf{y} \in \mathbb{Z}^m$ such that

$$\|{}^t A \mathbf{y}\|_{\mathbb{Z}} < |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} \gamma^{-1} T^{-1} \quad \text{and} \quad \|\mathbf{y}\| < |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} \gamma^{-1} \epsilon T^{\frac{n}{m}}.$$

If we substitute $\gamma^{-1} \epsilon T^{\frac{n}{m}} = X$, then we have $\gamma^{-1} T^{-1} = \gamma^{-1-\frac{m}{n}} \epsilon^{\frac{m}{n}} X^{-\frac{m}{n}}$, hence, ${}^t A$ is $c_2 \epsilon^{\frac{m}{n}}$ -Dirichlet improvable for some constant $c_2 > 0$.

Suppose that $A \in \mathbf{DI}_{c_1 \epsilon^{\frac{m}{n}}}(\mathbf{b})$ with some constant $c_1 > 0$ to be determined, that is, for all large enough $X > 1$, there exists $\mathbf{y} \in \mathbb{Z}^m$ such that

$$\|{}^t A \mathbf{y}\|_{\mathbb{Z}} < c_1 \epsilon^{\frac{m}{n}} |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} X^{-\frac{m}{n}} \quad \text{and} \quad \|\mathbf{y}\| < |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} X,$$

or equivalently,

$$|\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} > \gamma \max\left(\gamma^{-1} c_1^{-1} \epsilon^{-\frac{m}{n}} X^{\frac{m}{n}} \|{}^t A \mathbf{y}\|_{\mathbb{Z}}, \gamma^{-1} X^{-1} \|\mathbf{y}\|\right).$$

By Theorem 2.2 (1), there is no solution $\mathbf{q} \in \mathbb{Z}^n$ such that

$$\|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} \leq \gamma^{-1} X^{-1} \quad \text{and} \quad \|\mathbf{q}\| \leq \gamma^{-1} c_1^{-1} \epsilon^{-\frac{m}{n}} X^{\frac{m}{n}}.$$

By substituting $\gamma^{-1} c_1^{-1} \epsilon^{-\frac{m}{n}} X^{\frac{m}{n}} = T$ and taking $c_1 = \gamma^{-1-\frac{m}{n}}$, we have

$$\|A\mathbf{q} - \mathbf{b}\|_{\mathbb{Z}} \leq \epsilon^{-1} T^{-\frac{n}{m}} \quad \text{and} \quad \|\mathbf{q}\| \leq T.$$

Therefore, (A, \mathbf{b}) is $1/\epsilon$ -badly approximable. \square

3. DIMENSION ESTIMATES FOR $\mathbf{Bad}^b(\infty)$

In this section, we assume $m = 1$ and $n \geq 2$, and prove

Theorem 3.1. For any $b \in \mathbb{R} \setminus \mathbb{Z}$,

$$(3.1) \quad \dim_H \mathbf{Sing}(b) \geq \frac{n^2}{n+1}.$$

Moreover, if $b \in \mathbb{Q} \setminus \mathbb{Z}$, then for any $t > n$ and all small enough $\epsilon > 0$,

$$(3.2) \quad \dim_H \mathbf{DI}_{\epsilon}(b) \geq \frac{n^2}{n+1} + \epsilon^t.$$

Proof of Theorem 1.4. Combining Theorem 2.1, Theorem 3.1, and [CC16, Theorems 1.1 & 1.3], Theorem 1.4 follows. \square

In order to obtain the lower bound in Theorem 3.1, we basically follow the method of [CC16], but the set $\mathbf{Sing}(b)$ is smaller than the set of singular vectors as in Remark 1.2, hence we need to construct a certain smaller fractal structure than that of [CC16].

3.1. Self-similar coverings. We first recall the definition of self-similar structures and some related dimension results following [CC16].

We equip \mathbb{R}^n with the metric induced by the Euclidean norm $\|\cdot\|_2$. A *self-similar structure* on \mathbb{R}^n is a triple (J, σ, B) , where J is countable, $\sigma \subseteq J \times J$, and B is a map from J into the set of bounded subsets of \mathbb{R}^n . A σ -*admissible sequence* is a sequence $(x_k)_{k \in \mathbb{N}}$ in J such that for any k , $(x_k, x_{k+1}) \in \sigma$.

For a subset S of \mathbb{R}^n , a *self-similar covering* of S is a self-similar structure (J, σ, B) such that, for all θ in S , there exists a σ -admissible sequence $(x_k)_{k \in \mathbb{N}}$ in J such that

- $\lim_{k \rightarrow \infty} \text{diam } B(x_k) = 0$;
- $\bigcap_{k \in \mathbb{N}} B(x_k) = \{\theta\}$.

In this case, the self-similar structure (J, σ, B) is said to cover the subset S . Given $x \in J$, we denote

$$\sigma(x) = \{y \in J : (x, y) \in \sigma\}.$$

A triple (J, σ, B) is said to be a *strictly nested self-similar structure* on \mathbb{R}^n if it is a self-similar structure on \mathbb{R}^n such that

- for all $x \in J$, $\sigma(x)$ is finite, $B(x)$ is a nonempty compact subset of \mathbb{R}^n , and for all $y \in \sigma(x)$ we have $B(y) \subset B(x)$;
- for each σ -admissible sequence $(x_k)_{k \in \mathbb{N}}$, $\lim_{k \rightarrow \infty} \text{diam } B(x_k) = 0$;
- for each $x \in J$ and each $y \in \sigma(x)$, $\text{diam } B(y) < \text{diam } B(x)$.

We need the following dimension result.

Theorem 3.2. [CC16, Theorem 3.6] *Let S be a subset of \mathbb{R}^n . Suppose that there is a strictly nested self-similar structure (J, σ, B) that covers a subset of S , a subset $J_0 \subset J$ that contains a tail of any σ -admissible sequence, a function $\rho : J \rightarrow (0, 1)$, and two constants $c, s \geq 0$ such that*

- (1) *for each $x \in J_0$ and each $y \in \sigma(x)$, there are at most c points z in $\sigma(x) \setminus \{y\}$ such that*

$$d(B(y), B(z)) \leq \rho(x) \text{diam } B(x);$$

- (2) *for every $x \in J_0$,*

$$\sum_{y \in \sigma(x)} (\rho(y) \text{diam } B(y))^s \geq (c + 1) (\rho(x) \text{diam } B(x))^s;$$

- (3) *for all $\alpha \in J_0$ and all $y \in \sigma(x)$, $\rho(y) \text{diam } B(y) < \rho(x) \text{diam } B(x)$.*

Then S contains a subset of positive s -dimensional Hausdorff measure.

3.2. Fractal structure in [CC16]. In this subsection, we recall notation and the fractal structure given in [CC16]. Note that our ambient space is \mathbb{R}^n while the ambient space of [CC16] is \mathbb{R}^d . Therefore, d in [CC16] should be changed to n in our case. Since we do not consider the result in [CC16, Section 7], we only consider the fractal structure of [CC16, Section 6] in the case $x = y$.

Let

$$Q = \{(p_1, \dots, p_n, q) \in \mathbb{Z}^{n+1} : \gcd(p_1, \dots, p_n, q) = 1, q > 0\}.$$

Given $x = (p, q) = (p_1, \dots, p_n, q) \in Q$, we denote

$$|x| = q \quad \text{and} \quad \hat{x} = \frac{p}{q},$$

and define the *Farey lattice*

$$\Lambda_x = \mathbb{Z}^n + \mathbb{Z}\hat{x} = \pi_x(\mathbb{Z}^{n+1}),$$

where $\pi_x : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is the map given by $\pi_x(r, s) = r - s\hat{x}$ for $(r, s) \in \mathbb{R}^n \times \mathbb{R}$. Note that Λ_x is a lattice in \mathbb{R}^n with the covolume $\text{vol } \Lambda_x = |x|^{-1}$. Here and hereafter, $\text{vol } \Lambda$ stands for the covolume of the lattice Λ . For each $i = 1, \dots, n$, denote the i -th successive minimum of Λ_x by $\lambda_i(x)$ with respect to the Euclidean norm $\|\cdot\|_2$ on \mathbb{R}^n , and the normalized i -th successive minimum by $\hat{\lambda}_i(x) = |x|^{1/n} \lambda_i(x)$.

For each $x \in Q$, let Λ'_x be a codimension 1 sublattice of Λ_x with minimal covolume, H'_x be the real span of Λ'_x , and $H_x = \pi_x^{-1} H'_x$. Given $x \in Q$ and a primitive $\alpha \in \Lambda_x$, we let

$$\Lambda_{\alpha^\perp} = \pi_\alpha^\perp(\Lambda_x),$$

where π_α^\perp is the orthogonal projection of \mathbb{R}^n onto α^\perp , the subspace of vectors of \mathbb{R}^n orthogonal to α . For any $y \in Q$, let us denote

$$|x \wedge y| = |x||y|\|\hat{x} - \hat{y}\|_2.$$

For $y \in Q$ with $\pi_x(y) = \alpha$, the covolume of Λ_{α^\perp} is given by $\text{vol } \Lambda_{\alpha^\perp} = 1/|x \wedge y|$. As before, we denote the first successive minimum of Λ_{α^\perp} by $\lambda_1(\alpha)$ and its normalized version by $\hat{\lambda}_1(\alpha) = |y \wedge z|^{1/(n-1)} \lambda_1(\alpha)$. Given $\epsilon > 0$ and $x \in Q$, let

$$\Lambda_x(\epsilon) = \{\alpha \in \Lambda_x : \alpha \text{ is primitive and } \hat{\lambda}_1(\alpha) > \epsilon\}.$$

We fix a coset H' such that $\Lambda_x \cap H' \neq \emptyset$ and $\Lambda_x \setminus H'_x \subset \bigcup_{k \in \mathbb{Z}} kH'$ as in [CC16, Subsection 6.1], and denote by α_x^\perp the unique element of H' that is perpendicular to H'_x . Let $H'_x(k)$ be the coset $k\alpha_x^\perp + H'_x$ and let $\mathcal{C}(x)$ be the set of vectors in \mathbb{R}^n whose angle with α_x^\perp is at most $\tan^{-1} A_d$, where the constant A_d is in the proof of [CC16, Lemma 8.6]. Let

$$\mathcal{C}'_k(x) = \mathcal{C}(x) \cap H'_x(k) \quad \text{and} \quad \mathcal{C}_N(x) = \bigcup_{k=1}^N \mathcal{C}'_k(x).$$

For $N \in \mathbb{N}$, $\epsilon > 0$, and $x \in Q$, define

$$F_N(x, \epsilon) = \bigcup_{\alpha \in \Lambda_x(\epsilon) \cap \mathcal{C}_N(x)} \zeta(x, \alpha, \epsilon),$$

where $\zeta(x, \alpha, \epsilon) = \left\{ y \in Q : \pi_x(y) = \alpha, |y| \in \frac{|x \wedge y|^{n/(n-1)}}{\epsilon^{n/(n-1)}}(1, 2) \right\}$.

The following fractal structure was established in [CC16] to estimate the lower bound of the Hausdorff dimension of the ϵ -Dirichlet improvable set.

Proposition 3.3. [CC16, Proposition 6.6] *Let*

$$\sigma_{\epsilon, N}(x) = F_N(x, \epsilon), \quad B(x) = B\left(\hat{x}, \frac{\lambda_1(x)}{2|x|}\right), \quad Q_{\epsilon, N} = \bigcup_{x \in Q} \sigma_{\epsilon, N}(x).$$

The triple $(Q_{\epsilon, N}, \sigma_{\epsilon, N}, B)$ is a strictly nested self-similar structure covering a subset of the 2ϵ -Dirichlet improvable set provided ϵ is small enough.

Cheung and Chevallier were able to bound from below the Hausdorff dimension of this fractal structure by establishing spacing and local finiteness properties of this structure [CC16, Subsection 6.3] and controlling the distribution of Farey lattices with bounded distortion [CC16, Section 8].

3.3. Fractal structure for $\mathbf{Sing}(b)$. In this subsection, we fix $b \in \mathbb{R} \setminus \mathbb{Z}$, construct a certain fractal structure for $\mathbf{Sing}(b)$, and prove (3.1) of Theorem 3.1. For this, we modify the strategy of [CC16, Proposition 6.13] appropriately.

Fix any $\delta > 0$ and consider two sequences $(\epsilon_i)_{i \geq 0}$ and $(N_i)_{i \geq 0}$ given by

$$\epsilon_i = \eta 4^{-i} \quad \text{and} \quad N_i = \eta^{-1} 5^{\frac{\eta}{\delta} i}$$

for some constant $\eta > 0$ to be determined. Fix arbitrary $x_0 \in Q$ and choose $\eta > 0$ small enough so that the set

$$\tilde{F}_{N_0}(x_0, \epsilon_0) = \{y = (p, q) \in F_{N_0}(x_0, \epsilon_0) : |bq|_{\mathbb{Z}} > \eta 2^{-1}\}$$

is non-empty. It is possible due to $b \in \mathbb{R} \setminus \mathbb{Z}$. Pick $x_1 \in \tilde{F}_{N_0}(x_0, \epsilon_0)$ and define Q_i with $i \geq 1$ recursively by

$$Q_1 = \{x_1\}, \quad Q_{i+1} = \bigcup_{x \in Q_i} \tilde{F}_{N_i}(x, \epsilon_i),$$

where

$$\tilde{F}_{N_i}(x, \epsilon_i) = \left\{ y = (p, q) \in F_{N_i}(x, \epsilon_i) : |bq|_{\mathbb{Z}} > \eta 2^{-(i+1)} \right\}.$$

Let $Q' = \bigcup_{i \geq 1} Q_i$ be a disjoint union. For each $x \in Q_i$, define $\sigma'(x) = \tilde{F}_{N_i}(x, \epsilon_i)$ and $B(x) = B\left(\hat{x}, \frac{\lambda_1(x)}{2|x|}\right)$. Then we have

Proposition 3.4. *The triple (Q', σ', B) is a strictly nested self-similar structure covering a subset of $\mathbf{Sing}(b)$.*

Proof. It follows from Proposition 3.3 that (Q', σ', B) is a strictly nested self-similar structure. For each σ' -admissible sequence (x_i) , since $\bigcap_i B(x_i)$ is a single point θ , it is enough to show that $\theta \in \mathbf{Sing}(b)$.

For any large enough $X \geq 1$, there exists $i \geq 1$ such that $\eta^{-1}2^i|x_i| < X \leq \eta^{-1}2^{i+1}|x_{i+1}|$. As in the proof of [CC16, Proposition 6.6], it follows from [CC16, Lemma 4.2] that each $x_i = (p_i, q_i)$ is a best approximate to θ . Hence, it follows from [CC16, Lemma 4.3(iii)] that

$$\|q_i\theta - p_i\| < 2\|q_i\hat{x}_{i+1} - p_i\|.$$

Since $x_{i+1} \in \tilde{F}_{N_i}(x_i, \epsilon_i)$, we have $|x_{i+1}| > (\frac{|x_i \wedge x_{i+1}|}{\epsilon_i})^{\frac{n}{n-1}}$, hence

$$\|q_i\hat{x}_{i+1} - p_i\|_2 = \frac{|x_i \wedge x_{i+1}|}{|x_{i+1}|} < \epsilon_i|x_{i+1}|^{-\frac{1}{n}}.$$

It follows from $\|\cdot\| \leq \|\cdot\|_2$ that

$$\|q_i\theta - p_i\| < 2\epsilon_i|x_{i+1}|^{-\frac{1}{n}} \leq 2^{1+\frac{1}{n}}\epsilon_i(\eta^{-1}2^i)^{\frac{1}{n}}X^{-\frac{1}{n}}$$

Since $x_i \in \tilde{F}_{N_{i-1}}(x_{i-1}, \epsilon_{i-1})$, we have $|bq_i|_{\mathbb{Z}} > \eta 2^{-i}$, hence

$$\|q_i\theta - p_i\| < 2^{1+\frac{1}{n}}\epsilon_i(\eta^{-1}2^i)^{1+\frac{1}{n}}|bq_i|_{\mathbb{Z}}X^{-\frac{1}{n}} \quad \text{and} \quad |q_i| < \eta 2^{-i}X < |bq_i|_{\mathbb{Z}}X.$$

Since $\epsilon_i = \eta 4^{-i}$ and $n \geq 2$, $\epsilon_i 2^{(1+\frac{1}{n})i} \rightarrow 0$ as $i \rightarrow \infty$. Therefore, $\theta \in \mathbf{Sing}(b)$. \square

As in the proof of [CC16, Proposition 6.13], using [CC16, Lemmas 6.8 and 6.9], we can find constants $c_\rho > 0$ and $c > 0$ such that the function $\rho : Q' \rightarrow (0, 1)$ given by

$$\rho(x) = c_\rho \left(\frac{1}{N_i} \right)^{\frac{n+1}{n-1}} \epsilon_i^{\frac{2n}{n-1}+n} \quad \text{for each } x \in Q_i$$

has the property that for any $y \in \tilde{F}_{N_i}(x, \epsilon_i)$ there are at most c points $z \in \tilde{F}_{N_i}(x, \epsilon_i) \setminus \{y\}$ such that $d(B(y), B(z)) \leq \rho(x) \text{diam } B(x)$. Moreover, since $\rho(x)$ decreases with i , we have

$$\rho(y) \text{diam } B(y) < \rho(x) \text{diam } B(x)$$

for all $y \in \sigma'(x)$ when ϵ_i is small enough. Therefore, the conditions (1) and (3) of Theorem 3.2 hold.

In order to show the condition (2) of Theorem 3.2, we need the following proposition, which is a modification of [CC16, Proposition 6.10].

Proposition 3.5. *If ϵ_i is small enough, then for each $x \in Q_i$ and all real numbers s and t in $[0, 2n]$, we have*

$$\sum_{y \in \tilde{F}_{N_i}(x, \epsilon_i)} \frac{\hat{\lambda}_1(y)^s}{|y|^t} \gg S_1(N_i, t) \frac{\epsilon_i^{s+(t-1)\frac{n}{n-1}}}{\hat{\lambda}_n(x)^{(t-n)\frac{n}{n-1}}|x|^t},$$

where

$$S_1(N_i, t) = \sum_{k=1}^{N_i} \frac{1}{k^{1+(t-n)\frac{n}{n-1}}} \frac{\text{card}(\mathcal{C}'_k(x) \cap \Lambda_x(\epsilon_i))}{\text{card}(\mathcal{C}'_k(x) \cap \Lambda_x)}.$$

Proof. Observe that

$$\begin{aligned} \tilde{F}_{N_i}(x, \epsilon_i) &= \left\{ y = (p, q) \in F_{N_i}(x, \epsilon_i) : |bq|_{\mathbb{Z}} > \eta 2^{-(i+1)} \right\} \\ &= \bigcup_{\alpha \in \Lambda_x(\epsilon_i) \cap \mathcal{C}_{N_i}(x)} \zeta(x, \alpha, \epsilon_i, b), \end{aligned}$$

where $\zeta(x, \alpha, \epsilon_i, b) = \{y = (p, q) \in \zeta(x, \alpha, \epsilon_i) : |bq|_{\mathbb{Z}} > \eta 2^{-(i+1)}\}$. Therefore, it is enough to show

$$(3.3) \quad \text{card } \zeta(x, \alpha, \epsilon_i, b) \asymp \text{card } \zeta(x, \alpha, \epsilon_i).$$

Then, the same proof of [CC16, Proposition 6.10] works in this case.

In order to prove (3.3), write $x = (r, s)$ and fix a point $y = (p, q) \in \pi_x^{-1}(\alpha) \cap Q$. Since $\text{gcd}(r, s) = 1$, we have

$$\pi_x^{-1}(\alpha) \cap Q = \{(p + \ell r, q + \ell s) \cap Q : \ell \in \mathbb{Z}\}.$$

Since $|x \wedge y| = \|\alpha\|_2 |x|$,

$$\text{card } \zeta(x, \alpha, \epsilon_i) = \text{card } \{\ell \in \mathbb{Z} : (p + \ell r, q + \ell s) \in Q, \gamma < q + \ell s < 2\gamma\},$$

where $\gamma = \gamma_{x, \alpha, \epsilon_i} = \left(\frac{\|\alpha\|_2 |x|}{\epsilon_i}\right)^{\frac{n}{n-1}}$. It follows from the primitivity of α that $\text{gcd}(p + \ell r, q + \ell s) = 1$ for any $\ell \in \mathbb{Z}$. Hence, we have

$$\text{card } \zeta(x, \alpha, \epsilon_i) = \text{card } \{\ell \in \mathbb{Z} : \gamma < q + \ell s < 2\gamma\};$$

$$\text{card } \zeta(x, \alpha, \epsilon_i, b) = \text{card } \left\{ \ell \in \mathbb{Z} : \gamma < q + \ell s < 2\gamma, |b(q + \ell s)|_{\mathbb{Z}} > \eta 2^{-(i+1)} \right\}.$$

If $\ell \in \mathbb{Z}$ is such that $\gamma < q + \ell s < 2\gamma$ but $|b(q + \ell s)|_{\mathbb{Z}} \leq \eta 2^{-(i+1)}$, then it follows from $x \in Q_i$, hence $|bs|_{\mathbb{Z}} > \eta 2^{-i}$, that

$$|b(q + (\ell \pm 1)s)|_{\mathbb{Z}} \geq |bs|_{\mathbb{Z}} - |b(q + \ell s)|_{\mathbb{Z}} > \eta 2^{-i} - \eta 2^{-(i+1)} = \eta 2^{-(i+1)}.$$

Therefore, we have

$$\frac{1}{2} \text{card } \zeta(x, \alpha, \epsilon_i) \leq \text{card } \zeta(x, \alpha, \epsilon_i, b) \leq \text{card } \zeta(x, \alpha, \epsilon_i).$$

This proves (3.3). \square

We are now ready for the proof of (3.1) of Theorem 3.1.

Proof of (3.1) of Theorem 3.1. We will prove

$$\dim_H \mathbf{Sing}(b) \geq \frac{n^2}{n-1} - \delta.$$

Since δ is arbitrary, this implies (3.1) of Theorem 3.1. From the above observations and Proposition 3.4, it is enough to show the condition (2) with $s = \frac{n^2}{n+1} - \delta$ of Theorem 3.2 about (Q', σ', B) .

First, it follows from [CC16, Proposition 6.11] that if ϵ_i is small enough,

$$(3.4) \quad S_1(N_i, t) \gg \sum_{k=1}^{N_i} \frac{1}{k^{1+(t-n)\frac{n}{n-1}}}.$$

In fact, the condition $t > n$ was required. However, the same proof works for $t \leq n$ sufficiently near n .

Now, we fix any $x \in Q_i$ and it follows from the proof of [CC16, Corollary 6.12] that $\hat{\lambda}_1(x) \asymp \epsilon_{i-1}$ and $\hat{\lambda}_n(x)^{-1} \gg \epsilon_{i-1}^{n-1}$. Combining Proposition 3.5 and (3.4) with $t = \frac{n+1}{n}s = n - \frac{n+1}{n}\delta$, we have

$$\begin{aligned} \sum_{y \in \tilde{F}_{N_i}(x, \epsilon_i)} \frac{(\rho(y) \operatorname{diam} B(y))^s}{(\rho(x) \operatorname{diam} B(x))^s} &= \left(\frac{\rho(y)}{\rho(x)} \right)^s \frac{|x|^t}{\hat{\lambda}_1(x)^s} \sum_{y \in \tilde{F}_{N_i}(x, \epsilon_i)} \frac{\hat{\lambda}_1(y)^t}{|y|^s} \\ &\gg \left(\frac{\rho(y)}{\rho(x)} \right)^s \epsilon_i^{s+(t-1)\frac{n}{n-1}} \epsilon_{i-1}^{(t-n)n-s} \sum_{k=1}^{N_i} \frac{1}{k^{1+(t-n)\frac{n}{n-1}}} \\ &\gg \left(\frac{N_i}{N_{i+1}} \right)^{\frac{n+1}{n-1}s} \left(\frac{\epsilon_{i+1}}{\epsilon_i} \right)^{\left(\frac{2n}{n-1}+n\right)s} 4^{-i\left(n-\frac{n+1}{n-1}\delta-(n+1)\delta\right)} 5^{i\frac{n(n+1)}{n-1}} \\ &\gg (5/4)^{in}. \end{aligned}$$

Therefore, if i is large enough, then the sum exceeds $c + 1$. This proves the condition (2) with $s = \frac{n^2}{n+1} - \delta$ of Theorem 3.2. \square

3.4. Fractal structure for $\mathbf{DI}_\epsilon(b)$. Now we assume that $b \in \mathbb{Q} \setminus \mathbb{Z}$. Similar to the previous subsection, we will construct a certain fractal structure for $\mathbf{DI}_\epsilon(b)$ and prove (3.2) of Theorem 3.1.

Let $b = c/d$ with $\gcd(c, d) = 1$ and $d \geq 2$. The key property we will use is the following: for any $k \in \mathbb{Z}$,

$$(3.5) \quad |bk|_{\mathbb{Z}} < 1/d \iff |bk|_{\mathbb{Z}} = 0.$$

Given $\epsilon > 0$, $x \in Q$, and $N \in \mathbb{N}$, define

$$F_{N,b}(x, \epsilon) = \{y = (p, q) \in F_N(x, \epsilon) : |bq|_{\mathbb{Z}} \geq 1/d\},$$

and define

$$\sigma_{\epsilon, N, b}(x) = F_{N,b}(x, \epsilon), \quad Q_{\epsilon, N, b} = \bigcup_{x \in Q} \sigma_{\epsilon, N, b}(x), \quad B(x) = B\left(\hat{x}, \frac{\lambda_1(x)}{2|x|}\right).$$

Similar to Proposition 3.4, we have

Proposition 3.6. *The triple $(Q_{\epsilon, N, b}, \sigma_{\epsilon, N, b}, B)$ is a strictly nested self-similar structure covering a subset of $\mathbf{DI}_{2d^{1+1/n}\epsilon}(b)$ provided ϵ is small enough.*

Proof. It follows from Proposition 3.3 that $(Q_{\epsilon, N, b}, \sigma_{\epsilon, N, b}, B)$ is a strictly nested self-similar structure. For each $\sigma_{\epsilon, N, b}$ -admissible sequence (x_i) , since $\bigcap_i B(x_i)$ is a single point θ , it is enough to show that $\theta \in \mathbf{DI}_{2\epsilon}(b)$.

For any large enough $X \geq 1$, there exists $i \geq 1$ such that $d|x_i| < X \leq d|x_{k+1}|$. Following the same proof of Proposition 3.4, if we write $x_i = (p_i, q_i)$, it follows from $x_{i+1} \in F_{N,b}(x_i, \epsilon)$ that

$$\|q_i\theta - p_i\| < 2\epsilon|x_{i+1}|^{-\frac{1}{n}} \leq 2\epsilon d^{\frac{1}{n}} X^{-\frac{1}{n}}.$$

Since $x_i \in F_{N,b}(x_{i-1}, \epsilon)$, we have $|bq_i|_{\mathbb{Z}} \geq 1/d$, hence

$$\|q_i\theta - p_i\| < 2\epsilon d^{1+\frac{1}{n}} |bq_i|_{\mathbb{Z}} X^{-\frac{1}{n}} \quad \text{and} \quad |q_i| < \frac{1}{d} X \leq |bq_i|_{\mathbb{Z}} X.$$

Therefore, $\theta \in \mathbf{DI}_{2d^{1+1/n}\epsilon}(b)$. \square

As before, in order to show the condition (2) of Theorem 3.2, we need the following proposition.

Proposition 3.7. *If ϵ is small enough, then for any $N \in \mathbb{N}$, each $x \in Q_{\epsilon, N, b}$, and all real numbers s and t in $[0, 2n]$, we have*

$$\sum_{y \in F_{N,b}(x, \epsilon)} \frac{\widehat{\lambda}_1(y)^s}{|y|^t} \gg S_1(N, t) \frac{\epsilon^{s+(t-1)\frac{n}{n-1}}}{\widehat{\lambda}_n(x)^{(t-n)\frac{n}{n-1}} |x|^t},$$

where

$$S_1(N, t) = \sum_{k=1}^N \frac{1}{k^{1+(t-n)\frac{n}{n-1}}} \frac{\text{card}(\mathcal{C}'_k(x) \cap \Lambda_x(\epsilon))}{\text{card}(\mathcal{C}'_k(x) \cap \Lambda_x)}.$$

Proof. Similar to the proof of Proposition 3.5, since

$$\begin{aligned} F_{N,b}(x, \epsilon) &= \{y = (p, q) \in F_N(x, \epsilon) : |bq|_{\mathbb{Z}} \geq 1/d\} \\ &= \bigcup_{\alpha \in \Lambda_x(\epsilon) \cap \mathcal{C}_N(x)} \zeta(x, \alpha, \epsilon, b), \end{aligned}$$

where $\zeta(x, \alpha, \epsilon, b) = \{y = (p, q) \in \zeta(x, \alpha, \epsilon) : |bq|_{\mathbb{Z}} \geq 1/d\}$, it is enough to show that

$$(3.6) \quad \text{card} \zeta(x, \alpha, \epsilon, b) \asymp \text{card} \zeta(x, \alpha, \epsilon)$$

for all small enough $\epsilon > 0$. Then, the same proof of [CC16, Proposition 6.10] works in this case.

Following the same proof of Proposition 3.5, we have

$$\begin{aligned} \text{card} \zeta(x, \alpha, \epsilon) &= \text{card} \{\ell \in \mathbb{Z} : \gamma < q + \ell s < 2\gamma\}; \\ \text{card} \zeta(x, \alpha, \epsilon, b) &= \text{card} \{\ell \in \mathbb{Z} : \gamma < q + \ell s < 2\gamma, |b(q + \ell s)|_{\mathbb{Z}} \geq 1/d\}, \end{aligned}$$

where $x = (r, s)$, $y = (p, q) \in \pi_x^{-1}(\alpha) \cap Q$, and $\gamma = \gamma_{x, \alpha, \epsilon} = \left(\frac{\|\alpha\|_2 |x|}{\epsilon}\right)^{\frac{n}{n-1}}$. If $\ell \in \mathbb{Z}$ is such that $\gamma < q + \ell s < 2\gamma$ but $|b(q + \ell s)|_{\mathbb{Z}} < 1/d$, then $|b(q + \ell s)|_{\mathbb{Z}} = 0$ by (3.5). Thus, it follows from $x \in Q_{\epsilon, N, b}$, hence $|bs|_{\mathbb{Z}} \geq 1/d$, that

$$|b(q + (\ell \pm 1)s)|_{\mathbb{Z}} \geq |bs|_{\mathbb{Z}} - |b(q + \ell s)|_{\mathbb{Z}} \geq 1/d.$$

Therefore, we have

$$\frac{1}{2} \text{card} \zeta(x, \alpha, \epsilon) \leq \text{card} \zeta(x, \alpha, \epsilon, b) \leq \text{card} \zeta(x, \alpha, \epsilon).$$

This proves (3.6). \square

Proof of (3.2) of Theorem 3.1. Using Proposition 3.6 and Proposition 3.7, (3.2) of Theorem 3.1 follows by the same proof of [CC16, Corollary 6.12]. \square

4. DIMENSION ESTIMATES FOR $\mathbf{Bad}_A(\infty)$

In this section, we will prove Theorem 1.5. It is well-known that $A \in M_{m,n}(\mathbb{R})$ is singular if and only if ${}^tA \in M_{n,m}(\mathbb{R})$ is singular. This can be proved easily from the following transference theorem.

Theorem 4.1. [Cas57, Theorem II in Chapter V] *Let $0 < C < 1 \leq X$ be constants. Suppose that there is $\mathbf{q} \in \mathbb{Z}^n \setminus \{0\}$ such that*

$$\|A\mathbf{q}\|_{\mathbb{Z}} \leq C \quad \text{and} \quad \|\mathbf{q}\| \leq X.$$

Then there is $\mathbf{y} \in \mathbb{Z}^m \setminus \{0\}$ such that

$$\|{}^tA\mathbf{y}\|_{\mathbb{Z}} \leq D \quad \text{and} \quad \|\mathbf{y}\| \leq U,$$

where

$$D = (m+n-1)X^{\frac{1-m}{m+n-1}}C^{\frac{m}{m+n-1}}, \quad U = (m+n-1)X^{\frac{n}{m+n-1}}C^{\frac{1-n}{m+n-1}}.$$

We also obtain the following similar result for very singular matrices.

Proposition 4.2. *For $A \in M_{m,n}(\mathbb{R})$, A is very singular if and only if tA is very singular.*

Proof. It is enough to show one direction by the symmetry argument. Assume A is very singular, that is, $\widehat{w}(A) > \frac{n}{m}$. Choose $\delta > 0$ such that $\widehat{w}(A) > \frac{n}{m} + \delta$, that is, for all sufficiently large X there is $\mathbf{q} \in \mathbb{Z}^n$ such that

$$\|A\mathbf{q}\|_{\mathbb{Z}} < X^{-\frac{n}{m}-\delta} \quad \text{and} \quad 0 < \|\mathbf{q}\| < X.$$

Using Theorem 4.1, there is $\mathbf{y} \in \mathbb{Z}^m$ such that

$$\begin{aligned} \|{}^tA\mathbf{y}\|_{\mathbb{Z}} &\leq (m+n-1)X^{\frac{1-m}{m+n-1}-\frac{n}{m+n-1}-\delta\frac{m}{m+n-1}} = (m+n-1)X^{-1-\delta\frac{m}{m+n-1}}; \\ \|\mathbf{y}\| &\leq (m+n-1)X^{\frac{n}{m+n-1}-\frac{n}{m}\frac{1-n}{m+n-1}-\delta\frac{1-n}{m+n-1}} = (m+n-1)X^{\frac{n}{m}+\delta\frac{n-1}{m+n-1}}. \end{aligned}$$

By taking $Y = (m+n)X^{\frac{n}{m}+\delta\frac{n-1}{m+n-1}}$, we have

$$\|{}^tA\mathbf{y}\|_{\mathbb{Z}} < Y^{-\frac{m}{n}-\delta'} \quad \text{and} \quad 0 < \|\mathbf{y}\| < Y$$

for some small $\delta' > 0$. This proves that tA is very singular. \square

Now we recall the result in [Kim24]. Assume that the subgroup ${}^tAZ^m + \mathbb{Z}^n$ of \mathbb{R}^n has maximal rank $m+n$ over \mathbb{Z} . Following [BL05, Section 3], there exists a sequence of best approximations $(\mathbf{y}_k)_{k \geq 1}$ in \mathbb{Z}^m for tA . Denote $Y_k = \|\mathbf{y}_k\|$, $M_k = \|{}^tA\mathbf{y}_k\|_{\mathbb{Z}}$, and

$$\gamma_k = \max \left(\left(Y_k^{\frac{m}{n}} M_{k-1} \right)^{\frac{n}{m+n}}, \left(Y_{k+1}^{\frac{m}{n}} M_k \right)^{\frac{n}{m+n}} \right).$$

For any $\alpha > 0$, let

$$B_\alpha(\{\gamma_k\}) = \{\mathbf{b} \in [0, 1]^m : |\mathbf{b} \cdot \mathbf{y}_k|_{\mathbb{Z}} > \alpha\gamma_k \text{ for all large enough } k \geq 2\}.$$

Proposition 4.3. [Kim24, Proposition 5.1] *For any $\alpha > 0$,*

$$B_\alpha(\{\gamma_k\}) \subset \mathbf{Bad}_A\left(\frac{\alpha - n}{m}\right).$$

Using this proposition, we are able to prove Theorem 1.5.

Proof of Theorem 1.5. Since the theorem is trivial if $\widehat{w}(^tA) \leq \frac{m}{n}$, we may assume that $\widehat{w}(^tA) > \frac{m}{n}$, i.e. tA is very singular.

We first consider the case that the rank of the subgroup $^tAZ^m + \mathbb{Z}^n$ over \mathbb{Z} is strictly less than $m + n$. If so, we can choose a nonzero $\mathbf{y} \in \mathbb{Z}^m$ such that $^tA\mathbf{y} \in \mathbb{Z}^n$. Fix such nonzero $\mathbf{y} \in \mathbb{Z}^m$. Note that $\widehat{w}(^tA) = \infty$ in this case. Observe that if $|\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} > 0$, then tA is singular for \mathbf{b} . By Theorem 1.1, the set $\{\mathbf{b} \in \mathbb{R}^m : |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} > 0\}$ is contained in $\mathbf{Bad}_A(\infty)$. Hence, the complement $\mathbb{R}^m \setminus \mathbf{Bad}_A(\infty)$ is contained in the set $\{\mathbf{b} \in \mathbb{R}^m : |\mathbf{b} \cdot \mathbf{y}|_{\mathbb{Z}} = 0\}$, which is a countable union of $(m-1)$ -dimensional hyperplanes. This implies that $\dim_H(\mathbb{R}^m \setminus \mathbf{Bad}_A(\infty)) \leq m-1$.

Now assume that the subgroup $^tAZ^m + \mathbb{Z}^n$ has maximal rank $m+n$ over \mathbb{Z} . Since $\mathbf{b} \in \mathbf{Bad}_A(\infty)$ if and only if $\mathbf{b} + \mathbf{x} \in \mathbf{Bad}_A(\infty)$ for any $\mathbf{x} \in \mathbb{Z}^m$, by proposition 4.3, it is enough to show that for any $\alpha > 0$,

$$(4.1) \quad \dim_H([0, 1]^m \setminus B_\alpha(\{\gamma_k\})) \leq m - \frac{\widehat{w}(^tA) - \frac{m}{n}}{\widehat{w}(^tA) + 1}.$$

Observe that

$$[0, 1]^m \setminus B_\alpha(\{\gamma_k\}) = \limsup_{k \rightarrow \infty} \{\mathbf{b} \in [0, 1]^m : |\mathbf{b} \cdot \mathbf{y}_k|_{\mathbb{Z}} \leq \alpha \gamma_k\},$$

and for each k , the set $\{\mathbf{b} \in [0, 1]^m : |\mathbf{b} \cdot \mathbf{y}_k|_{\mathbb{Z}} \leq \alpha \gamma_k\}$ can be covered with $C_1 \frac{Y_k^m}{\gamma_k^{m-1}}$ balls of radius $C_2 \frac{\gamma_k}{Y_k}$ for some absolute constants $C_1, C_2 > 0$. Using Hausdorff-Cantelli Lemma [BD99, Lemma 3.10], it follows that for any $0 \leq s \leq m$, the s -dimensional Hausdorff measure

$$\mathcal{H}^s([0, 1]^m \setminus B_\alpha(\{\gamma_k\})) = 0 \quad \text{if} \quad \sum_{k \geq 2} \frac{Y_k^m}{\gamma_k^{m-1}} \left(\frac{\gamma_k}{Y_k}\right)^s < \infty.$$

Since tA is very singular, for any $0 < \delta < \widehat{w}(^tA) - \frac{m}{n}$, we have $Y_{k+1}^{\frac{m}{n} + \delta} M_k < 1$ for all sufficiently large k . Since $\gamma_k \leq Y_k^{-\frac{\delta n}{m+n}}$ for all sufficiently large k , it follows that

$$\sum_{k \geq 2} \frac{Y_k^m}{\gamma_k^{m-1}} \left(\frac{\gamma_k}{Y_k}\right)^s \leq \sum_{k \geq 2} Y_k^{m-s-(s-m+1)\frac{\delta n}{m+n}}$$

for any $m-1 \leq s \leq m$. For any $m - \frac{\delta n}{m+n+\delta n} < s < m$, since $m-s-(s-m+1)\frac{\delta n}{m+n} < 0$ and Y_k increases at least geometrically (see [BL05, Lemma 1]), it follows that

$$\sum_k Y_k^{m-s-(s-m+1)\frac{\delta n}{m+n}} < \infty.$$

Hence we have $\dim_H([0, 1]^m \setminus B_\alpha(\{\gamma_k\})) \leq s$. Taking $s \rightarrow m - \frac{\delta n}{m+n+\delta n}$ and $\delta \rightarrow \widehat{w}(tA) - \frac{m}{n}$, we finally have (4.1). \square

5. HOMOGENEOUS DYNAMICS

In this section, we discuss a relation between the infinitely badly approximable property and the dynamical property mentioned in Subsection 1.3. We first prove the following implication.

Proposition 5.1. *For any $x \in \mathcal{X}$,*

$$\Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty \implies \Delta_0(a_t \pi(x)) \rightarrow 0 \text{ as } t \rightarrow \infty.$$

Proof. Given a lattice Λ in \mathbb{R}^{m+n} , let us denote by $\lambda_j(\Lambda)$ be the j -th successive minimum of Λ , i.e. the infimum of λ such that the ball in \mathbb{R}^{m+n} of radius λ around 0 contains j independent vectors of Λ .

Given $x \in \mathcal{X}$ and $t \in \mathbb{R}$, let $\mathbf{v}_1, \dots, \mathbf{v}_{m+n}$ be independent vectors of the lattice $a_t \pi(x)$ satisfying $\|\mathbf{v}_i\| \leq \lambda_{m+n}(a_t \pi(x))$ for all $i = 1, \dots, m+n$. Since the parallelepiped $\Pi = \{\sum_i \alpha_i \mathbf{v}_i : \forall i, -1 \leq \alpha_i \leq 1\}$ contains a fundamental domain of the lattice $a_t \pi(x)$, there is a point \mathbf{w} of the grid $a_t x$ such that $\mathbf{w} + \Pi$ contains the origin 0. Hence, it follows that

$$\Delta(a_t x) \leq \|\mathbf{w}\| \leq \sum_{i=1}^{m+n} \|\mathbf{v}_i\| \leq (m+n) \lambda_{m+n}(a_t \pi(x)).$$

Since $\lambda_1(a_t \pi(x)) \cdots \lambda_{m+n}(a_t \pi(x)) \asymp 1$ by Minkowski's second theorem, we have $\Delta(a_t x) \ll \lambda_1(a_t \pi(x))^{-(m+n-1)}$. Therefore, if $\Delta(a_t x) \rightarrow \infty$ as $t \rightarrow \infty$, then $\Delta_0(a_t \pi(x)) \rightarrow 0$ as $t \rightarrow \infty$. \square

In order to prove Theorem 1.7, we need certain invariance properties for the divergence of the function Δ . Let us denote elements of $\text{ASL}_{m+n}(\mathbb{R})$ by $\langle g, \mathbf{v} \rangle$, where $g \in \text{SL}_{m+n}(\mathbb{R})$ and $\mathbf{v} \in \mathbb{R}^{m+n}$. Consider the action a_t on $\text{ASL}_{m+n}(\mathbb{R})$ by the left multiplication of $\langle a_t, 0 \rangle$ on $\text{ASL}_{m+n}(\mathbb{R})$. The *expanding horospherical subgroup* of $\text{ASL}_{m+n}(\mathbb{R})$ with respect to $(a_t)_{t \geq 0}$ is given by

$$H = \left\{ \left\langle \begin{pmatrix} I_m & A \\ 0 & I_n \end{pmatrix}, \begin{pmatrix} \mathbf{b} \\ 0 \end{pmatrix} \right\rangle : A \in M_{m,n}, \mathbf{b} \in \mathbb{R}^m \right\}.$$

On the other hand, the *nonexpanding horospherical subgroup* of $\text{ASL}_{m+n}(\mathbb{R})$ with respect to $(a_t)_{t \geq 0}$ is given by

$$\tilde{H} = \left\{ \left\langle \begin{pmatrix} P & 0 \\ R & Q \end{pmatrix}, \begin{pmatrix} 0 \\ \mathbf{c} \end{pmatrix} \right\rangle : P \in M_{m,m}, Q \in M_{n,n}, R \in M_{n,m}, \mathbf{c} \in \mathbb{R}^n \right\}.$$

Proposition 5.2. *For any $x \in \mathcal{X}$ and any $p \in \tilde{H}$,*

$$\Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty \iff \Delta(a_t p x) \rightarrow \infty \text{ as } t \rightarrow \infty.$$

Proof. The following is a key observation: for $p = \langle \begin{pmatrix} P & 0 \\ R & Q \end{pmatrix}, \begin{pmatrix} 0 \\ \mathbf{c} \end{pmatrix} \rangle \in \tilde{H}$,

$$a_t \left\langle \begin{pmatrix} P & 0 \\ R & Q \end{pmatrix}, \begin{pmatrix} 0 \\ \mathbf{c} \end{pmatrix} \right\rangle a_{-t} = \left\langle \begin{pmatrix} P & 0 \\ e^{-(\frac{1}{m} + \frac{1}{n})t} R & Q \end{pmatrix}, \begin{pmatrix} 0 \\ e^{-\frac{1}{n}t} \mathbf{c} \end{pmatrix} \right\rangle.$$

Therefore, for any $t \geq 0$ and any point \mathbf{v} of the grid x ,

$$\|a_t p \mathbf{v}\| = \|a_t p a_{-t} a_t \mathbf{v}\| \asymp \|a_t \mathbf{v}\|,$$

where the implied constant depends only on p . This concludes the proof of the proposition. \square

Proof of Theorem 1.7. Assume $m = 1$ and $n \geq 2$. By the upper bound (1.5), it is enough to show that

$$\dim_H \{x \in \mathcal{X} : \Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty\} \geq \dim_H \mathcal{X} - \frac{n}{n+1}.$$

Since the product map $\tilde{H} \times H \rightarrow \text{ASL}_{m+n}(\mathbb{R})$ is a local diffeomorphism, every element of a neighborhood of identity of $\text{ASL}_{m+n}(\mathbb{R})$ can be written as pu where $p \in \tilde{H}$ and $u \in H$. Therefore, using (1.4), Proposition 5.2, and Theorem 1.4, we have

$$\begin{aligned} \dim_H \{x \in \mathcal{X} : \Delta(a_t x) \rightarrow \infty \text{ as } t \rightarrow \infty\} &\geq \dim_H \mathbf{Bad}(\infty) + \dim_H \hat{H} \\ &= \dim_H \mathcal{X} - \frac{n}{n+1}. \end{aligned}$$

\square

REFERENCES

- [BD99] V. I. Bernik and M. M. Dodson, *Metric Diophantine approximation on manifolds*, Cambridge Tracts in Mathematics, vol. 137, Cambridge University Press, Cambridge, 1999. MR 1727177
- [BK15] R. Broderick and D. Kleinbock, *Dimension estimates for sets of uniformly badly approximable systems of linear forms*, Int. J. Number Theory **11** (2015), no. 7, 2037–2054. MR 3440444
- [BKLR21] Y. Bugeaud, D. H. Kim, S. Lim, and M. Rams, *Hausdorff dimension in inhomogeneous Diophantine approximation*, Int. Math. Res. Not. IMRN (2021), no. 3, 2108–2133. MR 4206606
- [BL05] Y. Bugeaud and M. Laurent, *On exponents of homogeneous and inhomogeneous Diophantine approximation*, Mosc. Math. J. **5** (2005), no. 4, 747–766, 972. MR 2266457
- [Cas57] J. W. S. Cassels, *An introduction to Diophantine approximation*, Cambridge Tracts in Mathematics and Mathematical Physics, vol. No. 45, Cambridge University Press, New York, 1957. MR 87708
- [CC16] Y. Cheung and N. Chevallier, *Hausdorff dimension of singular vectors*, Duke Math. J. **165** (2016), no. 12, 2273–2329. MR 3544282
- [Che11] Y. Cheung, *Hausdorff dimension of the set of singular pairs*, Ann. of Math. (2) **173** (2011), no. 1, 127–167. MR 2753601
- [Dan85] S. G. Dani, *Divergent trajectories of flows on homogeneous spaces and Diophantine approximation*, J. Reine Angew. Math. **359** (1985), 55–89. MR 794799
- [DFSU24] T. Das, L. Fishman, D. Simmons, and M. Urbański, *A variational principle in the parametric geometry of numbers*, Adv. Math. **437** (2024), Paper No. 109435, 130. MR 4671568
- [ET11] M. Einsiedler and J. Tseng, *Badly approximable systems of affine forms, fractals, and Schmidt games*, J. Reine Angew. Math. **660** (2011), 83–97. MR 2855820

- [Khi26] A. Khintchine, *Zur metrischen Theorie der diophantischen Approximationen*, Math. Z. **24** (1926), no. 1, 706–714. MR 1544787
- [Kim24] T. Kim, *On a Kurzweil type theorem via ubiquity*, Acta Arith. **213** (2024), no. 2, 181–191. MR 4742656
- [KK22] T. Kim and W. Kim, *Hausdorff measure of sets of Dirichlet non-improvable affine forms*, Adv. Math. **403** (2022), Paper No. 108353, 39. MR 4404030
- [KKL] T. Kim, W. Kim, and S. Lim, *Dimension estimates for badly approximable affine forms*, arXiv preprint, [arXiv:2111.15410](https://arxiv.org/abs/2111.15410).
- [KKLM17] S. Kadyrov, D. Kleinbock, E. Lindenstrauss, and G. A. Margulis, *Singular systems of linear forms and non-escape of mass in the space of lattices*, J. Anal. Math. **133** (2017), 253–277. MR 3736492
- [Kle99] D. Kleinbock, *Badly approximable systems of affine forms*, J. Number Theory **79** (1999), no. 1, 83–102. MR 1724255
- [MRS] N. Moshchevitin, A. Rao, and U. Shapira, *Badly approximable grids and k -divergent lattices*, arXiv preprint, [arXiv:2402.00196](https://arxiv.org/abs/2402.00196).
- [Sch69] W. M. Schmidt, *Badly approximable systems of linear forms*, J. Number Theory **1** (1969), 139–154. MR 248090
- [Sim18] D. Simmons, *A Hausdorff measure version of the Jarník-Schmidt theorem in Diophantine approximation*, Math. Proc. Cambridge Philos. Soc. **164** (2018), no. 3, 413–459. MR 3784262

THE EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL

Email address: taehyeong.kim@mail.huji.ac.il