

ON THE VARIETY OF TRITANGENTIAL PLANES TO A GENERAL K3 SURFACE OF DEGREE 6 AND GENUS 4 IN \mathbb{P}^4

CIRO CILIBERTO AND SANDRO VERRA

ABSTRACT. Let $S \subset \mathbb{P}^4$ be a general K3 surface of degree 6 and genus 4. In this paper we study the irreducible variety X_S of *tritangential planes* to S whose general point is a plane that intersects S in a curvilinear scheme of length six supported at three non collinear points. The variety X_S can be identified as the relevant part of the fixed locus of the so called *Beauville involution* defined on the Hilbert scheme $S[3]$ of 0-dimensional schemes of length three of S . In this paper we prove that: (a) X_S has dimension 3, is irreducible and smooth, except for 210 points that are at most of multiplicity 2 for X_S ; (b) X_S , in its natural embedding in the Grassmannian $\mathbb{G}(2, 4) \subset \mathbb{P}^9$ of planes in \mathbb{P}^4 , has degree 152.

CONTENTS

1.	Introduction	1
2.	The fixed locus of the Beauville involution for $g = 4$	3
3.	Irreducibility and local behaviour	4
	3.1. Irreducibility	4
	3.2. Regular and special planes	4
	3.3. Local behaviour	5
	3.4. Enumeration of the special planes	6
4.	The degree	9
	4.1. Schubert cycle computations	9
	4.2. The degree of \mathfrak{Z}_S	10
	References	16

1. INTRODUCTION

Let $S \subset \mathbb{P}^g$ be a smooth, irreducible, linearly normal, complex K3 surface of degree $2g - 2$ and genus g . In this paper we will assume that S is *general*, so that in particular $\text{Pic}(S) = \mathbb{Z}\langle \mathcal{O}_S(1) \rangle$. We will denote by $S[h]$ the Hilbert scheme of 0-dimensional subschemes of length h of S , that is a smooth, irreducible, projective variety of dimension $2h$.

There is a birational involution $\beta_S : S[g - 1] \dashrightarrow S[g - 1]$, called the *Beauville involution*, defined in the following way. It maps the general point $Z \in S[g - 1]$ to the scheme $Z' \in S[g - 1]$ such that the union of Z and Z' is the scheme theoretic intersection of S with the linear subspace of dimension $g - 2$ spanned by Z . The indeterminacy locus $\text{Ind}(\beta_S)$ of β_S consists of the set of those schemes $Z \in S[g - 1]$ such that the span of Z has dimension smaller than $g - 2$.

There is also a rational map $\text{sp}_S : S[g - 1] \dashrightarrow \mathbb{G}(g - 2, g)$ (where $\mathbb{G}(g - 2, g)$ is the Grassmannian of $(g - 2)$ -dimensional linear subspaces of \mathbb{P}^g) that maps the general $Z \in S[g - 1]$ to

the linear subspace of dimension $g - 2$ spanned by Z . The map sp_S is dominant, has degree $\binom{2g-2}{g-2}$, and it has the same indeterminacy locus $\text{Ind}(\beta_S)$ as β_S . Of course $\text{sp}_S(Z) = \text{sp}_S(\beta_S(Z))$, whenever $Z \in S[g - 1] \setminus \text{Ind}(\beta_S)$.

The involution β_S has been widely studied in the case $g = 3$, in which the indeterminacy locus is empty, mostly in relation with the locus of its fixed points, that identifies with the set of *bitangent lines* to a general quartic surface $S \subset \mathbb{P}^3$ (see, e.g., [5, 6, 15]). By contrast little is known about the involution β_S for $g > 3$. Here we will focus on the case $g = 4$, in which S is a general complete intersection surface of a smooth quadric \mathbb{Q} and a smooth cubic \mathbb{F} hypersurfaces in \mathbb{P}^4 (note that \mathbb{Q} is uniquely determined by S , whereas \mathbb{F} is not).

In this case the indeterminacy set $\text{Ind}(\beta_S)$ of β_S (and of sp_S) is the set of schemes $Z \in S[3]$ such that $\langle Z \rangle$ is a line: we call them *triples of collinear points*. The set $\text{Ind}(\beta_S)$ is easy to identify. Indeed $Z \in \text{Ind}(\beta_S)$ if and only if the line $\langle Z \rangle$ is contained in the unique quadric \mathbb{Q} containing S , so that there is a natural isomorphism between $\text{Ind}(\beta_S)$ and the Hilbert scheme of lines in \mathbb{Q} , that is isomorphic to \mathbb{P}^3 .

To resolve the indeterminacies of β_S (and of sp_S) one must blow-up $\text{Ind}(\beta_S)$ in $S[3]$. Let us consider $p_S : \widetilde{S[3]} \rightarrow S[3]$ this blow-up and let E_S be the exceptional divisor that sits over $\text{Ind}(\beta_S)$. Then E_S , that has dimension 5, is isomorphic to the set of pairs (Z, π) , with $Z \in \text{Ind}(\beta_S)$ and π a plane containing the line $\langle Z \rangle$.

The map sp_S extends to a morphism $\widetilde{\text{sp}}_S : \widetilde{S[3]} \rightarrow \mathbb{G}(2, 4)$ that acts as sp_S off E_S and on E_S it acts by sending a pair $(Z, \pi) \in E_S$ to the plane π .

The map β_S , in turn, extends to an involution morphism $\widetilde{\beta}_S : \widetilde{S[3]} \rightarrow \widetilde{S[3]}$, that coincides with β_S off E_S and on E_S works by mapping a pair $(Z, \pi) \in E_S$ to the pair $(Z', \pi) \in E_S$ such that the union of Z and Z' is the intersection scheme of S and π , so that $\widetilde{\beta}_S$ induces an involution on E_S .

In this paper we study the variety of fixed points of $\widetilde{\beta}_S$. This variety consists of two disjoint irreducible components (see §2 and §3.1). One component is the set Y_S of fixed points of the restriction of $\widetilde{\beta}_S$ on E_S , which can be easily described (see §2). The other component X_S , which is more important to us, has the property that any point $Z \in X_S$ is such that Z spans a plane and the general point Z of X_S consists of three distinct points and is such that the plane $\langle Z \rangle$ intersects S in a curvilinear scheme of length six supported on Z . We call such a plane a *3-tangential plane* to S at Z , and we will identify X_S with its image via sp_S , that is injective on X_S . Hence X_S can be seen as the closure in the Grassmannian $\mathbb{G}(2, 4)$ of the set of tritangential planes to S , and therefore it is called the *variety of tritangential planes* to S . This is the main character in this paper.

In §§3.3 and 3.4 we prove that X_S is smooth, except may be for the points of a 0-dimensional scheme of length 210, corresponding to the so called *special planes* for S , i.e., those tangent planes to S that intersect S uniquely at their tangency point with S . These points have multiplicity at most 2 for X_S . In order to compute the number 210 we make in Proposition 3.7 a more general computation, i.e., given a smooth, irreducible, projective surface in \mathbb{P}^4 , we compute the (expected) number of hyperplane sections of the surface with a triple point.

In §4 we compute the degree, equal to 152, of X_S considered as a subvariety of $\mathbb{G}(2, 4) \subset \mathbb{P}^9$. The main point of this computation is the computation of the degree of the scroll of (proper)

tritangent lines to a general projection in \mathbb{P}^3 of a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 .

We suspect that the variety X_S of tritangential planes a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 is of general type, but we have been unable to prove this.

It would be highly desirable to have a description as the one we make here for the fixed locus of β_S when S is a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 , also, more generally, for a general K3 surface S of degree $2g - 2$ and genus g in \mathbb{P}^g , for $g \geq 5$. This is however much harder. First of all it is in general more intricate the study of the indeterminacy locus $\text{Ind}(\beta_S)$ of β_S . The analogue of the variety of tritangential planes in the case of genus 4, is the closure X_S of the locus of $(g - 1)$ -*tangential* linear spaces of dimension $g - 2$ to S , i.e., of linear spaces of dimension $g - 2$ that intersect S in a curvilinear scheme of degree $2g - 2$ supported on a scheme Z consisting of $g - 1$ distinct points. It is very likely that this X_S is also irreducible and generically smooth, but studying its singular locus is definitely more complicated. Moreover the computation of the degree of X_S seems in general very hard, as well as the study of its birational structure, though we believe that X_S should always be of general type. We hope to come back on these topics in the future.

Acknowledgement: The authors are members of GNSAGA of the Istituto Nazionale di Alta Matematica ‘‘F. Severi’’.

2. THE FIXED LOCUS OF THE BEAUVILLE INVOLUTION FOR $g = 4$

Let $S \subset \mathbb{P}^4$ be a general K3 surface of degree 6 and genus 4. We want to study the set of fixed points of the Beauville involution $\widetilde{\beta}_S : \widetilde{S[3]} \rightarrow \widetilde{S[3]}$. First of all there is a set of fixed points on E_S . Precisely, $(Z, \pi) \in E_S$ is a fixed point for $\widetilde{\beta}_S$ if and only if the plane π is tangent along a line to the unique quadric \mathbb{Q} containing S , i.e., it intersects the quadric \mathbb{Q} along a line with multiplicity 2. For each line $\ell \subset \mathbb{Q}$ there is a unique plane π_ℓ that is tangent to \mathbb{Q} along ℓ , namely the intersection of all hyperplanes tangent to \mathbb{Q} at points of ℓ , hyperplanes that form a pencil with base locus the plane in question. Hence the set Y_S of fixed points of $\widetilde{\beta}_S$ along E_S is naturally isomorphic to the Hilbert scheme of lines in \mathbb{Q} , that is in turn isomorphic to \mathbb{P}^3 .

As mentioned in the Introduction, the remaining set of fixed points is denoted by X_S .

Proposition 2.1. *In the above set up one has $X_S \cap Y_S = \emptyset$.*

Proof. Suppose, by contradiction, that $X_S \cap Y_S \neq \emptyset$ and let $(Z, \pi) \in X_S \cap Y_S$. We can then find a flat family $\mathcal{Z} \rightarrow \mathbb{D}$, where \mathbb{D} is a disc, such that for $t \in \mathbb{D} \setminus \{0\}$ the fibre of $\mathcal{Z} \rightarrow \mathbb{D}$ over t is a scheme $Z_t \in S[3]$ such that $\pi_t := \langle Z_t \rangle$ is a 3-tangential plane to S at Z_t , whereas the fibre of $\mathcal{Z} \rightarrow \mathbb{D}$ over 0 is $Z_0 = Z$, and the flat limit of π_t , when t tends to 0, is $\pi_0 = \pi$.

Fix a general point $p \in \mathbb{P}^4$ and consider for all $t \in \mathbb{D}$, the hyperplane $H_t := \langle p, \pi_t \rangle$. Let C_t be the curve cut out by H_t on S . By the generality of p we may assume that the curve C_0 is smooth and therefore, by may be shrinking \mathbb{D} , we may assume that C_t is smooth for all $t \in \mathbb{D}$, so that C_t is a canonical curve of genus 4 for all $t \in \mathbb{D}$. Now for all $t \in \mathbb{D}$, $\mathcal{O}_{C_t}(Z_t)$ is a theta-characteristic on C_t . For $t \neq 0$, $\mathcal{O}_{C_t}(Z_t)$ is an odd theta-characteristic, whereas $\mathcal{O}_{C_0}(Z_0)$ is an even theta-characteristic, and this is impossible by the constancy of parity of theta-characteristics in flat families of curves (see [1, Appendix B, §33]). This proves the assertion. \square

By Proposition 2.1, any point $Z \in X_S$ is such that Z spans a plane, and the map sp_S is injective on X_S . Thus, in the rest of this paper, we will identify X_S with its image in $\mathbb{G}(2, 4)$.

3. IRREDUCIBILITY AND LOCAL BEHAVIOUR

3.1. Irreducibility. Let \mathcal{M}_g be the moduli space of curves of genus g and consider the variety \mathcal{S}_g^- of all pairs (C, θ) where C is a smooth irreducible curve of genus g and θ an odd theta-characteristic on C . We have the well known forgetful étale morphism $\mathcal{S}_g^- \rightarrow \mathcal{M}_g$ of degree $2^{g-1}(2^g - 1)$ and therefore all irreducible components of \mathcal{S}_g^- have dimension $3g - 3$. We will need the following result:

Theorem 3.1 (See [4], Lemma (6.3)). *The variety \mathcal{S}_g^- is irreducible for all $g \geq 2$. In particular the monodromy of the covering $\mathcal{S}_g^- \rightarrow \mathcal{M}_g$ is transitive for $g \geq 2$.*

Next we keep the set up of §2.

Theorem 3.2. *If S is a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 , then X_S is irreducible of dimension 3.*

Proof. Let us consider the variety \mathcal{X}_S of all pairs $(\pi, H) \in X_S \times (\mathbb{P}^4)^\vee$ such that $\pi \subset H$. We have the obvious projection

$$p_2 : (\pi, H) \in \mathcal{X}_S \longrightarrow H \in (\mathbb{P}^4)^\vee.$$

This is a finite morphism of degree 120, and this proves that \mathcal{X}_S has dimension 4. Look now at the other projection

$$p_1 : (\pi, H) \in \mathcal{X}_S \longrightarrow \pi \in X_S.$$

All fibres of p_1 are isomorphic to \mathbb{P}^1 . This shows that X_S has dimension 3.

To prove irreducibility, it suffices to show that \mathcal{X}_S is irreducible. To see this, notice that, since S is general, the general hyperplane section of S is a general curve of genus 4. From Theorem 3.1 it follows that the monodromy of p_2 is transitive, and this implies the irreducibility of \mathcal{X}_S , as wanted. \square

3.2. Regular and special planes. Let S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 , and let $\pi \in X_S$. Then π comes equipped with a 0-dimensional length 3 scheme Z such that the 0-dimensional length 6 intersection scheme Z' of π and S has the same support as Z and at every point of Z has length at least 2. If Z is curvilinear, we will say that π is *regular* for S . In this case the general hyperplane containing π cuts out on S , by Bertini's theorem, a smooth curve C , and the base point locus of the pencil of curves cut out on S by the pencil \mathcal{H}_π of hyperplanes containing π has base locus scheme the curvilinear 0-dimensional length 6 scheme Z' that has the same support as Z .

The plane $\pi \in X_S$ is called *special* for S if it is not regular. This is the case if and only if the 0-dimensional length 3 scheme Z is not curvilinear, hence there exists an intersection point p of π with S such that the defining ideal of Z on π is \mathfrak{m}_p^2 , where \mathfrak{m}_p is the maximal ideal of p on π . In particular π is tangent to S at p , the 0-dimensional length 6 intersection scheme Z' of π and S is supported at p and contains Z , and π intersects S only at p with intersection multiplicity 6.

Proposition 3.3. *Let S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 . Then a plane π of \mathbb{P}^4 is special for S if and only if there is a point $p \in S$ such that π is tangent to S at p and there is a hyperplane containing π that cuts out on S a curve with a triple point at p .*

Proof. Suppose that $\pi \in X_S$ is special. Then π is tangent to S at a point p and π intersects S only at p with intersection multiplicity 6. Consider the pencil of curves \mathcal{P}_π cut out on S by the pencil \mathcal{H}_π of hyperplanes containing π . The base locus scheme of \mathcal{P}_π is supported at p and all curves of \mathcal{P}_π are singular at p . Let C be a general curve in \mathcal{P}_π . It has a point of multiplicity exactly 2 at p , because otherwise two general curves of \mathcal{P}_π would have intersection multiplicity at least 9 at p , a contradiction. Let C' be another general curve in \mathcal{P}_π . Then C and C' have intersection multiplicity 6 at p . Then only the following two possibilities may occur:

- C and C' have both a node at p and the two branches of C at p are tangent to the two branches of C' at p ;
- C and C' have both a simple cusp at p with the same cuspidal tangent.

In either case, by imposing one single condition to the curves in \mathcal{P}_π we can obtain a curve C_0 in \mathcal{P}_π with multiplicity at least 3 at p . The multiplicity of C_0 at p has to be exactly 3 because otherwise C_0 would have intersection multiplicity at least 8 at p with the general curve C of \mathcal{P}_π , a contradiction.

Conversely, suppose we have a hyperplane H that cuts out on S a curve C_0 with a triple point p . Let π be the tangent plane to S at p , that is contained in H . Consider the pencil \mathcal{P}_π of curves cut out on S by the pencil \mathcal{H}_π of hyperplanes containing π . Then C_0 belongs to \mathcal{P}_π . The general curve C in \mathcal{P}_π is singular in p , so it must have a double point at p . Hence the base locus scheme of \mathcal{P}_π is supported at p , and this implies that π intersects S only at p with intersection multiplicity 6. This yields that π is special for S . \square

3.3. Local behaviour. Let again S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 . In this section we study the local structure of the points of X_S .

Theorem 3.4. *If $\pi \in X_S$ is regular, then it is a smooth point for X_S . If $\pi \in X_S$ is special then it is a point of multiplicity at most 2 for X_S .*

Proof. Let us consider again the variety \mathcal{X}_S introduced in the proof of Theorem 3.2, with the projection $p_2 : (\pi, H) \in \mathcal{X}_S \rightarrow H \in (\mathbb{P}^4)^\vee$, that is a finite morphism of degree 120. Let $\pi \in X_S$ be regular. Then we can choose a hyperplane H containing π such that the curve C cut out by H on S is smooth and π cuts out on C an odd theta-characteristic. The fibre $p_2^{-1}(H)$ consists of the points of the form (π', H) , with π' cutting out on C an odd theta-characteristic. This implies that the fibre $p_2^{-1}(H)$ consists of exactly 120 distinct points, hence p_2 is étale over H . This implies that \mathcal{X}_S is smooth at each point of the fibre $p_2^{-1}(H)$, in particular it is smooth at (π, H) . By looking at the projection $p_1 : (\pi, H) \in \mathcal{X}_S \rightarrow \pi \in X_S$, we see that \mathcal{X}_S is a \mathbb{P}^1 bundle over X_S . So that, being \mathcal{X}_S smooth at (π, H) , implies that X_S is smooth at π .

Let us assume next that $\pi \in X_S$ is special. Then we can choose a hyperplane H containing π such that the curve C cut out by H on S has a node or a simple cusp at a point p and it is otherwise smooth. Moreover the plane π intersects C only at p . Then the fibre $p_2^{-1}(H)$ consists of the points of the form (π', H) , with π' a plane whose intersection with C consists of 3 (proper or infinitely near) points each counted with multiplicity 2. Such planes are of two types:

- (i) planes π' that are tangent at three (proper or infinitely near) smooth points of C ;

(ii) planes π' passing through the double point p of C and tangent at two (proper or infinitely near) smooth points of C .

The planes of type (i) are exactly 64 (by [11, Cor. (2.7) and (2.13)]). As for the planes of type (ii), by projecting from p we get a smooth plane quartic C_0 and the planes in question project to bitangent lines to C_0 , which are 28. We claim that each of these planes has to be counted with multiplicity two. Let us give this for granted for the time being. Then the fibre $p_2^{-1}(H)$ contains the point (π, H) with multiplicity 2, and this implies that (π, H) is a point of multiplicity at most 2 for \mathcal{X}_S . Then, arguing as above, we see that π is a point of multiplicity at most 2 for X_S .

To finish our proof we have to prove the above claim. To do this, we note that C is a limit of a general nodal hyperplane section C' of S . Let H' be the hyperplane spanned by C' and let q be the node of C' . Then the fibre $p_2^{-1}(H')$ consists of the points of the form (π', H') , with:

- (i') π' a plane that is tangent at three (proper or infinitely near) smooth points of C' ;
- (ii') π' a plane passing through the node q of C and tangent at two (proper or infinitely near) smooth points of C .

The planes of type (i') are again 64. By projecting from q we get a general smooth plane quartic C'_0 and the planes of type (ii') project to bitangents to C'_0 , which are 28. By generality, we may assume that these bitangents are acted upon by a transitive monodromy. This shows that the planes of type (ii') have to be counted with the same multiplicity m . Since $64 + 28m = 120$, we see that $m = 2$, as wanted. \square

3.4. Enumeration of the special planes. To prove the next result we need a preliminary fact that we cite from [10]. Let \mathbb{D} be a disk and let $\mathcal{X} \rightarrow \mathbb{D}$ be a flat family of projective surfaces with smooth total space and smooth, irreducible general fiber \mathcal{X}_t , with $t \in \mathbb{D} \setminus \{0\}$, and whose special fiber $\mathcal{X}_0 = A \cup B$ consists of the union of two irreducible, reduced components A and B , intersecting transversally along a smooth curve $R = A \cap B$. Let \mathcal{L} be a line bundle on \mathcal{X} . Let $\mathcal{C} \rightarrow \mathbb{D}$ be a flat family of curves, with \mathcal{C} a section of \mathcal{L} on \mathcal{X} . We denote by C_t the fibre of $\mathcal{C} \rightarrow \mathbb{D}$ over $t \in \mathbb{D}$. We assume that, for $t \in \mathbb{D} \setminus \{0\}$, C_t is reduced, with a triple point p_t . We assume, moreover, that the curve C_t tends to the curve C_0 in such a way that p_t tends to a point $p \in R$. We will set $C_0 = C_A \cup C_B$ with C_A and C_B the components of C_0 on A and B respectively. The result we need is the following:

Proposition 3.5 ([10], §3). *In the above set up C_0 has at least the following singularity: both C_A and C_B have a double point at p with one branch tangent to R .*

The singularity of C_0 at p described in Proposition 3.5 will be called a *limit triple point*. Now we are in a position to prove the following:

Proposition 3.6. *Let S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 . There are only finitely many hyperplane sections of S having a triple point. In particular the set of special planes for S is finite.*

Proof. Let X be a general hypersurface of \mathbb{P}^4 of degree 3 with equation $f = 0$, let H, H' be general hyperplanes in \mathbb{P}^4 with equations $\ell = 0$ and $\ell' = 0$ respectively, and let Y be a general hypersurface of degree 2 in \mathbb{P}^4 with equation $g = 0$. We consider the threefold \mathcal{X}'' in $X \times \mathbb{A}^1 \subset \mathbb{P}^4 \times \mathbb{A}^1$ defined in $\mathbb{P}^4 \times \mathbb{A}^1$ by the equations $\{f = tg + \ell\ell' = 0, \text{ with } t \in \mathbb{A}^1\}$. Via the second projection $\mathcal{X}'' \rightarrow \mathbb{A}^1$ this is a flat family of surfaces, with smooth general fibre \mathcal{X}''_t ,

that is a general complete intersection surface of type $(2, 3)$ in \mathbb{P}^4 , and whose fibre over 0 is $\mathcal{X}_0'' = A \cup A'$ with $A = X \cap H$, $A' = X \cap H'$ two general cubic surfaces in \mathbb{P}^3 , and $A \cap A' = R$ is a smooth plane cubic curve, section of A and A' with the plane π with equations $\ell = \ell' = 0$.

We are interested in the singularities of \mathcal{X}'' in a neighborhood of the central fibre, i.e., we are interested in what happens if t belongs to a disc \mathbb{D} , centered at the origin of \mathbb{A}^1 . Thus we consider the flat family $\mathcal{X}' = \{f = tg + \ell\ell' = 0, \text{ with } t \in \mathbb{D}\} \rightarrow \mathbb{D}$. It is immediate to see that the singular locus of \mathcal{X}' coincides with the scheme with equation $t = f = g = \ell = \ell' = 0 \subset \mathcal{X}_0'$, that consists of six distinct points p_1, \dots, p_6 , located on the plane π and complete intersection there of the cubic R with the quadric Y . Moreover, \mathcal{X}' has points of type A_1 at p_1, \dots, p_6 . We resolve these singularities by blowing-up \mathcal{X}' at p_1, \dots, p_6 . One obtains a new family $\tilde{\mathcal{X}} \rightarrow \mathbb{D}$ with the same general fibre as $\mathcal{X}' \rightarrow \mathbb{D}$ and whose central fibre consists of eight components \tilde{A} and \tilde{A}' , the blow-ups of A and of A' along p_1, \dots, p_6 , plus the six exceptional divisors E_i over p_i , for $\leq i \leq 6$. Each of these exceptional divisors is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$. Now we can contract each of these exceptional divisors by contracting one of the two rulings of $\mathbb{P}^1 \times \mathbb{P}^1$. We choose to do this in the direction of \tilde{A}' . We thus obtain a new family of 3-folds $\mathcal{X} \rightarrow \mathbb{D}$, with \mathcal{X} smooth, with fiber $\mathcal{X}_t = \mathcal{X}'_t$ over $t \neq 0$, and whose central fiber $\mathcal{X}_0 = A \cup B$, where $B = \tilde{A}$. Moreover A and B intersect transversally along a curve that is isomorphic to R .

There is an obvious morphism $\mathcal{X} \rightarrow \mathbb{P}^4$, and we denote by \mathcal{L} the pull back of $\mathcal{O}_{\mathbb{P}^4}(1)$ via this morphism. The restriction \mathcal{L}_t of \mathcal{L} to \mathcal{X}_t , with $t \neq 0$, coincides with $\mathcal{O}_{\mathcal{X}_t}(1)$. The restriction \mathcal{L}_0 of \mathcal{L} to \mathcal{X}_0 maps $A \cup B$ to $A \cup A'$.

Suppose now, by contradiction, that for $t \in \mathbb{D}$ general, the surface \mathcal{X}_t has infinitely many hyperplane sections with a triple point. In particular, there would be a 1-dimensional family \mathcal{H}_t of such hyperplane sections of \mathcal{X}_t . Then also \mathcal{X}_0 would have a 1-dimensional family \mathcal{H}_0 of sections in \mathcal{L}_0 with a triple point or a limit triple point.

However, since A and A' are two general cubic surfaces in a \mathbb{P}^3 , they have no hyperplane section with a triple point (having a plane section with a triple point has codimension 1 in the moduli space of smooth cubics in \mathbb{P}^3 , see, e.g., [7, Thm. 1.3]). Hence there are no sections of \mathcal{L}_0 that have a triple point on A or B . So the triple points of the curves in \mathcal{H}_t for $t \in \mathbb{D} \setminus \{0\}$, would tend to points on the double curve R and would be limit triple points as described in Proposition 3.5. Now, if $p \in R = A \cap A'$ is a general point, there is a unique hyperplane section of $A \cup A'$ such that its two components C_A and $C_{A'}$ on A and A' respectively, have a double point at p . The hyperplane in question is the one containing the tangent planes to A and A' at p , that intersect along the tangent line to R at p . Then the two curves C_A and $C_{A'}$ do have a double point at p , but, if $p \in R$ is general, the branches of C_A and $C_{A'}$ are not tangent to R . This implies that if $p \in R = A \cap B$ is a general point, there is no section of \mathcal{L}_0 that has in p a limit triple point. In conclusion, in the limit, we cannot have a positive dimensional family of sections of \mathcal{L}_0 on \mathcal{X}_0 with a triple point or limit triple point, and this proves the assertion. \square

Next we want to prove that the set of special planes for a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 is non-empty and we want to compute their (expected) number. In order to do so, we need some more preliminaries.

Let V be a smooth, irreducible variety of dimension k endowed with a line bundle L . For any integer $n \geq 0$, let $J^n(V, L)$ be the n -th jet bundle relative to the pair (V, L) , that is defined in the following way (as a general reference on jet bundles see [14]). Consider $V \times V$ with the

two projections $p_i : V \times V \rightarrow V$, $1 \leq i \leq 2$, to the first and second factor. One defines

$$J^n(V, L) = p_{1*}(\mathcal{O}_{V \times V} / \mathcal{I}_{\Delta, V \times V}^n \otimes p_2^*(L))$$

where Δ is the diagonal in $V \times V$. Note that $J^0(V, L) = L$ and more generally $J^n(V, L)$ is a vector bundle of rank $\binom{n+k}{k}$. For all positive integers there is an exact sequence

$$(3.1) \quad 0 \longrightarrow L \otimes \text{Sym}^n(\Omega_V) \longrightarrow J^n(V, L) \longrightarrow J^{n-1}(V, L) \longrightarrow 0.$$

So, at a local level, we have

$$J^n(V, L) \sim L \otimes (1 \oplus \Omega_V \oplus \cdots \oplus \text{Sym}^n(\Omega_V))$$

and there is an evaluation map of vector bundles

$$\text{ev}_{V, L}^n : H^0(V, L) \otimes \mathcal{O}_V \longrightarrow J^n(V, L).$$

Let now $\Sigma \subset \mathbb{P}^4$ be a smooth, irreducible, projective, non-degenerate surface. We will denote by H_Σ the hyperplane bundle class of Σ . There is an *expected number* τ_Σ of hyperplane sections of Σ having a triple point. One has:

Proposition 3.7. *If $\Sigma \subset \mathbb{P}^4$ is a smooth, irreducible, projective, non-degenerate, linearly normal surface, we have*

$$(3.2) \quad \tau_\Sigma = 5K_\Sigma^2 + 20H_\Sigma \cdot K_\Sigma + 15H_\Sigma^2 + 5e(\Sigma)$$

where $e(\Sigma)$ is the Euler–Poincaré characteristic of Σ .

Proof. Consider the evaluation map

$$\text{ev}_{\Sigma, H_\Sigma}^2 : H^0(\Sigma, H_\Sigma) \otimes \mathcal{O}_\Sigma \longrightarrow J^2(\Sigma, H_\Sigma),$$

where $h^0(\Sigma, H_\Sigma) = 5$ and the rank of $J^2(\Sigma, H_\Sigma)$ is 6. The points $p \in \Sigma$ where $\text{ev}_{\Sigma, H_\Sigma}^2$ does not have maximal rank 5, are exactly the points p for which there is a section $s \in H^0(\Sigma, H_\Sigma)$ such that the curve C defined by $s = 0$ has a point of multiplicity at least 3 at p . So the required number τ_Σ is nothing but the second Chern class $c_2(J^2(\Sigma, H_\Sigma))$. Using the exact sequence (3.1), we have the identity of Chern polynomials

$$(3.3) \quad c_t(J^2(\Sigma, H_\Sigma)) = c_t(H_\Sigma) \cdot c_t(\Omega_\Sigma \otimes H_\Sigma) \cdot c_t(\text{Sym}^2(\Omega_\Sigma) \otimes H_\Sigma).$$

One has

$$c_t(H_\Sigma) = 1 + H_\Sigma t$$

$$c_t(\Omega_\Sigma \otimes H_\Sigma) = 1 + (K_\Sigma + 2H_\Sigma)t + (e(\Sigma) + K_\Sigma \cdot H_\Sigma + H_\Sigma^2)t^2$$

$$c_t(\text{Sym}^2(\Omega_\Sigma) \otimes H_\Sigma) = 1 + 3(K_\Sigma + H_\Sigma)t + (2K_\Sigma^2 + 4e(\Sigma) + 3H_\Sigma^2 + 6K_\Sigma \cdot H_\Sigma)t^2.$$

Plugging into (3.3) and computing, one checks that $\tau_\Sigma = c_2(J^2(\Sigma, H_\Sigma))$ is given by (3.2). \square

Remark 3.8. One can easily prove a more general result by letting Σ be not necessarily linearly normal and with improper double points. We do not dwell on this here.

Corollary 3.9. *Let S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 . Then $\tau_S = 210$, i.e., the expected number of special planes for S is 210.*

Proof. It follows from Proposition 3.7 by taking into account that $K_S = 0$, $H_S^2 = 6$ and $e(S) = 24$. \square

4. THE DEGREE

Let again S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 . In this section we will compute the degree of X_S as a subvariety of $\mathbb{G}(2, 4) \subset \mathbb{P}^9$. The result is the following:

Theorem 4.1. *If S is a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 , then the degree of X_S as a subvariety of in $\mathbb{G}(2, 4) \subset \mathbb{P}^9$ is 152.*

The proof consists of various steps that we will perform separately.

4.1. Schubert cycle computations. The Chow ring of $\mathbb{G}(2, 4)$ in dimension 3 is generated by two Schubert cycles, precisely

$$\begin{aligned} \mathbf{h} &= \{\text{planes contained in a fixed hyperplane}\}, \\ \mathbf{k} &= \{\text{planes intersecting a fixed plane in a line passing through a fixed point}\}. \end{aligned}$$

Hence in the Chow ring we have the relation

$$X_S = a\mathbf{h} + b\mathbf{k}$$

with a, b suitable integers. One has

$$\mathbf{h}^2 = \mathbf{k}^2 = 1, \quad \mathbf{h} \cdot \mathbf{k} = 0.$$

Hence

$$a = X_S \cdot \mathbf{h} = 120$$

because $X_S \cdot \mathbf{h}$ equals the number of tritangential planes to S contained in a general hyperplane, and this is the number of odd theta-characteristics of the curve of genus 4 general hyperplane section of S .

The number

$$b = X_S \cdot \mathbf{k}$$

is more complicated to compute, namely it is the number of (proper) *tritangent lines*¹ to the projection S' of S to a \mathbb{P}^3 from a general point, intersecting a general line of this \mathbb{P}^3 . The closure of the set of proper tritangent lines to S' is a ruled surface \mathfrak{Z}_S in \mathbb{P}^3 and b is the degree of \mathfrak{Z}_S .

Let $H_{\mathbb{G}}$ be the hyperplane class of $\mathbb{G}(2, 4)$, that is represented by the Schubert cycle

$$H_{\mathbb{G}} = \{\text{planes intersecting a fixed line}\}.$$

One has clearly $H_{\mathbb{G}}^3 \cdot \mathbf{h} = 1$ and $H_{\mathbb{G}}^3 \cdot \mathbf{k} = 2$. Hence:

Lemma 4.2. *One has*

$$\deg(X_S) = 120 + 2 \deg(\mathfrak{Z}_S).$$

¹A (proper) tritangent [resp. bitangent, resp. tangent] line ℓ to a surface Σ in \mathbb{P}^3 is such that ℓ is not contained in Σ and Σ cuts out on ℓ a divisor of the form $2D + D'$, with D, D' effective divisors, D of degree 3 [resp. of degree 2, resp. of degree 1] whose support lies in the smooth locus of Σ .

4.2. The degree of \mathfrak{Z}_S . To finish the proof of Theorem 4.1 we need to compute $\deg(\mathfrak{Z}_S)$. Before doing that we need the following preliminary information, which can be deduced from [13, Prop. 3.9]:

Proposition 4.3. *If $\tilde{S} \subset \mathbb{P}^3$ is a general sextic surface, the scroll of tritangent lines to \tilde{S} has degree 624.*

Next we can prove the following:

Proposition 4.4. *Let S be a general K3 surface of degree 6 and genus 4 in \mathbb{P}^4 . Then $\deg(\mathfrak{Z}_S) = 16$.*

Proof. Let S' be the projection of S from a general point to \mathbb{P}^3 . Note that S' is singular, having a canonical curve Γ of degree 6 and genus 4 of double points that are generically normal crossings with finitely many pinch points. The pull back $\tilde{\Gamma}$ of Γ on S is a smooth curve in $|\mathcal{O}_S(2)|$ and the projection map $\tilde{\Gamma} \rightarrow \Gamma$ is a double cover. Since $\tilde{\Gamma}$ has genus 13, by Riemann–Hurwitz formula $\tilde{\Gamma} \rightarrow \Gamma$ has 12 branch points, hence there are exactly 12 pinch points of S' along Γ .

Recall that \mathfrak{Z}_S is the scroll of (proper) tritangent lines to S' . Consider now a general surface \tilde{S} of degree 6 in \mathbb{P}^3 and consider the general specialization of \tilde{S} to S' . In this specialization the scroll $\mathfrak{Z}_{\tilde{S}}$ of tritangent lines to \tilde{S} specializes to the union of the following surfaces:

- (i) the scroll \mathfrak{Z}_S of (proper) tritangent lines to S' ;
- (ii) the scroll \mathfrak{Z}_1 of (proper) bitangent lines to S' intersecting Γ to be counted with a multiplicity m_1 ;
- (ii) the scroll \mathfrak{Z}_2 of (proper) tangent lines to S' that are also secant to Γ to be counted with a multiplicity m_2 ;
- (iii) the scroll \mathfrak{Z}_3 of trisecant lines to Γ to be counted with a multiplicity m_3 .

So we have

$$(4.1) \quad 624 = \deg(\mathfrak{Z}_{\tilde{S}}) = \deg(\mathfrak{Z}_S) + \sum_{i=1}^3 m_i \deg(\mathfrak{Z}_i).$$

We will compute m_i and $\deg(\mathfrak{Z}_i)$ for all $1 \leq i \leq 3$, and this will enable us to compute $\deg(\mathfrak{Z}_S)$.

Step 1: computation of $\deg(\mathfrak{Z}_1)$ and of the multiplicity m_1 .

The closure $\text{Bit}(S')$ of the set of (proper) bitangent lines to S' can be seen as a surface in $\mathbb{G}(1, 3) \subset \mathbb{P}^5$. The set $\text{Int}(\Gamma)$ of lines intersecting Γ can be seen as a hypersurface on $\mathbb{G}(1, 3)$, namely the *Cayley variety* of Γ , and it is well known that $\text{Int}(\Gamma)$ is cut out on $\mathbb{G}(1, 3)$ by a hypersurface of degree $6 = \deg(\Gamma)$ of \mathbb{P}^5 . Then \mathfrak{Z}_1 is contained in the intersection of $\text{Int}(\Gamma)$ with $\text{Bit}(S')$. More precisely we have

$$(4.2) \quad \text{Int}(\Gamma) \cdot \text{Bit}(S') = \mathfrak{Z}_1 + \mathfrak{D}_1 + \mathfrak{D}_2$$

where:

- \mathfrak{D}_1 is a perhaps non-reduced curve supported on the curve D_1 parametrizing the lines that intersect Γ at some point x , are tangent to a branch of S' passing through x and are also a proper tangent to S' at some point $y \neq x$;
- \mathfrak{D}_2 is a perhaps non-reduced curve supported on the irreducible curve D_2 parametrizing the

tangent lines to Γ .

We will look at D_1 and D_2 as a scrolls in \mathbb{P}^3 .

Claim 4.5. *The surface $\text{Bit}(S') \subset \mathbb{P}^5$ has degree 132, hence $\deg(\text{Int}(\Gamma) \cdot \text{Bit}(S')) = 792$.*

Proof of Claim 4.5. The Chow ring of $\mathbb{G}(1, 3)$ in dimension 2 is generated by the two Schubert cycles:

- P_1 the set of lines passing through a fixed point of \mathbb{P}^3 ;
 - P_2 the set of lines contained in a fixed plane of \mathbb{P}^3 ;
- and $P_i^2 = 1$ for $1 \leq i \leq 2$ whereas $P_1 \cdot P_2 = 0$.

We have the following relation in the Chow ring

$$\text{Bit}(S') = cP_1 + dP_2$$

with c, d integers, that we now will compute. The integer c is the number of (proper) bitangent lines to S' that pass through a general point of \mathbb{P}^3 . There is a formula that computes this number (see [9, p. 182] and [12] for a modern version), that gives $c = 60$. The integer d is the number of (proper) bitangent lines to S' contained in a general plane of \mathbb{P}^3 , i.e., the number of (proper) bitangent lines to a general plane section of S' that is a plane sextic curve with 6 nodes. This number is provided by Plücker formulae and it turns out that $d = 72$. Since P_1 and P_2 are two planes in $\mathbb{G}(1, 3)$, one has $\deg(\text{Bit}(S')) = 60 + 72 = 132$ as wanted. Then $\deg(\text{Int}(\Gamma) \cdot \text{Bit}(S')) = 6 \cdot 132 = 792$. \square

Claim 4.6. *One has $\deg(D_1) = 612$.*

Proof of Claim 4.6. Let ℓ be a general line of \mathbb{P}^3 . The degree of D_1 is the number of intersection points of ℓ with the scroll D_1 . To compute it we proceed as follows.

Consider the correspondence ω on ℓ defined in the following way. Take a general point $p \in \ell$. Consider all lines r passing through p , incident to Γ at a point x_r and such that r is tangent to a branch of S' passing through x_r . Such a line intersects S' off x_r at three further points $x_{r,1}, x_{r,2}, x_{r,3}$. Take the tangent planes $\pi_{r,1}, \pi_{r,2}, \pi_{r,3}$ to S' at $x_{r,1}, x_{r,2}, x_{r,3}$ respectively, and consider the points $p_{r,1}, p_{r,2}, p_{r,3}$ intersection of ℓ with $\pi_{r,1}, \pi_{r,2}, \pi_{r,3}$ respectively. The correspondence ω maps p to the divisor

$$E_p = \sum_r p_{r,1} + p_{r,2} + p_{r,3}$$

where the sum is taken over all lines r passing through p , incident to Γ at a point x_r and such that r is tangent to a branch of S' passing through x_r .

Let $[e, f]$ be the *indices* of ω , i.e., $e = \deg(E_p)$ and f is the degree of the divisor associated to a general point of ℓ in the inverse correspondence ω^{-1} . A point $z \in \ell$ sits in D_1 if and only if it is a *coincidence point* of ω , i.e., if and only if $\omega(z) = \omega^{-1}(z)$. The number of coincidence points of a correspondence of indices $[i, j]$ is clearly $i + j$, hence we have

$$(4.3) \quad \deg(D_1) = e + f.$$

So we have to compute e and f .

As for the computation of e we argue as follows. Take $p \in \ell$ as above. Consider the polar surface S'_p of S' with respect to p , which, by the generality of ℓ and p is a general polar of S' . Then S'_p contains Γ with multiplicity 1, and intersects S' in a further curve C_p of degree 18

that is well known to pass simply through the 12 pinch points of S' along Γ . The curve C_p is the closure of the set of smooth points z of S' such that the tangent plane to S' at z contains p . On S the curve C_p pulls back to a curve in $|\mathcal{O}_S(3)|$ and Γ to a curve in $|\mathcal{O}_S(2)|$. So they intersect in 36 points. Hence C_p and Γ intersect in 24 points off the 12 pinch points of S' on Γ . These points x are such that the line $r = \langle x, p \rangle$ is tangent to a branch of S' passing through x . So this proves that

$$(4.4) \quad e = 24 \cdot 3 = 72.$$

As for the computation of f we proceed as follows. Take a general point $q \in \ell$ and consider the curve C_q as above. Consider then the scroll T of all lines intersecting ℓ , Γ and C_q . Some properties of T are known by *Salmon–Cayley’s theorem* (see [2, p. 506–509]): T has degree 180, ℓ has multiplicity 72, Γ has multiplicity 18 and C_q has multiplicity 6 for T .

Let us consider the minimal desingularization $g : T' \rightarrow T$ of T and let $\tilde{\ell}$, $\tilde{\Gamma}$, \tilde{C}_q be the pull-back of ℓ , Γ and C_q on T' . It is clear that $\tilde{\ell}$, $\tilde{\Gamma}$, \tilde{C}_q are unisecant to the lines of the ruling, whose numerical class we denote by F . We will denote by H_T the pull back to T' of the hyperplane class of T .

Let us look at the pull-back on T' of the intersection of S' with T . We have

$$6H_T \sim 2\tilde{\Gamma} + \tilde{C}_q + A$$

where \sim stays for linear equivalence and A is a suitable curve. We have

$$6 = 6H_T \cdot F = 2\tilde{\Gamma} \cdot F + \tilde{C}_q \cdot F + A \cdot F = 2 + 1 + A \cdot F$$

hence $A \cdot F = 3$. Moreover, since $\tilde{\ell} \cdot \tilde{\Gamma} = \tilde{\ell} \cdot \tilde{C}_q = 0$, one has

$$A \cdot \tilde{\ell} = 6H_T \cdot \tilde{\ell} = 6 \cdot 72 = 432.$$

Furthermore we have

$$\tilde{\Gamma} \equiv \tilde{\ell} + \alpha F$$

where \equiv stays for numerical equivalence and α is a suitable integer. Then

$$6 \cdot 18 = H_T \cdot \tilde{\Gamma} = H_T \cdot \tilde{\ell} + \alpha = 72 + \alpha$$

hence $\alpha = 36$. So

$$A \cdot \tilde{\Gamma} = A \cdot \tilde{\ell} + 36(A \cdot F) = 432 + 3 \cdot 36 = 540.$$

Now notice that for the image z on Γ of each intersection point of A with $\tilde{\Gamma}$, the line of the ruling of T containing z is tangent to a branch of S' passing through z and intersects C_q in a point y such that the tangent plane to S' at y passes through q . This proves that

$$(4.5) \quad f = 540.$$

The assertion follows by (4.3), (4.4) and (4.5). \square

Claim 4.7. *One has $\deg(D_2) = 18$.*

Proof of Claim 4.7. This is an immediate consequence of Riemann–Hurwitz formula. \square

Claim 4.8. *The curve \mathfrak{D}_1 is reduced, so it coincides with D_1 . The curve \mathfrak{D}_2 is D_2 counted with multiplicity 4.*

Proof of Claim 4.8. Let us take $x \in S'$ a general point and let π_x be the tangent plane to S' at x . The section of S' with π_x is a curve C_x of degree 6 with six nodes, one at x and six more at the intersection points of π_x with Γ . Hence C_x has geometric genus 3, and the proper bitangent lines to S' passing through x correspond to the 12 ramification points of the g_4^1 pull-back on the normalization \tilde{C}_x of C_x of the series cut out on C_x by the lines of π_x passing through x .

Let us now see what happens when x tends to a general point ξ of Γ on a branch σ of S' passing through ξ . The plane π_x tends to the tangent plane $\pi_{\xi,\sigma}$ to the branch σ at ξ , that is also a tangent plane to Γ at ξ . This plane cuts out on S' a curve $C_{\xi,\sigma}$, still of geometric genus 3, that has an ordinary triple point at ξ and four more nodes at the further intersection points of $\pi_{\xi,\sigma}$ with Γ . Notice that the triple point of $C_{\xi,\sigma}$ at ξ has a distinguished branch γ that is the intersection of $\pi_{\xi,\sigma}$ with the other branch σ' of S' at ξ . If we remove from the triple point at ξ the branch γ , one is left with a node, that is cut out on the branch σ by the tangent plane $\pi_{\xi,\sigma}$ at ξ .

Now let us look at the limit, when x tends to ξ as indicated above, of the g_4^1 on the normalization \tilde{C}_x of C_x pull-back of the series cut out on C_x by the lines of π_x passing through x . The curve \tilde{C}_x flatly tends to the normalization $\tilde{C}_{\xi,\sigma}$ of $C_{\xi,\sigma}$. The g_4^1 on \tilde{C}_x tends to the g_3^1 pull-back on $\tilde{C}_{\xi,\sigma}$ of the linear series cut out on $C_{\xi,\sigma}$ by the lines of $\pi_{\xi,\sigma}$ passing through ξ , plus a fixed point, namely the point y of $\tilde{C}_{\xi,\sigma}$ corresponding to the branch γ of $C_{\xi,\sigma}$. It is then easily seen (see, e.g., [8, p. 179]) that the 12 ramification points of the g_4^1 tend to the 10 ramification points of the g_3^1 on $\tilde{C}_{\xi,\sigma}$ plus the divisor $2y$, and this implies the assertion. Indeed it is clear from the above argument that \mathfrak{D}_1 is reduced. As for \mathfrak{D}_2 , we see that the general point ξ of D_2 counts in \mathfrak{D}_2 with multiplicity two for each of the two branches of S' passing through ξ , and so it counts in \mathfrak{D}_2 with multiplicity 4. \square

Claim 4.9. *One has $\deg(\mathfrak{Z}_1) = 108$.*

Proof of Claim 4.9. The assertion follows by (4.2) and by Claims 4.6, 4.7 and 4.8. \square

Finally we have the:

Claim 4.10. *One has $m_1 = 2$.*

Proof of Claim 4.10. The problem is of a local nature and it boils down to the following. Let Σ be a surface in \mathbb{P}^3 with a double curve D along which it has generically normal crossings. One has to compute the multiplicity of a line intersecting Σ in a point of D as a point of the closure of the set of proper tangent lines to Σ . Since the problem is local we can argue as follows. Consider a smooth quadric Q in \mathbb{P}^3 that specializes to the union of two distinct planes intersecting along a line s . The family of tangent lines to Q is easily seen to be the intersection of $\mathbb{G}(1, 3)$ with a hypersurface of degree 2 in \mathbb{P}^5 . Under the above specialization this specializes to the set of lines intersecting the line s , which is the intersection of $\mathbb{G}(1, 3)$ with a hyperplane. Hence we see that this has to be counted with multiplicity 2. The assertion follows. \square

Step 2: computation of $\deg(\mathfrak{Z}_2)$ and of the multiplicity m_2 .

Claim 4.11. *One has $\deg(\mathfrak{Z}_2) = 90$.*

Proof of Claim 4.11. The set of secant lines to Γ can be considered as a surface $S(\Gamma)$ in $\mathbb{G}(1, 3) \subset \mathbb{P}^5$. The surface $S(\Gamma)$ is singular. Indeed Γ is contained in a unique smooth quadric

Q , that has two rulings of lines, and the lines of the two rulings are 3–secant to Γ . Hence the two rulings in question map on $\mathbb{G}(1, 3)$ to two irreducible conics that are sets of triple points for $S(\Gamma)$, and $S(\Gamma)$ is otherwise smooth. The normalization of $S(\Gamma)$ is smooth, it is isomorphic to the 2–fold symmetric product $\Gamma[2]$ of Γ and we have the normalization morphism $\nu : \Gamma[2] \rightarrow S(\Gamma)$. Let us denote by $H_{S(\Gamma)}$ the pull–back on $\Gamma[2]$ via ν of a hyperplane section of $S(\Gamma)$. Let r be a general line in \mathbb{P}^3 and consider the g_6^1 cut out on Γ by the planes containing r . Then we can consider $H_{S(\Gamma)}$ as consisting of all divisors of degree 2 contained in a divisor of this g_6^1 . It follows that

$$H_{S(\Gamma)}^2 = 21, \quad p_a(H_{S(\Gamma)}) = 22$$

(see [3, Prop. (1.3)]).

Take now a general line ℓ in \mathbb{P}^3 and consider the scroll Θ of lines that are secant to Γ and intersect ℓ . The surface Θ is singular. In particular, it is immediate that ℓ has multiplicity 6 and Γ has multiplicity 5 for Θ . We will consider the minimal desingularization $h : \Theta' \rightarrow \Theta$ of Θ and we will denote by $\tilde{\ell}$ and by $\tilde{\Gamma}$ the proper transform of ℓ and Γ on Θ' via h . We will also denote by F the numerical class of a line of the ruling of Θ' . Then it is clear that

$$F \cdot \tilde{\ell} = 1, \quad F \cdot \tilde{\Gamma} = 2.$$

Moreover, the ruling of Θ is birational to $H_{S(\Gamma)}$, hence its genus is 22, so that

$$(4.6) \quad p_a(\tilde{\ell}) = 22.$$

Let H_Θ be the pull–back on Θ' of a general plane section of Θ . It is clear that

$$H_\Theta^2 = H_{S(\Gamma)}^2 = 21.$$

By taking the pull–back on Θ' of a general plane section of Θ containing ℓ , we see that

$$H_\Theta \equiv \tilde{\ell} + 15F$$

and therefore

$$6 = H_\Theta \cdot \tilde{\ell} = \tilde{\ell}^2 + 15$$

hence

$$\tilde{\ell}^2 = -9.$$

One has

$$K_{\Theta'} \equiv -2\tilde{\ell} + \alpha F$$

with α a suitable integer, and, by (4.6), we have

$$42 = \tilde{\ell} \cdot (\tilde{\ell} + K_{\Theta'}) = \tilde{\ell}(-\tilde{\ell} + \alpha F) = 9 + \alpha$$

so that $\alpha = 33$.

Recall that Γ sits on a unique quadric Q . By pulling–back to Θ' the intersection of Q with Θ , we see that

$$2H_\Theta \equiv \tilde{\Gamma} + \beta F$$

with β a suitable integer. Intersecting with H_Θ we have

$$42 = 2H_\Theta^2 = H_\Theta \cdot \tilde{\Gamma} + \beta = 30 + \beta$$

so that $\beta = 12$. Hence

$$60 = 2H_\Theta \cdot \tilde{\Gamma} = \tilde{\Gamma}^2 + 12\tilde{\Gamma} \cdot F = \tilde{\Gamma}^2 + 24$$

thus

$$\tilde{\Gamma}^2 = 36.$$

By taking the pull-back to Θ' of the intersection of S' with Θ , we have

$$(4.7) \quad 6H_\Theta \sim 2\tilde{\Gamma} + A$$

where A is a suitable curve. Intersecting with F we see that $A \cdot F = 2$. Intersecting with $\tilde{\Gamma}$ we get

$$180 = 6H_\Theta \cdot \tilde{\Gamma} = 2\tilde{\Gamma}^2 + A \cdot \tilde{\Gamma} = 72 + A \cdot \tilde{\Gamma}$$

so that

$$A \cdot \tilde{\Gamma} = 108.$$

Now

$$2p_a(6H_\Theta) - 2 = 6H_\Theta \cdot (6H_\Theta + K_{\Theta'}) = 6H_\Theta \cdot (6H_\Theta - 2\tilde{\ell} + 33F) = 882$$

so that

$$p_a(6H_\Theta) = 442.$$

Similarly

$$2p_a(2\tilde{\Gamma}) - 2 = 2\tilde{\Gamma} \cdot (2\tilde{\Gamma} + K_{\Theta'}) = 2\tilde{\Gamma} \cdot (2\tilde{\Gamma} - 2\tilde{\ell} + 33F) = 276$$

so that

$$p_a(2\tilde{\Gamma}) = 139.$$

From (4.7) we deduce

$$442 = p_a(6H_\Theta) = p_a(2\tilde{\Gamma}) + p_a(A) + 2\tilde{\Gamma} \cdot A - 1 = 139 + p_a(A) + 216 - 1$$

so that

$$p_a(A) = 88.$$

Recall now that we have the 2:1 morphism $A \rightarrow \tilde{\ell}$ induced by the ruling of Θ' . By Hurwitz formula we see that the lines of the ruling tangent to A are exactly 90 and this is nothing else than the degree of \mathfrak{Z}_2 . \square

Claim 4.12. *One has $m_2 = 4$.*

Proof of Claim 4.12. The problem being local, the assertion follows from the argument in the proof of Claim 4.10. \square

Step 3: computation of $\deg(\mathfrak{Z}_3)$ and of the multiplicity m_3 .

Claim 4.13. *The image of \mathfrak{Z}_3 on $\mathbb{G}(1, 3)$ consists of two disjoint conics. Moreover $m_3 = 8$.*

Proof of Claim 4.13. Any trisecant line to Γ is a line of one of the two rulings of the unique smooth quadric Q containing Γ . The first assertion immediately follows. As for the second assertion, the problem being local, the assertion follows from the argument in the proof of Claim 4.10. \square

We are finally in a position to finish the proof of Proposition 4.4. Indeed, it follows right away from (4.1) and from Claims 4.9, 4.10, 4.11, 4.12, 4.13. \square

Finally we have:

Proof of Theorem 4.1. It follows from Proposition 4.4 and Lemma 4.2. \square

REFERENCES

- [1] E. Arbarello, M. Cornalba, Ph. Griffiths, J. Harris, *Geometry of Algebraic Curves, Vol. I*, Grundlehren der Math. Wissenschaften, **267**, Springer–Verlag, 1984.
- [2] L. Campedelli, *Geometria, Vol. II, Part II*, Cedam, Padova, 1948.
- [3] C. Ciliberto, *Sui sistemi di corrispondenze a valenza di una curva algebrica in sè*, Ricerche di Mat., **27**, (1978), 109–130.
- [4] M. Cornalba, *Moduli of curves and theta–characteristics*, in “Lectures on Riemann Surfaces”, ICTP, Trieste, Italy, 1987, Ed. M. Cornalba, X. Gomez–Mont, A. Verjovky, World Scientific, 1989, 560–589.
- [5] P. Corvaja, F. Zucconi, *Quartic surfaces, their bitangents and rational points*, Épijournal de Géométrie Algébrique, **7**, (2023).
- [6] P. Corvaja, F. Zucconi, *Bitangents to a quartic surface and infinitesimal Torelli*, to appear on Journal Diff. Geometry.
- [7] I. Coskun, A. Prendergast-Smith, *Eckardt loci on hypersurfaces*, Communications in Algebra, **43**(8), (2015), 3083–3101.
- [8] F. Enriques, O. Chisini, *Lezioni sulla teoria geometrica delle equazioni e delle funzioni algebriche, vol. I*, Zanichelli, Bologna, 1915.
- [9] F. Enriques, *Le superficie algebriche*, Zanichelli, Bologna, 1949.
- [10] C. Galati, *On the existence of curves with a triple point on a K3 surface*, Rend. Lincei Mat. Appl., **23** (3), (2012) 295–317.
- [11] J. Harris, *Theta–characteristics on algebraic curves*, Trans. of Amer. Math. Soc., **271** (2), (1982), 611–638.
- [12] B. Iversen, *Critical points of an algebraic function*, Invent. math., **12** (3), (1971), 210–224.
- [13] D. G. Markushevich, *Numerical invariants of families of lines on some Fano varieties*, Mathematics of the USSR-Sbornik, **44** (2), (1983), 239–260.
- [14] D. J. Saunders, *The Geometry of Jet Bundles*, Cambridge University Press, 1989.
- [15] G.E. Welters, *Abel–Jacobi isogenies for certain types of Fano threefolds*, Mathematical Centre Tracts 141, Mathematical Centre, 413 Kruislan, Amsterdam (1981), (available ir.cwi.nl pub Docworks PDF 600 dpi).

CIRO CILIBERTO, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00177 ROMA, ITALIA CILIBERTO@AXP.MAT.UNIROMA2.IT

SANDRO VERRA, DEPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ ROMA 3, LARGO SAN LEONARDO MURIALDO, 00146 ROMA, ITALIA SANDRO.VERRA@GMAIL.COM