

Tikhonov regularization of monotone operator flows not only ensures strong convergence of the trajectories but also speeds up the vanishing of the residuals

Radu Ioan Bot^{*}

Dang-Khoa Nguyen^{†‡}

August 21, 2024

Abstract

In the framework of real Hilbert spaces, we investigate first-order dynamical systems governed by monotone and continuous operators. It has been established that for these systems, only the ergodic trajectory converges to a zero of the operator. A notable example is the counterclockwise $\pi/2$ -rotation operator on \mathbb{R}^2 , which illustrates that general trajectory convergence cannot be expected. However, trajectory convergence is assured for operators with the stronger property of cocoercivity. For this class of operators, the trajectory's velocity and the operator values along the trajectory converge in norm to zero at a rate of $o(\frac{1}{\sqrt{t}})$ as $t \rightarrow +\infty$.

In this paper, we demonstrate that when the monotone operator flow is augmented with a Tikhonov regularization term, the resulting trajectory converges strongly to the element of the set of zeros with minimal norm. In addition, rates of convergence in norm for the trajectory's velocity and the operator along the trajectory can be derived in terms of the regularization function. In some particular cases, these rates of convergence can outperform the ones of the coercive operator flows and can be as fast as $\mathcal{O}(\frac{1}{t})$ as $t \rightarrow +\infty$. In this way, we emphasize a surprising acceleration feature of the Tikhonov regularization. Additionally, we explore these properties for monotone operator flows that incorporate time rescaling and an anchor point. For a specific choice of the Tikhonov regularization function, these flows are closely linked to second-order dynamical systems with a vanishing damping term. The convergence and convergence rate results we achieve for these systems complement recent findings for the Fast Optimistic Gradient Descent Ascent (OGDA) dynamics, leading to surprising outcomes.

When the monotone operator is defined as the identity minus a nonexpansive operator, the monotone equations transform into a fixed point problem. In such cases, explicitly discretizing the system with Tikhonov regularization, enhanced by an anchor point, leads to the Halpern fixed point iteration. Through convergence analysis, which can be viewed as the discrete counterpart of the continuous time analysis, we identify two regimes for the regularization sequence which ensure that the generated sequence of iterates converges strongly to the fixed point nearest to the anchor point. Furthermore, we establish a general theoretical framework that provides convergence rates for the vanishing of the discrete velocity and the fixed point residual. For certain regularization sequences, we derive specific convergence rates that align with those observed in continuous time.

Key Words. monotone equations, Tikhonov regularization, strong convergence, convergence rates, Lyapunov analysis,

AMS subject classification. 47J20, 47H05, 65K10, 65K15, 90C25

Contents

1	Introduction	2
2	First-order dynamical system with Tikhonov regularization	5
2.1	Convergence analysis and rates of convergence	6
2.2	First-order dynamical system with Tikhonov regularization and time rescaling	12
2.3	First-order dynamical system with anchor point	19

^{*}Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria, email: radu.bot@univie.ac.at. Research partially supported by FWF (Austrian Science Fund), projects W 1260 and P 34922-N.

[†]Faculty of Mathematics and Computer Science, University of Science, Ho Chi Minh City, Vietnam, email: ndkhoa@hcmus.edu.vn.

[‡]Vietnam National University, Ho Chi Minh City, Vietnam

3 From first-order dynamics with Tikhonov regularization to second-order dynamics with vanishing damping

4 Fixed point problem and discretization	21
4.1 Problem formulation and overview of existing numerical methods	21
4.2 Convergence results of the discretization	23
4.3 The averaged operator case	29
A Appendix	31
A.1 The cocoercive case	31
A.2 Auxiliary results used in the convergence analysis of the continuous time models	32
A.3 Auxiliary results used in the convergence analysis of the discrete time models	33

1 Introduction

Let \mathcal{H} be a real Hilbert space and $M: \mathcal{H} \rightarrow \mathcal{H}$ a continuous and monotone operator. We say that $M: \mathcal{H} \rightarrow \mathcal{H}$ is *monotone* if

$$\langle M(x) - M(y), x - y \rangle \geq 0 \quad \forall x, y \in \mathcal{H}.$$

Being monotone and continuous, M is also maximally monotone ([11]), which means that there is no other (set-valued) monotone operator with the graph in $\mathcal{H} \times \mathcal{H}$ being a strict superset of the graph of M . Therefore, the set $\text{Zer}M := \{x \in \mathcal{H} : M(x) = 0\}$ is convex and closed.

By assuming throughout this work that $\text{Zer}M$ is nonempty, we are interested in the investigation of continuous time models that solve the following equation

$$\text{Find } x \in \mathcal{H} \text{ such that } M(x) = 0. \tag{1.1}$$

A broad class of problems arising in applications can be reduced to the solving of a monotone equation in real Hilbert spaces such as (1.1).

Inspired by the gradient flow, we first attach to (1.1) the following dynamics

$$\dot{x}(t) + M(x(t)) = 0, \tag{1.2}$$

with starting time $t_0 \geq 0$ and initial condition $x(t_0) := x_0 \in \mathcal{H}$. We assume that it admits a unique strong global solution $x: [t_0, +\infty) \rightarrow \mathcal{H}$ (see [20, Appendix]) with the property that $t \mapsto M(x(t))$ is absolutely continuous on every compact interval $[t_0, T]$. If $M: \mathcal{H} \rightarrow \mathcal{H}$ is Lipschitz continuous, then this holds for every $x_0 \in \mathcal{H}$.

It has been long recognized (see, for instance, [9]) that only the ergodic trajectory $t \mapsto \frac{1}{t-t_0} \int_{t_0}^t x(s) ds$ converges weakly to a zero of M , while the trajectory $t \mapsto x(t)$ may not converge in general. To see this, one can consider the example ([24]) of the counterclockwise $\pi/2$ -rotation operator on \mathbb{R}^2 , $M: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $M(x_1, x_2) = (-x_2, x_1)$, which has $(0, 0)$ as its unique zero. For $t_0 = 0$ and $(x_1(0), x_2(0)) := r_0(\cos(\theta_0), \sin(\theta_0))$, for $r_0 > 0$ and $\theta_0 \in \mathbb{R}$, the solution trajectory of (1.2), $(x_1(t), x_2(t)) = r_0(\cos(\theta_0 - t), \sin(\theta_0 - t))$, is bounded, but does not have a limit point as $t \rightarrow +\infty$. On the other hand, the ergodic trajectory $t \mapsto \frac{1}{t} \int_0^t (x_1(s), x_2(s)) ds$ converges to $(0, 0)$ as $t \rightarrow +\infty$.

The situation changes drastically when $M: \mathcal{H} \rightarrow \mathcal{H}$ is a ρ -cocoercive operator for some $\rho > 0$, that is

$$\langle M(x) - M(y), x - y \rangle \geq \rho \|M(x) - M(y)\|^2 \quad \forall x, y \in \mathcal{H}.$$

A ρ -cocoercive operator is obviously monotone and $\frac{1}{\rho}$ -Lipschitz continuous. In this case, the solution trajectory $t \mapsto x(t)$ of (1.2) converges weakly to a zero of M , and it holds that $\|\dot{x}(t)\| = \|M(x(t))\| = o\left(\frac{1}{\sqrt{t}}\right)$ as $t \rightarrow +\infty$. According to [3, Theorem 11], this results holds true even for the trajectory of a cocoercive operator flow with small perturbations. For completeness we provide the proof of the statement for the system without perturbations in Theorem A.1 in the Appendix ((see, also, [13]).

In the following, we will mention some typical examples of cocoercive operators, which obviously also serve as examples of monotone operators.

For every convex and continuously differentiable function $f: \mathcal{H} \rightarrow \mathbb{R}$ it holds according to the Baillon-Haddad Theorem (see [11, Corollary 18.17]) that its gradient $\nabla f: \mathcal{H} \rightarrow \mathcal{H}$ is $\frac{1}{L}$ -cocoercive if and only if it is L -Lipschitz continuous. Further, an operator $T: \mathcal{H} \rightarrow \mathcal{H}$ is nonexpansive, meaning it is 1-Lipschitz continuous, if and only if $\text{Id} - T$ is $\frac{1}{2}$ -cocoercive, where $\text{Id}: \mathcal{H} \rightarrow \mathcal{H}$ denotes the identity mapping on \mathcal{H} . This emphasizes the strong connection between fixed point problems and equations governed by cocoercive operators. Finally, we

recall that the *Yoshida approximation* ${}^\rho M: \mathcal{H} \rightarrow \mathcal{H}$ with parameter $\rho > 0$ of a set-valued maximally monotone operator $M: \mathcal{H} \rightrightarrows \mathcal{H}$ is the single-valued operator defined as

$${}^\rho M: \mathcal{H} \rightarrow \mathcal{H}, \quad {}^\rho M := \frac{1}{\rho} (\text{Id} - J_{\rho M}),$$

where $J_{\rho M} := (\text{Id} + \rho M)^{-1}: \mathcal{H} \rightarrow \mathcal{H}$ denotes the *resolvent* of M with parameter $\rho > 0$. The operator ${}^\rho M$ is ρ -cocoercive ([11, Corollary 23.11]) and it holds $\text{Zer}{}^\rho M = \text{Zer} M := \{x \in \mathcal{H} : 0 \in M(x)\}$. This emphasizes the strong connection between monotone inclusions and equations governed by cocoercive operators.

We want to note that even in the case of a cocoercive operator, one cannot generally expect the solution trajectory of (1.2) to strongly converge to a zero of this operator. A counterexample in this sense was provided in [2] for a dynamical system governed by the gradient of a convex and continuously differentiable function with a Lipschitz continuous gradient.

In spite of being ubiquitous in applications, we will not focus in this paper on continuous time models for (1.1) driven by cocoercive operators, but rather by monotone and continuous ones, which constitute a larger class of operators. Operators such as

$$M: \mathcal{H} \times \mathcal{G} \rightarrow \mathcal{H} \times \mathcal{G}, \quad M(x, y) := (\nabla_x \Phi(x, y), -\nabla_y \Phi(x, y)),$$

where \mathcal{G} is another real Hilbert space and $\Phi: \mathcal{H} \times \mathcal{G} \rightarrow \mathcal{R}$ is a convex-concave function with a continuous gradient, are monotone and continuous but generally not coercive, as illustrated by the example of a nontrivial bilinear operator. For such operators, (1.1) is equivalent to the problem of finding the saddle points of the minimax problem

$$\min_{x \in \mathcal{H}} \max_{y \in \mathcal{G}} \Phi(x, y),$$

which is of great relevance in game theory and in convex optimization in the context of linear equality constraints.

If the operator $M := \nabla f$ is the (not necessarily Lipschitz continuous) gradient of a convex and continuously differentiable function $f: \mathcal{H} \rightarrow \mathbb{R}$, then it is monotone and continuous and the solutions of (1.1) are precisely the optimal solutions of the optimization problem

$$\min_{x \in \mathcal{H}} f(x). \tag{1.3}$$

The dynamical system (1.2) becomes in this case the gradient flow

$$\dot{x}(t) + \nabla f(x(t)) = 0, \tag{1.4}$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$. Whenever $\arg \min f \neq \emptyset$, that is, the set of minimizers of f is not empty, the generated trajectory converges weakly to an element of $\arg \min f$ and $f(x(t)) - \min f = o(\frac{1}{t})$ as $t \rightarrow +\infty$, where $\min f$ denotes the minimal value of f .

Relying on the previous works [49, 5], the second-order continuous time model

$$\ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \beta \nabla^2 f(x(t)) \dot{x}(t) + \nabla f(x(t)) = 0, \tag{1.5}$$

with $\alpha \geq 3$ and $\beta \geq 0$, initial time $t_0 > 0$ and initial conditions $x(t_0) := x_0 \in \mathcal{H}$ and $\dot{x}(t_0) := \dot{x}_0 \in \mathcal{H}$, was introduced in [8] in the context of minimizing a convex and twice differentiable function $f: \mathcal{H} \rightarrow \mathbb{R}$ with $\arg \min f \neq \emptyset$. The primary aim was to accelerate the convergence of the function values along the generated solution trajectory. Indeed, if $\alpha \geq 3$ and $\beta \geq 0$, then $f(x(t)) - \min f = \mathcal{O}(\frac{1}{t^2})$ as $t \rightarrow +\infty$, while if $\beta > 0$, then $\int_{t_0}^{+\infty} t^2 \|\nabla f(x(t))\|^2 dt < +\infty$. If $\alpha > 3$, then the trajectory converges weakly to an element of $\arg \min f$ and $f(x(t)) - \min f = o(\frac{1}{t^2})$ as $t \rightarrow +\infty$.

Recently, in [14], in the context of a monotone equation such as (1.1), a second-order dynamical system has been proposed that combines a vanishing damping term with the time derivative of the operator along the trajectory, which can be seen as an analogous of the Hessian-driven damping term in (1.5). This model exhibits fast convergence rates of convergence in norm for the trajectory's velocity and the operator along the trajectory, outperforming those of the cocoercive operator flow. This phenomenon is analogous to the one described above for optimization, despite the fact that the operator does not have to originate from a potential.

The precise formulation of this so-called Fast Optimistic Gradient Descent Ascent (OGDA) dynamics is

$$\ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \beta(t) \frac{d}{dt} M(x(t)) + \frac{1}{2} \left(\dot{\beta}(t) + \frac{\alpha \beta(t)}{t} \right) M(x(t)) = 0, \tag{1.6}$$

with initial time $t_0 > 0$, $\alpha \geq 2$, $\beta : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ a continuous differentiable and nondecreasing function, and initial conditions $x(t_0) = x_0 \in \mathcal{H}$ and $\dot{x}(t_0) = \dot{x}_0 \in \mathcal{H}$.

It has been shown that if

$$\sup_{t \geq t_0} \frac{t\dot{\beta}(t)}{\beta(t)} \leq \alpha - 2,$$

then

$$\|\dot{x}(t)\| = \mathcal{O}\left(\frac{1}{t}\right) \quad \text{and} \quad \|M(x(t))\| = \mathcal{O}\left(\frac{1}{t\beta(t)}\right) \quad \text{as } t \rightarrow +\infty,$$

and if

$$\sup_{t \geq t_0} \frac{t\dot{\beta}(t)}{\beta(t)} < \alpha - 2,$$

then

$$\|\dot{x}(t)\| = o\left(\frac{1}{t}\right) \quad \text{and} \quad \|M(x(t))\| = o\left(\frac{1}{t\beta(t)}\right) \quad \text{as } t \rightarrow +\infty,$$

and $x(t)$ converges weakly to a zero of M as $t \rightarrow +\infty$.

Explicit and implicit numerical algorithms, introduced as discrete time counterparts of (1.6) and sharing its convergence properties have also been provided in [14, 18].

In this paper, we consider the Tikhonov regularization of the dynamical system (1.2)

$$\dot{x}(t) + M(x(t)) + \varepsilon(t)x(t) = 0, \tag{1.7}$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$ and continuously differentiable regularization function $\varepsilon : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ satisfying

$$\lim_{t \rightarrow +\infty} \varepsilon(t) = 0 \quad \text{and} \quad \int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty.$$

It is well-known ([6, 24]) that the Tikhonov regularization ensures that the trajectory generated by the system converges strongly to the element of minimum norm in $\text{Zer}M$. We will demonstrate that, in addition, rates of convergence in norm for the trajectory's velocity and the operator along the trajectory can be derived in terms of $\varepsilon(t)$. In some particular cases, these rates of convergence can be as fast as $\mathcal{O}(\frac{1}{t})$ as $t \rightarrow +\infty$. In this way, we emphasize a surprising acceleration feature of the Tikhonov regularization.

Further, by using time rescaling techniques, we translate the strong convergence and the convergence rate statements to the more general dynamic

$$\dot{x}(t) + \beta(t)M(x(t)) + \delta(t)(x(t) - v) = 0, \tag{1.8}$$

with initial time $t_0 \geq 0$, $\beta, \delta : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ continuously differentiable functions satisfying

$$\lim_{t \rightarrow +\infty} \frac{\delta(t)}{\beta(t)} = 0, \quad \int_{t_0}^{+\infty} \delta(t) dt = +\infty, \quad \text{and} \quad \int_{t_0}^{+\infty} \beta(t) dt = +\infty.$$

initial conditions $x(t_0) := x_0 \in \mathcal{H}$ and anchor point $v \in \mathcal{H}$.

This dynamic, referred to as the continuous model of the Halpern method [31], was studied by Suh, Park, and Ryu in [50] in finite-dimensional spaces, in the case $\beta(t) := 1$ and $\varepsilon(t) := \frac{\alpha}{t^q}$ for every $t \geq t_0 > 0$, where $\alpha > 0$ and $0 < q \leq 1$. By relying on a very intricate Lyapunov analysis, the convergence of the trajectory and convergence rate for the norm of the operator along the trajectory has been provided. These results turn out to be particular cases of our more general approach, which is fundamentally different. In addition, our approach provides a rate of convergence for the velocity. In light of the above considerations, it is evident that the Halpern method is nothing else than a particular instance of the classical Tikhonov regularization approach.

The benefit of the time rescaling technique ([3, 4]) is that there is no need to develop a Lyapunov analysis for every single system, as it allows retrieving the convergence statements from the previously known results of other related dynamics.

Finally, we show the connection between a particular first-order dynamical system with Tikhonov regularization and anchor point of the type (1.8) and the second order dynamical system

$$\ddot{x}(t) + \frac{\alpha}{t}\dot{x}(t) + \beta(t)\frac{d}{dt}M(x(t)) + \left(\dot{\beta}(t) + \frac{\beta(t)}{t}\right)M(x(t)) = 0, \tag{1.9}$$

with initial time $t_0 > 0$, $\alpha \geq 2$, $\beta : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ a twice continuous differentiable and function, and initial conditions $x(t_0) = x_0 \in \mathcal{H}$ and $\dot{x}(t_0) = \dot{x}_0 \in \mathcal{H}$.

The dynamics (1.9) inherits from the first order dynamical system with Tikhonov regularization the strong convergence property of the trajectory and the fast rates of convergence in norm for the trajectory's velocity

and the operator along the trajectory. In addition, it coincides with (1.6) for $\beta(t) = \beta_0 t^{\alpha-2}$ with $\alpha \geq 2$ and $\beta_0 > 0$, and thus reveals the strong convergence of the trajectory for Fast OGDA in a setting where, according to [14], nothing could be said about the behaviour of the trajectory.

In the case when $M = \text{Id} - T$, for $T : \mathcal{H} \rightarrow \mathcal{H}$ a nonexpansive (1-Lipschitz continuous) operator, the explicit discretization of (1.7) with anchor point $v \in \mathcal{H}$ (or of the dynamic (1.8) in case $\beta \equiv 1$) leads to the Halpern fixed point iteration [31]. By performing a convergence analysis which can be seen as the discrete counterpart of the one provided for the continuous time system, we derive two different regimes for the regularization sequence, both of which guarantee that the generated sequence of iterates strongly converges to the fixed point of T (or the zero of M) that is closest to v . In addition, we provide a general theoretical framework that allows to derive convergence rates for the vanishing of the discrete velocity and of the fixed point residual. Again, for some instances of the regularization sequence, we are able to derive concrete convergence rates that actually match those in continuous time. Extensions to fixed point problems governed by averaged operators are given, which allow us to improve some statements concerning the asymptotic properties of some algorithms from the literature ([15], [36]).

2 First-order dynamical system with Tikhonov regularization

The starting point of our study is a first-order monotone operator flow with Tikhonov regularization.

Let $t_0 \geq 0$ and $\varepsilon : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ be given. We consider on $[t_0, +\infty)$ the following dynamical system

$$\dot{x}(t) + M(x(t)) + \varepsilon(t)x(t) = 0, \quad (2.1)$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$.

For the *regularization function* ε , the following conditions are imposed. These conditions are known to be necessary for the strong convergence of the trajectory to the minimum norm solution, as discussed in [6, 24].

The regularization function $\varepsilon : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ is required to be continuously differentiable and to satisfy

$$\lim_{t \rightarrow +\infty} \varepsilon(t) = 0 \quad \text{and} \quad \int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty. \quad (2.2)$$

The first-order dynamical system is assumed to admit a unique strong global solution $x : [t_0, +\infty) \rightarrow \mathcal{H}$, meaning that it is absolutely continuous on every compact interval $[t_0, T]$ (see [20, Appendix]) and it satisfies (2.1) almost everywhere, with the property that $t \mapsto M(x(t))$ is absolutely continuous on every compact interval $[t_0, T]$.

Remark 2.1. (i) The first condition in (2.2) is essential and inherently expected, as we anticipate the trajectory to converge towards a zero of the operator M . If we were to assume that $\varepsilon(t) \equiv \varepsilon_0 > 0$, then the trajectory $x(t)$ would indeed exhibit strong convergence, albeit towards the unique zero of the strongly monotone operator $M + \varepsilon_0 \text{Id}$.

(ii) In case $M \equiv 0$, the equation

$$\dot{x}(t) + \varepsilon(t)x(t) = 0$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$, has as its unique solution

$$x(t) = x_0 \exp\left(-\int_{t_0}^t \varepsilon(u) du\right).$$

We observe that $x(t)$ converges to 0 as $t \rightarrow +\infty$, representing the element of $\text{Zer}M$ with the minimum norm, if and only if $\int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty$. This property holds irrespective of the initial data x_0 and underscores the indispensability of the second condition in (2.2).

(iii) If $M : \mathcal{H} \rightarrow \mathcal{H}$ is Lipschitz continuous, then (see, for instance, [32, Proposition 6.2.1], [48, Theorem 54]) (2.1) has for every $x_0 \in \mathcal{H}$ a unique strong global solution $x : [t_0, +\infty) \rightarrow \mathcal{H}$. In this case, $t \mapsto M(x(t))$ is absolutely continuous on every compact interval $[t_0, T]$ and therefore almost everywhere differentiable on $[t_0, +\infty)$.

2.1 Convergence analysis and rates of convergence

Following Remark 2.1 (ii), we define

$$\gamma: [t_0, +\infty) \rightarrow \mathbb{R}_{++}, \quad \gamma(t) := \exp\left(\int_{t_0}^t \varepsilon(u) du\right), \quad (2.3)$$

such that for every $t \geq t_0$ it holds $\dot{\gamma}(t) = \varepsilon(t) \gamma(t)$. This function will naturally appear several times in the analysis. The conditions (2.2) imposed on ε ensure that γ is strictly increasing, thus $\gamma(t) \geq \gamma(t_0) = 1$ for every $t \geq t_0$, and that $\lim_{t \rightarrow +\infty} \gamma(t) = +\infty$.

Given a trajectory solution $x: [t_0, +\infty) \rightarrow \mathcal{H}$ of (2.1) and $\xi \in \text{Zer}M$, we define the functions

$$\varphi_\xi: [t_0, +\infty) \rightarrow \mathbb{R}_+, \quad \varphi_\xi(t) := \frac{1}{2} \|x(t) - \xi\|^2, \quad (2.4)$$

and

$$\psi: [t_0, +\infty) \rightarrow \mathbb{R}_+, \quad \psi(t) := \frac{1}{2} \|M(x(t)) + \varepsilon(t)x(t)\|^2. \quad (2.5)$$

These two objects are pivotal in our convergence analysis. Let us begin by establishing some initial estimates for them. In some of these estimates, we will utilize the quantity

$$D_0(x_0, v, \xi) := \max\{\|x_0 - \xi\|, \|v - \xi\|\}, \quad (2.6)$$

which is defined for a triple $(x_0, v, \xi) \in \mathcal{H} \times \mathcal{H} \times \mathcal{H}$.

Lemma 2.2. *Let $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be a trajectory solution of the first-order dynamical system with Tikhonov regularization (2.1) and $\xi \in \text{Zer}M$. Then, the following statements are true:*

(i) *for almost every $t \geq t_0$, it holds*

$$\dot{\varphi}_\xi(t) + 2\varepsilon(t)\varphi_\xi(t) \leq -\varepsilon(t)\langle x(t) - \xi, \xi \rangle \leq \varepsilon(t)\varphi_\xi(t) + \frac{\varepsilon(t)}{2} \|\xi\|^2; \quad (2.7)$$

(ii) *the trajectory is bounded, in particular, for every $t \geq t_0$, it holds*

$$\|x(t) - \xi\| \leq D_0(x_0, 0, \xi) \quad \text{and} \quad \|x(t)\| \leq 2D_0(x_0, 0, \xi); \quad (2.8)$$

(iii) *for almost every $t \geq t_0$, it holds*

$$\dot{\psi}(t) + 2\varepsilon(t)\psi(t) \leq -\frac{1}{2}\dot{\varepsilon}(t)\left(\frac{d}{dt}\|x(t)\|^2\right) = -\dot{\varepsilon}(t)\langle \dot{x}(t), x(t) \rangle; \quad (2.9)$$

(iv) *for almost every $t \geq t_0$, it holds*

$$\frac{d}{dt}(\gamma^2(t)\psi(t)) = \gamma^2(t)\dot{\psi}(t) + 2\varepsilon(t)\gamma^2(t)\psi(t) \leq -\frac{1}{2}\dot{\varepsilon}(t)\gamma^2(t)\left(\frac{d}{dt}\|x(t)\|^2\right) = -\dot{\varepsilon}(t)\gamma^2(t)\langle \dot{x}(t), x(t) \rangle. \quad (2.10)$$

Proof. (i) For almost every $t \geq t_0$, when we take the time derivative of φ_ξ , we obtain

$$\begin{aligned} \dot{\varphi}_\xi(t) &= \langle x(t) - \xi, \dot{x}(t) \rangle = -\langle x(t) - \xi, M(x(t)) \rangle - \varepsilon(t)\langle x(t) - \xi, x(t) \rangle \\ &= -\langle x(t) - \xi, M(x(t)) \rangle - \varepsilon(t)\langle x(t) - \xi, \xi \rangle - 2\varepsilon(t)\varphi_\xi(t) \\ &\leq -\varepsilon(t)\langle x(t) - \xi, \xi \rangle - 2\varepsilon(t)\varphi_\xi(t), \end{aligned} \quad (2.11)$$

where in the last inequality we used that $\langle x(t) - \xi, M(x(t)) \rangle \geq 0$, which holds due to the monotonicity of M . The second inequality in (2.7) follows from

$$-\varepsilon(t)\langle x(t) - \xi, \xi \rangle = \frac{\varepsilon(t)}{2} \left(\|x(t) - \xi\|^2 + \|\xi\|^2 - \|x(t)\|^2 \right) \leq \varepsilon(t)\varphi_\xi(t) + \frac{\varepsilon(t)}{2} \|\xi\|^2.$$

(ii) By invoking (2.7), it yields for almost every $t \geq t_0$

$$\frac{d}{dt}(\gamma(t)\varphi_\xi(t)) = \gamma(t)\dot{\varphi}_\xi(t) + \varepsilon(t)\gamma(t)\varphi_\xi(t) \leq \frac{\varepsilon(t)\gamma(t)}{2} \|\xi\|^2 = \dot{\gamma}(t) \frac{\|\xi\|^2}{2}.$$

After integration from t_0 to t , we obtain for every $t \geq t_0$

$$\gamma(t) \varphi_\xi(t) - \gamma(t_0) \varphi_\xi(t_0) \leq (\gamma(t) - \gamma(t_0)) \frac{\|\xi\|^2}{2},$$

which is equivalent to

$$\|x(t) - \xi\|^2 \leq \frac{1}{\gamma(t)} \|x_0 - \xi\|^2 + \left(1 - \frac{1}{\gamma(t)}\right) \|\xi\|^2.$$

The first statement in (2.8) is a direct consequence, as $\gamma(t) \geq \gamma(t_0) = 1$ for every $t \geq t_0$. The second statement in (2.8) follows by the triangle inequality.

(iii) Differentiating the function ψ with respect to t gives for almost every $t \geq t_0$

$$\begin{aligned} \dot{\psi}(t) &= \left\langle M(x(t)) + \varepsilon(t)x(t), \frac{d}{dt}(M(x(t)) + \varepsilon(t)x(t)) \right\rangle \\ &= - \left\langle \dot{x}(t), \frac{d}{dt}M(x(t)) + \dot{\varepsilon}(t)x(t) + \varepsilon(t)\dot{x}(t) \right\rangle \\ &= - \left\langle \dot{x}(t), \frac{d}{dt}M(x(t)) \right\rangle - \dot{\varepsilon}(t)\langle \dot{x}(t), x(t) \rangle - \varepsilon(t)\|\dot{x}(t)\|^2 \\ &\leq -\dot{\varepsilon}(t)\langle \dot{x}(t), x(t) \rangle - \varepsilon(t)\|\dot{x}(t)\|^2 \\ &= -\frac{1}{2}\dot{\varepsilon}(t) \left(\frac{d}{dt}\|x(t)\|^2 \right) - 2\varepsilon(t)\psi(t), \end{aligned} \tag{2.12}$$

where inequality (2.12) is a consequence of the monotonicity of M .

(iv) The statement follows from (2.9), by using that $\dot{\gamma}(t) = \gamma(t)\varepsilon(t)$ for every $t \geq t_0$. \square

Proposition 2.3. *Let $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the first-order dynamical system with Tikhonov regularization (2.1). If $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, then $x(t)$ converges strongly to the minimum norm solution $\xi_* := \text{P}_{\text{Zer}M}(0)$ as $t \rightarrow +\infty$.*

Proof. Invoking inequality (2.7) for $\xi_* := \text{P}_{\text{Zer}M}(0)$, we have for almost every $t \geq t_0$

$$\dot{\varphi}_{\xi_*}(t) + \varepsilon(t) \varphi_{\xi_*}(t) \leq \varepsilon(t) \langle x(t) - \xi_*, 0 - \xi_* \rangle.$$

By using Lemma A.5 and the boundedness of $x(\cdot)$, from here we obtain that

$$\limsup_{t \rightarrow +\infty} \varphi_{\xi_*}(t) \leq \limsup_{t \rightarrow +\infty} \langle x(t) - \xi_*, 0 - \xi_* \rangle. \tag{2.13}$$

We are going to show that, under the additional assumption $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, every weak cluster point of $x(\cdot)$ belongs to $\text{Zer}M$. Such cluster point always exists thanks to the boundedness of the trajectory.

If \hat{x} is arbitrary weak cluster point of $x(\cdot)$, then there exists a sequence $\{x(t_k)\}_{k \geq 0}$ such that $x(t_k)$ converges weakly to \hat{x} as $k \rightarrow +\infty$. On the other hand, $M(x(t_k))$ converges strongly to 0 as $k \rightarrow +\infty$. Since M is monotone and continuous, it is maximally monotone (see, for instance, [11, Corollary 20.28]). Therefore, its graph is sequentially closed in the weak \times strong topology of $\mathcal{H} \times \mathcal{H}$ (see [11, Proposition 20.38]). This means that $M(\hat{x}) = 0$.

This enables us to conclude from (2.13), by using the variational characterization of the projection, that $\limsup_{t \rightarrow +\infty} \varphi_{\xi_*}(t) \leq 0$, which is nothing else than $x(t) \rightarrow \xi_*$ as $t \rightarrow +\infty$. \square

According to the previous result, the condition (2.2) required for the regularization function, in combination with the assumption that $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, suffice to ensure the strong convergence of the trajectory $x(\cdot)$ towards the minimum norm solution of (1.1). In contrast, the following proposition demonstrates that assuming convergence rather than divergence for the integral of the regularization function results in the weak convergence of the trajectory $x(\cdot)$ to a solution of (1.1).

Proposition 2.4. *Let $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the first-order dynamical system with Tikhonov regularization (2.1). Regarding the regularization function, instead of (2.2), suppose that*

$$\lim_{t \rightarrow +\infty} \varepsilon(t) = 0 \quad \text{and} \quad \int_{t_0}^{+\infty} \varepsilon(t) dt < +\infty.$$

If $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, then $x(t)$ converges weakly to an element of $\text{Zer}M$ as $t \rightarrow +\infty$.

Proof. Observing that statement (i) in Lemma 2.2 holds regardless of condition (2.2), for every $\xi_* \in \text{Zer}M$ we have

$$\dot{\varphi}_{\xi_*}(t) \leq \frac{\varepsilon(t)}{2} \|\xi\|^2 \text{ for almost every } t \geq t_0.$$

In particular, according to Lemma A.2, the limit $\lim_{t \rightarrow +\infty} \varphi_{\xi_*}(t) \in \mathbb{R}$ exists. This yields that the first condition in Opial's Lemma (see Lemma A.3) is satisfied.

The demonstration that the second condition in Opial's Lemma is satisfied under the additional assumption $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$ has been presented in the proof of Proposition 2.3.

The trajectory $x(\cdot)$ converging weakly to a solution of (1.1) follows as a consequence of Lemma A.3. \square

In the following theorem, we present additional conditions on the regularization function that ensure $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, leading to the strong convergence of $x(\cdot)$ to the solution of (1.1) with minimal norm.

Theorem 2.5. *Let $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the first-order dynamical system with Tikhonov regularization (2.1). Regarding the regularization function, assume that, in addition to (2.2), it satisfies either*

$$\lim_{t \rightarrow +\infty} \frac{|\dot{\varepsilon}(t)|}{\varepsilon(t)} = 0. \quad (2.14)$$

or

$$\int_{t_0}^{+\infty} |\dot{\varepsilon}(t)| dt < +\infty. \quad (2.15)$$

Then, $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, and therefore the trajectory $x(t)$ converges strongly to the minimum norm solution $\xi_* := \text{P}_{\text{Zer}M}(0)$ as $t \rightarrow +\infty$.

Proof. According to Proposition 2.3, it is sufficient to show that each of the two conditions (2.14) and (2.15) leads to $\psi(t) \rightarrow 0$ as $t \rightarrow +\infty$, which will imply that $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$.

Let $\xi \in \text{Zer}M$. In the first scenario, we assume that (2.14) holds. Then there exists $t'_0 \geq t_0$ such that $t \mapsto \frac{|\dot{\varepsilon}(t)|}{\varepsilon(t)}$ is bounded on $[t'_0, +\infty)$. According to (2.9) and by using twice the Cauchy-Schwarz inequality and then Lemma 2.2 (ii), we have for almost every $t \geq t'_0$

$$\begin{aligned} \dot{\psi}(t) + \varepsilon(t)\psi(t) &\leq -\varepsilon(t)\psi(t) + |\dot{\varepsilon}(t)| \|\dot{x}(t)\| \|x(t)\| = -\frac{1}{2}\varepsilon(t)\|\dot{x}(t)\|^2 + |\dot{\varepsilon}(t)| \|\dot{x}(t)\| \|x(t)\| \\ &\leq \frac{1}{2} \frac{|\dot{\varepsilon}(t)|^2}{\varepsilon(t)} \|x(t)\|^2 \leq 2D_0(x_0, 0, \xi)^2 \frac{|\dot{\varepsilon}(t)|^2}{\varepsilon(t)} = \varepsilon(t) \left(2D_0(x_0, 0, \xi)^2 \frac{|\dot{\varepsilon}(t)|^2}{\varepsilon(t)^2} \right). \end{aligned}$$

By using Lemma A.5, we obtain $\lim_{t \rightarrow +\infty} \psi(t) = 0$.

In the second scenario, we assume that (2.15) holds. Using (2.10) and the Cauchy-Schwarz inequality, we obtain for almost every $t \geq t_0$

$$\begin{aligned} \frac{d}{dt}(\gamma^2(t)\psi(t)) &\leq -\dot{\varepsilon}(t)\gamma^2(t)\langle \dot{x}(t), x(t) \rangle \leq \frac{1}{2}|\dot{\varepsilon}(t)|\gamma^2(t)\|x(t)\|^2 + \frac{1}{2}|\dot{\varepsilon}(t)|\gamma^2(t)\|\dot{x}(t)\|^2 \\ &\leq 2D_0(x_0, 0, \xi)^2 |\dot{\varepsilon}(t)|\gamma^2(t) + |\dot{\varepsilon}(t)|\gamma^2(t)\psi(t). \end{aligned}$$

Let $s \geq t_0$. By integration, it yields for every $t \geq s$

$$\gamma^2(t)\psi(t) \leq \gamma^2(s)\psi(s) + 2D_0(x_0, 0, \xi)^2 \int_s^t |\dot{\varepsilon}(u)|\gamma^2(u) du + \int_s^t |\dot{\varepsilon}(u)|\gamma^2(u)\psi(u) du.$$

The function $t \mapsto \gamma^2(s)\psi(s) + 2D_0(x_0, 0, \xi)^2 \int_s^t |\dot{\varepsilon}(u)|\gamma^2(u) du$ is nondecreasing, therefore, according to Gronwall's Lemma (see Lemma A.6), we obtain for every $t \geq s$

$$\gamma^2(t)\psi(t) \leq \left(\gamma^2(s)\psi(s) + 2D_0(x_0, 0, \xi)^2 \int_s^t |\dot{\varepsilon}(u)|\gamma^2(u) du \right) \exp\left(\int_s^t |\dot{\varepsilon}(u)| du \right),$$

which, by using that γ is nondecreasing, implies that for every $t \geq s$

$$\begin{aligned} \psi(t) &\leq \left(\frac{\gamma^2(s)\psi(s)}{\gamma^2(t)} + 2D_0(x_0, 0, \xi)^2 \int_s^t |\dot{\varepsilon}(u)| \frac{\gamma^2(u)}{\gamma^2(t)} du \right) \exp\left(\int_s^t |\dot{\varepsilon}(u)| du \right) \\ &\leq \left(\frac{\gamma^2(s)\psi(s)}{\gamma^2(t)} + 2D_0(x_0, 0, \xi)^2 \int_s^t |\dot{\varepsilon}(u)| du \right) \exp\left(\int_s^t |\dot{\varepsilon}(u)| du \right). \end{aligned}$$

From here it yields that for every $s \geq t_0$

$$\limsup_{t \rightarrow +\infty} \psi(t) \leq 2D_0(x_0, 0, \xi)^2 \left(\int_s^{+\infty} |\dot{\varepsilon}(u)| du \right) \exp \left(\int_s^{+\infty} |\dot{\varepsilon}(u)| du \right). \quad (2.16)$$

Finally, letting $s \rightarrow +\infty$ in (2.16), we obtain $\lim_{t \rightarrow +\infty} \psi(t) = 0$.

To summarize, each condition (2.14) and (2.15) ensures that $\|M(x(t)) + \varepsilon(t)x(t)\| \rightarrow 0$ as $t \rightarrow +\infty$. Given the boundedness of the trajectory, this immediately implies $\|M(x(t))\| \rightarrow 0$ as $t \rightarrow +\infty$, thus concluding the proof. \square

Remark 2.6. (i) The condition (2.15) was introduced by Cominetti, Peypouquet, and Sorin in [24, Theorem 9]. The technique employed in proving $\lim_{t \rightarrow +\infty} \psi(t) = 0$ aligns with the approach used in this paper.

(ii) In [35] (also see [24, Proposition 5]), the strong convergence of the trajectory to the minimal norm solution has been demonstrated under assumptions regarding the regularization function which, in addition to (2.2), entail that $t \mapsto \varepsilon(t)$ is decreasing and satisfy the condition

$$\lim_{t \rightarrow +\infty} \frac{|\dot{\varepsilon}(t)|}{\varepsilon^2(t)} = 0. \quad (2.17)$$

The condition (2.14) in Theorem 2.5 represents a notable enhancement within this context. Specifically, it accommodates regularization functions of the form $\varepsilon(t) = \frac{\alpha}{t^q}$, where $0 < q \leq 1$ and $\alpha > 0$. This condition allows for the critical choice $q := 1$, for which (2.17) is not fulfilled.

While the previous theorem ensures the convergence of $\|M(x(t))\|$ to zero as $t \rightarrow +\infty$, it does not provide information about the speed of this convergence process or how it could be obtained. In the following result, we will introduce general conditions concerning the regularization function that enables the derivation of convergence rates for the operator norm and the velocity.

To this end, we define for $t_1 \geq t_0$ and a triple $(x_0, v, \xi) \in \mathcal{H} \times \mathcal{H} \times \mathcal{H}$ the following quantity

$$D_1(x_0, v, \xi, t_1) := \sqrt{\|M(x(t_1)) + \varepsilon(t_1)(x(t_1) - v)\|^2 + 8D_0^2(x_0, v, \xi)|\dot{\varepsilon}(t_1)|}. \quad (2.18)$$

Theorem 2.7. *Let $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the first-order dynamical system with Tikhonov regularization (2.1) and $\xi \in \text{Zer}M$. Then, the following statements are true:*

(i) *if there exists $t_1 \geq t_0$ such that*

$$t \mapsto \varepsilon(t) \text{ is twice differentiable and } t \mapsto \frac{d}{dt} (\dot{\varepsilon}(t) + \varepsilon^2(t)) \text{ preserves constant sign on } [t_1, +\infty), \quad (2.19)$$

then for every $t \geq t_1$ it holds

$$\begin{aligned} \psi(t) + \frac{\dot{\varepsilon}(t)}{2} \|x(t)\|^2 &\leq \frac{\gamma^2(t_1)}{2\gamma^2(t)} \left(\|M(x(t_1)) + \varepsilon(t_1)x(t_1)\|^2 + \dot{\varepsilon}(t_1) \|x(t_1)\|^2 \right) \\ &\quad + 2D_0(x_0, 0, \xi)^2 \left[\dot{\varepsilon}(t) - \frac{1}{\gamma^2(t)} \dot{\varepsilon}(t_1) \gamma^2(t_1) \right]_+, \end{aligned} \quad (2.20)$$

where $[r]_+ := \max\{r, 0\}$;

(ii) *if, in addition,*

$$t \mapsto \varepsilon(t) \text{ is nonincreasing on } [t_1, +\infty), \quad (2.21)$$

then for almost every $t \geq t_1$ it holds

$$\|\dot{x}(t)\| \leq D_1(x(t_1), 0, \xi, t_1) \max \left\{ \frac{\sqrt{|\dot{\varepsilon}(t)|}}{\sqrt{|\dot{\varepsilon}(t_1)|}}, \frac{\gamma(t_1)}{\gamma(t)} \right\}, \quad (2.22)$$

and for every $t \geq t_1$ it holds

$$\|M(x(t))\| \leq D_1(x(t_1), 0, \xi, t_1) \max \left\{ \frac{\sqrt{|\dot{\varepsilon}(t)|}}{\sqrt{|\dot{\varepsilon}(t_1)|}}, \frac{\gamma(t_1)}{\gamma(t)} \right\} + 2D_0(x_0, 0, \xi) \varepsilon(t). \quad (2.23)$$

Proof. For every $t > t_1 \geq t_0$, by integrating (2.10) from t_1 to t , we obtain

$$\begin{aligned} \gamma^2(t) \psi(t) - \gamma^2(t_1) \psi(t_1) &\leq -\frac{1}{2} \int_{t_1}^t \dot{\varepsilon}(u) \gamma^2(u) \left(\frac{d}{du} \|x(u)\|^2 \right) du \\ &= -\frac{1}{2} \dot{\varepsilon}(u) \gamma^2(u) \|x(u)\|^2 \Big|_{u=t_1}^{u=t} + \frac{1}{2} \int_{t_1}^t \frac{d}{du} (\dot{\varepsilon}(u) \gamma^2(u)) \|x(u)\|^2 du \end{aligned}$$

or, equivalently,

$$\psi(t) + \frac{\dot{\varepsilon}(t)}{2} \|x(t)\|^2 \leq \frac{\gamma^2(t_1)}{2\gamma^2(t)} \left(2\psi(t_1) + \dot{\varepsilon}(t_1) \|x(t_1)\|^2 \right) + \frac{1}{2\gamma^2(t)} \int_{t_1}^t \frac{d}{du} (\dot{\varepsilon}(u) \gamma^2(u)) \|x(u)\|^2 du. \quad (2.24)$$

Then, observe that for every $t \geq t_1$

$$\frac{d}{dt} (\dot{\varepsilon}(t) \gamma^2(t)) = \ddot{\varepsilon}(t) \gamma^2(t) + 2\dot{\varepsilon}(t) \gamma(t) \dot{\gamma}(t) = (\ddot{\varepsilon}(t) + 2\dot{\varepsilon}(t) \varepsilon(t)) \gamma^2(t) = \frac{d}{dt} (\dot{\varepsilon}(t) + \varepsilon^2(t)) \gamma^2(t), \quad (2.25)$$

which means $\frac{d}{dt} (\dot{\varepsilon}(t) \gamma^2(t))$ and $\frac{d}{dt} (\dot{\varepsilon}(t) + \varepsilon^2(t))$ share the same sign on $[t_1, +\infty)$.

The case $\frac{d}{dt} (\dot{\varepsilon}(t) + \varepsilon^2(t))$ is nonnegative on $[t_1, +\infty)$. Using Lemma 2.2 (ii), we deduce that for every $t \geq t_1$ it holds

$$\begin{aligned} 0 &\leq \int_{t_1}^t \frac{d}{du} (\dot{\varepsilon}(u) \gamma^2(u)) \|x(u)\|^2 du \leq 4D_0(x_0, 0, \xi)^2 \int_{t_1}^t \frac{d}{du} (\dot{\varepsilon}(u) \gamma^2(u)) du \\ &= 4D_0(x_0, 0, \xi)^2 (\dot{\varepsilon}(t) \gamma^2(t) - \dot{\varepsilon}(t_1) \gamma^2(t_1)). \end{aligned}$$

Plugging this inequality into (2.24), we obtain for every $t \geq t_1$

$$\psi(t) + \frac{\dot{\varepsilon}(t)}{2} \|x(t)\|^2 \leq \frac{\gamma^2(t_1)}{2\gamma^2(t)} \left(2\psi(t_1) + \dot{\varepsilon}(t_1) \|x(t_1)\|^2 \right) + 2D_0(x_0, 0, \xi)^2 \left[\dot{\varepsilon}(t) - \frac{1}{\gamma^2(t)} \dot{\varepsilon}(t_1) \gamma^2(t_1) \right]. \quad (2.26)$$

If, in addition, ε is nonincreasing on $[t_1, +\infty)$, then, invoking Lemma 2.2 (ii), we obtain for every $t \geq t_1$

$$\begin{aligned} \|M(x(t)) + \varepsilon(t)x(t)\|^2 &\leq \frac{\gamma^2(t_1)}{\gamma^2(t)} \left(\|M(x(t_1)) + \varepsilon(t_1)x(t_1)\|^2 + \dot{\varepsilon}(t_1) \|x(t_1)\|^2 - 4D_0(x_0, 0, \xi)^2 \dot{\varepsilon}(t_1) \right) \\ &\leq \frac{\gamma^2(t_1)}{\gamma^2(t)} D_1(x(t_1), 0, \xi, t_1)^2. \end{aligned}$$

The case $\frac{d}{dt} (\dot{\varepsilon}(t) + \varepsilon^2(t))$ is nonpositive on $[t_1, +\infty)$. We have for every $t \geq t_1$

$$\int_{t_1}^t \frac{d}{du} (\dot{\varepsilon}(u) \gamma^2(u)) \|x(u)\|^2 du \leq 0$$

and further, due to (2.24),

$$\psi(t) + \frac{\dot{\varepsilon}(t)}{2} \|x(t)\|^2 \leq \frac{\gamma^2(t_1)}{2\gamma^2(t)} \left(2\psi(t_1) + \dot{\varepsilon}(t_1) \|x(t_1)\|^2 \right). \quad (2.27)$$

The combination of (2.26) and (2.27) results in (2.20).

If, in addition, ε is nonincreasing on $[t_1, +\infty)$, then (2.25) yields for every $t \geq t_1$

$$\frac{1}{\gamma^2(t)} \leq \frac{1}{-\dot{\varepsilon}(t_1) \gamma^2(t_1)} (-\dot{\varepsilon}(t)) = \frac{|\dot{\varepsilon}(t)|}{|\dot{\varepsilon}(t_1)| \gamma^2(t_1)}$$

and therefore

$$\begin{aligned} \|M(x(t)) + \varepsilon(t)x(t)\|^2 &\leq -\dot{\varepsilon}(t) \|x(t)\|^2 + \frac{\gamma^2(t_1)}{\gamma^2(t)} \left(\|M(x(t_1)) + \varepsilon(t_1)x(t_1)\|^2 + \dot{\varepsilon}(t_1) \|x(t_1)\|^2 \right) \\ &\leq -\dot{\varepsilon}(t) 4D_0(x_0, 0, \xi)^2 + \frac{|\dot{\varepsilon}(t)|}{|\dot{\varepsilon}(t_1)|} \left(\|M(x(t_1)) + \varepsilon(t_1)x(t_1)\|^2 + 4D_0^2(x_0, v, \xi) |\dot{\varepsilon}(t_1)| \right) \\ &= \frac{|\dot{\varepsilon}(t)|}{|\dot{\varepsilon}(t_1)|} D_1(x(t_1), 0, \xi, t_1)^2. \end{aligned}$$

By combining the estimates we derived for $\|M(x(t)) + \varepsilon(t)x(t)\|$ in case where ε is nonincreasing, for every $t \geq t_1$, we obtain

$$\|M(x(t)) + \varepsilon(t)x(t)\| \leq \gamma(t_1) D_1(x(t_1), 0, \xi, t_1) \max \left\{ \frac{1}{\sqrt{|\dot{\varepsilon}(t_1)|} \gamma(t_1)} \sqrt{|\dot{\varepsilon}(t)|}, \frac{1}{\gamma(t)} \right\},$$

leading to the conclusion. \square

As an application of the theoretical considerations above, we consider dynamics generated by (2.1) for regularization functions of the form

$$\varepsilon: [t_0, +\infty) \rightarrow \mathbb{R}, \quad \varepsilon(t) := \frac{\alpha}{t^q}, \quad \text{with } t_0 > 0, 0 < q \leq 1 \text{ and } \alpha > 0.$$

For every $0 < q \leq 1$ and $\alpha > 0$, the regularization function ε evidently satisfies (2.2). Furthermore, it meets both conditions (2.14) and (2.15), ensuring the strong convergence of the trajectory $x(t)$ toward the minimum norm solution as $t \rightarrow +\infty$.

For every $t \geq t_0$ it holds $\dot{\varepsilon}(t) = -\frac{q\alpha}{t^{q+1}}$ and $\ddot{\varepsilon}(t) = \frac{q(q+1)\alpha}{t^{q+2}}$, therefore

$$\ddot{\varepsilon}(t) + 2\varepsilon(t)\dot{\varepsilon}(t) = \frac{q(q+1)\alpha}{t^{q+2}} - \frac{2q\alpha^2}{t^{2q+1}} = \frac{q\alpha}{t^{2q+1}} \left(\frac{q+1}{t^{1-q}} - 2\alpha \right). \quad (2.28)$$

In case $0 < q < 1$, there exists $t_1 := \max \left\{ t_0, \left(\frac{q+1}{2\alpha} \right)^{\frac{1}{1-q}} \right\}$ such that $\frac{d}{dt}(\dot{\varepsilon}(t) + \varepsilon^2(t)) \leq 0$ for every $t \geq t_1$.

According to (2.25), $\frac{\sqrt{|\dot{\varepsilon}(t)|}}{\sqrt{|\dot{\varepsilon}(t_1)|}} \geq \frac{\gamma(t_1)}{\gamma(t)}$ for every $t \geq t_1$. For every $t \geq t_0$ it holds

$$\gamma(t) = \exp \left(\int_{t_0}^t \frac{\alpha}{s^q} ds \right) = \exp \left(\frac{\alpha}{1-q} \left(t^{1-q} - t_0^{1-q} \right) \right),$$

and Theorem 2.7 and Theorem 2.5 lead to the following result.

Theorem 2.8 (the case $\varepsilon(t) := \frac{\alpha}{t^q}, 0 < q < 1$). *Let $t_0 > 0, 0 < q < 1, \alpha > 0$, and $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the dynamical system*

$$\dot{x}(t) + M(x(t)) + \frac{\alpha}{t^q}x(t) = 0, \quad (2.29)$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$, and $\xi \in \text{Zer}M$. Then, the following statements are true:

(i) for almost every $t \geq t_1 := \max \left\{ t_0, \left(\frac{q+1}{2\alpha} \right)^{\frac{1}{1-q}} \right\}$ it holds

$$\|\dot{x}(t)\| \leq D_1(x(t_1), 0, \xi, t_1) \left(\frac{t_1}{t} \right)^{\frac{q+1}{2}},$$

and for every $t \geq t_1 := \max \left\{ t_0, \left(\frac{q+1}{2\alpha} \right)^{\frac{1}{1-q}} \right\}$ it holds

$$\|M(x(t))\| \leq D_1(x(t_1), 0, \xi, t_1) \left(\frac{t_1}{t} \right)^{\frac{q+1}{2}} + \frac{2\alpha D_0(x_0, 0, \xi)}{t^q};$$

(ii) the trajectory $x(t)$ converges strongly to the minimum norm solution $\xi_* := P_{\text{Zer}M}(0)$ as $t \rightarrow +\infty$.

In case $q = 1$, if $\alpha \geq 1$, then $\frac{d}{dt}(\dot{\varepsilon}(t) + \varepsilon^2(t)) \leq 0$ for every $t \geq t_0$, and, if $\alpha < 1$, then $\frac{d}{dt}(\dot{\varepsilon}(t) + \varepsilon^2(t)) > 0$ for every $t \geq t_0$. For every $t \geq t_0$ it holds

$$\gamma(t) = \exp \left(\int_{t_0}^t \frac{\alpha}{t} dt \right) = \exp \left(\log \left(\left(\frac{t}{t_0} \right)^\alpha \right) \right) = \left(\frac{t}{t_0} \right)^\alpha,$$

and Theorem 2.7 and Theorem 2.5 lead to the following result.

Theorem 2.9 (the case $\varepsilon(t) := \frac{\alpha}{t}$). *Let $t_0 > 0, \alpha > 0$, and $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the dynamical system*

$$\dot{x}(t) + M(x(t)) + \frac{\alpha}{t}x(t) = 0, \quad (2.30)$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$, and $\xi \in \text{Zer}M$. Then, the following statements are true:

(i) for almost every $t \geq t_0$ it holds

$$\|\dot{x}(t)\| \leq D_1(x_0, 0, \xi, t_0) \left(\frac{t_0}{t} \right)^{\min\{\alpha, 1\}},$$

and for every $t \geq t_0$ it holds

$$\|M(x(t))\| \leq D_1(x_0, 0, \xi, t_0) \left(\frac{t_0}{t} \right)^{\min\{\alpha, 1\}} + \frac{2\alpha D_0(x_0, 0, \xi)}{t};$$

(ii) the trajectory $x(t)$ converges strongly to the minimum norm solution $\xi_* := P_{\text{Zer}M}(0)$ as $t \rightarrow +\infty$.

The above theorems suggest that the best rates of convergence occur when they are $\mathcal{O}(\frac{1}{t})$ as $t \rightarrow +\infty$, achieved when $q = 1$ and $\alpha \geq 1$. This represents a clear improvement concerning the convergence rates compared to those of the unregularized dynamical system, even in cases where the operator is cocoercive (see Theorem A.1). Should one aim to enhance the rates of convergence for both the velocity and the operator norm by opting for $q > 1$, it is important to note, as highlighted in Remark 2.1 (ii), that strong convergence of the trajectory to the minimum norm solution cannot be expected.

For $t \geq t_0 > 1$, we consider the regularization function $\varepsilon : [t_0, +\infty) \rightarrow \mathbb{R}, \varepsilon(t) = \frac{1}{t \log(t)}$. For every $t \geq t_0$ it holds

$$\dot{\varepsilon}(t) + \varepsilon^2(t) = \frac{-1}{t^2 \log(t)}, \quad \text{and} \quad \frac{d}{dt}(\dot{\varepsilon}(t) + \varepsilon^2(t)) = \frac{2}{t^3} > 0,$$

and $t \mapsto \varepsilon(t)$ is nonincreasing on $[t_0, +\infty)$. In addition, condition (2.14) as well as condition (2.15) is satisfied. According to (2.25), $\frac{\sqrt{|\dot{\varepsilon}(t)|}}{\sqrt{|\dot{\varepsilon}(t_0)|}} \leq \frac{\gamma(t_0)}{\gamma(t)}$ for every $t \geq t_0$. For every $t \geq t_0$ it holds

$$\gamma(t) = \exp\left(\int_{t_0}^t \frac{1}{s \log(s)} ds\right) = \exp(\log(\log(t)) - \log(\log(t_0))) = \frac{\log(t)}{\log(t_0)},$$

and Theorem 2.7 and Theorem 2.5 lead to the following result.

Theorem 2.10 (the case $\varepsilon(t) := \frac{1}{t \log(t)}$). *Let $t_0 > 1$ and $x : [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the dynamical system*

$$\dot{x}(t) + M(x(t)) + \frac{1}{t \log(t)} x(t) = 0, \quad (2.31)$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$, and $\xi \in \text{Zer}M$. Then, the following statements are true:

(i) for almost every $t \geq t_0$ it holds

$$\|\dot{x}(t)\| \leq D_1(x(t_0), 0, \xi, t_0) \frac{\log(t_0)}{\log(t)},$$

and for every $t \geq t_0$ it holds

$$\|M(x(t))\| \leq D_1(x(t_0), 0, \xi, t_0) \frac{\log(t_0)}{\log(t)} + \frac{2D_0(x_0, 0, \xi)}{t \log(t)};$$

(ii) the trajectory $x(t)$ converges strongly to the minimum norm solution $\xi_* := P_{\text{Zer}M}(0)$ as $t \rightarrow +\infty$.

We have seen that even if the regularization function converges to zero faster than $\frac{1}{t}$, while satisfying the standing assumptions (2.2), the rates of convergence for the velocity of the trajectory and the operator along the trajectory cannot be expected to be faster than $\mathcal{O}(\frac{1}{t})$ as $t \rightarrow +\infty$.

2.2 First-order dynamical system with Tikhonov regularization and time rescaling

In this subsection, we will extend the previous analysis to first-order dynamical systems that incorporate Tikhonov regularization and time rescaling, expressed through the presence of a time-dependent coefficient of the operator. With the introduction and analysis of this continuous dynamic, we take the first step in our attempt to demonstrate the close connection between first-order dynamical systems with Tikhonov regularization and second-order dynamical systems with a vanishing damping term associated with the monotone equation 1.1.

Let $s_0 \geq 0$ and $\beta, \delta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ be given functions. We consider on $[s_0, +\infty)$ the following dynamical system

$$\dot{z}(s) + \beta(s) M(z(s)) + \delta(s) z(s) = 0, \quad (2.32)$$

with initial condition $z(s_0) := z_0 \in \mathcal{H}$.

For the time rescaling function β and the regularization function δ , we consider the following assumptions.

The functions $\beta, \delta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ are required to be continuously differentiable and to satisfy

$$\lim_{s \rightarrow +\infty} \frac{\delta(s)}{\beta(s)} = 0, \quad \int_{s_0}^{+\infty} \delta(s) ds = +\infty, \quad \text{and} \quad \int_{s_0}^{+\infty} \beta(s) ds = +\infty. \quad (2.33)$$

The first-order dynamical system is assumed to admit a unique strong global solution $z : [s_0, +\infty) \rightarrow \mathcal{H}$ with the property that $s \mapsto M(x(s))$ is absolutely continuous on every compact interval $[s_0, S]$.

The following proposition will allow us to transfer the statements obtained for (2.1) in the previous section to (2.32).

Proposition 2.11. *Let $s_0, t_0 \geq 0$ and $\beta, \delta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}, \varepsilon : [t_0, +\infty) \rightarrow \mathbb{R}_{++}$ be continuously differentiable functions. Suppose that the time rescaling function satisfies*

$$\int_{s_0}^{+\infty} \beta(s) ds = +\infty.$$

Let $x : [t_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of

$$\dot{x}(t) + M(x(t)) + \varepsilon(t)x(t) = 0 \quad \text{with} \quad x(t_0) := z_0 \in \mathcal{H}, \quad (2.34)$$

and $z : [s_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of

$$\dot{z}(s) + \beta(s)M(z(s)) + \delta(s)z(s) = 0 \quad \text{with} \quad z(s_0) := z_0 \in \mathcal{H}. \quad (2.35)$$

Then the two trajectories are equivalent, subject to a time rescaling process. Precisely, there exist bijective functions $\tau : [s_0, +\infty) \rightarrow [t_0, +\infty)$ and $\sigma : [t_0, +\infty) \rightarrow [s_0, +\infty)$ such that $\sigma = \tau^{-1}$ and

- if $x(\cdot)$ is the trajectory solution of (2.34), then $z(s) := x(\tau(s))$ defines the trajectory solution of (2.35) with $\delta(s) := \varepsilon(\tau(s))\beta(s)$ for every $s \geq s_0$;
- if $z(\cdot)$ is the trajectory solution of (2.35), then $x(t) := z(\sigma(t))$ defines the trajectory solution of (2.34) with $\varepsilon(t) := \frac{\delta(\sigma(t))}{\beta(\sigma(t))}$ for every $t \geq t_0$.

Moreover,

$$\lim_{t \rightarrow +\infty} \varepsilon(t) = 0 \Leftrightarrow \lim_{s \rightarrow +\infty} \frac{\delta(s)}{\beta(s)} = 0 \quad \text{and} \quad \int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty \Leftrightarrow \int_{s_0}^{+\infty} \delta(s) ds = +\infty.$$

Proof. We define $\tau : [s_0, +\infty) \rightarrow [t_0, +\infty)$ for every $s \geq s_0$ as

$$\tau(s) = t_0 + \int_{s_0}^s \beta(u) du. \quad (2.36)$$

Since β is strictly positive and continuous, τ is a monotonically increasing and thus injective. Moreover, it holds $\tau(s_0) = t_0$ and $\lim_{s \rightarrow +\infty} \tau(s) = +\infty$, meaning that σ is also surjective and therefore, bijective. For clarity, we will denote its inverse τ^{-1} by $\sigma : [t_0, +\infty) \rightarrow [s_0, +\infty)$.

“ $x(t) \mapsto z(s)$ ”. Let $x(\cdot)$ be the trajectory solution of (2.34). We perform a change of time variable, setting $t := \tau(s)$ in the dynamics (2.34), and define for every $s \geq s_0$

$$z(s) := x(\tau(s)). \quad (2.37)$$

Then by the chain rule, we have for almost every $s \geq s_0$

$$\dot{z}(s) = \dot{x}(\tau(s)) \dot{\tau}(s) = \dot{x}(\tau(s)) \beta(s).$$

At the time $t = \tau(s)$, multiplying both sides of (2.34) by $\beta(s) > 0$ yields for almost every $s \geq s_0$

$$\dot{z}(s) + \beta(s)M(z(s)) + \varepsilon(\tau(s))\beta(s)z(s) = 0.$$

This is equivalent to (2.35) if for every $s \geq s_0$ we choose

$$\delta(s) := \varepsilon(\tau(s))\beta(s). \quad (2.38)$$

Suppose that $\lim_{t \rightarrow +\infty} \varepsilon(t) = 0$ and $\int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty$. It follows from (2.38) and the property that $\lim_{s \rightarrow +\infty} \tau(s) = +\infty$ that

$$\lim_{s \rightarrow +\infty} \frac{\delta(s)}{\beta(s)} = \lim_{s \rightarrow +\infty} \varepsilon(\tau(s)) = 0.$$

Furthermore, for every $s \geq s_0$, we have

$$\int_{s_0}^s \delta(u) du = \int_{s_0}^s \varepsilon(\tau(u))\beta(u) du = \int_{s_0}^s \varepsilon(\tau(u)) \dot{\tau}(u) du = \int_{\tau(s_0)}^{\tau(s)} \varepsilon(r) dr = \int_{t_0}^{\tau(s)} \varepsilon(r) dr \quad (2.39)$$

Taking the limit as s approaches $+\infty$, we obtain $\int_{s_0}^{+\infty} \delta(u) du = +\infty$.

" $x(t) \Leftarrow z(s)$ ". Let $z(\cdot)$ be the trajectory solution of (2.35). We substitute $s := \sigma(t)$ in the dynamic (2.35). By definition, for every $t \geq t_0$ it holds

$$\tau(\sigma(t)) = t$$

and differentiation with respect to t gives

$$\dot{\tau}(\sigma(t)) \dot{\sigma}(t) = \beta(\sigma(t)) \dot{\sigma}(t) = 1 \Leftrightarrow \dot{\sigma}(t) = \frac{1}{\beta(\sigma(t))}. \quad (2.40)$$

Therefore, if we set for every $t \geq t_0$

$$x(t) := z(\sigma(t)),$$

by the chain rule it yields for almost every $t \geq t_0$

$$\dot{x}(t) = \dot{z}(\sigma(t)) \dot{\sigma}(t) = \dot{z}(\sigma(t)) \cdot \frac{1}{\beta(\sigma(t))}.$$

Consequently, at the time $s = \sigma(t)$, dividing both sides of equation (2.35) by $\beta(\sigma(t)) > 0$, we deduce that for almost every $t \geq t_0$

$$\dot{x}(t) + M(x(t)) + \frac{\delta(\sigma(t))}{\beta(\sigma(t))} x(t) = 0. \quad (2.41)$$

By setting for every $t \geq t_0$

$$\varepsilon(t) := \frac{\delta(\sigma(t))}{\beta(\sigma(t))},$$

this is nothing else than (2.34).

Suppose that $\lim_{s \rightarrow +\infty} \frac{\delta(s)}{\beta(s)} = 0$ and $\int_{s_0}^{+\infty} \delta(s) ds = +\infty$. By following previous arguments, we have

$$\lim_{t \rightarrow +\infty} \varepsilon(t) = \lim_{t \rightarrow +\infty} \frac{\delta(\sigma(t))}{\beta(\sigma(t))} = 0.$$

Furthermore, for every $s \geq s_0$ it holds

$$\int_{t_0}^t \varepsilon(r) dr = \int_{t_0}^t \frac{\delta(\sigma(r))}{\beta(\sigma(r))} dr = \int_{t_0}^t \delta(\sigma(r)) \dot{\sigma}(r) dr = \int_{\sigma(t_0)}^{\sigma(t)} \delta(u) du = \int_{s_0}^{\sigma(t)} \delta(u) du,$$

thus, by taking the limit as t approaches $+\infty$, we conclude $\int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty$. \square

Proposition 2.11 allows us to transfer the convergence and convergence rate results stated in Theorem 2.5 and Theorem 2.7, respectively, to the dynamical system (2.32).

Theorem 2.12. *Let $z: [s_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the dynamical system (2.32). Regarding β and δ , assume that, in addition to (2.33), they satisfy either*

$$\lim_{s \rightarrow +\infty} \frac{1}{\beta(s)} \left| \frac{\dot{\delta}(s)}{\delta(s)} - \frac{\dot{\beta}(s)}{\beta(s)} \right| = 0. \quad (2.42)$$

or

$$\int_{s_0}^{+\infty} \left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right| ds < +\infty. \quad (2.43)$$

Then, the trajectory $z(s)$ converges strongly to the minimum norm solution $\xi_* := \mathbf{P}_{\text{Zer}M}(0)$ as $s \rightarrow +\infty$.

Proof. Given the trajectory solution $z(\cdot)$ of (2.32) and

$$\tau(s) = t_0 + \int_{s_0}^s \beta(u) du \quad \forall s \geq s_0,$$

where $t_0 \geq 0$, then, according to Proposition 2.11, $x(\cdot) := z(\tau^{-1}(\cdot))$ is the trajectory solution of (2.1) with

$$\varepsilon(t) := \frac{\delta(\tau^{-1}(t))}{\beta(\tau^{-1}(t))} \quad \text{for every } t \geq t_0.$$

For every $t \geq t_0$, we have

$$\dot{\varepsilon}(t) = \frac{1}{\beta(\tau^{-1}(t))} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)),$$

and thus it holds

$$\frac{|\dot{\varepsilon}(t)|}{\varepsilon(t)} = \frac{1}{\beta(\tau^{-1}(t))} \left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) \right| \frac{\beta(\tau^{-1}(t))}{\delta(\tau^{-1}(t))} = \frac{1}{\beta(\tau^{-1}(t))} \left| \frac{\dot{\delta}(\tau^{-1}(t))}{\delta(\tau^{-1}(t))} - \frac{\dot{\beta}(\tau^{-1}(t))}{\beta(\tau^{-1}(t))} \right|.$$

Therefore, if (2.42) is satisfied, then

$$\lim_{t \rightarrow +\infty} \frac{|\dot{\varepsilon}(t)|}{\varepsilon(t)} = 0.$$

On the other hand, by using a substitution argument,

$$\int_{t_0}^{+\infty} |\dot{\varepsilon}(t)| dt = \int_{t_0}^{+\infty} \frac{1}{\beta(\tau^{-1}(t))} \left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) \right| dt = \int_{s_0}^{+\infty} \left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right| ds.$$

Therefore, if (2.43) is satisfied, then

$$\int_{t_0}^{+\infty} |\dot{\varepsilon}(t)| dt < +\infty.$$

Invoking Theorem 2.5, the trajectory $x(t) = z(\tau^{-1}(t))$ converges strongly to the minimum norm solution $\xi_* := \mathbf{P}_{\text{Zer}M}(0)$ as $t \rightarrow +\infty$. This is nothing else than the trajectory $z(s)$ converges strongly to ξ_* as $s \rightarrow +\infty$. \square

Before deriving in a similar manner the rates of convergence for the trajectory solution of (2.32), we express the quantity introduced in (2.18) in terms of β and δ . For $s_1 \geq s_0$ and a triple $(z_0, v, \xi) \in \mathcal{H} \times \mathcal{H} \times \mathcal{H}$, we have

$$\begin{aligned} D_1(z(s_1), v, \xi, \tau(s_1)) &= \sqrt{\|M(x(\tau(s_1))) + \varepsilon(\tau(s_1))(x(\tau(s_1)) - v)\|^2 + 8D_0(z_0, v, \xi)^2 |\dot{\varepsilon}(\tau(s_1))|} \\ &= \frac{1}{\beta(s_1)} \sqrt{\|\beta(s_1)M(z(s_1)) + \delta(s_1)(z(s_1) - v)\|^2 + 8D_0(z_0, v, \xi)^2 \beta(s_1) \left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right|}. \end{aligned}$$

Theorem 2.13. *Let $z: [s_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of the dynamical system (2.32) and $\xi \in \text{Zer}M$. Then, the following statements are true:*

(i) *if there exists $s_1 \geq s_0$ such that*

$$s \mapsto \frac{\delta(s)}{\beta(s)} \text{ is twice differentiable and } s \mapsto \frac{d}{ds} \left(\frac{1}{\beta(s)} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) + \frac{\delta^2(s)}{\beta^2(s)} \right) \text{ preserves constant sign on } [s_1, +\infty), \quad (2.44)$$

then for every $s \geq s_1$ it holds

$$\begin{aligned} &\|\beta(s)M(x(\tau(s))) + \delta(s)x(\tau(s))\|^2 + \beta(s) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \|x(\tau(s))\|^2 \\ &\leq \frac{\beta^2(s)}{\beta^2(s_1)} \exp \left(-2 \int_{s_1}^s \delta(u) du \right) \left\{ \|\beta(s_1)M(z(s_1)) + \delta(s_1)z(s_1)\|^2 + \beta(s_1) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \|z(s_1)\|^2 \right\} \\ &\quad + 4D_0(z_0, 0, \xi)^2 \beta(s) \left\{ \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) - \frac{\beta(s)}{\beta(s_1)} \exp \left(-2 \int_{s_1}^s \delta(u) du \right) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right\}_+; \end{aligned} \quad (2.46)$$

(ii) *if, in addition,*

$$s \mapsto \frac{\delta(s)}{\beta(s)} \text{ is nonincreasing on } [s_1, +\infty), \quad (2.47)$$

then for almost every $s \geq s_1$ it holds

$$\|\dot{z}(s)\| \leq D_1(z(s_1), 0, \xi, \tau(s_1)) \max \left\{ \sqrt{\frac{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right|}{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right|}}, \beta(s) \exp \left(- \int_{s_1}^s \delta(u) du \right) \right\}, \quad (2.48)$$

and for every $s \geq s_1$ it holds

$$\begin{aligned} \|M(z(s))\| \leq & D_1(z(s_1), 0, \xi, \tau(s_1)) \max \left\{ \frac{\sqrt{\beta(s_1)} \sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right|}}{\sqrt{\beta(s)} \sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right|}}, \exp \left(- \int_{s_1}^s \delta(u) du \right) \right\} \\ & + 2D_0(z_0, 0, \xi) \frac{\delta(s)}{\beta(s)}. \end{aligned} \quad (2.49)$$

Proof. For $t_0 \geq 0$ and

$$\tau(s) = t_0 + \int_{s_0}^s \beta(u) du \quad \forall s \geq s_0,$$

we consider again $x(\cdot) := z(\tau^{-1}(\cdot))$ the trajectory solution of (2.1) with $\varepsilon(t) := \frac{\delta(\tau^{-1}(t))}{\beta(\tau^{-1}(t))}$ for every $t \geq t_0$.

(i) For every $t \geq t_0$ we have

$$\dot{\varepsilon}(t) = \frac{1}{\beta(\tau^{-1}(t))} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t))$$

and

$$\begin{aligned} \ddot{\varepsilon}(t) &= - \frac{\dot{\beta}(\tau^{-1}(t)) (\tau^{-1})'(t)}{\beta^2(\tau^{-1}(t))} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) + \frac{1}{\beta(\tau^{-1}(t))} \left[\frac{d^2}{ds^2} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) (\tau^{-1})'(t) \\ &= \frac{1}{\beta^2(\tau^{-1}(t))} \left\{ \left[\frac{d^2}{ds^2} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) - \frac{\dot{\beta}(\tau^{-1}(t))}{\beta(\tau^{-1}(t))} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) \right\}, \end{aligned}$$

therefore

$$\ddot{\varepsilon}(t) + 2\varepsilon(t) \dot{\varepsilon}(t) = \frac{1}{\beta^2(\tau^{-1}(t))} \left\{ \left[\frac{d^2}{ds^2} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) + \left(2\delta(\tau^{-1}(t)) - \frac{\dot{\beta}(\tau^{-1}(t))}{\beta(\tau^{-1}(t))} \right) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (\tau^{-1}(t)) \right\}.$$

For every $s \geq s_0$ it holds

$$\begin{aligned} & \frac{d}{ds} \left(\frac{1}{\beta(s)} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) + \frac{\delta^2(s)}{\beta^2(s)} \right) \\ &= - \frac{\dot{\beta}(s)}{\beta^2(s)} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) + \frac{1}{\beta(s)} \left[\frac{d^2}{ds^2} \left(\frac{\delta}{\beta} \right) \right] (s) + 2 \frac{\delta(s)}{\beta(s)} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \\ &= \frac{1}{\beta(s)} \left\{ \left[\frac{d^2}{ds^2} \left(\frac{\delta}{\beta} \right) \right] (s) + \left(2\delta(s) - \frac{\dot{\beta}(s)}{\beta(s)} \right) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right\}, \end{aligned}$$

therefore the function $t \mapsto \ddot{\varepsilon}(t) + 2\varepsilon(t) \dot{\varepsilon}(t) = \frac{d}{dt}(\dot{\varepsilon}(t) + \varepsilon^2(t))$ preserves constant sign on $[t_1, +\infty)$, where $t_1 := \tau(s_1)$.

According to Theorem 2.7 (i), the estimate (2.20) is satisfied for every $t \geq t_1$. By taking $t := \tau(s)$ in this inequality, we obtain for every $s \geq s_1$

$$\begin{aligned} & \frac{1}{2} \|M(x(\tau(s)) + \varepsilon(\tau(s))x(\tau(s)))\|^2 + \frac{\dot{\varepsilon}(\tau(s))}{2} \|x(\tau(s))\|^2 \\ & \leq \frac{\gamma^2(\tau(s_1))}{2\gamma^2(\tau(s))} \left(\|M(x(\tau(s_1)) + \varepsilon(\tau(s_1))x(\tau(s_1)))\|^2 + \dot{\varepsilon}(\tau(s_1)) \|x(\tau(s_1))\|^2 \right) \\ & \quad + 2D_0(x_0, 0, \xi)^2 \left[\dot{\varepsilon}(\tau(s)) - \frac{1}{\gamma^2(\tau(s))} \dot{\varepsilon}(\tau(s_1)) \gamma^2(\tau(s_1)) \right]_+. \end{aligned}$$

Multiplying both sides by $2[\dot{\tau}(s)]^2 = 2\beta^2(s) > 0$, we obtain for every $s \geq s_0$

$$\|\beta(s)M(x(\tau(s))) + \delta(s)x(\tau(s))\|^2 + \beta(s) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \|x(\tau(s))\|^2$$

$$\begin{aligned}
&\leq \frac{\beta^2(s) \gamma^2(\tau(s_1))}{\beta^2(s_1) \gamma^2(\tau(s))} \left\{ \|\beta(s_1)M(x(\tau(s_1))) + \delta(s_1)x(\tau(s_1))\|^2 + \beta(s_1) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \|x(\tau(s_1))\|^2 \right\} \\
&\quad + 4D_0(z_0, 0, \xi)^2 \beta(s) \left[\left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) - \frac{\beta(s)}{\gamma^2(\tau(s))} \frac{\gamma^2(\tau(s_1))}{\beta(s_1)} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right]_+ \\
&= \frac{\beta^2(s)}{\beta^2(s_1)} \exp \left(-2 \int_{s_1}^s \delta(u) du \right) \left\{ \|\beta(s_1)M(z(s_1)) + \delta(s_1)z(s_1)\|^2 + \beta(s_1) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \|z(s_1)\|^2 \right\} \\
&\quad + 4D_0(z_0, 0, \xi)^2 \beta(s) \left\{ \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) - \frac{\beta(s)}{\beta(s_1)} \exp \left(-2 \int_{s_1}^s \delta(u) du \right) \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right\}_+,
\end{aligned}$$

where, by recalling the calculation in (2.39), we used that

$$\gamma(\tau(s)) = \exp \left(\int_{t_0}^{\tau(s)} \varepsilon(r) dr \right) = \exp \left(\int_{s_0}^s \delta(u) du \right) \quad \forall s \geq s_0.$$

(ii) Assuming that (2.47) holds is equivalent with requiring that $t \mapsto \varepsilon(t)$ is nonincreasing on $t \geq t_1 = \tau(s_1)$. This means that we can make use of the two estimates provided in Theorem 2.7 (ii).

From (2.22), by taking $t := \tau(s)$, we obtain for almost every $s \geq s_1$ that

$$\begin{aligned}
\|\dot{z}(s)\| &= \|\dot{x}(\tau(s)) \dot{\tau}(s)\| = \beta(s) \|\dot{x}(\tau(s))\| \\
&\leq D_1(x(\tau(s_1)), 0, \xi, \tau(s_1)) \beta(s) \max \left\{ \frac{\sqrt{|\dot{\varepsilon}(\tau(s))|}}{\sqrt{|\dot{\varepsilon}(\tau(s_1))|}}, \frac{\gamma(\tau(s_1))}{\gamma(\tau(s))} \right\} \\
&= D_1(z(s_1), 0, \xi, \tau(s_1)) \max \left\{ \frac{\sqrt{\beta(s_1)\beta(s)} \sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right|}}{\sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right|}}, \beta(s) \exp \left(- \int_{s_1}^s \delta(u) du \right) \right\}.
\end{aligned}$$

From (2.23), by taking $t := \tau(s)$, we obtain that for every $s \geq s_1$ that

$$\begin{aligned}
\|M(z(s))\| &\leq D_1(z(s_1), 0, \xi, \tau(s_1)) \max \left\{ \frac{\sqrt{\beta(s_1)}}{\sqrt{\beta(s)}} \frac{\sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right|}}{\sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s_1) \right|}}, \exp \left(- \int_{s_1}^s \delta(u) du \right) \right\} \\
&\quad + 2D_0(z_0, 0, \xi) \frac{\delta(s)}{\beta(s)}.
\end{aligned}$$

□

To illustrate the two theoretical results of this subsection, we consider the case where, for $\alpha > 0$,

$$\delta(s) := \frac{\alpha}{s} \quad \forall s \geq s_0.$$

With this choice of the regularization function, we will later establish a connection between the first-order dynamical system and a second-order system with vanishing damping.

Theorem 2.14. *Let $s_0 > 0$, $\alpha > 0$, $\beta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ a twice differentiable function and $z : [s_0, +\infty) \rightarrow \mathcal{H}$ the trajectory solution of the dynamical system*

$$\dot{z}(s) + \beta(s)M(z(s)) + \frac{\alpha}{s}z(s) = 0, \quad (2.50)$$

with initial condition $z(s_0) := z_0 \in \mathcal{H}$. Suppose that there exists $s_1 \geq s_0$ such that

$$s \mapsto s\beta(s) \text{ is nondecreasing on } [s_1, +\infty) \quad \text{and} \quad \lim_{s \rightarrow +\infty} s\beta(s) = +\infty, \quad (2.51)$$

and

$$\frac{s\dot{\beta}(s)}{\beta(s)} \leq \alpha - 1 \quad \text{and} \quad \beta(s) \left(\dot{\beta}(s) + s\ddot{\beta}(s) \right) \geq s\dot{\beta}(s)^2 \quad \forall s \geq s_1. \quad (2.52)$$

Then, the following statements are true:

(i) it holds

$$\|\dot{z}(s)\| = \mathcal{O}\left(\frac{1}{s}\right) \quad \text{as } s \rightarrow +\infty, \quad (2.53)$$

$$\|M(z(s))\| = \mathcal{O}\left(\frac{1}{s\beta(s)}\right) \quad \text{as } s \rightarrow +\infty; \quad (2.54)$$

(ii) the trajectory $z(s)$ converges strongly to the minimum norm solution $\xi_* := \mathbf{P}_{\text{Zer}M}(0)$ as $s \rightarrow +\infty$.

Proof. It is straightforward to see that, by defining $\delta(s) := \frac{\alpha}{s}$, for every $s \geq s_0$,

$$\int_{s_0}^{+\infty} \delta(s) ds = +\infty \quad \text{and} \quad \int_{s_0}^{+\infty} \beta(s) ds = +\infty,$$

thus, the conditions (2.33) are satisfied.

For every $s \geq s_1$ it holds

$$\left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) = -\alpha \frac{\frac{d}{ds}(s\beta(s))}{s^2\beta(s)^2} = -\alpha \frac{\beta(s) + s\dot{\beta}(s)}{s^2\beta(s)^2} \leq 0,$$

thus, the function $s \mapsto \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s)$ is nonincreasing on $[s_1, +\infty)$.

Since even both conditions (2.42) and (2.43) are satisfied, Theorem 2.12 yields the strong convergence of $z(s)$ to the minimum norm solution as $s \rightarrow +\infty$.

In addition, for every $s \geq s_1$ it holds

$$\frac{1}{\beta(s)} \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) + \frac{\delta^2(s)}{\beta^2(s)} = -\alpha \frac{\beta(s) + s\dot{\beta}(s)}{s^2\beta(s)^3} + \frac{\alpha^2}{s^2\beta(s)^2} = \frac{\alpha}{s^2\beta(s)^2} \left(\alpha - 1 - \frac{s\dot{\beta}(s)}{\beta(s)} \right).$$

Since $s \mapsto \frac{1}{s\beta(s)}$ is nonincreasing on $[s_1, +\infty)$ and, due to (2.52), $s \mapsto \alpha - 1 - \frac{s\dot{\beta}(s)}{\beta(s)}$ is nonnegative and nonincreasing on $[s_1, +\infty)$, condition (2.44) is satisfied. This means that the rates of convergence for both the velocity and the operator norm can be deduced from the estimates given in Theorem 2.13 (ii).

To this end, we notice first that for every $s \geq s_1$ it holds

$$\sqrt{\beta(s)} \sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right|} = \sqrt{\alpha \frac{\frac{d}{ds}(s\beta(s))}{s^2\beta(s)}} \leq \frac{\alpha}{s}, \quad (2.55)$$

since, according to (2.52),

$$0 \leq \frac{\frac{d}{ds}(s\beta(s))}{\beta(s)} \leq \alpha \quad \forall s \geq s_1.$$

Invoking again (2.52), for every $s \geq s_1$ it holds

$$\frac{d}{ds} (s^{1-\alpha}\beta(s)) = s^{1-\alpha}\dot{\beta}(s) + (1-\alpha)s^{-\alpha}\beta(s) = s^{-\alpha} (s\dot{\beta}(s) + (1-\alpha)\beta(s)) \leq 0,$$

thus

$$\beta(s) \leq s_1^{1-\alpha}\beta(s_1)s^{\alpha-1}. \quad (2.56)$$

Since, for every $s \geq s_1$

$$\exp\left(-\int_{s_1}^s \delta(u) du\right) = \exp\left(-\int_{s_1}^s \frac{\alpha}{u} du\right) = \frac{s_1^\alpha}{s^\alpha}, \quad (2.57)$$

we obtain

$$\beta(s) \exp\left(-\int_{s_1}^s \delta(u) du\right) \leq \beta(s_1) \frac{s_1}{s}. \quad (2.58)$$

Substituting (2.55) and (2.58) into (2.48), and using (2.56), yields (2.53).

On the other hand, for every $s \geq s_1$ it holds

$$\frac{\sqrt{\left| \left[\frac{d}{ds} \left(\frac{\delta}{\beta} \right) \right] (s) \right|}}{\sqrt{\beta(s)}} = \sqrt{\alpha \frac{\frac{d}{ds}(s\beta(s))}{s^2\beta(s)^3}} \leq \frac{\alpha}{s\beta(s)}. \quad (2.59)$$

Substituting (2.59) and (2.57) into (2.43) yields (2.54). \square

The conditions (2.51) and (2.52) are fulfilled, for instance, for

$$\beta(s) = s^p, \quad \text{where } -1 < p \leq \alpha - 1.$$

In this case, Theorem 2.14 leads to the following result.

Theorem 2.15 (the case $\delta(s) := \frac{\alpha}{s}$ and $\beta(s) = s^p, p > -1$). *Let $s_0 > 0$, $\alpha > 0$, $-1 < p \leq \alpha - 1$, and $z: [s_0, +\infty) \rightarrow \mathcal{H}$ the trajectory solution of the dynamical system*

$$\dot{z}(s) + s^p M(z(s)) + \frac{\alpha}{s} z(s) = 0, \quad (2.60)$$

with initial condition $z(s_0) := z_0 \in \mathcal{H}$.

Then, the following statements are true:

(i) it holds

$$\|\dot{z}(s)\| = \mathcal{O}\left(\frac{1}{s}\right) \quad \text{as } s \rightarrow +\infty, \quad (2.61)$$

$$\|M(z(s))\| = \mathcal{O}\left(\frac{1}{s^{1+p}}\right) \quad \text{as } s \rightarrow +\infty; \quad (2.62)$$

(ii) the trajectory $z(s)$ converges strongly to the minimum norm solution $\xi_* := P_{\text{Zer}M}(0)$ as $s \rightarrow +\infty$.

2.3 First-order dynamical system with anchor point

Enhancing the dynamical systems (2.32) with an anchor point represents the final step in our attempt to establish a connection between first-order dynamical systems with Tikhonov regularization and second-order dynamical systems with a vanishing damping term.

Let $s_0 \geq 0$ and $\beta, \delta: [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ be given functions. We consider on $[s_0, +\infty)$ the following dynamical system

$$\dot{y}(s) + \beta(s) M(y(s)) + \delta(s) (y(s) - v) = 0, \quad (2.63)$$

with initial condition $y(s_0) := y_0 \in \mathcal{H}$ and anchor point $v \in \mathcal{H}$.

The convergence and convergence rate results derived for (2.32) can be transferred to (2.63). In particular, the rates of convergence are not affected by the presence of the anchor point. The only difference is the limit point to which the trajectory converges strongly, now being the closest point to v in the solution set $\text{Zer}M$.

Define $z: [s_0, +\infty) \rightarrow \mathcal{H}$ as

$$z(s) := y(s) - v \quad \forall s \geq s_0. \quad (2.64)$$

The main idea relies on the observation that $y(\cdot)$ is the trajectory solution of (2.63) if and only if $z(\cdot)$ is the trajectory solution of

$$\dot{z}(s) + \beta(s) M^v(z(s)) + \delta(s) z(s) = 0, \quad (2.65)$$

where $M^v: \mathcal{H} \rightarrow \mathcal{H}$ is the monotone operator defined as

$$M^v(z) = M(z + v) \quad \forall z \in \mathcal{H}. \quad (2.66)$$

The rates of convergence for the velocity and the operator norm follow from Theorem 2.13, since $\dot{z}(s) = \dot{y}(s)$ and $M^v(z(s)) = M(y(s))$ for every $s \geq s_0$.

If β and δ , in addition to (2.33), satisfy either (2.42) or (2.43), then the trajectory $y(s)$ converges strongly to $P_{\text{Zer}M}(v)$. Indeed, according to Theorem 2.12, $z(s) = y(s) - v$ converges strongly to $P_{\text{Zer}M^v}(0) = P_{\text{Zer}M}(v) - v$ (see [11, Proposition 29.1]) as $s \rightarrow +\infty$. In other words, $y(s)$ converges strongly to $P_{\text{Zer}M}(v)$ as $s \rightarrow +\infty$.

Remark 2.16. The dynamic (2.63) was also studied by Suh, Park, and Ryu in [50] in finite-dimensional spaces, in the case $\beta(s) := 1$ and $\varepsilon(s) := \frac{\alpha}{s^q}$ for every $s \geq s_0 > 0$, where $\alpha > 0$ and $0 < q \leq 1$. By relying on a very intricate Lyapunov analysis, the convergence of the trajectory and convergence rate for the operator norm has been provided.

These results are particular cases of our more general approach, see Theorems 2.8 and 2.9, which is fundamentally different, as it relies on the study of the two very simple functions φ and ψ , defined as in (2.4) and (2.5), respectively. In addition, our approach provides a rate of convergence for the velocity.

The dynamics (2.63) is referred to as the continuous model of the Halpern method. In light of the above considerations, it is evident that the Halpern method is nothing else than a particular instance of the classical Tikhonov regularization approach.

3 From first-order dynamics with Tikhonov regularization to second-order dynamics with vanishing damping term

In this section, we will demonstrate that, when applied to monotone equations of the type (1.1), first-order dynamics with a specific Tikhonov regularization term are intimately related to first-order dynamics with a vanishing damping term. The latter have been proved in [14] to accelerate the convergence rate of the operator norm with respect to the first-order monotone operator flow, while preserving the weak convergence of the trajectory to a zero of the operator.

Let $s_0 > 0$, $\alpha > 1$ and $\beta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ a continuous differentiable function. We consider on $[s_0, +\infty)$ the dynamical system

$$\ddot{z}(s) + \frac{\alpha}{s}\dot{z}(s) + \beta(s)\frac{d}{ds}M(z(s)) + \left(\dot{\beta}(s) + \frac{\beta(s)}{s}\right)M(z(s)) = 0, \quad (3.1)$$

with initial conditions $z(s_0) = z_0 \in \mathcal{H}$ and $\dot{z}(s_0) = \dot{z}_0 \in \mathcal{H}$.

We assume that it admits a unique strong global solution $z : [s_0, +\infty) \rightarrow \mathcal{H}$, meaning that z and $\cdot z$ are absolutely continuous on every compact interval $[s_0, S]$ and that z satisfies (3.1) almost everywhere, with the property that $s \mapsto M(z(s))$ is absolutely continuous on every compact interval $[s_0, S]$.

Proposition 3.1. *Let $s_0 > 0$ and $\alpha > 1$. Then $z : [s_0, +\infty) \rightarrow \mathcal{H}$ is the trajectory solution of (3.1) with the initial conditions $z(s_0) = z_0$ and $\dot{z}(s_0) = \dot{z}_0$ if and only if it is the trajectory solution of*

$$\dot{z}(s) + \beta(s)M(z(s)) + \frac{\alpha-1}{s}(z(s) - v) = 0, \quad (3.2)$$

with initial condition $z(s_0) = z_0$ and anchor point

$$v := z_0 + \frac{s_0}{\alpha-1}(\dot{z}_0 + \beta(s_0)M(z_0)).$$

Proof.

“second-order” \Leftrightarrow “first-order”. Let $z : [s_0, +\infty) \rightarrow \mathcal{H}$ be the trajectory solution of (3.1) with initial conditions $z(s_0) = z_0$ and $\dot{z}(s_0) = \dot{z}_0$. For almost every $s \geq s_0$ it holds

$$\frac{d}{ds}(s\dot{z}(s) + s\beta(s)M(z(s))) = s\ddot{z}(s) + \dot{z}(s) + s\beta(s)\frac{d}{ds}M(z(s)) + (s\dot{\beta}(s) + \beta(s))M(z(s)) = (1-\alpha)\dot{z}(s).$$

By integration from s_0 to s , we get for almost every $s \geq s_0$

$$s\dot{z}(s) - s_0\dot{z}_0 + s\beta(s)M(z(s)) - s_0\beta(s_0)M(z_0) = (1-\alpha)(z(s) - z_0)$$

or, equivalently,

$$\dot{z}(s) + \beta(s)M(z(s)) + \frac{\alpha-1}{s}(z(s) - v) = 0,$$

where $v := z_0 + \frac{s_0}{\alpha-1}(\dot{z}_0 + \beta(s_0)M(z_0))$.

“first-order” \Leftrightarrow “second-order”. Starting with the trajectory solution $z : [s_0, +\infty) \rightarrow \mathcal{H}$ of the first-order system (3.2) with initial condition $z(s_0) := z_0 \in \mathcal{H}$ and anchor point $v := z_0 + \frac{s_0}{\alpha-1}(\dot{z}_0 + \beta(s_0)M(z_0))$, it satisfies for almost every $s \geq s_0$

$$s\dot{z}(s) + s\beta(s)M(z(s)) + (\alpha-1)(z(s) - v) = 0.$$

In addition, $\dot{z}(s_0) = \dot{z}_0$. After differentiation, it yields for almost every $s \geq s_0$

$$s\ddot{z}(s) + \dot{z}(s) + s\beta(s)\frac{d}{ds}M(z(s)) + (s\dot{\beta}(s) + \beta(s))M(z(s)) + (\alpha-1)\dot{z}(s) = 0,$$

which is nothing else than (3.1). \square

Theorem 2.14 and the considerations in Subsection 2.3 lead to the following result.

Theorem 3.2. *Let $s_0 > 0$, $\alpha > 1$, $\beta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ a twice differentiable function and $z : [s_0, +\infty) \rightarrow \mathcal{H}$ the trajectory solution of the dynamical system (3.1) with initial conditions $z(s_0) = z_0$ and $\dot{z}(s_0) = \dot{z}_0$. Suppose that there exists $s_1 \geq s_0$ such that*

$$s \mapsto s\beta(s) \text{ is nondecreasing on } [s_1, +\infty) \text{ and } \lim_{s \rightarrow +\infty} s\beta(s) = +\infty,$$

and

$$\frac{s\dot{\beta}(s)}{\beta(s)} \leq \alpha - 2 \quad \text{and} \quad \beta(s) \left(\dot{\beta}(s) + s\ddot{\beta}(s) \right) \geq s\dot{\beta}(s)^2 \quad \forall s \geq s_1.$$

Then, the following statements are true:

(i) it holds

$$\|\dot{z}(s)\| = \mathcal{O}\left(\frac{1}{s}\right) \quad \text{as } s \rightarrow +\infty, \quad (3.3)$$

$$\|M(z(s))\| = \mathcal{O}\left(\frac{1}{s\beta(s)}\right) \quad \text{as } s \rightarrow +\infty; \quad (3.4)$$

(ii) the trajectory $z(s)$ converges strongly to $\mathbf{P}_{\text{Zer}M}\left(z_0 + \frac{s_0}{\alpha-1}(\dot{z}_0 + \beta(s_0)M(z_0))\right)$ as $s \rightarrow +\infty$.

Remark 3.3. In [14], for the Fast Optimistic Gradient Descent Ascent (OGDA) dynamics

$$\ddot{z}(s) + \frac{\alpha}{s}\dot{z}(s) + \beta(s)\frac{d}{ds}M(z(s)) + \frac{1}{2}\left(\dot{\beta}(s) + \frac{\alpha\beta(s)}{s}\right)M(z(s)) = 0 \quad (3.5)$$

with initial time $s_0 > 0$, $\alpha \geq 2$, $\beta : [s_0, +\infty) \rightarrow \mathbb{R}_{++}$ a continuous differentiable and nondecreasing function, and initial conditions $z(s_0) = z_0$ and $\dot{z}(s_0) = \dot{z}_0$, it has been shown that if

$$\sup_{s \geq s_0} \frac{s\dot{\beta}(s)}{\beta(s)} \leq \alpha - 2,$$

then

$$\|\dot{z}(s)\| = \mathcal{O}\left(\frac{1}{s}\right) \quad \text{and} \quad \|M(z(s))\| = \mathcal{O}\left(\frac{1}{s\beta(s)}\right) \quad \text{as } s \rightarrow +\infty,$$

without providing a statement regarding the convergence of the trajectory. On the other hand, if

$$\sup_{s \geq s_0} \frac{s\dot{\beta}(s)}{\beta(s)} < \alpha - 2,$$

then

$$\|\dot{z}(s)\| = o\left(\frac{1}{s}\right) \quad \text{and} \quad \|M(z(s))\| = o\left(\frac{1}{s\beta(s)}\right) \quad \text{as } s \rightarrow +\infty,$$

and $z(s)$ converges weakly to a zero of M as $s \rightarrow +\infty$.

If for every $s \geq s_0$ it holds

$$s\dot{\beta}(s) = (\alpha - 2)\beta(s),$$

that is, $\beta(s) = \beta_0 s^{\alpha-2}$ for some $\beta_0 > 0$, then the dynamical system (3.5) coincides with (3.1). According to Theorem 3.2, besides the rates of convergence for the velocity and the operator norm, we obtain in this borderline case that $z(s)$ converges strongly to $\mathbf{P}_{\text{Zer}M}\left(z_0 + \frac{s_0}{\alpha-1}(\dot{z}_0 + \beta(s_0)M(z_0))\right)$ as $s \rightarrow +\infty$.

This is particularly true when β is constant and $\alpha = 2$, which is of significant interest when deriving discrete explicit counterparts of (3.5).

4 Fixed point problem and discretization

In this section, we will introduce and analyze discrete time counterparts of the first-order dynamical system (2.1) in the context of identifying the fixed points of nonexpansive operators.

4.1 Problem formulation and overview of existing numerical methods

For $T : \mathcal{H} \rightarrow \mathcal{H}$ a *nonexpansive* (1-Lipschitz continuous) operator, i.e.,

$$\|T(x) - T(y)\| \leq \|x - y\| \quad \forall x, y \in \mathcal{H},$$

we consider the fixed point problem

$$\text{Find } x \in \mathcal{H} \text{ such that } x = T(x). \quad (4.1)$$

We denote the set of all *fixed points* of T by $\text{Fix}T := \{x \in \mathcal{H} : x = T(x)\}$ and assume throughout this section that this set is nonempty. The set $\text{Fix}T$ is convex and closed (see, for instance, [11, Corollary 4.24]).

There are many motivations for studying this particular class of problems. It has long been known that various monotone inclusions and convex optimization problems can be represented by fixed point formulations using nonexpansive operators. For a more detailed discussion of this connection, the reader is referred to [11, 27].

To establish the connection between (1.1) and (4.1), we observe that $\text{Fix}T = \text{Zer}(\text{Id} - T)$ and, since T is nonexpansive, $\text{Id} - T$ satisfies (see [11, Proposition 4.11]) the inequality

$$\langle x - y, (\text{Id} - T)(x) - (\text{Id} - T)(y) \rangle \geq \frac{1}{2} \|(\text{Id} - T)(x) - (\text{Id} - T)(y)\|^2 \geq 0 \quad \forall x, y \in \mathcal{H}. \quad (4.2)$$

This shows that $\text{Id} - T$ is a continuous and monotone operator. In other words, the fixed point problem (4.1) can be equivalently formulated as a monotone equation of type (1.1).

These premises allow to approach (4.1) by the dynamical system (2.30) for $M := \text{Id} - T$. This gives a convergence rate of $\mathcal{O}(\frac{1}{t})$ for $\|x(t) - T(x(t))\|$ and ensures the strong convergence of $x(t)$ to $\text{P}_{\text{Fix}T}(0)$ as $t \rightarrow +\infty$ (see also [16]). This is a significant improvement over both convergence statements for the dynamical system

$$\dot{x}(t) + (\text{Id} - T)(x(t)) = 0 \quad (4.3)$$

with the initial condition $x(t_0) = x_0$, as provided in Theorem A.1 (see, also, [13, Theorems 11 and 16]).

The most standard approach used to detect a fixed point of T is the *Banach-Picard iteration*, which reads for every $k \geq 0$

$$x_{k+1} := T(x_k), \quad (4.4)$$

where $x_0 \in \mathcal{H}$ is a given starting point. According to the Banach-Picard fixed point theorem, if T is a *contraction*, namely, T is Lipschitz continuous with modulus $\kappa \in [0, 1)$, then the sequence $\{x_k\}_{k \geq 0}$ generated by (4.4) converges strongly to the unique fixed point of T with a linear convergence rate. On the other hand, if T is merely nonexpansive, then Banach-Picard might diverge. To see this, it is enough to choose $T = -\text{Id}$ and $x_0 \neq 0$. A closer look at this example reveals that the Banach-Picard iteration not only fails to approach a fixed point of T , but also generates a sequence that does not satisfy the *asymptotic regularity property*. We say that the sequence $\{x_k\}_{k \geq 0}$ satisfies the asymptotic regularity property if the difference $x_k - T(x_k) \rightarrow 0$ as $k \rightarrow +\infty$. This property is crucial for guaranteeing the convergence of the iterates, as we will see later.

To overcome the restrictive contraction assumption on T , Krasnosel'skiĭ proposed in [37] applying the Banach-Picard iteration (4.4) to the operator $\frac{1}{2}\text{Id} + \frac{1}{2}T$ instead of T . Building on this idea, by iteratively considering a convex combination between Id and T , the so-called *Krasnosel'skiĭ-Mann iteration* emerges. Specifically, let $\{\alpha_k\}_{k \geq 0}$ be a sequence in $[0, 1]$. The Krasnosel'skiĭ-Mann iteration is defined for every $k \geq 0$ as

$$x_{k+1} := \alpha_k x_k + (1 - \alpha_k) T(x_k), \quad (4.5)$$

where, again, $x_0 \in \mathcal{H}$ is a given starting point.

A fundamental step in proving the convergence of the iterates of (4.5) is to demonstrate that $x_k - T(x_k) \rightarrow 0$ as $k \rightarrow +\infty$. This was first established by Browder and Petryshyn in [21] in the constant case $\alpha_k \equiv \alpha_0 \in (0, 1)$. The extension to nonconstant sequences was achieved by Groetsch in [28], who proved that if $\sum_{k \geq 0} \alpha_k (1 - \alpha_k) = +\infty$, then the asymptotic regularity property is satisfied. The weak convergence of the iterates has been studied in various settings in [11, 12, 28, 34, 46]. Techniques based on Tikhonov regularization to improve the convergence of the iterates from weak to strong have been recently explored in [15, 17].

It must be emphasized that the asymptotic regularity property does not automatically provide an explicit convergence rate. Indeed, deriving a precise rate of convergence is much more complex, and research on this subject is still limited compared to the extensive results on the convergence of iterates. The convergence of the Krasnosel'skiĭ-Mann iteration, expressed in terms of the fixed point residual $\|x_k - T(x_k)\|$, was proved to be $o\left(\frac{1}{\sqrt{k}}\right)$ by Baillon and Bruck in [10] for the case of a constant sequence $\{\alpha_k\}_{k \geq 0}$. For a nonconstant sequence, it was initially shown to be $\mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$ in [25, 39], and later on improved to $o\left(\frac{1}{\sqrt{k}}\right)$ in [26, 41, 19]. These results align with the statements in the continuous setting by Boţ and Csetnek ([13]) mentioned earlier. Contreras and Cominetti demonstrated in [23] that in the Banach space setting the lower bound of the Krasnosel'skiĭ-Mann iteration is $\mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$, whereas Maulén, Fierro, and Peypouquet proved in [42] that the rate of convergence of the fixed point residual of a general *inertial Krasnosel'skiĭ-Mann algorithm* is $o\left(\frac{1}{\sqrt{k}}\right)$.

In the following, we shall see that both Banach-Picard and Krasnosel'skiĭ-Mann iterations are obtained by temporal discretizations of the same dynamical system (4.3). The difference lies in the chosen step size. Specifically, we consider an explicit finite-difference scheme with step sizes $h_k > 0$

$$\frac{x_{k+1} - x_k}{h_k} + (\text{Id} - T)(x_k) = 0 \quad \forall k \geq 0,$$

which is equivalent to

$$x_{k+1} = x_k - h_k (\text{Id} - T)(x_k) = (1 - h_k)x_k + h_k T(x_k) \quad \forall k \geq 0. \quad (4.6)$$

From this, it is clear that Banach-Picard process is simply the scheme above with $h_k \equiv 1$, while taking $h_k = 1 - \alpha_k$ leads to the Krasnosel'skiĭ-Mann iteration.

By considering convex combinations with a fixed anchor point $v \in \mathcal{H}$, one obtains the *Halpern iteration* [31]

$$x_{k+1} := \alpha_k v + (1 - \alpha_k)T(x_k) \quad \forall k \geq 0. \quad (4.7)$$

The asymptotic regularity property of this iterative scheme has been studied in [55, 56]. This method has recently been gaining increasing attention [40, 45, 44, 50, 33]. One reason for its attractiveness, besides guaranteeing strong convergence of the iteration, is its acceleration property, which has been demonstrated recently. Sabach and Shtern proved in [47] for a general form of the Halpern iteration that the rate of convergence of the fixed point residual is of $\mathcal{O}(\frac{1}{k})$. Lieder obtained in [40] the same rate of convergence for the Halpern iteration (4.7) with $\alpha_k := \frac{1}{k+2}$ for every $k \geq 0$, while Park and Ryu showed in [44] that the $\mathcal{O}(\frac{1}{k})$ convergence rate is optimal within a specific family of fixed point iterations.

This rate of convergence is particularly attractive when compared to the closely related Krasnosel'skiĭ-Mann method. Building on these insights, researchers have started combining this anchoring technique with other methods to accelerate convergence. Notable contributions in this direction include works by Yoon and Ryu [58], Lee and Kim [38], and Tran-Dinh [53] and his coauthor Luo [54].

Recently, in [18] (see also [14]), explicit numerical methods with Nesterov momentum and correcting operator terms have been introduced. These methods exhibit a convergence rate for the fixed point residual of $\mathcal{O}(\frac{1}{k})$, while the sequence of iterates weakly converges to a fixed point of the operators. These numerical methods were derived through a subtle discretization of the Fast Optimistic Gradient Descent Ascent (OGDA) dynamics (1.6).

The main aim of this section is to show that the Halpern iteration serves as the discrete counterpart to the first-order dynamical system (2.1) with an anchor point $v \in \mathcal{H}$ (see (2.63)). Additionally, we will derive convergence rates for the vanishing of both the discrete velocity and the fixed point residual for various choices of the sequence $\{\alpha_k\}_{k \geq 0}$, and establish the strong convergence of the iterates to $\text{P}_{\text{Fix}T}(v)$. Our approach will utilize novel proof techniques that are inspired by those used in the continuous time analysis.

4.2 Convergence results of the discretization

To construct a numerical method, we consider the dynamics given by (2.1) where an anchor point $v \in \mathcal{H}$ is also added (or the dynamics (2.63) with $\beta \equiv 1$)

$$\dot{x}(t) + (\text{Id} - T)(x(t)) + \varepsilon(t)(x(t) - v) = 0,$$

with initial condition $x(t_0) = x_0 \in \mathcal{H}$.

The explicit finite-difference scheme with step sizes 1 gives

$$x_{k+1} - x_k + (\text{Id} - T)(x_k) + \varepsilon_k(x_{k+1} - v) = 0 \quad \forall k \geq 0$$

or, equivalently,

$$x_{k+1} := \frac{\varepsilon_k}{1 + \varepsilon_k}v + \frac{1}{1 + \varepsilon_k}T(x_k) \quad \forall k \geq 0.$$

Here $\{\varepsilon_k\}_{k \geq 0}$ is a positive, so-called *regularization sequence* which mimics the properties (2.2) of the regularization function $\varepsilon(t)$ in the continuous case, and is therefore required to fulfill

$$\lim_{k \rightarrow +\infty} \varepsilon_k = 0 \text{ and } \sum_{k \geq 0} \varepsilon_k = +\infty. \quad (4.8)$$

For convenience, we write for every $k \geq 0$

$$\alpha_k := \frac{\varepsilon_k}{1 + \varepsilon_k} \in (0, 1) \Leftrightarrow 1 - \alpha_k = \frac{1}{1 + \varepsilon_k}.$$

Then (4.8) translate into the following condition.

The sequence $\{\alpha_k\}_{k \geq 0} \subseteq (0, 1)$ satisfies

$$\lim_{k \rightarrow +\infty} \alpha_k = 0 \text{ and } \sum_{k \geq 0} \alpha_k = +\infty. \quad (4.9)$$

With this new regularization sequence, the update rule turns out to be nothing else but the Halpern fixed point iteration ([31]).

Let $v, x_0 \in \mathcal{H}$. For every $k \geq 0$ set

$$x_{k+1} := \alpha_k v + (1 - \alpha_k) T(x_k). \quad (4.10)$$

Next, we provide some preliminary estimates that will be useful in the convergence analysis of (4.10).

Lemma 4.1. *Let $\{x_k\}_{k \geq 0}$ be the sequences generated by the iterative scheme (4.10) and $\xi \in \text{Fix}T$. Then, the following inequalities are satisfied for every $k \geq 0$:*

$$\|x_k - \xi\| \leq D_0(x_0, v, \xi), \quad (4.11)$$

$$\|T(x_k) - v\| \leq 2D_0(x_0, v, \xi), \quad (4.12)$$

$$\begin{aligned} \frac{1}{2} \|x_{k+1} - \xi\|^2 &\leq \frac{1}{2} (1 - \alpha_k) \|x_k - \xi\|^2 + \alpha_k \langle x_k - \xi, v - \xi \rangle \\ &\quad + \alpha_k D_0(x_0, v, \xi) \left(\frac{3}{2} D_0(x_0, v, \xi) \alpha_k + \|x_k - T(x_k)\| \right). \end{aligned} \quad (4.13)$$

Proof. We use an induction argument to prove the first inequality. Obviously (4.11) holds when $k = 0$. Suppose that (4.11) holds for $k = k_0 \geq 0$. It follows from (4.10) and the triangle inequality that

$$\begin{aligned} \|x_{k_0+1} - \xi\| &= \|\alpha_{k_0} v + (1 - \alpha_{k_0}) T(x_{k_0}) - \xi\| \\ &\leq \alpha_{k_0} \|v - \xi\| + (1 - \alpha_{k_0}) \|T(x_{k_0}) - \xi\| \\ &= \alpha_{k_0} \|v - \xi\| + (1 - \alpha_{k_0}) \|T(x_{k_0}) - T(\xi)\| \\ &\leq \alpha_{k_0} \|v - \xi\| + (1 - \alpha_{k_0}) \|x_{k_0} - \xi\| \leq \max\{\|x_0 - \xi\|, \|v - \xi\|\} = D_0(x_0, v, \xi). \end{aligned}$$

This means that (4.11) is satisfied for $k = k_0 + 1$, so it is satisfied for every $k \geq 0$.

The inequality (4.12) follows immediately from this, since for every $k \geq 0$ we have

$$\begin{aligned} \|T(x_k) - v\| &\leq \|T(x_k) - T(\xi)\| + \|T(\xi) - v\| \\ &\leq \|x_k - \xi\| + \|T(\xi) - v\| = \|x_k - \xi\| + \|\xi - v\| \leq 2D_0(x_0, v, \xi). \end{aligned}$$

Now, let $k \geq 0$ be fixed. By the definition of x_{k+1} in (4.10), we have

$$\begin{aligned} \frac{1}{2} \|x_{k+1} - \xi\|^2 &= \frac{1}{2} \|\alpha_k v + (1 - \alpha_k) T(x_k) - \xi\|^2 = \frac{1}{2} \|\alpha_k (v - \xi) + (1 - \alpha_k) (T(x_k) - \xi)\|^2 \\ &= \frac{1}{2} (1 - \alpha_k)^2 \|T(x_k) - \xi\|^2 + (1 - \alpha_k) \alpha_k \langle T(x_k) - \xi, v - \xi \rangle + \frac{1}{2} \alpha_k^2 \|v - \xi\|^2 \\ &= \frac{1}{2} (1 - \alpha_k)^2 \|T(x_k) - T(\xi)\|^2 + (1 - \alpha_k) \alpha_k \langle x_k - \xi, v - \xi \rangle \\ &\quad - (1 - \alpha_k) \alpha_k \langle x_k - T(x_k), v - \xi \rangle + \frac{1}{2} \alpha_k^2 \|v - \xi\|^2. \end{aligned} \quad (4.14)$$

Using successively the nonexpansiveness of T , the Cauchy-Schwarz inequality, and the fact that $\alpha_k \in (0, 1)$, we infer

$$\begin{aligned} &\frac{1}{2} (1 - \alpha_k)^2 \|T(x_k) - T(\xi)\|^2 - \alpha_k^2 \langle x_k - \xi, v - \xi \rangle - (1 - \alpha_k) \alpha_k \langle x_k - T(x_k), v - \xi \rangle + \frac{1}{2} \alpha_k^2 \|v - \xi\|^2 \\ &\leq \frac{1}{2} (1 - \alpha_k)^2 \|x_k - \xi\|^2 + \alpha_k^2 \|x_k - \xi\| \|v - \xi\| + (1 - \alpha_k) \alpha_k \|x_k - T(x_k)\| \|v - \xi\| + \frac{1}{2} \alpha_k^2 \|v - \xi\|^2 \\ &\leq \frac{1}{2} (1 - \alpha_k) \|x_k - \xi\|^2 + \alpha_k^2 \|x_k - \xi\| \|v - \xi\| + \alpha_k \|x_k - T(x_k)\| \|v - \xi\| + \frac{1}{2} \alpha_k^2 \|v - \xi\|^2 \\ &\leq \frac{1}{2} (1 - \alpha_k) \|x_k - \xi\|^2 + \frac{3}{2} D_0(x_0, v, \xi)^2 \alpha_k^2 + D_0(x_0, v, \xi) \alpha_k \|x_k - T(x_k)\|, \end{aligned}$$

and the announced statement follows by plugging this estimate into (4.14). \square

In analogy to the analysis in the continuous time regime, we provide a result that allows us to derive convergence rates for the discrete velocity and the fixed point residual of the operator, as well as ensuring strong convergence for the sequence of iterates.

Proposition 4.2. *Let $\{x_k\}_{k \geq 0}$ be the sequences generated by the iterative scheme (4.10) and $\xi \in \text{Fix}T$. Then, the following statement are true:*

(i) for every $k \geq 1$ it holds

$$\|x_{k+1} - x_k\| \leq (1 - \alpha_k) \|x_k - x_{k-1}\| + 2D_0(x_0, v, \xi) |\alpha_k - \alpha_{k-1}|, \quad (4.15)$$

$$\|x_{k+1} - T(x_{k+1})\| \leq (1 - \alpha_k) \|x_{k+1} - x_k\| + 2D_0(x_0, v, \xi) \alpha_k; \quad (4.16)$$

(ii) if $\|x_k - T(x_k)\| \rightarrow 0$ as $k \rightarrow +\infty$, then $\{x_k\}_{k \geq 0}$ converges strongly to the fixed point of T closest to v , $\xi_* := \text{P}_{\text{Fix}T}(v)$, as $k \rightarrow +\infty$.

Proof. (i) Let $k \geq 1$ be fixed. According to (4.10), and by using the triangle inequality, we can derive that

$$\begin{aligned} \|x_{k+1} - x_k\| &= \|\alpha_k v + (1 - \alpha_k) T(x_k) - \alpha_{k-1} v - (1 - \alpha_{k-1}) T(x_{k-1})\| \\ &= \|(1 - \alpha_k) (T(x_k) - T(x_{k-1})) + (\alpha_k - \alpha_{k-1}) (v - T(x_{k-1}))\| \\ &\leq (1 - \alpha_k) \|T(x_k) - T(x_{k-1})\| + |\alpha_k - \alpha_{k-1}| \|T(x_{k-1}) - v\| \\ &\leq (1 - \alpha_k) \|x_k - x_{k-1}\| + 2D_0(x_0, v, \xi) |\alpha_k - \alpha_{k-1}|, \end{aligned}$$

which yields further

$$\begin{aligned} \|x_{k+1} - T(x_{k+1})\| &= \|\alpha_k v + (1 - \alpha_k) T(x_k) - T(x_{k+1})\| \\ &\leq (1 - \alpha_k) \|T(x_k) - T(x_{k+1})\| + \alpha_k \|v - T(x_{k+1})\| \\ &\leq (1 - \alpha_k) \|x_{k+1} - x_k\| + 2D_0(x_0, v, \xi) \alpha_k. \end{aligned}$$

(ii) In the view of (4.13), we will prove that

$$\limsup_{k \rightarrow +\infty} \langle x_k - \xi_*, v - \xi_* \rangle \leq 0. \quad (4.17)$$

Recall that the sequence of iterates $\{x_k\}_{k \geq 0}$ is bounded, see Lemma 4.1, therefore there exists a subsequence $\{x_{k_n}\}_{n \geq 0}$ of $\{x_k\}_{k \geq 0}$ that converges weakly to \hat{x} as $n \rightarrow +\infty$ and

$$\limsup_{k \rightarrow +\infty} \langle x_k - \xi_*, v - \xi_* \rangle = \lim_{n \rightarrow +\infty} \langle x_{k_n} - \xi_*, v - \xi_* \rangle = \langle \hat{x} - \xi_*, v - \xi_* \rangle.$$

Since, according to the assumption, $\|x_{k_n} - T(x_{k_n})\| \rightarrow 0$ as $n \rightarrow +\infty$, the Browder's demiclosedness principle ([11, Theorem 4.27]) yields $\hat{x} \in \text{Fix}T$. Therefore, since $\xi_* := \text{P}_{\text{Fix}T}(v)$, we have from the variational characterization of the projection that ([11, Theorem 3.16])

$$\langle \hat{x} - \xi_*, v - \xi_* \rangle \leq 0.$$

This yields (4.17), which allows us to deduce further that

$$\begin{aligned} &\limsup_{k \rightarrow +\infty} \left(\langle x_k - \xi_*, v - \xi_* \rangle + D_0(x_0, v, \xi) \left(\frac{3}{2} D_0(x_0, v, \xi) \alpha_k + \|x_k - T(x_k)\| \right) \right) \\ &= \limsup_{k \rightarrow +\infty} \langle x_k - \xi_*, v - \xi_* \rangle \leq 0. \end{aligned}$$

We are therefore in a position to use Lemma A.7 with

$$a_k := \frac{1}{2} \|x_k - \xi_*\|^2, \quad b_k := \langle x_k - \xi_*, v - \xi_* \rangle + D_0(x_0, v, \xi) \left(\frac{3}{2} D_0(x_0, v, \xi) \alpha_k + \|x_k - T(x_k)\| \right), \quad d_k := 0,$$

which yields the conclusion. \square

Relying on the estimates (4.15)-(4.16), we can formulate further conditions on the sequence $\{\alpha_k\}_{k \geq 0}$ which, in the light of Proposition 4.2(ii), lead to the strong convergence of the sequence generated by the Halpern iteration (4.10).

Theorem 4.3. Let $\{x_k\}_{k \geq 0}$ be the sequence generated by the iterative scheme (4.10). Assume that, in addition to (4.9), the sequence $\{\alpha_k\}_{k \geq 0}$ satisfies either

$$\lim_{k \rightarrow +\infty} \frac{|\alpha_{k+1} - \alpha_k|}{\alpha_{k+1}} = 0 \quad (4.18)$$

or

$$\sum_{k \geq 0} |\alpha_{k+1} - \alpha_k| < +\infty. \quad (4.19)$$

Then, $\|x_k - T(x_k)\| \rightarrow 0$ as $k \rightarrow +\infty$, and therefore the sequence $\{x_k\}_{k \geq 0}$ strongly converges to $\xi_* := P_{\text{Fix}T}(v)$ as $k \rightarrow +\infty$.

Proof. Let $\xi \in \text{Fix}T$. The statement follows by applying Lemma A.7 to the inequality (4.15) in two different ways, depending on the additional assumptions for $\{\alpha_k\}_{k \geq 0}$. Specifically, for every $k \geq 1$ we set $a_k := \|x_k - x_{k-1}\|$. If (4.19) holds, we choose $b_k := 0$ and $d_k := 2D_0(x_0, v, \xi) |\alpha_k - \alpha_{k-1}|$, while if (4.18) holds we choose $b_k := 2D_0(x_0, v, \xi) \frac{|\alpha_k - \alpha_{k-1}|}{\alpha_k}$ and $d_k := 0$, respectively. In both cases we can conclude that $\|x_k - x_{k-1}\| \rightarrow 0$ as $k \rightarrow +\infty$, which immediately gives $\|x_k - T(x_k)\| \rightarrow 0$ as $k \rightarrow +\infty$ due to the inequality (4.16) and the assumption $\lim_{k \rightarrow +\infty} \alpha_k = 0$. Proposition 4.2 (ii) gives the desired conclusion. \square

Remark 4.4. The condition (4.18) was proposed by Xu in [56], while (4.19) is due to Wittmann [55]. As pointed out in [57, Examples 3.1 and 3.2], these two conditions are not comparable in the sense that neither of them implies the other. Note also that the two conditions above can be interpreted as discrete versions of (2.14) and (2.15) and respectively.

To illustrate the theoretical results and to derive explicit convergence rates, we consider first the case $\alpha_k := \frac{\alpha-1}{(k+b)^q}$ for every $k \geq 0$, where $0 < q < 1$, $\alpha > 1$ and $b > 0$.

Theorem 4.5. Let $0 < q < 1$, $\alpha > 1$, $b > 0$ with $\alpha - 1 < b^q$, and $\xi \in \text{Fix}T$. For given $x_0, v \in \mathcal{H}$, let $\{x_k\}_{k \geq 0}$ be the sequence generated by

$$x_{k+1} := \frac{\alpha-1}{(k+b)^q} v + \left(1 - \frac{\alpha-1}{(k+b)^q}\right) T(x_k) \quad \forall k \geq 0.$$

Then, the following statement are true:

(i) for every $k \geq k_1 := \max \left\{1, \left\lceil \left(\frac{2}{\alpha-1}\right)^{\frac{1}{1-q}} - b \right\rceil\right\}$ it holds

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \frac{1}{k+b+1} \max \left\{ (k_1+b) \|x_{k_1} - x_{k_1-1}\|, \frac{4qD_0(x_0, v, \xi)(b+2)}{b} \right\}; \\ \|x_{k+1} - T(x_{k+1})\| &\leq \frac{1}{k+b+1} \max \left\{ (k_1+b) \|x_{k_1} - x_{k_1-1}\|, \frac{4qD_0(x_0, v, \xi)(b+2)}{b} \right\} + \frac{2(\alpha-1)D_0(x_0, v, \xi)}{(k+b)^q}; \end{aligned}$$

(ii) the sequence $\{x_k\}_{k \geq 0}$ converges strongly to $\xi_* := P_{\text{Fix}T}(v)$ as $k \rightarrow +\infty$.

Proof. Since $0 < q < 1$, the function $t \mapsto t^q$ is nondecreasing and concave on $(0, +\infty)$. We thus have $0 \leq (k+b)^q - (k+b-1)^q \leq q(k+b-1)^{q-1}$ for every $k \geq 1$. Therefore, according to (4.15) in Proposition 4.2 (i), we have for every $k \geq 1$ that

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \left(1 - \frac{\alpha-1}{(k+b)^q}\right) \|x_k - x_{k-1}\| + 2D_0(x_0, v, \xi) \left| \frac{\alpha-1}{(k+b)^q} - \frac{\alpha-1}{(k+b-1)^q} \right| \\ &\leq \left(1 - \frac{\alpha-1}{(k+b)^q}\right) \|x_k - x_{k-1}\| + \frac{2q(\alpha-1)D_0(x_0, v, \xi)}{(k+b)^q(k+b-1)}, \end{aligned}$$

and the statements follow from Lemma A.8 and (4.16), respectively. As $\|x_k - T(x_k)\| \rightarrow 0$ as $k \rightarrow +\infty$, the strong convergence of the sequence of iterates follows from Proposition 4.2 (ii). \square

There are several reasons for focusing on the case $q = 1$. One reason, of course, is to derive the discrete counterpart of the theorem 2.9 which gave the best rates of convergence in the continuous setting. Additionally, our approach in this critical case offers an alternative proof for the result by Lieder in [40], which corresponds to the scenario where $\alpha_k = \frac{1}{k+2}$ for every $k \geq 0$. Moreover, our method allows v to vary within \mathcal{H} instead of fixing it as x_0 .

Theorem 4.6. Let $\alpha > 1$ and $\xi \in \text{Fix}T$. For given $x_0, v \in \mathcal{H}$, let $\{x_k\}_{k \geq 0}$ be the sequence generated by

$$x_{k+1} := \frac{\alpha - 1}{k + \alpha} v + \left(1 - \frac{\alpha - 1}{k + \alpha}\right) T(x_k) \quad \forall k \geq 0.$$

Then, the following statements are true:

(i) for every $k \geq 1$ it holds

• if $\alpha > 2$, then

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \frac{1}{k + \alpha + 1} \max \left\{ (\alpha + 1) \|x_1 - x_0\|, \frac{2(\alpha - 1)(\alpha + 2) D_0(x_0, v, \xi)}{(\alpha - 2)\alpha} \right\}; \\ \|x_{k+1} - T(x_{k+1})\| &\leq \frac{1}{k + \alpha + 1} \max \left\{ (\alpha + 1) \|x_1 - x_0\|, \frac{2(\alpha - 1)(\alpha + 2) D_0(x_0, v, \xi)}{(\alpha - 2)\alpha} \right\} + \frac{2(\alpha - 1) D_0(x_0, v, \xi)}{k + \alpha}; \end{aligned}$$

• if $\alpha = 2$, then

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \frac{1}{k + 2} \sqrt{\frac{2(6 + \pi^2)}{3} D_0(x_0, v, \xi)^2 + \|v + T(x_0) - 2x_0\|^2 - \|T(x_0) - v\|^2}; \\ \|x_{k+1} - T(x_{k+1})\| &\leq \frac{1}{k + 2} \sqrt{\frac{2(6 + \pi^2)}{3} D_0(x_0, v, \xi)^2 + \|v + T(x_0) - 2x_0\|^2 - \|T(x_0) - v\|^2} + \frac{2D_0(x_0, v, \xi)}{k + 2}; \end{aligned}$$

• if $1 < \alpha < 2$, then

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \frac{1}{(k + \alpha + 1)^{\alpha - 1}} \max \left\{ (\alpha + 1)^{\alpha - 1} \|x_1 - x_0\|, \frac{8(\alpha^2 - 1)D_0(x_0, v, \xi)}{\alpha^2(2 - \alpha)(\alpha + 2)^{2 - \alpha}} \right\}; \\ \|x_{k+1} - T(x_{k+1})\| &\leq \frac{1}{(k + \alpha + 1)^{\alpha - 1}} \max \left\{ (\alpha + 1)^{\alpha - 1} \|x_1 - x_0\|, \frac{8(\alpha^2 - 1)D_0(x_0, v, \xi)}{\alpha^2(2 - \alpha)(\alpha + 2)^{2 - \alpha}} \right\} + \frac{2(\alpha - 1) D_0(x_0, v, \xi)}{k + \alpha}; \end{aligned}$$

(ii) the sequence $\{x_k\}_{k \geq 0}$ converges strongly to $\xi_* := \text{P}_{\text{Fix}T}(v)$ as $k \rightarrow +\infty$.

Proof. For every $k \geq 1$, we have from (4.15) in Proposition 4.2 (ii) that

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \left(1 - \frac{\alpha - 1}{k + \alpha}\right) \|x_k - x_{k-1}\| + 2D_0(x_0, v, \xi) \left| \frac{\alpha - 1}{k + \alpha} - \frac{\alpha - 1}{k + \alpha - 1} \right| \\ &\leq \left(1 - \frac{\alpha - 1}{k + \alpha}\right) \|x_k - x_{k-1}\| + \frac{2(\alpha - 1) D_0(x_0, v, \xi)}{(k + \alpha)(k + \alpha - 1)}. \end{aligned}$$

For $\alpha \neq 2$, the estimates for the discrete velocity and the fixed point residual follow from Lemma A.9 and (4.16).

For $\alpha = 2$, Lemma A.9 would give rates of convergence of $\mathcal{O}\left(\frac{\log k}{k}\right)$ as $k \rightarrow +\infty$. In the following we will see that we can get rid of the $\log k$ term by a different approach, inspired by the continuous time analysis.

To fix the setting, the iteration (4.10) is in this case for every $k \geq 0$

$$x_{k+1} := \frac{1}{k + 2} v + \frac{k + 1}{k + 2} T(x_k), \quad (4.20)$$

which also implies

$$T(x_k) - x_{k+1} = \frac{1}{k + 1} (x_{k+1} - v). \quad (4.21)$$

For every $k \geq 1$, we consider the difference

$$\begin{aligned} &\frac{1}{2} (k + 2)^2 \|x_{k+1} - x_k\|^2 - \frac{1}{2} (k + 1)^2 \|x_k - x_{k-1}\|^2 \\ &= \langle (k + 1)(x_k - x_{k-1}), (k + 2)(x_{k+1} - x_k) - (k + 1)(x_k - x_{k-1}) \rangle \\ &\quad + \frac{1}{2} \|(k + 2)(x_{k+1} - x_k) - (k + 1)(x_k - x_{k-1})\|^2. \end{aligned} \quad (4.22)$$

By making use of the formulas (4.20) and (4.21), respectively, we observe that for every $k \geq 1$

$$\begin{aligned} &(k + 2)(x_{k+1} - x_k) - (k + 1)(x_k - x_{k-1}) \\ &= v + (k + 1)T(x_k) - (k + 2)x_k - v - kT(x_{k-1}) + (k + 1)x_{k-1} \end{aligned}$$

$$\begin{aligned}
&= -(k+1) \left((x_k - T(x_k)) - (x_{k-1} - T(x_{k-1})) \right) - x_k + T(x_{k-1}) \\
&= -(k+1) \left((x_k - T(x_k)) - (x_{k-1} - T(x_{k-1})) \right) + \frac{1}{k} (x_k - v).
\end{aligned}$$

Plugging this relation into (4.22), we deduce that for every $k \geq 1$

$$\begin{aligned}
&\frac{1}{2} (k+2)^2 \|x_{k+1} - x_k\|^2 - \frac{1}{2} (k+1)^2 \|x_k - x_{k-1}\|^2 \\
&= \left\langle (k+1) (x_k - x_{k-1}), -(k+1) \left((x_k - T(x_k)) - (x_{k-1} - T(x_{k-1})) \right) + \frac{1}{k} (x_k - v) \right\rangle \\
&\quad + \frac{1}{2} \left\| -(k+1) \left((x_k - T(x_k)) - (x_{k-1} - T(x_{k-1})) \right) + \frac{1}{k} (x_k - v) \right\|^2 \\
&= -(k+1)^2 \langle x_k - x_{k-1}, (x_k - T(x_k)) - (x_{k-1} - T(x_{k-1})) \rangle + \frac{k+1}{k} \langle x_k - x_{k-1}, x_k - v \rangle \\
&\quad + \frac{1}{2} (k+1)^2 \|(x_k - T(x_k)) - (x_{k-1} - T(x_{k-1}))\|^2 \\
&\quad - \frac{k+1}{k} \langle (x_k - T(x_k)) - (x_{k-1} - T(x_{k-1})), x_k - v \rangle + \frac{1}{2k^2} \|x_k - v\|^2 \\
&\leq \frac{k+1}{k} \langle T(x_k) - T(x_{k-1}), x_k - v \rangle + \frac{1}{2k^2} \|x_k - v\|^2,
\end{aligned}$$

where the last inequality comes from the $\frac{1}{2}$ -cocoercivity of $\text{Id} - T$, see (4.2). Furthermore, by taking into consideration again the definition of iteration x_k , we have for every $k \geq 1$

$$\begin{aligned}
\frac{k+1}{k} \langle T(x_k) - T(x_{k-1}), x_k - v \rangle &= \langle T(x_k) - T(x_{k-1}), T(x_{k-1}) - v \rangle \\
&= -\frac{1}{2} \|T(x_k) - T(x_{k-1})\|^2 - \frac{1}{2} \|T(x_{k-1}) - v\|^2 + \frac{1}{2} \|T(x_k) - v\|^2 \\
&\leq -\frac{1}{2} \|T(x_{k-1}) - v\|^2 + \frac{1}{2} \|T(x_k) - v\|^2,
\end{aligned}$$

Therefore, by also utilizing Lemma 4.1, we conclude that for every $k \geq 1$

$$\begin{aligned}
\frac{1}{2} (k+2)^2 \|x_{k+1} - x_k\|^2 - \frac{1}{2} (k+1)^2 \|x_k - x_{k-1}\|^2 &\leq \frac{1}{2} \|T(x_k) - v\|^2 - \frac{1}{2} \|T(x_{k-1}) - v\|^2 + \frac{1}{2k^2} \|x_k - v\|^2 \\
&\leq \frac{1}{2} \|T(x_k) - v\|^2 - \frac{1}{2} \|T(x_{k-1}) - v\|^2 + \frac{2D_0(x_0, v, \xi)^2}{k^2}.
\end{aligned}$$

By telescoping, we obtain for every $k \geq 1$

$$\begin{aligned}
\frac{1}{2} (k+2)^2 \|x_{k+1} - x_k\|^2 - 2 \|x_1 - x_0\|^2 &\leq \frac{1}{2} \|T(x_k) - v\|^2 - \frac{1}{2} \|T(x_0) - v\|^2 + 2D_0(x_0, v, \xi)^2 \sum_{i=1}^k \frac{1}{i^2} \\
&\leq \frac{1}{2} \|T(x_k) - v\|^2 - \frac{1}{2} \|T(x_0) - v\|^2 + \frac{\pi^2 D_0(x_0, v, \xi)^2}{3}.
\end{aligned}$$

Using that $2(x_1 - x_0) = v + T(x_0) - 2x_0$, this yields for every $k \geq 1$

$$\begin{aligned}
(k+2)^2 \|x_{k+1} - x_k\|^2 &\leq \|T(x_k) - v\|^2 + \frac{2\pi^2 D_0(x_0, v, \xi)^2}{3} + \|v + T(x_0) - 2x_0\|^2 - \|T(x_0) - v\|^2 \\
&\leq \frac{2(6 + \pi^2)}{3} D_0(x_0, v, \xi)^2 + \|v + T(x_0) - 2x_0\|^2 - \|T(x_0) - v\|^2. \tag{4.23}
\end{aligned}$$

where last inequality follows from (4.12) in Lemma 4.1. This and (4.16) in Proposition Proposition 4.2 give the estimates in case $\alpha = 2$.

The strong convergence of the sequence $\{x_k\}_{k \geq 0}$ is a direct consequence of Proposition 4.2 (ii). \square

Remark 4.7. From the estimates given in the case of $\alpha = 2$, one can see that it is reasonable to choose $v := x_0$, as Lieder did in [40], since the last two terms in the expression under the square root can be simplified, and therefore the constant depends only on the distance from x_0 to the set $\text{Fix}T$.

4.3 The averaged operator case

The above investigations conducted for nonexpansive operators can be extended to θ -averaged operators $T: \mathcal{H} \rightarrow \mathcal{H}$ with $\theta \in (0, 1]$. This extension provides greater flexibility in the considered approaches.

A single-valued operator $T: \mathcal{H} \rightarrow \mathcal{H}$ is said to be θ -averaged if there exists a nonexpansive operator $N: \mathcal{H} \rightarrow \mathcal{H}$ such that

$$T = (1 - \theta) \text{Id} + \theta N.$$

In this case, for every $0 \leq \lambda \leq \frac{1}{\theta}$ the operator $T_\lambda := (1 - \lambda) \text{Id} + \lambda T$ is nonexpansive. Indeed, T_λ can be written as

$$T_\lambda = (1 - \lambda) \text{Id} + \lambda T = (1 - \lambda) \text{Id} + \lambda ((1 - \theta) \text{Id} + \theta N) = (1 - \lambda\theta) \text{Id} + \lambda\theta N,$$

which is obviously nonexpansive. Moreover, it holds $\|x - T(x)\| = \frac{1}{\lambda} \|x - T_\lambda(x)\|$ for every $x \in \mathcal{H}$ and $\text{Fix}T = \text{Fix}T_\lambda$. This shows that, in order to obtain an iterative method that detects the fixed points of T , one simply has to replace T by T_λ in the iterative scheme (4.10), without affecting the convergence results.

Let $v, x_0 \in \mathcal{H}$ and $0 < \lambda \leq \frac{1}{\theta}$. For every $k \geq 0$ set

$$x_{k+1} := \alpha_k v + (1 - \alpha_k) ((1 - \lambda) x_k + \lambda T(x_k)). \quad (4.24)$$

According to Theorem 4.3, the sequence $\{x_k\}_{k \geq 0}$ strongly converges to $\xi_* := \text{P}_{\text{Fix}T}(v)$ as $k \rightarrow +\infty$, provided that $\{\alpha_k\}_{k \geq 0}$ satisfies one of the conditions (4.18) or (4.19) in addition to (4.9). Convergence rates for the discrete velocity and the fixed point residual can be straightforwardly derived from Theorem 4.5 for $\alpha_k := \frac{\alpha-1}{(k+b)^q}$ for every $k \geq 0$, where $0 < q < 1$, $\alpha > 1$ and $b > 0$, and from Theorem 4.6 for $\alpha_k := \frac{\alpha-1}{k+\alpha}$ for every $k \geq 0$, where $\alpha > 1$.

Regarding the strong convergence of the iterates to $\xi_* := \text{P}_{\text{Fix}T}(v)$, we will see below that in the case $0 < \lambda < \frac{1}{\theta}$ this is guaranteed only by assuming that the sequence $\{\alpha_k\}_{k \geq 0}$ satisfies (4.9). The proof adapts an idea used in [22, 52].

Theorem 4.8. *Let $0 < \lambda < \frac{1}{\theta}$ and assume that the sequence $\{\alpha_k\}_{k \geq 0}$ satisfies only (4.9). For given $x_0, v \in \mathcal{H}$, let $\{x_k\}_{k \geq 0}$ be the sequence generated by (4.24). Then, the sequence $\{x_k\}_{k \geq 0}$ strongly converges to $\xi_* := \text{P}_{\text{Fix}T}(v)$ as $k \rightarrow +\infty$.*

Proof. By definition, there exists a nonexpansive operator $N: \mathcal{H} \rightarrow \mathcal{H}$ such that

$$(1 - \lambda) \text{Id} + \lambda T = (1 - \lambda) \text{Id} + \lambda ((1 - \theta) \text{Id} + \theta N) = (1 - \lambda\theta) \text{Id} + \lambda\theta N.$$

In particular, the iteration (4.24) becomes

$$x_{k+1} = \alpha_k v + (1 - \alpha_k) ((1 - \lambda\theta) x_k + \lambda\theta N(x_k)) \quad \forall k \geq 0. \quad (4.25)$$

From Lemma 4.1, we have for every $k \geq 0$

$$\|x_k - \xi_*\| \leq D_0(x_0, v, \xi_*),$$

which, by the triangle inequality and the fact that $\xi_* \in \text{Fix}T = \text{Fix}N$, leads to

$$\|N(x_k)\| \leq \|N(x_k) - N(\xi_*)\| + \|\xi_*\| \leq \|x_k - \xi_*\| + \|\xi_*\| \leq D_0(x_0, v, \xi_*) + \|\xi_*\|. \quad (4.26)$$

This shows that both sequences $\{x_k\}_{k \geq 0}$ and $\{N(x_k)\}_{k \geq 0}$ are bounded.

For every $k \geq 0$, we define

$$\begin{aligned} \theta_k &:= \alpha_k + (1 - \alpha_k) \lambda\theta \in (0, 1), \\ w_k &:= \frac{1}{\alpha_k + (1 - \alpha_k) \lambda\theta} (\alpha_k v + (1 - \alpha_k) \lambda\theta N(x_k)). \end{aligned}$$

Obviously, the sequence $\{w_k\}_{k \geq 0}$ is also bounded. Furthermore, $1 - \theta_k = (1 - \alpha_k)(1 - \lambda\theta)$ and $\lim_{k \rightarrow +\infty} \theta_k = \lambda\theta \in (0, 1)$. For every $k \geq 0$, we have

$$w_k = \frac{1}{\theta_k} (\alpha_k v + (1 - \alpha_k) \lambda\theta N(x_k)) = \frac{1}{\theta_k} (x_{k+1} - (1 - \theta_k) x_k)$$

or, equivalently,

$$x_{k+1} = (1 - \theta_k) x_k + \theta_k w_k.$$

Some simple calculations give for every $k \geq 0$

$$\begin{aligned} w_{k+1} - w_k &= \frac{1}{\theta_{k+1}} (\alpha_{k+1}v + (1 - \alpha_{k+1})\lambda\theta N(x_{k+1})) - \frac{1}{\theta_k} (\alpha_k v + (1 - \alpha_k)\lambda\theta N(x_k)) \\ &= \left(\frac{\alpha_{k+1}}{\theta_{k+1}} - \frac{\alpha_k}{\theta_k} \right) v + \frac{(1 - \alpha_{k+1})\lambda\theta}{\theta_{k+1}} (N(x_{k+1}) - N(x_k)) + \left(\frac{1 - \alpha_{k+1}}{\theta_{k+1}} - \frac{1 - \alpha_k}{\theta_k} \right) \lambda\theta N(x_k), \end{aligned}$$

therefore, by using that $\theta_k = \alpha_k + (1 - \alpha_k)\lambda\theta \geq (1 - \alpha_k)\lambda\theta$ and (4.26), we obtain the following estimates

$$\begin{aligned} \|w_{k+1} - w_k\| - \|x_{k+1} - x_k\| &\leq \left| \frac{\alpha_{k+1}}{\theta_{k+1}} - \frac{\alpha_k}{\theta_k} \right| \|v\| + \frac{(1 - \alpha_{k+1})\lambda\theta}{\theta_{k+1}} \|N(x_{k+1}) - N(x_k)\| \\ &\quad + \left| \left(\frac{1 - \alpha_{k+1}}{\theta_{k+1}} - \frac{1 - \alpha_k}{\theta_k} \right) \right| \lambda\theta \|N(x_k)\| - \|x_{k+1} - x_k\| \\ &\leq \left| \frac{\alpha_{k+1}}{\theta_{k+1}} - \frac{\alpha_k}{\theta_k} \right| \|v\| + \left(\frac{(1 - \alpha_{k+1})\lambda\theta}{\theta_{k+1}} - 1 \right) \|x_{k+1} - x_k\| \\ &\quad + \left| \left(\frac{1 - \alpha_{k+1}}{\theta_{k+1}} - \frac{1 - \alpha_k}{\theta_k} \right) \right| \lambda\theta \|N(x_k)\| \\ &\leq \left| \frac{\alpha_{k+1}}{\theta_{k+1}} - \frac{\alpha_k}{\theta_k} \right| \|v\| + \left| \left(\frac{1 - \alpha_{k+1}}{\theta_{k+1}} - \frac{1 - \alpha_k}{\theta_k} \right) \right| \lambda\theta (D_0(x_0, v, \xi_*) + \|\xi_*\|). \end{aligned}$$

This guarantees that $\limsup_{k \rightarrow +\infty} (\|w_{k+1} - w_k\| - \|x_{k+1} - x_k\|) \leq 0$, which, by using Lemma A.10, allows us to conclude that $\lim_{k \rightarrow +\infty} \|w_k - x_k\| = 0$. Consequently, $\lim_{k \rightarrow +\infty} \|x_{k+1} - x_k\| = \lim_{k \rightarrow +\infty} \theta_k \|w_k - x_k\| = 0$, which, by using (4.16), further implies $\lim_{k \rightarrow +\infty} \|x_k - T(x_k)\| = 0$. The conclusion follows from Proposition 4.2 (ii). \square

Remark 4.9. Bot, Csetnek, and Meier proposed in [15] for $z_0 \in \mathcal{H}$ the following iterative scheme

$$z_{k+1} := \beta_k z_k + \lambda(T(\beta_k z_k) - \beta_k z_k) \quad \forall k \geq 0, \quad (4.27)$$

with $\{\beta_k\}_{k \geq 0} \subseteq (0, 1]$, $\lim_{k \rightarrow +\infty} \beta_k = 1$, $\sum_{k \geq 0} (1 - \beta_k) = +\infty$ and $\sum_{k \geq 1} |\beta_k - \beta_{k-1}| < +\infty$, and $0 < \lambda \leq \frac{1}{\theta}$, designed with the aim of generating a strongly convergent sequence $\{z_k\}_{k \geq 0}$ to the fixed point of a θ -averaged operator $T : \mathcal{H} \rightarrow \mathcal{H}$ with minimum norm.

By setting $x_k := \beta_k z_k$ for every $k \geq 0$, the iterative scheme (4.27) can be rewritten as

$$x_{k+1} := \beta_{k+1} ((1 - \lambda)x_k + \lambda T(x_k)) \quad \forall k \geq 0, \quad (4.28)$$

which corresponds to (4.24) with $\alpha_k := 1 - \beta_{k+1}$ for every $k \geq 0$, and $v := 0$. Note that the strong convergence of $\{x_k\}_{k \geq 0}$ to $\text{P}_{\text{Fix}T}(0)$ is a direct consequence of Theorem 4.3 in the context of fulfilling condition (4.19). Since $\lim_{k \rightarrow +\infty} \beta_k = 1$, this says nothing more than that the sequence $\{x_k\}_{k \geq 0}$ strongly converges to $\text{P}_{\text{Fix}T}(0)$ as $k \rightarrow +\infty$. Furthermore, if $0 < \lambda \leq \frac{1}{\theta}$, then, according to Theorem 4.8, the same conclusion follows even without assuming that $\sum_{k \geq 1} |\beta_k - \beta_{k-1}| < +\infty$.

In addition, the rates of convergence for $\|x_k - T(x_k)\|$, as provided by the Halpern iteration, can be transferred to the discrete velocity and the fixed point residual expressed in terms of the original sequence $\{z_k\}_{k \geq 0}$, namely, by using the boundedness of the latter and that for every $k \geq 1$, the following holds

$$\|z_{k+1} - z_k\| \leq (1 - \beta_k) \|z_k\| + \lambda \|x_k - T(x_k)\| = \alpha_{k-1} \|z_k\| + \lambda \|x_k - T(x_k)\|$$

and

$$\begin{aligned} \|z_k - T(z_k)\| &\leq \|z_k - x_k\| + \|x_k - T(x_k)\| + \|T(x_k) - T(z_k)\| \\ &\leq 2 \|z_k - x_k\| + \|x_k - T(x_k)\| = (1 - \beta_k) \|z_k\| + \|x_k - T(x_k)\| \\ &= 2\alpha_{k-1} \|z_k\| + \|x_k - T(x_k)\|, \end{aligned}$$

respectively.

Remark 4.10. For $A : \mathcal{H} \rightrightarrows \mathcal{H}$ a (set-valued) maximally monotone operator, the problem of finding a zero of A , this is an element in the set $\text{Zer}A := \{x \in \mathcal{H} : 0 \in A(x)\}$, is for every $\eta > 0$ equivalent to the problem of finding a fixed point of the resolvent $J_{\eta A} : \mathcal{H} \rightarrow \mathcal{H}$, $J_{\eta A} = (\text{Id} + \eta A)^{-1}$, which is a single-valued and $\frac{1}{2}$ -averaged operator. The classical proximal point algorithm, defined for a given $x_0 \in \mathcal{H}$ as

$$x_{k+1} = J_{\eta A}(x_k) \quad \forall k \geq 0,$$

produces a sequence $\{x_k\}_{k \geq 0}$ that weakly converges to a zero of A as $k \rightarrow +\infty$. The fixed point residual $\|J_{\eta A}(x_k) - x_k\|$ vanishes with a rate of convergence of $\mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$ as $k \rightarrow +\infty$.

Gu and Yang have shown in [30] that for $\mathcal{H} := \mathbb{R}^N$, this rate of convergence can be improved to

$$\|J_{\eta A}(x_k) - x_k\| = \begin{cases} \mathcal{O}\left(\frac{1}{k}\right), & \text{if } N = 1, \\ \mathcal{O}\left(\frac{1}{\sqrt{(1 + \frac{1}{k})^k k}}\right), & \text{if } N \geq 2. \end{cases} \quad \text{as } k \rightarrow +\infty.$$

In the setting of finite-dimensional Hilbert spaces, Kim introduced in [36] (see also [44]) the following iterative scheme, which, for given $y_0 = x_{-1} = x_0 \in \mathcal{H}$, for every $k \geq 0$, reads

$$\begin{aligned} y_{k+1} &:= J_{\eta A}(x_k), \\ x_{k+1} &:= y_{k+1} + \frac{k}{k+2}(y_{k+1} - y_k) - \frac{k}{k+2}(y_k - x_{k-1}). \end{aligned} \quad (4.29)$$

Using the performance estimation approach, he proved that this method has a fixed point residual convergence rate of $\mathcal{O}\left(\frac{1}{k}\right)$ as $k \rightarrow +\infty$.

One can note that, for every $k \geq 1$, the second update formula in (4.29) can be equivalently written as

$$\begin{aligned} (k+2)x_{k+1} &= 2(k+1)y_{k+1} - 2ky_k + kx_{k-1}, \\ (k+1)x_k &= 2ky_k - 2(k-1)y_{k-1} + (k-1)x_{k-2}, \\ &\vdots \\ 3x_2 &= 4y_3 - 2y_1 + x_0, \\ 2x_1 &= 2y_1. \end{aligned}$$

Summing these relations, we obtain that for every $k \geq 0$

$$(k+2)x_{k+1} + (k+1)x_k = 2(k+1)y_{k+1} + x_0.$$

In the view of the first equation in (4.29), it yields that the sequence $\{x_k\}_{k \geq 0}$ generated by Kim's algorithm fulfills

$$x_{k+1} = \frac{1}{k+2}x_0 + \frac{k+1}{k+2}(2J_{\eta A}(x_k) - x_k) \quad \forall k \geq 0.$$

Consequently, according to Theorem 4.3 applied to averaged operators, even in infinite-dimensional Hilbert spaces, the sequence $\{x_k\}_{k \geq 0}$ strongly converges to $\text{P}_{\text{Zer}A}(x_0)$ as $k \rightarrow +\infty$. In addition, from Theorem 4.6 in case $\alpha = 2$ it follows that $\|x_{k+1} - x_k\|$ and $\|J_{\eta A}(x_k) - x_k\|$ converge to zero with a rate of convergence of $\mathcal{O}\left(\frac{1}{k}\right)$ as $k \rightarrow +\infty$.

A Appendix

In the appendix, we discuss the asymptotic behaviour of the trajectory generated by (1.2) in case M is an cocoercive operator (see [3, Theorem 11] for a general formulation), and outline additional results that play an essential role in the convergence of the continuous time models.

A.1 The cocoercive case

Theorem A.1. *Let $M: \mathcal{H} \rightarrow \mathcal{H}$ be a ρ -cocoercive operator for some $\rho > 0$, that is*

$$\langle M(x) - M(y), x - y \rangle \geq \rho \|M(x) - M(y)\|^2 \quad \forall x, y \in \mathcal{H},$$

with $\text{Zer}M \neq \emptyset$, and $x: [t_0, +\infty) \rightarrow \mathcal{H}$ be the solution trajectory of

$$\dot{x}(t) + M(x(t)) = 0,$$

with initial condition $x(t_0) := x_0 \in \mathcal{H}$. Then, the following statements are true:

- (i) *it holds $\|M(x(t))\| = o\left(\frac{1}{\sqrt{t}}\right)$ as $t \rightarrow +\infty$;*

(ii) the solution trajectory $x(t)$ converges weakly to an element of $\text{Zer}M$ as $t \rightarrow +\infty$.

Proof. Let $\xi \in \text{Zer}M$. For every $t \geq t_0$, we consider

$$\mathcal{E}(t) := \frac{1}{2} \|x(t) - \xi\|^2.$$

Then, for almost every $t \geq t_0$ it holds

$$\frac{d}{dt}\mathcal{E}(t) = \langle x(t) - \xi, \dot{x}(t) \rangle = -\langle x(t) - \xi, M(x(t)) \rangle.$$

By integration, we get for every $t \geq t_0$

$$\frac{1}{2} \|x(t) - \xi\|^2 \leq \frac{1}{2} \|x(t_0) - \xi\|^2 + \int_{t_0}^t \langle x(s) - \xi, M(x(s)) - M(\xi) \rangle ds \leq \frac{1}{2} \|x(t_0) - \xi\|^2.$$

This gives the boundedness of the trajectory. On the other hand, by invoking the ρ -cocoercivity of M and that $M(\xi) = 0$, and by letting t converge to infinity, we obtain

$$\rho \int_{t_0}^{+\infty} \|M(x(s))\|^2 ds \leq \int_{t_0}^{+\infty} \langle x(s) - \xi, M(x(s)) \rangle du \leq \frac{1}{2} \|x(t_0) - \xi\|^2 < +\infty. \quad (\text{A.1})$$

Therefore, $\liminf_{t \rightarrow +\infty} t \|M(x(t))\|^2 = 0$. By computing the time derivative at almost every $t \geq t_0$ and by making use of the monotonicity of M , we get

$$\begin{aligned} \frac{d}{dt} \left(\frac{1}{2} t \|M(x(t))\|^2 \right) &= \frac{1}{2} \|M(x(t))\|^2 + t \left\langle M(x(t)), \frac{d}{dt}(M(x(t))) \right\rangle \\ &= \frac{1}{2} \|M(x(t))\|^2 - t \left\langle \dot{x}(t), \frac{d}{dt}(M(x(t))) \right\rangle \\ &\leq \frac{1}{2} \|M(x(t))\|^2. \end{aligned}$$

The right-hand side of the above inequality belongs to $\mathbb{L}^1([t_0, +\infty); \mathcal{H})$ thanks to (A.1). According to Lemma A.2, $\lim_{t \rightarrow +\infty} t \|M(x(t))\|^2 \in \mathbb{R}$ exists, and therefore it must be equal to 0. In other words,

$$\|M(x(t))\| = o\left(\frac{1}{\sqrt{t}}\right) \text{ as } t \rightarrow +\infty,$$

which means that statement (i) is proven.

Using that for almost every $t \geq t_0$ it holds

$$\frac{d}{dt} \left(\frac{1}{2} \|x(t) - \xi\|^2 \right) = \langle x(t) - \xi, \dot{x}(t) \rangle = -\langle x(t) - \xi, M(x(t)) \rangle \leq 0,$$

thanks to Lemma A.2, we obtain that the limit $\lim_{t \rightarrow +\infty} \|x(t) - \xi\| \in \mathbb{R}$ exists. This shows that the first condition in Opial's Lemma (see Lemma A.3) is satisfied.

Using that $\lim_{t \rightarrow +\infty} M(x(t)) = 0$ and the sequential closedness of the graph of M in the weak \times strong topology of $\mathcal{H} \times \mathcal{H}$, it yields that every weak limit point of the trajectory $x(t)$ belongs to $\text{Zer}M$. This shows that the second condition in Opial's Lemma is also satisfied.

Consequently, according to Lemma A.3, the trajectory $x(t)$ converges weakly to an element in $\text{Zer}M$ as $t \rightarrow +\infty$, which proves (ii). \square

A.2 Auxiliary results used in the convergence analysis of the continuous time models

The following result and its proof can be referenced in [1, Lemma 5.1].

Lemma A.2. *Let $h: [t_0, +\infty) \rightarrow \mathbb{R}$ be a locally absolutely continuous function that is bounded from below, and $g \in \mathbb{L}^1([t_0, +\infty); \mathbb{R})$ be such that*

$$\dot{h}(t) \leq g(t) \quad \text{for almost every } t \geq t_0.$$

Then the limit $\lim_{t \rightarrow +\infty} h(t) \in \mathbb{R}$ exists.

The continuous form of Opial's Lemma ([43]) serves as the primary tool utilized to establish the weak convergence of the trajectory.

Lemma A.3. *Let \mathcal{S} be a nonempty subset of \mathcal{H} and $x: [t_0, +\infty) \rightarrow \mathcal{H}$. Assume that*

- (i) *for every $\xi \in \mathcal{S}$, $\lim_{t \rightarrow +\infty} \|x(t) - \xi\|$ exists;*
- (ii) *every weak sequential cluster point of the trajectory $x(t)$ as $t \rightarrow +\infty$ belongs to \mathcal{S} .*

Then $x(t)$ converges weakly to a point in \mathcal{S} as $t \rightarrow +\infty$.

Lemma A.4. [7, Lemma A.4] *Take $t_0 > 0$, and let $h \in \mathbb{L}^1([t_0, +\infty))$ be nonnegative and continuous. Consider an nondecreasing function $\gamma: [t_0, +\infty) \rightarrow \mathbb{R}_+$ such that $\lim_{t \rightarrow +\infty} \gamma(t) = +\infty$. Then,*

$$\lim_{t \rightarrow +\infty} \frac{1}{\gamma(t)} \int_{t_0}^t \gamma(r) h(r) = 0.$$

The following result, provided in [24, Lemma 1], is employed in proving the strong convergence of the trajectory.

Lemma A.5. *Let $h: [t_0, +\infty) \rightarrow \mathbb{R}$ be a locally absolutely continuous function, $g: [t_0, +\infty) \rightarrow \mathbb{R}$ a bounded function and $\varepsilon: [t_0, +\infty) \rightarrow \mathbb{R}_+$ a locally integrable function such that*

$$\dot{h}(t) + \varepsilon(t) h(t) \leq \varepsilon(t) g(t) \quad \text{for almost every } t \geq t_0.$$

Then h is also bounded and, if $\int_{t_0}^{+\infty} \varepsilon(t) dt = +\infty$, then $\limsup_{t \rightarrow +\infty} h(t) \leq \limsup_{t \rightarrow +\infty} g(t)$.

Gronwall's Lemma ([29]) in integral form for continuous functions will be utilized to establish the strong convergence of the trajectory to the minimal solution under a certain condition.

Lemma A.6. *Let $h, g: [t_0, +\infty) \rightarrow \mathbb{R}$ be continuous function and $k: [t_0, +\infty) \rightarrow \mathbb{R}$ a function with the property that its negative part is integrable on every closed and bounded subinterval of $[t_0, +\infty)$. If h is non-negative and*

$$h(t) \leq k(t) + \int_{t_0}^t g(u) h(u) du \quad \text{for every } t \geq t_0,$$

then

$$h(t) \leq k(t) + \int_{t_0}^t k(u) g(u) \exp\left(\int_u^t g(r) dr\right) du \quad \text{for every } t \geq t_0.$$

If, in addition, k is nondecreasing, then

$$h(t) \leq k(t) \exp\left(\int_{t_0}^t g(u) du\right) \quad \text{for every } t \geq t_0.$$

A.3 Auxiliary results used in the convergence analysis of the discrete time models

The following result by Xu (see [56, Lemma 2.5]) and will play a crucial role in the proof of the strong convergence of the sequence of iterates.

Lemma A.7. *Let $\{a_k\}_{k \geq 0}$ be a sequence of nonnegative real numbers satisfying for every $k \geq 0$*

$$a_{k+1} \leq (1 - \alpha_k) a_k + \alpha_k b_k + d_k, \tag{A.2}$$

where $\{\alpha_k\}_{k \geq 0}$, $\{b_k\}_{k \geq 0}$, and $\{d_k\}_{k \geq 0}$ are sequences of real numbers with the properties

- (i) $\{\alpha_k\}_{k \geq 0} \subseteq [0, 1]$ and $\sum_{k \geq 0} \alpha_k = +\infty$;
- (ii) $\limsup_{k \rightarrow +\infty} b_k \leq 0$;
- (iii) $\sum_{k \geq 0} d_k < +\infty$.

Then, $\lim_{k \rightarrow +\infty} a_k = 0$.

The following two lemmas are used to derive the convergence rates for the discrete time models.

Lemma A.8. Let $\{\xi_k\}_{k \geq 1}$ be a sequence of nonnegative real numbers satisfying

$$\xi_{k+1} \leq \left(1 - \frac{c}{(k+b)^q}\right) \xi_k + \frac{d}{(k+b)^q (k+b-1)} \quad \forall k \geq 1, \quad (\text{A.3})$$

where $0 < q < 1$, $0 < c < b^q$, and $d \geq 0$. Then, for every $k \geq k_1 := \max\left\{1, \left[\left(\frac{2}{c}\right)^{\frac{1}{1-q}} - b\right]\right\}$ it holds

$$\xi_{k+1} \leq \frac{1}{k+b+1} \max\left\{(k_1+b)\xi_{k_1}, \frac{2d(b+2)}{cb}\right\}.$$

Proof. Multiplying both sides of (A.3) by $k+b+1 > 0$, we get for every $k \geq 1$

$$\begin{aligned} (k+b+1)\xi_{k+1} &\leq (k+b+1) \left(1 - \frac{c}{(k+b)^q}\right) \xi_k + \frac{d}{(k+b)^q} \frac{k+b+1}{k+b-1} \\ &\leq (k+b+1) \left(1 - \frac{c}{(k+b)^q}\right) \xi_k + \frac{d}{(k+b)^q} \frac{b+2}{b}. \end{aligned} \quad (\text{A.4})$$

Since $0 < q < 1$, we have

$$\begin{aligned} (k+b+1) \left(1 - \frac{c}{(k+b)^q}\right) &= k+b+1 - c(k+b)^{1-q} - \frac{c}{(k+b)^q} \\ &\leq k+b - \frac{c}{2}(k+b)^{1-q} + 1 - \frac{c}{2}(k+b)^{1-q} \\ &\leq k+b - \frac{c}{2}(k+b)^{1-q} \\ &= \left(1 - \frac{c}{2(k+b)^q}\right) (k+b), \end{aligned}$$

where the second inequality holds for $k \geq k_1 := \max\left\{1, \left[\left(\frac{2}{c}\right)^{\frac{1}{1-q}} - b\right]\right\}$. Under these premises, we have for every $k \geq k_1$

$$\begin{aligned} (k+b+1)\xi_{k+1} &\leq \left(1 - \frac{c}{2(k+b)^q}\right) (k+b)\xi_k + \frac{d}{(k+b)^q} \frac{b+2}{b} \\ &= \left(1 - \frac{c}{2(k+b)^q}\right) (k+b)\xi_k + \frac{c}{2(k+b)^q} \frac{2d(b+2)}{cb}. \end{aligned}$$

By induction, it follows that for every $k \geq k_1$

$$(k+b+1)\xi_{k+1} \leq \max\left\{(k_1+b)\xi_{k_1}, \frac{2d(b+2)}{cb}\right\}.$$

□

Lemma A.9. Let $\{\xi_k\}_{k \geq 1}$ be a sequence of nonnegative real numbers satisfying

$$\xi_{k+1} \leq \left(1 - \frac{c}{k+b}\right) \xi_k + \frac{d}{(k+b)(k+b-1)} \quad \forall k \geq 1, \quad (\text{A.5})$$

where $0 < c < b$ and $d \geq 0$. Then, the following statements are true for every $k \geq 1$:

(i) if $c > 1$, then

$$\xi_{k+1} \leq \frac{1}{k+b+1} \max\left\{(b+1)\xi_1, \frac{d(b+2)}{(c-1)b}\right\};$$

(ii) if $c = 1$, then

$$\xi_{k+1} \leq \frac{(1+b)\xi_1 + 3d}{k+b+1} + 3d \frac{\log k}{k+b+1};$$

(iii) if $0 < c < 1$, then

$$\xi_{k+1} \leq \frac{1}{(k+b+1)^c} \left((b+1)^c \xi_1 + \frac{d(b+1)(b+3-c)}{b^2(b+2)^{1-c}(1-c)} \right).$$

Proof. Multiplying both sides of (A.5) by $k + b + 1 > 0$, we get for every $k \geq 1$

$$\begin{aligned} (k + b + 1) \xi_{k+1} &\leq (k + b + 1) \left(1 - \frac{c}{k + b}\right) \xi_k + \frac{d(k + b + 1)}{(k + b)(k + b - 1)} \\ &\leq (k + b + 1) \left(1 - \frac{c}{k + b}\right) \xi_k + \frac{d(b + 2)}{(k + b)b}. \end{aligned}$$

Observing that

$$(k + b + 1) \left(1 - \frac{c}{k + b}\right) = (k + b) \left(1 + \frac{1}{k + b}\right) \left(1 - \frac{c}{k + b}\right) = (k + b) \left(1 - \frac{c - 1}{k + b} - \frac{c}{(k + b)^2}\right),$$

we obtain for every $k \geq 1$

$$(k + b + 1) \xi_{k+1} \leq \left(1 - \frac{c - 1}{k + b} - \frac{c}{(k + b)^2}\right) (k + b) \xi_k + \frac{d(b + 2)}{(k + b)b}. \quad (\text{A.6})$$

(i) *The case $c > 1$.* From (A.6) we deduce that

$$(k + b + 1) \xi_{k+1} \leq \left(1 - \frac{c - 1}{k + b}\right) (k + b) \xi_k + \frac{(b + 2)d}{b(k + b)} = \left(1 - \frac{c - 1}{k + b}\right) (k + b) \xi_k + \frac{c - 1}{k + b} \frac{d(b + 2)}{(c - 1)b} \quad \forall k \geq 1,$$

and the conclusion follows, once again, by induction arguments.

(ii) *The case $c = 1$.* The inequality (A.6) now implies that

$$(k + b + 1) \xi_{k+1} \leq (k + b) \xi_k + \frac{3d}{k} \quad \forall k \geq 1.$$

For every integer $K \geq 1$, by summing up the inequality above for $k = 1, \dots, K$, we obtain

$$\begin{aligned} (K + b + 1) \xi_{K+1} &\leq (1 + b) \xi_1 + 3d \sum_{k=1}^K \frac{1}{k} \leq (1 + b) \xi_1 + 3d \left(1 + \int_1^K \frac{1}{t} dt\right) \\ &= (1 + b) \xi_1 + 3d(1 + \log K) \\ &\leq (1 + b) \xi_1 + 3d + 3d \log K. \end{aligned}$$

This ensures that for every $k \geq 1$

$$\xi_{k+1} \leq \frac{(1 + b) \xi_1 + 3d}{k + b + 1} + 3d \frac{\log k}{k + b + 1}.$$

(iii) *The case $0 < c < 1$.* Multiplying both sides of (A.5) by $(k + b + 1)^c$, we get for every $k \geq 1$

$$(k + b + 1)^c \xi_{k+1} \leq (k + b + 1)^c \left(1 - \frac{c}{k + b}\right) \xi_k + \frac{d(k + b + 1)^c}{(k + b)(k + b - 1)}. \quad (\text{A.7})$$

Using that $0 \leq (t + 1)^c - t^c \leq ct^{c-1}$ for every $t \geq 0$, we obtain

$$(k + b + 1)^c \leq (k + b)^c + c(k + b)^{c-1} = (k + b)^c \left(1 + \frac{c}{k + b}\right) \quad \forall k \geq 1.$$

Under these premises, from (A.7) we obtain for every $k \geq 1$

$$\begin{aligned} (k + b + 1)^c \xi_{k+1} &\leq (k + b)^c \left(1 + \frac{c}{k + b}\right) \left(1 - \frac{c}{k + b}\right) \xi_k + \frac{(b + 2)^2 d}{b(b + 1)(k + b + 1)^{2-c}} \\ &\leq (k + b)^c \xi_k + \frac{d(b + 1)(b + 2)}{b^2(k + b + 1)^{2-c}}. \end{aligned}$$

Once again, by the telescoping sum argument, we obtain for every $k \geq 1$

$$(k + b + 1)^c \xi_{k+1} \leq (b + 1)^c \xi_1 + \frac{d(b + 1)(b + 2)}{b^2} \sum_{i=1}^k \frac{1}{(i + b + 1)^{2-c}}$$

$$\begin{aligned}
&\leq (b+1)^c \xi_1 + \frac{d(b+1)(b+2)}{b^2} \left(\frac{1}{(b+2)^{2-c}} + \int_1^k \frac{1}{(x+b+1)^{2-c}} dx \right) \\
&\leq (b+1)^c \xi_1 + \frac{d(b+1)(b+2)}{b^2} \left(\frac{1}{(b+2)^{2-c}} + \frac{1}{(b+2)^{1-c}(1-c)} \right) \\
&= (b+1)^c \xi_1 + \frac{d(b+1)(b+3-c)}{b^2(b+2)^{1-c}(1-c)}.
\end{aligned}$$

□

The following result is [51, Lemma 2.2] and plays an important role in the proof of the strong convergence of the Halpern iteration in the case of averaged operators.

Lemma A.10. *Let $\{x_k\}_{k \geq 0}$ and $\{w_k\}_{k \geq 0}$ be bounded sequences in \mathcal{H} and $\{\delta_k\}_{k \geq 0}$ be a sequence in $[0, 1]$ with $0 < \liminf_{k \rightarrow +\infty} \delta_k \leq \limsup_{k \rightarrow +\infty} \delta_k < 1$. If $x_{k+1} = (1 - \delta_k)x_k + \delta_k w_k$ for every $k \geq 0$ and*

$$\limsup_{k \rightarrow +\infty} (\|w_{k+1} - w_k\| - \|y_{k+1} - x_k\|) \leq 0,$$

then $\lim_{k \rightarrow +\infty} \|w_k - y_k\| = 0$.

Acknowledgements. The authors are thankful to Ernő Robert Csetnek (University of Vienna) for comments and suggestions which improved the quality of the manuscript.

References

- [1] **B. Abbas, H. Attouch, B.F. Svaiter.** *Newton-like dynamics and forward-backward methods for structured monotone inclusions in Hilbert spaces.* Journal of Optimization Theory and Applications 161(2), 331–360 (2014)
- [2] **H. Attouch, J.-B. Baillon.** *Weak versus strong convergence of a regularized Newton dynamic for maximal monotone operators.* Vietnam Journal of Mathematics 46, 177–195 (2018)
- [3] **H. Attouch, R. I. Boţ, D.-K. Nguyen.** *Fast convex optimization via time scale and averaging of the steepest descent.* [arXiv:2208.08260](https://arxiv.org/abs/2208.08260)
- [4] **H. Attouch, R. I. Boţ, D.-K. Nguyen.** *Fast convex optimization via closed-loop time scaling of gradient dynamics.* [arXiv:2301.00701](https://arxiv.org/abs/2301.00701)
- [5] **H. Attouch, Z. Chbani, J. Peypouquet, P. Redont.** *Fast convergence of inertial dynamics and algorithms with asymptotic vanishing viscosity.* Mathematical Programming 168(1), 123–175 (2018)
- [6] **H. Attouch, R. Cominetti.** *A dynamical approach to convex minimization coupling approximation with the steepest descent method.* Journal of Differential Equations 128(2), 519–540 (1996)
- [7] **H. Attouch, J. Peypouquet.** *Convergence of inertial dynamics and proximal algorithms governed by maximally monotone operators.* Mathematical Programming 174(1-2), 391–432 (2019)
- [8] **H. Attouch, J. Peypouquet, P. Redont.** *Fast convex optimization via inertial dynamics with Hessian driven damping.* Journal of Differential Equations 261(10), 5734–5783 (2016)
- [9] **J.-B. Baillon, H. Brézis.** *Une remarque sur le comportement asymptotique des semigroupes non linéaires,* Houston Journal of Mathematics 2(1), 5–7 (1976)
- [10] **J.-B. Baillon, R. E. Bruck.** *The rate of asymptotic regularity is $o(1/n)$.* In: Theory and Applications of Nonlinear Operators of Accretive and Monotone Types, Lecture Notes in Pure and Applied Mathematics 178, 51–81, Dekker, New York (1996)
- [11] **H. H. Bauschke, P. L. Combettes.** *Convex Analysis and Monotone Operator Theory in Hilbert Spaces.* CMS Books in Mathematics, Springer, New York (2017)
- [12] **J. Borwein, S. Reich, I. Shafrir.** *Krasnoselski-Mann iterations in normed spaces.* Canadian Mathematical Bulletin 35(1), 21–28 (1992)
- [13] **R. I. Boţ, E. R. Csetnek.** *A dynamical system associated with the fixed points set of a nonexpansive operator.* Journal of Dynamics and Differential Equations 29, 155–168 (2017)

- [14] **R. I. Boř, E. R. Csetnek, D.-K. Nguyen.** *Fast Optimistic Gradient Descent Ascent (OGDA) method in continuous and discrete time.* Foundations of Computational Mathematics, DOI: 10.1007/s10208-023-09636-5 (2023)
- [15] **R. I. Boř, E. R. Csetnek, D. Meier.** *Inducing strong convergence into the asymptotic behaviour of proximal splitting algorithms in Hilbert spaces.* Optimization Methods and Software 34(3), 489–514 (2019)
- [16] **R. I. Boř, S.-M. Grad, D. Meier, M. Staudigl.** *Inducing strong convergence of trajectories in dynamical systems associated to monotone inclusions with composite structure.* Advances in Nonlinear Analysis 10(1), 450–476 (2021)
- [17] **R. I. Boř, D. Meier.** *A strongly convergent Krasnosel’skii-Mann-type algorithm for finding a common fixed point of a countably infinite family of nonexpansive operators in Hilbert spaces.* Journal of Computational and Applied Mathematics 395, Article number: 113589 (2021)
- [18] **R. I. Boř, D.-K. Nguyen.** *Fast Krasnosel’skii-Mann algorithm with a convergence rate of the fixed point iteration of $o(1/k)$.* SIAM Journal on Numerical Analysis 61(6), 2813–2843 (2023)
- [19] **M. Bravo, R. Cominetti.** *Sharp convergence rates for averaged nonexpansive maps.* Israel Journal of Mathematics 227, 163–188 (2018)
- [20] **H. Brézis.** *Opérateurs Maximaux Monotones dans les Espaces de Hilbert et Équations d’Évolution.* Lecture Notes 5, North Holland, Amsterdam (1972)
- [21] **F. E. Browder, W. V. Petryshyn.** *The solution by iteration of nonlinear functional equations in Banach spaces.* Bulletin of the American Mathematical Society 72, 571–575 (1966)
- [22] **C. E. Chidume, C. O. Chidume.** *Iterative approximation of fixed points of nonexpansive mappings.* Journal of Mathematical Analysis and Applications 318(1), 288–295 (2006)
- [23] **J. P. Contreras, R. Cominetti.** *Optimal error bounds for nonexpansive fixed-point iterations in normed spaces.* Mathematical Programming 199, 343–374 (2023)
- [24] **R. Cominetti, J. Peypouquet, S. Sorin.** *Strong asymptotic convergence of evolution equations governed by maximal monotone operators with Tikhonov regularization.* Journal of Differential Equations 245(12), 3753–3763 (2008)
- [25] **R. Cominetti, J. A. Soto, J. Vaisman.** *On the rate of convergence of Krasnosel’skii-Mann iterations and their connection with sums of Bernoullis.* Israel Journal of Mathematics 199(2), 757–772 (2014)
- [26] **D. Davis, W. Yin.** *Convergence rate analysis of several splitting schemes.* In: **R. Glowinski, S. J. Osher, W. Yin,** Splitting Methods in Communication, Imaging, Science, and Engineering, 115–163. Springer (2016)
- [27] **Q.-L. Dong, Y. J. Cho, S. He, P. M. Pardalos, T. M. Rassias.** *The Krasnosel’skii-Mann Iterative Method: Recent Progress and Applications.* SpringerBriefs in Optimization, Springer, Cham (2022)
- [28] **C. W. Groetsch.** *A note on segmenting Mann iterates.* Journal of Mathematical Analysis and Applications 40, 369–372 (1972)
- [29] **T.H. Gronwall** *Note on the derivatives with respect to a parameter of the solutions of a system of differential equations.* Annals of Mathematics 20(2), 292–296 (1919)
- [30] **G. Gu, J. Yang.** *Tight sublinear convergence rate of the proximal point algorithm for maximal monotone inclusion problems.* SIAM Journal on Optimization 30(3), 1905–1921 (2020)
- [31] **B. Halpern.** *Fixed points of nonexpanding maps.* Bulletin of the American Mathematical Society 73(6), 957–961 (1967)
- [32] **A. Haraux.** *Systèmes Dynamiques Dissipatifs et Applications,* Recherches en Mathématiques Appliquées 17, Masson, Paris (1991)
- [33] **S. He, H.-K. Xu, Q.-L. Dong, N. Mei.** *Convergence analysis of the Halpern iteration with adaptive anchoring parameters.* Mathematics of Computation 93, 327–345 (2024).
- [34] **S. Ishikawa.** *Fixed points and iteration of a nonexpansive mapping in a Banach space.* Proceedings of the American Mathematical Society 59(1), 65–71 (1976)

- [35] **M.M. Israel Jr., S. Reich.** *Asymptotic behavior of solutions of a nonlinear evolution equation.* Journal of Mathematical Analysis and Applications 83(1), 43–53 (1981)
- [36] **D. Kim.** *Accelerated proximal point method for maximally monotone operators.* Mathematical Programming 190, 57–87 (2021)
- [37] **M. A. Krasnosel’skiĭ.** *Two remarks on the method of successive approximations.* Uspekhi Matematicheskikh Nauk 10, 123–127 (1955)
- [38] **S. Lee, D. Kim.** *Fast extra gradient methods for smooth structured nonconvex-nonconcave minimax problems.* Advances in Neural Information Processing Systems 34 (2021)
- [39] **J. Liang, J. Fadili, G. Peyré.** *Convergence rates with inexact non-expansive operators.* Mathematical Programming 159(1–2), 403–434 (2016)
- [40] **F. Lieder.** *On the convergence rate of the Halpern-iteration.* Optimization Letters 15(2), 405–418 (2021)
- [41] **S.-Y. Matsushita.** *On the convergence rate of the Krasnosel’skiĭ-Mann iteration.* Bulletin of the Australian Mathematical Society 96(1), 162–170 (2017)
- [42] **J. J. Maulén, I. Fierro, J. Peypouquet.** *Inertial Krasnoselskii-Mann iterations.* [arXiv:2210.03791](https://arxiv.org/abs/2210.03791)
- [43] **Z. Opial.** *Weak convergence of the sequence of successive approximations for nonexpansive mappings.* Bulletin of the American Mathematical Society 73, 591–597 (1967)
- [44] **J. Park, E. K. Ryu.** *Exact optimal accelerated complexity for fixed-point iterations.* Proceedings of the 39th International Conference on Machine Learning, PMLR 162 (2022)
- [45] **H. Qi, H.-K. Xu.** *Convergence of Halpern’s iteration method with applications in optimization.* Numerical Functional Analysis and Optimization 42:15, 1839–1854 (2021)
- [46] **S. Reich.** *Weak convergence theorems for nonexpansive mappings in Banach spaces.* Journal of Mathematical Analysis and Applications 67, 274–276 (1979)
- [47] **S. Sabach, S. Shtern.** *A first order method for solving convex bilevel optimization problems.* SIAM Journal on Optimization 27(2), 640–660 (2017)
- [48] **E.D. Sontag.** *Mathematical Control Theory. Deterministic Finite-Dimensional Systems.* Texts in Applied Mathematics 6, Springer, New York, 1998
- [49] **W. Su, S. Boyd, E. Candès.** *A differential equation for modeling Nesterov’s accelerated gradient method: theory and insights.* Journal of Machine Learning Research 17(153), 1–43 (2016)
- [50] **J. J. Suh, J. Park, E. K. Ryu.** *Continuous-time Analysis of Anchor Acceleration.* Advances in Neural Information Processing Systems 36 (2023)
- [51] **T. Suzuki.** *Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces.* Fixed Point Theory and Applications 2005(1) 103–123 (2005)
- [52] **T. Suzuki.** *A sufficient and necessary condition for Halpern-type strong convergence to fixed points of nonexpansive mappings.* Proceedings of the American Mathematical Society 135(01), 99–106 (2006).
- [53] **Q. Tran-Dinh.** *From Halpern’s fixed-point iterations to Nesterov’s accelerated interpretations for root-finding problems.* Computational Optimization and Applications 87, 181–218 (2024)
- [54] **Q. Tran-Dinh, Y. Luo.** *Halpern-type accelerated and splitting algorithms for monotone inclusions.* [arXiv:2110.08150](https://arxiv.org/abs/2110.08150)
- [55] **R. Wittmann.** *Approximation of fixed points of nonexpansive mappings.* Archiv der Mathematik 58(5), 486–491 (1992)
- [56] **H.-K. Xu.** *Iterative Algorithms for Nonlinear Operators.* Journal of the London Mathematical Society 66(1), 240–256 (2002)
- [57] **H.-K. Xu.** *An Iterative Approach to Quadratic Optimization.* Journal of Optimization Theory and Applications 116(3), 659–678 (2003)
- [58] **T. H. Yoon, E. K. Ryu.** *Accelerated algorithms for smooth convex-concave minimax problems with $\mathcal{O}(1/k^2)$ rate on squared gradient norm.* Proceedings of the 38th International Conference on Machine Learning, PMLR 139, 12098–12109 (2021)