Collapse of a quantum vortex in an attractive two-dimensional Bose gas

Sambit Banerjee,^{1,*} Kai Zhou,^{1,2,*} Shiva Kant Tiwari,^{3,*} Hikaru Tamura,^{1,†} Rongjie Li,¹

Panayotis Kevrekidis,⁴ Simeon I. Mistakidis,⁵ Valentin Walther,^{1,3,6} and Chen-Lung Hung^{1,6,‡}

¹Department of Physics and Astronomy, Purdue University, West Lafayette, IN 47907, USA

²School of the Gifted Young, University of Science and Technology of China, Hefei 230026, China

³Department of Chemistry, Purdue University, West Lafayette, IN 47907, USA

⁴Department of Mathematics and Statistics, University of Massachusetts Amherst, Amherst, Massachusetts 01003-9305, USA

⁵Department of Physics, Missouri University of Science and Technology, Rolla, MO 65409, USA

⁶Purdue Quantum Science and Engineering Institute,

Purdue University, West Lafayette, IN 47907, USA

(Dated: June 4, 2024)

We experimentally and numerically study the collapse dynamics of a quantum vortex in a twodimensional atomic superfluid following a fast interaction ramp from repulsion to attraction. We find the conditions and time scales for a superfluid vortex to radially converge into a quasi-stationary density profile, demonstrating the first spontaneous formation of a vortex soliton in an atomic Bose gas. We record an emergent universal dynamics of an azimuthal modulational instability, which amplifies initial density perturbations and leads to the eventual splitting of a vortex soliton or direct fragmentation of a superfluid into disordered, but roughly circular arrays of Townes solitonlike wavepackets. Our study sets the stage for exploring universal out-of-equilibrium dynamics of vortex quantum matter quenched to attractive interactions.

Vortices are prevalent fundamental excitations in nonlinear fields [1]. Probing vortex dynamics has played a pivotal role in studies of condensed matter, quantum gases, and nonlinear optics, from developing better understandings of superconductivity and superfluidity [2– 4] to finding new applications using angular momentum carrying optical beams [5].

Generally, the stability of vortices relies on the nature of the nonlinearity involved. As vortices are described by a field with integer multiples of 2π -phase winding around a phase singularity, which necessitates a zero amplitude defect at the vortex core, a self-defocusing (repulsive) nonlinearity can smoothen and stabilize the wave away from the defect. With a self-focusing (attractive) interaction, on the other hand, such waveforms become unstable against wave collapse. Nevertheless, it has been shown to be possible to embed vorticity in a stationary state, called a vortex soliton [6–8], where the self-focusing effect balances the wave dispersion of a ring-shaped waveform with phase winding.

However, even without radial wave collapse, a vortex soliton is still unstable against a pattern-forming instability [9], with which any azimuthal wave perturbations beyond a critical length scale can self-amplify. This instability leads to the growth of modulations in discrete angular modes and can eventually fragment a ring-shaped vortex soliton into a circular array of solitary waves in a distinct angular pattern like a 'necklace' [9–11]. To-date, experimental studies of solitons with vorticity focus on engineered optical vortices in nonlinear media [7, 8, 10– 19] and photonic lattices [20-22].

In quantum gas experiments, to our knowledge, vortex dynamics with attractive interactions has remained completely unexplored. Despite existing discussions on the formation and stability of vortex solitons in Bose-Einstein condensates [23–25], a successful demonstration of a collapsing vortex has remained elusive. This is hindered by state preparation in attractive Bose gases that have strong tendencies to collapse [26–30]. Whether a many-body vortex soliton can form spontaneously also remains an open question.

Recently, optical box-confined atomic quantum gases have emerged as an excellent platform [31] for studying wave collapse [30, 32] and pattern-forming instabilities [32–36]. This is due to the ability to remove undesired trap length scales and the flexibility to adjust the box boundary conditions. It thus becomes possible to access intrinsically unstable many-body states through nonequilibrium quench dynamics. By using a magnetic Feshbach resonance to quench the atomic interaction from repulsion to attraction, self-trapped, yet unstable fundamental solitons in two dimensions (2D)-the Townes solitons [37]-were found, surprisingly, following a universal wave collapse dynamics of a modulational instability in 2D [32, 38].

Watching vortices collapse following an interaction quench in a quantum gas can open pathways to uncover facinating self-patterned structures or vortex solitons. Collapse dynamics of vortices can exhibit new universal scaling behaviors and time scales distinct from those of wave collapse in vortex-free, non-rotating Bose gases [32]. Azimuthal modulational instability in a quantum vortex, when seeded by zero-temperature quantum fluctuations, can serve as a 'quantum parametric amplifier' [39, 40]. This may lead to macroscopic quantum entanglement and even many-body fragmentation [41–47] in the angu-

^{*} These authors contributed equally to this work.

[†] Present address: Insitute for Molecular Science, Okazaki, Aichi 444-8585, Japan

[‡] clhung@purdue.edu

lar momentum states that can have applications in quantum metrology [48, 49].

In this article, we study, for the first time, the collapse dynamics of a single vortex in attractive Bose gases confined in a quasi-2D box, and report the spontaneous formation of vortex solitons and self-patterned Townes soliton 'necklaces' from out-of-equilibrium dynamics. We create a single vortex nearly deterministically in a circular 2D superfluid, followed by ramping the atomic interaction to the weakly attractive regime and performing 2D time-of-flight (TOF) imaging. We observe clear radial convergence to a vortex soliton density profile and record a new universal scaling behavior and time scale of an azimuthal modulational instability that either splits a vortex soliton or directly fragments a superfluid into disordered, but roughly circular arrays of Townes solitonlike wavepackets. These dynamics are reproduced by the appropriate 2D Gross-Pitaevskii equation (GPE) incorporating a three-body loss term and initial random noise simulating zero-temperature phonon fluctuations. Our results pave the way for revealing universal collapse dynamics of vortex quantum matter.

The starting point of the experiment is a homogeneous 2D superfluid of $\approx 3.8 \times 10^4$ cesium atoms at low temperature T < 8 nK and trapped in a circular box (Appendix A), where the out-of-plane atomic motion




FIG. 1. Creation of vortex solitons and Townes soliton necklaces in 2D Bose gases. (a) Experimental scheme. A single vortex is created at the center of a circular 2D superfluid, followed by simultaneous ramp-down of the in-plane confinement and the interaction parameter to $g_f < 0$ for a TOF time τ in the 2D plane. (b) Density image taken after the ramp at $\tau = 5$ ms, $g_f \approx -0.0077$, and averaged over 4 shots. (c) Single-shot images with variable interaction g_f and time τ . Ring-like and necklace-like solitonic structures are observed at sufficiently long TOF times. Image size: (b) $(56\mu m)^2$ and (c) $(65\mu m)^2$. All images use the same color scale.

is frozen in the ground state with an oscillator length $l_z \approx 265$ nm. The initial radius of the 2D superfluid is $r_i \approx 21 \ \mu\text{m}$ and the surface density $n_i \approx 25 \ \mu\text{m}^{-2}$. The interaction strength $g = \sqrt{8\pi}a_s/l_z$ is controlled by a magnetic Feshbach resonance that tunes the s-wave scattering length a_s , setting the initial value to $g_i \approx 0.1$.

We create a single vortex near deterministically at the center of the 2D superfluid using a scheme demonstrated in Refs. [50–52]. The procedure begins with a repulsive circular beam of an initial radius of $\approx 4 \ \mu m$ and a potential height $\approx k_B \times 47$ nK, slowly sweeping across the superfluid while splitting into two beams of the same diameter at a 65° separation angle. Here, k_B is the Boltzmann constant. The beams move approximately 12 μ m in 400 ms, corresponding to a linear speed of $\approx 0.04c$, where $c = \hbar \sqrt{n_i g_i} / m \approx 0.76$ mm/s is the bulk sound speed, $\hbar = h/2\pi$ is the reduced Planck constant and m is the atomic mass. Ideally, this procedure creates a density defect with a 2π -phase winding around each beam center but with opposite circularity, forming a vortex dipole. We arrange the potential sweep such that one beam ends up at the center of the superfluid while the other one stops near the box boundary [51], as illustrated in Fig. 1(a). After each beam reaches its final position, the beam size is slowly shrunk down to zero in another 490 ms. From single-shot absorption images, shown in Fig. 1(b), we observe a single vortex at the superfluid center with nearly 90% probability. Near the edge of the 2D gas, we observe the second vortex for less than 10%of the experimental repetitions (Appendix B).

To induce collapse dynamics following the vortex preparation procedure, we hold the gas in the box for another 16 ms and then quickly ramp the interaction strength in 5 ms to a variable attractive value $g_f < 0$, as schematically depicted in Fig. 1(a), while simultaneously ramping off (in 3 ms) the in-plane box confinement. The ramp speed is fast compared to subsequent vortex dynamics at $g = g_f$, but slow enough to avoid creating additional excitations near the edge of the 2D gas, such as ring dark solitons [34] that can be generated following a quick potential ramp-down, or bright solitonic density peaks due to a fast interaction reduction. We then allow the gas to freely evolve in the 2D plane for a variable TOF time τ , followed by absorption imaging to record the 2D density distribution.

Figure 1(c) shows single-shot images of samples held at different interaction strengths. For weak attraction $g_f \approx -0.0017$ (top row), the vortex core soon expands to a larger size and the disk-like density structure evolves into a ring-shaped profile for an extended period of time. At longer hold times, the ring structure becomes distorted and splits, manifesting the presence of the azimuthal modulational instability [53]. At more negative values of g_f , the 2D gas directly fragments into circular arrays of density blobs instead of forming a single ring-shaped structure, as shown in the second and the third rows of Fig. 1(c). This peculiar multi-ring fragmentation indicates fast wave collapse, both radi-




FIG. 2. Evolution of the radial density profiles. Experiment: (a, b) Radially averaged density profiles (filled symbols) for (a) $g_f \approx -0.0017$ and (b) $g_f \approx -0.0077$ measured at the indicated time τ . Shaded band in (a) shows the stationary vortex soliton density profile rescaled to match the peak densities measured at $\tau = 37 \sim 77$ ms. (c) Variance V (filled symbols) of the measured density profiles relative to a reference profile recorded at time $\tau_{\rm ref}$ and normalized by the mean value evaluated near the reference time. Grey dashed line marks the value V = 1. Inset depicts an averaged density image taken at $\tau_r = 49$ ms when the radial profile converges. Simulation: (d, e) Radial density profiles evaluated from the GPE simulations (solid curves) using experimental parameters as in (a) and (b), respectively. Filled circles illustrate a density profile from (a) at time $\tau > \tau_r$. (f) Normalized variance V of simulated density profiles. Arrow indicates the onset of a weak radial convergence plateau and the inset presents the corresponding averaged density image.

ally and azimuthally. One expects the blobs to form around the size of an interaction length $\xi = \pi/k_{\rm MI}$, where $k_{\rm MI} = \sqrt{2n_i |g_f|}$ is the most unstable wavenumber with the largest imaginary Bogoliubov dispersion [32]. At longer TOF times, the circular arrays appear to collide with each other in the radial direction and collapse, leaving behind a disordered array of disjoint blobs whose density profiles approach those of Townes solitons (see Appendix C). We dub this disordered array a Townes soliton necklace.

The multi-ring collapse dynamics can be analyzed through radially averaged density profiles, where we set the vortex core as the origin; see Fig. 2. At $g_f \approx -0.0077$ (Fig. 2(b)), two density peaks appear at $\tau \approx 25$ ms, corresponding to the formation of two approximately circular arrays of blobs as shown in Fig. 1(c). However, these peaks soon collapse. The remaining broad peak observed at $\tau \gtrsim 37$ ms comes from the radial average of a disordered Townes soliton necklace and the collision remnant.

Remarkably, the radial profile of the ring-shaped structure observed at the weaker interaction $g_f \approx -0.0017$ remains nearly stationary for more than 40 ms from $\tau \gtrsim 37$ ms; see Fig. 2(a). This suggests that the vortex evolves into a self-trapped vortex soliton. The peak density is found at a fixed radial position at $r \approx 10 \ \mu m$, agreeing fairly well with a stationary-state solution of the GPE [24], $n(r) = n_p \left| \phi_{vs} \left(\sqrt{n_p |g_f|} r \right) \right|^2$, where n_p is the peak density and ϕ_{vs} is the radial part of an ideal vortex wavefunction $\psi(R,\theta) = \phi_{\rm vs}(R)e^{is\theta}$ with $s = \pm 1$. We solve the radial wavefunction using the 2D GPE in a scale-invariant form $-\frac{1}{2}\left(\frac{\partial^2}{\partial R^2} + \frac{1}{R}\frac{\partial}{\partial R} - \frac{s^2}{R^2}\right)\phi_{\rm vs} - |\phi_{\rm vs}|^2\phi_{\rm vs} = \tilde{\mu}\phi_{\rm vs}$, where $R = \sqrt{n_p |g_f|}r$ is the rescaled radial coordinate and $\tilde{\mu} = -0.3247$ is the scaled chemical potential. We note this is a unique, scale-invariant vortex solution in 2D, with n_p fixing the physical size of the wavefunction, and no free parameters are used in the comparison; see shaded band in Fig. 2(a). The measured quasi-stationary density profile nevertheless deviates from the ideal solution near the low-density wings, potentially due to very slow collapse dynamics of atoms dispersed during the ramp-down of the box confinement.

We can quantify exactly when a vortex soliton forms by comparing the radial density profiles $n(r, \tau)$ to a reference profile obtained at a long TOF time $\tau_{\rm ref}$, chosen to be 4 ms after the last point of each curve in Fig. 2(c). We compute the time-dependent variance $V(\tau) \equiv \int [n(r, \tau) - n(r, \tau_{\rm ref})]^2 dr$ normalized by the mean value evaluated near time $\tau_{\rm ref}$. If the density radially converges to a quasi-stationary profile, V can either settle around unity as it is limited by technical noise, or plateau at some intermediate time followed by slow reduction due to residual radial evolutions. Indeed, the density profiles of $g_f \approx -0.0017$ satisfy this criterion, with $V \approx 1$ after time $\tau_r \approx 49$ ms. For more attractive samples, V decreases monotonically at all times, implying the absence of a quasi-stationary intermediate state.




FIG. 3. Dynamics of azimuthal modulational instability and universal scaling behavior. (a) The angular power spectrum S_l at $g_f \approx -0.0017$ for different TOF times τ . Single-shot images at the top are recorded at $\tau = 77$ ms, representing soliton splitting into an angular mode at l = 3 and 4, respectively. (b) S_l at $g_f \approx -0.0077$. Single-shot images are recorded at $\tau = 53$ ms, evincing dominant mode(s) l = 7and $9 \sim 11$, respectively. Vertical dashed lines in (a) and (b) indicate the peak position $l_{\rm p}$ identified using the last seven TOF time points. (c) l_p versus g_f (filled circles). Error bars represent the standard error of the mean. Blue dashed line marks $l_p = \bar{r}k_{\rm MI}$, evaluated using the mean radii of the samples. Red solid line is a fit. Green dotted line marks the most unstable mode of a vortex soliton. Shaded band shows l_{p} and the standard error evaluated from the GPE simulations. (d) Time evolution of the averaged power spectrum \overline{S} (color symbols). Insets present single-shot density images taken at the saturation time of \bar{S} , $\tau_{\theta} = 17$ and 37 ms, for $g_f \approx -0.011$ and -0.0047, respectively. Color scales are the same as in (a). (e) Scaled power spectra of (d) (symbols) and simulations (shaded bands, same color) versus rescaled time. The amplitude of numerical spectra is reduced by 2 times for comparing with experiments. Data plotted in (d) and (e) use the same labeled color symbols.

We can fully reproduce key signatures of the radial collapse dynamics in the time evolutions of the 2D GPE (Appendix D), see Figs. 2(d-f). To properly model atom losses during the collapse dynamics, a nonlinear three-body loss term is incorporated in the GPE simulation. To identify soliton formation time scales, in Fig. 2(f) we find that the variance V of simulated density profiles weakly converges for interactions $|g_f| \leq 0.0025$ at $\tau \geq 40$ ms. After a short-lived plateau, V slowly reduces due to the residual evolution in the low-density wing, as seen in Fig. 2(d).

The insets in Fig. 2(c) and (f) compare experimental and numerical density images (each averaged over ~ 10 shots) measured at $\tau = \tau_r$ when the radial density profiles converge. Both suggest that the vortex solitons already suffer significant azimuthal distortions by this time. We attribute this to the presence of an azimuthal modulational instability, which amplifies initial density perturbations in our samples.

We now ask how and when the azimuthal instability manifests in the vortex collapse dynamics. To analyze the dynamics, we evaluate the azimuthal number distribution $N_{\theta} = \int_{0}^{r_{i}} n(r,\theta)rdr$ and calculate the angular power spectrum $S_{l} = \langle |N_{l}|^{2} \rangle / N$, where $l \in \mathbb{N}$ labels discrete angular modes, $N_{l} = \int_{0}^{2\pi} N_{\theta} e^{-il\theta} d\theta$, N is the total atom number, and $\langle \cdots \rangle$ denotes ensemble averaging.

In Figs. 3(a) and (b), we present the time evolution of measured spectra S_l , revealing dynamic competition within a band of unstable angular modes. Each spectrum shows significant growth initially at some larger angular modes near l = 6 and 11, respectively, which is succeeded by a band of lower l modes that ultimately culminate in a distinct peak. The peak is broader for more attractive interactions, as in Fig. 3(b), where smaller-scale perturbations also become unstable. The prevailing peak in the power spectrum reveals the most unstable l-mode(s) that could most frequently fragment a ring structure into lpieces–a signature of pattern-forming instability dynamics. In the top panels of Figs. 3(a) and (b), we show single-shot density images to illustrate the ensuing fragmentation caused by different modes.

In Fig. 3(c), we extract the peak position l_p measured at later times in the spectra (Appendix E) and compare it with theoretical expectations. Overall, the peak locations follow reasonably well the prediction from a variational calculation $l_p \approx \bar{r} k_{\rm MI} = \pi \bar{r} / \xi$ [53]. Here the peak azimuthal mode number is determined by the most unstable wavenumber $k_{\rm MI}$ in the Bogoliubov modes and an effective radius $\bar{r} = \sqrt{\int \frac{n(r)^2}{n_i} r dr / \int \frac{n(r)}{r^2} r dr} \approx 7.5 \ \mu {\rm m}$, evaluated using the density profiles near the fragmentation time (Appendix F). We also fit l_p using a variable $\bar{r}.$ The fitted $\bar{r}\approx 10~\mu{\rm m}$ appears to reflect the peak radii of actual samples as shown in Fig. 2. Meanwhile, we find qualitatively similar power spectra from the GPE data and very good agreement with the measured l_p . The overall trend reflects that the interaction length ξ is a dominant length scale fragmenting our disk-shaped atomic samples. Starting with an ideal vortex soliton wavefunction, on the other hand, the expected most unstable mode l = 2 is independent of the values of (q_f, n_p) (Appendix G) [23, 25].

Surprisingly, we find a new universal scaling behavior and a universal time scale when the azimuthal modulational instability fragments a sample. To see this, we first calculate \bar{S} , the mean power spectrum of S_l averaged over a wide band $(l = 1 \sim 30)$, and monitor its time evolution in Fig. 3(d). For every interaction except $g_f \approx -0.0017$, the growth of \bar{S} saturates, indicating when the azimuthal perturbations cease to grow and the samples fragment. The insets are representative singleshot density images taken at the onset of saturation (τ_{θ}) , showing single- and double-ring fragmentation patterns.




FIG. 4. Soliton formation timescales versus rescaled initial sample size. Formation of vortex solitons (VS) at $\tilde{\tau}_r = \gamma \tau_r \lesssim$ 1 is observed for a sample with $r_i/\xi \lesssim 2$ (red circle). Samples with $r_i/\xi \gtrsim 3$ feature direct fragmentation to Townes solitons (TS) at nearly identical times $\tilde{\tau}_{\theta} = \gamma \tau_{\theta} \approx 2$ (blue triangles). Blue dashed line marks the mean value. Shaded bands indicate the rescaled time and uncertainty for radial convergence (red) as well as azimuthal fragmentation (blue) identified from the GPE simulations.

Notably, the scaled spectra $(\gamma/\gamma_i)^{\alpha}\bar{S}$ versus the rescaled time $\tilde{\tau} = \gamma \tau$ in Fig. 3(e) clearly exhibit a self-similar behavior, where $\alpha = 0.5$ is an empirical scaling exponent and $\hbar \gamma_{(i)} = \hbar^2 n_i |g_{f(i)}|/m$ is the (initial) interaction energy. This behavior suggests that, regardless of the radial dynamics, azimuthal perturbations exhibit a universal growth curve with a peculiar interaction scaling and fragmentation occurs at a nearly fixed time $\tilde{\tau}_{\theta} \approx 2$ despite manifesting by different angular patterns. The power spectra from the GPE simulations feature the same scaling behavior and peak at $\tilde{\tau}_{\theta} \approx 2.5$. We note that a universal scaling behavior of $\alpha = 1$, predicted by a linear analysis of the modulational instability, and wave fragmentation at $\tilde{\tau} \approx 0.8$ was observed in vortex-free 2D gases [32]. Our observation suggests that the presence of a single vortex significantly delays (azimuthal) wave fragmentation.

Finally, we summarize the fate of a single vortex following a fast ramp to attractive interactions. In Fig. 4, we present the identified soliton formation time scales versus the initial sample radius r_i normalized by the interaction length scale ξ . Based on the scale-invariant nature of weakly interacting 2D Bose gases [38, 54–56], we expect that the observed dynamics can be replicated in samples of different initial sizes and final interaction strengths within the same range $1 \leq r_i/\xi \leq 5$, provided that the weak three-body loss dynamics does not sig-

nificantly break the scale-invariance. For samples with $r_i/\xi \lesssim 2$, we observe the formation of a quasi-stationary, ring-shaped density profile at $\tilde{\tau} = \tilde{\tau}_r \lesssim 1$, which we identify as a vortex soliton. Perturbations in the vortex soliton grow continuously due to the azimuthal modulational instability until approaching $\tilde{\tau} \approx 2$ where the density ring nearly splits into a circular array of bright solitons with distinct angular patterns (Fig. 3(a)). Soliton splitting speeds up when the interaction length approaches the sample initial size $\xi \approx r_i$, as confirmed by GPE simulations (Appendix D). On the other hand, for larger size or more attractive samples with $r_i/\xi \gtrsim 3$, more than one density ring forms in the radial direction. These samples all fragment azimuthally at $\tilde{\tau} = \tilde{\tau}_{\theta} \approx 2$. The multiring structure eventually collapses, creating a disordered Townes soliton necklace. We note that starting with different initial interaction energy $\hbar \gamma_i$ and temperature T can change the amplitudes of initial density fluctuations that seed the instability. Our samples have suppressed thermal fluctuations due to $\eta = \hbar \gamma_i / k_B T \gtrsim 1$. When increasing $\eta > 1$, we expect $\tilde{\tau}_{\theta}, \tilde{\tau}_r$ to remain roughly unchanged, as the instability remains primarily seeded by quantum fluctuations (small initial perturbations from the vortex preparation are also present as seen in our GPE simulations). For samples with lower $\eta < 1$, however, increased initial thermal fluctuations and density perturbations may lead to early fragmentation, making a vortex soliton shorter-lived with smaller $\tilde{\tau}_{\theta} < 2$.

Our observed quench dynamics point towards a universal behavior and extended time scales to create selftrapped vortex matter from out-of-equilibrium dynamics. We note that collapse of vortices and soliton formation can also be investigated in two-component Bose gases with an effective attractive interaction [56, 57]. Further manipulation or studies of the azimuthal modulational instability of a vortex soliton can be carried out by adding a ring trap for radial confinement or by pinning the vortex core with a repulsive beam. Another nontrivial extension of this work would be to study the quench dynamics of a vortex lattice in a superfluid [58, 59]. By tuning the vortex density to reach a lattice constant $\gtrsim 2\xi$, roughly corresponding to $r_i/\xi \gtrsim 1$ for the size of individual vortex puddles in the lattice, one may find novel vortex soliton array formations and can study the subsequent collision and collapse of these solitonic structures. While our work is based on weakly attractive Bose gases, our experimental method can be adapted to embed vortices in a quantum droplet [60–63], where wave collapse from a strong mean-field attraction is prevented by an effective repulsion from quantum fluctuations [64–66], and a self-trapped state with vorticity will exhibit different instability behaviors [67–69].

- L. M. Pismen, Vortices in nonlinear fields: from liquid crystals to superfluids, from non-equilibrium patterns to cosmic strings, Vol. 100 (Oxford University Press, 1999).
- [2] M. Salomaa and G. Volovik, Quantized vortices in superfluid he 3, Reviews of Modern Physics 59, 533 (1987).
- [3] G. Blatter, M. V. Feigel'man, V. B. Geshkenbein, A. I.

Larkin, and V. M. Vinokur, Vortices in high-temperature superconductors, Reviews of Modern Physics **66**, 1125 (1994).

- [4] M. W. Zwierlein, J. R. Abo-Shaeer, A. Schirotzek, C. H. Schunck, and W. Ketterle, Vortices and superfluidity in a strongly interacting fermi gas, Nature 435, 1047 (2005).
- [5] Y. Shen, X. Wang, Z. Xie, C. Min, X. Fu, Q. Liu, M. Gong, and X. Yuan, Optical vortices 30 years on: Oam manipulation from topological charge to multiple singularities, Light: Science & Applications 8, 90 (2019).
- [6] V. Kruglov, Y. A. Logvin, and V. Volkov, The theory of spiral laser beams in nonlinear media, Journal of Modern Optics 39, 2277 (1992).
- [7] A. S. Desyatnikov, Y. S. Kivshar, and L. Torner, Optical vortices and vortex solitons, Progress in Optics 47, 291 (2005).
- [8] B. A. Malomed, Vortex solitons: Old results and new perspectives, Physica D: Nonlinear Phenomena **399**, 108 (2019).
- [9] W. J. Firth and D. V. Skryabin, Optical solitons carrying orbital angular momentum, Physical Review Letters 79, 2450 (1997).
- [10] D. V. Petrov, L. Torner, J. Martorell, R. Vilaseca, J. P. Torres, and C. Cojocaru, Observation of azimuthal modulational instability and formation of patterns of optical solitons in a quadratic nonlinear crystal, Optics Letters 23, 1444 (1998).
- [11] M. S. Bigelow, P. Zerom, and R. W. Boyd, Breakup of ring beams carrying orbital angular momentum in sodium vapor, Physical Review Letters 92, 083902 (2004).
- [12] G. Swartzlander Jr and C. Law, Optical vortex solitons observed in kerr nonlinear media, Physical Review Letters 69, 2503 (1992).
- [13] G. Duree, M. Morin, G. Salamo, M. Segev, B. Crosignani, P. Di Porto, E. Sharp, and A. Yariv, Dark photorefractive spatial solitons and photorefractive vortex solitons, Physical Review Letters 74, 1978 (1995).
- [14] Z. Chen, M.-f. Shih, M. Segev, D. W. Wilson, R. E. Muller, and P. D. Maker, Steady-state vortex-screening solitons formed in biased photorefractive media, Optics Letters 22, 1751 (1997).
- [15] M. Soljačić, S. Sears, and M. Segev, Self-trapping of "necklace" beams in self-focusing kerr media, Physical Review Letters 81, 4851 (1998).
- [16] C.-C. Jeng, M.-F. Shih, K. Motzek, and Y. Kivshar, Partially incoherent optical vortices in self-focusing nonlinear media, Physical Review Letters 92, 043904 (2004).
- [17] A. S. Reyna, G. Boudebs, B. A. Malomed, and C. B. de Araújo, Robust self-trapping of vortex beams in a saturable optical medium, Physical Review A 93, 013840 (2016).
- [18] A. S. Reyna, H. T. Baltar, E. Bergmann, A. M. Amaral, E. L. Falcão-Filho, P.-F. Brevet, B. A. Malomed, and C. B. de Araújo, Observation and analysis of creation, decay, and regeneration of annular soliton clusters in a lossy cubic-quintic optical medium, Physical Review A 102, 033523 (2020).
- [19] L. T. Vuong, T. D. Grow, A. Ishaaya, A. L. Gaeta, G. W. 't Hooft, E. R. Eliel, and G. Fibich, Collapse of optical vortices, Physical Review Letters 96, 133901 (2006).
- [20] J. W. Fleischer, G. Bartal, O. Cohen, O. Manela, M. Segev, J. Hudock, and D. N. Christodoulides, Observation of vortex-ring "discrete" solitons in 2d photonic

lattices, Physical Review Letters 92, 123904 (2004).

- [21] D. N. Neshev, T. J. Alexander, E. A. Ostrovskaya, Y. S. Kivshar, H. Martin, I. Makasyuk, and Z. Chen, Observation of discrete vortex solitons in optically induced photonic lattices, Physical Review Letters **92**, 123903 (2004).
- [22] G. Bartal, O. Manela, O. Cohen, J. W. Fleischer, and M. Segev, Observation of second-band vortex solitons in 2d photonic lattices, Physical Review Letters 95, 053904 (2005).
- [23] H. Saito and M. Ueda, Split instability of a vortex in an attractive bose-einstein condensate, Physical Review Letters 89, 190402 (2002).
- [24] L. Carr and C. W. Clark, Vortices in attractive boseeinstein condensates in two dimensions, Physical Review Letters 97, 010403 (2006).
- [25] D. Mihalache, D. Mazilu, B. A. Malomed, and F. Lederer, Vortex stability in nearly-two-dimensional boseeinstein condensates with attraction, Physical Review A 73, 043615 (2006).
- [26] C. Sackett, H. Stoof, and R. Hulet, Growth and collapse of a bose-einstein condensate with attractive interactions, Physical review letters 80, 2031 (1998).
- [27] J. M. Gerton, D. Strekalov, I. Prodan, and R. G. Hulet, Direct observation of growth and collapse of a bose– einstein condensate with attractive interactions, Nature 408, 692 (2000).
- [28] E. A. Donley, N. R. Claussen, S. L. Cornish, J. L. Roberts, E. A. Cornell, and C. E. Wieman, Dynamics of collapsing and exploding bose–einstein condensates, Nature **412**, 295 (2001).
- [29] J. L. Roberts, N. R. Claussen, S. L. Cornish, E. A. Donley, E. A. Cornell, and C. E. Wieman, Controlled collapse of a bose-einstein condensate, Physical Review Letters 86, 4211 (2001).
- [30] C. Eigen, A. L. Gaunt, A. Suleymanzade, N. Navon, Z. Hadzibabic, and R. P. Smith, Observation of weak collapse in a bose-einstein condensate, Physical Review X 6, 041058 (2016).
- [31] N. Navon, R. P. Smith, and Z. Hadzibabic, Quantum gases in optical boxes, Nature Physics 17, 1334 (2021).
- [32] C.-A. Chen and C.-L. Hung, Observation of universal quench dynamics and townes soliton formation from modulational instability in two-dimensional bose gases, Physical Review Letters 125, 250401 (2020).
- [33] Z. Zhang, K.-X. Yao, L. Feng, J. Hu, and C. Chin, Pattern formation in a driven bose–einstein condensate, Nature Physics 16, 652 (2020).
- [34] H. Tamura, C.-A. Chen, and C.-L. Hung, Observation of self-patterned defect formation in atomic superfluids– from ring dark solitons to vortex dipole necklaces, Physical Review X 13, 031029 (2023).
- [35] D. Hernandez-Rajkov, N. Grani, F. Scazza, G. Del Pace, W. Kwon, M. Inguscio, K. Xhani, C. Fort, M. Modugno, F. Marino, *et al.*, Universality of the superfluid kelvinhelmholtz instability by single-vortex tracking, arXiv preprint arXiv:2303.12631 (2023).
- [36] N. Liebster, M. Sparn, E. Kath, K. Fujii, S. Görlitz, T. Enss, H. Strobel, and M. K. Oberthaler, Emergence of crystalline steady state in a driven superfluid, (2023), arXiv:2309.03792 [cond-mat.quant-gas].
- [37] R. Y. Chiao, E. Garmire, and C. H. Townes, Self-trapping of optical beams, Physical Review Letters 13, 479 (1964).
- [38] C.-A. Chen and C.-L. Hung, Observation of scale invariance in two-dimensional matter-wave townes solitons,

Physical Review Letters 127, 023604 (2021).

- [39] J. Esteve, C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler, Squeezing and entanglement in a bose– einstein condensate, Nature 455, 1216 (2008).
- [40] C.-A. Chen, S. Khlebnikov, and C.-L. Hung, Observation of quasiparticle pair production and quantum entanglement in atomic quantum gases quenched to an attractive interaction, Physical Review Letters 127, 060404 (2021).
- [41] P. Nozieres and D. Saint James, Particle vs. pair condensation in attractive bose liquids, Journal de Physique 43, 1133 (1982).
- [42] N. Wilkin, J. Gunn, and R. Smith, Do attractive bosons condense?, Physical Review Letters 80, 2265 (1998).
- [43] R. Spekkens and J. Sipe, Spatial fragmentation of a boseeinstein condensate in a double-well potential, Physical Review A 59, 3868 (1999).
- [44] T.-L. Ho and S. K. Yip, Fragmented and single condensate ground states of spin-1 bose gas, Physical review letters 84, 4031 (2000).
- [45] E. J. Mueller, T.-L. Ho, M. Ueda, and G. Baym, Fragmentation of bose-einstein condensates, Physical Review A 74, 033612 (2006).
- [46] J. Nguyen, M. Tsatsos, D. Luo, A. Lode, G. D. Telles, V. S. Bagnato, and R. Hulet, Parametric excitation of a bose-einstein condensate: From faraday waves to granulation, Physical Review X 9, 011052 (2019).
- [47] B. Evrard, A. Qu, J. Dalibard, and F. Gerbier, Observation of fragmentation of a spinor bose-einstein condensate, Science **373**, 1340 (2021).
- [48] X.-W. Luo, C. Zhang, and S. Du, Quantum squeezing and sensing with pseudo-anti-parity-time symmetry, Physical Review Letters 128, 173602 (2022).
- [49] L. Pezze, A. Smerzi, M. K. Oberthaler, R. Schmied, and P. Treutlein, Quantum metrology with nonclassical states of atomic ensembles, Reviews of Modern Physics 90, 035005 (2018).
- [50] E. Samson, K. Wilson, Z. Newman, and B. Anderson, Deterministic creation, pinning, and manipulation of quantized vortices in a bose-einstein condensate, Physical Review A 93, 023603 (2016).
- [51] B. Gertjerenken, P. G. Kevrekidis, R. Carretero-Gonzalez, and B. Anderson, Generating and manipulating quantized vortices on-demand in a bose-einstein condensate: A numerical study, Physical Review A 93, 023604 (2016).
- [52] W. J. Kwon, G. Del Pace, K. Xhani, L. Galantucci, A. Muzi Falconi, M. Inguscio, F. Scazza, and G. Roati, Sound emission and annihilations in a programmable quantum vortex collider, Nature **600**, 64 (2021).
- [53] R. M. Caplan, Q. Hoq, R. Carretero-González, and P. G. Kevrekidis, Azimuthal modulational instability of vortices in the nonlinear schrödinger equation, Optics Communications 282, 1399 (2009).
- [54] C.-L. Hung, X. Zhang, N. Gemelke, and C. Chin, Observation of scale invariance and universality in twodimensional bose gases, Nature 470, 236 (2011).
- [55] T. Yefsah, R. Desbuquois, L. Chomaz, K. J. Günter, and J. Dalibard, Exploring the thermodynamics of a two-dimensional bose gas, Physical Review Letters 107, 130401 (2011).
- [56] B. Bakkali-Hassani, C. Maury, Y.-Q. Zou, É. Le Cerf, R. Saint-Jalm, P. C. M. Castilho, S. Nascimbene, J. Dalibard, and J. Beugnon, Realization of a townes soliton in

a two-component planar bose gas, Physical Review Letters **127**, 023603 (2021).

- [57] A. Romero-Ros, G. Katsimiga, S. Mistakidis, S. Mossman, G. Biondini, P. Schmelcher, P. Engels, and P. Kevrekidis, Experimental realization of the peregrine soliton in repulsive two-component bose-einstein condensates, Physical Review Letters **132**, 033402 (2024).
- [58] K. W. Madison, F. Chevy, V. Bretin, and J. Dalibard, Stationary states of a rotating bose-einstein condensate: Routes to vortex nucleation, Physical Review Letters 86, 4443 (2001).
- [59] J. R. Abo-Shaeer, C. Raman, J. M. Vogels, and W. Ketterle, Observation of vortex lattices in bose-einstein condensates, Science 292, 476 (2001).
- [60] I. Ferrier-Barbut, H. Kadau, M. Schmitt, M. Wenzel, and T. Pfau, Observation of quantum droplets in a strongly dipolar bose gas, Physical review letters **116**, 215301 (2016).
- [61] L. Chomaz, S. Baier, D. Petter, M. Mark, F. Wächtler, L. Santos, and F. Ferlaino, Quantum-fluctuation-driven crossover from a dilute bose-einstein condensate to a macrodroplet in a dipolar quantum fluid, Physical Review X 6, 041039 (2016).
- [62] G. Semeghini, G. Ferioli, L. Masi, C. Mazzinghi, L. Wolswijk, F. Minardi, M. Modugno, G. Modugno, M. Inguscio, and M. Fattori, Self-bound quantum droplets of atomic mixtures in free space, Physical review letters 120, 235301 (2018).
- [63] C. Cabrera, L. Tanzi, J. Sanz, B. Naylor, P. Thomas, P. Cheiney, and L. Tarruell, Quantum liquid droplets in a mixture of bose-einstein condensates, Science **359**, 301 (2018).
- [64] A. Bulgac, Dilute quantum droplets, Physical review letters 89, 050402 (2002).
- [65] D. Petrov, Quantum mechanical stabilization of a collapsing bose-bose mixture, Physical review letters 115, 155302 (2015).
- [66] P. Naidon and D. Petrov, Mixed bubbles in bose-bose mixtures, Physical Review Letters 126, 115301 (2021).
- [67] Y. Li, Z. Chen, Z. Luo, C. Huang, H. Tan, W. Pang, and B. A. Malomed, Two-dimensional vortex quantum droplets, Physical Review A 98, 063602 (2018).
- [68] Y. V. Kartashov, B. A. Malomed, L. Tarruell, and L. Torner, Three-dimensional droplets of swirling superfluids, Physical Review A 98, 013612 (2018).
- [69] L. Dong and Y. V. Kartashov, Rotating multidimensional quantum droplets, Physical Review Letters **126**, 244101 (2021).
- [70] S. K. Tiwari and S. Wüster, Tracking rydberg atoms with bose-einstein condensates, Phys. Rev. A 99, 043616 (2019).
- [71] G. R. Dennis, J. J. Hope, and M. T. Johnsson, (2012), http://www.xmds.org/.
- [72] G. R. Dennis, J. J. Hope, and M. T. Johnsson, XMDS2: Fast, scalable simulation of coupled stochastic partial differential equations, Comput. Phys. Comm. 184, 201 (2013).
- [73] P. B. Blakie, A. Bradley, M. Davis, R. Ballagh, and C. Gardiner, Dynamics and statistical mechanics of ultracold bose gases using c-field techniques, Advances in Physics 57, 363 (2008).
- [74] T. Kraemer, M. Mark, P. Waldburger, J. G. Danzl, C. Chin, B. Engeser, A. D. Lange, K. Pilch, A. Jaakkola, H.-C. Nägerl, *et al.*, Evidence for efimov quantum states

in an ultracold gas of caesium atoms, Nature ${\bf 440},\,315$ (2006).

- [75] R. M. Caplan, R. Carretero-González, P. G. Kevrekidis, and B. A. Malomed, Existence, stability, and scattering of bright vortices in the cubic-quintic nonlinear schrödinger equation, Mathematics and Computers in Simulation 82, 1150 (2012).
- [76] M. Edwards, R. Dodd, C. W. Clark, and K. Burnett, Zero-temperature, mean-field theory of atomic boseeinstein condensates, Journal of research of the National Institute of Standards and Technology 101, 553 (1996).

ACKNOWLEDGMENTS

We thank Qiyu Liang, Cheng Chin, Moti Segev, and Ron Ziv for fruitful discussions. This work is supported in part by the NSF (Grant # PHY-1848316), the AFOSR (FA9550-22-1-0327), and the DOE QuantISED program through the Fermilab Quantum Consortium. P.G.K. acknowledges support by the NSF under the awards PHY-2110030 and DMS-2204702. R. Li acknowledges the Rolf Scharenberg Graduate Summer Research Fellowship.

Appendix A: The box potential

The 2D homogeneous Bose gases are confined in a flat optical box potential. The in-plane confinement is provided by a repulsive ring beam, created by a blue-detuned 780 nm light reflecting off a digital mirror device (DMD) and projected through a microscope objective (numerical aperture N.A. = 0.6). The same DMD is also used to dynamically project additional repulsive potentials to create a single vortex at the center of the 2D superfluid. The out-of-plane (vertical) confinement of the 2D gas is provided by a single node of a repulsive optical lattice potential, formed using another 780 nm light.

We note that, without compensation, a very weak inplane trap corrugation is present in the box due to the confining lattice beam. The potential variation is estimated to be $\Delta U \leq k_{\rm B} \times 2$ nK. While it imposes no visible effect on the initial 2D superfluid, a weakly attractive sample could eventually be localized in the local potential minima created by this corrugation, forming linear density stripes. To mitigate this effect, we ramp on a compensating potential pattern at $\tau = 2$ ms using the DMD with a matching periodicity ($\approx 18 \ \mu$ m) to minimize the density stipes observed after long TOF times. It is to be noted that this compensation is not perfect (estimate residual $\Delta U < k_{\rm B} \times 0.2$ nK) and could be partially responsible for seeding the azimuthal modulational instability.

Appendix B: Deterministic vortex creation and imaging

We use a dynamic repulsive potential patterned by the DMD to deterministically create vortices in the 2D superfluids. While we have performed GPE simulations using the so-called chopstick method [50, 51] as depicted in Fig. 1(a) and in Appendix D, we have experimentally searched for the proper conditions for deterministic vortex creation and pinning, which largely depends on the potential height and the sweep speed of the chopsticks, as well as the time over which we ramp off the size of the potential. We have used a potential height of $V_0 \approx k_B \times 47$ nK optimized for our sample density. The sweep speed and ramp-off time are optimized so that the vortices are pinned to the chopstick beam centers with high probability. We observed that a faster sweep generates more than one randomly positioned density defect. Similarly, a faster ramp-off of the chopstick beam size either leads to a lower vortex creation probability or results in vortex unpinning. We have also optimized the size of the chopstick beams and the angle of divergence, with respect to the sample size, to reach higher vortex generation probability. Detailed systematic studies of our deterministic 2D vortex creation will be presented elsewhere.

Figure 5 illustrates sample images of vortices created in 2D superfluids. Figure 5(a) shows a single vortex imaged in-situ in the repulsive interaction regime. In order to clearly resolve the vortex core, we slowly ramp the interaction from $g \approx 0.1$ down to 0.0056 in 40 ms to enlarge the core size before imaging. Figure 5(b) presents another image of a vortex prepared under the same interaction ramp procedure but with an additional 40 ms of 2D TOF time at the same interaction $g \approx 0.0056$. The vortices remain stable under a variety of 2D TOF conditions, including long TOF under a strong repulsive interaction. To clearly image the vortex core, however, we always need to ramp the interaction down to a weaker value prior to imaging.

Figure 5(c) shows an in situ image of a vortex dipole created symmetrically about the trap center. Figure 5(d) depicts another image of a vortex dipole of a different orientation, with the second vortex created near the edge of the superfluid at the lower side of the image. This is the configuration used for the main experiment. The probability of observing a second vortex near the edge, as shown in (d), is $\leq 10\%$. Although not demonstrated in this study, bringing the second chopstick beam out of the box can in principle completely anihilate the second vortex [51].




FIG. 5. Single-shot images of (a) a single vortex imaged insitu after an interaction ramp from $g \approx 0.1$ to 0.0056, (b) a single vortex imaged after a 40 ms 2D TOF at $g \approx 0.0056$, (c) and (d) vortex dipoles imaged in-situ.

Appendix C: Formation of Townes solitons

Figure 1(c) shows that, at long enough TOF times, the gas fragments into a ring of disjoint density blobs. To test whether these blobs are Townes solitons, we numerically solve the GPE with zero winding number s = 0 to obtain the steady-state solution $\phi_{TS}(R)$. The density profile of Townes solitons must have the form [37, 38]

$$n(r) = n_p \left| \phi_{TS} \left(\sqrt{n_p \left| g_f \right|} r \right) \right|^2 \,. \tag{C1}$$

In Fig. 6, we randomly pick some isolated and round blobs to fit their radially averaged density profiles with




FIG. 6. Comparison of density blobs with the density profile of Townes solitons. Each panel shows a radially averaged density profile (filled circles) from the density image shown in the inset. The images in the top row are randomly selected from samples of interaction strength $g_f \approx -0.011$ and TOF time $\tau = 49$ ms, while the images in the bottom row are from samples of $g_f \approx -0.0077$ and $\tau = 53$ ms. Cyan dots in the images mark the center used for radial averaging. Solid lines are fits using Eq. (C1).

Eq. (C1), using the peak density n_p as the only free parameter. Indeed, we find reasonable agreement between the density profiles and those of Townes solitons except near the tails at $r \gtrsim 4 \ \mu m$. The deviation may be attributed to collapsed remnant gas in the background, dispersed blobs that failed to form Townes solitons, or the close proximity with other waves.

Appendix D: Extended mean-field analyses of radial wave collapse and the azimuthal modulational instability

We numerially evaluate the vortex collapse dynamics via the time-dependent GPE emulating the experimental procedures, including vortex generation via the chopstick potential and the interaction ramp-down followed by the 2D TOF.

We begin with the time-dependent 2D GPE (without the three-body loss term),

$$i\hbar\frac{\partial}{\partial t}\psi(\mathbf{r}) = \left(-\frac{\hbar^2}{2m}\nabla^2 + W(\mathbf{r}) + g|\psi(\mathbf{r})|^2 + V(\mathbf{r})\right)\psi(\mathbf{r}), \tag{D1}$$

where $\psi(\mathbf{r})$ is the condensate wavefunction, $W(\mathbf{r}) = W_0[1 + \operatorname{erf}((r - r_t)/\sigma_t)]/2$ denotes the circular confining potential, $W_0 = k_{\rm B} \times 47$ nK is the potential height, $r_t = 26 \ \mu {\rm m}$ is the radius, and $\sigma_t = 4.8 \ \mu {\rm m}$ the width of the confining potential. Here, $g = g_{3D}/(\sqrt{2\pi}l_z)$ refers to the effective 2D interaction strength obtained by considering a tight confinement along the z-direction [70], where $g_{3D} = 4\pi\hbar^2 a_s/m$, $a_s = 100a_0$, and a_0 is the Bohr radius. The last line in Eq. (D1) is the chopstick potential $V(\mathbf{r}) = V_0[1 + \text{erf}((|\mathbf{r} - \mathbf{r}_0| - r_c)/\sigma_c)]/2$, which is initially centered around $\mathbf{r}_0 = (x_0, y_0) = (-10.0 \ \mu\text{m}, -7.0 \ \mu\text{m})$ with $V_0 = W_0$, $r_c = 3.5 \ \mu\text{m}$ the radius, and $\sigma_c = 3.0 \ \mu\text{m}$.

Using a high-level computing language XMDS [71, 72], which invokes an adaptive step-size 8/9'th order Runge-Kutta method for time-stepping, and the Fast-Fourier-Transform for calculating the Laplacian, we numerically solve Eq. (D1) in the imaginary time evolution and obtain the ground state with the surface density $n_i = |\psi|^2 \approx$ $25 \, \mu \mathrm{m}^{-2}$ in the bulk. The density falls to half of this value at the radial distance $r_i \approx 21 \ \mu m$ as in the experiment. After finding the ground state, we move and split the chopstick potentials at a separation angle 65° to their final position within 400 ms while simultaneously solving the GPE in real-time. The final position of the first chopstick is $(x_f, y_f) = (0, 0)$, whereas we move the second chopstick out of the box. The size of the chopsticks is reduced to zero at their final positions within an additional 500 ms, which results in only a single vortex at the center of the wavefunction.

Following the vortex preparation sequence, we modify the mean-field wavefunction $\psi(\mathbf{r})$ by seeding noise in the form of the Bogoliubov modes,

$$\eta(\mathbf{r}) = \psi(\mathbf{r}) + \frac{1}{\sqrt{2}} \sum_{\mathbf{k}} [\alpha_{\mathbf{k}} u_k(\mathbf{r}) - \alpha_{\mathbf{k}}^* v_k^*(\mathbf{r})], \qquad (D2)$$

where $\alpha_{\mathbf{k}}$ is a random complex Gaussian noise satisfying the relation $\overline{\alpha_{\mathbf{k}}\alpha_{\mathbf{k}'}} = 0$, $\overline{\alpha_{\mathbf{k}}\alpha_{\mathbf{k}'}^*} = \delta_{\mathbf{k}\mathbf{k}'}$, and $\overline{\cdots}$ is a stochastic average. Here, $u_k(\mathbf{r})$ and $v_k(\mathbf{r})$ are the Bogoliubov modes assuming a homogeneous 2D condensate density. This procedure corresponds to seeding on-average 0.5 phonons per **k**-mode in the modified wavefunction, simulating fluctuations at zero temperature [73].

We then perform the GPE calculations in which we phenomenologically incorporate a three-body loss term $-i\hbar\zeta K_3|\psi(\mathbf{r})|^4\psi(\mathbf{r})/2$ in Eq. (D1) with a constant loss rate $K_3 = 1.2 \times 10^{-26} \text{ cm}^6/\text{s}$. This rate is around 10 times larger than a previously measured value using thermal gases prepared at $a_s \approx 0$ [74]. The factor $\zeta = 1/(6\sqrt{3\pi l_z^2})$ takes into account Bose symmetrization and integration along the z-axis. We evolve the GPE for 6 ms by inserting the wavefunction of Eq. (D2) and then slowly ramp down the trap potential height W_0 to zero within 3 ms while simultaneously ramping the interaction g to an attractive value $g_f < 0$ within 5 ms as mentioned in the main text. We then continue to evolve the GPE for additional time τ at each final interaction strength. Ten different initial random noise samples are considered for each time evolution.

Figure 7 shows the evolution of the density $n(\mathbf{r}, \tau) = |\eta(\mathbf{r}, \tau)|^2$, in a single realization, for three different in-




FIG. 7. Dynamics of the density evolution, $n(\mathbf{r}, \tau) = |\eta(\mathbf{r}, \tau)|^2$, obtained from the GPE calculation with a variable interaction strength g_f and different hold time τ as in Fig. 1(c). Image size: $(75 \ \mu \text{m})^2$.




FIG. 8. Atom loss for three different values of g_f . The black solid lines are obtained from the simulations, and the black circles are from the experiment.

teraction strengths. The simulation employs 256×256 spatial grid points for a box of 64 μ m in length. The first row of panels in Fig. 7 are snapshots of the density for $g \approx -0.0017$ $(a_s = -1.65a_0)$. Enlargement of the density defect surrounded by an enhanced ring structure similar to the experiment can be clearly seen (see Fig. 1). Wave fragmentation and enhanced density blob features appear to be more prominent as the interaction strength becomes more attractive, as shown in the last two rows of panels in Fig. 7. Inclusion of the three-body loss term suggests delay of the wavefunction collapse and soliton patterns are observed in the final snapshots. The corresponding atom loss for these three interaction strengths are presented in Fig. 8 together with atom loss measured in the experiments. One can see excellent match between the simulation and experimental data for higher interaction strengths. A slightly larger atom loss occurs in the experiment for lower interaction strengths. We attribute this to imperfect parameter matching for different experimental conditions such as slightly higher initial density or deviation of the three-body loss coefficients. Notice that within the GPE simulations, we have assumed that all parameters are fixed except for g_f .

In Fig. 2, we numerically evaluate the time evolution of the radial density profiles averaged over ten realizations of initial random noise. To identify the time scales for radial convergence in the density profiles, we adopt the procedure for evaluating the variance V as described in the main text. In Fig. 2(f), for comparing with Fig. 2(c), we




FIG. 9. Angular power spectra S_l evaluated from the timedependet GPE with $g_f \approx (a)$ -0.0017, (b) -0.0047, (c) -0.0077, and (d) -0.011, respectively. White dashed lines mark the peak l_p . (e) Averaged power spectra \bar{S} versus time and (f) rescaled power spectra versus rescaled time obtained at the interaction strengths $g_f \approx$ -0.011, -0.0077, -0.0047, -0.003, -0.0025, -0.0017, -0.001, and -0.0005 (solid curves, from dark to light gray), respectively.

have added to each $V(\tau)$ a constant offset obtained from the experiment to account for the contributions stemming from the technical noise.

To analyze the azimuthal modulational instability and compare with the experimental data, we evaluate the angular power spectra as shown in Fig. 9, from which we identify the peak position of the power spectrum S_l as described in Appendix E. The averaged power spectrum \bar{S} is evaluated within the same range $(l = 1 \sim 30)$ described in the main text. The results are shown in Fig. 9(e). Rescaled spectra $(\gamma/\gamma_i)^{\alpha}\bar{S}$ versus rescaled time $\tilde{\tau} = \gamma \tau$ are shown in Fig. 9(f) with an empirical scaling exponent $\alpha = 0.5$. The rescaled spectra are compared with the experimental data in Fig. 3(e), both exhibiting a remarkable universal behavior before the fragmentation time $\tilde{\tau} \lesssim 2$. We attribute the nonuniversal curves at $|g_f| \lesssim 0.001$ to finite size effects when r_i approaches ξ .

Appendix E: Determination of the peak position in the power spectrum S_l

In experiments and in numerical simulations, the angular power spectrum S_l developes a distinct peak at each interaction g_f . To determine the most unstable angular mode, we identify the mode number l of the maximum S_l measured at seven longest TOF times for each g_f (or at times $\tau \leq 100$ ms for GPE simulations) and then compute their mean value l_p and the standard error. The results are provided in Fig. 3(c). In Fig. 10, we compare l_p (black dashed line) with the power spectra measured experimentally for three TOF times.




FIG. 10. Angular power spectrum S_l experimentally measured at three longest TOF times with (a) $g_f \approx -0.0017$ and (b) $g_f \approx -0.0077$, respectively. Red circles represent the mean of the power spectra and the shaded band indicates the corresponding standard deviation of the mean. Black dashed line indicates l_p determined from the mean of individually identified peak-l in each spectrum.

Appendix F: Variational analysis of the azimuthal modulational instability

When the interaction strength of a homogeneous, non-rotating 2D superfluid is quenched to a negative value, a modulational instability will manifest, with selfamplifying density waves to fragment a sample into many density blobs [32]. The most unstable wavenumber is $k_{\rm MI} = \sqrt{2n_i |g_f|}$, where n_i is the initial atomic density and g_f is the attractive interaction strength. For a circular sample carrying one quantum vortex at the center, the instability may be analyzed in discrete angular modes. However, for a sample whose initial radius is several times larger than the characteristic interaction length $\xi = \pi/k_{\rm MI}$, one may still expect a local manifestation of the modulational instability, as already hinted in the multi-ring fragmentation observed in Fig. 1(c). If the instability shares the same length scale as the modulational instability, the most unstable angular mode can be expressed as $l_{\rm MI} = \bar{r}k_{\rm MI} = \bar{r}\sqrt{2n_i|g_f|}$, where \bar{r} is a mean radial position of the sample.

Our intuitive picture can be supported by a variational analysis detailed in [53, 75], which assumes a wavefunction of the form $\psi(r,\theta) = \phi(r)A(\theta,t)$ with a fixed radial function $\phi(r) = \sqrt{n(r)}$ and derives the nonlinear Schrödinger equation for $A(\theta,t)$ using a variational method. By assuming a vortex state of $A(\theta,t) = e^{i(\pm \theta + \mu t)}$ with small perturbations in discrete *l* angular modes and performing analyses for the azimuthal modulational instability, one finds the eigenfrequencies of the *l*-th mode to be

$$\omega_l = \pm \gamma_{\rm va} \sqrt{l^2 (l^2 - 2l_p^2)}, \qquad (F1)$$

where
$$\gamma_{\rm va} = \frac{\hbar}{m} \frac{\int \frac{n(r)}{r^2} r dr}{\int n(r) r dr}$$
, $l_p = \sqrt{2|g_f|\mathcal{C}}$, and $\mathcal{C} = \frac{\int n(r)^2 r dr}{\int \frac{n(r)}{r^2} r dr}$. According to Eq. (F1), ω_l becomes purely imaginary when $l < \sqrt{2}l_n$ and the modes are unstable.

The nearest integer from l_p is the most unstable mode with the largest imaginary frequency. We may carry out the radial integrations by using experimentally measured radial density profiles for n(r) in the integrant. We may also link the most unstable mode l_p with the most unstable wavenumber $k_{\rm MI}$ in the modulational instability as

$$l_p = \bar{r}_v \sqrt{2|g_f|n_i} = \bar{r}_v k_{\rm MI} \,, \tag{F2}$$

and define the effective radius for the azimuthal modulational instability as $\bar{r}_v \equiv \sqrt{C/n_i}$. While the bove variational analysis is most accurate with a fixed radial density profile, in Fig. 3(c) we estimate l_p using the density profile at the saturation time τ_{θ} of \bar{S} when the azimuthal fragmentation takes place.




FIG. 11. Azimuthal modulational instability of a vortex soliton. $\gamma_l = |\text{Im}(\omega_l)|$ denotes the growth rate at the *l*-th mode. It is only non-zero for l = 1, 2, 3, and the most unstable mode is l = 2.

Appendix G: Azimuthal modulational instability of a vortex soliton

The azimuthal modulational instability of a vortex soliton can be directly analyzed in a Bogoliubov analysis [23, 76]. We do this by introducing small perturbations to the scale-invariant stationary-state solution of a vortex soliton,

$$\psi(R,\theta,\tilde{\tau}) = \left\{ \phi_{\rm vs}(R) + \sum_{l} \left[u_l(R) e^{-i\omega_l \tilde{\tau} + il\theta} + v_l^*(R) e^{i\omega_l^* \tilde{\tau} - il\theta} \right] \right\} e^{-i\tilde{\mu}\tilde{\tau} \pm i\theta},\tag{G1}$$

where $\phi_{\rm vs}(R)$ is the radial wavefunction, $R = \sqrt{n_p |g_f|} r$ is the rescaled radial coordinate, $\tilde{\tau} = \gamma_p \tau$ is the rescaled time with $\gamma_p = \hbar n_p |g_f|/m$, and $\tilde{\mu} = \mu/(\hbar \gamma_p)$ is the scaled chemical potential as in the main text. Here, u_l and v_l^* are the Bogoliubov coefficients for the *l*-th angular mode,

l

and ω_l is the corresponding eigenvalue. If the imaginary part of ω_l is non-zero, the mode becomes unstable with an exponential growth rate $\gamma_l = |\text{Im}(\omega_l)|$. Substituting Eq. (G1) into the scale-invariant 2D GPE and gathering all the terms evolving in the form of $e^{-i\omega_l \tilde{\tau}}$ and $e^{i\omega_l^* \tilde{\tau}}$, we obtain the Bogoliubov equations in the matrix form

$$\begin{pmatrix} -\frac{1}{2} \begin{bmatrix} \frac{\partial^2}{\partial R^2} + \frac{1}{R} \frac{\partial}{\partial R} - \frac{(s+l)^2}{R^2} \end{bmatrix} - (2|\phi_{\rm vs}|^2 + \tilde{\mu}) & -\phi_{\rm vs}^2 \\ \phi_{\rm vs}^{*2} & \frac{1}{2} \begin{bmatrix} \frac{\partial^2}{\partial R^2} + \frac{1}{R} \frac{\partial}{\partial R} - \frac{(s-l)^2}{R^2} \end{bmatrix} + (2|\phi_{\rm vs}|^2 + \tilde{\mu}) \end{pmatrix} \begin{pmatrix} u_l \\ v_l \end{pmatrix} = \omega_l \begin{pmatrix} u_l \\ v_l \end{pmatrix}, \quad (G2)$$

and further discretize the coordinate R and wavefunctions (u_l, v_l, ϕ_{vs}) with 500 grid points of step size $\Delta R = 0.03$. Here, $\left(\frac{\partial^2}{\partial R^2}\right)_{ij} = (\delta_{i,j-1} - 2\delta_{i,j} + \delta_{i,j+1})/\Delta R^2$ and $\left(\frac{1}{R}\frac{\partial}{\partial R}\right)_{ij} = (-\delta_{i,j-1} + \delta_{i,j+1})/(2\Delta RR_i)$, where i(j) is the grid index and $\delta_{i,j}$ is the Kronecker delta. We calculate the growth rate γ_l by solving the above eigenvalue problem. The result is shown in Fig. 11, where the most unstable mode is at l = 2, corresponding to the blue dashed line in Fig. 3(c). Our result is consistent with the split instability discussed in Refs. [23, 25].