

PLANNING FOR GOLD: SAMPLE SPLITTING FOR VALID POWERFUL DESIGN OF OBSERVATIONAL STUDIES

WILLIAM BEKERMAN^{*1}, ABHINANDAN DALAL^{*1}, CARLO DEL NINNO², AND DYLAN S. SMALL¹

¹*Department of Statistics and Data Science, University of Pennsylvania, Philadelphia, PA, USA*

²*Retired, Washington DC, USA*

ABSTRACT. Observational studies are valuable tools for inferring causal effects in the absence of controlled experiments. However, these studies may be biased due to the presence of some relevant, unmeasured set of covariates. The design of an observational study has a prominent effect on its sensitivity to hidden biases, and the best design may not be apparent without examining the data. One approach to facilitate a data-inspired design is to split the sample into a planning sample for choosing the design and an analysis sample for making inferences. We devise a powerful and flexible method for selecting outcomes in the planning sample when an unknown number of outcomes are affected by the treatment. We investigate the theoretical properties of our method and conduct extensive simulations that demonstrate pronounced benefits, especially at higher levels of allowance for unmeasured confounding. Finally, we demonstrate our method in an observational study of the multi-dimensional impacts of a devastating flood in Bangladesh.

1. INTRODUCTION

Observational studies have widespread presence in areas ranging from epidemiology to economics and are often employed by researchers when randomized experiments are either unethical or infeasible. Unlike a randomized experiment, however, a researcher does not control the assignment of treatment to the subjects. Even after matching on a potentially wide array of observed covariates, a researcher is unlikely to be absolutely confident about the exhaustiveness of her set of confounders and unmeasured covariates may serve to nullify causal claims between treatment and outcome (Fewell et al. [2007], Rosenbaum [2010]). Under these circumstances, the design of an observational study plays a crucial role in its robustness and validity.

Rosenbaum [2010] argues that a methodical design strengthens the power and transparency of observational analyses. He writes:

The quality and strength of evidence provided by an observational study is determined largely by its design. Excellent methods of analysis will not salvage

E-mail addresses: `bekerman@wharton.upenn.edu`.

Key words and phrases. Sensitivity analysis, confounding bias, multiple hypothesis testing.

**Denotes equal contribution.*

a poorly designed study ... An observational study that begins by examining outcomes is a formless, undisciplined investigation ...

The design of an observational study may encompass various considerations, including what protocol to use, the level of unconfoundedness at which to control, choice of test statistics, and outcomes for which to test. To best formulate such a design, a glimpse of the data is often advantageous and randomly splitting the sample into a smaller planning sample for choosing the design and a larger analysis sample for making inferences can help maintain an honest and transparent inference process, while not sacrificing validity. Although other strategies exist to conduct valid statistical inference after making data-driven decisions, such as the simultaneous confidence intervals proposed by Berk et al. [2013], these procedures are not necessarily well-posed for addressing causal questions (Small [2018]). Sample splitting on the other hand is a computationally efficient and intuitive technique with many useful qualities (Heller et al. [2009], Rosenbaum [2010], Small [2024]). Data splitting facilitates an informed choice of statistical techniques and methods, as well as outcomes for which to test, after looking at the data without the risk of p -hacking. In practice, a researcher would aim to be deliberate in how she allocates the power of her data analysis, preferring to select outcomes which would have a reasonable chance of being significant in the analysis sample, even after accounting for potential confounding biases. The goal of this paper is to develop a powerful method for selecting outcomes to test on the analysis sample from the planning sample data. As a motivating example, we consider the wide-ranging effects of the 1998 floods in Bangladesh, which endangered the health and lives of millions through food shortages, diminished purchasing power for essential goods, and the potential spread of water-borne illnesses (del Ninno et al. [2001]). We employ split-sampling to establish hypotheses likely to be less sensitive to unmeasured biases, helping researchers to identify the specific impacts of treatment; in this case, evaluating whether these floods reduced food availability, decreased access to sanitary water, or worsened illness intensity, to highlight a few possibilities, among residents in Bangladesh following these disasters.

Seeking to improve design under concerns of potential unmeasured confounding, Heller et al. [2009] utilize sample splitting to compare design sensitivities (Rosenbaum [2004]). They demonstrate that sample splitting may reduce power, but leaves design sensitivity unchanged. The most salient rebuttal against the use of data splitting has been the loss of power resulting from

the decrease in sample size. Cox [1975] utilized sample splitting in randomized trials to choose several hypotheses. Similar approaches have been employed by Wasserman and Roeder [2006], weighting hypotheses based on estimated signal strengths, and Rubin et al. [2006], estimating optimal cutoffs in a multiple testing problem. Unfortunately, the gain in power using clever techniques does not generally compensate the loss due to the reduced sample size in randomized experiments. Yet Heller et al. [2009] show that sample splitting can be powerful in the context of observational studies. Although their work constitutes an illuminating proof of concept, their approach of choosing the smallest p -value is largely heuristic-based and is limited to either one specific outcome or one general outcome of interest which can manifest itself through various avenues. In his defense of the use of protocols for observational studies, Small [2024] (Section 6.1) raises the choice of how many and which outcomes to select in a planning sample as an open problem.

In this article, we provide an adaptive procedure to select outcomes to test in an observational study. Our method is based on the sensitivity value, coined first by Zhao [2018] but frequently reported in observational studies since Cornfield et al. [1959]. Defined as the smallest level of unmeasured confounding $\tilde{\Gamma}^l$ required in sample to invalidate the significance of a causal effect, the sensitivity value is a sample-based random variable that captures the trade-off between finite sample efficiency and bias insensitivity more succinctly than the design sensitivity. In more detail discussed in the subsequent sections, our strategy has the advantage of dealing with the behavior of a finite sample, as it is based on the random variable (sensitivity value) rather than its asymptotic limit parameter (design sensitivity). In particular, our method computes the sensitivity value of each outcome on the planning sample and estimates its corresponding variability to construct predictive intervals on the analysis sample. Intuitively, it uses the information contained in the planning sample to predict whether or not each outcome will have a reasonable chance of proving significant in the latter sample, allowing us to make an informed choice about how many and which outcomes to carry over for analysis. Our approach demonstrates competitive power compared to the full sample Bonferroni correction in observational studies, while allowing for the full benefits of data splitting to be reaped. Simulation results suggest that benefits are more pronounced with increasing wariness about unmeasured confounding; that is, increasing Γ -levels at which to control.

We outline our article as follows: in Section 2, we provide the notation we will use in the article and provide background on treatment assignment and sensitivity analysis for observational

studies. We discuss methods for outcome selection and provide some theoretical arguments in Sections 3 and 4. In Sections 5 and 6, we conduct simulation studies and an application of the proposed methods to study the specific household-level effects of Bangladesh flooding. We conclude the article with a brief discussion in Section 7. An extension to multiple controls, technical proofs, additional design considerations, and details of the data application can be found in the Appendix.

2. NOTATION AND REVIEW

2.1. Treatment Assignments and Treatment Effects. Consider an observational study with I matched sets, $i = 1, \dots, I$, where the i^{th} set contains $n_i \geq 2$ subjects, $j = 1, \dots, n_i$, one of which is treated with $Z_{ij} = 1$, while the others are controls with $Z_{ij} = 0$, such that $\sum_{j=1}^{n_i} Z_{ij} = 1$ and $N = \sum_{i=1}^I n_i$ subjects in total. Sets are matched for observed covariates $x_{ij} = x_{ij'}$ for all i, j, j' , but a researcher may be concerned about an unobserved covariate possibly leading to $u_{ij} \neq u_{ij'}$ for some i, j, j' . There are L outcomes in which the researcher may be interested. Let $r_{T_{ij}}^l$ and $r_{C_{ij}}^l$ denote the potential outcomes for the l^{th} outcome for the j^{th} subject in the i^{th} matched set when assigned to treatment and control, respectively, for $l = 1, \dots, L$. The researcher gets to observe $R_{ij}^l = Z_{ij}r_{T_{ij}}^l + (1 - Z_{ij})r_{C_{ij}}^l$; hence, the individual treatment effect $r_{T_{ij}}^l - r_{C_{ij}}^l$ is not calculable Neyman [1923], Rubin [1974]. Take $\mathcal{F} = \{(x_{ij}, u_{ij}, r_{T_{ij}}^l, r_{C_{ij}}^l) \mid i = 1, \dots, I; j = 1, \dots, n_i; l = 1, \dots, L\}$ and \mathcal{Z} be the event $\sum_{j=1}^{n_i} Z_{ij} = 1$. Let $\mathbf{R}^l = (R_{11}^l, R_{12}^l, \dots, R_{In_I}^l)^T$, $\mathbf{R} = (\mathbf{R}^1, \dots, \mathbf{R}^L)^T$, and $\mathbf{Z} = (Z_{11}, Z_{12}, \dots, Z_{In_I})^T$. We use Ω to denote the support of \mathbf{Z} .

In a randomized experiment, a subject within each matched set i is randomly selected to receive treatment with probability $1/n_i$. Treatment assignment is independent across distinct matched sets in this setting, ensuring that $\mathbb{P}(z) = 1/|\Omega|$ for all $z \in \Omega$, where $|\cdot|$ refers to the size of a finite set (Fisher [1935]). The sharp null posits $H_0^l : r_{T_{ij}}^l = r_{C_{ij}}^l$ for all i, j . An observational study seeks to emulate a randomized experiment (Cochran [1965], Rosenbaum [2010] Section 1.2) by matching on the observed covariates so that any departure of the observed outcomes from the null distribution can be ascribed directly to the treatment. However, in an observational study, failure to control for some relevant, unobserved confounder can undermine the attribution of a causal effect. Thus, a researcher often looks to account for some level of bias due to unmeasured confounding.

2.2. Sensitivity Analysis. In order to evaluate how the conclusions of a study would change under different levels of unmeasured confounding, a typical strategy is to perform sensitivity analyses of the results. One common framework is the Rosenbaum sensitivity model (Rosenbaum [2002]), parameterized by a single parameter Γ , which quantifies the amount of departure of the observed treatment exposure in a matched study from random assignment. A researcher often calculates a set of p -values $[p_\Gamma, \bar{p}_\Gamma]$ in these analyses, whereas $p_1 = \bar{p}_1$. If these p -values remain significant at larger values of Γ , then an analysis is more robust to possible violations of the unconfoundedness assumption.

The Rosenbaum sensitivity model presupposes that the odds of receiving treatment in a matched set with the same values of \mathbf{x} is given by

$$(2.1) \quad \frac{1}{\Gamma} \leq \frac{\mathbb{P}(Z_{ij} = 1 | \mathcal{F}, \mathcal{Z})}{\mathbb{P}(Z_{ik} = 1 | \mathcal{F}, \mathcal{Z})} \leq \Gamma, \quad 1 \leq j, k \leq n_i, 1 \leq i \leq I$$

with independent assignment across different matched sets. A value of $\Gamma = 1$ yields random assignment, whereas any value $\Gamma \geq 1$ implies an unknown but restricted departure from random assignment due to some unobserved covariate. In the subsequent discussion, we focus on the special case of matched pairs; that is, $n_i = 2$. This condition is relaxed in the Appendix.

Let $Y_i^l = (Z_{i1} - Z_{i2})(R_{i1}^l - R_{i2}^l)$ denote the treatment-control difference in the i^{th} matched set for the l^{th} outcome. A popular choice of statistics for testing the null are the signed score statistics

$$(2.2) \quad T^l(Z, R) = \frac{\sum_{i=1}^I \text{sgn}(Y_i^l) q_i^l}{\sum_{i=1}^I q_i^l}$$

where $\text{sgn}(y) = \mathbb{1}(y \geq 0)$ and q_i^l is some function of $|Y_i^l|$ such that $q_i^l = 0$ if $Y_i^l = 0$. The statistic in Equation 2.2 has been normalized by $\sum_{i=1}^I q_i$ following Zhao [2018] so that it remains confined between zero and one. For instance, $q_i^l = \mathbb{1}(|Y_i^l| > 0)$ yields the sign-statistics and $q_i^l = \text{rank}(|Y_i^l|)$ gives the well-known Wilcoxon signed rank statistics. Under the null hypothesis and assuming random treatment assignment (i.e., $\Gamma = 1$ in Equation 2.1), conditional on \mathcal{F} and \mathcal{Z} , $\text{sgn}(Y_i^l)$ are independent and identically distributed Bernoulli(1/2) random variables and q_i^l 's are fixed constants, from which the null distribution of the statistics can be derived. However, the null hypothesis when $\Gamma > 1$ is composite so one seeks to maximize the p -value subject to Equation 2.1.

Rosenbaum [2002] showed that one can obtain $\underline{p}_\Gamma^l := \mathbb{P}(T_\Gamma^l \geq t | \mathcal{F}, \mathcal{Z}) \leq \mathbb{P}(T^l \geq t | \mathcal{F}, \mathcal{Z}) \leq \mathbb{P}(T_\Gamma^l \geq t | \mathcal{F}, \mathcal{Z}) =: \bar{p}_\Gamma^l$, where T_Γ^l is the sum of I independent random variables taking the value q_i^l with probability $\Gamma/(1 + \Gamma)$ and zero with probability $1/(1 + \Gamma)$, whereas $T_\Gamma^l = T_{1/\Gamma}^l$. A standard sensitivity analysis computes the plausible p -values \underline{p}_Γ^l and \bar{p}_Γ^l for the statistic in Equation 2.2 under Model 2.1 using the previous result. Indeed, for large sample size I , T_Γ^l satisfies a central-limit theorem (Sidak et al. [1999]). Conditional on \mathcal{F} and \mathcal{Z} , we have

$$(2.3) \quad \frac{\sqrt{I} \left(\bar{T}_\Gamma^l - \frac{\Gamma}{1+\Gamma} \right)}{\sqrt{\frac{\Gamma}{(1+\Gamma)^2} (\sigma_q^l)^2}} \xrightarrow{d} \mathcal{N}(0, 1), \text{ when } (\sigma_q^l)^2 := \lim_{I \rightarrow \infty} (\sigma_{q,I}^l)^2 = \lim_{I \rightarrow \infty} \frac{\sum_{i=1}^n (q_i^l)^2 / I}{(\sum_{i=1}^n q_i^l / I)^2} < \infty$$

Another common practice in observational studies is to report the sensitivity value (Zhao [2018]), defined as $\Gamma_\alpha^*(\mathbf{Z}, \mathbf{R}^l) = \inf\{\Gamma > 0 : \bar{p}_\Gamma^l(\mathbf{Z}, \mathbf{R}^l) \geq \alpha\}$. Put simply, the sensitivity value is the smallest level of confounding required to render the p -value insignificant. While a typical sensitivity analysis is conducted for $\Gamma \geq 1$, the model is well-defined for any positive Γ , and defining it as before removes the mass at one. Although the name “*sensitivity value*” is relatively new, it has been widely reported in observational studies to underpin the robustness of a study to unmeasured biases. Before proceeding to the next section, it is important to substantiate the difference between sensitivity value and the more commonly used design sensitivity $\tilde{\Gamma}^l$. Design sensitivity is defined as the parameter $\tilde{\Gamma}^l$ such that H_{0l} is rejected under true causal effect and no unmeasured confounding with probability one when $\Gamma < \tilde{\Gamma}^l$ and with probability zero when $\Gamma > \tilde{\Gamma}^l$ as $I \rightarrow \infty$ (Rosenbaum [2004]). Sensitivity value, on the other hand, is a deterministic function of the data, and is hence a random variable instead of an asymptotic concept. In particular, design sensitivity is the stochastic limit of the sensitivity value as $I \rightarrow \infty$ and is insensitive to the sample size.

3. METHOD

3.1. Principles of Outcome Selection. A researcher may use sample splitting to help design an observational study. We describe this setup precisely in Framework 1.

Framework 1. (*Sample splitting*). Consider the dataset randomly partitioned into planning and analysis sets, and a design choice based on the planning sample, encoded by $\mathcal{G}_{\text{planning}}$. Let there be L possible outcomes to test, of which $\mathcal{G}_{\text{planning}}$ decides to test a subset S . A potential choice for the level at which outcome l is tested is $\alpha_l \mathbb{1}(l \in S)$, where $\sum_{l \in S} \alpha_l = \alpha$, $l \in \{1, \dots, L\}$.

Proposition 1. *Framework 1 satisfies:*

$$\mathbb{P}(\exists l \in \{1, \dots, L\} : H_{0l} \text{ is true and rejected in the analysis sample} \mid \mathcal{G}_{\text{planning}}) \leq \alpha$$

Proof: $\mathbb{P}(\exists l : H_{0l} \text{ is false and rejected} \mid \mathcal{G}_{\text{planning}}) \leq \sum_{l=1}^L \mathbb{P}(H_{0l} \text{ is rejected} \mid \mathcal{G}_{\text{planning}}, H_{0l} \text{ is true})$
 $\leq \sum_{l=1}^L \alpha_l \mathbb{1}(l \in S) = \alpha.$ ■

Remark: Proposition 1 implies that the family-wise error rate (FWER) of Framework 1 is controlled at level α .

Rosenbaum [2004] prefers designs with higher design sensitivity, all else being equal. However, design sensitivity is a complicated function of outcome signal strength, study design, and statistics used, and is not easily computed for several choices of test statistics. Moreover, the outcome signal strength is usually unknown and must be estimated in to obtain an estimator for design sensitivity.

One approach is to estimate the design sensitivity via the sensitivity value in the planning sample, as this is known to be consistent for design sensitivity (Zhao [2018]). A researcher could then select those outcomes for which the sensitivity value is larger than control level Γ_{con} , which is the level of bias the researcher aims to account for in the analysis sample. A natural implementation is thus to directly test for the outcomes at some α and Γ_{con} , which is delineated in Method 1.

Method 1. (Naïve). *Consider Framework 1. To control for bias at level Γ_{con} in the analysis sample, test for those outcomes $H_{0l} : l \in S_{\text{plan}}^{\text{Naïve}}(\Gamma_{\text{con}}, \alpha)$, where*

$$S_{\text{plan}}^{\text{Naïve}}(\Gamma_{\text{con}}, \alpha) = \{l \in \{1, \dots, L\} : \bar{p}_{\Gamma_{\text{con}}}^l(\mathbf{Z}, \mathbf{R}^l) \leq \alpha\}$$

Method 1 is a practical implementation of Rosenbaum’s principle of choosing designs based on maximizing design sensitivity (Rosenbaum [2004], Rosenbaum [2010]). In some settings, such as observational studies with moderate to large sample sizes and high levels of control for bias due to unmeasured confounding, this strategy can be useful in selecting outcomes prior to data analysis (Heller et al. [2009]). As an example, consider a high-dimensional regime with $N = 1000$ subjects randomly assigned to treatment or control with probability 1/2 and $L = 500$ outcomes of possible interest. Suppose that we wish to maintain considerable control for potential confounding biases in the analysis sample, leading us to set $\Gamma_{\text{con}} = 3.5$. All control

potential outcomes $r_{C_{ij}}^l$ are generated independently from the standard Normal distribution and treated potential outcomes are taken as:

$$r_{T_{ij}}^l = \begin{cases} r_{C_{ij}}^l + 1, & \text{if } l \in \{1, \dots, 10\}; \\ r_{C_{ij}}^l, & \text{if } l \in \{11, \dots, 500\}. \end{cases}$$

Using the Wilcoxon signed rank statistic and a planning sample proportion of 0.20, the power for the Naïve method is approximately 0.24 while the power for Bonferroni correction without data splitting is around 0.16 when averaged across 1,000 simulations at $\alpha = 0.05$.

Unfortunately, there also exist numerous situations which do not lend themselves well to this approach for the design of a study. For instance, the Naïve method often performs poorly compared to full-sample techniques in lower-dimensional settings and when controlling for small values of Γ_{con} . Despite its intuitive appeal and concise implementation, these shortcomings warrant the development of a more sophisticated technique to select outcomes and enhance design. The procedure we introduce next will consider the sensitivity value and its estimated variability for each outcome on the planning sample to construct predictive intervals on the analysis sample and render a more robust selection of outcomes.

3.2. Sensitivity Value Based Outcome Selection. In a data splitting regime such as Framework 1, a researcher prioritizes the meaningful filtering of outcomes so that those tested in the analysis sample have a reasonable chance of turning out to be significant. As discussed in Section 2.2, the sensitivity value is a deterministic function of the data instead of an asymptotic concept like design sensitivity, so we can expect outcome selection using the sensitivity value to be more purposeful in finite samples, especially when working with small or moderate sample sizes.

Under $\Gamma_{\alpha}^*(\mathbf{Z}, \mathbf{R}^l)$, we define the transformed sensitivity value $\kappa_{\alpha}^l(\mathbf{Z}, \mathbf{R}^l) = \frac{\Gamma_{\alpha}^*(\mathbf{Z}, \mathbf{R}^l)}{1 + \Gamma_{\alpha}^*(\mathbf{Z}, \mathbf{R}^l)}$. In a data splitting framework, we partition $(\mathbf{Z}, \mathbf{R}) = \cup_{k \in \{\text{planning}, \text{analysis}\}} (\mathbf{Z}_k, \mathbf{R}_k)$ and modify the notation accordingly. Now, suppose that the statistic T^l , as defined in Equation 2.2, satisfies

$$(3.1) \quad \frac{\sqrt{I}(T^l - \mu_F^l)}{\sigma_F^l} \xrightarrow{d} \mathcal{N}(0, 1).$$

This assumption is valid under weak regularity conditions (Hettmansperger [1984]). Using Results 2.3 and 3.1, Zhao [2018] showed that one may obtain

$$(3.2) \quad \sqrt{I}(\kappa_{\alpha}^l - \mu_F^l) \xrightarrow{d} \mathcal{N}(-\sigma_q^l \Phi^{-1}(1 - \alpha) \sqrt{\mu_F^l(1 - \mu_F^l)}, (\sigma_F^l)^2),$$

where μ_F^l is related to Rosenbaum's design sensitivity as $\mu_F^l = \tilde{\Gamma}^l / (1 + \tilde{\Gamma}^l)$, Φ denotes the standard Normal distribution function, and subscript F signifies dependence on the distribution of outcomes.

Suppose we test outcome l in the analysis sample at level α_l . We know that outcome l will be deemed significant if $\kappa_{\alpha_l}^l(\mathbf{Z}_{\text{analysis}}, \mathbf{R}_{\text{analysis}}^l) > \kappa_{\text{con}}$, where $\kappa_{\text{con}} = \Gamma_{\text{con}} / (1 + \Gamma_{\text{con}})$ is the transformed bias level at which the researcher would like to control. If we can create a predictive interval for $\kappa_{\alpha_l}^l(\mathbf{Z}_{\text{analysis}}, \mathbf{R}_{\text{analysis}}^l)$, abbreviated as $\kappa_{\alpha_l}^l$, based on the data from the planning sample (containing rI matched pairs, $r \in (0, 1)$), then we should be able to see whether that interval lies towards the right of κ_{con} and select the outcomes to test accordingly.

Unfortunately, $\kappa_{\alpha_l}^l$ is a function of the analysis sample and is hence independent of the planning sample. However, Equation 3.2 tells us that the sensitivity value at level α_{plan} in the planning sample has the same centering and scaling as the sensitivity value in the analysis sample. We can thus obtain a predictive interval for $\kappa_{\alpha_l}^l$ based on $\kappa_{\alpha_{\text{plan}}}^l(\mathbf{Z}_{\text{plan}}, \mathbf{R}_{\text{plan}}^l)$, abbreviated as $\kappa_{\alpha_{\text{plan}}}^l$, which we delineate in Theorem 3.1.

Theorem 3.1. *Assume conditions 2.3 and 3.1 hold. Then, we have*

$$\sqrt{I} \left(\kappa_{\alpha_l}^l - \kappa_{\alpha_{\text{plan}}}^l \right) + \sqrt{\kappa_{\alpha_{\text{plan}}}^l (1 - \kappa_{\alpha_{\text{plan}}}^l)} \sigma_q^l \left[\frac{\Phi^{-1}(1 - \alpha_{\text{plan}})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1 - r}} \right] \xrightarrow{d} \mathcal{N} \left(0, \frac{(\sigma_F^l)^2}{r(1 - r)} \right)$$

We need one final detail to derive a confidence set for $\kappa_{\alpha_l}^l$. The astute reader will note that σ_F^l is still unknown in Theorem 3.1 and cannot be estimated directly since $\kappa_{\alpha_{\text{plan}}}^l$ can be only obtained once in the planning sample. To mitigate this issue, we take bootstrap samples of the matched pairs in $\mathcal{G}_{\text{plan}}$ and compute the sensitivity values $\{\kappa_{\alpha_{\text{plan}}, b}^l\}_{1 \leq b \leq B}$. We discuss the potential benefits that arise from our use of the bootstrap in Subsection 4.1. Then, we can estimate $(\hat{\sigma}_F^l)^2 = I \widehat{\text{Var}}(\kappa_{\alpha_{\text{plan}}, b}^l)$, where $\widehat{\text{Var}}(\cdot)$ denotes the sample variance. Together, this constitutes our Sens-Val Method 2.

Method 2. (Sens-Val). *Consider Framework 1. To control for bias at level Γ_{con} in the analysis sample, for each outcome l :*

- Compute $\kappa_{\alpha_{\text{plan}}}^l$, the sensitivity value for the l^{th} outcome at level α_{plan} .
- Take B bootstrap samples of I_{plan} matched pairs from $(\mathbf{Z}_{\text{plan}}, \mathbf{R}_{\text{plan}}^l)$ and calculate sensitivity values $\{\kappa_{\alpha_{\text{plan}}, b}^l\}_{1 \leq b \leq B}$. Compute $(\hat{\sigma}_F^l)^2 = I \widehat{\text{Var}}(\kappa_{\alpha_{\text{plan}}, b}^l)$.

- Test for the outcomes $H_{0l} : l \in S_{plan}^{Sens-Val}(\Gamma_{con}, \alpha_{plan})$, where

$$S_{plan}^{Sens-Val}(\Gamma_{con}, \alpha_{plan}) = \left\{ l \in [L] : \kappa_{\alpha_{plan}}^l + \frac{\sqrt{\kappa_{\alpha_{plan}}^l (1 - \kappa_{\alpha_{plan}}^l)} \sigma_q^l}{\sqrt{I}} \left[\frac{\Phi^{-1}(1 - \alpha_{plan})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1 - r}} \right] \right. \\ \left. > \kappa_{con} - \frac{\hat{\sigma}_F^l \Phi^{-1}(1 - \alpha_{coverage})}{\sqrt{Ir(1 - r)}} \right\}$$

where $[L] = \{1, \dots, L\}$, and $\{\alpha_l\}_1^L$ and $\alpha_{coverage}$ are hyper-parameters to be determined by the researcher. See Subsection 3.3 for more details.

We present a corollary to Theorem 3.1 below, which follows from the consistency of $\hat{\sigma}_F^l$.

Corollary 3.1. *If $\hat{\sigma}_F^l$ is consistent for σ_F^l , then*

$$\mathbb{P} \left(\kappa_{\alpha_l}^l \geq \kappa_{\alpha_{plan}}^l + \frac{\sqrt{\kappa_{\alpha_{plan}}^l (1 - \kappa_{\alpha_{plan}}^l)} \sigma_q^l}{\sqrt{I}} \left[\frac{\Phi^{-1}(1 - \alpha_{plan})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1 - r}} \right] \right. \\ \left. + \frac{\hat{\sigma}_F^l \Phi^{-1}(1 - \alpha_{coverage})}{\sqrt{Ir(1 - r)}} \right) = (1 - \alpha_{coverage}) + o(1)$$

Corollary 3.1 provides a $(1 - \alpha_{coverage})$ predictive interval for the sensitivity value of the l^{th} outcome when tested on the analysis sample at level α_l . We use one-sided confidence regions in Method 2 in order to maximize the probability that true causal effects are detected by our method. Since a signal cannot be detected if it is not tested on the analysis sample, we intend to be reasonably cautious in filtering out outcomes during the planning stage of our procedure, the validity of which is ensured by Proposition 1.

The Sens-Val method often demonstrates competitive or superior performance compared to the Naïve approach and full-sample alternatives. Consider a simple setup with $N = 200$ subjects randomly assigned to treatment or control with probability 1/2 and $L = 20$ outcomes of possible interest. Suppose that all control potential outcomes $r_{C_{ij}}^l$ are generated independently from the standard Normal distribution and treated potential outcomes are taken as:

$$r_{T_{ij}}^l = \begin{cases} r_{C_{ij}}^l + \frac{3}{4}, & \text{if } l \in \{1, \dots, 5\}; \\ r_{C_{ij}}^l, & \text{if } l \in \{6, \dots, 20\}. \end{cases}$$

Again, we consider the Wilcoxon signed rank statistic and a planning sample proportion of 0.20. While the power for Sens-Val is approximately 0.648, Bonferroni correction without data

splitting is around 0.583, and the Naïve method is approximately 0.275 when averaged across 1,000 simulations and controlled at levels $\alpha = 0.05$ and $\Gamma_{\text{con}} = 1.5$.

Recall that the aforementioned Wilcoxon signed rank statistic corresponds to $q_i = \text{rank}(|Y_i|)$. Thus, we know $\sum_i q_i = I(I+1)/2$ and $\sum_i q_i^2 = I(I+1)(2I+1)/6$. By Hettmansperger [1984], we then have $\mu_F = P(Y_1+Y_2 > 0)$ and $\sigma_F^2 = 4 [P(Y_1+Y_2 > 0) - P(Y_1+Y_2 > 0, Y_1+Y_3 > 0)]^2$. In our previous setup, Wilcoxon's test has mean $\mu_F = \Phi(3\sqrt{2}/4) \approx 0.856$, which corresponds to design sensitivity $\tilde{\Gamma}^l = \mu_F/(1 - \mu_F) \approx 5.924$, for $l \in \{1, \dots, 5\}$. As we have fixed $\Gamma_{\text{con}} = 1.5$, testing outcomes l with large design sensitivity is equivalent to testing the true signals. Imagine an oracle that tests only these true outcomes. The average power of this oracle when applied to the previous setting is roughly 0.659, which is only slightly higher, if far less realistic, than Sens-Val.

3.3. Hyperparameter Search. In the previous experiment, we set $\{\alpha_l\}_1^L = \alpha/L$ and $\alpha_{\text{coverage}} = \alpha$ for the Sens-Val procedure. Our theoretical results presume that α_l is known *a priori* and the sensitivity values for each l are calculated under this belief. However, if the researcher uses a Bonferroni correction, i.e., $\alpha_l = \alpha \mathbb{1}(l \in S_{\text{plan}}^{\text{Sens-Val}})/|S_{\text{plan}}^{\text{Sens-Val}}|$, then α_l s are implicit and the outcomes are generally not selected optimally for the criteria they would have to face in the analysis sample. In this regard, we can implement a dynamic α_l framework as follows:

- Set $\alpha_l^0 = \alpha/L$, for each l .
- For $n \geq 1$, compute $S_{\text{plan}}^{\text{Sens-Val}}(\Gamma_{\text{con}}, \alpha_{\text{plan}})^{n-1}$ at $\alpha_l = \alpha_l^{n-1}$.
- Update $\alpha_l^n = \alpha/|S_{\text{plan}}^{\text{Sens-Val}}(\Gamma_{\text{con}}, \alpha_{\text{plan}})^{n-1}|$.
- Continue till convergence of $\{\alpha_l^n\}_{n \geq 1}$ for all $1 \leq l \leq L$, and report $S_{\text{plan}}^{\text{Sens-Val}}(\Gamma_{\text{con}}, \alpha_{\text{plan}})$ as $S_{\text{plan}}^{\text{Sens-Val}}(\Gamma_{\text{con}}, \alpha_{\text{plan}})$ at the converged values of $\{\alpha_l^n\}_{n \geq 1}$ for each l .

Although this procedure does not necessarily have the same statistical guarantees with respect to the coverage of the sensitivity values in the analysis sample, we find the dynamic α_l framework tends to demonstrate slightly enhanced performance in simulations, particularly in settings with a sparse number of true signals. Intuitively, we are selecting outcomes based on criteria as stringent as they would actually face in the analysis sample, and the selection bias resulting from the dynamic choice of α_l s turns out to be insignificant. We can safely do this since Proposition 1 ensures FWER control in the analysis sample.

The Sens-Val method is relatively stable with the choice of α_{plan} , while the researcher can tune α_{coverage} based on how liberal or stringent she wishes to be. We set $\alpha_{\text{plan}} = \alpha_{\text{coverage}} = \alpha$

throughout the paper and conduct simulations and data-analysis with dynamic α 's moving forward.

4. THEORETICAL PROPERTIES

In this section, we consider further theoretical rationale behind Method 2. Although our primary objective is to justify the performance of Sens-Val with respect to selection of outcomes, this section contains results that may be of independent theoretical interest. We begin in Subsection 4.1 by deriving a novel Edgeworth expansion for the sensitivity value and discussing how it can be related to a bootstrap distribution leading to a higher order accuracy under standard assumptions. While previous work has demonstrated accuracy up to $O(1/\sqrt{I})$, we consider a new expansion of the transformed sensitivity value which is proven accurate up to $O(1/I)$. This complements Corollary 3.1 by demonstrating the reliability of Method 2. In Subsection 4.2, we demonstrate that our filter is powerful in selecting important outcomes under local alternatives. This suggests that even if the strength of true signals is barely above the detection threshold, Method 2 has a good chance of selecting these outcomes. Along with Framework 1, these results demonstrate that our approach is valid, accurate, and reasonably powerful, thereby illustrating why it is our recommended procedure for choosing outcomes in a planning sample.

4.1. Edgeworth Expansion and Estimation of Unknown Parameters. One of the contributions of Method 2 is the use of the bootstrap for estimation of unknown parameter σ_F^l . This allows us to construct a predictive interval for the sensitivity value in the analysis sample based only on the information contained in the planning sample, yet we must rely on the accuracy of the bootstrap procedure in order to do so.

In his work, Zhao [2018] derives an expression which relates the transformed sensitivity value κ_α^l to observed statistic T^l with accuracy up to $O(1/\sqrt{I})$. This yields a central limit theorem for the sensitivity value, although the limiting asymptotic distribution remains dependent on unknown parameters μ_F^l and σ_F^l . Employing commonly-used techniques in the bootstrap literature, we consider a new expansion of the transformed sensitivity value which is accurate up to $O(1/I)$.

Proposition 2. *Suppose the conditions of Equation 2.3 and Equation 3.1 holds, and additionally $c_{q,I}^l = \frac{\frac{1}{n} \sum_{i=1}^I (q_i^l)^3}{\left(\frac{1}{n} \sum_{i=1}^n q_i^l\right)^3}$ is such that $c_q^l = \lim_{I \rightarrow \infty} c_{q,I}^l < \infty$. Define $T_{alt}^l = \frac{\sqrt{I}(T^l - \mu_F^l)}{\sigma_F^l}$. Then,*

$$(i) \quad V^l := \frac{\sqrt{I}(\kappa_\alpha^l - \mu_F^l)}{\sigma_F^l} + \frac{z_\alpha \sigma_{q,I}^l}{\sigma_F^l} \sqrt{\kappa_\alpha^l (1 - \kappa_\alpha^l)} = T_{alt}^l - (2\mu_F^l - 1) \frac{c_1}{\sqrt{I}} + O_p\left(\frac{1}{I}\right)$$

$$(ii) \quad W^l := \frac{\sqrt{I}(\kappa_\alpha^l - \mu_F^l)}{\sigma_F^l} \text{ admits the expression}$$

$$W^l = T_{alt}^l \left[1 + \frac{\sigma_{q,I}^l z_\alpha (2\mu_F^l - 1)}{2\sqrt{I} \sqrt{\mu_F^l (1 - \mu_F^l)}} \right] - \frac{\sigma_{q,I}^l z_\alpha}{\sigma_F^l} \sqrt{\mu_F^l (1 - \mu_F^l)} - \frac{2\mu_F^l - 1}{\sqrt{I}} \left[c_1 + \frac{(\sigma_{q,I}^l)^2}{\sigma_F^l} z_\alpha^2 \right] + O_p\left(\frac{1}{I}\right)$$

where $c_1 = c_{q,I}^l g(z_\alpha) / \sigma_F^l$ and $g(x) = (x^2 - 1)/6$.

It can be seen from Proposition 2(ii) that if $I \rightarrow \infty$, one recovers the asymptotic result given as 3.2. Proposition 2 allows us to consider the Edgeworth expansions for V^l and W^l based on that for T_{alt}^l , assuming that one exists as in Equation 4.1:

$$(4.1) \quad \mathbb{P}(T_{alt}^l \leq x) = \Phi(x) + \frac{\phi(x) b^l(x, F)}{\sqrt{I}} + O\left(\frac{1}{I}\right) \text{ and } \mathbb{P}_*(T_{alt}^l \leq x) = \Phi(x) + \frac{b^l(x, F_I)}{\sqrt{I}} + O\left(\frac{1}{I}\right),$$

where $b(\cdot, F)$ is a polynomial in x that depends on the parameters of the distribution F , F_I denotes the empirical distribution of T_{alt}^l , and Φ and ϕ denote the standard Normal distribution function and density, respectively. Expansions of this form have been derived and discussed in Albers et al. [1976], where the coefficients of b^l are generally associated with skewness of the distribution of T_{alt}^l .

Theorem 4.1. *Suppose the conditions of Proposition 2 and Equation 4.1 hold. Then,*

$$(i) \quad \mathbb{P}(V^l \leq v) = \Phi(v) + \frac{1}{\sqrt{I}} \phi(v) [(2\mu_F^l - 1)c_1 + b^l(v, F)] + O\left(\frac{1}{I}\right)$$

$$\begin{aligned}
(ii) \mathbb{P}(W^l \leq w) &= \Phi \left(w + \frac{\sigma_{q,I}^l z_\alpha}{\sigma_F^l} \sqrt{\mu_F^l (1 - \mu_F^l)} \right) \\
&+ \frac{1}{\sqrt{I}} \phi \left(w + \frac{\sigma_{q,I}^l z_\alpha}{\sigma_F^l} \sqrt{\mu_F^l (1 - \mu_F^l)} \right) \left[(2\mu_F^l - 1) \left(c_1 + \frac{(\sigma^l)_{q,I}^2 z_\alpha^2}{2\sigma_F^l} \right) \right. \\
&\quad \left. - \left(w + \frac{\sigma_{q,I}^l z_\alpha}{\sigma_F^l} \sqrt{\mu_F^l (1 - \mu_F^l)} \right) \left(\frac{(2\mu_F^l - 1)\sigma_q^l z_\alpha}{2\sqrt{\mu_F^l (1 - \mu_F^l)}} \right) + b^l \left(w + \frac{\sigma_{q,I}^l z_\alpha}{\sigma_F^l}, F \right) \right] \\
&+ O\left(\frac{1}{I}\right)
\end{aligned}$$

Analogous expressions hold for the bootstrapped version with \mathbb{P} replaced by \mathbb{P}_* , μ_F and F replaced by κ_α and empirical distribution \hat{F} , respectively.

Theorem 4.1 reveals that the difference between the distribution for V^l and its bootstrapped version is $O(1/I)$. This is because

$$\mathbb{P}(V^l \leq v) - \mathbb{P}_*(V^l \leq v) = \frac{1}{\sqrt{I}} [2c_1(\mu_F^l - \kappa_\alpha) + b^l(v, F) - b^l(v, \hat{F})] + O(1/I),$$

of which $\mu_F^l - \kappa_\alpha$ and $b^l(v, F) - b^l(v, \hat{F})$ are both $O(1/\sqrt{I})$, rendering the approximation $O(1/I)$ consistent. This stands in contrast to the central limit theorem approximation, which only provides accuracy up to $O(1/\sqrt{I})$. Therefore, not only does the bootstrap-based procedure work as intended, but it can also bring about enhanced accuracy compared to the usual Normal approximation. Although Method 2 is described utilizing the Normal approximation combined with the bootstrap estimate of σ_F^l —a choice driven by its already adequate performance and to help ensure clarity on part of the reader—using the quantiles of the bootstrap distribution directly could further improve accuracy. This idea is elaborated upon in Section 7.

Theorem 4.1 also allows us to consider the accuracy of the bootstrap distribution in general and that of its moments. Wasserman [2006] discusses how the accuracy of the bootstrap of a functional relates to its Hadamard differentiability and one can observe that Method 2 is essentially a version of the non-parametric delta method. With respect to estimation of moments like σ_F^l , Bickel and Freedman [1981] uses the concept of von-Mises functionals to demonstrate the accuracy of plug-in bootstrap U-statistic-based functionals. Indeed, since $(\sigma_F^l)^2$ is the asymptotic limit of $\mathbb{E}[(\kappa_\alpha^l - (\kappa_\alpha^l)')^2]/2$, where κ_α^l and $(\kappa_\alpha^l)'$ are *iid* copies of the transformed sensitivity values, the discussion in Section 2 of Bickel and Freedman [1981] allows us to infer consistency of $\sigma_{F_{\text{boot}}}^l$.

4.2. Local Asymptotic Power. We consider the quality of the filter outlined by Method 2 in terms of the probability that it selects a true outcome when its corresponding signal is small. Although data splitting has been proposed to aim at raising the design sensitivity (Heller et al. [2009]), there is only so much a practitioner can achieve if the true causal effect lies close to the design sensitivity. In this regard, we consider the behavior of our approach when a true effect lies only marginally above the design sensitivity.

We consider a location family such that the pairwise difference in the treated-control pair is obtained by a distribution from the family $\mathcal{F}^l = \{F_\theta^l : F_\theta^l \text{ admits a density of form } f_l(\cdot - \theta)\}$, where θ quantifies the treatment effect. We denote $\mu_{F_\theta^l}^l$ as $\mu(\theta)$ and the same for $\sigma(\theta)$, suppressing the l superscript as it would be clear from the context. We take θ_0 such that $\mu(\theta_0) = \tilde{\Gamma}/(1 + \tilde{\Gamma})$ and $\mu(\cdot)$ to be differentiable at θ_0 . To consider the effect of moderate sample sizes without the asymptotic degeneracy with sample size, we consider the class of local alternatives $\theta_I = \theta_0 + h/\sqrt{I_{\text{plan}}} = \theta_0 + h/(r\sqrt{I})$. Although we typically do not expect a true effect to depend on the sample size, this exercise allows us to control the degeneracy of large samples. Suppose 3.1 also holds under this local alternative with θ_I given as above. This is not directly implied by 3.1, as now the distribution parameters also change with the sample size I . Nonetheless, the local version of 3.1 can hold under contiguity arguments, which we discuss in Proposition 3. This version also allows us to obtain Theorem 4.2.

Proposition 3. *Suppose the statistic T^l in Equation 2.2 be such that q_i^l is generated by $q_i^l = \psi(\text{rank}(|Y_i^l|)/(I+1))$ for some differentiable function ψ on $[0, 1]$, such that $0 < \int_0^1 \psi(u) du < \infty$, $\int_0^1 \psi(u)^2 < \infty$ and ψ' is bounded. Also, let $F_\theta^l \in \mathcal{F}^l$ is in a location family that admits a positive continuously differentiable density. Then, Equation 3.1 can be strengthened to obtain,*

$$(4.2) \quad \frac{\sqrt{I}(T^l - \mu^l(\theta_I))}{\sigma^l(\theta_I)} \xrightarrow{d} \mathcal{N}(0, 1)$$

with $\theta_I = \theta_0 + h/\sqrt{I}$, $h > 0$.

Theorem 4.2. *Suppose that Equation 4.2 holds and $\mu(\theta)$, $\sigma(\theta)$ are differentiable and continuous, respectively, at point θ_0 . Then, assuming $\theta_I = \theta_0 + h/\sqrt{I_{\text{plan}}}$, where θ_0 is such that $\mu(\theta_0) = \Gamma_{\text{con}}/(1 + \Gamma_{\text{con}})$, the local power, i.e., the probability that a true outcome with location parameter θ_I is selected via Method 2, converges to*

$$1 - \Phi \left(-\frac{z_{\alpha_{\text{coverage}}}}{\sqrt{1-r}} - h \frac{(\mu^l)'(\theta_0)}{\sigma^l(\theta_0)} + z_{\alpha_{\text{plan}}} \frac{\sqrt{\Gamma_{\text{con}}}\sigma_q}{(1 + \Gamma_{\text{con}})\sigma(\theta_0)} \frac{\sqrt{r}}{\sqrt{1-r}} \right).$$

Theorem 4.2 reveals that Method 2 is powerful even under local alternatives. For instance, let us consider the setting where we use the Wilcoxon signed rank statistic and control for potential unmeasured confounding bias at level $\Gamma_{\text{con}} = 2$, implying a transformed design sensitivity of $2/3$. Under the sequence of alternatives given as $\theta = \Phi^{-1}(2/3)/\sqrt{2} + h/\sqrt{I_{\text{plan}}}$, where $h = 0.20$ and $r = 0.20$, Method 2 would have a local power of 0.997, based on the calculations of $\mu(\theta)$ and $\sigma(\theta)$ given by Zhao [2018].

While the second term in the expression given by Theorem 4.2 suggests possible type I error inflation in the planning stage, we note that the primary objective in constructing our filter is establishing its considerable power. Our procedure is formulated intentionally to diminish the probability that true signals are removed prematurely from the pipeline, while still being able to reduce the number of outcomes examined in the analysis stage effectively. Since the validity of our entire procedure is ensured by Proposition 1 and Corollary 3.1 guarantees asymptotic coverage for the sensitivity values in the analysis sample, the substantial power of Method 2 under local alternatives motivates it as our recommended choice when sample splitting.

5. SIMULATION STUDIES

We conduct simulation studies to assess the performance of the Sens-Val procedure (Method 2). We also implement the Naïve approach (Method 1) and standard Bonferroni correction for purpose of comparison. These experiments are intended to examine the efficacy of our procedure in both large samples and for our data application. In particular, we emulate the setting proposed by Yadlowsky et al. [2022] and show the potential of Sens-Val compared to alternatives. We then assess the effectiveness of each method with simulated data that has the same dimensions as the Bangladesh flooding dataset.

The purpose of our first simulation study is to assess the performance of our sensitivity value-based split-sampling framework compared to a Bonferroni correction alternative that does not partition the data. We take a total number of subjects $N = 5000$ and number of outcomes $L = 250$, from which we assume there are five outcomes that constitute “true” signals in our data. We choose $d = 20$ covariates and generate $X \sim \text{Uniform}[0, 1]^d$. Then, conditional on $X = x$, we take $U \sim \mathcal{N}(0, (1 + \sin(3x_1)/2)^2)$. Amongst our five “true” signals, we set our potential outcomes $r_{C_{ij}}^l = \beta^\top x + \varepsilon$ and $r_{T_{ij}}^l = \tau + r_{C_{ij}}^l$, while the remaining potential outcomes are generated as: $r_{C_{ij}}^l = r_{T_{ij}}^l = \beta^\top x + U + \varepsilon$, where $\beta \sim \mathcal{N}(1, 1)$, $\mu \sim \mathcal{N}(0, 1)$, and $\varepsilon \sim \mathcal{N}(0, 1/4)$ are independent.

Seeking to incorporate the variability characteristic of observational data, we draw the proportion of subjects who are assigned treatment from $\text{Beta}(10, 10)$. We then draw the treatment assignment according to $Z \sim \text{Bernoulli}\left(\frac{\exp(\alpha_0 + x^\top \mu - \log(\Gamma_{\text{data}})1\{u > 0\})}{1 + \exp(\alpha_0 + x^\top \mu - \log(\Gamma_{\text{data}})1\{u > 0\})}\right)$, where α_0 is a constant to calibrate the overall treatment assignment ratio close to the desired proportion. Note that for a valid sensitivity analysis, we must have $\Gamma_{\text{con}} \geq \Gamma_{\text{data}}$. We consider two setups, one when there is no unmeasured confounding (NUC) in any of the outcomes (unknown to the researcher), where we set $\Gamma_{\text{data}} = 1$, and the other where only the false outcomes suffer from unmeasured confounding (UC) but not the true outcome, setting $\Gamma_{\text{data}} = \Gamma_{\text{con}}$. Note that, even in the latter situation, we do not include unmeasured confounding in the true potential outcomes, akin to the Rosenbaum favorable setting (Rosenbaum [2010], Section 14.2). Propensity score pair matching with a caliper is implemented to balance covariate distributions between treated and control units. 20% of our total matched pairs are randomly selected to be used in the planning sample, while the remaining data are assigned to the analysis sample. We vary both τ and Γ_{con} across experiments to evaluate the performance of the methods, all of which consider a possible positive treatment effect for each outcome l . All experiments take 250 bootstrap samples and the average power over all the true outcomes across 1000 simulations to produce a univariate measurement of power.

In Figure 1, we display plots showing the results of Sens-Val, the Naïve approach, and Bonferroni correction in the NUC and UC settings at multiple values of Γ_{con} . We observe that Sens-Val demonstrates competitive performance across values of Γ_{con} and τ , with increasing benefits at increasing levels of wariness for potential unmeasured confounding biases.

In the second set of experiments, we investigate the behavior of the discussed methods on simulated data with the same dimensions as the Bangladesh flooding dataset. To match the real data, we take a total number of $N = 757$ subjects, $d = 33$ covariates, and $L = 93$ outcomes of interest, from which we assume there are five outcomes that constitute “true” signals. The remaining variables are generated the same as the previous simulation. In particular, unmeasured confounding U , potential outcomes, and treatment assignment are taken the same as in the large-sample setting. However, the proportion of subjects who are assigned treatment is taken equal to 0.71, consistent with the actual ratio of treated to control subjects in the data.

In Figure 2, we display plots showing the results of Sens-Val, the Naïve approach, and Bonferroni correction in the NUC and UC settings at multiple values of Γ_{con} . We observe that, despite maintaining just 80% of the sample for inference, Sens-Val demonstrates competitive




FIGURE 1. Simulation results in the large sample setting. (A) Results for the NUC setting. (B) Results for the UC setting.

or superior power across values of Γ_{con} and τ , with increasing benefits at increasing levels of wariness for potential unmeasured confounding biases. These results show that the Sens-Val procedure can be quite useful for data of moderate sample size and suggest that it should demonstrate strong performance when applied to study the effects of the Bangladesh floods.

6. APPLICATION TO THE IMPACTS OF FLOODING IN BANGLADESH

6.1. Data Description and Preliminary Examinations. Finally, we apply each of the methods to the Bangladesh flooding dataset. In the study by del Ninno et al. [2001], a detailed household survey was conducted among 757 rural households in seven flood-affected *thanas*. The data was collected at three points in time over a period of a year between November 1998 and December 1999. We focus our analysis on the immediate impacts of the flood on food consumption, food security, health, and nutrition at the household level.




FIGURE 2. Simulation results in the data-inspired setting. (A) Results for the NUC setting. (B) Results for the UC setting.

Using the first round of household-level data, which was collected between the third week of November and the third week of December 1998, we gather covariates related to household composition, education, assets, and previous flood exposures in order to identify and match similar households. Households and *thanas* were classified by the Water Board to be not affected, moderately affected, severely affected, or very severely affected by the floods, depending on the level and depth of the floodwater. We thus define our treatment to be any of moderate, severe, or very severe flood exposures, rendering our controls to be those matched non-exposed households. We conduct propensity score pair matching with a caliper and obtain satisfactory covariate balance, as shown in the Appendix. This yields 215 matched pairs to comprise our sample. We fix our planning sample proportion equal to 0.20 and partition the data into planning and analysis samples as in Framework 1. Seeking to investigate the effects of the flood, we collect flood-specific outcomes, as well as those related to food consumption and food security,

gender discrimination, illness, nutritional status of children and women, and economic status that were examined by del Ninno et al. [2001].

Akin to previous observational studies such as Zhang et al. [2011], statisticians on our team conducted a comprehensive examination of the planning sample and followed up directly with the subject matter expert in the group (del Ninno) to discuss findings and formulate an analysis plan that reflected both his domain knowledge and our statistical judgement. For example, we initially considered outcomes related to both illness and nutritional status in the first round of data, which was collected shortly after flooding affected the region. However, the subject matter expert advised that malnutrition and growth stunting is often due to repeated illnesses and may not be detectable so soon following a disaster. To supplement his guidance, we found on the planning sample little discrepancy between matched households related to the presence of common forms of under-nutrition. On the other hand, we found credible evidence that the intensity of disease, as measured by its duration with correction for extreme values (Bhargava [1994]), may have increased with flood exposure. This motivated us to continue to analyze the impacts of the flood on illness in the first round of household-level data, while turning our focus to the third round of data, collected one year after the first round, to assess its long-term effects on nutrition. Moreover, the planning sample also revealed that several of the binary outcomes that report forms of under-nutrition are very sparse and likely too rare to study even with the much larger analysis sample. Accordingly, we restrict our analysis to only include the variables used to construct them, namely body mass index, height-for-age, weight-for-height, and weight-for-age z-scores. Removing variables with missing proportions of at least 75%, we obtain a total of 93 relevant outcomes, details of which are discussed in the Appendix.

6.2. Data Analysis. We run our Sens-Val procedure and compare it to the previous alternatives at multiple levels of control for potential unmeasured confounding Γ_{con} . Both sample splitting-based methods exploit the planning sample to estimate the sign of the average treatment effect for each outcome of interest, which is then used to formulate an alternative hypothesis and test the significance of that outcome. Since this is done only using the planning sample, this is a valid strategy which can supplement or supplant domain knowledge when desired. As standard Bonferroni correction does not enjoy the benefits of a planning sample, it conducts two-sided hypothesis tests for each outcome of interest.

For binary outcomes, sensitivity analyses and sensitivity value calculations are conducted using McNemar’s test statistic (Rosenbaum [1987], Hsu and Small [2013]), while remaining outcomes rely on Wilcoxon’s signed-rank statistic with a Normal approximation (Rosenbaum [2010]). We go through each outcome and, if either household in a matched pair has a missing value for that particular outcome, that matched pair is omitted from the sensitivity analyses. This produces valid inference under the assumption that the potential outcomes and treatment assignment are independent of the missingness mechanism conditional on the covariates (MAR; Rubin [1976]).

We set our α -level equal to 0.10 for sake of comparison with the results obtained by del Ninno et al. [2001]. The previous work uncovered some notable trends in its explorations of the data, such as a rise in food prices, illness, and credit usage, especially for food, and decreases in food production among flood-affected households. However, the authors focused primarily on discovery and did not seek to control the FWER of their analyses. We intend to provide confirmatory evidence for these findings while accounting for the testing of multiple hypotheses in the form of multiplicity correction. We vary Γ_{con} over the values in the set $\{1.00, 1.25, 1.50, 2.00, 2.50, 3.00, 4.00, 6.00\}$ and present our results in Table 1. A more detailed description of the outcomes we reject is provided in the Appendix.

As expected, all procedures reject the largest number of outcomes at small levels of Γ_{con} and the fewest number at larger levels of Γ_{con} . At $\Gamma_{\text{con}} = 1$, our results are very similar to those identified by del Ninno et al. [2001]. In particular, all three procedures identify significant differences in `credit.val` and `food.credit.val.pc`, between exposed and non-exposed households. As discussed in the report, borrowing was one of the primary coping mechanisms employed by households in response to the floods. While the proportion of households that took out any loan increased substantially during the flood period, this was especially pronounced for the purchasing of food. Specifically, the percentage of households who took out a loan for the purpose of buying food was only 7% before the flooding, but increased to almost 16% by the final month of flooding. This rise in borrowing can likely be attributed to the reduced availability of food due to the loss in food production among affected households, which was uncovered by our method and Bonferroni correction at $\Gamma_{\text{con}} = 1$, along with the growth in food price index, which was detected by the Sens-Val and Naïve methods until $\Gamma_{\text{con}} = 1.5$. Likewise, all approaches discover a significant increase in budget share used to buy pulses, such as beans,

TABLE 1. Application of proposed methods to specific household-level effects of 1998 Bangladesh floods. Cells marked with “✓” denote that the particular variable in the row was rejected by the method in the column at the given level of Γ_{con} . The shaded column reflects the results of Sens-Val Method 2.

Method No. Tested	$\Gamma_{\text{con}} = 1.00$			$\Gamma_{\text{con}} = 1.25$			$\Gamma_{\text{con}} = 1.50$			$\Gamma_{\text{con}} = 2.00$		
	Bonf. 93	Naïve 27	S-V 82	Bonf. 93	Naïve 15	S-V 22	Bonf. 93	Naïve 10	S-V 13	Bonf. 93	Naïve 6	S-V 7
ft.water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
repair.cost	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
days.water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
days.away.home	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
tubewell.cooking	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
tubewell.washing	✓	✓	✓	✓	✓	✓			✓			
food.price.idx	✓	✓	✓	✓	✓	✓		✓	✓			
food.credit.val.pc	✓	✓	✓	✓	✓	✓		✓	✓			
credit.val	✓	✓	✓	✓	✓	✓						
days.ill	✓	✓	✓		✓	✓						
pulses.budget	✓	✓	✓									
rice.consumed.pc	✓		✓									
food.produced.val	✓		✓									
food.produced.val.pc			✓									

Method No. Tested	$\Gamma_{\text{con}} = 2.50$			$\Gamma_{\text{con}} = 3.00$			$\Gamma_{\text{con}} = 4.00$			$\Gamma_{\text{con}} = 6.00$		
	Bonf. 93	Naïve 4	S-V 6	Bonf. 93	Naïve 4	S-V 4	Bonf. 93	Naïve 4	S-V 4	Bonf. 93	Naïve 4	S-V 4
ft.water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
repair.cost	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
days.water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
days.away.home	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓
tubewell.cooking			✓									
tubewell.washing												
food.price.idx												
food.credit.val.pc												
credit.val												
days.ill												
pulses.budget												
rice.consumed.pc												
food.produced.val												
food.produced.val.pc												

lentils, and peas, which was also alluded to in the report. Sens-Val and Bonferroni correction also identify a decrease in rice consumption as an effect of the flooding.

Moreover, the Sens-Val procedure uncovers the discrepancy in the proportion of households that receive their washing water from tubewells until $\Gamma_{\text{con}} = 1.5$ and the proportion of households that receive their cooking water from tubewells until $\Gamma_{\text{con}} = 2.5$. However, the Naïve method and full-sample Bonferroni correction fail to detect these effects at the same levels of allowance for potential biases, further demonstrating the robustness and power of our new technique. Since tubewells are the only source of clean water accessible to the vast majority of rural

inhabitants in Bangladesh (del Ninno et al. [2001]), residents were thus more likely to fall ill due to poor water quality and sanitation, which has been heavily documented following floods and other natural disasters (CDC et al. [2000], Du et al. [2010]). Indeed, both the Sens-Val and Naïve methods find that households exposed to the floods experienced a larger number of days with illness compared to their matched counterparts until $\Gamma_{\text{con}} = 1.5$. Although we were able to identify this augmented illness intensity among exposed households, no significant evidence of long-term nutritional detriments was discovered. This result is partially in-line with a later work by del Ninno and Lundberg [2005] which found that many of these metrics had recovered one full year after the floods.

Our Sens-Val approach consistently selects more outcomes in the planning sample to test compared to the Naïve procedure. Far from being a bug, this was a primary reason for our introduction of a more nuanced methodology. The Naïve method often suffers from lesser power due to filtering out promising outcomes in the planning stage, despite their potential to prove significant in the subsequent part of the pipeline. On the other hand, our method rejects at least as many outcomes as those rejected by Bonferroni correction for each value of Γ_{con} . Our approach also demonstrates enhanced power at large values of Γ_{con} . In particular, Bonferroni correction fails to recover any outcomes for $\Gamma_{\text{con}} \geq 16$ and the Naïve method fails to do so for $\Gamma_{\text{con}} \geq 20$. Yet Sens-Val still rejects `ft.water`, `repair.cost`, and `days.water` at $\Gamma_{\text{con}} = 38$. These outcomes are all quite intuitively related to our treatment exposure and it is thus encouraging to see our Sens-Val method identifying their significance even when accounting for very large potential biases.

7. DISCUSSION

In this work, we introduce a data-adaptive method to identify the outcomes to select from a planning sample when an unknown number of outcomes are affected by the treatment in an observational study. Our procedure relies on the sensitivity value, a widely-used but only recently formalized concept that is particularly suited for use in finite samples. We consider both the sensitivity value and its corresponding variability for each outcome in the planning sample to construct predictive intervals on the analysis sample and render a robust selection of hypotheses that have a reasonable chance of being rejected. Our approach is valid, accurate, computationally efficient, and powerful in both theory and practice, especially at higher levels of allowance for unmeasured confounding. Along with helping to improve the sensitivity of

observational studies to potential unmeasured biases, our technique allows for researchers to procure the many advantages of sample splitting with limited drawbacks. In practice, our new approach enabled us to identify the multi-dimensional effects of the 1998 Bangladesh floods with higher resolution than other strategies.

A random split-sampling framework can facilitate the development of a data-informed design while not introducing a significant cost on computational complexity or a loss of efficiency due to reduced sample size. Accordingly, our method can be adapted to consider other aspects of design for powerful analysis. Examples are provided in the Appendix. Sample splitting can also yield unforeseen insights and help motivate new ideas. For instance, in our observational study of the multi-dimensional impacts of Bangladesh floods, the statisticians on our team first conducted a thorough examination of the planning sample and followed up directly with a subject matter expert to discuss their findings and modify the analysis plan accordingly. Likewise, we can utilize the planning sample to inform the direction of the alternative hypothesis we wish to test for certain variables, especially when our understanding of the impact of treatment is limited or highly variable. While standard statistical practice would compel a researcher to obtain additional independent data to address these points, data splitting provides this flexibility based only on a planning sample.

One possible direction to extend Method 2 is inspired by the results in Section 4.1. Since the Edgeworth expansion delivers a confidence interval more accurate than the classical central limit theorem, one could obtain prediction intervals that are accurate up to $O(1/I)$. Indeed, one can easily verify that $\sqrt{r}V_{\text{plan}}^l - \sqrt{1-r}V_{\text{analysis}}^l$ equals the statistic that we use in Method 2, up to $O(1/I)$, where the sub-scripts denote the counterparts of V^l from the planning and analysis samples. By virtue of the independence of these samples, the Edgeworth expansion of this linear combination can be obtained, albeit by tedious calculation, to render a form similar to that in Theorem 4.1. The approximate quantiles of that distribution can be obtained by estimating the coefficients of the $1/\sqrt{I}$ -term, which in turn can be estimated at \sqrt{I} -rate via bootstrapping from the planning sample, yielding a $O(1/I)$ approximation of the quantiles and a corresponding coverage probability of $(1 - \alpha_{\text{coverage}}) + O(1/I)$, improving upon Corollary 3.1. However, we do not pursue this direction, as doing so would complicate the method while offering only limited insights into its working. Also, the effectiveness of our filter is based on the principle of encapsulating the first order bias and variability of the sensitivity value, and

it is unlikely that higher order bias corrections would be of substantial practical value beyond its theoretical insights.

Although our method has been presented as a means for securing FWER control, thereby rendering the Bonferroni procedure a clear baseline for comparison, our strategy is highly flexible and can be adapted easily to alternative criteria. For instance, the false discovery rate (FDR) is another metric for which control is often sought when conducting multiple comparisons. The FDR can be controlled during the variable selection process by techniques such as the Benjamini–Hochberg procedure (Benjamini and Hochberg [1995]) and model-X knockoffs (Candes et al. [2018]), the latter of which inspired the title of this article. One future direction is to modify the model-X knockoffs framework to establish FDR control when working with such a bias-concerned testing procedure.

Another possible direction could be to adapt the proposed method to the framework of cross-screening (Zhao et al. [2018]) to further promote detection power. Cross-screening randomly splits in half some data, uses each half to plan and execute a study on the other half, and then reports the more favorable results, adjusting for multiple comparisons using Bonferroni correction. While the primary focus of this paper is to describe our procedure using traditional data splitting, whereas the data is divided and a partition of the data is regarded as a pilot sample, it could be advantageous to employ our design filter in conjunction with the cross-screening framework, especially when there are many hypotheses of potential interest.

8. REPRODUCIBILITY

Bangladesh flooding dataset is publicly available at <https://doi.org/10.7910/DVN/UJA9N5>. Codes for implementing our method, reproducing numerical experiments, and conducting real data analysis is available at <https://github.com/WillBekerman/planning-for-gold>.

9. ACKNOWLEDGEMENTS

The authors thank Ting Ye for her helpful ideas and constructive suggestions.

10. FUNDING

William Bekerman was partially supported by the National Science Foundation Graduate Research Fellowship. Dylan S. Small was partially supported by the National Institutes of Health grant 1R01AG065276-01.

REFERENCES

- Willem Albers, Peter J Bickel, and Willem R van Zwet. Asymptotic expansions for the power of distribution free tests in the one-sample problem. *The Annals of Statistics*, pages 108–156, 1976.
- Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(1):289–300, 1995.
- Richard Berk, Lawrence Brown, Andreas Buja, Kai Zhang, and Linda Zhao. Valid post-selection inference. *The Annals of Statistics*, pages 802–837, 2013.
- Alok Bhargava. Modelling the health of Filipino children. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 157(3):417–432, 1994.
- Peter J Bickel and David A Freedman. Some asymptotic theory for the bootstrap. *The Annals of Statistics*, pages 1196–1217, 1981.
- Emmanuel Candès, Yingying Fan, Lucas Janson, and Jinchi Lv. Panning for gold: ‘Model-X’ knockoffs for high dimensional controlled variable selection. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80(3):551–577, 2018.
- CDC et al. Morbidity and mortality associated with Hurricane Floyd–North Carolina, September–October 1999. *MMWR. Morbidity and mortality weekly report*, 49(17):369–372, 2000.
- William G Cochran. The planning of observational studies of human populations. *Journal of the Royal Statistical Society. Series A (General)*, 128(2):234–266, 1965.
- Jerome Cornfield, William Haenszel, E Cuyler Hammond, Abraham M Lilienfeld, Michael B Shimkin, and Ernst L Wynder. Smoking and lung cancer: recent evidence and a discussion of some questions. *Journal of the National Cancer Institute*, 22(1):173–203, 1959.
- David R Cox. A note on data-splitting for the evaluation of significance levels. *Biometrika*, 62(2):441–444, 1975.
- Anirban DasGupta. *The Bootstrap*, pages 461–497. Springer New York, New York, NY, 2008. ISBN 978-0-387-75971-5.
- Carlo del Ninno and Mattias Lundberg. Treading water: the long-term impact of the 1998 flood on nutrition in Bangladesh. *Economics & Human Biology*, 3(1):67–96, 2005.

- Carlo del Ninno, Paul A Dorosh, Lisa C Smith, and Dilip K Roy. *The 1998 Floods in Bangladesh: Disaster Impacts, Household Coping Strategies, and Response*, volume 122. International Food Policy Research Institute, 2001.
- Weiwei Du, Gerard Joseph FitzGerald, Michele Clark, and Xiang-Yu Hou. Health impacts of floods. *Prehospital and disaster medicine*, 25(3):265–272, 2010.
- Zoe Fewell, George Davey Smith, and Jonathan AC Sterne. The impact of residual and unmeasured confounding in epidemiologic studies: a simulation study. *American Journal of Epidemiology*, 166(6):646–655, 2007.
- Ronald A Fisher. *The Design of Experiments*. Oliver and Boyd, Edinburgh, 1935.
- Joseph L Gastwirth, Abba M Krieger, and Paul R Rosenbaum. Asymptotic separability in sensitivity analysis. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 62(3):545–555, 2000.
- Ruth Heller, Paul R Rosenbaum, and Dylan S Small. Split samples and design sensitivity in observational studies. *Journal of the American Statistical Association*, 104(487):1090–1101, 2009.
- Thomas P Hettmansperger. *Statistical Inference Based on Ranks*. John Wiley & Sons, 1984.
- Jesse Y Hsu and Dylan S Small. Calibrating sensitivity analyses to observed covariates in observational studies. *Biometrics*, 69(4):803–811, 2013.
- Jesse Y Hsu, Dylan S Small, and Paul R Rosenbaum. Effect modification and design sensitivity in observational studies. *Journal of the American Statistical Association*, 108(501):135–148, 2013.
- Jerzy Neyman. On the application of probability theory to agricultural experiments. essay on principles. *Annals of Agricultural Sciences*, pages 1–51, 1923.
- John Winsor Pratt and Jean Dickinson Gibbons. *Concepts of nonparametric theory*. Springer, 1981.
- Paul R Rosenbaum. Sensitivity analysis for certain permutation inferences in matched observational studies. *Biometrika*, 74(1):13–26, 1987.
- Paul R Rosenbaum. Signed rank statistics for coherent predictions. *Biometrics*, pages 556–566, 1997.
- Paul R Rosenbaum. *Observational Studies*. Springer Series in Statistics. Springer, 2002.
- Paul R Rosenbaum. Design sensitivity in observational studies. *Biometrika*, 91(1):153–164, 2004.

- Paul R Rosenbaum. Sensitivity analysis for M-estimates, tests, and confidence intervals in matched observational studies. *Biometrics*, 63(2):456–464, 2007.
- Paul R Rosenbaum. *Design of Observational Studies*. Springer, 2010.
- Paul R Rosenbaum. A new U-statistic with superior design sensitivity in matched observational studies. *Biometrics*, 67(3):1017–1027, 2011.
- Paul R Rosenbaum. Weighted M-statistics with superior design sensitivity in matched observational studies with multiple controls. *Journal of the American Statistical Association*, 109(507):1145–1158, 2014.
- Paul R Rosenbaum. Combining planned and discovered comparisons in observational studies. *Biostatistics*, 21(3):384–399, 2020.
- Daniel Rubin, Sandrine Dudoit, and Mark Van der Laan. A method to increase the power of multiple testing procedures through sample splitting. *Statistical Applications in Genetics and Molecular Biology*, 5(1), 2006.
- Donald B Rubin. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology*, 66(5):688, 1974.
- Donald B Rubin. Inference and missing data. *Biometrika*, 63(3):581–592, 1976.
- Walter Rudin et al. *Principles of Mathematical Analysis*, volume 3. McGraw-Hill New York, 1976.
- Henry Scheffé. A method for judging all contrasts in the analysis of variance. *Biometrika*, 40(1-2):87–110, 1953.
- Zbynek Sidak, Pranab K Sen, and Jaroslav Hajek. *Theory of Rank Tests*. Elsevier, 1999.
- Dylan S Small. Larry Brown: Remembrance and connections of his work to observational studies. *Observational Studies*, 4(1):250–259, 2018.
- Dylan S. Small. Protocols for observational studies: Methods and open problems. *arXiv preprint 2403.19807 [stat.ME]*, 2024.
- Aad W Van der Vaart. *Asymptotic Statistics*, volume 3. Cambridge university press, 2000.
- Larry Wasserman. *All of nonparametric statistics*. Springer Science & Business Media, 2006.
- Larry Wasserman and Kathryn Roeder. Weighted hypothesis testing. *arXiv preprint math/0604172*, 2006.
- Steve Yadlowsky, Hongseok Namkoong, Sanjay Basu, John Duchi, and Lu Tian. Bounds on the conditional and average treatment effect with unobserved confounding factors. *The Annals of Statistics*, pages 2587–2615, 2022.

- Ting Ye, Dylan S Small, and Paul R Rosenbaum. Dimensions, power and factors in an observational study of behavioral problems after physical abuse of children. *The Annals of Applied Statistics*, 16(4):2732–2754, 2022.
- Kai Zhang, Dylan S Small, Scott Lorch, Sindhu Srinivas, and Paul R Rosenbaum. Using split samples and evidence factors in an observational study of neonatal outcomes. *Journal of the American Statistical Association*, 106(494):511–524, 2011.
- Qingyuan Zhao. On sensitivity value of pair-matched observational studies. *Journal of the American Statistical Association*, 114(526):713–722, 2018.
- Qingyuan Zhao, Dylan S Small, and Paul R Rosenbaum. Cross-screening in observational studies that test many hypotheses. *Journal of the American Statistical Association*, 113(523):1070–1084, 2018.

APPENDIX A. MULTIPLE CONTROLS

When $n_i \geq 2$, with there being one treated and $n_i - 1$ control units in the i th matched set, $i = 1(1)I$, one might still be interested in the treated-minus-control difference Y_{ij}^l for the l th outcome. For instance, if $n_1 = 3$, one can have two treatment-minus-control differences, Y_{11} and Y_{12} in the first matched set, such that if the first unit is treated, then $Z_{11} = 1$, $R_{11}^l - R_{12}^l$ and $R_{11}^l - R_{13}^l$, while if the second unit is treated, then $Z_{12} = 1$, the treated-minus-control differences are $R_{12}^l - R_{11}^l$, $R_{12}^l - R_{13}^l$, and so on. In general, if J_i is the unit treated in the i th matched set with $J_i = \sum_{j=1}^{n_i} jZ_{ij}$, then the $n_i - 1$ treated-minus-control differences in set i for the l th outcome are $Y_{ik}^l = R_{i,J_i}^l - R_{ik}^l$, with $1 \leq k \leq n_i, k \neq J_i$.

Now, one can look the treated-minus-control differences and compare them with the control-minus-control differences in the matched set, and thus assign a significant departure from the null distribution under no-unmeasured confounding to be an evidence in favor of the alternative. Define $N_{i,-j} = \{1 \leq k \leq n_i, k \neq j\}$, such that $N_{i,-J_i}$ are the control units in the i th matched set. Thus, one can consider the generalized test-statistic $T_g^l(\mathbf{Z}, \mathbf{R}^l) = \frac{\sum_{i=1}^I \sum_{k \neq J_i} \text{sgn}(Y_{ik}^l) q_{ik}^l}{\sum_{i=1}^I \sum_{k \neq J_i} q_{ik}^l}$, where q_{ik}^l are again appropriately chosen functions of the treated-minus-control differences and control-minus-control differences, and the statistic has been normalized. One may refer to Rosenbaum [2007] Section 4.2 for a detailed discussion. Unlike the matched pairs setting, there does not exist a bounding random variable T_{Γ}^l in the context of Model 2.1 that holds in finite samples. However, one can obtain a random variable $T_{\Gamma,g}^l$ such that, conditional on \mathcal{F} and \mathcal{Z} ,

$$(A.1) \quad \frac{\sqrt{I}(T_{\Gamma,g}^l - a_{\Gamma}^l)}{b_{\Gamma}^l} \sim \mathcal{N}(0, 1) \quad \text{and} \quad \mathbb{P}(T_{l,g} \geq t | \mathcal{F}, \mathcal{Z}) \leq \mathbb{P}(T_{\Gamma,g}^l \geq t | \mathcal{F}, \mathcal{Z}) \text{ as } I \rightarrow \infty,$$

for some appropriate choice of a_{Γ}^l and b_{Γ}^l (Gastwirth et al. [2000], Rosenbaum [2007]) when $n_i \leq p$ for all i and regularity conditions on q_i s. Now, suppose one conducts the sensitivity analysis using Equation A.1, then the sensitivity value would satisfy

$$(A.2) \quad T_g^l - a_{\Gamma_{\alpha}^*}^l(\mathbf{Z}, \mathbf{R}^l) = \Phi^{-1}(1 - \alpha_l) \frac{b_{\Gamma_{\alpha}^*}^l(\mathbf{Z}, \mathbf{R}^l)}{\sqrt{I}} + o_p\left(\frac{1}{\sqrt{I}}\right).$$

Then, suppose the true distribution of T_g^l , akin to Assumption 3.1, admits an asymptotic distribution of the form

$$(A.3) \quad \sqrt{I}(T_g^l - \mu_{Fg}^l) \xrightarrow{d} \mathcal{N}(0, (\sigma_{Fg}^l)^2).$$

Then, we can have a result analogous to 3.2.

Theorem A.1. *Suppose Equations A.1 and A.3 hold. Also, suppose that the mapping $\Gamma \mapsto a_\Gamma^l$ is continuously differentiable at a neighborhood of $\tilde{\Gamma}^l$ and a has non-zero derivative at $\tilde{\Gamma}^l$, where $\tilde{\Gamma}^l$ is the design sensitivity of the l th outcome. Then,*

$$\sqrt{I}(\Gamma_\alpha^*(\mathbf{Z}, \mathbf{R}^l) - \tilde{\Gamma}^l) \xrightarrow{d} \mathcal{N}(-\Phi^{-1}(1 - \alpha_l)\mu_{\tilde{\Gamma}^l}^g, (\sigma_{\tilde{\Gamma}^l}^g)^2)$$

for some constants $\mu_{\tilde{\Gamma}^l}^g, \sigma_{\tilde{\Gamma}^l}^g$.

We can use the same strategy as in Section 3.2 using Theorem A.1, but the expressions for $\mu_{\tilde{\Gamma}^l}^{l,g}$ are again fairly complicated. However, this time we can estimate both the mean and variance of the standardized asymptotic distribution using the bootstrap.

Method 3. (Sens-Val-Multiple-Controls). *Consider Framework 1. To control for bias at level Γ_{con} in the analysis sample when using multiple controls in matched sets, for each outcome l :*

- Compute $\Gamma_{\alpha_{plan}}^*(\mathbf{Z}_{plan}, \mathbf{R}_{plan}^l)$, the sensitivity value for the l^{th} outcome at level α_{plan} .
- Take B bootstrap samples of I_{plan} matched sets from $(\mathbf{Z}_{plan}, \mathbf{R}_{plan}^l)$ and calculate sensitivity values $\{\Gamma_{\alpha_{plan},b}^*(\mathbf{Z}_{plan,b}, \mathbf{R}_{plan,b}^l)\}_{1 \leq b \leq B}$. Compute

$$(\hat{\sigma}_{F,g}^l)^2 = I \widehat{\text{Var}}(\Gamma_{\alpha_{plan},b}^*(\mathbf{Z}_{plan,b}, \mathbf{R}_{plan,b}^l)).$$

Also, compute $\Gamma_{\alpha_{plan},boot}^*(\mathbf{Z}_{plan}, \mathbf{R}_{plan}^l)$, the sample mean of the bootstrap samples, and obtain

$$\hat{\mu}_{\tilde{\Gamma}^l}^{l,g} = \sqrt{I}(\Gamma_{\alpha_{plan}}^*(\mathbf{Z}_{plan}, \mathbf{R}_{plan}^l) - \Gamma_{\alpha_{plan},boot}^*(\mathbf{Z}_{plan}, \mathbf{R}_{plan}^l))$$

- Test for the outcomes $H_{0l} : l \in S_{plan}^{\text{Sens-Val-Multiple-Controls}}(\Gamma_{con}, \alpha_{plan})$, is given by

$$\left\{ l \in [L] : \Gamma_{\alpha_{plan}}^*(\mathbf{Z}, \mathbf{R}_{plan}^l) + \frac{\hat{\mu}_{\tilde{\Gamma}^l}^{l,g}}{\sqrt{I}} \left[\frac{\Phi^{-1}(1 - \alpha_{plan})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1-r}} \right] > \kappa_{con} - \frac{\hat{\sigma}_{F,g}^l \Phi^{-1}(1 - \alpha_{coverage})}{\sqrt{I r (1-r)}} \right\}$$

where $[L] = \{1, \dots, L\}$, and $\{\alpha_l\}_1^L$ and $\alpha_{coverage}$ are hyper-parameters to be determined by the researcher.

Theorem A.2. *If $\hat{\mu}_F^{l,g}$ and $\hat{\sigma}_F^{l,g}$ are consistent for their corresponding parameters and the conditions for Theorem A.1 hold, then*

$$\mathbb{P} \left(\Gamma_{\alpha_l}^* (\mathbf{Z}_{analysis}, \mathbf{R}_{analysis}^l) \geq \Gamma_{\alpha_{plan}}^* (\mathbf{Z}_{plan}, \mathbf{R}_{plan}^l) + \frac{\hat{\mu}_F^{l,g}}{\sqrt{I}} \left[\frac{\Phi^{-1}(1 - \alpha_{plan})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1-r}} \right] + \frac{\hat{\sigma}_F^{l,g} \Phi^{-1}(1 - \alpha_{coverage})}{\sqrt{Ir(1-r)}} \right) = (1 - \alpha_{coverage}) + o(1)$$

The proof of Theorem A.2 is *mutatis mutandis* identical to the proofs of Theorem 3.1 and Corollary 3.1, and is thus omitted. Theorem A.2 provides a way to select outcomes from the planning sample, even under choice of multiple controls.

APPENDIX B. PROOFS OF THEOREMS

B.1. **Theorem 3.1.** We suppress the superscript l in the proof for ease of notation. Using Equation 3.2, after sufficient enriching of the underlying probability space, we have

$$\begin{aligned}\kappa_{\alpha_{\text{plan}}} - \mu_F &= -\frac{1}{\sqrt{I_{\text{plan}}}}\sigma_q\Phi^{-1}(1 - \alpha_{\text{plan}})\sqrt{\mu_F(1 - \mu_F)} + \frac{Z_1\sigma_F}{\sqrt{I_{\text{plan}}}} + o_p\left(\frac{1}{\sqrt{I_{\text{plan}}}}\right) \\ \kappa_{\alpha_l} - \mu_F &= -\frac{1}{\sqrt{I_{\text{analysis}}}}\sigma_q\Phi^{-1}(1 - \alpha_l)\sqrt{\mu_F(1 - \mu_F)} + \frac{Z_2\sigma_F}{\sqrt{I_{\text{analysis}}}} + o_p\left(\frac{1}{\sqrt{I_{\text{analysis}}}}\right),\end{aligned}$$

where Z_1 and Z_2 are *independent* standard Normals by Framework 1. Subtracting one from the other and noting that $I_{\text{plan}} = rI$, $I_{\text{analysis}} = (1 - r)I$, we obtain

$$\kappa_{\alpha_l} - \kappa_{\alpha_{\text{plan}}} = \frac{\sqrt{\mu_F(1 - \mu_F)}\sigma_q}{\sqrt{I}} \left[\frac{\Phi^{-1}(1 - \alpha_{\text{plan}})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1 - r}} \right] - \frac{\sigma_F}{\sqrt{I}} \left(\frac{Z_1}{\sqrt{r}} - \frac{Z_2}{\sqrt{1 - r}} \right) + o_p\left(\frac{1}{\sqrt{I}}\right)$$

Finally, noting that $\kappa_{\alpha_{\text{plan}}} = \mu_F + o_p(1)$, we find that

$$\begin{aligned}\kappa_{\alpha_l} &= \kappa_{\alpha_{\text{plan}}} + \frac{\sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})}\sigma_q}{\sqrt{I}} \left[\frac{\Phi^{-1}(1 - \alpha_{\text{plan}})}{\sqrt{r}} - \frac{\Phi^{-1}(1 - \alpha_l)}{\sqrt{1 - r}} \right] - \frac{\sigma_F}{\sqrt{I}} \left(\frac{Z_1}{\sqrt{r}} - \frac{Z_2}{\sqrt{1 - r}} \right) \\ &\quad + o_p\left(\frac{1}{\sqrt{I}}\right)\end{aligned}$$

Since $Z_1/\sqrt{r} - Z_2/\sqrt{1 - r}$ has a $\mathcal{N}(0, 1/r(r - r))$ distribution, our proof is complete. \blacksquare

B.2. **Proposition 2.** We suppress the subscript l as understood from context. Referring to the distribution of $T_\kappa := \sqrt{I}(\bar{T}_\Gamma - \kappa)/\sqrt{\kappa(1 - \kappa)\sigma_{q,I}}$ as $G_{I,\kappa}$, where $\kappa = \Gamma/(1 + \Gamma)$, if $c_{q,I} = \frac{\frac{1}{n}\sum_{i=1}^I q_i^3}{\left(\frac{1}{n}\sum_{i=1}^n q_i\right)^3}$ is such that $c_q = \lim_{I \rightarrow \infty} c_{q,I} < \infty$, the quantiles of $G_{I,\kappa}$ admit a Cornish-Fisher expansion (DasGupta [2008]) of the form

$$G_{I,\kappa}^{-1}(t) = \Phi^{-1}(t) + \frac{\kappa_3}{\kappa_2^{3/2}}g(\Phi^{-1}(t)) + O\left(\frac{1}{I}\right)$$

where κ_r is the r -th cumulant of T_κ , $r \geq 1$, and $g(x) = \frac{1}{6}(x^2 - 1)$. Now, by the scaling chosen for T_κ , we see that $\kappa_2 = \text{Var}(T_\kappa) = 1$ and a quick calculation reveals

$$\kappa_3 = \mathbb{E}[T_\kappa^3] = \frac{1}{\sqrt{I}} \frac{2\kappa - 1}{\sqrt{\kappa(1 - \kappa)\sigma_{q,I}}} c_{q,I} = O(1) \quad (\text{by assumption})$$

Now the transformed sensitivity value κ_α satisfies

$$\begin{aligned} \frac{\sqrt{I}(T - \kappa_\alpha)}{\sqrt{\kappa_\alpha(1 - \kappa_\alpha)}\sigma_{q,I}} &= G_{I,\kappa_\alpha}^{-1}(1 - \alpha) = z_\alpha + \frac{1}{\sqrt{I}} \frac{2\kappa_\alpha - 1}{\sqrt{\kappa_\alpha(1 - \kappa_\alpha)}\sigma_{q,I}} c_{q,Ig}(z_\alpha) + O_p\left(\frac{1}{I}\right) \\ \text{(B.1)} \quad \Rightarrow \quad \frac{\sqrt{I}(T - \mu_F)}{\sigma_F} &= \frac{\sqrt{I}(\kappa_\alpha - \mu_F)}{\sigma_F} + z_\alpha \frac{\sigma_{q,I}}{\sigma_F} \sqrt{\kappa_\alpha(1 - \kappa_\alpha)} + \frac{2\kappa_\alpha - 1}{\sqrt{I}} \frac{c_{q,Ig}(z_\alpha)}{\sigma_F} + O_p\left(\frac{1}{I}\right) \end{aligned}$$

Finally, noting that $\kappa_\alpha = \mu_F + O_p(1/\sqrt{I})$ we thus obtain from Equation B.1

$$V = T_{\text{alt}} - (2\mu_F - 1) \frac{c_1}{\sqrt{I}} + O_p\left(\frac{1}{I}\right)$$

as claimed in Proposition 2 (i).

Next, we prove (ii) and continue from Equation B.1. Using the variables defined in Proposition 2, we have

$$\begin{aligned} T_{\text{alt}} &= W + \frac{\sigma_{q,I}z_\alpha}{\sigma_F} \sqrt{\left(\mu_F + \frac{W\sigma_F}{\sqrt{I}}\right) \left(1 - \mu_F - \frac{W\sigma_F}{\sqrt{I}}\right) + \left(\frac{2W\sigma_F}{\sqrt{I}} + 2\mu_F - 1\right) \frac{c_1}{\sqrt{I}}} + O_p\left(\frac{1}{I}\right) \\ &= W + \frac{\sigma_{q,I}z_\alpha}{\sigma_F} \sqrt{\mu_F(1 - \mu_F) - \frac{W\sigma_F}{\sqrt{I}}(2\mu_F - 1) + (2\mu_F - 1) \frac{c_1}{\sqrt{I}}} + O_p\left(\frac{1}{I}\right) \end{aligned}$$

Next, using $\sqrt{a+x} - \sqrt{a} = \frac{x}{2\sqrt{a}} + O(x^2)$ we obtain

$$T_{\text{alt}} = W \left[1 - \frac{\sigma_{q,I}z_\alpha(2\mu_F - 1)}{2\sqrt{I}\sqrt{\mu_F(1 - \mu_F)}} \right] + \frac{\sigma_{q,I}z_\alpha}{\sigma_F} \sqrt{\mu_F(1 - \mu_F)} + (2\mu_F - 1) \frac{c_1}{\sqrt{I}} + O_p\left(\frac{1}{I}\right).$$

Continuing with the trail of equalities

$$\begin{aligned} W &= \left[1 - \frac{\sigma_{q,I}z_\alpha(2\mu_F - 1)}{2\sqrt{I}\sqrt{\mu_F(1 - \mu_F)}} \right]^{-1} \left[T_{\text{alt}} - \frac{\sigma_{q,I}z_\alpha}{\sigma_F} \sqrt{\mu_F(1 - \mu_F)} - (2\mu_F - 1) \frac{c_1}{\sqrt{I}} \right] + O_p\left(\frac{1}{I}\right) \\ &= \left[1 + \frac{\sigma_{q,I}z_\alpha(2\mu_F - 1)}{2\sqrt{I}\sqrt{\mu_F(1 - \mu_F)}} + O_p\left(\frac{1}{I}\right) \right] \left[T_{\text{alt}} - \frac{\sigma_{q,I}z_\alpha}{\sigma_F} \sqrt{\mu_F(1 - \mu_F)} - (2\mu_F - 1) \frac{c_1}{\sqrt{I}} \right] \\ &\quad + O_p\left(\frac{1}{I}\right) \\ \therefore W &= T_{\text{alt}} \left[1 + \frac{\sigma_{q,I}z_\alpha(2\mu_F - 1)}{2\sqrt{I}\sqrt{\mu_F(1 - \mu_F)}} \right] - \frac{\sigma_{q,I}z_\alpha}{\sigma_F} \sqrt{\mu_F(1 - \mu_F)} - \frac{2\mu_F - 1}{\sqrt{I}} \left[c_1 + \frac{\sigma_{q,I}^2 z_\alpha^2}{2\sigma_F} \right] + O_p\left(\frac{1}{I}\right), \end{aligned}$$

concluding the proof. \blacksquare

B.3. Theorem 4.1. We provide a proof of Theorem 4.1 (i) as (ii) follows similarly. We suppress the superscript l and prove the version for \mathbb{P} , with the version for \mathbb{P}_* being analogous. From Proposition 2 and Equation 4.1 we obtain

$$\begin{aligned}\mathbb{P}(V \leq v) &= \mathbb{P}\left(T_{\text{alt}} \leq v + (2\mu_F - 1)\frac{c_1}{\sqrt{I}} + O_p\left(\frac{1}{I}\right)\right) \\ &= \Phi\left(v + (2\mu_F - 1)\frac{c_1}{\sqrt{I}} + O\left(\frac{1}{I}\right)\right) \\ &\quad + \frac{1}{\sqrt{I}}\phi\left(v + (2\mu_F - 1)\frac{c_1}{\sqrt{I}} + O\left(\frac{1}{I}\right)\right)b\left(v + (2\mu_F - 1)\frac{c_1}{\sqrt{I}} + O\left(\frac{1}{I}\right), F\right) + O\left(\frac{1}{I}\right)\end{aligned}$$

Using $f(x + O(1/\sqrt{I})) = f(x) + O(1/\sqrt{I})$, for any differentiable function f and $\Phi(a + x/\sqrt{I}) = \Phi(a) + \phi(a)x/\sqrt{I} + O(1/I)$ we deduce

$$\mathbb{P}(V \leq v) = \Phi(v) + \frac{1}{\sqrt{I}}[(2\mu_F - 1)c_1 + b(v, F)]\phi(v) + O\left(\frac{1}{I}\right),$$

completing the proof. ■

B.4. Proposition 3. We suppress the superscript l throughout the proof.

$$T_I = \frac{\sum_{i=1}^I \psi(IG_I(|Y_i|/(I+1))\text{sgn}(Y_i)}{\sum_{i=1}^I \psi(IG_I(|Y_i|/(I+1))}$$

where G_I is the empirical CDF of $I|Y_i|/(I+1)$. We have

$$\begin{aligned}\sqrt{I}T_I &= \frac{\frac{1}{\sqrt{I}}\sum_{i=1}^I \psi(IG_I(|Y_i|/(I+1))\text{sgn}(Y_i)}{\frac{1}{I}\sum_{i=1}^I \psi(IG_I(|Y_i|/(I+1))} = \frac{1}{\sqrt{I}\|\psi\|_1} \sum_{i=1}^I \psi(IG_I(|Y_i|/(I+1))\text{sgn}(Y_i) + o_p(1) \\ &= \frac{1}{\sqrt{I}\|\psi\|_1} \sum_{i=1}^I \psi(G(|Y_i|)\text{sgn}(Y_i) + \frac{1}{\sqrt{I}\|\psi\|_1} \sum_{i=1}^I \left[\psi\left(\frac{I}{I+1}G_I(|Y_i|)\right) - \psi(G(|Y_i|))\right] \text{sgn}(Y_i) \\ &\quad + o_p(1)\end{aligned}$$

where G is the CDF of $|Y_i|$. Call the second term $D_I/\|\psi\|_1$. Applying Hoeffding's inequality, we have

$$\begin{aligned}
\mathbb{P}(|D_I| > \varepsilon) &= \mathbb{E} \left[\mathbb{P} \left(\frac{1}{\sqrt{I}} \sum_{i=1}^I \left[\psi \left(\frac{I}{I+1} G_I(|Y_i|) \right) - \psi(G(|Y_i|)) \right] \text{sgn}(Y_i) > \varepsilon \mid |Y_1|, \dots, |Y_I| \right) \right] \\
&\leq \mathbb{E} \left[2 \exp \left(- \frac{I\varepsilon^2}{2 \sum_{i=1}^I \left[\psi \left(\frac{I}{I+1} G_I(|Y_i|) \right) - \psi(G(|Y_i|)) \right]^2} \right) \right] \\
\text{(B.2)} \quad &\leq \mathbb{E} \left[2 \exp \left(- \frac{I\varepsilon^2}{2B^2 \sum_{i=1}^I \left[\frac{I}{I+1} G_I(|Y_i|) - G(|Y_i|) \right]^2} \right) \right]
\end{aligned}$$

where $B = \sup_{x \in (0,1)} \psi'(x) < \infty$. Now, we observe

$$\begin{aligned}
\frac{1}{I} \sum_{i=1}^I \left[\frac{I}{I+1} G_I(|Y_i|) - G(|Y_i|) \right]^2 &\leq \frac{1}{I} \sum_{i=1}^I 2 \left[\frac{I^2}{(I+1)^2} (G_I(|Y_i|) - G(|Y_i|))^2 + \frac{1}{(I+1)^2} G(|Y_i|)^2 \right] \\
&\leq \frac{2I^2}{(I+1)^2} \|G_I - G\|_\infty + \frac{2}{(I+1)^2} \frac{1}{I} \sum_{i=1}^I G(|Y_i|)^2
\end{aligned}$$

which converges to 0 by Glivenko-Cantelli theorem and the weak law of large numbers since $\mathbb{E}[G(|Y|)^2] < \infty$. Noting that $e^{-x} \leq 1$ for all $x \geq 0$, by bounded convergence theorem on Equation B.2, we obtain $D_I = o_p(1)$.

Thus, we have

$$\frac{\sqrt{I}}{\sigma(\theta_0)} (T_I - \mu(\theta_0)) = \frac{1}{\sqrt{I}} \frac{\sum_{i=1}^I U_i (\text{sgn}(Y_i) - \mu(\theta_0))}{\sigma(\theta_0) \|\psi\|_1} + o_p(1),$$

where $U_i = \psi(G(|Y_i|))$ and G is the CDF of $|Y_i|$, thereby allowing T_I to have an asymptotic linear expansion at θ_0 . Also, as the location family has a continuously differentiable positive density, the Fisher information derived at θ_0 is $I = \int \frac{f'(x - \theta_0)^2}{f(x - \theta_0)} dx = \int \frac{f'(x)^2}{f(x)} dx$ is constant and thus continuous at θ_0 . Hence, the family is quadratic mean differentiable (Van der Vaart [2000], Theorem 7.2), leading to $\sqrt{I}(T_I - \mu(\theta_0))/\sigma(\theta_0)$ and the log-likelihood ratio at θ_I vs θ_0

satisfying

$$\begin{bmatrix} \sqrt{I}(T_I - \mu(\theta_0))/\sigma(\theta_0) \\ \log \frac{\prod_{i=1}^I f_{\theta_0+h/\sqrt{I}}(Y_i)}{\prod_{i=1}^I f_{\theta_0}(Y_i)} \end{bmatrix} = \frac{1}{\sqrt{I}} \begin{bmatrix} \sum_{i=1}^I \frac{U_i(\text{sgn}(Y_i) - \mu(\theta_0))}{\sigma(\theta_0)\|\psi\|_1} \\ h \sum_{i=1}^I \frac{f'_{\theta_0}(Y_i)}{f_{\theta_0}(Y_i)} \end{bmatrix} + \begin{bmatrix} 0 \\ -h^2 I/2 \end{bmatrix} + \begin{bmatrix} o_p(1) \\ o_p(1) \end{bmatrix}$$

and thus converging to a bivariate normal distribution. By Le-Cam's third lemma (Van der Vaart [2000], Section 6.2), our proof is complete. \blacksquare

B.5. **Theorem 4.2.** An outcome l is selected by Method 2 if its corresponding

$$\begin{aligned} & \kappa_{\alpha_{\text{plan}}} + \frac{\sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})}\sigma_q z_{\alpha_{\text{plan}}}}{\sqrt{Ir}} - \frac{\sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})}\sigma_q z_{\alpha_l}}{\sqrt{I(1-r)}} > \kappa_{\text{con}} - \frac{\hat{\sigma}(\hat{\theta}_I)z_{\alpha_{\text{coverage}}}}{\sqrt{Ir(1-r)}} \\ \Leftrightarrow & \sqrt{Ir} \left(\kappa_{\alpha_{\text{plan}}} + \frac{\sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})}\sigma_q z_{\alpha_{\text{plan}}}}{\sqrt{Ir}} - \frac{\sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})}\sigma_q z_{\alpha_l}}{\sqrt{I(1-r)}} \right) > \\ & \sqrt{Ir}\kappa_{\text{con}} - \frac{\hat{\sigma}(\hat{\theta}_I)z_{\alpha_{\text{coverage}}}}{\sqrt{1-r}} \\ \Leftrightarrow & \sqrt{I_a} \frac{(\kappa_{\alpha_{\text{plan}}} - \mu(\theta_I))}{\sigma(\theta_I)} + \sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})} \frac{\sigma_q}{\sigma(\theta_I)} z_{\alpha_{\text{plan}}} - \frac{\sqrt{\kappa_{\alpha_{\text{plan}}}(1 - \kappa_{\alpha_{\text{plan}}})}\sqrt{r}}{\sqrt{(1-r)}} \frac{\sigma_q}{\sigma(\theta_I)} z_{\alpha_l} > \\ & \sqrt{I_a} \frac{(\kappa_c - \mu(\theta_I))}{\sigma(\theta_I)} - \frac{\hat{\sigma}(\hat{\theta}_I)z_{\alpha_{\text{coverage}}}}{\sigma(\theta_I)\sqrt{1-r}} \end{aligned}$$

the probability of which, under θ_I and as $I \rightarrow \infty$, is given by

$$\pi_I(\theta_I) \rightarrow 1 - \Phi \left(-\frac{z_{\alpha_{\text{coverage}}}}{\sqrt{1-r}} - h \frac{\mu'(\theta_0)}{\sigma(\theta_0)} + z_{\alpha_{\text{plan}}} \frac{\sqrt{\kappa(1-\kappa)}\sigma_q}{\sigma(\theta_0)} \frac{\sqrt{r}}{\sqrt{1-r}} \right)$$

as intended. \blacksquare

B.6. **Theorem A.2.** We suppress the l -superscript throughout the proof and drop mention of \mathbf{Z} and \mathbf{R}^l . If the sensitivity analysis is computed via Equation A.1, then Γ_α^* satisfies Equation A.2, i.e.,

$$(B.3) \quad T - a_{\Gamma_\alpha^*} = \Phi^{-1}(1 - \alpha) \frac{b_{\Gamma_\alpha^*}}{\sqrt{I}} + o_p \left(\frac{1}{\sqrt{I}} \right).$$

Let f be the mapping $f : \Gamma \mapsto a_\Gamma$, which is continuously differentiable at a neighborhood of $\tilde{\Gamma}$ with a non-zero derivative at $\tilde{\Gamma}$ by assumption. By inverse function theorem (Rudin et al. [1976], Theorem 9.24), f is invertible at neighborhood of $\tilde{\Gamma}$ and the inverse also admits a derivative.

Since $T = \mu_{F,g} + o_p(1)$, rejection of the null hypothesis occurs w.p. 1 as $I \rightarrow \infty$ if Γ is such that $a_\Gamma < \mu_{F,g}$. On the other hand, rejection of the null hypothesis occurs w.p. 0 as $I \rightarrow \infty$ if Γ is such that $a_\Gamma > \mu_{F,g}$. Thus, by definition, $a_{\tilde{\Gamma}} = \mu_{F,g}$ implies $\tilde{\Gamma} = f^{-1}(\mu_{F,g})$, where the inverse is well-defined in a neighborhood of $\mu_{F,g}$ by the inverse function theorem.

Since $T \xrightarrow{P} \mu_{F,g}$, we obtain $a_{\Gamma_\alpha^*} \xrightarrow{P} \mu_{F,g}$ from Equation B.3 and deduce $\Gamma_\alpha^* \xrightarrow{P} \tilde{\Gamma}$ by continuous mapping theorem. Using Equation A.2 and Assumption A.3, we can obtain

$$\frac{\sqrt{I}(a_{n,\Gamma_\alpha^*} - \mu_{F,g})}{\sigma_{F,g}} = \frac{\sqrt{I}(T - \mu_{F,g})}{\sigma_{F,g}} - \Phi^{-1}(1 - \alpha) \frac{b_{\Gamma_\alpha^*}}{\sigma_{F,g}} + o_p(1)$$

Replacing $b_{\Gamma_\alpha^*}$ by $b_{\tilde{\Gamma}}$ and using continuous mapping theorem, we have

$$\frac{\sqrt{I}(a_{\Gamma_\alpha^*} - \mu_{F,g})}{\sigma_{F,g}} \xrightarrow{d} \mathcal{N}\left(-\Phi^{-1}(1 - \alpha) \frac{b_{\tilde{\Gamma}}}{\sigma_{F,g}}, 1\right)$$

Thus, by the inverse-function theorem and delta method using the function f^{-1} in a neighborhood of $\mu_{F,g}$ we conclude

$$\sqrt{I}(\Gamma_\alpha^* - \tilde{\Gamma}) \xrightarrow{d} \mathcal{N}\left(-\Phi^{-1}(1 - \alpha)b_{\tilde{\Gamma}}(f^{-1})'(\mu_{F,g}), \sigma_{F,g}^2((f^{-1})'(\mu_{F,g}))^2\right)$$

■

APPENDIX C. ADDITIONAL DISCUSSIONS

C.1. Further Design Considerations.

C.1.1. *Choice of Statistics.* Rosenbaum advocates for catering design with the intent of maximizing design sensitivity. Zhao [2018] applies this idea to the selection of test statistics with the sensitivity value. He considers the generalized score statistic where $q_i^l = \psi^l(\text{rank}(Y_i^l))/(I + 1)$ and obtains a formula for the asymptotic mean and variance in Equation 3.2 based on the outcome density and score function ψ^l which can vary by outcome. However, his analysis relies only on the mean, as the variance parameter is quite involved to calculate. It also requires prior knowledge of the outcome distributions.

Due to our use of split-sampling Framework 1, we can bypass excess considerations and directly compare test statistics on the planning sample. For instance, suppose the l^{th} outcome has $\{\psi_k^l\}_{1 \leq k \leq p_l}$ candidate choices to be used. Although we have framed Sens-Val as a procedure for outcome selection, one can redefine an outcome as its (l, k) pair, $1 \leq k \leq p_l$, $1 \leq l \leq L$, and choose the (outcome, statistic) pair suggested by the method. In case of a tie among the statistics, one can simply choose the statistic with the higher sensitivity value. Such a choice bypasses the theoretical considerations of Zhao [2018], circumvents dealing with variances σ_F^l , and provides a data-driven means for selection of outcomes and test statistics. Moreover, it is driven by the quantiles of the asymptotic distribution of the sensitivity value, adhering to the suggestions by Zhao [2018] and augmenting it with the possibility that quantiles used in planning and analysis samples may differ.

We illustrate the potential benefits of this idea as follows. Consider a setting with $N = 200$ subjects randomly assigned to treatment or control with probability $1/2$ and $L = 20$ possible outcomes, among which there exist four true signals. Suppose the control potential outcomes $r_{C_{ij}}^l$ are all generated independently from either the standard Normal distribution, standard logistic distribution, or a t -distribution with four degrees of freedom. Treated units amongst the true signals equal their control potential outcome plus a constant treatment effect of τ , which is specified for each distribution. This setup allows us to assess the performance of our method for various types of data. Using a planning sample proportion of 0.20, we evaluate the average power of our Sens-Val method across 100 simulations for various elements in the class of U -statistics proposed by Rosenbaum [2011, 2014], which are indexed by parameters $(m, \underline{m}, \overline{m})$.

For a fixed outcome l , we let $h(y)$ denote a function of m variables that quantifies the positive differences among the order statistics between $|y|_{(\underline{m})}$ and $|y|_{(\overline{m})}$. The corresponding U -statistic can then be taken as $T = \binom{I}{m}^{-1} \sum_{|\mathcal{I}|=m} h(Y_{\mathcal{I}})$ and written as signed score form 2.2. In the absence of ties:

$$d_i^l = \binom{I}{m}^{-1} \sum_{k=\underline{m}}^{\overline{m}} \binom{\text{rank}(|Y_i^l|) - 1}{k-1} \binom{I - \text{rank}(|Y_i^l|)}{m-k} \approx I^{-1} \sum_{k=\underline{m}}^{\overline{m}} k \binom{m}{k} p^{k-1} (1-p)^{m-k},$$

where $p = \text{rank}(|Y_i^l|)/I$ (Rosenbaum [2011], Section 3.1). Note that the choice $(m, \underline{m}, \overline{m}) = (2, 2, 2)$ closely approximates Wilcoxon's signed rank statistic (Pratt and Gibbons [1981]).

We observe the simulated average power among these statistics in Table 2 below. In terms of performance, the best statistic to use often changes based on the simulation setting and data generating process. This shows that it can often be beneficial in terms of detection power to compare statistics to use during the planning stage.

TABLE 2. Simulated average power with $N = 200$ subjects of a 0.05 level analysis for several values of Γ . The power is the proportion of simulations where the test correctly rejects the null hypothesis when it is false, performed at the given value of Γ . Each situation is replicated 100 times. Scores above the horizontal line are monotone increasing scores, while those below are descending scores.

$(m, \underline{m}, \overline{m})$	Normal(0,1) $\tau = 1$			Logistic(0,1) $\tau = 1.5$			t with 4 d.f. $\tau = 1.5$		
	$\Gamma = 1.5$	$\Gamma = 2.5$	$\Gamma = 3.5$	$\Gamma = 1.5$	$\Gamma = 2.5$	$\Gamma = 3.5$	$\Gamma = 1.5$	$\Gamma = 2.5$	$\Gamma = 3.5$
(2,2,2)	0.855	0.565	0.150	0.648	0.200	0.048	0.390	0.113	0.013
(5,4,5)	0.843	0.555	0.183	0.620	0.215	0.055	0.368	0.110	0.015
(8,7,8)	0.778	0.463	0.165	0.495	0.160	0.043	0.243	0.068	0.015
(8,5,8)	0.860	0.573	0.193	0.643	0.225	0.058	0.413	0.133	0.018
(20,16,20)	0.753	0.445	0.180	0.463	0.170	0.033	0.225	0.058	0.013
(20,12,20)	0.848	0.575	0.223	0.618	0.223	0.058	0.398	0.130	0.020
(8,7,7)	0.723	0.438	0.178	0.500	0.173	0.045	0.380	0.108	0.025
(8,5,7)	0.753	0.423	0.115	0.563	0.168	0.040	0.515	0.128	0.023
(20,16,19)	0.653	0.368	0.170	0.425	0.153	0.053	0.260	0.078	0.025
(20,12,19)	0.760	0.488	0.173	0.548	0.185	0.050	0.483	0.143	0.023

C.1.2. *Choice of Sub-Populations.* Zhao [2018] considers the choice of sub-populations to be an important aspect of the study design and explains the trade-off of using sub-populations instead of all subgroups observed by Hsu et al. [2013]. While this explains the phenomenon based on the difference of design sensitivities (and sub-group proportions), the ignorance of design-sensitivity may leave a researcher no better off, unless she employs a pilot sample to

estimate the design sensitivities. As noted before, estimation of design sensitivity is not only challenging, but also not quite the goal of the researcher who has to run the tests on the analysis sample. Sens-Val solves the problem more pragmatically – one may consider the effect on all possible sub-populations as different outcomes, and use our method to select the sub-populations which have a reasonable chance to be significant in the analysis sample.

We demonstrate this advantage with a simulation. Consider a setting with $N = 2000$ subjects randomly assigned to treatment or control with probability $1/2$ and $L = 10$ possible outcomes, among which there exist three true signals. We use a planning sample proportion of 0.20 to divide control/treated pairs into discrete planning and analysis samples. We then take $K = 5$ discrete sub-populations, where control/treated pairs within the planning and analysis samples are assigned to sub-population k with probability $\pi_k = (0.10, 0.15, 0.20, 0.20, 0.35)$. Suppose the control potential outcomes $r_{C_{ij}}^l$ are all generated independently from the standard Normal distribution and treated potential outcomes are taken as:

$$r_{T_{ij}}^l = \begin{cases} r_{C_{ij}}^l + \frac{2}{3}, & \text{if } l = 1 \text{ or } (l = 2 \text{ and } k \in 1, 2, 3) \text{ or } (l = 3 \text{ and } k \in 1, 2); \\ r_{C_{ij}}^l + \frac{1}{3}, & \text{if } l = 3 \text{ and } k \in 3, 4; \\ r_{C_{ij}}^l, & \text{otherwise.} \end{cases}$$

The Sens-Val method selects any promising sub-populations in the planning sample to test in the latter stage of the pipeline. We conduct our Sens-Val method and Bonferroni correction across 1,000 simulations at $\alpha = 0.05$ and $\Gamma_{\text{com}} = 2$, and display our simulated performance in Table 3 below. We see that Sens-Val does a very good job choosing sub-populations during the planning stage, which yields notable benefits in terms of detection power.

C.1.3. Coherence. An effect is said to be coherent if the observed association between the exposure and outcomes are compatible with the mechanism thought to produce the effects. Rosenbaum [1997] considers a linear combination of the signed-rank statistics to create a coherent statistic. Ye et al. [2022] extends this idea to all linear combinations using the ideas of Scheffé [1953] and examines the impacts of such choices on the power of the study. Rosenbaum [2020], meanwhile, uses an *a priori* guess for the choice of linear combinations obtained from previous data, thereby circumventing the need for multiplicity corrections. We concur that coherence is an important consideration in the design of a study, as a coherent mechanism is

TABLE 3. Simulation results whereas each situation is replicated 1,000 times. The left-hand-side shows the proportion of simulations in which an outcome/sub-population pair is selected by Sens-Val in the planning sample. The right-hand-side gives the frequency for which each method rejects a certain outcome, along with an average true positive rate for each method. Bolded numbers depict scenarios with non-zero treatment effects.

Outcome	Proportion Tested					Proportion Rejected	
	Sub-Population					Method	
	1	2	3	4	5	Sens-Val	Bonferroni
1	0.999	1.000	1.000	1.000	1.000	0.923	0.917
2	1.000	1.000	1.000	0.022	0.001	0.639	0.001
3	0.999	0.999	0.297	0.470	0.007	0.161	0.000
4	0.009	0.006	0.004	0.010	0.005	0.000	0.000
5	0.013	0.008	0.002	0.014	0.000	0.000	0.000
6	0.015	0.008	0.004	0.013	0.007	0.000	0.000
7	0.010	0.007	0.009	0.015	0.007	0.000	0.000
8	0.016	0.009	0.004	0.013	0.007	0.000	0.000
9	0.014	0.004	0.005	0.012	0.005	0.000	0.000
10	0.007	0.009	0.007	0.017	0.005	0.000	0.000
See that ■ denotes $\tau = \frac{2}{3}$ while ■ denotes $\tau = \frac{1}{3}$.						True Positive Rate	
						0.574	0.306

less likely to suffer from unmeasured biases. Method 2 can easily be morphed to impose conditions on the outcomes to test. For example, if an exposure affects an outcome solely through a mediator, the criteria in Method 2 can be adjusted to include the outcome only if both the impact of exposure on the mediator and the mediator’s effect on the outcome meet the specified thresholds. Such restrictions, if known, would be helpful to increase the sensitivity value of the true outcomes while filtering out null or weak signals to enhance discovery.

C.1.4. *Weighting of Outcomes.* Since smaller effects can often arise due to small unattenuated biases, while larger effects can be rendered non-causal only by biases of large magnitudes, one might be encouraged to use weighted statistics. Rosenbaum [2014] considers such an approach using weighted M-statistics for superior design sensitivity, but are likely to be driven by the data. In the classical inference literature, Wasserman and Roeder [2006] consider choosing α_l s in Framework 1 using estimated signal strengths from the planning sample. While the primary concern there is bias rather than efficiency, it may still be useful in a finite sample study tasked with managing both efficiency and bias. Method 2 used in conjunction with the dynamic α_l framework might lend itself to a middle-ground approach, where the statistics and α_l s are

suitably driven by the data to obtain enhanced power under bias and moderate sample size. As long as α_{lS} are suitably normalized to control the FWER in the analysis sample, one can experiment via Method 2 with different choices of the superior statistics based on Rosenbaum [2014] and the level thresholds based on signal strengths Wasserman and Roeder [2006] to optimize their power.

C.2. Computational Considerations. One may contrast Method 2 with the implementations of Heller et al. [2009]. In the vanilla version, they choose one outcome to test in the planning sample according to the smallest p -value, which is quite limited by its applicability. They also introduce a technique in which they consider all possible subsets of the outcomes and select that which has the min-max p -value. However, this approach is computationally infeasible for even a moderate number of outcomes, requiring the consideration of 2^L subsets of outcomes in the planning sample. On the other hand, Method 2 is linear in the number of outcomes and is thus much more usable in practice. Moreover, due to potential correlations among the outcomes of interest, the min-max p -value may have complicated nuances attached to it, the theoretical properties of which are poorly understood. However, our proposed Method 2 has simple theoretical properties (3.1) and is robust to potential correlational complications since it relies only on the marginal structure of the outcomes.

We also note that the bootstrap samples need not be recalculated for each α^n when employing the dynamic α framework. As the bootstrap samples are used only to obtain an estimate of σ_F^l , which is devoid of α from its definition via Equation 3.1, $\hat{\sigma}_F^l(\alpha)$ and $\hat{\sigma}_F^l(\alpha^*)$ are likely to have an $o_p(1)$ difference, leading to an $o_p(1/\sqrt{I_{\text{plan}}})$ difference in the odds of the l^{th} outcome being selected. Hence, one can safely do away with implementing the bootstrap repetitively without any additional asymptotic cost.

APPENDIX D. DETAILS OF DATA APPLICATION




FIGURE 3. A Love plot displaying the absolute standardized mean differences across covariates.

TABLE 4. Description of rejected outcomes, in accordance with the results in Table 1. Among the 93 total outcomes, only 14 are rejected by any method at the evaluated levels of Γ_{con} .

Outcome	Definition
ft.water	Feet of water in household due to flood
repair.cost	Cost of repairing flood damage
days.water	Number of days with water in household
days.away.home	Number of days out of home due to flood
tubewell.cooking	Household obtains cooking water from tubewell
tubewell.washing	Household obtains washing water from tubewell
food.price.idx	Food price index
food.credit.val.pc	Per capita value of credit purchases for food
credit.val	Total household value of credit purchases
days.ill	Average number of days with illness
pulses.budget	Share of total household expenditure used to purchase pulses
rice.consumed.pc	Amount of rice consumed per capita
food.produced.val	Total household value of food produced
food.produced.val.pc	Per capita value of food produced

TABLE 5. All of the 93 outcomes examined in the data application. Outcomes are grouped into flood-specific, food consumption and security, health, nutrition, and economic categories. Although many of the outcomes can be grouped into more than one category, we list them under a single group for simplicity.

Category	Specific Outcomes
Flood-Specific	days.away.home, days.water, ft.water
Food Consumption & Security	amt.fish.female, amt.fish.male, amt.milk.meat.egg.female, amt.milk.meat.egg.male, calories.pc, calories.per.adult, cereals.consumed.pc, cereals.budget, egg.consumed.pc, egg.budget, fish.consumed.pc, fish.budget, food.price.idx, fruit.consumed.pc, fruit.budget, icrs.fish.female, icrs.fish.male, meat.consumed.pc, meat.budget, milk.consumed.pc, milk.budget, oil.consumed.pc, oil.budget, protein.pc, protein.per.adult, pulses.consumed.pc, pulses.budget, rice.consumed.pc, rice.budget, veg.consumed.pc, veg.budget, wheat.consumed.pc, wheat.budget
Health	any.ill.0-5yrs, any.ill.6-18yrs, any.ill.fever, any.ill.female, any.ill.gastrointest, any.ill.male, any.ill.over18yrs, any.ill.respiratory, days.ill, distance.cooking, distance.drinking, distance.washing, liters.water.collection, sanitary.latrine, time.water.collection, tubewell.cooking, tubewell.drinking, tubewell.washing
Nutrition	rd3.bmi, rd3.bmi.adolesc.women, rd3.height.on.age, rd3.height.on.age.preschool, rd3.weight.on.age, rd3.weight.on.age.preschool, rd3.weight.on.height, rd3.weight.on.height.preschool
Economic	adult.equivalent.tot.expend, any.electricity, credit.val, food.consumed.val, food.credit.val.pc, food.expend.pc, food.purchased.val, food.purchased.val.pc, food.produced.val, food.produced.val.pc, food.received.val, food.received.val.pc, marriage.funerals.expend, nonfood.expend, nonfood.expend.incl.repairs, nonfood.expend.pc, repair.cost, share.food.expend, share.food.expend.incl.repairs, share.nonfood.expend, share.nonfood.expend.incl.repairs, tot.expend, tot.expend.incl.repairs, tot.expend.pc, val.post.agric.equip, val.post.domestic, val.post.house, val.post.livestock, val.post.transport, val.post.valuables, val.post.water