EXISTENCE, UNIQUENESS AND ALGEBRAIC REDUCTIBILITY EXPERIMENTS OF RANS EQUATIONS WITH PERIODIC BOUNDARY CONDITIONS

A PREPRINT

C. V. Valencia, S. Velasco and M. Romero de Terreros Department of Physics and Mathematics Universidad Iberoamericana Ciudad de México

> Prolongación Paseo de la Reforma 880, Mexico City —01219, MEXICO carla.valencia@ibero.mx

> > June 4, 2024

ABSTRACT

In the present work, we show the existence and uniqueness of the two-dimensional turbulent vorticity term under conditions given on the weak solutions to the classical incompressible Navier-Stokes problem in a compact domain R with periodic boundary conditions and initial conditions in Sobolev space $H^m(R)$. To achieve this, we derive a pressureless incompressible system and calculate kinetic energy quantities on the turbulent vorticity from the properties of the classical curl. Finally, since the system obtained is algebraic, we experimented with the possibilities of reducing it by applying the Rosenfeld-Gröbner algorithm.

Keywords Turbulence theory, Vortex dynamics

Introduction

It is a well-known practice to use the rotational operator on the Navier-Stokes equations to derive pressureless systems. However, we have not come across similar approaches for the Navier-Stokes equations that take into consideration velocities for the *two-scale decomposition of the flow*, as outlined in the *Reynolds-Averaged Navier-Stokes* (RANS) equations statement before the time average hypothesis is considered. The system of equations obtained includes non-linear differential polynomials, the factorization of which corresponds to limit cases that could also be obtained from scenarios with small parameters. We look for terms that persist during different algebraic simplifications, to estimate the maximum time of existence and uniqueness of a classical solution, and to recommend using algorithms from the theory of differential algebra to investigate the multiplicity of formal solutions beyond that maximum time. It is our belief that this method gives a unique perspective on exploring causal terms in various turbulent scenarios.

Olga Ladyzhenskaya, stated the: "Problem 1. Do the Navier-Stokes equations together with the initial and boundary conditions actually give a deterministic description of the dynamics of an incompressible fluid?" [19, p. 252]. However, there are inaccuracies in the description of flows given by ideal models and real fluids [13, 14]. In particular, there is *turbulence*, and study of fluid dynamics, especially *turbulent flow*, is pivotal in engineering applications. The Reynolds-averaged Navier-Stokes (RANS) equations remain central in this field and offer valuable insights to increasingly complex problems relating to the flow of fluids.

In this article, we lay the existence and uniqueness foundations for our search for sufficient conditions to establish well-posed problems in the context of the algebraically simplified cases of the *Reynolds-Averaged Navier-Stokes* equations (RANS). Our far-reaching objective is to establish the conditions required for the existence and uniqueness of periodic solutions, which are empirically observed, and for which, even recently, no mathematical proofs have been reported, such as the case of a fluid flowing past a cylinder and having Reynolds numbers between 50 and 250 [25]. Moreover, we hope that the identification of the terms that prevail in the limit when the viscosity tends to zero will allow us to propose explanations for why this periodicity is lost for Reynolds numbers greater than 250 [25]. The results




Figure 1: $R = [0, L_1] \times [0, L_2] \times [0, L_3]$

obtained in this research were presented, without detailed proofs, at *the Engineering Mechanics Institute Conference* and Probabilistic Mechanics & Reliability Conference (EMI/PMC 2024).

In particular, we see value in examining the theoretical implications of the issues associated with the RANS approach to gain insight into the dynamics of the Earth's Atmosphere. The Intergovernmental Panel on Climate Change (IPCC) states that "Atmospheric models include representations of (...) turbulence, convection and gravity waves that are not fully represented by grid-scale dynamics" [28, p. 277]. Both the general existence and uniqueness problem and the specific need to address this Issues point us towards the broader question at hand: Do the limit models derived from Reynolds-Averaged Navier-Stokes Equations' problems could give a better understanding of the dynamics of a turbulent fluid in the Atmosphere? We would like to highlight that there are very few fundamental results on local and global existence and uniqueness of solutions for a two scale decomposition of the flow, as it appears on the RANS equations [23, p. 126] and LES initial statements of the problem [26, p. 11], [21, 22].

We analyze the new system of equations with both the classical energy method, and as a system of differential polynomials through the Rosenfeld-Gröbner algorithm [7], [6], and [1]. In recent years, the field of differential algebra has been extending its reach, offering a promising alternative approach for gaining ground against complex differential systems. We propose analyzing fluid dynamics theory from an algebraic standpoint. This is achieved through the implementation of Rosenfeld-Gröbner algorithms, a method pioneered by Francois Boulier. Boulier's extensive work in differential algebra provides the main framework for our investigation [3]. We aim to obtain decomposed systems of from the original system of equations.

1 Problem Statement

Here, we present the general form obtained by applying the rotational to the Navier-Stokes equations where the velocity is split as a linear combination of two parts, $v + \tilde{v}$, where $v, \tilde{v} \in C^{\infty}(D)$ are infinitely differentiable functions on $R = [0, L_1] \times [0, L_2] \times [0, L_3]$, for fixed $L_1, L_2, L_3 > 0$, so that the results can be extended to *starshaped* real domains (where any ray with an origin in a fixed point has a unique common point with the topological boundary of the domain) and distributions in the Sobolev spaces $W_p^k(R)$, with $p \in [1, \infty)$, thanks to the fact that, in particular, $C^{\infty}(R)$ is dense in $W_p^k(R)$ if R is a rectangular prism.

Remark 1. While a necessary and sufficient condition for a distribution in a Sobolev space to be the limit of a succession of analytic functions is unknown it is possible to assume that the domain is starshaped with respect to a point to say that the space $C^{\infty}(\bar{R})$ of infinitely differentiable functions on a domain D is dense in the Sobolev space $W_p^k(R)$ of distributions on R with derivatives of order $0 \le l \le k$ in the Lebesgue space $L_p(R)$, for $p \in [1, \infty)$. Therefore, we can just assume that the domain is a rectangular prism R and v is an element of the space $C^{\infty}(\bar{R})$ that satisfies both the Leibinz rule for the product derivative, the Gauss-Green formula [31, p. 699], and the generalisation of Schwarz's Theorem [31, p. 280] to exchange the order of the generalised second partial derivatives with respect to the spatial and temporal independent variables [29, 30].




Figure 2: Periodic boundary conditions at ∂R

Theorem 1. Let $p, v, \tilde{v} \in C^1([0,T), C^{\infty}(\bar{R}))$, for some $0 < T < \infty$, with periodic conditions for v and \tilde{v} at ∂R , where ∂R denotes the topological boundary of R. This is, $v(x_1, x_2, x_3, t) = v(x_1 + L_1, x_2 + L_2, x_3 + L_3, t)$, and $\tilde{v}(x_1, x_2, x_3, t) = \tilde{v}(x_1 + L_1, x_2 + L_2, x_3 + L_3, t)$, $\forall (x_1, x_2, x_3) \in \partial R$. If v, \tilde{v} , and p verify the following system of equations:

$$\begin{cases} \nabla \cdot \boldsymbol{v} = 0, \\ \nabla \cdot \tilde{\boldsymbol{v}} = 0, \\ \frac{D(\boldsymbol{v} + \tilde{\boldsymbol{v}})}{Dt} = \nu \Delta(\boldsymbol{v} + \tilde{\boldsymbol{v}}) - \nabla p. \end{cases}$$
(1)

Then, for at least one point $x^0 = (x_1^0, x_2^0, x_3^0) \in R$, v and \tilde{v} satisfy:

$$\frac{\partial \boldsymbol{\omega}}{\partial t} + \frac{\partial \tilde{\boldsymbol{\omega}}}{\partial t} + (\boldsymbol{v} \cdot \nabla)\boldsymbol{\omega} + (\tilde{\boldsymbol{v}} \cdot \nabla)\tilde{\boldsymbol{\omega}} + (\boldsymbol{v} \cdot \nabla)\tilde{\boldsymbol{\omega}} + (\tilde{\boldsymbol{v}} \cdot \nabla)\boldsymbol{\omega} - \nu\Delta(\boldsymbol{\omega} + \tilde{\boldsymbol{\omega}}) = 0,$$
(2)

where Δ denotes the Laplacian operator, $\boldsymbol{\omega} = \nabla \times \boldsymbol{v}$, and $\tilde{\boldsymbol{\omega}} = \nabla \times \tilde{\boldsymbol{v}}$.

Proof. In order to obtain Eq. (10), we will integrate each term on both sides of the Conservation of Momentum Law of Eq. (1) over the entire physical domain R:

$$\iiint_R \left(\frac{D(\boldsymbol{v}+\tilde{\boldsymbol{v}})}{Dt}\right) d\boldsymbol{x} = \iiint_R (\nu \Delta(\boldsymbol{v}+\tilde{\boldsymbol{v}}) - \nabla p) d\boldsymbol{x},$$

where $dx = dx_1 dx_2 dx_3$ is the volume differential in \mathbb{R}^3 . By the monotone convergence theorem [31], if:

$$\nabla \times \left(\iiint_R \left(\frac{D(\boldsymbol{v} + \tilde{\boldsymbol{v}})}{Dt} \right) d\boldsymbol{x} \right) = \nabla \times \left(\iiint_R \left(\nu \Delta(\boldsymbol{v} + \tilde{\boldsymbol{v}}) - \nabla p \right) d\boldsymbol{x} \right),$$

then,

$$\iiint_{R} \left(\nabla \times \frac{D(\boldsymbol{v} + \tilde{\boldsymbol{v}})}{Dt} \right) d\boldsymbol{x} = \iiint_{R} \left(\nabla \times \left(\nu \Delta (\boldsymbol{v} + \tilde{\boldsymbol{v}}) - \nabla p \right) \right) d\boldsymbol{x}.$$
(3)

Given that $v, \tilde{v} \in C^{\infty}(D)$ and the Schwarz theorem, we can exchange the order of derivation. Because both v and \tilde{v} have zero divergence and periodic conditions, by direct calculations and integration by parts, the left hand side of Eq. (3) becomes:

$$\begin{split} \iiint_{R} \left(\nabla \times \frac{D(\boldsymbol{v} + \tilde{\boldsymbol{v}})}{Dt} \right) d\boldsymbol{x} &= \iiint_{R} \left(\mathbf{A_{1}} + \mathbf{B_{1}} + \mathbf{M_{1}} + \mathbf{M_{2}} + \mathbf{M_{3}} + \mathbf{M_{4}} \right) d\boldsymbol{x}, \\ \mathbf{A_{1}} &= \frac{\partial}{\partial t} \left(\nabla \times \boldsymbol{v} \right), \qquad \mathbf{B_{1}} = \frac{\partial}{\partial t} \left(\nabla \times \tilde{\boldsymbol{v}} \right). \end{split}$$

where:

$$\partial t$$
 ∂t ∂t

$$\mathbf{M_2} + \mathbf{M_3} = (\boldsymbol{v} \cdot \nabla) \left(\nabla \times \tilde{\boldsymbol{v}} \right) + \left(\tilde{\boldsymbol{v}} \cdot \nabla \right) \left(\nabla \times \boldsymbol{v} \right)$$

and

$$\mathbf{M_4} = \left(\mathbf{\tilde{v}} \cdot \nabla \right) \left(\nabla \times \mathbf{\tilde{v}} \right).$$

On the right hand,

$$\nabla \times \left(\nu \Delta (\boldsymbol{v} + \tilde{\boldsymbol{v}}) - \nabla p\right) = \nu \Delta \left(\left(\nabla \times \boldsymbol{v}\right) + \left(\nabla \times \tilde{\boldsymbol{v}}\right) \right).$$

We can remove the triple integral through the Intermediate Value Theorem [32, p. 124], knowing there is at least one $x_0 \in R$ such that, putting together the expressions for the left and right hand sides, we obtain Eq. (10).

2 Existence and Uniqueness Theorems

In light of the problem statement above, the system that arises is algebraic. This allows for the classical energy method to be applied in order to obtain estimates and quantities that are unaffected by viscosity, and only dependant on the initial and boundary conditions.

Remark 2. If $\mathbf{f} = (f_1, f_2, f_3)$, $\mathbf{g} = (g_1, g_2, g_3) \in L^2(R; \mathbb{R}^3)$, we denote its inner product as: $(\mathbf{f}, \mathbf{g})_0 = \iiint_R (\mathbf{f} \cdot \mathbf{g}) d\mathbf{x} = \iiint_R (f_1 g_1 + f_2 g_2 + f_3 g_3) d\mathbf{x}$. Similarly, $(f, f)_0 = ||f||_0^2 = \iiint_R (f_1^2 + f_2^2 + f_3^2) d\mathbf{x}$. Additionally, let ∂_{α}^k be a differential operator that express the partial derivative of absolute order k with order α for the different spatial variables: $\partial_{\alpha}^k = \frac{\partial^k}{\partial x^{\alpha_1} \partial x^{\alpha_2} \partial x^{\alpha_3}}$, where $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^3$ and $\alpha_1 + \alpha_2 + \alpha_3 = k$.

Lemma 1. Let $f, g \in C^1([0,T), C^{\infty}(\overline{R}))$, for some $0 < T < \infty$ satisfy periodic conditions at ∂R , such that $\nabla \cdot f = 0$. In that case, for $k \in \mathbb{N}$, $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^3$ and $\alpha_1 + \alpha_2 + \alpha_3 = k$, we have that:

$$\begin{cases} (\frac{\partial \boldsymbol{f}}{\partial t}, \boldsymbol{f})_0 = \frac{1}{2} \frac{\partial}{\partial t} (\boldsymbol{f}, \boldsymbol{f})_0 = \frac{1}{2} \frac{\partial}{\partial t} \|\boldsymbol{f}\|_0^2; \\ ((\boldsymbol{f} \cdot \nabla) \boldsymbol{f}, \boldsymbol{f})_0 = 0; \\ ((\boldsymbol{f} \cdot \nabla) \boldsymbol{g}, \boldsymbol{f})_0 = -((\boldsymbol{f} \cdot \nabla) \boldsymbol{f}, \boldsymbol{g})_0; \\ ((\boldsymbol{f} \cdot \nabla) \boldsymbol{f}, \boldsymbol{g})_0 = -((\boldsymbol{f} \cdot \nabla) \boldsymbol{g}, \boldsymbol{f})_0; \\ (\partial_{\alpha}^k \Delta \boldsymbol{f}, \partial_{\alpha}^k \boldsymbol{g})_0 = -(\nabla \partial_{\alpha}^k \boldsymbol{f}, \nabla \partial_{\alpha}^k \boldsymbol{g})_0. \end{cases}$$
(4)

Proof. Direct calculation, integration by parts, and applying the Schwarz theorem to interchange the order of the partial derivatives lead to each of the above identities. \Box

Lemma 2. Assume the same conditions as the Theorem 1. Therefore, using the energy method, we find that:

$$\frac{1}{2} \frac{\partial}{\partial t} \left\| \partial_{\alpha}^{k} \tilde{\boldsymbol{w}} \right\|_{0}^{2} + \left(\partial_{\alpha}^{k} \left(\frac{\partial \boldsymbol{\omega}}{\partial t} \right), \partial_{\alpha}^{k} \tilde{\boldsymbol{\omega}} \right)_{0} + \left(\left(\partial_{\alpha}^{k} \left[(\boldsymbol{v} \cdot \nabla) \boldsymbol{\omega} \right], \partial_{\alpha}^{k} \tilde{\boldsymbol{\omega}} \right)_{0} + \left(\left(\partial_{\alpha}^{k} \left[(\boldsymbol{v} \cdot \nabla) \boldsymbol{\omega} \right], \partial_{\alpha}^{k} \tilde{\boldsymbol{\omega}} \right)_{0} + \left(\partial_{\alpha}^{k} \left[(\boldsymbol{v} \cdot \nabla) \boldsymbol{\omega} \right], \partial_{\alpha}^{k} \tilde{\boldsymbol{\omega}} \right)_{0} + \left(\left(\partial_{\alpha}^{k} \left[(\boldsymbol{v} \cdot \nabla) \boldsymbol{\omega} \right], \partial_{\alpha}^{k} \tilde{\boldsymbol{\omega}} \right)_{0} + \nu \left(\partial_{\alpha}^{k} \nabla \boldsymbol{\omega}, \partial_{\alpha}^{k} \nabla \boldsymbol{\omega} \right)_{0} + \nu \| \partial_{\alpha}^{k} \nabla \boldsymbol{\omega} \|_{0}^{2} = 0, \quad (5)$$

 $\forall k \in \mathbb{N} \text{ and } \forall \alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^3 \text{ such that } k = \alpha_1 + \alpha_2 + \alpha_3.$

Proof. Consider the Eq. (1), and apply the differential operator ∂_{α}^{k} to each of its terms:

$$\partial_{\alpha}^{k} \left[\frac{\partial \boldsymbol{\omega}}{\partial t} + \frac{\partial \tilde{\boldsymbol{\omega}}}{\partial t} + (\boldsymbol{v} \cdot \nabla)\boldsymbol{\omega} + (\tilde{\boldsymbol{v}} \cdot \nabla)\tilde{\boldsymbol{\omega}} + (\boldsymbol{v} \cdot \nabla)\tilde{\boldsymbol{\omega}} + (\tilde{\boldsymbol{v}} \cdot \nabla)\boldsymbol{\omega} - \nu\Delta(\boldsymbol{\omega} + \tilde{\boldsymbol{\omega}}) \right] = 0.$$
(6)

Now, we apply the inner product in the Lebesgue space $L_2(R;\mathbb{R}^3)$ of each side of Eq. (6) with $\partial_{\alpha}^k \tilde{\omega}$. According to the properties established in Eq. (4), we reach Eq. (5).

Corollary 1. If k = 0:

$$\frac{1}{2}\frac{\partial}{\partial t}\left\|\tilde{\boldsymbol{\omega}}\right\|_{0}^{2} + \left(\frac{\partial\boldsymbol{\omega}}{\partial t},\,\tilde{\boldsymbol{\omega}}\right)_{0} + \left((\boldsymbol{v}\cdot\nabla)\boldsymbol{\omega},\,\tilde{\boldsymbol{\omega}}\right)_{0} + \left((\tilde{\boldsymbol{v}}\cdot\nabla)\tilde{\boldsymbol{\omega}},\,\tilde{\boldsymbol{\omega}}\right)_{0} + \left((\boldsymbol{v}\cdot\nabla)\boldsymbol{\omega},\,\tilde{\boldsymbol{\omega}}\right)_{0} + \nu\left(\nabla\boldsymbol{\omega},\nabla\tilde{\boldsymbol{\omega}}\right)_{0} + \nu\left\|\nabla\tilde{\boldsymbol{\omega}}\right\|_{0}^{2} = 0. \quad (7)$$

In particular, for planar fields $\mathbf{v} = (v_1, v_2, 0)$, $\tilde{\mathbf{v}} = (\tilde{v}_1, \tilde{v}_2, 0)$:

$$\frac{1}{2}\frac{\partial}{\partial t}\left\|\tilde{\boldsymbol{\omega}}\right\|_{0}^{2} + \left(\frac{\partial\boldsymbol{\omega}}{\partial t}, \, \tilde{\boldsymbol{\omega}}\right)_{0} = 0.$$
(8)

Proof. Here, we can recall the relationships between the inner product and the norm in the Lebesgue Space of Eq. (4) for R and periodic boundary conditions for both v and \tilde{v} over ∂R to deduce Eq. (7). Moreover, if v and \tilde{v} are two dimensional, by definition of ω and $\tilde{\omega}$, the components $\omega_1, \omega_2, \tilde{\omega}_1, \tilde{\omega}_2 = 0$, and $\omega_3, \tilde{\omega}_3$ are independent of x_3 , so that the terms: $\|\nabla \omega\|_0^2 = 0$ and $(\nabla \omega, \nabla \tilde{\omega})_0 = 0$.

Theorem 2. Let $p, v, \tilde{v} \in C^1([0,T), C^{\infty}(\bar{R}))$, for some $0 < T < \infty$, with initial conditions $v_0 = v(\cdot, 0), \tilde{v}_0 = \tilde{v}(\cdot, 0) \in V^m(R)$, for $m \ge 2$, and periodic conditions for v and \tilde{v} at ∂R . Assume that (v, \tilde{v}, p) is a solution of:

$$\begin{cases} \nabla \cdot \boldsymbol{v} = 0, \\ \frac{D\boldsymbol{v}}{Dt} = \nu \Delta \boldsymbol{v} - \nabla p, \\ \nabla \cdot \tilde{\boldsymbol{v}} = 0, \\ \frac{D(\boldsymbol{v} + \tilde{\boldsymbol{v}})}{Dt} = \nu \Delta (\boldsymbol{v} + \tilde{\boldsymbol{v}}) - \nabla p. \end{cases}$$
(9)

Additionally, if the dimension n = 2, $R = (0, L_1) \times (0, L - 2)$, there exist a unique weak solution $\tilde{\boldsymbol{\omega}} = \nabla \times \tilde{\boldsymbol{v}}$ such that:

$$\frac{\partial \boldsymbol{\omega}}{\partial t} + \frac{\partial \boldsymbol{\omega}}{\partial t} + (\boldsymbol{v} \cdot \nabla)\boldsymbol{\omega} + (\boldsymbol{\tilde{v}} \cdot \nabla)\boldsymbol{\tilde{\omega}} + (\boldsymbol{v} \cdot \nabla)\boldsymbol{\tilde{\omega}} + (\boldsymbol{\tilde{v}} \cdot \nabla)\boldsymbol{\omega} - \nu\Delta(\boldsymbol{\omega} + \boldsymbol{\tilde{\omega}}) = 0,$$
(10)

for a finite time T < 0.

Proof. First, we recall the Picard Theorem for Banach Spaces, and consider the subspace of divergence free distributions $V^m(R)$ of the Sobolev space $H^m(R) = W^{k,2}(R)$, whose elements f, and weak derivatives $\partial_{\alpha}^k f$ have finite kinetic energy for $0 \le k \le m$. Second, we notice that the Conservation of Volume and Momentum laws corresponding to v have a unique weak solution $v \in C^1([0,T); C^k(\mathbb{R}^n))$, for a finite time T < 0, and whose vorticity $\omega = \nabla \times v$ satisfies a non-linear parabolic partial equation with a Maximum Principle, and the classical energy constraints. Finally, we extend the application of the Gagliardo-Niremberg inequalities to the convective terms of the form $(f \cdot \nabla g)$, and apply the Grönwall lemma to verify uniqueness.

Let \mathcal{B} is a Banach space with a distance $\|.\|_B$, and $U \subset \mathcal{B}$ is an open subset in \mathcal{B} , then $F: U \to \mathcal{B}$ is a Lipschitz operator if there is a magnitude L such that

$$\|F(x) - F(y)\|_{\mathcal{B}} \le L \|x - y\|_{\mathcal{B}}, \quad \forall x, y \in U.$$

Assume a fixed initial condition $\mathbf{f}(0) = \mathbf{f}_0 \in C^{\infty}(\bar{\mathcal{R}})$ of $\mathbf{f}(t) \in \mathcal{B}$. Let $U = B_{\mathbf{f}_0}^R \subset \mathcal{B}$ an open ball with center in \mathbf{f}_0 and radius R. If $F: U \to \mathcal{B}$ is a bounded, continuous and a Lipschitz operator in U, then there is a unique generalized solution $\mathbf{f} \in C^1([0,T]; U)$ to:

$$\frac{\partial \boldsymbol{f}}{\partial t} = F\left(\boldsymbol{f}\left(t\right)\right),\tag{11}$$

for that initial condition, and for $t \in [0, T)$, where $T = \min\left(\frac{R}{M}, \frac{1}{T}\right)$ and M > 0 is a bound of F is on U [4].

Let $\mathcal{B} = H^k(R)$. This is, \boldsymbol{f} and $\partial_{\alpha}^m \boldsymbol{f}$ belong to the Lebesgue space $L^2(R)$ for all $0 \le m \le k$. $H^k(R)$ is a Banach space [30]. In particular, $V^k(R)$ is the subspace of $H^k(R)$ of divergence free fields. The norm is indicated as: $\|\cdot\|_{\mathcal{B}} = \|\cdot\|_m$, where $(\boldsymbol{f}, \boldsymbol{g})_m = \sum_{0 \le k \le m} \iiint_R (D^k \boldsymbol{f} \cdot D^k \boldsymbol{g}) d\boldsymbol{x}$ is an inner product on $H^m(R)$, and $\|\boldsymbol{f}\|_m = (\boldsymbol{f}, \boldsymbol{f})_m$.

This result has been applied to show that the incompressible Navier-Stokes equations of system (9) with an initial condition $v_0 \in V^m(R)$ have a weak unique solution $v \in V^m(R)$ for a finite time T > 0 [4].

We define the operator $F: H^m(R) \to H^m(R)$:

$$F(\tilde{\boldsymbol{\omega}}) = -\frac{\partial \boldsymbol{\omega}}{\partial t} - (\boldsymbol{v} \cdot \nabla)\boldsymbol{\omega} - (\tilde{\boldsymbol{v}} \cdot \nabla)\tilde{\boldsymbol{\omega}} - (\boldsymbol{v} \cdot \nabla)\tilde{\boldsymbol{\omega}} - (\tilde{\boldsymbol{v}} \cdot \nabla)\boldsymbol{\omega} + \nu\Delta(\boldsymbol{\omega} + \tilde{\boldsymbol{\omega}})$$

with:

$$\begin{cases} F_{1}(\tilde{\boldsymbol{\omega}}) = -\frac{\partial \boldsymbol{\omega}}{\partial t}, \\ F_{2}(\boldsymbol{v}, \boldsymbol{\omega}) = -(\boldsymbol{v} \cdot \nabla)\boldsymbol{\omega}, \\ F_{3}(\tilde{\boldsymbol{v}}, \tilde{\boldsymbol{\omega}}) = -(\tilde{\boldsymbol{v}} \cdot \nabla)\tilde{\boldsymbol{\omega}}, \\ F_{4}(\boldsymbol{v}, \tilde{\boldsymbol{\omega}}) = -(\boldsymbol{v} \cdot \nabla)\tilde{\boldsymbol{\omega}}, \\ F_{5}(\tilde{\boldsymbol{v}}, \boldsymbol{\omega}) = -(\tilde{\boldsymbol{v}} \cdot \nabla)\boldsymbol{\omega}, \\ F_{5}(\tilde{\boldsymbol{v}}, \boldsymbol{\omega}) = -(\tilde{\boldsymbol{v}} \cdot \nabla)\boldsymbol{\omega}, \\ F_{6}(\boldsymbol{\omega}) = \nu\Delta\boldsymbol{\omega}, \\ F_{7}(\tilde{\boldsymbol{\omega}}) = \nu\Delta\tilde{\boldsymbol{\omega}}. \end{cases}$$
(12)

The first operator, F_1 , is determined by the parabolic equation satisfied by ω , where $\frac{\partial \omega}{\partial t} + (v \cdot \nabla \omega) = \nu \Delta \omega$, that is naturally reduced to an equation for the third component ω_3 of $\omega = (\omega_1, \omega_2, \omega_3)$. The last two Laplacian operators, F_6 and F_7 , are bounded and Lipschitz operators in $V^m(R)$. The second operator, F_2 inherits this properties from the classical case, where one can apply the Gagliardo-Niremberg inequalities. As far as we know, there are no improvements of the Gagliardo-Niremberg classical inequalities as it appear in [3, 4, 26], and [5, p. 81].

Model	Compressible	Incompressible	Stokes	Euler	Stationary
Busemann [9]	Irreducible	N/A	N/A	N/A	N/A
Prandtl [1]	N/A	Reducible	N/A	N/A	N/A
Navier-Stokes 3D	Irreducible	Irreducible	Reducible	Irreducible	Irreducible
RANS [10]	Irreducible	Irreducible	Reducible	Irreducible	Irreducible
ω -RANS	Irreducible	Irreducible	Reducible	Irreducible	Irreducible
ψ -NS	N/A	Reducible	N/A	N/A	Reducible

Figure 3: Experimental results

3 Experiments on multiplicity

Our empirical experiments suggest that the 3D Navier-Stokes equations as well as the 3D Reynolds-averaged Navier-Stokes equations (RANS) are irreducible as it is shown in the Figure 3 above. After attempting to utilize Boulier's algorithm on either Navier-Stokes or Reynolds-averaged Navier-Stokes (RANS) equations, the algorithm fails to converge unless the dimension is n = 2 and the problem is restated in terms of the *stream-function*. The complexity of the convective derivative has proved to be a major challenge in the path towards solving complex differential systems.

4 Conclusions

If we consider the existence and uniqueness of a classical velocity in the Navier-Stokes incompressible equations over a two dimensional domain for a finite time, then, we might think about the associated fluctuation properties as a consequence the original classical variables. The symmetry showed in the vorticity equations show that the fluctuation itself depends on the original one. Moreover, the application of Differential Algebra methods through the algorithm developed by Boulier and Lemaire allow new possible reductions and comparisons in the turbulent regime.

From the existence and uniqueness point of view, we will review the Maximum Principles available for quasi-linear parabolic equations (For example, those given by Oleinik in 1963.) to see if the result might be extended to n = 3. We would like to see the impact of different classical vorticity ω frequencies over $\tilde{\omega}$. In this direction, we will experiment with the algorithm to count rational solutions over differential polynomial systems; in particular, for periodic solutions. We greatly appreciate the support of the Universidad Iberoamericana Ciudad de México in the framework of the 18th call for scientific, humanistic and technological research 2023; in particular, the Direction of Research and Graduate Studies. We deeply appreciate the constant support of our families, friends and colleagues, without whom any journey would only be a dream.

Declarations of interest: none

References

- [1] F. Boulier, D. Lazard, F. Ollivier and M. Petitot, Computing representations for radicals of finitely generated differential ideals, *AAECC* (2009) **20** : 1, p.73-1219
- [2] C. V. Valencia-Negrete, *Ecuaciones de Navier-Stokes y Euler en* R^2 y R^3 , Instituto Politécnico Nacional (2014)
- [3] R. A. Adams, Sobolev Spaces, Academic Press (1975)
- [4] A. Majda and A. L. Bertozzi, Vorticity and Incompressible Flow, Cambridge University Press (2002)
- [5] H. Tanabe, *Functional Analytic Methods for Partial Differential Equations*, Monographs and textbooks in pure and applied Mathematics 204, Marcel Dekker Inc., (1996)
- [6] F. Boulier and F. Lemaire, Computing canonical representatives of regular differential ideals, ISSAC (2000)
- [7] F. Boulier, D. Lazard, F. Ollivier, and M. Petitot, Representation for the radical of a finitely generated differential ideal, *ISSAC* (1995)
- [8] J. F. Ritt, *Differential Algebra*, Dover Publications Inc., (1950)
- [9] E., R. Kolchin, Differential Algebra and Algebraic Groups, Academic Press, (1973)
- [10] H. A. Harrington and R. A. van Gorder, Reduction of dimension for nonlinear dynamical systems, *Nonlinear Dynamics* (2017) 88 p.715-734
- [11] B. Buchberger, Bruno Buchberger's PhD thesis 1965: An algorithm for finding the basis elements of the residue class ring of a zero dimensional polynomial ideal, *Journal of symbolic computation* (2006) **41** : 3 4, p.475-511

- [12] B. Buchberger, What is a Gröbner Basis, *Notices of the AMS* (2005) November 52 : 10, p.1199-1200
- [13] O. Darrigol, Worlds of Flow: A History of Hydrodynamics from the Bernoullis to Prandtl, Oxford University Press (2005)
- [14] O. Darrigol and U. Frisch, From Newton's mechanics to Euler's equations, *Physica D: Nonlinear Phenomena* (2008) 237: 14, p.1855-1869
- [15] U. Frisch, B. Hasslacher and Y. Pomeau, Lattice-Gas Automata for the Navier-Stokes Equation, *Physical Review Letters* (1986) 56 : 14, p. 1505-1508
- [16] J. Hardy, O. de Pazzis and Y. Pomeau, Molecular dynamics of a classical lattice gas: Transport properties and time correlation functions, *Phys. Rev. A* (1976) 13: 5 p. 1949-1961
- [17] S. Heinz Unified turbulence models for LES and RANS, FDF and PDF simulations, *Theoretical and Computational Fluid Dynamics* (2006) 21: 2 p. 99-118
- [18] T. Krüger, H. Kusumaatmaja, A. Kuzmin, O. Shardt, G. Silva and E. M. Viggen, *The Lattice Boltzmann Method Principles and Practice*, (2016) Springer
- [19] O. A. Ladyzhenskaya, Sixth problem of the millennium: Navier-Stokes equations, existence and smoothness, *Russian Mathematical Surveys* (2003) 58 : 2 p. 251–286
- [20] L. Saint-Raymond, *Hydrodynamic Limits of the Boltzmann Equation*, Lecture Notes in Mathematics, Springer (2009)
- [21] J. E. Marsden and S. Shkoller, Global well-posedness for the Lagrangian averaged Navier–Stokes (LANS– α) equations on bounded domains, *Physical review letters* (2001) **359** : 1784, p.1449-1468
- [22] J. E. Marsden and S. Shkoller, The Anisotropic Lagrangian Averaged Euler and Navier-Stokes Equations, *Arch. Rational Mech. Anal.*, (2003) **166** : 1 p.27-46
- [23] O. Reynolds, IV. On the dynamical theory of incompressible viscous fluids and the determination of the criterion, *Philosophical Transactions of the Royal Society of London. (A.)*, (1895) **186**, p. 123–164
- [24] O. Reynolds, An Experimental Investigation of the Circumstances Which Determine Whether the Motion of Water Shall Be Direct or Sinuous, and of the Law of Resistance in Parallel Channels, *Philosophical Transactions of the Royal Society of London*, (1883) 174, p. 935-982
- [25] L. R. Scott and R. Durst, Chaotic dynamics of two-dimensional flows around a cylinder, *Physics of Fluids* (2024) 36: 2, p.024118
- [26] L. Berselli, I. Traian and L. William, Mathematics of Large Eddy Simulation of Turbulent Flows, Springer (2006)
- [27] G. R. McNamara and G. Zanetti, Use of the Boltzmann Equation to Simulate Lattice-Gas Automata, *Physical review letters*, (1988) 61 : 12 p. 2332-2335
- [28] Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change by Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.), IPCC, Cambridge University Press. In Press (2021)
- [29] V. G. Maz'ya and S. V. Poborchi, S. V., Differentiable functions on bad domains, World Scientific (1997)
- [30] V. Maz'ya, Sobolev spaces, Springer (2013)
- [31] N. Piskunov, Differential and Integral Calculus, Mir Publishers (1969)
- [32] V. A. Zorich, Mathematical Analysis, 2, Springer (2002)