

Tau Data-Based Evaluations of the hadronic vacuum polarization contribution to the muon g-2

Pere Masjuan^{1,2}, Alejandro Miranda² and Pablo Roig^{3*}

1 Grup de Física Teòrica, Departament de Física, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain.

2 Institut de Física d'Altes Energies (IFAE) and The Barcelona Institute of Science and Technology (BIST), Campus UAB, 08193 Bellaterra (Barcelona), Spain.

3 Departamento de Física, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apdo. Postal 14-740, 07000 Ciudad de México, México.

* pablo.roig@cinvestav.mx



The 17th International Workshop on Tau Lepton Physics
Louisville, USA, 4-8 December 2023
doi:[10.21468/SciPostPhysProc.3](https://doi.org/10.21468/SciPostPhysProc.3)?

Abstract

We review the τ data-driven computation of Euclidean windows for the hadronic vacuum polarization contribution to the muon anomalous magnetic moment (a_μ), which agree with the lattice results, making the difference of the e^+e^- data-driven methods with them more intriguing. This conundrum needs to be solved by next year, when the final FNAL a_μ measurement will be published, in order to extract firm conclusions on possible new physics effects.

1 Introduction

The first two publications from the FNAL Muon g-2 Collaboration [1, 2], together with the BNL final result [3] yield the current experimental world average

$$a_\mu^{\text{exp}} = 116592059(22) \times 10^{-11}, \quad (1)$$

with an impressive accuracy of 0.19 ppm.

Although there is intense work, within the Muon g-2 Theory Initiative, to update the corresponding SM prediction, its recommended value is still the one in the White Paper [4]

$$a_\mu^{\text{SM}} = 116591810(43) \times 10^{-11}, \quad (2)$$

which is based on Refs. [5–39] (see also, e.g. [40–84]). The 5.1σ deviation between Eqs. (1) and (2) needs to be taken cautiously, as sketched below.

The BMW collaboration [85] achieved a very precise lattice QCD evaluation of the hadronic vacuum polarization contribution (HVP) which dominates the error of the SM prediction, and

challenged the WP number four years ago. According to this result, the discrepancy with respect to Eq. (1) would be reduced to 1.5σ . There is not yet another lattice QCD computation of similar accuracy which can be compared with the BMW number. Since Ref. [67], it has become standard to compare data-driven and lattice QCD evaluations of a_μ^{HVP} in three different windows in Euclidean time, which we pioneered using τ decay input in Ref. [60] (see also [74]), on which this contribution is fundamentally based. Remarkably, in the so-called intermediate window (to be introduced and discussed below), τ -based evaluations [60, 74] and the Mainz CLS [65], ETMC [66] and RBC/UKQCD [61] collaborations agree closely with BMW [85], with competing precision.

At the beginning of 2023, the CMD-3 Collaboration [86, 87] released their measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$, which disagreed substantially with previous data [88–93, 95–100] though was in line with the BMW result. a_μ^{HVP} based on CMD-3 alone would yield agreement between the SM prediction and Eq. (1) within one sigma.

In this puzzling situation we insist [41, 60] on the value of data-driven determinations of a_μ^{HVP} using τ input for the dominant di-pion channel, giving $\sim 73(85)\%$ of the whole value(uncertainty). An additional independent determination of a_μ^{HVP} will come soon from the space-like measurement in MUonE [101, 102] and will help to settle this question. After the end of the FNAL muon g-2 experiment, the J-PARC one will start operating, with the added value of being affected by completely different systematics [103]. The community hopes that this effort will further strength a_μ^{exp} and increase its precision, motivating to continue the improvement of a_μ^{SM} beyond 2025.

2 a_μ^{HVP} in the SM

The e^+e^- data-driven evaluation of the leading order HVP contribution to the muon g-2 is computed from

$$a_\mu^{\text{HVP, LO}} = \frac{1}{4\pi^3} \int_{s_{\text{thr}}}^\infty ds K(s) \sigma_{e^+e^- \rightarrow \text{hadrons}(\gamma)}^0(s), \quad (3)$$

which uses the measurement of the bare hadronic cross-section (specified by the upper-index 0). We note that both the $1/s$ behaviour of the cross-section (away from resonance peaks) and the $K(s)$ kernel enhance the low-energy contributions, making the precise evaluation of the $\pi^+\pi^-$ cut critical. Currently, the discrepancy between the BaBar [98] and KLOE [97] results has been very much aggravated by the CMD-3 data [86], jeopardizing this data-driven evaluation at the level of precision currently needed.

In this context, it is worth to resurrect the corresponding evaluation using τ data, that was once very useful in combination with the e^+e^- measurements to improve the accuracy of the SM prediction [104], and has been used fruitfully since then [105–118].

This requires isospin-breaking (IB) corrections to relate the measured di-pion τ decay spectrum to the needed bare cross-section

$$\sigma_{\pi\pi(\gamma)}^0 = \left[\frac{K_\sigma(s)}{K_\Gamma(s)} \frac{d\Gamma_{\pi\pi(\gamma)}}{ds} \right] \frac{R_{\text{IB}}(s)}{S_{\text{EW}}}, \quad (4)$$

where the first factor includes kinematic dependencies and global constants and S_{EW} corresponds to the universal short-distance radiative correction (see its updated value in [60]).

The most difficult corrections to evaluate are inside $R_{\text{IB}}(s)$ [106, 112]

$$R_{\text{IB}}(s) = \frac{\text{FSR}(s)}{G_{\text{EM}}(s)} \frac{\beta_{\pi^+\pi^-}^3(s)}{\beta_{\pi^-\pi^0}^3(s)} \left| \frac{F_V(s)}{f_+(s)} \right|^2, \quad (5)$$

where the easy part (for the needed precision) corresponds to the phase-space correction coming from the ratio of β functions and the final state radiative correction, FSR. It is much more difficult to quantify the long-distance process-dependent electromagnetic corrections entering the $G_{\text{EM}}(s)$ [106] and the difference between the neutral (F_V) and charged (f_+) di-pion form factors.

The S_{EW} contribution gives $\Delta a_\mu = -119.6 \times 10^{-11}$, consistent with earlier determinations and with a negligible error. The PS correction induces $\Delta a_\mu = -74.5 \times 10^{-11}$, as in previous works. The FSR correction yields $\Delta a_\mu = +45.5(4.6) \times 10^{-11}$, in accord with [112].

We evaluate [41] the structure-dependent IB corrections using Chiral Perturbation Theory [119–121] extended with Resonance fields and accounting for short-distance QCD constraints [122–126], using a large- N_C expansion [127].

We find that IB in the form factors induces the shift $+40.9(48.9) \times 10^{-11}$ or $+77.6(24.0) \times 10^{-11}$, depending on the input that we use. These results agree with earlier evaluations [106, 112], all with sizable uncertainties. For the G_{EM} effect, we obtain $(-15.9^{+5.7}_{-16.0}) \times 10^{-11}$, which agrees with Refs. [106, 112].

Fig. 1 shows the $\pi\pi$ contribution to $a_\mu^{\text{HVP, LO}}$ around the ρ peak, found using either $\sigma(e^+e^- \rightarrow \text{hadrons})$ (top, with average -without the CMD-3 point- in yellow) or the $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ spectrum (bottom, with mean in green that agrees with CMD-3). Tau data yields a $\sim 10 \times 10^{-10}$ larger value.




Figure 1: The $\pi\pi(\gamma)$ contribution to $a_\mu^{\text{HVP, LO}}$ around the ρ peak, obtained from the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ cross section (top) and di-pion τ decays (bottom).

In our recent paper [60] we translate these results into the window quantities introduced in Ref. [14], as used in [67]. The corresponding weights in Euclidean time or energy (the latter reproduced here as Fig. 2) are shown in Fig. 1 of [67]. It is important to clarify that the names can be misleading, as the intermediate window (of simply 'the window') receives significant contributions from both low and high energies. Similarly, the short-distance window is still mildly sensitive to hadronization, and the tail of the long-distance window enters

the perturbative regime. This observation needs to be taken into account when interpreting results in the different windows, with respect to the expected theory uncertainty. It is also crucial that the contributions of these windows to a_μ^{HVP} scale as $\sim 1 : 10 : 25$, respectively, so that the relative accuracy needed varies substantially between them.




Figure 2: Normalized weights of the short-distance (*SD*), intermediate (*win*, also *int*) and long-distance (*LD*) windows in \sqrt{s} [67].

Our main results for the three different window contributions to a_μ^{HVP} are plotted in Fig. 3, where the consistency among the different τ measurements is evident. In the *SD* and *int* windows, e^+e^- (from Ref. [67]) and τ data-based values are at odds.

Our τ -based $\pi\pi$ contribution to $a_\mu^{\text{HVP LO}}$ is supplemented with that from the remaining modes in order to compare it directly with the full evaluations. For this we considered two approaches, as explained in detail in Ref. [60], and took the difference between them as the uncertainty associated to this procedure. In this way, we found the results shown in Fig. 4. There is a clear trend of τ -based evaluations agreeing with the lattice results, while the e^+e^- ones differ markedly with both.

3 Conclusion

We have reviewed our τ -based evaluation of the dominant di-pion contribution to $a_\mu^{\text{HVP LO}}$ [41, 60]. Interestingly, it agrees nicely with the lattice QCD computations in the so-called intermediate window (as well as in the other ones), with the e^+e^- -based results differing clearly with both. In light of these findings and the CMD-3 data (which is compatible with the lattice and τ -based results), the consistency of the different $e^+e^- \rightarrow \pi^-\pi^-$ data seem puzzling, and further work appears needed to clarify this situation. This would be extremely important for interpreting the final FNAL measurement of a_μ , concluding on possible new physics contributions through its comparison with the SM prediction, which is not clear enough at the moment.

Acknowledgements

It is our pleasure to thank Swagato and his crew for this excellent workshop.




Figure 3: Window quantities (*SD* top, *int* medium, and *LD* bottom) for the 2π contribution below 1.0 GeV to a_μ^{HVP} according to our reference results. The blue band displays the mean from τ data, with the e^+e^- result from [67].

Funding information The work of P. M. has been supported by the European Union's Horizon 2020 Research and Innovation Programme under grant 824093 (H2020-INFRAIA-2018-1), the Ministerio de Ciencia e Innovación under grant PID2020-112965GB-I00, and by the Secretaria d'Universitats i Recerca del Departament d'Empresa i Coneixement de la Generalitat de Catalunya under grant 2021 SGR 00649. IFAE is partially funded by the CERCA program of the Generalitat de Catalunya. A. M. is also supported by MICINN with funding from European Union NextGenerationEU (PRTR-C17.I1) and by Generalitat de Catalunya. P. R. thanks partial funding from Conahcyt (México).




Figure 4: Comparison of the total intermediate window contribution to $a_\mu^{\text{HVP LO}}$ according to lattice QCD, e^+e^- and τ data-driven evaluations. The blue band corresponds to the weighted average of the lattice results excluding RBC/UKQCD 2018 [14] and ETMC 2021 [52] (superseded by refs. [61] and [66], respectively).

References

- [1] B. Abi *et al.* [Muon g-2], “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm,” Phys. Rev. Lett. **126** (2021) no.14, 141801, doi:[10.1103/PhysRevLett.126.141801](https://doi.org/10.1103/PhysRevLett.126.141801).
- [2] D. P. Aguillard *et al.* [Muon g-2], “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm,” Phys. Rev. Lett. **131** (2023) no.16, 161802, doi:[10.1103/PhysRevLett.131.161802](https://doi.org/10.1103/PhysRevLett.131.161802).
- [3] G. W. Bennett *et al.* [Muon g-2], “Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL,” Phys. Rev. D **73** (2006), 072003, doi:[10.1103/PhysRevD.73.072003](https://doi.org/10.1103/PhysRevD.73.072003).
- [4] T. Aoyama *et al.* “The anomalous magnetic moment of the muon in the Standard Model,” Phys. Rept. **887** (2020), 1-166, doi:[10.1016/j.physrep.2020.07.006](https://doi.org/10.1016/j.physrep.2020.07.006).
- [5] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, “Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon $g - 2$ and $\alpha(m_Z^2)$ using newest hadronic cross-section data,” Eur. Phys. J. C **77** (2017) no.12, 827, doi:[10.1140/epjc/s10052-017-5161-6](https://doi.org/10.1140/epjc/s10052-017-5161-6).
- [6] A. Keshavarzi, D. Nomura and T. Teubner, “Muon $g - 2$ and $\alpha(M_Z^2)$: a new data-based analysis,” Phys. Rev. D **97** (2018) no.11, 114025, doi:[10.1103/PhysRevD.97.114025](https://doi.org/10.1103/PhysRevD.97.114025).
- [7] G. Colangelo, M. Hoferichter and P. Stoffer, “Two-pion contribution to hadronic vacuum polarization,” JHEP **02** (2019), 006, doi:[10.1007/JHEP02\(2019\)006](https://doi.org/10.1007/JHEP02(2019)006).
- [8] M. Hoferichter, B. L. Hoid and B. Kubis, “Three-pion contribution to hadronic vacuum polarization,” JHEP **08** (2019), 137, doi:[10.1007/JHEP08\(2019\)137](https://doi.org/10.1007/JHEP08(2019)137).
- [9] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, “A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(m_Z^2)$,” Eur. Phys. J. C **80** (2020) no.3, 241 [erratum: Eur. Phys. J. C **80** (2020) no.5, 410], doi:[10.1140/epjc/s10052-020-7792-2](https://doi.org/10.1140/epjc/s10052-020-7792-2).

- [10] A. Keshavarzi, D. Nomura and T. Teubner, “ $g - 2$ of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium,” Phys. Rev. D **101** (2020) no.1, 014029, doi:[10.1103/PhysRevD.101.014029](https://doi.org/10.1103/PhysRevD.101.014029).
- [11] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, “Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order,” Phys. Lett. B **734** (2014), 144-147, doi:[10.1016/j.physletb.2014.05.043](https://doi.org/10.1016/j.physletb.2014.05.043).
- [12] B. Chakraborty *et al.* [Fermilab Lattice, LATTICE-HPQCD and MILC], “Strong-Isospin-Breaking Correction to the Muon Anomalous Magnetic Moment from Lattice QCD at the Physical Point,” Phys. Rev. Lett. **120** (2018) no.15, 152001, doi:[10.1103/PhysRevLett.120.152001](https://doi.org/10.1103/PhysRevLett.120.152001).
- [13] S. Borsanyi *et al.* [Budapest-Marseille-Wuppertal], “Hadronic vacuum polarization contribution to the anomalous magnetic moments of leptons from first principles,” Phys. Rev. Lett. **121** (2018) no.2, 022002, doi:[10.1103/PhysRevLett.121.022002](https://doi.org/10.1103/PhysRevLett.121.022002).
- [14] T. Blum *et al.* [RBC and UKQCD], “Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment,” Phys. Rev. Lett. **121** (2018) no.2, 022003, doi:[10.1103/PhysRevLett.121.022003](https://doi.org/10.1103/PhysRevLett.121.022003).
- [15] D. Giusti, V. Lubicz, G. Martinelli, F. Sanfilippo and S. Simula, “Electromagnetic and strong isospin-breaking corrections to the muon $g - 2$ from Lattice QCD+QED,” Phys. Rev. D **99** (2019) no.11, 114502, doi:[10.1103/PhysRevD.99.114502](https://doi.org/10.1103/PhysRevD.99.114502).
- [16] E. Shintani *et al.* [PACS], “Hadronic vacuum polarization contribution to the muon $g - 2$ with 2+1 flavor lattice QCD on a larger than $(10 \text{ fm})^4$ lattice at the physical point,” Phys. Rev. D **100** (2019) no.3, 034517, doi:[10.1103/PhysRevD.100.034517](https://doi.org/10.1103/PhysRevD.100.034517).
- [17] C. T. H. Davies *et al.* [Fermilab Lattice, LATTICE-HPQCD and MILC], “Hadronic-vacuum-polarization contribution to the muon’s anomalous magnetic moment from four-flavor lattice QCD,” Phys. Rev. D **101** (2020) no.3, 034512, doi:[10.1103/PhysRevD.101.034512](https://doi.org/10.1103/PhysRevD.101.034512).
- [18] A. Gérardin, M. Cè, G. von Hippel, B. Hörrz, H. B. Meyer, D. Mohler, K. Ottnad, J. Wilhelm and H. Wittig, “The leading hadronic contribution to $(g - 2)_\mu$ from lattice QCD with $N_f = 2 + 1$ flavours of $O(a)$ improved Wilson quarks,” Phys. Rev. D **100** (2019) no.1, 014510, doi:[10.1103/PhysRevD.100.014510](https://doi.org/10.1103/PhysRevD.100.014510).
- [19] C. Aubin, T. Blum, C. Tu, M. Golterman, C. Jung and S. Peris, “Light quark vacuum polarization at the physical point and contribution to the muon $g - 2$,” Phys. Rev. D **101** (2020) no.1, 014503, doi:[10.1103/PhysRevD.101.014503](https://doi.org/10.1103/PhysRevD.101.014503).
- [20] D. Giusti and S. Simula, “Lepton anomalous magnetic moments in Lattice QCD+QED,” PoS LATTICE**2019** (2019), 104, doi:[10.22323/1.363.0104](https://doi.org/10.22323/1.363.0104).
- [21] K. Melnikov and A. Vainshtein, “Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment revisited,” Phys. Rev. D **70** (2004), 113006, doi:[10.1103/PhysRevD.70.113006](https://doi.org/10.1103/PhysRevD.70.113006).
- [22] P. Masjuan and P. Sánchez-Puertas, “Pseudoscalar-pole contribution to the $(g_\mu - 2)$: a rational approach,” Phys. Rev. D **95** (2017) no.5, 054026, doi:[10.1103/PhysRevD.95.054026](https://doi.org/10.1103/PhysRevD.95.054026).

- [23] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, “Dispersion relation for hadronic light-by-light scattering: two-pion contributions,” JHEP **04** (2017), 161, doi:[10.1007/JHEP04\(2017\)161](https://doi.org/10.1007/JHEP04(2017)161).
- [24] M. Hoferichter, B. L. Hoid, B. Kubis, S. Leupold and S. P Schneider, “Dispersion relation for hadronic light-by-light scattering: pion pole,” JHEP **10** (2018), 141, doi:[10.1007/JHEP10\(2018\)141](https://doi.org/10.1007/JHEP10(2018)141).
- [25] A. Gérardin, H. B. Meyer and A. Nyffeler, “Lattice calculation of the pion transition form factor with $N_f = 2 + 1$ Wilson quarks,” Phys. Rev. D **100** (2019) no.3, 034520, doi:[10.1103/PhysRevD.100.034520](https://doi.org/10.1103/PhysRevD.100.034520).
- [26] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, “Short-distance constraints for the HLbL contribution to the muon anomalous magnetic moment,” Phys. Lett. B **798** (2019), 134994, doi:[10.1016/j.physletb.2019.134994](https://doi.org/10.1016/j.physletb.2019.134994).
- [27] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, “Longitudinal short-distance constraints for the hadronic light-by-light contribution to $(g-2)_\mu$ with large- N_c Regge models,” JHEP **03** (2020), 101, doi:[10.1007/JHEP03\(2020\)101](https://doi.org/10.1007/JHEP03(2020)101).
- [28] V. Pauk and M. Vanderhaeghen, “Single meson contributions to the muon’s anomalous magnetic moment,” Eur. Phys. J. C **74** (2014) no.8, 3008, doi:[10.1140/epjc/s10052-014-3008-y](https://doi.org/10.1140/epjc/s10052-014-3008-y).
- [29] I. Danilkin and M. Vanderhaeghen, “Light-by-light scattering sum rules in light of new data,” Phys. Rev. D **95** (2017) no.1, 014019, doi:[10.1103/PhysRevD.95.014019](https://doi.org/10.1103/PhysRevD.95.014019).
- [30] F. Jegerlehner, “The Anomalous Magnetic Moment of the Muon,” Springer Tracts Mod. Phys. **274** (2017), pp.1-693, Springer, 2017, doi:[10.1007/978-3-319-63577-4](https://doi.org/10.1007/978-3-319-63577-4).
- [31] M. Knecht, S. Narison, A. Rabemananjara and D. Rabetiarivony, “Scalar meson contributions to a_μ from hadronic light-by-light scattering,” Phys. Lett. B **787** (2018), 111-123, doi:[10.1016/j.physletb.2018.10.048](https://doi.org/10.1016/j.physletb.2018.10.048).
- [32] G. Eichmann, C. S. Fischer and R. Williams, “Kaon-box contribution to the anomalous magnetic moment of the muon,” Phys. Rev. D **101** (2020) no.5, 054015, doi:[10.1103/PhysRevD.101.054015](https://doi.org/10.1103/PhysRevD.101.054015).
- [33] P. Roig and P. Sánchez-Puertas, “Axial-vector exchange contribution to the hadronic light-by-light piece of the muon anomalous magnetic moment,” Phys. Rev. D **101** (2020) no.7, 074019, doi:[10.1103/PhysRevD.101.074019](https://doi.org/10.1103/PhysRevD.101.074019).
- [34] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, “Remarks on higher-order hadronic corrections to the muon g-2,” Phys. Lett. B **735** (2014), 90-91, doi:[10.1016/j.physletb.2014.06.012](https://doi.org/10.1016/j.physletb.2014.06.012).
- [35] T. Blum, N. Christ, M. Hayakawa, T. Izubuchi, L. Jin, C. Jung and C. Lehner, “Hadronic Light-by-Light Scattering Contribution to the Muon Anomalous Magnetic Moment from Lattice QCD,” Phys. Rev. Lett. **124** (2020) no.13, 132002, doi:[10.1103/PhysRevLett.124.132002](https://doi.org/10.1103/PhysRevLett.124.132002).
- [36] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, “Complete Tenth-Order QED Contribution to the Muon g-2,” Phys. Rev. Lett. **109** (2012), 111808, doi:[10.1103/PhysRevLett.109.111808](https://doi.org/10.1103/PhysRevLett.109.111808).

- [37] T. Aoyama, T. Kinoshita and M. Nio, “Theory of the Anomalous Magnetic Moment of the Electron,” Atoms **7** (2019) no.1, 28, doi:[10.3390/atoms7010028](https://doi.org/10.3390/atoms7010028).
- [38] A. Czarnecki, W. J. Marciano and A. Vainshtein, “Refinements in electroweak contributions to the muon anomalous magnetic moment,” Phys. Rev. D **67** (2003), 073006 [erratum: Phys. Rev. D **73** (2006), 119901], doi:[10.1103/PhysRevD.67.073006](https://doi.org/10.1103/PhysRevD.67.073006).
- [39] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, “The electroweak contributions to $(g - 2)_\mu$ after the Higgs boson mass measurement,” Phys. Rev. D **88** (2013), 053005, doi:[10.1103/PhysRevD.88.053005](https://doi.org/10.1103/PhysRevD.88.053005).
- [40] G. Colangelo *et al.* “Prospects for precise predictions of a_μ in the Standard Model,” [arXiv:2203.15810 [hep-ph]], contribution to Snowmass.
- [41] J. A. Miranda and P. Roig, “New τ -based evaluation of the hadronic contribution to the vacuum polarization piece of the muon anomalous magnetic moment,” Phys. Rev. D **102** (2020), 114017, doi:[10.1103/PhysRevD.102.114017](https://doi.org/10.1103/PhysRevD.102.114017).
- [42] P. Masjuan, P. Roig and P. Sánchez-Puertas, “The interplay of transverse degrees of freedom and axial-vector mesons with short-distance constraints in $g-2$,” J. Phys. G **49** (2022) no.1, 015002, doi:[10.1088/1361-6471/ac3892](https://doi.org/10.1088/1361-6471/ac3892).
- [43] J. Lüdtke and M. Procura, “Effects of longitudinal short-distance constraints on the hadronic light-by-light contribution to the muon $g - 2$,” Eur. Phys. J. C **80** (2020) no.12, 1108, doi:[10.1140/epjc/s10052-020-08611-6](https://doi.org/10.1140/epjc/s10052-020-08611-6).
- [44] J. Bijnens, N. Hermansson-Truedsson, L. Laub and A. Rodríguez-Sánchez, “The two-loop perturbative correction to the $(g - 2)_\mu$ HLbL at short distances,” JHEP **04** (2021), 240, doi:[10.1007/JHEP04\(2021\)240](https://doi.org/10.1007/JHEP04(2021)240).
- [45] L. Cappiello, O. Catà and G. D’Ambrosio, “Scalar resonances in the hadronic light-by-light contribution to the muon $(g-2)$,” Phys. Rev. D **105** (2022) no.5, 056020, doi:[10.1103/PhysRevD.105.056020](https://doi.org/10.1103/PhysRevD.105.056020).
- [46] E. H. Chao, R. J. Hudspith, A. Gérardin, J. R. Green, H. B. Meyer and K. Ott nad, “Hadronic light-by-light contribution to $(g - 2)_\mu$ from lattice QCD: a complete calculation,” Eur. Phys. J. C **81** (2021) no.7, 651, doi:[10.1140/epjc/s10052-021-09455-4](https://doi.org/10.1140/epjc/s10052-021-09455-4).
- [47] G. Colangelo, M. Hoferichter, B. Kubis, M. Niehus and J. R. de Elvira, “Chiral extrapolation of hadronic vacuum polarization,” Phys. Lett. B **825** (2022), 136852, doi:[10.1016/j.physletb.2021.136852](https://doi.org/10.1016/j.physletb.2021.136852).
- [48] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, “Short-distance constraints for the longitudinal component of the hadronic light-by-light amplitude: an update,” Eur. Phys. J. C **81** (2021) no.8, 702, doi:[10.1140/epjc/s10052-021-09513-x](https://doi.org/10.1140/epjc/s10052-021-09513-x).
- [49] I. Danilkin, M. Hoferichter and P. Stoffer, “A dispersive estimate of scalar contributions to hadronic light-by-light scattering,” Phys. Lett. B **820** (2021), 136502, doi:[10.1016/j.physletb.2021.136502](https://doi.org/10.1016/j.physletb.2021.136502).
- [50] M. Hoferichter and T. Teubner, “Mixed Leptonic and Hadronic Corrections to the Anomalous Magnetic Moment of the Muon,” Phys. Rev. Lett. **128** (2022) no.11, 112002, doi:[10.1103/PhysRevLett.128.112002](https://doi.org/10.1103/PhysRevLett.128.112002).

- [51] J. Leutgeb and A. Rebhan, “Hadronic light-by-light contribution to the muon g-2 from holographic QCD with massive pions,” Phys. Rev. D **104** (2021) no.9, 094017, doi:[10.1103/PhysRevD.104.094017](https://doi.org/10.1103/PhysRevD.104.094017).
- [52] D. Giusti and S. Simula, “Window contributions to the muon hadronic vacuum polarization with twisted-mass fermions,” PoS **LATTICE2021** (2022), 189, doi:[10.22323/1.396.0189](https://doi.org/10.22323/1.396.0189).
- [53] Á. Miramontes, A. Bashir, K. Raya and P. Roig, “Pion and Kaon box contribution to $a\mu_{HLbL}$,” Phys. Rev. D **105** (2022) no.7, 074013, doi:[10.1103/PhysRevD.105.074013](https://doi.org/10.1103/PhysRevD.105.074013).
- [54] V. Biloshytskyi, E. H. Chao, A. Gérardin, J. R. Green, F. Hagelstein, H. B. Meyer, J. Parrino and V. Pascalutsa, “Forward light-by-light scattering and electromagnetic correction to hadronic vacuum polarization,” JHEP **03** (2023), 194, doi:[10.1007/JHEP03\(2023\)194](https://doi.org/10.1007/JHEP03(2023)194).
- [55] G. Colangelo, M. Hoferichter, B. Kubis and P. Stoffer, “Isospin-breaking effects in the two-pion contribution to hadronic vacuum polarization,” JHEP **10** (2022), 032, doi:[10.1007/JHEP10\(2022\)032](https://doi.org/10.1007/JHEP10(2022)032).
- [56] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, “Constraints on the hadronic light-by-light in the Melnikov-Vainshtein regime,” JHEP **02** (2023), 167, doi:[10.1007/JHEP02\(2023\)167](https://doi.org/10.1007/JHEP02(2023)167).
- [57] D. Boito, M. Golterman, K. Maltman and S. Peris, “Evaluation of the three-flavor quark-disconnected contribution to the muon anomalous magnetic moment from experimental data,” Phys. Rev. D **105** (2022) no.9, 093003, doi:[10.1103/PhysRevD.105.093003](https://doi.org/10.1103/PhysRevD.105.093003).
- [58] D. Boito, M. Golterman, K. Maltman and S. Peris, “Data-based determination of the isospin-limit light-quark-connected contribution to the anomalous magnetic moment of the muon,” Phys. Rev. D **107** (2023) no.7, 074001, doi:[10.1103/PhysRevD.107.074001](https://doi.org/10.1103/PhysRevD.107.074001).
- [59] D. Stamen, D. Hariharan, M. Hoferichter, B. Kubis and P. Stoffer, “Kaon electromagnetic form factors in dispersion theory,” Eur. Phys. J. C **82** (2022) no.5, 432 doi:[10.1140/epjc/s10052-022-10348-3](https://doi.org/10.1140/epjc/s10052-022-10348-3).
- [60] P. Masjuan, A. Miranda and P. Roig, “ τ data-driven evaluation of Euclidean windows for the hadronic vacuum polarization,” Phys. Lett. B **850** (2024), 138492, doi:[10.1016/j.physletb.2024.138492](https://doi.org/10.1016/j.physletb.2024.138492).
- [61] T. Blum *et al.* [RBC and UKQCD], “Update of Euclidean windows of the hadronic vacuum polarization,” Phys. Rev. D **108** (2023) no.5, 054507, doi:[10.1103/PhysRevD.108.054507](https://doi.org/10.1103/PhysRevD.108.054507).
- [62] J. Lüdtke, M. Procura and P. Stoffer, “Dispersion relations for hadronic light-by-light scattering in triangle kinematics,” JHEP **04** (2023), 125, doi:[10.1007/JHEP04\(2023\)125](https://doi.org/10.1007/JHEP04(2023)125).
- [63] M. Davier, D. Díaz-Calderón, B. Malaescu, A. Pich, A. Rodríguez-Sánchez and Z. Zhang, “The Euclidean Adler function and its interplay with $\Delta\alpha_{\text{QED}}^{\text{had}}$ and α_s ,” JHEP **04** (2023), 067, doi:[10.1007/JHEP04\(2023\)067](https://doi.org/10.1007/JHEP04(2023)067).
- [64] S. J. Wang, Z. Fang and L. Y. Dai, “Two body final states production in electron-positron annihilation and their contributions to $(g-2)_\mu$,” JHEP **07** (2023), 037, doi:[10.1007/JHEP07\(2023\)037](https://doi.org/10.1007/JHEP07(2023)037).

- [65] M. Cè *et al.* “Window observable for the hadronic vacuum polarization contribution to the muon $g-2$ from lattice QCD,” Phys. Rev. D **106** (2022) no.11, 114502, doi:[10.1103/PhysRevD.106.114502](https://doi.org/10.1103/PhysRevD.106.114502).
- [66] C. Alexandrou *et al.* [Extended Twisted Mass], “Lattice calculation of the short and intermediate time-distance hadronic vacuum polarization contributions to the muon magnetic moment using twisted-mass fermions,” Phys. Rev. D **107** (2023) no.7, 074506, doi:[10.1103/PhysRevD.107.074506](https://doi.org/10.1103/PhysRevD.107.074506).
- [67] G. Colangelo, A. X. El-Khadra, M. Hoferichter, A. Keshavarzi, C. Lehner, P. Stoffer and T. Teubner, “Data-driven evaluations of Euclidean windows to scrutinize hadronic vacuum polarization,” Phys. Lett. B **833** (2022), 137313, doi:[10.1016/j.physletb.2022.137313](https://doi.org/10.1016/j.physletb.2022.137313).
- [68] G. Benton, D. Boito, M. Golterman, A. Keshavarzi, K. Maltman and S. Peris, “Data-Driven Determination of the Light-Quark Connected Component of the Intermediate-Window Contribution to the Muon $g-2$,” Phys. Rev. Lett. **131** (2023) no.25, 251803, doi:[10.1103/PhysRevLett.131.251803](https://doi.org/10.1103/PhysRevLett.131.251803).
- [69] C. Lehner and A. S. Meyer, “Consistency of hadronic vacuum polarization between lattice QCD and the R-ratio,” Phys. Rev. D **101** (2020), 074515, doi:[10.1103/PhysRevD.101.074515](https://doi.org/10.1103/PhysRevD.101.074515).
- [70] G. Wang *et al.* [chiQCD], “Muon $g-2$ with overlap valence fermions,” Phys. Rev. D **107** (2023) no.3, 034513, doi:[10.1103/PhysRevD.107.034513](https://doi.org/10.1103/PhysRevD.107.034513).
- [71] C. Aubin, T. Blum, M. Golterman and S. Peris, “Muon anomalous magnetic moment with staggered fermions: Is the lattice spacing small enough?,” Phys. Rev. D **106** (2022) no.5, 054503, doi:[10.1103/PhysRevD.106.054503](https://doi.org/10.1103/PhysRevD.106.054503).
- [72] A. Bazavov *et al.* [Fermilab Lattice, HPQCD, and MILC], “Light-quark connected intermediate-window contributions to the muon $g-2$ hadronic vacuum polarization from lattice QCD,” Phys. Rev. D **107** (2023) no.11, 114514, doi:[10.1103/PhysRevD.107.114514](https://doi.org/10.1103/PhysRevD.107.114514).
- [73] M. Davier, Z. Fodor, A. Gerardin, L. Lellouch, B. Malaescu, F. M. Stokes, K. K. Szabo, B. C. Toth, L. Varnhorst and Z. Zhang, “Hadronic vacuum polarization: comparing lattice QCD and data-driven results in systematically improvable ways,” Phys. Rev. D **109** (2024) no.7, 076019, doi:[10.1103/PhysRevD.109.076019](https://doi.org/10.1103/PhysRevD.109.076019).
- [74] M. Davier, A. Hoecker, A. M. Lutz, B. Malaescu and Z. Zhang, “Tensions in $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ measurements: the new landscape of data-driven hadronic vacuum polarization predictions for the muon $g-2$,” [arXiv:2312.02053 [hep-ph]].
- [75] M. Hoferichter, G. Colangelo, B. L. Hoid, B. Kubis, J. R. de Elvira, D. Schuh, D. Stamen and P. Stoffer, “Phenomenological Estimate of Isospin Breaking in Hadronic Vacuum Polarization,” Phys. Rev. Lett. **131** (2023) no.16, 161905, doi:[10.1103/PhysRevLett.131.161905](https://doi.org/10.1103/PhysRevLett.131.161905).
- [76] J. Bijnens, N. Hermansson-Truedsson, L. Laub and A. Rodríguez-Sánchez, “Short-distance HLbL contributions to the muon anomalous magnetic moment beyond perturbation theory,” JHEP **10** (2020), 203 doi:[10.1007/JHEP10\(2020\)203](https://doi.org/10.1007/JHEP10(2020)203).
- [77] E. H. Chao, A. Gérardin, J. R. Green, R. J. Hudspith and H. B. Meyer, “Hadronic light-by-light contribution to $(g-2)_\mu$ from lattice QCD with SU(3) flavor symmetry,” Eur. Phys. J. C **80** (2020) no.9, 869 doi:[10.1140/epjc/s10052-020-08444-3](https://doi.org/10.1140/epjc/s10052-020-08444-3).

- [78] G. Colangelo, M. Hoferichter and P. Stoffer, “Constraints on the two-pion contribution to hadronic vacuum polarization,” Phys. Lett. B **814** (2021), 136073 doi:[10.1016/j.physletb.2021.136073](https://doi.org/10.1016/j.physletb.2021.136073).
- [79] B. L. Hoid, M. Hoferichter and B. Kubis, “Hadronic vacuum polarization and vector-meson resonance parameters from $e^+e^- \rightarrow \pi^0\gamma$,” Eur. Phys. J. C **80** (2020) no.10, 988, doi:[10.1140/epjc/s10052-020-08550-2](https://doi.org/10.1140/epjc/s10052-020-08550-2).
- [80] A. Miranda, P. Roig and P. Sánchez-Puertas, “Axial-vector exchange contribution to the hyperfine splitting,” Phys. Rev. D **105** (2022) no.1, 016017, doi:[10.1103/PhysRevD.105.016017](https://doi.org/10.1103/PhysRevD.105.016017).
- [81] R. Escribano, J. A. Miranda and P. Roig, “Radiative corrections to the $\tau^- \rightarrow (P_1 P_2)^- \nu_\tau (P_{1,2} = \pi, K)$ decays,” Phys. Rev. D **109** (2024) no.5, 053003, doi:[10.1103/PhysRevD.109.053003](https://doi.org/10.1103/PhysRevD.109.053003).
- [82] P. Roig, A. Guevara and G. López Castro, “ $VV'P$ form factors in resonance chiral theory and the $\pi - \eta - \eta'$ light-by-light contribution to the muon $g - 2$,” Phys. Rev. D **89** (2014) no.7, 073016, doi:[10.1103/PhysRevD.89.073016](https://doi.org/10.1103/PhysRevD.89.073016).
- [83] A. Guevara, P. Roig and J. J. Sanz-Cillero, “Pseudoscalar pole light-by-light contributions to the muon ($g - 2$) in Resonance Chiral Theory,” JHEP **06** (2018), 160, doi:[10.1007/JHEP06\(2018\)160](https://doi.org/10.1007/JHEP06(2018)160).
- [84] K. Raya, A. Bashir and P. Roig, “Contribution of neutral pseudoscalar mesons to a_μ^{HLbL} within a Schwinger-Dyson equations approach to QCD,” Phys. Rev. D **101** (2020) no.7, 074021, doi:[10.1103/PhysRevD.101.074021](https://doi.org/10.1103/PhysRevD.101.074021).
- [85] S. Borsanyi *et al.* “Leading hadronic contribution to the muon magnetic moment from lattice QCD,” Nature **593** (2021) no.7857, 51-55, doi:[10.1038/s41586-021-03418-1](https://doi.org/10.1038/s41586-021-03418-1).
- [86] F. V. Ignatov *et al.* [CMD-3], “Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section from threshold to 1.2 GeV with the CMD-3 detector,” [arXiv:2302.08834 [hep-ex]].
- [87] F. V. Ignatov *et al.* [CMD-3], “Measurement of the pion formfactor with CMD-3 detector and its implication to the hadronic contribution to muon ($g-2$),” [arXiv:2309.12910 [hep-ex]].
- [88] R. R. Akhmetshin *et al.* [CMD-2], “Reanalysis of hadronic cross-section measurements at CMD-2,” Phys. Lett. B **578** (2004), 285-289, doi:[10.1016/j.physletb.2003.10.108](https://doi.org/10.1016/j.physletb.2003.10.108).
- [89] V. M. Aul’chenko *et al.* [CMD-2], “Measurement of the pion form-factor in the range 1.04-GeV to 1.38-GeV with the CMD-2 detector,” JETP Lett. **82** (2005), 743-747, doi:[10.1134/1.2175241](https://doi.org/10.1134/1.2175241).
- [90] T. K. Pedlar *et al.* [CLEO], “Precision measurements of the timelike electromagnetic form-factors of pion, kaon, and proton,” Phys. Rev. Lett. **95** (2005), 261803, doi:[10.1103/PhysRevLett.95.261803](https://doi.org/10.1103/PhysRevLett.95.261803).
- [91] V. M. Aul’chenko *et al.* “Measurement of the e+ e- to pi+ pi- cross section with the CMD-2 detector in the 370 - 520-MeV c.m. energy range,” JETP Lett. **84** (2006), 413-417, doi:[10.1134/S0021364006200021](https://doi.org/10.1134/S0021364006200021).
- [92] R. R. Akhmetshin *et al.* [CMD-2], “High-statistics measurement of the pion form factor in the rho-meson energy range with the CMD-2 detector,” Phys. Lett. B **648** (2007), 28-38, doi:[10.1016/j.physletb.2007.01.073](https://doi.org/10.1016/j.physletb.2007.01.073).

- [93] M. N. Achasov *et al.* “Update of the $e^+e^- \rightarrow \pi^+\pi^-$ cross-section measured by SND detector in the energy region $400 \text{ MeV} < E < 1000 \text{ MeV}$,” *J. Exp. Theor. Phys.* **103** (2006), 380-384, doi:[10.1134/S106377610609007X](https://doi.org/10.1134/S106377610609007X).
- [94] F. Ambrosino *et al.* [KLOE], “Measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ and the dipion contribution to the muon anomaly with the KLOE detector,” *Phys. Lett. B* **670** (2009), 285-291, doi:[10.1016/j.physletb.2008.10.060](https://doi.org/10.1016/j.physletb.2008.10.060).
- [95] F. Ambrosino *et al.* [KLOE], “Measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$ from threshold to 0.85 GeV^2 using Initial State Radiation with the KLOE detector,” *Phys. Lett. B* **700** (2011), 102-110, doi:[10.1016/j.physletb.2011.04.055](https://doi.org/10.1016/j.physletb.2011.04.055).
- [96] D. Babusci *et al.* [KLOE], “Precision measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma)/\sigma(e^+e^- \rightarrow \mu^+\mu^-\gamma)$ and determination of the $\pi^+\pi^-$ contribution to the muon anomaly with the KLOE detector,” *Phys. Lett. B* **720** (2013), 336-343, doi:[10.1016/j.physletb.2013.02.029](https://doi.org/10.1016/j.physletb.2013.02.029).
- [97] A. Anastasi *et al.* [KLOE-2], “Combination of KLOE $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ measurements and determination of $a_\mu^{\pi^+\pi^-}$ in the energy range $0.10 < s < 0.95 \text{ GeV}^2$,” *JHEP* **03** (2018), 173, doi:[10.1007/JHEP03\(2018\)173](https://doi.org/10.1007/JHEP03(2018)173).
- [98] J. P. Lees *et al.* [BaBar], “Precise Measurement of the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ Cross Section with the Initial-State Radiation Method at BABAR,” *Phys. Rev. D* **86** (2012), 032013, doi:[10.1103/PhysRevD.86.032013](https://doi.org/10.1103/PhysRevD.86.032013).
- [99] M. Ablikim *et al.* [BESIII], “Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section between 600 and 900 MeV using initial state radiation,” *Phys. Lett. B* **753** (2016), 629-638 [erratum: *Phys. Lett. B* **812** (2021), 135982], doi:[10.1016/j.physletb.2015.11.043](https://doi.org/10.1016/j.physletb.2015.11.043).
- [100] M. N. Achasov *et al.* [SND], “Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ process cross section with the SND detector at the VEPP-2000 collider in the energy region $0.525 < \sqrt{s} < 0.883 \text{ GeV}$,” *JHEP* **01** (2021), 113, doi:[10.1007/JHEP01\(2021\)113](https://doi.org/10.1007/JHEP01(2021)113).
- [101] P. Banerjee *et al.* “Theory for muon-electron scattering @ 10 ppm: A report of the MUonE theory initiative,” *Eur. Phys. J. C* **80** (2020) no.6, 591. doi:[10.1140/epjc/s10052-020-8138-9](https://doi.org/10.1140/epjc/s10052-020-8138-9).
- [102] G. Abbiendi, “Status of the MUonE experiment,” *Phys. Scripta* **97** (2022) no.5, 054007, doi:[10.1088/1402-4896/ac6297](https://doi.org/10.1088/1402-4896/ac6297).
- [103] H. Iinuma [J-PARC muon g-2/EDM], “New approach to the muon g-2 and EDM experiment at J-PARC,” *J. Phys. Conf. Ser.* **295** (2011), 012032 doi:[10.1088/1742-6596/295/1/012032](https://doi.org/10.1088/1742-6596/295/1/012032).
- [104] R. Alemany, M. Davier and A. Höcker, “Improved determination of the hadronic contribution to the muon (g-2) and to alpha (M(z)) using new data from hadronic tau decays,” *Eur. Phys. J. C* **2** (1998), 123-135, doi:[10.1007/s100520050127](https://doi.org/10.1007/s100520050127).
- [105] S. Narison, “Muon and tau anomalies updated,” *Phys. Lett. B* **513** (2001), 53-70 [erratum: *Phys. Lett. B* **526** (2002), 414-415], doi:[10.1016/S0370-2693\(01\)00743-2](https://doi.org/10.1016/S0370-2693(01)00743-2).
- [106] V. Cirigliano, G. Ecker and H. Neufeld, “Isospin violation and the magnetic moment of the muon,” *Phys. Lett. B* **513** (2001), 361-370, doi:[10.1016/S0370-2693\(01\)00764-X](https://doi.org/10.1016/S0370-2693(01)00764-X).
- [107] V. Cirigliano, G. Ecker and H. Neufeld, “Radiative tau decay and the magnetic moment of the muon,” *JHEP* **08** (2002), 002, doi:[10.1088/1126-6708/2002/08/002](https://doi.org/10.1088/1126-6708/2002/08/002).

- [108] M. Davier, S. Eidelman, A. Höcker and Z. Zhang, “Confronting spectral functions from e+ e- annihilation and tau decays: Consequences for the muon magnetic moment,” Eur. Phys. J. C **27** (2003), 497-521, doi:[10.1140/epjc/s2003-01136-2](https://doi.org/10.1140/epjc/s2003-01136-2).
- [109] M. Davier, S. Eidelman, A. Höcker and Z. Zhang, “Updated estimate of the muon magnetic moment using revised results from e+ e- annihilation,” Eur. Phys. J. C **31** (2003), 503-510, doi:[10.1140/epjc/s2003-01362-6](https://doi.org/10.1140/epjc/s2003-01362-6).
- [110] K. Maltman, “Constraints on the I=1 hadronic tau decay and e+e- \rightarrow hadrons data sets and implications for the muon (g-2),” Phys. Lett. B **633** (2006), 512-518, doi:[10.1016/j.physletb.2005.12.032](https://doi.org/10.1016/j.physletb.2005.12.032).
- [111] K. Maltman and C. E. Wolfe, “Isospin breaking in the relation between the tau- \rightarrow nu(tau) pi- pi0 and e+ e- \rightarrow pi+ pi- versions of $|F(\pi)(s)|^{**2}$ and implications for (g-2)(mu),” Phys. Rev. D **73** (2006), 013004, doi:[10.1103/PhysRevD.73.013004](https://doi.org/10.1103/PhysRevD.73.013004).
- [112] M. Davier, A. Hoecker, G. López Castro, B. Malaescu, X. H. Mo, G. Toledo Sánchez, P. Wang, C. Z. Yuan and Z. Zhang, “The Discrepancy Between tau and e+e- Spectral Functions Revisited and the Consequences for the Muon Magnetic Anomaly,” Eur. Phys. J. C **66** (2010), 127-136, doi:[10.1140/epjc/s10052-009-1219-4](https://doi.org/10.1140/epjc/s10052-009-1219-4).
- [113] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, “Reevaluation of the Hadronic Contributions to the Muon g-2 and to alpha(MZ),” Eur. Phys. J. C **71** (2011), 1515 [erratum: Eur. Phys. J. C **72** (2012), 1874], doi:[10.1140/epjc/s10052-012-1874-8](https://doi.org/10.1140/epjc/s10052-012-1874-8).
- [114] M. Benayoun, P. David, L. DelBuono and F. Jegerlehner, “Upgraded Breaking Of The HLS Model: A Full Solution to the $\tau^-e^+e^-$ and ϕ Decay Issues And Its Consequences On g-2 VMD Estimates,” Eur. Phys. J. C **72** (2012), 1848, doi:[10.1140/epjc/s10052-011-1848-2](https://doi.org/10.1140/epjc/s10052-011-1848-2).
- [115] M. Davier, A. Höcker, B. Malaescu, C. Z. Yuan and Z. Zhang, “Update of the ALEPH non-strange spectral functions from hadronic τ decays,” Eur. Phys. J. C **74** (2014) no.3, 2803, doi:[10.1140/epjc/s10052-014-2803-9](https://doi.org/10.1140/epjc/s10052-014-2803-9).
- [116] M. Bruno, T. Izubuchi, C. Lehner and A. Meyer, “On isospin breaking in τ decays for $(g-2)_\mu$ from Lattice QCD,” PoS **LATTICE2018** (2018), 135, doi:[10.22323/1.334.0135](https://doi.org/10.22323/1.334.0135).
- [117] S. Narison, “QCD parameters and SM-high precisions from e+e- \rightarrow Hadrons,” Nucl. Phys. A **1039** (2023), 122744, doi:[10.1016/j.nuclphysa.2023.122744](https://doi.org/10.1016/j.nuclphysa.2023.122744)
- [118] L. Esparza-Arellano, A. Rojas and G. Toledo, “Sizing the double pole resonant enhancement in $e^+e^- \rightarrow \pi^0\pi^0\gamma$ cross section and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau\gamma$ decay,” Phys. Rev. D **108** (2023) no.11, 116020, doi:[10.1103/PhysRevD.108.116020](https://doi.org/10.1103/PhysRevD.108.116020).
- [119] S. Weinberg, “Phenomenological Lagrangians,” Physica A **96** (1979) no.1-2, 327-340, doi:[10.1016/0378-4371\(79\)90223-1](https://doi.org/10.1016/0378-4371(79)90223-1).
- [120] J. Gasser and H. Leutwyler, “Chiral Perturbation Theory to One Loop,” Annals Phys. **158** (1984), 142, doi:[10.1016/0003-4916\(84\)90242-2](https://doi.org/10.1016/0003-4916(84)90242-2).
- [121] J. Gasser and H. Leutwyler, “Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark,” Nucl. Phys. B **250** (1985), 465-516, doi:[10.1016/0550-3213\(85\)90492-4](https://doi.org/10.1016/0550-3213(85)90492-4).
- [122] G. Ecker, J. Gasser, A. Pich and E. de Rafael, “The Role of Resonances in Chiral Perturbation Theory,” Nucl. Phys. B **321** (1989), 311-342, doi:[10.1016/0550-3213\(89\)90346-5](https://doi.org/10.1016/0550-3213(89)90346-5).

- [123] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, “Chiral Lagrangians for Massive Spin 1 Fields,” Phys. Lett. B **223** (1989), 425-432, doi:[10.1016/0370-2693\(89\)91627-4](https://doi.org/10.1016/0370-2693(89)91627-4).
- [124] V. Cirigliano, G. Ecker, M. Eidemuller, R. Kaiser, A. Pich and J. Portolés, “Towards a consistent estimate of the chiral low-energy constants,” Nucl. Phys. B **753** (2006), 139-177, doi:[10.1016/j.nuclphysb.2006.07.010](https://doi.org/10.1016/j.nuclphysb.2006.07.010).
- [125] K. Kampf and J. Novotny, “Resonance saturation in the odd-intrinsic parity sector of low-energy QCD,” Phys. Rev. D **84** (2011), 014036, doi:[10.1103/PhysRevD.84.014036](https://doi.org/10.1103/PhysRevD.84.014036).
- [126] P. Roig and J. J. Sanz Cillero, “Consistent high-energy constraints in the anomalous QCD sector,” Phys. Lett. B **733** (2014), 158-163, doi:[10.1016/j.physletb.2014.04.034](https://doi.org/10.1016/j.physletb.2014.04.034).
- [127] G. ’t Hooft, “A Planar Diagram Theory for Strong Interactions,” Nucl. Phys. B **72** (1974), 461 doi:[10.1016/0550-3213\(74\)90154-0](https://doi.org/10.1016/0550-3213(74)90154-0).