

Relations in Twisted Quantum K -Rings

Irit Huq-Kuruvilla

June 4, 2024

Abstract

We introduce twisted quantum K -rings, defined via twisted K -theoretic Gromov-Witten invariants. We develop a toolkit for computing relations by adapting some results about ordinary quantum K rings to our setting, and discuss some applications, including Ruan-Zhang's quantum K -theory with level structure, and complete intersections inside projective space.

In addition, we formulate a ring-theoretic abelian/non-abelian correspondence conjecture, relating the quantum K -ring of a GIT quotient $X//G$ to a certain twist of the quantum K -ring of $X//T$, the quotient by the maximal torus. We prove this conjecture for the case of Grassmanians, and use this to give another proof of the Whitney relations of Mihalcea-Gu-Sharpe-Zhou.

1 Introduction

The quantum K -ring (introduced by Givental and Lee) of smooth projective variety X is a deformation of its topological K -ring involving K -theoretic Gromov-Witten invariants of X . These invariants are defined as certain holomorphic Euler characteristics on Kontsevich's moduli space of stable maps $\overline{M}_{0,n,d}(X)$.

It is a K -theoretic counterpart to the quantum cohomology ring introduced by Witten and Kontsevich. Many recent developments in physics (including those of [15], [21]) have resulted in connections to quantum K -theory, and predictions for relations in quantum K -rings. However, quantum K -rings are in general much harder to compute than their cohomological counterparts, so many physical predictions go untested.

It was observed by Givental that one can calculate K -theoretic Gromov-Witten invariants of the quintic by doing intersection theory of $\overline{M}_{0,n,d}(\mathbb{P}^4)$, with the virtual fundamental class modified by the Euler class of an index bundle, corresponding to the restriction of $\mathcal{O}(5)$ to the source curve. This introduced the theory of twisted Gromov-Witten invariants, fully developed for cohomology theory in [3]. In general, twisted invariants can be computed from their untwisted counterparts. Other examples twisted invariants can be used to compute Gromov-Witten invariants of cotangent bundles, and to define an "abelian/non-abelian correspondence", relating Gromov-Witten invariants of a GIT quotient $X//G$ to those of the quotient $X//T$, where T is the maximal torus in G .

The K -theoretic counterpart of twisted invariants were introduced in [20], and generalizations were developed in [7] and [13]. In this paper, we will consider the most common type of twisting (type I), and develop a theory of quantum K -rings based on these invariants.

In full generality, given $V \in K^*(X)$ and a multiplicative K -theoretic characteristic class C , twisted K -theoretic Gromov-Witten invariants with twisting $C(V)$ are defined as holomorphic Euler characteristics on $\overline{M}_{0,n,d}(X)$ with respect to a twisted virtual structure sheaf:

$$\mathcal{O}^{tw} := \mathcal{O}^{vir} \times C(ft_*ev_{n+1}^*V)$$

As it turns out, type-I twisted invariants satisfy very similar properties to their untwisted counterparts, and as such, many results on quantum K -rings pass unchanged to the twisted case, in particular, we show the following:

Theorem. *Type I twisted invariants satisfy analogues of the Kontsevich-Manin axioms, and they are the structure constants to a ring, the (big or small) twisted quantum K ring of X .*

Furthermore we introduce the twisted J function $\mathcal{J}^{tw}(q, t)$, a certain kind of generating function for twisted invariants, and we prove the following theorem (which was proven by Iritani-Milanov-Tonita in [14] in the untwisted case):

Theorem. *If $K^*(X)$ is generated by line bundles whose 1st Chern classes are effective curve classes, then a polynomial difference/differential operator in q, t annihilating \mathcal{J}^{tw} yields a relation in the twisted quantum K -ring of X .*

Using the above theorem in general is difficult, as the relations are obtained in terms of operators $A_{i,com}$ which arise as solutions to a Lax-type equation. We give some conditions on which these operators can be simply interpreted in terms of multiplication by classes in $K^*(X)$, generalizing some results of [1]. The results we adapt to this setting form a robust toolkit for computing many examples of quantum K -rings.

To demonstrate the effectiveness of these results, we calculate relations in the (small) quantum K -rings for complete intersections in \mathbb{P}^N , as well as Ruan-Zhang's quantum K -theory with level structure, and we give a stable-map analogue of the quasimap quantum K -ring for $T^*\mathbb{P}^N$ introduced in [19].

We also introduce an abelian/non-abelian correspondence conjecture for small quantum K -rings. Our setting will be some reductive group G acting on a vector space X . The classical abelian/non-abelian correspondence proven by Martin constructs a ring map $sp : H^*(X//T)^W \rightarrow H^*(X//G)$, and identifies the Poincaré pairing on the target with the Poincaré pairing on the source modified by the Euler class of the vector bundle $V^M := \sum_{\alpha \in \text{roots}(G)} L_\alpha$. The analogous statement in K -theory was proven by Harada-Landweber [12].

Conjecture. The map $x \mapsto \lim_{\lambda \rightarrow 1} sp(x)$, extended to Novikov variables, is a surjective ring homomorphism from $QK^{tw}(X//T)$ to $QK^*(X//G)$, where the twisting is determined by $Eu_\lambda(V_M)$, where Eu_λ is the \mathbb{C}^* -equivariant K -theoretic Euler class.

We note that this is not a K -theoretic version of the correspondence for quantum cohomology rings conjectured by Ciocan-Fontanine–Kim–Sabbah in [2], that conjecture relates untwisted rings. There is a likely a K -theoretic counterpart to that conjecture, however the version we work with matches up well with the kind of formulas that appear in physics for relations in quantum K -rings, as we will see for the Grassmanian. We show this conjecture is a consequence of other abelian/non-abelian-type conjectures for J -functions, and as a result, using theorems from Givental-Yan in [8] on the J -function of the Grassmanian, we prove the conjecture for Grassmanians. Using this, we reprove the presentation for $QK^*(Gr(k, n))$ from [11]. Our arguments in principle extend to all partial flag varieties, where a presentation for the quantum K -ring is otherwise not known is not known, however due to the details involved, we save that case for a future work.

2 Twisted K -Theoretic Gromov-witten Invariants

K -theoretic Gromov-Witten invariants are defined as certain holomorphic Euler characteristics on the Kontsevich moduli space of stable maps $X_{g,n,d}(X)$ (this will be shortened to $X_{g,n,d}$ for the remainder of this text). Given Laurent polynomials $f_i(q)$ with coefficients in $K^*(X)$, the associated Gromov-Witten invariant is written in correlator notation as $\langle f_1, \dots, f_n \rangle_{g,n,d}$ and is defined to be:

$$\chi(X_{g,n,d}; \mathcal{O}^{vir} \otimes \prod_i ev_i^* f_i(L_i))$$

Here the notation $ev_i^* f_i(L_i)$ means to pull back the coefficients of f_i by ev_i , and evaluate the resulting polynomial at $q = L_i$.

A *twisting* is a modification of the virtual structure sheaf \mathcal{O}^{vir} of the form $C(ft_*\alpha)$, where C is some invertible multiplicative characteristic class in K -theory, expressible via Adams operations as:

$$C(E) = e^{\sum_{k \neq 0} \Psi^k(s_k E)},$$

ft is the forgetting map from the universal curve $X_{g,n+1,d}$, and α can be one of three types of elements:

- Type I: $\alpha = ev_{n+1}^* V$, where $V \in K^*(X)$ (introduced in [20]).
- Type II: $\alpha = ev_{n+1}^* f(L_{n+1})$, where f is a Laurent polynomial with coefficients in $K^*(X)$ satisfying $f(1) = 0$. (The other cases are handled by splitting into type II and type I components) (introduced in [7]).
- Type III: $\alpha = i_* f(L_+, L_-)$, where f is a symmetric bivariate Laurent polynomial, i is the inclusion of \mathcal{Z} , the codimension-2 locus of nodes in the universal family, and L_+, L_- are the cotangent lines to each branch at the node (introduced in [13]).

Twistings of different kinds can be taken in succession, and theorems relating twisted invariants of each type to their untwisted counterparts were proven in [7] and [13]. This paper will only be concerned with twistings of type I.

2.1 Examples of Twistings of Type I

The rest of this text will be concerned with twistings of type I. These encompass many important geometric situations. All of these twistings will be some variation of an Euler class, so we include some facts about K -theoretic Euler classes here:

The K -theoretic Euler class Eu is defined on line bundles L by $Eu(L) = 1 - L^{-1}$. We can write this as $e^{\sum_{k < 0} \Psi^k(L)/k}$, using the Taylor series for the logarithm. If we let \mathbb{C}^* act by scaling on L with equivariant parameter λ , this class becomes invertible. By multiplicativity and the splitting principle, the expression $e^{\sum_{k < 0} \Psi^k(V)/k}$ computes the Euler class for any bundle V .

2.1.1 Vector Bundles

Let V be a vector bundle over X . Since V is not compact, we cannot define Gromov-Witten invariants of the total space of V directly using the theory of stable maps.

However, we can take the following approach: Given the standard \mathbb{C}^* -action scaling the fibers of V , we work equivariantly. In the hypothetical moduli space $V_{g,n,d}$, the \mathbb{C}^* fixed locus is exactly $X_{g,n,d}$, corresponding to curves that land in X . Given such a map $f : C \rightarrow X$, the normal bundle would be f^*V . Thus the normal sheaf to $X_{g,n,d}$ is $ft_* ev_{n+1}^* V$. Thus equivariant holomorphic Euler characteristics on $V_{g,n,d}$ could be defined on $X_{g,n,d}$ via heuristically emulating the approach of fixed point localization, which involves dividing by the equivariant Euler class of the normal sheaf. So the stable map Gromov-Witten invariants of V are defined as the Gromov-Witten invariants of X , twisted by the class $Eu^{-1}(ev_* ft_* V)$.

2.1.2 Complete Intersections

If V was an ample vector bundle, its sections define a complete intersection $Y \in X$. If V is furthermore convex, $ft_* ev^*(V)$ is a bundle over $X_{0,n,d}$ whose zero locus is $Y_{0,n,d}$. So we can compute K -theoretic Gromov-Witten invariants of Y by inserting the Euler class of $ft_* ev^*(V)$ into X . To make it invertible, we use the equivariant Euler class as before.

2.1.3 Abelian/non-Abelian Correspondence

As stated in the introduction, our setting will be some reductive group G acting on a vector space X . The classical Abelian/non-Abelian correspondence proven by Martin relates $H^*(X//G)$ with $H^*(X//T)^W$, and uses the vector bundle $V^M := \sum_{\alpha \in \text{roots}(G)} L_\alpha$. The analogous statement in K -theory reads as follows:

Theorem 2.1 (Harada-Landweber, [12]). • $K^*(X//G) \cong \frac{K^*(X//T)^W}{\text{ann}(Eu(V^M))}$. We refer to this isomorphism by sp

- For $\alpha \in K^*(X//T)$, $\frac{1}{|W|}\chi(X//T; Eu(V^M)\alpha) = \chi(X//T; sp(\alpha))$.

Remark. This is an equivalent formulation to the one given in Harada-Landweber, but it doesn't correspond exactly to their main theorem. Our formulation is closer to Martin's, but in K -theory rather than cohomology. Their proof applies to equivariant contexts as well, so we can work in equivariant K -theory with respect to a torus action compatible with G

The quantum version of this correspondence relates the K -theoretic Gromov-Witten theory of $X//G$ with Gromov-Witten theory of $X//T$ twisted by $Eu_\lambda(V^M)$, the S^1 -equivariant Euler class of V^M . We will outline the precise statements in a later section of this text.

2.1.4 Level Structures

Quantum K -theory with level structure was introduced by Ruan and Zhang, and corresponds to twisting K -theoretic Gromov-Witten invariants twisted by the class $det(ft_* ev_{n+1}^* V)^{-\ell}$, here V is some vector bundle on X and ℓ is a positive integer known as the *Ruan-Zhang level*. While the characteristic class given by the determinant is invertible, it cannot be expressed convergently as $e^{\sum_k s_k \Psi^k(L)}$, so to include it into the framework of twisted invariants we use the following approach:

First observe that for V a vector bundle with nonvanishing Euler class, we have:

Proposition 2.2.

$$\frac{Eu(V)}{Eu(V^*)} = (-1)^{rk(E)} det(E)^{-1}$$

Proof. For V a line bundle, this is $\frac{1-L^{-1}}{1-L} = -L^{-1}$. Using the splitting principle, write $V = \bigoplus_{i=1}^{rk(V)} L_i$, which yields:

$$\frac{Eu(V)}{Eu(V^*)} = \prod_i \frac{1-L_i^{-1}}{1-L_i} = \prod_i -L_i^{-1} = -1^{rk(V)} det(V)^{-1}$$

□

So if we introduce two variables Y_1 and Y_2 , the class $C(V) := (-1)^{rk(\ell V)} \frac{\Lambda_{Y_1}(\ell V)}{\Lambda_{Y_2}(\ell V^*)}$ is an invertible multiplicative characteristic class whose limit as $Y_1, Y_2 \rightarrow 1$ becomes $-1^{\ell \times rk(V)} det(V)^{-\ell}$. Replacing V with $-V$ corrects for the sign issues, but in practice it is often useful to use the above formulation, and then make ad hoc corrections to account for the sign.

Remark. In fact, Ruan and Zhang show directly that the properties we use in the next section also hold for the determinant, with no extra variables necessary. So in the contexts that appear in this work, the failure of det to be expressible in standard form can actually be ignored.

3 Basic Properties of Type I Theories

Given that many of the above examples represent Gromov-Witten theories of different spaces, it is not particularly surprising that type I twisted K -theoretic Gromov-Witten invariants satisfy essentially the same set of properties as their untwisted counterparts. Before stating and proving these properties we introduce some definitions and notation.

Given a twisting $C(V)$, the classical K -ring is just $K^*(X)$, with Poincare pairing modified by $C(V)$. More precisely, given some basis ϕ_α for $K^*(X)$, the Poincare pairing is defined by

$$(\phi_\alpha, \phi_\beta) = \chi(X; \phi_\alpha \phi_\beta C(V))$$

. We denote this metric by g , with matrix coefficients $g_{\alpha\beta}$, and inverse matrix $g^{\alpha\beta}$.

Let F be the exponential generating function of the genus-0 K -theoretic Gromov-Witten invariants twisted by $C(V)$, i.e:

$$F := \sum_{n,d} \frac{Q^d}{n!} \langle t, \dots, t \rangle_{0,n,d}^{tw}$$

Here t stands for $\sum_{\alpha} t_{\alpha} \phi_{\alpha}$. Differentiation of F with respect to t_{α} corresponds to insertion of ϕ_{α} in one of the inputs, we denote $\frac{\partial}{\partial t_{\alpha}} F$ by F_{α} (and similarly for other functions of t).

The twisted quantum K -product will be based on the function

$$G := \frac{1}{2} g_{\alpha\beta} t_{\alpha} t_{\beta} + F$$

We first define a deformation of the Poincare pairing, the quantum pairing is defined by:

$$((\phi_{\alpha}, \phi_{\beta})) := G_{\alpha\beta}$$

$H_{g,n,d}(\alpha_1, \dots, \alpha_n)$ denotes the K -class $\prod_{i=1}^n ev_i^* \alpha_i(L_i) \otimes \mathcal{O}_{g,n,d}^{tw}$, i.e. the contents of a twisted K -theoretic Gromov-Witten invariant before the holomorphic Euler characteristic is taken.

For $V \in K^*(X)$, if we denote $V_{g,n,d}$ to be $ft_* ev_{n+1}^* V$, then the following are of the theory with twisting determined by V and a multiplicative characteristic class C (these results go back to Coates in [3]):

Theorem 3.1 ([3]).

- $ft^* \mathcal{O}_{g,n,d}^{tw} = \mathcal{O}_{g,n+1,d}^{tw}$
- The restriction of $\mathcal{O}_{g,n,d}^{tw}$ to the preimage of the gluing map $X_{g_1, n_1+1, d_1} \times_{\Delta} X_{g_2, n_2+1, d_2}$ is:

$$\frac{\mathcal{O}_{g_1, n_1+1, d_1}^{tw} \mathcal{O}_{g_2, n_2+1, d_2}^{tw}}{ev_{\Delta}^* C(V)} \rightarrow X_{g,n,d}$$

- The restriction of $\mathcal{O}_{g,n,d}^{tw}$ to the preimage of the gluing map $X_{g-1, n+1, d} \otimes_{X \times X} \Delta \rightarrow X_{g,n,d}$ is $\frac{\mathcal{O}_{g-1, n+2, d}^{tw}}{ev_{\Delta}^* (C(V))}$.

Property i) here implies the following:

Lemma 3.2. *If ft denotes the map forgetting the $n+1$ st marked point, then:*

$$ft^* \mathcal{O}_{g,n,d}^{tw} = \mathcal{O}_{g,n+1,d}^{tw}$$

These facts are sufficient to show that a type-I twisted theory satisfies analogues of the K -theoretic Kontsevich-Manin axioms, which we state below:

Theorem 3.3. *Twisted K -theoretic Gromov-Witten invariants satisfy the following set of axioms:*

- S_n covariance: For $\sigma \in S_n$ acting on $X_{g,n,d}$: $\sigma(H_{g,n,d}(\alpha_1, \dots, \alpha_n)) = H_{g,n,d}(\alpha_{\sigma(1)}, \dots, \alpha_{\sigma(n)})$.
- Fundamental class: For nondescendant inputs α_i , $\langle \alpha_1, \dots, \alpha_n, 1 \rangle_{g, n+1, d}^{tw} = \langle \alpha_1, \dots, \alpha_n \rangle_{g, n, d}^{tw}$
- Mapping to a point: $\mathcal{O}_{g,n,0}^{tw, vir} \cong \mathcal{O}_{\overline{M}_{g,n}} \boxtimes C(V) \Lambda_{-1}(T_X)$
- Splitting: Fix g_1, g_2 and n_1, n_2 such that $g = g_1 + g_2, n = n_1 + n_2$. Let

$$\Phi: \bigcup_{d_1+d_2=d} X_{g_1, n_1+1, d_1} \times X_{g_2, n_2+1, d_2} \rightarrow X_{g,n,d}$$

be the contraction map which glues the last marked point of $\overline{M}_{g_1, n_1+1}$ to the first marked point of $\overline{M}_{g_2, n_2+1}$. Then

$$\begin{aligned} & \Phi^* \sum_{k,d} Q^d \frac{1}{k!} ft_*^k H_{g, n+k, d}^X(\gamma_1, \dots, \gamma_n, t, \dots, t) \\ &= \sum_{\alpha, \beta} \left(\sum_{k_1, d_1} Q^{d_1} \frac{1}{k_1!} ft_*^{k_1} H_{g_1, n_1+k_1+1, d_1}^X(\gamma_1, \dots, \gamma_{n_1}, t, \dots, t, \phi_\alpha) \right) \\ & \quad G^{\alpha\beta}(t) \left(\sum_{k_2, d_2} Q^{d_2} \frac{1}{k_2!} ft_*^{k_2} H_{g_2, n_2+k_2+1, d_2}^X(\phi_\beta, \gamma_{n_1+1}, \dots, \gamma_n, t, \dots, t) \right) \end{aligned}$$

where the notation ft^k stands for the forgetful map which forget the additional k marked points.

- Genus reduction: Let

$$\Phi : X_{g-1, n+2, d} \rightarrow X_{g, n, d}$$

be the contraction map which glues the last two marked points. Then

$$\begin{aligned} & \Phi^* \sum_{k,d} Q^d \frac{1}{k!} ft_*^k H_{g, n+k, d}^X(\gamma_1, \dots, \gamma_n, t, \dots, t) \\ &= \sum_{\alpha, \beta} \left(\sum_{k,d} Q^d \frac{1}{k!} ft_*^k H_{g+1, n+k+2, d}^X(\gamma_1, \dots, \gamma_n, t, \dots, t, \phi_\alpha, \phi_\beta) \right) G^{\alpha\beta}(t) \end{aligned}$$

The proofs of the first three axioms are identical to those for the untwisted theory in [16]. We prove the latter two in the following section.

Remark. Our statements use $H_{g, n, d}$, classes in $K(X_{g, n, d})$, the axioms in [16] use $I_{g, n, d}$, which are the pushforwards to $\overline{M}_{g, n}$ of $H_{g, n, d}$, all proofs are equivalent once translated.

3.1 Splitting and Genus Reduction Axioms

We first recall the statement and proof for the splitting axiom for untwisted theories, and then cover the necessary changes for the twisted version.

The preimage of the divisor D is the locus in $X_{g, n+k, d}$ consisting of two curves of genus g_1 and g_2 , each containing n_1, n_2 marked points respectively, with $n_1 + n_2 = n$. The curves are connected by a chain of m \mathbb{P}^1 s. The k additional marked points are distributed on each irreducible component, such that there is at least one marked point in each curve in the chain. The degrees are chosen to add up to d . Call $m+1$ the depth of such a curve. Let $M_{s, k_\alpha, d_\alpha}$ be the space of curves of depth at least s , with a fixed distribution of degrees and marked points.

The map $\Phi : \bigcup M_{1, k_\alpha, d_\alpha} \rightarrow D$ gives D the structure of a divisor with normal crossings, so the structure sheaf of D can be obtained from the structure sheaf of each $M_{1, \dots}$ via the inclusion-exclusion principle. However, the intersection of any two depth-1 strata is a unique depth-2 stratum, etc. So if \mathcal{O}_m denotes the virtual structure sheaf of the union of the depth m strata, we have: The intersection of the loci of depth 1 is the union of the loci of depth 2, etc. So we can calculate the structure sheaf \mathcal{O}_D in terms of these by the inclusion-exclusion principle. This is a (virtual) normal crossing divisor, so if we call D_m the set of curve configurations of depth m , we have:

$$\mathcal{O}_D = \sum_m (-1)^m \sum_{C \in D_m} \mathcal{O}_C$$

Since the map ft is (virtually) traverse to the inclusion of D , we can calculate $ft_*^k I(\dots)$ by first restricting to $\bigcup_{C \in D_1} C$, and then pushing forward.

This means for untwisted invariants, we obtain:

$$\begin{aligned} \Phi^* ft_*^k H_{g,n+k,d}^X(\gamma_1, \dots, \gamma_n, t \dots, t) = \\ \sum_{\alpha, \beta, k_1+k_2=k} I_{g_1, n_1+k_1+1, d_1}(\gamma_1, \dots, \gamma_n, t \dots, \phi_\alpha) g^{\alpha\beta} H_{0, k_2+2, d_2}(\phi_\beta, t, \dots, \phi_k) g^{kl} H_{g_2, n_2+k_2+1, d_2} - \\ \sum_{\alpha, \beta, \nu, \mu, d_i, k_i} H_{g_1, n_1+k_1+1, d_1}(\gamma_1, \dots, \gamma_n, t \dots, \phi_\alpha) g^{\alpha\beta} H_{0, k_2+2, d_2}(\phi_\beta, t, \dots, \phi_\nu) g^{\nu\mu} H_{g_2, n_2+k_2+1, d_2}(\phi_\mu \gamma_1, \dots, \gamma_n, t \dots) \\ + \dots \end{aligned} \tag{3.1}$$

A shorthand way of writing this is using the quantum pairing. Since the matrix inverse of $1 + F$ is $1 - F + F^2 + \dots$, since $G_{\alpha\beta} = g_{\alpha\beta} + F_{\alpha\beta}$ be the matrix of the quantum pairing, we can sum the above expression over k to yield:

$$\langle t, \dots, t \rangle_{0,n,d}^\Phi = \langle t, \dots, \phi^\alpha \rangle_{n_1+1, g_1, d_1} G^{\alpha\beta} \langle \phi^\beta, t, \dots \rangle_{n_2+1, g_2, d_2} \tag{3.2}$$

The twisted version works essentially the same way. Using the restriction theorem for twisting class cited earlier, the same formula is true, replacing all correlators by their twisted counterparts. The difference is that, by Theorem 3.1, there is a factor of $\frac{1}{C(V)}$ for each node in the stratum. We can remove this factor by using the twisted Poincare pairing rather than the untwisted one, since the extra factor of $C(V)$ is cancelled by the $\frac{1}{C(V)}$ appearing at each node, yielding the desired result of:

$$\langle t, \dots, t \rangle_{0,n,d}^{tw, \Phi} = \langle t, \dots, \phi^\alpha \rangle_{n_1+1, g_1, d_1}^{tw} G^{\alpha\beta, tw} \langle \phi^\beta, t, \dots \rangle_{n_2+1, g_2, d_2}^{tw}$$

The proof of the genus reduction axiom is identical.

3.2 String, Dilaton, and WDVV Equations

We also have the following additional properties:

Theorem 3.4.

- *String Equation:* If \mathcal{H} denotes the Hodge bundle, then (for $g > 1$):

$$ft_* \mathcal{O}_{g,n,d}^{tw} \left(\frac{1}{1-qH} \prod_i \frac{1}{1-q_i L_i} \right) = * \mathcal{O}_{g,n,d}^{tw} \left(\frac{1}{1-qH} (1 + H^* + \sum_i \frac{q_i}{1-q_i} \prod_i \frac{1}{1-q_i L_i} \right)$$

For $g = 0$. the same equation is true with H^* set to 0, which leads to the following, perhaps more familiar, expression in terms of correlators:

$$\langle a_1(L), \dots, a_n(L), 1 \rangle_{0, n+1, d}^{tw} = \sum_i \langle a_1(L), \dots, \frac{a_i(L) - a_i(1)}{L-1}, a_n(L) \rangle_{0, n, d}^{tw}$$

- *Dilaton Equation:* With the same setup as for the string equation:

$$ft_* \mathcal{O}_{g,n,d}^{tw} \left(\frac{1}{1-qH} \prod_i \frac{1}{1-q_i L_i} \right) L_{n+1} = * \mathcal{O}_{g,n,d}^{tw} \left(\frac{1}{1-qH} (1 + H + \sum_i \frac{q_i}{1-q_i} \prod_i \frac{1}{1-q_i L_i} \right)$$

As consequence $ft_*(1 + L_{n+1}) = H + H^* + n - 2$.

In genus 0, this yields the following relation among correlators:

$$\langle a_1, \dots, a_n, L_{n+1} - 1 \rangle_{g, n+1, d} = (n-2) \langle a_1, \dots, a_n \rangle_{0, n, d}$$

Both of these proofs are identical to the ones in [16].

Theorem 3.5 (WDVV Identity for K -Theory).

$$\sum_{d_1, d_2, n_1, n_2, \alpha, \beta} \langle A, B, t, \dots, \phi_\alpha \rangle_{0, n_1+3, d_1} g^{\alpha\beta} \langle C, D, t, \dots, \phi_\beta \rangle_{0, n_2+3, d_2} -$$

$$\sum \langle A, B, t, \dots, \phi_\alpha \rangle_{0, n_1+3, d_1} g^{\alpha\beta} \langle \phi_\beta, t, \dots, \phi_\gamma \rangle_{0, n_2+2, d+2} g^{\gamma\delta} \langle C, D, t, \dots, \phi_\delta \rangle_{0, n_3+3, d_3} + \sum \dots$$

is symmetric in A, B, C, D .

There is a contraction map $ct : X_{0, n+4, d} \rightarrow \overline{M}_{0,4} \cong \mathbb{P}^1$, which forgets the map to X , all marked points but the first 4, and stabilizes the resulting curve by contracting unstable components. The points $0, 1, \infty \in \mathbb{P}^1$ are the boundary points of $\overline{M}_{0,4}$, parametrized by cross-ratio. They correspond to 2-component curves where the 4th marked point shares a component with the first, second, and third marked points respectively. Using the splitting axiom, the expression above is equivalent to restricting to a boundary point of $\overline{M}_{0,4}$, and calculating the resulting Gromov-Witten invariant. Permuting A, B, C, D chooses a different boundary point.

4 Twisted Quantum Products

For a compact Kahler X , the quantum K -ring of X is a deformation of the ordinary K -ring, using genus-0 K -theoretic Gromov-Witten invariants as structure constants. The same construction can be applied to (type-I) twisted invariants.

The quantum product is defined via structure constants, and is determined by the formula:

$$((\phi_\alpha * \phi_\beta, \phi_\gamma)) = G_{\alpha\beta\gamma}$$

Solving for the quantum product gives:

$$\phi_\alpha * \phi_\beta = \sum_{\gamma} G_{\alpha\beta\gamma} \sum_{\delta} G^{\delta\gamma} \phi_\delta$$

We now verify some basic facts about the quantum product:

Theorem 4.1. *The quantum product is associative.*

Proof. This is identical to the proof in the untwisted case, but we recount it for the sake of completeness.

We can write the coefficients $G^{\alpha\beta}$ explicitly in the following way. Call the matrices with coefficients $G_{\alpha\beta}, F_{\alpha\beta}$ \mathcal{G} and \mathcal{F} respectively:

We have: $\mathcal{G} = g + \mathcal{F}$, where g is the matrix of the twisted Poincare pairing.

Using the fact that g is invertible and $(1 + M)^{-1} = 1 - M + M^2 - \dots$, as in the proof of the splitting axiom, we have:

$$\mathcal{G}^{-1} = (1 - g^{-1}\mathcal{F} + g^{-1}\mathcal{F}g^{-1}\mathcal{F} + \dots)g^{-1}$$

So a particular matrix coefficient looks like:

$$G^{\alpha\beta} = g^{\alpha\beta} - \sum_{\delta, \gamma} g^{\alpha\gamma} F_{\gamma\delta} g^{\delta\beta} + \dots$$

The statement of associativity is equivalent to the symmetry in $\alpha, \beta, \gamma, \delta$ of:

$$\sum_{\mu, \nu} G_{\alpha\beta\mu} G^{\mu\nu} G_{\nu\gamma\delta}$$

Using the expression for $G^{\mu\nu}$ calculated previously, and the fact that third derivatives of G agrees with those of F . we can rewrite this as:

$$\sum_{\mu\nu} F_{\alpha\beta\mu} g^{\mu\nu} F_{\nu\gamma\delta} - F_{\alpha\beta\mu} \sum_{\mu,\nu,\zeta,\eta} g^{\mu\zeta} F_{\zeta\eta} g^{\eta\nu} F_{\nu\gamma\delta} + \dots$$

The symmetry of this expression is precisely the generating function version of the WDVV identity for twisted K -theoretic invariants. □

Theorem 4.2. *The quantum product has unit 1*

Proof. This is equivalent to the statement

$$G_{\alpha,\beta,1} = G_{\alpha,\beta}$$

The left hand side is equal to:

$$\sum_{n,d} \frac{1}{n!} Q^d \langle \phi_\alpha, \phi_\beta, 1, t, \dots, t \rangle_{0,n+3,d}^{tw}$$

The right hand side is:

$$g_{\alpha\beta} + \sum_{n,d} \frac{1}{n!} Q^d \langle \phi_\alpha, \phi_\beta, t, \dots, t \rangle_{0,n+2,d}^{tw}$$

Using the string equation, we can equate all but one term on the left hand side with a term on the right hand side by observing that:

$$\frac{1}{n!} Q^d \langle \phi_\alpha, \phi_\beta, 1, t, \dots, t \rangle_{0,n+3,d}^{tw} = \frac{1}{n!} Q^d \langle \phi_\alpha, \phi_\beta, t, \dots, t \rangle_{0,n+2,d}^{tw}$$

This handles all terms except $\langle \phi_\alpha, \phi_\beta, 1 \rangle_{0,3,0}$, since there is no forgetful map to apply here. However $X_{0,3,0}$ is just X , so this correlator is equal to $\langle \phi_\alpha, \phi_\beta \rangle = g_{\alpha\beta}$, identifying it with the constant term (in both t and Q) on the right hand side. □

5 q -difference Modules

Having established some basic properties of twisted Gromov-Witten invariants, we now illustrate the connection with q -difference equations, which will be the main tool we use to compute ring relations. Many of the results here have proofs that are completely identical to their untwisted counterparts, so we only include proofs that are particularly illustrative, or are different in the twisted case.

5.1 Quantum Connection

As in the untwisted case, and for the exact same reasons, quantum K -theory determines a Frobenius structure on $K(X)$

Theorem 5.1. *$K(X) \otimes \mathbb{C}$, equipped with the metric G and the quantum product $\partial_{ijk}G$, has the structure of a Frobenius manifold.*

In particular, the Dubrovin connection:

$$\nabla_q := d - \frac{1}{1-q} (e_i^*) dt$$

is flat

Proof. This is completely a formal consequence of the WDVV and string equations, see [16] for details. \square

Theorem 5.2. *The twisted quantum differential equation:*

$$\nabla_q S = 0$$

has fundamental solution matrix $S \in \text{End}(K(X))[q, Q]$ given by:

$$S_{ij} = g_{ij} + \sum_{n,d} \frac{Q^d}{n!} \langle \phi_i, t, \dots, t, \frac{\phi_j}{1 - qL} \rangle_{0,n,d}^{tw}$$

Proof. This is also a completely formal consequence of WDVV and string equations, just refer to [16]. \square

Theorem 5.3. *Regarding S as an endomorphism valued function in $\text{End}(K^*(X))(q)[[Q, t]]$, it is invertible, and S^{-1} is the adjoint of S with respect to the twisted pairing, and is a fundamental solution to:*

$$(1 - q)\partial_\alpha T = T(\phi_\alpha * Q)$$

Remark. By Kawasaki's Riemann-Roch Theorem, each entry of S is a rational function of q with poles at roots of unity and vanishes at ∞ .

Definition 5.4. The twisted big J -function, denoted $\mathcal{J}^{tw}(q, t)$ is defined by:

$$\mathcal{J}^{tw} = (1 - q)S^{-1}\mathbf{1}$$

Actually expanding this yields:

$$\mathcal{J}^{tw} = (1 - q) + t + \sum_d \sum_n \frac{Q^d}{n!} \langle t, \dots, t, \frac{\phi^\alpha}{1 - qL} \rangle_{0,n+1,d} \phi_\alpha$$

The value at $t = 0$, denoted J^{tw} is the twisted small J -function.

5.2 The J -function and Symplectic Loop Space

The proper setting to think about generating functions for quantum K -theoretic invariants is within the context of Givental's symplectic loop space \mathcal{K} . We give a brief overview of the theory below:

The symplectic loop space \mathcal{K} is a $K^*(X)[Q]$ -module consisting of (Q -adic completions of) rational functions in q with coefficients in $K^*(X)$, with poles allowed only at $0, \infty$, or roots of unity. It has a symplectic form given by:

$$\Omega(f, g) = -\text{Res}_{0, \infty}(f(q), g(q^{-1})) \frac{dq}{q}$$

Here Res_S denotes the sum of the residues at values in the set S , and $(,)$ denotes the K -theoretic Poincaré pairing.

\mathcal{K} is also equipped with a polarization into two Lagrangian subspaces:

- $\mathcal{K}^+ :=$ Laurent polynomials in q
- $\mathcal{K}^- := \{f | f(\infty) = 0, f(0) \neq \infty\}$

As a consequence of the Kawasaki-Riemann-Roch theorems for stacks the $Q^{d>0}$ terms of the big J -function lie in \mathcal{K}^- , its projection to \mathcal{K}^+ is $1 - q + t$.

Theorem 5.5 (Givental-Tonita). *The range of the big J -function is an overruled Lagrangian cone (denoted \mathcal{L}) in \mathcal{K} . This means it is a cone that has the following properties:*

- Tangent spaces are invariant with respect to multiplication by $1 - q$
- For $T = T_p \mathcal{L}$, $(1 - q)T \subset \mathcal{L}$
- The tangent spaces for all $p \in (1 - q)T$ are equal to T .

5.3 Relations and Shift Operators

We now make the assumption that we have a basis p_i of $H_2(X)$ corresponding to the Novikov variables Q_i , and there exist a set of line bundles P_i with $c_1(P_i) = -p_i$.

\mathcal{L} has the structure of a q -difference module in the following way:

Theorem 5.6 (Givental-Tonita). *Let p_i be some basis for $H^2(X)$ consisting of first Chern classes of line bundles P_i . \mathcal{L} and each of its tangent spaces is invariant under the algebra of finite difference operators in Q_i generated by P_i, Q^d (for d in the Mori cone of X), where P_i acts as $P_i q^{Q_i \partial_{Q_i}}$*

Givental showed in [7] that essentially the same things holds in the context of twisted invariants of type I, the graph of \mathcal{J}^{tw} is a Lagrangian cone in the space \mathcal{K}^{tw} defined the same way as \mathcal{K} but using the twisted Poincare pairing, and the ruling spaces have the same difference module structure.

Theorem 5.7. $S^{-1}(K^*(X)[q^\pm, Q, t])$ is a tangent space to the twisted Lagrangian cone \mathcal{L}^{tw}

This is a twisted version of a corresponding theorem of Givental-Tonita and the proof is the same, it is a direct consequence of the string and dilaton equations.

By the invariance property, $S^{-1}(K^*(X)[q^\pm, Q, t])$ is thus invariant under action of $P_i q^{\partial_{Q_i}}$.

This means the operator $A_i := S^{-1} P_i q^{\partial_{Q_i}} S$ is a polynomial in q , and lies in $End(K(X))[q, Q, t]$.

We define the q -shift operator \mathcal{A}_i as $A_i q^{\partial_{Q_i}}$, it satisfies the following properties (again the proofs are the same as the proofs of the untwisted versions):

- $S \circ P_i q^{\partial_{Q_i}} = \mathcal{A}_i \circ S$
- $P_i q^{\partial_{Q_i}} \circ T = T \circ \mathcal{A}_i$
- $[\nabla_{q,\alpha}, \nabla_{q,\beta}] = [\mathcal{A}_i, \mathcal{A}_j] = [\nabla_{q,\alpha}, A_i] = 0$
- Given $s_1, s_2 \in \mathcal{K}^+$:
- $\partial_\alpha G(\overline{s_1}, s_2) = G(\frac{\nabla_{q,\alpha}}{1-q} s_1, s_2) + G(\overline{s_1}, \text{frac} \nabla_{q,\alpha} 1 - q s_2)$
- $q^{Q_i \partial_{Q_i}} G(\overline{s_1}, s_2) = G(\overline{\mathcal{A}_i^{-1} s_1}, \mathcal{A}_i s_2)$

Proofs: Entirely formal consequences of string and divisor equations and the quantum differential equation for S . As a result, we have the following theorem:

Theorem 5.8. • $(1-q)\partial_\alpha A_i = A_i(q^{Q_i \partial_{Q_i}} \phi_\alpha * Q) - (\phi_\alpha * Q) A_i$

- $G(\phi_\alpha, A_i \phi_\beta) = q^{Q_i \partial_{Q_i}} G(\overline{A_i} \phi_\alpha, \phi_\beta)$
- Let $A_{i,com}$ denote $A_i|_{q=1}$, then $A_{i,com}$ commutes with quantum multiplication.
- $A_i = P_i * Q + \sum_{d|d_i > 0} \sum_{d=0}^{d_i-1} Q^d A_{i,d,k}(t) (1-q)^k Q^d$ Where $A_{i,d,k}(t)$ does not depend on t_0 . All expansions here are in $End(K(X)) \otimes Q[[q^\pm]][[Q, t]]$.

It is sometimes more convenient to work not with S and \mathcal{J}^{tw} but with their logarithmic-modified counterparts \tilde{S} and $\tilde{\mathcal{J}}^{tw}$, obtained by replacing S with $S(\prod_i P_i^{\frac{\ln(Q_i)}{\ln(q)}})$. (This results in left multiplying J by $\prod_i P_i^{-\frac{\ln(Q_i)}{\ln(q)}}$). This expression is defined by rewriting P_i as $(1 - (1 - P_i))$ and using the binomial expansion. The result converges as a power series in Q_i since $(1 - P_i)$ is nilpotent.

The purpose of this is to remove the factors of P_i from theorems about \mathcal{A}_i , due to the observation:

$$q^{Q_i \partial_{Q_i}} \left(\prod_i P_i^{\frac{\ln(Q_i)}{\ln(q)}} \right) = P_i \left(\prod_i P_i^{\frac{\ln(Q_i)}{\ln(q)}} \right)$$

Applying this to the properties written earlier yields:

Theorem 5.9.

- $\tilde{S} \circ q^{Q_i \partial_{Q_i}} = \mathcal{A}_i \circ \tilde{S}$
- $q^{Q_i \partial_{Q_i}} \tilde{S}^{-1} = \tilde{S}^{-1} \circ \mathcal{A}_i$

Combining 5.2 and 5.3 yield a twisted version of Iritani-Milanov-Tonita's theorem on relations in big quantum K -theory:

Theorem 5.10. *Given some polynomial difference/differential operator $D(q, Q, q^{Q_i \partial_{Q_i}}, \partial_{t_\alpha})$, we have:*

$D\tilde{\mathcal{J}} = 0$ implies that the following relation holds in the big quantum K -ring:

$$D(1, Q, A_{i,com}, \phi_\alpha * Q) = 0 \in QK_{big}^*(X)$$

Proof. We apply D to $\tilde{J}^{tw} = (1 - q)\tilde{S}^{-1}\mathbf{1}$:

If $D = (1 - q)\partial_\alpha$, then $D\tilde{J}^{tw} = (1 - q)\tilde{S}^{-1}(\phi_\alpha)$.

If $D = q^{Q_i \partial_{Q_i}}$, then $D\tilde{J}^{tw} = (1 - q)\tilde{S}^{-1}(\mathcal{A}_i(\mathbf{1}))$.

Since the two kinds of operators commute, for a general D if $D\tilde{J}^{tw} = 0$, then $\tilde{S}^{-1}D(q, Q, \mathcal{A}_i, \phi_\alpha * Q)\mathbf{1} = 0$

Multiplying both sides by \tilde{S} and setting $q = 1$ gives the desired result. □

The same theorem holds for the small J -function.

Theorem 5.11. *Given some polynomial difference operator $D(q, Q, q^{Q_i \partial_{Q_i}})$, we have:*

$DJ^{tw} = 0$ implies that the following relation holds in the small quantum K -ring:

$$D(1, Q, A_{i,com}|_{t=0}) = 0 \in QK^*(X)$$

If we replace J^{tw} with $\mathcal{J}^{tw}|_{t=t_0}$, the same theorem holds for twisted quantum K -theory with bulk shift t_0 .

Proof. We note that this is not strictly a corollary of the previous theorem but rather a consequence of its proof. We can set $t = 0$ (or some other value) in each line of the proof and it remains true. (We remark that $\tilde{S}_{t=0}$ is invertible since we can specialize $\tilde{S}\tilde{S}^{-1} = I$ to $t = 0$, and the RHS does not depend on t). □

5.4 Calculating the Shift Operators

In practice, applying the theorem above only yields useful relations when the \mathcal{A}_i s (or their limits A_i) can be calculated directly. As a consequence of the theorems of Iritani, Milanov, and Tonita (which all generalize to this setting), there is a recursive procedure for calculating them, given the small J -function, if the K -ring is generated by divisors.

However, the following approach, used in the case of G/P by Anderson-Chen-Tseng-Iritani[1] is often more practical:

Theorem 5.12 (Quantum Triviality Theorem). *Given some polynomial $f(P_i)$, if $\frac{1}{1-q}J^{tw}$ has the property that all terms of positive Q -degree in $f(q^{Q_i \partial_{Q_i}})J$ vanish at $q = \infty$, then $f(A_{i,com})|_{t=0}$ acts as quantum multiplication by $f(P_i)$ in the small twisted quantum K -ring of X .*

Proof. We make the simplification that $f = P_i$ here to save notation, but the proof is essentially identical in the case of nontrivial f .

$$P_i q^{Q_i \partial_{Q_i}} T(1_X) = T(\mathcal{A}_i q^{Q_i \partial_{Q_i}}(1_X))$$

The left hand side is $P_i q^{Q_i \partial_{Q_i}} \frac{1}{1-q} \mathcal{J}_X^{tw}$. We first restrict to $t = 0$, then expand at $q = \infty$, since $T = I +$ terms vanishing at $q = \infty$, the expansion becomes:

$$P_i q^{Q_i \partial_{Q_i}} \frac{1}{1-q} \mathcal{J}_X^{tw} = c_0 + \sum_d c_d Q^d + \dots$$

For some terms c_d given by the expansion of A_i at $q = 1, t = 0$, with c_0 being the operator of (small) quantum multiplication by P_i .

Clearly, if the left-hand side has no $Q^{>0}$ terms that do not vanish at infinity, then neither does the right hand side, hence $c_d = 0$ for all $d \neq 0$. □

What this means is that provided we know the small J -function for some theory, provided the q -degrees satisfy the above conditions, we can convert difference operators annihilating the J -function to relations without having to solve the quantum Lax equation.

In practice, finding these operators is often not difficult. However, if we wish to compute a presentation for a (twisted) quantum K -ring, we need some way of telling when we have found enough relations.

Proposition 5.13. *If we know a set I of relations holds in $QK^{tw}(X)$, and $I|_{Q=0}$ form a complete set of relations for $K^*(X)$, then I form a complete set of relations for $QK^{tw}(X)$.*

Proof. Since $WK^{tw}(X)$ is complete with respect to Q , this follows from a general result in commutative algebra, Proposition A.3 in [11]. □

We note that we should only expect to get a full set of relations when P_i generate $K^*(X)$, as q -difference operators can only give us relations in terms of functions of P_i .

In addition, finding ring relations using these methods requires knowing the small J -function, the next section gives some details on cases where it can be calculated.

5.5 S_n -invariant Theory

In a series of papers, Givental introduced a generalization of K -theoretic Gromov-Witten invariants that involve the S_n -action on the marked points. Generating functions for such invariants generally behave better with respect to twisting and fixed-point localization.

Theorems about cohomological Gromov-Witten invariants often find natural generalizations in this setting, rather than in the "ordinary" quantum K -theory we have considered so far.

The main way this affects the current topic of this paper is that, for many spaces, it is difficult to find values for the (twisted or untwisted) big J -function. However, it is much easier if we instead use the symmetrized big J function, denoted \mathcal{J}^{sym} , defined in the following way:

$$\mathcal{J}^{sym} = (1-q) + t + \sum_d \sum_n \frac{Q^d}{n!} \langle t, \dots, t, \frac{\phi^\alpha}{1-qL} \rangle_{0,n+1,d}^{sym} \phi_\alpha$$

The correlator $\langle \rangle^{sym}$ means we take pushforwards from the spaces $X_{0,n+1,d}/S_n$. These invariants are a special case of the *permutation-equivariant K -theoretic Gromov-Witten invariants* introduced in [5]. The important thing for us is that at $t = 0$, the S_n -action is irrelevant and we recover the usual small J -function. Twistings are defined for this theory in exactly the same way.

We do not consider any ring structure involving the S_n -invariant correlators in this paper.

For the examples of twistings we have covered, we can construct values of \mathcal{J}^{sym} :

Theorem 5.14. *Let V be a vector bundle that splits into a sum of r line bundles V_i . Let $J := \sum_d Q_d J_d$. If $(1-q)J$ is a value of \mathcal{J}^{sym} , then:*

- $(1 - q) \sum_d Q^d J_d \prod_{i=1}^r \frac{\prod_{\ell \leq (c_1(V_i), d)} (1 - V_i^{-1} q^\ell)}{\prod_{\ell < 0} (1 - \lambda V_i^{-1} q^\ell)}$ is a value of the symmetrized big J -function of the theory twisted by $Eu_\lambda(V)$.
- $(1 - q) \sum_d Q^d J_d \prod_{i=1}^r \frac{\prod_{\ell < (c_1(V_i), d)} (1 - \lambda V_i q^\ell)}{\prod_{\ell < 0} (1 - \lambda V_i^{-1} q^\ell)}$ is a value of the symmetrized big J -function of the theory twisted by $\frac{1}{Eu_\lambda(V^*)}$.
- $(1 - q) \sum_d Q^d J_d \prod_i (V_i^{-(d_i, c_1(V_i))} q^{(d_i, c_1(V_i) + 1)})^\ell$ is a value of the symmetrized big J -function of the theory with level structure given by V, ℓ .

If these values turn out to be the small J -function, which can be tested by verifying that their projection to K_+ is equal to $1 - q$, then we can calculate relations in the associated twisted quantum K -rings via finding appropriate q -difference operators.

Remark. There is a ring structure involving symmetrized invariants (or more general S_n -equivariant ones), but we do not discuss it in this work.

6 Reconstruction for Twisted Theories

Another fact about genus-0 twisted quantum K -theoretic invariants is that, in nice situations, they are completely determined by the invariants for $n = 1$. In particular, if $K^*(X)$ is generated by line bundles. This can be seen at the level of Lagrangian cones from twisted versions of results in [14] and [6] (the proofs are identical to the ones in the untwisted case). However, for use in the following section, we will use the result in terms of correlators due to Lee-Pandharipande [17].

Lemma 6.1 (Lee-Pandharipande). *The following relations hold in $Pic(X_{0,n,d})$, for ϕ a line bundle on X :*

$$ev_i^*(\phi) = ev_j^*(\phi) + \langle d, \phi \rangle L_j - \sum_{d_1 + d_2 = d} \langle d_1, \phi \rangle D_{i, d_1 | j, d_2} \quad (6.1)$$

$$L_1 + L_2 = D_{i | j} \quad (6.2)$$

Noting that a relationship between divisors and line bundles in the Picard group is equivalent to the same relation between their structure sheaves and Euler classes in K -theory, the lemma has the following K -theoretic interpretation:

Lemma 6.2. *The following relations hold in $K^*(X_{0,n,d})$, for ϕ a line bundle on X :*

$$(1 - ev_i^*(\phi)^{-1}) \mathcal{O}_{0,n,d}^{vir} = (1 - ev_j^*(\phi)^{-1} L_j^{-\langle d, \phi \rangle}) \mathcal{O}_{0,n,d}^{vir} - \sum_{d_1 + d_2 = d} \langle d_1, \phi \rangle \mathcal{O}_{D_{i, d_1 | j, d_2}}^{vir} \quad (6.3)$$

$$1 - L_i^{-1} L_j^{-1} = \mathcal{O}_{D_{i | j}}^{vir} \quad (6.4)$$

By applying these relations, coupled with the splitting axiom of the virtual structure sheaf, any n -pointed degree d invariant whose inputs are written in terms of line bundles on X can be expressed in terms of invariants of lower degree and/or fewer marked points. For details on how this reconstruction looks concretely, see [17]

Lemma 6.3. *Given an arbitrary twisting of type 1, denote the modified virtual structure sheaf by $\mathcal{O}^{vir, tw}$, then the following relations hold in $K^*(X_{0,n,d})$, for ϕ a line bundle on X :*

$$(1 - ev_i^*(\phi)) \mathcal{O}_{0,n,d}^{vir, tw} = (1 - ev_j^*(\phi)^{-1} L_j^{-\langle d, \phi \rangle}) \mathcal{O}_{0,n,d}^{vir, tw} - \sum_{d_1 + d_2 = d} ev_{node}^* C(V) \langle d_1, \phi \rangle \mathcal{O}_{D_{i, d_1 | j, d_2}}^{vir, tw} \quad (6.5)$$

$$(1 - L_i^{-1} L_j^{-1}) \mathcal{O}_{0,n,d}^{vir, tw} = ev_{node}^* C(V) \mathcal{O}_{D_{i | j}}^{vir, tw} \quad (6.6)$$

Both of these lemmas follow from their untwisted counterparts paired with the restriction formula for twisting classes.

This allows us to prove the following theorem:

Theorem 6.4. *By applying the lemma sequentially, all twisted Gromov-Witten invariants can be reconstructed from the one-pointed descendant invariants.*

Proof: Choose a generating set ϕ_i of line bundles. We simultaneously induct on d , the ϕ_i -degree of an input, and the absolute value of the L_i -degree.

For our base cases, when the ϕ_i and L_i degrees are 0, the input is 1, and we can lower the number of marked points by virtue of the string equation.

If the i th input is a monomial, it is $P(\phi_i)L_i^a$ for ϕ_i some line bundle, we can use the first relation to write the correlator in terms of one with a lower the ϕ_i -degree of P , and ones with lower degree and number of marked points.

If we have two different inputs with descendants with negative degrees, we can use the second relation to remove one L_i^{-1}, L_j^{-1} from each of them, and calculate the invariant via the splitting axiom.

For an input with positive L_i degree, we can use a rearranged version of the first relation to lower the degree.

Remark. This procedure is often not practical to actually carry out in practice.

7 Quantum Abelian-Non/Abelian Correspondence

Our setting is the same as in the section discussing the classical abelian/non-Abelian correspondence. Let X, Y be $X//G$ and $X//T$ respectively. The correspondence relates invariants of X to twisted invariants of Y , with twisting given by $Eu_\lambda(V)$, where V is Martin's bundle.

Let the small J -function of the twisted theory of Y be denoted J_Y^{tw} , let the corresponding Lagrangian cone be denoted \mathcal{L}_Y^{tw} . We introduce the following morphisms:

- Let sp denote the usual specialization map $K^*(Y)^W \rightarrow K^*(X)$
- Let ν_Q denote the specialization map on Novikov variables.
- Let sp_Q denote ${}^o nu_Q$, which maps $K^*(Y)[[Q]]$ to $K^*(X)[[Q]]$.

We will give 4 different conjectures that go under the name ‘‘abelian/non-abelian correspondence’’:

Conjecture 7.1. sp_Q maps a formal germ $N \subset \mathcal{L}_Y^{tw}$ to \mathcal{L}_X .

Conjecture 7.2. sp_Q maps a formal germ $N \subset \mathcal{L}_Y^{tw, sym}$ to \mathcal{L}_X^{sym} .

Conjecture 7.3. $\lim_{\lambda \rightarrow 1} sp_Q(J_Y^{tw} = J_X$

Conjecture 7.4. $\langle a_1, \dots, a_n \rangle_{g, n, d}^X = \lim_{\lambda \rightarrow 1} \sum_{d' \rightarrow d} \langle \tilde{a}_1, \tilde{a}_2, \dots \rangle_{g, n, d'}^{tw, Y}$

Conjecture 7.5. The map $\phi : x \mapsto \lim_{y \rightarrow 1} sp_Q(x)$ is a surjective ring homomorphism from $QK^{tw}(Y)^W$ to $QK^*(X)$.

We will give some relationships between the conjectures below:

Theorem 7.6. *If $(0, J_Y^{tw}) \in N$, then Conjecture 7.1 or Conjecture 7.2 implies Conjecture 7.3*

Proof. Obvious □

Theorem 7.7. *Conjecture 7.4 implies Conjecture 7.5.*

Proof. First, we show that sp (and sp_Q) respects the quantum pairing in the following sense.

$$\lim_{\lambda \rightarrow 1} \frac{1}{|W|} \nu_Q(((a, b))^{Y, tw}) = ((sp(a), sp(b)))^X$$

The Q^d coefficient of the left hand side is equal to:

$$\lim_{\lambda \rightarrow 1} \frac{1}{|W|} \sum_{d' \rightarrow d} \langle a, b \rangle_{0,2,d'}^{tw} Q^d$$

This equal to the RHS by Conjecture 7.4.

Next, we show that ϕ is a ring homomorphism. Let a, b be Weyl-invariant classes in $K^*(Y)$. For ψ_i a basis in $K^*(Y)$, we have the following:

$$a \circ_Y b = \sum_{d \in H^2(X//T), i} \langle a, b, \psi_i \rangle_{0,3,d}^{tw} \psi_i^* Q^d$$

Observe that W induces an action on $\coprod_{d' \rightarrow d} Y_{g,n,d'}$. Since W acts on Y by automorphisms, it induces isomorphisms from $Y_{g,n,d}$ and Y_{g,n,d^w} , which preserve the twisting class, we conclude that

$$\langle a^w, b^w, c^w \rangle_{0,3,d^w}^{tw} = \langle a, b, c \rangle_{0,3,d}^{tw},$$

this means $a \circ b$ is W -invariant, so applying sp_Q is possible.

For $t \in K^*(X)$, with a lift $\tilde{t} \in K^*(Y)$, we have:

$$((\phi(a \circ_Y b), t)) = \lim_{\lambda \rightarrow 1} ((sp_Q(a \circ_Y b), t)) = \lim_{\lambda \rightarrow 1} \frac{1}{|W|} \nu_Q(((a \circ_Y b, \tilde{t}))^{tw}) = \lim_{\lambda \rightarrow 1} \sum_{d \rightarrow d', d' \in H_2(X)} Q^d \langle a, b, \tilde{t} \rangle_{0,3,d'}^{tw} = ((\phi(a) \circ_X \phi(b), t))$$

Thus $\phi(a \circ_Y b) = \phi(a) \circ_X \phi(b)$. surjectivity of ϕ is obvious. Since sp is a map of Q -modules (if we interpret Q as the Novikov variables of $X//T$), the kernel of sp is the ideal generated by the kernel of sp on $K^*(X)$, which is generated by $Eu(V^M)$ and kernel of sp on the Novikov variables. □

Theorem 7.8. *Conjecture 7.3 implies Conjecture 7.4 for $n = 1$*

Proof. The correspondence for descendent invariants is equivalent to the following statement for all $t \in K^*(X)$:

$$\left\langle \frac{t}{1-q} \right\rangle_{0,1,d} = \frac{1}{|W|} \sum_{d' \rightarrow d} \left\langle \frac{\tilde{t}}{1-q} \right\rangle_{0,1,d'}^{tw}$$

The left hand side is (J_X, t) , the right-hand side is $\lim_{y \rightarrow 1} sp_Q(J_Y^{tw}, \tilde{t})$, these are equal by the hypothesis, and the classical abelian/non-abelian correspondence. □

Theorem 7.9. *If $K^*(X)$ and $K^*(Y)$ are generated by line bundles, then Conjecture Conjecture 7.3 implies Conjecture Conjecture 7.4 for all values of n .*

Proof. This is a consequence of the twisted reconstruction theorem. We induct on n, d and the degree of any line bundles appearing in the inputs, as well as the degree of descendants.

Beginning with $\sum_{d' \rightarrow d} \langle \tilde{a}_1, \dots \rangle_{0, n, d}$ we can use the reconstruction relations to reduce the degrees of the inputs, n , and d , and apply the induction hypothesis.

The nontrivial thing to check is applying the 3rd relation and using the splitting axiom. What we need to prove is that (assuming the induction hypothesis):

$$\langle a_1, \dots \rangle_{0, n, d}^{\Phi} = \sum_{d' \rightarrow d} \langle \tilde{a}_1, \dots \rangle_{0, n, d'}^{tw, \Phi}$$

Where Φ is the contraction map appearing in the proof of the splitting axiom.

$$\sum_{d' \rightarrow d, n_1 + n_2 = n, d_1 + d_2 = d'} \sum_{i, j} \langle \tilde{a}_1, \dots, \phi_i \rangle_{0, n_1 + 1, d_1} g^{ij} \langle \phi_j, \tilde{a}_{n_1 + 1}, \dots \rangle_{0, n_2 + 1, d_2}$$

Similarly, the left hand side is:

$$\sum_{i, j, d_1 + d_2 = d, n_1 + n_2 = n} \langle a_1, \dots, a_{n_1}, \phi_i \rangle_{0, n_1 + 1, d_1} g^{ij} \langle a_1, \dots, a_{n_1}, \phi_j \rangle_{0, n_2 + 1, d_2}.$$

First we note that if $d_1 + d_2 = d'$, then $sp(d_1) + sp(d_2) = d$, hence for any $w_1, w_2 \in W$, $sp(d_1^{w_1}) + sp(d_2^{w_2}) = d$. So we can rewrite the right hand side as:

$$\sum_{n_1 + n_2 = n, d_1 + d_2 = d} \sum_{i, j} \left(\sum_{d'_1 \rightarrow d_1} \langle \tilde{a}_1, \dots, \phi_i \rangle_{0, n_1 + 1, d'_1} \right) g^{ij} \left(\sum_{d'_2 \rightarrow d_2} \langle \phi_j, \tilde{a}_{n_1 + 1}, \dots \rangle_{0, n_2 + 1, d'_2} \right)$$

We can write the LHS as $T_1(\sum g_{ij} \phi_i \otimes \phi_j)$, where

$$T_1 = \sum_{d_1 + d_2 = d, n_1 + n_2 = n} \langle a_1, \dots, \cdot \rangle_{0, n_1 + 1, d_1} \otimes \langle \cdot, a_{n_1 + 1}, \dots \rangle_{0, n_2 + 1, d_2}.$$

Similarly we can write the left hand side as $T_2(\sum_{i, j} g^{ij, tw} \psi_i \otimes \psi_j)$, where T_2 is the tensor

$$T_2 := \frac{1}{|W|} \sum_{n_1 + n_2 = n, d_1 + d_2 = d} \left(\sum_{d'_1 \rightarrow d_1} \langle a_1, \dots, \cdot \rangle_{0, n_1 + 1, d'_1} \right) \otimes \left(\sum_{d'_2 \rightarrow d_2} \langle \cdot, a_{n_1 + 1}, \dots \rangle_{0, n_2 + 1, d'_2} \right)$$

The induction hypothesis allows us to relate the tensors T_1 and T_2 , as linear functions on $K^*(X) \otimes K^*(X)$ and $(K^*(Y) \otimes K^*(Y))^W$ respectively. Applying it yields:

$$sp^* T_1 = \frac{1}{|W|} \lim_{\lambda \rightarrow 1} T_2$$

The class $\sum_{i, j} g^{ij, tw} \psi_i \otimes \psi_j$ is the Poincare dual to the diagonal in $X \times X$. The class $\sum_{i, j} g^{ij, tw} \psi_i \otimes \psi_j$ is the twisted Poincare dual to the diagonal in $Y \times Y$.

Since sp identifies the structure sheaves of the diagonals of X and Y , we have: $\lim_{\lambda \rightarrow 1} PD^{tw}(\Delta_Y) = \frac{1}{|W|} sp^* PD(\Delta_X)$. The factor $\frac{1}{|W|}$ comes from the difference between the Poincare pairings. Thus the two quantities are equal, as desired. \square

Remark. The conditions fails for X a Grassmanian or flag variety, but we remedy this in the following way: We work equivariantly with respect to the usual torus action, then after localizing at the ideal generated by $1 - \Lambda_i$, $K^*(X)$ is generated by line bundles, and the argument above goes through.

7.1 Twistings

A G representation E induces a vector bundles E_X and E_Y on X and Y respectively. Their classes in K -theory are related by the classical abelian/non-abelian correspondence.

One can make the exact same conjectures relating invariants of X twisted by $C(E_X)$ to twisted invariants of Y , further twisted by $C(E_Y)$, and the same equivalences hold.

Theorem 7.10. *Theorems 7.8 and 7.9 also apply in the setting of twisted invariants.*

There is also a direct relationship between the conjecture for Lagrangian cones and twisted Lagrangian cones.

Theorem 7.11. *For any choice of C, E , Conjecture 7.2 and Conjecture 7.1 for untwisted invariants imply Conjecture 7.2, Conjecture 7.1 for twisted invariants.*

This is a consequence of Givental’s Quantum Adams–Riemann–Roch theorem, which relates the Lagrangian cones of the twisted theory to the untwisted theory via a symplectomorphism Δ . It applies in both the ordinary and S_n -invariant settings. Let Δ_X, Δ_Y be the maps coming from the twistings on X, Y , respectively. We observe that $sp_Q(\Delta_Y) = \Delta_X$, thus proving the theorem. (The same result for cohomology is theorem in [4]).

8 Summary: Toolkit for Computing Quantum K -rings

In view of the results of the previous sections, we have the following “toolkit” for calculating relations in small quantum K -rings (twisted or not).

1. Interpret the ring as a twisted quantum K -ring of a simpler space if possible. (i.e. move from $X//G$ to $X//T$, or consider the Gromov–Witten invariants of a complete intersection as twisted Gromov–Witten invariants of the ambient space).
2. Determine the small J -function of the theory. (In many cases, this is doable by using Givental’s quantum Riemann–Roch theorem to get some value of \mathcal{J}^{sym} , if it is the value at $t = 0$ we are done.)
3. Find a bound for the q -degrees of each term in J
4. Find difference equations annihilating J
5. Convert these into relations, using the degree bound and quantum triviality theorem (if possible) to simplify functions of $A_{i,com}|_{t=0}$.

9 Applications

9.1 Level Structures of Projective Space

We first consider $X = \mathbb{P}^N$ with the level structure twisting determined by the bundle $P := \mathcal{O}(-1)$ and the integer ℓ . The small J -function of this theory, for $-1 < \ell \leq N + 1$, was calculated by Givental–Yan in [8] and is equal to:

$$J^\ell := (1 - q) \sum_d Q^d \frac{P^\ell q^{\binom{d}{2}}}{\prod_i \prod_{j=1}^d (1 - Pq^j / \Lambda_i)}$$

Now, we can find

Proposition 9.1. \tilde{J}^ℓ satisfies the following difference equation:

$$\prod_i (1 - \Lambda_i^{-1} q^{Q\partial_Q}) \tilde{J}^\ell = Q q^{\ell Q\partial_Q} J_{1,N+1}^\ell$$

Proof.

$$\begin{aligned} \prod_i (1 - \Lambda_i^{-1} q^{Q\partial_Q}) P^{\frac{\ln(Q)}{\ln(q)}} J_{1,N+1}^\ell &= (1-q) \prod_i (1 - q^d P/\Lambda_i) P^{\frac{\ln(Q)}{\ln(q)}} \sum_{d \geq 0} \frac{Q^d P^{\ell d} q^{\ell \binom{d}{2}}}{\prod_i \prod_{m=1}^d (1 - q^m P/\Lambda_i)} \\ &= (1-q) P^{\frac{\ln(Q)}{\ln(q)}} \sum_{d \geq 1} \frac{Q^d P^{\ell d} q^{\ell \binom{d}{2}}}{\prod_i \prod_{m=1}^{d-1} (1 - q^m P/\Lambda_i)} = (1-q) P^{\frac{\ln(Q)}{\ln(q)}} \sum_{d \geq 1} q^{\ell(d-1)} P^{\ell d} \frac{Q^d P^{\ell(d-1)} q^{\ell \binom{d-1}{2}}}{\prod_i \prod_{m=1}^{d-1} (1 - q^m P/\Lambda_i)} \quad (9.1) \\ &= (1-q) Q q^{\ell Q\partial_Q} P^{\frac{\ln(Q)}{\ln(q)}} \sum_{d \geq 1} \frac{Q^{d-1} P^{\ell(d-1)} q^{\ell \binom{d-1}{2}}}{\prod_i \prod_{m=1}^{d-1} (1 - q^m P/\Lambda_i)} = Q q^{\ell Q\partial_Q} \tilde{J}_{1,N+1} \end{aligned}$$

□

Writing J^ℓ as $(1-q) \sum Q^d J_d$, we have that $\deg(J_d) = \ell \binom{d}{2} - (N+1) \binom{d+1}{2}$, this is always strictly less than $-d$, so by the quantum triviality theorem $A_{com}|_{t=0} = P$. So we can conclude:

Corollary 9.2. The quantum K -ring with level structure of \mathbb{P}^N with respect to the bundle P and the level ℓ is determined by the relation:

$$\prod_i (1 - P/\Lambda_i) = Q P^\ell \quad (9.2)$$

Remark. The physical derivation of the quantum K -ring of projective space, done in e.g. [11], relies on a specific choice of parameters called the Chern-Simons levels. Ordinary quantum K -theory comes from choosing the parameter \tilde{k}_1 (in the notation of [11]) to be $-\frac{N+1}{2}$. The prediction corresponding to $\tilde{k}_1 = \ell - \frac{N+1}{2}$ are exactly the relations we have computed here for level ℓ .

9.2 Complete Intersections in Projective Space

If $P = O(1)$, the J -function of \mathbb{P}^N is:

$$\sum_{d \geq 0} \frac{Q^d}{\prod_{m=1}^d (1 - P q^m)^{N+1}}$$

Given some complete intersection Y of hypersurfaces of degree ℓ_1, \dots, ℓ_r , we can apply Givental's quantum Lefschetz to get that the following function I_Y is a value of the J -function for invariants twisted by the Euler class of $\bigoplus O(\ell_i)$:

$$I_Y : \sum_{d \geq 0} \frac{Q^d}{\prod_{m=1}^d (1 - P q^m)^{N+1}} \prod_i \prod_{0 \leq b \leq \ell_i, d} (1 - \lambda P^{\ell_i} q^b)$$

I_Y has no poles at $q = 0$, and $I_Y = (1-q) \sum_d Q^d I_d$, with $\deg(I_d) = \sum_i \binom{\ell_i d}{2} - (N+1) \binom{d+1}{2}$.

Proposition 9.3. If $\sum_i \ell_i^2 \leq N+1$, then I_Y is the small J -function of this twisted theory, i.e. it is the pushforward to X of J_Y , furthermore all $Q^{\neq 0}$ terms of $P q^{Q\partial_Q} \frac{I_Y}{1-q}$ vanish at $q = \infty$.

Proof. First we note that I_Y has no poles at $q = 0$, and if we write $I_Y = (1-q) \sum_d Q^d I_d$, then the remaining two conditions are both checked by bounds on $\deg(I_d) = \sum_i \binom{\ell_i d}{2} - (N+1) \binom{d+1}{2}$. We want to check that

$\deg(I_d) < -1$ so I_Y is the small J -function, and that $\deg(I_d) < -d$ so the required terms vanish at infinity. We have:

$$\deg(I_d) = \frac{d}{2} \left(d \left(\sum_i \ell_i^2 - (N+1) \right) + \sum_i \ell_i - (N+1) \right)$$

Our desired result is equivalent to the inequality

$$d \left(\sum_i \ell_i^2 - (N+1) \right) + \sum_i \ell_i - (N+1) < -2 \quad (9.3)$$

By hypothesis, $\sum_i \ell_i^2 \leq N+1$, it is sufficient to check it for $d=1$, i.e. to show:

$$\sum_i (\ell_i^2 - (N+1)) + \left(\sum_i \ell_i - (N+1) \right) < -2 \quad (9.4)$$

Equivalently:

$$\sum_i 2\ell_i^2 + \sum_i (\ell_i - \ell_i^2) < 2N \quad (9.5)$$

However we observe that unless at most one $\ell_i = 2$, and all the others are equal to 1, we have:

$$\sum_i (\ell_i - \ell_i^2) < -2 \quad (9.6)$$

Thus when (9.6) is true:

$$\sum_i 2\ell_i^2 + \sum_i (\ell_i - \ell_i^2) \leq 2N + 2 - \sum_i (\ell_i - \ell_i^2) < 2N \quad (9.7)$$

In those exceptional cases, we can check by hand that (9.5) is satisfied, noting that $r < N$, so it is impossible to have both $\sum_i \ell_i^2 - (N+1)$ and $\sum_i \ell_i - (N+1)$ lie in the set $\{0, -1\}$. \square

Theorem 9.4. *If $\sum_i \ell_i^2 \leq N+1$, then the restriction to Y of the following expression holds in $QK^*(Y)$:*

$$(1-P)^{N+1-r} = \prod_i \frac{1-P^{\ell_i}}{1-P} Q$$

Proof. We have the equation:

$$(1-Pq^{Q\partial Q})^{N+1} I_Y = \prod_i (1-\lambda P_i^{\ell_i} q^{\ell_i Q\partial Q}) Q I_Y \quad (9.8)$$

After normalizing, this corresponds to the equation for \tilde{I}_Y (we omit this step in subsequent examples):

$$(1-q^{Q\partial Q})^{N+1} \tilde{I}_Y = \prod_i (1-\lambda q^{\ell_i Q\partial Q}) Q \tilde{I}_Y \quad (9.9)$$

Which gives us the relation in the twisted quantum K -theory of \mathbb{P}^N , since I_Y is the small J -function of this theory and $A_{com}|_{t=0} = P$ by the quantum triviality theorem:

$$(1-P)^{N+1} = \prod_i (1-\lambda P^{\ell_i}) Q,$$

equivalently:

$$(1 - P)^{N+1-r} = \prod_i \frac{1 - \lambda P^{\ell_i}}{1 - P} Q$$

Taking non-equivariant limits and pulling back to $QK^*(Y)$ gives the relation:

$$(1 - P)^{N+1-r} = \prod_i \frac{1 - P^{\ell_i}}{1 - P} Q$$

□

Remark. In the case that $r = 1$, i.e. when Y is a hypersurface, these results are mathematical proofs of the physical predictions made in section 5 of [10].

9.3 Quasimaps to $T^*\mathbb{P}^N$

An alternative formulation of quantum K -theory has been developed using the quasimap compactifications instead of the stable map ones. In particular, there have been many results relating relations in these quasimap rings to integrable systems. The quasimap K -ring of $T^*\mathbb{P}^N$ is determined by the relation ([19]):

$$P^{N+1} \hbar^{(-N-1)/2} \prod_i (\Lambda_i^{-1}) Q = \prod_{i=1}^{N+1} \frac{1 - P/\Lambda_i}{1 - \hbar P/\Lambda_i}$$

An important question to ask is whether there is a relationship between the quasimap quantum ring and some version of the stable map one. We will give a partial answer this for the case of $T^*\mathbb{P}^{N-1}$.

The quasimap I -function that describes the invariants $T^*\mathbb{P}^{N-1}$ has the following form:

$$\hat{I}^T = \sum_{0 \leq d} Q^d \prod_{j=1}^{N+1} \frac{\prod_{m=0}^{d-1} 1 - q^m y X / \Lambda_j}{\prod_{m=0}^d 1 - q^m X / \Lambda_j}$$

It was shown in [18] that $(1 - q)\hat{I}^T$ is the small J function for a particular set of twisted invariants of \mathbb{P}^{N-1} . The twisting is the inverse Euler class of the tangent bundle, followed by the determinant of the cotangent bundle. (Equivalently, it's the stable map theory corresponding to $T^*\mathbb{P}^N$ with level structure determined by $T^*\mathbb{P}^N$ and $\ell = 1$).

Using the twisted version of the Iritani-Milanov-Tonita theorem, we can calculate the quantum K -ring with respect to this twisting, and compare it to the quasimap relations.

Let $I^T = \prod_j 1 - y \frac{X}{\Lambda_j} q^{Q\partial_Q} \hat{I}^T$. The action of this operator replaces the $d - 1$ in the product with a d . If we denote the operator taking \hat{I}^T to I^T by D , we have:

$$\prod_{j=1}^{N+1} \frac{1 - \frac{X}{\Lambda_j} q^{Q\partial_Q}}{1 - y \frac{X}{\Lambda_j} q^{Q\partial_Q}} D \hat{I}^T = Q D \hat{I}^T \quad (9.10)$$

Since the q -degree of the Q^d th term is at most $-d$, we have that the operator $q^{Q\partial_Q}$ gets converted into multiplication by P_i via the quantum triviality theorem. This yields the relation (after dividing both sides by $D|_{q=1}$):

$$\prod_{j=1}^{N+1} (1 - X/\Lambda_j) = Q \prod_{j=1}^{N+1} (1 - yX/\Lambda_j) \quad (9.11)$$

We can rewrite this as:

$$\prod_{j=1}^{N+1} \frac{(1 - X/\Lambda_j)}{1 - yX/\Lambda_j} = Q$$

We can multiply both sides by $\prod_j Xy/\Lambda_j$ to get:

$$\prod_{j=1}^{N+1} \frac{(1 - X/\Lambda_j)}{\Lambda_j/yX - 1} = Q \prod_j \frac{X}{\Lambda_j} y^N$$

Replacing y with \hbar^{-1} , this is similar to what is obtained in the quasimap case, up to a factor of $\hbar^{(N+1)/2}$. We leave the explanation behind this discrepancy, and some more detailed comparisons between the quasimap and stable map rings to a future work.

9.4 Grassmanians

We will use the ring-theoretic abelian/non-abelian correspondence to recover some relations for the case $X = Gr(k, n)$, the Grassmanian of k -planes in \mathbb{C}^n .

Recall the Grassmanian can be written as the GIT quotient $Hom(\mathbb{C}^k, \mathbb{C}^n)//GL(k)$. The corresponding abelian quotient Y is a product of k copies of \mathbb{P}^{n-1} s.

The classical abelian/non-abelian correspondence takes the following form in this case. $K^*(Y) \cong \prod_{i=1}^n \frac{\mathbb{C}[P]}{\prod_i (1 - P_i/\Lambda_i)}$, where P_i is the tautological bundle of the i th projective space.

The ring map $K^*(Y)^W \rightarrow K^*(Gr(k, n))$ is given by sending symmetric functions of the P_i to the same symmetric functions of the Chern roots of the tautological bundle V . We will thus abusively also write $V = \sum_i P_i$.

Givental-Yan proved Conjecture 7.2 for Grassmanians, in particular they established the following:

Theorem 9.5. [8] *The small J -function of the Grassmanian is equal to:*

$$(1 - q) \sum_{d_1, \dots, d_n \leq 0} \frac{Q^{\sum d_i}}{\prod_{i=1}^n \prod_{j=1}^n \prod_{m=1}^{d_i} (1 - q^m \frac{P_i}{\Lambda_j})} \prod_{i \neq j} \frac{\prod_{m=-\infty}^{d_i - d_j} (1 - q^m \frac{P_i}{P_j})}{\prod_{m=-\infty}^0 (1 - q^m \frac{P_i}{P_j})}$$

It is the image under sp_Q of J_Y^{tw} , which is a value of the big S_n -equivariant J -function of the twisted theory of Y :

$$J_Y^{tw}(q) := (1 - q) \sum_{d_1, \dots, d_n \leq 0} \frac{\prod_i Q_i^{d_i}}{\prod_{i=1}^n \prod_{j=1}^n \prod_{m=1}^{d_i} (1 - q^m \frac{P_i}{\Lambda_j})} \prod_{i \neq j} \frac{\prod_{m=-\infty}^{d_i - d_j} (1 - q^m \Lambda_0 \frac{P_i}{P_j})}{\prod_{m=-\infty}^0 (1 - q^m \Lambda_0 \frac{P_i}{P_j})}$$

In fact, J_Y^{tw} is actually the small J -function of this theory. To prove this, it is sufficient to show that its projection to K_+ is equal to $1 - q$, so it corresponds to the input $t = 0$. This means the terms for $d \neq 0$ must be reduced rational functions, i.e. with no poles at $q = 0, \infty$ and with negative q -degree outside the Q^0 term.

It is easy to see this since if we write J_Y^{tw} as $(1 - q) \sum_d Q^d J_d$, J_d can only have q^{-1} terms in the denominator, so there are no poles at 0. In terms of degree, we have:

$$deg(J_d) = \sum_{i,j} \binom{d_i - d_j + 1}{2} - n \sum_i \binom{d_i + 1}{2}$$

We can rewrite this as

$$\sum_i \left(\sum_j \binom{d_i - d_j + 1}{2} - n \binom{d_i + 1}{2} \right)$$

Each term $\sum_j \binom{d_i - d_j + 1}{2} - n \binom{d_i + 1}{2}$ must be strictly negative unless $d_i = 0$ since $N > k$ and $\binom{d_i - d_j}{2} < \binom{d_i + 1}{2}$. In fact we can say each term contributes degree at most $k - n$, hence $\deg(J_d) < -1$.

Thus, we can apply Conjecture 7.5 in this case.

In fact, for the purpose of using the quantum triviality theorem, we will prove two stronger bounds on $\deg(J_d)$

Lemma 9.6.

- $\deg(J_d) < (k - n) \max(d_i)$
- $\deg(J_d) < \sum_i d_i$

Proof. From before we have:

$$\deg(J_d) = \sum_i \left(\sum_j \binom{d_i - d_j + 1}{2} - n \binom{d_i + 1}{2} \right)$$

Each summand is less than $(k - n) \max(d_i)$, which establishes both inequalities. □

This gives us the following corollaries as a result of the quantum triviality theorem:

Corollary 9.7.

- $A_{i,com}|_{t=0} = P_i$
- Letting $e_\ell(P)$ denote the ℓ th symmetric polynomial applied to P_1, \dots, P_k , the expression $e_\ell(P)$ is equal to its classical product, i.e. $\wedge^\ell(\bigoplus_i P_i)$.
- The same is true for $h_\ell(P)$ with $\ell \leq n - k$, where h_ℓ denotes the complete homogenous symmetric polynomial.

We can now find the relations in $QK^{tw}(Y)$.

Theorem 9.8. *The following relations are true in $QK^{tw}(Y)$:*

$$F_\lambda(Q_i, P_i) = 0$$

Where

$$F_\lambda(x, t) := t^k e_n(\Lambda) \prod_j \frac{(P_j - \lambda t)}{(t - \lambda P_j)} - \prod_a (\Lambda_a - t) \prod_j P_j$$

Proof. J_Y^{tw} satisfies the following q -difference equations:

$$\prod_{j \neq i} (1 - \lambda q P_j P_i^{-1} q^{Q_j \partial_{Q_j} - Q_i \partial_{Q_i}}) \prod_a (1 - P_i q^{Q_i \partial_{Q_i}} / \Lambda_a) \mathcal{J} = Q_i \prod_{j \neq i} (1 - \lambda q P_i P_j^{-1} q^{Q_i \partial_{Q_i} - Q_j \partial_{Q_j}}) \mathcal{J}$$

These translate to the relations (using the fact that $A_{i,com} = P_i$ by the quantum triviality theorem):

$$\prod_a (1 - P_i / \Lambda_a) = Q_i \prod_{j \neq i} \frac{1 - \lambda P_i / P_j}{1 - \lambda P_j / P_i}$$

Since

$$\frac{1 - \lambda P_i/P_j}{1 - \lambda P_j/P_i} = \frac{P_i(P_j - \lambda P_i)}{P_j(P_i - \lambda P_j)},$$

the above relation is equivalent to:

$$\prod_j P_j(P_i - \lambda P_j) \prod_a (1 - P_i/\Lambda_a) = P_i^k Q_i \prod_j (P_j - \lambda P_i)$$

Multiplying by $\prod_i \Lambda_i$ gives the desired result. □

We have established that elements of $QK^{tw}(Y)^W|_{Q_i=Q}$ have well-defined limits as λ approaches 1. Call the ring thus obtained S , which maps to $QK^*(X)$ via sp .

If we denote F as $\lim_{\lambda \rightarrow 1} F_\lambda(Q, t)$, then symmetric combinations of $F(P_i) = 0$ become relations in S . As before, we will denote the set of P_i collectively by P , with the notation that $f(P) = f(P_1, \dots, P_k)$ for f any symmetric function. Similarly, we will denote the equivariant parameters collectively by Λ .

Calculating the limit explicitly yields:

$$F = \sum_{\ell=0}^{n-k} (-1)^\ell t [e_k(P) e_\ell(\Lambda) + Q e_n(\Lambda) e_{\ell-n+k}(0)]$$

Remark. The equations $F(P_i) = 0$ are exactly the Bethe Ansatz equations obtained from physical models of $QK^*(X)$ in [11].

We will now use Vieta's formulas to explicitly compute relations in S . The algebraic manipulations done here are identical to those in [9] for the case of Grassmanians. However, in our situation, we can give each relation a concrete mathematical meaning since we are working in $QK^{tw}(Y)$. Denote the entire set of roots of F by w , and denote the subset of these roots not in P by \bar{P} , as consequence:

$$e_\ell(w) = \sum_{i=0}^{n-k} e_{\ell-i}(P) e_i(\bar{P}) \tag{9.12}$$

Denote the t^ℓ coefficient of F by F_ℓ , we have that:

$$F_{n-\ell} = (-1)^\ell (e_k(P) e_\ell(\Lambda))$$

Thus, applying Vieta's formula to F gives:

$$e_\ell(w) = (-1)^\ell \frac{e_k(P) e_\ell(T)}{e_k(P)} \tag{9.13}$$

Combining the (9.12) and (9.13) gives:

$$e_\ell(w) = e_\ell(T) + Q e_\ell(\bar{P}) e_{\ell-n+k}(0) \tag{9.14}$$

Using (9.14) to replace $e_\ell(w)$ gives us:

$$e_\ell(T) + Q e_\ell(\bar{P}) e_{\ell-n+k}(0) = \sum_{i=0}^{n-k} e_{\ell-i}(P) e_i(\bar{P}) \tag{9.15}$$

We would like to use (9.15) to solve for $e_\ell(\bar{P})$, to do this, first note that (9.15) is the y^ℓ term in the equation:

$$\left(\sum_{i=1}^k y^i e_i(P)\right)\left(\sum_{j=1}^k y^j e_j(\bar{P})\right) = \sum_{s=0}^n y^s e_s(\Lambda) + Qe^{n-k} e_{n-k}(\Lambda) \quad (9.16)$$

Since the inverse of the series $\sum y^i e_i(S)$ is $\sum (-1)^i y^i h_i(S)$, where h_i are the total homogenous symmetric polynomials, we can rewrite (9.16) as:

$$\left(\sum_{j=1}^k y^j e_j(\bar{P})\right) = \left(\sum_{i=1}^{\infty} (-y^i) h_i(P)\right)\left(\sum_{s=0}^n y^s e_s(\Lambda) + Qe^{n-k} e_{n-k}(\Lambda)\right) \quad (9.17)$$

Extracting the degree ℓ part yields:

$$e_{\ell}(\bar{P}) = \sum_{i=0}^{n-k} (-1)^i e_{\ell-i}(\Lambda) h_i(P) + Qe_{n-k}(\bar{X}) e_{\ell-n+k}(0) \quad (9.18)$$

Now, having established these relations, we can specialize them into $QK^*(X)$, noting that $e_{\ell}(P_i)$ and $h_{\ell}(P_i)$ can be interpreted as their classical products. Since specialization sends P_i to the i th Chern root of the tautological bundle S , we have the following:

$$sp(e_{\ell}(\bar{P})) = \begin{cases} \wedge^{\ell}(\mathbb{C}^n/S) & \ell < n-k \\ \frac{1}{1-Q} \wedge^{\ell}(\mathbb{C}^n/S) & \ell = n-k \end{cases} \quad (9.19)$$

$$sp(e_{\ell}(P)) = \wedge^{\ell} S \quad (9.20)$$

Thus specializing the relation (9.15) yields:

$$\sum_{r=0}^{n-k} \wedge^{\ell-r} \mathcal{S} \times_Q \wedge^r(\mathbb{C}^n/S) = \wedge^{\ell} \mathbb{C}^n - \frac{q}{1-q} \det(\mathbb{C}^n/S) \times_Q (\wedge^{\ell-n+k} \mathcal{S} - \mathcal{O}\delta_{\ell, n-k}). \quad (9.21)$$

Adding factors of y , this yields the Whitney relations:

$$\Lambda_y(\mathcal{S}) \times_Q \Lambda_y(\mathbb{C}^n/S) = \Lambda_y(\mathbb{C}^n) - y^{n-k} \frac{q}{1-q} \det(\mathbb{C}^n/S) \times_Q (\Lambda_y(\mathcal{S}) - 1). \quad (9.22)$$

These are deformations of the usual Whitney relations, which generate $K^*(X)$, hence this is a presentation for $QK^*(X)$.

References

- [1] David Anderson, Linda Chen, and Hsian-Hua Tseng. On the finiteness of quantum k-theory of a homogeneous space. *International Mathematics Research Notices*, 2022(2):1313–1349, 2022.
- [2] Ionuț Ciocan-Fontanine, Bumsig Kim, and Claude Sabbah. The abelian/nonabelian correspondence and frobenius manifolds. *Inventiones mathematicae*, 171(2):301–343, 2008.
- [3] Tom Coates. *Riemann–Roch theorems in Gromov–Witten theory*. PhD thesis, UC Berkeley, 2003.
- [4] Tom Coates, Wendelin Lutz, and Qaasim Shafi. The abelian/nonabelian correspondence and gromov–witten invariants of blow-ups. *Forum of Mathematics, Sigma*, 10, 2021.
- [5] Alexander Givental. Permutation-equivariant quantum k-theory i. definitions. elementary k-theory of $\overline{M}_{0,n}/s_n$. *arXiv preprint arXiv:1508.02690*, 2015.
- [6] Alexander Givental. Explicit reconstruction in quantum cohomology and k-theory. In *Annales de la Faculté des sciences de Toulouse: Mathématiques*, volume 25, pages 419–432, 2016.
- [7] Alexander Givental. Permutation-equivariant quantum K-theory XI: Quantum Adams-Riemann-Roch. 2017.
- [8] Alexander Givental, Xiaohan Yan, et al. Quantum k-theory of grassmannians and non-abelian localization. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications*, 17:018, 2021.
- [9] Wei Gu, Leonardo Mihalcea, Eric Sharpe, Weihong Xu, Hao Zhang, and Hao Zou. Quantum k theory rings of partial flag manifolds. *Journal of Geometry and Physics*, 198:105127, 2024.
- [10] Wei Gu, Leonardo Mihalcea, Eric Sharpe, and Hao Zou. Quantum k theory of symplectic grassmannians. *Journal of Geometry and Physics*, 177:104548, 2022.
- [11] Wei Gu, Leonardo C Mihalcea, Eric Sharpe, and Hao Zou. Quantum k theory of grassmannians, wilson line operators, and schur bundles. *arXiv preprint arXiv:2208.01091*, 2022.
- [12] Megumi Harada and Gregory D Landweber. The k-theory of abelian versus nonabelian symplectic quotients. *arXiv preprint math/0601294*, 2006.
- [13] Irit Huq-Kuruvilla. Multiplicative quantum cobordism theory, 2021.
- [14] Hiroshi Iritani, Todor Milanov, and Valentin Tonita. Reconstruction and convergence in quantum k-theory via difference equations. *International Mathematics Research Notices*, 2015(11):2887–2937, 2015.
- [15] Hans Jockers and Peter Mayr. A 3d gauge theory/quantum k-theory correspondence. *arXiv preprint arXiv:1808.02040*, 2018.
- [16] Y-P Lee et al. Quantum K-theory I: Foundations. *Duke Mathematical Journal*, 121(3):389–424, 2004.
- [17] Y.-P. Lee and R. Pandharipande. A reconstruction theorem in quantum cohomology and quantum k-theory. *American Journal of Mathematics*, 126(6):1367–1379, 2004.
- [18] Henry Liu. Self-duality in quantum k-theory. *arXiv: Algebraic Geometry*, 2019.
- [19] Petr P. Pushkar, Andrey V. Smirnov, and Anton M. Zeitlin. Baxter q-operator from quantum k-theory. *Advances in Mathematics*, 360:106919, 2020.
- [20] Valentin Tonita. Twisted k-theoretic gromov–witten invariants. *Mathematische Annalen*, 372(1):489–526, 2018.
- [21] Kazushi Ueda and Yutaka Yoshida. 3d n=2 chern-simons-matter theory, bethe ansatz, and quantum k-theory of grassmannians. *Journal of High Energy Physics*, 2020(8):1–44, 2020.