

Elliptic Quantum Toroidal Algebra $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$, Vertex Operators and L -operators

Hitoshi Konno[†] and Andrey Smirnov^{*}

[†]*Department of Mathematics, Tokyo University of Marine Science and Technology,
Etchujima, Koto, Tokyo 135-8533, Japan*

hkonno0@kaiyodai.ac.jp

^{*}*Department of Mathematics, University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA*

asmirnov@unc.edu

Abstract

We propose new vertex operators, both the type I and the type II dual, of the elliptic quantum toroidal algebra $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ by combining representations of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ and the notions of the elliptic stable envelopes for the instanton moduli space $\mathcal{M}(n, r)$. The vertex operators reproduce the shuffle product formula of the elliptic stable envelopes by their composition. We also show that the vacuum expectation value of a composition of the vertex operators gives a correct formula of the K-theoretic vertex function for $\mathcal{M}(n, r)$. We then derive exchange relations among the vertex operators and construct a L -operator satisfying the $RLL = LLR^*$ relation with R and R^* being elliptic dynamical instanton R -matrices defined as transition matrices of the elliptic stable envelopes. Assuming a universal form of L , we define a comultiplication Δ in terms of it. It turns out that the new vertex operators are intertwining operators of the $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ -modules w.r.t Δ .

1 Introduction

The aim of this paper is to present a new formulation of the vertex operators of the elliptic quantum toroidal algebra (EQTA) $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ by combining its representations and the notions of the elliptic stable envelopes for the instanton moduli space $\mathcal{M}(n, r)$.

The EQTA $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ is an elliptic quantum group associated with the toroidal algebra of type \mathfrak{gl}_1 formulated in [32]. Here parameters t_1 and t_2 correspond to q^{-1} and t in [32], respectively. It is defined by generators and relations in the same scheme as the elliptic quantum group (EQG) $U_{q, p}(\widehat{\mathfrak{g}})$ associated with the affine Lie algebra $\widehat{\mathfrak{g}}$ [7, 19, 23, 31] and equipped with a Hopf algebroid structure given by the Drinfeld comultiplication, i.e. a comultiplication formula in terms of the Drinfeld generators or their generating functions called the elliptic currents.

Two types of vertex operators, type I and type II dual, have been constructed as intertwining operators of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ -modules and shown to give a realization of the generating function of the affine quiver W -algebra associated with the Jordan quiver $\Gamma(\widehat{A}_0)$ [20]. At the same time, the elliptic currents of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ gives the screening currents of it. In addition, it has been shown that these vertex operators realize the refined topological vertices [15, 16] relevant to calculate the instanton partition functions of the 5d and 6d lift of the 4d $\mathcal{N} = 2^*$ gauge theory [39, 40]. However their relations to the elliptic stable envelopes and to the vertex functions of the corresponding quiver variety were missing.

The notion of stable envelopes was initiated in [35, 41]. They provide a good basis of equivariant cohomology and K-theory of Nakajima quiver varieties X [37, 38] and has nice applications to enumerative geometry, geometric representation theory and quantum integrable systems. In particular, the transition matrices between stable envelopes defined for different chambers give the R -matrices satisfying the Yang-Baxter equation (YBE). Regarding such R -matrix as universal one i.e. a L -operator which generate a quantum group, the transition property can be understood as a Gauss decomposition of the L -operator in terms of the stable envelopes. Hence the stable envelopes provide a new geometric formulation of quantum groups as well as quantum integrable systems associated with the quiver varieties. The elliptic version of stable envelopes (ESE) [1] for the equivariant elliptic cohomology $E_T(X)$ is more fascinating. They depend on both the equivariant and the Kähler parameters and provide the elliptic dynamical R -matrix as their transition matrix. There these two parameters play a role of the spectral and the dynamical parameters, respectively.

An typical example is the case $X = T^*fl$, the cotangent bundle to the partial flag variety. The ESE's for $E_T(T^*fl)$ can be identified with the elliptic weight functions of the A type appearing in the elliptic hypergeometric integral solutions of the elliptic q -KZ equation [28, 29, 43] and their transition matrices coincide with the known elliptic dynamical R -matrices of type A [12, 17, 18, 24]. In [28, 29], such identification was carried out by using the vertex operators defined as intertwining operators of the $U_{q, p}(\widehat{\mathfrak{sl}}_N)$ -modules w.r.t the standard comultiplication given in terms of the L -operator instead of the Drinfeld comultiplication. Basically such vertex operators are operator valued integrals, the so called screened vertex operators, possessing a basic elliptic weight function i.e. elliptic stable envelope, as an integration kernel. Then a composition of them produces more general ESE corresponding to a general configuration in T^*fl via the shuffle product formula [29]. Such structure can indeed be regarded as an inheritance from the L -operator of $U_{q, p}(\widehat{\mathfrak{sl}}_N)$, whose Gauss components are given by the so called half currents, i.e. integrals of the elliptic currents of $U_{q, p}(\widehat{\mathfrak{sl}}_N)$ with the same basic elliptic weight function as an

integration kernel [22]. This is consistent to a picture that the stable envelope gives a half of the Gauss decomposition of the L -operator.

In addition, for the same case $X = T^*fl$, it was shown that the vertex operators yield a K-theoretic vertex functions [33, 41, 42] as a vacuum expectation value of a composition of the vertex operators [30]. The vertex function is a generating function of the counting of quasi maps from X to \mathbb{P}^1 . It is given by a certain q -hypergeometric integral containing the elliptic weight function inside the integral and satisfies the q -KZ equation. The monodromy matrix of it is then given by a transition matrix of the weight functions. In [1], such elliptic weight functions are called the pole subtraction elliptic stable envelopes.

Motivated by such nice relationship between representation theory of $U_{q,p}(\widehat{\mathfrak{g}})$ and the elliptic stable envelopes, we in this paper propose new vertex operators of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ by combining representations of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ [32] and the elliptic stable envelopes for the equivariant elliptic cohomology $E_T(\mathcal{M}(n, r))$ of the instanton moduli space $\mathcal{M}(n, r)$ constructed in [4, 44]. We construct both the type I and the type II dual vertex operators as screened vertex operators whose integration kernel is given by the ESE for $E_T(\mathcal{M}(n, r))$. We then make several checks on their consistency. One of them is the shuffle product formula of ESE's. We derive it by considering a composition of the vertex operators and check its consistency with the one obtained geometrically in [2]. We also construct K-theoretic vertex functions for $\mathcal{M}(n, r)$ as expectation value of compositions of the vertex operators and show their coincidence with the formula obtained in [4, 45]. Moreover, thus obtained vertex operators, in particular for the Hilbert scheme $\text{Hilb}^n(\mathbb{C}^2)$, allow us to construct a L -operator $L^+(u)$ on a certain representation, which satisfies the $RLL = LLR^*$ relation. Here R and R^* are elliptic dynamical R -matrices given as transition matrices of ESE's for $E_T(\text{Hilb}^n(\mathbb{C}^2))$ appearing in the type I and the type II dual vertex operators, respectively. Hence they should be the elliptic instanton R -matrices. We then consider a universal form $\mathcal{L}^+(u)$ of $L^+(u)$ and show that these vertex operators are intertwining operators of the $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ -modules w.r.t the standard comultiplication defined by $\mathcal{L}^+(u)$.

This paper is organized as follows. Sec.2 is a review of the elliptic quantum toroidal algebra $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ including the Hopf algebroid structure and representations of the level $(1, N)$ and $(0, 1)$. In Sec.3, we summarize known results on the ESE for $E_T(\mathcal{M}(n, r))$. In Sec.4, we propose a new construction of the type I and the type II dual vertex operators of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ and check the shuffle product formula of ESE's. Sec.5 is devoted to a study of the K-theoretic vertex functions for $\mathcal{M}(n, r)$. In Sec.6, we derive exchange relations of the vertex operators. Finally, in Sec.7, we construct a L -operator $L^+(u)$, which satisfies the $RLL = LLR^*$ relation. The exchange relations between the vertex operators and $L^+(u)$ are given. Defining a new comultiplication

Δ , we show that our vertex operators are intertwining operators of the $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ -modules w.r.t. Δ . For comparison, we list in Appendix A a summary of the vertex operators w.r.t. the Drinfeld comultiplication Δ^D obtained in [32].

2 Elliptic Quantum Toroidal Algebra $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$

We review the elliptic quantum toroidal algebra $\mathcal{U}_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ introduced in [32]. Parameters $t_1, t_2, \hbar = t_1 t_2$ in this paper correspond to $q^{-1}, t, t/q$ in [32], respectively. We assume p, t_1, t_2 being generic complex number and $|p| < 1$.

2.1 Definition

Let us consider the Heisenberg algebras generated by $c, \Lambda_0, c^\perp, \Lambda_0^\perp, P, Q$ satisfying the commutation relations

$$[c, \Lambda_0] = 1 = [c^\perp, \Lambda_0^\perp] = [P, Q], \quad (2.1)$$

the others are zero. We set $\gamma = \hbar^{c/2}, C = \hbar^{c^\perp/2}, \mathfrak{z}^* = \hbar^P$ and $\mathfrak{z} = \mathfrak{z}^* C^2 = \hbar^{P+c^\perp}$. We call \mathfrak{z}^* the dynamical parameter. Let \mathbb{F} be the field of meromorphic functions of \mathfrak{z} and \mathfrak{z}^* . We have

$$g(\mathfrak{z}, \mathfrak{z}^*) e^{\Lambda_0^\perp - Q} = e^{\Lambda_0^\perp - Q} g(\mathfrak{z}, \mathfrak{z}^* \hbar^{-1}), \quad g(\mathfrak{z}, \mathfrak{z}^*) e^{-\Lambda_0^\perp} = e^{-\Lambda_0^\perp} g(\mathfrak{z} \hbar^{-1}, \mathfrak{z}^*) \quad \forall g(\mathfrak{z}, \mathfrak{z}^*) \in \mathbb{F}.$$

Set

$$\begin{aligned} \kappa_m &= -(1 - t_1^m)(1 - t_2^m)(1 - \hbar^{-m}), \\ G^\pm(z) &= (1 - t_1^{\mp 1} z)(1 - t_2^{\mp 1} z)(1 - \hbar^{\pm 1} z). \end{aligned}$$

Definition 2.1. *The elliptic quantum toroidal algebra $\mathcal{U} = U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ is a topological associative algebra over $\mathbb{F}[[p]]$ generated by $\alpha_m, x_n^\pm, (m \in \mathbb{Z} \setminus \{0\}, n \in \mathbb{Z})$ and $C, \gamma^{1/2}$. Let $x^\pm(z), \psi^\pm(z)$ be the following generating functions¹.*

$$\begin{aligned} x^\pm(z) &:= \sum_{n \in \mathbb{Z}} x_n^\pm z^{-n}, \\ \psi^+(z) &:= C^{-1} \exp \left(- \sum_{m>0} \frac{p^m}{1-p^m} \alpha_{-m} (\gamma^{-1/2} z)^m \right) \exp \left(\sum_{m>0} \frac{1}{1-p^m} \alpha_m (\gamma^{-1/2} z)^{-m} \right), \\ \psi^-(z) &:= C \exp \left(- \sum_{m>0} \frac{1}{1-p^m} \alpha_{-m} (\gamma^{1/2} z)^m \right) \exp \left(\sum_{m>0} \frac{p^m}{1-p^m} \alpha_m (\gamma^{1/2} z)^{-m} \right). \end{aligned}$$

¹Our C and $x^\pm(z)$ here are $\psi_0^- = (\psi_0^+)^{-1}$ and $x^\pm(\gamma^{1/2} z)$ in [32], respectively.

We call them the elliptic currents. The defining relations are given by

$$C, \gamma^{1/2} : \text{central}, \quad (2.2)$$

$$[\alpha_m, \alpha_n] = -\frac{\kappa_m}{m}(\gamma^m - \gamma^{-m})\gamma^{-m} \frac{1-p^m}{1-p^{*m}} \delta_{m+n,0}, \quad (2.3)$$

$$[\alpha_m, x^+(z)] = -\frac{\kappa_m}{m} \frac{1-p^m}{1-p^{*m}} \gamma^{-m} z^m x^+(z) \quad (m \neq 0). \quad (2.4)$$

$$[\alpha_m, x^-(z)] = \frac{\kappa_m}{m} z^m x^-(z) \quad (m \neq 0), \quad (2.5)$$

$$[x^+(z), x^-(w)] = -\frac{(1-t_1)(1-t_2)}{(1-\hbar)} \left(\delta(\gamma^{-1}z/w)\psi^+(\gamma^{1/2}w) - \delta(\gamma z/w)\psi^-(\gamma^{-1/2}w) \right) \quad (2.6)$$

$$z^3 G^+(w/z)g(w/z; p^*)x^+(z)x^+(w) = -w^3 G^+(z/w)g(z/w; p^*)x^+(w)x^+(z), \quad (2.7)$$

$$z^3 G^-(w/z)g(w/z; p)^{-1}x^-(z)x^-(w) = -w^3 G^-(z/w)g(z/w; p)^{-1}x^-(w)x^-(z), \quad (2.8)$$

$$g(w/z; p^*)g(u/w; p^*)g(u/z; p^*) \left(\frac{w}{u} + \frac{w}{z} - \frac{z}{w} - \frac{u}{w} \right) x^+(z)x^+(w)x^+(u) \\ + \text{permutations in } z, w, u = 0, \quad (2.9)$$

$$g(w/z; p)^{-1}g(u/w; p)^{-1}g(u/z; p)^{-1} \left(\frac{w}{u} + \frac{w}{z} - \frac{z}{w} - \frac{u}{w} \right) x^-(z)x^-(w)x^-(u) \\ + \text{permutations in } z, w, u = 0, \quad (2.10)$$

where we set $p^* = p\gamma^{-2}$ and

$$g(z; s) = \exp \left(\sum_{m>0} \frac{\kappa_m}{m} \frac{s^m}{1-s^m} z^m \right) \in \mathbb{C}[[z]] \quad (2.11)$$

for $s = p, p^*$.

We treat these relations as formal Laurent series in z, w and u . All the coefficients in z, w, u are well defined in the p -adic topology [13, 27]. The algebra $\mathcal{U} = U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ is an extension of the one given in [32] by introducing the dynamical parameter \mathfrak{z}^* and the field \mathbb{F} .

It is also convenient to set

$$\alpha'_m = \frac{1-p^{*m}}{1-p^m} \gamma^m \alpha_m \quad (m \in \mathbb{Z}_{\neq 0}).$$

We have

$$[\alpha'_m, \alpha'_n] = -\frac{\kappa_m}{m}(\gamma^m - \gamma^{-m})\gamma^m \frac{1-p^{*m}}{1-p^m} \delta_{m+n,0}. \quad (2.12)$$

We have the following duality.

Proposition 2.2.

$$U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}}) \cong U_{t_1^{-1}, t_2^{-1}, p^*}(\mathfrak{gl}_{1, \text{tor}})$$

by

$$x_n^+ \mapsto x_n^-, \quad x_n^- \mapsto x_n^+, \quad \alpha_m \mapsto \alpha'_m, \quad \gamma^{1/2} \mapsto \gamma^{-1/2}.$$

Note that in $U_{t_1^{-1}, t_2^{-1}, p^*}(\mathfrak{gl}_{1, tor})$, one has

$$(p^*)^* = p^*(\gamma^{-1})^{-2} = p.$$

Remark. In [32], it has been shown that the elliptic currents $x^+(z)$ and $x^-(z)$ in $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ are screening currents of the Jordan quiver W -algebra $W_{p, p^*}(\Gamma(\hat{A}_0))$ [20]. The latter W -algebra is self-dual under the exchange between $x^+(z)$ and $x^-(z)$. The duality (2.2) is a generalization of the Feigin-Frenkel duality of the (deformed) W -algebras to the affine quiver W -algebras [8, 14].

Remark. For \mathcal{U} -modules, on which the central element $\gamma^{1/2}$ takes a complex value, we regard $p, p^* = p\gamma^{-2}$ as a generic complex number with $|p| < 1, |p^*| < 1$. Then we have

$$g(z; p) = \frac{(pt_1^{-1}z; p)_\infty (pt_2^{-1}z; p)_\infty (p\hbar z; p)_\infty}{(pt_1z; p)_\infty (pt_2z; p)_\infty (p\hbar^{-1}z; p)_\infty},$$

where we set

$$(z; p)_\infty = \prod_{n=0}^{\infty} (1 - zp^n) \quad |z| < 1.$$

Then one can rewrite, for example, (2.7)-(2.8) in the sense of analytic continuation as follows.

$$\begin{aligned} x^+(z)x^+(w) &= g_{\theta^*}(w/z)x^+(w)x^+(z), \\ x^-(z)x^-(w) &= g_\theta(z/w)x^-(w)x^-(z). \end{aligned}$$

Here we set

$$\theta(z) = -z^{-1/2}(z; p)_\infty (p/z; p)_\infty, \quad \theta^*(z) = \theta(z)|_{p \rightarrow p^*}, \quad (2.13)$$

$$g_\theta(z) = \frac{\theta(t_1z)\theta(t_2z)\theta(\hbar^{-1}z)}{\theta(t_1^{-1}z)\theta(t_2^{-1}z)\theta(\hbar z)}, \quad g_{\theta^*}(z) = g_\theta(z)|_{\theta \rightarrow \theta^*}. \quad (2.14)$$

Note that

$$g_\theta(z^{-1}) = g_\theta(z)^{-1}, \quad g_\theta(pz) = g_\theta(z).$$

2.2 Hopf algebroid structure

For $F(\mathfrak{z}, p) \in \mathbb{F}[[p]]$, let $\tilde{\otimes}$ denote the usual tensor product with the following extra condition [6, 21, 26]

$$F(\mathfrak{z}^*, p^*)a\tilde{\otimes}b = a\tilde{\otimes}F(\mathfrak{z}, p)b. \quad (2.15)$$

Here $F(\mathfrak{z}^*, p^*)$ denotes the same F with replacing \mathfrak{z} and p by \mathfrak{z}^* and $p^* = p\gamma^{-2}$, respectively.

Define two moment maps $\mu_l, \mu_r : \mathbb{F}[[p]] \rightarrow \mathcal{U}$ by

$$\mu_l(F(\mathfrak{z}, p)) = F(\mathfrak{z}, p), \quad \mu_r(F(\mathfrak{z}, p)) = F(\mathfrak{z}^*, p^*).$$

Let $\gamma_{(1)} = \gamma \tilde{\otimes} 1, \gamma_{(2)} = 1 \tilde{\otimes} \gamma$ and $p_{(i)}^* = p\gamma_{(i)}^{-2}$ ($i = 1, 2$). Let us define two algebra homomorphisms $\Delta^D : \mathcal{U} \rightarrow \mathcal{U} \tilde{\otimes} \mathcal{U}$ and $\varepsilon^D : \mathcal{U} \rightarrow \mathbb{C}$ by

$$\Delta^D(\gamma^{\pm 1/2}) = \gamma^{\pm 1/2} \tilde{\otimes} \gamma^{\pm 1/2}, \quad (2.16)$$

$$\Delta^D(\psi^\pm(z)) = \psi^\pm(\gamma_{(2)}^{\mp 1/2} z) \tilde{\otimes} \psi^\pm(\gamma_{(1)}^{\pm 1/2} z), \quad (2.17)$$

$$\Delta^D(x^+(z)) = 1 \tilde{\otimes} x^+(\gamma_{(1)}^{-1/2} z) + x^+(\gamma_{(2)}^{1/2} z) \tilde{\otimes} \psi^-(\gamma_{(1)}^{-1/2} z), \quad (2.18)$$

$$\Delta^D(x^-(z)) = x^-(\gamma_{(2)}^{-1/2} z) \tilde{\otimes} 1 + \psi^+(\gamma_{(2)}^{-1/2} z) \tilde{\otimes} x^-(\gamma_{(1)}^{1/2} z), \quad (2.19)$$

$$\Delta^D(\mu_l(F(\mathfrak{z}, p))) = \mu_l(F(\mathfrak{z}, p)) \tilde{\otimes} 1, \quad \Delta^D(\mu_r(F(\mathfrak{z}, p))) = 1 \tilde{\otimes} \mu_r(F(\mathfrak{z}, p)), \quad (2.20)$$

$$\varepsilon^D(\gamma^{1/2}) = \varepsilon^D(C) = 1, \quad \varepsilon^D(\psi^\pm(z)) = 1, \quad \varepsilon^D(x^\pm(z)) = 0, \quad (2.21)$$

$$\varepsilon^D(\mu_l(F(\mathfrak{z}, p))) = \varepsilon^D(\mu_r(F(\mathfrak{z}, p))) = F(\mathfrak{z}, p). \quad (2.22)$$

Δ^D is called the Drinfeld comultiplication. Then we have

Proposition 2.3. *The maps ε^D and Δ^D satisfy*

$$(\Delta^D \tilde{\otimes} \text{id}) \circ \Delta^D = (\text{id} \tilde{\otimes} \Delta^D) \circ \Delta^D, \quad (2.23)$$

$$(\varepsilon^D \tilde{\otimes} \text{id}) \circ \Delta^D = \text{id} = (\text{id} \tilde{\otimes} \varepsilon^D) \circ \Delta^D. \quad (2.24)$$

We also define an algebra anti-homomorphism $S^D : \mathcal{U} \rightarrow \mathcal{U}$ by

$$S^D(\gamma^{1/2}) = \gamma^{-1/2},$$

$$S^D(\psi^\pm(z)) = \psi^\pm(z)^{-1},$$

$$S^D(x^+(z)) = -x^+(z)\psi^-(z)^{-1},$$

$$S^D(x^-(z)) = -\psi^+(z)x^-(z),$$

$$S^D(\mu_l(F(\mathfrak{z}, p))) = \mu_r(F(\mathfrak{z}, p)), \quad S^D(\mu_r(F(\mathfrak{z}, p))) = \mu_l(F(\mathfrak{z}, p)).$$

Then one can check the following.

Proposition 2.4.

$$m \circ (\text{id} \tilde{\otimes} S^D) \circ \Delta^D(a) = \mu_l(\varepsilon^D(a)1) \quad \forall a \in \mathcal{U},$$

$$m \circ (S^D \tilde{\otimes} \text{id}) \circ \Delta^D(a) = \mu_r(\varepsilon^D(a)1).$$

These Propositions indicate that $(\mathcal{U}, \Delta^D, \varepsilon^D, \mu_l, \mu_r, S^D)$ is a Hopf algebroid [6, 21, 25, 26].

In Sec.7, we introduce a new comultiplication Δ defined by the L -operator.

2.3 Representations of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$

Let \mathcal{V} be a \mathcal{U} -module. For $(k, l) \in \mathbb{C}^2$, we say that \mathcal{V} has level (k, l) , if the central elements γ and C act as

$$\gamma \cdot \xi = \hbar^{k/2} \xi, \quad C \cdot \xi = \hbar^{l/2} \xi \quad \forall \xi \in \mathcal{V}.$$

In the rest of this paper, we regard $p, p^* = p\gamma^{-2}$ as a generic complex number with $|p|, |p^*| < 1$.

2.3.1 The level- $(1, N)$ representation

Let us introduce one more Heisenberg algebra generated by h, α with

$$[h, \alpha] = 1.$$

Define for $v \in \mathbb{C}^\times$

$$|0\rangle_v^{(1, N)} := v^\alpha e^{\Lambda_0} e^{N\Lambda_0^\perp} \mathbf{1}. \quad (2.25)$$

We assume $\gamma^{1/2} \cdot \mathbf{1} = C \cdot \mathbf{1} = e^{\pm h} \cdot \mathbf{1} = \mathbf{1}$ and $e^Q \cdot \mathbf{1} = e^Q \mathbf{1}$. One has

$$e^{\pm h} x^{\pm c^\perp} \cdot |0\rangle_v^{(1, N)} = v^{\pm 1} x^{\pm N} |0\rangle_v^{(1, N)}, \quad \gamma \cdot |0\rangle_v^{(1, N)} = \hbar^{1/2} |0\rangle_v^{(1, N)}, \quad C^{\pm 1} \cdot |0\rangle_v^{(1, N)} = \hbar^{\pm N/2} |0\rangle_v^{(1, N)}.$$

Let $\tilde{\mathcal{F}}_v^{(1, N)} = \mathbb{C}[\alpha_{-m} \ (m > 0)]|0\rangle_v^{(1, N)}$ be a Fock space on which the Heisenberg subalgebra $\{\alpha_m \ (m \in \mathbb{Z}_{\neq 0})\}$ acts as

$$\begin{aligned} \alpha_{-m} \cdot |0\rangle_v^{(1, N)} &= 0, \\ \alpha_{-m} \cdot \xi &= \alpha_{-m} \xi, \quad \alpha_m \cdot \xi = -\frac{\kappa_m}{m} (1 - \hbar^{-m}) \frac{1 - p^m}{1 - p^{*m}} \frac{\partial}{\partial \alpha_{-m}} \xi \end{aligned}$$

for $m > 0$, $\xi \in \tilde{\mathcal{F}}_v^{(1, N)}$. Note that $p^* = p\hbar^{-1}$ on $\tilde{\mathcal{F}}_v^{(1, N)}$. We set

$$\mathcal{F}_v^{(1, N)} = \bigoplus_{m \in \mathbb{Z}} \tilde{\mathcal{F}}_v^{(1, N)} \otimes e^{mQ}.$$

Theorem 2.5. *The following assignment gives a level $(1, N)$ representation of \mathcal{U} on $\mathcal{F}_v^{(1, N)}$.*

$$x^+(z) = e^{h(z^{-1}\hbar^{1/2})^{c^\perp}} \exp \left\{ -\sum_{n>0} \frac{\hbar^{n/2}}{1 - \hbar^n} \alpha_{-n} z^n \right\} \exp \left\{ \sum_{n>0} \frac{\hbar^{n/2}}{1 - \hbar^n} \alpha_n z^{-n} \right\}, \quad (2.26)$$

$$x^-(z) = e^{-h(z^{-1}\hbar^{1/2})^{-c^\perp}} \exp \left\{ \sum_{n>0} \frac{\hbar^{n/2}}{1 - \hbar^n} \alpha'_{-n} z^n \right\} \exp \left\{ -\sum_{n>0} \frac{\hbar^{n/2}}{1 - \hbar^n} \alpha'_n z^{-n} \right\}, \quad (2.27)$$

$$\psi^+(\hbar^{1/4}z) = \hbar^{-c^\perp/2} \exp \left\{ -\sum_{n>0} \frac{p^n}{1 - p^n} \alpha_{-n} z^n \right\} \exp \left\{ \sum_{n>0} \frac{1}{1 - p^n} \alpha_n z^{-n} \right\}, \quad (2.28)$$

$$\psi^-(\hbar^{-1/4}z) = \hbar^{c^\perp/2} \exp \left\{ -\sum_{n>0} \frac{1}{1 - p^n} \alpha_{-n} z^n \right\} \exp \left\{ \sum_{n>0} \frac{p^n}{1 - p^n} \alpha_n z^{-n} \right\}. \quad (2.29)$$

This is a reformulation of the same representation given in [32] by introducing the zero-modes operators $c, \Lambda_0, c^\perp, \Lambda_0^\perp$ and h, α .

2.3.2 The level-(0,1) representation

We next consider the level-(0,1) representation of \mathcal{U} called the q -Fock representation.

For $u \in \mathbb{C}^\times$, let \mathcal{F}_u be a vector space spanned by $|\lambda\rangle_u$ ($\lambda \in \mathcal{P}$), where

$$\mathcal{P} = \{\lambda = (\lambda_1, \lambda_2, \dots) \mid \lambda_i \geq \lambda_{i+1}, \lambda_i \in \mathbb{Z}_{\geq 0}, \lambda_l = 0 \text{ for sufficiently large } l\}. \quad (2.30)$$

We denote by $\ell(\lambda)$ the length of $\lambda \in \mathcal{P}$ i.e. $\lambda_{\ell(\lambda)} > 0$ and $\lambda_{\ell(\lambda)+1} = 0$. We also set $|\lambda| = \sum_{i \geq 1} \lambda_i$ and denote by λ' the conjugate of λ .

Theorem 2.6. *The following action gives a level (0,1) representation of \mathcal{U} on \mathcal{F}_u . We denote this by $\mathcal{F}_u^{(0,1)}$.*

$$\gamma^{1/2}|\lambda\rangle_u = |\lambda\rangle_u, \quad (2.31)$$

$$x^+(z)|\lambda\rangle_u = a^+(p) \sum_{i=1}^{\ell(\lambda)+1} A_{\lambda,i}^+(p) \delta(u_i/z) |\lambda + \mathbf{1}_i\rangle_u, \quad (2.32)$$

$$x^-(z)|\lambda\rangle_u = a^-(p) \sum_{i=1}^{\ell(\lambda)} A_{\lambda,i}^-(p) \delta(t_1 u_i/z) |\lambda - \mathbf{1}_i\rangle_u, \quad (2.33)$$

$$\psi^+(z)|\lambda\rangle_u = B_\lambda^+(u/z; p) |\lambda\rangle_u, \quad (2.34)$$

$$\psi^-(z)|\lambda\rangle_u = B_\lambda^-(z/u; p) |\lambda\rangle_u, \quad (2.35)$$

where we set

$$a^+(p) = (1-t) \frac{(p\hbar; p)_\infty (p/t_2; p)_\infty}{(p; p)_\infty (p/q; p)_\infty}, \quad (2.36)$$

$$a^-(p) = (1-t^{-1}) \frac{(p/\hbar; p)_\infty (pt_2; p)_\infty}{(p; p)_\infty (pq; p)_\infty}, \quad (2.37)$$

$$A_{\lambda,i}^+(p) = \prod_{j=1}^{i-1} \frac{\theta(t_2 u_i/u_j) \theta(\hbar^{-1} u_i/u_j)}{\theta(t_1^{-1} u_i/u_j) \theta(u_i/u_j)}, \quad (2.38)$$

$$A_{\lambda,i}^-(p) = \prod_{j=i+1}^{\ell(\lambda)} \frac{\theta(\hbar^{-1} u_j/u_i)}{\theta(u_j/u_i)} \prod_{j=i+1}^{\ell(\lambda)+1} \frac{\theta(t_2 u_j/u_i)}{\theta(t_1^{-1} u_j/u_i)}, \quad (2.39)$$

$$B_\lambda^+(u/z; p) = \prod_{i=1}^{\ell(\lambda)} \frac{\theta(t_2^{-1} u_i/z)}{\theta(t_1 u_i/z)} \prod_{i=1}^{\ell(\lambda)+1} \frac{\theta(\hbar u_i/z)}{\theta(u_i/z)}, \quad (2.40)$$

$$B_\lambda^-(z/u; p) = \prod_{i=1}^{\ell(\lambda)} \frac{\theta(t_2 z/u_i)}{\theta(t_1^{-1} z/u_i)} \prod_{i=1}^{\ell(\lambda)+1} \frac{\theta(\hbar^{-1} z/u_i)}{\theta(z/u_i)}. \quad (2.41)$$

Note that we use a slightly different notation (2.13) for the theta function from the one in [32].

Remark. In [32] it is conjectured that the level-(0,1) representation in Theorem 2.6 gives a geometric action of \mathcal{U} on the equivariant elliptic cohomology of the Hilbert schemes $\bigoplus_n E_T(\text{Hilb}^n(\mathbb{C}^2))$ by identifying $|\lambda\rangle_u$ with the fixed point class $[\lambda]$ in $\bigoplus_n E_T(\text{Hilb}^n(\mathbb{C}^2))$. The class $[\lambda]$ with $|\lambda| = n$ should be realized in terms of the elliptic stable envelopes $\text{Stab}_{\mathfrak{e}}$ for $E_T(\text{Hilb}^n(\mathbb{C}^2))$ constructed in [44] (see Sec.3.2 below) by

$$[\lambda] = \sum_{\substack{\mu \in \mathcal{P} \\ |\mu| = n}} \text{Stab}_{\mathfrak{e}}^{-1}(\mu)|_{\lambda} \text{Stab}_{\mathfrak{e}}(\mu)$$

in the similar way to the action of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on the equivariant elliptic cohomology of the cotangent bundle to partial flag varieties [29].

3 Elliptic Stable Envelopes for $E_T(\mathcal{M}(n, r))$

In this section we summarize basic facts on the instanton moduli space $\mathcal{M}(n, r)$ and the elliptic stable envelopes for the equivariant elliptic cohomology $E_T(\mathcal{M}(n, r))$ following [1, 4, 44].

3.1 The instanton moduli space $\mathcal{M}(n, r)$

Let $\mathcal{M}(n, r)$ be the moduli space of framed rank r torsion free sheaves \mathcal{S} on \mathbb{P}^2 with $c_2(\mathcal{S}) = n$. A framing of a sheaf \mathcal{S} means a choice of isomorphism

$$\mathcal{S}|_{\mathbb{P}} \rightarrow \mathcal{O}_{L_{\infty}}^{\oplus r},$$

where $L_{\infty} \subset \mathbb{P}^2$ is chosen as a line at infinity of a plane in \mathbb{P}^2 . The space $\mathcal{M}(n, r)$ is isomorphic to the Nakajima quiver variety associated to the quiver consisting of one vertex and one loop with the vertex dimension n and the framing dimension r , and known as the Atiyah-Drinfeld-Hitchin-Manin (ADHM) instanton moduli space. One has a natural action of $G = GL(r) \times GL(2)$ on $\mathcal{M}(n, r)$. Let T be the maximal torus of G and set

$$A = T \cap GL(r).$$

The parameters t_1, t_2 are identified with the generators of the character group of T/A . Note also that the rank 1 case is isomorphic to the Hilbert scheme of n -points on \mathbb{C}^2 .

$$\mathcal{M}(n, 1) \cong \text{Hilb}^n(\mathbb{C}^2).$$

We also have the following isomorphism for the fixed points w.r.t A .

$$\mathcal{M}(n, r)^A = \coprod_{n_1 + \dots + n_r = n} \text{Hilb}^{n_1}(\mathbb{C}^2) \times \dots \times \text{Hilb}^{n_r}(\mathbb{C}^2).$$

Let us consider the case $r = 1$, the Hilbert scheme $\mathcal{H}_n = \text{Hilb}^n(\mathbb{C}^2)$, $A = \mathbb{C}^\times$. We denote the coordinate on A by u such that

$$t_1 = u\hbar^{1/2}, \quad t_2 = u^{-1}\hbar^{1/2}.$$

The fixed points $\mathcal{H}_n^T = \mathcal{H}_n^A$ is a finite set labeled by partitions of n . Let

$$\mathcal{P}_n = \{ \lambda \in \mathcal{P} \mid |\lambda| = n \}. \quad (3.1)$$

We regard $\lambda \in \mathcal{P}_n$ as a Young diagram with n boxes. For a box $\square = (i, j) \in \lambda$, we define

$$\begin{aligned} c_\square &:= i - j, & h_\square &:= i + j - 2, \\ \rho_\square &= c_\square - \epsilon h_\square \end{aligned}$$

with $0 < \epsilon \ll 1$. We introduce a canonical ordering on the n boxes of λ by

$$a < b \Leftrightarrow \rho_a < \rho_b \quad a, b \in \lambda$$

and define a bijection $\iota : \lambda \rightarrow [1, n]$ if $a \in \lambda$ is the $\iota(a)$ -th box in this order. In the following we often denote the box a by $\square_{\iota(a)}$ or simply $\iota(a)$.

Let us consider the following presentation of the equivariant K-theory of \mathcal{H}_n .

$$\text{K}_T(\mathcal{H}_n) = \mathbb{Z}[x_1^\pm, \dots, x_n^\pm, t_1^\pm, t_2^\pm]^{\mathfrak{S}_n} / R,$$

where \mathfrak{S}_n denotes the symmetrization in x_a 's, and R the ideal of Laurent polynomials vanishing at all fixed points in \mathcal{H}_n^T . We often loosely use x_a as $x_{\iota(a)}$ for $a \in \lambda$ and vice versa. For $\square = (i, j) \in \lambda$, we set

$$\varphi_\square^\lambda = t_1^{-(j-1)} t_2^{-(i-1)} \in \text{K}_T(pt).$$

The restriction of a K-theory class $f(x_1, \dots, x_n, t_1, t_2)$ to a fixed point labeled by $\lambda \in \mathcal{P}_n$ is given by

$$i_\lambda^* f(x_1, \dots, x_n, t_1, t_2) = f(\varphi_{\square_1}^\lambda, \dots, \varphi_{\square_n}^\lambda, t_1, t_2).$$

Here $i_\lambda : \lambda \rightarrow \mathcal{H}_n$ denotes the canonical inclusion of a fixed point.

Let \mathcal{V} be the rank n tautological bundle on \mathcal{H}_n . We present \mathcal{V} as

$$\mathcal{V} = x_1 + \dots + x_n$$

regarding x_1, \dots, x_n as the Chern roots of \mathcal{V} . The line bundle

$$\mathcal{O}(1) = \det \mathcal{V} = x_1 \cdots x_n$$

generates the Picard group $\text{Pic}(\mathcal{H}_n) = \mathbb{Z}$.

In general, fixed points in $\mathcal{M}(n, r)$ are labeled by a r -tuple partition $\boldsymbol{\lambda} = (\lambda^{(1)}, \dots, \lambda^{(r)})$ with $|\boldsymbol{\lambda}| = \sum_{i=1}^r |\lambda^{(i)}| = n$. We denote the Chern roots of the rank n tautological bundle \mathcal{V} on $\mathcal{M}(n, r)$ by $\mathbf{x} = (x_1, \dots, x_n)$ and a coordinate of $A = (\mathbb{C}^\times)^r$ by $\mathbf{u} = (u_1, \dots, u_r)$. We define a canonical ordering on the n boxes of $\boldsymbol{\lambda}$ by extending the one for each partition $\lambda^{(i)}$ with adding the following condition, for $i < j$

$$a < b \Leftrightarrow a \in \lambda^{(i)}, b \in \lambda^{(j)}.$$

Note that this additional condition is realized as a condition $\rho_a < \rho_b$ if one extends the definition of ρ_a to

$$\rho_a := c_a - \epsilon h_a + u_i \quad \text{for } a \in \lambda^{(i)} \quad (3.2)$$

and assume $|u_1| \ll \dots \ll |u_r|$. In the following we use this convention for $\boldsymbol{\lambda}$.

3.2 Elliptic stable envelopes

The elliptic stable envelopes for $E_T(\mathcal{H}_n)$ was constructed in [44]. Let $T^{1/2} \in K_T(\mathcal{H}_n)$ be a polarization of \mathcal{H}_n satisfying

$$T\mathcal{H}_n = T^{1/2} + \hbar(T^{1/2})^\vee \in K_T(\mathcal{H}_n).$$

The term $\hbar(T^{1/2})^\vee$ is called the opposite polarization. For $\lambda \in \mathcal{P}_n$, let us set

$$S_\lambda^{Ell}(x_1, \dots, x_n, u) := \frac{\prod_{\substack{a, b \in \lambda \\ \rho_a + 1 < \rho_b}} \theta(t_1 x_a / x_b) \prod_{\substack{a, b \in \lambda \\ \rho_a + 1 > \rho_b}} \theta(t_2 x_b / x_a) \prod_{\substack{a \in \lambda \\ \rho_a \leq 0}} \theta(x_a / u) \prod_{\substack{a \in \lambda \\ \rho_a > 0}} \theta(\hbar u / x_a)}{\prod_{\substack{a, b \in \lambda \\ \rho_a < \rho_b}} \theta(x_a / x_b) \theta(\hbar x_a / x_b)}. \quad (3.3)$$

Let \mathbf{t} be a λ -tree, see Definition 1 in Section 4.2 of [44]. Namely, \mathbf{t} is a rooted tree in a Young diagram λ with vertices corresponding to boxes of λ , edges connecting only adjacent boxes and the root at the box $(1, 1) \in \lambda$. Let us set (formula (54) in [44]):

$$W_{T^{1/2}}^{Ell}(\mathbf{t}; x_1, \dots, x_n, u, \mathfrak{z}) = (-1)^{\kappa_{\mathbf{t}}} \phi\left(\frac{x_r}{u}, \mathfrak{z}^n (t_1 t_2)^{\nu_r}\right) \prod_{e \in \mathbf{t}} \phi\left(\frac{x_{h(e)} \varphi_{t(e)}^\lambda}{x_{t(e)} \varphi_{h(e)}^\lambda}, \mathfrak{z}^{w_e} (t_1 t_2)^{\nu_e}\right)$$

with

$$\phi(x, y) = \frac{\theta(xy)}{\theta(x)\theta(y)},$$

where the product runs over the edges of the tree \mathbf{t} and $h(e) \in \lambda$, $t(e) \in \lambda$ denote the head and tail box of the edge e . And, $\kappa_{\mathbf{t}}, w_e, \nu_e \in \mathbb{Z}$ are certain integers computed from the tree in a

combinatorial way, we refer to Sections 4.2-4.5 in [44] for definitions of these integers. We also identify the dynamical parameter \mathfrak{z} with the Kähler parameter in $E_T(\mathcal{H}_n)$ [1] in the same way as in [29]. Finally, let Υ_λ be the set of λ -trees without L-shaped subgraphs, see section 4.6 in [44]. Then, we have:

Theorem 3.1. [44] *The elliptic stable envelope of a fixed point $\lambda \in \mathcal{H}_n^A$ is given by*

$$\text{Stab}_{\mathfrak{C}, T^{1/2}}(\lambda; \mathfrak{z}) = \text{Sym} \left(S_\lambda^{Ell}(x_1, \dots, x_n, u) \sum_{\delta \in \Upsilon_\lambda} W_{T^{1/2}}^{Ell}(\mathfrak{t}_\delta; x_1, \dots, x_n, u; \mathfrak{z}) \right), \quad (3.4)$$

where the symbol Sym stands for symmetrization over x_1, \dots, x_n . The chamber \mathfrak{C} is taken as a stability condition $t_1/t_2 > 0$.

The elliptic stable envelope for $E_T(\mathcal{M}(n, r))$ is constructed by taking the shuffle product [2] of those for the Hilbert schemes. Let λ', λ'' be two partitions with $|\lambda'| = n', |\lambda''| = n''$ and consider the elliptic stable envelopes $\text{Stab}_{T^{1/2}, \mathfrak{C}'}(\lambda'; \mathfrak{z})$ and $\text{Stab}_{T^{1/2}, \mathfrak{C}''}(\lambda''; \mathfrak{z})$ for $E_T(\mathcal{H}_{n'})$ and $E_T(\mathcal{H}_{n''})$ with the equivariant parameters u_1, u_2 , respectively. Here one takes $\mathfrak{C}' = \mathfrak{C}''$ as $t_1/t_2 > 0$. We take the canonical ordering on the $n' + n''$ boxes in the double partition (λ', λ'') as defined in Sec.3.1. Let us consider $\text{Stab}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z})$ given by

$$\begin{aligned} & \text{Stab}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z}) \\ & := \text{Sym}_{\{x_a\}_{a \in (\lambda', \lambda'')}} \left(\prod_{a \in \lambda', b \in \lambda''} \frac{\theta(t_1 x'_a/x''_b) \theta(t_2 x'_a/x''_b)}{\theta(x'_a/x''_b) \theta(\hbar x'_a/x''_b)} \prod_{b \in \lambda''} \theta(\hbar u_1/x''_b) \prod_{a \in \lambda'} \theta(x'_a/u_2) \right. \\ & \quad \left. \times \text{Stab}_{\mathfrak{C}', T^{1/2}}(\lambda'; \mathfrak{z} \hbar^{-1}) \text{Stab}_{\mathfrak{C}'', T^{1/2}}(\lambda''; \mathfrak{z}) \right). \quad (3.5) \end{aligned}$$

Here x'_a ($a \in \lambda'$) and x''_b ($b \in \lambda''$) are the Chern roots for the tautological bundle on $\mathcal{H}_{n'}$ and $\mathcal{H}_{n''}$, respectively, and we set $\{x_a\}_{a \in (\lambda', \lambda'')} = \{x'_a\}_{a \in \lambda'} \cup \{x''_b\}_{b \in \lambda''}$. Then one finds that $\text{Stab}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z})$ is the elliptic stable envelope for $E_T(\mathcal{M}(n' + n'', 2))$ with the chamber \mathfrak{C} given by $|u_1| \ll |u_2|$. The formula (3.5) is called the shuffle product [2]. By abuse of notation we use $T^{1/2}$ in $\text{Stab}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z})$ as $T^{1/2}\mathcal{M}$ i.e. the polarization satisfying

$$T\mathcal{M} = T^{1/2}\mathcal{M} + \hbar(T^{1/2}\mathcal{M})^\vee \in K_T(\mathcal{M})$$

for $\mathcal{M} = \mathcal{M}(n' + n'', 2)$. The shuffle product keeps such assignment of polarization consistent through a construction of the elliptic stable envelopes. See the remark below. In the following, we use the same convention.

By repeating the shuffle product, one obtains the elliptic stable envelopes for $E_T(\mathcal{M}(n, r))$. Let $\lambda^{(i)}$ ($i = 1, \dots, r$) be partitions with $|\lambda^{(i)}| = n_i$ and consider a r -tuple partition $\boldsymbol{\lambda} = (\lambda^{(1)}, \dots, \lambda^{(r)})$. We set $|\boldsymbol{\lambda}| = \sum_{i=1}^r n_i = n$.

Proposition 3.2. *The following gives the elliptic stable envelopes for $E_T(\mathcal{M}(n, r))$ with the chamber \mathfrak{C} being $|u_1| \ll \cdots \ll |u_r|$.*

$$\text{Stab}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) = \text{Sym}_{\mathbf{x}} \left(S_{\boldsymbol{\lambda}}^{\text{Ell}}(\mathbf{x}, \mathbf{u}) \sum_{\mathbf{t}} W_{T^{1/2}}^{\text{Ell}}(\mathbf{t}; \mathbf{x}, \mathbf{u}; \mathfrak{z}) \right),$$

where we set

$$\begin{aligned} S_{\boldsymbol{\lambda}}^{\text{Ell}}(\mathbf{x}, \mathbf{u}) &:= \frac{\prod_{\substack{a, b \in \lambda \\ \rho_a + 1 < \rho_b}} \theta(t_1 x_a / x_b) \prod_{\substack{a, b \in \lambda \\ \rho_a + 1 > \rho_b}} \theta(t_2 x_b / x_a)}{\prod_{\substack{a, b \in \lambda \\ \rho_a < \rho_b}} \theta(x_a / x_b) \theta(\hbar x_a / x_b)} \prod_{i=1}^r \left[\prod_{\substack{a \in \lambda \\ \rho_a > \rho_{r_i}}} \theta(\hbar u_i / x_a) \prod_{\substack{a \in \lambda \\ \rho_a \leq \rho_{r_i}}} \theta(x_a / u_i) \right], \\ \sum_{\mathbf{t}} W_{T^{1/2}}^{\text{Ell}}(\mathbf{t}; \mathbf{x}, \mathbf{u}; \mathfrak{z}) &:= \sum_{\delta^{(1)} \in \Upsilon_{\lambda^{(1)}}, \dots, \delta^{(r)} \in \Upsilon_{\lambda^{(r)}}} W_{T^{1/2}}^{\text{Ell}}(\mathbf{t}_{\delta^{(1)}}^{(1)}; \mathbf{x}^{(1)}, u_1; \mathfrak{z} \hbar^{-r+1}) W_{T^{1/2}}^{\text{Ell}}(\mathbf{t}_{\delta^{(2)}}^{(2)}; \mathbf{x}^{(2)}, u_2; \mathfrak{z} \hbar^{-r+2}) \cdots \\ &\quad \cdots \times W_{T^{1/2}}^{\text{Ell}}(\mathbf{t}_{\delta^{(r)}}^{(r)}; \mathbf{x}^{(r)}, u_r; \mathfrak{z}) \end{aligned}$$

with r_i being the root box of $\lambda^{(i)}$, $\mathbf{x}^{(i)} = (x_1^{(i)}, \dots, x_{n_i}^{(i)})$ being the Chern roots associated to the Hilbert scheme \mathcal{H}_{n_i} and $\mathbf{t} := (\mathbf{t}_{\delta^{(1)}}^{(1)}, \dots, \mathbf{t}_{\delta^{(r)}}^{(r)})$.

Remark. A more general formula for the elliptic stable envelope of affine A -type quivers is obtained in [3]. For $\mathcal{M}(n, r)$, this result coincides with Proposition 3.2. We note that the dynamical shifts of \mathfrak{z} is controlled in [3] via the positive part of polarization bundle $\det T_{\rho > 0, \lambda}^{1/2}$, see formula (13) and Definition 4.2.1 in [3]. For ρ defined by (3.2) and the chamber $|u_1| \ll \cdots \ll |u_r|$, these shifts are given by substitutions $\mathfrak{z} \rightarrow \mathfrak{z} \hbar^{-r+i}$ in the functions $W_{T^{1/2}}^{\text{Ell}}$, exactly as in the formulas of Proposition 3.2.

The shuffle product formula for $\text{Stab}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z})$'s is obtained by repeating (3.5).

Corollary 3.3. [2] *Let $\boldsymbol{\lambda}'$ and $\boldsymbol{\lambda}''$ be r' -tuple and r'' -tuple partitions with $|\boldsymbol{\lambda}'| = n'$, $|\boldsymbol{\lambda}''| = n''$, respectively. Consider the elliptic stable envelopes $\text{Stab}_{\mathfrak{C}', T^{1/2}}(\boldsymbol{\lambda}'; \mathfrak{z})$ and $\text{Stab}_{\mathfrak{C}'', T^{1/2}}(\boldsymbol{\lambda}''; \mathfrak{z})$ for $E_T(\mathcal{M}(n', r'))$ and $E_T(\mathcal{M}(n'', r''))$ with the equivariant parameters $\mathbf{u}' = (u'_1, \dots, u'_{r'})$, $\mathbf{u}'' = (u''_1, \dots, u''_{r''})$, respectively. Let \mathfrak{C}' and \mathfrak{C}'' be chambers $|u'_1| \ll \cdots \ll |u'_{r'}|$ and $|u''_1| \ll \cdots \ll |u''_{r''}|$, respectively with a common stability condition $t_1/t_2 > 0$. Then the following shuffle product gives the elliptic stable envelope $\text{Stab}_{\mathfrak{C}, T^{1/2}}((\boldsymbol{\lambda}', \boldsymbol{\lambda}''); \mathfrak{z})$ for $E_T(\mathcal{M}(n' + n'', r' + r''))$ with the chamber*

\mathfrak{C} given by $|u'_1| \ll \cdots \ll |u'_{r'}| \ll |u''_1| \ll \cdots \ll |u''_{r''}|$ with the stability condition $t_1/t_2 > 0$.

$$\begin{aligned} & \text{Stab}_{\mathfrak{C}, T^{1/2}}((\boldsymbol{\lambda}', \boldsymbol{\lambda}''); \mathfrak{z}) \\ &= \text{Sym}_{\{x_a\}_{a \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'')}} \left(\prod_{a \in \boldsymbol{\lambda}', b \in \boldsymbol{\lambda}''} \frac{\theta(t_1 x'_a/x'_b) \theta(t_2 x'_a/x''_b)}{\theta(x'_a/x''_b) \theta(\hbar x'_a/x''_b)} \prod_{i=1}^{r'} \prod_{b \in \boldsymbol{\lambda}''} \theta(\hbar u'_i/x''_b) \prod_{j=1}^{r''} \prod_{a \in \boldsymbol{\lambda}'} \theta(x'_a/u''_j) \right. \\ & \quad \left. \times \text{Stab}_{\mathfrak{C}', T^{1/2}}(\boldsymbol{\lambda}'; \mathfrak{z} \hbar^{-r''}) \text{Stab}_{\mathfrak{C}'', T^{1/2}}(\boldsymbol{\lambda}''; \mathfrak{z}) \right). \end{aligned} \quad (3.6)$$

Here x'_a ($a \in \boldsymbol{\lambda}'$) and x''_b ($b \in \boldsymbol{\lambda}''$) are the Chern roots for the tautological bundle on $\mathcal{M}(n', r')$ and $\mathcal{M}(n'', r'')$, respectively, and we set $\{x_a\}_{a \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'')} = \{x'_a\}_{a \in \boldsymbol{\lambda}'} \cup \{x''_b\}_{b \in \boldsymbol{\lambda}''}$.

In the next sections, we use $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z})$ defined by

$$\text{Stab}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) = \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) (-)^{\varepsilon(\boldsymbol{\lambda}, r)} \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a \neq \rho_b}} \frac{\theta(t_1 x_a/x_b)}{\theta(x_a/x_b)} \prod_{i=1}^r \prod_{a \in \boldsymbol{\lambda}} \theta(x_a/u_i), \quad (3.7)$$

where

$$\varepsilon(\boldsymbol{\lambda}, r) = \sum_{i=1}^r \sum_{\substack{a \in \boldsymbol{\lambda} \\ \rho_a > \rho_{r_i}}} 1.$$

The difference between Stab and $\widehat{\text{Stab}}$ is essentially the Thom class of $T^{1/2} = T^{1/2} \mathcal{M}(n, r)$ i.e.

$$\Theta(T^{1/2}) = \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a \neq \rho_b}} \frac{\theta(t_1 x_a/x_b)}{\theta(x_a/x_b)} \prod_{i=1}^r \prod_{a \in \boldsymbol{\lambda}} \theta(x_a/u_i).$$

Explicitly, $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z})$ is given by

$$\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) := \text{Sym}_{\mathbf{x}} \left(\widehat{S}_{\boldsymbol{\lambda}}^{Ell}(\mathbf{x}, \mathbf{u}) \sum_{\mathbf{t}} W_{T^{1/2}}^{Ell}(\mathbf{t}; \mathbf{x}, \mathbf{u}; \mathfrak{z}) \right) \quad (3.8)$$

with

$$\begin{aligned} \widehat{S}_{\boldsymbol{\lambda}}^{Ell}(\mathbf{x}, \mathbf{u}) &:= \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a < \rho_b}} \frac{\theta(x_b/x_a) \theta(t_2 x_a/x_b)}{\theta(t_1 x_b/x_a) \theta(\hbar x_a/x_b)} \prod_{i=1}^r \prod_{\substack{a \in \boldsymbol{\lambda} \\ \rho_a > \rho_{r_i}}} \left(-\frac{\theta(\hbar u_i/x_a)}{\theta(x_a/u_i)} \right) \\ &\quad \times \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ c_a = c_b \\ h_a > h_b}} \frac{\theta(t_2 x_b/x_a)}{\theta(t_1 x_a/x_b)} \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ c_a + 1 = c_b \\ h_a < h_b}} \frac{\theta(t_2 x_b/x_a)}{\theta(t_1 x_a/x_b)}. \end{aligned} \quad (3.9)$$

Proposition 3.4. *The shuffle product formula for $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z})$'s is given by*

$$\begin{aligned} & \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}((\boldsymbol{\lambda}', \boldsymbol{\lambda}''); \mathfrak{z}) \\ &= \text{Sym}_{\{x_a\}_{a \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'')}} \left(\prod_{\substack{a \in \boldsymbol{\lambda}', b \in \boldsymbol{\lambda}'' \\ \rho_a \neq \rho_b}} \frac{\theta(x_b''/x_a')\theta(t_2x_a'/x_b'')}{\theta(t_1x_b''/x_a')\theta(\hbar x_a'/x_b'')} \prod_{i=1}^{r'} \prod_{b \in \boldsymbol{\lambda}''} \left(-\frac{\theta(\hbar u_i'/x_b'')}{\theta(x_b''/u_i')} \right) \right. \\ & \quad \left. \times \widehat{\text{Stab}}_{\mathfrak{C}', T^{1/2}}(\boldsymbol{\lambda}'; \mathfrak{z}\hbar^{-r''}) \widehat{\text{Stab}}_{\mathfrak{C}'', T^{1/2}}(\boldsymbol{\lambda}''; \mathfrak{z}) \right). \end{aligned} \quad (3.10)$$

Proof. The statement follows from (3.6) and (3.7) by noting that the following factor is symmetric in x_a , $a \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'')$.

$$(-)^{\varepsilon((\boldsymbol{\lambda}', \boldsymbol{\lambda}''), r' + r'')} \prod_{\substack{a, b \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'') \\ \rho_a \neq \rho_b}} \frac{\theta(t_1x_a/x_b)}{\theta(x_a/x_b)} \prod_{i=1}^{r'+r''} \prod_{a \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'')} \theta(x_a/u_i).$$

□

In the following sections, we also use the elliptic stable envelopes for $E_T(\mathcal{M}(n, r))$ with the opposite polarization $T_{opp}^{1/2} = \hbar(T^{1/2})^\vee$, the elliptic nome p^* and the Kähler parameter \mathfrak{z}^{*-1} :

$$\text{Stab}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\boldsymbol{\lambda}; \mathfrak{z}^{*-1}) := \text{Stab}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) \Big|_{T^{1/2} \mapsto T_{opp}^{1/2}, p \mapsto p^*, \mathfrak{z} \mapsto \mathfrak{z}^{*-1}} \quad (3.11)$$

and its hatted version defined by

$$\text{Stab}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\boldsymbol{\lambda}; \mathfrak{z}^{*-1}) = \widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\boldsymbol{\lambda}; \mathfrak{z}^{*-1}) (-)^{\varepsilon^*(\boldsymbol{\lambda}, r)} \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a \neq \rho_b}} \frac{\theta^*(t_2x_a/x_b)}{\theta^*(\hbar x_a/x_b)} \prod_{i=1}^r \prod_{a \in \boldsymbol{\lambda}} \theta^*(\hbar u_i/x_a). \quad (3.12)$$

Here we set

$$\varepsilon^*(\boldsymbol{\lambda}, r) = \sum_{i=1}^r \sum_{\substack{a \in \boldsymbol{\lambda} \\ \rho_a \leq \rho r_i}} 1.$$

Note that for $T_{opp}^{1/2} = \hbar(T^{1/2}\mathcal{M}(n, r))^\vee$, we have

$$\Theta(T_{opp}^{1/2}) = \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a \neq \rho_b}} \frac{\theta^*(t_2x_a/x_b)}{\theta^*(\hbar x_a/x_b)} \prod_{i=1}^r \prod_{a \in \boldsymbol{\lambda}} \theta^*(\hbar u_i/x_a).$$

Proposition 3.5. *The shuffle product formula for $\widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\boldsymbol{\lambda}; \mathfrak{z}^{*-1})$'s is given by*

$$\begin{aligned} & \widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*((\boldsymbol{\lambda}', \boldsymbol{\lambda}''); \mathfrak{z}^{*-1}) \\ &= \text{Sym}_{\{x_a\}_{a \in (\boldsymbol{\lambda}', \boldsymbol{\lambda}'')}} \left(\prod_{a \in \boldsymbol{\lambda}', b \in \boldsymbol{\lambda}''} \frac{\theta^*(t_1x_a'/x_b'')\theta^*(\hbar x_b''/x_a')}{\theta^*(x_a'/x_b'')\theta^*(t_2x_b''/x_a')} \prod_{i=1}^{r''} \prod_{a \in \boldsymbol{\lambda}'} \left(-\frac{\theta^*(x_a'/u_i'')}{\theta^*(\hbar u_i''/x_a')} \right) \right. \\ & \quad \left. \times \widehat{\text{Stab}}_{\mathfrak{C}'', T_{opp}^{1/2}}^*(\boldsymbol{\lambda}''; \mathfrak{z}^{*-1}) \widehat{\text{Stab}}_{\mathfrak{C}', T_{opp}^{1/2}}^*(\boldsymbol{\lambda}'; \mathfrak{z}^{*-1}\hbar^{-r''}) \right). \end{aligned} \quad (3.13)$$

Proof. The statement follows from (3.12) and the fact that $\text{Stab}_{\mathfrak{e}, T_{opp}^{1/2}}^*(\boldsymbol{\lambda}; \mathfrak{z}^{*-1})$ satisfies the same shuffle product formula as (3.6) except for replacing all θ with θ^* . \square

4 Vertex Operators of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$

In this section, we introduce new vertex operators of type I and the type II dual by combining representations of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ in Sec.2.3 and the elliptic stable envelopes for $E_T(\mathcal{M}(n, r))$. We show that they yield correct shuffle product formula of the elliptic stable envelopes (Sec.4.3), the K-theoretic vertex functions (Sec.5), exchange relations (Sec.6) and L -operator (Sec.7).

4.1 OPE of the elliptic currents

Let us consider the level $(1, N)$ representation given in Sec.2.3.1, on which $p^* = p\hbar^{-1}$. For the elliptic currents $x^+(u), x^-(v)$, one gets the following operator product expansion (OPE).

$$x^+(u)x^+(v) = \langle x^+(u)x^+(v) \rangle : x^+(u)x^+(v) :, \quad (4.1)$$

$$x^-(u)x^-(v) = \langle x^-(u)x^-(v) \rangle : x^-(u)x^-(v) :, \quad (4.2)$$

with coefficients

$$\begin{aligned} \langle x^+(u)x^+(v) \rangle &= \frac{(v/u, v/\hbar u, p^* t_2 v/u, p^* t_1 v/u; p^*)_\infty}{(v/t_1 u, v/t_2 u, p^* \hbar v/u, p^* v/u; p^*)_\infty}, \\ \langle x^-(u)x^-(v) \rangle &= \frac{(v/u, \hbar v/u, pv/t_1 u, pv/t_2 u; p)_\infty}{(t_2 v/u, t_1 v/u, pv/\hbar u, pv/u; p)_\infty}. \end{aligned}$$

Here we set

$$(a_1, \dots, a_M; p)_\infty = \prod_{i=1}^M (a_i; p)_\infty.$$

Divide the OPE coefficient into two parts as follows.

$$\langle x^\pm(u)x^\pm(v) \rangle = \langle x^\pm(u)x^\pm(v) \rangle^{sym} \langle x^\pm(u)x^\pm(v) \rangle^{non-sym}.$$

The first factor is symmetric in u, v

$$\langle x^+(u)x^+(v) \rangle^{sym} = t_1^{-1} \frac{(p^* t_2 v/u, p^* t_2 u/v, v/u, u/v; p^*)_\infty}{(p^* \hbar v/u, p^* \hbar u/v, v/t_1 u, u/t_1 v; p^*)_\infty}, \quad (4.3)$$

$$\langle x^-(u)x^-(v) \rangle^{sym} = t_1^{-1} \frac{(pt_1^{-1} v/u, pt_1^{-1} u/v, \hbar v/u, \hbar u/v; p)_\infty}{(pv/u, pu/v, t_2 v/u, t_2 u/v; p)_\infty}, \quad (4.4)$$

whereas the second is not

$$\langle x^+(u)x^+(v) \rangle^{non-sym} = \frac{\theta^*(t_1 v/u)\theta^*(\hbar u/v)}{\theta^*(v/u)\theta^*(t_2 u/v)}, \quad (4.5)$$

$$\langle x^-(u)x^-(v) \rangle^{non-sym} = \frac{\theta(v/u)\theta(t_2 u/v)}{\theta(t_1 v/u)\theta(\hbar u/v)}. \quad (4.6)$$

Remark. There is an ambiguity in the division. For example another choice is as follows.

$$\langle x^+(u)x^+(v) \rangle_0^{sym} = \frac{(v/u, u/v, v/\hbar u, u/\hbar v; p^*)_\infty}{(v/t_1 u, u/t_1 v, v/t_2 u, u/t_2 v; p^*)_\infty}, \quad (4.7)$$

$$\langle x^-(u)x^-(v) \rangle_0^{sym} = \frac{(v/u, u/v, \hbar v/u, \hbar u/v; p)_\infty}{(t_1 v/u, t_1 u/v, t_2 v/u, t_2 u/v; p)_\infty}, \quad (4.8)$$

$$\langle x^+(u)x^+(v) \rangle_0^{non-sym} = \frac{\theta^*(t_2 v/u)\theta^*(t_1 v/u)}{\theta^*(v/u)\theta^*(\hbar v/u)}, \quad (4.9)$$

$$\langle x^-(u)x^-(v) \rangle_0^{non-sym} = \frac{\theta(t_1 u/v)\theta(t_2 u/v)}{\theta(u/v)\theta(\hbar u/v)}. \quad (4.10)$$

This non-symmetric parts directly give a part of factors in $S_\lambda^{Ell}(\mathbf{x}, u)$ (3.3) and naturally yield the shuffle product formula (3.6). However the symmetric parts lead to slightly different expressions for the vertex functions from (5.4) and (5.5).

4.2 Vertex operators for $\mathcal{M}(n, r)$

We start from the vertex operators corresponding to the case $r = 1$ i.e. $\text{Hilb}^n(\mathbb{C}^2)$. Let us define

$$\Phi_\theta(u) := (-u)^{-\alpha} e^{-\Lambda_0^\perp} \tilde{\Phi}_\theta(u), \quad (4.11)$$

$$\Psi_\theta^*(u) := (-u)^\alpha e^{\Lambda_0^\perp - Q} \tilde{\Psi}_\theta^*(u), \quad (4.12)$$

where $\tilde{\Phi}_\theta(u)$ and $\tilde{\Psi}_\theta^*(u)$ are given by

$$\tilde{\Phi}_\theta(u) = \exp \left\{ - \sum_{m>0} \frac{1}{\kappa_m} \alpha'_{-m} (\hbar^{1/2} u)^m \right\} \exp \left\{ \sum_{m>0} \frac{1}{\kappa_m} \alpha'_m (\hbar^{1/2} u)^{-m} \right\}, \quad (4.13)$$

$$\tilde{\Psi}_\theta^*(u) = \exp \left\{ \sum_{m>0} \frac{1}{\kappa_m} \alpha_{-m} (\hbar^{1/2} u)^m \right\} \exp \left\{ - \sum_{m>0} \frac{1}{\kappa_m} \alpha_m (\hbar^{1/2} u)^{-m} \right\}. \quad (4.14)$$

One can show the following commutation relations.

Proposition 4.1.

$$\Psi_\theta^*(u)x^+(v) = -\frac{\theta^*(v/u)}{\theta^*(\hbar u/v)} x^+(v)\Psi_\theta^*(u), \quad (4.15)$$

$$\Phi_\theta(u)x^-(v) = -\frac{\theta(\hbar u/v)}{\theta(v/u)} x^-(v)\Phi_\theta(u), \quad (4.16)$$

$$x^+(v)\Phi_\theta(u) = \Phi_\theta(u)x^+(v), \quad x^-(v)\Psi_\theta^*(u) = \Psi_\theta^*(u)x^-(v), \quad (4.17)$$

$$\mathfrak{z}\Phi_\theta(u) = \hbar^{-1}\Phi_\theta(u)\mathfrak{z}, \quad \Psi_\theta^*(u)\mathfrak{z}^* = \hbar\mathfrak{z}^*\Psi_\theta^*(u), \quad (4.18)$$

$$[x^\pm(x), \mathfrak{z}] = [\Psi_\theta^*(u), \mathfrak{z}] = 0, \quad [x^\pm(x), \mathfrak{z}^*] = [\Phi_\theta(u), \mathfrak{z}^*] = 0. \quad (4.19)$$

Definition 4.2. We define the type I $\Phi(u)$ and the type II dual $\Psi^*(u)$ vertex operators to be the following linear maps.

$$\begin{aligned} \Phi(u) &: \mathcal{F}_v^{(1,N)} \rightarrow \mathcal{F}_u^{(0,1)} \tilde{\otimes} \mathcal{F}_{-v/u}^{(1,N-1)}, \\ \Psi^*(u) &: \mathcal{F}_v^{(1,N)} \tilde{\otimes} \mathcal{F}_u^{(0,1)} \rightarrow \mathcal{F}_{-vu}^{(1,N+1)} \end{aligned}$$

with components defined by

$$\Phi(u) = \sum_{\lambda \in \mathcal{P}} |\lambda\rangle_u \tilde{\otimes} \Phi_\lambda(u), \quad (4.20)$$

$$\Phi_\lambda(u) = \int_{\mathcal{C}} \prod_{a \in \lambda} dx_a : \prod_{a \in \lambda} x^-(x_a) : \Phi_\emptyset(u) \prod_{\rho_a < \rho_b} \langle x^-(x_a) x^-(x_b) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\lambda; \mathfrak{z}),$$

and

$$\Psi_\lambda^*(u)\xi = \Psi^*(u)(\xi \tilde{\otimes} |\lambda\rangle_u), \quad \forall \xi \in \mathcal{F}_v^{(1, N)}, \quad (4.21)$$

$$\Psi_\lambda^*(u) = \int_{\mathcal{C}^*} \prod_{a \in \lambda} dx_a \widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\lambda; \mathfrak{z}^{*-1}) : \prod_{a \in \lambda} x^+(x_a) : \Psi_\emptyset^*(u) \prod_{\rho_a < \rho_b} \langle x^+(x_a) x^+(x_b) \rangle^{sym}.$$

The integration cycles $\mathcal{C}, \mathcal{C}^*$ are chosen appropriately depending on the situation of the application. We call $\Phi_\lambda(u)$ (resp. $\Psi_\lambda^*(u)$) with $\lambda \in \mathcal{P}_n$ the type I (resp. type II dual) vertex operator for $\text{Hilb}^n(\mathbb{C}^2)$. The total operators $\Phi(u)$ and $\Psi^*(u)$ are then the vertex operators for $\bigoplus_n \text{Hilb}^n(\mathbb{C}^2)$.

From the following Proposition one can deduce a representation theoretical origin of a part of factors in $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\lambda; \mathfrak{z})$ or $\widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\lambda; \mathfrak{z}^{*-1})$.

Proposition 4.3.

$$\begin{aligned} \Phi_\lambda(u) &= \int_{\mathcal{C}} \prod_{a \in \lambda} dx_a \prod_{\rho_a \leq 0} x^-(x_a) \cdot \Phi_\emptyset(u) \cdot \prod_{\rho_a > 0} x^-(x_a) \\ &\quad \times \prod_{\substack{c_a = c_b \\ h_a > h_b}} \frac{\theta(t_2 x_b / x_a)}{\theta(t_1 x_a / x_b)} \prod_{\substack{c_a + 1 = c_b \\ h_a < h_b}} \frac{\theta(t_2 x_b / x_a)}{\theta(t_1 x_a / x_b)} \sum_{\delta \in \Upsilon_\lambda} W_{T^{1/2}}^{Ell}(\mathbf{t}_\delta; x_1, \dots, x_{|\lambda|}, u; \mathfrak{z}), \\ \Psi_\lambda^*(u) &= \int_{\mathcal{C}^*} \prod_{a \in \lambda} dx_a \sum_{\delta \in \Upsilon_\lambda} W_{T_{opp}^{1/2}}^{Ell*}(\mathbf{t}_\delta; x_1, \dots, x_n, u; \mathfrak{z}^{*-1}) \times \prod_{\rho_a > 0} x^+(x_a) \cdot \Psi_\emptyset^*(u) \cdot \prod_{\rho_a \leq 0} x^+(x_a) \\ &\quad \times \prod_{\substack{c_a = c_b \\ h_a > h_b}} \frac{\theta^*(t_2 x_b / x_a)}{\theta^*(t_1 x_a / x_b)} \prod_{\substack{c_a + 1 = c_b \\ h_a < h_b}} \frac{\theta^*(t_2 x_b / x_a)}{\theta^*(t_1 x_a / x_b)}. \end{aligned}$$

Here $\prod_{a \in \lambda} x^-(x_a)$ (resp. $\prod_{a \in \lambda} x^+(x_a)$) denotes a product in the increasing (resp. decreasing) order of $a \in \lambda$ from left to right.

Proof. Use (4.15), (4.16) and

$$\begin{aligned} \prod_{a \in \lambda} x^+(x_a) &= \prod_{\substack{a, b \in \lambda \\ \rho_a > \rho_b}} \langle x^+(x_a) x^+(x_b) \rangle : \prod_{a \in \lambda} x^+(x_a) :, \\ \prod_{a \in \lambda} x^-(x_a) &= \prod_{\substack{a, b \in \lambda \\ \rho_a < \rho_b}} \langle x^-(x_a) x^-(x_b) \rangle : \prod_{a \in \lambda} x^-(x_a) :. \end{aligned}$$

Note that the normal ordered products : $\prod_{a \in \lambda} x^\pm(x_a)$: are symmetric in x_a $a \in \lambda$. \square

Remark. The vertex operators (4.20) and (4.21) should be compared with (3.21) and (3.23) in [28], where $\varphi_\mu, \varphi_\mu^*$ are the basic elliptic stable envelopes for the type A linear quiver variety.

4.3 Shuffle product formula

We show that the vertex operators reproduce the shuffle product formula for the elliptic stable envelopes.

4.3.1 Type I vertex operator

Let λ', λ'' be two partitions with $|\lambda'| = n', |\lambda''| = n''$. Let us consider the following composition of the two type I vertex operators for the Hilbert schemes.

$$\begin{aligned} & \Phi_{\lambda'}(u_1) \Phi_{\lambda''}(u_2) \\ &= \int_{\mathcal{C}} \prod_{a \in \lambda'} dx'_a \int_{\mathcal{C}} \prod_{b \in \lambda''} dx''_b : \prod_{a \in \lambda'} x^-(x'_a) : \Phi_\emptyset(u_1) \prod_{\rho_a < \rho_b} \langle x^-(x'_a) x^-(x'_b) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}', T^{1/2}}(\lambda'; \mathfrak{z}) \\ & \quad \times : \prod_{b \in \lambda''} x^-(x''_b) : \Phi_\emptyset(u_2) \prod_{\rho_c < \rho_d} \langle x^-(x''_c) x^-(x''_d) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}'', T^{1/2}}(\lambda''; \mathfrak{z}). \end{aligned}$$

Here the chambers $\mathfrak{C}', \mathfrak{C}''$ are the same and taken as the stability condition $t_1/t_2 > 0$. We also assume $|u_1| \ll |u_2|$. In a similar way to the type A linear quiver case studied in [28], let us arrange the order of the elements in the integrand as follows.

1. Move $\widehat{\text{Stab}}_{\mathfrak{C}', T^{1/2}}(\lambda'; \mathfrak{z})$ to the right of all operators.
2. Move : $\prod_{b \in \lambda''} x^-(x''_b)$: to the left of $\Phi_\emptyset(u_1)$ by using the formula (4.16).
3. Make : $\prod_{a \in \lambda'} x^-(x'_a) :: \prod_{b \in \lambda''} x^-(x''_b)$: totally normal ordered product by the formula

$$: \prod_{a \in \lambda'} x^-(x'_a) :: \prod_{b \in \lambda''} x^-(x''_b) := \prod_{a \in \lambda', b \in \lambda''} \langle x^-(x'_a) x^-(x''_b) \rangle : \prod_{a \in (\lambda', \lambda'')} x^-(x_a) : .$$

Here we set $\{x_a\}_{a \in (\lambda', \lambda'')} = \{x'_a\}_{a \in \lambda'} \cup \{x''_b\}_{b \in \lambda''}$. We define the order of boxes in the different partitions by $\rho_a < \rho_b$ for $a \in \lambda', b \in \lambda''$.

4. Divide $\langle x^-(x'_a) x^-(x''_b) \rangle$ into the symmetric part (4.4) and the non-symmetric part (4.6).
5. Symmetrize the integrand over $\{x_a\}_{a \in (\lambda', \lambda'')}$.

One then obtains

$$\begin{aligned} \Phi_{\lambda'}(u_1)\Phi_{\lambda''}(u_2) &= \int_{\mathcal{C} \times \mathcal{C}} \prod_{a \in (\lambda', \lambda'')} dx_a : \prod_{a \in (\lambda', \lambda'')} x^-(x_a) : \Phi_{\emptyset}(u_1)\Phi_{\emptyset}(u_2) \\ &\times \prod_{\substack{a, b \in (\lambda', \lambda'') \\ \rho_a < \rho_b}} \langle x^-(x_a)x^-(x_b) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z}), \end{aligned} \quad (4.22)$$

where

$$\begin{aligned} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z}) &= \text{Sym}_{\{x_a\}_{a \in (\lambda', \lambda'')}} \left(\prod_{a \in \lambda', b \in \lambda''} \frac{\theta(x_b''/x_a')\theta(t_2 x_a'/x_b'')}{\theta(t_1 x_b''/x_a')\theta(\hbar x_a'/x_b'')} \prod_{b \in \lambda''} \left(-\frac{\theta(\hbar u_1/x_b'')}{\theta(x_b''/u_1)} \right) \right. \\ &\quad \left. \times \widehat{\text{Stab}}_{\mathfrak{C}', T^{1/2}}(\lambda'; \mathfrak{z}\hbar^{-1}) \widehat{\text{Stab}}_{\mathfrak{C}'', T^{1/2}}(\lambda''; \mathfrak{z}) \right) \end{aligned}$$

Comparing this with (3.10), one finds that $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z})$ coincides with the hatted version of the elliptic stable envelope for $E_T(\mathcal{M}(n' + n'', 2))$ given in (3.5). We thus regard the composition $\Phi_{\lambda'}(u_1)\Phi_{\lambda''}(u_2)$ as a vertex operator for $\mathcal{M}(n' + n'', 2)$.

By repeating such consideration, one obtains a vertex operator for $\bigoplus_n E_T(\mathcal{M}(n, r))$ by the following composition.

$$(\text{id} \otimes \cdots \otimes \text{id} \otimes \tilde{\Phi}(u_1)) \circ \cdots \circ (\text{id} \otimes \tilde{\Phi}(u_{r-1})) \circ \tilde{\Phi}(u_r) : \mathcal{F}_v^{(1, N+r)} \rightarrow \mathcal{F}_{u_r}^{(0,1)} \otimes \cdots \otimes \mathcal{F}_{u_1}^{(0,1)} \otimes \mathcal{F}_{(-)^r v/u_1 \cdots u_r}^{(1, N)}.$$

Its components $\Phi_{\lambda}(u_1, \dots, u_r)$ are defined by

$$\begin{aligned} &(\text{id} \otimes \cdots \otimes \text{id} \otimes \tilde{\Phi}(u_1)) \circ \cdots \circ (\text{id} \otimes \tilde{\Phi}(u_{r-1})) \circ \tilde{\Phi}(u_r) \\ &= \sum_{n \in \mathbb{Z}_{\geq 0}} \sum_{\substack{\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)}) \\ |\lambda| = n}} |\lambda^{(r)}\rangle_{u_r} \otimes \cdots \otimes |\lambda^{(1)}\rangle_{u_1} \otimes \tilde{\Phi}_{\lambda}(u_1, \dots, u_r). \end{aligned} \quad (4.23)$$

One finds

$$\begin{aligned} \Phi_{\lambda}(u_1, \dots, u_r) &= \Phi_{\lambda^{(1)}}(u_1) \cdots \Phi_{\lambda^{(r)}}(u_r) \\ &= \int_{\mathcal{C}^r} \prod_{a \in \lambda} dx_a : \prod_{a \in \lambda} x^-(x_a) : \Phi_{\emptyset}(u_1) \cdots \Phi_{\emptyset}(u_r) \prod_{\substack{a, b \in \lambda \\ \rho_a < \rho_b}} \langle x^-(x_a)x^-(x_b) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\lambda; \mathfrak{z}), \end{aligned} \quad (4.24)$$

where $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\lambda; \mathfrak{z})$ is the hatted version of the elliptic stable envelope for $E_T(\mathcal{M}(n, r))$ given in (3.8) satisfying the shuffle product formula (3.10). We hence regard $\Phi_{\lambda}(u_1, \dots, u_r)$ as the type I vertex operator for $\mathcal{M}(n, r)$. Then the total one (4.23) is the vertex operator for $\bigoplus_n \mathcal{M}(n, r)$.

4.3.2 Type II dual vertex operator

Composing the type II dual vertices, one gets

$$\Psi^*(u_r) \circ (\Psi^*(u_{r-1}) \widetilde{\otimes} \text{id}) \circ \cdots \circ (\Psi^*(u_1) \widetilde{\otimes} \text{id} \widetilde{\otimes} \cdots \widetilde{\otimes} \text{id}) : \mathcal{F}_v^{(1,N)} \widetilde{\otimes} \mathcal{F}_{u_1}^{(0,1)} \widetilde{\otimes} \cdots \widetilde{\otimes} \mathcal{F}_{u_r}^{(0,1)} \rightarrow \mathcal{F}_{(-)^r v u_1 \cdots u_r}^{(1,N+r)}. \quad (4.25)$$

We define its components $\Psi_{\lambda}^*(u_1, \dots, u_r)$ labeled by a r -tuple partition $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ with $|\lambda| = n$ by

$$\begin{aligned} \Psi_{\lambda}^*(u_1, \dots, u_r) \xi &= \Psi^*(u_r) \circ (\Psi^*(u_{r-1}) \widetilde{\otimes} \text{id}) \circ \cdots \circ (\Psi^*(u_1) \widetilde{\otimes} \text{id} \widetilde{\otimes} \cdots \widetilde{\otimes} \text{id}) (\xi \widetilde{\otimes} |\lambda^{(1)}\rangle_{u_1} \widetilde{\otimes} \cdots \widetilde{\otimes} |\lambda^{(r)}\rangle_{u_r}) \\ &= \Psi_{\lambda^{(r)}}^*(u_r) \Psi_{\lambda^{(r-1)}}^*(u_{r-1}) \cdots \Psi_{\lambda^{(1)}}^*(u_1) \xi, \quad \forall \xi \in \mathcal{F}_v^{(1,N)}. \end{aligned}$$

Similarly to the type I vertex (4.24), we obtain the following expression.

$$\begin{aligned} &\Psi_{\lambda}^*(u_1, \dots, u_r) \\ &= \int_{\mathcal{C}^{*r}} \prod_{a \in \lambda} dx_a \widehat{\text{Stab}}_{\mathcal{C}, T_{opp}^{1/2}}^*(\lambda; \mathfrak{z}^{*-1}) : \prod_{a \in \lambda} x^+(x_a) : \Psi_{\emptyset}^*(u_r) \cdots \Psi_{\emptyset}^*(u_1) \\ &\quad \times \prod_{\substack{a, b \in \lambda \\ \rho_a < \rho_b}} \langle x^+(x_a) x^+(x_b) \rangle^{sym}. \end{aligned} \quad (4.26)$$

Here $\widehat{\text{Stab}}_{\mathcal{C}, T_{opp}^{1/2}}^*(\lambda; \mathfrak{z}^{*-1})$ is given in (3.12). We hence regard $\Psi_{\lambda}^*(u_1, \dots, u_r)$ as the type II dual vertex operator for $\mathcal{M}(n, r)$. The total one (4.25) is the vertex operator for $\bigoplus_n \mathcal{M}(n, r)$.

The shuffle product formula for $\widehat{\text{Stab}}_{\mathcal{C}, T_{opp}^{1/2}}^*(\lambda; \mathfrak{z}^{*-1})$ is given by (3.13). This is reproduced by considering a composition $\Psi_{\lambda''}^*(u''_1, \dots, u''_{r''}) \Psi_{\lambda'}^*(u'_1, \dots, u'_{r'})$ and making the following arrangements of the operators in the integrand.

1. Move $\widehat{\text{Stab}}_{\mathcal{C}, T_{opp}^{1/2}}^*(\lambda'; \mathfrak{z}^{*-1})$ to the left of all operators.
2. Move $:\prod_{a \in \lambda'} x^+(x'_a):$ to the left of $\Psi_{\emptyset}^*(u''_{r''}) \cdots \Psi_{\emptyset}^*(u''_1)$ by using the formula (4.15).
3. Make $:\prod_{b \in \lambda''} x^+(x''_b): :: \prod_{a \in \lambda'} x^+(x'_a):$ totally normal ordered product by the formula

$$:\prod_{b \in \lambda''} x^+(x''_b): :: \prod_{a \in \lambda'} x^+(x'_a) := \prod_{a \in \lambda', b \in \lambda''} \langle x^+(x''_b) x^+(x'_a) \rangle : \prod_{a \in (\lambda', \lambda'')} x^+(x_a) : .$$

Here we set $\{x_a\}_{a \in (\lambda', \lambda'')} = \{x'_a\}_{a \in \lambda'} \cup \{x''_b\}_{b \in \lambda''}$. We define the order of boxes in the different partitions by $\rho_a < \rho_b$ for $a \in \lambda', b \in \lambda''$.

4. Divide $\langle x^+(x''_b) x^+(x'_a) \rangle$ into the symmetric part (4.3) and the non-symmetric part (4.5).
5. Symmetrize the integrand over $\{x_a\}_{a \in (\lambda', \lambda'')}$.

5 The K-Theoretic Vertex functions for $\mathcal{M}(n, r)$

In this section, we calculate vacuum expectation values of the vertex operators constructed in the last section and show that they give K-theoretic vertex functions for $\mathcal{M}(n, r)$.

5.1 The type I case

Let λ, μ be two partitions with $|\lambda| = |\mu| = n$. There is a bijection ς from boxes in λ to those in μ defined by $\varsigma(a) = b \in \mu$ for $a \in \lambda$ if $\iota(a) = \iota(b)$. Here ι is defined in Sec.3.1. For a box $a = (i, j) \in \mu$, we set $\varphi_a^\mu = t_1^{-(j-1)} t_2^{-(i-1)}$ as before. For the Chern root x_a ($a \in \lambda$) we take the Jackson integral

$$\int_0^{\varphi_{\varsigma(a)}^\mu} d_p x_a f(x_a) = (1-p) \varphi_{\varsigma(a)}^\mu \sum_{d \in \mathbb{N}} f(\varphi_{\varsigma(a)}^\mu p^d) p^d$$

in the vertex operators (4.20) for $\text{Hilb}^n(\mathbb{C}^2)$. We have

$$\begin{aligned} \Phi_\lambda(u) &= \prod_{a \in \lambda} \left(\int_0^{\varphi_{\varsigma(a)}^\mu} d_p x_a \right) : \prod_{a \in \lambda} x^-(x_a) : \Phi_\emptyset(u) \prod_{\substack{a, b \in \lambda \\ \rho_a < \rho_b}} \langle x^-(x_a) x^-(x_b) \rangle^{sym} \\ &\quad \times \widehat{\text{Stab}}_{\mathfrak{e}, T^{1/2}}(\lambda; \mathfrak{z}). \end{aligned} \quad (5.1)$$

Let $|0\rangle_v^{(1, N)}$ be the vacuum state (2.25) in $\mathcal{F}_v^{(1, N)}$ and ${}^{(1, N)}\langle 0|$ be the dual state satisfying

$${}^{(1, N)}\langle 0| |0\rangle_v^{(1, N)} = 1.$$

Let us consider the vacuum expectation value of $\Phi_\lambda(u)$. One only needs to take a normal ordering of the operators:

$$: \prod_{a \in \lambda} x^-(x_a) : \Phi_\emptyset(u) = \prod_{a \in \lambda} \left(-\frac{\hbar^{1/2} u}{x_a} \right) \frac{(pu/x_a; p)_\infty}{(\hbar u/x_a; p)_\infty} : \prod_{a \in \lambda} x^-(x_a) \cdot \Phi_\emptyset(u) : .$$

Here the normal ordering of the zero-modes operators is given by

$$\prod_{a \in \lambda} e^{-h} \left(\hbar^{1/2}/x_a \right)^{-c^\perp} \cdot (-u)^{-\alpha} e^{-\Lambda_0^\perp} = \prod_{a \in \lambda} \left(-\frac{\hbar^{1/2} u}{x_a} \right) \cdot (-u)^{-\alpha} e^{-\Lambda_0^\perp} \prod_{a \in \lambda} e^{-h} \left(\hbar^{1/2}/x_a \right)^{-c^\perp} .$$

Then using (4.4), one obtains

$$\begin{aligned} & {}^{(1, N-1)}_{-v/u} \langle 0 | \Phi_\lambda(u) | 0 \rangle_v^{(1, N)} \\ &= (-u/v)^n \prod_{a \in \lambda} \left(\int_0^{\varphi_{\varsigma(a)}^\mu} d_p x_a \right) \prod_{a \in \lambda} \left(\frac{x_a}{\hbar^{1/2}} \right)^{N-1} \frac{(pu/x_a; p)_\infty}{(\hbar u/x_a; p)_\infty} \\ &\quad \times \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} t_1^{-1} \frac{(pt_1^{-1} x_a/x_b; p)_\infty (\hbar x_a/x_b; p)_\infty}{(px_a/x_b; p)_\infty (t_2 x_a/x_b; p)_\infty} \widehat{\text{Stab}}_{\mathfrak{e}, T^{1/2}}(\lambda; \mathfrak{z}). \end{aligned} \quad (5.2)$$

To evaluate the Jackson integrals, we use the following Proposition.

Proposition 5.1. For the elliptic stable envelope $\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z})$ for $\text{Hilb}^n(\mathbb{C}^2)$, one has

$$\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{x_a = \varphi_{\zeta(a)}^\mu p^{d_a}} = \mathfrak{z}^{-\sum_{a \in \lambda} d_a} \times \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{\mu},$$

where $\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{\mu}$ stands for the elliptic stable envelope restricted to the fixed point μ i.e.

$$\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{\mu} := \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{x_a = \varphi_{\zeta(a)}^\mu},$$

and is independent from $\mathbf{d} = (d_a)_{a \in \lambda}$.

Proof. We recall that the quasiperiods of the elliptic stable envelope $\text{Stab}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z})$ in variables x_a are controlled by the universal line bundle on $E_T(\text{Hilb}^n(\mathbb{C}^2))$. For instance, see Section 2 in [44] for the definitions. Namely, these quasiperiods are the same as for the function

$$\phi\left(\prod_{a \in \lambda} x_a, \mathfrak{z}\right) \times \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} \frac{\theta(t_1 x_a / x_b)}{\theta(x_a / x_b)} \prod_{a \in \lambda} \theta(x_a / u).$$

see Section 3.7 in [44]. Comparing this function with (3.7) we find that $\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z})$ has the same quasiperiods as the function $\phi(\prod_{a \in \lambda} x_a, \mathfrak{z})$. Since $\phi(xp, y) = y^{-1} \phi(x, y)$, $\phi(x, yp) = x^{-1} \phi(x, y)$, we obtain

$$\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{x_a \rightarrow px_a} = \mathfrak{z}^{-1} \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}),$$

and therefore

$$\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{x_a = \varphi_{\zeta(a)}^\mu p^{d_a}} = \mathfrak{z}^{-\sum_{a \in \lambda} d_a} \times \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{x_a = \varphi_{\zeta(a)}^\mu} = \mathfrak{z}^{-\sum_{a \in \lambda} d_a} \times \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{\mu}.$$

□

Remark. In [45], the following function is used in the vertex function instead of $\widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z})$.

$$\mathbf{e}(\mathbf{x}, \mathfrak{z}) = \exp \left\{ -\frac{1}{\log p} \sum_{a \in \lambda} \log x_a \log \mathfrak{z} \right\}.$$

This yields the same effect as $\widehat{\text{Stab}}_{\mathfrak{E}}(\lambda; \mathfrak{z})$ in Proposition 5.1.

$$\mathbf{e}(\mathbf{x}, \mathfrak{z}) \Big|_{x_a = \varphi_{\zeta(a)}^\mu p^{d_a}} = \mathfrak{z}^{-\sum_{a \in \lambda} d_a} \times \mathbf{e}(\mathbf{x}, \mathfrak{z}) \Big|_{x_a = \varphi_{\zeta(a)}^\mu}.$$

From Proposition 5.1, one obtains

$$\begin{aligned} & \langle 0 | \Phi_\lambda(u) | 0 \rangle_v^{(1, N)} \\ &= \left(-\frac{(1-p)u}{v(\hbar^{1/2})^{N-1}} \right)^n \sum_{\mathbf{d} \in \mathbb{N}^n} (p^{-N} \mathfrak{z})^{-\sum_{a \in \lambda} d_a} \prod_{a \in \lambda} (\varphi_{\zeta(a)}^\mu)^N \frac{(p^{-d_a} pu / \varphi_{\zeta(a)}^\mu; p)_\infty}{(p^{-d_a} \hbar u / \varphi_{\zeta(a)}^\mu; p)_\infty} \\ & \times \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} t_1^{-1} \frac{(p^{d_a - d_b} p t_1^{-1} \varphi_{\zeta(a)}^\mu / \varphi_{\zeta(b)}^\mu; p)_\infty (p^{d_a - d_b} \hbar \varphi_{\zeta(a)}^\mu / \varphi_{\zeta(b)}^\mu; p)_\infty}{(p^{d_a - d_b} p \varphi_{\zeta(a)}^\mu / \varphi_{\zeta(b)}^\mu; p)_\infty (p^{d_a - d_b} t_2 \varphi_{\zeta(a)}^\mu / \varphi_{\zeta(b)}^\mu; p)_\infty} \times \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}}(\lambda; \mathfrak{z}) \Big|_{\mu}. \end{aligned} \quad (5.3)$$

Let \mathcal{N}_μ be the specialization of the integrand in (5.2) to $x_a = \varphi_{\varsigma(a)}^\mu$ ($a \in \lambda$) i.e. the $\mathbf{d} = \mathbf{0}$ term in (5.3). Then the K-theoretic vertex function $V_\lambda^\mu(u, z)$ for $\text{Hilb}^n(\mathbb{C}^2)$ is obtained as follows.

$$\begin{aligned} V_\lambda^\mu(u, z) &= \frac{1}{\mathcal{N}_\mu^{(1, N-1)_{-v/u}}} \langle 0 | \Phi_\lambda(u) | 0 \rangle_v^{(1, N)} \\ &= \sum_{\mathbf{d} \in \mathbb{N}^n} (\hbar p^{-N-1} \mathfrak{z})^{-\sum_a d_a} \prod_{a \in \lambda} \frac{(\varphi_{\varsigma(a)}^\mu / u; p)_{d_a}}{(p \hbar^{-1} \varphi_{\varsigma(a)}^\mu / u; p)_{d_a}} \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} \frac{(p \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b} (t_2 \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b}}{(p t_1^{-1} \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b} (\hbar \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b}}. \end{aligned}$$

Here we used for $d \in \mathbb{Z}$

$$(z; p)_d = \frac{(z; p)_\infty}{(p^d z; p)_\infty}$$

and the formula for $d > 0$

$$(z; p)_{-d} = (-z)^{-d} p^{\frac{p(p-1)}{2}} \frac{1}{(p z^{-1}; p)_d}.$$

In order to compare this with the known formula, one needs a shift $u \mapsto \hbar^{-1}u$.

$$\begin{aligned} V_\lambda^\mu(\hbar^{-1}u, \mathfrak{z}) &= \sum_{\mathbf{d} \in \mathbb{N}^n} (\hbar p^{-N-1} \mathfrak{z})^{-\sum_a d_a} \prod_{a \in \lambda} \frac{(\hbar \varphi_{\varsigma(a)}^\mu / u; p)_{d_a}}{(p \varphi_{\varsigma(a)}^\mu / u; p)_{d_a}} \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} \frac{(p \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b} (t_2 \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b}}{(p t_1^{-1} \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b} (\hbar \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b}}. \end{aligned}$$

The special case $\mu = \lambda$, hence $\varsigma = \text{id}$, of this coincides with the formula obtained geometrically in [45].

In the same way, the vertex function for $\mathcal{M}(n, r)$ is obtained from the vertex operator (4.24). We use the following formula obtained by combining Proposition 5.1 and the shuffle product formula (3.10).

$$\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) \Big|_{x_a = \varphi_{\varsigma(a)}^\mu p^{d_a}} = \left(\hbar^{-(r-1)} \mathfrak{z} \right)^{-\sum_{a \in \lambda} d_a} \times \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) \Big|_{\boldsymbol{\mu}},$$

where $\boldsymbol{\lambda}, \boldsymbol{\mu}$ are r -tuple partitions with $|\boldsymbol{\lambda}| = |\boldsymbol{\mu}| = n$, and the bijection $\varsigma : \boldsymbol{\lambda} \rightarrow \boldsymbol{\mu}$ is defined in the same way as above. One obtains the following result.

$$\begin{aligned} V_\lambda^\mu(u_1/\hbar, \dots, u_r/\hbar, \mathfrak{z}) &:= \frac{1}{\mathcal{N}_\mu^{(1, N-r)_{-v/u_1 \dots u_r}}} \langle 0 | \Phi_\lambda(u_1/\hbar, \dots, u_r/\hbar) | 0 \rangle_v^{(1, N)} \\ &= \sum_{\mathbf{d} \in \mathbb{N}^n} \prod_{i=1}^r (\hbar^{-(r-2)} p^{r-N-1} \mathfrak{z})^{-\sum_{a \in \lambda^{(i)}} d_a} \prod_{i=1}^r \prod_{a \in \lambda} \frac{(\hbar \varphi_{\varsigma(a)}^\mu / u_i; p)_{d_a}}{(p \varphi_{\varsigma(a)}^\mu / u_i; p)_{d_a}} \\ &\quad \times \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} \frac{(p \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b} (t_2 \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b}}{(p t_1^{-1} \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b} (\hbar \varphi_{\varsigma(a)}^\mu / \varphi_{\varsigma(b)}^\mu; p)_{d_a - d_b}}. \end{aligned} \tag{5.4}$$

Here $\mathbf{d} = (d_a)$, $a \in \lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ and

$$\begin{aligned} \mathcal{N}_\mu &= \prod_{1 \leq i < j \leq r} \langle \Phi_\emptyset(u_i) \Phi_\emptyset(u_j) \rangle \times \prod_{a \in \lambda} \frac{(1-p)\varphi_\zeta^\mu}{v} \left(\frac{\hbar^{1/2}}{\varphi_\zeta^\mu} \right)^{-N} \\ &\times \prod_{i=1}^r \prod_{a \in \lambda} \left(-\frac{\hbar^{-1/2} u_i (p u_i / \varphi_\zeta^\mu; p)_\infty}{\varphi_\zeta^\mu (\hbar u_i / \varphi_\zeta^\mu; p)_\infty} \right) \times \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} t_1^{-1} \frac{(p t_1^{-1} \varphi_\zeta^\mu / \varphi_\zeta^\mu; p)_\infty (\hbar \varphi_\zeta^\mu / \varphi_\zeta^\mu; p)_\infty}{(p \varphi_\zeta^\mu / \varphi_\zeta^\mu; p)_\infty (t_2 \varphi_\zeta^\mu / \varphi_\zeta^\mu; p)_\infty} \\ &\times \widehat{\text{Stab}}_{\mathfrak{E}, T^{1/2}(\lambda; \mathfrak{z})} \Big|_\mu. \end{aligned}$$

5.2 The type II dual case

Similarly to the type I case, one obtains the K-theoretic vertex function from the type II dual vertex operator (4.26). By using the OPE formula

$$\begin{aligned} \Psi_{\lambda^{(r)}}^*(u_r) \cdots \Psi_{\lambda^{(1)}}^*(u_1) &= \prod_{1 \leq i < j \leq r} \langle \Psi_\emptyset^*(u_j) \Psi_\emptyset^*(u_i) \rangle \prod_{i=1}^r \prod_{a \in \lambda} \left(-\frac{\hbar^{1/2} u_i (\hbar p^* u_i / x_a; p^*)_\infty}{x_a (u_i / x_a; p^*)_\infty} \right) \\ &\times \prod_{i=1}^r \left((-u_i)^\alpha e^{\Lambda_\emptyset^\perp - Q} \right) \prod_{a \in \lambda} e^h \left(\frac{\hbar^{1/2}}{x_a} \right)^{c^\perp} : \prod_{a \in \lambda} \tilde{x}^+(x_a) \prod_{i=1}^r \tilde{\Psi}_\emptyset^*(u_i) :, \end{aligned}$$

one obtains the following formula.

$$\begin{aligned} V_\lambda^{*\mu}(u_1, \dots, u_r, \mathfrak{z}^{*-1}) &:= \frac{1}{\mathcal{N}_\mu^*} (-)^{r \nu u_1 \cdots u_r} \langle 0 | \Psi_\lambda^*(u_1, \dots, u_r) | 0 \rangle_v^{(1, N)} \\ &= \sum_{\mathbf{d} \in \mathbb{N}^n} \prod_{i=1}^r (\hbar^{-r} p^{*N-1} \mathfrak{z}^{*-1})^{-\sum_{a \in \lambda^{(i)}} d_a} \prod_{i=1}^r \prod_{a \in \lambda} \frac{(\hbar^{-1} \varphi_\zeta^\mu / u_i; p^*)_{d_a}}{(p^* \varphi_\zeta^\mu / u_i; p^*)_{d_a}} \\ &\times \prod_{\substack{a, b \in \lambda \\ \rho_a \neq \rho_b}} \frac{(p^* \varphi_\zeta^\mu / \varphi_\zeta^\mu; p^*)_{d_a - d_b} (t_2^{-1} \varphi_\zeta^\mu / \varphi_\zeta^\mu; p^*)_{d_a - d_b}}{(p^* t_1 \varphi_\zeta^\mu / \varphi_\zeta^\mu; p^*)_{d_a - d_b} (\hbar^{-1} \varphi_\zeta^\mu / \varphi_\zeta^\mu; p^*)_{d_a - d_b}}. \end{aligned} \quad (5.5)$$

Here we used

$$\widehat{\text{Stab}}_{\mathfrak{E}, T_{opp}^{1/2}(\lambda; \mathfrak{z}^{*-1})} \Big|_{x_a = \varphi_\zeta^\mu p^{*d_a}} = \mathfrak{z}^{*\sum_{a \in \lambda} d_a} \times \widehat{\text{Stab}}_{\mathfrak{E}, T_{opp}^{1/2}(\lambda; \mathfrak{z}^{*-1})} \Big|_\mu.$$

Comparing (5.5) with (5.4), one finds a duality compatible to Proposition 2.2 under a certain tuning of the Kähler parameter \mathfrak{z}^* .

6 Exchange Relations of the Vertex Operators

We derive exchange relations of the vertex operators for $\text{Hilb}^n(\mathbb{C}^2)$. We show that their coefficients are given by the elliptic dynamical instanton R -matrices, which arise as the transition matrices of the elliptic stable envelopes. These results can be easily extended to the case $\mathcal{M}(n, r)$.

6.1 Type I vertex operator

Let denote by $\Phi_{\lambda'}^{(\mu')}(u_1)$ the type I vertex operator in (5.1) and consider the composition

$$\begin{aligned} \Phi_{\lambda'}^{(\mu')}(u_1)\Phi_{\lambda''}^{(\mu'')}(u_2) &= \prod_{a \in \lambda'} \left(\int_0^{\varphi_{\zeta}^{\mu'}(a)} d_p x'_a \right) \prod_{b \in \lambda''} \left(\int_0^{\varphi_{\zeta}^{\mu''}(b)} d_p x''_b \right) : \prod_{a \in (\lambda', \lambda'')} x^-(x_a) : \Phi_{\emptyset}(u_1)\Phi_{\emptyset}(u_2) \\ &\times \prod_{\substack{a, b \in (\lambda', \lambda'') \\ \rho_a < \rho_b}} \langle x^-(x_a)x^-(x_b) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}((\lambda', \lambda''); \mathfrak{z}), \end{aligned} \quad (6.1)$$

where the chamber \mathfrak{C} is given by $u_1 \ll u_2$ with the stability condition $t_1/t_2 > 0$. Similarly we have the opposite order of composition

$$\begin{aligned} \Phi_{\lambda''}^{(\mu'')}(u_2)\Phi_{\lambda'}^{(\mu')}(u_1) &= \prod_{b \in \lambda''} \left(\int_0^{\varphi_{\zeta}^{\mu''}(b)} d_p x''_b \right) \prod_{a \in \lambda'} \left(\int_0^{\varphi_{\zeta}^{\mu'}(a)} d_p x'_a \right) : \prod_{a \in (\lambda', \lambda'')} x^-(x_a) : \Phi_{\emptyset}(u_2)\Phi_{\emptyset}(u_1) \\ &\times \prod_{\substack{a, b \in (\lambda', \lambda'') \\ \rho_a < \rho_b}} \langle x^-(x_a)x^-(x_b) \rangle^{sym} \widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T^{1/2}}((\lambda'', \lambda'); \mathfrak{z}), \end{aligned} \quad (6.2)$$

where the chamber $\bar{\mathfrak{C}}$ is given by $u_1 \gg u_2$ with the stability condition $t_1/t_2 > 0$.

Let $(\mathcal{P}^2)_n = \{\alpha = (\alpha', \alpha'') \in \mathcal{P} \times \mathcal{P} \mid |\alpha| = |\alpha'| + |\alpha''| = n\}$. For $\alpha = (\alpha', \alpha'')$, let $\bar{\alpha} = (\alpha'', \alpha')$. We define the elliptic dynamical instanton R -matrix $R_{T^{1/2}}(u_1, u_2; \mathfrak{z}) \in \text{End}_{\mathbb{F}}(\mathcal{F}_{u_1}^{(0,1)} \tilde{\otimes} \mathcal{F}_{u_2}^{(0,1)})$ as the following transition matrix of the elliptic stable envelopes. For $\alpha, \beta, \gamma \in (\mathcal{P}^2)_n$,

$$R_{T^{1/2}}(u_1, u_2; \mathfrak{z})|\beta'\rangle_{u_1} \tilde{\otimes} |\beta''\rangle_{u_2} = \sum_{\alpha \in (\mathcal{P}^2)_n} R_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\alpha}^{\beta} |\alpha'\rangle_{u_1} \tilde{\otimes} |\alpha''\rangle_{u_2}, \quad (6.3)$$

$$R_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\alpha}^{\beta} = \mu(u_1/u_2) \bar{R}_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\alpha}^{\beta}, \quad (6.4)$$

$$\widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T^{1/2}}(\bar{\alpha}; \mathfrak{z})|_{\bar{\gamma}} = \sum_{\beta \in (\mathcal{P}^2)_n} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\beta; \mathfrak{z})|_{\gamma} \bar{R}_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\alpha}^{\beta}. \quad (6.5)$$

Here $\mu(u)$ is a scalar function defined by

$$\mu(u_1/u_2)\Phi_{\emptyset}(u_1)\Phi_{\emptyset}(u_2) = \Phi_{\emptyset}(u_2)\Phi_{\emptyset}(u_1). \quad (6.6)$$

It is explicitly calculated by using (4.11) as

$$\mu(u) = \frac{\Gamma(\hbar u; t_1, t_2, p)}{\Gamma(pu; t_1, t_2, p)},$$

where $\Gamma(z; t_1, t_2, p)$ denotes the triple Gamma function defined by

$$\begin{aligned} \Gamma(z; t_1, t_2, p) &= (z; t_1, t_2, p)_{\infty} (t_1 t_2 p / z; t_1, t_2, p)_{\infty}, \\ (z; t_1, t_2, p)_{\infty} &= \prod_{m_1, m_2, m_3=0}^{\infty} (1 - z t_1^{m_1} t_2^{m_2} p^{m_3}). \end{aligned}$$

By definition, the $R_{T^{1/2}}(u_1, u_2; \mathfrak{z})$ preserves the weight measured by c^\perp , i.e. the total number of the framing dimensions :

$$[R_{T^{1/2}}(u_1, u_2; \mathfrak{z}), c^{\perp(1)} + c^{\perp(2)}] = 0. \quad (6.7)$$

We also assume the property

$$R_{T^{1/2}}(u_1, u_2; \mathfrak{z} \hbar^{c^{\perp(1)} + c^{\perp(2)}}) = R_{T^{1/2}}(u_1, u_2; \mathfrak{z}). \quad (6.8)$$

See Remark below (6.14). Note also that $\widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\alpha}; \mathfrak{z})$ is depend on u_1, u_2 only through the chamber $\mathfrak{C} : u_1 \ll u_2$ essentially. Hence for any $a \in \mathbb{C}^\times$

$$R_{T^{1/2}}(au_1, au_2; \mathfrak{z}) = R_{T^{1/2}}(u_1, u_2; \mathfrak{z}). \quad (6.9)$$

Proposition 6.1. *The type I vertex operators satisfy the following exchange relation.*

$$\Phi_{\omega''}^{(\mu'')}(u_2) \Phi_{\omega'}^{(\mu')}(u_1) = \sum_{\boldsymbol{\lambda} = (\lambda', \lambda'') \in (\mathcal{P}^2)_n} R_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\omega}^{\boldsymbol{\lambda}} \Phi_{\lambda'}^{(\mu')}(u_1) \Phi_{\lambda''}^{(\mu'')}(u_2),$$

where $\omega = (\omega', \omega'') \in (\mathcal{P}^2)_n$.

Proof. Let us consider the RHS. By using (6.1), (6.6) and moving $\bar{R}_{\mathfrak{C}\bar{\mathfrak{C}}}(u_1, u_2; \mathfrak{z})_{\omega}^{\boldsymbol{\lambda}}$ to the right end, one obtains

$$\begin{aligned} & \sum_{\boldsymbol{\lambda} = (\lambda', \lambda'')} \prod_{a \in \lambda'} \left(\int_0^{\varphi_{\zeta(a)}^{\mu'}} d_p x_a \right) \prod_{a \in \lambda''} \left(\int_0^{\varphi_{\zeta(a)}^{\mu''}} d_p x_a \right) : \prod_{a \in \boldsymbol{\lambda}} x^-(x_a) : \Phi_{\emptyset}(u_2) \Phi_{\emptyset}(u_1) \\ & \times \prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a < \rho_b}} \langle x^-(x_a) x^-(x_b) \rangle^{sym} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) \bar{R}_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\omega}^{\boldsymbol{\lambda}}. \end{aligned}$$

Here we used (6.8). Note that : $\prod_{a \in \boldsymbol{\lambda}} x^-(x_a) :$ and $\prod_{\substack{a, b \in \boldsymbol{\lambda} \\ \rho_a < \rho_b}} \langle x^-(x_a) x^-(x_b) \rangle^{sym}$ are symmetric in x_a ($a \in \boldsymbol{\lambda}$) so that they are independent from a choice $\boldsymbol{\lambda} \in (\mathcal{P}^2)_n$. Evaluating the Jackson integral and using Proposition 5.1 and (6.5), one obtains

$$\begin{aligned} & \sum_{\mathbf{d} \in \mathbb{N}^n} \mathfrak{z}^{-\sum_{a \in \bullet} d_a} : \prod_{a \in \bullet} x^-(x_a) : \Big|_{x_a = \varphi_{\zeta(a)}^{\mu} p^{d_a}} \Phi_{\emptyset}(u_2) \Phi_{\emptyset}(u_1) \prod_{\substack{a, b \in \bullet \\ \rho_a < \rho_b}} \langle x^-(x_a) x^-(x_b) \rangle^{sym} \Big|_{x_a = \varphi_{\zeta(a)}^{\mu} p^{d_a}} \\ & \times \sum_{\boldsymbol{\lambda}} \widehat{\text{Stab}}_{\mathfrak{C}, T^{1/2}}(\boldsymbol{\lambda}; \mathfrak{z}) \Big|_{\boldsymbol{\mu}} \bar{R}_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\omega}^{\boldsymbol{\lambda}} \\ = & \sum_{\mathbf{d} \in \mathbb{N}^n} \mathfrak{z}^{-\sum_{a \in \bullet} d_a} : \prod_{a \in \bullet} x^-(x_a) : \Big|_{x_a = \varphi_{\zeta(a)}^{\mu} p^{d_a}} \Phi_{\emptyset}(u_2) \Phi_{\emptyset}(u_1) \\ & \times \prod_{\substack{a, b \in \bullet \\ \rho_a < \rho_b}} \langle x^-(x_a) x^-(x_b) \rangle^{sym} \Big|_{x_a = \varphi_{\zeta(a)}^{\mu} p^{d_a}} \times \widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T^{1/2}}(\bar{\boldsymbol{\omega}}; \mathfrak{z}) \Big|_{\bar{\boldsymbol{\mu}}} \\ = & \Phi_{\omega''}^{(\mu'')}(u_2) \Phi_{\omega'}^{(\mu')}(u_1). \end{aligned}$$

Here \bullet denotes any partition in $(\mathcal{P}^2)_n$. We also set $\{x_a = \varphi_{\zeta(a)}^{\mu} p^{d_a}\} := \{x'_a = \varphi_{\zeta(a)}^{\mu'} p^{d_a} (a \in \lambda')\} \cup \{x''_b = \varphi_{\zeta(b)}^{\mu''} p^{d_b} (b \in \lambda'')\}$. \square

6.1.1 Type II

From Sec.4.3.2 one has

$$\begin{aligned} \Psi_{\lambda''}^{*(\mu'')} (u_2) \Psi_{\lambda'}^{*(\mu')} (u_1) &= \prod_{a \in \lambda''} \left(\int_0^{\varphi_{\zeta(a)}^{\mu''}} d_p x''_a \right) \prod_{a \in \lambda'} \left(\int_0^{\varphi_{\zeta(a)}^{\mu'}} d_p x'_a \right) \widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*((\lambda', \lambda''); \mathfrak{z}^{*-1}) \\ &\times : \prod_{a \in (\lambda', \lambda'')} x^+(x_a) : \Psi_{\emptyset}^*(u_2) \Psi_{\emptyset}^*(u_1) \prod_{\substack{a, b \in (\lambda', \lambda'') \\ \rho_a < \rho_b}} \langle x^+(x_a) x^+(x_b) \rangle^{sym}, \end{aligned} \quad (6.10)$$

where the chamber \mathfrak{C} is given by $u_1 \ll u_2$ with the stability condition $t_1/t_2 > 0$. Hence $\rho_a < \rho_b$ for $a \in \lambda', b \in \lambda''$ due to (3.2). Similarly

$$\begin{aligned} \Psi_{\lambda'}^{*(\mu')} (u_1) \Psi_{\lambda''}^{*(\mu'')} (u_2) &= \prod_{a \in \lambda'} \left(\int_0^{\varphi_{\zeta(a)}^{\mu'}} d_p x'_a \right) \prod_{a \in \lambda''} \left(\int_0^{\varphi_{\zeta(a)}^{\mu''}} d_p x''_a \right) \widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T_{opp}^{1/2}}^*((\lambda'', \lambda'); \mathfrak{z}^{*-1}) \\ &\times : \prod_{a \in (\lambda', \lambda'')} x^+(x_a) : \Psi_{\emptyset}^*(u_1) \Psi_{\emptyset}^*(u_2) \prod_{\substack{a, b \in (\lambda', \lambda'') \\ \rho_a < \rho_b}} \langle x^+(x_a) x^+(x_b) \rangle^{sym}, \end{aligned} \quad (6.11)$$

where the chamber $\bar{\mathfrak{C}}$ is given by $u_1 \gg u_2$ with the stability condition $t_1/t_2 > 0$. Hence $\rho_a < \rho_b$ for $a \in \lambda'', b \in \lambda'$.

Let us define $R_{T_{opp}^{1/2}}^*(u_1, u_2; \mathfrak{z}^{*-1}) \in \text{End}_{\mathbb{F}}(\mathcal{F}_{u_1}^{(0,1)} \tilde{\otimes} \mathcal{F}_{u_2}^{(0,1)})$ as the following transition matrix of the elliptic stable envelopes $\widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\bullet; \mathfrak{z}^{*-1})$. For $\alpha, \beta, \gamma \in (\mathcal{P}^2)_n$,

$$R_{T_{opp}^{1/2}}^*(u_1, u_2; \mathfrak{z}^{*-1})_{\alpha}^{\beta} = \mu^*(u_1/u_2) \bar{R}_{T_{opp}^{1/2}}^*(u_1, u_2; \mathfrak{z}^{*-1})_{\alpha}^{\beta}, \quad (6.12)$$

$$\widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T_{opp}^{1/2}}^*(\bar{\alpha}; \mathfrak{z}^{*-1})|_{\bar{\gamma}} = \sum_{\beta \in (\mathcal{P}^2)_n} \widehat{\text{Stab}}_{\mathfrak{C}, T_{opp}^{1/2}}^*(\beta; \mathfrak{z}^{*-1})|_{\gamma} \bar{R}_{T_{opp}^{1/2}}^*(u_1, u_2; \mathfrak{z}^{*-1})_{\alpha}^{\beta}, \quad (6.13)$$

where $\mu^*(u)$ is a scalar function satisfying

$$\Psi_{\emptyset}^*(u_1) \Psi_{\emptyset}^*(u_2) = \mu^*(u_1/u_2) \Psi_{\emptyset}^*(u_2) \Psi_{\emptyset}^*(u_1). \quad (6.14)$$

Explicitly it is given by

$$\mu^*(u) = \frac{\Gamma(p^* \hbar u; t_1, t_2, p^*)}{\Gamma(u; t_1, t_2, p^*)}.$$

Remark. The $R_{T_{opp}^{1/2}}^*$ too satisfies (6.7) and (6.9). The relation between $R_{T_{1/2}}$ and $R_{T_{opp}^{1/2}}$ is obtained from Proposition 3.4 in [1], which states

$$\text{Stab}_{\bar{\mathfrak{C}}, T_{opp}^{1/2}}(\mathfrak{z}^{-1})^{\vee} \circ \text{Stab}_{\mathfrak{C}, T_{1/2}}(\mathfrak{z}) = 1.$$

Here \vee means transpose defined geometrically in [1]. We identify it as

$$\left(\widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T_{opp}^{1/2}}(\mathfrak{z}^{-1})^{\vee} \right)_{\alpha\gamma} = \widehat{\text{Stab}}_{\bar{\mathfrak{C}}, T_{opp}^{1/2}}(\bar{\alpha}; \mathfrak{z}^{-1})|_{\gamma}.$$

Applying this to (6.5) and (6.13) and noting that for any polarization, $R_{\bullet}^*(; \mathfrak{z}^*)$ is the same $R_{\bullet}(\cdot; \mathfrak{z})$ except for replacing p and \mathfrak{z} by p^* and \mathfrak{z}^* , respectively, one obtains the following relation.

Proposition 6.2.

$$R_{T^{1/2}}(u_1, u_2; \mathfrak{z}) = {}^t R_{T_{opp}^{1/2}}(u_1, u_2; \mathfrak{z}^{-1}). \quad (6.15)$$

The statement is valid for any Nakajima quiver variety X . The other explicit examples are all elliptic dynamical R -matrices associated with untwisted affine Lie algebras $\widehat{\mathfrak{g}}$ [24]. In the next section, we also show that $R_{T^{1/2}}(u_1, u_2; \mathfrak{z})$ and ${}^t R_{T_{opp}^{1/2}}^*(u_1, u_2; \mathfrak{z}^{*-1})$ satisfy the same dynamical Yang-Baxter equation under the assumption (6.8).

In the similar way to Proposition 6.1, one can prove the following statement.

Proposition 6.3.

$$\Psi_{\omega'}^{*(\mu')}(u_1) \Psi_{\omega''}^{*(\mu'')}(u_2) = \sum_{\lambda=(\lambda', \lambda'') \in (\mathcal{P}^2)_n} \Psi_{\lambda'}^{*(\mu'')}(u_2) \Psi_{\lambda'}^{*(\mu')}(u_1) R_{T^{1/2}}^*(u_1, u_2; \mathfrak{z}^*) \omega_{\lambda},$$

where $\omega = (\omega', \omega'') \in (\mathcal{P}^2)_n$.

Proof. Start from the RHS with (6.10) replacing $R_{T^{1/2}}^*(; \mathfrak{z}^*)$ with ${}^t R_{T_{opp}^{1/2}}^*(; \mathfrak{z}^{*-1})$ by (6.15), move $R_{T_{opp}^{1/2}}^*$ to the left end. Then use (6.13) and (6.11). \square

Finally, the type I and the type II dual vertex operators exchange by a scalar function.

Proposition 6.4. *In the level $(1, N)$ representation, one has*

$$\begin{aligned} \Phi_{\lambda}(u) \Psi_{\mu}^*(v) &= \chi(u/v) \Psi_{\mu}^*(v) \Phi_{\lambda}(u) \quad \forall \lambda, \mu \in \mathcal{P}_n, \\ \chi(u) &= \frac{1}{\Gamma(\hbar^{1/2}u; t_1, t_2)} = \Gamma(\hbar^{1/2}/u; t_1, t_2). \end{aligned}$$

Here $\Gamma(z; t_1, t_2)$ denotes the elliptic Gamma function given by

$$\Gamma(z; t_1, t_2) = \frac{(t_1 t_2 / z; t_1, t_2)_{\infty}}{(z; t_1, t_2)_{\infty}}, \quad (z; t_1, t_2)_{\infty} = \prod_{m_1, m_2=0}^{\infty} (1 - z t_1^{m_1} t_2^{m_2}).$$

Proof. From (4.13), (4.14), (4.11) and (4.12), one has

$$\Phi_{\emptyset}(u) \Psi_{\emptyset}^*(v) = \chi(u/v) \Psi_{\emptyset}^*(v) \Phi_{\emptyset}(u)$$

Then the statement follows from

$$[\Phi_{\emptyset}(u), x^+(z)] = 0 = [\Psi_{\emptyset}^*(u), x^-(z)].$$

\square

Note $\gamma = \hbar^{1/2}$ and $p^* = p\hbar^{-1}$ in the level $(1, N)$ representation and the formula²

$$\mu^*(u) = \left. \frac{\chi(\hbar^{1/2}/u)}{\chi(\hbar^{1/2}u)} \mu(u) \right|_{p \rightarrow p^*}. \quad (6.16)$$

²In this occasion let us correct a misprint in [22]. On page 431, the formula in (6.54) should read $\mu^*(v) = \frac{\chi(\frac{1}{2}-v)}{\chi(\frac{1}{2}+v)} \mu(v)|_{r \rightarrow r^*}$.

6.2 Dynamical Yang-Baxter equation

From the associativity of the composition of the type I and type II dual vertex operators, one can derive the dynamical Yang-Baxter equation for the elliptic R -matrices in (6.4) and (6.12).

For the type I vertex operator, using Proposition 6.1 for $\Phi_{\omega'''}^{(\mu''')}(u_3)\Phi_{\omega''}^{(\mu'')}(u_2)\Phi_{\omega'}^{(\mu')}(u_1)$ and assuming that $\Phi(u)$ is invertible, one obtains

$$\begin{aligned} & R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z}\hbar^{c^\perp(3)})R_{T^{1/2}}^{(13)}(u_1, u_3; \mathfrak{z})R_{T^{1/2}}^{(23)}(u_2, u_3; \mathfrak{z}\hbar^{c^\perp(1)}) \\ &= R_{T^{1/2}}^{(23)}(u_2, u_3; \mathfrak{z})R_{T^{1/2}}^{(13)}(u_1, u_3; \mathfrak{z}\hbar^{c^\perp(2)})R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z}), \end{aligned} \quad (6.17)$$

or

$$\begin{aligned} & R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z}\hbar^{-c^\perp(1)-c^\perp(2)})R_{T^{1/2}}^{(13)}(u_1, u_3; \mathfrak{z}\hbar^{-c^\perp(1)-c^\perp(2)-c^\perp(3)})R_{T^{1/2}}^{(23)}(u_2, u_3; \mathfrak{z}\hbar^{-c^\perp(2)-c^\perp(3)}) \\ &= R_{T^{1/2}}^{(23)}(u_2, u_3; \mathfrak{z}\hbar^{-c^\perp(1)-c^\perp(2)-c^\perp(3)})R_{T^{1/2}}^{(13)}(u_1, u_3; \mathfrak{z}\hbar^{-c^\perp(1)-c^\perp(3)})R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z}\hbar^{-c^\perp(1)-c^\perp(2)-c^\perp(3)}), \end{aligned} \quad (6.18)$$

depending on whether one puts operator $\Phi_{\bullet}^{(\mu''')}(u_1)\Phi_{\bullet}^{(\mu'')}(u_2)\Phi_{\bullet}^{(\mu')}(u_3)$ on the RHS of the coefficient “ RRR ” or on the other side. Here $R_{T^{1/2}}^{(ij)}(u_i, u_j; \mathfrak{z})$ are operators acting on $\mathcal{F}_{u_1}^{(0,1)} \tilde{\otimes} \mathcal{F}_{u_1}^{(0,1)} \tilde{\otimes} \mathcal{F}_{u_3}^{(0,1)}$ by

$$R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z})|\lambda'\rangle_{u_1} \tilde{\otimes} |\lambda''\rangle_{u_2} \tilde{\otimes} |\lambda'''\rangle_{u_3} = \sum_{\mu', \mu''} R_{T^{1/2}}(u_1, u_2; \mathfrak{z})_{\mu', \mu''}^{\lambda', \lambda''} |\mu'\rangle_{u_1} \tilde{\otimes} |\mu''\rangle_{u_2} \tilde{\otimes} |\lambda'''\rangle_{u_3}$$

etc. and $c^\perp(3) = 1 \tilde{\otimes} 1 \tilde{\otimes} c^\perp$ etc.. Note also $\hbar^{c^\perp}|\lambda\rangle_u = \hbar|\lambda\rangle_u$. The above two relations should be equivalent. In particular, if one assumes (6.8), one can rewrite (6.18) as

$$\begin{aligned} & R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z})R_{T^{1/2}}^{(13)}(u_1, u_3; \mathfrak{z}\hbar^{-c^\perp(2)})R_{T^{1/2}}^{(23)}(u_2, u_3; \mathfrak{z}) \\ &= R_{T^{1/2}}^{(23)}(u_2, u_3; \mathfrak{z}\hbar^{-c^\perp(1)})R_{T^{1/2}}^{(13)}(u_1, u_3; \mathfrak{z})R_{T^{1/2}}^{(12)}(u_1, u_2; \mathfrak{z}\hbar^{-c^\perp(3)}). \end{aligned} \quad (6.19)$$

By the same argument using Proposition 6.3 for the type II dual vertex operators, one finds that $R_{T^{1/2}}^*(u_1, u_2; \mathfrak{z}^*)$ satisfies (6.19) directly and (6.17) indirectly by applying the same assumption (6.8).

7 L -operator of $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$

For the elliptic dynamical instanton R -matrices introduced in the last section, we construct the L -operator L^+ on $\mathcal{F}_{\bullet}^{(1, N)}$ satisfying the $RLL = LLR^*$ relation. We then consider its universal form \mathcal{L}^+ and define the standard comultiplication Δ in terms of it and show that the vertex operators in Sec.4.2 are intertwining operators of the $U_{t_1, t_2, p}(\mathfrak{gl}_{1, tor})$ -modules w.r.t. Δ .

7.1 L -operator on $\mathcal{F}_\bullet^{(1,N)}$

Let $\sigma^{op} : \xi \tilde{\otimes} \eta \rightarrow \eta \tilde{\otimes} \xi$ and consider the following composition of the type I and type II vertex operators.

$$\mathcal{F}_u^{(0,1)} \tilde{\otimes} \mathcal{F}_{\hbar^{1/2}v}^{(1,N)} \xrightarrow{\sigma^{op}} \mathcal{F}_{\hbar^{1/2}v}^{(1,N)} \tilde{\otimes} \mathcal{F}_u^{(0,1)} \xrightarrow{\Phi(\hbar^{1/2}u) \tilde{\otimes} \text{id}} \mathcal{F}_{\hbar^{1/2}u}^{(0,1)} \tilde{\otimes} \mathcal{F}_{-v/u}^{(1,N-1)} \tilde{\otimes} \mathcal{F}_u^{(0,1)} \xrightarrow{\text{id} \tilde{\otimes} \Psi^*(u)} \mathcal{F}_{\hbar^{1/2}u}^{(0,1)} \tilde{\otimes} \mathcal{F}_v^{(1,N)}.$$

Hence we have the operator

$$L^+(u) := g(\text{id} \tilde{\otimes} \Psi^*(u)) \circ (\Phi(\hbar^{1/2}u) \tilde{\otimes} \text{id}) \sigma^{op} : \mathcal{F}_u^{(0,1)} \tilde{\otimes} \mathcal{F}_{\hbar^{1/2}v}^{(1,N)} \rightarrow \mathcal{F}_{\hbar^{1/2}u}^{(0,1)} \tilde{\otimes} \mathcal{F}_v^{(1,N)}$$

for $N \in \mathbb{Z}$, $v \in \mathbb{C}^\times$. Here we set

$$g = (\hbar; t_1, t_2)_\infty.$$

Define the components of $L^+(u)$ by

$$L^+(u) \cdot |\nu\rangle_u \tilde{\otimes} \xi = \sum_{\mu} |\mu\rangle_{\hbar^{1/2}u} \tilde{\otimes} L_{\mu\nu}^+(u) \xi,$$

for $|\mu\rangle_u \tilde{\otimes} \xi \in \mathcal{F}_u^{(0,1)} \tilde{\otimes} \mathcal{F}_{\hbar^{1/2}v}^{(1,N)}$. On the other hand one obtains

$$\begin{aligned} L^+(u) \cdot |\nu\rangle_u \tilde{\otimes} \xi &= g(\text{id} \tilde{\otimes} \Psi^*(u)) \circ (\Phi(\hbar^{1/2}u) \tilde{\otimes} \text{id}) \cdot \xi \tilde{\otimes} |\nu\rangle_u \\ &= g(\text{id} \tilde{\otimes} \Psi^*(u)) \cdot \sum_{\mu} |\mu\rangle_{\hbar^{1/2}u} \tilde{\otimes} \Phi_{\mu}(\hbar^{1/2}u) \xi \tilde{\otimes} |\nu\rangle_u \\ &= g \sum_{\mu} |\mu\rangle_{\hbar^{1/2}u} \tilde{\otimes} \Psi_{\nu}^*(u) \Phi_{\mu}(\hbar^{1/2}u) \xi. \end{aligned}$$

Hence

$$L_{\mu\nu}^+(u) = g \Psi_{\nu}^*(u) \Phi_{\mu}(\hbar^{1/2}u). \quad (7.1)$$

In particular, setting $k_{\emptyset}^+(u) = L_{\emptyset\emptyset}^+(u)$, one has

$$\begin{aligned} k_{\emptyset}^+(u) &= g \Psi_{\emptyset}^*(u) \Phi_{\emptyset}(\hbar^{1/2}u) \\ &= \hbar^{-\Lambda_0/2} e^{-Q} : \exp \left\{ - \sum_{m \neq 0} \frac{p^m}{(1-t_1^m)(1-t_2^m)(1-p^{*m})} \alpha_m(\hbar^{1/2}u)^{-m} \right\} : . \end{aligned}$$

By using (2.3), one obtains the following commutation relation.

Proposition 7.1.

$$k_{\emptyset}^+(u) k_{\emptyset}^+(v) = \rho(u/v) k_{\emptyset}^+(v) k_{\emptyset}^+(u), \quad (7.2)$$

$$\rho(u) = \frac{\rho^{+*}(u)}{\rho^+(u)}, \quad \rho^+(u) = \frac{\Gamma(\hbar u; t_1, t_2, p)}{\Gamma(u; t_1, t_2, p)}, \quad \rho^{+*}(u) = \rho^+(u) \Big|_{p \rightarrow p^*}. \quad (7.3)$$

Let us consider the following elliptic dynamical R -matrices.

$$R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\alpha}^{\beta} := \rho^+(u/v) \bar{R}_{T^{1/2}}(u, v; \mathfrak{z})_{\alpha}^{\beta}, \quad (7.4)$$

where $\bar{R}_{T^{1/2}}$ is given in (6.5) and $R_{T^{1/2}}^{+*} = R_{T^{1/2}}^+|_{p \rightarrow p^*}$.

Proposition 7.2. *The L^+ operator satisfies the following relation.*

$$\sum_{\mu', \nu'} R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} L_{\mu'\mu''}^+(u) L_{\nu'\nu''}^+(v) = \sum_{\mu', \nu'} L_{\nu\nu'}^+(v) L_{\mu\mu'}^+(u) R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''} \quad (7.5)$$

Proof. From (6.9) and Propositions 6.1, 6.3, 6.4, one has

$$\begin{aligned} & \sum_{\mu', \nu'} R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} L_{\mu'\mu''}^+(u) L_{\nu'\nu''}^+(v) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g^2 \sum_{\mu', \nu'} R_{T^{1/2}}(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} \Psi_{\mu''}^*(u) \Phi_{\mu'}(\hbar^{1/2}u) \Psi_{\nu''}^*(v) \Phi_{\nu'}(\hbar^{1/2}v) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g^2 \chi(\hbar^{1/2}u/v) \sum_{\mu', \nu'} R_{T^{1/2}}(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} \Psi_{\mu''}^*(u) \Psi_{\nu''}^*(v) \Phi_{\mu'}(\hbar^{1/2}u) \Phi_{\nu'}(\hbar^{1/2}v) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g^2 \chi(\hbar^{1/2}u/v) \Psi_{\mu''}^*(u) \Psi_{\nu''}^*(v) \Phi_{\nu}(\hbar^{1/2}v) \Phi_{\mu}(\hbar^{1/2}u) \quad (7.6) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g^2 \chi(\hbar^{1/2}u/v) \sum_{\mu', \nu'} \Psi_{\nu'}^*(v) \Psi_{\mu'}^*(u) \Phi_{\nu}(\hbar^{1/2}v) \Phi_{\mu}(\hbar^{1/2}u) R_{T^{1/2}}^*(u, v; \mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''} \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g^2 \chi(\hbar^{1/2}u/v) \chi(\hbar^{1/2}v/u)^{-1} \sum_{\mu', \nu'} \Psi_{\nu'}^*(v) \Phi_{\nu}(\hbar^{1/2}v) \Psi_{\mu'}^*(u) \Phi_{\mu}(\hbar^{1/2}u) R_{T^{1/2}}^*(u, v; \mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''} \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} \chi(\hbar^{1/2}u/v) \chi(\hbar^{1/2}v/u)^{-1} \sum_{\mu', \nu'} L_{\nu\nu'}^+(v) L_{\mu\mu'}^+(u) \frac{\mu^*(u/v)}{\rho^{+*}(u/v)} R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''} \\ &= \sum_{\mu', \nu'} L_{\nu\nu'}^+(v) L_{\mu\mu'}^+(u) R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''}. \end{aligned}$$

In the last equality we used

$$\frac{\rho^+(u)}{\rho^{+*}(u)} = \frac{\chi(\hbar^{1/2}/u)}{\chi(\hbar^{1/2}u)} \frac{\mu(u)}{\mu^*(u)}.$$

□

7.2 Intertwining relations

We next derive the exchange relation between the vertex operators and $L^+(u)$. They turns out to be the intertwining relations w.r.t. the standard comultiplication defined in terms of a universal form of $L^+(u)$.

Proposition 7.3. *The type I and the type II vertex operators satisfy the following relations.*

$$\Phi_\nu(\hbar^{1/2}v)L_{\mu\mu''}^+(u) = \sum_{\mu'\nu'} R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} L_{\mu'\mu''}^+(u)\Phi_{\nu'}(\hbar^{1/2}v), \quad (7.7)$$

$$L_{\mu\mu''}^+(u)\Psi_{\nu''}^*(v) = \sum_{\mu'\nu'} \Psi_{\nu'}^*(v)L_{\mu\mu'}^+(u)R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^*)_{\mu'\nu''}^{\mu''\nu''}. \quad (7.8)$$

Proof. From the derivation of the first half of (7.6), one has

$$\begin{aligned} & g \sum_{\mu', \nu'} R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} \Psi_{\mu''}^*(u)\Phi_{\mu'}(\hbar^{1/2}u)\Psi_{\nu''}^*(\tilde{v})\Phi_{\nu'}(\hbar^{1/2}v) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g\chi(\hbar^{1/2}u/\tilde{v})\Psi_{\mu''}^*(u)\Psi_{\nu''}^*(\tilde{v})\Phi_\nu(\hbar^{1/2}v)\Phi_\mu(\hbar^{1/2}u). \end{aligned}$$

Make the following manipulation.

$$\begin{aligned} \text{LHS} &= \sum_{\mu', \nu'} R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} L_{\mu'\mu''}^+(u)\chi(\hbar^{1/2}v/\tilde{v})^{-1}\Phi_{\nu'}(\hbar^{1/2}v)\Psi_{\nu''}^*(\tilde{v}), \\ \text{RHS} &= \frac{\rho^+(u/v)}{\mu(u/v)} g\chi(\hbar^{1/2}u/\tilde{v})\chi(\hbar^{1/2}v/\tilde{v})^{-1}\Psi_{\mu''}^*(u)\Phi_\nu(\hbar^{1/2}v)\Psi_{\nu''}^*(\tilde{v})\Phi_\mu(\hbar^{1/2}u) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} g\chi(\hbar^{1/2}u/\tilde{v})\chi(\hbar^{1/2}v/\tilde{v})^{-1}\chi(\hbar^{1/2}v/u)^{-1}\chi(\hbar^{1/2}u/\tilde{v})^{-1} \\ &\quad \times \Phi_\nu(\hbar^{1/2}v)\Psi_{\mu''}^*(u)\Phi_\mu(\hbar^{1/2}u)\Psi_{\nu''}^*(\tilde{v}) \\ &= \frac{\rho^+(u/v)}{\mu(u/v)} \chi(\hbar^{1/2}v/\tilde{v})^{-1}\chi(\hbar^{1/2}v/u)^{-1}\Phi_\nu(\hbar^{1/2}v)L_{\mu\mu''}^+(u)\Psi_{\nu''}^*(\tilde{v}). \end{aligned}$$

Therefore we have

$$\sum_{\mu', \nu'} R_{T^{1/2}}^+(u, v; \mathfrak{z})_{\mu\nu}^{\mu'\nu'} L_{\mu'\mu''}^+(u)\Phi_{\nu'}(\hbar^{1/2}v) = \Phi_\nu(\hbar^{1/2}v)L_{\mu\mu''}^+(u).$$

Here we used

$$\rho^+(u/v) = \mu(u/v)\chi(\hbar^{1/2}v/u).$$

Similarly, from the derivation of the second half of (7.6), one has

$$\begin{aligned} & g \sum_{\mu', \nu'} \Psi_{\nu'}^*(v)\Phi_\nu(\hbar^{1/2}\tilde{v})\Psi_{\mu'}^*(u)\Phi_\mu(\hbar^{1/2}u)R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^*)_{\mu'\nu''}^{\mu''\nu''} \\ &= \frac{\rho^{+*}(u/v)}{\mu^*(u/v)} g\chi(\hbar^{1/2}v/u)\Psi_{\mu''}^*(u)\Psi_{\nu''}^*(v)\Phi_\nu(\hbar^{1/2}\tilde{v})\Phi_\mu(\hbar^{1/2}u) \end{aligned}$$

for $\tilde{v} = v$. Then one obtains

$$\begin{aligned}
\text{LHS} &= \chi(\hbar^{1/2}\tilde{v}/v)^{-1}\Phi_\nu(\hbar^{1/2}\tilde{v})\sum_{\mu',\nu'}\Psi_{\nu'}^*(v)L_{\mu\mu'}^+(u)R_{T^{1/2}}^{+*}(u,v;\mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''}, \\
\text{RHS} &= \frac{\rho^{+*}(u/v)}{\mu^*(u/v)}g\chi(\hbar^{1/2}v/u)\chi(\hbar^{1/2}\tilde{v}/u)^{-1}\chi(\hbar^{1/2}\tilde{v}/v)^{-1}\Phi_\nu(\hbar^{1/2}\tilde{v})\Psi_{\mu''}^*(u)\Psi_{\nu''}^*(v)\Phi_\mu(\hbar^{1/2}u) \\
&= \frac{\rho^{+*}(u/v)}{\mu^*(u/v)}g\chi(\hbar^{1/2}v/u)\chi(\hbar^{1/2}\tilde{v}/u)^{-1}\chi(\hbar^{1/2}\tilde{v}/v)^{-1}\chi(\hbar^{1/2}u/v)^{-1} \\
&\quad \times \Phi_\nu(\hbar^{1/2}\tilde{v})\Psi_{\mu''}^*(u)\Phi_\mu(\hbar^{1/2}u)\Psi_{\nu''}^*(v) \\
&= \frac{\rho^{+*}(u/v)}{\mu^*(u/v)}\chi(\hbar^{1/2}v/u)\chi(\hbar^{1/2}\tilde{v}/u)^{-1}\chi(\hbar^{1/2}\tilde{v}/v)^{-1}\chi(\hbar^{1/2}u/v)^{-1}\Phi_\nu(\hbar^{1/2}\tilde{v})L_{\mu\mu''}^+(u)\Psi_{\nu''}^*(v).
\end{aligned}$$

Therefore we have

$$\begin{aligned}
&\sum_{\mu',\nu'}\Psi_{\nu'}^*(v)L_{\mu\mu'}^+(u)R_{T^{1/2}}^{+*}(u,v;\mathfrak{z}^*)_{\mu'\nu'}^{\mu''\nu''} \\
&= \frac{\rho^{+*}(u/v)}{\mu^*(u/v)}\chi(\hbar^{1/2}v/u)\chi(\hbar^{1/2}\tilde{v}/u)^{-1}\chi(\hbar^{1/2}u/v)^{-1}L_{\mu\mu''}^+(u)\Psi_{\nu''}^*(v).
\end{aligned}$$

After taking the limit $\tilde{v} \rightarrow v$, use

$$\begin{aligned}
\mu^*(u) &= \mu(u)\Big|_{p \rightarrow p^*} \times \chi(\hbar^{1/2}/u)\chi(\hbar^{1/2}u)^{-1}, \\
\rho^{+*}(u) &= \mu(u)\Big|_{p \rightarrow p^*} \times \chi(\hbar^{1/2}/u).
\end{aligned}$$

One gets the desired result. \square

Finally let us consider a coalgebra structure which allows us to interpret the relations in Proposition 7.3 as intertwining relations on the modules of $\mathcal{U} = U_{t_1,t_2,p}(\mathfrak{gl}_{1,tor})$. Let $\mathcal{L}^+(u) \in \text{End}_{\mathbb{F}}(\mathcal{F}_u^{(0,1)}) \otimes \tilde{\mathcal{U}}$ be the universal L -operator with

$$\mathcal{L}^+(u) \cdot |\nu\rangle_u \otimes \tilde{\eta} = \sum_{\mu \in \mathcal{P}} |\mu\rangle_u \otimes \tilde{\mathcal{L}}_{\mu\nu}^+(u)\eta, \quad \eta \in \mathcal{U}.$$

We assume $\mathcal{L}_{\mu\nu}^+(u)$ satisfies the RLL -relation (7.5). Let us further set $\mathcal{L}^+(u) = \mathcal{L}^+(u;\mathfrak{z}^*)e^{-c^\perp} \otimes \tilde{\mathcal{Q}}$ with

$$\mathcal{L}^+(u;\mathfrak{z}^*)e^{-c^\perp} \otimes \tilde{\mathcal{Q}} \cdot |\nu\rangle_u \otimes \tilde{\eta} = \sum_{\mu \in \mathcal{P}} |\mu\rangle_u \otimes \tilde{\mathcal{L}}_{\mu\nu}^+(u;\mathfrak{z}^*)e^{-Q}\eta.$$

Then due to (2.1), one obtains the following dynamical RLL -relation.

$$\begin{aligned}
&R_{T^{1/2}}^{+(12)}(u,v;\mathfrak{z}^*\hbar^{c^\perp(3)})\mathcal{L}^{+(1)}(u;\mathfrak{z}^*)\mathcal{L}^{+(2)}(v;\mathfrak{z}^*\hbar^{c^\perp(1)}) \\
&= \mathcal{L}^{+(2)}(v;\mathfrak{z}^*)\mathcal{L}^{+(1)}(u;\mathfrak{z}^*\hbar^{c^\perp(2)})R_{T^{1/2}}^{+*(12)}(u,v;\mathfrak{z}^*).
\end{aligned} \tag{7.9}$$

We hence assume $e^{\pm Q} \cdot |\lambda\rangle_u = |\lambda\rangle_u$ and

$$\begin{aligned}\mathcal{L}_{\mu\nu}^+(u) \cdot |\lambda\rangle_v &= \mathcal{L}_{\mu\nu}^+(u; \mathfrak{z}^*) \cdot |\lambda\rangle_v = \sum_{\lambda'} R_{T^{1/2}}^+(u, v; \mathfrak{z}^*)_{\mu\lambda'}^{\nu\lambda} |\lambda'\rangle_v, & |\lambda\rangle_v \in \mathcal{F}_v^{(0,1)}, \\ \mathcal{L}_{\mu\nu}^+(u) \cdot \xi &= L_{\mu\nu}^+(u)\xi, & \xi \in \mathcal{F}_{\bullet}^{(1,N)}.\end{aligned}$$

Define a new comultiplication Δ by

$$\Delta(\mathcal{L}_{\mu\nu}^+(u)) = \sum_{\lambda} \mathcal{L}_{\mu\lambda}^+(u\gamma^{(2)}) \tilde{\otimes} \mathcal{L}_{\lambda\nu}^+(u), \quad (7.10)$$

$$\Delta(g(\mathfrak{z})) = g(\mathfrak{z}) \tilde{\otimes} 1, \quad \Delta(g(\mathfrak{z}^*)) = 1 \tilde{\otimes} g(\mathfrak{z}), \quad \forall g(\mathfrak{z}), g(\mathfrak{z}^*) \in \mathbb{F}, \quad (7.11)$$

$$\Delta(\gamma^{1/2}) = \gamma^{1/2} \tilde{\otimes} \gamma^{1/2}, \quad \Delta(C) = C \tilde{\otimes} C. \quad (7.12)$$

We call this the standard comultiplication. Defining further a counit ε and an antipode S in terms of $\mathcal{L}^+(u)$ in the same way as in Sec.4.2 of [27], one can show that $(U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}}), \Delta, \varepsilon, \mu_l, \mu_r, S)$ is a Hopf algebroid.

Proposition 7.4. *Relations in Proposition 7.3 are the intertwining relations w.r.t. Δ for $\Phi(u)$ and $\Psi^*(u)$, respectively i.e.*

$$\Phi(v)\mathcal{L}_{\mu\nu}^+(u) = \Delta(\mathcal{L}_{\mu\nu}^+(u))\Phi(v), \quad (7.13)$$

$$\mathcal{L}_{\mu\nu}^+(u)\Psi^*(v)(\xi \tilde{\otimes} |\lambda\rangle_v) = \Psi^*(v)\Delta(\mathcal{L}_{\mu\nu}^+(u))(\xi \tilde{\otimes} |\lambda\rangle_v), \quad \xi \in \mathcal{F}_{\bullet}^{(1,N)}, |\lambda\rangle_v \in \mathcal{F}_v^{(0,1)}. \quad (7.14)$$

Proof. From (7.13), one has

$$\begin{aligned}\text{LHS} \cdot \xi &= \sum_{\lambda} |\lambda\rangle_v \tilde{\otimes} \Phi_{\lambda}(v) L_{\mu\nu}^+(u)\xi, \\ \text{RHS} \cdot \xi &= \sum_{\mu'} \mathcal{L}_{\mu\mu'}^+(\gamma^{(2)}u) \tilde{\otimes} \mathcal{L}_{\mu'\nu}^+(u) \cdot \sum_{\lambda'} |\lambda'\rangle_v \tilde{\otimes} \Phi_{\lambda'}(v)\xi \\ &= \sum_{\mu', \lambda', \lambda} R_{T^{1/2}}^+(\hbar^{1/2}u, v; \mathfrak{z}^*)_{\mu\lambda}^{\mu'\lambda'} |\lambda\rangle_v \tilde{\otimes} L_{\mu'\nu}^+(u)\Phi_{\lambda'}(v)\xi \\ &= \sum_{\lambda} |\lambda\rangle_v \tilde{\otimes} \sum_{\mu', \lambda'} R_{T^{1/2}}^+(\hbar^{1/2}u, v; \mathfrak{z}^*)_{\mu\lambda}^{\mu'\lambda'} L_{\mu'\nu}^+(u)\Phi_{\lambda'}(v)\xi.\end{aligned}$$

In the last equality we used (2.15). Similarly, from (7.14), one obtains

$$\begin{aligned}\text{LHS} &= L_{\mu\nu}^+(u)\Psi_{\lambda}^*(v)\xi, \\ \text{RHS} &= \Psi^*(v) \cdot \sum_{\mu'} \mathcal{L}_{\mu\mu'}^+(\gamma^{(2)}u) \tilde{\otimes} \mathcal{L}_{\mu'\nu}^+(u) \cdot (\xi \tilde{\otimes} |\lambda\rangle_v) \\ &= \Psi^*(v) \cdot \sum_{\mu', \lambda'} L_{\mu\mu'}^+(u)\xi \tilde{\otimes} R_{T^{1/2}}^+(u, v; \mathfrak{z}^*)_{\mu\lambda}^{\mu'\lambda'} |\lambda'\rangle_v\end{aligned}$$

$$\begin{aligned}
&= \Psi^*(v) \cdot \sum_{\mu', \lambda'} R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^* \hbar^{-c^\perp(3)})_{\mu' \lambda'}^{\nu \lambda} L_{\mu \mu'}^+(u) \xi \tilde{\otimes} |\lambda'\rangle_v \\
&= \Psi^*(v) \cdot \sum_{\mu', \lambda'} L_{\mu \mu'}^+(u) R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^* \hbar^{-c^\perp(1) - c^\perp(3)})_{\mu' \lambda'}^{\nu \lambda} \xi \tilde{\otimes} |\lambda'\rangle_v \\
&= \sum_{\mu', \lambda'} \Psi_{\lambda'}^*(v) L_{\mu \mu'}^+(u) R_{T^{1/2}}^{+*}(u, v; \mathfrak{z}^*)_{\mu' \lambda'}^{\nu \lambda} \xi.
\end{aligned}$$

To obtain the fourth line, we used $\mathfrak{z}^* = \mathfrak{z} \hbar^{-c^\perp}$ and (2.15). Note also $p^* = p \hbar^{-1}$ on $\mathcal{F}_\bullet^{(1, N)}$. \square

Remark. The $\gamma^{(2)}$ -shift in (7.10) and the resultant intertwining relation (7.7) is consistent to those in [18, 27] for elliptic quantum groups associated with affine Lie algebras. In particular, in [25, 28] the comultiplication formula without the $\gamma^{(2)}$ -shift [27] was considered in a derivation of the vertex operators of $U_{q, p}(\widehat{\mathfrak{sl}}_N)$. There, instead, the components of the vertex operators was defined with the shift.

Acknowledgements

The authors would like to thank Tatsuyuki Hikita, Taro Kimura, Vitaly Tarasov and Yutaka Yoshida for useful discussions. H. Konno is supported by the Grant-in-Aid for Scientific Research (C) 23K03029 JSPS, Japan. Work of A. Smirnov is supported in part by the NSF under grant DMS - 2054527.

A Vertex Operators w.r.t. Δ^D

For a comparison of the vertex operators presented in this paper with those obtained in the previous paper [32], we summarize the results on the latter. They, both the type I and the type II dual vertex operators, were constructed as intertwining operators of the same $U_{t_1, t_2, p}(\mathfrak{gl}_{1, \text{tor}})$ -modules but w.r.t the Drinfeld comultiplication Δ^D given in Sec.2.2.

The type I $\Phi^D(u)$ and type II dual $\Psi^{D*}(u)$ vertex operators w.r.t. Δ^D , are the intertwining operators defined by

$$\begin{aligned}
\Phi^D(u) &: \mathcal{F}_v^{(1, N)} \rightarrow \mathcal{F}_u^{(0, 1)} \tilde{\otimes} \mathcal{F}_{-v/u}^{(1, N-1)}, \\
\Psi^{D*}(u) &: \mathcal{F}_v^{(1, N)} \tilde{\otimes} \mathcal{F}_u^{(0, 1)} \rightarrow \mathcal{F}_{-vu}^{(1, N+1)}
\end{aligned}$$

satisfying the intertwining relations

$$\Delta^D(x) \Phi^D(u) = \Phi^D(u) x, \tag{A.1}$$

$$x \Psi^{D*}(v) = \Psi^{D*}(v) \Delta^D(x) \quad \forall x \in \mathcal{U}. \tag{A.2}$$

Their components are defined by

$$\Phi^D(u)\xi = \sum_{\lambda \in \mathcal{P}^+} |\lambda\rangle'_u \tilde{\otimes} \Phi_\lambda^D(u)\xi, \quad (\text{A.3})$$

$$\Psi_\lambda^{D*}(u)\xi = \Psi^{D*}(u) (\xi \tilde{\otimes} |\lambda\rangle_u) \quad \forall \xi \in \mathcal{F}_v^{(1,N)}, \quad (\text{A.4})$$

where we set

$$|\lambda\rangle'_u = \frac{c_\lambda(p)}{c'_\lambda(p)} |\lambda\rangle_u, \quad (\text{A.5})$$

$$c_\lambda(p) = (-t_2)^{|\lambda|} t_1^{-n(\lambda')} t_2^{n(\lambda)} \prod_{\square \in \lambda} \theta(t_1^{-a(\square)} t_2^{\ell(\square)+1}), \quad (\text{A.6})$$

$$c'_\lambda(p) = (-t_1)^{-|\lambda|} t_1^{-n(\lambda')} t_2^{n(\lambda)} \prod_{\square \in \lambda} \theta(t_1^{-a(\square)-1} t_2^{\ell(\square)}) \quad (\text{A.7})$$

with $a(\square) = \lambda_i - j$, $\ell(\square) = \lambda'_j - i$, $n(\lambda) = \sum_{i \geq 1} (i-1)\lambda_i$, $n(\lambda') = \sum_{i \geq 1} (i-1)\lambda'_i$ for $\square = (i, j) \in \lambda$.

Solving the intertwining relations by using the representations in Theorem 2.5 and Theorem 2.6, one obtains the following formula.

Theorem A.1. [32]

$$\Phi_\lambda^D(u) = \frac{t_1^{-n(\lambda')-|\lambda|} N_\lambda(p)}{c_\lambda} : \tilde{\Phi}_\emptyset(u) \prod_{\square \in \lambda} x^-(\varphi_\square^\lambda u) : (-u)^{-\Lambda_0} e^{-\Lambda_0^\perp}, \quad (\text{A.8})$$

$$\Psi_\lambda^{D*}(u) = \frac{t_1^{-n(\lambda')}}{c_\lambda N'_\lambda(p^*)} : \tilde{\Psi}_\emptyset^*(u) \prod_{\square \in \lambda} x^+(\varphi_\square^\lambda u) : (-u)^{\Lambda_0} e^{\Lambda_0^\perp}, \quad (\text{A.9})$$

where $\tilde{\Phi}_\emptyset(u)$ and $\tilde{\Psi}_\emptyset^*(u)$ are given in (4.13) and (4.14), respectively. We set also $\varphi_\square^\lambda = t_1^{-j+1} t_2^{-i+1}$ for $\square = (i, j) \in \lambda$ and

$$c_\lambda = \prod_{\square \in \lambda} (1 - t_1^{-a(\square)} t_2^{\ell(\square)+1}), \quad c'_\lambda = \prod_{\square \in \lambda} (1 - t_1^{-a(\square)+1} t_2^{\ell(\square)}),$$

$$N_\lambda(p) = \prod_{\square \in \lambda} \frac{(pt_1^{-a(\square)-1} t_2^{\ell(\square)}; p)_\infty}{(pq^{a(\square)} t^{\ell(\square)+1}; p)_\infty}, \quad N'_\lambda(p) = \prod_{\square \in \lambda} \frac{(pt_1^{a(\square)} t_2^{-\ell(\square)-1}; p)_\infty}{(pq^{-a(\square)-1} t^{-\ell(\square)}; p)_\infty}.$$

Remark. The formulas (A.8) and (A.9) are reformulation of $\Phi_\lambda(u)$ and $\Psi_\lambda^*(u)$ given in [32] by introducing the zero mode operators c, Λ_0 and c^\perp, Λ_0^\perp . The combinatorial factors c_λ, c'_λ are the standard objects in the Macdonald theory [34] and related to an elliptic analogue of them $c_\lambda(p), c'_\lambda(p)$ as follows [32].

$$\frac{c'_\lambda N_\lambda(p)}{c_\lambda N'_\lambda(p)} = \frac{c'_\lambda(p)}{c_\lambda(p)}. \quad (\text{A.10})$$

References

- [1] M.Aganagic and A.Okounkov, Elliptic Stable Envelopes, *J. Amer. Math. Soc.* **34** (2021), no. 1, 79–133.
- [2] T.M.Botta, Shuffle Products for Elliptic Stable Envelopes of Nakajima Varieties, (2022), arXiv:2104.00976
- [3] H.Dinkins, Elliptic Stable Envelopes of Affine Type A Quiver Varieties, *Int.Math.Res.Not.* (2023), no.15, 13426–13476, arXiv:2107.09569
- [4] H.Dinkins, Exotic Quantum Difference Equations and Integral Solutions, PhD Thesis, University of North Carolina at Chapel Hill, 2022, arXiv:2205.01596
- [5] V.G. Drinfeld, A New Realization of Yangians and Quantized Affine Algebras. *Soviet Math. Dokl.* **36** (1988) 212-216.
- [6] P.Etingof and A.Varchenko, Solutions of the Quantum Dynamical Yang-Baxter Equation and Dynamical Quantum Groups, *Comm.Math.Phys.* **196** (1998) 591–640; Exchange Dynamical Quantum Groups, *Comm.Math.Phys.* **205** (1999) 19–52.
- [7] R.M.Farghly, H.Konno and K.Oshima, Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{g}})$ and Quantum Z -algebras, *Alg.Rep.Theory* **18** (2014) 103–135. DOI 10.1007/s10468-014-9483-x.
- [8] B.Feigin and E.Frenkel, Duality in W -Algebras, *Int.Math.Res.Not.* (1991) no.6, 75–82.
- [9] B. Feigin, M.Jimbo, T.Miwa and E.Mukhin, Quantum Toroidal \mathfrak{gl}_1 : Plane Partitions, *Kyoto J.Math.* **52** (2012) 621–659.
- [10] B. Feigin, M.Jimbo, T.Miwa and E.Mukhin, Quantum Toroidal \mathfrak{gl}_1 and Bethe Ansatz, *J.Phys.A* **48** (2015) 244001, 27pp.
- [11] B.Feigin and A.Tsybaliuk, Heisenberg action in the equivariant K -theory of Hilbert schemes via shuffle algebra, *Kyoto J.Math.* **51** (2011) 831–854.
- [12] G. Felder, Elliptic Quantum Groups, *Proc. ICMP Paris-1994* (1995), 211–218.
- [13] O.Foda, K.Iohara, M.Jimbo, R.Kedem, T.Miwa, H.Yan, Notes on Highest Weight Modules of the Elliptic Algebra $\mathcal{A}_{q,p}(\widehat{sl}_2)$. *Quantum field theory, integrable models and beyond* (Kyoto, 1994). *Progr. Theoret. Phys. Suppl.* **118** (1995) 1-34.

- [14] E.Frenkel and N.Reshetikhin, Deformation of W -Algebras Associated to Simple Lie Algebras, *Commun. Math. Phys.* **197** (1998) 1–31. q-alg/9708006.
- [15] T.Hollowood, A.Iqbal and C.Vafa, Matrix Models, Geometric Engineering and Elliptic Genera, *J.High Energy Phys.* (2008) 069.
- [16] A.Iqbal, C.Kozcaz and C.Vafa, The Refined Topological Vertex, *J.High Energy Phys.* (2009) 069, 58pp.
- [17] M. Jimbo, T. Miwa and M. Okado. Solvable Lattice Models Related to the Vector Representation of Classical Simple Lie Algebras, *Comm. Math. Phys.*, **116** (1988) 507–525.
- [18] M. Jimbo, H. Konno, S. Odake and J. Shiraishi, Quasi-Hopf Twistors for Elliptic Quantum Groups, *Transformation Groups* **4** (1999) 303–327.
- [19] M. Jimbo, H. Konno, S. Odake and J. Shiraishi, Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{sl}}_2)$: Drinfeld Currents and Vertex Operators, *Comm. Math. Phys.* **199** (1999) 605–647.
- [20] T.Kimura and V.Pestun, Quiver W -algebras, *Lett.Math.Phys.* **108** (2018), 1351–1381.
- [21] E. Koelink and H. Rosengren, Harmonic Analysis on the $SU(2)$ Dynamical Quantum Group, *Acta.Appl.Math.*, **69**, 2001, 163–220.
- [22] T. Kojima and H. Konno, The Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ and the Drinfeld Realization of the Elliptic Quantum Group $\mathcal{B}_{q,\lambda}(\widehat{\mathfrak{sl}}_N)$, *Comm.Math.Phys.***239** (2003) 405-447.
- [23] H. Konno, An Elliptic Algebra $U_{q,p}(\widehat{\mathfrak{sl}}_2)$ and the Fusion RSOS Models, *Comm. Math. Phys.* **195** (1998) 373–403.
- [24] H. Konno, Dynamical R Matrices of Elliptic Quantum Groups and Connection Matrices for the q -KZ Equations, *SIGMA* **2** (2006) Paper 091.
- [25] H. Konno, Elliptic Quantum Group $U_{q,p}(\widehat{\mathfrak{sl}}_2)$ and Vertex Operators, *J.Phys. A* **41** (2008) 194012 (12pp).
- [26] H. Konno, Elliptic Quantum Group $U_{q,p}(\widehat{\mathfrak{sl}}_2)$, Hopf Algebroid Structure and Elliptic Hypergeometric Series, *J. Geom. Phys.* **59** (2009) 1485-1511.
- [27] H.Konno, Elliptic Quantum Groups $U_{q,p}(\widehat{\mathfrak{gl}}_N)$ and $E_{q,p}(\widehat{\mathfrak{gl}}_N)$, *Adv.Studies in Pure Math.* **76** (2018) 347-417.

- [28] H.Konno, Elliptic Weight Functions and Elliptic q -KZ Equation, *Journal of Integrable Systems* **2** (2017) 1-43. doi: 10.1093/integr/xyx011.
- [29] H.Konno, Elliptic Stable Envelopes and Finite-Dimensional Representations of Elliptic Quantum Group, *Journal of Integrable Systems* **3** (2018) 1-43. doi: 10.1093/integr/xyy012.
- [30] H.Konno, Elliptic Quantum Groups, *the Encyclopedia of Mathematical Physics 2nd edition*, Elsevier, to appear.
- [31] H.Konno, Elliptic Quantum Groups, Representations and Related Geometry, *Springer Briefs in Mathematical Physics* **37** (2020) Springer.
- [32] H.Konno and K.Oshima, Elliptic Quantum Toroidal Algebra $U_{q,t,p}(\mathfrak{gl}_{1,tor})$ and Affine Quiver Gauge Theories, *Lett.Math.Phys.* **113** (2023) Paper No.32, 64pp.
- [33] P. Koroteev, P. Pushkar, A. Smirnov and Zeitlin, Quantum K-theory of Quiver Varieties and Many-body Systems, arXiv:1705.10419.
- [34] I.Macdonald, Symmetric Functions and Hall Polynomials, 2nd ed., Oxford Univ. Press, New York, 1995.
- [35] D.Maulik and A.Okounkov, Quantum Groups and Quantum Cohomology, *Astérisque* **408** (2019), Societe mathematique de France. arXiv:1211.1287
- [36] K.Miki, A (q, γ) Analogue of the $W_{1+\infty}$ Algebra, *J.Math.Phys.* **48** (2007) 123520, 35 pages.
- [37] H. Nakajima, Instantons on ALE Spaces, Quiver Varieties and Kac-Moody Algebras, *Duke Math. J.* **76** (1994) 365–416.
- [38] H. Nakajima, Quiver Varieties and Kac-Moody Algebras, *Duke Math. J.* **91** (1998) 515–560.
- [39] N.Nekrasov, Seiberg-Witten Prepotential From Instanton Counting, *Adv.Theor.Math.Phys.* **7** (2004), 831–864.
- [40] N.Nekrasov, BPS/CFT Correspondence: Non-perturbative Dyson-Schwinger Equations and qq -Characters, *J.High Energy Phys.* **03** (2016), 181.
- [41] A.Okounkov, Lectures on K-theoretic Computations in Enumerative Geometry. (2015), arXiv: 1512.07363.
- [42] A.Okounkov and A.Smirnov, Quantum Difference Equation for Nakajima Varieties, *Invent.Math.* **229** (2022), 1203–1299.

- [43] R. Rimányi, V. Tarasov and A. Varchenko, Elliptic and K -theoretic Stable Envelopes and Newton Polytopes, *Selecta Math.* **25** (2019), Paper No.16.
- [44] A. Smirnov, Elliptic stable envelope for Hilbert scheme of points in the plane, *Selecta Math.* **26** (2020) Paper No.3, 57pp.
- [45] A. Smirnov and H.Dinkins, Characters of Tangent Spaces at Torus Fixed Points and 3d-Mirror Symmetry, *Lett. Math. Phys.* **110** (2020) 2337–2352.