

# DETOUR MONOPHONIC VERTEX COVER PEBBLING NUMBER (DMVCPN) OF SOME STANDARD GRAPHS

K. CHRISTY RANI<sup>1</sup>, I. DHIVVIYANANDAM<sup>2</sup>

**ABSTRACT.** Let  $G$  be a connected graph with vertex set  $V(G)$  and edge set  $E(G)$ . Pebbling shift is a deletion of two pebbles from a vertex and a placement of one pebble to a neighbouring vertex. The vertex cover set,  $D_{vc}$  for graph  $G$  is the subset of  $V(G)$  such that every edge in  $G$  has minimum one end in  $D_{vc}$ . A detour monophonic path is considered to be a longest chordless path between two non adjacent vertices  $x$  and  $y$ . A detour monophonic vertex cover pebbling number,  $\mu_{vc}(G)$ , is a minimum number of pebbles require to cover all the vertices of the vertex cover set of  $G$  with at least one pebble each on them after the transformation of pebbles by using detour monophonic paths. We determine the detour monophonic vertex cover pebbling number (DMVCPN) of Cycle, Path, Fan and Wheel graphs.

**Keywords:** detour monophonic pebbling, vertex cover set, detour monophonic vertex cover pebbling.

**AMS Subject Classification:** 05C12, 05C25, 05C38, 05C76, 05C99

## 1. Introduction

Graph pebbling is an interesting area emerged a few decades ago and enhanced Graph theory. Graph pebbling has produced results in Combinatorial Group Theory and Combinatorial Number Theory [4], [5]. It is widely applied in network optimization model for transportation and for transmitting information through its medium to military troops, etc. A pebbling move is an extraction of two pebbles from one vertex and placing one pebble on the adjacent vertex and eliminating the other pebble. A vertex cover of a graph  $G$  is studied in [9], which defines the vertex cover as a set  $D_{vc} \subset V(G)$  such that for each edge  $uv \in E(G)$ , at least one of  $u$

---

<sup>1</sup> Department of Mathematics, Bon Secours College for Women (Autonomous), Thanjavur-613006, Tamil Nadu, India.

e-mail: christy.agnes@gmail.com; <https://orcid.org/0000-0002-2837-9388>.

<sup>2</sup> Department of Mathematics, North Bengal St. Xavier's College, Rajganj-735134, West Bengal, India.

e-mail: divyanasj@gmail.com; <https://orcid.org/0000-0002-3805-6638>.

or  $v$  is in  $D_{vc}$ . A detour monophonic path from  $x$  to  $y$  is the longest path with no chords studied in [10]. In this article we define the detour monophonic vertex cover pebbling number,  $\mu_{vc}(G)$ , which is a minimum number of pebbles require to cover all the vertices of the vertex cover set  $D_{vc}$  of  $G$  and the corresponding edges of  $G$  with at least one pebble each on them after the transformation of pebbles by using detour monophonic paths. We also investigate the detour monophonic vertex cover pebbling number of standard graphs. To prove the worst condition, we use the stacking theorem. It states that when the initial configuration  $D$  is placed on a single vertex  $v$  such that the distance of  $v = \dot{s}$  is maximum and hence  $\dot{s} = \sum_{u \in V_G} 2^{dist(u,v)}$ . Applying this for every vertex  $v \in V(G)$ , we determine  $\mu_{vc}(G)$ . We refer to [1] and [2] for basic terminologies of graph theory. Throughout this paper, we refer  $G$  to be a simple connected graph.

**Definition 1.1.** [10] *In a graph  $G$ , the smallest number  $\mu(G, x)$  for  $x \in G$  such that minimum single pebble can be shifted to  $x$  using a monophonic path by the successive pebbling shifts to place  $\mu(G, x)$  pebbles on the vertices of  $G$ . The maximum  $\mu(G, x)$  over  $v \in V(G)$  is the monophonic pebbling number denoted by  $\mu(G)$ .*

**Definition 1.2.** [8] *Any vertex  $v \in V(G)$  is called a source vertex when  $D(u, v)$  is maximum.*

**Notation 1.1.** *In this article we use*

- a.  $D_{vc}$  for the vertex cover set,
- b.  $\dot{s}$  to denote the source vertex,
- c.  $D_m$  for the detour monophonic distance and
- d.  $\mu_{vc}(G)$  to denote the detour monophonic vertex cover pebbling number - DMVCPN.

## 2. Motivation

Motivated by the concepts of the cover pebbling number [3] and detour monophonic number [10] and monophonic pebbling number [6] with vast applicatoins, we have discussed the detour monophonic vertex cover pebbling number for some standard graphs such as  $C_n (n \geq 3)$ ,  $P_n$ ,  $F_n$  and  $W_n$ . Detour monophonic vertex cover pebbling is applied in deciding the equal distribution of goods on every customer by using the detour monophonic path. This is also applied in the network transmission of the information from one node to the other.

**Theorem 2.1.** *Let  $P_n$  be a path of order  $n$  ( $n$  is odd and  $n \geq 3$ ).*

*Then  $\mu_{vc}(P_n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}$ .*

*Proof.* Let  $P_n$  be a path of order  $n$ , where  $n$  is odd. Let  $D_{vc}$  be the vertex cover set. Then,  $D_{vc} = \{v_2, v_4, v_6, \dots, v_{n-1}\}$ . Let the initial configuration be  $\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1} - 1$  pebbles. Without loss of generality let  $\dot{s} = v_1$ . The detour monophonic distance from  $v_1$  to any other vertex is at most  $n - 2$ . In order to cover the farthest vertex  $v_{n-1}$  using the detour monophonic path we require  $2^{n-2}$  pebbles. Similarly, to cover the remaining vertices of  $D_{vc}$ , we require the following pebble distributions.

$2^1 + 2^3 + 2^5 + 2^7 + \dots + 2^{n-2}$ . That sums upto  $\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}$ . But our initial configuration of pebbles is  $\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1} - 1$  which is a contradiction. Therefore,

$$\mu_{vc}(P_n) \geq \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}.$$

Now we prove the sufficient condition by using  $\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}$  pebbles on the vertices of  $P_n$ .

**Case 1:** Let  $s = v_1$  or  $v_n$ .

Without loss of generality, let  $v_n$  be the source vertex. Then  $D^*_{vc} = \{v_{n-1}, v_{n-3}, v_{n-5}, \dots, v_2\}$ .

	$v_2$	$v_4$	$v_6$	$\dots$	$v_{n-5}$	$v_{n-3}$	$v_{n-1}$
$v_1$	1	3	5	$\dots$	$n-6$	$n-4$	$n-2$
$v_n$	$n-2$	$n-4$	$n-6$	$\dots$	5	3	1

TABLE 1. Detour monophonic distance from end vertices to  $V(D^*_{vc})$ .

In order to cover the vertices of  $P_n$ , it suffices to cover the vertices of vertex cover set  $V(D^*_{vc})$ . From Table 1, we require  $2^{n-2}$ ,  $2^{n-4}$  and  $2^{n-6}$  pebbles from  $v_n$  to cover  $v_2$ ,  $v_4$  and  $v_6$  respectively. Proceeding in the same way we use  $2^5$ ,  $2^3$  and 2 pebbles to cover the vertices  $v_{n-5}$ ,  $v_{n-3}$  and  $v_{n-1}$ . Thus the following series of pebbles guarantees covering all the vertices of  $D^*_{vc}$ .

$$\begin{aligned} & \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{n-5}) + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1}) \\ &= 2^{n-2} + 2^{n-4} + 2^{n-6} + \dots + 2^5 + 2^3 + 2^1. \end{aligned}$$

Thus the number of pebbles used to cover  $V(D^*_{vc})$  is,

$$\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}.$$

**Case 2:** When the middle vertex is the source vertex.

Let  $s = v_i$ ,  $i = \lfloor \frac{n}{2} \rfloor + 1$ .

**Subcase 2.1:** If  $n \equiv 1 \pmod{4}$  and then  $D^1_{vc} = \{v_2, v_4, v_6, \dots, v_{\lfloor \frac{n}{2} \rfloor - 2}, v_{\lfloor \frac{n}{2} \rfloor}, v_{\lfloor \frac{n}{2} \rfloor + 2}, v_{\lfloor \frac{n}{2} \rfloor + 4}, \dots, v_{n-1}\}$ .

	$v_2$	$v_4$	$v_6$	$\dots$	$v_{\lfloor \frac{n}{2} \rfloor - 2}$	$v_{\lfloor \frac{n}{2} \rfloor}$	$v_{\lfloor \frac{n}{2} \rfloor + 2}$	$v_{\lfloor \frac{n}{2} \rfloor + 4}$	$\dots$	$v_{n-3}$	$v_{n-1}$
$v_{\lfloor \frac{n}{2} \rfloor + 1}$	$\lfloor \frac{n}{2} \rfloor - 1$	$\lfloor \frac{n}{2} \rfloor - 3$	$\lfloor \frac{n}{2} \rfloor - 5$	$\dots$	3	1	1	3	$\dots$	$\lfloor \frac{n}{2} \rfloor - 3$	$\lfloor \frac{n}{2} \rfloor - 1$

TABLE 2. Detour monophonic distance from the middle vertex of  $P_n$  to  $V(D^1_{vc})$ , where  $n \equiv 1 \pmod{4}$ .

From Table 2, we use  $2^{\lfloor \frac{n}{2} \rfloor - 1}$ ,  $2^{\lfloor \frac{n}{2} \rfloor - 3}$  and  $2^{\lfloor \frac{n}{2} \rfloor - 5}$  pebbles from  $v_{\lfloor \frac{n}{2} \rfloor + 1}$  to cover  $v_2$ ,  $v_4$  and  $v_6$ . We continue in this manner and cover the vertices of  $D^1_{vc}$  with the following pebble distribution.

$$\begin{aligned} & \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor - 2}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor + 2}) \\ &+ \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor + 4}) + \dots + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1}). \\ &= 2^{\lfloor \frac{n}{2} \rfloor - 1} + 2^{\lfloor \frac{n}{2} \rfloor - 3} + 2^{\lfloor \frac{n}{2} \rfloor - 5} + \dots + 2^3 + 2^1 + 2^1 + 2^3 + \dots + 2^{\lfloor \frac{n}{2} \rfloor - 3} + 2^{\lfloor \frac{n}{2} \rfloor - 1} \end{aligned}$$

$$= 2(2^1 + 2^3 + 2^5 + 2^7 + \dots + 2^{\lfloor \frac{n}{2} \rfloor - 1}).$$

Thus in order to cover the vertices of  $D_{vc}^1$  we used

$$2\left(\sum_{j=1}^{\lfloor \frac{n}{4} \rfloor} 2^{2(j)-1}\right) \text{ pebbles which is less than the total number of pebbles available.}$$

**Subcase 2.2:** Let  $n \equiv 3(\text{mod } 4)$ . Without loss of generality  $D_{vc}^2 = \{v_2, v_4, v_6, \dots, v_{\lfloor \frac{n}{2} \rfloor - 1}, v_{\lfloor \frac{n}{2} \rfloor + 1}, v_{\lfloor \frac{n}{2} \rfloor + 3}, \dots, v_{n-1}\}$ .

	$v_2$	$v_4$	$v_6$	$\dots$	$v_{\lfloor \frac{n}{2} \rfloor - 1}$	$v_{\lfloor \frac{n}{2} \rfloor + 1}$	$v_{\lfloor \frac{n}{2} \rfloor + 3}$	$\dots$	$v_{n-3}$	$v_{n-1}$
$v_{\lfloor \frac{n}{2} \rfloor + 1}$	$\lfloor \frac{n}{2} \rfloor - 1$	$\lfloor \frac{n}{2} \rfloor - 3$	$\lfloor \frac{n}{2} \rfloor - 5$	$\dots$	2	0	2	$\dots$	$\lfloor \frac{n}{2} \rfloor - 3$	$\lfloor \frac{n}{2} \rfloor - 1$

TABLE 3. Detour monophonic distance from the middle vertex of  $P_n$  to  $V(D_{vc}^2)$ , where  $n \equiv 3(\text{mod } 4)$ .

From Table 3, we use  $2^{\lfloor \frac{n}{2} \rfloor - 1}$ ,  $2^{\lfloor \frac{n}{2} \rfloor - 3}$  and  $2^{\lfloor \frac{n}{2} \rfloor - 5}$  pebbles to cover  $v_2$ ,  $v_4$  and  $v_6$ . Similarly we cover the vertices of  $D_{vc}^2$  with the following pebble distributions.

$$\begin{aligned} & \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor - 1}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor + 1}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor + 3}) + \\ & \dots + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1}). \\ & = 2^{\lfloor \frac{n}{2} \rfloor - 1} + 2^{\lfloor \frac{n}{2} \rfloor - 3} + 2^{\lfloor \frac{n}{2} \rfloor - 5} + \dots + 2^2 + 2^0 + 2^2 + \dots + 2^{\lfloor \frac{n}{2} \rfloor - 3} + 2^{\lfloor \frac{n}{2} \rfloor - 1} \\ & = 2^0 + 2(2^2 + 2^4 + 2^6 + \dots + 2^{\lfloor \frac{n}{2} \rfloor - 1}). \end{aligned}$$

Thus, we used  $2\left(\sum_{k=1}^{\lfloor \frac{n}{4} \rfloor} 2^k\right) + 1$  pebbles, which is less than  $\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}$ .

**Case 3:** Consider  $s$  is an internal vertex. Let  $s = v_i$ , where  $2 \leq i \leq n - 1$ ,  $i \neq \lfloor \frac{n}{2} \rfloor + 1$ .

**Subcase 3.1:** When  $i$  is even.

Let  $V(D_{vc}^3) = \{v_2, v_4, v_6, \dots, v_{i-2}, v_i, v_{i+2}, \dots, v_{n-3}, v_{n-1}\}$ .

	$v_2$	$v_4$	$v_6$	$\dots$	$v_{i-2}$	$v_i$	$v_{i+2}$	$\dots$	$v_{n-3}$	$v_{n-1}$
$v_i$	$ i - 2 $	$ i - 4 $	$ i - 6 $	$\dots$	2	0	2	$\dots$	$n - (i + 3)$	$n - (i + 1)$

TABLE 4. Detour monophonic distance from the internal vertex of  $P_n$  to  $V(D_{vc}^3)$ .

When  $i = 2$ , we use  $2^0, 2^2, 2^6, \dots, 2^{n-5}$  and  $2^{n-3}$  pebbles to cover  $v_2, v_4, v_6, \dots, v_{n-3}$  and  $v_{n-5}$  respectively. when  $i = 4$ , we use  $2^2, 2^0, 2^2, \dots, 2^{n-7}$  and  $2^{n-5}$  pebbles to cover  $v_2, v_4, v_6, \dots, v_{n-3}$  and  $v_{n-5}$  respectively. Continuing this process, when  $i$  is even, we cover the vertices of  $D_{vc}^3$  with the following pebble distribution.

$$\mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{i-2}) + \mu_{vc}(G, v_i) + \mu_{vc}(G, v_{i+2}) + \dots + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1}).$$

From Table 4, we can sum up as follows.

$$\sum_{l=1}^{\frac{i}{2}-1} 2^{2(l)} + \sum_{m=0}^{\frac{n-(i+1)}{2}} 2^{2(m)}.$$

	$v_2$	$v_4$	$v_6$	$\cdots$	$v_{i-3}$	$v_{i-1}$	$v_{i+1}$	$v_{i+3}$	$\cdots$	$v_{n-3}$	$v_{n-1}$
$v_i$	$ i-2 $	$i-4$	$i-6$	$\cdots$	3	1	1	3	$\cdots$	$n-(i+3)$	$n-(i+1)$

TABLE 5. Detour monophonic distance from the internal vertices of  $P_n$  to  $V(D_{vc}^4)$ .

**Subcase 3.2:** When  $i$  is odd.

Consider  $V(D_{vc}^4) = \{v_2, v_4, v_6, \cdots, v_{i-3}, v_{i-1}, v_{i+1}, v_{i+3}, \cdots, v_{n-3}, v_{n-1}\}$ .

From Table 5, when  $i$  is odd, we cover the vertices of  $D_{vc}^4$  with the following pebble distribution.

$$\mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \cdots + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{i-1}) + \mu_{vc}(G, v_{i+1}) + \mu_{vc}(G, v_{i+3}) + \cdots + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1}).$$

$= 2^1 + 2^3 + \cdots + 2^{i-4} + 2^{i-2} + 2^1 + 2^3 + \cdots + 2^{n-(i+3)} + 2^{n-(i+1)}$ . Thus, we have utilized

$\sum_{l=1}^{\lfloor \frac{i}{2} \rfloor} 2^{2(l)-1} + \sum_{m=0}^{\lfloor \frac{n-(i+1)}{2} \rfloor} 2^{2(m)+1}$  pebbles, which is less than the total number of pebbles available. Hence,

$$\mu_{vc}(P_n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}.$$

□

**Theorem 2.2.** Let  $P_n$  be a path of order  $n$  ( $n$  is even and  $n \geq 2$ ).

Then  $\mu_{vc}(P_n) = \sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)}$ .

*Proof.* Let  $P_n$  be a path of order  $n$ , where  $n$  is even. Let  $D_{vc}$  be the vertex cover set. Then,  $D_{vc} = \{v_1, v_3, v_5, \cdots, v_{n-1}\}$ . Let the initial configuration be  $\sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)} - 1$  and  $\dot{s} = v_1$ . The detour monophonic distance from  $v_1$  to any other vertex of  $P_n$  is at most  $n-2$ . Then using  $2^{n-2}$  pebbles we could cover  $v_{n-1}$ . Similarly to cover  $v_{n-3}$ , we require  $2^{n-4}$  pebbles. In a similar way we can cover the remaining vertices of  $D_{vc}$  using the following pebble distributions.

$2^2 + 2^4 + 2^6 + \cdots + 2^{n-4} + 2^{n-2}$  that sums upto  $\sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)} - 1$ . Now, there exists a vertex  $v_1$  uncovered. Thus we fail to cover all the vertices of  $D_{vc}$ . Hence,

$$\mu_{vc}(P_n) \geq \sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)}.$$

Now we prove the sufficient condition with  $\sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)}$  pebbles on the vertices of  $P_n$ .

**Case 1:**

Assume the end vertex to be the source vertex.

Let  $\dot{s} = v_i$ , where  $i \in \{1, n\}$ .

Without loss of generality let  $\dot{s} = v_1$  and  $D_{vc}^1 = \{v_1, v_3, v_5, \cdots, v_{n-3}, v_{n-1}\}$ .

	$v_1$	$v_3$	$v_5$	$v_7$	$v_9$	$\cdots$	$v_{n-7}$	$v_{n-5}$	$v_{n-3}$	$v_{n-1}$
$v_1$	0	2	4	6	8	$\cdots$	$n-8$	$n-6$	$n-4$	$n-2$

TABLE 6. Detour monophonic distance from end vertex to  $V(D_{vc}^1)$ .

We make use of  $2^0, 2^2, 2^4$  pebbles to cover the vertices  $v_1, v_3$  and  $v_5$  respectively. Proceeding in the same way we use  $2^{n-4}$  and  $2^{n-2}$  pebbles to cover the vertices  $v_{n-3}$  and  $v_{n-1}$ . Thus, from Table 6 the following series of pebble distribution guarantees covering of all the vertices of  $D_{vc}^1$ .

$$\begin{aligned} & \mu_{vc}(G, v_1) + \mu_{vc}(G, v_3) + \mu_{vc}(G, v_5) + \mu_{vc}(G, v_7) + \cdots + \mu_{vc}(G, v_{n-7}) + \mu_{vc}(G, v_{n-5}) + \\ & \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1}) \\ &= 2^0 + 2^2 + 2^4 + \cdots + 2^{n-2}. \\ &= \sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)}. \end{aligned}$$

By symmetry, we can cover the vertices of  $D_{vc}^1$ , when the source vertex is  $v_n$ .

### Case 2:

Let the middle vertex be the source vertex. Then there exist two subcases.

#### Subcase 2.1:

When  $n$  is even and  $n \equiv 0 \pmod{4}$ .

Assume  $\dot{s} = v_i$ .

If  $i = \frac{n}{2}$ , then  $D_{vc}^2 = \{v_1, v_3, v_5, \dots, v_{\frac{n}{2}-1}, v_{\frac{n}{2}+1}, \dots, v_{n-3}, v_{n-1}\}$ .

If  $i = \frac{n}{2} + 1$ , then  $D_{vc}^{2*} = \{v_2, v_4, v_6, \dots, v_{\frac{n}{2}-1}, v_{\frac{n}{2}+1}, \dots, v_n\}$ .

Without loss of generality, consider the vertex cover set  $D_{vc}^2$ .

	$v_1$	$v_3$	$v_5$	$\cdots$	$v_{\frac{n}{2}-3}$	$v_{\frac{n}{2}-1}$	$v_{\frac{n}{2}+1}$	$v_{\frac{n}{2}+3}$	$\cdots$	$v_{n-3}$	$v_{n-1}$
$v_i$	$\frac{n}{2} - 1$	$\frac{n}{2} - 3$	$\frac{n}{2} - 5$	$\cdots$	3	1	1	3	$\cdots$	$\frac{n}{2} - 3$	$\frac{n}{2} - 1$

TABLE 7. Detour monophonic distance from middle vertex to  $V(D_{vc}^2)$ .

Now, from Table 7 the following pebble distribution covers the vertices of  $D_{vc}^2$ .

$$\begin{aligned} & 2^{\frac{n}{2}-1} + 2^{\frac{n}{2}-3} + 2^{\frac{n}{2}-5} + \cdots + 2^3 + 2^1 + 2^1 + 2^3 + \cdots + 2^{\frac{n}{2}-3} + 2^{\frac{n}{2}-1} \\ &= 2(2^1 + 2^3 + 2^5 + \cdots + 2^{\frac{n}{2}-1}). \end{aligned}$$

Similarly, we can cover the vertices of  $D_{vc}^{2*}$ . Thus in this case to cover the vertices of the vertex cover set, we use

$$2\left(\sum_{j=0}^{\frac{n}{4}} 2^{2(j)-1}\right) \text{ pebbles, where } n \equiv 0 \pmod{4}.$$

#### Subcase 2.2:

When  $n$  is even and  $n \equiv 2 \pmod{4}$ .

Let  $i = \frac{n}{2}$ . Then  $\dot{s} = v_{\frac{n}{2}}$ .

As we consider the odd middle vertex, the vertex cover set also includes all the odd vertices of  $P_n$ . Thus,  $D_{vc}^3 = \{v_1, v_3, v_5, \dots, v_{\frac{n}{2}}, v_{\frac{n}{2}+2}, \dots, v_{n-1}\}$ . Let  $i = \frac{n}{2} + 1$  and  $\dot{s} = v_{\frac{n}{2}+1}$ . When we consider the even middle vertex, the vertex cover set consists of all the even vertices of  $P_n$ . Hence  $D_{vc}^{3*} = \{v_2, v_4, v_6, \dots, v_{\frac{n}{2}-1}, v_{\frac{n}{2}+1}, v_{\frac{n}{2}+3}, \dots, v_n\}$ .

Without loss of generality, assume the vertex cover set  $D_{vc}^{3*}$ .

	$v_2$	$v_4$	$v_6$	$\cdots$	$v_{\frac{n}{2}-3}$	$v_{\frac{n}{2}-1}$	$v_{\frac{n}{2}+1}$	$v_{\frac{n}{2}+3}$	$v_{\frac{n}{2}+5}$	$\cdots$	$v_{n-2}$	$v_n$
$v_i$	$\frac{n}{2} - 1$	$\frac{n}{2} - 3$	$\frac{n}{2} - 5$	$\cdots$	4	2	0	2	4	$\cdots$	$\frac{n}{2} - 3$	$\frac{n}{2} - 1$

TABLE 8. Detour monophonic distance from middle vertex to  $V(D_{vc}^{3*})$ .

Thus, from Table 8 we cover the vertices of  $D_{vc}^{3*}$  with the following pebble distribution.

$$2^{\frac{n}{2}-1} + 2^{\frac{n}{2}-3} + 2^{\frac{n}{2}-5} + \dots + 2^4 + 2^2 + 2^0 + 2^2 + 2^4 + \dots + 2^{\frac{n}{2}-3} + 2^{\frac{n}{2}-1}.$$

$$= 2^{2\lfloor \frac{n}{4} \rfloor} + \dots + 2^4 + 2^2 + 2^0 + 2^2 + 2^4 + \dots + 2^{2\lfloor \frac{n}{4} \rfloor}.$$

In this manner we can cover the vertices of  $D_{vc}^3$ . Hence, the pebble distribution is summed up as follows.

$$1 + 2\left(\sum_{k=1}^{\lfloor \frac{n}{4} \rfloor} 2^{2(k)}\right), \text{ where } n \equiv 2 \pmod{4} \text{ and } \dot{s} \text{ is a middle vertex.}$$

**Case 3:** Consider  $\dot{s}$  to be an internal vertex. Let  $\dot{s} = v_i$ , where  $2 \leq i \leq n-1$ ,  $i \neq \frac{n}{2}, \frac{n}{2} + 1$ .

**Subcase 3.1:**

If  $v_i$  is even and  $i < \frac{n}{2}$ . Then  $D_{vc}^4 = \{v_1, v_3, v_5, \dots, v_{i-1}, v_{i+1}, \dots, v_{n-1}\}$ .

	$v_1$	$v_3$	$v_5$	$\dots$	$v_{i-3}$	$v_{i-1}$	$v_{i+1}$	$v_{i+3}$	$\dots$	$v_{n-1}$
$v_i$	$i-1$	$i-3$	$i-5$	$\dots$	3	1	1	3	$\dots$	$n-(i+1)$

TABLE 9. Detour monophonic distance from  $v_i$  to  $V(D_{vc}^4)$ .

From Table 9 to cover the vertices of  $D_{vc}^4$ , we require the following pebble series. The vertices  $v_1$  and  $v_{n-1}$  are covered with  $2^{i-1}$  and  $2^{n-(i+1)}$  pebbles. Thus the required pebble distribution results as follows.

$$\mu_{vc}(G, v_1) + \mu_{vc}(G, v_3) + \dots + \mu_{vc}(G, v_{i-3}) + \mu_{vc}(G, v_{i-1}) + \mu_{vc}(G, v_{i+1}) + \mu_{vc}(G, v_{i+3}) + \dots + \mu_{vc}(G, v_{n-1}).$$

$$= 2^{i-1} + 2^{i-3} + \dots + 2^3 + 2^1 + 2^1 + 2^3 + \dots + 2^{n-(i+1)}$$

$$= \sum_{j=1}^{\lfloor \frac{i}{2} \rfloor} 2^{2(j)-1} + \sum_{k=1}^{\lfloor \frac{n-(i+1)}{2} \rfloor} 2^{2(k)-1}.$$

When  $i > \frac{n}{2} + 1$ , we obtain the reversal result using the similar pebble distribution. Hence the number of pebbles employed in the process of covering  $D_{vc}^4$  is,

$$\sum_{j=1}^{\lfloor \frac{n-(i+1)}{2} \rfloor} 2^{2(j)-1} + \sum_{k=1}^{\lfloor \frac{i}{2} \rfloor} 2^{2(k)-1}.$$

**Subcase 3.2:**

If  $v_i$  is odd and  $i < \frac{n}{2}$ . Then  $D_{vc}^5 = \{v_1, v_3, v_5, \dots, v_i, v_{i+2}, \dots, v_{n-1}\}$ .

	$v_1$	$v_3$	$v_5$	$\dots$	$v_{i-4}$	$v_{i-2}$	$v_i$	$v_{i+2}$	$v_{i+4}$	$\dots$	$v_{n-1}$
$v_i$	$i-1$	$i-3$	$i-5$	$\dots$	4	2	0	2	4	$\dots$	$n-(i+1)$

TABLE 10. Detour monophonic distance from  $v_i$  to  $V(D_{vc}^5)$ .

The vertex  $v_1$  is covered with  $2^{i-1}$  pebbles and  $v_{n-1}$  is covered with  $2^{n-(i+1)}$  pebbles. Thus, from Table 10 to cover the vertices of  $D_{vc}^5$ , we require the following pebble distributions.

$$\mu_{vc}(G, v_1) + \dots + \mu_{vc}(G, v_{i-4}) + \mu_{vc}(G, v_{i-2}) + \mu_{vc}(G, v_i) + \mu_{vc}(G, v_{i+2}) + \mu_{vc}(G, v_{i+4}) + \dots + \mu_{vc}(G, v_{n-1}).$$

$$= 2^{i-1} + \dots + 2^4 + 2^2 + 2^0 + 2^2 + 2^4 + \dots + 2^{n-(i+1)}$$

$$= \sum_{l=0}^{\lfloor \frac{i}{2} \rfloor} 2^{2(l)} + \sum_{m=1}^{\lfloor \frac{n-(i+1)}{2} \rfloor} 2^{2(m)}.$$

If  $i > \frac{n}{2} + 1$ , we obtain the reversal result using the similar pebble distribution. Hence,

$$\sum_{l=1}^{\lfloor \frac{n-(i+1)}{2} \rfloor} 2^{2(l)} + \sum_{m=0}^{\lfloor \frac{i}{2} \rfloor} 2^{2(m)}.$$

Thus in each case, we have used the pebbles which sums up less than the considered pebbles. Therefore,  $\mu_{vc}(P_n) = \sum_{i=0}^{\frac{n}{2}-1} 2^{2(i)}$ .  $\square$

**Theorem 2.3.** For Fan  $F_n$  ( $n$  is even),  $\mu_{vc}(F_n) = \sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2$ .

*Proof.* Let  $F_n = P_{n-1} + v_0$  be a fan of order  $n$  ( $n$  is even) with  $V(F_n) = \{v_0, v_1, v_2, \dots, v_{n-1}\}$ . Let  $D_{vc}$  be the vertex cover set. Then,  $D_{vc} = \{v_0, v_2, v_4, v_6, \dots, v_{n-2}\}$ . Consider the following configuration of pebbles on the vertices of fan  $F_n$ .

Let the initial configuration be  $\sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 1$  pebbles. Assume  $v_1$  to be the source vertex. The detour monophonic distance from  $v_1$  to any other vertex is at most  $n-2$ . Then using  $2^{n-3}$  pebbles we could place a pebble on  $v_{n-2}$ . Then we place  $2^{n-5}$  pebbles on  $v_{n-4}$ . In a similar way, we distribute the pebbles on the vertices of  $D_{vc}$ . Now, there remains only one pebble on  $v_2$  leaving  $v_1$  pebbleless. Thus, we fail to cover all the vertices of  $D_{vc}$ . Therefore,

$$\mu_{vc}(F_n) \geq \sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2.$$

Now we prove the sufficient condition by using  $\sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2$  pebbles on the vertices of  $F_n$ .

**Case 1:** Let  $\dot{s} = v_1$  or  $v_{n-1}$ .

Without loss of generality, let  $v_n$  be the source vertex. Then  $D_{vc}^1 = \{v_0, v_2, v_4, v_6, \dots, v_{n-6}, v_{n-4}, v_{n-2}\}$ .

	$v_0$	$v_2$	$v_4$	$v_6$	$\dots$	$v_{n-6}$	$v_{n-4}$	$v_{n-2}$
$v_1$	1	1	3	5	$\dots$	$n-7$	$n-5$	$n-3$
$v_{n-1}$	1	$n-3$	$n-5$	$n-7$	$\dots$	5	3	1

TABLE 11. Detour monophonic distance from  $v_1$  and  $v_{n-1}$  to  $V(D_{vc}^1)$ .

From Table 11, we use 2 pebbles to cover  $v_0$ . Also, we use  $2^{n-3}$ ,  $2^{n-5}$ ,  $2^{n-7}$  pebbles to cover  $v_0$ ,  $v_2$ ,  $v_4$  and  $v_6$  respectively. Proceeding in the same way we need  $2^5$ ,  $2^3$  and 2 pebbles to cover the vertices  $v_{n-6}$ ,  $v_{n-4}$  and  $v_{n-2}$ . Thus, the following series of pebbles guarantees covering all the vertices of  $D_{vc}^1$ .

$$\begin{aligned} & \mu_{vc}(G, v_0) + \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{n-6}) + \mu_{vc}(G, v_{n-4}) + \\ & \mu_{vc}(G, v_{n-2}) \\ & = 2^1 + 2^{n-3} + 2^{n-5} + 2^{n-7} + \dots + 2^5 + 2^3 + 2^1. \end{aligned}$$

Thus,  $\sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2$  pebbles are used to cover  $D_{vc}^1$ .

**Case 2:**

Consider  $\dot{s} = v_i$ ,  $i = \frac{n}{2}$ .

**Subcase 2.1:** If  $n \equiv 2 \pmod{4}$  and then  $D_{vc}^2 = \{v_0, v_2, v_4, v_6, \dots, v_{\frac{n}{2}-3}, v_{\frac{n}{2}-1}, v_{\frac{n}{2}+1},$

$v_{\frac{n}{2}+3}, \dots, v_{n-4}, v_{n-2}\}$ .

	$v_0$	$v_2$	$v_4$	$v_6$	$\dots$	$v_{\frac{n}{2}-3}$	$v_{\frac{n}{2}-1}$	$v_{\frac{n}{2}+1}$	$v_{\frac{n}{2}+3}$	$\dots$	$v_{n-4}$	$v_{n-2}$
$v_{\frac{n}{2}}$	1	$\frac{n}{2}-2$	$\frac{n}{2}-4$	$\frac{n}{2}-6$	$\dots$	3	1	1	3	$\dots$	$\frac{n}{2}-4$	$\frac{n}{2}-2$

TABLE 12. Detour monophonic distance from  $v_{\frac{n}{2}}$  of  $F_n$  to  $V(D_{vc}^2)$ , where  $n \equiv 2(\text{mod } 4)$ .

From Table 12, we use  $2^{\frac{n}{2}-2}$ ,  $2^{\frac{n}{2}-4}$  pebbles to cover  $v_2$ ,  $v_{n-2}$  and  $v_4$ ,  $v_{n-4}$  respectively. Obviously,  $v_0$  is covered with 2 pebbles. We continue in this manner and cover the vertices of  $D_{vc}^2$  with the following pebble distribution.

$$\mu_{vc}(G, v_0) + \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{\frac{n}{2}-3}) + \mu_{vc}(G, v_{\frac{n}{2}-1}) + \mu_{vc}(G, v_{\frac{n}{2}+1}) + \mu_{vc}(G, v_{\frac{n}{2}+3}) + \dots + \mu_{vc}(G, v_{n-4}) + \mu_{vc}(G, v_{n-2}).$$

$$= 2^1 + 2^{\frac{n}{2}-2} + 2^{\frac{n}{2}-4} + 2^{\frac{n}{2}-6} + \dots + 2^3 + 2^1 + 2^1 + 2^3 + \dots + 2^{\frac{n}{2}-4} + 2^{\frac{n}{2}-2}.$$

Thus in order to cover the vertices of  $D_{vc}^2$  we used

$$2\left(\sum_{j=1}^{\lfloor \frac{n}{4} \rfloor} 2^{2(j-1)}\right) + 2 \text{ pebbles, which is less than the total number of pebbles available.}$$

**Subcase 2.2:** Let  $n \equiv 0(\text{mod } 4)$ . Without loss of generality  $D_{vc}^3 = \{v_0, v_2, v_4, v_6, \dots, v_{\frac{n}{2}-2}, v_{\frac{n}{2}}, v_{\frac{n}{2}+2}, \dots, v_{n-4}, v_{n-2}\}$ .

	$v_0$	$v_2$	$v_4$	$v_6$	$\dots$	$v_{\frac{n}{2}-2}$	$v_{\frac{n}{2}}$	$v_{\frac{n}{2}+2}$	$\dots$	$v_{n-4}$	$v_{n-2}$
$v_{\frac{n}{2}}$	1	$\frac{n}{2}-2$	$\frac{n}{2}-4$	$\frac{n}{2}-6$	$\dots$	2	0	2	$\dots$	$\frac{n}{2}-4$	$\frac{n}{2}-2$

TABLE 13. Detour monophonic distance from the middle vertex of  $F_n$  to  $V(D_{vc}^3)$ , where  $n \equiv 0(\text{mod } 4)$ .

From Table 13,  $v_0$  is covered with 2 pebbles. We use  $2^{\frac{n}{2}-2}$ ,  $2^{\frac{n}{2}-4}$  and  $2^{\frac{n}{2}-6}$  pebbles to cover  $v_2$ ,  $v_4$  and  $v_6$ . Similarly we cover the vertices of  $D_{vc}^3$  with the following pebble distributions.

$$\mu_{vc}(G, v_0) + \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \dots + \mu_{vc}(G, v_{\frac{n}{2}-2}) + \mu_{vc}(G, v_{\frac{n}{2}}) + \mu_{vc}(G, v_{\frac{n}{2}+2}) + \dots + \mu_{vc}(G, v_{n-4}) + \mu_{vc}(G, v_{n-2}).$$

$$= 2^0 + 2(2^2 + 2^4 + 2^6 + \dots + 2^{\frac{n}{4}-1}) + 2^1.$$

Thus we used  $2\left(\sum_{k=1}^{\frac{n}{4}-1} 2^k\right) + 3$ , which is less than  $\sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2$ .

**Case 3:** Consider  $\dot{s}$  is an internal vertex. Let  $\dot{s} = v_i$ , where  $2 \leq i \leq n-2$ ,  $i \neq \lfloor \frac{n-1}{2} \rfloor + 1$ .

**Subcase 3.1:** When  $i$  is even.

Let  $V(D_{vc}^4) = \{v_0, v_2, v_4, v_6, \dots, v_{i-2}, v_i, v_{i+2}, \dots, v_{n-4}, v_{n-2}\}$ .

From Table 14, when  $i = 2$ , we use  $2, 2^0, 2^2, 2^6, \dots, 2^{n-6}$  and  $2^{n-4}$  pebbles to

	$v_0$	$v_2$	$v_4$	$v_6$	$\cdots$	$v_{i-2}$	$v_i$	$v_{i+2}$	$\cdots$	$v_{n-4}$	$v_{n-2}$
$v_i$	1	$ i-2 $	$ i-4 $	$ i-6 $	$\cdots$	2	0	2	$\cdots$	$n-(i+4)$	$n-(i+2)$

TABLE 14. Detour monophonic distance from  $v_i$  of  $F_n$  to  $V(D_{vc}^4)$ .

cover  $v_0, v_2, v_4, v_6, \dots, v_{n-4}$  and  $v_{n-6}$  respectively. when  $i = 4$ , we use  $2^0, 2^2, 2^0, 2^2, \dots, 2^{n-8}$  and  $2^{n-5}$  pebbles to cover  $v_2, v_4, v_6, \dots, v_{n-4}$  and  $v_{n-6}$  respectively. Continuing this process, when  $i$  is even, we cover the vertices of  $D_{vc}^4$  with the following pebble distribution.

$$\begin{aligned} & \mu_{vc}(G, v_0) + \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \cdots + \mu_{vc}(G, v_{i-2}) + \mu_{vc}(G, v_i) + \\ & \mu_{vc}(G, v_{i+2}) + \cdots + \mu_{vc}(G, v_{n-4}) + \mu_{vc}(G, v_{n-2}). \\ & = \sum_{l=1}^{\frac{i}{2}-1} 2^{2(l)} + \sum_{m=0}^{\frac{n-(i+2)}{2}} 2^{2(m)} + 2. \end{aligned}$$

**Subcase 3.2:** When  $i$  is odd.

Consider  $V(D_{vc}^5) = \{v_0, v_2, v_4, v_6, \dots, v_{i-3}, v_{i-1}, v_{i+1}, v_{i+3}, \dots, v_{n-4}, v_{n-2}\}$ .

	$v_0$	$v_2$	$v_4$	$v_6$	$\cdots$	$v_{i-3}$	$v_{i-1}$	$v_{i+1}$	$v_{i+3}$	$\cdots$	$v_{n-4}$	$v_{n-2}$
$v_i$	1	$ i-2 $	$ i-4 $	$ i-6 $	$\cdots$	3	1	1	3	$\cdots$	$n-(i+4)$	$n-(i+2)$

TABLE 15. Detour monophonic distance from  $v_i$  of  $F_n$  to  $V(D_{vc}^5)$ .

When  $i$  is odd, we cover the vertices of  $D_{vc}^5$  with the following pebble distribution.  
 $\mu_{vc}(G, v_0) + \mu_{vc}(G, v_{i-1}) + \mu_{vc}(G, v_{i-3}) + \mu_{vc}(G, v_4) + \mu_{vc}(G, v_6) + \cdots + \mu_{vc}(G, v_{n-6}) +$   
 $\mu_{vc}(G, v_{i-1}) + \mu_{vc}(G, v_{i+1}) + \mu_{vc}(G, v_{i+3}) + \cdots + \mu_{vc}(G, v_{n-4}) + \mu_{vc}(G, v_{n-2}).$   
 $= 2^1 + 2^1 + 2^3 + \cdots + 2^{i-4} + 2^{i-2} + 2^1 + 2^3 + \cdots + 2^{n-(i+4)} + 2^{n-(i+2)}$ . Thus, we have ended up with

$\sum_{l=1}^{\lfloor \frac{i}{2} \rfloor} 2^{2(l)-1} + \sum_{m=0}^{\lfloor \frac{n-(i+2)}{2} \rfloor} 2^{2(m)+1} + 2$  pebbles which is less than the total number of pebbles available.

#### Case 4:

Consider the hub vertex as the source vertex. Then  $\dot{s} = v_0$  and

$$D_{vc}^6 = \{v_{n-2}, v_{n-4}, v_{n-6}, \dots, v_2, v_0\}.$$

The detour monophonic distance from any other vertex of  $F_n$  to  $v_0$  is 1. Thus in order to cover the vertices of  $D_{vc}^6$ , we have the pebble distribution that sums up as follows.

$$\begin{aligned} & \mu_{vc}(G, v_0) + \mu_{vc}(G, v_2) + \mu_{vc}(G, v_4) + \cdots + \mu_{vc}(G, v_{n-6}) + \mu_{vc}(G, v_{n-4}) + \mu_{vc}(G, v_{n-2}) \\ & = 2^0 + 2^1 + 2^1 + \cdots + 2^1 + 2^1 + 2^1 \end{aligned}$$

$2 \lfloor \frac{n-1}{2} \rfloor + 1$  which is less than the considered number of pebbles.

Hence,

$$\mu_{vc}(F_n) = \sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2, \text{ when } n \text{ is even.}$$

□

**Theorem 2.4.** For Fan  $F_n$  ( $n$  is odd),  $\mu_{vc}(F_n) = \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)} + 2$ .

*Proof.* The proof is similar to that of Theorems 2.2 and 2.3.

□

**Theorem 2.5.** For Cycle  $C_n (n \geq 3)$ ,

$$\mu_{vc}(C_n) = \begin{cases} 2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2(i)}\right) + 2^{\frac{n}{2}} + 1, & \text{if } n \text{ is even and } n \equiv 0(\text{mod } 4) \\ 2\left(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\frac{n}{2}-1} 2^{2(j)}\right) + 1, & \text{if } n \text{ is even and } n \equiv 2(\text{mod } 4) \\ \sum_{k=\lfloor \frac{n}{2} \rfloor + 1}^{n-2} 2^k + 3, & \text{if } n \text{ is odd.} \end{cases}$$

*Proof.* Let  $C_n$  be an even cycle of order  $n (n \geq 3)$ .

**Case 1:**

When  $n \equiv 0(\text{mod } 4)$ .

Consider the detour monophonic vertex cover set  $D_{vc}^1 = \{v_{i+2(j)} | i = 1, 2, 3, \dots, n; j = 0, 1, 2, \dots, \frac{n}{4} - 1, \frac{n}{4}, \frac{n}{4} + 1, \dots, \frac{n}{2} - 2, \frac{n}{2} - 1\}$ , the indices run over modulus  $n$ . Thus  $|D_{vc}^1| = \frac{n}{2}$ .

Let  $\dot{s} = v_1$  and the initial configuration be  $2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2i}\right) + 2^{\frac{n}{2}}$ . In order to cover the vertices  $v_{(n-1)}$  we require  $2^{n-2}$  pebbles and to cover  $v_{(n-3)}$  we require  $2^{n-4}$  pebbles respectively using the detour monophonic path. Similar process can be carried on to cover the remaining vertices of  $D_{vc}^1$  using the following series of pebble distribution that sums up

$$\begin{aligned} & 2^{n-2} + 2^{n-4} + \dots + 2^{\frac{n}{2}+2} + 2^{\frac{n}{2}} + 2^{\frac{n}{2}+2} + \dots + 2^{n-2} + 2^{n-4} \\ &= 2(2^{n-2} + 2^{n-4} + \dots + 2^{\frac{n}{2}+2}) + 2^{\frac{n}{2}} \\ &= 2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2i}\right) + 2^{\frac{n}{2}}. \end{aligned}$$

Thus we can cover the vertices  $v_3$  and  $v_5$  with  $2^{n-2}$  and  $2^{n-4}$  pebbles and we are left with no pebble to cover the source vertex  $v_1$ . Therefore, the detour monophonic vertex cover number  $\mu_{vc}(C_n) \geq 2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2(i)}\right) + 2^{\frac{n}{2}} + 1$ .

	$v_i$	$v_{i+2}$	$v_{i+4}$	$\dots$	$v_{\frac{n}{2}+i-2}$	$v_{\frac{n}{2}+i}$	$v_{\frac{n}{2}+i+2}$	$\dots$	$v_{n+i-4}$	$v_{n+i-2}$
$v_i$	0	$n-2$	$n-4$	$\dots$	$\frac{n}{2}+2$	$\frac{n}{2}$	$\frac{n}{2}+2$	$\dots$	$n-4$	$n-2$

TABLE 16. Detour monophonic distance from  $v_i$  to  $V(D_{vc}^1)$ .

Obviously  $v_i$  is covered with one pebble. From Table 16, in order to cover the vertices  $v_{i+2}$  and  $v_{n+i-2}$  we require  $2^{n-2}$  pebbles each and to cover  $v_{i+4}$  and  $v_{n+i-4}$  we require  $2^{n-4}$  pebbles each. Similarly to cover  $v_{\frac{n}{2}+i-2}$  and  $v_{\frac{n}{2}+i+2}$ , we require  $2^{\frac{n}{2}+2}$  pebbles. We need  $2^{\frac{n}{2}}$  pebbles to cover the vertex  $v_{i+\frac{n}{2}}$ . Thus in order to cover the vertices in  $D_{vc}^1$  we distribute the pebbles as follows.  $\mu_{vc}(G, v_i) + \mu_{vc}(G, v_{i+2}) + \mu_{vc}(G, v_{i+4}) + \dots + \mu_{vc}(G, v_{\frac{n}{2}+i-2}) + \mu_{vc}(G, v_{\frac{n}{2}+i}) + \mu_{vc}(G, v_{\frac{n}{2}+i+2}) + \dots + \mu_{vc}(G, v_{n+i-4}) + \mu_{vc}(G, v_{n+i-2})$ .

$$\begin{aligned} &= 2^0 + 2^{n-2} + 2^{n-4} + \dots + 2^{\frac{n}{2}+2} + 2^{\frac{n}{2}} + 2^{\frac{n}{2}+2} + \dots + 2^{n-4} + 2^{n-2} \\ &= 1 + 2(2^{n-2} + 2^{n-4} + \dots + 2^{\frac{n}{2}+2}) + 2^{\frac{n}{2}} \\ &= 2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2i}\right) + 2^{\frac{n}{2}} + 1. \end{aligned}$$

Since all the vertices are of same detour monophonic distance, the number of pebbles required to place on any  $v_i \in V(D_{vc}^1)$  is  $2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2(i)}\right) + 2^{\frac{n}{2}} + 1$ .

Thus,  $\mu_{vc}(C_n) = 2\left(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2i}\right) + 2^{\frac{n}{2}} + 1$ .

**Case 2:**

When  $n$  is even and  $n \equiv 2 \pmod{4}$ , let  $D_{vc}^2 = \{v_{i+2j} | i = 1, 2, 3, \dots, n; j = 0, 1, 2, \dots, \lfloor \frac{n}{4} - 1 \rfloor, \lfloor \frac{n}{4} \rfloor, \lfloor \frac{n}{4} + 1 \rfloor, \dots, \frac{n}{2} - 2, \frac{n}{2} - 1\}$  be the vertex cover set, where the indices run over modulus  $n$ . Thus,  $|D_{vc}^2| = \frac{n}{2}$ . Now consider the configuration which engages  $2(\sum_{j=\lfloor \frac{n}{4} \rfloor+1}^{\frac{n}{2}-1} 2^{2(j)})$  pebbles to cover the vertices of  $C_n$ . To cover the vertices  $v_{i+2}$  and  $v_{i+n-2}$ , we need  $2^{n-2}$  pebbles. Similarly to cover the vertices  $v_{i+4}$  and  $v_{i+n-4}$ , we need  $2^{n-4}$  pebbles. We use  $2^{\frac{n}{2}+1}$  pebbles to cover  $v_{i+2(\lfloor \frac{n}{4} \rfloor)}$  and  $v_{i+2(\lfloor \frac{n}{4} \rfloor+1)}$ . Obviously, we require a pebble to cover  $\dot{s}$ . Thus,  $\mu_{vc}(C_n) \geq 2(\sum_{j=\lfloor \frac{n}{4} \rfloor+1}^{\frac{n}{2}-1} 2^{2(j)}) + 1$ .

	$v_i$	$v_{i+2}$	$v_{i+4}$	$\dots$	$v_{\frac{n}{2}+i-3}$	$v_{\frac{n}{2}+i-1}$	$v_{\frac{n}{2}+i+1}$	$v_{\frac{n}{2}+i+3}$	$\dots$	$v_{n+i-4}$	$v_{n+i-2}$
$v_i$	0	$n-2$	$n-4$	$\dots$	$\frac{n}{2}+3$	$\frac{n}{2}+1$	$\frac{n}{2}+1$	$\frac{n}{2}+3$	$\dots$	$n-4$	$n-2$

TABLE 17. Detour monophonic distance from  $v_i$  to  $V(D_{vc}^2)$ .

Clearly only one pebble suffices to cover the vertex  $v_i (i = 1, 2, 3, \dots, n)$ . From Table 17, all the vertices of  $D_{vc}^2$  are covered by the following series of pebble distribution.

$$\begin{aligned} & \mu_{vc}(G, v_i) + \mu_{vc}(G, v_{i+2}) + \mu_{vc}(G, v_{i+4}) + \dots + \mu_{vc}(G, v_{\frac{n}{2}+i-3}) + \mu_{vc}(G, v_{\frac{n}{2}+i-1}) + \\ & \mu_{vc}(G, v_{\frac{n}{2}+i+1}) + \mu_{vc}(G, v_{\frac{n}{2}+i+3}) + \dots + \mu_{vc}(G, v_{n+i-4}) + \mu_{vc}(G, v_{n+i-2}). \\ & = 2^0 + 2^{n-2} + 2^{n-4} + \dots + 2^{\frac{n}{2}+1} + 2^{\frac{n}{2}+1} + \dots + 2^{n-2} + 2^{n-4} \\ & 1 + 2(2^{n-2} + 2^{n-4} + \dots + 2^{2(\frac{n}{2}-1)} + 2^{2(\frac{n}{2}-2)} + \dots + 2^{2(\lfloor \frac{n}{4} \rfloor+1)}) \end{aligned}$$

Thus,  $2(\sum_{j=\lfloor \frac{n}{4} \rfloor+1}^{\frac{n}{2}-1} 2^{2(j)}) + 1$ .

Since all the vertices of  $C_n$  are at equi distance, the result holds for any  $v_i \in V(C_n)$ , when  $n$  is even and  $n \equiv 2 \pmod{4}$ . Hence,  $\mu_{vc}(C_n) = 2(\sum_{j=\lfloor \frac{n}{4} \rfloor+1}^{\frac{n}{2}-1} 2^{2(j)}) + 1$ .

**Case 3:** When  $n$  is odd, assume the detour monophonic cover set  $D_{vc}^3 = \{v_{i+j} | i = 1, 2, 3, \dots, n; j = 0, 1, 2, \dots, n-1\}$ , where all the indices run over modulus  $n$  and  $|D_{vc}^3| = \lfloor \frac{n}{2} \rfloor + 1$ . Without loss of generality we fix  $\dot{s} = v_1$ .

Let the initial configuration be  $\sum_{k=\lfloor \frac{n}{2} \rfloor+1}^{n-2} 2^k + 2$ . The vertices  $v_3, v_5$  and  $v_7$  are covered with  $2^{n-2}, 2^{n-4}$  and  $2^{n-5}$  pebbles respectively. To cover the vertices  $v_{\lfloor \frac{n}{2} \rfloor-2}, v_{\lfloor \frac{n}{2} \rfloor}$  and  $v_{\lfloor \frac{n}{2} \rfloor+2}$ , we have to use  $2^{\lfloor \frac{n}{2} \rfloor+4}, 2^{\lfloor \frac{n}{2} \rfloor+2}$  and  $2^{\lfloor \frac{n}{2} \rfloor+1}$  pebbles on the corresponding vertices. Continuing in this manner we are left with  $v_1$  uncovered. Hence, the monophonic vertex cover pebbling number

$$\mu_{vc}(C_n) \geq \sum_{k=\lfloor \frac{n}{2} \rfloor+1}^{n-2} 2^k + 3.$$

If  $n$  is odd and  $n \equiv 1 \pmod{4}$ , then  $D_{vc}^3 = \{v_1, v_3, v_5, \dots, v_{\lfloor \frac{n}{2} \rfloor-1}, v_{\lfloor \frac{n}{2} \rfloor+1}, v_{\lfloor \frac{n}{2} \rfloor+3}, v_{\lfloor \frac{n}{2} \rfloor+5}, \dots, v_{n-4}, v_{n-2}, v_n\}$ . Without loss of generality consider  $v_1$  to be the source vertex.

If  $n$  is odd and  $n \equiv 3 \pmod{4}$ , then  $D_{vc}^{3*} = \{v_1, v_3, v_5, \dots, v_{\lfloor \frac{n}{2} \rfloor-2}, v_{\lfloor \frac{n}{2} \rfloor}, v_{\lfloor \frac{n}{2} \rfloor+2}, v_{\lfloor \frac{n}{2} \rfloor+4}, \dots, v_{n-4}, v_{n-2}, v_n\}$ . Without loss of generality let  $v_1$  be the source vertex.

	$v_1$	$v_3$	$v_5$	$\cdots$	$v_{\lfloor \frac{n}{2} \rfloor - 1}$	$v_{\lfloor \frac{n}{2} \rfloor + 1}$	$v_{\lfloor \frac{n}{2} \rfloor + 3}$	$v_{\lfloor \frac{n}{2} \rfloor + 5}$	$\cdots$	$v_{n-4}$	$v_{n-2}$	$v_n$
$v_1$	0	$n-2$	$n-4$	$\cdots$	$\lfloor \frac{n}{2} \rfloor + 3$	$\lfloor \frac{n}{2} \rfloor + 1$	$\lfloor \frac{n}{2} \rfloor + 2$	$\lfloor \frac{n}{2} \rfloor + 4$	$\cdots$	$n-5$	$n-3$	1

TABLE 18. Detour monophonic distance from  $v_1$  to  $V(D_{vc}^3)$ .

	$v_1$	$v_3$	$v_5$	$\cdots$	$v_{\lfloor \frac{n}{2} \rfloor - 2}$	$v_{\lfloor \frac{n}{2} \rfloor}$	$v_{\lfloor \frac{n}{2} \rfloor + 2}$	$v_{\lfloor \frac{n}{2} \rfloor + 4}$	$\cdots$	$v_{n-4}$	$v_{n-2}$	$v_n$
$v_1$	0	$n-2$	$n-4$	$\cdots$	$\lfloor \frac{n}{2} \rfloor + 4$	$\lfloor \frac{n}{2} \rfloor + 2$	$\lfloor \frac{n}{2} \rfloor + 1$	$\lfloor \frac{n}{2} \rfloor + 3$	$\cdots$	$n-5$	$n-3$	1

TABLE 19. Detour monophonic distance from  $v_1$  to  $V(D_{vc}^{3*})$ .

In order to cover the vertices of  $C_n$ , it suffices to cover the vertices of vertex cover set. From Tables 18 and 19, the pebbles are distributed sums up as follows.

$$= 2^0 + 2^1 + 2^{\lfloor \frac{n}{2} \rfloor + 1} + 2^{\lfloor \frac{n}{2} \rfloor + 2} + 2^{\lfloor \frac{n}{2} \rfloor + 3} + 2^{\lfloor \frac{n}{2} \rfloor + 4} + \cdots + 2^{n-5} + 2^{n-4} + 2^{n-3} + 2^{n-2}$$

$$= \sum_{k=\lfloor \frac{n}{2} \rfloor + 1}^{n-2} 2^k + 3.$$

$$\text{Thus, } \mu_{vc}(C_n) = \sum_{k=\lfloor \frac{n}{2} \rfloor + 1}^{n-2} 2^k + 3.$$

Since the vertices of  $C_n$  are equi distanced, the result is true for any  $v_i \in \{v_i | 2 \leq i \leq n\}$ , when  $n$  is odd.

Hence proved. □

**Theorem 2.6.** For a Wheel  $W_n$ ,

$$\mu_{vc}(W_n) = \begin{cases} \sum_{i=\frac{n}{2}}^{n-3} 2^i + 5, & \text{if } n \text{ is even} \\ 2\left(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}\right) + 2^{\lfloor \frac{n}{2} \rfloor} + 3, & \text{if } n \text{ is odd and } n \equiv 1 \pmod{4} \\ 2\left(\sum_{k=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(k)}\right) + 3, & \text{if } n \text{ is odd and } n \equiv 3 \pmod{4} \end{cases}$$

*Proof.* Let  $W_n = C_{n-1} + v_0$  be a wheel of order  $n$  with  $V(W_n) = \{v_i | 0 \leq i \leq n-1\}$  with  $|V(W_n)| = n$ .

**Case 1:** When  $n$  is even, assume the detour monophonic cover set  $D'_{vc} = \{v_0, v_{i+j} | i = 1, 2, 3, \dots, n-1; j = 0, 1, 2, \dots, n-2\}$ , where all the indices run over modulus  $n$  and  $|D'_{vc}| = \frac{n}{2} + 1$ . Without loss of generality consider  $\dot{s} = v_1$ . Let the initial configuration be  $\sum_{i=\frac{n}{2}}^{n-3} 2^i + 4$  pebbles assigned to cover the vertices of  $W_n$ . Since the detour monophonic distance between  $v_0$  and the source vertex is 1, we can cover 2 pebbles. The vertices  $v_3, v_5$  and  $v_7$  are covered with  $2^{n-3}, 2^{n-5}$  and  $2^{n-7}$  pebbles respectively. Proceeding in this manner we cover the all vertices of  $D'_{vc}$  except the source vertex with  $\sum_{i=\frac{n}{2}}^{n-3} 2^i + 4$  pebbles. Thus, continuing in this manner we are left with  $v_1$  uncovered. Hence,

$$\mu_{vc}(W_n) \geq \sum_{i=\frac{n}{2}}^{n-3} 2^i + 5.$$

Now, consider  $n$  is even and  $n \equiv 0 \pmod{4}$ .

Let the vertex cover set  $V(D_{vc}^1) = \{v_0, v_1, v_3, v_5, \dots, v_{\frac{n}{2}-3}, v_{\frac{n}{2}-1}, v_{\frac{n}{2}+1}, v_{\frac{n}{2}+3}, \dots, v_{n-3}, v_{n-1}\}$ , where  $|D_{vc}^1| = \frac{n}{2} + 1$ . Fix  $\dot{s} = v_1$ .

	$v_0$	$v_1$	$v_3$	$v_5$	$\cdots$	$v_{\frac{n}{2}-3}$	$v_{\frac{n}{2}-1}$	$v_{\frac{n}{2}+1}$	$v_{\frac{n}{2}+3}$	$\cdots$	$v_{n-3}$	$v_{n-1}$
$v_1$	1	0	$n-3$	$n-5$	$\cdots$	$\frac{n}{2}+3$	$\frac{n}{2}+1$	$\frac{n}{2}$	$\frac{n}{2}+2$	$\cdots$	$n-4$	1

TABLE 20. Detour monophonic distance from  $v_1$  to  $V(D_{vc}^1)$ .

Let  $n$  is even and  $n \equiv 2(\text{mod } 4)$ . Then the vertex cover set  $V(D_{vc}^{1*}) = \{v_0, v_1, v_3, v_5, \cdots, v_{\frac{n}{2}-2}, v_{\frac{n}{2}}, v_{\frac{n}{2}+2}, v_{\frac{n}{2}+4}, \cdots, v_{n-3}, v_{n-1}\}$  and  $|D_{vc}^{1*}| = \frac{n}{2} + 1$ . Without loss of generality take  $\dot{s} = v_1$ .

	$v_0$	$v_1$	$v_3$	$v_5$	$\cdots$	$v_{\frac{n}{2}-2}$	$v_{\frac{n}{2}}$	$v_{\frac{n}{2}+2}$	$v_{\frac{n}{2}+4}$	$\cdots$	$v_{n-3}$	$v_{n-1}$
$v_1$	1	0	$n-3$	$n-5$	$\cdots$	$\frac{n}{2}+2$	$\frac{n}{2}$	$\frac{n}{2}+1$	$\frac{n}{2}+3$	$\cdots$	$n-4$	1

TABLE 21. Detour monophonic distance from  $v_1$  to  $V(D_{vc}^{1*})$ .

From Tables 20 and 21, in order to cover the vertices of  $W_n$ , it suffices to cover the vertices of the corresponding vertex cover set. Thus, the pebble distribution with either of the above vertex cover sets is given by,

$$2^1 + 2^0 + 2^{\frac{n}{2}} + 2^{\frac{n}{2}+1} + 2^{\frac{n}{2}+2} + 2^{\frac{n}{2}+3} + \cdots + 2^{n-5} + 2^{n-4} + 2^{n-3}, \text{ which sums up into } \sum_{i=\frac{n}{2}}^{n-3} 2^i + 5.$$

The result is true for all  $v_i (2 \leq i \leq n-1)$ .

Let the source vertex be the hub. Then  $\dot{s} = v_0$ . Without loss of generality let  $D_{vc}^4 = \{v_0, v_1, v_3, \cdots, v_{n-5}, v_{n-3}, v_{n-1}\}$ .

The detour monophonic distance from any other vertex of  $W_n$  to  $v_0$  is 1. Thus in order to cover the vertices of  $D_{vc}^4$ , we have the pebble distribution that sums up as follows.  $\mu_{vc}(G, v_0) + \mu_{vc}(G, v_1) + \mu_{vc}(G, v_3) + \cdots + \mu_{vc}(G, v_{n-5}) + \mu_{vc}(G, v_{n-3}) + \mu_{vc}(G, v_{n-1})$   
 $= 2^0 + 2^1 + 2^1 + \cdots + 2^1 + 2^1 + 2^1$   
 $2(\frac{n}{2}) + 1 = n + 1$  which is less than the considered number of pebbles.

Thus,  $\mu_{vc}(W_n) = \sum_{i=\frac{n}{2}}^{n-3} 2^i + 5$ . Hence proved.

**Case 2:** When  $W_n$  be an odd wheel of order  $n$ .

Let  $(n \geq 3)$  and  $n \equiv 1(\text{mod } 4)$ .

Consider the detour monophonic vertex cover set  $D_{vc}'' = \{v_0, v_{2(k)-1} | k = 1, 2, 3, \cdots, n-2\}$  and thus  $|D_{vc}''| = \lfloor \frac{n}{2} \rfloor$ .

Let  $\dot{s} = v_1$  and the initial configuration be  $2(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}) + 2^{\lfloor \frac{n}{2} \rfloor} + 1$ . In order to cover the vertices  $v_{(n-3)}$  we require  $2^{n-3}$  pebbles and to cover  $v_{(n-5)}$  we require  $2^{n-5}$  pebbles respectively using the detour monophonic path. Similar process can be carried on to cover the remaining vertices of  $D_{vc}^1$  using the following series of pebble distribution that sums up,

$$2^1 + 2^{n-3} + 2^{n-5} + \cdots + 2^{\lfloor \frac{n}{2} \rfloor + 2} + 2^{\lfloor \frac{n}{2} \rfloor} + 2^{\lfloor \frac{n}{2} \rfloor + 2} + \cdots + 2^{n-5} + 2^{n-3}$$

$$= 2 + 2(2^{n-3} + 2^{n-5} + \cdots + 2^{\lfloor \frac{n}{2} \rfloor + 2}) + 2^{\lfloor \frac{n}{2} \rfloor}$$

$$= 2\left(\sum_{i=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)}\right) + 2^{\lfloor \frac{n}{2} \rfloor} + 2.$$

Thus we can cover the vertices  $v_3$  and  $v_5$  with  $2^{n-3}$  and  $2^{n-5}$  pebbles and we are left with no pebble to cover the source vertex  $v_1$ . Therefore, the detour monophonic vertex cover number  $\mu_{vc}(W_n) \geq 2\left(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}\right) + 2^{\lfloor \frac{n}{2} \rfloor} + 3$ .

Now let  $\dot{s} = v_i (1 \leq i \leq n-1)$  and  $D_{vc}^2 = \{v_0, v_{i+2}, v_{i+4}, \dots, v_{\lfloor \frac{n}{2} \rfloor + i - 2}, v_{\lfloor \frac{n}{2} \rfloor + i}, v_{\lfloor \frac{n}{2} \rfloor + i + 2}, \dots, v_{n+i-5}, v_{n+i-3}\}$ , where the indices run over modulus  $n$ .

	$v_0$	$v_i$	$v_{i+2}$	$v_{i+4}$	$\dots$	$v_{\lfloor \frac{n}{2} \rfloor + i - 2}$	$v_{\lfloor \frac{n}{2} \rfloor + i}$	$v_{\lfloor \frac{n}{2} \rfloor + i + 2}$	$\dots$	$v_{n+i-5}$	$v_{n+i-3}$
$v_i$	1	0	$n-3$	$n-5$	$\dots$	$\lfloor \frac{n}{2} \rfloor + 2$	$\lfloor \frac{n}{2} \rfloor$	$\lfloor \frac{n}{2} \rfloor + 2$	$\dots$	$n-5$	$n-3$

TABLE 22. Detour monophonic distance from  $v_i$  to  $V(D_{vc}^2)$ .

Obviously  $v_i$  is covered with one pebble. From Table 22,  $D_m(v_0, v_i) = 1$  and so  $v_0$  is covered with two pebbles. In order to cover the vertices  $v_{i+2}$  and  $v_{n+i-3}$  we require  $2^{n-3}$  pebbles each and to cover  $v_{i+4}$  and  $v_{n+i-5}$  we require  $2^{n-5}$  pebbles each. Similarly to cover  $v_{\lfloor \frac{n}{2} \rfloor + i - 2}$  and  $v_{\lfloor \frac{n}{2} \rfloor + i + 2}$ , we require  $2^{\lfloor \frac{n}{2} \rfloor + 2}$  pebbles. We need  $2^{\lfloor \frac{n}{2} \rfloor}$  pebbles to cover the vertex  $v_{\lfloor \frac{n}{2} \rfloor + i}$ . Thus in order to cover the vertices in  $D_{vc}^2$  we distribute the pebbles as follows.  $\mu_{vc}(G, v_0) + \mu_{vc}(G, v_i) + \mu_{vc}(G, v_{i+2}) + \mu_{vc}(G, v_{i+4}) + \dots + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor + i - 2}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor + i}) + \mu_{vc}(G, v_{i + \frac{n+4}{2}}) + \dots + \mu_{vc}(G, v_{n+i-5}) + \mu_{vc}(G, v_{n+i-3})$ .

$$= 2^1 + 2^0 + 2^{n-3} + 2^{n-5} + \dots + 2^{\lfloor \frac{n}{2} \rfloor + 2} + 2^{\lfloor \frac{n}{2} \rfloor} + 2^{\lfloor \frac{n}{2} \rfloor + 2} + \dots + 2^{n-5} + 2^{n-3}$$

$$= 2 + 1 + 2(2^{n-3} + 2^{n-5} + \dots + 2^{\lfloor \frac{n}{2} \rfloor + 2}) + 2^{\lfloor \frac{n}{2} \rfloor}$$

$$= 2\left(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}\right) + 2^{\lfloor \frac{n}{2} \rfloor} + 3.$$

Since all the vertices are of same detour monophonic distance, the number of pebbles required to cover from any other vertex of  $W_n$  except the hub is  $2\left(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}\right) + 2^{\lfloor \frac{n}{2} \rfloor} + 3$ .

Assume the source vertex is the hub.

Then  $\dot{s} = v_0$ . Without loss of generality let  $D_{*vc} = \{v_0, v_1, v_3, \dots, v_{n-6}, v_{n-4}, v_{n-2}\}$ .

Since  $D_m(x, v_0) = 1$ , where  $x$  is any vertex of  $W_n$ , Thus in order to cover the vertices of  $D_{*vc}$ , we have the pebble distribution that sums up as follows.  $\mu_{vc}(G, v_0) + \mu_{vc}(G, v_1) + \mu_{vc}(G, v_3) + \dots + \mu_{vc}(G, v_{n-6}) + \mu_{vc}(G, v_{n-4}) + \mu_{vc}(G, v_{n-2})$

$$= 2^0 + 2^1 + 2^1 + \dots + 2^1 + 2^1 + 2^1$$

$$= 2\lfloor \frac{n}{2} \rfloor + 1$$

which is less than the considered number of pebbles.

Thus,  $\mu_{vc}(W_n) = 2\left(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}\right) + 2^{\lfloor \frac{n}{2} \rfloor} + 3$ .

Hence proved.

### Case 3:

When  $n$  is odd and  $n \equiv 3 \pmod{4}$ , let  $D_{vc}^3 = \{v_{i+2j} | i = 1, 2, 3, \dots, n; j = 0, 1, 2, \dots, \lfloor \frac{n}{4} - 1 \rfloor, \lfloor \frac{n}{4} \rfloor, \lfloor \frac{n}{4} + 1 \rfloor, \dots, \frac{n}{2} - 2, \frac{n}{2} - 1\}$  be the vertex cover set, where

the indices run over modulus  $n$ . Thus,  $|D_{vc}^3| = \frac{n}{2}$ . Now consider the configuration which engages  $2(\sum_k 2^{2(k)}) + 2$  pebbles to cover the vertices of  $W_n$ . To cover the vertices  $v_{i+2}$  and  $v_{i+n-2}$ , we need  $2^{n-2}$  pebbles. Similarly to cover the vertices  $v_{i+4}$  and  $v_{i+n-4}$ , we need  $2^{n-4}$  pebbles. We use  $2^{\frac{n}{2}+1}$  pebbles to cover  $v_{i+2(\lfloor \frac{n}{4} \rfloor)}$  and  $v_{i+2(\lfloor \frac{n}{4} \rfloor+1)}$ . Obviously, we require a pebble to cover  $\dot{s} = v_i (i = 1, 2, 3, \dots, n-1)$ .

Thus,  $\mu_{vc}(W_n) \geq 2(\sum_{k=\lfloor \frac{n}{4} \rfloor+1}^{\lfloor \frac{n}{2} \rfloor-1} 2^{2(k)}) + 3$ .

	$v_i$	$v_{i+2}$	$v_{i+4}$	$\dots$	$v_{\lfloor \frac{n}{2} \rfloor+i-3}$	$v_{\lfloor \frac{n}{2} \rfloor+i-1}$	$v_{\lfloor \frac{n}{2} \rfloor+i+1}$	$v_{\lfloor \frac{n}{2} \rfloor+i+3}$	$\dots$	$v_{n+i-5}$	$v_{n+i-3}$
$v_i$	0	$n-3$	$n-5$	$\dots$	$\lfloor \frac{n}{2} \rfloor + 3$	$\lfloor \frac{n}{2} \rfloor + 1$	$\lfloor \frac{n}{2} \rfloor + 1$	$\lfloor \frac{n}{2} \rfloor + 3$	$\dots$	$n-5$	$n-3$

TABLE 23. Detour monophonic distance from  $v_i$  to  $V(D_{vc}^3)$ .

Clearly only one pebble suffices to cover the vertex  $v_i (i = 1, 2, 3, \dots, n)$ . From Table 23, all the vertices of  $D_{vc}^2$  are covered by the following series of pebble distribution.

$$\begin{aligned} & \mu_{vc}(G, v_0) + \mu_{vc}(G, v_i) + \mu_{vc}(G, v_{i+2}) + \mu_{vc}(G, v_{i+4}) + \dots + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor+i-3}) + \\ & \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor+i-1}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor+i+1}) + \mu_{vc}(G, v_{\lfloor \frac{n}{2} \rfloor+i+3}) + \dots + \mu_{vc}(G, v_{n+i-5}) + \mu_{vc}(G, v_{n+i-3}). \\ & = 2^1 + 2^0 + 2^{n-3} + 2^{n-5} + \dots + 2^{\lfloor \frac{n}{2} \rfloor+3} + 2^{\lfloor \frac{n}{2} \rfloor+1} + 2^{\lfloor \frac{n}{2} \rfloor+1} + 2^{\lfloor \frac{n}{2} \rfloor+3} + \dots + 2^{n-5} + 2^{n-3} \\ & = 2(\sum_{k=\lfloor \frac{n}{4} \rfloor+1}^{\lfloor \frac{n}{2} \rfloor-1} 2^{2(k)}) + 3. \end{aligned}$$

Since all the vertices except  $v_0$  of  $W_n$  are at equi distance, the result holds for any  $v_i \in V(W_n) (2 \leq i \leq n-1)$ , when  $n$  is odd and  $n \equiv 3 \pmod{4}$ .

When the source vertex is the hub, the result follows from the previous discussion of this Theorem and the configuration of pebbles is  $2\lfloor \frac{n}{2} \rfloor + 1$  which is less than the considered number of pebbles.

Hence,

$$\mu_{vc}(W_n) = 2(\sum_{k=\lfloor \frac{n}{4} \rfloor+1}^{\lfloor \frac{n}{2} \rfloor-1} 2^{2(k)}) + 3.$$

□

### 3. Algorithm to find DMVCPN

**Step 1:** Find the vertex cover set  $D_{vc}$  with minimum cardinality for a graph  $G$ , which covers all the edges of  $G$ .

**Step 2:** Fix the source vertex  $\dot{s}$ .

**Step 3:** Find the detour monophonic distance  $D_m(\dot{s}, v)$ , where  $v \in V(G)$ .

**Step 4:** By distributing the pebbles place one pebble each to cover the vertices of  $D_{vc}$  such that the number of pebbles used is minimum for all configuration of pebbles.

**Step 5:** Repeat Step 4 for various source vertices until reaching the minimal number of pebbles that cover all the vertices of  $D_{vc}$  and so the vertices of  $G$ .

**Step 6:** End the process if each vertex of  $G$  is considered to be  $\dot{s}$ .

### 4. Various Pebbling Numbers of $G$

The various graph pebbling numbers displayed in the Table 24 exhibit that DMVCPN of a graph is distinct.

Standard Graph $G$	Detour pebbling number $f^*(G)$ [7]	Monophonic pebbling number $\mu(G)$ [6]	Detour monophonic vertex cover pebbling number $\mu_{vc}(G)$
$P_n$	$2^{n-1}$	$2^{n-1}$	$\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(i)+1}$ , $n$ is odd $\sum_{j=0}^{\frac{n}{2}-1} 2^{2(j)}$ , $n$ is even
$C_n$	$2^{n-1}$	$2^{n-2} + 1$	$\begin{cases} 2(\sum_{i=\frac{n}{4}+1}^{\frac{n}{2}-1} 2^{2(i)}) + 2^{\frac{n}{2}} + 1, & \text{if } n \text{ is even and } n \equiv 0(mod 4) \\ 2(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\frac{n}{2}-1} 2^{2(j)}) + 1, & \text{if } n \text{ is even and } n \equiv 2(mod 4) \\ \sum_{k=\lfloor \frac{n}{2} \rfloor + 1}^{n-2} 2^k + 3, & \text{if } n \text{ is odd} \end{cases}$
$F_n$	$2^{n-1}$	$2^{n-2} + 1$	$\sum_{j=0}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)} + 2$ , $n$ is odd $\sum_{i=0}^{\frac{n}{2}-2} 2^{2(i)+1} + 2$ , $n$ is even
$W_n$	$2^n$	$2^{n-2} + 2$	$\begin{cases} \sum_{i=\frac{n}{2}}^{n-3} 2^i + 5, & \text{if } n \text{ is even} \\ 2(\sum_{j=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(j)}) + 2^{\lfloor \frac{n}{2} \rfloor} + 3, & \text{if } n \text{ is odd and } n \equiv 1(mod 4) \\ 2(\sum_{k=\lfloor \frac{n}{4} \rfloor + 1}^{\lfloor \frac{n}{2} \rfloor - 1} 2^{2(k)}) + 3, & \text{if } n \text{ is odd and } n \equiv 3(mod 4) \end{cases}$

TABLE 24. Various pebbling numbers for some standard graphs

### 5. Conclusion

In this article, we have combined cover pebbling number and detour monophonic number to obtain an interesting variate of graph theory, the detour monophonic vertex cover pebbling number the interesting areas of graph theory. We have studied and determined  $\mu_{vc}(G)$  for some standard graphs. To evaluate the specific values of  $\mu_{vc}(G)$  for other classes of well-known graphs is an open area of research.

### REFERENCES

- [1] Bondy J.A. and Murty U.S.R., *Graph theory with applications*, London: Macmillan, Vol. 290 (1977).
- [2] Chartrand G., *Introductory Graph theory*, Dover books on Mathematics, (1985).
- [3] Crull, B., Cundiff, T., Feltman, P., Hurlbert, G.H., Pudwell, L., Szaniszlo, Z. and Tuza, Z., *The cover pebbling number of graphs*, Discrete mathematics, 296(1),(2005), pp.15-23. DOI: 10.1016/j.disc.2005.03.009
- [4] Daoud S. N. and Mohamed K., *The complexity of some families of cycle-related graphs*, Journal of Taibah University for Science II (2017), pp: 205-228. DOI: 10.1016/ j.jtusci.2016.04.002
- [5] Hurlbert G. *Graph pebbling*, Handbook of graph theory, (2013), pp:1428-1449. DOI:10.1016/j.disc.2005.03.009
- [6] Lourdusamy A., Dhivvianandam I., and Iammal S.K., *Monophonic pebbling number of some standard graphs*, Journal of Applied Mathematics Informatics, 21 (2022), pp: 77-83 DOI: 10.14317/jami.2024.077
- [7] Lourdusamy A. and Nellainayaki S.S., *Detour pebbling on path related graphs*, Advances in Mathematics: Scientific Journal, 10(4), (2021), pp.2017-2024. DOI:10.37418/amsj.10.4.16
- [8] Michael E. S. and Imelda S. A., *The cover pebbling number of the join of some graphs*, Applied Mathematical Science 8(86), (2014), pp.4275-4283. DOI: 10.12988/ams.2014.45377

- [9] Pushpam P. R. L. and Suseendran C., *Secure vertex cover of a graph*, Discrete Mathematics, Algorithms and Applications, 9(2), (2017), pp: 1750026. DOI:10.1142/S1793830917500264
- [10] Titus P. and Ganesamoorthy K., *Upper detour monophonic number of a graph*, Electronic Notes in Discrete Mathematics, 53 (2016), pp.331-342. DOI: 10.1016/j.endm.2016.05.028