# An analogue of Girstmair's formula in function fields

By D. Shiomi

# 1 Introduction

Suppose that p is an odd prime and g > 1 is a primitive root modulo p. Let M be a number field contained in the p-th cyclotomic field. In [2, 3], Girstmair found a surprising relation between the relative class number of M and the digits of 1/p in base g. In particular, he obtained

**Theorem 1.1.** Suppose that  $p \ge 7$  and  $p \equiv 3 \pmod{4}$ . Then,

$$(g+1)h(-p) = \sum_{k=1}^{p-1} (-1)^k x_k$$

where h(-p) is the class number of  $\mathbb{Q}(\sqrt{-p})$ , and  $1/p = \sum_{k=1}^{\infty} x_k g^{-k}$  is the digit expansion of 1/p in base g.

After his works, this type of formulas has been further studied by several authors (cf. [5, 6, 7, 8]).

In this paper, we consider an analogue of Girstmair's formula in function fields. Let  $\mathbb{F}_q$  be the finite field with q elements of characteristic p. Let  $K = \mathbb{F}_q(T)$  be the rational function field over  $\mathbb{F}_q$ , and put  $\mathbb{A} = \mathbb{F}_q[T]$ . Suppose that  $P \in \mathbb{A}$  is monic irreducible and  $G \in \mathbb{A}$  is a primitive root modulo P. Let Lbe an extension field of K contained in the P-th cyclotomic function field. We denote by  $h_L$  the divisor class number of L, i.e., the order of the divisor class group of L of degree 0.

The goal of this paper is to give relations between the plus and minus parts of  $h_L$  and the digits of 1/P in base G (see Theorem 3.1 and 3.2). As a consequence, we prove the following theorem, which can be regarded as an analogue of Theorem 1.1.

**Theorem 1.2.** Suppose that gcd(2,q) = 1,  $d = \deg P$ , and  $\deg G \ge \deg P$ . Let

<sup>2020</sup> Mathematics Subject Classification: 11R29, 11R60, 11T55

Key words and Phrases: Digits, Class numbers, Quadratic function fields, Cyclotomic function fields

 $L = K(\sqrt{(-1)^d P})$ . Let  $1/P = \sum_{k=1}^{\infty} H_k/G^k$  be the digit expansion of 1/P in base G. Then,

$$h_L = \begin{cases} \sum_{k=1}^r (-1)^k \deg H_k & \text{if } d \text{ is even,} \\ (-\eta_G) \sum_{k=1}^r (-1)^k \varepsilon_k, & \text{if } d \text{ is odd.} \end{cases}$$

Here,  $r = (q^d - 1)/(q - 1)$  and

$$\eta_G = \begin{cases} 1 & \text{if } \delta(G) \in (\mathbb{F}_q^{\times})^2, \\ -1 & \text{if } \delta(G) \notin (\mathbb{F}_q^{\times})^2, \end{cases} \qquad \varepsilon_k = \begin{cases} 1 & \text{if } \delta(H_k) \in (\mathbb{F}_q^{\times})^2, \\ -1 & \text{if } \delta(H_k) \notin (\mathbb{F}_q^{\times})^2, \end{cases}$$

where  $\delta(I)$  is the leading coefficient of  $I \in \mathbb{A}$  if  $I \neq 0$ , and  $\delta(0) = 0$ .

Remark 1.1. Assume the notation in Theorem 1.2. Consider

$$S_{P,G} = \sum_{k=1}^{q^{d}-1} (-1)^{k} H_{k},$$

which is analogous to the alternating sum in Theorem 1.1. However, this sum seems to have no connection with the divisor class number of  $K(\sqrt{(-1)^d P})$ . In fact, we have  $S_{P,G} = 0$  by considering the case  $\alpha = -1$  in Corollary 2.1 (see Section 2.1).

This paper is organized as follows. In Section 2.1, we study about some properties of digits of rational functions over finite fields. In Section 2.2, we review basic facts about cyclotomic function fields and some formulas for divisor class numbers. In Section 3, we give the main results of this paper and some examples.

# 2 Preparations

#### 2.1 The digit expansion

Let  $v_{\infty}$  be the valuation of K satisfying with  $v_{\infty}(T) = -1$ . Then

$$v_{\infty}(F_1/F_2) = \deg F_2 - \deg F_1 \quad (F_1, F_2 \in \mathbb{A} \setminus \{0\}).$$

We denote by  $K_{\infty}$  the completion of K by  $v_{\infty}$ .

**Proposition 2.1.** Let  $G \in \mathbb{A}$  with deg  $G \ge 1$ . Define

$$S_G = \{I \in \mathbb{A} \mid \deg I < \deg G\} \cup \{0\}.$$

Any rational function  $f \in K$  can be uniquely expressed as

$$f = \sum_{k=0}^{\infty} \frac{H_k}{G^k} \quad in \ K_{\infty} \tag{1}$$

where  $H_0 \in \mathbb{A}$  and  $H_k \in S_G$   $(k \ge 1)$ . Furthermore,  $v_{\infty}(f) > 0$  if and only if  $H_0 = 0$ . We call the expression (1) the digit expansion of f in base G.

*Proof.* Suppose  $f = F_1/F_2$   $(F_1, F_2 \in \mathbb{A})$ . Let  $H_0, R \in \mathbb{A}$  with  $F_1 = H_0F_2 + R$  and deg  $R < \deg F_2$ . Putting  $f_1 = R/F_2$ , then we have

$$f = H_0 + f_1$$
 and  $v_{\infty}(f_1) > 0$ .

For  $k \geq 1$ , we inductively determine  $H_k \in \mathbb{A}$  and  $f_{k+1} \in K$  satisfying with

$$Gf_k = H_k + f_{k+1}$$
 and  $v_{\infty}(f_{k+1}) > 0.$ 

Then  $H_k \in S_G$   $(k \ge 1)$ , and

$$f = H_0 + \frac{H_1}{G} + \dots + \frac{H_k}{G^k} + \frac{f_{k+1}}{G^k}$$

By  $\lim_{k\to\infty} v_{\infty}(f_{k+1}/G^k) = \infty$ , we get (1).

We next prove the uniqueness of (1). Assume that

$$\sum_{k=0}^{\infty} \frac{H_k}{G^k} = \sum_{k=0}^{\infty} \frac{I_k}{G^k},$$

where  $H_0, I_0 \in \mathbb{A}$  and  $H_k, I_k \in S_G$   $(k \ge 1)$ . Suppose that there exists an integer  $k_0 \ge 0$  such that  $H_k = I_k$   $(k < k_0)$  and  $H_{k_0} \ne I_{k_0}$ . Then

$$H_{k_0} - I_{k_0} = \sum_{k=1}^{\infty} \frac{I_{k+k_0} - H_{k+k_0}}{G^k}.$$

Since  $v_{\infty}((I_{k+k_0} - H_{k+k_0})/G^k) > 0$  for  $k \ge 1$ , we have  $v_{\infty}(H_{k_0} - I_{k_0}) > 0$ . This contradicts  $H_{k_0} - I_{k_0} \in \mathbb{A} \setminus \{0\}$ . Thus  $H_k = I_k$  for  $k \ge 0$ .

Finally, we see that

$$v_{\infty}(f) > 0 \iff v_{\infty}(H_0) > 0 \iff H_0 = 0$$
 in (1)

since  $v_{\infty}(H_k/G^k) > 0$  for  $k \ge 1$ .

For  $M \in \mathbb{A}$ , let  $\mathcal{R}_M = \mathbb{A}/M\mathbb{A}$ .

**Proposition 2.2.** Let  $G, M \in \mathbb{A}$  with  $\deg G \geq 1$ ,  $\deg M \geq 1$ , and  $\gcd(G, M) = 1$ . Let g be the order of  $G \mod M$  in  $\mathcal{R}_M^{\times}$ . Suppose that  $1/M = \sum_{k=1}^{\infty} H_k/G^k$  is the digit expansion of 1/M in base G. For  $k \geq 0$ , we choose  $G_k \in \mathbb{A}$  with  $\deg G_k < \deg M$  and  $G_k \equiv G^k \pmod{M}$ .

(1) For  $k \ge 1$ , we have

$$H_k = \frac{GG_{k-1} - G_k}{M}.$$
(2)

In particular, we obtain  $H_k \neq 0$  (k = 1, 2...) when deg  $G \ge \deg M$ .

(2) The sequence  $\{H_k\}_{k>1}$  is purely periodic, and its period l is equal to g.

*Proof.* (1) Put  $I_k = (GG_{k-1} - G_k)/M$  for  $k \ge 1$ . Since deg $(GG_{k-1} - G_k) < \deg G + \deg M$  and  $GG_{k-1} \equiv G_k \pmod{M}$ , we have  $I_k \in S_G$ . We see that

$$\sum_{k=1}^{\infty} \frac{I_k}{G^k} = \frac{1}{M} \left\{ \sum_{k=1}^{\infty} \frac{G_{k-1}}{G^{k-1}} - \sum_{k=1}^{\infty} \frac{G_k}{G^k} \right\} = \frac{1}{M} = \sum_{k=1}^{\infty} \frac{H_k}{G^k}.$$

By the uniqueness of the digit expansion, we have  $I_k = H_k$  for  $k \ge 1$ .

(2) By assertion (1), the sequence  $\{H_k\}_{k\geq 1}$  is purely periodic, and  $l \leq g$ . On the other hand, we have

$$\frac{1}{M} = \sum_{k=1}^{l} \frac{H_k}{G^k} + \frac{1}{G^l} \cdot \frac{1}{M}$$

Multiply both sides by  $G^l M$  and consider modulo M. Then  $G^l \equiv 1 \pmod{M}$ . We thus get  $g \leq l$ .

**Remark 2.1.** Equality (2) plays an important role in the proof of our main results. For the number field version of this equality, see [2], [7], [8].

**Corollary 2.1.** Assume the notation in Proposition 2.2. Let  $\alpha \in \mathbb{F}_q^{\times}$  of order t. If  $gcd(M, G(\alpha G - 1)) = 1$  and  $t \mid g$ , then

$$\sum_{k=1}^{g} \alpha^k H_k = 0.$$

Proof. Put

$$U = \sum_{k=1}^{g} \alpha^{k-1} G_{k-1} \left( = \sum_{k=1}^{g} \alpha^{k} G_{k} \right).$$

By  $gcd(M, G(\alpha G - 1)) = 1$  and  $t \mid g$ , we have

$$U \equiv \sum_{k=1}^{g} \alpha^{k-1} G^{k-1} = \frac{(\alpha G)^g - 1}{\alpha G - 1} \equiv 0 \pmod{M}.$$

This gives U = 0 since deg U < deg M. Therefore, by (2), we obtain

$$\sum_{k=1}^{g} \alpha^k H_k = \frac{(\alpha G - 1)U}{M} = 0.$$

**Remark 2.2.** Considering the case  $\alpha = 1$  in Corollary 2.1, we obtain

$$\sum_{k=1}^{g} H_k = 0$$

This result was discovered by Rudnick [10].

#### 2.2 Cyclotomic function fields and divisor class numbers

In this subsection, we review basic facts about cyclotomic function fields and divisor class numbers. For details, see [1, 4, 9].

Let  $K^{ac}$  be an algebraic closure of K. We define the action

$$I * x = I(\varphi + \mu)(x) \quad (x \in K^{ac}, \ I \in \mathbb{A}),$$

where  $\varphi$  and  $\mu$  are the  $\mathbb{F}_q$ -linear maps defined by

$$\varphi: K^{ac} \to K^{ac} \quad x \mapsto x^q, \qquad \mu: K^{ac} \to K^{ac} \quad x \mapsto Tx.$$

By the above action,  $K^{ac}$  becomes an A-module. Suppose that  $P \in \mathbb{A}$  is monic irreducible of degree d. Put

$$\Lambda_P = \{ x \in K^{ac} \mid P * x = 0 \}.$$

This is a cyclic A-submodule of  $K^{ac}$ . Fix a generator  $\lambda_P$  of  $\Lambda_P$ . Then we have the following isomorphism of A-modules

$$\mathcal{R}_P \longrightarrow \Lambda_P \quad (\overline{I} \to I * \lambda_P),$$

where  $\mathcal{R}_P = \mathbb{A}/P\mathbb{A}$ , and  $\overline{I} = I \mod P \in \mathcal{R}_P$ . Let  $K_P = K(\Lambda_P)$ . Then  $K_P/K$  is a Galois extension, and we have the following isomorphism

$$\mathcal{R}_P^{\times} \to \operatorname{Gal}(K_P/K) \ (\overline{I} \mapsto \sigma_I),$$

where  $\operatorname{Gal}(K_P/K)$  is the Galois group of  $K_P/K$ , and  $\sigma_I$  is the isomorphism given by  $\sigma_I(\lambda_P) = I * \lambda_P$ . Since  $\mathcal{R}_P$  is a finite field of order  $q^d$ , the extension  $K_P/K$  is cyclic of degree  $q^d - 1$ . The field  $K_P$  is called the *P*-th cyclotomic function field.

Let L be an intermediate field of  $K_P/K$ . We denote by  $H_L$  the subgroup of  $\mathcal{R}_P^{\times}$  corresponding to L. The field  $L^+ = L \cap K_{\infty}$  is called the maximal real subfield of L. By Theorem 12.14 in [9], we have  $H_{L^+} = H_L \mathbb{F}_q^{\times}$ .

The next lemma is well-known to the experts. But we give a proof for the reader's convenience.

**Lemma 2.1.** Suppose that q is odd, and  $P \in \mathbb{A}$  is monic irreducible of degree d. Then  $L = K\left(\sqrt{(-1)^d P}\right)$  is contained in  $K_P$ . Furthermore,

$$L^{+} = \begin{cases} L & \text{if } d \text{ is even,} \\ K & \text{if } d \text{ is odd.} \end{cases}$$
(3)

*Proof.* By the discussion of the proof of Lemma 16.13 in [9], we can take an element  $\pi \in K_P$  satisfying with  $\pi^{q-1} = (-1)^r P$ , where  $r = (q^d - 1)/(q - 1)$ . Hence L is contained in  $K_P$  since  $r \equiv d \pmod{2}$  and q is odd.

We note that  $K_P/K$  is cyclic and  $[K_P^+:K] = r \equiv d \pmod{2}$ . Therefore,

$$L = L^+ \iff L \subseteq K_P^+ \iff d \equiv 0 \pmod{2}.$$

This implies (3).

Let  $X_P$  be the group of Dirichlet characters modulo P. For an intermediate field L of  $K_P/K$ , we set

$$X_L = \{ \chi \in X_P \mid \chi(\overline{I}) = 1 \ (\overline{I} \in H_L) \},\$$
  
$$X_L^+ = \{ \chi \in X_P \mid \chi(\overline{I}) = 1 \ (\overline{I} \in H_{L^+}) \},\$$

and  $X_L^- = X_L \setminus X_L^+$ . We denote by  $h_L$  (resp.  $h_L^+$ ) the divisor class number of L (resp.  $L^+$ ). By a similar argument as in the proof of Theorem 2 in [1], we have

$$h_L = \prod_{\substack{\chi \in X_L^+ \\ \chi \neq 1_P}} \left\{ -\sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) \deg I \right\} \cdot \prod_{\chi \in X_L^-} \left\{ \sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) \right\}, \quad (4)$$

$$h_L^+ = \prod_{\substack{\chi \in X_L^+ \\ \chi \neq 1_P}} \left\{ -\sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) \deg I \right\}.$$
 (5)

Here,  $\mathbb{M}$  is the set of all monic polynomials of  $\mathbb{A}$ , and  $1_P$  is the trivial character of  $X_P$ . The rational number  $h_L^- = h_L/h_L^+$  is called the relative divisor class number of L. By (4) and (5), we have

$$h_L^- = \prod_{\chi \in X_L^-} \left\{ \sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) \right\}.$$
 (6)

The right-hand side of (6) is an algebraic integer. Hence the rational number  $h_L^-$  is an integer.

### 3 Main results

Suppose that  $P \in \mathbb{A}$  is monic irreducible and  $G \in \mathbb{A}$  is a primitive root modulo P. Let  $d = \deg P$ ,  $e = \deg G$ ,  $r = (q^d - 1)/(q - 1)$ . Let  $1/P = \sum_{k=1}^{\infty} H_k/G^k$  be the digit expansion of 1/P in base G. Choose  $G_k \in \mathbb{A}$   $(0 \le k \le q^d - 1)$  with  $\deg G_k < d$  and  $G_k \equiv G^k \pmod{P}$ .

**Lemma 3.1.** For  $0 \le s \le d-1$ , we have

$$\left\{ \begin{bmatrix} \overline{I} \end{bmatrix} \mid \begin{array}{c} I \in \mathbb{M}, \\ \deg I = s \end{array} \right\} = \left\{ \begin{bmatrix} \overline{G_k} \end{bmatrix} \mid \begin{array}{c} 0 \le k \le r - 1, \\ \deg G_k = s \end{array} \right\},$$

where  $[\overline{I}]$  is the equivalent class of  $\overline{I} \in \mathcal{R}_P$  in  $\mathcal{R}_P^{\times}/\mathbb{F}_q^{\times}$ .

*Proof.* Since G is a primitive root of modulo P, we have

$$\overline{G_k} = (\overline{G})^k \in \mathbb{F}_q^{\times} \iff r \mid k.$$

Hence {  $\overline{G_k} \mid 0 \leq k \leq r-1$  } is a complete residue system of  $\mathcal{R}_P^{\times}/\mathbb{F}_q^{\times}$ . This implies Lemma 3.1.

We first consider the plus-part of divisor class numbers. We put

$$F^{(+)}(u) = \sum_{k=1}^{r} (\deg H_k) u^{k-1}$$

In the above definition, we understand  $\deg H_k$  as 0 if  $H_k = 0$ . Let

$$X_P^+ = \{ \chi \in X_P \mid \chi(a) = 1 \text{ for all } a \in \mathbb{F}_q^{\times} \}.$$

Then we have

**Proposition 3.1.** For  $\chi \in X_P^+$ , we have

$$F^{(+)}(\zeta) = \sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} (\deg I - (d-e))\chi(I),$$

where  $\zeta = \chi(G)$ .

*Proof.* By (2), we have  $GG_{k-1} = H_k P + G_k$ . It follows that

$$\deg H_k = \begin{cases} \deg G_{k-1} - (d-e) & \text{if } \deg G_{k-1} \ge d-e, \\ 0 & \text{if } \deg G_{k-1} < d-e. \end{cases}$$

Define Deg :  $\mathcal{R}_P^{\times} \to \mathbb{Z}$  ( $\overline{I} \mapsto \deg r_I$ ), where  $r_I$  is the remainder of I divided by P. It is easy to check that Deg and  $\chi$  are both functions on  $\mathcal{R}_P^{\times}/\mathbb{F}_q^{\times}$ . Therefore, by Lemma 3.1, we have

$$\sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} (\deg I - (d-e))\chi(I)$$
  
= 
$$\sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} (\operatorname{Deg}([\overline{I}]) - (d-e))\chi([\overline{I}])$$
  
= 
$$\sum_{\substack{I \leq k \leq r \\ d-e \leq \deg G_{k-1}}} (\operatorname{Deg}([\overline{G_{k-1}}]) - (d-e))\chi([\overline{G_{k-1}}])$$
  
= 
$$\sum_{k=1}^{r} (\deg H_k)\zeta^{k-1}.$$

We thus get Proposition 3.1.

Corollary 3.1.

$$\sum_{k=1}^{r} \deg H_k = \begin{cases} q^{d-e} s_1(e) & \text{if } e < d, \\ (e-d)s_0(d) + s_1(d) & \text{if } e \ge d, \end{cases}$$

where  $s_0(j) = \sum_{i=0}^{j-1} q^i$  and  $s_1(j) = \sum_{i=0}^{j-1} i q^i$ .

*Proof.* We consider the case  $\chi = 1_P$  in Proposition 3.1. Assume that e < d. Since the number of monic polynomials of degree i in  $\mathbb{A}$  is equal to  $q^i$ , we have

$$\sum_{k=1}^{r} \deg H_k = \sum_{i=0}^{e-1} \sum_{\substack{I \in \mathbb{M} \\ \deg I = d-e+i}} (\deg I - (d-e))$$
$$= \sum_{i=0}^{e-1} iq^{d-e+i}$$
$$= q^{d-e} s_1(e).$$

We next assume that  $e \ge d$ . Then

$$\sum_{k=1}^{r} \deg H_k = \sum_{i=0}^{d-1} \sum_{I \in \mathbb{M} \atop \deg I=i} (\deg I - (d-e))$$
$$= \sum_{i=0}^{d-1} (i + (e-d))q^i$$
$$= s_1(d) + (e-d)s_0(d).$$

**Corollary 3.2.** If  $\chi \in X_P^+ \setminus \{1_P\}$  and  $e \ge d$ , then

$$F^{(+)}(\zeta) = \sum_{I \in \mathbb{M} \atop \deg I < d} \chi(I) \deg I,$$

where  $\zeta = \chi(G)$ .

*Proof.* Since  $\chi(\mathbb{F}_q^{\times}) = \{1\}$ , we have

$$\sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) = \frac{1}{q-1} \sum_{I \in \mathbb{A} \\ \deg I < d} \chi(I) = 0.$$

Therefore, by Proposition 3.1, we obtain

$$F^{(+)}(\zeta) = \sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I)(\deg I - (d - e)) = \sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) \deg I.$$

**Theorem 3.1.** Let L be an intermediate field of  $K_P/K$ . Assume that  $m = [L^+:K] > 1$  and  $e \ge d$ . Then,

$$(-1)^{m-1}h_L^+ = \prod_{\substack{\zeta \in U_L^+ \\ \zeta \neq 1}} F^{(+)}(\zeta),$$

where  $U_L^+$  is the set of all m-th roots of unity.

*Proof.* Since  $X_L^+$  is a cyclic group of order m, the map  $X_L^+ \longrightarrow U_L^+$   $(\chi \mapsto \chi(G))$  is a group isomorphism. By (6) and Corollary 3.2, we have

$$(-1)^{m-1}h_L^+ = \prod_{\substack{\chi \in X_L^+ \\ \chi \neq 1_P}} \left\{ \sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I) \deg I \right\}$$
$$= \prod_{\substack{\zeta \in U_L^+ \\ \zeta \neq 1}} F^{(+)}(\zeta).$$

Now we prove the even case of Theorem 1.2.

*Proof.* By Lemma 2.1, the field L is contained in  $K_P$  and  $L = L^+$ . Therefore, by Theorem 3.1, we obtain

$$h_L = h_L^+ = -F^{(+)}(-1) = \sum_{k=1}^r (-1)^k \deg H_k.$$

**Example 3.1.** Suppose that q = 3,  $P = T^2 + 1$ ,  $G = T^2 + T + 2$ , and  $L = K(\sqrt{P})$ . Then P is irreducible in  $\mathbb{F}_3[T]$ , and G is a primitive root modulo P. Let  $1/P = \sum_{k=1}^{\infty} H_k/G^k$  be the digit expansion of 1/P in base G. Then  $r = (q^2 - 1)/(q - 1) = 4$ , and

$$H_1 = 1$$
,  $H_2 = T + 2$ ,  $H_3 = 2T + 2$ ,  $H_4 = 2T$ .

By Theorem 1.2, we have

$$h_L = (\deg H_2 + \deg H_4) - (\deg H_1 + \deg H_3) = 1.$$

We next calculate the divisor class number of  $K_P^+$ . We see that  $[K_P^+ : K] = 4$ , and

$$F^{(+)}(u) = \sum_{k=1}^{4} (\deg H_k) u^{k-1} = u + u^2 + u^3, \quad U_{K_P}^+ = \{\zeta \in \mathbb{C} \mid \zeta^4 = 1\}.$$

By Theorem 3.1, we have

$$h_{K_P}^+ = (-1)^{[K_P^+:K]-1} \prod_{\substack{\zeta \in U_{K_P}^+ \\ \zeta \neq 1}} F^+(\zeta) = 1.$$

**Example 3.2.** Suppose that q = 2,  $P = T^3 + T + 1$ ,  $G = T^3$ . Then P is irreducible in  $\mathbb{F}_2[T]$ , and G is a primitive root modulo P. Let  $1/P = \sum_{k=1}^{\infty} H_k/G^k$  be the digit expansion of 1/P in base G. Then  $r = (q^3 - 1)/(q - 1) = 7$ , and

$$H_1 = 1, \quad H_2 = T + 1, \quad H_3 = T^2, \quad H_4 = T^2 + 1,$$
  
 $H_5 = T^2 + T, \quad H_6 = T, \quad H_7 = T^2 + T + 1.$ 

We see that  $[K_P^+:K] = 7$ , and

$$F^{(+)}(u) = \sum_{k=1}^{7} (\deg H_k) u^{k-1} = u + 2u^2 + 2u^3 + 2u^4 + u^5 + 2u^6,$$
$$U^+_{K_P} = \{\zeta \in \mathbb{C} \mid \zeta^7 = 1\}.$$

By Theorem 3.1, we have

$$h_{K_P}^+ = (-1)^{[K_P^+:K]-1} \prod_{\substack{\zeta \in U_{K_P}^+ \\ \zeta \neq 1}} F^+(\zeta) = 71.$$

We next consider the minus-part of divisor class numbers. Denote by  $\widehat{\mathbb{F}_q^{\times}}$  the character group of  $\mathbb{F}_q^{\times}$ . Recall that  $\delta(I)$  is the leading coefficient of  $I \in \mathbb{A}$  if  $I \neq 0$ , and  $\delta(0) = 0$ . Suppose that  $\lambda \in \widehat{\mathbb{F}_q^{\times}}$ . We define

$$F^{(\lambda)}(u) = \lambda(\delta(G)) \sum_{k=1}^{r} \bar{\lambda}(\delta(H_k)) u^{k-1},$$

where  $\bar{\lambda}$  is the complex conjugate of  $\lambda$ . In the above definition, we understand  $\bar{\lambda}(\delta(H_k))$  as 0 if  $\delta(H_k) = 0$ . We set

$$X_P^{(\lambda)} = \{ \chi \in X_P \mid \chi|_{\mathbb{F}_q^{\times}} = \lambda \},\$$

where  $\chi|_{\mathbb{F}_q^{\times}}$  is the restriction to  $\mathbb{F}_q^{\times}$  of  $\chi$ .

**Proposition 3.2.** For  $\chi \in X_P^{(\lambda)}$ , we have

$$F^{(\lambda)}(\zeta) = \sum_{I \in \mathbb{M} \atop d-e \leq \deg I < d} \chi(I),$$

where  $\zeta = \chi(G)$ .

*Proof.* Define  $H_{\chi} : \mathcal{R}_P^{\times} \to \mathbb{C}$  by

$$H_{\chi}(\overline{I}) = \overline{\lambda}(\delta(r_I))\chi(r_I),$$

where  $r_I$  is the remainder of I divided by P. It is easy to check that  $H_{\lambda}$  is a function on  $\mathcal{R}_P^{\times}/\mathbb{F}_q^{\times}$ . By Lemma 3.1, we have

$$\sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} \chi(I) = \sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} \overline{\lambda}(\delta(r_I))\chi(r_I)$$
$$= \sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} H_{\chi}([\overline{I}])$$
$$= \sum_{\substack{I \leq k \leq r \\ d-e \leq \deg G_{k-1}}} H_{\chi}([\overline{G_{k-1}}])$$
$$= \sum_{\substack{1 \leq k \leq r \\ d-e \leq \deg G_{k-1}}} \overline{\lambda}(\delta(G_{k-1}))\zeta^{k-1}$$

By  $GG_{k-1} = H_k P + G_k$ , we have

$$\delta(H_k) = \begin{cases} \delta(G)\delta(G_{k-1}) & \text{if } \deg G_{k-1} \ge d-e, \\ 0 & \text{if } \deg G_{k-1} < d-e. \end{cases}$$

Therefore,

$$\sum_{\substack{I \in \mathbb{M} \\ d-e \leq \deg I < d}} \chi(I) = \lambda(\delta(G)) \sum_{k=1}^{r} \overline{\lambda}(\delta(H_k)) \zeta^{k-1} = F^{(\lambda)}(\zeta).$$

•

**Corollary 3.3.** Assume that  $\chi \in X_P^{(\lambda)}$  and  $e \ge d$ . Then,

$$F^{(\lambda)}(\zeta) = \sum_{\substack{I \in \mathbb{M} \\ \deg I < d}} \chi(I).$$

Proof. This follows immediately from Proposition 3.2.

Let L be an intermediate field of  $K_P/K$ . Put

$$l = [L:K], m = [L^+:K], n = [L:L^+].$$

Since  $X_P$  is cyclic, we have

$$X_L = \{ \chi \in X_P \mid \chi^l = 1_P \}, \quad X_L^+ = \{ \chi \in X_P \mid \chi^m = 1_P \}.$$

We set

$$Y_L = \{ \lambda \in \widehat{\mathbb{F}_q^{\times}} \mid \lambda^n = \lambda_0 \}, \quad U_n = \{ \zeta \in \mathbb{C} \mid \zeta^n = 1 \},$$

where  $\lambda_0$  is the trivial character of  $\mathbb{F}_q^{\times}$ . Then the maps

$$\varphi: X_L/X_L^+ \longrightarrow Y_L \ (\chi X_L^+ \mapsto \chi|_{\mathbb{F}_q^{\times}}),$$
$$\psi: X_L/X_L^+ \longrightarrow U_n \ (\chi X_L^+ \mapsto \chi(G)^m)$$

are both group isomorphisms. For  $\lambda \in Y_L$ , we put  $\alpha_{\lambda} = \psi \circ \varphi^{-1}(\lambda)$ .

**Theorem 3.2.** Let L be an intermediate field of  $K_P/K$ . Assume that  $L \neq L^+$ and  $e \geq d$ . Then

$$h_L^- = \prod_{\substack{\lambda \in Y_L \\ \lambda \neq \lambda_0}} \prod_{\zeta \in U_L^{(\lambda)}} F^{(\lambda)}(\zeta),$$

where  $U_L^{(\lambda)} = \{ \zeta \in \mathbb{C} \mid \zeta^m = \alpha_\lambda \}.$ 

*Proof.* For  $\lambda \in Y_L$ , we set

$$X_L^{(\lambda)} = \{ \chi \in X_L \mid \chi|_{\mathbb{F}_q^{\times}} = \lambda \}.$$

Then,

$$X_L^- = X_L \setminus X_L^+ = \bigcup_{\substack{\lambda \in Y_L \\ \lambda \neq \lambda_0}} X_L^{(\lambda)}.$$

By the definition of  $\alpha_{\lambda}$ , we have

$$X_L^{(\lambda)} = \{ \chi \in X_L \mid \chi(G)^m = \alpha_\lambda \}.$$

This implies  $U_L^{(\lambda)} = \{\chi(G) \mid \chi \in X_L^{(\lambda)}\}$ . Therefore, by (6) and Corollary 3.3, we have

$$\begin{split} h_L^- &= \prod_{\chi \in X_L^-} \left\{ \sum_{I \in \mathbb{M} \atop \deg I < d} \chi(I) \right\} \\ &= \prod_{\substack{\lambda \in Y_L \\ \lambda \neq \lambda_0}} \prod_{\chi \in X_L^{(\lambda)}} \left\{ \sum_{I \in \mathbb{M} \atop \deg I < d} \chi(I) \right\} \\ &= \prod_{\substack{\lambda \in Y_L \\ \lambda \neq \lambda_0}} \prod_{\zeta \in U_L^{(\lambda)}} F^{(\lambda)}(\zeta). \end{split}$$

We prove the odd case of Theorem 1.2.

*Proof.* By Lemma 2.1, the field L is contained in  $K_P$  and  $L^+ = K$ . Suppose that  $\lambda$  is the quadratic character of  $\mathbb{F}_q^{\times}$ . Then  $Y_L$  is generated by  $\lambda$ . Note that  $\alpha_{\lambda} = -1$  and

$$U_L^{(\lambda)} = \{-1\}, \quad \lambda(\delta(G)) = \eta_G, \quad \overline{\lambda}(\delta(H_k)) = \lambda(\delta(H_k)) = \varepsilon_k.$$

Therefore, by Theorem 3.2, we obtain

$$h_L = h_L^- = F^{(\lambda)}(-1) = (-\eta_G) \sum_{k=1}^r (-1)^k \varepsilon_k.$$

**Example 3.3.** Suppose that q = 3,  $P = T^3 + 2T + 2$ ,  $G = T^3 + T + 2$ , and  $L = K(\sqrt{-P})$ . Then P is irreducible in  $\mathbb{F}_3[T]$ , and G is a primitive root modulo P. Let  $1/P = \sum_{k=1}^{\infty} H_k/G^k$  be the digit expansion of 1/P in base G. Then  $r = (q^3 - 1)/(q - 1) = 13$ ,  $\eta_G = 1$ , and

$$\begin{array}{ll} H_1=1, & H_2=2T, & H_3=T^2+2, & H_4=2T+2, & H_5=T^2+T+2, \\ H_6=2T^2+2T, & H_7=T^2+2T, & H_8=T^2+T+1, & H_9=2T^2, \\ H_{10}=2T+1, & H_{11}=T^2+2T+2, & H_{12}=T^2+2T+1, & H_{13}=T^2+1. \end{array}$$

Let  $\varepsilon_k = 1$  or -1 (k = 1, 2, ..., 13) according to  $\delta(H_k) = 1$  or  $\delta(H_k) = 2$ . By Theorem 1.2, we have

$$h_L = (-\eta_G) \sum_{k=1}^{13} (-1)^k \varepsilon_k = 7.$$

We next calculate the relative class number of  $K_P$ . Let  $\lambda$  be the quadratic character of  $\mathbb{F}_3^{\times}$ . Then  $[K_P^+:K] = 13$ ,  $\alpha_{\lambda} = -1$ , and

$$F^{(\lambda)}(u) = \lambda(\delta(G)) \sum_{k=1}^{13} \overline{\lambda}(\delta(H_k)) u^{k-1}$$
  
= 1 - u + u^2 - u^3 + u^4 - u^5 + u^6 + u^7 - u^8 - u^9 + u^{10} + u^{11} + u^{12},  
$$U_{K_P}^{(\lambda)} = \{\zeta \in \mathbb{C} \mid \zeta^{13} = -1\}.$$

Therefore, by Theorem 3.2, we have

$$h_{K_P}^- = \prod_{\zeta \in U_{K_P}^{(\lambda)}} F^{(\lambda)}(\zeta) = 774144 = 2^{12} \cdot 3^3 \cdot 7.$$

## References

- S. Galovich, M. Rosen, The Class Number of Cyclotomic Function Fields, J. Number Theory 13 (1981), 363-375.
- [2] K. Girstmair, A "popular" class number formula, Amer. Math. Monthly 101 (1994), 997-1001.
- [3] K. Girstmair, The digits of 1/p in connection with class number factors, Acta Arith. 67 (1994), 381-386.
- [4] D. R. Hayes, Explicit class field theory for rational function fields, Trans. Amer. Math. Soc. 189 (1974), 77–91.
- [5] M. Hirabayashi, Generalizations of Girstmair's formulas, Abh. Math. Sem. Univ. Hamburg 75 (2005), 83-95.
- [6] C. Kalyan, K. Krishnarjun, On some symmetries of the base n expansion of 1/m: the class number connection, Pac. J. Math. **319**, No. 1, 39-53 (2022).
- [7] Y. Mizuno, A certain character twisted average value of the digits of rational numbers and the class numbers of imaginary quadratic fields, Acta Arith. 208, No. 3, 215-233 (2023).
- [8] R. Murty, R. Thangadurai, The class number of  $\mathbb{Q}(\sqrt{-p})$  and digits of 1/p, Proc. Amer. Math. Soc. **139** (2011), 1277-1289.
- [9] M. Rosen, Number Theory in Function Fields, Springer-Verlag, Berlin, 2002.
- [10] Z. Rudnick, On the sum of digits of 1/M in  $\mathbb{F}_q[x]$ , arXiv :2108.10142 (2021).

Daisuke Shiomi Department of Science Faculty of Science, Yamagata University Kojirakawa-machi 1-4-12 Yamagata 990-8560, Japan shiomi@sci.kj.yamagata-u.ac.jp