

Classification of generalized Alexander quandles

Akihiro Higashitani, Seiichi Kamada, Jin Kosaka and Hirotake Kurihara

Abstract

The aim of this paper is to provide a new characterization of isomorphism classes of generalized Alexander quandles in terms of the underlying groups and their automorphisms. This extends the previous result [4, Theorem 1.4]. Additionally, we compute the number of generalized Alexander quandles up to quandle isomorphism arising from groups up to order 127 and their group automorphisms.

1 Introduction

1.1 Motivation

Quandles [6, 7] are algebraic systems equipped with three axioms that correspond to Reidemeister moves in knot theory. According to Joyce, any quandle can be expressed by a disjoint union of homogeneous quandles ([6, Theorem 7.2]). Here, we say that a quandle Q is *homogeneous* if the automorphism group of Q acts transitively on Q . Furthermore, it has been shown that any homogeneous quandle can be constructed by a so-called *quandle triplet* (G, H, ψ) ([6, Theorem 7.1]).

The primary focus of this paper is a special kind of homogeneous quandles called *generalized Alexander quandles*, which are quandles arising from groups. Alexander quandles, which are generalized Alexander quandles arising from abelian groups, have been extensively studied in the context of knot theory. Finite Alexander quandles have been classified up to quandle isomorphism by Nelson [8].

Recently, the first named and fourth named authors established a kind of characterization theorem of finite generalized Alexander quandles of finite groups under certain assumptions ($P1$) and ($P2$) (detailed in Section 3.) The main objective of this paper is to extend the result of [4].

1.2 Main results

The following is the main theorem of this paper:

Theorem 1.1 (Main Result). *Let G and G' be groups with their identity elements e and e' , respectively. Let $\psi \in \text{Aut}(G)$, $\psi' \in \text{Aut}(G')$ and let*

$$Q = (Q(G, \psi), *), Q' = (Q(G', \psi'), *'), P = P(Q) \text{ and } P' = P(Q').$$

2020 Mathematics subject classification: Primary: 20N02, Secondary: 20B05, 53C35.
Key words and phrases: generalized Alexander quandles, finite groups, automorphisms.

Then the following conditions (1) and (2) are equivalent:

- (1) Q and Q' are isomorphic as quandles.
- (2) There exists a group isomorphism $h : P \rightarrow P'$ such that the following two conditions (A) and (B) are satisfied:
 - (A) $h \circ \psi|_P = \psi'|_{P'} \circ h$;
 - (B) there exist representatives $A \subset G$ and $A' \subset G'$ of G/P and G'/P' , respectively, and a bijection $k : A \rightarrow A'$ between A and A' such that $h(e * a) = e' *' k(a)$ holds for any $a \in A$.

The notation used in this paper will be explained in Sections 2 and 3. The proof of Theorem 1.1 will be given in Section 4.

Remark 1.1. The statement of Theorem 1.1 does not assume the finiteness of groups.

By [4], generalized Alexander quandles of finite groups up to order 15 are completely classified. An observation of those of order 16 is also made, but the main result of [4] is not applicable for two cases. By Theorem 1.1, we can completely classify not only all generalized Alexander quandles of finite groups of order 16, but also those with orders at most 127 (with some additional exceptional treatments). See Section 5.

Acknowledgements

The authors are grateful to S. Nelson and H. Tamaru for useful comments on an earlier draft of the paper. The first named author is partially supported by JSPS KAKENHI Grant Number JP20K03513. The second named author is supported by JP19H01788 and JP23H05437. The fourth named author is supported by JP20K03623.

2 Preliminaries

In this section, we prepare notation and recall notions on quandles and groups used throughout the paper.

2.1 Definition of quandles

First, let us recall a definition of quandle. A non-empty set Q equipped with a binary operation $*$ satisfying the following three axioms is called a *quandle*:

- (Q1) $x * x = x$ for any $x \in Q$;
- (Q2) for any $x \in Q$, the map defined by $s_x : Q \ni y \mapsto y * x \in Q$ is bijective;
- (Q3) for any $x, y, z \in Q$, we have $(x * y) * z = (x * z) * (y * z)$.

The map appearing in (Q2) is called a *point symmetry*.

2.2 Homogeneous quandles and generalized Alexander quandles

Let G be a group and let ψ be an automorphism of G . Moreover, we take a subgroup H of G satisfying $\psi(h) = h$ for any $h \in H$. We define a binary operation $*$ on the set of left cosets G/H defined by $xH * yH := y\psi(y^{-1}x)H$. Then this binary operation becomes well-defined and $(G/H, *)$ gives us a quandle. The triplet (G, H, ψ) providing a new quandle $(G/H, *)$ is called a *quandle triplet*, and we denote this quandle by $Q(G, H, \psi)$. We notice that for any group G with its identity element e and any automorphism ψ of G , the triplet $(G, \{e\}, \psi)$ always becomes a quandle triplet.

Definition 2.1. Let G be a group and let ψ be its automorphism. Considering a binary operation $*$ on G defined by $x * y := y\psi(y^{-1}x)$, we obtain a quandle $(G, *)$, denoted by $Q(G, \psi)$ or $(Q(G, \psi), *)$. We call this quandle $Q(G, \psi)$ a *generalized Alexander quandle* (or a *principal quandle*, see [2]).

In the case where G is abelian, we call $Q(G, \psi)$ an *Alexander quandle*, which has been studied especially in the context of knot theory. Generalized Alexander quandles can be regarded as a generalization of it.

2.3 Notation and terminologies on quandles and groups

In what follows, we use the notation $(Q, *)$ for quandles if we would like to emphasize the equipped binary operation; otherwise, we simply write Q .

Let $(Q, *)$ and $(Q', *')$ be quandles. A map $f : Q \rightarrow Q'$ is said to be a *quandle homomorphism* if $f(x*y) = f(x)*'f(y)$ holds for any $x, y \in Q$. In addition to this, if f is bijective, then we call f a *quandle isomorphism*. If there exists a quandle isomorphism between quandles Q and Q' , then we say that Q and Q' are isomorphic as quandles, which we denote by $Q \cong Q'$.

- Let $\text{Aut}(Q)$ be the group consisting of all quandle automorphisms on Q , which we call the *automorphism group* of Q . Note that the point symmetry s_x for each $x \in Q$ belongs to $\text{Aut}(Q)$.
- Let $\text{Inn}(Q)$ be the subgroup of $\text{Aut}(Q)$ generated by $\{s_x : x \in Q\}$, which we call the *inner automorphism group* of Q . More explicitly, we have $\text{Inn}(Q) = \{s_{x_n}^{\pm 1} \circ s_{x_{n-1}}^{\pm 1} \circ \cdots \circ s_{x_1}^{\pm 1} : n \in \mathbb{Z}_{>0}, x_1, \dots, x_n \in Q\}$. In the case where Q is finite, we have $\text{Inn}(Q) = \{\text{id}_Q, s_{x_n} \circ s_{x_{n-1}} \circ \cdots \circ s_{x_1} : n \in \mathbb{Z}_{>0}, x_1, \dots, x_n \in Q\}$.

We say that a quandle Q is *homogeneous* if the automorphism group $\text{Aut}(Q)$ of Q acts on Q transitively. According to [6, Section 7], the following one-to-one correspondence is known:

$$\{\text{homogeneous quandles}\} / \cong \xleftrightarrow{1:1} \{Q(G, H, \psi) : (G, H, \psi) \text{ is a quandle triplet}\} / \cong,$$

where $/ \cong$ stands for that we consider this correspondence up to quandle isomorphism.

Remark 2.1. Let Q and Q' be homogeneous quandles which are isomorphic. Then, for any $x \in Q$ and $x' \in Q'$, we can always find a quandle isomorphism $f : Q \rightarrow Q'$ with $f(x) = x'$. In fact, even if $f(x) = x'' \neq x'$, since $\text{Aut}(Q')$ acts on Q' transitively, we can

find $g \in \text{Aut}(Q')$ with $g(x'') = x'$, so we may take an isomorphism $g \circ f$ between Q and Q' instead of f .

Therefore, for any generalized Alexander quandles $Q(G, \psi)$ and $Q(G', \psi')$ which are isomorphic, we may assume the existence of a quandle isomorphism $f : Q(G, \psi) \rightarrow Q(G', \psi')$ with $f(e) = e'$, where G and G' are groups with their identity elements e and e' , respectively.

We also collect some terminologies and notation concerning with groups. Let G be a group.

- We denote the automorphism group of G by $\text{Aut}(G)$. Remark that this notation is the same as that for quandles.
- Given $\psi \in \text{Aut}(G)$, let $\text{Fix}(\psi, G) = \{g \in G : \psi(g) = g\}$.
- Let N, H be groups and let $\sigma : H \rightarrow \text{Aut}(N)$, $h \mapsto \sigma_h$ be a group homomorphism. The (outer) *semidirect product* $N \rtimes_{\sigma} H$ of N and H with respect to σ is a group which is the direct product $N \times H$ as a set equipped with the operation defined by $(n_1, h_1)(n_2, h_2) = (n_1\sigma_{h_1}(n_2), h_1h_2)$.

2.4 Problem

Given a group G , let $\mathcal{Q}(G)$ denote the set of all generalized Alexander quandles $Q(G, \psi)$ with $\psi \in \text{Aut}(G)$ up to quandle isomorphism. Namely,

$$\mathcal{Q}(G) = \{Q(G, \psi) : \psi \in \text{Aut}(G)\} / \cong .$$

Problem 2.1 ([5, Problem 3.5]). *Determine $\mathcal{Q}(G)$ for a given group G .*

This problem has some certain partial answers in terms of the automorphism group of groups. For example, [8, Corollary 2.2] claims that $\mathcal{Q}(G)$ can be characterized in terms of just numerical information in the case where G is finite and abelian. Moreover, it is proved in [4, Theorem 1.2] that if G is a finite simple group, then $\mathcal{Q}(G)$ one-to-one corresponds to the conjugacy classes of $\text{Aut}(G)$. Using this, we can also get this one-to-one correspondence in the case of symmetric groups ([4, Corollary 1.3]), though it is not simple. Theorem 1.1 asserts a new kind of complete solution of Problem 2.1.

3 Previous results on generalized Alexander quandles

In this section, we recall some results on generalized Alexander quandles, especially from [4].

Let G be a group, ψ an automorphism of G , and $Q = Q(G, \psi)$ the generalized Alexander quandle. We define $P = P(Q)$ as the orbit of the identity element e in Q under the action of $\text{Inn}(Q)$, i.e.,

$$P = P(Q) = \{s_{x_n}^{\pm 1} \circ s_{x_{n-1}}^{\pm 1} \circ \cdots \circ s_{x_1}^{\pm 1}(e) : n \in \mathbb{Z}_{>0}, x_1, \dots, x_n \in Q\}.$$

Then P enjoys a special property. In particular, the following statements hold:

Proposition 3.1 ([4, Propositions 3.2 and 3.3]). *The subset P of G is a subquandle of Q and a normal subgroup of G .*

Theorem 3.1 ([4, Theorem 3.10], [9, Lemma 2.4]). *Let G and G' be groups with their identity elements e and e' , respectively. Let $\psi \in \text{Aut}(G)$, $\psi' \in \text{Aut}(G')$ and let*

$$Q = Q(G, \psi), Q' = Q(G', \psi'), P = P(Q) \text{ and } P' = P(Q').$$

Assume that there is a quandle isomorphism $f : Q \rightarrow Q'$ with $f(e) = e'$. Then the following hold:

- (i) $f \circ \psi = \psi' \circ f$;
- (ii) $f|_P : P \rightarrow P'$ is a group isomorphism;
- (iii) $f|_P : Q(P, \psi|_P) \rightarrow Q(P', \psi'|_{P'})$ is a quandle isomorphism;
- (iv) $f(xP) = f(x)P'$ holds for any $x \in G$.

Remark 3.1. In [4, Theorem 3.10], the finiteness of groups is assumed, but Proposition 3.1 and Theorem 3.1 do not need this assumption. In fact, even in the case of infinite groups, we see that Proposition 3.1 and the statements of (i), (iii) and (iv) of Theorem 3.1 hold by almost the same proofs with those given in [4]. Moreover, the statement (ii) for (non-necessarily finite) groups is nothing but [9, Lemma 2.4].

The following proposition is useful for study of generalized Alexander quandles up to isomorphism. It asserts that a conjugacy class of $\text{Aut}(G)$ determines an element of $Q(G)$.

Proposition 3.2 ([4, Proposition 3.5]). *Let $\psi, \psi' \in \text{Aut}(G)$ and let $Q = Q(G, \psi)$ and $Q' = Q(G, \psi')$. If ψ and ψ' are conjugate in $\text{Aut}(G)$, then $Q \cong Q'$ holds.*

Now, we recall the main result of [4]. For this, we have to introduce more notions. Given a group G and its automorphism ψ , let $Q = Q(G, \psi)$ and $P = P(Q)$. We can consider a generalized Alexander quandle $Q(P, \psi|_P)$. Let P^2 be $P(Q(P, \psi|_P))$. By Proposition 3.1, P^2 is a normal subgroup of P .

We consider the following two properties (P1) and (P2).

(P1) P^2 is a normal subgroup of G .

(P2) $P^2 = \{s_p(e) : p \in P\}$ holds.

These two properties are independent each other and become invariants on generalized Alexander quandles (see [4, Proposition 3.16]).

Theorem 3.2 ([4, Theorem 1.4]). *Let G and G' be finite groups, $\psi \in \text{Aut}(G)$, $\psi' \in \text{Aut}(G')$ and let*

$$Q = (Q(G, \psi), *), Q' = (Q(G', \psi'), *'), P = P(Q) \text{ and } P' = P(Q').$$

Assume that Q and Q' satisfy (P1) and (P2). Then the following conditions (1) and (2) are equivalent:

(1) $Q \cong Q'$.

(2) *The following three conditions are satisfied:*

(i) $|G| = |G'|$;

(ii) $|\text{Fix}(\psi, G)| = |\text{Fix}(\psi', G')|$;

(iii) *there exists a group isomorphism $h : P \rightarrow P'$ such that the following (A) and (B) are satisfied:*

(A) $h \circ \psi|_P = \psi'|_{P'} \circ h$;

(B) *for any $a \in G$, there exists $a' \in G'$ such that $h(e * a) = e' *' a'$.*

Note that the finiteness of the groups G and G' is necessary in this theorem since we have to take the orders of the groups into account, while Theorem 1.1 does require neither the finiteness condition on G and G' nor the conditions (P1) and (P2) on Q and Q' .

4 Proof of Theorem 1.1

This section is devoted to proving Theorem 1.1.

(1) \Rightarrow (2):

Let f be a quandle isomorphism $f : Q \rightarrow Q'$ with $f(e) = e'$. (See Remark 2.1.) Let h be the restriction of f into P , i.e., $h = f|_P : P \rightarrow P'$. Then, by Theorem 3.1 (i) and (ii), we see that the condition (A) is satisfied.

Take an arbitrary representative $A \subset G$ of G/P , let $A' = f(A)$ and let k be the restriction of f into A , i.e., $k = f|_A : A \rightarrow A'$. Then it follows from the bijectivity of f (i.e., k) together with Theorem 3.1 (iv) that A' also becomes a representative of G'/P' . Since $e * a$ is an element of P , we see that $h(e * a) = f(e * a) = f(e) *' f(a) = e' *' k(a)$ holds for any $a \in A$. Hence, (B) is also satisfied.

(2) \Rightarrow (1):

Let $h : P \rightarrow P'$ and $k : A \rightarrow A'$ be a group isomorphism and a bijection between representatives A and A' , respectively, satisfying the conditions (A) and (B). Given any $x \in G$, we can uniquely choose $a_x \in A$ and $p_x \in P$ such that $x = a_x p_x$. Since P is a normal subgroup of G (see Proposition 3.1), we know that $a_x p_x a_x^{-1} \in P$ holds.

Now, we define a map $f : G \rightarrow G'$ by setting

$$f(x) := h(a_x p_x a_x^{-1}) k(a_x) \in G'$$

for any $x = a_x p_x \in G$. The bijectivity of f follows from that of h and k . In what follows, we prove that f is a quandle homomorphism.

Since $h(a_x p_x a_x^{-1}) \in P'$ and P' is a normal subgroup of G' , $k(a_x)^{-1} h(a_x p_x a_x^{-1}) k(a_x)$ belongs to P' . Hence the coset decomposition of $f(x) = a'_{f(x)} p'_{f(x)}$ becomes $a'_{f(x)} = k(a_x)$ and $p'_{f(x)} = k(a_x)^{-1} h(a_x p_x a_x^{-1}) k(a_x)$.

For any $x = a_x p_x$ and $y = a_y p_y$ in G , we have

$$x * y = a_y p_y \psi((a_y p_y)^{-1} a_x p_x) = a_x \cdot a_x^{-1} a_y p_y \psi((a_x^{-1} a_y p_y)^{-1}) \psi(p_x).$$

Since we have $a_x^{-1}a_y p_y \psi((a_x^{-1}a_y p_y)^{-1}) = e * (a_x^{-1}a_y p_y) \in P$ and $\psi(p_x) \in P$, the coset decomposition of $x * y = a_{x*y} p_{x*y}$ becomes

$$a_{x*y} = a_x$$

and

$$p_{x*y} = a_x^{-1}a_y p_y \psi((a_x^{-1}a_y p_y)^{-1})\psi(p_x).$$

Thus, we have the following:

$$\begin{aligned} f(x * y) &= h(a_{x*y} p_{x*y} a_{x*y}^{-1})k(a_{x*y}) \\ &= h(a_x \cdot a_x^{-1}a_y p_y \psi((a_x^{-1}a_y p_y)^{-1})\psi(p_x) \cdot a_x^{-1})k(a_x) \\ &= h(a_y p_y \psi(p_y^{-1}a_y^{-1}) \psi(a_x p_x a_x^{-1}) k(a_x)) \\ &= h(a_y p_y a_y^{-1} \cdot a_y \psi(p_y^{-1}a_y^{-1}) (a_x \psi(p_x^{-1}a_x^{-1}))^{-1}) k(a_x) \\ &= h(a_y p_y a_y^{-1}) h(a_y \psi(p_y^{-1}a_y^{-1})) h(a_x \psi(p_x^{-1}a_x^{-1}))^{-1} k(a_x) \\ &= h(a_y p_y a_y^{-1}) h(a_y \psi(a_y^{-1})) \cdot \psi(a_y p_y^{-1} a_y^{-1}) h(\psi(a_x p_x a_x^{-1}) \cdot \psi(a_x) a_x^{-1}) k(a_x) \\ &= h(a_y p_y a_y^{-1}) h(a_y \psi(a_y^{-1})) h \circ \psi(a_y p_y^{-1} a_y^{-1} a_x p_x a_x^{-1}) h(\psi(a_x) a_x^{-1}) k(a_x). \end{aligned}$$

On the other hand, for the coset decomposition of $f(x) *' f(y) = a'_{f(x)*' f(y)} p'_{f(x)*' f(y)}$, we also have $a'_{f(x)*' f(y)} = a'_{f(x)} = k(a_x)$ and

$$\begin{aligned} p'_{f(x)*' f(y)} &= a'_{f(x)}^{-1} a'_{f(y)} p'_{f(y)} \psi'((a'_{f(x)}^{-1} a'_{f(y)} p'_{f(y)})^{-1}) \psi'(p'_{f(x)}) \\ &= k(a_x)^{-1} k(a_y) \cdot k(a_y)^{-1} h(a_y p_y a_y^{-1}) k(a_y) \psi'((k(a_x)^{-1} k(a_y) \cdot k(a_y)^{-1} h(a_y p_y a_y^{-1}) k(a_y))^{-1}) \\ &\quad \cdot \psi'(k(a_x)^{-1} h(a_x p_x a_x^{-1}) k(a_x)) \\ &= k(a_x)^{-1} h(a_y p_y a_y^{-1}) k(a_y) \psi'(k(a_y)^{-1} h(a_y p_y^{-1} a_y^{-1}) h(a_x p_x a_x^{-1}) k(a_x)). \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} f(x) *' f(y) &= a'_{f(x)*' f(y)} p'_{f(x)*' f(y)} \\ &= h(a_y p_y a_y^{-1}) k(a_y) \psi'(k(a_y)^{-1} h(a_y p_y^{-1} a_y^{-1}) h(a_x p_x a_x^{-1}) k(a_x)) \\ &= h(a_y p_y a_y^{-1}) k(a_y) \psi'(k(a_y)^{-1}) \psi'(h(a_y p_y^{-1} a_y^{-1} a_x p_x a_x^{-1})) \psi'(k(a_x)) \\ &= h(a_y p_y a_y^{-1}) k(a_y) \psi'(k(a_y)^{-1}) h \circ \psi(a_y p_y^{-1} a_y^{-1} a_x p_x a_x^{-1}) \psi'(k(a_x)) \quad (\text{by (A)}). \end{aligned}$$

We consider $k(a_y) \psi'(k(a_y)^{-1})$ and $\psi'(k(a_x))$ in the above equation. By (B), we have

$$k(a_y) \psi'(k(a_y)^{-1}) = e' *' k(a_y) = h(e * a_y) = h(a_y \psi(a_y^{-1}))$$

and

$$\begin{aligned} \psi'(k(a_x)) &= (k(a_x) \psi'(k(a_x)^{-1}))^{-1} k(a_x) = (e' *' k(a_x))^{-1} k(a_x) = h(e * a_x)^{-1} k(a_x) \\ &= h(a_x \psi(a_x^{-1}))^{-1} k(a_x) = h(\psi(a_x) a_x^{-1}) k(a_x). \end{aligned}$$

Thus, we have

$$f(x) *' f(y) = h(a_y p_y a_y^{-1}) h(a_y \psi(a_y^{-1})) h \circ \psi(a_y p_y^{-1} a_y^{-1} a_x p_x a_x^{-1}) h(\psi(a_x) a_x^{-1}) k(a_x) = f(x*y).$$

Therefore, we conclude that f is a quandle isomorphism, as required.

5 Classification of generalized Alexander quandles arising from finite groups with small order

In this section, we apply Theorem 1.1 to obtain the complete classification of generalized Alexander quandles arising from finite groups with small orders.

Given a positive integer n , let

$$\mathcal{Q}_{\text{GAQ}}(n) := \{Q(G, \psi) : G \text{ is a group of order } n, \psi \in \text{Aut}(G)\} / \cong.$$

What we will work in this section is precisely to determine $|\mathcal{Q}_{\text{GAQ}}(n)|$ for $n \leq 127$.

5.1 The case $n \leq 15$

In [4, Section 6], classification of generalized Alexander quandles arising from finite groups G is discussed. For example, the cases where $|G| = p, 2p$ or p^2 are investigated theoretically, where p is a prime. Moreover, it follows from [4, Corollary 1.7] that any generalized Alexander quandles arising from cyclic groups of order $2n$ can be realized as those arising from dihedral groups. By additional case-by-case discussions, we can eventually get the complete classification of $\mathcal{Q}_{\text{GAQ}}(n)$ for $n \leq 15$.

5.2 The case $n = 16$

Most parts of $\mathcal{Q}_{\text{GAQ}}(16)$ are also discussed in [4, Section 6], although two cases $\{Q_1, Q'_1\}$ and $\{Q_2, Q'_2\}$ are remained to be open. In this subsection, we figure these cases out by using Theorem 1.1.

Let Q_8 be the quaternion group $\langle \bar{e}, \mathbf{i}, \mathbf{j}, \mathbf{k} : \bar{e}^2 = e, \mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{ijk} = \bar{e} \rangle$. We denote the elements $e, \mathbf{i}, \mathbf{j}, \mathbf{k}, \bar{e}, \bar{e}\mathbf{i}, \bar{e}\mathbf{j}, \bar{e}\mathbf{k}$ by $1, \mathbf{i}, \mathbf{j}, \mathbf{k}, -1, -\mathbf{i}, -\mathbf{j}, -\mathbf{k}$, respectively, so that $Q_8 = \{1, \mathbf{i}, \mathbf{j}, \mathbf{k}, -1, -\mathbf{i}, -\mathbf{j}, -\mathbf{k}\}$.

Let $\mathbb{Z}/2\mathbb{Z} = \{\bar{0}, \bar{1}\}$ be the group of order 2. Let $\sigma : \mathbb{Z}/2\mathbb{Z} \rightarrow \text{Aut}(Q_8)$ be defined by $i \mapsto \sigma_i$, where $\sigma_{\bar{0}} = \text{id}_{Q_8}$ and $\sigma_{\bar{1}} : (\mathbf{i}, \mathbf{j}, \mathbf{k}) \mapsto (\mathbf{i}, -\mathbf{j}, -\mathbf{k})$. Then σ_i can be written as the inner automorphism $\sigma_i(a) = \mathbf{i}^i a \mathbf{i}^{-i}$ for $a \in Q_8$. Note that $\mathbf{i}^i a \mathbf{i}^{-i}$ is well-defined on $\mathbb{Z}/2\mathbb{Z}$. We define the groups G and G' as follows:

$$G = Q_8 \times \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad G' = Q_8 \rtimes_{\sigma} \mathbb{Z}/2\mathbb{Z}.$$

Let ψ be the group automorphism of Q_8 defined by $\psi(\mathbf{i}) = \mathbf{j}$ and $\psi(\mathbf{j}) = \mathbf{k}$, i.e., ψ cyclically permutes $\mathbf{i}, \mathbf{j}, \mathbf{k}$. Moreover, we take the following maps ψ_1 on G and ψ'_1 on G' , respectively:

$$\psi_1(a, i) = (\psi(a), i) \quad \text{and} \quad \psi'_1(a, i) = (\psi(a\mathbf{i}^i)\mathbf{i}^{-i}, i).$$

Note that we have

$$\psi(a\mathbf{i}^i)\mathbf{i}^{-i} = \begin{cases} \psi(a) & \text{if } i = \bar{0}, \\ \psi(a)\mathbf{k} & \text{if } i = \bar{1}. \end{cases}$$

Obviously, ψ_1 becomes a group automorphism of G of order 3. Now, we check that ψ'_1 becomes a group automorphism of G' of order 3. Note that the injectivity of ψ'_1 is

straightforward. So ψ'_1 is bijective. For checking that ψ'_1 is a group homomorphism, we see that

$$\psi'_1((a_1, i_1) \cdot (a_2, i_2)) = \psi'_1(a_1 \sigma_{i_1}(a_2), i_1 + i_2) = (\psi(a_1 \sigma_{i_1}(a_2) \mathbf{i}^{i_1+i_2}) \mathbf{i}^{-(i_1+i_2)}, i_1 + i_2).$$

On the other hand, we have

$$\begin{aligned} \psi'_1(a_1, i_1) \cdot \psi'_1(a_2, i_2) &= (\psi(a_1 \mathbf{i}^{i_1}) \mathbf{i}^{-i_1}, i_1) \cdot (\psi(a_2 \mathbf{i}^{i_2}) \mathbf{i}^{-i_2}, i_2) \\ &= (\psi(a_1 \mathbf{i}^{i_1}) \mathbf{i}^{-i_1} \sigma_{i_1}(\psi(a_2 \mathbf{i}^{i_2}) \mathbf{i}^{-i_2}), i_1 + i_2) \\ &= (\psi(a_1 \mathbf{i}^{i_1}) \mathbf{i}^{-i_1} \mathbf{i}^{i_1} \psi(a_2 \mathbf{i}^{i_2}) \mathbf{i}^{-i_2} \mathbf{i}^{-i_1}, i_1 + i_2) \\ &= (\psi(a_1 \mathbf{i}^{i_1}) \psi(a_2 \mathbf{i}^{i_2}) \mathbf{i}^{-(i_1+i_2)}, i_1 + i_2) \\ &= (\psi(a_1 \mathbf{i}^{i_1} a_2 \mathbf{i}^{i_2}) \mathbf{i}^{-(i_1+i_2)}, i_1 + i_2) \\ &= (\psi(a_1 \mathbf{i}^{i_1} a_2 \mathbf{i}^{-i_1} \mathbf{i}^{i_1+i_2}) \mathbf{i}^{-(i_1+i_2)}, i_1 + i_2) \\ &= (\psi(a_1 \sigma_{i_1}(a_2) \mathbf{i}^{i_1+i_2}) \mathbf{i}^{-(i_1+i_2)}, i_1 + i_2). \end{aligned}$$

These imply $\psi'_1(a_1, i_1) \cdot \psi'_1(a_2, i_2) = \psi'_1((a_1, i_1) \cdot (a_2, i_2))$. We can also check that the order of ψ'_1 is 3. In fact,

$$\psi_1'^3(a, i) = \psi'_1 \circ \psi'_1(\psi(a \mathbf{i}^i) \mathbf{i}^{-i}, i) = \psi'_1(\psi^2(a \mathbf{i}^i) \mathbf{i}^{-i}, i) = (\psi^3(a \mathbf{i}^i) \mathbf{i}^{-i}, i) = (a \mathbf{i}^i \mathbf{i}^{-i}, i) = (a, i).$$

Let $Q_1 = (Q(G, \psi_1), *)$ and $Q'_1 = (Q(G', \psi'_1), *')$ be the generalized Alexander quandles.

For Q_1 , since $(1, \bar{0}) * (a, i) = (a, i) \psi_1(a^{-1}, i) = (a, i) (\psi(a^{-1}), i) = (a \psi(a^{-1}), \bar{0})$, we have $P(Q_1) \subset Q_8 \times \{\bar{0}\}$. Moreover, since we know $\mathbf{j} \psi(\mathbf{j}^{-1}) = -\mathbf{i}$ and $\mathbf{k} \psi(\mathbf{k}^{-1}) = -\mathbf{j}$ and Q_8 is generated by $-\mathbf{i}$ and $-\mathbf{j}$, we have $P(Q_1) = Q_8 \times \{\bar{0}\}$. In a similar way, we can also obtain $P(Q'_1) = Q_8 \times \{\bar{0}\}$. Note that $P(Q'_1)$ is not only $Q_8 \times \{\bar{0}\}$ as a set, but has the group structure $Q_8 \times \{\bar{0}\}$ as a direct product. Thus both of the coset decompositions of $G/P(Q_1)$ and $G'/P(Q'_1)$ are $Q_8 \times \{\bar{0}\} \sqcup Q_8 \times \{\bar{1}\}$. Since neither of these two quandles Q_1 and Q'_1 satisfy (P2), we cannot use Theorem 3.2 to determine whether Q_1 and Q'_1 are isomorphic. In the following, we use Theorem 1.1 to show that Q_1 and Q'_1 are isomorphic.

Now we construct the two maps h and k in Theorem 1.1 for Q_1 and Q'_1 . Since $P(Q_1) = P(Q'_1) = Q_8 \times \{\bar{0}\}$ as sets, we can define the map $h: P(Q_1) \rightarrow P(Q'_1)$ to be the identity map. Obviously, the condition (A) of Theorem 1.1 can be verified. We take $A = \{(1, \bar{0}), (1, \bar{1})\}$ and $A' = \{(1, \bar{0}), (-\mathbf{i}, \bar{1})\}$ as the coset representatives of $G/P(Q_1)$ and $G'/P(Q'_1)$, respectively. Then we define the map $k: A \rightarrow A'$ by $k(1, \bar{0}) = (1, \bar{0})$ and $k(1, \bar{1}) = (-\mathbf{i}, \bar{1})$. Now we check the condition (B) of Theorem 1.1. For $(1, \bar{0}) \in A$, we have

$$h((1, \bar{0}) * (1, \bar{0})) = h(1, \bar{0}) = (1, \bar{0}) = (1, \bar{0}) *' k((1, \bar{0})).$$

For $(1, \bar{1}) \in A$, we have

$$h((1, \bar{0}) * (1, \bar{1})) = h((1, \bar{0})) = (1, \bar{0})$$

and

$$(1, \bar{0}) *' k(1, \bar{1}) = (1, \bar{0}) *' (-\mathbf{i}, \bar{1}) = (-\mathbf{i}, \bar{1}) \psi'_1((-\mathbf{i}, \bar{1})^{-1}) = (-\mathbf{i}, \bar{1}) (\mathbf{i}, \bar{1}) = (-\mathbf{i}^2, \bar{0}) = (1, \bar{0}).$$

Hence the condition (B) of Theorem 1.1 is also verified. Therefore, by Theorem 1.1, we have $Q_1 \cong Q'_1$.

Let $\tau: Q_8 \times \mathbb{Z}/2\mathbb{Z} \rightarrow Q_8 \times \mathbb{Z}/2\mathbb{Z}$ be the map defined by $\tau(a, i) = ((-1)^i a, i)$. We can check that τ is both of group automorphisms of G and G' . Let $\psi_2 = \tau \circ \psi_1$ and $\psi'_2 = \tau \circ \psi'_1$. Then both of the orders ψ_2 and ψ'_2 are 6.

Let Q_2 and Q'_2 be the generalized Alexander quandles $(Q(G, \psi_2), *)$ and $(Q(G', \psi'_2), *')$, respectively.

We construct h and k in Theorem 1.1 to show $Q_2 \cong Q'_2$.

We can check $P(Q_2) = P(Q'_2) = Q_8 \times \{\bar{0}\}$. Let $h: P(Q_2) \rightarrow P(Q'_2)$ be the identity map on $Q_8 \times \{\bar{0}\}$. The condition (A) of Theorem 1.1 is satisfied. We take $A = \{(1, \bar{0}), (1, \bar{1})\}$ and $A' = \{(1, \bar{0}), (\mathbf{i}, \bar{1})\}$ as the coset representatives of $G/P(Q_2)$ and $G'/P(Q'_2)$, respectively. Then we define the map $k: A \rightarrow A'$ by $k(1, \bar{0}) = (1, \bar{0})$ and $k(1, \bar{1}) = (\mathbf{i}, \bar{1})$. Now we verify the condition (B) of Theorem 1.1. For $(1, \bar{0}) \in A$, we have

$$h((1, \bar{0}) * (1, \bar{0})) = h(1, \bar{0}) = (1, \bar{0}) = (1, \bar{0}) *' k((1, \bar{0})).$$

For $(1, \bar{1}) \in A$, we have

$$h((1, \bar{0}) * (1, \bar{1})) = h((1, \bar{1})\psi_2((1, \bar{1})^{-1})) = h((1, \bar{1})\psi_2((1, \bar{1}))) = h((-1, \bar{1} + \bar{1})) = (-1, \bar{0})$$

and

$$(1, \bar{0}) *' k(1, \bar{1}) = (1, \bar{0}) *' (\mathbf{i}, \bar{1}) = (\mathbf{i}, \bar{1})\psi'_2((\mathbf{i}, \bar{1})^{-1}) = (\mathbf{i}, \bar{1})(\mathbf{i}, \bar{1}) = (-1, \bar{0}).$$

Hence the condition (B) of Theorem 1.1 is satisfied. Therefore, by Theorem 1.1, we have $Q_2 \cong Q'_2$.

Remark 5.1. For a quandle $Q = (Q, *)$ and a positive integer n , we can construct a quandle $Q^{(n)} = (Q, *^{(n)})$ by $x *^{(n)} y = s_y^n(x)$ where s_y is the point symmetry at y for the quandle $(Q, *)$. If two quandles Q and Q' are isomorphic, then $Q^{(n)}$ and $Q'^{(n)}$ are isomorphic (see [5, Section 4]). For a generalized Alexander quandle $Q = Q(G, \psi)$, we have $Q^{(n)} = Q(G, \psi^n)$. For the quandles Q_1, Q'_1, Q_2 and Q'_2 above, we see that $Q_2^{(2)} \cong Q_1$ and $Q_2'^{(2)} \cong Q'_1$. Thus, there is no need to verify $Q_1 \cong Q'_1$ once $Q_2 \cong Q'_2$ is verified. However, for the sake of clarity of argument, we verified $Q_1 \cong Q'_1$ at first.

5.3 Complete list of $|\mathcal{Q}_{\text{GAQ}}(n)|$ with $n \leq 127$

Similar to the case of order 16, there are some pairs of groups G, G' and their automorphisms ψ, ψ' such that we cannot use Theorem 3.2 to determine whether $Q(G, \psi)$ and $Q(G', \psi')$ are isomorphic. (For example, we encounter such cases in order 24 and 32.) By using Theorem 1.1, we can classify $|\mathcal{Q}_{\text{GAQ}}(n)|$ for small n 's. Table 1 is the list of $|\mathcal{Q}_{\text{GAQ}}(n)|$ with $n \leq 127$. These computations have been performed by using GAP [3] and the packages SmallGrp [1] and Rig [10]. We have made code available at the following URL:

https://github.com/Kurihara190/Classification_of_Generalized_Alexander_Quandles

Remark 5.2. The reason why the case $n = 128$ is not mentioned is that the number of nonisomorphic groups of order 128 is significantly larger compared to orders up to 127. Consequently, given the computational capacity of our current computer, the search would require an impractical amount of time to complete the calculations. However, Theorem 1.1 is valid for any order, so it may be possible to classify generalized Alexander quandles with orders higher than 128 in the future, provided that computational capacity and efficiency are improved.

n	1	2	3	4	5	6	7	8	9	10	11
$ Q_{\text{GAQ}}(n) $	1	1	2	3	4	3	6	9	11	5	10
n	12	13	14	15	16	17	18	19	20	21	22
$ Q_{\text{GAQ}}(n) $	11	12	7	8	29	16	17	18	15	13	11
n	23	24	25	26	27	28	29	30	31	32	33
$ Q_{\text{GAQ}}(n) $	22	32	39	13	51	20	28	15	30	87	20
n	34	35	36	37	38	39	40	41	42	43	44
$ Q_{\text{GAQ}}(n) $	17	24	64	36	19	25	45	40	23	42	32
n	45	46	47	48	49	50	51	52	53	54	55
$ Q_{\text{GAQ}}(n) $	44	23	46	114	83	49	32	39	52	87	41
n	56	57	58	59	60	61	62	63	64	65	66
$ Q_{\text{GAQ}}(n) $	66	37	29	58	60	60	31	69	363	48	33
n	67	68	69	70	71	72	73	74	75	76	77
$ Q_{\text{GAQ}}(n) $	66	51	44	35	70	205	72	37	84	56	60
n	78	79	80	81	82	83	84	85	86	87	88
$ Q_{\text{GAQ}}(n) $	41	78	158	299	41	82	79	64	43	56	97
n	89	90	91	92	93	94	95	96	97	98	99
$ Q_{\text{GAQ}}(n) $	88	85	72	68	61	47	72	378	96	97	110
n	100	101	102	103	104	105	106	107	108	109	110
$ Q_{\text{GAQ}}(n) $	154	100	51	102	117	52	53	106	342	108	57
n	111	112	113	114	115	116	117	118	119	120	121
$ Q_{\text{GAQ}}(n) $	73	204	112	59	88	87	135	59	96	177	219
n	122	123	124	125	126	127					
$ Q_{\text{GAQ}}(n) $	61	80	92	297	127	126					

Table 1: List of $|Q_{\text{GAQ}}(n)|$ with $n \leq 127$

References

- [1] H.U. Besche, B. Eick, E. O'Brien and M. Horn, *Small-Grp, The GAP Small Groups Library*. Available at <https://gap-packages.github.io/smallgrp/>.
- [2] M. Bonatto, *Principal and doubly homogeneous quandles*, *Monatsh. Math.* **191** (2020), no. 4, 691–717.

- [3] GAP - Groups, Algorithms, and Programming, Version 4.12.2 (2022), <http://www.gap-system.org>.
- [4] A. Higashitani and H. Kurihara, *Generalized Alexander quandles of finite groups and their characterizations*, to appear in *European J. Math.*, also available at arXiv:2210.16763.
- [5] A. Higashitani and H. Kurihara, *Homogeneous quandles arising from automorphisms of symmetric groups*, *Comm. Algebra* **51** (2023), no. 4, 1413–1430.
- [6] D. Joyce, *A classifying invariant of knots, the knot quandle*, *J. Pure Appl. Algebra* **23** (1982), no. 1, 37–65.
- [7] S.V. Matveev, *Distributive groupoids in knot theory*, *Mat. Sb.* **161** (1982), no. 1, 78–88.
- [8] S. Nelson, *Classification of finite Alexander quandles*, *Proceedings of the Spring Topology and Dynamical Systems Conference*, *Topology Proc.* **27** (2003), no. 1, 245–258.
- [9] Y. Taniguchi, *Good involutions of generalized Alexander quandles*, *J. Knot Theory Ramifications* **32** (2023), no. 12, Paper No. 2350081, 7 pp.
- [10] L. Vendramin, *Rig, a GAP package for racks, quandles and Nichols algebras*. Available at <https://github.com/gap-packages/rig/>.

Author’s addresses.

Akihiro Higashitani:

Department of Pure and Applied Mathematics, Osaka University, Suita, Osaka 565-0871, Japan

E-mail address: higashitani@ist.osaka-u.ac.jp

Seiichi Kamada:

Department of Mathematics, Osaka University, Toyonaka, Osaka 560-0043, Japan

E-mail address: kamada@math.sci.osaka-u.ac.jp

Jin Kosaka:

Department of Mathematics, Osaka University, Toyonaka, Osaka 560-0043, Japan

E-mail address: kosakajin123@gmail.com

Hirotake Kurihara:

Department of Applied Science, Yamaguchi University, Ube 755-8611, Japan

E-mail address: kurihara-hiro@yamaguchi-u.ac.jp