

# CHARACTERISATION OF DISTAL ACTIONS OF AUTOMORPHISMS ON THE SPACE OF ONE-PARAMETER SUBGROUPS OF LIE GROUPS

DEBAMITA CHATTERJEE AND RIDDHI SHAH

**ABSTRACT.** For a connected Lie group  $G$  and an automorphism  $T$  of  $G$ , we consider the action of  $T$  on  $\text{Sub}_G$ , the compact space of closed subgroups of  $G$  endowed with the Chabauty topology. We study the action of  $T$  on  $\text{Sub}_G^p$ , the closure in  $\text{Sub}_G$  of the set of closed one-parameter subgroups of  $G$ . We relate the distality of the  $T$ -action on  $\text{Sub}_G^p$  with that of the  $T$ -action on  $G$  and characterise the same in terms of compactness of the closed subgroup generated by  $T$  in  $\text{Aut}(G)$  when  $T$  acts distally on the maximal central torus and  $G$  is not a vector group. We extend these results to the action of subgroups  $\mathcal{H}$  of  $\text{Aut}(G)$  and equate the distal action of any closed subgroup  $\mathcal{H}$  on  $\text{Sub}_G^p$  with that of every element in  $\mathcal{H}$ . Moreover, we show that a connected Lie group  $G$  acts distally on  $\text{Sub}_G^p$  by conjugation if and only if  $G$  is either compact or it is isomorphic to a direct product of a compact group and a vector group. Our results generalise some results proven by Shah and Yadav.

*2020 Mathematics Subject Classification.* Primary 37B05; Secondary 22E15, 22D45

*Keywords.* Distal actions of automorphisms, Lie groups, one-parameter subgroups, Chabauty topology.

## 1. INTRODUCTION

For a Hausdorff topological space, a homeomorphism  $T$  of  $X$  is said to be *distal* (equivalently,  $T$  acts *distally* on  $X$ ) if for any pair of distinct elements  $x, y \in X$ , the closure of the double orbit  $\{(T^n(x), T^n(y)) \mid n \in \mathbb{Z}\}$  in  $X \times X$  does not intersect the diagonal, i.e. for  $x, y \in X$  with  $x \neq y$ ,  $\overline{\{(T^n(x), T^n(y)) \mid n \in \mathbb{Z}\}} \cap \{(a, a) \mid a \in X\} = \emptyset$ . David Hilbert introduced the notion of distality to study non-ergodic actions on compact spaces (see Ellis [16] and Moore [29]). Distal actions on compact spaces, Lie groups as well as locally compact groups have

---

*Date:* May 30, 2024.

been studied by many mathematicians in different contexts (see Abels [1, 2], Choudhuri et al [10], Ellis [16], Furstenberg [17], Palit-Shah [30], Palit-Prajapati-Shah [31], Raja-Shah [33, 34], Shah [36], Shah-Yadav [38, 39, 40], and references cited therein).

Let  $G$  be a locally compact (Hausdorff) group with the identity  $e$  and let  $\text{Sub}_G$  denote the set of all closed subgroups of  $G$  endowed with the Chabauty topology. Note that  $\text{Sub}_G$  is compact and Hausdorff, and it is metrisable if  $G$  is second countable [6]. Let  $\text{Aut}(G)$  denote the group of all (bicontinuous) automorphisms of  $G$ . There is a canonical group action of  $\text{Aut}(G)$  on  $\text{Sub}_G$ ; namely,  $(T, H) \mapsto T(H)$ ,  $T \in \text{Aut}(G)$ ,  $H \in \text{Sub}_G$ . Equivalently, we have a homomorphism from  $\text{Aut}(G)$  to  $\text{Homeo}(\text{Sub}_G)$ . As the image of  $\text{Aut}(G)$  under this action is a large subclass of homeomorphisms of  $\text{Sub}_G$ , it is important to study the dynamics of the action of this special subclass in terms of distality. In [40], Shah and Yadav have studied the action of automorphisms of a connected Lie groups  $G$  on  $\text{Sub}_G$  extensively and have shown in particular that for a large class of  $G$  without the compact central subgroup of positive dimension, an automorphism  $T$  of  $G$  acts distally on  $\text{Sub}_G^a$ , the space of closed abelian subgroups of  $G$  if and only if  $T$  generates a relatively compact subgroup of  $\text{Aut}(G)$ . A similar characterisation also holds for  $T$  belonging to the class (NC), i.e. those  $T$  for which the closure of the  $T$ -orbit of any discrete cyclic group does not contain the trivial group  $\{e\}$ . Note that  $\text{Sub}_G^a$  is very large in many groups, e.g. it is the same as  $\text{Sub}_G$  if  $G$  is abelian. It also contains all discrete cyclic subgroups. Shah, with Palit in [30], and with Palit and Prajapati in [31], has studied distal actions of automorphisms of a discrete group  $G$  on  $\text{Sub}_G^a$ , where  $G$  is either polycyclic or a lattice in a connected Lie group.

Here, we would like to consider a closed subset of  $\text{Sub}_G^a$  containing smallest closed connected subgroups of  $G$ ; namely, the closure in  $\text{Sub}_G$  of the set of all closed one-parameter subgroups of  $G$ ; we denote this closed set by  $\text{Sub}_G^p$ . Note that  $\text{Sub}_G^p$  is invariant under the action of  $\text{Aut}(G)$ . We would like to study the dynamics of the actions of automorphisms on  $\text{Sub}_G^p$  and characterise them in terms of distality. In the case of  $\mathbb{R}^n$ ,  $n \geq 2$ ,  $\text{Sub}_{\mathbb{R}^n}^p$  is homeomorphic to  $\mathbb{R}\mathbb{P}^{n-1} \sqcup \{\{0\}\}$ . It is indeed much smaller than  $\text{Sub}_{\mathbb{R}^n}^a = \text{Sub}_{\mathbb{R}^n}$ . In general for a Lie group  $G$ ,  $\text{Sub}_G^p$  does not contain any non-trivial discrete subgroup as it can not be a limit point of closed one-parameter subgroups (cf. Lemma

2.2). Theorem 4.1 of Shah and Yadav in [40] mainly deals with the action of automorphisms on discrete cyclic subgroups. Here, we are concentrating on the action of  $T$  on  $\text{Sub}_G^p$ , the smallest compact subset of  $\text{Sub}_G$  which contains all closed (connected) one-parameter subgroups of  $G$ . and get a characterisation for the distal action for a large class of connected non-abelian Lie groups  $G$ .

**Theorem 1.1.** *Let  $G$  be a connected Lie group without any non-trivial compact connected central subgroup and let  $T$  be an automorphism of  $G$ . Then the following hold:*

- (1) *If  $G$  is abelian, then  $G$  is isomorphic to  $\mathbb{R}^d$  for some  $d \in \mathbb{N}$ , and  $T$  acts distally on  $\text{Sub}_G^p$  if and only if  $T \in \mathcal{KD}$ , where  $\mathcal{K}$  is a compact subgroup of  $\text{GL}(d, \mathbb{R})$  and  $\mathcal{D}$  is the center of  $\text{GL}(d, \mathbb{R})$ .*
- (2) *If  $G$  is not abelian, then  $T$  acts distally on  $\text{Sub}_G^p$  if and only if  $T \in \mathcal{K}$ , a compact subgroup of  $\text{Aut}(G)$ .*

Note that the condition on the center in Theorem 1.1 holds if  $G$  is simply connected, or more generally its nilradical is simply connected. It also holds for  $G/M$  for any connected Lie group  $G$ , where  $M$  is the maximal central torus in  $G$  (the maximal compact connected central subgroup in  $G$ ). The result in case of non-abelian groups  $G$  in (2) as above gives a similar characterisation as in [40], even though the space  $\text{Sub}_G^p$  is much smaller than  $\text{Sub}_G^a$ . In case  $G$  as above is abelian, then  $G = \mathbb{R}^d$ , a vector group, and  $\text{Sub}_G^p$  is homeomorphic to  $\mathbb{RP}^{d-1} \sqcup \{0\}$  as noted earlier, and the Theorem 1.1 (1) shows that a larger class of  $T$  act distally on  $\text{Sub}_G^p$ .

In [40], it is shown for a connected Lie group without any compact central subgroup of positive dimension that if  $T$  acts distally on  $\text{Sub}_G^a$ , then it acts distally on  $G$  (more generally, see Theorem 3.6 and Corollary 3.7 of [40]). We generalise this for Lie groups which are not vector groups; as in the later case, Theorem 1.1 (1) shows that it is possible to have automorphisms which act distally on  $\text{Sub}_G^p$  but not on  $G$ .

**Theorem 1.2.** *Let  $G$  be a connected Lie group which is not a vector group and let  $T \in \text{Aut}(G)$  be such that it acts distally on the largest compact connected central subgroup of  $G$ . If  $T$  acts distally on  $\text{Sub}_G^p$ , then  $T$  acts distally on  $G$ .*

An automorphism  $T$  of a connected Lie group  $G$  is said to be unipotent if all the eigenvalues of  $dT$  are equal to 1, where  $dT$  is the corresponding Lie algebra automorphism of the Lie algebra of  $G$ . It is known that  $T$  acts distally on  $G$  if and only if all the eigenvalues of  $dT$  have absolute value 1 (cf. [1]). Any unipotent automorphism acts distally on  $G$  and if it belongs to a compact subgroup of  $\text{Aut}(G)$ , then it is trivial. Theorem 4.3 of [40] shows that no non-trivial unipotent automorphism acts distally on  $\text{Sub}_G^a$ ; generalising this statement we prove the following.

**Theorem 1.3.** *Let  $G$  be a connected Lie group and let  $T \in \text{Aut}(G)$  be unipotent. Then  $T$  acts distally on  $\text{Sub}_G^p$  if and only if  $T = \text{Id}$ , the identity map.*

Theorem 1.3 illustrates in particular that the converse of Theorem 1.2 does not hold as the non-trivial unipotent automorphisms do not act distally on  $\text{Sub}_G^p$ .

Let  $\text{Sub}_G^c$  denote the set of all closed discrete cyclic subgroups of  $G$ . It is invariant under the action of  $\text{Aut}(G)$  and so is its closure in  $\text{Sub}_G$ . In [30] and [31], distal actions of automorphisms of  $\text{Sub}_\Gamma^c$  have been characterised, where  $\Gamma$  is a certain type of discrete group, e.g. a discrete polycyclic group or a lattice in a connected Lie group. In a Lie group  $G$ , any closed one-parameter subgroup  $H = \{x_t\}_{t \in \mathbb{R}}$  is the limit of  $\{H_n\}_{n \in \mathbb{N}}$  as  $n \rightarrow \infty$ , where  $H_n = \{x_{m/n} \mid m \in \mathbb{Z}\}$ . Here,  $H_n$  is a closed discrete cyclic group generated by  $x_{1/n}$ . If  $H$  is compact, we may assume that  $x_1 = e$ , the identity in  $G$ . Thus  $\text{Sub}_G^p \subset \overline{\text{Sub}_G^c}$  (the closure of  $\text{Sub}_G^c$  in  $\text{Sub}_G$ ) and the latter is contained in  $\text{Sub}_G^a$ . We should note that  $\overline{\text{Sub}_G^c}$  is much bigger than  $\text{Sub}_G^p$  as the latter contains all discrete cyclic subgroups, and these do not belong to  $\text{Sub}_G^p$  (see Lemma 2.2). It is easy to see that Theorems 1.1 (2), 1.2 and 1.3 remain valid if  $\text{Sub}_G^p$  is replaced by  $\overline{\text{Sub}_G^c}$  in the respective statement.

The following generalises a part of Theorem 4.1 of [40] and also Theorem 1.1 (2).

**Theorem 1.4.** *Let  $G$  be a connected Lie group such that it is not a vector group. Let  $T \in \text{Aut}(G)$  be such that it acts distally on the largest compact connected central subgroup of  $G$ . Then the following are equivalent:*

- (1)  $T$  acts distally on  $\text{Sub}_G^p$ .
- (2)  $T$  acts distally on  $\overline{\text{Sub}_G^c}$ .

- (3)  $T$  acts distally on  $\text{Sub}_G^a$ .
- (4)  $T$  acts distally on  $\text{Sub}_G$ .
- (5)  $T$  is contained in a compact subgroup of  $\text{Aut}(G)$ .

We may note that in case  $G$  is a vector group, Theorem 1.1(1) shows that (1) and (5) above are not equivalent; however, in this case, statements (2 – 5) above are equivalent by Theorem 4.1 of [40] as (2) above implies that  $T \in (\text{NC})$ , i.e.  $\{e\}$  is not a limit point of  $\{T^n(H)\}_{n \in \mathbb{Z}}$  for any closed discrete cyclic subgroup  $H$  of  $G$ .

Recall that a semigroup  $\Gamma$  of homeomorphisms of a (Hausdorff) topological space  $X$  acts distally on  $X$  if  $\overline{\{(\gamma(x), \gamma(y)) \mid \gamma \in \Gamma\}} \cap \{(d, d) \mid d \in X\} = \emptyset$ . Note that a homeomorphism  $T$  of  $X$  acts distally on  $X$  if and only if the cyclic group  $\{T^n \mid n \in \mathbb{Z}\}$  acts distally on  $X$ . For a Lie group  $G$ , any compact subgroup of  $\text{Aut}(G)$  acts distally on  $\text{Sub}_G$  (see e.g. Lemma 2.2 in [39]). It follows from Theorem 1.1 of [2] that  $\mathcal{H}$  acts distally on  $G$  if and only if  $\mathcal{H}$  acts distally on the Lie algebra of  $G$ , and these statements are equivalent to the statement that every element of  $\mathcal{H}$  acts distally on  $G$  (see also [11]). The following result generalises Theorems 1.1 and 1.4 to the action of any subgroup  $\mathcal{H}$  of  $\text{Aut}(G)$  and, if  $\mathcal{H}$  is closed, then it equates the distal action of  $\mathcal{H}$  on  $\text{Sub}_G^p$  with that of every element of  $\mathcal{H}$ . Let  $\overline{\mathcal{H}}$  denote the closure of  $\mathcal{H}$  in  $\text{Aut}(G)$ . It is easy to see using Abel's results in [1, 2] that  $\mathcal{H}$  acts distally on  $G$  if and only if so does  $\overline{\mathcal{H}}$ . It follows from Ellis' result in [16] that  $\mathcal{H}$  acts distally on any compact  $\mathcal{H}$ -invariant subspace of  $\text{Sub}_G$  if and only if so does  $\overline{\mathcal{H}}$ .

**Theorem 1.5.** *Let  $G$  be a connected Lie group. Let  $\mathcal{H}$  be a subgroup of  $\text{Aut}(G)$  such that it acts distally on the largest compact connected central subgroup of  $G$ . Consider the following statements:*

- (1) Every element of  $\overline{\mathcal{H}}$  acts distally on  $\text{Sub}_G^p$ .
- (2)  $\mathcal{H}$  acts distally on  $\text{Sub}_G^p$ .
- (3)  $\mathcal{H}$  acts distally on  $\overline{\text{Sub}_G^c}$ .
- (4)  $\mathcal{H}$  acts distally on  $\text{Sub}_G^a$ .
- (5)  $\mathcal{H}$  acts distally on  $\text{Sub}_G$ .
- (6)  $\overline{\mathcal{H}}$  is a compact group.

*Then the following hold:*

- (a) (1) and (2) are equivalent.
- (b) (3 – 6) are equivalent.
- (c) If  $G$  is not a vector group, then (1 – 6) are equivalent.

- (d) If  $G = \mathbb{R}^d$ , for some  $d \in \mathbb{N}$ , then each of (1) and (2) is equivalent to the following:
- (7)  $\mathcal{H} \subset \mathcal{KD}$ , where  $\mathcal{K}$  is a compact subgroup of  $\mathrm{GL}(d, \mathbb{R})$  and  $\mathcal{D}$  is the center of  $\mathrm{GL}(d, \mathbb{R})$ .

Note that in Theorem 1.5, one can also have equivalent statements similar to (1 – 2), where  $\mathrm{Sub}_G^p$  is replaced by any of  $\overline{\mathrm{Sub}}_G^c$ ,  $\mathrm{Sub}_G^a$  or  $\mathrm{Sub}_G$  for the action of a subgroup  $\mathcal{H}$ . Statement (6) in the theorem implies that  $\mathcal{H}$  acts distally on  $G$ , and hence the assertion (c) above implies that Theorem 1.2 remains valid for the action of a subgroup of  $\mathrm{Aut}(G)$  instead of an automorphism. We may note here that there is a subgroup  $\mathcal{H}$  of  $\mathrm{SL}(4, \mathbb{C})$  in which every element of  $\mathcal{H}$  is contained in a compact subgroup of  $\mathrm{SL}(4, \mathbb{C})$  but the closure of  $\mathcal{H}$  in  $\mathrm{SL}(4, \mathbb{C})$  is connected and contains unipotent elements (cf. [5]). Hence, one can not replace  $\overline{\mathcal{H}}$  by  $\mathcal{H}$  in (1) of Theorem 1.5.

The condition that  $T$  (resp. a subgroup  $\mathcal{H}$  of  $\mathrm{Aut}(G)$ ) acts distally on the maximal compact connected central subgroup (maximal central torus) of  $G$  is satisfied easily if  $T \in \mathrm{Aut}(G)^0$ , (resp.  $\mathcal{H} \subset \mathrm{Aut}(G)^0$ ), as  $\mathrm{Aut}(G)^0$  acts trivially on any compact central subgroup. It is also satisfied if  $T^n \in \mathrm{Aut}(G)^0$  for some  $n \in \mathbb{N}$  (resp.  $\mathcal{H}$  or  $\mathcal{H}\mathrm{Aut}(G)^0$  has finitely many connected components). Here,  $\mathrm{Aut}(G)^0$  denotes the connected component of identity in  $\mathrm{Aut}(G)$ .

Let  $\mathrm{Inn}(G)$  denote the group of inner automorphisms of  $G$ . It is a (not necessarily closed) normal subgroup of  $\mathrm{Aut}(G)$ . The following corollaries generalises the first part of Corollaries 4.5 [40].

**Corollary 1.6.** *Let  $G$  be a connected Lie group. Then the following are equivalent:*

- (1) *Every inner automorphism of  $G$  acts distally on  $\mathrm{Sub}_G^p$ .*
- (2) *Every inner automorphism of  $G$  acts distally on  $\mathrm{Sub}_G$ .*
- (3)  *$\mathrm{Inn}(G)$  acts distally on  $\mathrm{Sub}_G^p$ .*
- (4)  *$\mathrm{Inn}(G)$  acts distally on  $\mathrm{Sub}_G$ .*
- (5)  *$G$  is compact or  $G = \mathbb{R}^n \times K$ , for some  $n \in \mathbb{N}$  and the maximal compact normal subgroup  $K$  of  $G$ .*

In Corollary 1.6, one can also write equivalent statements involving the action on  $\overline{\mathrm{Sub}}_G^c$ .

Note that  $\mathrm{Sub}_G^p$  is contained in a larger compact set  $\mathrm{Sub}_G^{co}$ , which is the closure of the set of all closed connected subgroups of  $G$ . Hence it is easy to show that Theorems 1.1–1.3 are all valid if  $\mathrm{Sub}_G^p$  is replaced

by  $\text{Sub}_G^{\text{co}}$ . In Theorems 1.4 and 1.5 and Corollary 1.6, the action of the given automorphism or a subgroup of  $\text{Aut}(G)$  is distal on  $\text{Sub}_G^{\text{co}}$  if and only if it is so on  $\text{Sub}_G^p$ .

Ellis [16] has shown that a semigroup  $\Gamma$  of homeomorphisms of a compact space  $X$  acts distally on  $X$  if and only if the closure of  $\Gamma$  in  $X^X$  is a group. Note that minimal distal actions on compact spaces are characterised by Furstenberg [17]. For many groups  $G$ , compact spaces  $\text{Sub}_G$ ,  $\text{Sub}_G^a$  and  $\text{Sub}_G^c$  are identified (see Baik-Clavier [3, 4], Bridson et al [8], Hamrouni and Kadri [19], Kloeckner [27] and also Pourezza and Habbard [32]); one can also identify  $\text{Sub}_G^p$  for some groups. The action of  $\text{Aut}(G)$  on some subspaces of  $\text{Sub}_G$  for the 3-dimensional Heisenberg group has also been described in detail [8]. The space of  $G$ -invariant measures on  $\text{Sub}_G$  and in particular, ergodic  $G$ -invariant measures on  $\text{Sub}_G$  for some  $G$  have also been studied extensively (see Gelandar [18] and the references cited therein). Since automorphisms of  $G$  corresponds to a large subclass of homeomorphisms of  $\text{Sub}_G$  it is of significant interest to study the dynamics of the action of  $\text{Aut}(G)$  on closed (compact) invariant subspaces of  $\text{Sub}_G$ .

Throughout,  $G$  is a connected Lie group unless specified otherwise. Let  $R$  denote the radical of  $G$ , i.e. the maximal connected solvable normal subgroup of  $G$  and let  $N$  denote the nilradical of  $G$ , i.e. the maximal connected nilpotent normal subgroup of  $G$ . Both the radical and the nilradical of  $G$  are characteristic in  $G$ . Recall that  $G$  is said to be semisimple if its radical is trivial. In particular,  $G/R$  is semisimple. For a closed subgroup  $H$  of  $G$ , let  $H^0$  denote the connected component of the identity  $e$  in  $H$ ,  $[H, H]$  denote the commutator subgroup of  $H$ , i.e.  $[H, H]$  is generated by  $\{xy, x^{-1}y^{-1} \mid x, y \in H\}$ . Note that  $H^0$ ,  $[H, H]$  and its closure  $\overline{[H, H]}$  are characteristic in  $H$ .

In §2, we study general properties of  $\text{Sub}_G$  and prove some elementary results on the structure of  $\text{Sub}_G^p$ . In §3, we discuss some preliminary results on the action of  $\text{Aut}(G)$  on  $G$  and  $\text{Sub}_G^p$  and prove main results stated above.

## 2. ACTIONS OF AUTOMORPHISMS OF $G$ ON $\text{Sub}_G$ AND $\text{Sub}_G^p$

In this section, we discuss some basic properties of Chabauty topology on the space  $\text{Sub}_G$  of closed subgroups of  $G$ , the structure of  $\text{Aut}(G)$  and its action on  $\text{Sub}_G$ . We also discuss some elementary results about the structure of  $\text{Sub}_G$  and  $\text{Sub}_G^p$ .

Let  $G$  be a locally compact Hausdorff topological group. For a compact set  $K$  and an open set  $U$  in  $G$ , we set  $U_1(K) = \{H \in \text{Sub}_G \mid H \cap K = \emptyset\}$  and  $U_2(U) = \{H \in \text{Sub}_G \mid H \cap U \neq \emptyset\}$ . Here,  $\{U_1(K) \mid K \text{ is compact}\} \cup \{U_2(U) \mid U \text{ is open}\}$  is a sub-basis of Chabauty topology on  $\text{Sub}_G$ . As mentioned earlier,  $\text{Sub}_G$  is compact and Hausdorff, and if  $G$  is second countable, then  $\text{Sub}_G$  is metrisable (cf. [6], Lemma E.1.1). In particular, for a connected Lie group and its closed subgroups  $G$ ,  $\text{Sub}_G$  is metrisable.

**Lemma 2.1.** *Let  $G$  be a locally compact second countable group. A sequence  $\{H_n\} \subset \text{Sub}_G$  converges to  $H \in \text{Sub}_G$  if and only if the following hold:*

- (I) *For  $g \in G$ , if there exists a subsequence  $\{H_{n_k}\}$  of  $\{H_n\}$  with  $h_k \in H_{n_k}$ ,  $k \in \mathbb{N}$ , such that  $h_k \rightarrow g$  in  $G$ , then  $g \in H$ .*
- (II) *For every  $h \in H$ , there exists a sequence  $\{h_n\}_{n \in \mathbb{N}}$  such that  $h_n \in H_n$ ,  $n \in \mathbb{N}$ , and  $h_n \rightarrow h$ .*

The following lemma is well-known. We include a short proof for the sake of completeness.

**Lemma 2.2.** *Let  $G$  be a second countable Lie group. Suppose  $\{H_n\}_{n \in \mathbb{N}}$  in  $\text{Sub}_G$  is such that it converges to  $H \in \text{Sub}_G$ . If  $H$  is discrete, then  $H_n$  is discrete for all large  $n$ . In particular, the set of discrete subgroups of  $G$  is open in  $\text{Sub}_G$ .*

*Proof.* It is enough to show that  $H_n^0$  is trivial for all large  $n$ . As  $G^0$  is open in  $G$ , we get that  $H_n \cap G^0 \rightarrow H \cap G^0$  and the latter is discrete. As  $H_n^0 \subset G^0$ , we may replace  $H_n$  by  $H_n \cap G^0$ ,  $H$  by  $H \cap G^0$ , and  $G$  by  $G^0$  and assume that  $G$  is connected. As  $G$  is a Lie group, there exists a neighbourhood  $U$  of the identity  $e$  in  $G$  such that the only subgroup contained in  $U$  is the trivial subgroup  $\{e\}$ . As  $H$  is discrete, there exists a neighbourhood  $W$  of the identity  $e$  in  $G$  such that  $H \cap W = \{e\}$ . Since  $G$  is locally compact and Hausdorff, there exist an open set  $V$  and a compact set  $K$  such that  $e \in V \subset K \subset U \cap W$ . As  $G$  is connected,  $K \cap V^c$  is a nonempty compact set and

$$H \cap (K \cap V^c) \subset H \cap W \cap V^c = \emptyset,$$

and hence  $H \in U_1(K \cap V^c)$  which is an open set in  $\text{Sub}_G$ . As  $H_n \rightarrow H$  in  $\text{Sub}_G$ , we get that for all large  $n$ ,  $H_n \in U_1(K \cap V^c)$ , i.e.  $H_n \cap (K \cap V^c) = \emptyset$ , and hence  $H_n \subset V \sqcup K^c$ . Thus for all large  $n$ ,  $H_n^0 \subset V \subset U$ , and it



follows that  $H_n^0 = \{e\}$ , and hence  $H_n$  is discrete. The second assertion easily follows from the first.  $\square$

Recall that an automorphism of  $G$  is a homomorphism and a homeomorphism, and  $\text{Aut}(G)$  denotes the group of all automorphisms of  $G$ . There is a natural group action of  $\text{Aut}(G)$  on  $\text{Sub}_G$  defined as follows:

$$\text{Aut}(G) \times \text{Sub}_G \rightarrow \text{Sub}_G, (T, H) \mapsto T(H); T \in \text{Aut}(G), H \in \text{Sub}_G.$$

It is easy to see that the map  $H \mapsto T(H)$  defines a homeomorphism of  $\text{Sub}_G$  for every  $T \in \text{Aut}(G)$ , and the corresponding map is  $\phi : \text{Aut}(G) \rightarrow \text{Homeo}(\text{Sub}_G)$ , where  $\phi(T)(H) = T(H)$ . It is easy to check that  $\phi$  is a group homomorphism.

Recall that for a locally compact Hausdorff group  $G$ , if  $\text{Aut}(G)$  is equipped with the compact-open topology, then

$$\text{Aut}(G) \times G \rightarrow G, (T, x) \mapsto T(x), T \in \text{Aut}(G), x \in G,$$

is continuous. If we consider the modified compact-open topology on  $\text{Aut}(G)$ , it is a topological group [42]. In Lemma 2.4 of [40], it is proved that the action of the group  $\text{Aut}(G)$  on  $\text{Sub}_G$  is continuous, implying that  $\phi$  is continuous. Note that for a connected Lie group  $G$ , the compact-open topology and the modified compact-open topology on  $\text{Aut}(G)$  are same. In particular, every  $T$  which generates a relatively compact group in  $\text{Aut}(G)$  acts distally on  $\text{Sub}_G$  (see e.g. [30], Lemma 4.2).

Let  $G$  be a connected Lie group with the Lie algebra  $\mathfrak{g}$ . There is an exponential map  $\exp : \mathfrak{g} \rightarrow G$  which is a continuous map and its restriction to a small open neighbourhood of 0 in  $\mathfrak{g}$  is a homeomorphism onto an open neighbourhood of the identity  $e$  in  $G$ . Given  $T \in \text{Aut}(G)$ , there exists a corresponding Lie algebra automorphism  $dT : \mathfrak{g} \rightarrow \mathfrak{g}$  such that  $\exp \circ dT = T \circ \exp$ . The map  $d : \text{Aut}(G) \rightarrow \text{Aut}(\mathfrak{g})$ ,  $T \mapsto dT$  is an injective continuous homomorphism and its image is closed, and in fact,  $d$  is a homeomorphism onto its image (cf. [9], see also [21]) It is known that  $\text{Aut}(\mathfrak{g})$  is a closed subgroup of  $\text{GL}(\mathfrak{g})$ , and we can view  $\text{Aut}(G)$  as a closed subgroup of  $\text{GL}(\mathfrak{g})$ . The topology on  $\text{Aut}(G)$  when considered as a subspace topology of  $\text{GL}(\mathfrak{g})$  coincides with the (modified) compact-open topology. We refer the reader to [22] for the basic structure theory of Lie groups.

Let  $G$  be a connected Lie group. A (continuous real) one-parameter subgroup in  $G$  is the image of a continuous homomorphism from  $\mathbb{R}$  to

$G$ . Any one-parameter subgroup is of the form  $\{\exp tX\}_{t \in \mathbb{R}}$  for some  $X$  in the Lie algebra  $\mathfrak{g}$  of  $G$ . Moreover, if  $H$  is a one-parameter subgroup in  $G$ , then so is  $T(H)$  for any  $T \in \text{Aut}(G)$ . Let

$$\mathcal{P}_1(G) := \{A \mid A \text{ is a closed one-parameter subgroup of } G\}.$$

Then  $\mathcal{P}_1(G) \subset \text{Sub}_G$  and it is invariant under the action of  $\text{Aut}(G)$ . Let  $\text{Sub}_G^p$  denote the closure (in  $\text{Sub}_G$ ) of the set  $\mathcal{P}_1(G)$  of all closed one-parameter subgroups in  $G$ , i.e.

$$\text{Sub}_G^p := \overline{\mathcal{P}_1(G)}.$$

Then  $\text{Sub}_G^p$  is compact and it is also invariant under the action of  $\text{Aut}(G)$ .

Note that if any one-parameter subgroup in a locally compact group is not closed, then its closure is compact and isomorphic to an  $n$  dimensional torus  $\mathbb{T}^n = (\mathbb{S}^1)^n$  for some  $n \in \mathbb{N}$  with  $n \geq 2$ . If it is closed and bounded (compact), then it is isomorphic to  $\mathbb{S}^1$ . If it is unbounded (not relatively compact), then it is closed and isomorphic to  $\mathbb{R}$ .

For brevity, for a set  $\{x_t\}_{t \in \mathbb{R}}$  which may be a continuous one-parameter set or a one-parameter subgroup, we may just write  $\{x_t\}$ , i.e. we may omit the parameter set if there is no ambiguity.

Now we prove some elementary results about the structure of  $\text{Sub}_G^p$ . The following lemma shows that if  $G$  contains any non-trivial compact connected abelian subgroup and the dimension of  $G$  is at least 2, then  $\mathcal{P}_1(G)$  is not closed in  $\text{Sub}_G$ .

**Lemma 2.3.** *Let  $G$  be a connected Lie group and let  $\text{Sub}_G^p$  be the closure in  $\text{Sub}_G$  of all closed one-parameter subgroups. Then the following holds:*

- (1) *All compact connected abelian subgroups of  $G$  belong to  $\text{Sub}_G^p$ .*
- (2) *If  $\{x_t\} \in \mathcal{P}_1(G) \subset \text{Sub}_G^p$  and if  $K$  is a compact connected abelian subgroup of  $G$  which is centralised by each  $x_t$ . Then  $\{x_t\}K \in \text{Sub}_G^p$ .*

*Proof.* Let  $K$  be a compact connected abelian subgroup. Then the set  $F$  of all finite order elements in  $K$  is dense in  $K$  and each such element of  $F$  is contained in a closed one-parameter subgroup of  $K$ . Since  $K$  is monothetic (i.e. it has a dense cyclic subgroup), we get that  $K$  is a limit of a sequence of compact one-parameter subgroups of  $K$  in  $\text{Sub}_K$ . Therefore,  $K \in \text{Sub}_G^p$ . Thus (1) holds.

Let  $\{x_t\} \in \mathcal{P}_1(G) \subset \text{Sub}_G^p$  be such that each  $x_t$  centralises  $K$ . If  $\{x_t\}$  is compact, then  $\{x_t\}K$  is a compact connected abelian subgroup of  $G$  and hence  $\{x_t\}K \in \text{Sub}_G^p$  from (1). Now suppose  $\{x_t\}$  is isomorphic to  $\mathbb{R}$ . Let  $k \in K$  be such that it generates a dense cyclic subgroup in  $K$ . Since  $K$  is exponential, there exists a one-parameter subgroup  $\{k_t\}$ , which is dense in  $K$  and  $k = k_1$ . Let  $\{n_m\}$  be such that  $k^{n_m} \rightarrow k$ . Let  $A_m = \{a_t(m) = x_{t/n_m^2} k_t\}_{t \in \mathbb{R}}$ . Then each  $A_m$  is a closed one-parameter subgroup in  $G$ , and passing to a subsequence if necessary, we get that  $A_m \rightarrow A$  for some subgroup  $A \in \text{Sub}_G^p$ . Note that  $\{x_t\} \times K$  is a closed subgroup of  $G$  and it contains each  $A_m$ . Therefore,  $A \subset \{x_t\} \times K$ . Since  $a_{n_m}(m) = x_{1/n_m} k_{n_m} = x_{1/n_m} k^{n_m} \rightarrow k$ , we get that  $k \in A$ . Therefore,  $K \subset A$ . For a fixed  $t$ , let  $b_m = a_{tn_m^2}(m) = x_t k_{tn_m^2} \in A_m$ ,  $m \in \mathbb{N}$ . Passing to a subsequence of  $\{b_m\}$  if necessary, we get that  $b_m \rightarrow x_t h$  for some  $h \in K$ , and hence,  $x_t \in A$ . Now  $\{x_t\} \subset A$ . Hence,  $A = \{x_t\} \times K$  and (2) holds.  $\square$

The following lemma relates the set  $\text{Sub}_{G/K}^p$  and a compact subset of  $\text{Sub}_G^p$  which is invariant under the action of  $\text{Aut}(G)$ .

**Lemma 2.4.** *Let  $G$  be a connected Lie group,  $K$  be a compact connected central subgroup of  $G$  and let  $\pi : G \rightarrow G/K$  be the natural projection. Then the following hold:*

- (1)  $\{HK \mid H \in \text{Sub}_G\}$  is a closed (compact) subset of  $\text{Sub}_G$  and  $\{HK \mid H \in \text{Sub}_G^p\}$  is a closed (compact) subset of  $\text{Sub}_G^p$ , and they are  $T$ -invariant if  $T(K) = K$ , where  $T \in \text{Aut}(G)$ .
- (2)  $\mathcal{P}_1(G/K) = \{\pi(A) \mid A \in \mathcal{P}_1(G)\}$ .
- (3)  $\text{Sub}_{G/K}^p = \{\pi(H) \mid H \in \text{Sub}_G^p\}$ .
- (4)  $\text{Sub}_{G/K}^p$  is homeomorphic to  $\{HK \mid H \in \text{Sub}_G^p\}$ , and the latter is the same set as  $\{H \in \text{Sub}_G^p \mid HK = KH = H\}$ .

*Proof.* Let  $K$  and  $\pi$  be as above. Since  $K$  is compact and central in  $G$ ,  $HK$  is a closed subgroup of  $G$  for any closed subgroup  $H$  of  $G$ . Also, if  $H_n K \rightarrow L$  in  $\text{Sub}_G$ , then since  $\text{Sub}_G$  and  $K$  are compact, it is easy to see that for any limit point  $H$  of  $\{H_n\}$ ,  $L = HK$ . Therefore,  $\{HK \mid H \in \text{Sub}_G\}$  is closed. Now if  $H \in \text{Sub}_G^p$ , then there exists  $\{A_n\} \subset \mathcal{P}_1(G)$ , such that  $A_n \rightarrow H$ . By Lemma 2.3 (2),  $A_n K \in \text{Sub}_G^p$ . As  $A_n K \rightarrow HK$  and  $\text{Sub}_G^p$  is closed, we get that  $HK \in \text{Sub}_G^p$ . Now arguing as above, it is easy to show that  $\{HK \mid H \in \text{Sub}_G^p\}$  is a closed subset of  $\text{Sub}_G^p$ . Moreover, if  $T \in \text{Aut}(G)$  with  $T(K) = K$ , then

$T(HK) = T(H)K$  for any  $H \in \text{Sub}_G$ . Therefore, the sets mentioned in (1) are invariant under the action of  $\text{Aut}(G)$  if  $T(K) = K$ . Thus (1) holds.

If  $A \in \mathcal{P}_1(G)$ , then  $AK$  is closed and  $\pi(A) \in \mathcal{P}_1(G/K)$ . Conversely, suppose  $B \in \mathcal{P}_1(G/K)$ . It is easy to see that  $B_1 := \pi^{-1}(B)$  is a closed connected nilpotent Lie subgroup of  $G$ , and any one-parameter subgroup of  $B_1$ , which not contained in  $K$ , projects onto  $B$ . Let  $U$  be an open neighbourhood of the identity  $e$  in  $B_1$  such that every  $x \in U$  is contained in a one-parameter subgroup. As  $B_1$  is connected,  $U \neq K$  and  $U \setminus K$  is open. Suppose  $B_1$  is compact. Being connected and nilpotent,  $B_1$  is abelian. Since the set of roots of unity in  $B_1$  is dense and  $U \setminus K$  is open in  $B_1$ , we can choose  $x \in U \setminus K$  such that  $x^n = e$  for some  $n \in \mathbb{N}$ . If  $B_1$  is unbounded, then let  $x \in U \setminus K$  be any element. Let  $\{x_t\} \subset B_1$  be a one-parameter subgroup in  $B_1$  such that  $x_1 = x$ . Then  $\pi(\{x_t\}) = \{x_t\}K/K = B$ . If  $B_1$  is compact, then  $\{x_t\}$  is closed as  $x_1^n = e$ . If  $B_1$  is unbounded, then so is  $\{x_t\}$ , and hence it is closed. Therefore,  $\mathcal{P}_1(G/K) = \{\pi(A) \mid A \in \mathcal{P}_1(G)\}$ , i.e. (2) holds.

Let  $H \in \text{Sub}_G^p$ . Then  $A_n \rightarrow H$  for some  $\{A_n\} \subset \mathcal{P}_1(G)$ . By (2), we have that  $\{\pi(A_n)\} \subset \mathcal{P}_1(G/K)$ , and as  $K$  is compact,  $\pi(A_n) \rightarrow \pi(H)$ . Therefore,  $\pi(H) \in \text{Sub}_{G/K}^p$ . Conversely, let  $B \in \text{Sub}_{G/K}^p$ . There exist  $B_n \in \mathcal{P}_1(G/K)$ ,  $n \in \mathbb{N}$ , such that  $B_n \rightarrow B$ . By (2), there exist  $A_n \in \mathcal{P}_1(G)$  such that  $\pi(A_n) = B_n$ ,  $n \in \mathbb{N}$ . Let  $A$  be a limit point of  $\{A_n\}$  in  $\text{Sub}_G$ . Then  $A \in \text{Sub}_G^p$  and  $B = \pi(A)$ . Thus (3) holds.

Let  $\mathcal{S}(K) = \{H \in \text{Sub}_G^p \mid HK = KH = H\}$ . It is easy to show that  $\mathcal{S}(K)$  is closed in  $\text{Sub}_G^p$ , and hence it is compact. Observe that the set  $\mathcal{S}(K)$  is same as the set  $\{HK \mid H \in \text{Sub}_G^p\}$ . For  $H_1, H_2 \in \text{Sub}_G$ , if  $\pi(H_1) = \pi(H_2)$ , then  $H_1K = H_2K$ . Therefore,  $H \mapsto \pi(H)$  from  $\mathcal{S}(K)$  to  $\text{Sub}_{G/K}^p$  is injective, and it easily follows from (3) that it is surjective. Also if  $H_n \rightarrow H$  in  $\text{Sub}_G$ , then  $\pi(H_n) \rightarrow \pi(H)$  in  $\text{Sub}_{G/K}$ . Since  $\mathcal{S}(K)$  is compact and  $\text{Sub}_{G/K}^p$  is Hausdorff, we have that the map induced by  $\pi$  from  $\mathcal{S}(K)$  to  $\text{Sub}_{G/K}^p$  is a homeomorphism. Thus (4) holds.  $\square$

### 3. CHARACTERISATION OF DISTAL ACTIONS ON THE SPACE OF ONE-PARAMETER SUBGROUPS

In this section we discuss the action of an automorphism as well as that of a subgroup of automorphisms of a connected Lie group  $G$  on  $\text{Sub}_G^p$  and prove results stated in the introduction. We first set some notations and discuss some preliminary results which are useful.

For convenience, we define a class  $\mathcal{C}'$  of Lie groups as follows:

**Definition 3.1.** *A connected Lie group  $G$  is said to be in class  $\mathcal{C}'$  if  $G$  has no compact central subgroup of positive dimension (i.e. if  $G$  has no non-trivial compact connected central subgroup).*

The notation is inspired by the fact that the class  $\mathcal{C}'$  is a subset of the class  $\mathcal{C}$  defined by Dani and McCrudden in [13]. The latter consists of connected Lie groups which admit a (real) linear representation with discrete kernel. Proposition 2.5 of [13] shows that  $G$  is in class  $\mathcal{C}$  if and only if for the radical  $R$  of  $G$ , the intersection of  $\overline{[R, R]}$  with the center of  $G$  has no non-trivial compact subgroup. Therefore class  $\mathcal{C}'$  is a subset of class  $\mathcal{C}$ . Note that any connected Lie group  $G$  admits a maximal compact connected central subgroup  $M$ , and  $G/M$  has no non-trivial compact central subgroup and it belongs to class  $\mathcal{C}'$ .

Recall that for a connected Lie group  $G$  with the Lie algebra  $\mathfrak{g}$ , there is an exponential map  $\exp : \mathfrak{g} \rightarrow G$ , and for any  $T \in \text{Aut}(G)$ , there is a Lie algebra automorphism  $dT : \mathfrak{g} \rightarrow \mathfrak{g}$  such that  $\exp((dT)(X)) = T(\exp X)$ ,  $X \in \mathfrak{g}$ , and  $T \mapsto dT$  is a homomorphism and homeomorphism from  $\text{Aut}(G)$  onto its image in  $\text{GL}(\mathfrak{g})$  which is closed; the image consists of Lie algebra automorphisms of  $\mathfrak{g}$ . If  $G$  is in class  $\mathcal{C}'$ , then  $\text{Aut}(G)$  is almost algebraic as a subgroup of  $\text{GL}(\mathfrak{g})$ , it is closed and has finitely many connected components (cf. [12]). If  $H$  is a closed connected normal subgroup of  $G$  with the Lie algebra  $\mathfrak{h}$  which is a Lie subalgebra of  $\mathfrak{g}$  and it is an ideal in  $\mathfrak{g}$ . Moreover, the Lie algebra of  $G/H$  is isomorphic to  $\mathfrak{g}/\mathfrak{h}$ . Also, if  $H$  is  $T$ -invariant and if  $\overline{T}$  denotes the automorphism of  $G/H$  induced by  $T$ . Then  $d\overline{T} = \overline{dT}$ , which is the Lie algebra automorphism on  $\mathfrak{g}/\mathfrak{h}$  induced by  $dT$ . Also  $dT|_H$  is the same as  $(dT)|_{\mathfrak{h}}$ . Note that the eigenvalues of  $d\overline{T}$  (as well as those of  $dT|_H$ ) are eigenvalues of  $dT$ .

It is known that  $T$  acts distally on  $G$  if and only if  $dT$  acts distally on  $\mathfrak{g}$ , and the latter statement is equivalent to the statement that all the eigenvalues of  $dT$  have absolute value 1 (cf [2]). It is also known that if  $G$  admits an automorphism  $T$  such that  $dT$  has no eigenvalue of absolute value 1 (equivalently, if  $T$  is expansive; see [37], see also [7]), then  $G$  must be nilpotent (this is well-known, see e.g. [37]). We will need the following lemma which is slightly stronger. The lemma may be known, but we give a short proof for the sake of completeness.

**Lemma 3.2.** *Let  $G$  be a connected non-nilpotent Lie group and let  $T$  be an automorphism of  $G$ . Then at least one eigenvalue of  $dT$  is a root of unity.*

*Proof.* Let  $M$  be the largest compact connected central subgroup of  $G$ . Then  $G/M$  is in class  $\mathcal{C}'$ . As  $G$  is not nilpotent,  $G/M$  is also not nilpotent. Since  $M$  is characteristic in  $G$ , we may replace  $G$  by  $G/M$  and assume that  $G \in \mathcal{C}'$ , i.e. it has no compact central subgroup of positive dimension. If  $T$  has finite order, then  $(dT)^n = \text{Id}$  for some  $n \in \mathbb{N}$ , and the assertion follows in this case. Now suppose  $T$  does not have finite order.

Since  $G \in \mathcal{C}'$ ,  $\text{Aut}(G)$  is almost algebraic (cf. [12]), and hence it has finitely many connected components. Now we may replace  $T$  by its suitable power and assume that  $T \in \text{Aut}(G)^0$ , the connected component of the identity in  $\text{Aut}(G)$ , and that  $T$  is contained in a non-trivial one-parameter subgroup (say)  $\{T_t\}$  of  $\text{Aut}(G)^0$  as  $T = T_1$ . We will show that 1 is an eigenvalue of  $dT$ .

Suppose  $G$  is solvable. Let  $H = \{T_t\} \times G$ . Then  $H$  is solvable and the nilradical of  $H$ , being contained in  $G$ , is the same as the nilradical  $N$  of  $G$ . Therefore,  $H/N$  is abelian. Thus  $T$  acts trivially on  $G/N$ . As  $G$  is not nilpotent,  $G \neq N$ , and hence 1 is an eigenvalue  $dT$ .

Now suppose  $G$  is not solvable. Let  $R$  be the radical of  $G$ . Then  $R$  is characteristic in  $G$ , and let  $\overline{T}_t$  be the automorphisms of  $G/R$  induced by  $T_t$  for each  $t$ . It is enough to prove that 1 is an eigenvalue of  $d\overline{T}$ . Hence we may replace  $G$  by  $G/R$  and  $\{T_t\}$  by  $\{\overline{T}_t\}$  and assume that  $G$  is semisimple. Now  $\text{Aut}(G)^0$  consists of inner automorphisms of  $G$  and  $T_t = \text{inn}(x_t)$ ,  $t \in \mathbb{R}$ , for some one-parameter subgroup  $\{x_t\} \subset G$ . If  $x_1 = e$ , then  $T = T_1 = \text{Id}$ . Now suppose  $x_1 \neq e$ . Then  $\{x_t\}$  is non-trivial. Let  $X \in \mathfrak{g}$  be such that  $\exp tX = x_t$ ,  $t \in \mathbb{R}$ . As  $T(x_t) = x_t$ ,  $t \in \mathbb{R}$ , we have that  $dT(X) = X$  and 1 is an eigenvalue of  $dT$ .  $\square$

**Remark 3.3.** *The proof above also shows that for a connected solvable Lie group  $G$  with the nilradical  $N$ , for any  $T \in \text{Aut}(G)$ ,  $\overline{T}$  has finite order, where  $\overline{T}$  is the automorphism of  $G/N$  induced by  $T$ . In particular, all the eigenvalues of  $d\overline{T}$  on the Lie algebra of  $G/N$  are roots of unity.*

Note that the largest compact central subgroup and its connected component of the identity in a connected Lie group are characteristic.

We note the following useful proposition which will enable us to work on Lie groups in class  $\mathcal{C}'$ .

**Proposition 3.4.** *Let  $G$  be a connected Lie group and let  $T \in \text{Aut}(G)$ . Let  $K$  be a  $T$ -invariant compact connected central subgroup of  $G$  and let  $\bar{T}$  denote the automorphism of  $G/K$  induced by  $T$ . If  $T$  acts distally on  $\text{Sub}_G^p$ , then  $\bar{T}$  acts distally on  $\text{Sub}_{G/K}^p$ .*

*Proof.* Let  $T \in \text{Aut}(G)$  and let  $\pi : G \rightarrow G/K$  be the natural projection. Note that  $\bar{T}(\pi(H)) = \pi(T(H))$  for any  $H \in \text{Sub}_G$  and  $\text{Sub}_{G/K}^p$  is  $\bar{T}$  invariant. Let  $\mathcal{S}(K) = \{H \in \text{Sub}_G^p \mid HK = H\}$  which is a compact  $T$ -invariant subset of  $\text{Sub}_G^p$  and homeomorphic to  $\text{Sub}_{G/K}^p$  (cf. Lemma 2.4). Suppose  $T$  acts distally on  $\text{Sub}_G^p$ . Then the  $T$ -action on  $\mathcal{S}(K)$  is distal. Let  $L_1, L_2 \in \text{Sub}_{G/K}^p$  be such that for some  $\{n_k\} \subset \mathbb{Z}$ ,  $\bar{T}^{n_k}(L_i) \rightarrow L$ , for some  $L \in \text{Sub}_{G/K}^p$ ,  $i = 1, 2$ . Then there exist  $H_1, H_2 \in \mathcal{S}(K)$ , such that  $\pi(H_i) = L_i$  and  $T^{n_k}(H_i) \rightarrow \pi^{-1}(L)$  in  $\mathcal{S}(K)$ ,  $i = 1, 2$  (cf. Lemma 2.4(4)). As  $T$  acts distally on  $\mathcal{S}(K)$ , we get that  $H_1 = H_2$ , and hence that  $L_1 = L_2$ . Thus  $\bar{T}$  acts distally on  $\text{Sub}_{G/K}^p$ .  $\square$

We also need the following elementary lemma.

**Lemma 3.5.** *Let  $G$  be a simply connected nilpotent Lie group with the Lie algebra  $\mathfrak{g}$  and let  $T \in \text{Aut}(G)$ . Then the set  $\mathcal{P}_1(G)$  consisting of all closed one-parameter subgroups of  $G$  is closed in  $\text{Sub}_G$  and is the same as  $\text{Sub}_G^p$ . Moreover,  $T$  acts distally on  $\text{Sub}_G^p$  if and only if  $dT$  acts distally on  $\text{Sub}_{\mathfrak{g}}^p$ .*

*Proof.* In a simply connected nilpotent group, all one-parameter subgroups are closed and there is a one-to-one correspondence between non-trivial (closed) one-parameter subgroups of  $G$  and one-dimensional subspaces of  $\mathfrak{g}$ . The correspondence is via the exponential map  $\exp : \mathfrak{g} \rightarrow G$  with  $\log$  as its inverse. Namely, for  $\{x_t\}$ ,  $x_t = \exp tX$ , for some  $X \in \mathfrak{g}$ , and  $\log x_t = tX$ ,  $t \in \mathbb{R}$ . Suppose  $\{H_n\} \subset \mathcal{P}_1(G)$  is such that  $H_n \rightarrow H$  in  $\text{Sub}_G$ . Then  $\log H_n \in \mathcal{P}_1(\mathfrak{g})$  and using Lemma 2.1 and the continuity of the exponential map and its inverse, it is easy to show that  $\{\log H_n\}$  converges to (say)  $B$  in  $\text{Sub}_{\mathfrak{g}}^p$ . and since  $\mathfrak{g}$  is isomorphic to  $\mathbb{R}^n$  for some  $n$ ,  $\text{Sub}_{\mathfrak{g}}^p = \mathcal{P}_1(\mathfrak{g})$  which is closed and we have that  $B \in \mathcal{P}_1(\mathfrak{g})$ . Then  $H = \exp B \in \mathcal{P}_1(G)$  and, in particular, we get that  $\mathcal{P}_1(G)$  is closed in  $\text{Sub}_G$ , i.e.  $\mathcal{P}_1(G) = \text{Sub}_G^p$ . In particular,  $H_n \rightarrow H$  in  $\text{Sub}_G^p$  implies that  $\log H_n \rightarrow \log H$  in  $\text{Sub}_{\mathfrak{g}}^p$ .

Conversely, if  $L_n \rightarrow L$  in  $\mathcal{P}_1(\mathfrak{g})$ , using Lemma 2.1 and the fact that  $\exp$  is a homeomorphism from  $\mathfrak{g}$  to  $G$ , it is easy to see that  $\exp L_n \rightarrow \exp L$  in  $\mathcal{P}_1(G)$ . Now the exponential map defines a homeomorphism from  $\mathcal{P}_1(\mathfrak{g})$  to  $\mathcal{P}_1(G)$  and  $T$  (resp.  $dT$ ) defines a homeomorphism of  $\mathcal{P}_1(G)$  (resp.  $\mathcal{P}_1(\mathfrak{g})$ ) such that  $\exp \circ dT = T \circ \exp$ . Therefore, it is easy to see that  $T$  acts distally on  $\mathcal{P}_1(G) = \text{Sub}_G^p$  if and only if  $dT$  acts distally on  $\mathcal{P}_1(\mathfrak{g}) = \text{Sub}_{\mathfrak{g}}^p$ .  $\square$

*Proof of Theorem 1.1 (1) and (2) for nilpotent groups.* Let  $G$  be a (non-trivial) connected Lie group in class  $\mathcal{C}'$ , i.e.  $G$  does not have any non-trivial compact connected central subgroup. Let  $T \in \text{Aut}(G)$ .

(1) : Suppose  $G$  is a connected abelian group in class  $\mathcal{C}'$ , i.e.  $G$  does not have any non-trivial compact subgroup, and it is a vector group isomorphic to  $\mathbb{R}^d$ , for some  $d \in \mathbb{N}$ . Note that  $\mathcal{D} = \{r \text{ Id} \mid r \in \mathbb{R} \setminus \{0\}\}$ , where  $\text{Id}$  is the identity matrix in  $\text{GL}(d, \mathbb{R})$ . If  $d = 1$ , then  $\text{GL}(1, \mathbb{R})$  is abelian, and the assertion follows trivially. Now suppose  $d \geq 2$ . Then  $\text{Sub}_G^p$  is isomorphic to  $\mathbb{R}\mathbb{P}^{d-1} \sqcup \{\{e\}\}$  and  $T \in \text{GL}(d, \mathbb{R})$ . Both  $\mathbb{R}\mathbb{P}^{d-1}$  and  $\{e\}$  are  $T$  invariant. Hence  $T$  acts distally on  $\text{Sub}_{\mathbb{R}^d}^p$  if and only if  $T$  acts distally on  $\mathbb{R}\mathbb{P}^{d-1}$ . One can define an action of  $T$  on  $\mathbb{S}^{d-1}$ , the boundary of unit sphere in  $\mathbb{R}^d$ , as  $\overline{T}(x) = T(x)/\|x\|$ , where  $\|\cdot\|$  denotes the usual norm on  $\mathbb{R}^d$ . Here,  $\overline{T}$  is a homeomorphism and  $\overline{T}(-x) = -\overline{T}(x)$ . As  $\mathbb{R}\mathbb{P}^{d-1}$  is a quotient space of  $\mathbb{S}^{d-1}$ , the action of  $T$  on  $\mathbb{R}\mathbb{P}^{d-1}$  is the same as the action of  $\overline{T}$  on  $\mathbb{S}^{d-1}/\sim$ , where  $\sim$  is the equivalence relation which identifies  $x$  with  $-x$  for each  $x \in \mathbb{S}^{d-1}$ . It is easy to see that  $T$  acts distally on  $\mathbb{R}\mathbb{P}^{d-1}$  if and only if  $\overline{T}$  acts distally on  $\mathbb{S}^{d-1}$ . The latter statement is equivalent to the following:  $T \in \mathcal{KD}$ , where  $\mathcal{K}$  is a compact subgroup of  $\text{GL}(d, \mathbb{R})$  and  $\mathcal{D}$  is the center of  $\text{GL}(d, \mathbb{R})$ , (cf. [38], Theorem 1).

Conversely, for  $G = \mathbb{R}^d$  and  $T \in \mathcal{KD}$ , with a compact set  $\mathcal{K} \subset \text{GL}(d, \mathbb{R})$  and the center  $\mathcal{D}$  of  $\text{GL}(d, \mathbb{R})$ , the action of  $T$  on  $\text{Sub}_G^p$  is same as that of  $S$  on  $\text{Sub}_G^p$  for some  $S \in \mathcal{K}$  since  $\mathcal{D}$  acts trivially on  $\text{Sub}_G^p$ . As the compact subgroups of  $\text{Aut}(G)$  act distally on any closed invariant subspace of  $\text{Sub}_G^p$  ( $\text{Sub}_G^p$  in this case), we have that  $T$  acts distally on  $\text{Sub}_G^p$ . This completes the proof of Theorem 1.1 (1).

(2) : Suppose  $G$  belongs to the class  $\mathcal{C}'$  and it is non-abelian and nilpotent. Then  $G$  is simply connected as  $G$  has no compact central subgroup of positive dimension. Suppose  $T$  acts distally on  $\text{Sub}_G^p$ . We want to show that  $T$  is contained in a compact subgroup of  $\text{Aut}(G)$ . Since



$G$  belongs to the class  $\mathcal{C}'$ ,  $\text{Aut}(G)$  is almost algebraic as a subgroup of  $\text{GL}(\mathfrak{g})$  (cf. [12]).

As  $G$  is simply connected and nilpotent, by Lemma 3.5,  $dT$  acts distally on  $\text{Sub}_{\mathfrak{g}}^p$ . Then from (1) above, we have that  $dT \in \mathcal{K}\{r\text{Id} \mid r \neq 0\}$  for some compact subgroup  $\mathcal{K}$  of  $\text{GL}(\mathfrak{g})$  and  $\text{Id}$  is the identity map on  $\mathfrak{g}$ . As  $dT^n = \psi \circ r\text{Id}$  for some  $r \in \mathbb{R} \setminus \{0\}$  and  $\psi \in \mathcal{K}$ . If  $r = \pm 1$ , then  $dT^2 \in \mathcal{K}$  and  $T$  generates a relatively compact subgroup of  $\text{Aut}(G)$ . If possible suppose  $r \neq \pm 1$ . Since  $\text{Aut}(G)$  is almost algebraic, there exists  $n \in \mathbb{N}$  such that  $dT^n$  is contained in a connected almost algebraic subgroup  $\mathbb{R} \times \mathbb{T}^k$  of  $\text{Aut}(G)^0$  (which is identified with its image in  $\text{GL}(\mathfrak{g})$ ), where  $\mathbb{T}^k = (\mathbb{S}^1)^k$  for some  $k \in \mathbb{N}$ . Therefore,  $\psi^n \circ r^n \text{Id} \in \mathbb{R} \times \mathbb{T}^k \subset \text{Aut}(G)^0$ . Hence both  $\psi^n$  and  $r^n \text{Id} \in \text{Aut}(G)^0$ . But  $r^n \text{Id}$  is not a Lie algebra automorphism of  $\mathfrak{g}$  as  $r^n \neq 1$ , for if  $r^n[X, Y] = r^n \text{Id}[X, Y] = [r^n \text{Id}(X), r^n \text{Id}(Y)] = r^{2n}[X, Y]$ , then  $[X, Y] = 0$  for all  $X, Y \in \mathfrak{g}$ , which leads to a contradiction as  $G$  is not abelian. Thus,  $r = \pm 1$  and  $T$  is contained in a compact subgroup of  $\text{Aut}(G)$ . The converse statement is obvious. Thus (2) holds if  $G$  is nilpotent.  $\square$

We will continue with the proof of Theorem 1.1 after proving Theorems 1.2 and 1.3 and some elementary results.

For a Lie group  $G$  and  $T \in \text{Aut}(G)$ , let  $C(T)$  denote the contraction group of  $T$  and it is defined as

$$C(T) := \{x \in G \mid T^n(x) \rightarrow e \text{ as } n \rightarrow \infty\}.$$

Note that  $C(T)$  is a connected nilpotent subgroup of  $G$  and it is simply connected if it is closed in  $G$  (see [41], see also [20]). Both  $C(T)$  and  $C(T^{-1})$  are trivial if and only if  $T$  acts distally on  $G$  (cf. [34], Theorem 4.1). For a connected Lie group  $G$ ,  $C(T)$  is trivial if and only if all the eigenvalues of  $dT$  have absolute value greater than or equal to 1 ([34, 2]).

We need the following lemma which can be proven easily by using Theorem 1.1 of [14], Theorem 2.4 and Corollary 2.7 of [20]. Note that the largest compact connected central subgroup of  $G$  is characteristic in  $G$ . The lemma generalises a part of the statement in Proposition 4.3 of [34].

**Lemma 3.6.** *Let  $G$  be a connected Lie group and let  $T \in \text{Aut}(G)$ . Then the following are equivalent:*

- (1)  $T$  acts distally on the largest compact connected central subgroup of  $G$ .
- (2) The contraction groups  $C(T)$  and  $C(T^{-1})$  are closed, simply connected and nilpotent.
- (3)  $T$  acts distally on every compact  $T$ -invariant subgroup of  $G$ .

*Proof.* Let  $M$  be the largest compact connected central subgroup of  $G$ . Let  $\pi : G \rightarrow G/M$  be the natural projection and let  $T' \in \text{Aut}(G/M)$  be the automorphism induced by  $T$  on  $G/M$ . As  $G/M$  does not have any non-trivial compact connected central subgroup, using Theorem 1.1 of [14], one can show that  $C(T')$  is closed (see the proof of Proposition 4.3 of [34] for more details). Hence  $\pi^{-1}(C(T')) = C_M(T) := \{x \in G \mid T^n(x)M \rightarrow M \text{ in } G/M\}$  is closed. Suppose (1) holds, i.e.  $T$  acts distally on  $M$ . Then  $C(T) \cap M = C(T|_M) = \{e\}$ . As  $C_M(T)$  is closed, by Corollary 2.7 of [20],  $C(T)$  is closed. Similarly,  $C(T^{-1})$  is closed. The rest of the assertion in (2) follows from Corollary 2.4 in [41]. This proves (1)  $\implies$  (2).

Now suppose (2) holds, i.e.  $C(T)$  is closed. Let  $K$  be any compact  $T$ -invariant subgroup of  $G$ . Then  $C(T|_K) = C(T) \cap K$  is a compact subgroup of  $C(T)$ . Hence  $C(T|_K)$  is trivial (cf. [20], Corollary 2.5). Similarly,  $C((T|_K)^{-1})$  is trivial and  $T$  acts distally on  $K$  (cf. [25]). Thus (3) holds. (3)  $\implies$  (1) is obvious.  $\square$

*Proof of Theorem 1.2.* Let  $G$  be a connected Lie group which is not a vector group. Let  $M$  be the largest compact connected central subgroup of  $G$ . Suppose  $T \in \text{Aut}(G)$  be such that it acts distally on  $M$ . Suppose  $T$  acts distally on  $\text{Sub}_G^p$ . Then we want to show that  $T$  acts distally on  $G$ .

**Step 1:** Suppose  $G$  is simply connected and nilpotent. Since  $G$  is not a vector group,  $G$  is non-abelian and by the proof of Theorem 1.1 (2) for the nilpotent case,  $T$  is contained in a compact subgroup of  $\text{Aut}(G)$ . Hence  $T$  acts distally on  $G$ .

**Step 2:** Now suppose  $G$  is not a simply connected nilpotent group. By Theorem 4.1 of [34], it is enough to prove that the contraction groups  $C(T)$  and  $C(T^{-1})$  are trivial. As  $T$  acts distally on the maximal compact connected central subgroup  $M$ , by Lemma 3.6, both  $C(T)$  and  $C(T^{-1})$  are closed, simply connected and nilpotent. Suppose  $C(T)$  is non-trivial. If possible, suppose  $C(T)$  is non-abelian. As it is  $T$ -invariant and  $T|_{C(T)}$  acts distally on  $\text{Sub}_{C(T)}^p$ , we get from Step 1 that

$T|_{C(T)}$  acts distally on  $C(T)$ , a contradiction. Now suppose  $C(T)$  is abelian. Then  $C(T)$  is isomorphic to  $\mathbb{R}^d$  as  $C(T)$  is simply connected. By Theorem 1.1 (1),  $T_1 = T|_{C(T)} = \phi \circ \alpha I$  for some  $\phi$  contained in a compact subgroup of  $\mathrm{GL}(d, \mathbb{R})$  and  $\alpha \in \mathbb{R}$ ,  $0 < |\alpha| < 1$ ; here  $I$  denotes the identity matrix in  $\mathrm{GL}(d, \mathbb{R})$ . We may replace  $T$  by  $T^2$  (and  $T_1$  by  $T_1^2$ ), and assume that  $\alpha > 0$ . Here,  $G \neq C(T)$  as  $G$  is not a vector group.

Suppose  $G$  is nilpotent but not simply connected. Then  $M$  as above is non-trivial. Since  $T$  acts distally on  $M$ , by Lemma 2.5 of [2], we have that  $T^k|_M$  is unipotent for some  $k \in \mathbb{N}$ , i.e.  $dT$  has an eigenvalue which is a root of unity. If  $G$  is not nilpotent, by Lemma 3.2,  $dT$  has an eigenvalue which is a root of unity. We may replace  $T$  by a suitable power and assume that 1 is an eigenvalue of  $dT$ .

Let  $X \in \mathfrak{g}$ ,  $X \neq 0$  be such that  $dT(X) = X$  and let  $A = \{\exp tX\}$ , a one-parameter subgroup, it is either closed or its closure  $\bar{A}$  is compact. Since  $T|_A = \mathrm{Id}$ , we have that  $T|_{\bar{A}} = \mathrm{Id}$  and it follows that  $\bar{A}$  normalises  $C(T)$  and  $\bar{A} \cap C(T) = \{e\}$ . Let  $H = \overline{A \times C(T)}$ . Then  $H$  is solvable. As  $A$  normalises  $C(T)$ , so does  $H$ . Now  $H/C(T)$  is connected and abelian as it is the closure of the image of the one-parameter subgroup  $A$  in it. Then  $H/C(T)$  is either isomorphic to  $\mathbb{R}$  or it is compact. In the first case it follows that  $H = A \times C(T)$ . In the second case, since  $C(T)$  is a vector group,  $H = K \times C(T)$  for some (maximal) compact subgroup  $K \subset H$  (cf. [24], Lemma 3.7). Here,  $H$  is a closed  $T$ -invariant subgroup of  $G$  and  $T(h) \in hC(T)$  for all  $h \in H$ . If  $A$  (resp.  $\bar{A}$ ) is compact, then  $H = A \times C(T)$  (resp.  $H = \bar{A} \times C(T)$ ). Note that  $C(T)$  is isomorphic to  $\mathbb{R}^d$  and  $T_1$ , the restriction of  $T$  to  $C(T)$ , is as described earlier.

**Step 3:** Suppose  $H/C(T)$  is isomorphic to  $\mathbb{R}$ . Then so is  $A$  and  $H = A \times C(T)$ . Let  $U, V$  be open neighbourhoods of the identity  $e$  in  $H$  such that  $V^2 \subset U$  and every element of  $U$  is contained in a one-parameter subgroup. Let  $x = ac$  for some  $a \in A \cap V$  and  $c \in C(T) \cap V$  with  $c \neq e$ . Then  $x \in V^2 \subset U$  and  $T(x) = xy$  for  $y = c^{-1}T(c) \in C(T)$  and  $y \neq e$  as  $c \in C(T)$  and  $c \neq e$ . Now  $x$  is contained in a one-parameter subgroup (say)  $\{x_t\}$  in  $H$  as  $x = x_1$ . Then  $T(x_t) = x_t y_t$ ,  $y_t \in C(T)$ ,  $t \in \mathbb{R}$  and  $y_1 = y \neq e$ . Here,  $\{x_t\}$  is closed as  $H$  is simply connected and solvable. As  $\mathrm{Sub}_H^p$  is compact, we get that  $T^{n_k}(\{x_t\}) \rightarrow B$  for some  $B \in \mathrm{Sub}_H^p$  and some unbounded sequence  $\{n_k\} \subset \mathbb{N}$ . By Lemma 2.2,  $B \neq \{e\}$ . Let  $b \in B \setminus \{e\}$ . Then  $T^{n_k}(x_{t_k}) \rightarrow b$  for some sequence  $\{t_k\} \subset \mathbb{R}$ . Here,  $\{x_t\}C(T)/C(T) = H/C(T)$  is isomorphic to  $\mathbb{R}$ . Since

$x_{t_k}C(T) \rightarrow bC(T)$  in  $H/C(T)$ , we get that  $\{t_k\}$  is bounded. Passing to a subsequence if necessary, we may assume that  $t_k \rightarrow t_0$ . Then  $x_{t_k} \rightarrow x_{t_0}$  and  $y_{t_k} \rightarrow y_{t_0}$ . As all the eigenvalues of  $T|_{C(T)} = T_1 = \phi \circ \alpha I$  have absolute value  $\alpha$  with  $0 < \alpha < 1$ , we get that  $I - T_1$  is invertible in  $\text{GL}(d, \mathbb{R})$ , where  $I$  is the identity matrix in  $\text{GL}(d, \mathbb{R})$ . Using the addition as the group operation in  $\mathbb{R}^d$ , and the fact that  $T_1^n(y) \rightarrow 0$  for all  $y \in \mathbb{R}^d$ , we get that

$$T^{n_k}(x_{t_k}) = x_{t_k} \left( \sum_{i=0}^{n_k-1} T_1^i(y_{t_k}) \right) \rightarrow x_{t_0}(I - T_1)^{-1}(y_{t_0}) = b$$

(as  $n_k \rightarrow \infty$ ), where  $T_1^0 = I$ , the identity in  $\text{GL}(d, \mathbb{R})$ . Note that

$$I + T_1(I - T_1)^{-1} = (I - T_1)(I - T_1)^{-1} + T_1(I - T_1)^{-1} = (I - T_1)^{-1}.$$

Therefore,

$$T(b) = x_{t_0}y_{t_0}T_1(I - T_1)^{-1}(y_{t_0}) = x_{t_0}(I + T_1(I - T_1)^{-1})(y_{t_0}) = b.$$

Thus  $T(B) = B$ . Now we get that  $T^{n_k}(T(\{x_t\})) \rightarrow T(B) = B$  and hence that  $T(\{x_t\}) = \{x_t\}$  since  $T$  acts distally on  $\text{Sub}_G^p$ . As  $T$  is an automorphism, we have that  $T(x_t) = x_{st}$  for some  $s \in \mathbb{R} \setminus \{0\}$ , and  $xy = x_1y_1 = x_s$ . Therefore,  $x_{s-1} = y_1 = y \neq e$  and  $s \neq 1$ . Then  $x_{s-1} \in C(T)$ , and  $\{x_t\}C(T)/C(T)$  is compact. This leads to a contradiction as  $H = \{x_t\}C(T)$  and  $H/C(T)$  is isomorphic to  $\mathbb{R}$ .

**Step 4:** Now we have that  $H/C(T)$  is compact and  $H = K \times C(T)$ , where  $K$  is a maximal compact subgroup of  $H$ . We show that  $T(x) = x$  for all  $x \in K$ . We know that  $T(x) \in xC(T)$  for all  $x \in K$ . Since  $K$  is connected and abelian, the set  $F$  of roots of unity in  $K$  is dense in  $K$  and it is enough to show that  $T(x) = x$  for all  $x \in F$ . If possible, suppose there exists  $x \in F$  such that  $T(x) \neq x$ . Let  $m \in \mathbb{N}$  be the order of  $x$ , i.e.  $x^m = e$  and  $m \neq 1$ . Let  $\{x_t\}$  be a one-parameter subgroup in  $K$  such that  $x_1 = x$ . As  $x_1^m = e$ ,  $\{x_t\}$  is closed and isomorphic to  $\mathbb{S}^1$  and  $T(x_t) = x_t y_t$ , where  $y_t \in C(T)$ ,  $t \in \mathbb{R}$  and  $y_1 = x^{-1}T(x) \neq e$ . There exists an unbounded sequence  $\{n_k\} \subset \mathbb{N}$  such that  $T^{n_k}(\{x_t\}) \rightarrow B$  for some  $B \subset H$ . Then  $B \neq \{e\}$  as  $x_1 \neq e$  and  $\{x_t\}$  is connected (cf. Lemma 2.2). Let  $b \in B$  be such that  $b \neq e$ . Then  $T^{n_k}(x_{t_k}) \rightarrow b$  for some  $\{t_k\} \subset \mathbb{R}$ . Here,  $\{x_t\}$  is periodic i.e.  $x_{mn} = e$  and  $x_{m+t} = x_t$  for all  $n \in \mathbb{Z}$  and  $t \in \mathbb{R}$ , where  $m$  is the order of  $x_1 = x$ . Hence we can choose  $\{t_k\}$  such that it is bounded. Then  $\{x_{t_k}\}$ , and hence,  $\{y_{t_k}\}$  is bounded. Now arguing as in Step 3, we get that  $T(b) = b$ . Therefore,  $T(B) = B$ ,

and hence  $T^{n_k}(T(\{x_t\})) \rightarrow B$ . Since  $T$  acts distally on  $\text{Sub}_G^p$ , we get that  $T(\{x_t\}) = \{x_t\}$ . As  $T$  is an automorphism and  $\{x_t\}$  is isomorphic to  $\mathbb{S}^1$ , we get that  $T(x_t) = x_t$  or  $T(x_t) = x_t^{-1} = x_{-t}$ . In either case, we get that  $y_1 \in K \cap C(T) = \{e\}$ . This leads to a contradiction. Hence  $T(x) = x$ , for all  $x \in F$ , and hence for all  $x \in K$ .

**Step 5:** We now show that  $K$  centralises  $C(T)$  and  $H = K \times C(T)$ . We have that  $T|_K = \text{Id}$ . Let  $K_c = c^{-1}Kc$ ,  $c \in C(T)$ . Then  $K_c$  is a compact connected abelian group and  $H = K_c \times C(T)$  and  $T(x) \in xC(T)$  for all  $x \in K_c$ . Therefore, arguing as in Step 4 for  $K_c$  instead of  $K$ , we get that  $T|_{K_c} = \text{Id}$ , i.e.  $c^{-1}xc = T(c^{-1}xc) = T(c^{-1})xT(c)$  for all  $c \in C(T)$  and  $x \in K$ . Thus  $cT(c^{-1}) = (I - T_1)(c)$  centralises  $K$ ,  $c \in C(T)$ , where  $T_1 = T|_{C(T)}$ . As  $I - T_1$  is invertible in  $GL(d, \mathbb{R})$ , we get that  $C(T)$  centralises  $K$ . Thus  $H = K \times C(T)$ .

**Step 6:** Let  $\psi, S : H \rightarrow H$  be such that  $S|_K = \text{Id}$ ,  $S|_{C(T)} = \alpha \text{Id}$ ,  $\psi|_K = \text{Id}$  and  $\psi|_{C(T)} = \phi$ , where  $\alpha$  and  $\phi$  are as in Step 2. Now  $T|_H = S\psi = \psi S$ , and  $\psi$  is contained in a compact subgroup of  $\text{Aut}(H)$ . Therefore,  $S$  acts distally on  $\text{Sub}_H^p$ , and  $C(S) = C(T)$ . Using that  $C(T) = C(S) \neq \{e\}$ , we show that  $S$  does not act distally on  $\text{Sub}_H^p$ , which would lead to a contradiction.

Let  $\{z_t = x_t c_t\}$  be a closed one-parameter subgroup, where  $\{x_t\} \subset K$  is isomorphic to  $\mathbb{S}^1$  and  $\{c_t\} \subset C(S) = C(T)$  is a non-trivial (closed) one-parameter subgroup such that  $S(c_t) = c_{\alpha t}$ ,  $t \in \mathbb{R}$  (In additive notation on  $\mathbb{R}^d$ , we choose  $c_t = tc$ ,  $t \in \mathbb{R}$ , for some nonzero  $c \in \mathbb{R}^d$ ). Let  $\{n_k\}$  be an unbounded sequence in  $\mathbb{N}$  such that  $S^{n_k}(\{z_t\}) \rightarrow C$  for some  $C$  in  $\text{Sub}_H^p$ . Note that  $\{z_t\} \subset \{x_t\} \times \{c_t\}$  and the latter is  $S$ -invariant and isomorphic to  $\mathbb{S}^1 \times \mathbb{R}$ . So  $C \subset \{x_t\} \times \{c_t\}$ . For a fixed  $t \in \mathbb{R}$ , as  $c_t \in C(S) = C(T)$ , we get that

$$S^{n_k}(z_t) = x_t S^{n_k}(c_t) \rightarrow x_t,$$

as  $n_k \rightarrow \infty$ . Therefore,  $\{x_t\} \subset C$ . Again for a fixed  $t \in \mathbb{R}$ ,

$$S^{n_k}(z_{\alpha^{-n_k} t}) = x_{\alpha^{-n_k} t} S^{n_k}(c_{\alpha^{-n_k} t}) = x_{\alpha^{-n_k} t} c_t \rightarrow x_s c_t$$

along a subsequence of  $\{n_k\}$ , for some  $s \in \mathbb{R}$ . Therefore,  $x_s c_t$ , and hence,  $c_t$  belongs to  $C$ . Thus  $\{c_t\} \subset C$  and  $C = \{x_t\} \times \{c_t\}$ , and hence  $C$  is  $S$ -invariant. This implies that  $S^{n_k}(S(\{z_t\})) \rightarrow C$ , and hence that  $S(\{z_t\}) = \{z_t\}$ . As  $\{z_t\}$  is isomorphic to  $\mathbb{R}$ , there exists  $\beta \in \mathbb{R} \setminus \{0\}$

such that for all  $t \in \mathbb{R}$ ,

$$S(z_t) = z_{\beta t} = x_{\beta t} c_{\beta t} \quad \text{and} \quad S(x_t) = S(x_t c_t) = x_t c_{\alpha t}.$$

As  $\{x_t\} \times \{c_t\}$  is a direct product and  $\{c_t\}$  is isomorphic to  $\mathbb{R}$ , it follows that  $\beta = \alpha$  and  $x_t = x_{\alpha t}$ , i.e.  $x_{(1-\alpha)t} = e$  for all  $t \in \mathbb{R}$ . As  $\alpha \neq 1$ ,  $x_t = e$  for each  $t$ , a contradiction. Hence,  $C(S) = C(T)$  is trivial.

Similarly, we can show that  $C(T^{-1})$  is trivial. Thus  $T$  acts distally on  $G$ .  $\square$

The following corollary which extends Theorem 1.2 is an easy consequence of the theorem in view of Theorem 1.1 of [2]. We will omit the proof.

**Corollary 3.7.** *Let  $G$  be a connected Lie group which is not a vector group and let  $H$  be a subgroup of  $\text{Aut}(G)$  such that it acts distally on the largest compact connected central subgroup of  $G$ . If  $H$  acts distally on  $\text{Sub}_G^p$ , then  $H$  acts distally on  $G$ .*

Before proving Theorem 1.3, we first prove the following proposition which is a special case of the theorem.

**Proposition 3.8.** *Let  $G$  be a connected Lie group,  $T \in \text{Aut}(G)$  and let  $L$  be a closed connected nilpotent normal  $T$ -invariant subgroup. Suppose  $T$  acts trivially on both  $L$  and  $G/L$ . If  $T$  acts distally on  $\text{Sub}_G^p$ , then  $T = \text{Id}$ .*

*Proof.* Since  $T$  acts trivially on  $G/L$  and  $L$ , we have that  $T$  is distal. In fact, all the eigenvalues of  $T$  are equal to 1 and  $T$  is unipotent. Suppose  $T$  acts distally on  $\text{Sub}_G^p$ . Let  $U$  be a neighbourhood of the identity  $e$  such that every element of  $U$  is contained in a one-parameter subgroup. Since  $G$  is connected,  $U$  generates  $G$  and it is enough to show that  $T(x) = x$  for all  $x \in U$ .

**Step 1:** If possible, suppose  $T(x) \neq x$  for some  $x \in U$ . There exists a one-parameter subgroup  $\{x_t\}$  in  $G$  such that  $x_1 = x$ . Let  $H = \overline{\{x_t\}L}$ . Then  $H$  is connected solvable and  $H/L$  is abelian. Let  $\pi : G \rightarrow G/L$  be the natural projection. Then  $\pi(H) = \overline{\pi(\{x_t\})}$ . If  $\pi(\{x_t\})$  is unbounded, it is closed and it is isomorphic to  $\mathbb{R}$  and in this case  $\{x_t\}$  is also unbounded and closed. Moreover  $H = \{x_t\}L$ .

Now suppose  $\pi(\{x_t\})$  is relatively compact. As  $H$  is solvable, there exists a maximal compact (connected, abelian) subgroup  $K$  in  $H$  such that  $H = KL$ . As  $T(x) \neq x$  and  $T|_L = \text{Id}$ , we have that  $T(h) \neq h$  for

some  $h \in K$ , and hence  $T(h) \neq h$  for some finite order element  $h$  in  $K$ . There exists a closed (compact) one-parameter subgroup  $\{h_t\}$  in  $K$  such that  $h_1 = h$ . We may replace  $\{x_t\}$  by  $\{h_t\}$  and assume also in this case that  $H = \{x_t\}L$ .

Now we have that  $H = \{x_t\}L$  is closed, where either  $\pi(\{x_t\})$  (as well as  $\{x_t\}$ ) is isomorphic to  $\mathbb{R}$  or  $\{x_t\}$  is isomorphic to  $\mathbb{S}^1$  and  $T(x_1) \neq x_1$ . As  $T$  acts trivially on  $G/L$ , we have that  $T(H) = H$  and  $T(x_t) = x_t y_t$ , for some  $y_t \in L$ ,  $t \in \mathbb{R}$ , where  $y_1 \neq e$ .

**Step 2:** Let  $A := \{x_t\}$  and let  $\{n_k\} \subset \mathbb{N}$  be an unbounded sequence such that  $T^{n_k}(A) \rightarrow B$  for some  $B \in \text{Sub}_H^p$ . By Lemma 2.2,  $B$  is not discrete as  $A$  is connected and non-trivial. Let  $b \in B$  be such that  $b \neq e$ . There exists  $t_k \in \mathbb{R}$ ,  $k \in \mathbb{N}$ , such that

$$T^{n_k}(x_{t_k}) = x_{t_k} y_{t_k}^{n_k} \rightarrow b.$$

As  $y_t \in L$ ,  $t \in \mathbb{R}$ , we get that  $\pi(x_{t_k}) \rightarrow \pi(b)$ . If  $H/L$  is isomorphic to  $\mathbb{R}$ , then we get that the sequence  $\{t_k\}$  is bounded. In the second case where  $\{x_t\}$  is compact, we can choose  $\{t_k\}$  to be bounded. Passing to a subsequence if necessary, we may assume that  $t_k \rightarrow t_0$ . Therefore,  $x_{t_k} \rightarrow x_{t_0}$  and  $y_{t_k} \rightarrow y_{t_0}$  and  $b = x_{t_0} y'$ , where  $y_{t_k}^{n_k} \rightarrow y'$  in  $L$ .

**Step 3:** We first assume that  $L$  is simply connected. We show that  $y_{t_0} = e$  in this case. Let  $\mathfrak{l}$  be the Lie algebra of  $L$ . Since  $L$  is simply connected and nilpotent,  $\exp : \mathfrak{l} \rightarrow L$  is a homeomorphism with  $\log$  as its inverse. Let  $y_{t_k} = \exp Y_k$  for some  $Y_k \in \mathfrak{l}$ . Then  $y_{t_k}^{n_k} = \exp(n_k Y_k)$  and as  $y_{t_k}^{n_k} \rightarrow y'$ , we have that  $n_k Y_k \rightarrow \log y'$  in  $\mathfrak{l}$ . As  $\mathfrak{l}$  is isomorphic to  $\mathbb{R}^d$  for some  $d \in \mathbb{N}$ , it follows that  $Y_k \rightarrow 0$  in  $\mathfrak{l}$ , and hence  $y_{t_k} \rightarrow e$ , i.e.  $y_{t_0} = e$ . Now

$$T(b) = T(x_{t_0})y' = x_{t_0} y_{t_0} y' = x_{t_0} y' = b.$$

Thus  $T(B) = B$ . As  $T$  acts distally on  $\text{Sub}_G^p$ , we have that  $T(A) = A$ . Since  $A$  is a one-parameter subgroup in  $G$  and  $T$  is unipotent, we get that  $T(x_t) = x_t$ ,  $t \in \mathbb{R}$ . This leads to a contradiction as  $T(x_1) \neq x_1$ . Therefore,  $T(x) = x$  for all  $x \in U$ , and hence, for all  $x \in G$ . Thus  $T = \text{Id}$  if  $L$  is simply connected.

**Step 4:** Suppose that  $L$  is not simply connected. Let  $M$  be the maximal compact (normal) subgroup of  $L$ . Then  $M$  is central in  $G$  and  $L/M$  is simply connected. Let  $\varrho : G \rightarrow G/M$  be the natural projection and let  $\bar{T}$  be the automorphism on  $G/M$  induced  $T$ . Then  $\bar{T}$  acts trivially on both  $\varrho(L)$  and  $\varrho(G)/\varrho(L)$ . By Proposition 3.4,  $\bar{T}$  acts distally

on  $\text{Sub}_{G/M}^p$ . From Step 3, we get that  $\overline{T}$  acts trivially on  $G/M$ , i.e.  $T(g) \in gM$ ,  $g \in G$ . In particular, for  $\{x_t\}$  as above,  $T(x_t) = x_t y_t$ , where  $y_t \in M$ , for all  $t \in \mathbb{R}$ , and  $y_1 \neq e$ . Since  $M$  is central in  $G$ ,  $y_t x_s = x_s y_t$ ,  $t, s \in \mathbb{R}$ , and  $\{y_t\}$  is a one-parameter subgroup of the compact central subgroup  $M$ . Note that  $\{y_t\}$  is closed if  $\{x_t\}$  is compact.

As  $\text{Sub}_G^p$  is compact, there exists an unbounded sequence  $\{n_k\}$  in  $\mathbb{N}$  such that  $T^{n_k}(\{x_t\}) \rightarrow C$  for some  $C \in \text{Sub}_G^p$ . We show that  $C = \{x_t\} \times \overline{\{y_t\}}$ . As  $\{x_t\} \subset \{x_t\} \times \overline{\{y_t\}}$ , and the latter is  $T$ -invariant, we get that  $C \subset \{x_t\} \times \overline{\{y_t\}}$ . For a fixed  $t \in \mathbb{R}$ ,

$$T^{n_k}(x_{t/n_k}) = x_{t/n_k} y_{t/n_k}^{n_k} = x_{t/n_k} y_t \rightarrow y_t$$

as  $n_k \rightarrow \infty$ . Therefore,  $y_t \in C$  for each  $t$  and  $\overline{\{y_t\}} \subset C$  as  $C$  is closed. Also,  $T^{n_k}(x_t) = x_t y_t^{n_k} \rightarrow x_t y$  along a subsequence of  $\{n_k\}$  for some  $y \in \overline{\{y_t\}} \subset C$ . Hence,  $x_t \in C$ ,  $t \in \mathbb{R}$ . Thus  $\{x_t\} \times \overline{\{y_t\}} \subset C$ . Now  $T(C) = C$ , and we get that  $T^{n_k}(T(\{x_t\})) \rightarrow C$ . as  $T$  acts distally on  $\text{Sub}_G^p$ , we have that  $T(\{x_t\}) = (\{x_t\})$ . Since  $T$  is unipotent, we have that  $T(x_t) = x_t$  for each  $t$ . But  $T(x_1) = x_1 y_1 \neq x_1$ . This leads to a contradiction. Therefore,  $T(x) = x$  for each  $x \in U$ , and hence, for each  $x \in G$ . Thus  $T = \text{Id}$ .  $\square$

*Proof of Theorem 1.3.* Let  $G$  be a connected Lie group and let  $T \in \text{Aut}(G)$ . Suppose  $T$  is unipotent and it acts distally on  $\text{Sub}_G^p$ . We show that  $T = \text{Id}$ .

**Step 1:** Suppose  $G$  is nilpotent. If possible, suppose  $T \neq \text{Id}$ . Since  $T$  is unipotent, by Proposition 3.10 of [40], there exists an increasing sequence of closed connected normal  $T$ -invariant subgroups  $\{e\} = G_0 \subset G_1 \subset \dots \subset G_n = G$ ,  $n \geq 2$ , such that  $T$  acts trivially on each successive quotient group  $G_i/G_{i-1}$ ,  $1 \leq i \leq n$ , and  $T$  does not act trivially on  $G_{i+1}/G_{i-1}$ ,  $1 \leq i \leq n-1$ . Then  $G_2 \neq G_1$ . As  $T$  acts distally on  $\text{Sub}_G^p$ , it acts distally on  $\text{Sub}_{G_2}^p$ . Now by Proposition 3.8,  $T|_{G_2} = \text{Id}$ , which leads to a contradiction. Therefore,  $T = \text{Id}$  when  $G$  is nilpotent.

**Step 2:** Suppose  $G$  is not nilpotent. Let  $N$  be the nilradical of  $G$ . Then  $T|_N = \text{Id}$ . Suppose  $G$  is solvable. As noted in Remark 3.3, we have that the automorphism induced by  $T$  on  $G/N$  is a finite order automorphism, and it is trivial, since it is unipotent. Now by Proposition 3.8,  $T = \text{Id}$ .

**Step 3:** Suppose  $G$  is not solvable. Let  $R$  be the the radical of  $G$ . Then as  $T$  acts distally on  $\text{Sub}_R^p$ , we have from from Step 2 that  $T|_R = \text{Id}$ .



We first assume that  $G$  belongs to class  $\mathcal{C}'$ , i.e.  $G$  has no compact subgroup of positive dimension. (For this case, we will follow the proof of Theorem 4.1 in [40] to some extent and use Step 1 instead of Proposition 4.2 of [40]). Now  $\text{Aut}(G)$  is almost algebraic and it has finitely many connected components (see [12]). Thus some power of  $T$  belongs to a one-parameter subgroup of  $\text{Aut}(G)^0$ . Since  $T$  is unipotent, we get that  $T$  itself belongs to a one-parameter subgroup  $\{T_t\}$  consisting of unipotent elements as  $T = T_1$ . Also, since  $\{T^n\}_{n \in \mathbb{Z}}$  is co-compact in  $\{T_t\}$ , it is easy to see that each  $T_t$  also acts distally on  $\text{Sub}_G^p$ .

Let  $G' = G/R$  and let  $T'_t$  be the automorphism of  $G'$  induced by  $T_t$ ,  $t \in \mathbb{R}$ . Here,  $T'_t$  is also unipotent. Note that  $G'$  is semisimple and  $\text{Aut}(G')^0$  is the group of inner automorphisms of  $G'$ . There exists a one-parameter subgroup  $\{u_t\}$  consisting of Ad-unipotent elements in  $G'$ , such that  $T'_t = \text{inn}(u_t)$ , the inner automorphisms by  $u_t$ ,  $t \in \mathbb{R}$ . As shown in [40] we can choose a Iwasawa decomposition  $G' = KAU$  such that  $\{u_t\} \subset U$ , and  $U$  (resp.  $AU$ ) is a closed simply connected nilpotent (resp. solvable) group. Let  $\pi : G \rightarrow G' = G/R$  denote the natural projection. Then  $B := \pi^{-1}(AU)$  is a closed connected solvable group,  $T_t(B) = B$  and  $T_t$  acts distally on  $\text{Sub}_B^p$ . This implies from Step 2, that  $T_t|_B = \text{Id}$ . In particular,  $\text{Inn}(u_t)$  acts trivially on  $AU$ , i.e.  $\{u_t\}$  is contained in the center of  $AU$ . Let  $\mathcal{A}$  (resp.  $\mathcal{U}$ ) be the Lie algebra of  $A$  (resp.  $U$ ). Then  $u_t = \exp tX$ ,  $t \in \mathbb{R}$ , for some  $X \in \mathcal{U}$ , and hence  $X$  centralises  $\mathcal{A}$ . Using the restricted root system and Lemma 6.50 of [28], it follows that the centraliser of  $\mathcal{A}$  in  $\mathcal{AU}$  is contained in  $\mathcal{A}$ . Therefore,  $X \in \mathcal{A} \cap \mathcal{U} = \{0\}$ . This implies that for each  $t$ ,  $u_t = e$ , and hence that  $T'_t = \text{Id}$ .

Let  $G = SR$  be a Levi decomposition, where  $S$  is a Levi subgroup (maximal connected semisimple subgroup) of  $G$ . Let  $\pi_1 : G \rightarrow G/N$  be the natural projection. Then  $\pi_1(S)$  and  $\pi_1(R)$  centralise each other. Since  $T = T_1$  and  $T'_1 = \text{Id}$  on  $G/R$ , we have that  $T(x) \in xR$ ,  $x \in G$ . Let  $x, y \in S$ . Then  $\pi_1(T(xyx^{-1}y^{-1})) = \pi_1(xyx^{-1}y^{-1})$ . Since  $[S, S] = S$ , we get that  $T(x) \in xN$  for all  $x \in S$ . As  $N$  is closed, we get that  $T(x) \in xN$  for all  $x \in \overline{SN}$ . As  $T$  acts distally on  $\text{Sub}_{\overline{SN}}^p$  and  $T|_N = \text{Id}$ , by Proposition 3.8, we get that  $T(x) = x$  for all  $x \in S$ . Since  $T|_R = \text{Id}$ . We get that  $T = \text{Id}$ .

We have shown that under the conditions stated in the proposition,  $T = \text{Id}$  if  $G$  is either solvable or if  $G$  belongs to class  $\mathcal{C}'$ .

**Step 4:** Let  $M$  be the largest compact connected central subgroup of  $G$ . Suppose  $M$  is non-trivial. Then  $G/M$  belongs to class  $\mathcal{C}'$ ,  $M$  is  $T$ -invariant and by Proposition 3.4,  $T$  acts distally on  $\text{Sub}_{G/M}^p$ . Since the automorphism of  $G/M$  induced by  $T$  is also unipotent, we get from Step 3 that  $T$  acts trivially on  $G/M$ . That is,  $T(x) \in xM$  for all  $x \in G$ . As  $M$  is abelian, by Step 1,  $T$  acts trivially on  $M$ . Now by Proposition 3.8,  $T = \text{Id}$ .

The converse statement in the theorem is obvious.  $\square$

*Proof of Theorem 1.1(2) for non-nilpotent case.* Let  $G$  be a connected Lie group in class  $\mathcal{C}'$ . Then  $\text{Aut}(G)$  is almost algebraic as a subgroup of  $\text{GL}(\mathfrak{g})$  (via the correspondence  $T \leftrightarrow dT$ ) (cf. [12]). Suppose  $T \in \text{Aut}(G)$  acts distally on  $\text{Sub}_G^p$ . Suppose  $G$  is not nilpotent. In particular, it is not a vector group. By Theorem 1.2,  $T$  acts distally on  $G$ . Then for some  $n \in \mathbb{N}$ ,  $T^n$  is contained in a connected abelian almost algebraic subgroup of  $\text{Aut}(G)$  and it has the form  $\mathcal{K} \times \mathcal{U}$ , where  $\mathcal{K}, \mathcal{U} \subset \text{Aut}(G)$ ,  $\mathcal{K}$  is an abelian compact connected group and  $\mathcal{U}$  is an abelian unipotent connected group; i.e. it consists of unipotent elements (cf. [1], Corollaries 2.3 and 2.5). Now  $T^n = S \circ \psi = \psi \circ S$ , where  $S \in \mathcal{U}$  is unipotent and  $\psi \in \mathcal{K}$ . Hence,  $S$  acts distally on  $\text{Sub}_G^p$  (cf. [39], Lemma 2.2). By Theorem 1.3,  $S = \text{Id}$ . Thus  $T^n = \psi \in \mathcal{K}$  and hence  $T$  is contained in a compact subgroup of  $\text{Aut}(G)$ . The converse is obvious (cf. [30], Lemma 4.2).  $\square$

Now we want to prove Theorem 1.4. We need an elementary result about certain automorphisms keeping a maximal compact abelian subgroups (maximal tori) invariant. Maximal compact subgroups in a connected Lie group  $G$  are connected and conjugate to each other (see [24]). Any maximal compact subgroup of a connected solvable Lie group is abelian. Any maximal compact abelian subgroup of  $G$  is connected (in fact, it is a Cartan subgroup of the maximal compact group in which it is contained). A maximal compact (connected) abelian subgroup of a connected Lie group  $G$  is called a maximal torus and maximal tori are conjugate to each other. Any compact abelian subgroup is contained in a maximal torus. If  $A$  is a maximal torus of  $G$ , then so is  $T(A)$ , for any  $T \in \text{Aut}(G)$ , and  $T(A)$  is a conjugate of  $A$ .

In [15], any subgroup of  $\text{Aut}(G)$  which fixes elements of a maximal torus are shown to be almost algebraic. Given any automorphism  $T$  of  $G$ , it need not keep any torus invariant. However, this is the case

if  $T$  is contained in a compact connected abelian subgroup of  $\text{Aut}(G)$  as shown in the following elementary lemma, which also holds if we replace the radical  $R$  by any closed connected characteristic subgroup of  $G$ .

**Lemma 3.9.** *Let  $G$  be a connected Lie group. If  $\mathcal{K}$  is a compact connected abelian subgroup of  $\text{Aut}(G)$ , then there exists a maximal torus of  $G$  (resp. of the radical of  $G$ ) which is invariant under every automorphism of  $G$  which belongs to  $\mathcal{K}$ .*

*Proof.* Let  $A$  be any maximal torus in a connected Lie group  $G$ . For any closed connected normal subgroup  $L$  of  $G$ , since the maximal tori of  $G$  (resp.  $L$ ) are conjugate to each other in  $G$  (resp.  $L$ ), we get that  $A$  contains a maximal torus of  $L$  and hence,  $A \cap L$  is a maximal torus of  $L$ .

Let  $\mathcal{K}$  be a compact connected abelian subgroup of  $\text{Aut}(G)$  and let  $H = \mathcal{K} \rtimes G$ ; where the action of  $\mathcal{K}$  on  $G$  is by automorphisms. Then  $H$  is a connected Lie group and  $G$  is normal in  $H$ . Let  $R$  be the radical of  $G$ . Then  $R$  is characteristic in  $G$  and hence normal in  $H$ . Let  $A$  be a maximal torus in  $H$  containing  $\mathcal{K}$ . Arguing as above, we get that  $A \cap G$  (resp.  $A \cap R$ ) is a maximal torus in  $G$  (resp.  $R$ ). As  $A \cap G$  (resp.  $A \cap R$ ) is normal in  $A$  and  $\mathcal{K} \subset A$ , we get that  $A \cap G$  (resp.  $A \cap R$ ) is invariant under every automorphism which belongs to  $\mathcal{K}$ .  $\square$

*Proof of Theorem 1.4.* Let  $G$  be a connected Lie group and let  $T \in \text{Aut}(G)$ . The statements (5)  $\implies$  (4)  $\implies$  (3)  $\implies$  (2)  $\implies$  (1) are obvious. Now it is enough to show that (1)  $\implies$  (5) when  $G$  is not a vector group and  $T$  acts distally on the largest compact connected central subgroup (say)  $M$  of  $G$ . Suppose (1) holds, i.e.  $T$  acts distally on  $\text{Sub}_G^p$ . Then we want to show that  $T$  is contained in a compact subgroup of  $\text{Aut}(G)$ . If  $M$  is trivial, then  $G$  belongs to the class  $\mathcal{C}'$  and the assertion follows from Theorem 1.1 (2). Now suppose  $M$  is non-trivial. By Theorem 1.2,  $T$  acts distally on  $G$ . Now  $\overline{T}$  also acts distally on  $G/M$  (cf. [33], Theorem 3.1).

Suppose  $G = M$ . Since  $T$  acts distally on  $M$ , by Lemma 2.5 of [2],  $T^n$  is unipotent for some  $n \in \mathbb{N}$ . As  $T$  acts distally on  $\text{Sub}_G^p$ , so does  $T^n$ , and by Theorem 1.3,  $T^n = \text{Id}$ . Hence the assertion holds when  $G$  is a compact and abelian.

Now suppose  $M$  is non-trivial and  $G \neq M$ . We show that there exists a maximal torus of  $G$  which is  $T^k$ -invariant for some  $k \in \mathbb{N}$ . Let

$\pi : G \rightarrow G/M$  be the natural projection. By Proposition 3.4,  $\overline{T}$  acts distally on  $\text{Sub}_{G/M}^p$ , where  $\overline{T}$  is the automorphism of  $G/M$  induced by  $T$ .

Suppose  $G/M$  is a vector group. Then by Theorem 1.1 (1),  $\overline{T} = \psi \circ \alpha \text{Id}$ , for some  $\alpha \in \mathbb{R} \setminus \{0\}$  and  $\psi$  generates a relatively compact subgroup in  $\text{Aut}(G/M)$ . Since  $\overline{T}$  acts distally on  $G/M$ , we get that  $|\alpha| = 1$ , and hence  $\overline{T}$  is contained in a compact subgroup of  $\text{Aut}(G/M)$ . Now suppose  $G/M$  is not a vector group. As  $G/M$  belongs to the class  $\mathcal{C}'$ , by Theorem 1.1 (2),  $\overline{T}$  is contained in a compact subgroup of  $\text{Aut}(G/M)$ . So in either case we have that  $\overline{T}$  is contained in a compact subgroup of  $\text{Aut}(G/M)$ .

Let  $\mathcal{K}$  be the connected component of the identity in the closed subgroup generated by  $\overline{T}$  in  $\text{Aut}(G/M)$ , then  $(\overline{T})^k \in \mathcal{K}$  for some  $k \in \mathbb{N}$ . Now by Lemma 3.9,  $\overline{T}$  keeps a maximal torus (say),  $A$  of  $G/M$  invariant. Let  $K = \pi^{-1}(A)$ . Then  $K$  is a maximal torus of  $G$  and  $T^k(K) = K$ .

Note that  $K$  is compact, connected and abelian. Moreover, as  $T$  acts distally on  $G$ , so does  $T^k$  and the restriction of  $T^k$  to  $K$  is also distal. Arguing as above for  $K$  instead of  $M$ , we get that for some  $m \in \mathbb{N}$ ,  $T^m|_K$  is unipotent, and as it acts distally on  $\text{Sub}_K^p$ , it acts trivially on  $K$  (by Theorem 1.3). Replacing  $T$  by  $T^m$ , we may assume that  $T$  fixes every element of  $K$ , i.e.  $T \in F_K(G) = \{\psi \in \text{Aut}(G) \mid \psi(x) = x \text{ for all } x \in K\}$ . We know that  $F_K(G)$  is almost algebraic (cf. [15]). As  $T$  is distal and  $T \in F_K(G)$ , for some  $n \in \mathbb{N}$ ,  $T^n$  is contained in a connected almost algebraic subgroup  $\mathcal{B} \times \mathcal{U}$ , where  $\mathcal{B}$  is a compact group and  $\mathcal{U}$  consists of unipotent elements (cf. [1], Corollaries 2.3 and 2.5). Now we have that  $T = \phi \circ S = S \circ \phi$  such that  $\phi \in \mathcal{B}$  and  $S \in \mathcal{U}$ . As  $\mathcal{B}$  is compact, we have that  $S$  acts distally on  $\text{Sub}_G^p$ . By Theorem 1.3,  $S = \text{Id}$ . Thus  $T = \phi \in \mathcal{B}$  and (5) holds.  $\square$

Using Theorems 1.1 and 1.4 and some known results, we prove Theorem 1.5.

*Proof of Theorem 1.5.* Let  $G$  be a connected Lie group and let  $\mathcal{H}$  be a subgroup of  $\text{Aut}(G)$ . Suppose  $\mathcal{H}$  acts distally on the largest compact connected central subgroup of  $G$ . Then so does  $\overline{\mathcal{H}}$ , which is a closed subgroup of  $\text{Aut}(G)$ . Note that  $\mathcal{H}$  acts distally on a compact  $\mathcal{H}$ -invariant subspace of  $\text{Sub}_G$  if and only if so does  $\overline{\mathcal{H}}$  (both the above statements follow from [16]). Also (7) holds for  $\mathcal{H}$  if and only if it holds

when  $\mathcal{H}$  is replaced by  $\overline{\mathcal{H}}$ , as  $\mathcal{KD}$  is closed in  $\mathrm{GL}(d, \mathbb{R})$  for  $\mathcal{K}, \mathcal{D}$  as in (7). Therefore to prove the assertions, we can assume that  $\mathcal{H}$  is closed. Note that every compact subgroup of  $\mathrm{Aut}(G)$  acts distally on  $\mathrm{Sub}_G$  (see e.g. [39], Lemma 2.2). Now (6)  $\implies$  (5)  $\implies$  (4)  $\implies$  (3)  $\implies$  (2)  $\implies$  (1) are obvious.

First suppose that  $G$  is not a vector group. We show that (1)  $\implies$  (6). As in the hypothesis, Suppose (1) holds, i.e. every  $T \in \mathcal{H}$  acts distally on  $\mathrm{Sub}_G^p$ . By Theorem 1.4, every  $T \in \mathcal{H}$  is contained in a compact subgroup of  $\mathrm{Aut}(G)$ . Recall that  $\mathrm{Aut}(G)$  is identified with a closed subgroup of  $\mathrm{GL}(\mathfrak{g})$  via the map  $T \rightarrow dT$ , where  $\mathfrak{g}$  is the Lie algebra of  $G$  and it is a  $d$ -dimensional vector space for some  $d \in \mathbb{N}$  (cf. [9], see also [21]). Therefore,  $\mathcal{H}$  is also a closed subgroup of  $\mathrm{GL}(\mathfrak{g})$  and the latter is isomorphic to  $\mathrm{GL}(d, \mathbb{R})$ , and we get by Proposition 2 of [38] that  $\mathcal{H}$  is compact. Thus (6) holds and (1 – 6) are equivalent. Therefore, statements (a), (b) and (c) hold in case  $G$  is not a vector group.

Now suppose  $G = \mathbb{R}^d$  for some  $d \in \mathbb{N}$ . Suppose (3) holds, i.e.  $\mathcal{H}$  acts distally on  $\overline{\mathrm{Sub}_G^c}$ . Then every  $T \in (\mathrm{NC})$ . Since  $G$  belongs to the class  $\mathcal{C}'$ , by Theorem 4.1 of [40],  $T$  is contained in a compact subgroup of  $\mathrm{Aut}(G)$ . Since  $\mathcal{H}$  is closed, arguing as above using Proposition 2 of [38], we get that  $\mathcal{H}$  is compact. Hence (6) holds. Thus (3 – 6) are equivalent and hence (c) holds when  $G$  is a vector group. Now we show that (1), (2) and (7) are equivalent. Suppose (7) holds. Note that  $\mathcal{D} = \{r \mathrm{Id} \in \mathrm{GL}(d, \mathbb{R}) \mid r \in \mathbb{R} \setminus \{0\}\}$ . Therefore,  $\mathcal{D}$  acts trivially on  $\mathrm{Sub}_G^p$  and the action of  $\mathcal{KD}$  on  $\mathrm{Sub}_G^p$  is same as that of the compact group  $\mathcal{K}$  on  $\mathrm{Sub}_G^p$ . Hence the action of  $\mathcal{KD}$ , and hence, the action of  $\mathcal{H}$  on  $\mathrm{Sub}_G^p$  is distal. Thus (7)  $\implies$  (2). We have already noted that (2)  $\implies$  (1).

Now we show that (1)  $\implies$  (7). Suppose every  $T \in \mathcal{H}$  acts distally on  $\mathrm{Sub}_G^p$ . Let  $T \in \mathcal{H}$  be fixed. By Theorem 1.1 (1),  $T \in \mathcal{BD}$ , for some compact group  $\mathcal{B} \subset \mathrm{GL}(d, \mathbb{R})$ . Let  $\mathcal{H}_1 := \overline{[\mathcal{H}, \mathcal{H}]}$ . Then  $\mathcal{H}_1 \subset \mathcal{H}$  as the latter is closed. Let  $T \in \mathcal{H}_1$ . Then  $\det T = 1$ . From above we have that  $T = \phi \circ r \mathrm{Id}$  for some  $\phi \in \mathcal{B}$  and some  $r \in \mathbb{R} \setminus \{0\}$ . Since  $\mathcal{B}$  is a compact group,  $\det \phi = \pm 1$ , and hence  $r = \pm 1$ . Therefore,  $T$  generates a relatively compact group, for every  $T \in \mathcal{H}_1$ . As  $\mathcal{H}_1$  is a closed subgroup of  $\mathrm{GL}(d, \mathbb{R})$  by Proposition 2 of [38],  $\mathcal{H}_1$  is compact. Now  $\mathcal{H}_1$  is a compact normal subgroup of  $\mathcal{H}$ . Let  $N(\mathcal{H}_1)$  be the normaliser of  $\mathcal{H}_1$  in  $\mathrm{GL}(d, \mathbb{R})$ . Then  $N(\mathcal{H}_1)$  is a closed algebraic subgroup of  $\mathrm{GL}(d, \mathbb{R})$

and  $\mathcal{H}, \mathcal{D} \subset N(\mathcal{H}_1)$ . Moreover,  $N(\mathcal{H}_1) \subset N(\mathcal{H}_1\mathcal{D})$ . Being algebraic,  $N(\mathcal{H}_1)$  has finitely many connected components, and hence so does  $N(\mathcal{H}_1)/\mathcal{H}_1\mathcal{D}$ . In particular any closed abelian subgroup of  $N(\mathcal{H}_1)/\mathcal{H}_1\mathcal{D}$  is compactly generated (cf. [23]) Let  $\rho : N(\mathcal{H}_1) \rightarrow N(\mathcal{H}_1)/\mathcal{H}_1\mathcal{D}$  be the natural projection. Then  $\rho(\mathcal{H})$  is abelian, and hence so is  $\overline{\rho(\mathcal{H})}$ . Then  $\overline{\rho(\mathcal{H})}$  is compactly generated, and it has a unique compact subgroup, say  $C$ . Now let  $T \in \mathcal{H}$ . Then  $T = \psi \circ r \text{Id}$  for some  $\psi$  which generates a relatively compact group and  $r \text{Id} \in \mathcal{D}$ . Then  $\rho(T) = \rho(\psi)$  also generates a relatively compact group in  $\overline{\rho(\mathcal{H})}$ , and hence it is contained in  $C$ . Thus  $\rho(\mathcal{H}) \subset C$  and  $\overline{\rho(\mathcal{H})}$  is compact. As  $\mathcal{H}_1$  is compact, it follows that  $\overline{\mathcal{H}\mathcal{D}}/\mathcal{D}$  is compact. Since  $\mathcal{D}$  has two connected components, we get that  $\overline{\mathcal{H}\mathcal{D}}/\mathcal{D}^0$  is also compact and it is a Lie group. As  $\mathcal{D}^0$  is a vector group which is central in  $\overline{\mathcal{H}\mathcal{D}}$ , by Lemma 3.7 of [24],  $\overline{\mathcal{H}\mathcal{D}} = \mathcal{K} \times \mathcal{D}^0$ , where  $\mathcal{K}$  is the maximal compact subgroup of  $\overline{\mathcal{H}\mathcal{D}}$ . In particular,  $\mathcal{H} \subset \mathcal{K}\mathcal{D}^0$ . Thus (7) holds. Therefore, if  $G$  is a vector group, (1), (2), and (7) are equivalent and (a) holds and (d) holds. This completes the proof.  $\square$

Now we state two corollaries which are easy to deduce using results proven above and a result in [1].

**Corollary 3.10.** *Let  $G$  be a connected Lie group and let  $\mathcal{H}$  be a subgroup of  $\text{Aut}(G)$ . Then  $\mathcal{H}$  acts distally on both  $G$  and  $\text{Sub}_G^p$  if and only if  $\overline{\mathcal{H}}$  is compact.*

Corollary 3.10 is a direct consequence of Theorem 1.5 in case  $G$  is not a vector group. If  $G$  is a vector group, then as  $\mathcal{H}$  acts distally on  $G$ , the eigenvalues of every element in  $\mathcal{H}$  have absolute value 1 (cf. [1]), and this together with the statement (7) in Theorem 1.5 is equivalent to the statement that  $\overline{\mathcal{H}}$  is compact. We omit the details.

The following corollary is an extension of Corollary 4.6 of [40] for a particular class of Lie groups. It can be proven easily using Theorems 1.1 and 1.5 and Proposition 3.4. We omit the proof.

**Corollary 3.11.** *Let  $G$  be a connected Lie group such that  $G/M$  is not a vector group, where  $M$  is the maximal compact connected central subgroup of  $G$ . Let  $T \in \text{Aut}(G)$  (resp.  $\mathcal{H}$  be a subgroup of  $\text{Aut}(G)$ ). If  $T$  (resp.  $\mathcal{H}$ ) acts distally on  $\text{Sub}_G^p$ , then  $T$  (resp.  $\mathcal{H}$ ) has bounded orbits; i.e.  $\{T^n(x) \mid n \in \mathbb{Z}\}$  (resp.  $\mathcal{H}x$ ) is relatively compact in  $G$ , for every  $x \in G$ .*

Now we prove Corollary 1.6. The statement in the corollary which requires a proof is (1)  $\implies$  (5) and can be proven along the lines of proof of (1)  $\implies$  (5) of Corollary 4.5 in [40]; instead of Corollary 4.3 and Theorem 4.3 there, one has to use Theorems 1.2 and 1.3. We will give a proof here for the sake of completeness.

*Proof of Corollary 1.6.* Let  $G$  be a connected Lie group. The statement in (5) implies that  $\text{Inn}(G)$  is compact, and hence it follows that (5)  $\implies$  (4). Also (4)  $\implies$  (3)  $\implies$  (1) and (4)  $\implies$  (2)  $\implies$  (1) are obvious. It is enough to show that (1)  $\implies$  (5).

Suppose (1) holds; i.e. every inner automorphism of  $G$  acts distally on  $\text{Sub}_G^p$ . If  $G$  is either compact or a vector group, then (5) holds trivially. Now suppose  $G$  neither compact nor a vector group. Since every inner automorphism acts trivially on the center of  $G$ , we have by Theorem 1.2 that it acts distally on  $G$ . Therefore,  $G$  is distal (cf. [35]), i.e. the action of  $\text{Inn}(G)$  on  $G$  is distal. It follows from Theorem 9 of [35] and Corollary 2.1 (ii) of [26] that  $G/R$  is compact, where  $R$  is the radical of  $G$ . Let  $N$  be the nilradical of  $G$  and let  $x \in N$ . Then  $\text{inn}(x)$  is unipotent, and by Theorem 1.3,  $\text{inn}(x) = \text{Id}$ . This implies that  $x$  is central in  $G$  for every  $x \in N$ . Then  $N$  is abelian and central in  $G$ . Now for the radical  $R$ , we know that  $[R, R] \subset N$  is central in  $G$ , and hence  $R$  is nilpotent and it is same as  $N$  and it is abelian and central in  $G$ . Now  $G/N$  is compact, where  $N$  is central in  $G$ , i.e.  $N = \mathbb{R}^n \times M$ , for some  $n \in \mathbb{N}$ , where  $M$  is compact. Here,  $V = \mathbb{R}^n$  is a vector subgroup which is central. By Lemma 3.7 of [24],  $G = K \times \mathbb{R}^n$ , where  $K$  is the maximal compact subgroup of  $G$ . Thus (5) holds.  $\square$

The distal actions of automorphisms  $T$  of a connected Lie group  $G$  on  $\text{Sub}_G$  and its closed  $T$ -invariant subspaces were studied if  $G$  is in class  $\mathcal{C}'$  or  $T$  is unipotent; here we have studied a more general case when  $T$  acts distally on the maximal central torus of  $G$ . We have also studied the action of subgroups of  $\text{Aut}(G)$  on  $\text{Sub}_G$  and  $\text{Sub}_G^p$  for a connected Lie group  $G$  under a condition that the action on the maximal central torus is distal. It would be interesting to characterise general automorphisms of  $G$  which act distally on  $\text{Sub}_G$  or its closed invariant subspace for any connected Lie group  $G$  without the condition involving the action on the central torus.

**Acknowledgement.** The authors would like to acknowledge the support in part by the International Centre for Theoretical Sciences (ICTS)

during a visit for participating in the program - ICTS Ergodic Theory and Dynamical Systems (code: ICTS/etds-2022/12).

## REFERENCES

- [1] H. Abels. Distal affine transformation groups. *J. Reine Angew. Math.* **299/300** (1978), 294–300.
- [2] H. Abels. Distal automorphism groups of Lie groups. *J. Reine Angew. Math.* **329** (1981), 82–87.
- [3] H. Baik and L. Clavier. The space of geometric limits of one-generator closed subgroups of  $\mathrm{PSL}_2(\mathbb{R})$ . *Algebr. Geom. Topol.* **13** (2013), 549–576.
- [4] H. Baik and L. Clavier. The space of geometric limits of abelian subgroups of  $\mathrm{PSL}_2(\mathbb{C})$ . *Hiroshima Math. J.* **46** (2016), 1–36.
- [5] H. Bass. Groups of integral representation type, *Pacific J. Math.* **86** (1980), 15–51.
- [6] R. Benerdetti, C. Petronio. *Lecture on Hyperbolic Geometry*. Universitext, Springer, Berlin, 1992.
- [7] S. Bhattacharya. Expansiveness of algebraic actions on connected groups. *Transac. Amer. Math. Soc.* **356** (2004), 4687–4700.
- [8] M. R. Bridson, P. de la Harpe, and V. Kleptsyn, The Chabauty space of closed subgroups of the three-dimensional Heisenberg group. *Pacific J. Math.* **240** (2009), 1–48.
- [9] C. Chevalley, *Theory of Lie groups*, Princeton University Press, 1946.
- [10] M. Choudhuri, G. Faraco and A. K. Yadav. On the distality and expansivity of certain maps on spheres. *Dyn. Syst.* **39** (2024), 166–181.
- [11] J. P. Conze and Y. Guivarc’h. *Remarques sur la distalité dans les espaces vectoriels*. C. R. Acad. Sci. Paris Ser. A **278** (1974), 1083–1086.
- [12] S. G. Dani. On automorphism groups of connected Lie groups. *Manuscripta Math.* **74** (1992), 445–452.
- [13] S. G. Dani and M. McCrudden. Convolution roots and embeddings of probability measures on Lie groups, *Adv. Math.* **209** (2007), 198–211.
- [14] S. G. Dani and R. Shah. Contractible measures and Levy’s measures on Lie groups. In: Probability on Algebraic Structures (Ed. G. Budzban P. Feinsilver and A. Mukherjea), *Contemporary Math.* **261** (2000), 3–13.
- [15] S. G. Dani and R. Shah. On the algebraicity of certain subgroups of automorphism groups of connected Lie groups. preprint.
- [16] R. Ellis. Distal transformation Groups. *Pacific J. Math.* **8** (1958), 401–405.
- [17] H. Furstenberg. The structure of distal flows. *Amer. J. Math.* **85** (1963), 477–515.
- [18] T. Gelander, A lecture on invariant random subgroups. In *New Directions in Locally Compact Groups*, pp. 186–204, London Math. Soc. Lecture Note Ser. 447, Cambridge Univ. Press, Cambridge, 2018.
- [19] H. Hamrouni and B. Kadri, On the compact space of closed subgroups of locally compact groups. *J. Lie Theory* **24** (2014), 715–723.



- [20] W. Hazod and E. Siebert. Automorphisms on a Lie group contracting modulo a compact subgroup and applications to semistable convolution semigroups, *J. Theoret. Probab.* **1** (1988), 211–225.
- [21] G. Hochschild. The automorphism group of a Lie group. *Trans. Amer. Math. Soc.* **72** (1952), 209–216.
- [22] G. Hochschild. *The Structure of Lie Groups*. Holden-Day, San Francisco, 1965.
- [23] K. H. Hofmann and K.-H. Neeb, The compact generation of closed subgroups of locally compact groups. *J. Group Theory* **12** (2009), 555–559.
- [24] K. Iwasawa. On Some Types of Topological Groups. *Ann. Math.* **50** (1949), 507–558.
- [25] W. Jaworski. Contraction groups, ergodicity and distal properties of automorphisms on compact groups. *Illinois J. Math.* **56** (2012), 1023–1084.
- [26] J. Jenkins. Growth of connected locally compact groups. *J. Functional Analysis* **12** (1973), 113–127.
- [27] B. Kloeckner. The space of closed subgroups of  $\mathbb{R}^n$  is stratified and simply connected. *J. Topology.* **2** (2009), 570–588.
- [28] A. W. Knap. *Lie Groups Beyond an Introduction*. Second Edition. Progress in Mathematics 140. (Birkhäuser Boston Inc., 2002).
- [29] C. C. Moore. Distal affine transformation groups. *Amer. J. Math.* **90** (1968), 733–751.
- [30] R. Palit and R. Shah. Distal actions of automorphisms of nilpotent groups  $G$  on  $\text{Sub}_G$  and applications to lattice in Lie groups. *Glasgow Math. J.* **63** (2021) 343–362.
- [31] R. Palit, M. B. Prajapati and R. Shah. Dynamics of actions of automorphisms of discrete groups  $G$  on  $\text{Sub}_G$  and applications to lattices in Lie groups. *Groups Geom. Dyn.* **17** (2023), 185–213.
- [32] I. Poureza and J. Hubbard. The space of closed subgroups of  $\mathbb{R}^2$ . *Topology* **18** (1979), 143–146.
- [33] C. R. E. Raja and R. Shah, Distal actions and shifted convolution property, *Israel J. Math.* **177** (2010), 301–318.
- [34] C. R. E. Raja and R. Shah, Some properties of distal actions on locally compact groups, *Ergodic Theory and Dynam. Sys.* **39** (2019) 1340–1360.
- [35] J. Rosenblatt. A distal property of groups and the growth of connected locally compact groups. *Mathematika* **26** (1979), 94–98.
- [36] R. Shah. Orbits of distal actions on locally compact groups. *J. Lie Theory* **22** (2010), 586–599.
- [37] R. Shah. Expansive automorphisms on connected locally compact groups. *New York J. Math.* **26** (1920), 285–302.
- [38] R. Shah and A. K. Yadav. Dynamics of certain distal actions on spheres. *Real Analysis Exchange* **44** (2019), 77–88.
- [39] R. Shah and A. K. Yadav. Distally of certain actions on  $p$ -adic spheres. *J. Australian Math. Soc.* **109** (2020), 250–261.
- [40] R. Shah and A. K. Yadav. Distal action of Automorphisms of Lie groups  $G$  on  $\text{Sub}_G$ . *Math. Proc. Camb. Phil. Soc.* **173** (2022), 457–478.

- [41] E. Siebert. Contractive automorphisms on locally compact groups. *Math. Z.* **191** (1986), 73–90.
- [42] M. Stroppel. *Locally Compact Groups*. EMS Textbooks in Mathematics (European Math. Soc. - EMS, 2006).

DEBAMITA CHATTERJEE, SCHOOL OF PHYSICAL SCIENCES, JAWAHARLAL NEHRU UNIVERSITY, NEW DELHI 110 067

*Email address:* debamitachatterjee1994@gmail.com

RIDDHI SHAH, SCHOOL OF PHYSICAL SCIENCES, JAWAHARLAL NEHRU UNIVERSITY, NEW DELHI 110 067

*Email address:* rshah@jnu.ac.in, riddhi.kausti@gmail.com