

# ON THE STRONGLY SUBDIFFERENTIABLE POINTS IN LIPSCHITZ-FREE SPACES

CHRISTIAN COBOLLO, SHELDON DANTAS, PETR HÁJEK, AND MINGU JUNG

*Dedicated to Professor William B. Johnson on the occasion of his 80th birthday.*

**ABSTRACT.** In this paper, we present some sufficient conditions on a metric space  $M$  for which every elementary molecule is a strongly subdifferentiable (SSD, for short) point in the Lipschitz-free space  $\mathcal{F}(M)$  over  $M$ . Our main result reads as follows: if  $(M, d)$  is a metric space and  $\gamma > 0$ , then there exists a metric  $d_\gamma$  in  $M$  such that every finitely supported element in  $\mathcal{F}(M, d_\gamma)$  is an SSD point. Consequently, the subset of all SSD points in  $\mathcal{F}(M, d_\gamma)$  is dense.

## 1. INTRODUCTION

We focus our attention on a concept of differentiability of the norm of a Banach space, which is called *strong subdifferentiability* (SSD, for short). To put it simply, studying when the norm of a Banach space is strongly subdifferentiable is essentially the same as investigating what is lacking for a Gâteaux differentiable norm to become a Fréchet differentiable norm. In fact, SSD is a strictly weaker concept than Fréchet differentiability (indeed, every point of a finite-dimensional Banach space is SSD). More precisely, an element  $x \in X$  is an SSD point if the limit of the expression  $\frac{\|x+th\|-\|x\|}{t}$  exists uniformly in  $h$  as  $h \rightarrow 0^+$ . To give the interested reader a sense of what is known about SSD in classical Banach spaces, we present some relevant results in this area. In what follows, we say that a Banach space  $X$  is SSD when *every* point of its unit sphere is an SSD point. Every uniformly smooth Banach space is SSD since its norm is uniformly Fréchet differentiable on the unit sphere, and if a dual space  $X^*$  is SSD, then  $X$  must be reflexive (see [16, Theorem 3.3]) which implies in particular that neither  $\ell_1$  nor  $\ell_\infty$  are SSD. On the other hand, if  $X$  is a predual of a Banach space with the  $w^*$ -Kadec-Klee property, then  $X$  is SSD (see [12, Proposition 2.6]); hence  $c_0$  is an example of a non-reflexive SSD Banach space. It is well-known that a Banach space admitting a Fréchet differentiable norm is an Asplund space. In fact, it turns out that the same implication holds for a Banach space admitting an SSD renorming (see [16, 21]). However, one of the important distinctions between Fréchet differentiability and SSD consists in the fact that the set of points of Fréchet differentiable points is always  $G_\delta$  while the set of SSD points is an  $F_{\sigma\delta}$  set (cf. [21, Proposition 5]). In particular, if the set of Fréchet differentiable points is dense, then it is automatically  $G_\delta$

---

2020 *Mathematics Subject Classification.* Primary 46B20; Secondary 49J50.

*Key words and phrases.* Strong subdifferentiability; Lipschitz-free spaces; Fréchet differentiability; Gâteaux differentiability; uniformly discrete spaces.

dense, which is not true for SSD points. Indeed, there is an example of Banach space, where the set of SSD points is both dense and meager (e.g., SSD points in  $\ell_1$  are nothing but finitely supported elements as observed in [18, Example 1.1] and [15, Theorem 6]); so it cannot be residual. Let us also mention that there are spaces whose norms are nowhere SSD except at the origin (see [21, Theorem 1] and [13, Thm III.1.9 and Prop.III.4.5]), and for every space with a fundamental biorthogonal system, there exists a dense subspace with an SSD norm (we send the reader to [9, 10] for more information in this line of research).

For a comprehensive study of SSD, we recommend consulting [13, 16]. Additionally, we suggest the following references [2, 7, 8, 11, 15, 18, 19, 21, 22] for insights into this theory across various Banach spaces and [14] for the basic theory of Banach spaces.

Here, our focus lies in exploring the presence of SSD points within Lipschitz-free spaces over certain metric spaces  $M$ . It is important to note that the existence of Fréchet differentiable points in  $\mathcal{F}(M)$  relies on the boundedness and uniform discreteness of  $M$  (see [4] and, more generally, [25, Theorem 4.3]). We will observe a significant change in this context concerning SSD points (cf. Corollary 3.17). On a different note, Aliaga and Rueda Zoca demonstrated that within a bounded uniformly discrete metric space  $M$ , a finitely supported element  $\mu \in S_{\mathcal{F}(M)}$  is Fréchet differentiable if and only if it is Gâteaux differentiable. They obtained such an equivalence by leveraging a geometric condition on the pairs  $(x_i, y_i)$  involved in an optimal representation of  $\mu = \sum_{i=1}^n \lambda_i m_{x_i, y_i}$  (cf. [1, Theorem 3.5]). A similar result holds true for SSD and Fréchet differentiable points (cf. Proposition 3.2) when focusing on the Lipschitz-free space over  $\cup_n [x_n, x_{n+1}]$ , where the molecule is given by  $\mu = \sum_n a_n m_{x_n, x_{n+1}}$  for some  $\{x_n\}_n \subseteq M$  containing 0 and  $\sum_n a_n = 1$  with  $a_n > 0$ .

As we have already mentioned some of our results, let us now present the contents of this manuscript in a more systematic manner. We present the necessary background in the next section. The third (and final) section of the paper is divided into three smaller parts, where we study the following topics. First, we exhibit examples of when SSD points can be obtained in  $\mathcal{F}(M)$  for certain metric spaces  $M$ . Specifically, as we briefly mentioned earlier, we prove that for any metric space  $M$ , a molecule  $\mu = \sum_n a_n m_{x_n, x_{n+1}}$  with  $\sum_n a_n = 1$  is an SSD point in  $\mathcal{F}(\cup_n [x_n, x_{n+1}])$  if and only if it is Fréchet differentiable in this space. Next, we observe that a Lipschitz-free space  $\mathcal{F}(M)$  is SSD if and only if  $\mathcal{F}(M)$  is finite-dimensional (cf. Proposition 3.5). In Subsection 3.2, we define locally uniformly non-aligned metric spaces. This allows us to prove that every elementary molecule is an SSD point in  $\mathcal{F}(M)$ , provided  $M$  is a metric space of this kind. Finally, we conclude the paper by presenting our main result. For any metric space  $(M, d)$  and  $\gamma > 0$ , we can construct a metric  $d_\gamma$  on  $M$  such that every finitely supported element in  $\mathcal{F}(M, d_\gamma)$  is an SSD point. Therefore, the set of all SSD points in  $\mathcal{F}(M, d_\gamma)$  turns out to be dense in  $\mathcal{F}(M, d_\gamma)$ .

## 2. PRELIMINARIES

This section is dedicated to introducing the main tools we will be using throughout the paper. We start by presenting the definitions and main properties we will be using about Lipschitz-free spaces, and then we proceed to discuss SSD points, which form the central

concept of the paper. Throughout the paper, we consider only *real* Banach spaces and denote by  $B_X$  and  $S_X$  the unit ball and unit sphere of a Banach space  $X$ , respectively.

**2.1. Lipschitz-free spaces.** A *pointed metric space* is just a metric space  $M$  in which we distinguish an element, called 0. Given a pointed metric space  $M$  with distinguished point 0, we write  $\text{Lip}_0(M)$  to denote the Banach space of all Lipschitz functions  $f : M \rightarrow \mathbb{R}$  which vanish at the distinguished point 0. This space is endowed with the Lip-norm

$$\|f\|_L := \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)} : x, y \in M, x \neq y \right\}.$$

The canonical isometric embedding of  $M$  into  $\text{Lip}_0(M)^*$  is denoted by  $\delta$  and defined by  $\langle f, \delta(x) \rangle = f(x)$  for all  $x \in M$  and all  $f \in \text{Lip}_0(M)$ . We denote by  $\mathcal{F}(M)$  the norm-closed linear space of  $\delta(M)$  in the dual space  $\text{Lip}_0(M)^*$ , which is usually called the *Lipschitz-free space over  $M$* .

An *elementary molecule* in  $\mathcal{F}(M)$  is an element of the form

$$m_{x,y} := \frac{\delta(x) - \delta(y)}{d(x, y)}$$

where  $x, y \in M$  are such that  $x \neq y$ . A *molecule* in  $\mathcal{F}(M)$  is understood to be a finite sum of elementary molecules. When  $\mu$  is a molecule, we have that it has an optimal representation [26, Proposition 3.16]. In other words, we can write  $\mu$  as follows

$$\mu = \sum_{i=1}^n \lambda_i m_{x_i, y_i}, \quad \text{where } \lambda_i > 0 \quad \text{with} \quad \|\mu\| = \sum_{i=1}^n \lambda_i.$$

It is easy to check and well known that for every  $\mu \in S_{\mathcal{F}(M)}$  and  $\varepsilon > 0$ , there exist  $x_n \neq y_n \in M$  and  $(a_n) \in \ell_1$  with  $a_n > 0$  such that

$$\mu = \sum_n a_n m_{x_n, y_n}$$

and  $\sum_n a_n < 1 + \varepsilon$ . We send the reader to the survey [20] and the monograph [26] for a complete background.

Finally, recall that a metric space  $M$  is called *uniformly discrete* when  $\inf\{d(x, y) : x \neq y \in M\} > 0$ .

**2.2. Strongly subdifferentiable points.** The norm  $\|\cdot\|$  of a Banach space  $X$  is *strongly subdifferentiable* (SSD, for short) at  $x \in S_X$  when the one-sided limit

$$\lim_{t \rightarrow 0^+} \frac{\|x + th\| - 1}{t}$$

exists uniformly in  $h \in B_X$ . For simplicity, we shall say that  $x \in S_X$  is an SSD point if the norm of  $X$  is SSD at  $x$ . When every  $x \in S_X$  is an SSD point,  $X$  will be said to be SSD.

One can realize immediately that a norm of a Banach space  $X$  is Fréchet differentiable at a point  $x \in S_X$  if and only if it is both Gâteaux and SSD at this point. We have the following characterization due to Franchetti and Payá for SSD points (we will be using such a characterization without any explicit reference throughout). A point  $x \in S_X$  is an SSD

point if and only if  $x$  strongly exposes the set  $D(x) := \{x^* \in S_{X^*} : x^*(x) = 1\}$ , i.e., given  $\varepsilon > 0$ , there exists  $\eta(\varepsilon, x) > 0$  such that whenever  $x^*(x) > 1 - \eta(\varepsilon, x)$  for some  $x^* \in B_{X^*}$ , we have that  $\text{dist}(x^*, D(x)) < \varepsilon$  (see [16, Theorem 1.2]).

### 3. MAIN RESULTS

In this section, we start by presenting some introductory results as examples and remarks on  $\mathcal{F}(M)$  for some metric spaces  $M$ .

**3.1. Some examples and remarks.** Recall that, for a Banach space  $X$ , a point  $x \in S_X$  is a point of Gâteaux differentiability if and only if there exists a unique  $f \in S_{X^*}$  such that  $f(x) = 1$ . Let us recall also that given two points  $x, y \in M$ , we define the metric segment between them as the set

$$[x, y] := \{z \in M : d(x, z) + d(y, z) = d(x, y)\}.$$

The next result states that an element, which is a limit in norm of a convex series of elementary molecules, is a Gâteaux differentiable point in the Lipschitz-free over the union of segments determined by the representation of the given element. Although the proof is straightforward, we include it for completeness.

**Proposition 3.1.** *Let  $M$  be a pointed metric space and  $I$  be an index set. If  $\mu = \sum_{n \in I} a_n m_{x_n, x_{n+1}} \in S_{\mathcal{F}(M)}$  for some  $\{x_n : n \in I\} \subseteq M$  containing 0 and  $a_n > 0$  with  $\sum_{n \in I} a_n = 1$ , then  $\mu$  is a Gâteaux differentiable point in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$ .*

*Proof.* If  $f \in S_{\text{Lip}_0(M)}$  satisfies that  $\langle f, \mu \rangle = 1$ , then by using a convex argument, we have that  $\langle f, m_{x_n, x_{n+1}} \rangle = 1$  for every  $n \in I$ . For simplicity, say  $I = \{n_1, n_2, \dots\}$  and  $x_{n_1} = 0$ . As  $f$  attains its Lipschitz constant between  $x_{n_1} = 0$  and  $x_{n_2}$ , the value of  $f$  on the segment  $[x_{n_1}, x_{n_2}]$  is determined, i.e.,  $f(t) = d(x_{n_1}, t)$  for  $t \in [x_{n_1}, x_{n_2}]$ . From the fixed value of  $f(x_{n_2}) = d(x_{n_1}, x_{n_2})$ , we observe that the value of  $f$  is also determined on  $[x_{n_2}, x_{n_3}]$ . In this way, the value of  $f$  is uniquely determined on the entire segment  $[x_n, x_{n+1}]$  for every  $n \in I$ , which proves that the molecule  $\mu$  must be a point of Gâteaux differentiability in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$  as we wanted.  $\square$

**Proposition 3.2.** *Let  $M$  be a pointed metric space and  $I$  be an index set. Let  $\mu = \sum_{n \in I} a_n m_{x_n, x_{n+1}} \in S_{\mathcal{F}(M)}$  for some  $\{x_n : n \in I\} \subseteq M$  containing 0 and  $a_n > 0$  with  $\sum_{n \in I} a_n = 1$ . Consider the following statements.*

- (1)  $\mu$  is an SSD point in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$ .
- (2)  $\mu$  is a Fréchet differentiable point in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$ .
- (3)  $\cup_{n \in I} [x_n, x_{n+1}]$  is uniformly discrete and bounded.

Then (1)  $\Leftrightarrow$  (2)  $\Rightarrow$  (3).

*Proof.* Notice from Proposition 3.1 that the element  $\mu$  is a Gâteaux differentiable point in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$ . Thus,  $\mu$  is a SSD point in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$  if and only if  $\mu$  is a Fréchet differentiable point in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$ . This proves that (1) and (2) are equivalent. Now, suppose that  $\mu$  is Fréchet differentiable in  $\mathcal{F}(\cup_{n \in I} [x_n, x_{n+1}])$ . Then, the set  $\cup_{n \in I} [x_n, x_{n+1}]$  must be uniformly discrete and bounded due to [4, Theorem 2.4].  $\square$

Implication (3)  $\Rightarrow$  (2) in Proposition 3.2 is *not* true in general.

**Example 3.3.** Consider the equilateral metric space  $M = \mathbb{N} \cup \{0\}$ , i.e.,  $d(0, n) = 1$  for every  $n \in \mathbb{N}$  and  $d(n, m) = 2$  for all  $n \neq m \in \mathbb{N}$ . The operator  $\ell_1$  to  $\mathcal{F}(M)$  given by  $\delta(e_n) := \delta(n)$  for every  $n \in \mathbb{N}$  is a linear surjective isometry. Note that  $\cup_{n \in \mathbb{N}} [n-1, n] = M$  is uniformly discrete and bounded while  $\mu = \sum_{n=1}^{\infty} a_n m_{n-1, n}$  is not an SSD point in  $\mathcal{F}(M)$  since  $\mathcal{F}(M) = \ell_1$  does not have Fréchet differentiable points [21].

*Remark 3.4.* Let us observe that Proposition 3.1 *does not* hold true for the more general case where  $\mu$  is given as  $\sum_n a_n m_{x_n, y_n}$ , with  $\sum_n a_n = 1$  and  $x_n \neq y_n \in M$ ,  $n \in I$ . Indeed, consider the metric space  $M = \{0, 1, 2, 3\}$  in  $\mathbb{R}$ . Define the molecule  $\mu := \frac{1}{2}m_{1,0} + \frac{1}{2}m_{3,2}$ . Then, the Lipschitz functions  $f := (0, 1, 2, 3)$  and  $g := (0, 1, 0, 1)$  both attain their norms at  $\mu$ . So,  $\mu$  cannot be a Gâteaux differentiable point. We thank Ramón J. Aliaga for pointing this out.

The next result implies that unless the cardinality of  $M$  is finite, the set of SSD points in  $\mathcal{F}(M)$  should differ from the unit sphere, as this would imply that the space  $\mathcal{F}(M)$  is an Asplund space.

**Proposition 3.5.** *Let  $M$  be a pointed metric space. The following are equivalent.*

- (a)  $\mathcal{F}(M)$  admits a renorming which is SSD.
- (b) Every renorming of  $\mathcal{F}(M)$  is SSD.
- (c)  $\mathcal{F}(M)$  is finite dimensional.

*Proof.* Let us prove that (a) implies (c). Note that a Lipschitz-free space  $\mathcal{F}(M)$  is an Asplund space only when it is finite-dimensional (that is, the cardinality of  $M$  is finite) as any infinite-dimensional Lipschitz-free space contains an isomorphic copy of  $\ell_1$  (see [23, Proposition 3]). For (c) implying (b), notice that every finite-dimensional Banach space is SSD (as a consequence of Dini's theorem).  $\square$

**3.2. Locally uniformly non-aligned metric spaces.** In this subsection, our aim is to introduce a metric space  $M$  where *every* elementary molecule exhibits the property of being an SSD point in  $\mathcal{F}(M)$ . For any  $x, y, z \in M$ , the *Gromov product of  $x, y$  with respect to  $z$*  is defined as

$$G_z(x, y) := d(x, z) + d(z, y) - d(x, y)$$

(some authors consider the Gromov product as the above value divided by 2—see, for instance, [5, 6]; in our context, this would not make much of a difference).

**Definition 3.6.** Let  $M$  be a pointed metric space and  $x, y \in M$  with  $x \neq y$ . We say that the pair  $(x, y)$  satisfies *property (G)* if there exists  $\eta := \eta(x, y) > 0$  such that  $G_z(x, y) > \eta$  for every  $z \in M \setminus \{x, y\}$ . The metric space  $M$  is said to be *locally uniformly non-aligned* if every pair  $(x, y)$  of points in  $M$  satisfies property (G).

If  $(x, y)$  satisfies property (G), then it is clear that  $[x, y] = \{x, y\}$ . Let us mention that  $m_{x,y}$  is an extreme point of  $B_{\mathcal{F}(M)}$  if and only if  $G_z(x, y) > 0$  for every  $z \in M \setminus \{x, y\}$  (see [3, Theorem 1.1]). Notice also that a locally uniformly non-aligned metric space does




FIGURE 1. Example 3.7

not need to be uniformly discrete (for instance, take  $M = \{(1, 0, \dots, 0, \alpha_n, 0, 0, \dots) : n \geq 2\} \cup \{0\} \subseteq c_0$ , where  $(\alpha_n)$  is a sequence of positive real numbers converging to 0).

**Example 3.7.** Let  $M = \{x = 0, y, z_1, z_2, \dots\}$  be such that  $d(x, y) = d(z_n, z_m) = 1$ ,  $d(x, z_n) = \frac{1}{2}$  and  $d(y, z_n) = \frac{1}{2} + \varepsilon_n$ , where  $(\varepsilon_n) \subseteq \mathbb{R}^+$  such that  $\varepsilon_n \rightarrow 0$  as  $n \rightarrow \infty$  (see Figure 1). It is clear that  $G_{z_n}(x, y) = \varepsilon_n \rightarrow 0$  as  $n \rightarrow \infty$ ; so  $(x, y)$  fails to have property (G).

To present a characterization for property (G) of a pair of points  $(x, y)$ , we introduce the following notion.

**Definition 3.8.** Let  $M$  be a pointed metric space and  $x \neq y \in M$ . We say that  $f \in S_{\text{Lip}_0(M)}$  *peaks at*  $(x, y)$  if  $f(m_{x,y}) = 1$  and

$$\sup\{|f(m_{p,q})| : (p, q) \neq (x, y)\} < \gamma < 1 \text{ for some } \gamma > 0.$$

Let us point out that our Definition 3.8 is different from the one introduced in [17, Definition 5.2], while these two concepts coincide when the given points  $x$  and  $y$  are isolated points. For  $x \neq y \in M$ , recall also the auxiliary Lipschitz function  $f_{x,y} \in S_{\text{Lip}_0(M)}$  given by

$$f_{x,y}(z) := \frac{d(x, y)}{2} \frac{d(z, y) - d(z, x)}{d(z, y) + d(z, x)}$$

(see [17, Lemma 3.6] and [24]).

**Proposition 3.9.** *Let  $M$  be a pointed metric space and  $x \neq y \in M$  be given. The following are equivalent.*

- (1) *The pair  $(x, y)$  satisfies property (G).*
- (2) *The function  $f_{x,y}$  peaks at  $(x, y)$ .*
- (3) *There exists  $f \in S_{\text{Lip}_0(M)}$  which peaks at  $(x, y)$ .*

*Proof.* (1)  $\Leftrightarrow$  (2): Suppose that  $f_{x,y}$  peaks at  $(x, y)$  (witnessed by  $0 < \gamma < 1$ ) and let  $z \in M \setminus \{x, y\}$  be given. Notice first that  $f_{x,y}(x) = \frac{d(x,y)}{2}$ . Then,

$$|f_{x,y}(m_{x,z})| = \frac{d(x,y)}{2d(x,z)} \left| 1 - \left( \frac{d(z,y) - d(z,x)}{d(z,y) + d(z,x)} \right) \right| = \frac{d(x,y)}{d(z,y) + d(z,x)} < \gamma.$$

This implies that

$$G_z(x, y) = d(x, z) + d(z, y) - d(x, y) > \frac{1}{\gamma}d(x, y) - d(x, y) = \left( \frac{1}{\gamma} - 1 \right) d(x, y).$$

If we define  $\eta := \left( \frac{1}{\gamma} - 1 \right) d(x, y) > 0$ , we are done. Conversely, let  $\eta := \eta(x, y)$  be the one given by property (G) for the pair  $(x, y)$ . It is clear that  $\langle f_{x,y}, m_{x,y} \rangle = 1$ . Now, for any pair  $(p, q) \neq (x, y)$ , we get

$$\langle f_{x,y}, m_{p,q} \rangle \leq \frac{d(x, y)}{\max\{d(x, p) + d(p, y), d(x, q) + d(q, y)\}} \leq \frac{d(x, y)}{d(x, y) + \eta} < 1;$$

hence  $f_{x,y}$  peaks at  $(x, y)$ .

It remains to show that (3)  $\Rightarrow$  (1). Let  $f \in S_{\text{Lip}_0(M)}$  peak at  $(x, y)$  with constant  $0 < \gamma < 1$  and let  $z \in M \setminus \{x, y\}$ . Observe that

$$\begin{aligned} G_z(x, y) &= d(x, z) + d(z, y) - d(x, y) > \frac{1}{\gamma}(f(x) - f(z)) + \frac{1}{\gamma}(f(z) - f(y)) - d(x, y) \\ &= \frac{1}{\gamma}(f(x) - f(y)) - d(x, y) \\ &= \left( \frac{1}{\gamma} - 1 \right) d(x, y) \end{aligned}$$

for every  $z \in M \setminus \{x, y\}$ . This shows that  $(x, y)$  satisfies property (G).  $\square$

Now we are ready to prove our main interest in introducing the aforementioned property.

**Theorem 3.10.** *Let  $M$  be a pointed metric space and  $x, y \in M$  with  $x \neq y$ . If  $(x, y)$  has property (G), then the elementary molecule  $m_{x,y}$  is an SSD point in  $\mathcal{F}(M)$ .*

*Proof.* By Proposition 3.9, there exists  $f \in S_{\text{Lip}_0(M)}$  such that  $f$  peaks at  $(x, y)$  with constant  $0 < \gamma < 1$ . Let  $\varepsilon > 0$  and take  $g \in S_{\text{Lip}_0(M)}$  to be such that  $\langle g, m_{x,y} \rangle > 1 - \gamma_\varepsilon$ , where  $\gamma_\varepsilon > 0$  satisfies the following condition

$$0 < \gamma_\varepsilon < \frac{\varepsilon}{4 - \varepsilon}(1 - \gamma).$$

Put  $h := \left(1 - \frac{\varepsilon}{4}\right)g + \frac{\varepsilon}{4}f$  and observe that

$$\langle h, m_{x,y} \rangle > \left(1 - \frac{\varepsilon}{4}\right)(1 - \gamma_\varepsilon) + \frac{\varepsilon}{4}.$$

By the choice of  $\gamma_\varepsilon$ , we have that for  $(p, q) \neq (x, y)$ ,

$$|\langle h, m_{p,q} \rangle| \leq \left(1 - \frac{\varepsilon}{4}\right) + \frac{\varepsilon}{4}\gamma < \langle h, m_{x,y} \rangle.$$

It follows that  $\|h\| = \langle h, m_{x,y} \rangle > \left(1 - \frac{\varepsilon}{4}\right)(1 - \gamma_\varepsilon) + \frac{\varepsilon}{4}$ . Finally, considering  $\hat{h} := \frac{h}{\|h\|}$ , we get

$$\|\hat{h} - g\| \leq (1 - \|h\|) + \frac{\varepsilon}{2} < 1 - \left(1 - \frac{\varepsilon}{4}\right)(1 - \gamma_\varepsilon) + \frac{\varepsilon}{4}.$$

Now it is enough to take small enough  $\varepsilon$  and  $\gamma_\varepsilon$ .  $\square$

As a direct consequence of Proposition 3.9 and Theorem 3.10, we have the following corollary.

**Corollary 3.11.** *Let  $M$  be a pointed metric space. If  $M$  is locally uniformly non-aligned, then every elementary molecule is an SSD point in  $\mathcal{F}(M)$ .*

As a matter of fact, the idea of the proof of Theorem 3.10 will be important in a more refined way when we prove our main result, Theorem 3.16.

*Remark 3.12.* The sufficient condition of having property (G) for a pair of points does not characterize SSD elementary molecules. In order to see this, let us consider the following examples.

- (1) Let  $M = \mathbb{N} \cup \{0\}$ . Then the linear operator from  $\mathcal{F}(M)$  to  $\ell_1$  sending  $\delta(n)$  to  $\sum_{k=1}^n e_k$  for  $n \in \mathbb{N}$  is a surjective isometry. In this case, every elementary molecule  $m_{x,y}$  with  $x \neq y \in M$  is an SSD point in  $\mathcal{F}(M)$ . However, it is easy to check that  $m_{x,y}$  satisfies property (G) if and only if  $|x - y| = 1$ .
- (2) Consider  $\mathfrak{C}$  to be the usual middle third Cantor set with the usual distance. The set  $\mathfrak{C}$  may be written as  $\mathfrak{C} = [0, 1] \setminus (\bigcup_{n=1}^{\infty} I_n)$ , where the sequence of intervals  $I_n := ]a_n, b_n[$  is an enumeration of the middle thirds we are taking on each step (in particular,  $I_1 = ]\frac{1}{3}, \frac{2}{3}[$ ,  $I_2 = ]\frac{1}{9}, \frac{2}{9}[$ ,  $I_3 = ]\frac{7}{9}, \frac{8}{9}[$ , etc.). It is well known that  $\mathcal{F}(\mathfrak{C})$  is isometrically isomorphic to  $\ell_1$ . Observe that  $m_{a_n, b_m}$  is an SSD point in  $\mathcal{F}(\mathfrak{C})$  if and only if  $n = m$ , while  $m_{x,y}$  fails to have property (G) for any  $x \neq y \in \mathfrak{C}$ .

*Remark 3.13.* Recall the metric space  $M$  and the points  $x \neq y \in M$  from Example 3.7. It was previously observed that  $(x, y)$  fails property (G) (equivalently, thanks to Proposition 3.9, there is no  $f \in S_{\text{Lip}_0(M)}$  that peaks at  $(x, y)$ ). However, as we observed in Remark 3.12, the fact that  $(x, y)$  lacks property (G) does not provide any information about whether  $m_{x,y}$  is an SSD point in  $\mathcal{F}(M)$ .

We claim that, in this case,  $m_{x,y}$  is indeed an SSD point in  $\mathcal{F}(M)$ . To this end, let  $\varepsilon > 0$  be given and let us find  $n_0 \in \mathbb{N}$  such that  $0 < \varepsilon_n < \varepsilon$  for all  $n > n_0$ . We consider  $M_0 := \{x = 0, y, z_1, \dots, z_{n_0}\}$ . Take  $0 < \gamma < \varepsilon$  sufficiently small so that whenever  $g \in S_{\text{Lip}_0(M_0)}$  satisfies that  $g(m_{x,y}) > 1 - \gamma$ , there exists  $h \in S_{\text{Lip}_0(M_0)}$  so that  $h(m_{x,y}) = 1$  and  $\|h - g\| < \varepsilon$ . Now, suppose that  $f \in S_{\text{Lip}_0(M)}$  satisfies that  $f(m_{x,y}) > 1 - \gamma$ . Then

$$\begin{aligned} 1 = d(x, y) &< \frac{f(x) - f(z_n) + f(z_n) - f(y)}{1 - \gamma} &\leq \frac{d(x, z_n) + d(z_n, y)}{1 - \gamma} \\ &= \frac{1 + \varepsilon_n}{1 - \gamma} = 1 + \frac{\varepsilon_n + \gamma}{1 - \gamma}. \end{aligned}$$



Thus, we get that  $d(z_n, y) - (f(z_n) - f(y))$  and  $d(x, z_n) - (f(x) - f(z_n))$  are smaller than or equal to  $\varepsilon_n + \gamma$ . This yields that

$$f(m_{x,z_n}) \geq 1 - (2\varepsilon_n + 2\gamma) \quad \text{and} \quad f(m_{z_n,y}) \geq 1 - \frac{2\varepsilon_n + 2\gamma}{1 + 2\varepsilon_n}. \quad (3.1)$$

Letting  $\tilde{f} := \|f|_{M_0}\|^{-1}f|_{M_0}$ , we have  $\tilde{f}(m_{x,y}) > 1 - \gamma$ . Thus, there exists  $h \in S_{\text{Lip}_0(M_0)}$  such that  $h(m_{x,y}) = 1$  and  $\|h - \tilde{f}\| < \varepsilon$ . Extend  $h$  to a norm one Lipschitz map on  $M$  and denote it again by  $h$ . As before, we observe that

$$h(m_{x,z_n}) \geq 1 - 2\varepsilon_n \quad \text{and} \quad h(m_{z_n,y}) \geq 1 - \frac{2\varepsilon_n}{1 + 2\varepsilon_n}. \quad (3.2)$$

Finally, we estimate  $\|h - f\|$  by dividing some cases. We use (3.1) and (3.2).

(1)  $|(h - f)(m_{x,y})| < \gamma < \varepsilon$ .

(2) If  $1 \leq n \leq n_0$ , then

$$|(h - f)(m_{x,z_n})| \leq \|h|_{M_0} - \tilde{f}\| + \|\tilde{f}\| - 1 < \varepsilon + \gamma < 2\varepsilon.$$

Similarly,  $|(h - f)(m_{z_n,y})| < 2\varepsilon$ .

(3) If  $n > n_0$ , then

$$|(h - f)(m_{x,z_n})| \leq 2\varepsilon_n + 2\gamma < 2\varepsilon + 2\varepsilon = 4\varepsilon.$$

Similarly,  $|(h - f)(m_{z_n,y})| \leq \frac{2\varepsilon_n + 2\gamma}{1 + 2\varepsilon_n} < 4\varepsilon$ .

(4) Let  $i, j \in \mathbb{N}$ . If  $1 \leq i, j \leq n_0$ , then  $|(h - f)(m_{z_i,z_j})| \leq \|h|_{M_0} - f|_{M_0}\| < 2\varepsilon$ . Now suppose that  $1 \leq i \leq n_0$  and  $j > n_0$ . Let us recall that  $d(z_i, z_j) = 1$  and  $d(x, z_i) = d(x, z_j) = 1/2$ . Using this and the previous cases, we get that

$$\begin{aligned} |(h - f)(m_{z_i,z_j})| &\leq d(x, z_i)|(h - f)(m_{x,z_i})| + d(x, z_j)|(h - f)(m_{x,z_j})| \\ &< \frac{2\varepsilon}{2} + \frac{4\varepsilon}{2} = 3\varepsilon. \end{aligned}$$

Similarly for  $i, j > n_0$ , we have  $|(h - f)(m_{z_i,z_j})| < 4\varepsilon$ .

Therefore, by using (1)-(4), we observe that  $h \in S_{\text{Lip}_0(M)}$  is such that  $h(m_{x,y}) = 1$  and satisfies  $\|h - f\| \leq 4\varepsilon$ . This shows that the elementary molecule  $m_{x,y}$  is an SSD point in  $\mathcal{F}(M)$  as we wanted to show.

Let us now conclude this subsection by considering some relations between property (G), uniformly Gromov rotundness (see [5, 6]), and uniformly concaveness. Let  $M$  be a pointed metric space and  $A$  a set of elementary molecules in  $M$ . Recall that  $A$  is *uniformly Gromov rotund* if there exists  $\varepsilon = \varepsilon(A) > 0$  such that

$$G_z(x, y) > \varepsilon \min\{d(x, z), d(y, z)\}$$

whenever  $m_{x,y} \in A$  and  $z \in M \setminus \{x, y\}$ . For simplicity, let us say that the pair  $(x, y)$  is uniformly Gromov rotund if the singleton  $\{m_{x,y}\}$  is uniformly Gromov rotund.

Also, recall that a metric space  $M$  is called *uniformly concave* if for every  $x \neq y \in M$  and  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $G_z(x, y) > \delta$  for every  $z \in M$  satisfying  $\min\{d(x, z), d(y, z)\} \geq \varepsilon$  (see [26, Definition 3.3]). Let us say that a pair of points  $(x, y)$  is

uniformly concave if for any  $\varepsilon > 0$ , there exists  $\delta = \delta(x, y, \varepsilon) > 0$  such that  $G_z(x, y) > \delta$  for every  $z \in M$  satisfying  $\min\{d(x, z), d(y, z)\} \geq \varepsilon$ .

**Proposition 3.14.** *Let  $M$  be a pointed metric space. Let  $x, y \in M$  be such that  $x \neq y$ .*

- (1) *If the pair  $(x, y)$  satisfies property (G), then it is uniformly Gromov rotund.*
- (2) *If the pair  $(x, y)$  is uniformly Gromov rotund, then it is uniformly concave.*
- (3) *If  $x$  and  $y$  are isolated points in  $M$ , then  $(x, y)$  satisfies property (G) if and only if  $(x, y)$  is uniformly Gromov rotund if and only if  $(x, y)$  is uniformly concave.*

*Proof.* (1): Suppose that  $M$  is any metric space and that  $(x, y)$  satisfies property (G). Then, there exists  $\eta = \eta(x, y) > 0$  such that  $G_z(x, y) > \eta$  for every  $z \in M \setminus \{x, y\}$ . We have two cases. If  $d(z, y) > d(x, y)$  or  $d(x, z) > d(x, y)$ , then we have that  $G_z(x, y) > d(x, z)$  or  $G_z(x, y) > d(y, z)$ . In any case, we have that  $G_z(x, y) > \min\{d(x, z), d(y, z)\}$ . On the other hand, if  $d(z, y)$  and  $d(x, z)$  are both  $\leq d(x, y)$ , then we have that

$$G_z(x, y) > \eta \geq \frac{\eta}{d(x, y)} \max\{d(x, z), d(z, y)\}.$$

Putting together these cases, we get that

$$G_z(x, y) > \min \left\{ 1, \frac{\eta}{d(x, y)} \right\} \min\{d(x, z), d(z, y)\}$$

for every  $z \in M \setminus \{x, y\}$ . This shows that  $\varepsilon = \varepsilon(\{x, y\}) := \min \left\{ 1, \frac{\eta}{d(x, y)} \right\}$  does the job.

(2): This is obvious by definition.

(3): Suppose now that  $x, y \in M$  are isolated and  $(x, y)$  is uniformly concave. Take  $r_x, r_y > 0$  so that  $B(x, r_x) = \{x\}$  and  $B(y, r_y) = \{y\}$ , respectively. Considering  $\varepsilon(x, y) := \min\{r_x, r_y\}$ , we can find  $\delta = \delta(x, y, \varepsilon(x, y)) > 0$  such that  $G_z(x, y) > \delta$  for every  $z \in M \setminus \{x, y\}$  (noting that  $\min\{d(x, z), d(y, z)\} \geq \min\{r_x, r_y\}$ ). This proves that  $(x, y)$  has property (G).  $\square$

*Remark 3.15.* The converse of (1) in Proposition 3.14 is *not* true in general. Indeed, consider  $M = \{x = 0, y, z_1, z_2, \dots\}$  endowed with the metric given by  $d(x, y) = 1$ ,  $d(x, z_n) = \frac{1}{2n}$ ,  $d(y, z_n) = 1 - \frac{1}{4n}$  and  $d(z_n, z_m) = \frac{1}{2n} + \frac{1}{2m}$  for every  $n, m \in \mathbb{N}$  with  $n \neq m$ . On the one hand, we have that

$$G_{z_n}(x, y) = \frac{1}{2n} + 1 - \frac{1}{4n} - 1 > \varepsilon \min\{d(x, z_n), d(y, z_n)\}$$

for every  $n \in \mathbb{N}$ , whenever  $0 < \varepsilon < \frac{1}{2}$ . This shows that the pair  $(x, y)$  is uniformly Gromov rotund, while  $G_{z_n}(x, y) \rightarrow 0$  as  $n \rightarrow \infty$ ; so  $(x, y)$  fails property (G).

**3.3. The subset of SSD points is dense.** Let  $M$  be a pointed metric space with metric  $d$ . One natural way of producing a locally uniformly non-aligned metric space from  $M$  is to consider the metric space  $(M, d_\gamma)$  with  $\gamma > 0$ , where  $d_\gamma : M \times M \rightarrow \mathbb{R}$  is given by  $d_\gamma(x, y) := d(x, y) + \gamma$  for every  $x, y \in M$  with  $x \neq y$  and  $d_\gamma(x, x) := 0$  for every  $x \in M$ . It is obvious from Corollary 3.11 that every elementary molecule in  $\mathcal{F}(M, d_\gamma)$  is an SSD

point. We have the following result, which will lead to the conclusion that every molecule in  $\mathcal{F}(M, d_\gamma)$  is indeed an SSD point.

**Theorem 3.16.** *Let  $(M, d)$  be a pointed metric space and  $\gamma > 0$ . Then every element of*

$$\left\{ \sum_{i=1}^n \lambda_i \left( \frac{\delta(x_i) - \delta(y_i)}{d(x_i, y_i) + \gamma} \right) : \exists f \in S_{\text{Lip}_0(M, d)} : f(m_{x_i, y_i}) = 1, \lambda_i > 0, \sum_{i=1}^n \lambda_i = 1 \text{ and } n \in \mathbb{N} \right\}$$

*is an SSD point in  $\mathcal{F}(M, d_\gamma)$ .*

*Proof.* Let  $\mu \in \mathcal{F}(M, d_\gamma)$  be an element of the following form

$$\mu = \sum_{i=1}^n \lambda_i \left( \frac{\delta(x_i) - \delta(y_i)}{d(x_i, y_i) + \gamma} \right)$$

where there exists  $f \in S_{\text{Lip}_0(M, d)}$  such that  $f(x_i) - f(y_i) = d(x_i, y_i)$  for every  $i = 1, \dots, n$  and  $\sum_{i=1}^n \lambda_i = 1$  with  $\lambda_i > 0$ . Let us notice that the sets  $\{x_i\}_{i=1}^n$  and  $\{y_i\}_{i=1}^n$  are disjoint. Indeed, let  $g \in S_{\text{Lip}_0(M, d_\gamma)}$  be such that  $g(x_i) - g(y_i) = d(x_i, y_i) + \gamma$  for every  $i = 1, \dots, n$ . Assuming for simplicity that  $x_1 = y_2$ , we have

$$\begin{aligned} g(x_2) &= g(y_2) + d(x_2, y_2) + \gamma = g(y_1) + d(x_1, y_1) + d(x_2, x_1) + 2\gamma \\ &\geq g(y_1) + d(y_1, x_2) + 2\gamma. \end{aligned}$$

This yields a contradiction since  $g(m_{x_2, y_1})$  should be smaller than or equal to 1. Setting  $N := \{0, x_1, y_1, \dots, x_n, y_n\} \subseteq M$ , it is clear that  $f|_N \in S_{\text{Lip}_0(N, d)}$  satisfies that  $|f|_N| \leq \beta := \max\{d(0, q) : q \in N\}$ . Applying the McShane extension, we can extend  $f|_N$  to  $(M, d)$  whose values are contained in  $[-\beta, \beta]$  without increasing its norm. We still denote this extension by  $f$ .

Now let us define  $f_\gamma \in \text{Lip}_0(M, d_\gamma)$  as

$$f_\gamma(z) := \begin{cases} f(x_i) + \gamma, & \text{if } z = x_i, \forall i = 1, \dots, n, \\ f(y_i), & \text{if } z = y_i, \forall i = 1, \dots, n, \\ f(z) + \frac{\gamma}{2}, & \text{if } z \notin N. \end{cases}$$

Observe that  $\|f_\gamma\|_{\text{Lip}_0(M, d_\gamma)} = 1$ . As a matter of fact, for  $1 \leq i, j \leq n$  and  $p \notin N$  we have

- (1)  $|f_\gamma(x_i) - f_\gamma(y_j)| = |f(x_i) + \gamma - f(y_j)| \leq d(x_i, y_j) + \gamma;$
- (2)  $|f_\gamma(x_i) - f_\gamma(p)| = |f(x_i) + \gamma - (f(p) + \frac{\gamma}{2})| \leq d(x_i, p) + \frac{\gamma}{2};$
- (3)  $|f_\gamma(y_j) - f_\gamma(p)| = |f(y_j) - (f(p) + \frac{\gamma}{2})| \leq d(y_j, p) + \frac{\gamma}{2}.$

Thus, we have

$$\langle f_\gamma, \mu \rangle = \sum_{i=1}^n \langle f_\gamma, m_{x_i, y_i} \rangle = \sum_{i=1}^n \frac{\lambda_i}{d(x_i, y_i) + \gamma} (f(x_i) + \gamma - f(y_i)) = \sum_{i=1}^n \lambda_i = 1.$$

As it is clear that  $\|\mu\|_{\mathcal{F}(M, d_\gamma)} \leq 1$ , we actually obtain that  $\|\mu\|_{\mathcal{F}(M, d_\gamma)} = 1$ . Let us note that

$$|f_\gamma(p)| \leq \beta + \frac{\gamma}{2} \text{ for every } p \notin N. \quad (3.3)$$

We also consider  $\xi : \mathbb{R}_{\geq 0} \rightarrow [0, 1]$  to be defined as  $\xi(t) = 1$  for  $0 \leq t \leq \beta$ ,

$$\xi(t) = -\frac{1}{T-\beta}(t-T) \quad \text{for } \beta < t \leq T$$

and  $\xi(t) = 0$  for all  $t \geq T$ , where  $T > 0$  is sufficiently large (which will be specified later). Now, we define  $G_\gamma \in \text{Lip}_0(M, d_\gamma)$  given by

$$G_\gamma(z) := f_\gamma(z) \xi(d(0, z))$$

for every  $z \in M$ . By using the definitions of  $\xi$  and  $f_\gamma$ , we have immediately that  $\langle G_\gamma, \mu \rangle = 1$ .

**Claim A:**  $\sup\{|\langle G_\gamma, m_{p,q} \rangle| : p \notin N \text{ or } q \notin N\} < 1$ .

In order to prove the claim, we consider different cases. Let  $p \notin N$  or  $q \notin N$  with  $p \neq q$ . If  $d(0, p), d(0, q) \leq \beta$ , then

$$\begin{aligned} |\langle G_\gamma, m_{p,q} \rangle| &= \frac{|f_\gamma(p)\xi(d(0, p)) - f_\gamma(q)\xi(d(0, q))|}{d(p, q) + \gamma} \\ &= \frac{|f_\gamma(p) - f_\gamma(q)|}{d(p, q) + \gamma} \\ &\stackrel{(3.3)}{\leq} \frac{d(p, q) + \frac{\gamma}{2}}{d(p, q) + \gamma} \leq \frac{2\beta + \frac{\gamma}{2}}{2\beta + \gamma} < 1. \end{aligned}$$

Otherwise, we may (and we do) assume without loss of generality that  $d(0, q) > \beta$  (in particular,  $q \notin N$ ). In this case, we have that

$$\begin{aligned} |\langle G_\gamma, m_{p,q} \rangle| &\leq \xi(d(0, p)) \frac{|f_\gamma(p) - f_\gamma(q)|}{d(p, q) + \gamma} + |f_\gamma(q)| \frac{|\xi(d(0, p)) - \xi(d(0, q))|}{d(p, q) + \gamma} \\ &\leq \frac{|f_\gamma(p) - f_\gamma(q) - \frac{\gamma}{2}|}{d(p, q) + \gamma} + |f_\gamma(q)| \frac{|\xi(d(0, p)) - \xi(d(0, q))| |d(0, p) - d(0, q)|}{|d(0, p) - d(0, q)| d(p, q) + \gamma} \\ &\stackrel{(3.3)}{\leq} \frac{|f(p) - f(q)| + \frac{\gamma}{2}}{d(p, q) + \gamma} + \left(\beta + \frac{\gamma}{2}\right) \left(\frac{1}{T-\beta}\right) \frac{d(p, q)}{d(p, q) + \gamma} =: (I) \end{aligned}$$

We shall see that (I) is strictly smaller than 1 uniformly for  $p$  and  $q$ , provided that  $T$  is sufficiently large. For simplicity, let us set

$$T_0 := \frac{\gamma}{4} \left( \frac{T-\beta}{\beta + \frac{\gamma}{2}} \right).$$

At this moment, say  $T > 0$  is large enough so that

$$\frac{2\beta + \frac{3\gamma}{2}}{T_0 + \gamma} + \left(\beta + \frac{\gamma}{2}\right) \left(\frac{1}{T-\beta}\right) < \frac{3}{4}. \quad (3.4)$$

We consider two cases. Suppose first that  $d(p, q) < T_0$ , which is equivalent to

$$\frac{\beta + \frac{\gamma}{2}}{T-\beta} < \frac{\gamma}{4} \frac{1}{d(p, q)}. \quad (3.5)$$

By (3.5), we get that

$$(I) \leq \frac{d(p, q) + \frac{\gamma}{2}}{d(p, q) + \gamma} + \left(\beta + \frac{\gamma}{2}\right) \left(\frac{1}{T - \beta}\right) \frac{d(p, q)}{d(p, q) + \gamma} < \frac{d(p, q) + \frac{3\gamma}{4}}{d(p, q) + \gamma} < \frac{T_0 + \frac{3\gamma}{4}}{T_0 + \gamma} < 1.$$

On the other hand, if  $d(p, q) \geq T_0$ , then by (3.4) we get that

$$(I) \leq \frac{2\beta + \frac{3\gamma}{2}}{T_0 + \gamma} + \left(\beta + \frac{\gamma}{2}\right) \left(\frac{1}{T - \beta}\right) < \frac{3}{4}.$$

This completes the proof of the claim. Thus, we can find  $K > 0$  sufficiently large such that the following holds true

$$\sup\{|\langle G_\gamma, m_{p,q} \rangle| : p \notin N \text{ or } q \notin N\} < \frac{K + \frac{\gamma}{2}}{K + \gamma} < 1.$$

Let us now move towards the end of the proof. Let  $\varepsilon > 0$  be given and  $g \in S_{\text{Lip}_0(M, d_\gamma)}$  be such that  $\langle g, \mu \rangle > 1 - \rho$ , where  $\rho = \rho(\varepsilon, \mu) > 0$  is small enough such that<sup>1</sup>

$$(1 - \sqrt{\varepsilon}) + \sqrt{\varepsilon} \left(\frac{K + \frac{\gamma}{2}}{K + \gamma}\right) + \frac{\beta\varepsilon}{\gamma} < (1 - \sqrt{\varepsilon})(1 - \rho) + \sqrt{\varepsilon}. \quad (3.6)$$

**Claim B:** There exists  $\psi \in S_{\text{Lip}_0(M, d_\gamma)}$  such that  $\langle \psi, \mu \rangle = 1$  and  $\|\psi - g\| \approx 0$ .

First, define  $h : (M, d_\gamma) \rightarrow \mathbb{R}$  by

$$h := (1 - \sqrt{\varepsilon})g + \sqrt{\varepsilon}G_\gamma.$$

Then, we have that  $\langle h, \mu \rangle > (1 - \sqrt{\varepsilon})(1 - \rho) + \sqrt{\varepsilon}$  and

$$|\langle h, m_{p,q} \rangle| \leq (1 - \sqrt{\varepsilon}) + \sqrt{\varepsilon} \left(\frac{K + \frac{\gamma}{2}}{K + \gamma}\right)$$

for every  $p \notin N$  or  $q \notin N$ . This implies that

$$\begin{aligned} \langle h, \mu \rangle &> (1 - \sqrt{\varepsilon})(1 - \rho) + \sqrt{\varepsilon} \\ &\stackrel{(3.6)}{>} (1 - \sqrt{\varepsilon}) + \sqrt{\varepsilon} \left(\frac{K + \frac{\gamma}{2}}{K + \gamma}\right) + \frac{\beta\varepsilon}{\gamma} \\ &> \sup\{|\langle h, m_{p,q} \rangle| : p \notin N \text{ or } q \notin N\}. \end{aligned}$$

In particular, this shows that  $\|h\| = \|h|_N\|$  with  $\langle h|_N, \mu \rangle > (1 - \sqrt{\varepsilon})(1 - \rho) + \sqrt{\varepsilon}$ . Consider  $\mu$  as an element of the finite-dimensional space  $\mathcal{F}(N, d_\gamma)$ , and recall that the norm of a finite-dimensional normed space is SSD. Thus, by letting  $\rho = \rho(\varepsilon, \mu)$  even smaller if

<sup>1</sup>Let us notice that (3.6) is indeed possible to be considered since

$$(1 - \sqrt{\varepsilon}) + \sqrt{\varepsilon} \left(\frac{K + \frac{\gamma}{2}}{K + \gamma}\right) + \frac{\beta\varepsilon}{\gamma} = 1 - \sqrt{\varepsilon} \left(1 - \frac{K + \frac{\gamma}{2}}{K + \gamma} - \frac{\beta\sqrt{\varepsilon}}{\gamma}\right)$$

is smaller than 1 for a sufficiently small  $\varepsilon > 0$ , and then we can take  $\rho = \rho(\varepsilon, \mu) > 0$  small enough in order to satisfy (3.6).

necessary, we can find a Lipschitz function  $\varphi \in \text{Lip}_0(N, d_\gamma)$  such that  $\|\varphi\| = \langle \varphi, \mu \rangle = \|h\|$  and  $\|\varphi - h|_N\|_{\text{Lip}_0(N, d_\gamma)} < \varepsilon$ . Now, we define  $\psi : (M, d_\gamma) \rightarrow \mathbb{R}$  by

$$\psi(p) := \begin{cases} \varphi(p), & \forall p \in N, \\ h(p), & \forall p \in M \setminus N \end{cases}$$

which will serve as the required element for our claim.

Notice that  $\langle \psi, \mu \rangle = \langle \varphi, \mu \rangle = \|\varphi\|$ . Moreover,  $\|\psi - h\|_{\text{Lip}_0(M, d_\gamma)}$  can be estimated as follows. If  $p, q \in N$ , then

$$\frac{|(\psi - h)(p) - (\psi - h)(q)|}{d(p, q) + \gamma} = \frac{|(\varphi - h|_N)(p) - (\varphi - h|_N)(q)|}{d(p, q) + \gamma} < \varepsilon.$$

On the other hand, if  $p \in M \setminus N$  and  $q \in N$ , then

$$\begin{aligned} \frac{|(\psi - h)(p) - (\psi - h)(q)|}{d(p, q) + \gamma} &= \frac{|(h - h)(p) - (\varphi - h)(q)|}{d(p, q) + \gamma} \\ &\leq \|h - h|_N\|_{\text{Lip}_0(N, d_\gamma)} \frac{\beta}{\gamma} < \frac{\beta\varepsilon}{\gamma}. \end{aligned}$$

Finally, the case when  $p, q \in M \setminus N$  is trivial. Consequently, we get that

$$\|\psi - h\|_{\text{Lip}_0(M, d_\gamma)} \leq \|\psi - h\|_{\text{Lip}_0(M, d_\gamma)} + \|h - g\|_{\text{Lip}_0(M, d_\gamma)} < \max\left\{\varepsilon, \frac{\beta\varepsilon}{\gamma}\right\} + 2\sqrt{\varepsilon}.$$

It remains to prove that  $\psi$  attains its norm at  $\mu$ . To this end, observe that if  $p, q \in M \setminus N$ , then

$$|\langle \psi, m_{p,q} \rangle| = |\langle h, m_{p,q} \rangle| \leq \|h\| = \|\varphi\| = \langle \psi, \mu \rangle.$$

If  $p, q \in N$ , then

$$|\langle \psi, m_{p,q} \rangle| = |\langle \varphi, m_{p,q} \rangle| \leq \|\varphi\| = \langle \psi, \mu \rangle.$$

Finally, if  $p \in N$  and  $q \in M \setminus N$ , then

$$\begin{aligned} |\langle \psi, m_{p,q} \rangle| &= \frac{|\psi(p) - \psi(q)|}{d(p, q) + \gamma} \leq \frac{|\varphi(p) - h(p)| + |h(p) - h(q)|}{d(p, q) + \gamma} \\ &\leq \frac{\|\varphi - h|_N\|_{\text{Lip}_0(N, d_\gamma)} \beta}{\gamma} + |\langle h, m_{p,q} \rangle| \\ &< \frac{\beta\varepsilon}{\gamma} + (1 - \sqrt{\varepsilon}) + \sqrt{\varepsilon} \left( \frac{K + \frac{\gamma}{2}}{K + \gamma} \right) \\ &\stackrel{(3.6)}{<} (1 - \sqrt{\varepsilon})(1 - \rho) + \sqrt{\varepsilon} \\ &< \|h\| = \|\varphi\| = \langle \psi, \mu \rangle. \end{aligned}$$

This completes the proof of the claim. Therefore, we conclude that  $\mu$  is an SSD point in  $\mathcal{F}(M, d_\gamma)$  as we wanted to prove.  $\square$

Now we are ready to prove the denseness of the SSD points in  $\mathcal{F}(M, d_\gamma)$  for every  $\gamma > 0$ . This is a consequence of Theorem 3.16.

**Corollary 3.17.** *Let  $(M, d)$  be a pointed metric space and  $\gamma > 0$ . Then every molecule in  $\mathcal{F}(M, d_\gamma)$  is an SSD point. In particular, the set of SSD points in  $\mathcal{F}(M, d_\gamma)$  is dense.*

*Proof.* Let  $\gamma > 0$  be arbitrary. For simplicity, let us put  $\tilde{d} := d_\gamma$  (then,  $\tilde{d}_\gamma = d_{2\gamma}$ ). Let  $\mu \in \mathcal{F}(M, \tilde{d}_\gamma)$  be a norm-one finitely supported element having an optimal representation and write

$$\mu = \sum_{i=1}^n \lambda_i \frac{\delta(x_i) - \delta(y_i)}{\tilde{d}(x_i, y_i) + \gamma} \in \mathcal{F}(M, \tilde{d}_\gamma)$$

with  $\sum_{i=1}^n \lambda_i = \|\mu\|_{\mathcal{F}(M, \tilde{d}_\gamma)} = 1$ . Take  $f \in S_{\text{Lip}_0(M, \tilde{d}_\gamma)}$  to be such that

$$f(x_i) - f(y_i) = \tilde{d}(x_i, y_i) + \gamma, \quad \forall i = 1, \dots, n.$$

Arguing as in the proof of Theorem 3.16, observe that the sets  $\{x_i\}_{i=1}^n$  and  $\{y_i\}_{i=1}^n$  are disjoint. Now, we consider the finite set  $N := \{x_1, y_1, \dots, x_n, y_n\}$  and assume, without loss of generality, that  $0 \in \{y_1, \dots, y_n\}$ .

Define  $\tilde{f} : N \rightarrow \mathbb{R}$  by  $\tilde{f}(x_i) := f(x_i) - \gamma$  and  $\tilde{f}(y_i) := f(y_i)$  for every  $i = 1, \dots, n$ . We claim that  $\tilde{f}$  is 1-Lipschitz in  $\text{Lip}_0(N, \tilde{d})$ . For this, let  $i, j \in \{1, \dots, n\}$ . We split the argument into two cases. Suppose first that  $\tilde{f}(x_i) - \tilde{f}(y_j) \geq 0$ . Then, we have that

$$|\tilde{f}(x_i) - \tilde{f}(y_j)| = f(x_i) - f(y_j) - \gamma \leq \tilde{d}(x_i, y_j)$$

and we are done. Let us suppose now that  $\tilde{f}(x_i) - \tilde{f}(y_j) < 0$ . Then

$$\begin{aligned} |\tilde{f}(x_i) - \tilde{f}(y_j)| &= \tilde{f}(y_j) - \tilde{f}(x_j) + (\tilde{f}(x_j) - \tilde{f}(x_i)) = -\tilde{d}(y_j, x_j) + f(x_j) - f(x_i) \\ &\leq -\tilde{d}(y_j, x_j) + \tilde{d}(x_j, x_i) + \gamma \stackrel{(*)}{\leq} \tilde{d}(x_i, y_j), \end{aligned}$$

where the last inequality  $(*)$  holds by the definition of  $\tilde{d} = d_\gamma$ .

According to McShane's theorem, we can extend  $\tilde{f}$  to a norm one element in  $\text{Lip}_0(M, \tilde{d})$ . Let us denote this extension once again by  $\tilde{f}$ . By Theorem 3.16, applied to the metric space  $(M, \tilde{d})$  and  $\mu \in \mathcal{F}(M, \tilde{d}_\gamma)$ , we can conclude that  $\mu$  is an SSD point in  $\mathcal{F}(M, \tilde{d}_\gamma)$  (noting that  $\tilde{f}(x_i) - \tilde{f}(y_i) = \tilde{d}(x_i, y_i)$  for every  $i = 1, \dots, n$ ). In other words, given  $\mu \in \mathcal{F}(M, d_{2\gamma})$  with an optimal representation, we proved that  $\mu$  is an SSD point in  $\mathcal{F}(M, d_{2\gamma})$ . This shows that the set of all SSD points in  $\mathcal{F}(M, d_{2\gamma})$  is dense in  $\mathcal{F}(M, d_{2\gamma})$  as we wanted.  $\square$

Note that if  $(M, d)$  is uniformly discrete, then the identity map between  $(M, d)$  and  $(M, d_\gamma)$  turns to be bi-Lipschitz for every  $\gamma > 0$ . In this regard, the following result is a straightforward consequence of Corollary 3.17.

**Corollary 3.18.** *Let  $M$  be a uniformly discrete metric space and  $\varepsilon > 0$  be given. Then there exists a metric space  $N$  and a bi-Lipschitz map  $\phi : M \rightarrow N$  with  $1 \leq \|\phi\| \leq 1 + \varepsilon$  such that the set of SSD points in  $\mathcal{F}(N)$  is dense.*

**Acknowledgements:** This paper was done while the first, the second and the fourth authors were visiting the Department of Mathematics of the Faculty of Electrical Engineering at the Czech Technical University in Prague. They are thankful for the support they have

received there. They would like also to thank Ramón J. Aliaga, Rubén Medina, Rafael Payá and Andrés Quilis for fruitful conversations and for providing relevant references about the topic of the paper.

### Funding information:

*Christian Cobollo* was supported by

- (1) The grants PID2021-122126NB-C33, PID2019-105011GB-I00 and PID2022-139449NB-I00 funded by MICIU/AEI/10.13039/501100011033 and by ERDF/EU,
- (2) Generalitat Valenciana (through Project PROMETEU/2021/070 and the predoctoral contract CIACIF/2021/378).

*Sheldon Dantas* was supported by

- (1) The Spanish AEI Project PID2019 - 106529GB - I00 / AEI / 10.13039/501100011033,
- (2) Generalitat Valenciana project CIGE/2022/97 and
- (3) The grant PID2021-122126NB-C33 funded by MICIU/AEI/10.13039/501100011033 and by ERDF/EU

*Petr Hájek* was supported by

- (1) GA23-04776S and
- (2) SGS22/053/OHK3/1T/13 of CTU in Prague.

*Mingu Jung* was supported by June E Huh Center for Mathematical Challenges (HP086601) at Korea Institute for Advanced Study.

### REFERENCES

- [1] R.J. ALIAGA AND A. RUEDA-ZOCA, *Points of differentiability of the norm in Lipschitz-free spaces*. J. Math. Anal. Appl. 489 (2020), no.2, 124171, 17 pp.
- [2] A. APARICIO, F. OCAÑA, R. PAYÁ, AND A. RODRÍGUEZ, *A non-smooth extension of Fréchet-differentiability of the norm with applications to numerical ranges*. Glasgow Math. J. **28** (1986), pp. 121–137.
- [3] R.J. ALIAGA AND E. PERNECKÁ, *Supports and extreme points in Lipschitz-free spaces*. Rev. Mat. Iberoam. 36 (2020), no. 7, pp. 2073–2089.
- [4] J.B. GUERRERO, G. LÓPEZ PÉREZ AND A. RUEDA-ZOCA, *Octahedrality in Lipschitz-free Banach spaces*, Proc. Roy. Soc. Edinburgh Sect. A. 148, **3** (2018), 447–460
- [5] M.R. BRIDSON AND A. HAEFLIGER, *Metric Spaces of Non-Positive Curvature*, Fundamental Principles of Mathematical Sciences, vol.319, Springer-Verlag, Berlin, 1999.
- [6] B. CASCALES, R. CHICLANA, L.C. GARCÍA-LIROLA, M. MARTÍN AND A. RUEDA ZOCA, *On strongly norm attaining Lipschitz maps*. J. Funct. Anal. **277** (2019), 1677–1717
- [7] M.D. CONTRERAS, *Strong subdifferentiability in spaces of vector-valued continuous functions*. Quart. J. Math. Oxford Ser. **47** (1996), pp. 147–155.
- [8] M.D. CONTRERAS AND R. PAYÁ, *On upper semicontinuity of duality mappings*. Proc. Amer. Math. Soc. **121** (1994), pp. 451–459.  
*C\*-algebras that are I-rings*. J. Math. Anal. Appl. **198** (1996), no. 1, 227–236.
- [9] S. DANTAS, P. HÁJEK, AND T. RUSSO, *Smooth norms in dense subspaces of Banach spaces*, J. Math. Anal. Appl. **487** (2020), 123963.



- [10] S. DANTAS, P. HÁJEK, AND T. RUSSO, *Smooth and polyhedral norms via fundamental biorthogonal systems*, Int. Math. Res. Not. IMRN, (2022); rnac211, [doi.org/10.1093/imrn/rnac211](https://doi.org/10.1093/imrn/rnac211).
- [11] S. DANTAS, M. JUNG, M. MAZZITELLI AND J.T. RODRÍGUEZ, *On the strong subdifferentiability of homogeneous polynomials and (symmetric) tensor products*. Accepted by Publicacions Matemàtiques. Available in <https://arxiv.org/abs/2209.01767>
- [12] S. DANTAS, S.K. KIM, H.J. LEE AND M. MAZZITELLI, *Local Bishop-Phelps-Bollobás properties*. J. Math. Anal. Appl. **468** (2018), pp. 304–323.
- [13] R. Deville, G. Godefroy, and V. Zizler, *Smoothness and Renormings in Banach Spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics, **64**. Longman, Essex, 1993.
- [14] M. FABIAN, P. HABALA, P. HÁJEK, V. MONTESINOS, AND V. ZIZLER, *Banach Space Theory: The Basis for Linear and Nonlinear Analysis*, CMS Books in Mathematics, Springer New York, 2010.
- [15] C. FRANCHETTI, *Lipschitz maps and the geometry of the unit ball in normed spaces*. Arch. Math. **46** (1986), pp. 76–84.
- [16] C. FRANCHETTI AND R. PAYÁ, *Banach spaces with strongly subdifferentiable norm*. Boll. Uni. Mat. Ital. **VII-B** (1993), pp. 45–70.
- [17] L.C. GARCÍA-LIROLA, A. PROCHÁZKA AND A. RUEDA ZOCA, *A characterisation of the Daugavet property in spaces of Lipschitz functions*, J. Math. Anal. Appl. **464**, 1 (2018), 473–492
- [18] J.R. GILES, D.A. GREGORY, AND B. SIMS, *Geometrical implications of upper semicontinuity of the duality mapping of a Banach space*. Pacific J. Math. **79** (1978), pp. 9–109.
- [19] G. GODEFROY, *Some applications of Simons' inequality*. Serdica Math. J. **26** (2000), pp. 59–78.
- [20] G. GODEFROY, *A survey on Lipschitz-free Banach spaces*, Comment. Math. **55** (2) (2015) 89–118.
- [21] G. GODEFROY, V. MONTESINOS, AND V. ZIZLER, *Strong subdifferentiability of norms and geometry of Banach spaces*. Comment. Math. Univ. Carolin. **36** (1995), pp. 493–502.
- [22] D.A. GREGORY, *Upper semi-continuity of subdifferential mappings*. Can. Math. Bull. **23** (1980), pp. 11–19.
- [23] P. HÁJEK AND M. NOVOTNÝ, *Some remarks on the structure of Lipschitz-free spaces*, Bull. Belg. Math. Soc. Simon Stevin **24** (2017), 283–304
- [24] Y. IVAKHNO, V. KADETS AND D. WERNER, *Corrigendum to: The Daugavet property for spaces of Lipschitz functions*, Math. Scand. **104** (2009) 319. Available in the following link: <http://page.mi.fu-berlin.de/werner99/preprints/daugalip-corr.pdf>
- [25] A. PROCHÁZKA AND A. RUEDA ZOCA, *A characterisation of octahedrality in Lipschitz-free spaces*, Ann. Inst. Fourier (Grenoble) **68** (2) (2018) 569–588.
- [26] N. WEAVER, *Lipschitz Algebras*, 2nd ed., World Scientific Publishing Co., River Edge, NJ, 2018.

(Cobollo) UNIVERSITAT POLITÈCNICA DE VALÈNCIA. INSTITUTO UNIVERSITARIO DE MATEMÀTICA PURA Y APLICADA, CAMINO DE VERA, S/N 46022 VALENCIA, SPAIN

ORCID: 0000-0002-5901-5798

*Email address:* `chcogo@upv.es`

(Dantas) DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVERSIDAD DE VALENCIA, DOCTOR MOLINER 50, 46100 BURJASOT (VALENCIA), SPAIN.

ORCID: 0000-0001-8117-3760

*Email address:* `sheldon.dantas@uv.es`

(Hájek) DEPARTMENT OF MATHEMATICS, FACULTY OF ELECTRICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, TECHNICKÁ 2, 166 27 PRAGUE 6, CZECH REPUBLIC

*Email address:* `hajekpe8@fel.cvut.cz`

(Jung) JUNE E HUH CENTER FOR MATHEMATICAL CHALLENGES, KOREA INSTITUTE FOR ADVANCED STUDY, 02455 SEOUL, REPUBLIC OF KOREA

ORCID: 0000-0003-2240-2855

*Email address:* `jmingoo@kias.re.kr`