

Meson modes of probe $D7$ -branes in Klebanov-Strassler background

M. Chemtob*

Université Paris-Saclay, CNRS, CEA, Institut de physique théorique, 91191, Gif-sur-Yvette, France[†]

(Dated: June 4, 2024)

We examine the properties of 4-d bosonic and fermionic (mesons and mesinos) modes on $D7$ -branes wrapped over Kuperstein 4-cycle of the Klebanov-Strassler background. The wave equations are derived in a convenient parameterization of the warped deformed conifold metric. The mass spectra, wave functions and couplings of bosonic modes are evaluated by means of semi-classical tools. We explore the possible existence of fermionic zero modes.

CONTENTS

I. Introduction	1
II. Type II b string theory on warped deformed conifold with Dp -branes	2
A. Flavoured Klebanov-Strassler background	2
B. Dp -brane action	4
III. Bosonic sector of $D7$ -branes	6
A. Classical embedding in the warped deformed conifold	6
B. Meson scalar modes from brane moduli fields	8
C. Meson modes from $D7$ -brane gauge sector	13
D. Properties of meson modes	15
1. Parameter space	15
2. Mass spectra of scalar and vector mesons	18
3. Interactions of scalar mesons	21
IV. Fermionic sector of $D7$ -branes	22
A. Dimensional reduction of spinor fields	23
B. Fermionic zero modes	26
V. Summary and conclusions	27
A. Embedding of $D7$ -branes in deformed conifold	28
1. Geometrical properties of 4-cycle embedding	28
2. Reduced bosonic action	30
3. Reduced Dirac operator	33
4. Undeformed conifold limit	34
References	35

I. INTRODUCTION

Branes and fluxes are necessary ingredients [1] in building solutions of superstring theories dual to confining gauge theories that include flavour degrees of freedom [2–7] and/or chiral fermions [8–12]. The pioneering applications to hadronic physics considered configurations of Dirichlet $D3/D7$, $D4/D6$ -branes [5, 13–15] or $D4/D8$ -branes [16–18] embedded in their own backgrounds. (See the reviews [19–21].) For superbranes embedded in curved spacetimes [22], useful help came from the constraints imposed by κ -symmetry [23, 24] for internal space manifolds of calibration type supporting magnetized or instantonic backgrounds [25–28]. The Klebanov-Strassler solution [29] for type II b supergravity on the deformed conifold offered an ideal setting for this task. The addressed issues included the classification

* marc.chemtob@ipht.fr

[†] Supported by Direction de la Recherche Fondamentale et aux Energies Alternatives Saclay

of holomorphic 4-cycles [30, 31], stability of supersymmetric embeddings [32], mesons spectroscopy [4, 33–36], chiral symmetry breaking [37–39], duality correspondence to Klebanov-Witten gauge theory with quark flavours [40, 41], supersymmetry breaking and its mediation [42], back-reaction on background geometry [43, 44], ultraviolet instabilities [45–47], and connection to the Randall-Sundrum mechanism [48].

In this work we pursue this research thread by extending our previous work on the Kaluza-Klein theory for Klebanov-Strassler solution [49] to the flavoured background including probe $D7$ -branes. Our analysis follows the Giddings-Kachru-Polchinski (GKP) approach [50] to flux compactification on $M_4 \times X_6$ spacetimes, involving a conic Calabi-Yau orientifold X_6 with an attached warped deformed conifold throat \mathcal{C}_6 [51]. We introduce in this background $D7$ -branes that extend over Minkowski spacetime M_4 and wrap over Kuperstein [34] 4-cycle $\Sigma_4 \subset \mathcal{C}_6$. The supergravity solution with 5-flux $N = MK$ and 3-fluxes (M, K) and a N_f $D7$ -brane stack is dual to the flavoured version of Klebanov-Witten [52] supersymmetric field theory. This is the 4-d gauge theory on $(N D3 + M D5)$ -branes near the conifold apex with N_f conjugate pairs of quark flavours Q, \tilde{Q} , which decouples from gravity in the 't Hooft-Veneziano limit, $N_f \ll N \gg 1$. Along with the glueball modes from fluctuations of the supergravity multiplet (closed strings) fields in \mathcal{C}_6 , the theory comprises meson modes from fluctuations of $((D3, D7) + (D7, D3))$ open strings fields in Σ_4 .

The bosonic (scalar and vector) and fermionic (spinor) modes are examined within the streamlined approach [34, 46, 47] using the deformed conifold parameterization as a foliation by Kuperstein 4-cycle leaves [42]. The wave equations are solved by means of the semi-classical JWKB (Jeffreys-Wentzel-Kramers-Brillouin) method. The information on the modes masses and wave functions is used to compute the self-couplings of meson modes along with their couplings to graviton modes. An attempt is also made to test the existence of fermionic zero modes.

The contents of this paper are organized into four sections. The formalism for the flavoured Klebanov-Strassler background is presented in Section II. Subsection II A reviews the supergravity-gauge theory dual descriptions and Subsection II B discusses a simplified version of the $D7$ -brane world volume action which will be used in later sections to compute the mesons properties.

Section III discusses the Kaluza-Klein reduction of the bosonic action for $D7$ -branes wrapped on Kuperstein 4-cycle of the warped deformed conifold. We derive the wave equations for scalar and vector meson modes and present numerical results for the modes wave functions, mass spectra, self-couplings and couplings to bulk graviton modes, all evaluated by means of the JWKB method. Subsection III A discusses algebraic properties of the 4-cycle embedding in the conifold. Subsections III B and III C discuss the reduced action for scalar moduli and gauge vector boson fields on $D7$ -branes. Subsection III D starts with a presentation of the free parameters and next presents results of numerical applications for the mass spectra and interactions of scalar and vector meson modes.

Section IV focuses on the fermionic action of $D7$ -branes in the approach of [6]. Subsection IV A discusses the reduced action and Kaluza-Klein decomposition of spinor fields and Subsection IV B the zero modes wave functions. Section V summarizes the main conclusions. Useful tools and intermediate results are presented in Appendix A. Subsection A 1 discusses geometrical properties of the 4-cycle embedding, Subsection A 2 the $D7$ -brane bosonic action, Subsection A 3 the Dirac operator restriction to Σ_4 and Subsection A 4 collects intermediate results appropriate to the singular conifold limit.

II. TYPE $II b$ STRING THEORY ON WARPED DEFORMED CONIFOLD WITH Dp -BRANES

We consider the GKP [50] flux compactification of type $II b$ superstring theory on the 10-d target spacetime orientifold $M_{10} = M_4 \times X_6$ with the warped deformed conifold \mathcal{C}_6 glued at a conic singularity in the moduli space of X_6 . The classical background in \mathcal{C}_6 is described by Klebanov-Strassler solution [29] for type $II b$ supergravity theory. We introduce in this background probe $D7$ -branes wrapped over an holomorphic 4-cycle of \mathcal{C}_6 preserving $\mathcal{N} = 1$ supersymmetry in M_4 . At low energies and small curvatures relative to the string scale and up to back-reaction effects on the curved background that we neglect, the solution provides a strong coupling description (in the sense of the anti-de Sitter conformal field theory (AdS/CFT) or the gravity-gauge duality) of the flavoured Klebanov-Witten gauge theory [41, 52]. We shall begin the discussion with a brief review of the Klebanov-Strassler background [29]. A simplified version of the bosonic and fermionic action of Dp -branes is presented next in view of later applications to the Kaluza-Klein reduction of the $D7$ -brane action.

A. Flavoured Klebanov-Strassler background

The deformed conifold \mathcal{C}_6 is commonly defined as the locus of the quadratic algebraic equation in C^4 ,

$$Det(W) = - \sum_{a=1}^4 w_a^2 = -z_1 z_2 + z_3 z_4 = -\epsilon^2, [W = \begin{pmatrix} w^3 + iw^4 & w^1 - iw^2 \\ w^1 + iw^2 & -w^3 + iw^4 \end{pmatrix} \equiv \begin{pmatrix} z_3 & z_1 \\ z_2 & z_4 \end{pmatrix}] \quad (\text{II.1})$$

where $\epsilon \in C$ is the complex structure modulus and w_a and z_a , $[a = 1, \dots, 4]$ are linearly related sets of complex coordinates, $(w_1, -iw_2) = \frac{1}{2}(z_1 \pm z_2)$, $(w_3, iw_4) = \frac{1}{2}(z_3 \mp z_4)$. The conifold admits the isometry group $SO(4) \simeq SU(2)_1 \times SU(2)_2$ acting on the coordinates matrix W by left and right multiplication, $W \rightarrow g_1 W g_2^\dagger$. Its intersections with the spherical manifolds,

$$\frac{1}{2}Tr(W^\dagger W) = \sum_a |w_a|^2 = \sum_a |z_a|^2 = |\epsilon|^2 \cosh \tau, \quad (\text{II.2})$$

labelled by the parameter $\tau \in R$, define radial sections of the conifold isomorphic to the compact coset space $T^{1,1} \simeq SU(2)_1 \times SU(2)_2 / U(1)$. The conifold parameterization combining the radial coordinate $\tau \in [0, \infty]$ with the five independent angle coordinates $\Theta^\alpha = (\theta_1, \phi_1, \theta_2, \phi_2, \psi)$ of $T^{1,1}$ [22],

$$\begin{aligned} w_1 &= \epsilon [\cosh S \cos \theta_+ \cos \phi_+ + i \sinh S \cos \theta_- \sin \phi_+], & w_2 &= \epsilon [-\cosh S \cos \theta_+ \sin \phi_+ + i \sinh S \cos \theta_- \cos \phi_+], \\ w_3 &= \epsilon [-\cosh S \sin \theta_+ \cos \phi_- + i \sinh S \sin \theta_- \sin \phi_-], & w_4 &= \epsilon [-\cosh S \sin \theta_+ \sin \phi_- - i \sinh S \sin \theta_- \cos \phi_-], \\ [S &= \frac{\tau + i\psi}{2}, \theta_\pm = \frac{1}{2}(\theta_1 \pm \theta_2), \phi_\pm = \frac{1}{2}(\phi_1 \pm \phi_2), (\theta \in [0, \pi], \phi_i \in [0, 2\pi], \psi \in [0, 4\pi])] \end{aligned} \quad (\text{II.3})$$

is encoded in the representation of the matrix W ,

$$W = g_1 W_0(\tau) g_2^\dagger, [W_0(\tau) = \overline{\text{diag}}(e^{\tau/2}, e^{-\tau/2}), \varphi_1 - \varphi_2 = 0, \psi = \varphi_1 + \varphi_2] \quad (\text{II.4})$$

with the left and right action of $g_i(\theta_i, \phi_i, \varphi_i) \in SU(2)_i$, $[i = 1, 2]$ on the anti-diagonal matrix $W_0(\tau)$. The Klebanov-Strassler solution of type *II b* supergravity [29] on $M_4 \times C_6$ involves 3-fluxes, $\hat{l}_s^2 \int_{B,A} (H_3, F_3) = (-K, M)$, $[N = MK, \hat{l}_s = 2\pi\sqrt{\alpha'}]$ across the pair of Poincaré dual 3-cycles B, A of C_6 , with the resulting 5-flux $\hat{l}_s^4 \int_{T^{1,1}} F_5 = N$ sourcing a warped spacetime of classical metric,

$$\begin{aligned} ds_{10}^2 &= h^{-\frac{1}{2}}(\tau) d\bar{s}_4^2 + h^{\frac{1}{2}}(\tau) d\bar{s}^2(C_6), [h(\tau) \equiv e^{-4A(\tau)} = 2^{2/3} (g_s M \alpha')^2 \epsilon^{-8/3} I(\tau), \\ d\bar{s}_6^2(C_6) &= \frac{\epsilon^{4/3} K(\tau)}{2} \left(\frac{1}{3K^3(\tau)} (d\tau^2 + (g^{(5)})^2) + \cosh^2\left(\frac{\tau}{2}\right) ((g^{(3)})^2 + (g^{(4)})^2) + \sinh^2\left(\frac{\tau}{2}\right) ((g^{(1)})^2 + (g^{(2)})^2) \right), \\ K(\tau) &= \frac{(\sinh(2\tau) - 2\tau)^{1/3}}{2^{\frac{1}{3}} \sinh \tau}, I(\tau) = \int_\tau^\infty dx \frac{x \coth x - 1}{\sinh^2 x} (\sinh(2x) - 2x)^{\frac{1}{3}} \end{aligned} \quad (\text{II.5})$$

asymptotic to the direct product metric of the spacetime $AdS_5 \times T^{1,1}$, describing the background for $ND3+MD5$ -brane stacks embedded near the conifold apex. We follow same notations as [49, 53] where $h(\tau)$ is the warp profile, $K(\tau), I(\tau)$ are auxiliary radial functions, $g^{(a)}(\Theta)$ is the diagonal basis of left-invariant 1-forms on $T^{1,1}$ and $g_s, \alpha' = 1/m_s^2$ denote the string theory coupling constant and scale parameters. The classical solutions for the 2- and 3-form fields are quoted in Eq. (A.18) below.

In the limit $N \gg 1$, fixed $g_s N$, the dual gauge theory on the regular $ND3$ - and fractional $MD5$ -branes is the Klebanov-Witten 4-d $\mathcal{N} = 1$ supersymmetric theory with local symmetry group $\mathcal{G} = \mathcal{G}_1 \times \mathcal{G}_2 = SU(N+M) \times SU(N)$ [52] of coupling constants $g_{1,2}$ and global (flavour) symmetry group $G = SU(2) \times SU(2) \times U(1)_b \times Z_2^r, [Z_2^r \subset U(1)_r]$. The small curvature regime $g_s N \gg 1$, where classical supergravity theory is valid, corresponds to 't Hooft limit $\lambda = g_{1,2}^2 N \gg 1$ of the gauge theory. The gauge sector consists of vector supermultiplets $V_{1,2}$ in the adjoint representations of $\mathcal{G}_{1,2}$ and the (matter) flavour sector consists of two pairs of chiral supermultiplets in the bi-fundamental representations, $A_i \sim (N+M, \bar{N}), B_i \sim (\bar{N}+M, N)$, $[i = 1, 2]$ of \mathcal{G} and the representations $(2, 1)_{1, \frac{1}{2}}, (1, 2)_{-1, \frac{1}{2}}$ of G . The renormalization group evolution features a cascading type flow towards the infrared through self-similar Seiberg type dualities $SU(N+M) \times SU(N) \rightarrow SU(N-M) \times SU(N)$. For $N = kM$, $[k \in Z]$ the successive duality steps $\delta k = -1$ end into the confining pure gauge theory $SU(M)$ [54]. In the presence of the interaction superpotential $W_{KW} = \frac{h}{2} \epsilon^{ik} \epsilon^{jl} Tr(A_i B_j A_k B_l)$, the renormalization group flow (for $N \gg M$) drives the theory to a superconformal (infrared) fixed submanifold $h = h(g_1, g_2)$ in the space of coupling parameters g_1, g_2, h [55]. The fields and superpotential are assigned in the limit $M \rightarrow 0$ the scaling dimensions $\Delta(\phi) = d(\phi) + \gamma(\phi)/2$ and (unbroken) R-symmetry charges R (including quantum contributions),

$$R(A_i) = 2\Delta(A_i)/3 = 1/2, R(B_i) = 2\Delta(B_i)/3 = 1/2, \Delta(\lambda) = 3/2, R(\lambda) = 1, R(W) = 2\Delta(W)/3 = 2. \quad (\text{II.6})$$

The probe N_f $D7$ -branes wrapped over a 4-cycle $\Sigma_4 \subset C_6$, add modes of open $F1$ -strings stretched between the $N_f D7$ - and $(ND3+MD5)$ -branes. Massless quark modes are present if Σ_4 touches the conifold apex where the $D3$ -branes are located. The gauge dual theory is then the flavoured Klebanov-Witten gauge theory with N_f conjugate pairs of quarks and anti-quarks Q, \tilde{Q} [3] in bi-fundamental representations of the colour and flavour groups. One

must consider holomorphic and topologically trivial Σ_4 in order to preserve $\mathcal{N} = 1$ 4-d supersymmetry along with $D3$ -charge conservation and C_4 -tadpole cancellation. We shall specialize to the Kuperstein 4-cycle [34], $w_4 = \mu$, sub-locus of the conifold $\sum_{a=1}^4 w_a^2 = \epsilon^2$, where the complex parameter μ (of same energy dimension $E^{-3/2}$ as ϵ) sets the minimal classical radial distance between the apexes of \mathcal{C}_6 and Σ_4 . Note that our ϵ^2 , μ correspond to the ϵ , $\mu/\sqrt{2}$ of [41].

In full generality, the $D7$ -branes stack can be split into left and right sub-stacks, $N_f = N_{f_L} + N_{f_R}$. The gauge theory is then described by an horizontal type quiver diagram with Q_L , \tilde{Q}_L in bi-fundamentals of $SU(N_{f_L}) \times SU(N + M)$ and Q_R , \tilde{Q}_R in bi-fundamentals of $SU(N) \times SU(N_{f_R})$. (For comparison, the Ouyang embedding [33], $z_4 = \mu$, involves instead the vertical type quiver diagram with Q_L , \tilde{Q}_R charged under $SU(N + M)$ and Q_R , \tilde{Q}_L charged under $SU(N)$.) The dual theory of gauge symmetry group $\mathcal{G} = SU(N_{f_L}) \times SU(N + M) \times SU(N) \times SU(N_{f_R})$ includes, in addition to A_i , B_i , a quark flavour sector with left and right (conjugate pairs) of quarks supermultiplets in the \mathcal{G} group representations

$$Q_L \sim (N_{f_L}, \overline{N + M}, 1, 1), \tilde{Q}_L \sim (\bar{N}_{f_L}, N + M, 1, 1), Q_R \sim (1, 1, \bar{N}, N_{f_R}), \tilde{Q}_R \sim (1, 1, N, \bar{N}_{f_R}). \quad (\text{II.7})$$

The total quartic order superpotential with (implicit) traces over the gauge and quark flavour quantum numbers,

$$\begin{aligned} W = & h(A_1 B_1 A_2 B_2 - A_1 B_2 A_2 B_1) - \eta_L \sqrt{h} \tilde{Q}_L (A^i B_i - \frac{\mu}{\sqrt{h}}) Q_L + \frac{\eta_L^2}{2} (\tilde{Q}_L Q_L)^2 \\ & - \eta_R \sqrt{h} \tilde{Q}_R (B^i A_i - \frac{\mu}{\sqrt{h}}) Q_R - \frac{\eta_R^2}{2} (\tilde{Q}_R Q_R)^2, \end{aligned} \quad (\text{II.8})$$

drives the theory (in the quarks massless limit $\mu \rightarrow 0$) to a superconformal fixed submanifold on which $\Delta(Q_{L,R}) = 3/4$, $R(Q_{L,R}) = 1/2$. One can always rescale the superfields $Q_{L,R}$ so as to set $\eta_{L,R} = \sqrt{h}$ in the superpotential W which then depends on the overall coupling constant h and the quarks bare mass parameter $m_Q \simeq \mu/\sqrt{h}$. The correspondence between the dimensional parameters on the string theory side ϵ , μ and the dynamical and bare quark mass scales Λ_{ir} , m_Q on the gauge theory side, $\epsilon^{2/3} \simeq \alpha' \Lambda_{ir}$ and $\mu^{2/3} \simeq \alpha' m_Q$, is set via dimensional analysis.

The structure of the superpotential in Eq. (II.8) is formally similar to that of the $SU(N_c)$, $\mathcal{N} = 2$ supersymmetric gauge theory [56] with a massive adjoint superfield ϕ coupled to massive $N_f(Q, \tilde{Q})$ fundamental quarks hypermultiplets. The decoupling of ϕ along the Higgs branch, labelled by the scalar field component VEV of ϕ , deforms the $\mathcal{N} = 2$ theory to the $\mathcal{N} = 1$ theory and changes the superpotential as

$$W_{\mathcal{N}=2} = \frac{s}{2} \text{Tr}(\phi^2) + \text{Tr}(\phi \tilde{Q} Q) + \text{Tr}(m Q \tilde{Q}) \implies W_{\mathcal{N}=1} = -\frac{1}{2s} \text{Tr}((\tilde{Q} Q)^2) + \text{Tr}(m Q \tilde{Q}). \quad (\text{II.9})$$

An even closer analog [41] to the flavoured gauge theory is provided by the type II b supergravity on the (flat orbifold) spacetime $M_4 \times (C_{1,2}^2/Z_2) \times C_3$ (coordinates $(x^\mu, w_{1,2}, w_3)$) with N $D3$ -branes at the fixed point $w_{1,2,3} = 0$ and N_f $D7$ -branes over $M_4 \times (C_{1,2}^2/Z_2)$ at $w_3 = m$. The 4-d supersymmetric gauge theory $SU(N) \times SU(N_f)$, $\mathcal{N} = 2$ consists (in $\mathcal{N} = 1$ language) of one vector and three complex chiral supermultiplets $\Phi_{i=1,2,3}$, adjoints of $SU(N)$ (descending from the supermultiplet of the $\mathcal{N} = 4$ theory on N $D3$ -branes) and N_f quark flavour hypermultiplets $Q = \text{diag}(Q_L, Q_R)$, $\tilde{Q} = \text{diag}(\tilde{Q}_L, \tilde{Q}_R)$. The total superpotential, including mass terms for Φ_3 and quarks,

$$W = -\text{Trace}[[\Phi_1, \Phi_2] \Phi_3 + \frac{s}{2} \Phi_3^2 + \tilde{Q} \Phi_3 Q - \mu(\tilde{Q}_L Q_L + \tilde{Q}_R Q_R)], \quad (\text{II.10})$$

yields, after the Higgs field $\Phi_3 = \text{diag}(\phi_3, -\tilde{\phi}_3)$ is integrated out, the $\mathcal{N} = 1$ flavoured Klebanov-Witten gauge theory superpotential in Eq. (II.8), with the anti-diagonal matrix $\Phi_{1,2} = \overline{\text{diag}}(A_{1,2}, \pm B^{2,1})$ and $m \simeq (B^i A_i - \mu)$.

In order that the moduli space of vacua of the supergravity theory matches that of the flavoured gauge theory, one may need to activate classical fields in the $D7$ -branes gauge theory [40, 41, 57]. The case where no classical gauge fields are present corresponds to having only Q_R -type quarks coupled to the lower rank gauge theory factor $SU(N)$. We shall specialize here to this case which then involves the single species of quarks $Q \sim (1, \bar{N}, N_f)$, $\tilde{Q}(1, N, \bar{N}_f)$ of $SU(N + M) \times SU(N) \otimes SU(N_f)$ with the total interaction superpotential $W = W_{KW} - h[\tilde{Q}(B^i A_i - \frac{\mu}{\sqrt{h}})Q + \frac{1}{2}(\tilde{Q}Q)^2]$.

B. Dp -brane action

The dimensional reduction of the $D7$ -brane action on Kuperstein 4-cycle gives rise to an effective field theory in M_4 with meson modes added to the glueball modes from the supergravity multiplet. The geometrical tools needed in

applying the Kaluza-Klein approach for submanifolds embedded in curved manifolds, are well documented [58, 59]. To prepare the ground for this discussion, we introduce here general notations that will be used throughout the text. The target spacetime $M_{10} = M_4 \times X_6$ is parameterized by world coordinates X^M with a metric tensor g_{MN} describing the (diffeomorphism invariant) proper distances squared,

$$ds^2(M_{10}) = g_{MN}dX^M dX^N, [X^M = (X^\mu, X^m), M = 0, \dots, 9, \mu = 0, \dots, 3, m = 1, \dots, 6]. \quad (\text{II.11})$$

The coordinates X^A for the (flat) tangent spaces of (constant) metric tensor $\eta_{AB} = \text{diag}(-1, +1, \dots, +1)$ are related to the X^M by the transformations $X^A = e_M^A X^M$, $X^M = e_A^M X^A$ involving the components e_M^A , e_A^M of the dual bases of (local inertial vielbein) frame vectors (e^A , e_A) which satisfy the properties

$$ds^2(M_{10}) = \sum_{A=0}^9 (e^A)^2, [e^A = e_M^A dX^M, e_A = e_A^M \partial_M, \eta_{AB} e_M^A e_M^B = g_{MN}]. \quad (\text{II.12})$$

Similar definitions are adopted for the world and flat bases of Dirac gamma matrices obeying the Clifford-Dirac algebra,

$$\{\Gamma_A, \Gamma_B\} = 2\eta_{AB}, \{\Gamma_M, \Gamma_N\} = 2g_{MN}, [\Gamma^A = e_M^A \Gamma^M, \Gamma^M = e_A^M \Gamma^A]. \quad (\text{II.13})$$

The world volume M_{p+1} of parallel Dp -branes in M_{10} is parameterized by the intrinsic coordinates $\xi^\alpha = (\xi^\mu, \xi^r)$, $[\alpha = 0, \dots, p, \mu = 0, \dots, 3, r = 1, \dots, p-3]$. Its embedding along the directions of M_4 and the $(p-3)$ -cycle $\Sigma_{p-3} \subset C_6$ splits up the coordinates into tangential and normal subsets, $X^M = (X^\alpha, X^u) = (X^\mu, X^r, X^u)$, $[\alpha = 0, \dots, p, \mu = 0, \dots, 3, r = 1, \dots, p-3, u = 1, \dots, 9-p]$ regarded as fields $X^M(\xi^\alpha)$ on the world volume M_{p+1} . On the N_f Dp -branes stack, the coordinates fields are promoted to $N_f \times N_f$ matrices in the adjoint representation of $U(N_f)$ with the X^α/α' identified to gauge fields A^α and the X^u to moduli fields. We shall restrict consideration to diagonal matrices in the Abelian subgroup $U(1)^{N_f}$ corresponding to N_f copies of single Dp -branes.

The $D7$ -brane action includes the familiar Dirac-Born-Infeld and Chern-Simons bosonic terms, built from the embedding coordinates X^M , the brane pull-backs of bulk fields B_2 , C_r and the brane 1-form gauge field $A_1 = A_\alpha d\xi^\alpha$. The fermionic action is constructed as an expansion in powers of the pull-back of the spinor doublet field Θ of 10-d type II b supergravity [60]. We restrict hereafter to the quadratic order action in Θ coupled to the NSNS 2-form and RR 4-form background fields and the brane gauge field strength $F_2 = dA_1$. The combined bosonic and fermionic action for $Dp/\bar{D}p$ -branes in Einstein frame

$$\begin{aligned} S(Dp/\bar{D}p) &= -\tau_p \int_{M_{p+1}} d^{p+1} \xi e^{(p-3)\phi/4} (-\text{Det}(\gamma + e^{-\phi/2} \mathcal{F}))^{1/2} \\ &\times [1 - i \bar{\Theta} P_{\mp}^{Dp} ((\mathcal{M}^{-1})^{\alpha\beta} \Gamma_\beta D_\alpha + \frac{i}{8} \gamma^{\alpha\beta} \Gamma_\beta \mathcal{H}_5 \Gamma_\alpha \otimes \sigma_2) \Theta] \pm \mu_p \int_{M_{p+1}} [\sum_q C_q \wedge e^{\mathcal{F}_2}]_{p+1}, \\ [\tau_p = \mu_p = \frac{2\pi}{\hat{l}_s^{p+1}}, \mathcal{F}_2 = B_2 + F_2, F_2 = dA(\xi), F_{q+1} = dC_q, \Theta(\xi) = \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}, \bar{\Theta} = \Theta^\dagger \Gamma^0, \\ \mathcal{M}_{\alpha\beta} = \gamma_{\alpha\beta} + e^{-\phi/2} \mathcal{F}_{\alpha\beta} \Gamma_{(10)} \otimes \sigma_3, P_{\pm}^{Dp}(\mathcal{F}) = \frac{1}{2}(1 \pm \Gamma_{Dp}(\mathcal{F}))] \end{aligned} \quad (\text{II.14})$$

is expressed here in same notations as [6, 49] with $\tau_p = 2\pi/\hat{l}_s^{p+1}$ and $g_{Dp}^2 = 4\pi g_s \hat{l}_s^{p-3}$ denoting the brane tension and gauge theory coupling constant. The action is built from the covariant derivatives along M_8 : $D_\alpha = \nabla_\alpha + A_\alpha$, $[\nabla_\alpha = \partial_\alpha + \omega_\alpha]$ and the pull-back transforms of the bulk spacetime metric and Dirac matrix fields g_{MN} , Γ_M , the axio-dilaton scalar fields (ϕ , C_0) and the q -form fields B_2 , $C_{2,4}$, $[F_5 = dC_4]$

$$\begin{pmatrix} \gamma_{\alpha\beta} \\ B_{\alpha\beta} \end{pmatrix} = \partial_\alpha X^M(\xi) \partial_\beta X^N(\xi) \begin{pmatrix} g_{MN}(X(\xi)) \\ B_{MN}(X(\xi)) \end{pmatrix}, \Gamma_\alpha = \partial_\alpha X^M \Gamma_M = E_A^M \partial_\alpha X^A \Gamma_A. \quad (\text{II.15})$$

The self-dual field strength 5-form has the classical value $F_5^{cl} = \partial_\tau h^{-1}(\tau) d\tau \wedge \text{vol}(M_4)$. The (anticommuting, Grassmannian) bulk spinor field, $\Theta(X(\xi)) = (\theta_1(X(\xi)), \theta_2(X(\xi)))$, of Dirac conjugate, $\bar{\Theta} = \Theta^\dagger \Gamma^0 = \Theta^T \Gamma^0$, consists of a pair of Majorana-Weyl spinor fields $\theta_i(X) \in 16$ of $SO(9,1)$ doublet in the Pauli matrices $\sigma_{1,2,3}$ space. The Weyl spinor fields of fixed 10-d chirality, $\Gamma_{(10)} \theta_i = \theta_i$, $[\Gamma_{(10)} = \Gamma^{01\dots 9}]$ satisfy the Majorana reality condition,

$$B\theta_i = \theta_i^*, [B\Gamma_M B^* = \Gamma_M^*, B^* B = 1] \implies \bar{\theta}_i = \theta_i^T C, [\Gamma_A^T = -C\Gamma_A C^{-1}, C = C^\dagger = C^{-1}] \quad (\text{II.16})$$

where the matrices B and C , which transform the Dirac matrices to their complex conjugate and transpose versions, respectively, are related (up to convention dependent choices of relative signs) as, $C = B^T \Gamma_0 = -B^T \Gamma^0$, $B^* = \Gamma_0 C =$

$\Gamma_0 C^{-1}$. A comprehensive presentation of tools for supersymmetry and Dirac matrices in various dimensions can be found in [61].

Finally, we hope that no confusion should arise between spinor indices of gamma matrices $(\Gamma_M)_{\alpha\beta}$ and the induced brane metric components $\gamma_{\alpha\beta}$ in the effective metric $\mathcal{M}_{\alpha\beta}$ and likewise in our use of same symbols for 1-forms and differentials in the cotangent and tangent spaces. Note also that the transition from our Einstein frame to the alternative Einstein frame in [42] is realized by replacing $g_{MN} \rightarrow g_{MN}/g_s^{1/2}$, $\sqrt{\gamma} \rightarrow \sqrt{\gamma}/g_s^2$, $\mathcal{F} \rightarrow g_s^{1/2}\mathcal{F}$, $\tau_p \rightarrow \tau_p/g_s^{(p+1)/4}$.

III. BOSONIC SECTOR OF D7-BRANES

We examine in this section the bosonic modes on D7-branes wrapped over Kuperstein 4-cycle [34] of the deformed conifold. After discussing properties of the classical embedding, we derive the wave equations for wave functions of mode fields in M_4 arising from fluctuations of the scalar moduli and gauge vector brane fields decomposed on harmonics of radial sections of the 4-cycle compact base of geometry S^3/Z_2 . We apply the semi-classical JWKB method to evaluate the mass spectra and wave functions of meson modes. This information is used to compute their self couplings and boundary couplings to graviton modes. (The fermionic action will be examined in Section IV.)

A. Classical embedding in the warped deformed conifold

The Kuperstein 4-cycle [34] $\Sigma_4 : w_4 = \mu$, is invariant under the diagonal group $SO(3)_D \subset (SU(2)_L \times SO(4))/Z_2$ of the conifold isometry group which comprises rotations of the complex coordinates $w_{1,2,3}$. We see from Eq. (II.3) that this cycle intersects the fixed- τ subloci of the conifold, $\theta_- = 0$, $\phi_- = 0 \implies \theta_1 = \theta_2 = \theta$, $\phi_1 = \phi_2 = \phi$, generated by the restricted form of the coordinates matrix $W = g_1 W_0(\tau) g_2^\dagger$, [$g_1 = g_2$] [49]. The initial study [34] invoked this observation to parameterize the deformed conifold by combining the coordinates τ , θ , ϕ with the angles γ , δ of the $SU(2)$ matrix, $S(\gamma, \delta) = g_1 W_0(\tau) g_1^\dagger$. Instead, we shall consider the alternative parameterization of the deformed conifold as a foliation by Σ_4 leaves [42],

$$\begin{aligned} w_1 &= i\eta(\chi)(\hat{C}_\rho c_\phi c_\theta + i\hat{S}_\rho s_\phi), \quad w_2 = i\eta(\chi)(-\hat{C}_\rho s_\phi c_\theta + i\hat{S}_\rho c_\phi), \quad w_3 = -i\eta(\chi)\hat{C}_\rho s_\theta, \quad w_4 = (\mu + \chi), \\ [c_\phi &= \cos \phi, \quad c_\theta = \cos \theta, \quad s_\phi = \sin \phi, \quad s_\theta = \sin \theta, \quad \hat{C}_\rho = \cosh\left(\frac{\rho + i\gamma}{2}\right), \quad \hat{S}_\rho = \sinh\left(\frac{\rho + i\gamma}{2}\right), \\ \chi \in C, \quad \rho \in [0, \infty], \quad \alpha^k &= (\theta \in [0, \pi], \quad \phi \in [0, 2\pi], \quad \gamma \in [0, 4\pi]). \end{aligned} \quad (\text{III.1})$$

The D7-brane splits up the spacetime $M_4 \times \mathcal{C}_6$ coordinates $X^M = (x^\mu, y^m)$ into 8 longitudinal real coordinates $X^\alpha = (X^\mu, X^a = (\rho, \alpha^k))$, [$\alpha = 0, \dots, 7$, $a = 1, \dots, 4$, $k = 1, 2, 3$], and the single transversal complex coordinate, χ . The bulk bosonic and fermionic coordinates, X^M and Θ^α , are regarded as brane fields on the world volume M_8 of intrinsic coordinates ξ^α . The invariance of the theory under coordinates diffeomorphisms can be used to choose the so-called static gauge, $X^\alpha(\xi) = \xi^\alpha$, with the complex modulus field $\chi(\xi^\alpha) = \chi(\rho, \alpha^m)$ describing the brane location along the transversal directions. Upon substituting Eq. (III.1) into Eqs. (II.1) and (II.2) for the conifold and its radial fixed- τ sections, one obtains the pair of matching equations linking the conifold and 4-cycle coordinates,

$$\sum_a w_a^2 = \epsilon^2 = (\mu + \chi)^2 - \eta^2(\chi), \quad \sum_a |w_a|^2 = |\epsilon|^2 \cosh \tau = \hat{r}^3 = |\eta(\chi)|^2 \cosh \rho + |\mu + \chi|^2, \quad [\hat{r}^3 = \left(\frac{2}{3}\right)^{3/2} r^3] \quad (\text{III.2})$$

where the first equation defines the complex function $\eta(\chi) = \pm(\mu^2 - \epsilon^2 + 2\mu\chi + \chi^2)^{1/2}$ and the second defines the 1-to-1 correspondence between the radial variables ρ and τ at fixed χ , with $|\mu|$ measuring the distance between the apexes of \mathcal{C}_6 and Σ_4 . To lighten the formalism, we introduce in the sequel the dimensionless complex parameter $L = \mu/\epsilon$ along with the auxiliary complex functions $Y(\chi) = 1/X(\chi) = \eta_\chi/\mu_\chi$, [$\eta_\chi \equiv \eta(\chi)$, $\mu_\chi \equiv \mu + \chi$] which satisfy the relations

$$\eta_\chi = \frac{\mu_\chi}{X(\chi)} = \mu_\chi Y(\chi) \implies d\eta_\chi = X d\chi = \frac{d\chi}{Y}, \quad [X(\chi) \equiv \frac{1}{Y(\chi)} = \frac{1}{(1 - \epsilon^2/\mu_\chi^2)^{1/2}} = \left(1 - \frac{1}{(L + \chi/\epsilon)^2}\right)^{-1/2}] \quad (\text{III.3})$$

For real values of the parameters μ , ϵ and the variable χ , one finds that $\eta_\chi \simeq \chi/X(\chi) \rightarrow \chi(1 - \epsilon^2/\chi^2)^{1/2}$ at $\chi \gg \mu$ and $\eta_\chi \rightarrow \mu(1 - \epsilon^2/\mu^2)^{1/2}$ at $\chi \ll \mu$. In the parameters interval, $\mu < \epsilon$, these functions develop imaginary parts with $\eta_\chi \rightarrow \pm i|\epsilon/\mu|$ for values of the coordinate variable in the ranges $\epsilon > \chi \gg \mu$ and $\chi \ll \mu < \epsilon$. The matching

equations in Eq. (III.2) can be combined into the χ -dependent functional relation, linking the radial variables τ and ρ ,

$$\cosh \tau = \left| \frac{\mu_X}{\epsilon} \right|^2 (|Y|^2 \cosh \rho + 1) = \left| \frac{\mu_X}{\epsilon^2} - 1 \right| \cosh \rho + \left| \frac{\mu_X^2}{\epsilon^2} \right|. \quad (\text{III.4})$$

Since the classical solution for the brane modulus field is expected to coincide with the 4-cycle locus, $\chi(\rho, \alpha^m) = 0$, as we show below, then at the on-shell value $\chi = 0$ the change of radial variable from $\rho \rightarrow \tau$ and vice versa can be written as,

$$\cosh \tau = |L|^2 (|1 - L^{-2}| \cosh \rho + 1), \quad \cosh \rho = |1 - L^{-2}|^{-1} \left(\frac{\cosh \tau}{|L|^2} - 1 \right), \quad \left[\frac{d\tau}{d\rho} = |L^2 - 1| \frac{\sinh \rho}{\sinh \tau} \right] \quad (\text{III.5})$$

where $L^2 = \mu^2/\epsilon^2$ and $Y \equiv 1/X = (1 - L^{-2})^{1/2}$ are now regarded as complex parameters. Note that $\lim_{\rho \rightarrow 0} d^2\tau/d\rho^2 \simeq \cosh \rho / \sinh \rho$ diverges as $1/\rho$. Along the real L^2 -axis, X is pure imaginary inside the segment $0 < L^2 < 1$ (with $X \simeq \pm iL$ for small L^2), and real elsewhere. At large $|L^2|$, the limits $X \rightarrow 1$, $\cosh \rho \rightarrow (\cosh \tau / |L|^2 - 1)$ show that the radial region $\tau < \tau_0$, $[\cosh \tau_0 = |L|^2]$ is forbidden. The minimal radial distance τ_{min} from the conifold apex (at $\tau = 0$) to the 4-cycle apex (at $\rho \rightarrow \rho_{min} = 0$),

$$\cosh \tau_{min} = \left| \frac{\mu}{\epsilon} \right|^2 \left(\frac{1}{|X|^2} + 1 \right) = |L|^2 (1 + |1 - \frac{1}{L^2}|), \quad \left[\frac{d\tau}{d\rho} \Big|_{\tau_{min}} \simeq \frac{|L^2 - 1| \rho}{(4|L^2 - 1| - 1)^{1/2}} \right] \quad (\text{III.6})$$

measures the extent to which Σ_4 protrudes inside \mathcal{C}_6 . Inside the three contiguous intervals along the real axis of the L^2 -plane this can be expressed by the formulas

$$\tau_{min} = [\cosh^{-1}(-2L^2 + 1), 0, \cosh^{-1}(2L^2 - 1)], \quad [L^2 < 0, 0 < L^2 < 1, L^2 > 1] \quad (\text{III.7})$$

which can be rewritten in a more familiar way using the elementary relationship, $\cosh^{-1} A = \ln(A + \sqrt{A^2 - 1})$. For $L^2 > 1$: $\cosh \tau_{min} = 2L^2 - 1$, $\sinh \tau_{min} = 2L(L^2 - 1)^{1/2}$. Inside the real axis segment $0 < L^2 \leq 1$, we see that $\tau_{min} = 0$ saturates the lower bound, $\tau \geq 0$, and that its value elsewhere, $\tau_{min} = \ln(2|L|^2 - 1 + 2|L|(|L|^2 \mp 1)^{1/2})$, grows as $\tau_{min} \simeq 2 \ln(|L|)$ at large $|L^2|$. Note that $\tau_{min}(L)$ is a regular function of the variable real L except for a discontinuous derivative, $d\tau_{min}/dL \simeq 2\sqrt{2}L/(L^2 - 1)^{1/2}$, near $L = 1$, forecasting a possible cusp behaviour there.

The undeformed conifold limit $\epsilon \rightarrow 0$, $\tau \rightarrow \infty$ must be taken at fixed r and fixed μ . In the change of variable $\tau \rightarrow r$, the residual dependence on ϵ gets absorbed inside the product $\epsilon L \rightarrow \mu$. The minimal distance to the conifold apex sets at the conic radial variable value,

$$\hat{r}_{min}^3 \equiv \left(\frac{2}{3}\right)^{3/2} r_{min}^3 = |\epsilon|^2 \cosh \tau_{min} = (|\mu^2 - \epsilon^2| + |\mu|^2) \simeq 2|\mu|^2. \quad (\text{III.8})$$

With the definition of the conic radial variable r in Eq. (III.2), the radial warping profile function reads $h(\tau) \rightarrow h(r) \simeq (\mathcal{R}/r)^4$. Our rescaled conic variable \hat{r} coincides with the r of [42] while our $h(r) = (4/9)h(\hat{r})$ is (4/9) times their $h(r)$.

Having dealt with the algebraic properties of the 4-cycle embedding, we now focus on its geometrical properties. The unwarped metric is evaluated by substituting the parameterization in Eq. (III.1) into the defining formula [51],

$$d\tilde{s}^2(\mathcal{C}_6) = \frac{1}{2} \frac{\partial \mathcal{F}}{\partial \hat{r}^3} Tr(dW^\dagger dW) + \frac{1}{4} \frac{\partial^2 \mathcal{F}}{\partial (\hat{r}^3)^2} |Tr(W^\dagger dW)|^2 = (2F_1 \delta_{ab} + 4F_2 \bar{w}_a w_b) dw^a \bar{d}w^b, \quad (\text{III.9})$$

where the coefficient functions $F_1(\rho)$, $F_2(\rho)$ correspond to the single and double trace terms for the differential dW of the coordinates matrix W in Eq. (II.1). To take advantage of the isometry group $SO(3) = SU(2)/Z_2 \sim S^3/Z_2$ of the 4-cycle fixed- ρ sections, it is useful to trade the angular 1-forms $d\alpha^m = (d\theta, d\phi, d\gamma)$ for the left invariant 1-forms \hat{h}_k , $[k = 1, 2, 3]$ of $SO(3)$, satisfying Maurer-Cartan relations, $d\hat{h}_i = \frac{1}{4} \epsilon_{ijk} \hat{h}_j \wedge \hat{h}_k$. The resulting basis of left invariant 1-forms [42] is found to be given by the (angle dependent) linear combinations of angular differential forms $d\alpha_m$,

$$\begin{aligned} \hat{h}_1 &= -2(\cos \frac{\gamma}{2} d\theta + \sin \frac{\gamma}{2} \sin \theta d\phi), \quad \hat{h}_2 = 2(-\sin \frac{\gamma}{2} d\theta + \cos \frac{\gamma}{2} \sin \theta d\phi), \quad \hat{h}_3 = 2(\frac{d\gamma}{2} + \cos \theta d\phi), \\ [d\theta &= -\frac{1}{2}(\cos \frac{\gamma}{2} \hat{h}_1 + \sin \frac{\gamma}{2} \hat{h}_2), \quad d\phi = \frac{1}{2 \sin \theta}(-\sin \frac{\gamma}{2} \hat{h}_1 + \cos \frac{\gamma}{2} \hat{h}_2), \quad d\gamma = \hat{h}_3 - \cot \theta (-\sin \frac{\gamma}{2} \hat{h}_1 + \cos \frac{\gamma}{2} \hat{h}_2)] \end{aligned} \quad (\text{III.10})$$

Note that our sign conventions for the w_a in Eq. (III.1) and for the above \hat{h}_i differ from those of [42] by the replacements $\theta \rightarrow -\theta$ and $\phi \rightarrow -\phi$ and $w_3 \rightarrow -w_3$. One can now use the above relations to express the conifold metric as a diagonal quadratic form in \hat{h}_i (up to cross terms $d\chi(d\rho - i\hat{h}_3)$) weighted by ρ -dependent functions,

$$d\tilde{s}^2(\mathcal{C}_6) = F_1 \left[2(d\bar{\chi} d\chi + d\bar{\eta} d\eta \cosh \rho) + \frac{1}{4} \bar{\eta}_\chi \eta_\chi (\hat{h}_1^2 - \hat{h}_2^2 + (2d\rho^2 + \hat{h}_1^2 + \hat{h}_2^2 + 2\hat{h}_3^2) \cosh \rho) \right]$$

$$\begin{aligned}
& +d\eta(d\rho - i\hat{h}_3)\bar{\eta}\sinh\rho + d\bar{\eta}(d\rho + i\hat{h}_3)\eta\sinh\rho + R\left[2\mu_\chi d\bar{\chi} + (2\cosh\rho d\bar{\eta} + \bar{\eta}_\chi\sinh\rho)(d\rho - i\hat{h}_3)\right]^2, \\
[F_1 = \frac{1}{2}\frac{\partial\mathcal{F}}{\partial\hat{\rho}^2} = \frac{\epsilon^{-2/3}K(\tau)}{2}, R \equiv \frac{F_2}{F_1} = -\frac{1}{2\epsilon^2\sinh^2\tau}(\cosh\tau - \frac{2}{3K^3(\tau)})]. \tag{III.11}
\end{aligned}$$

The factor of $R = F_2/F_1$ in the last term inside brackets has the expected perfect square form. We rewrite for convenience the expressions for the unwarped metric tensor components,

$$\begin{aligned}
\tilde{g}_{\rho\rho} = \tilde{g}_{\hat{h}_3\hat{h}_3} = \frac{F_1}{2}|\eta_\chi|^2 A_1, \quad [\tilde{g}_{\hat{h}_1\hat{h}_1}, \tilde{g}_{\hat{h}_2\hat{h}_2}] = \frac{F_1}{4}|\eta_\chi|^2(\pm 1 + C_\rho), \quad \tilde{g}_{\chi\bar{\chi}} = F_1 A_3, \quad \tilde{g}_{\rho\chi} = i\tilde{g}_{\hat{h}_3\chi} = \frac{1}{2}F_1\bar{\eta}_\chi S_\rho A_2, \\
[A_1 = C_\rho + 2\eta_\chi\bar{\eta}_\chi R S_\rho^2, \quad A_2 = X + 2\bar{\mu}_\chi\eta_\chi R(1 + C_\rho\frac{X}{X}), \quad A_3 = 1 + C_\rho X\bar{X} + 2\mu_\chi\bar{\mu}_\chi R|1 + C_\rho\frac{X}{X}|^2] \tag{III.12}
\end{aligned}$$

where we introduced the useful auxiliary functions A_1, A_2, A_3 and the abbreviations, $C_\rho = \cosh\rho, S_\rho = \sinh\rho$. For $0 < L^2 < 1, X$ becomes complex and A_2 pure imaginary. The restriction to Σ_4 at constant χ removes the derivatives $d\chi, d\eta$ and reduces the metric to a diagonal quadratic form in the S^3 tangent vectors $\hat{h}_{1,2,3}$ with ρ -dependent coefficients,

$$\begin{aligned}
d\tilde{s}^2(\mathcal{C}_6)|_{\Sigma_4} &= \bar{\eta}_\chi\eta_\chi[\frac{1}{4}F_1(\tau)(\hat{h}_1^2 - \hat{h}_2^2 + (2d\rho^2 + \hat{h}_1^2 + \hat{h}_2^2 + 2\hat{h}_3^2)\cosh\rho) + R(\tau)(d\rho^2 + \hat{h}_3^2)\sinh^2\rho] \\
&= \frac{|\eta_\chi|^2 K(\tau)}{2\epsilon^{2/3}}[K_2(\rho)(d\rho^2 + \hat{h}_3^2) + \cosh^2\frac{\rho}{2}\hat{h}_1^2 + \sinh^2\frac{\rho}{2}\hat{h}_2^2], \\
[K_2(\rho) &= \cosh(\rho) + 2|\eta_\chi|^2\sinh^2(\rho)R(\tau) = \cosh\rho - \frac{|\eta_\chi|^2\sinh^2\rho}{|\epsilon|^2\sinh^2\tau}(\cosh\tau - \frac{2}{3K^3(\tau)})]. \tag{III.13}
\end{aligned}$$

One must not confuse the c-number frame vectors $\hat{h}_k = \hat{h}_k^m d\alpha^m$ above with the corresponding differential 1-forms $h^{\hat{k}} = h_m^{\hat{k}} d\alpha^m$. These quantities admit similar decompositions on the angles differentials, $h_m^{\hat{k}} d\alpha^m$, but satisfy different c-number and wedge product composition laws, $\hat{h}_k\hat{h}_l$ and $h^{\hat{k}}\wedge h^{\hat{l}}$. The basis of derivative operators $h_{\hat{k}} \equiv \partial_{h_{\hat{k}}} = h_{\hat{k}}^m\partial/\partial\alpha^m$ is dual to the basis of 1-forms $h^{\hat{k}}$, as made explicit by the expressions of components $h_m^{\hat{k}}, h_k^m$ in Eqs. (A.5). In most parts of the text we choose ρ as the independent radial variable. The transformation from $\tau \rightarrow \rho$ via the change of variables in Eqs. (III.5) must then be accompanied by the substitutions, $g_{\rho\rho} \rightarrow \dot{\tau}^2 g_{\tau\tau}, \partial_\rho \rightarrow \dot{\tau}\partial_\tau, Det(\gamma) \rightarrow \dot{\tau}^2 Det(\gamma), [\dot{\tau} \equiv \partial\tau/\partial\rho]$. More details on the formalism are provided in Appendix A 1.

B. Meson scalar modes from brane moduli fields

The bosonic brane action in Eq. (II.14) depends on the moduli field χ and its spacetime derivatives through the bulk fields pull-back transforms. The induced metric on the brane

$$d\tilde{s}^2(D7) = \gamma_{\alpha\beta}d\xi^\alpha d\xi^\beta = \gamma_{\mu\nu}dx^\mu dx^\nu + \gamma_{\rho\rho}d\rho^2 + \gamma_{h_i h_j}\hat{h}_i\hat{h}_j + 2(\gamma_{\mu\rho}dx^\mu d\rho + \gamma_{\mu h_i}dx^\mu \hat{h}_i + \gamma_{\rho h_i}d\rho \hat{h}_i), \tag{III.14}$$

can be calculated from the bulk metric in Eq. (A.11) either using the definition, $\gamma_{\alpha\beta} = \partial_{\xi^\alpha}X^M\partial_{\xi^\beta}X^N g_{MN}$, or substituting in $d\tilde{s}^2(M_{10}) = g_{MN}dX^M dX^N$ the decomposition of the spacetime coordinates differentials, $dX^M = \sum_\alpha(\partial X^M/\partial\xi^\alpha)d\xi^\alpha$ in $d\tilde{s}^2(M_{10}) = g_{MN}dX^M dX^N$. In the static gauge, $\xi^\alpha = X^\alpha$, the induced metric acquires in addition to the bulk metric restriction $\gamma_{\alpha\beta}^{(0)} = g_{\alpha\beta}$ the extra part $\alpha_{\alpha\beta} = \gamma_{\alpha\beta} - \gamma_{\alpha\beta}^{(0)}$ comprising linear and quadratic order terms in $d\chi(x, \rho, \hat{h}_a)$. The unwarped induced metric components $\tilde{\gamma}_{\alpha\beta}$ (distinguished by tilde symbols) are related to the warped ones as, $\gamma_{\mu\nu} = h^{-1/2}\tilde{\gamma}_{\mu\nu}, \gamma_{\mu a} = h^{1/2}\tilde{\gamma}_{\mu a}, \gamma_{ab} = h^{1/2}\tilde{\gamma}_{ab}$. The general formulas for the induced metric including the dependence on $\eta_\chi = \mu_\chi Y = \mu_\chi/X$ and $\partial_\mu\chi, \partial_\rho\chi, \partial_{h_a}\chi$ are given in Eqs. (A.14). We select from these results the following formulas for the induced metric components along M_4 and radial direction of Σ_4 , that will be of direct relevance to us in the applications,

$$\begin{aligned}
\tilde{\gamma}_{\mu\nu} &= \tilde{g}_{\mu\nu} + 2F_1 h(\tau)A_3\partial_\mu\chi\partial_\nu\bar{\chi}, \quad \tilde{\gamma}_{\rho\rho} = \tilde{g}_{\rho\rho} + F_1[(\bar{\eta}_\chi\partial_\rho\chi S_\rho A_2 + H. c.) + 2\partial_\rho\chi\partial_\rho\bar{\chi}A_3], \\
\tilde{\gamma}_{\mu\rho} &= \frac{1}{2}F_1(\bar{\eta}_\chi\partial_\mu\chi S_\rho A_2 + 2\partial_\rho\bar{\chi}\partial_\mu\chi A_3 + H. c.), \quad [C_\rho = \cosh\rho, S_\rho = \sinh\rho]. \tag{III.15}
\end{aligned}$$

The auxiliary functions A_1, A_2, A_3 were defined in Eq. (III.12) and the terms linear in $\partial_\mu\chi, \partial_\rho\chi$ originate from the off-diagonal $d\rho d\chi$ components of the conifold metric. The pull-back transforms to Σ_4 of the classical solutions

for the Kähler and potential 2-forms J_2 , B_2 and the complex field strength 3-form G_3 are quoted in Eq. (A.18) of Appendix A 2 where one can find complementary details on the formalism.

The determinant factor in the $D7$ -brane action can be evaluated as a power expansion in field derivatives $\partial_{\mu,\rho,h_i}\chi$ using Eq. (A.15). Since the leading order part of the induced metric is diagonal, $g_{\alpha\beta} \propto g_{\alpha\alpha}\delta_{\alpha\beta}$, one can make use of the simplified formula

$$\frac{Det^{1/2}(g+\alpha)}{Det^{1/2}(g)} = 1 + \frac{1}{2} \sum_{\alpha} \frac{\alpha_{\alpha\alpha}}{g_{\alpha\alpha}} - \frac{1}{2} \sum_{\alpha,\beta} \frac{\alpha_{\alpha\beta}\alpha_{\beta\alpha}}{g_{\alpha\alpha}g_{\beta\beta}} + \frac{1}{8} \left(\sum_{\alpha} \frac{\alpha_{\alpha\alpha}}{g_{\alpha\alpha}} \right)^2 + O(\alpha^3). \quad (\text{III.16})$$

We ignore momentarily the contributions from B_2^{cl} and, instead of the general intricate formula up to $O(\partial_{\mu,\rho,h_i}\chi)^2$ in Eq. (A.16), quote below the approximate expression retaining the dependence on η_χ but neglecting contributions of $O(\epsilon/\mu)$ (relative to unity),

$$\begin{aligned} S_B^{(2)}(D7) &= -\mu_7 \int d^4x \int d\rho \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 e^\phi \sqrt{-\tilde{\gamma}^{(0)}} \left(1 + \frac{T_2}{|\eta_\chi|^2 T_3} (\eta_\chi \partial_\rho \bar{\chi} + \bar{\eta}_\chi \partial_\rho \chi) \right. \\ &\quad \left. + F_1 h(\tau) \left(T_1 - \frac{T_2^2}{T_3} \right) \tilde{g}^{\mu\nu} \partial_\mu \chi \partial_\nu \bar{\chi} + \frac{2}{|\eta_\chi|^2 T_3} \left(T_1 - \frac{T_2^2}{T_3} \right) \partial_\rho \chi \partial_\rho \bar{\chi} \right), \\ [T_1 &= (1 + C_\rho)T, \quad T_2 = S_\rho T, \quad T_3 = C_\rho + 2|\eta_\chi|^2 R S_\rho^2, \quad T = (1 + 2|\eta_\chi|^2 R(1 + C_\rho)), \\ F_1 &= \frac{1}{2} \epsilon^{-2/3} K(\tau), \quad R = \frac{F_2}{F_1} = \frac{1}{2\epsilon^2 \sinh^2 \tau} \left(\frac{2}{3K^3(\tau)} - \cosh \tau \right) = \frac{K'(\tau)}{2\epsilon^2 K(\tau) \sinh^2 \tau}, \\ V(\rho) &\equiv \sqrt{-\tilde{\gamma}^{(0)}} = \frac{F_1^2}{16} \sinh(2\rho) |\eta_\chi|^4 (1 + 2R|\eta_\chi|^2 \frac{\sinh^2 \rho}{\cosh \rho}) = \frac{|\eta_\chi|^4}{8} F_1^2(\tau) T_3(\rho) \sinh(\rho). \end{aligned} \quad (\text{III.17})$$

The radial integral measure, $V(\rho) = (-\tilde{\gamma}^{(0)})^{1/2}$, includes the contributions from the sub-determinants of the induced metric along M_4 and Σ_4 . Only the real quadratic order terms $|\partial_{\mu,\rho}\chi|^2$ are present while the complex terms $(\partial_{\mu,\rho}\chi)^2$ cancel out. The field components in the decompositions, $\chi = \phi_1 + i\phi_2 = |\chi|e^{i\Phi}$, are thus associated to decoupled degenerate modes in leading order of the derivatives expansion. (Note that the first and second terms in the combination $(T_1 - T_2^2/T_3)|\partial_{\mu,\rho}\chi|^2$ originate from the determinant correction terms, $\alpha_{\alpha\alpha}g^{\alpha\alpha}$ and $\alpha_{\alpha\beta}\alpha_{\beta\alpha}g^{\alpha\alpha}g^{\beta\beta}$ in Eq. (III.16), respectively, and that the second term inside parentheses is missing in the results of [42].)

In the limit $|\mu/\epsilon| \gg 1$: $Y \equiv 1/X = \eta_\chi/\mu_\chi = (1 - (L + \chi/\epsilon)^{-2})^{1/2} \rightarrow (1 - (L)^{-2})^{1/2} \simeq 1$. The results prior to taking the limit of large L^2 are obtained from Eq. (A.16) by replacing $(T_1 - T_2^2/T_3) \rightarrow (A_3 - |A_2|^2 S_\rho^2/T_3)$ in Eq. (III.17). The correspondence between the auxiliary functions $A_{1,2,3}$ in Eq. (III.15) and the real functions $T, T_{1,2,3}$ is given by $A_1 \rightarrow T_3, A_2 \rightarrow T, A_3 \rightarrow T_1, A_2 \sinh \rho \rightarrow T_2$. In order to describe accurately the off shell dependence on $\chi(\rho) \leq \mu$ of the fluctuation field derivative in the wave equation, $\partial_\rho \chi / \eta_\chi \simeq \partial_\rho \chi / (\mu + \chi(\rho))$, it is useful to retain the dependence on χ in η_χ .

We are now in a position to discuss the $D7$ -brane stability. The brane effective scalar potential is expected to vanish at its minimum due to the preserved supersymmetry. This is verified upon combining the $O((d\chi)^0)$ contribution from the scalar potential $V(\rho)$ with the linear order (tadpole) term $O(\chi)$ from integration by parts of the ρ -integral in Eq. (III.17), and expanding in powers of χ ,

$$\begin{aligned} V(\rho) &= V_0(\rho) - \left(\frac{\chi}{\eta_\chi} + \frac{\bar{\chi}}{\bar{\eta}_\chi} \right) \partial_\rho \left(\frac{V_0(\rho) T_2}{T_3} \right) \\ &\simeq V_0(\rho) + \left(\frac{\chi}{\eta_\chi} + \frac{\bar{\chi}}{\bar{\eta}_\chi} \right) \frac{|\mu|^4}{8} [2F_1^2 S_\rho (C_\rho + 3R|\mu|^2 S_\rho^2) - \partial_\rho (F_1^2 S_\rho^2 (1 + 2R|\mu|^2 (1 + C_\rho)))] + O(\chi^2), \end{aligned} \quad (\text{III.18})$$

where $V_0(\rho) = V(\rho)|_{\chi=0}$. The verification is easier in the singular conifold limit where the net effective potential admits the expansion in powers of χ ,

$$\begin{aligned} V(\rho) &= \mu_7 e^\phi [\sqrt{-\gamma^{(0)}} + \sqrt{-\gamma^{(0)}} \gamma^{ab} \gamma_{\chi a} \partial_b \chi] = \mu_7 e^\phi [V_0(\rho) - \left(\frac{\chi}{\eta} + \frac{\bar{\chi}}{\bar{\eta}} \right) \frac{\partial}{\partial \rho} \left(\frac{V_0(\rho) T_2}{T_3} \right)_{\chi=0}] \\ &\simeq \mu_7 e^\phi \left[\frac{|\mu + \chi|^{8/3}}{96} - \frac{4}{3} \frac{|\mu|^{8/3}}{96} \left(\frac{\chi}{\mu} + \frac{\bar{\chi}}{\bar{\mu}} \right) \right] (2r_\chi^3 - 1) \left(\frac{r_\chi^3 - 2}{r_\chi} \right)^{1/2} + O(\chi^2), \end{aligned} \quad (\text{III.19})$$

showing explicitly how the linear order terms in $V(\rho) \propto |\mu|^{8/3} \chi$ cancel out after integrating by parts the $O(\chi)$ source term. Since the finite term V_0 can always be removed through a constant shift of the action, the net brane potential of $O(\chi^2)$ is minimized at the stable vacuum (on shell) value for constant $\chi(\rho) = 0$. Whether the cancellation still

holds upon switching on the background 2-form field B_2^{cl} is verified by combining the Born-Infeld and Chern-Simons actions and replacing the metric determinant $\tilde{\gamma}^{(0)} \rightarrow \tilde{q}$, as detailed in Appendix A 2,

$$V_0(\rho) + V_0^{CS}(\rho) = \mu_7 [e^\phi \sqrt{-\tilde{q}} - \frac{1}{2} C_4 \wedge B^2] = \mu_7 [e^\phi (\sqrt{-\tilde{\gamma}^{(0)}}) + e^{-\phi} \frac{Pf(B)}{h(\tau)} - \frac{Pf(B)}{h(\tau)}] = \mu_7 e^\phi V_0(\rho). \quad (\text{III.20})$$

We now return to the diagonal terms in the reduced $D7$ -brane action in Eq. (III.17) and express the quadratic order in $|d\chi|$ in terms of the effective brane inverse metric components $G^{\mu\nu}$, $G^{\rho\rho}$,

$$\begin{aligned} \delta S_B^{(2)}(D7) &= -\mu_7 \int d^4x \int d\rho \hat{h}_1 \wedge \hat{h}_1 \wedge \hat{h}_3 e^\phi V_0(\rho) (G^{\mu\nu} \partial_\mu \chi \partial_\nu \bar{\chi} + G^{\rho\rho} \partial_\rho \chi \partial_\rho \bar{\chi}), \\ [G^{\mu\nu} = \tilde{g}^{\mu\nu} Q(\rho) = \tilde{g}^{\mu\nu} F_1(\rho) h(\tau) A(\rho), \quad G^{\rho\rho} = \frac{2P(\rho)}{|\eta_\chi|^2} = \frac{2A(\rho)}{|\eta_\chi|^2 T_3}, \quad A(\rho) = (A_3 - \frac{|A_2|^2 S_\rho^2}{A_1})]. \end{aligned} \quad (\text{III.21})$$

The Kaluza-Klein decomposition of the complex moduli field

$$\chi(\xi) = \sum_m \chi^{(m)}(x) e^{ik_m \cdot x} \Psi_m(\rho, \alpha) = \sum_{m,j} \chi^{(m)}(x) e^{ik_m \cdot x} f_{m,j}(\rho) \Phi_{m_l, m_r}^j(\alpha), \quad (\text{III.22})$$

introduces mode fields $\chi^{(m)}(x)$ in M_4 , labelled by the radial and angular excitation indices k_m , whose wave functions $\Psi_m(\rho, \alpha)$, $[\alpha = (\theta, \phi, \gamma)]$ are linear combinations with radial coefficient functions $f_{m,j}(\rho)$ of the $S^3 = SO(4)/SO(3)$ scalar harmonic functions $\Phi_{m_l, m_r}^j(\alpha) \sim \mathcal{Y}_{m_l, m_r}^{l/2, l/2}(\alpha)$, $[j = \frac{1}{2} = 0, 1/2, \dots, (m_l, m_r) \in [-j, \dots, j]]$.

Substituting the decomposition in Eq. (III.22) into the perturbed action in Eq. (III.21) yields a coupled system of second order differential equations of Sturm-Liouville type for the radial wave functions $f_{m,j}(\rho)$. We shall restrict our study to the (angle independent) singlet modes described by radial wave functions $f_m(\rho) = f_{m,j=0}(\rho)$. The volume integral over the compact base $\int \hat{h}_1 \wedge \hat{h}_1 \wedge \hat{h}_3 = 64\pi^2$ can then be absorbed into the overall constant factor including the brane tension, $C = 64\pi^2 \mu_7$. One convenient way to derive the wave equations is by adding and subtracting the diagonal terms $\omega_m^2 \delta_{mn}$ so as to separate the quadratic order action for the $\chi^{(m)}(x)$ into kinetic and mass terms and on-shell constraint terms,

$$\begin{aligned} \delta S_B^{(2)}(D7)(\chi) &= C \int d^4x d\rho e^\phi \sqrt{-\tilde{\gamma}^{(0)}} \sum_{m,n} \bar{\chi}^{(m)}(x) f_m^\dagger(\rho) [Q(\rho) (\tilde{\nabla}_4^2 - \omega_m^2 \delta_{mn}) \\ &+ (Q(\rho) \omega_m^2 \delta_{mn} + \frac{1}{\sqrt{-\tilde{\gamma}^{(0)}}} \partial_\rho \mathcal{G} \partial_\rho)] \chi^{(n)}(x) f_n(\rho), \\ [Q(\rho) = F_1(\rho) h(\tau) A(\rho), \quad \mathcal{G} = G^{\rho\rho} \sqrt{-\tilde{\gamma}^{(0)}} = \frac{2\sqrt{-\tilde{\gamma}^{(0)}}}{|\eta_\chi|^2} P(\rho), \quad P(\rho) = \frac{A(\rho)}{T_3}, \quad \omega_m^2 = -k_m^2]. \end{aligned} \quad (\text{III.23})$$

The classical field B_2^{cl} can be included at this stage by simply replacing $A(\rho) \rightarrow [A_3 - (|A_2 S_\rho|^2 / (A_1 R_S))]$, as explained near Eqs. (A.24) or (A.27) in Appendix A 2. The rescaled fields $\chi^{(m)}(x) \rightarrow \chi^{(m)}(x) / \sqrt{C}$ with canonical kinetic actions, $\delta S^{(2)} = \int d^4x \bar{\chi}^{(m)}(x) (\tilde{\nabla}_4^2 - \omega_m^2) \chi^{(m)}(x)$, are associated to orthonormal modes of squared masses ω_m^2 and wave functions $f_m(\rho)$ satisfying the wave equations and normalization conditions,

$$\begin{aligned} 0 &= \sqrt{-\tilde{\gamma}^{(0)}} (Q(\rho) \omega_m^2 + \frac{1}{\sqrt{-\tilde{\gamma}^{(0)}}} \partial_\rho \mathcal{G} \partial_\rho) f_m(\rho) = \mathcal{G} (\frac{Q(\rho)}{G^{\rho\rho}} \omega_m^2 - \frac{1}{\mathcal{G}} \partial_\rho \mathcal{G} \partial_\rho) f_m(\rho), \\ [\int d\rho \sqrt{-\tilde{\gamma}^{(0)}} Q(\rho) f_m^\dagger(\rho) f_{n'}(\rho) = \delta_{mm'}]. \end{aligned} \quad (\text{III.24})$$

One could have reached the above results also by requiring the stationary condition $\partial S_B(D7) / \partial \bar{\chi}^{(m)} = 0$, subject to the mass shell constraint, $\tilde{\nabla}_4^2 \rightarrow \omega_m^2$. The wave equation can be transformed via the wave function rescaling $f_m = \tilde{f}_m / \mathcal{G}^{1/2}(\rho)$ to the Schrodinger type form

$$\begin{aligned} (\partial_\rho^2 - V_{eff}) \tilde{f}_m(\rho) &= 0, \quad [V_{eff} = \mathcal{G}_1 - \frac{\omega_m^2 Q(\rho)}{G^{\rho\rho}} = \mathcal{G}_1 - \hat{\omega}_m^2 L^2 \frac{\hat{Q}(\rho)}{2P(\rho)}, \quad \mathcal{G}_1 = \frac{(\mathcal{G}^{1/2})''}{\mathcal{G}^{1/2}}, \quad P(\rho) = \frac{1}{2} |\eta_\chi|^2 G^{\rho\rho}, \\ \omega_m^2 = \frac{\epsilon^{4/3} \hat{\omega}_m^2}{(g_s M \alpha')^2}, \quad Q(\rho) = \frac{(g_s M \alpha')^2}{\epsilon^{10/3}} \hat{Q}(\rho), \quad \hat{Q}(\rho) = \frac{1}{2^{1/3}} K(\tau) I(\tau) A(\rho)] \end{aligned} \quad (\text{III.25})$$

where the prime denotes the $\partial/\partial\rho$ derivative and the normalization integral for wave functions is changed to, $\int d\rho \sqrt{-\tilde{\gamma}^{(0)}} (Q(\rho)/\mathcal{G}) \tilde{f}_m^\dagger(\rho) \tilde{f}_{m'}(\rho) = \delta_{mm'}$. The dimensionless effective potential V_{eff} depends on the ratio parameter $L = \mu/\epsilon$ and the rescaled masses $\hat{\omega}_m^2 = \epsilon^{-4/3} (g_s M \alpha')^2 \omega_m^2$. Note the naive energy dimensions, $[f_m(\rho)] =$

$E^{-3/2}$, $[\tilde{f}_m(\rho)] = E^0$, $[\mu] = [\epsilon] = E^{-2/3}$ and the fact that $A(\rho)$ cancels out in the mass dependent term, $\omega_m^2 Q(\rho)V(\rho)/\mathcal{G} = \omega_m^2 F_1 h|\eta|^2 T_3/2$, but not in the term \mathcal{G}_1 .

The discrete mass spectrum for the ground state and radially excited modes is evaluated by applying the semi-classical Bohr-Sommerfeld quantization rule,

$$(n - \frac{1}{2} + \frac{\delta_W}{4})\pi = \int_{\rho_{min}}^{\rho_0} d\rho (-V_{eff}(\rho))^{1/2}, \quad [n = 1, 2, \dots, V_{eff}(\rho_0) = 0, \rho_{min} = 0] \quad (\text{III.26})$$

with corresponding wave functions in the inner well region before the turning point ρ_0 of form

$$\tilde{f}_m(\rho) = \frac{C_m}{(-V_{eff}(\rho))^{1/4}} \sin(\frac{\pi}{4} + \int_{\rho}^{\rho_0} d\rho' (-V_{eff}(\rho'))^{1/2}). \quad (\text{III.27})$$

The barrier parameter δ_W in the phase integral is set to 0 if the potential slopes at the turning point near threshold $\rho_{min} = 0$ and to 1 if the potential is finite there which amounts to impose a hard wall forcing the wave function to vanish at the origin. At large ρ , the two terms in the effective potential V_{eff} vanish exponentially. Near the origin $\rho = \rho_{min} = 0$, the second (mass dependent) term is regular but the first (mass independent) term diverges as $\mathcal{G}_1 \rightarrow -\frac{1}{4\rho^2}$. The quantization rule in Eq. (III.26) is thus invalidated due to the divergent contribution to the phase integral near the origin diverges, $\int_{\rho_{min}} d\rho (-V_{eff}(\rho))^{1/2} = 1/(4\rho_{min})$. The divergence is still present if one used τ as the independent radial variable [36] in place of ρ . Analyzing the limit in the case $L^2 > 1$, in terms of the limited expansions in powers of $\tau - \tau_{min}$, one finds that the contribution to the phase integral near the origin diverges logarithmically,

$$\begin{aligned} \rho &= 2\alpha(\tau - \tau_{min})^{1/2}, \quad \frac{d\tau}{d\rho} = \alpha^{-1}(\tau - \tau_{min})^{1/2}, \quad \mathcal{G}_1 = -\frac{1}{(4\alpha)^2(\tau - \tau_{min})}, \quad [\alpha = \frac{L^{1/2}}{(L^2 - 1)^{1/4}}] \\ \implies \int_{\tau_{min}} \frac{d\tau}{d\tau/d\rho} (-\mathcal{G}_1)^{1/2} &= \int_{\tau_{min}} d\tau \frac{1}{4(\tau - \tau_{min})} = -\frac{1}{4} \ln(\tau - \tau_{min}). \end{aligned} \quad (\text{III.28})$$

Since the divergence is mode independent, one could consider the artificial cure of imposing an infrared cutoff $\rho \geq \rho'_0$. However, the natural way to remove the singularity is by choosing an appropriate boundary condition on the radial wave functions. Proceeding along same lines as the modified semi-classical approach of [63, 64]. we are led to consider the wave function rescaling and change of radial variable, $f_m = (\rho/\mathcal{G})^{1/2} \hat{f}_m$, $\rho = e^y$, $[y \in (-\infty, \infty)]$ transforming Eq. (III.25) to the equivalent wave equation,

$$0 = e^{-3y/2} (\partial_y^2 - W_{eff}) \hat{f}_m, \quad [W_{eff} = \frac{1}{4} + e^{2y} V_{eff} = \rho^2 (\hat{\mathcal{G}}_1 - \omega_m^2 \frac{Q(\rho)V_0(\rho)}{\mathcal{G}}), \quad \hat{\mathcal{G}}_1 = \mathcal{G}_1 + \frac{1}{4\rho^2}] \quad (\text{III.29})$$

where the modified effective potential W_{eff} is everywhere regular, thanks to the shift transformation $\mathcal{G}_1 \rightarrow \hat{\mathcal{G}}_1$ with $\hat{\mathcal{G}}_1(0) = 0$. (For the sake of completeness, we note the alternative choice for the radial variable made in [47] $e^y = e^\rho - 1$. We also mention the similar kind of singular wave equation encountered in the analysis of scalar glueballs in [62], where taking advantage of the supersymmetric type structure of the wave operator as a product of first order operators, $Q_1 Q_2 w = m^2 w$, the authors replaced the initial wave equation by the regular one, $Q_2 Q_1 \hat{w} = m^2 \hat{w}$, retaining the same mass spectrum.) Substituting the power expansion near $\rho \rightarrow 0$ of the wave function f_m into the wave equation,

$$0 = (\partial_\rho^2 - V_{eff}) \tilde{f}_m \simeq (\partial_\rho^2 + \frac{1}{4\rho^2}) (\rho^{1/2} \hat{f}_m), \quad [\tilde{f}_m \equiv f_m \sqrt{\mathcal{G}} \equiv \rho^{1/2} \hat{f}_m = \rho^{1/2} (x_0 + x_1 \rho + x_2 \rho^2 + \dots)] \quad (\text{III.30})$$

requires imposing $x_1 = 0$ while leaving x_0, x_2 arbitrary. It follows that $f_m(0) \neq 0$, $\partial_\rho f_m(0) = 0$ are allowed boundary conditions at the origin. The quantization rule for the modified wave equation in Eq. (III.29),

$$\begin{aligned} (n - \frac{1}{2} + \frac{\delta_W}{4})\pi &= \int_{-\infty}^{y_0} dy (-W_{eff}(y))^{1/2} = \int_0^{\rho_0} d\rho (-\hat{V}_{eff}(\rho))^{1/2}, \quad [\hat{V}_{eff}(\rho_0) = 0, n = 1, 2, \dots, \\ \hat{V}_{eff}(\rho) &= (\mathcal{G}_1 + \frac{1}{4\rho^2}) - \omega_m^2 \frac{Q(\rho)}{\mathcal{G}^{\rho\rho}} = \hat{\mathcal{G}}_1 - \omega_m^2 \frac{Q(\rho)V_0(\rho)}{\mathcal{G}}, \quad G^{\rho\rho} = \frac{\mathcal{G}}{V_0(\rho)}] \end{aligned} \quad (\text{III.31})$$

is seen to take the standard form when expressed in terms of the auxiliary potential, $\hat{V}_{eff}(\rho) = W_{eff}(y)/\rho^2$. We can then compute the mass spectra by applying the standard numerical procedure [65], used in our earlier work [49], to the modified potential $\hat{V}_{eff}(\rho)$ and evaluate the rescaled wave functions using

$$\hat{f}_m(\rho) = \frac{C_m}{\rho^{1/2} (-\hat{V}_{eff}(\rho))^{1/4}} \sin(\frac{\pi}{4} + \int_{\rho}^{\rho_0} d\rho' (-\hat{V}_{eff}(\rho'))^{1/2}), \quad [\hat{V}_{eff} = \frac{W_{eff}}{\rho^2}]. \quad (\text{III.32})$$

We now examine the couplings for scalar modes $\chi^{(m)}(x)$ inferred from expansion of the action in powers of χ of the overall factor in the effective potential

$$V_{eff}(\rho) \simeq V_0(\rho)|\mu + \chi|^4 = V_0(\rho)(1 + (2\chi/\mu + (\chi/\mu)^2 + H. c.) + 4|\chi/\mu|^4 + \dots), [V_0(\rho) = \frac{|\mu|^2}{8} F_1^2 S_\rho A_1(\rho)] \quad \text{III.33}$$

This is seen to contribute mass terms to the meson modes, replacing in the wave equation, $\hat{\omega}_m^2 \rightarrow \hat{\omega}_m'^2 = \hat{\omega}_m^2 - 4/(|L|^2 \hat{Q}(\rho))$, $[\hat{Q}(\rho) = 2^{-1/3} K(\tau) I(\tau) (A_3 - |A_2 S_\rho|^2 / A_1)]$. The correction to the squared mass parameter has a negative sign and is sizeable at small L where $\hat{Q}(\rho) \rightarrow 0$ at large ρ . The kinetic energy term in Eq. (III.23) produces couplings of meson modes to massive gravitons. These are derived by substituting $\tilde{g}_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} + h_{\mu\nu}$ and decomposing the graviton field into modes $h_{\mu\nu}^{(m)}(x)$,

$$h_{\mu\nu}(X) = \frac{1}{\sqrt{\lambda_2}} \sum_m N_m h_{\mu\nu}^{(m)}(x) R_m(\tau) \Phi_m(\Theta), [N_m = \frac{2^{13/6} \sqrt{3} \epsilon^{-2/3}}{g_s M \alpha' J_{(m)}}, \frac{1}{\sqrt{\lambda_2}} = \frac{2\sqrt{V_W}}{M_\star} = \frac{2(2\pi)^3 \eta_W^3 \mathcal{R}^3}{M_\star},$$

$$\eta_W = \frac{L_W}{\mathcal{R}}, J_{(m)} = (\int d\tau \frac{I(\tau)}{K^2(\tau)} B_m^\dagger B_m(\tau))^{1/2}, R_m = \frac{B_m}{\tilde{G}^{1/2}}] \quad \text{III.34}$$

in same notations as [53]. Starting from the kinetic action,

$$\delta S_{B,1}^{(2)}(D7) = - \int_{M_4} d^4x \int d\rho W(\rho) (\tilde{g}^{\mu\nu} - h^{\mu\nu}) \partial_\mu \chi^\dagger \partial_\nu \chi,$$

$$[W(\rho) = C' V_0(\rho) Q(\rho) = \frac{C' |\mu \chi|^4}{8} F_1^3 h(\tau) S_\rho T_3 A(\rho), C' = C e^\phi = 64\pi^2 \mu_7 g_s = \frac{g_s m_s^8}{2\pi^5}] \quad \text{III.35}$$

and expanding the factor $|\mu \chi|^4$ in powers of χ , as detailed in Eq. (III.33), yields interactions for graviton modes coupled to canonically normalized mesons mode fields, $\chi^{(m)} \rightarrow \chi^{(m)}/(\int d\rho W_0 f_m^2)^{1/2}$, represented by operators of dimensions 4, 5, 6,

$$\delta S_{B,1}^{(2)}(D7) = + \int_{M_4} d^4x \left(- \tilde{g}^{\mu\nu} \partial_\mu \chi^{(m)\dagger} \partial_\nu \chi^{(m)}(x) + \frac{N_p}{\sqrt{\lambda_2}} h^{(p)\mu\nu}(x) \partial_\mu \chi^{(m)\dagger} \partial_\nu \chi^{(n)}(x) \int d\rho W_0(\rho) \frac{f_m f_n(\rho) R_p(\tau)}{\prod_{x=m,n} (\int d\rho W_0 f_x^2)^{1/2}} \right.$$

$$\left. + h^{(p)\mu\nu}(x) \partial_\mu \chi^{(m)\dagger} \partial_\nu \chi^{(n)}(x) \chi^{(l)}(x) \int d\rho W_1(\rho) \frac{f_m f_n f_l(\rho) R_p(\tau)}{\prod_{x=m,n,l} (\int d\rho W_0 f_x^2)^{1/2}} \right),$$

$$[W_0 = \frac{C' |\mu|^4}{8} F_1^3 h(\tau) S_\rho (T_1 T_3 - T_2^2), W_1 = \frac{4W_0}{\mu}, W_2 = \frac{6W_0}{\mu^2}]. \quad \text{III.36}$$

The kinetic energy term in Eq. (III.35) for the $D7$ -brane action also contributes self couplings for the $\chi^{(m)}(x)$ represented by operators of dimension 5

$$\delta S_{B,2}^{(2)}(D7) = - \int_{M_4} d^4x \tilde{g}^{\mu\nu} \partial_\mu \chi^{(m)\dagger} \partial_\nu \chi^{(n)}(x) \chi^{(p)}(x) \frac{\int d\rho W_1 f_m f_n f_p}{\prod_{x=m,n,p} (\int d\rho W_0 f_x^2)^{1/2}}, \quad \text{III.37}$$

where W_1 is specified in Eq. (III.36). Substituting $\chi \rightarrow \chi^{(m)}$ in the radial derivative part of the $D7$ -brane action,

$$\delta S_{B,3}^{(2)}(D7) = - \int_{M_4} d^4x \int d\rho S(\rho) \partial_\rho \chi^\dagger \partial_\rho \chi,$$

$$[S(\rho) = \frac{C' |\mu \chi|^2}{4} F_1^2 S_\rho X(\rho) = S_0 |1 + \frac{\chi}{\mu}|^2 = S_0 + (\frac{\chi}{\mu} + H. c.) S_1 + |\frac{\chi}{\mu}|^2 S_2] \quad \text{III.38}$$

yields, via the factor $|\mu \chi|^2$, local self couplings represented by operators of dimensions 2, 3, 4,

$$\delta S_{B,2}^{(2)}(D7) = - \int d^4x \chi^{(m)\dagger} \chi^{(n)}(x) \int d\rho [S_0(\rho) \partial_\rho f_m^\dagger \partial_\rho f_n + (\frac{\chi^{(p)}(x)}{\mu} S_1(\rho) \partial_\rho f_m^\dagger \partial_\rho f_n f_p + H. c.)$$

$$+ \chi^{(p)}(x) \chi^{(q)\dagger}(x) \frac{S_2(\rho)}{|\mu|^2} \partial_\rho f_m \partial_\rho f_n f_p f_q^\dagger], \quad \text{III.39}$$

where we absorbed the normalization factors inside the wave functions by substituting, $f_x/(\int d\rho W_0 f_x^2)^{1/2} \rightarrow f_x$.

C. Meson modes from $D7$ -brane gauge sector

We here examine the mesons modes descending from the $D7$ -brane gauge connection 1-form field, $A = A_\alpha d\xi^\alpha$. We shall include from the outset the 2-form classical field B_2 inside the brane effective metric tensor, $q_{\alpha\beta} = \gamma_{\alpha\beta} + e^{-\phi/2} \mathcal{F}_{\alpha\beta}^{cl}$. The contributions from the Born-Infeld and Chern-Simons parts in the Einstein frame action

$$S_B(D7) = \mu_7 \int d^8\xi [-e^\phi Det^{1/2}(-(\gamma + e^{-\phi/2}\mathcal{F})) + (C_8 + C_6 \wedge \mathcal{F} + \frac{1}{2}C_4 \wedge \mathcal{F}^2)], \quad (\text{III.40})$$

depend on the 2-form combination of the gauge field strength $F = dA$ and NSNS sector B_2 potential, $\mathcal{F} = B + F$, $F = dA$, and the RR sector $C_{8,6,4}$ -form potentials [66]. In the present classical background solution, C_8 is absent, C_6 vanishes identically, $F_7 \equiv dC_6 = \star_{10}F_3 - C_4 \wedge H_3 = 0$ [34], and C_4 takes the simple form

$$C_4(\tau) = \int_0^\tau d\tau' \tilde{F}_5(\tau') = [4\epsilon^{-8/3}(g_s M\alpha')^2 \int_0^\tau d\tau' (l(\tau')/(K(\tau') \sinh(\tau'))^2)]^{-1} vol(M_4) = vol(M_4)/h(\tau) \quad (\text{III.41})$$

The determinant of the matrix q separates into two factors associated to the two diagonal blocks in the M_4 and Σ_4 vector spaces,

$$\begin{aligned} Det^{1/2}(-(\gamma + e^{-\phi/2}\mathcal{F})) &= Det^{1/2}(-q)_{\alpha\beta} Det^{1/2}(1 + e^{-\phi/2} q^{\alpha\beta} F_{\beta\alpha}) \\ &= Det^{1/2}(-\tilde{\gamma})_{\mu\nu} Det^{1/2}(\tilde{\gamma} + e^{-\phi/2} h^{-1/2} B)_{ab} Det^{1/2}(1 + e^{-\phi} q^{-1} F)_{\alpha\beta}, \\ [q_{\mu\nu} &= h^{-1/2} \tilde{\gamma}_{\mu\nu} + e^{-\phi/2} B_{\mu\nu}, \quad q_{ab} = h^{1/2} \tilde{\gamma}_{ab} + \mathcal{F}_{ab}, \\ Det^{1/2}(-\tilde{q})_{\alpha\beta} &= Det^{1/2}(-\tilde{\gamma})_{\mu\nu} Det^{1/2}(\tilde{q})_{ab} = \sqrt{-\tilde{\gamma}^{(0)}} (Det^{1/2}(\tilde{\gamma}) + e^{-\phi} h^{-1}(\tau) Pf(B))_{ab} \end{aligned} \quad (\text{III.42})$$

where we have ignored in the final line entry above the components $B_{\mu\nu}$ along M_4 . The expansions of the determinant $Det^{1/2}(1 + e^{-\phi/2} q^{-1} F)$ and exponential factor e^F in powers of F yields the 4-d gauge action of quadratic order in F

$$\delta S_B^{(2)}(D7) = \mu_7 \int d^4x \wedge d\rho \wedge \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 [-\sqrt{-\tilde{q}}(e^\phi - \frac{1}{4} Tr(q^{-1} F q^{-1} F) + \frac{1}{8} (Tr(\tilde{q}^{-1} F))^2) + \frac{1}{2} C_4 \wedge \mathcal{F} \wedge \mathcal{F}] \quad (\text{III.43})$$

The Kaluza-Klein ansatz for the gauge connection components, $A_\alpha(X) = (A_\mu, A_a) = (A_\mu, A_\rho, A_k)$,

$$A_\mu(X) = \sum_{m,j} a_\mu^{(m)}(x) a_{m,j}(\rho) \Phi^j(\alpha), \quad A_{[\rho,k]}(X) = \sum_{m,j} a_{[\rho,k]}^{(m)}(x) g_{m,j}(\rho) \Phi_{[\rho,k]}^j(\alpha), \quad (\text{III.44})$$

introduces fields in M_4 belonging to a single tower of vector modes and three towers of scalar (gauge invariant) modes $[a_\mu^{(m)}(x), a_\rho^{(m)}(x), a_k^{(m)}(x)]$. The wave functions are given by linear combinations of radial functions $[a_{m,j}(\rho), g_{m,j}(\rho)]$ times scalar and vector harmonics $\Phi^j(\alpha), \Phi_k^j(\alpha)$ of the compact base manifold S^3/Z_2 . Upon removing the classical part $-\mu_7 \int d^4x \sqrt{-\tilde{q}} e^\phi$ and using the relations, $q^{\mu\nu} = h^{1/2} \tilde{g}^{\mu\nu}$, $q^{\rho\rho} = h^{-1/2} \tilde{q}^{\rho\rho}$, $\sqrt{-q} = \sqrt{-\tilde{q}}$, the reduced action for vector modes $a_\mu^{(m)}$ in Lorentz gauge $\partial_\mu a^{(m)\mu} = 0$ becomes

$$\begin{aligned} \delta S_B^{(2)}(a_\mu) &= +\frac{\mu_7}{4} \int d^4x \int d\rho \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 e^\phi \sqrt{-q} (q^{\alpha\beta} F_{\beta\gamma} q^{\gamma\delta} F_{\delta\alpha} - \frac{1}{2} (q^{\alpha\beta} F_{\beta\alpha})^2) \\ &= -\frac{\mu_7}{2} \int d^4x \int d\rho \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 e^\phi \sqrt{-\tilde{q}} [\tilde{g}^{\mu\nu} \tilde{g}^{\lambda\rho} h(\tau) \partial_\mu A_\lambda \partial_\nu A_\rho + \tilde{g}^{\mu\nu} \tilde{q}^{\rho\rho} \partial_\rho A_\mu \partial_\rho A_\nu]. \end{aligned} \quad (\text{III.45})$$

Selecting in Eq. (III.44) only the singlet modes $A_\mu \rightarrow a_\mu^{(m)}(x) a_m(\rho)$ of squared masses $\tilde{\nabla}_4^2 \rightarrow \omega_m^2$, and separating the terms in the sum over modes into kinetic and on-shell constraint parts (by subtracting and adding the mass terms $\omega_m^2 |a_\mu^{(m)}|^2$) yields

$$\begin{aligned} \delta S_B^{(2)}(a_\mu) &= +\frac{C}{2} \int d^4x \int d\rho \sqrt{-\tilde{q}} \tilde{g}^{\mu\nu} a_\mu^{(m)\dagger}(x) a_m^\dagger(\rho) [h(\tau) (\tilde{\nabla}_4^2 - \omega_m^2) \\ &+ (h(\tau) \omega_m^2 + \frac{1}{\sqrt{-\tilde{q}}} \partial_\rho \tilde{q}^{\rho\rho} \sqrt{-\tilde{q}} \partial_\rho)] a_\nu^{(m)}(x) a_m(\rho), \quad [C = 64\pi^2 \mu_7 = \frac{1}{2\pi^5 l_8^8}]. \end{aligned} \quad (\text{III.46})$$

One can now deduce by inspection the equations for the modes wave functions and their normalization conditions,

$$0 = [\frac{\omega_m^2 |\eta_\chi|^2 Q(\rho)}{2P(\rho)} + \frac{1}{\mathcal{G}_a} \partial_\rho \mathcal{G}_a \partial_\rho] a_m(\rho) = [\frac{1}{2^{4/3}} \hat{\omega}_m^2 L^2 K(\tau) I(\tau) R_S + \frac{1}{\mathcal{G}_a} \partial_\rho \mathcal{G}_a \partial_\rho] a_m(\rho),$$

$$\begin{aligned} [\omega_m^2 = \epsilon^{4/3}(g_s M \alpha')^{-2} \hat{\omega}_m^2, \quad Q = h(\tau), \quad P(\rho) = \frac{|\eta_\chi|^2 \tilde{q}^{\rho\rho}}{2} = \frac{|\eta_\chi|^2 \tilde{\gamma}^{\rho\rho}}{2R_S}, \\ \mathcal{G}_a(\rho) = \sqrt{-\tilde{q}} \tilde{q}^{\rho\rho}, \quad R_S = \frac{\sqrt{-\tilde{q}}}{\sqrt{-\tilde{\gamma}}}, \quad \tilde{\gamma}_{\rho\rho} = \frac{1}{\tilde{\gamma}^{\rho\rho}} \frac{1}{2} F_1 |\eta_\chi|^2 A_1, \quad \int d\rho \sqrt{-\tilde{q}} Q(\rho) a_m^\dagger(\rho) a_{m'}(\rho) = \delta_{mm'}] \quad (\text{III.47}) \end{aligned}$$

where only the first term involving the mass parameter ω_m depends on the classical field B_2^c , unlike the second term, $\mathcal{G}_a(\rho) = \sqrt{-\tilde{q}} \tilde{q}^{\rho\rho} = \sqrt{-\tilde{\gamma}} \tilde{\gamma}^{\rho\rho}$, as inferred from Eq. (A.23). The wave functions rescaling $a_m(\rho) \rightarrow \tilde{a}_m(\rho)/\mathcal{G}_a^{1/2}$ transforms the wave equations to the Schrödinger form, $(\partial_\rho^2 - V_{eff})\tilde{a}_m(\rho) = 0$,

$$V_{eff}^{(m)}(\rho) = \mathcal{G}_{a,1} - \frac{Q\omega_m^2}{\tilde{q}^{\rho\rho}} = \mathcal{G}_{a,1} - |\eta_\chi|^2 \frac{Q}{2P} \omega_m^2 = \mathcal{G}_{a,1} - \frac{\hat{\omega}_m^2 R_S}{2^{4/3}} |L|^2 I(\tau) K(\tau) A_1(\rho), \quad [\mathcal{G}_{a,1} = \frac{(\mathcal{G}_a^{1/2}(\rho))''}{\mathcal{G}_a^{1/2}(\rho)}] \quad (\text{III.48})$$

The dimensionless effective potential $V_{eff}^{(m)}$ is formally similar to that of χ modes in Eq. (III.25) but with different auxiliary functions $Q(\rho)$, $P(\rho)$. The mass independent contribution again diverges near the origin, $\mathcal{G}_a \sim 1/\rho$, $\mathcal{G}_{a,1} \sim -1/(4\rho^2)$, so the semi-classical method must be modified by changing the independent variable $\rho \rightarrow y = \ln \rho$ and the wave functions $a_m(\rho) \rightarrow (\rho/\mathcal{G}_a)^{1/2} \hat{a}_m(\rho)$, as in Eq. (III.29). One can then evaluate the masses by applying the familiar quantization rule to the modified effective potential, similarly to Eq. (III.31),

$$\hat{V}_{eff}^{(m)}(\rho) = (\mathcal{G}_{a,1} + \frac{1}{4\rho^2}) - \omega_m^2 \frac{Q(\rho)}{\tilde{q}^{\rho\rho}} = \hat{\mathcal{G}}_{a,1} - \omega_m^2 \frac{Q(\rho)\sqrt{-\tilde{q}}}{\mathcal{G}_a}, \quad [\hat{\mathcal{G}}_{a,1} = \mathcal{G}_{a,1} + \frac{1}{4\rho^2}]. \quad (\text{III.49})$$

The coupling of gravitons to pairs of brane modes can be obtained by substituting in Eq. (III.45) for the reduced action the decomposition on singlet modes, $a_\mu^{(m)}$, $[a_\mu^{(m)} k^\mu = 0]$. The resulting 4-d effective gravitational action is given in same notations as Eq. (III.34) by

$$\delta S_{B,1}^{(2)}(D7) = + \int d^4 x \lambda_{mnp}^h h^{(p)\mu\nu} \partial_\lambda a_\mu^{(m)} \partial^\lambda a_\nu^{(n)}, \quad [\lambda_{mnp}^h = \frac{C'}{2\sqrt{\lambda_2}} \frac{\int d\rho V(\rho) R_S h(\tau) \tilde{a}_m^\dagger \tilde{a}_n(\rho) R_p(\tau)}{(\prod_{x=m,n} \int d\rho' W_A(\rho') |\tilde{a}_x|^2)^{1/2}}]. \quad (\text{III.50})$$

We consider next the components of the gauge field 1-form $A = A_\alpha d\xi^\alpha = A_\mu dx^\mu + A_\rho d\rho + A_{\hat{k}} h^{\hat{k}}$ along the radial and angular directions of the tangent space of Σ_4 in $M_4 \times \Sigma_4$ and settle again to the static gauge. Note that these are linearly related to the tangent frame vielbeins components, $A = A_{\hat{\rho}} e^{\hat{\rho}} + A_{\hat{\epsilon}^k} e^{\hat{\epsilon}^k} = A_{\hat{\rho}} \eta^{\hat{\rho}} d\rho + A_{\hat{k}} \eta^{\hat{k}} h_m^{\hat{k}} d\alpha^m$, as $A_\rho = A_{\hat{\rho}} \eta^{\hat{\rho}}$, $A_{\hat{k}} = A_{\hat{\epsilon}^k} \eta^{\hat{\epsilon}^k}$, $A_m = A_{\hat{k}} h_m^{\hat{k}}$. Substituting the decomposition for the field strength 2-form $F = dA = \frac{1}{2} F_{\alpha\beta} d\xi^\alpha \wedge d\xi^\beta$,

$$\begin{aligned} F = \frac{1}{2!} F_{\nu\mu} dx^\nu \wedge dx^\mu + F_{\rho\mu} d\rho \wedge dx^\mu + F_{\rho\hat{k}} d\rho \wedge h^{\hat{k}} + \frac{1}{2!} F_{\hat{k}\hat{l}} h^{\hat{k}} \wedge h^{\hat{l}}, \\ [F_{\nu\mu} = 2\partial_{[\nu} A_{\mu]}, \quad F_{\rho\nu} = 2\partial_{[\rho} A_{\nu]}, \quad F_{\rho\hat{k}} = 2\partial_{[\rho} A_{\hat{k}]}, \quad F_{\hat{k}\hat{l}} = 2\partial_{[\hat{k}} A_{\hat{l}]} + \frac{1}{2} \epsilon_{mkl} A_{\hat{m}}] \quad (\text{III.51}) \end{aligned}$$

yields the diagonal form for the reduced action,

$$\delta S_{B,2}^{(2)}(D7) = -\frac{\mu_7}{2} \int d^8 \xi e^\phi \sqrt{-\tilde{q}} [q^{\mu\mu} q^{\rho\rho} F_{\rho\mu}^2 + q^{\mu\mu} q^{\hat{k}\hat{k}} F_{\mu\hat{k}}^2 + q^{\hat{k}\hat{k}} q^{\rho\rho} F_{\rho\hat{k}}^2 + q^{\hat{l}\hat{l}} q^{\hat{m}\hat{m}} F_{\hat{l}\hat{m}}^2 + q^{\hat{l}\hat{l}} q^{\hat{m}\hat{m}}]. \quad (\text{III.52})$$

We consider the $U(1)$ theory in the gauge $A_{\hat{\rho}} = 0$, substitute the Kaluza-Klein decompositions of the fields $A_{\hat{k}}$ in Eq. (III.51) analogous to those in Eqs. (III.44) and restrict again to the (purely radial) singlet modes $A_{\hat{k}}(\rho)$, hence ignoring angular derivatives. The contributions to the reduced $D7$ -brane action from the symmetric part of the metric tensor $q^{(\alpha\beta)}$ in Eq. (III.43) takes the diagonal quadratic form,

$$\delta S_{B,2}^{(2)}(D7) = -\frac{C}{2} \int d^4 x d\rho [\sqrt{-\tilde{q}} (q^{\mu\nu} q^{\hat{k}\hat{k}} \partial_\mu A_{\hat{k}} \partial_\nu A_{\hat{k}} + q^{\hat{\rho}\hat{\rho}} q^{\hat{k}\hat{k}} \partial_\rho A_{\hat{k}} \partial_\rho A_{\hat{k}} + \frac{1}{2} q^{\hat{l}\hat{l}} q^{\hat{m}\hat{m}} A_{\hat{k}}^2) + \frac{1}{h(\tau)} A_{\hat{k}} \partial_\rho A_{\hat{k}}] \quad (\text{III.53})$$

where the third term inside parentheses is summed over cyclic permutations of the indices $(\hat{k}, \hat{l}, \hat{m})$ running over the basis $[\hat{3}, \hat{1}, \hat{2}]$. The Chern-Simons contribution in the last term above is expressed through integration by parts (and neglect of boundary contributions) by the effective mass term, $-(C/4) \partial_\rho (h^{-1}(\tau)) A_{\hat{k}}^2$. After integration by parts and substitution of the expression for the inverse effective metric tensor, one finds the action for singlet fields,

$$\delta S_{B,2}^{(2)}(D7) = -\frac{C}{2} \int d^4 x d\rho \mathcal{G}_k [-\frac{\sqrt{-\tilde{q}}}{\mathcal{G}_k (\eta^{\hat{k}})^2 R_S} A_{\hat{k}} \tilde{\nabla}_4^2 A_{\hat{k}} - \frac{1}{\mathcal{G}_k} A_{\hat{k}} \partial_\rho \mathcal{G}_k \partial_\rho A_{\hat{k}}]$$

$$+ \frac{\sqrt{-\tilde{q}}}{2h(\tau)\mathcal{G}_k R_S^2} \frac{1}{(\eta^{\hat{l}}\eta^{\hat{m}})^2} A_{\hat{k}}^2 + \frac{1}{2\mathcal{G}_k h(\tau)} \partial_\rho(A_{\hat{k}}^2)], [\mathcal{G}_k = \frac{\sqrt{-\tilde{q}}}{h(\tau)(\eta^{\hat{\rho}}\eta^{\hat{k}})^2 R_S^2}, R_S = \frac{\sqrt{-\tilde{q}}}{\sqrt{-\tilde{\gamma}}}. \quad (\text{III.54})$$

The wave functions of singlet modes $a_{\hat{k},m}(\rho)$ satisfy the wave equations

$$\begin{aligned} 0 &= [\omega_m^2 \frac{\sqrt{-\tilde{q}}}{\mathcal{G}_k |\eta^{\hat{k}}|^2 R_S} + \frac{1}{\mathcal{G}_k} \partial_\rho \mathcal{G}_k \partial_\rho - \frac{\sqrt{-\tilde{q}}}{2h(\tau)\mathcal{G}_k R_S^2 (\eta^{\hat{l}}\eta^{\hat{m}})^2} + \frac{\partial_\rho h^{-1}(\tau)}{2\mathcal{G}_k} \partial_\rho] a_{\hat{k},m} \\ &= [\omega_m^2 Q R_S + \frac{1}{\mathcal{G}_k} \partial_\rho \mathcal{G}_k \partial_\rho - \frac{1}{2} (\frac{\eta^{\hat{\rho}}\eta^{\hat{k}}}{\eta^{\hat{l}}\eta^{\hat{m}}})^2 + \frac{\partial_\rho h^{-1}(\tau)}{2\mathcal{G}_k} \partial_\rho] a_{\hat{k},m}, [Q = h(\tau)(\eta^{\hat{\rho}})^2]. \end{aligned} \quad (\text{III.55})$$

where the summation over cyclic indices $\hat{k}, \hat{l}, \hat{m}$ in the third term inside the brackets yield $-(\eta^{\hat{\rho}}\eta^{\hat{k}}/(\eta^{\hat{l}}\eta^{\hat{m}}))^2$, for the modes $a_{\hat{3},m}$. The rescaled wave functions $\tilde{a}_{\hat{k},m} = a_{\hat{k},m} \mathcal{G}_k^{1/2}$ satisfy the Schrödinger type wave equations,

$$(\partial_\rho^2 - V_{eff}^{\hat{k}}) \tilde{a}_{\hat{k},m} = 0, [V_{eff}^{\hat{k}} = -Q\omega_m^2 R_S + \mathcal{G}_{k,1} + \frac{1}{2} (\frac{\eta^{\hat{l}}\eta^{\hat{m}}}{\eta^{\hat{\rho}}\eta^{\hat{k}}})^2 - \frac{\partial_\rho(h^{-1}(\tau))}{2\mathcal{G}_k}, \mathcal{G}_{k,1} = \frac{(\mathcal{G}_k^{1/2})''}{\mathcal{G}_k^{1/2}}]. \quad (\text{III.56})$$

The singularity near $\rho = 0$ in $V_{eff}(\rho)$ from $\mathcal{G}_{k,1}$ is again removed by the changes of variable and wave functions, $\rho = e^y$, $a_{\hat{k},m} = (\rho/\mathcal{G}_k)^{1/2} \hat{a}_{\hat{k},m}$. The resulting Schrödinger wave equation with the regular effective potential $W_{eff}(y) = \frac{1}{4} + e^{2y} V_{eff}$ and the quantization rule for the modes potential \hat{V}_{eff} and masses ω_m^2 ,

$$\begin{aligned} (\partial_y^2 - W_{eff}^{\hat{k}}) \hat{a}_{\hat{k},m} &= 0, [\hat{V}_{eff}^{\hat{k}} \equiv \frac{W_{eff}^{\hat{k}}}{\rho^2} = -Q\omega_m^2 R_S + \hat{\mathcal{G}}_{k,1} + \frac{1}{2} (\frac{\eta^{\hat{l}}\eta^{\hat{m}}}{\eta^{\hat{\rho}}\eta^{\hat{k}}})^2 - \frac{\partial_\rho(h^{-1}(\tau))}{\mathcal{G}_k}, \hat{\mathcal{G}}_{k,1} = \mathcal{G}_{k,1} + \frac{1}{4\rho^2}, \\ (n - 1/2 + \delta_W/4)\pi &= \int_{-\infty}^{y_0} dy (-W_{eff}^{\hat{k}}(y))^{1/2} = \int_0^{\rho_0} d\rho (-\hat{V}_{eff}^{\hat{k}}(\rho))^{1/2}, \hat{V}_{eff}^{\hat{k}} \equiv \frac{W_{eff}^{\hat{k}}}{\rho^2} = \frac{1}{4\rho^2} + V_{eff}^{\hat{k}}] \end{aligned} \quad (\text{III.57})$$

are derived in a similar fashion as Eq. (III.29).

We turn finally to the contributions from the antisymmetric part of the effective metric of quadratic order in $q^{[\alpha\beta]} = h^{-1/2}(\tau) \tilde{q}^{[\alpha\beta]}$ and linear order $q^{[\alpha\beta]} q^{(\delta\delta)}$, given in the notations of Eq. (A.23) by $\tilde{q}^{[\alpha\beta]} = (B^{-1})_{\alpha\beta}/R_A$, $\tilde{q}^{(\alpha\beta)} = (\tilde{\gamma})^{\alpha\beta}/R_S$. Retaining for simplicity the leading contributions at large μ which are included in the metric components, $q^{[\rho\hat{h}_2]}$, $q^{[\hat{h}_1\hat{h}_3]}$, adds the following part to the reduced action,

$$\begin{aligned} \delta S_{B,3}^{(2)}(D7) &= -\mu_7 \int d^8 \xi \sqrt{-\tilde{q}} e^\phi q^{[\rho\hat{h}_2]} q^{[\hat{h}_3\hat{h}_1]} (F_{\hat{h}_2\rho} F_{\hat{h}_1\hat{h}_3} + F_{\hat{h}_3\rho} F_{\hat{h}_2\hat{h}_1} + F_{\hat{h}_1\rho} F_{\hat{h}_3\hat{h}_2}), \\ &\simeq -\frac{C}{2} \int d^4 x \int d\rho \sqrt{-\tilde{q}} q^{[\rho\hat{h}_2]} q^{[\hat{h}_3\hat{h}_1]} A_{\hat{k}} \partial_\rho A_{\hat{k}}, \end{aligned} \quad (\text{III.58})$$

where the second entry is obtained by restricting to the singlet modes.

D. Properties of meson modes

1. Parameter space

The data needed to evaluate the properties of meson modes consists of three sets of inputs: (1) the string theory and internal manifold volume parameters, g_s , $\alpha' = 1/m_s^2$ and $V_W = (2\pi L_W)^{1/6}$, (2) the Klebanov-Strassler throat parameters ϵ , μ , associated to the confinement scale and the quarks bare masses of the dual gauge theory, $\Lambda_{ir} \sim \epsilon^{2/3}/\alpha'$ and $m_Q \sim \mu^{2/3}/\alpha'$ and (3) the throat ultraviolet cutoff τ_{uv} . To ease the contact with phenomenology, we trade m_s for the Planck mass M_\star and ϵ for the mass hierarchy warping factor $w = m_{eff}/M_\star = w_s m_s/M_\star$, using the relations

$$m_s = (\frac{\pi M_\star^2}{L_W^6})^{1/8} = M_\star \sqrt{\frac{\pi}{\mathcal{V}}}, \quad m_s^{3/2} \epsilon = (\frac{2^{1/3} a_0^{1/2} g_s M}{\pi})^{3/4} (w \mathcal{V}^{1/3})^{3/2} = (2^{1/3} a_0^{1/2} g_s M)^{3/4} (\frac{w_s}{\mathcal{V}^{1/6}})^{3/2}, \quad (\text{III.59})$$

where the dependence of ϵ on the compactification manifold X_6 in string units, $\mathcal{V} \equiv e^{6u} = V_W/\hat{l}_s^6 = (L_W/l_s)^6$, results from our definition for the warping factor [49], $w_s = e^u h^{-1/4}(0)$. (Note that the power index change from

$\mathcal{V}^{-1/4} \rightarrow \mathcal{V}^{1/2}$ upon going from $w_s \rightarrow w$.) The dimensionless mesons masses $\hat{\omega}_m$, depending on the parameters ratio L ,

$$\hat{\omega}_m = \epsilon^{-2/3}(g_s M \alpha') \omega_m = \mu^{-2/3}(g_s M \alpha') L^{2/3} \omega_m = (g_s M)^{1/2} \mathcal{V}^{1/6} \omega_m / (2^{1/6} a_0^{1/4} M_* w), \quad [L \equiv \mu/\epsilon \simeq (\frac{m_Q}{\Lambda_{ir}})] \quad (\text{III.60})$$

are defined with the same rescaling as for glueball masses \hat{E}_m .

The complex fields χ and $A_1 + iA_2$ belong to chiral supermultiplets of $\mathcal{N} = 1$ supersymmetry and the real fields A_μ and A_3 to vector supermultiplets. The components in each supermultiplet have equal masses but not necessarily same radial wave functions [41]. The gravity-gauge holographic correspondence between bulk and boundary theories in AdS_d and M_{d-1} spacetimes relates the mass ω of normalizable modes to the scaling dimensions Δ of operators of same spin s and quantum numbers) by $\omega^2 = (\Delta - s)(\Delta - d + s)$. For bulk spacetimes asymptotic to $AdS_d \times X_{10-d}$, one can use the asymptotic expansions near the boundary for the non-normalizable (NN) and normalizable (N) classical fields of fixed mass,

$$\delta\Phi_{\omega,s}(x,r) \sim r^s (c_{NN} r^{\Delta-d} + c_N r^{-\Delta}), \quad [(\frac{2}{\epsilon^2})^{1/3} \hat{r} \simeq e^{\tau/3} \simeq L^{3/2} e^{\rho/3}] \quad (\text{III.61})$$

to determine the scaling dimensions Δ from the radial profiles and hence identify the structure of the dual gauge theory composite operators in the quark superfields. The following table shows the correspondence between the superspace $\theta, \bar{\theta}$ components of vector and chiral bilinear quark operators and the asymptotic radial wave functions $\Phi_{\omega,s}(x,r)$ of the modes $A_3^{(m)}$ and $A_\mu^{(m)}$ part of the same vector supermultiplet and the modes $A_{1,2}^{(m)}$ and $\chi^{(m)}$ part of the same chiral supermultiplet.

Operator O	$O_{\theta\bar{\theta}}^V = q^\dagger \partial_\mu q - \bar{q}^\dagger \partial_\mu \bar{q}$	$O_{\theta\bar{\theta}0}^V = q^\dagger q - \bar{q}^\dagger \bar{q}$	$(\bar{Q}Q)_{\theta^0}$	$(\bar{Q}Q)_{\theta^2}$
Δ, s	3, 1	2, 0	3/2, 0	5/2, 0
AdS Field	A_μ	A_3	(A_1, A_2)	χ
$\lim_{r \rightarrow \infty} \delta\Phi_{\Delta,s}(x,r)$	$c_{NN} r^0 + c_N r^{-2}$	$c_{NN} r^{-2} + c_N r^{-2} \ln r$	$c_{NN} r^{-5/2} + c_N r^{-3/2}$	$c_{NN} r^{-3/2} + c_N r^{-5/2}$

The independent (non-normalizable and normalizable) solutions for the classical bulk fields perturbations $\delta\Phi_{\omega,s}(x,r) = (r^{-3/2} \chi, A_a, A_\mu)(x,r)$ have asymptotic series expansions at large radial distances $r^{3/2}/\mu \gg 1$ whose leading terms are assigned the constant coefficients c_{NN} , c_N . The correspondence to the gauge field theory bare action associates c_{NN} to the source perturbation, $\delta L = c_{NN} O_{\Delta,s}(x)$, and c_N to the perturbed operator VEV, $c_N \sim \langle O_{\Delta,s} \rangle$. (Note that the radial variable rescaling factor for χ is necessary to transform it to a canonically normalized field and that the components c_{NN} and c_N are leading and subleading for χ , A_3 , A_μ with the reverse holding for $A_{1,2}$.) The fields $A_3^{(m)}$ and $A_\mu^{(m)}$ are part of a vector supermultiplet whose dual counterpart is the flavor $U(N_f)$ current superfield operator, $O^V = Q^\dagger e^V Q - \bar{Q}^\dagger e^{-V} \bar{Q}$, $[Q = q + \theta\psi_q + \theta^2 F_q]$. Since current conservation entails non-renormalization (or vanishing anomalous dimension), one is led to assign the unperturbed scaling dimension, $\Delta(O_{\theta\bar{\theta}0}^V) = 2$. The chiral operators are renormalized with their dimensions determined from the superconformal symmetry by the additive rule, $\Delta(Q^n)_{\theta^0} = 3n/4$. The fields $A_{1,2}^{(m)}$ and $\chi^{(m)}$ thus correspond to the superspace components of the chiral superfield operator $\bar{Q}Q$, $[\Delta(\bar{Q}Q)_{\theta^0} = 3/2]$.

In order to gather existing information on the supergravity parameter μ , we consider first the possibility [36] that the mesons in the flavoured Klebanov-Strassler background correspond to the $q - \bar{q}$ quarkonia of QCD. To pursue with this comparison one must specify the matrix μ in the quarks flavour space of dimension N_f . We specialize to the simplified case of an $U(1)^{N_f}$ group for which $L = \mu/\epsilon$ is a diagonal matrix with entries related to the bare quark masses and the dynamical scale as, $L_i \equiv \mu_i/\epsilon = (m_{Q_i}/\Lambda_{ir,N_f})^{3/2} \simeq (m_Q/\Lambda_{ir})^{3/2}$, assuming equality of the dynamical scales for the pure and flavoured gauge theories. The extension to a non-Abelian flavour symmetry group involves tools whose implementation exceeds the scope of the present study [74, 75]. The lattice simulations of 3-flavour QCD using chiral perturbation theory assign the following values to the up, down and strange quark masses, $\bar{m}_q = (m_u + m_d)/2 = (3.39 \pm 0.04) \text{ MeV}$, $m_s = (92.09 \pm 0.7) \text{ MeV}$ for the (renormalization scheme dependent) scale $\Lambda = 200 \text{ MeV}$. The experimental data for heavy flavour quarkonia set the charm and bottom quarks masses at $m_c = (1.280 \pm 0.025) \text{ GeV}$, $m_b = (4.18 \pm 0.03) \text{ GeV}$. The resulting values for L_i for the two choices $\Lambda = 200 - 770 \text{ MeV}$, utilized in lattice simulations and holographic quark models, are displayed in the following table. Note that the estimates for L_i in the two line entries differ by a constant rescaling factor.

Λ (MeV)	$L_{u,d}$	L_s	L_c	L_b
200	2.21×10^{-3}	0.312	16.2	95.5
770	3.4×10^{-4}	0.0413	2.14	12.6

We postpone a further discussion of this application to Subsection (III D 2) below and proceed now to the cosmological type constraints on the parameter μ . The analyses [69, 70] of the inflation model of KKLT [68] are concerned with the slow-roll of a $D3$ -brane in Klebanov-Strassler throat attracted to a $\bar{D}3$ -brane located near the warped deformed conifold tip. The potential energy arises from three sources: (1) The superpotentials $W_{flux}(S)$ and $W_{np}(\rho)$ from the $\int F_3 \sim M$, $\int H_3 \sim -K$ fluxes [50, 77, 78] and the non-perturbative Euclidean $\mathcal{E}3$ -branes ($n_f = 1$) or gaugino condensation on n_f $D7$ -branes, necessary to stabilize the axio-dilaton and deformed conifold complex structure and Kahler volume moduli τ , S , ρ . (2) The $D3 - \bar{D}3$ -branes attractive Coulomb potential V_{Coul} . (3) The $N_{\bar{D}3}$ $\bar{D}3$ -branes potential energy $V(\bar{D}3) = 2e^{4A(\tau)}\tau_{D3}$ necessary to produce a de Sitter metastable vacuum with spontaneously broken supersymmetry. We quote for definiteness schematic expressions for these three contributions

$$\begin{aligned} W_{flux}(S) &= \hat{t}_s^2 M_*^8 S \left(\frac{M}{2\pi i} \left(\ln \frac{\Lambda_{uv}^3}{S} + 1 \right) + i \frac{K}{g_s} \right), \quad W_{np}(\rho) = A_0 f^{1/n_F}(w_a) e^{-a\rho}, \\ V_{Coul} &= 2N\mu_3 w^4 \left(1 - \frac{4}{9N} \left(\frac{w\mathcal{R}}{r} \right)^4 \right), \quad V(\bar{D}3) \simeq \frac{S^{4/3}}{(g_s M)^2}, \end{aligned} \quad (\text{III.62})$$

but refrain from quoting the Kahler potentials for the moduli superfields [76] which are significant inputs in this problem [69, 70]. The $\bar{D}3$ -branes play a critical rôle regarding the supersymmetry breaking mechanism and the adverse possibility that their back-reaction on the metric and 3-form fluxes produces non-physical singularities. Favourable answers to both issues were reported in the extensive literature on this subject. Firstly, it appears that the de Sitter uplifting can be described in terms a linear realizations of a (global or local) supersymmetry coupled to a nilpotent superfield [72]. Secondly, it seems that a proper account of the constraints from warping and off-shell effects on the scalar potential of the modulus S can remove singularities [73].

There are two possible (angular and radial) inflation scenarios depending on whether the $D3$ -brane moves along an angle direction of the S^3 at the conifold tip or along the radial τ direction. Both were analyzed for n_f $D7$ -branes embedded in Kuperstein 4-cycle $f(w) = w_4/\mu - 1$ and in Ouyang 4-cycle [33], $f(w) = z_4/\mu - 1 = (-w_3 + iw_4)/\mu - 1$. For angular inflation, the condition that the $\bar{D}3$ -brane mass term from compactification effects [79], $m_{comp}^2 \simeq w^{3.28}/(g_s M \alpha')$, is dominated by the mass term from moduli stabilization m_{infl}^2 , yields the bound on μ [80]

$$m_{infl}^2 \simeq w^2 \epsilon / (\mu n_f g_s M \alpha') \gg m_{comp}^2 \implies L n_f \equiv \mu n_f / \epsilon \ll w^{-1.28}. \quad (\text{III.63})$$

The alternative analysis using the large volume limit $\mathfrak{R}(\rho) \gg 1$ [81] finds a comparable bound for both Ouyang and Kuperstein embeddings, $n_f L \equiv n_f \mu / \epsilon \ll (g_s M \alpha' / \epsilon^{4/3})^{2/3} \approx w_s^{-4/3}$ (setting for convenience $w^{-0.32} \rightarrow w^{-1/3}$).

The status of the radial inflation scenario is inconclusive due to insufficient cancellations in the radial scalar potential $V(D3 - \bar{D}3)$ to guarantee slow-roll. An independent information on μ can still be inferred from the annihilation of $\bar{D}3$ -branes with 3-fluxes [82] in the adverse case where this process occurs before the $\bar{D}3$ -branes tunneling to a metastable supersymmetry breaking vacuum. The condition that moduli stabilization dominates over brane-flux contributions sets the upper bound [83], $\ln(\mu/\sqrt{\alpha'}) \ll 2\mathcal{V}^{1/3}/(3M)(\epsilon^{2/3}/\sqrt{\alpha'})$.

Useful constraints on μ are provided by the holographic gauge mediation model [42] in which the deformed non-supersymmetric Klebanov-Strassler background (in the large radius limit) is used as a hidden sector with a spontaneously broken supersymmetry set by the scale parameter \mathcal{S} [84]. The observable sector is the grand unified $SU(5)$ gauge theory on a n_f $D7$ -brane stack ($n_f = 5$) localized near the ultraviolet boundary of the throat. The massive gauginos of $D7$ -branes in the throat with broken supersymmetry mediate soft contributions to other brane modes via their coupling with pairs of messenger meson modes $\chi^{(m)} \sim \bar{Q} O_m Q$. The bound on the gauginos mass for low energy supersymmetry breaking, where $\alpha(\Lambda_\mu)$ is the n_f $D7$ -brane running gauge coupling constant at the scale $\Lambda_\mu \simeq \mu^{2/3}/\alpha'$, yields the condition [42]

$$m_\lambda \simeq \frac{\alpha(\Lambda_\mu) n_f \mathcal{S} \alpha'^3}{4\pi \mu^2} \frac{1}{(4\pi g_s N)^{1/2}} \simeq 10^{+1} \times \frac{\mathcal{S} \alpha'^3}{\mu^2} \times 10^{-2} > O(100) \text{ GeV} \implies \frac{\mathcal{S} \alpha'^3}{\mu^2} > O(1) \text{ TeV}, \quad (\text{III.64})$$

which translates in the case of an intermediate scale breaking of supersymmetry, $\mathcal{S}^{1/4} \sim 10^9 \text{ GeV}$, into the conditions on μ and L ,

$$\mu^{2/3} \leq \frac{10^{11} \text{ GeV}}{m_s^2} \implies L = \frac{\mu}{\epsilon} \leq 10^{-21/2} \frac{\mathcal{V}}{(w_s(\pi g_s M)^{1/2})^{3/2}} \left(\frac{\mathcal{S}}{10^{36} \text{ GeV}^4} \right)^{3/2}. \quad (\text{III.65})$$

The scale evolution towards the ultraviolet of the n_f $D7$ -brane running gauge coupling in this model [42] provides an additional constraint on μ . As the floating energy scale exceeds the mesons mass, $\Lambda > \Lambda_\mu \sim \mu^{2/3}/\alpha'$, $\alpha(\Lambda)$ picks up positive logarithmic contributions from 1-loops of virtual pairs of messengers that could produce a Landau pole (LP)

$\alpha(\Lambda) \rightarrow \infty$ invalidating the model. The flow of $x(\Lambda) = 2\pi/\alpha(\Lambda)$ towards the ultraviolet is expected to proceed through pairs of self-similar Seiberg dualities of the Klebanow-Witten gauge theory, $SU(N) \rightarrow SU(N+M) \rightarrow SU(N+2M)$ in steps $\delta t = \delta \ln Q = 2\pi/(3g_s M)\delta k$, $[t = \ln Q]$ with the $D7$ -brane gauge theory beta function $\beta_k(x) = -3kM/2$, $[k \in Z_+]$. The integrated increase of the $SU(N)$ gauge group rank from $N_\mu = k_\mu M \rightarrow N_{uv} = k_{uv}M$ induces the growth of the $SU(n_f)$ gauge theory coupling constant,

$$\Delta x = 2\pi(\alpha^{-1}(\Lambda_{uv}) - \alpha^{-1}(\Lambda_\mu)) = -\sum_{k_\mu}^{k_{uv}} \delta t \beta_k \simeq -\frac{\pi}{2g_s}(k_{uv}^2 - k_\mu^2) \implies (\alpha^{-1}(\Lambda_\mu) - \alpha^{-1}(\Lambda_{uv})) \simeq \frac{1}{4g_s}(k_{uv}^2 - k_\mu^2) \quad (\text{E.66})$$

For an initial coupling constant at the GUT value, $\alpha(\Lambda_\mu) \simeq 1/25$, the Landau pôle $\alpha(\Lambda_{LP}) \simeq \infty$ occurs at $1/\alpha(\Lambda_\mu) \simeq (k_{LP}^2 - k_\mu^2)/(4g_s)$, corresponding (for $g_s \simeq 1/4$) to $k_{LP} \leq 5$. The resulting upper bound on the ratio of mass scales (for $8\pi/3M \sim 1$),

$$\ln \frac{\Lambda_{LP}}{\Lambda_\mu} = \Delta t = t_{LP} - t_\mu = \frac{2\pi}{3g_s M}(k_{LP} - k_\mu) = \frac{8\pi\alpha^{-1}(\Lambda_\mu)}{3M(k_{LP} + k_\mu)} \simeq \frac{25}{k_{LP}} \geq 5 \implies \Lambda_\mu/\Lambda_{LP} \leq e^{-5} \simeq 10^{-2.2} \quad (\text{E.67})$$

implies a problematic narrow window of scales from $\Lambda_\mu \rightarrow \Lambda_{uv}$. For the values $w_s \sim 10^{-4}$, $\Lambda_\mu = \mu^{2/3}m_s^2$, $\Lambda_{LP} = m_s$: $\Lambda_\mu/\Lambda_{LP} < 10^{-2} \implies L < 10^{-3}w_s^{-3/2}/\mathcal{V}^{1/4}$. As discussed in [42], one can widen this bound via the conifold orbifolding. For the orbifold \mathcal{C}_6/Z_Q , the flux parameter gets reduced, $M \rightarrow M/Q$, hence $\beta_k \rightarrow \beta_k/Q$, but the scale evolution $\delta t/\delta k = 2\pi/(3g_s M)$ is unaffected. The modified bound for $Q = 5$ reads, $1/\alpha(\Lambda_\mu) \simeq (k_{LP}^2 - k_\mu^2)/(4g_s Q) \implies \Lambda_\mu/\Lambda_{LP} \leq e^{-5Q} \simeq 10^{-10}$.

We conclude from the above comparisons that the conditions on $L = \mu/\epsilon$ favour values $L = O(1)$ in hadronic physics applications and larger values $L > O(10^2)$ in particle physics applications which feature a sensitive dependence on the string theory and compactification parameters α' , ϵ , \mathcal{V} .

2. Mass spectra of scalar and vector mesons

Our applications were all developed within the semi-classical approach using ‘Mathematica’ numerical tools. The predictions for the mass spectra of the radial excitations of scalar and vector mesons are listed in Table I and displayed as a function of real $L = \sqrt{L^2} \in [0, 100]$ in Fig. 1. It is clearly seen that scalar modes are significantly lighter than vector modes. The masses are nearly constant for $L \in (0, 1)$ with a weak inflection point at $L = 1$ and a slow growth until $L = O(10)$ beyond which they follow the asymptotic power law regimes $\hat{\omega}_1 \simeq (0.8 - 0.6) L^{2/3}$ for scalars and $\hat{\omega}_1 \simeq (1. - 0.7) L^{2/3}$ for vectors. The masses grow with the radial quantum number as $\hat{\omega}_m \propto n$, as expected for low curvature throats and previous results for $\mathcal{N} = 2$ backgrounds [20]. This contrasts with the growth law $\hat{\omega}_m \propto n^{1/2}$ in strongly coupled gauge theories [85] and the mixed law $\hat{\omega}_n \sim (a + bn)$ in analyses for multidimensional fields spaces [62, 86, 87, 91]. The mass ratios $[\hat{\omega}_2/\hat{\omega}_1, \hat{\omega}_3/\hat{\omega}_1]$ are approximately independent of L , ranging inside $[2, 3]$ for scalars and $[1.7, 2.4]$ for vectors.

The mass gap, defined as the ratio of the mesons mass $\omega_{mes} \simeq \mu^{2/3}m_s^2/(g_s M)$ (at $L \gg 1$) to the open string mass (in flat spacetime) $m_Q \simeq \mu^{2/3}T(F1) \sim \mu^{2/3}/(2\pi\alpha')$, sets typically at $\omega_{mes}/m_Q \sim 1/(g_s M)$. Note this is smaller than the ratio in the undeformed conifold case [35], $M_{mes} \sim m_q/\sqrt{g_s N}$. If one used instead the $F1$ -string tension near the conifold apex, $T(F1) \simeq w_s m_s^2 \mathcal{V}^{-1/3}$, a substantially larger mass gap would result, $\omega_{mes}/m_Q(F1) \simeq (\epsilon^{2/3}m_s)^{-2} \simeq \mathcal{V}^{1/3}/(g_s M w_s^2)$.

For completeness, we compare in the table below our semi-classical predictions for the reduced masses of (ground and first radial excitation) scalar and vector mesons $\hat{\omega}_m(\chi)$, $\hat{\omega}_m(A_\mu)$, $[m = 1, 2]$ with those of other authors which made use of the shooting technique. Four cases **A**, **B**, **C**, **D** are considered in the successive columns. The results refer to $\omega_m/m_{gb} = 2^{1/3}\hat{\omega}_m$ in cases **A**, **B** and to $\lambda_m = \omega_m L_{eff}^2/\mu^{2/3} = (81 \ln(r_\mu/r_0)\hat{\omega}_m^2/(8L^{4/3}))^{1/2} \rightarrow 5.13\hat{\omega}_m/L^{2/3}$ in cases **C**, **D**. In all cases but case **B**, we set $B_2^l = 0$. The comparisons reflect on the numerical evaluation scheme used but also on the parameter choices, since the previous studies considered the limit $\mu = 0$ in **A** and the limit $\epsilon = 0$ in **C**, **D**. The results show some disparity but they roughly agree within a factor 2.

	A ($\mu \simeq 0$) $\rightarrow L \simeq 0.1$ [34]	B ($L = 1.5$) [36]	C ($\mu \gg 1$) $\rightarrow L = 50$ [41]	D ($\mu = 10, \epsilon = 0$) $\rightarrow L = 100$ [42]
	$2^{1/3}\hat{\omega}_m(\chi)$ $2^{1/3}\hat{\omega}_m(A_\mu)$	$2^{1/3}\hat{\omega}_m(A_\mu)$	$\lambda_m(\chi)$ $\lambda_m(A_\mu)$	$\lambda_m(A_\mu)$
Shooting	(3.38, 4.92) (4.32, 5.8)	(1.62, 3.04)	(1.89, 4.39) (2.56, 4.97)	(1.89, 3.73)
JWKB	(1.37, 2.70) (1.85, 3.13)	(2.03, 3.39)	(3.08, 6.16) (4.09, 7.11)	(3.83, 6.69)

TABLE I: List of masses $\hat{\omega}_{1,2,3}^2$ of the scalar and vector ground state and two first radially excited mesons at a set of real positive values of $L = \mu/\epsilon \in [0, 100]$.

L	0.1	0.2	0.4	0.7	0.8	0.9	0.95	1.5	3	5	10	20	50	100	
$\chi^{(m)}$	$\hat{\omega}_1^2$	1.187	1.181	1.161	1.124	1.131	1.273	1.367	1.620	3.221	5.408	11.19	23.84	67.07	149.6
	$\hat{\omega}_2^2$	4.610	4.595	4.535	4.367	4.290	4.241	4.350	6.307	12.70	21.40	44.51	95.17	268.7	600.2
	$\hat{\omega}_3^2$	10.25	10.22	10.09	9.72	9.516	9.26	9.170	14.03	28.31	47.73	99.41	212.7	600.76	1366.3
$A_\mu^{(m)}$	$\hat{\omega}_1^2$	2.161	2.146	2.206	2.316	2.519	2.648	2.673	3.276	5.938	9.75	19.90	42.01	117.2	260.5
	$\hat{\omega}_2^2$	6.19	6.13	6.13	6.10	6.12	6.37	6.68	8.807	17.14	28.64	59.18	125.9	354.2	790.4
	$\hat{\omega}_3^2$	12.39	12.27	12.20	11.99	11.86	11.80	12.20	17.40	34.4	57.87	120.0	256.17	722.3	1614.0

The results in Fig. 1 and Table II show that the classical field B_2^{cl} has a negligible effect at $L < 1$ but contributes significantly at $L > 5$, where masses are reduced by a factor $\sqrt{2}$ while mass splittings between radial excitations are enhanced. We have also evaluated the mass spectra in the region of the parameter space $L^2 < 0$. The results in Table III show minor changes relative to those for $L^2 > 0$ (in Table II). The variation in $|L| < 1$ is weaker but the growth $\hat{\omega}_m \sim |L|^{2/3}$ is stronger for $|L| \gg 1$ and more effective for χ^m relative to A_μ^m .

Before addressing the holography correspondence [36] to the QCD (Quantum Chromodynamics) hadrons, it is useful to recall that the space inversion and charge conjugation parities assigned to meson modes are linked to the choice of boundary conditions near the origin $\rho = 0$, $\Omega = 0$. The transformations under P , C of the 4-d fields




FIG. 1: The scalar $\chi^{(m)}$ (left) and the vector $A_\mu^{(m)}$ (right) mesons masses $\hat{\omega}_m$ are plotted as a function of the parameter $L = \mu/\epsilon \in [0 \rightarrow 100]$. The curves from bottom to top refer to the ground state and first two radially excited modes $n = 1, 2, 3$. The upper panels (a), (b) refer to the case $B_2^{cl} = 0$ (ignoring the NSNS field) and those in the lower panels (c), (d) to the case $B_2^{cl} \neq 0$ for the full background solution. We set $g_s = 1$ in numerical calculations.

TABLE II: *Scalar and vector mesons masses of the ground state and first radially excited mesons including the contributions fom NSNS B_2^{cl} solution evaluated for a list of real values of $L^2 > 0$.*

L	0.1	0.2	0.4	0.7	0.8	0.9	0.95	1.5	3	5	10	20	50	100
$\chi^{(m)} \hat{\omega}_1^2$	1.20	1.19	1.14	1.41	1.77	2.60	3.05	2.26	3.31	4.37	7.76	13.4	31.1	65.4
$\hat{\omega}_2^2$	4.75	3.18	4.95	4.95	5.70	6.53	8.57	7.56	9.83	14.6	30.3	51.7	106.6	245.5
$A_\mu^{(m)} \hat{\omega}_1^2$	2.15	2.14	2.10	2.14	2.32	2.44	2.47	2.62	3.84	5.48	9.45	17.18	40.41	80.09
$\hat{\omega}_2^2$	6.17	6.13	5.88	5.71	5.69	5.93	6.245	7.278	11.68	17.21	30.62	57.12	137.8	277.5

TABLE III: *Scalar and vector mesons masses of ground state and first radially excited modes with NSNS B_2^{cl} solution for a list of real values of the ratio parameter $L^2 = \mu^2/\epsilon^2 < 0$.*

$-iL$	0.1	0.2	0.4	0.7	0.8	0.9	0.95	1.5	3	5	10	20	50	100
$\chi^{(m)} \hat{\omega}_1^2$	1.21	1.22	1.25	1.35	1.39	1.40	1.42	2.83	4.51	6.58	11.5	21.2	50.3	100.2
$\hat{\omega}_2^2$	4.77	4.43	5.36	5.86	5.84	5.85	5.96	11.75	19.17	28.5	51.2	96.2	233.3	470.3
$A_\mu^{(m)} \hat{\omega}_1^2$	2.16	2.17	2.26	2.35	2.40	2.46	2.49	2.86	4.00	5.58	9.50	17.2	40.4	80.0
$\hat{\omega}_2^2$	6.20	6.25	6.54	6.83	7.01	7.20	7.30	8.51	12.3	17.5	30.8	57.2	137.8	277.5

$f_m(\rho)\chi^{(m)}(x)$ and $a_m(\rho)a_\mu^{(m)}(x)$ follow from the induced action of these symmetry operators on the (transversal) spatial and (longitudinal) gauge coordinates $\chi \sim X^{8+i9}$, $A_\mu \sim X_\mu$ in the $D7$ -brane theory, $P : [\chi \rightarrow -\chi, A_\mu \rightarrow -A_\mu]$ and $C : [\chi \sim \chi, A_\mu \rightarrow -A_\mu^T]$, as discussed in [17, 38, 62, 92]. One can choose, for instance, radial and angular wave functions $f_m(\rho, \Omega)$, $a_m(\rho, \Omega)$ that are even or odd under the reflections, $\rho \rightarrow -\rho$, $\Omega \rightarrow -\Omega$. The angular dependence (on Ω) can be ignored for singlet modes. For radial wave functions $f_m \sim \hat{f}_m$, $a_m \sim \hat{a}_m$ even under $\rho \rightarrow -\rho$, hence obeying the boundary conditions at $\rho = 0$: $f_m(0) \neq 0$, $\partial_\rho f_m(0) = 0$ and $a_m(0) \neq 0$, $\partial_\rho a_m(0) = 0$, the assigned J^{PC} quantum numbers are 0^{-+} for scalar mesons and 1^{--} for vector mesons.

It is safe to restrict our study to S -wave quarkonia and their radial excitations, ignoring the 0^{++} scalar hadrons which exhibit large decay widths attributed to mixing with glueballs. We focus on the natural parity vector and pseudoscalar hadrons of spacetime quantum numbers $J^{PC} = 1^{--}$, 0^{-+} , and their radial excitations and restrict to the electric charge neutral members of $SU(3)_{fl}$ multiplets. We also discard the $\pi^0(140)$ -meson, since spontaneously broken chiral symmetry is definitely absent in the present model. We thus consider the masses (in MeV units) of (1) the ground and radially excited vector mesons of u, d flavours for ρ : [770, 1450, 2100] and ω : [782, 1420, 1650] and of the mixed (u, d, s) mixed flavours for ϕ : [1020, 1680, 2170] and (2) the excited pseudoscalar mesons of (u, d) flavours, π : [1300, 1800] and of (u, d, s) mixed flavours, η : [958, 1295, 1405, 1475]. We include in our test the heavy flavours hadrons of same quantum numbers.

The comparison of holographic mesons to QCD hadrons reveals two discouraging features given that scalar mesons are lighter than vector mesons and the masses for mesons with light quark flavours change little in the relevant parameter range $L_i = \mu_i/\epsilon < 1$. With the choice $L_i \simeq (m_{q_i}/\Lambda)^{3/2} = [0.1, 0.2, 10, 100]$ for the (q, s, c, b) flavours, fitting the predicted vector meson ground state mass $\omega_1 \equiv \Lambda_{ir}\hat{\omega}_1/(g_s M)$, [$\hat{\omega}_1 = \sqrt{2.16}$] to the ρ^0 -hadron mass determines $\Lambda_{ir}/(g_s M) = 524$ MeV. The resulting mass predictions for the ground state mesons [(770), 767, 2336, 8454] MeV compare poorly to those of the lightest 1^{--} hadrons, $m[\rho, \phi, J/\psi, \Upsilon] = [(770), 1015, 3097, 9460]$. The predicted scalar meson mass $\omega_1(0^{-+}) = 629$ MeV also misses the observed value by a factor 2. We turn next to the ratios of mesons masses where one expects a weaker sensitivity to the parameters. The predicted ratios of the first and second radially

TABLE IV: *Comparison of ratios for the ground state and first or second radially excited modes between mesons and hadrons for $L = [0.1, 0.2, 10, 100]$ assigned to $[(u, d), s, c, b]$ quark flavours. No data is available for the second radial modes of the η_c, η_b hadrons.*

1^{--}	$\frac{\omega_2}{\omega_1}$	ρ	ϕ	J/Ψ	Υ	$\frac{\omega_3}{\omega_1}$	ρ	ϕ	J/Ψ	Υ
Hadrons		1.88	1.65	1.19	1.06		2.73	2.12	1.21	1.08
Mesons		1.69	1.69	1.72	1.74		2.39	2.39	2.45	2.49
0^{-+}	$\frac{\omega_2}{\omega_1}$	π	η	η_c	η_b	$\frac{\omega_3}{\omega_1}$	π	η	η_c	η_b
Hadrons		1.39	1.35	1.22	1.04		—	1.47	—	—
Mesons		1.97	1.97	1.99	2.0		2.94	2.94	2.98	3.02

excited meson modes relative to the ground state mode are compared to those of quarkonia hadrons in Table IV. In spite of the improved agreement, we fail to account for the uniform decrease of hadronic mass ratios upon going from light to heavy flavours.

Two main drawbacks of supersymmetric holography models for hadrons are the presence of fermionic superpartners and the absence of a spontaneous breaking of chiral symmetry $U(N_f)_L \times U(N_f)_R \rightarrow U(N_f)_V$ (excluding light Nambu-Goldstone pseudoscalar mesons). To estimate the uncertainties one could examine from the gauge theory side how soft supersymmetry breaking effects affect the transition to the non-supersymmetric theory. Unfortunately, the contributions from F-term components of spurion superfields at small energy scales are understood moderately for the gauginos and scalars mixing [88, 89] and poorly for the scalars masses. On the other hand, relaxing the supersymmetry constraints on the supergravity side (holomorphic 4-cycles with anti-self dual, primitive gauge fields, $\mathcal{F} = -\star_{S^4} \mathcal{F}$, $\mathcal{F} \wedge J = 0$) [38] entails starting anew from U -shaped embedding of $D7 - \bar{D}7$ -branes in the ultraviolet joining together towards the infrared [17] which also require activating gauge and dilaton-axion fields [5, 37–39, 90]. We here consider the non-supersymmetric deformed Klebanov-Strassler background embedding a smeared distribution of $p(D3 - \bar{D}3)$ -branes near the conifold apex which is dual to the $SU(N - p + M) \times SU(N - p)$ gauge theory in a metastable supersymmetry breaking vacuum [84]. The dilaton, 2-form and energy-momentum tensor fields $O_+ \sim e^{-\phi}$, $O_- \sim e^{-\phi} \int_{S^2} (B_2 + iC_2)$, $\Theta_{\mu\nu}$ acquire finite VEVs in the deformed solution of size set by the warped vacuum energy scale, $\mathcal{S} = \frac{p}{N} (w_s m_s)^4$. The corrections to the vector mesons masses in this background were actually evaluated in [42]

$$\omega_m^2 \rightarrow \omega_m^2 \left(1 - \frac{\mathcal{S} \alpha'^4 r_m}{\mu^{8/3}}\right) = \omega_m^2 \left(1 - \frac{p \mathcal{V}^{2/3} r_m}{N (g_s M)^2 L^{8/3}}\right), \quad (\text{III.68})$$

where the constant coefficients $r_m = \delta\beta_m / \lambda_m^2$ take numerical values of $O(10)$ increasing with the radial excitation. The negative sign correction reduces the masses of light relative to heavy flavour mesons, owing to the factor $\mu^{-8/3}$, while the factor $r_m = O(10)$ reduce the ratios $\omega_2^2 / \omega_1^2 \propto (r_2 - r_1)^{1/2}$ upon going from light to heavy modes. Both features have the potential of removing the disagreements found in the above comparison using Table IV.

3. Interactions of scalar mesons

The tree level couplings of modes are obtained by expanding the $D7$ -brane action in powers of field fluctuations and evaluating coupling constants in terms of overlap integrals of wave functions. It is useful to first have a look on the radial profiles of the effective potentials and wave functions which are displayed in Fig. 2 for scalars and Fig. 3 for vectors. The plots of the potentials $W_{eff}(\rho)$ in panels (a) feature 'well' regions getting deeper and broader with increasing radial quantum number n (or mass) with monotonic growth beyond the classical turning points ρ_0 . The potentials for vector modes are deeper and narrower than those of scalar modes.

The semi-classical wave functions $\hat{f}_m(\rho)$ and $\hat{a}_m(\rho)$ in panels (b) are peaked at the 4-cycle tip and decrease with long tails beyond the turning points. The previously announced boundary conditions near $\rho = 0$ are visible on these plots. One sees the expected oscillatory behaviour at large ρ for radially excited modes with exponentially decreasing tails beyond the turning points. The vector modes have smaller amplitudes. The radial profiles of the effective potentials $\hat{V}_{eff} = W_{eff} / \rho^2$, displayed in panels (c), shows that the L -dependence significantly saturates beyond $L \sim 10$. The smooth dependence of the predicted masses on L is a consequence of the slow variation of the parameters X and τ_{min} and of the effective potentials turning points ρ_0 , as illustrated in the following table.

L	τ_{min}	X	$\rho_0(\chi)$	$\rho_0(A_\mu)$
0. \rightarrow 0.9	0	0.1 \rightarrow 2.0	6.7 \rightarrow 9.6	5.4 \rightarrow 8.2
1.5 \rightarrow 100	1.9 \rightarrow 10	1.3 \rightarrow 1.0	7.2 \rightarrow 6.0	6.1 \rightarrow 4.8

For imaginary parameter values $L = i(0.1 \rightarrow 0.9)$ the variations are $\tau_{min} \simeq (0.2 \rightarrow 1)$, $X \simeq (0.1 \rightarrow 1.7)$. We also note that the predictions depend weakly on the auxiliary ultraviolet cutoff parameter values as long as one chooses $\tau_{uv} < 25$ or $\rho_{uv} < 15$. For example, increasing $\rho_{uv} = 10 \rightarrow 12$ changes the mass at $L = 100$ from $\omega_m^2 = 152 \rightarrow 149$. The τ -profiles are similar to the ρ -profiles except for the successive thresholds at $\tau_{min} = \tau(\rho = 0)$ which grow with increasing L .

The glueball-meson coupling constants λ_Δ^A of dimension $\Delta = 5, 6$ and the mesons self couplings λ_Δ^S of dimension $\Delta = 2, 3, 4$ are described by the Lagrangians

$$L_{EFF} = \lambda_{mnp}^A h^{(m)\mu\nu} \partial_\mu \chi^{(n)} \partial_\nu \bar{\chi}^{(p)} + \lambda_{mnp}^A h^{(m)\mu\nu} \partial_\mu \chi^{(n)} \partial_\nu \bar{\chi}^{(p)} \chi^{(l)}$$




FIG. 2: Plots versus ρ of the effective potentials $W_{eff}(\rho)$, $\hat{V}_{eff}(\rho) = W_{eff}(\rho)/\rho^2$ and wave functions $f_n(\rho)$ (in units of $\epsilon = 1$, $g_s M \alpha' = 1$) of the scalar meson mode fields $\chi^{(n)}(x)$ at fixed values of $L = \mu/\epsilon$ in the case $B_2^{cl} = 0$.

$$+\lambda_{mn}^S \chi^{(m)} \bar{\chi}^{(n)} + \lambda_{mnp}^S \chi^{(m)} \bar{\chi}^{(n)} \chi^{(p)} + \lambda_{mnpq}^S \chi^{(m)} \bar{\chi}^{(n)} \chi^{(p)} \bar{\chi}^{(q)}, \quad (\text{III.69})$$

using Eqs. (III.36), (III.39) and (III.50). We associate to λ_{Δ}^A , λ_{Δ}^S the reduced dimensionless coupling constants $\hat{\lambda}_{\Delta}^A = \lambda_{\Delta}^A/\mathcal{R}_{\Delta}^A$, $\hat{\lambda}_{\Delta}^S = \lambda_{\Delta}^S/\mathcal{R}_{\Delta}^S$ by factoring out the dependence on parameters in the coefficients \mathcal{R}_{Δ}^A , \mathcal{R}_{Δ}^S (which differ from one coupling type to the other and may include powers of L). Our numerical predictions for the reduced coupling constants $\hat{\lambda}_{\Delta}^{A,S}$ are listed in Table V for 3 values of $L = [0.5, 1.5, 3]$ where the expressions for the coefficients \mathcal{R}_{Δ} are displayed in the second line. We see that the gravitational couplings are strongly suppressed for $L > 3$, owing to the negligible overlap due to the widened distance between glueballs and mesons. The reduced coupling constants are seen to be $O(1)$ with the exception of A_6 . Inside the interval $L \in [0, 3]$ the gravitons-mesons coupling constants decrease by a factor 5, while the derivative mesons couplings A_6 and the non-derivative reduced couplings S_2, S_3, S_4 increase by a factor < 2 . However, upon taking the L -dependence in \mathcal{R} into account, one finds that also $\lambda_{3,4}^S$ decrease slowly with increasing $L > 1$.

The term $S_2 \simeq \lambda_2^S |\chi|^2$ corresponds to a mode independent tree level contribution to the scalar meson masses coming on top to the squared masses ω_m^2 . The mass shifts, $\delta m_{\chi}^2 = \mathcal{R}_2^S \hat{\lambda}_2^S \simeq (M_{\star} w)^2 / (g_s M \mathcal{V}^{1/3})$ are typically smaller than the masses $\omega_m^2 = m_s^4 \epsilon^{4/3} \hat{\omega}_m^2 / (g_s M)^4 \simeq (M_{\star} w)^2 \hat{\omega}_m^2 / (g_s M \mathcal{V}^{1/3})$ and are strongly suppressed at large L since $\hat{\omega}_m^2 \propto L^{4/3}$.

IV. FERMIONIC SECTOR OF D7-BRANES

We examine in this section the fermionic modes on $D7$ -branes wrapped over the Kuperstein 4-cycle Σ_4 of the Klebanov-Strassler background. The towers of 4-d spinor fields from the Kaluza-Klein decomposition of the 10-d bispinor field $\Theta(X)$ on $M_4 \times \mathcal{C}_6$ are examined for the (linearized) wave equations derived in [6]. The Dirac operator in the simplified action in Eq. (II.14) (dropping the contributions from classical form fields except the F_5 -form) is utilized in a truncated version ignoring the twisting effects from the normal bundle and the extrinsic curvature of Σ_4 [58, 96]. Motivated by the similarity of the present model to that of non-compact conic manifolds of G_2 holonomy [10], we also address the possible existence of normalizable massless fermion modes.

TABLE V: Reduced coupling constants $\hat{\lambda}_\Delta = (\hat{\lambda}^{A_5^n}, \hat{\lambda}^{A_6}, \hat{\lambda}^{S_n})$ for the dimension $\Delta = 5$ operators $A_5^{(n)}(h\partial\bar{\chi}\partial\chi)$ in the ground and first two radial modes $n = (0, 1, 2)$, the dimension $\Delta = 6$ operator $A_6(h\chi\partial\bar{\chi}\partial\chi)$ and the dimension $\Delta = 2, 3, 4$ operators $S_2(\chi^2)$, $S_3(\chi^3)$, $S_4(\chi^4)$ in the ground state mode. The ratios of the physical coupling constants to the reduced ones, $\mathcal{R}_\Delta^A = [\lambda_{5,6}^A/\hat{\lambda}_{5,6}^A, \mathcal{R}_\Delta^S = \lambda_{2,3,4}^S/\hat{\lambda}_{2,3,4}^S]$, including the dependence on free parameters, appear in the second line entry. The next lines entries display the reduced coupling constants at the values $L = \mu/\epsilon = (0.5, 1.5, 3)$.

	A_5^0	A_5^1	A_5^2	A_6	S_2	S_3	S_4
$\mathcal{R}_\Delta = \frac{\lambda_\Delta}{\hat{\lambda}_\Delta}$	$\frac{0.787 \cdot 10^4 \rho^{1/4}}{(g_s M^3)^{1/2} M_* w}$			$\frac{0.899 \cdot 10^6 (\rho/g_s^5)^{1/2}}{M^3 L (M_* w)^2}$	$\frac{0.754 (M_* w)^2}{g_s M \rho^{1/2}}$	$\frac{42.9 M_* w}{g_s^3 M^{5/2} L \rho^{1/4}}$	$\frac{612.}{g_s^5 M^4 L^2}$
L= 0.5	0.171	0.181	0.182	28.78	0.954	8.741	2.052
L= 1.5	0.106	0.101	0.112	47.98	0.873	3.029	0.619
L= 3	0.043	0.053	0.045	62.31	3.044	11.88	3.519

A. Dimensional reduction of spinor fields

The embedding of the brane world volume $M_8 \subset M_{10} = M_4 \times \mathcal{C}_6 = M_4 \times \Sigma_4 \times N_2$ involves the consecutive splittings of dimensions $10 \rightarrow 4 + 6$ and $6 \rightarrow 4 + 2$ for the curved and flat coordinates of the M_{10} spacetime,

$$\begin{aligned} X^M &= (X^\mu, X^m) = (X^\mu, X^r, X^u), [M = 0, \dots, 9, \mu = 0, \dots, 3, m = 1, \dots, 6, r = 1, \dots, 4, u = 1, 2] \\ X^a &= (X^\mu, X^m) = (X^\mu, X^a, X^i), [a = 0, \dots, 9, a = 1, \dots, 4, i = 1, 2] \end{aligned} \quad (\text{IV.1})$$

where the coordinates X^r , X^a , [$r, a = 1, \dots, 4$] indices are ordered as $(\rho, \hat{h}_3, \hat{h}_1, \hat{h}_2)$. Similar splittings occur for the curved and flat (constant) Dirac gamma matrices Γ^M and Γ^A acting in the 32 dimensional spinor of the tangent spacetime group $SO(9, 1)$. The Dirac matrices representations adapted to the brane consist of direct products of Dirac matrices acting on the 4 and 8 spinors of the $SO(1, 3) \times SO(6)$ group and on the 4, 4, 2 spinors of the $SO(1, 3) \times SO(4) \times SO(2)$ group,

$$\Gamma^\mu = \gamma^\mu \otimes 1_4 \otimes 1_2, \quad \Gamma^a = \gamma_{(4)} \otimes \tilde{\gamma}^a \otimes 1_2, \quad \Gamma^r = \gamma_{(4)} \otimes \tilde{\gamma}_{(4)} \otimes \hat{\gamma}^r, \quad [\Gamma_{(4)} = i\Gamma^{0123} = \gamma_{(4)} \otimes 1_8,$$




FIG. 3: Plots versus ρ of the effective potentials $W_{eff}(\rho)$, $\hat{V}_{eff}(\rho) = W_{eff}(\rho)/\rho^2$ and wave functions $a_n(\rho)$ (in units of $\epsilon = 1$, $g_s M \alpha' = 1$) of the vector fields modes $A_\mu^{(n)}$ at fixed values of $L = \mu/\epsilon$ in the case $B_2^c = 0$.

$$\Gamma_{(6)} = -i\Gamma^{\underline{456789}} = 1_{(4)} \otimes \tilde{\gamma}_{(6)}, \quad \Gamma_{\Sigma_4} = \Gamma^{\underline{4567}} = 1_{(4)} \otimes \tilde{\gamma}_{(4)}, \quad \Gamma_{N_2} = -i\Gamma^{\underline{89}} = 1_{(4)} \otimes 1_{(4)} \otimes \tau_3, \quad \Gamma_{(8)} = -i\Gamma^{01\dots 6} \quad (\text{IV.2})$$

The 10-d chirality matrix $\Gamma_{(10)}$, $[\Gamma_{(10)}^2 = 1]$ factors into the chirality matrices of the submanifolds, M_4 , \mathcal{C}_6 and Σ_4 , N_2 , $\Gamma_{(10)} = \Gamma_{(4)}\Gamma_{(6)} = \Gamma_{(4)}\Gamma_{\Sigma_4}\Gamma_{N_2} = \gamma_{(4)} \otimes \tilde{\gamma}_{(6)} = \gamma_{(4)} \otimes \tilde{\gamma}_{(4)} \otimes \hat{\gamma}_{(2)}$. We also specialize below to the matrix representations,

$$\begin{aligned} \gamma^\mu &= \begin{pmatrix} 0 & -\bar{\sigma}^\mu \\ \sigma^\mu & 0 \end{pmatrix} = \begin{pmatrix} 0 & [-1, \sigma_{xyz}] \\ [1, \sigma_{xyz}] & 0 \end{pmatrix}, \quad \tilde{\gamma}^a = \begin{pmatrix} 0 & [-i, \sigma_{zxy}] \\ [i, \sigma_{zxy}] & 0 \end{pmatrix}, \quad \hat{\gamma}^r = (\tau_1, \tau_2), \\ [\mu = 0, \dots, 3, a = 1, \dots, 4, B = \Gamma^2\Gamma^4\Gamma^7\Gamma^9] &= \begin{pmatrix} 0 & -\sigma_2 \\ \sigma_2 & 0 \end{pmatrix} \otimes \begin{pmatrix} -i\sigma_2 & 0 \\ 0 & -i\sigma_2 \end{pmatrix} \otimes \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \\ \gamma_{(4)} &= \text{diag}(1_2, -1_2), \quad \tilde{\gamma}_{(6)} = \text{diag}(1_8, -1_8), \quad \tilde{\gamma}_{(4)} = \text{diag}(1_2, -1_2), \quad \hat{\gamma}_{(2)} = \text{diag}(1, -1). \end{aligned} \quad (\text{IV.3})$$

Note that our choice of the $SO(6)$ gamma matrices representations differs from [6] but coincides with [9]. The fermionic brane action of quadratic order in the bi-spinor fields, $\Theta = (\theta_1 \theta_2)^T$, involves the extended Dirac operator in M_4 , Σ_4 including the classical 5-form [6],

$$\begin{aligned} \gamma^{\alpha\beta}(\Gamma_\alpha D_\beta + \frac{i}{8}\Gamma_\alpha \mathcal{H}_5 \Gamma_\beta \otimes \sigma_2) &= \Gamma^\mu \mathcal{D}_\mu + \Gamma^a \mathcal{D}_a, \quad [\mathcal{D}_\mu = \nabla_\mu - \frac{1}{4}\Gamma_\mu \hat{\theta}_{\Sigma_4}(\ln h(\tau)) P_+^{O3}, \\ \mathcal{D}_a &= \nabla_a + \frac{1}{8}\partial_a(\ln h(\tau)) - \frac{1}{4}\hat{\theta}_{\Sigma_4}(\ln h(\tau))\Gamma_a P_+^{O3}, \quad P_\pm^{O3} = \frac{1}{2}(1 \pm \Gamma_{(6)} \otimes \sigma_2)]. \end{aligned} \quad (\text{IV.4})$$

The quantities adapted to the unwarped metric (independent of the warp profile $h(\tau)$) are introduced via the replacements

$$\tilde{D}_{M_4} = \Gamma^\mu \mathcal{D}_\mu \rightarrow h^{1/4}(\tau) \mathcal{D}_{M_4}, \quad \tilde{D}_{\mathcal{C}_6} = \Gamma^m \mathcal{D}_m \rightarrow h^{-1/4}(\tau) \mathcal{D}_{\mathcal{C}_6}, \quad (\text{IV.5})$$

where the tilde symbols were suppressed, for simplicity. The projections on Σ_4 of the covariant derivatives of \mathcal{C}_6 comprise both the Lorentz spin connection ω_a and the gauge connection A_a , with $\nabla_{\mathcal{C}_6} = \Gamma^a \nabla_a$, $[\nabla_a = \partial_a + \frac{1}{4}\omega_a^{\hat{a}\hat{b}}\Gamma_{\hat{a}\hat{b}} + A_a]$ but ignore the topological twisting terms from the R-symmetry group $SO(2)$ [96] to be discussed at the end of this subsection. Note that the pull-back transformations of bulk fields do not affect scalar quantities such as $\hat{\theta}\phi(X)$, $\hat{\theta}h(\tau)$, \mathcal{H} and that the cross terms $(d\rho + i\hat{h}_3)d\bar{\chi} + H$. c. in the conifold metric $g^{ab}\Gamma_a D_b \rightarrow g^{\rho\chi}\Gamma_\rho \partial_\chi$ drop out in the kinetic action. The fermionic $D7$ -brane

$$\begin{aligned} \delta S_F^{(2)}(D7) &= i\tau_7 \int d^7\xi \sqrt{-\gamma} e^\phi \bar{\Theta} P_-^{D7} [h^{1/4} \tilde{D}_{M_4} + h^{-1/4} \tilde{D}_{\mathcal{C}_6} + \hat{\theta}_{\Sigma_4}(\ln h(\tau)) (\frac{1}{8} - \frac{1}{2} P_+^{O3})] \Theta, \\ [P_-^{D7} &= (1 - \Gamma_{(8)} \otimes \sigma_2)/2, \quad \Gamma_{(8)} = -\Gamma_{(4)}\Gamma_{\Sigma_4}] \end{aligned} \quad (\text{IV.6})$$

includes the projection operator P_-^{D7} ensuring κ -symmetry. The Kaluza-Klein ansatz for the component Dirac spinors involves direct products of the 4-d and 6-d Weyl spinors, $\theta_{i,4}^{(m)}(x)$ and $\theta_{i,6}^{(m)}(y)$ of same chiralities ± 1 ,

$$\theta_i(X) = \sum_m [\theta_{i,4}^{(m)}(x) \otimes \theta_{i,6}^{(m)}(y) + (B_4 \theta_{i,4}^{(m)})^* \otimes (B_6 \theta_{i,6}^{(m)})^*], \quad (\text{IV.7})$$

where the linear combinations ensure that the 10-d Majorana-Weyl spinors $\theta_i(X)$ satisfy the reality condition which was expressed in terms of the factorized form of the charge conjugation matrix, $B = B_4 \otimes B_6$, $[B_4 \gamma^\mu B_4^* = \gamma^{\mu*}, B_6 \tilde{\gamma}^m B_6^* = -\tilde{\gamma}^{m*}]$ in Eq. (IV.3) for the gamma matrices. The summation over the modes index label m involves products of (same chirality) 4-d and 6-d Dirac spinors to which are added the products of opposite chirality spinors. Consider specifically the basis $\lambda_{1,2,3,4} = [(+, +, +), (+, -, -), (-, -, +), (-, +, -)]$ of the + chirality spinors in M_{10} : $\theta_i \in 32$, $[\Gamma_{10} \theta_i = \theta_i]$ formed from direct products of the \pm -chiralities spinors 4, 4, 2 of $M_4 \times \Sigma_4 \times N_2$. The projection on Majorana-Weyl spinors selects the basis of spinors, $\lambda_1 + \lambda_4 = \lambda_1 + (B\lambda_1)^*$ and $\lambda_2 + \lambda_3 = \lambda_2 + (B\lambda_2)^*$, of chiralities +1 and -1 in Σ_4 , but mixed chiralities in M_4 , N_2 , as seen on their representations in terms of the 2, 2, 1 Weyl spinors ξ , η , ζ of M_4 , Σ_4 , N_2 ,

$$\begin{aligned} \lambda_1 + \lambda_4 &= \begin{pmatrix} \xi \\ 0 \end{pmatrix} \begin{pmatrix} \eta \\ 0 \end{pmatrix} \begin{pmatrix} \zeta \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ -\sigma_2 \xi^* \end{pmatrix} \begin{pmatrix} -i\sigma_2 \eta^* \\ 0 \end{pmatrix} \begin{pmatrix} 0 \\ -i\zeta^* \end{pmatrix}, \\ \lambda_2 + \lambda_3 &= \begin{pmatrix} \xi \\ 0 \end{pmatrix} \begin{pmatrix} 0 \\ \eta \end{pmatrix} \begin{pmatrix} 0 \\ \zeta \end{pmatrix} + \begin{pmatrix} 0 \\ -\sigma_2 \xi^* \end{pmatrix} \begin{pmatrix} 0 \\ -i\sigma_2 \eta^* \end{pmatrix} \begin{pmatrix} i\zeta^* \\ 0 \end{pmatrix}. \end{aligned} \quad (\text{IV.8})$$

Substitution of Eq. (IV.7) into Eq. (IV.6) yields sums over pairs of square and cross matrix elements, $\overline{(\theta + (B\theta)^*)} \mathcal{D}(\theta + (B\theta)^*) = (S_1 + S_2) + (C_1 + C_2)$, which by virtue of the relations $\overline{(B\theta)^*} \mathcal{D}(B\theta)^* = -\eta \bar{\theta} \mathcal{D} \theta$, $\bar{\theta} \mathcal{D}(B\theta)^* = -(\theta^T C \mathcal{D}$

$\theta)^*$, $(B\theta)^* \mathcal{D}\bar{\theta} = \theta^T C \mathcal{D}\theta$, $[B^T = C\Gamma^0, B^* = \Gamma_0 C]$ and the identities $\Gamma_0 \mathcal{D}\Gamma_0 = \mathcal{D}^\dagger$, $\Gamma_a^\dagger = \Gamma_0 \Gamma_a \Gamma_0$ satisfy $S_1 + S_2 = 2S_1$ and $C_1 + C_2 = C_1 - C_1^* = 0$.

The calculations are facilitated by selecting a suitable gauge for the local fermionic κ -symmetry. For the simple choice, $\Theta \equiv (\theta_1, \theta_2) = (\theta, 0)$, the wave equations for modes of mass ω_m , subject to the mass shell conditions, $\Gamma_{(4)} \mathcal{D}_{M_4} \Theta_4 = -\omega_m (B_4 \Theta_4(x))^*$, are derived from the stationarity property of the action in the form

$$\Gamma_{(4)} [\mathcal{D}_{\mathcal{C}_6} - \frac{1}{8} \tilde{\phi}_{\Sigma_4}(\ln h)(1 + 2\Gamma_{\Sigma_4})] \theta_6^{(m)}(y) = h^{1/2}(y) \omega_m (B_6 \theta_6^{(m)}(y))^*, \quad (\text{IV.9})$$

where the $\Gamma_{(4)}$ factor ensures that the external and internal space Dirac operators commute. The 6-d massless ($\omega_m = 0$) zero modes of chirality $\Sigma_4 = \pm 1$ have the explicit dependence on the warp profile $h^{3/8}(\tau)$ and $h^{-1/8}(\tau)$, as verified from the wave equations

$$[\mathcal{D}_{\mathcal{C}_6} - \frac{1}{8} \begin{pmatrix} 3 \\ -1 \end{pmatrix} \tilde{\phi}_{\Sigma_4}(\ln h)] \theta_{6,\pm}^{(0)}(y) = 0 \implies \mathcal{D}_{\mathcal{C}_6} \psi_{\pm} = 0, [\theta_{6,+}^{(0)}(y) = h^{3/8} \psi_+(y), \theta_{6,-}^{(0)}(y) = h^{-1/8} \psi_-(y)] (\text{IV.10})$$

Note that if the classical F_5 -term were ignored, all modes would have the same warp profile factor, $\theta_{6,\pm} = h^{1/8} \psi_{\pm}$. The kinetic energy action for zero modes becomes

$$\delta S_F^{(2)}(D7) = i\tau_7 \int_{M_4} d^4x \sqrt{-\hat{g}_4} e^\phi \sum_r \int_{\Sigma_4} V_0(\rho) [(\bar{\theta}_{4,+}^{(r)} \tilde{D}_4 \theta_{4,+}^{(r)}) h(\tau) (\psi_+^\dagger \psi_+(\rho)) + (\bar{\theta}_{4,-}^{(r)} \tilde{D}_4 \theta_{4,-}^{(r)}) (\psi_-^\dagger \psi_-(\rho))] (\text{IV.11})$$

The correspondence between fermionic and bosonic superpartner modes is more transparent in the alternative κ -symmetry gauge fixing condition [6, 96], $P_-^{Dp} \Theta = 0 \implies \Theta = (\theta, -i\Gamma_{(8)} \theta)^T$. Using the relations $P_\pm^{O3} \Theta_{6,\pm} = \Theta_{6,\pm}$, $P_\mp^{O3} \Theta_{6,\pm} = 0$ in the extended Dirac equation $[\mathcal{D}_{M_4} + \mathcal{D}_{\mathcal{C}_6} + \tilde{\phi}_{\Sigma_4}(\ln h)(1/8 - P_+^{O3}/2)] \Theta = 0$, yields the wave equations,

$$-h^{1/2}(\tau) \omega_m (B_6 \Theta_6^{(m)})^* + \Gamma_{(4)} (\mathcal{D}_{\mathcal{C}_6} - \frac{1}{8} \tilde{\phi}_{\Sigma_4}(\ln h)(1 + 2\Gamma_{\Sigma_4} \otimes \sigma_2)) \Theta_6^{(m)} = 0, [\Theta_6^{(m)} = (\theta_{1,6}^{(m)}, \theta_{2,6}^{(m)})]. \quad (\text{IV.12})$$

The zero modes of Σ_4 -chirality ± 1 have the same warp profile dependence, $\Theta_{6,\pm}^0 = h^{[3/8, -1/8]} \Psi_{\pm}$, and the correspondence to the bosonic superpartner modes of same warp profile dependence associates Ψ_+ to gaugino and modulino and Ψ_- to wilsonini. Recall that for compact Calabi-Yau manifolds [6], the $(SO(2)_R$ singlet) gaugino wave function are built from the covariantly constant spinor η^{CY} . If Σ_4 is a complex manifold of Kähler metric, $\gamma_{AB} dW^A dW^B$ in a suitable basis of holomorphic coordinates W^A , one can then build the modulini and wilsonini wave functions by acting with linear combinations of Dirac matrices [97], $\eta_m = m_{AB} \Gamma^{W^A W^B} \eta^{CY}$, $\eta_W = W_A \Gamma^{W^A} \eta^{CY}$. The independent harmonic 2- and 1-forms m_{AB} , W_A on Σ_4 (of Betti numbers $b_2(\Sigma_4) = h^{2,0}$ and $b_1(\Sigma_4) = 2h^{1,0}$) also serve to build up the wave functions of bosonic superpartner modes, $\chi \sim X^{8,9}$ and $A_{\hat{a}} \sim X^a$. The wave functions for gaugino and modulini $\theta_{6,+} \sim (\lambda_2, \lambda_3)$ and wilsonini $\theta_{6,-} \sim (\lambda_1, \lambda_4)$ are of form [6],

$$\theta_{6,+} = \frac{h^{3/8}}{\sqrt{2}} \left[\left[\begin{pmatrix} i\eta^{CY} \\ \eta^{CY} \end{pmatrix} + \dots \right], \left[\begin{pmatrix} i\eta_m \\ \eta_m \end{pmatrix} + \dots \right] \right], \quad \theta_{6,-} = \frac{h^{-1/8}}{\sqrt{2}} \left[\left[\begin{pmatrix} -i\eta_W \\ \eta_W \end{pmatrix} + \dots \right], \right], \quad (\text{IV.13})$$

where the central dots refer to the terms projecting on Majorana spinors. While the above properties need not apply to non-compact manifolds, normalizable massless fermions might still exist in the conifold case. This is found to be the case for Killing spinors in supergravity backgrounds [31] and also for the massless axino [99] superpartner of the axion mode [98] induced through the spontaneous breaking of the baryon charge symmetry $U(1)_b$ in Klebanov-Strassler background. For completeness, we also recall that fermionic zero modes arise typically in 4-d gauge theories with instantons [11] or magnetic backgrounds [6, 9, 12] and in theories with extra dimensions with D -branes intersecting at points D -branes [7], or instanton (Euclidean) branes wrapped on divisors of the compactification manifold [93–95]. In theories with anomalous global symmetries the numbers of massless fermions are determined via the Atiyah-Singer index or the holomorphic (arithmetic) index, $\chi(D_{6,4}) = \sum_p (-1)^p h^{(0,p)}$.

Before analyzing further the issue of fermionic zero modes, we review briefly the geometrical approach [96] to embed superbranes in curved manifolds. (Generalized versions exist for superspacetimes and superworld volumes [58]). One assumes here a linear relationship between the bulk spacetime vielbeins $e^a = e_M^a dX^M$ and the tangential and normal vielbeins of the brane world volume $\epsilon^a = \epsilon_m^a d\xi^m$, $\epsilon^i = \epsilon_m^i d\xi^m$, involving the local Lorentz transformations $u_{\underline{b}}^{\underline{a}}(\xi) \in SO(1,9)$,

$$\epsilon^a = u_{\underline{b}}^a e^{\underline{b}}, \quad \epsilon^i = u_{\underline{b}}^i e^{\underline{b}}, \quad [u_{\underline{a}}^{\underline{b}} = (u_{\underline{a}}^{\underline{b}}, u_{\underline{a}}^{\underline{c}}) = (u_{\underline{a}}^{\underline{b}}, u_{\underline{c}}^{\underline{b}}) = \begin{pmatrix} u_{\underline{a}}^{\underline{b}} & u_{\underline{a}}^{\underline{c}} \\ u_{\underline{c}}^{\underline{b}} & u_{\underline{c}}^{\underline{c}} \end{pmatrix}, \quad \epsilon^a \cdot \epsilon^i = 0] \quad (\text{IV.14})$$

which align $(p+1)$ of the bulk tangent space basis onto the world volume tangent space basis. Similar relations hold between the bulk gamma matrices and the gamma matrices, $\Gamma^a = \Gamma^{\underline{a}}u_{\underline{a}}^a$, $\Gamma^i = \Gamma^{\underline{a}}u_{\underline{a}}^i$ along tangential and normal directions. Imposing the embedding equations $\epsilon^i \equiv u_{\underline{b}}^i e^{\underline{b}} = 0$, which imply that the $(9-p)$ normal vielbeins are annihilated by the Lorentz transformation aligning the ϵ^a along the world volume, makes u become auxiliary (non-dynamical fields). The 10-d metric and the world volume embedding $\partial_m X^M$ determine the tangential sub-block $u_{\underline{b}}^{\underline{a}} = e_{\underline{M}}^{\underline{a}} \partial_m X^M \epsilon_b^m$ (reducing to $e_i^{\underline{a}} \partial_m X^i \epsilon_b^m$ in the static gauge) while the orthogonality conditions $u_{\underline{a}}^i u_{\underline{b},i} = \eta_{\underline{ab}} - u_{\underline{a}}^i u_{\underline{b}}^j \eta_{ab}$ determine the normal sub-blocks. With these definitions and a suitable choice for the gamma matrices, one can express after some work the covariant derivatives $D_m = \partial_m + \Omega_m$ involving the bulk spin connection Ω_m in terms of tangential and normal induced spin connections ω_m^{ab} , A_m^{ij} for the reduced spinors of $SO(1,p)$ and $SO(9-p)$,

$$D_x \equiv (e_m^{\underline{c}} \partial_{\underline{c}} - \frac{1}{4} \Omega_m^{\underline{ab}} \Gamma_{\underline{ab}}) = \mathcal{D}_m - \frac{1}{4} u^{ca} \partial_m u_{\underline{c}}^b \Gamma_{\underline{ab}} - \frac{1}{2} K_m^{bi} \Gamma_{bi}, \quad [\mathcal{D}_m = e_m^{\underline{c}} \partial_{\underline{c}} - \frac{1}{4} \Omega_m^{\underline{ab}} \Gamma_{\underline{ab}} - \frac{1}{4} A_m^{ij} \Gamma_{ij},$$

$$\omega^{bc} = u_{\underline{b}}^b \Omega^{\underline{bc}} u_{\underline{c}}^c - u^{\underline{ab}} du_{\underline{a}}^c, \quad A^{ij} = u_{\underline{b}}^i \Omega^{\underline{bc}} u_{\underline{c}}^j - u^{\underline{ai}} du_{\underline{a}}^j, \quad K_b^i = -D(e_{\underline{b}}^{\underline{a}}) u_{\underline{a}}^i = -(d(e_{\underline{b}}^{\underline{a}}) + u_{\underline{b}}^{\underline{a}} \Omega_{\underline{a}}^{\underline{b}}) u_{\underline{a}}^i] \quad (\text{IV.15})$$

where $K^i = d\xi^b K_b^i$ denotes the world volume extrinsic curvature (second fundamental form).

B. Fermionic zero modes

We discuss in this subsection the existence of massless fermions on $D7$ -branes in the Klebanov-Strassler background, within the simplified description using the Dirac operator restriction on Σ_4 , ignoring twisting terms,

$$\tilde{D}|_{\Sigma_4} = \Gamma^m (\partial_m + \frac{1}{4} \omega_m^{\hat{a}\hat{b}} \Gamma_{\hat{a}\hat{b}}), \quad [\Gamma_m = e_m^{\hat{a}} \Gamma_{\hat{a}}, \quad \Gamma^m = e_{\hat{a}}^m \Gamma^{\hat{a}}, \quad \Gamma_{\hat{a}} = \delta_{\hat{a}\hat{b}} \Gamma^{\hat{b}}, \quad \{\Gamma_{\hat{a}}, \Gamma_{\hat{b}}\} = 2\delta_{\hat{a}\hat{b}}] \quad (\text{IV.16})$$

where the indices m and \hat{a} refer to curved and flat directions and the spin connection is evaluated from Eq.(A.29) in the basis of vielbeins defined in Eq. (A.32). Thanks to the $SO(3)$ isometry of Kuperstein 4-cycle, the general formulas for the Dirac operator in Eqs.(A.30) and (A.31) of Appendix A 3 simplify to

$$\tilde{D}|_{\Sigma_4} = \frac{1}{\eta^{\hat{\rho}}} \Gamma^{\hat{\rho}} (\partial_{\hat{\rho}} + \frac{1}{2} \sum_{\hat{k}} \partial_{\hat{\rho}} (\ln \eta^{\hat{k}}) + X(\rho) \Gamma_{\Sigma_4} + \eta^{\hat{\rho}} \Gamma_{\hat{\rho}} \Gamma^{\hat{k}} e_{\hat{k}}^m \partial_m) = \eta^{\hat{\rho}} \Gamma^{\hat{\rho}} (\partial_{\hat{\rho}} + \frac{1}{2} \partial_{\hat{\rho}} \ln D + X(\rho) \Gamma_{\Sigma_4} + \eta^{\hat{\rho}} \Gamma_{\hat{k}} \Gamma^{\hat{\rho}\hat{k}} \nabla_{\hat{h}_k}),$$

$$[X(\rho) = \frac{(\eta^{\hat{\rho}})^2}{8D} (\frac{N_1}{\eta^{\hat{1}}} + \frac{N_2}{\eta^{\hat{2}}} + \frac{N_3}{\eta^{\hat{3}}}), \quad D = \eta^{\hat{1}} \eta^{\hat{2}} \eta^{\hat{3}}, \quad N_1 = -\eta^{\hat{1}} + \eta^{\hat{2}} + \eta^{\hat{3}}, \quad N_2 = \eta^{\hat{1}} - \eta^{\hat{2}} + \eta^{\hat{3}}, \quad N_3 = \eta^{\hat{1}} + \eta^{\hat{2}} - \eta^{\hat{3}}] \quad (\text{IV.17})$$

The successive contributions refer to the radial derivative term, two potential terms (proportional to the unit matrix and chirality matrix $\Gamma_{\Sigma_4} = \Gamma^{\hat{\rho}\hat{1}\hat{2}\hat{3}}$) and the angular term $\tilde{D}(S^3)$. The latter term drops out for singlet modes, satisfying the conditions $e_{\hat{k}}^m \partial_{\alpha^m} \psi(\rho, \alpha) = 0$, to which we restrict hereafter. In the gamma matrices representation of Σ_4 ,

$$\Gamma^{\hat{a}} = \gamma_{(4)} \times \tilde{\gamma}^{\hat{a}}, \quad \tilde{\gamma}_{(4)} = \tilde{\gamma}^{\hat{\rho}\hat{1}\hat{2}\hat{3}} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad [\hat{a} = (\hat{\rho}, \hat{k}), \quad \tilde{\gamma}^{\hat{\rho}} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \tilde{\gamma}^{\hat{1},\hat{2},\hat{3}} = \begin{pmatrix} 0 & \sigma_{z,x,y} \\ \sigma_{z,x,y} & 0 \end{pmatrix}] \quad (\text{IV.18})$$

quoted from Eq. (IV.2), the chirality matrix diagonal $\Gamma_{\Sigma_4} = \text{diag}(1_2, -1_2)$ and the 4-spinors split up into a decoupled pair of 2-spinors of ± 1 chirality, $\psi = (\psi_+, \psi_-)$. The solutions $\psi_{\pm}(\rho)$ of the decoupled radial wave equations,

$$(\partial_{\hat{\rho}} + \frac{1}{2} \sum_{\hat{k}} \partial_{\hat{\rho}} (\ln \eta^{\hat{k}}) \pm X(\rho)) \psi_{\pm}(\rho) = 0 \implies \psi_{\pm}(\rho) = C_{\pm} \psi_0(\rho) e^{\mp \int_{\rho_C}^{\rho} d\rho' X(\rho')} \xi_{\pm}, \quad [\psi_0(\rho) = \frac{1}{\prod_{\hat{k}} (\xi^{\hat{k}})^{1/2}}] \quad (\text{IV.19})$$

are expressed in terms of the 2-d constant spinors ξ_{\pm} are 2-d constant spinors and the constant factors C_{\pm} , along with the boundary values ρ_C in the definite integrals, are set through the wave functions normalizations. For modes of canonical kinetic actions in Eq. (IV.11), the normalization conditions, $1 = C_+^2 \int d\rho \sqrt{-\gamma^{(0)}} h(\tau) |\psi_+|^2$, $1 = C_-^2 \int d\rho \sqrt{-\gamma^{(0)}} |\psi_-|^2$, determine C as a function of ρ_C . The existence of normalizable zero modes is thus dictated by the behaviour of radial wave functions at small and large distances.

We display in Fig. 4 the solutions for the zero modes radial wave functions. The zeroth-order factor $\psi_0(\tau)$ is regular at infinity but diverges at threshold. Both solutions ψ_{\pm} slope to zero at large τ , with ψ_+ converging much more rapidly than ψ_- . Near the threshold ($\tau = \tau_{min} \simeq 2.58$ for $L = 1.5$), ψ_- vanishes with a rapid slope and ψ_+ diverges. Since the normalization integrals diverge for both ψ_+ and ψ_- , whether the warp profile is taken into account or not, it follows that the existence of zero modes is ruled out. Only if one imposed an ultraviolet radial cutoff τ_{uv} could the normalizable mode ψ_- be present.

Similar conclusions are reached in the undeformed conifold case. The radial Dirac operator (for singlet modes)

$$\begin{aligned} \mathcal{D} &= \eta_{\hat{\rho}} \Gamma^{\hat{\rho}} (\partial_{\rho} + \frac{1}{2} \partial_{\rho} (\ln D) + X(\rho) \Gamma_{\Sigma_4}), \quad [D(\rho) = \eta^1 \eta^2 \eta^3 = \frac{|\mu|^2}{16\sqrt{3}} ((2r_{\chi}^3 - 1)(r_{\chi}^3 - 2))^{1/2}, \\ \frac{X(\rho)}{dr_{\chi}/d\rho} &= \frac{(2r_{\chi}^3 - 1)^{1/2}}{\sqrt{6}r_{\chi}^{5/2}} + \frac{1}{12r_{\chi}} \left[3 + \frac{2\sqrt{6}(r_{\chi}^3 + 1)(2r_{\chi}^3 - 1)^{1/2}}{(r_{\chi}^3 - 2)^{3/2}} \right] \end{aligned} \quad (\text{IV.20})$$

admits the zero modes solutions

$$\psi_{\pm}(r_{\chi}) = C_{\pm} \psi_0(r_{\chi}) e^{\mp \int_{\rho_C}^{\rho} d\rho X(\rho)}, \quad [\psi_0(r_{\chi}) = \frac{4 \cdot 3^{1/4}}{|\mu|((r_{\chi}^3 - 2)(2r_{\chi}^3 - 1))^{1/4}}] \quad (\text{IV.21})$$

where the indefinite integral in the exponential is expressed by the analytic formula,

$$\begin{aligned} \int^{r_{\chi}} d\rho X(\rho) &= \frac{1}{36} [-6\sqrt{6}(\frac{2r_{\chi}^3 - 1}{r_{\chi}^3 - 2})^{1/2} - 4\sqrt{6}(\frac{2r_{\chi}^3 - 1}{r_{\chi}^3})^{1/2} \\ &+ 8\sqrt{3}(\text{Arcsinh}(\sqrt{\frac{2}{3}}(r_{\chi}^3 - 2)^{1/2}) + \text{Arcsinh}((2r_{\chi}^3 - 1)^{1/2})) + 2\sqrt{3}\text{Arctanh}((\frac{r_{\chi}^3 - 2}{2(2r_{\chi}^3 - 1)})^{1/2}) + 9 \log r_{\chi}] \end{aligned} \quad (\text{IV.22})$$

The series expansions near threshold $z \equiv (r - 2^{1/3}) \rightarrow 0$ and the boundary $r_{\chi} \rightarrow \infty$ of the various factors in the integrands of the normalization integrals, $N(\psi_{\pm}) = \int_{\rho_C}^{\rho} d\rho \sqrt{-\gamma^{(0)}} [\psi_{\pm}^2 h(r), \psi_{\pm}^2]$ are displayed in the table below for the parameter value $\mu = 1.5$. The values of the constant coefficients C_{\mp} are correlated with the value chosen for the radial integral lower bound ρ_C .

	Limit $\frac{dr_{\chi}}{d\rho}$	ψ_0	$e^{\int_{\rho_C}^{\rho} d\rho X(\rho)}$	ψ_{\mp}	$N(\psi_{\pm}^2)$	$N(\psi_{\pm}^2 h(r))$
$z \rightarrow 0$	$(2^{1/6}/\sqrt{3})z^{1/2}$	$1.8z^{-1/4}$	$e^{\mp 0.32z^{-1/2}}$	$1.8z^{-1/4}e^{\mp 0.32z^{-1/2}}$		
$r_{\chi} \rightarrow \infty$	$r_{\chi}/3$	$2.95r_{\chi}^{-3/2}$	$r_{\chi}^{(1/4+2/\sqrt{3})}$	$\begin{pmatrix} C_- r_{\chi}^{-5/4+2/\sqrt{3}} \\ C_+ r_{\chi}^{-7/4-2/\sqrt{3}} \end{pmatrix}$	$r_{\chi}^{3/2+4/\sqrt{3}}$	$r_{\chi}^{-7/2-4/\sqrt{3}}$

The normalization integral for ψ_+ converges at large radius but diverges near threshold, while that for ψ_- diverges at large radius but converges near threshold, as verified from the limited expansions above. The radial wave functions displayed in Fig.4 are qualitatively similar to those in the deformed conifold case. The solution ψ_- vanishes at threshold $r_{\chi}^{\min} \simeq (2\mu^2)^{1/3} \simeq 1.65$, and jumps rapidly to a plateau that slopes very slowly to zero. This is explained by our choice for the arbitrary parameter ρ_C which contributes a large constant part to the wave function, $\psi(r_{\chi})$ that is compensated by the normalization factor C_- . We thus conclude that also the undeformed conifold does not support singlet zero modes but could allow one ψ_- if an ultraviolet cutoff were imposed.

V. SUMMARY AND CONCLUSIONS

We examined in this work the Kaluza-Klein theory for $D7$ -branes embedded in Kuperstein 4-cycle of Klebanov-Strassler background. The mass spectra and wave functions of ground state and radially excited meson modes, singlet under the base manifold isometry group, were evaluated using semi-classical tools. The reduced masses $\hat{\omega}_m$ (in same units $\epsilon^{2/3}(g_s M \alpha')/\epsilon^{2/3}$ as for glueball masses \hat{E}_m) exhibit a nearly uniform dependence on the ratio $L = \mu/\epsilon \in R$, being nearly constant in $L \in [0, 1]$ and growing monotonically with the power law $\hat{\omega}_m \sim L^{2/3}$ at $L > 5$. Similar results hold for $L^2 < 0$ with no qualitative differences are expected for complex values of L . The classical 2-form B_2^l gives contributions resulting in lighter modes changing the power law growth from $\omega_m^2 \approx 0.5 L^{4/3} \rightarrow \omega_m^2 \approx (0.4 - 0.3) L^{4/3}$. The meson modes are more strongly bound than glueball modes with typical ratios $\hat{\omega}_n^2/\hat{E}_n^2 \simeq 1/3$ for $L \in [0, 1]$. Scalar modes are always lighter than vectors modes. The mass splittings between radial excitations depend weakly on L . The mass splittings for radial excitations are independent of L .

The coupling constants of interactions between meson and glueball modes were evaluated from the overlap integrals of meson and glueball wave functions. The gravitational couplings of meson modes decrease rapidly beyond $L > 3$ while their self couplings increase slowly with L . The existence of normalizable fermionic massless modes is ruled out unless one imposes an ultraviolet cutoff on radial distances.

Our test of duality identifying the holographic meson modes to the hadronic quarkonia of light u, d, s and heavy c, b flavour was inconclusive partly due to the uncertain gauge-gravity duality correspondence relations between parameters. Improvements are expected if supersymmetry breaking effects are taken into account.




FIG. 4: Plots of the fermionic zero mode wave functions (in units $\epsilon = 1$, $g_s M \alpha' = 1$) on Kuperstein 4-cycle as a function of the deformed conifold radial variable τ for $L = 1.5$, $[\tau \geq \tau_{\min} = 2.58]$ (left panel) and the undeformed conifold radial variable r_χ for $\mu = 1.5$, $[r_\chi \geq (r_\chi)_{\min} = 1.65]$ (right panel). The (red) curves in long dashes refer to the prefactor $\psi_0(\tau)$ and the (green and blue) curves in shorter dashes refer to the unnormalized radial wave functions, $\psi_\pm(\rho) = C_\pm \psi_0 e^{\mp \int_{\rho_C}^{\rho} d\rho X(\rho)}$ of chiralities ± 1 . The values of the normalization constant C_\pm depend on the choice of the arbitrary lower limit ρ_B affecting the wave functions and the normalization integrals.

Appendix A: Embedding of $D7$ -branes in deformed conifold

We review in this appendix useful tools for computing the reduced $D7$ -brane action on Kuperstein submanifold [34] Σ_4 : $w_4 \equiv \frac{1}{2i} \text{Tr}(W) = \mu$ of the deformed conifold in Eq. (III.1), where the complex parameter $\mu \in C$, $[|\mu| = L^{2/3}]$ measures the minimal distance from its apex to the conifold apex. Helpful details complementing the discussion in the text are provided in the following four subsections. We start with geometrical properties of the 4-cycle embedding, discuss in turn the reduced bosonic and fermionic actions of $D7$ -branes and finally conclude with intermediate results in the undeformed conifold limit.

1. Geometrical properties of 4-cycle embedding

The metric in the spacetime $M_4 \times C_6$ is expressed in a convenient form by introducing the basis of cotangent space frame vectors, $e^{\hat{\mu}}$, $(e^{\hat{\rho}}, e^{\hat{k}}, e^{\hat{\chi}})$, $[\mu = 0, 1, 2, 3, k = 1, 2, 3]$ and the dual orthogonal basis of tangent space frame vectors $e_{\hat{\mu}}$, $(e_{\hat{\rho}}, e_{\hat{k}}, e_{\hat{\chi}})$, such that the sets of components of these two bases, $(e_{\hat{\nu}}^{\hat{\mu}}, e_{\hat{\rho}}^{\hat{\rho}}, e_{\hat{m}}^{\hat{k}})$ and $(e_{\hat{\mu}}^{\hat{\nu}}, e_{\hat{\rho}}^{\hat{\rho}}, e_{\hat{k}}^{\hat{m}})$ describe mutually inverse matrices. (The present notation for covariant and contravariant vectors is also used for differential forms and derivatives.) The warped metric consists of 4-d and 6-d parts, $ds_{10}^2 = h^{-1/2}(\tau) d\tilde{s}_4^2 + h^{1/2}(\tau) d\tilde{s}_6^2$, defined as

$$\begin{aligned} d\tilde{s}_4^2 &= \tilde{g}_{\mu\nu} dx^\mu dx^\nu = \eta_{\hat{\mu}\hat{\nu}} e^{\hat{\mu}} e^{\hat{\nu}} = \eta_{\hat{\mu}\hat{\nu}} e_{\hat{\mu}}^{\hat{\mu}} e_{\hat{\nu}}^{\hat{\nu}} dx^\mu dx^\nu, \\ d\tilde{s}_6^2 &= \tilde{g}_{\rho\rho} d\rho^2 + \tilde{g}_{\hat{h}_i \hat{h}_j}(\rho) \hat{h}_i \hat{h}_j + (\tilde{g}_{\chi\bar{\chi}} d\chi d\bar{\chi} + \tilde{g}_{\rho\bar{\chi}} d\rho d\bar{\chi} + \tilde{g}_{\hat{h}_i \bar{\chi}} \hat{h}_i d\bar{\chi} + H. c.) = (e^{\hat{\rho}})^2 + (e^{\hat{h}_i})^2 + 2e^{\hat{\chi}}(e^{\hat{\chi}})^*. \end{aligned} \quad (\text{A.1})$$

It proves convenient to separate out the radial variable dependence of the frame vectors along the angular directions of the 4-cycle manifold Σ_4 , $e^{\hat{k}} = \eta^{\hat{k}}(\rho) h^{\hat{k}}$, $e_{\hat{k}} = \eta_{\hat{k}}(\rho) h_{\hat{k}}$, where $h_{\hat{k}}^{\hat{k}}$, $h_{\hat{k}}^{\hat{m}}$ denote the components of base manifold S^3/Z_2 vielbeins, depending only on the angle variables α^m . The non-vanishing components of the C_6 metric in Eq. (III.12) take the equivalent forms

$$\begin{aligned} \tilde{g}_{\mu\nu} &= \eta_{\mu\nu}, \quad \tilde{g}_{\rho\rho} \equiv (\eta^{\hat{\rho}})^2 = \tilde{g}_{\hat{h}_3 \hat{h}_3} \equiv (\eta^{\hat{3}})^2 = \frac{F_1}{2} |\eta_\chi|^2 A_1, \quad \tilde{g}_{\hat{h}_1 \hat{h}_1} \equiv (\eta^{\hat{1}})^2 = \frac{F_1}{4} |\eta_\chi|^2 (1 + C_\rho), \\ \tilde{g}_{\hat{h}_2 \hat{h}_2} &\equiv (\eta^{\hat{2}})^2 = \frac{F_1}{4} |\eta_\chi|^2 (-1 + C_\rho), \quad \tilde{g}_{\rho\chi} = (\tilde{g}_{\rho\bar{\chi}})^* = i\tilde{g}_{\hat{h}_3 \chi} = -i(\tilde{g}_{\hat{h}_3 \bar{\chi}})^* = \frac{1}{2} F_1 \bar{\eta}_\chi S_\rho A_2, \quad \tilde{g}_{\chi\bar{\chi}} = \tilde{g}_{\bar{\chi}\chi} = F_1 A_3, \\ [A_1 &= (C_\rho + 2|\eta_\chi|^2 R S_\rho^2), \quad A_2 = X(1 + 2|\eta_\chi|^2 R (\frac{\bar{X}}{X} + C_\rho)), \quad A_3 = 1 + X\bar{X}(C_\rho + 2|\eta_\chi|^2 R |1 + C_\rho \frac{\bar{X}}{X}|^2)] \end{aligned} \quad (\text{A.2})$$

where $C_\rho = \cosh \rho$, $S_\rho = \sinh \rho$. The cross terms $\bar{\eta}_\chi d\bar{\chi}(d\rho - i\hat{h}_3) + H. c.$ signal that C_6 is not a direct product manifold of Σ_4 times its complementary sub-manifold N_2 in C_6 . It is still possible to express the metric as a diagonal quadratic form in the basis of vielbein vectors,

$$e^{\hat{\rho}} = (\tilde{g}_{\rho\rho} - 2\frac{|\tilde{g}_{\rho\chi}|^2}{\tilde{g}_{\chi\bar{\chi}}})^{1/2} d\rho, \quad e^{\hat{h}_3} = (\tilde{g}_{\hat{h}_3 \hat{h}_3} - 2\frac{|\tilde{g}_{\hat{h}_3 \chi}|^2}{\tilde{g}_{\chi\bar{\chi}}})^{1/2} \hat{h}_3,$$

$$e^{\hat{\chi}} = \tilde{g}_{\chi\bar{\chi}}^{1/2}(d\chi + \frac{\tilde{g}_{\rho\bar{\chi}}}{\tilde{g}_{\chi\bar{\chi}}}(d\rho - i\hat{h}_3)), \quad e^{\hat{h}_1} = \tilde{g}_{h_1 h_1}^{1/2} \hat{h}_1, \quad e^{\hat{h}_2} = \tilde{g}_{h_2 h_2}^{1/2} \hat{h}_2. \quad (\text{A.3})$$

in same notations as Eq. (A.1). The complex vielbein e^χ splits up into two real vielbeins, $e^{\hat{\phi}_1} = \tilde{g}_{\chi\bar{\chi}}^{1/2}(d\phi_1 + (\tilde{g}_{\rho\bar{\chi}}/\tilde{g}_{\chi\bar{\chi}})d\rho)$, $e^{\hat{\phi}_2} = \tilde{g}_{\chi\bar{\chi}}^{1/2}(d\phi_2 - \tilde{g}_{\rho\bar{\chi}}/\tilde{g}_{\chi\bar{\chi}}\hat{h}_3)$, $[e^{\hat{\chi}}e^{\bar{\chi}} = (e^{\hat{\phi}_1})^2 + (e^{\hat{\phi}_2})^2]$, corresponding to the pair of real moduli fields in $\chi = \phi_1 + i\phi_2$, $\bar{\chi} = \phi_1 - i\phi_2$. The restriction $\mathcal{C}_6 \rightarrow \Sigma_4$ leads to a metric and scalar Laplacian for Σ_4 of diagonal form in the coordinates (ρ, \hat{h}_i) consistently with the underlying $SO(3)$ isometry group,

$$d\tilde{s}^2(\mathcal{C}_6)|_{\Sigma_4} = (e^{\hat{\rho}})^2 d\rho^2 + \delta_{ij} e^{\hat{i}} e^{\hat{j}} d\alpha^m d\alpha^n = (e^{\hat{\rho}})^2 d\rho^2 + \eta_{\hat{k}}^2(\rho) \delta_{mn} h_{\hat{k}}^m h_{\hat{k}}^n, \quad \nabla^2|_{\Sigma_4} = e_{\hat{\rho}}^\rho \partial_\rho e_{\hat{\rho}}^\rho \partial_\rho + e_i^m \frac{\partial}{\partial \alpha^m} e_i^n \partial_{\alpha^n}, \\ [e^{\hat{\rho}} = e_{\hat{\rho}}^\rho d\rho, \quad e^{\hat{k}} = e_{\hat{l}}^{\hat{k}} \hat{h}_l = e_{\hat{m}}^{\hat{k}} d\alpha^m = \eta^{\hat{k}}(\rho) h_{\hat{m}}^{\hat{k}}(\alpha) d\alpha^m, \quad e_{\hat{\rho}} = e_{\hat{\rho}}^\rho \partial_\rho, \quad e_{\hat{k}} = e_{\hat{m}}^{\hat{k}} \nabla_m = e_{\hat{k}}^m \partial_{\alpha^m}]. \quad (\text{A.4})$$

The components of the dual bases of frame vectors $h_{\hat{m}}^{\hat{k}}$, $h_{\hat{k}}^m$ are expressed via the (row-column) matrix notation for the matrices A , B in terms of the Euler angle coordinates $\alpha^m = (\theta, \phi, \gamma/2 \in [0, 2\pi])$, describing the (Lens space) orbifold S^3/Z_2 as the Hopf map onto the fibre $S^1(\gamma/2)$ over $S^2(\theta, \phi)$,

$$h_{\hat{m}}^{\hat{k}} = A_{km} \equiv \begin{pmatrix} -2c_\gamma & -2s_\gamma s_\theta & 0 \\ -2s_\gamma & 2c_\gamma s_\theta & 0 \\ 0 & 2c_\theta & 1 \end{pmatrix}, \quad h_{\hat{k}}^m = B_{mk} = \frac{1}{2} \begin{pmatrix} -c_\gamma & -s_\gamma & 0 \\ -\frac{s_\gamma}{s_\theta} & \frac{c_\gamma}{s_\theta} & 0 \\ \frac{2c_\theta s_\gamma}{s_\theta} & -\frac{2c_\theta c_\gamma}{s_\theta} & 1 \end{pmatrix}, \quad (\text{A.5})$$

where $[c_\gamma, s_\gamma] = [\cos \gamma/2, \sin \gamma/2]$, $[c_\theta, s_\theta] = [\cos \theta, \sin \theta]$. The resulting formulas for the metric tensor,

$$\tilde{g}_{mn}^0 = \sum_{\hat{k}} h_{\hat{m}}^{\hat{k}} h_{\hat{n}}^{\hat{k}} = \begin{pmatrix} 4 & 0 & 0 \\ 0 & 4 & 2c_\theta \\ 0 & 2c_\theta & 1 \end{pmatrix}, \quad \tilde{g}^{0,mn} = \sum_{\hat{k}} h_{\hat{k}}^m h_{\hat{k}}^n = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/s_\theta^2 & -2c_\theta/s_\theta^2 \\ 0 & 2c_\theta/s_\theta^2 & 1/s_\theta^2 \end{pmatrix} \\ \implies ds^2(S^3/Z_2) = \tilde{g}_{mn}^0 d\alpha^m d\alpha^n = 4(d\theta^2 + s_\theta^2 d\phi^2 + (\frac{d\gamma}{2} + c_\theta d\phi)^2), \quad (\text{A.6})$$

give the expected results for the volume integral of the (unit radius) orbifold [100], $Vol(S^3/Z_2) = 2^{-3} \int \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 = 8\pi^2 = Vol(S^3)/2$. The geometry of Σ_4 at large radial distances $1 \ll \rho \sim \hat{r}/\mu^{2/3}$ is that of a real cone over a squashed S^3/Z_2 (in agreement with Eq. (A.38)), while that near $\rho \rightarrow 0$ is that of a real cone over the direct product of S^2 times a collapsed S^1 ,

$$\lim_{\rho \rightarrow \infty} d\tilde{s}^2(\mathcal{C}_6)|_{\Sigma_4} = \frac{3}{2} d\hat{r}^2 + \frac{\hat{r}^2}{6} (\hat{h}_1^2 + \hat{h}_2^2 + \frac{4}{3} \hat{h}_3^2), \quad \lim_{\rho \rightarrow 0} ds^2(\Sigma_4) = \frac{|\eta_\chi|^2}{2} F_1^2(\tau) (d\rho^2 + \hat{h}_3^2 + \hat{h}_2^2 + \frac{\rho^2}{4} \hat{h}_1^2). \quad (\text{A.7})$$

We now consider the harmonics of the base manifold $S^3 = SO(4)/SO(3)$. The basis of scalar harmonics $\mathcal{Y}_{m_l, m_r}^{l/2, l/2}(\alpha)$, $[j \equiv l/2 = 0, 1/2, \dots, (m_l, m_r) \in [-j, \dots, j]]$, running over the irreducible representations of $SO(4) \sim SU(2)_l \times SU(2)_r$, consists of eigenfunctions of the $S^3 \sim SU(2)$ scalar Laplacian $-\nabla^2(S^3) \rightarrow l(l+2) = 4j(j+1)$, $[j = l/2]$ of equal angular momentas $j = l/2$, $r = l/2$ of the left and right acting subgroups $SU(2)_{l,r}$. The left magnetic quantum number m_l labels the states of $SU(2)_l$ while the right one m_r labels the representations part of the harmonic basis, with the multiplicities $D_3(l) = (l+1)^2 = (2j+2)^2$. The harmonic basis in the Euler angles parameterization, $Y_{m,n}^j(\alpha_E)$, $[\alpha_E = (\theta, \phi, \gamma)]$ and that in the hyperspherical angles parameterization, $Y_{j,L,M}(\alpha_S)$, $[\alpha_S = (\chi, \theta, \phi)]$ involving Wigner functions [101] and products of Jacobi polynomials times spherical harmonics [102], respectively are linearly related via the Clebsch-Gordan coefficients [103] (enforcing the conditions $L \leq 2j$, $M = m+r$),

$$\mathcal{Y}_{m,r}^j(\alpha_E) = \sum_{L,M} \langle L, M | jm, jr \rangle Y_{j,l,M}(\alpha_S), \\ \left[\mathcal{Y}_{m,r}^j(\alpha_E) = D_{m,r}^{\frac{l}{2}, \frac{l}{2}}(\theta, \phi, \gamma), \quad [j \equiv \frac{l}{2} = 0, \frac{1}{2}, \dots, (m, r) \in (-j, \dots, +j)] \right. \\ \left. Y_{j,L,M}(\alpha_S) = \phi_{jL}(\chi) Y_{LM}(\theta, \phi), \quad [j = 0, 1/2, 1, \dots, L = 0, 1, \dots, 2j] \right]. \quad (\text{A.8})$$

The allowed functions on S^3/Z_2 , invariant under the action of Z_2 , are restricted for the Euler angles harmonics by the projection condition $r = 0 \pmod{1}$.

The harmonic decomposition for the gauge field components A_μ along M_4 utilizes the scalar harmonics while that for the longitudinal and transversal components along Σ_4 , defined by the respective conditions, $\epsilon^{ijk} \nabla_j A_k^\parallel = 0$, $\nabla^k A_k^\perp = 0$, utilizes the bases of scalar and vector harmonics,

$$A_k^\parallel = \nabla_k \mathcal{Y}^{\frac{l}{2}, \frac{l}{2}}(\alpha), \quad [E_l = -l(l+2) + 2, \quad D_l = (l+1)^2]$$

$$A_k^\perp = \mathcal{Y}_{m,r}^{l,\pm} = [\mathcal{Y}_{m,r}^{\frac{l+1}{2}, \frac{l-1}{2}}(\alpha), \mathcal{Y}_{m,r}^{\frac{l-1}{2}, \frac{l+1}{2}}(\alpha)], [E_l = -l(l+2) + 1, D_l = 2l(l+2)] \quad (\text{A.9})$$

with the above indicated values of the eigenvalues and degeneracies [67] E_l and D_l .

The spinor harmonics of S^3 can be constructed recursively [104] in terms of the basis of 2×2 Dirac matrices, $\Gamma^1 = -\sigma_2, \Gamma^2 = \sigma_1, \Gamma^3 = -i\Gamma^1\Gamma^2 = \sigma_3$. The Dirac operator harmonic eigenfunctions of angular momentum $n, 0 = (\not{D} (S^3) \mp i(n+3/2))\psi_{\pm, nl_1}(\chi, \Omega_2)$, are given by linear combinations of S^2 spinor harmonics $\chi_{ll_1}^{(\pm)}(\Omega_2)$, eigenfunctions of the Dirac operator of angular momentum $l, (\Gamma^3 \not{D}_{S^2} \mp (l+1))\chi_{ll_1}^{(\pm)}(\Omega_2) = 0$. The latter are in turn expressed in terms of S^1 spinor harmonics, eigenfunctions of Dirac operator of angular momentum $l_1, (\not{D}_{S^1} \mp i(l_1+1/2))\chi_{l_1}^{(\pm)}(\phi) = 0$,

$$\begin{aligned} \psi_{\pm, nl_1}(\chi, \Omega_2) &= \phi_{nl}(\chi)\chi_{\pm, ll_1}^{(-)}(\Omega_2) \pm i\psi_{nl}(\chi)\chi_{\pm, ll_1}^{(+)}(\Omega_2), \\ [\chi_{\pm, ll_1}^{(-)}(\Omega_2) &= \begin{pmatrix} \phi_{ll_1}(\theta) \\ \pm i\psi_{ll_1}(\theta) \end{pmatrix} \chi_{l_1}^{(-)}(\phi), \chi_{\pm, ll_1}^{(+)} = \begin{pmatrix} i\psi_{ll_1}(\theta) \\ \pm \phi_{ll_1}(\theta) \end{pmatrix} \chi_{l_1}^{(+)}(\phi), \chi_{l_1}^{(\pm)}(\phi) = e^{\pm i(l_1+1/2)\phi}] \end{aligned} \quad (\text{A.10})$$

where the \pm lower indices in ψ_{\pm, ll_1} and χ_{\pm, ll_1} label the independent solutions while the upper indices (\pm) refer to the S^2 chiralities. The integer angular momenta for the harmonics of S^3, S^2, S^1 satisfy $n \geq l \geq l_1, [n = 0, 1, \dots]$ with degeneracies $D_3(n) = (n+1)(n+2), D_2(l) = 2(l+1), D_1(l_1) = 1$ and the scalar harmonic functions ϕ_{nl}, ψ_{nl} are expressed in terms of Jacobi polynomials.

2. Reduced bosonic action

The induced metric on the $D7$ -brane world volume of intrinsic coordinates ξ^α is the pull-back transform of the target spacetime metric through the directional derivatives of the coordinates $X^M(\xi)$, defining the world volume embedding in spacetime,

$$ds^2(D7) = \gamma_{\alpha\beta} d\xi^\alpha d\xi^\beta, [\gamma_{\alpha\beta} = \frac{\partial X^M}{\partial \xi^\alpha} \frac{\partial X^N}{\partial \xi^\beta} g_{MN}]. \quad (\text{A.11})$$

The invariance under coordinates diffeomorphisms allows choosing the static gauge in which the intrinsic coordinates are identified to subset of spacetime coordinates tangential to the world volume, $\xi^\alpha = X^\alpha(\xi) = [\rho, (\hat{h}_3, \hat{h}_1, \hat{h}_2)]$. The remaining coordinates along transversal directions $X^u(\xi)$ are interpreted as brane moduli fields $\chi(\xi)$ and are assigned classical values (VEVs) describing the brane embedding in spacetime. The induced metric

$$ds^2(D7) = \gamma_{\mu\nu} dx^{\mu\nu} + \gamma_{\rho\rho} d\rho^2 + 2\gamma_{h_i h_j} \hat{h}_i \hat{h}_j + 2\gamma_{\rho h_i} \hat{h}_i d\rho, \quad (\text{A.12})$$

can also be directly evaluated by substituting the decomposition $d\chi = \partial_\mu \chi dx^\mu + \partial_\rho \chi d\rho + \partial_{h_i} \chi h_i$ into Eq. (A.1) for the spacetime metric. We consider for definiteness the induced warped metric. This consists of two additive parts, $\gamma_{\alpha\beta}(\chi) \equiv \gamma_{\alpha\beta}^{(0)} + \alpha_{\alpha\beta} = g_{\alpha\beta} + \alpha_{\alpha\beta}$, the first coinciding with the spacetime metric restriction to the brane world volume and the second involving directional derivatives of the scalar moduli fields $\partial_{\mu, \rho, \hat{h}_i} \chi$. Both parts are functions of χ and μ_χ , the first $g_{\alpha\beta}$ being of diagonal form in the present case and the second $\alpha_{\alpha\beta}$ of non-diagonal form,

$$\begin{aligned} \gamma_{\mu\nu} &= \gamma_{\mu\nu}^{(0)} + \alpha_{\mu\nu} = h^{-1/2} \tilde{g}_{\mu\nu} + 2h^{1/2} \tilde{g}_{\chi\bar{\chi}} \partial_\mu \chi \partial_\nu \bar{\chi}, \\ \gamma_{\alpha\nu} &= \gamma_{\nu\alpha} = \alpha_{\nu\alpha} = h^{1/2} (\tilde{g}_{\alpha\chi} \partial_\nu \chi + \tilde{g}_{\alpha\bar{\chi}} \partial_\nu \bar{\chi} + \tilde{g}_{\chi\bar{\chi}} (\partial_\nu \chi \partial_\alpha \bar{\chi} + \partial_\nu \bar{\chi} \partial_\alpha \chi)), [\alpha = \rho, \hat{h}_3], \\ \gamma_{\rho\rho} &= \gamma_{\rho\rho}^{(0)} + \alpha_{\rho\rho} = h^{1/2} (\tilde{g}_{\rho\rho} + 2\tilde{g}_{\rho\chi} \partial_\rho \chi + 2\tilde{g}_{\rho\bar{\chi}} \partial_\rho \bar{\chi} + 2\tilde{g}_{\chi\bar{\chi}} \partial_\rho \chi \partial_\rho \bar{\chi}), \\ \gamma_{\rho h_i} &= h^{1/2} \tilde{g}_{\rho h_i} + h^{1/2} (2\tilde{g}_{h_i \chi} \partial_\rho \chi + 2\tilde{g}_{h_i \bar{\chi}} \partial_\rho \bar{\chi}), \\ \gamma_{\hat{h}_i \hat{h}_j} &= h^{1/2} (\tilde{g}_{\hat{h}_i \hat{h}_j} + 2\tilde{g}_{\chi h_i} \partial_{h_j} \chi + 2\tilde{g}_{\bar{\chi} h_j} \partial_{h_j} \bar{\chi} + 2\tilde{g}_{\chi\bar{\chi}} \partial_{h_i} \chi \partial_{h_j} \bar{\chi}), [\tilde{g}_{\rho h_i} = 0, \tilde{g}_{\hat{h}_i \hat{h}_j} = 0]. \end{aligned} \quad (\text{A.13})$$

The linear and quadratic order contributions in $d\chi$ and η_χ are given by the explicit formulas for the warped metric components

$$\begin{aligned} \alpha_{\mu\nu} &= 2F_1 h^{1/2} (\tau) A_3 \partial_\mu \chi \partial_\nu \bar{\chi}, \alpha_{\rho\rho} = F_1 h^{1/2} (2A_3 \partial_\rho \bar{\chi} \partial_\rho \chi + \bar{A}_2 \eta_\chi S_\rho \partial_\rho \bar{\chi} + A_2 S_\rho \bar{\eta}_\chi \partial_\rho \chi), \\ \alpha_{h_3 h_3} &= F_1 h^{1/2} (2A_3 \partial_{h_3} \chi \partial_{h_3} \bar{\chi} + i\bar{A}_2 \eta_\chi S_\rho \partial_{h_3} \bar{\chi} - iA_2 S_\rho \bar{\eta}_\chi \partial_{h_3} \chi), \\ \alpha_{h_2 h_2} &= 2A_3 F_1 h^{1/2} \partial_{h_1} \chi \partial_{h_1} \bar{\chi}, \alpha_{h_1 h_1} = 2A_3 F_1 h^{1/2} \partial_{h_2} \chi \partial_{h_2} \bar{\chi}, \\ \alpha_{\mu\rho} &= \frac{1}{2} F_1 h^{1/2} (2\bar{A}_3 \partial_\rho \bar{\chi} \partial_\mu \chi + 2A_3 \partial_\mu \bar{\chi} \partial_\rho \chi + \bar{A}_2 S_\rho \partial_\mu \bar{\chi} \eta_\chi + A_2 S_\rho \bar{\eta}_\chi \partial_\mu \chi), \\ \alpha_{\mu h_3} &= \frac{i}{2} F_1 h^{1/2} S_\rho (\bar{A}_2 \partial_\mu \bar{\chi} \eta_\chi - A_2 \bar{\eta}_\chi \partial_\mu \chi), \alpha_{\mu h_1} = A_3 F_1 h^{1/2} \partial_{h_1} \bar{\chi} \partial_\mu \chi, \alpha_{\mu h_2} = A_3 F_1 h^{1/2} \partial_{h_2} \bar{\chi} \partial_\mu \chi, \end{aligned}$$

$$\begin{aligned}
\alpha_{\rho h_3} &= \frac{i}{2} F_1 h^{1/2} S_\rho (\bar{A}_2 \partial_\rho \bar{\chi} \eta_\chi - A_2 \bar{\eta}_\chi \partial_\rho \chi), \quad \alpha_{\rho h_1} = A_3 F_1 h^{1/2} \partial_{h_1} \bar{\chi} \partial_\rho \chi, \quad \alpha_{\rho h_2} = A_3 F_1 h^{1/2} \partial_{h_2} \bar{\chi} \partial_\rho \chi, \\
\alpha_{h_3 h_1} &= A_3 F_1 h^{1/2} \partial_{h_3} \chi \partial_{h_1} \bar{\chi} - \frac{i}{2} A_2 h^{1/2} S_\rho \bar{\eta}_\chi F_1 \partial_{h_1} \chi, \quad \alpha_{h_1 h_2} = A_3 F_1 h^{1/2} \partial_{h_1} \chi \partial_{h_2} \bar{\chi}, \\
\alpha_{h_3 h_2} &= A_3 F_1 h^{1/2} \partial_{h_3} \chi \partial_{h_2} \bar{\chi} - \frac{i}{2} A_2 F_1 h^{1/2} S_\rho \bar{\eta}_\chi \partial_{h_2} \chi, \\
[\eta_\chi &= \frac{\mu_\chi}{X}, \quad A_1 = C_\rho + 2\eta_\chi \bar{\eta}_\chi S_\rho^2 R, \quad A_2 = X + 2\bar{\mu}_\chi \eta_\chi (1 + C_\rho \frac{X}{\bar{X}}) R, \quad A_3 = 1 + C_\rho X \bar{X} + 2\mu_\chi \bar{\mu}_\chi R |1 + C_\rho \frac{X}{\bar{X}}|^2] \quad (\text{A.14})
\end{aligned}$$

The action for the Kaluza-Klein modes of χ depends on the 4-d Minkowski spacetime derivatives $\partial_\mu \chi$ as well as the radial and angle derivatives, $\partial_\rho \chi$ and $\partial_{h_i} \chi$. For the singlet modes under the 4-cycle isometry group (as opposed to charged modes) only the radial derivatives are relevant. The square root of the induced metric determinant is evaluated as a power expansion in the fields derivatives by means of the familiar formula

$$\text{Det}^{1/2}(-(\gamma^{(0)} + \alpha)) = \text{Det}^{1/2}(-\gamma^{(0)}) \left(1 + \frac{1}{2} \text{Tr}(\gamma^{(0)-1} \alpha) - \frac{1}{4} \text{Tr}(\gamma^{(0)-1} \alpha \gamma^{(0)-1} \alpha) + \frac{1}{8} (\text{Tr}(\gamma^{(0)-1} \alpha))^2 + \dots \right) \quad (\text{A.15})$$

The $D7$ -brane bosonic action up to $O((d\chi)^4)$ is written below with the linear, quadratic, cubic and quartic order (square and cross) terms in $d\chi$ appearing in succession,

$$\begin{aligned}
S_B(D7) &= -\mu_7 \int d^4 x \sqrt{g_4^{(0)}} \int d\rho \sqrt{-\gamma^{(0)}} e^\phi \int \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 \\
&\times \left[1 + \left(\frac{S_\rho}{|\eta_\chi|^2 T_3} [A_2 \bar{\eta}_\chi (\partial_\rho \chi - i \partial_{h_3} \chi) - F_1 h A_3 \bar{A}_2 \eta_\chi (\partial_\mu \bar{\chi})^2 (\partial_\rho \chi + i \partial_{h_3} \chi)] + H. c. \right) \right. \\
&+ F_1 h(\tau) \left(A_3 - \frac{|A_2|^2 S_\rho^2}{T_3} \right) \hat{g}^{\mu\nu} \partial_\mu \bar{\chi} \partial_\nu \chi + \frac{2}{|\eta_\chi|^2 T_3} \left(A_3 - \frac{|A_2|^2 S_\rho^2}{T_3} \right) (\partial_\rho \bar{\chi} \partial_\rho \chi + \partial_{h_3} \bar{\chi} \partial_{h_3} \chi) \\
&+ \frac{4}{(1 + C_\rho) |\eta_\chi|^2} \left(A_3 - \frac{|A_2|^2 S_\rho^2}{T_3} \right) \partial_{h_1} \bar{\chi} \partial_{h_1} \chi + \frac{4}{(-1 + C_\rho) |\eta_\chi|^2} \left(A_3 - \frac{|A_2|^2 S_\rho^2}{T_3} \right) \partial_{h_2} \bar{\chi} \partial_{h_2} \chi + \left(\frac{2i |A_2|^2 S_\rho^2}{|\eta_\chi|^2 T_3^2} \partial_\rho \chi \partial_{h_3} \bar{\chi} + H. c. \right) \\
&+ \frac{1}{2 |\eta_\chi|^4 T_3^2} ((\partial_\rho \chi)^2 + (\partial_{h_3} \chi)^2) \left((4A_2 A_3 \bar{\eta}_\chi S_\rho (-\partial_\rho \chi + i \partial_{h_3} \chi) + H. c.) - 4A_3^2 ((\partial_\rho \bar{\chi})^2 + (\partial_{h_3} \bar{\chi})^2) \right) \\
&- \frac{1}{|\eta_\chi|^4} (\partial_{h_1} \chi)^2 \left[\frac{4A_2 A_3 \bar{\eta}_\chi S_\rho}{(1 + C_\rho) T_3} \partial_\rho \bar{\chi} + \frac{4A_3^2}{(1 + C_\rho) T_3} (\partial_\rho \bar{\chi})^2 + \frac{8A_3^2}{(1 + C_\rho)^2} (\partial_{h_1} \bar{\chi})^2 \right] \\
&- \frac{1}{|\eta_\chi|^4} (\partial_{h_2} \chi)^2 \left[\frac{4A_2 A_3 \bar{\eta}_\chi S_\rho}{(-1 + C_\rho) T_3} \partial_\rho \bar{\chi} + \frac{4A_3^2}{(-1 + C_\rho) T_3} (\partial_\rho \bar{\chi})^2 + \frac{8A_3^2}{(-1 + C_\rho)^2} (\partial_{h_2} \bar{\chi})^2 \right]. \quad (\text{A.16})
\end{aligned}$$

The cross angular terms, $(\partial_{h_i} \bar{\chi} \partial_{h_j} \chi)$, $[i \neq j]$ are absent owing the symmetry under the isometry group $SO(3) \sim SU(2)/Z_2$. Note also the absence at quadratic order of complex terms mixing the real and imaginary field components, $(\partial_\mu \chi)^2$, $(\partial_\rho \chi)^2$, $(\partial_{h_j} \chi)^2$.

The pull-backs of the Kähler, NSNS 2-form and complex 3-form fields J , B_2 , G_3 in Einstein frame are conveniently evaluated by substituting the expressions in Eq. (III.1) for the complex coordinates $(w_a) \in C^4$ in the $SO(4)$ invariant expressions of the Klebanov-Strassler classical solutions [66],

$$\begin{aligned}
\bullet J &= i \frac{K(\tau)}{\epsilon^{2/3}} \left[\delta^{ab} - \frac{1}{\epsilon^2 \sinh^2 \tau} \left(\cosh \tau - \frac{2}{3K^3(\tau)} \right) \bar{w}^a w^b \right] dw_a \wedge d\bar{w}_b, \\
\bullet B_2^{cl} &= ib(\tau) \epsilon_{abcd} w^a \bar{w}^b dw^c \wedge d\bar{w}^d, \quad [b(\tau) = \frac{g_s M \alpha'}{2|\epsilon|^4} \frac{(-1 + \tau \coth \tau)}{\sinh^2 \tau}] \\
\bullet G_3 &= G_{(2,1)} = F_3 - ie^{-\phi} H_3 = g_1(\tau) \epsilon_{abcd} w^a \bar{w}^b dw^c \wedge d\bar{w}^d \wedge \bar{w}_e dw^e + g_2(\tau) \epsilon_{abcd} w^a \bar{w}^b dw^c \wedge dw^d \wedge w_e d\bar{w}^e, \\
[g_1(\tau) &= \frac{M \alpha'}{2\epsilon^6 \sinh^4 \tau} \frac{\sinh 2\tau - 2\tau}{\sinh \tau}, \quad g_2(\tau) = \frac{M \alpha'}{2\epsilon^6 \sinh^4 \tau} 2(1 - \tau \coth \tau)]. \quad (\text{A.17})
\end{aligned}$$

The resulting formulas for the differential forms restriction to Σ_4 read

$$\begin{aligned}
\bullet J|_{\Sigma_4} &= \frac{|\eta_\chi|^2}{2\epsilon^{2/3}} K(\tau) \left[\left(\cosh \rho - \frac{|\eta_\chi|^2}{\epsilon^2 \sinh^2 \tau} \left(\cosh \tau - \frac{2}{3K^3(\tau)} \right) \sinh^2 \rho \right) d\rho \wedge \hat{h}_3 + \frac{1}{2} \sinh \rho \hat{h}_1 \wedge \hat{h}_2 \right] \\
&= \frac{|\eta_\chi|^2}{2\epsilon^{2/3}} K(\tau) [K_2(\tau) d\rho \wedge \hat{h}_3 + \frac{1}{2} \sinh \rho \hat{h}_1 \wedge \hat{h}_2], \\
\bullet B_2^{cl}|_{\Sigma_4} &= |\mu_\chi^2 Y|^2 b(\tau) \left[\frac{i}{2} (Y - \bar{Y}) \sinh^2 \frac{\rho}{2} \left(\cosh \frac{\rho}{2} d\rho \wedge \hat{h}_1 + \sinh \frac{\rho}{2} \hat{h}_3 \wedge \hat{h}_2 \right) \right. \\
&- \left. \frac{1}{2} (Y + \bar{Y}) \cosh^2 \frac{\rho}{2} \left(\sinh \frac{\rho}{2} d\rho \wedge \hat{h}_2 - \cosh \frac{\rho}{2} \hat{h}_3 \wedge \hat{h}_1 \right) \right],
\end{aligned}$$

$$\begin{aligned}
\bullet G_3|_{\Sigma_4} &= \frac{|\mu_\chi^2 Y|^2}{4} \sinh \rho [g_1(\tau) |\mu_\chi|^2 |Y|^2 [(Y - \bar{Y}) \sinh^2 \frac{\rho}{2} (\cosh \frac{\rho}{2} d\rho \wedge \hat{h}_1 + \sinh \frac{\rho}{2} \hat{h}_3 \wedge \hat{h}_2) \\
&+ i(Y + \bar{Y}) \cosh^2 \frac{\rho}{2} (\sinh \frac{\rho}{2} d\rho \wedge \hat{h}_2 - \cosh \frac{\rho}{2} \hat{h}_3 \wedge \hat{h}_1)] \wedge (d\rho + i\hat{h}_3) \\
&+ g_2(\tau) \mu_\chi^2 Y^2 (Y + \bar{Y} \cosh \rho) [\cosh \frac{\rho}{2} (d\rho + i\hat{h}_3) \wedge \hat{h}_1 + i \sinh \frac{\rho}{2} (d\rho + i\hat{h}_3) \wedge \hat{h}_2] \wedge (d\rho - i\hat{h}_3)]. \tag{A.18}
\end{aligned}$$

We see that $\hat{J} \equiv J|_{\Sigma_4}/2$ and $\mathcal{F} = B_2^{cl}|_{\Sigma_4} + F_2$ obey (for $F_2 = 0$ and the value $\theta = 0$ of the arbitrary parameter θ) the conditions set by the κ -symmetry [25],

$$\frac{1}{2}(\hat{J} \wedge \hat{J})|_{\Sigma_4} = \text{vol}(\Sigma_4), \quad (\hat{J} \wedge \mathcal{F})|_{\Sigma_4} = \frac{1}{2} \tan \theta (\hat{J} \wedge \mathcal{F} - \mathcal{F} \wedge \mathcal{F})|_{\Sigma_4}, \tag{A.19}$$

as expected [42] in the case of a calibrated (holomorphic, supersymmetry preserving) 4-cycle.

The pull-backs of the background solutions also include extra contributions depending on $d\chi$. The Kähler 2-form and the 3-form solutions acquire the correction terms,

$$\begin{aligned}
\delta J &= iK(\tau) \epsilon^{-2/3} \left[\frac{1}{2} (-\bar{\eta}_\chi X S_\rho (d\rho - i\hat{h}_3) \wedge d\chi + \eta_\chi \bar{X} S_\rho (d\rho + i\hat{h}_3) \wedge d\bar{\chi}) + (1 + X \bar{X} C_\rho) d\chi \wedge d\bar{\chi} \right] \\
&+ i \frac{K'(\tau) \epsilon^{-8/3}}{\sinh(\tau)} \left[-\frac{1}{2} (\mu_\chi \bar{\mu}_\chi^2 (1 + C_\rho) S_\rho (d\rho - i\hat{h}_3) \wedge d\chi - H. c.) + |\mu_\chi|^2 (1 + C_\rho)^2 d\chi \wedge d\bar{\chi} \right], \\
\delta G_3 &= g_1(\tau) [i|\mu_\chi|^4 \cosh^2 \frac{\rho}{2} (\sinh \frac{\rho}{2} d\rho \wedge \hat{h}_2 - \cosh \frac{\rho}{2} \hat{h}_3 \wedge \hat{h}_1)] \wedge ((\bar{\mu}_\chi + C_\rho \bar{\eta}_\chi) d\chi + \frac{|\eta_\chi|^2}{2} S_\rho (d\rho + i\hat{h}_3)) \\
&+ g_2(\tau) \left[-\frac{1}{2} |\eta_\chi|^2 \mu_\chi (X - Y) \frac{S_\rho^2}{\sinh \frac{\rho}{2}} \hat{h}_1 \wedge d\chi - i|\eta_\chi|^2 \mu_\chi (X - Y) \sinh \frac{\rho}{2} S_\rho^2 \hat{h}_2 \wedge d\chi \right] \wedge ((\mu_\chi + C_\rho \eta_\chi) d\chi + \frac{|\eta_\chi|^2}{2} S_\rho (d\rho - i\hat{h}_3)). \tag{A.20}
\end{aligned}$$

The additional terms in $\delta B_{\alpha\beta} = B_{\alpha\beta} - B_{\alpha\beta}|_{\Sigma_4} = 2B_{\alpha\chi} \partial_\beta \chi + 2B_{\alpha\bar{\chi}} \partial_\beta \bar{\chi} + 2B_{\chi\bar{\chi}} \partial_\alpha \chi \partial_\beta \bar{\chi}$ are found to vanish in the limit $\mu \gg \epsilon$ (in which $X \simeq 1/Y \rightarrow 1$) and hence will not be quoted.

We now explain how to include the contributions from the classical fields B_2^{cl} , F_2^{cl} components along Σ_4 present in (Einstein frame) Klebanov-Strassler solution. The dependence enters through the (unwarped) effective metric $\tilde{q}_{\alpha\beta} = \tilde{\gamma}_{\alpha\beta} + e^{-\phi/2} h^{-1/2} \mathcal{F}_{\alpha\beta}$, $[\mathcal{F}_{\alpha\beta} = B_{\alpha\beta} + F_{\alpha\beta}]$ where the index $\alpha = (\mu = 0, 1, 2, 3, a = 1, 2, 3, 4)$ labels directions along M_4 and Σ_4 in the ordering convention adopted for the tangent space basis of Σ_4 : $(d\rho, \hat{h}_3, \hat{h}_1, \hat{h}_2)$. The matrix \tilde{q} is block diagonal with one sub-block $\tilde{g}_{\mu\nu}$ in M_4 while the other sub-block $\tilde{q} = \tilde{\gamma} + b$, $[b = e^{-\phi/2} h^{-1/2} B_2^{cl}]$ along Σ_4 is the sum of a symmetric (diagonal) matrix $\tilde{\gamma}_{ab}$ and an antisymmetric (anti-diagonal) matrix b_{ab} ,

$$\begin{aligned}
\tilde{\gamma} &= \text{diag}(\gamma_1, \gamma_2, \gamma_3, \gamma_4), \quad b = \overline{\text{diag}}(b_1, b_2, -b_2, -b_1), \quad [\gamma_{a=1, \dots, 4} = \frac{1}{2} F_1 |\eta|^2 [A_1, A_1, \frac{1}{2}(1 + C_\rho), \frac{1}{2}(-1 + C_\rho)], \\
b_{[1,2]} &= -e^{-\phi/2} h^{-1/2} b_0(\tau) [\sinh \frac{\rho}{2}, -\cosh \frac{\rho}{2}], \quad b_0(\tau) = \frac{|\eta_\chi|^4 |X|^2}{4} (1 + C_\rho) b(\tau), \quad b(\tau) = \frac{g_s M \alpha'}{2\epsilon^4} \frac{(-1 + \tau \coth \tau)}{\sinh^2 \tau}. \tag{A.21}
\end{aligned}$$

In the convenient matrix notation in the tangent space of Σ_4 , \tilde{q} and its inverse \tilde{q}^{-1} are both given by a sum of diagonal and anti-diagonal (antisymmetric) matrices,

$$\begin{aligned}
\tilde{q} = \tilde{\gamma} + b &= \begin{pmatrix} \gamma_1 & 0 & 0 & b_1 \\ 0 & \gamma_2 & b_2 & 0 \\ 0 & -b_2 & \gamma_3 & 0 \\ -b_1 & 0 & 0 & \gamma_4 \end{pmatrix} \implies \tilde{q}^{-1} = (\tilde{\gamma} + b)^{-1} = \begin{pmatrix} g_1 & 0 & 0 & -\beta_1 \\ 0 & g_2 & -\beta_2 & 0 \\ 0 & \beta_2 & g_3 & 0 \\ \beta_1 & 0 & 0 & g_4 \end{pmatrix}, \\
[g_a]_{a=1,4} &= \frac{1}{\gamma_i R_i}, \quad R_1 = R_4 = 1 + \frac{b_1^2}{\gamma_1 \gamma_4}, \quad R_2 = R_3 = 1 + \frac{b_2^2}{\gamma_2 \gamma_3}, \quad \beta_i|_{i=1,2} = \frac{1}{b_i \rho_i}, \quad \rho_1 = 1 + \frac{\gamma_1 \gamma_4}{b_1^2}, \quad \rho_2 = 1 + \frac{\gamma_2 \gamma_3}{b_2^2}, \\
\tilde{q} \equiv \text{Det}(\tilde{q}) &= (b_2^2 + \gamma_2 \gamma_3)(b_1^2 + \gamma_1 \gamma_4) \tilde{\gamma} + b + b_1^2 \gamma_2 \gamma_3 + b_2^2 \gamma_1 \gamma_4, \\
\tilde{\gamma} \equiv \text{Det}(\tilde{\gamma}) &= \gamma_1 \gamma_2 \gamma_3 \gamma_4, \quad b \equiv \text{Det}(b) = (Pf(b))^2 = (b_1 b_2)^2. \tag{A.22}
\end{aligned}$$

Substituting the expressions in Eq. (A.21) for the matrices entries, γ_a , b_i and g_a , β_i , $[a = 1, \dots, 4, i = 1, 2]$ specifies the parameters, $R_1 = R_2 = 1 + \frac{\sqrt{b}}{\sqrt{\gamma}}$, $\rho_1 = \rho_2 = 1 + \frac{\sqrt{\tilde{q}}}{\sqrt{b}}$, hence yielding the simplified formulas for the matrix determinant, $\sqrt{\tilde{q}} = \sqrt{\tilde{\gamma}} + \sqrt{b}$, and for the inverse matrix entries,

$$\begin{aligned}
g_a &= \frac{1}{R_S \gamma_a} \implies \tilde{q}^{\rho\rho} = \frac{1}{R_S \gamma_1}, \quad \tilde{q}^{\hat{h}_3 \hat{h}_3} = \frac{1}{R_S \gamma_2}, \quad \tilde{q}^{\hat{h}_1 \hat{h}_1} = \frac{1}{R_S \gamma_3}, \quad \tilde{q}^{\hat{h}_2 \hat{h}_2} = \frac{1}{R_S \gamma_4}, \quad [R_S = \frac{\sqrt{\tilde{q}}}{\sqrt{\tilde{\gamma}}}] \\
\beta_i &= -\frac{1}{R_A b_i} \implies \tilde{q}^{[\rho \hat{h}_2]} = -\frac{1}{R_A b_1}, \quad \tilde{q}^{[\hat{h}_3 \hat{h}_1]} = -\frac{1}{R_A b_2}, \quad [R_A = \frac{\sqrt{\tilde{q}}}{\sqrt{b}}]. \tag{A.23}
\end{aligned}$$

The effective metric and its inverse along $M_4 \oplus \Sigma_4$ can be represented in a general (basis independent) way by block diagonal matrices $\tilde{q}_{\alpha\beta} = [\tilde{\gamma}_{\mu\nu}, \tilde{q}_{ab}]$, $\tilde{q}^{\alpha\beta} = [\tilde{\gamma}^{\mu\nu}, \tilde{q}^{ab}]$, satisfying the relations [42],

$$\begin{aligned} Det(\tilde{q}) &= (\sqrt{Det(\tilde{\gamma})} + h^{-1}\sqrt{Det(B)})^2, \quad \tilde{q}^{(\alpha\beta)} = \frac{\sqrt{\tilde{\gamma}}}{\sqrt{\tilde{q}}}\gamma^{\alpha\beta} = \frac{\gamma^{\alpha\beta}}{R_S}, \quad \tilde{q}^{[\alpha\beta]} = -\frac{\sqrt{b}}{\sqrt{-\tilde{q}}}B^{\alpha\beta} = -\frac{B^{\alpha\beta}}{R_A}, \\ [\sqrt{\tilde{\gamma}} = \frac{1}{2}(\frac{1}{2}F_1|\eta|^2)^2 A_1 \sinh \rho, \quad \sqrt{Det(B)} = Pf(B) = \frac{1}{2}b_0^2 \sinh \rho, \quad R_S = 1 + \frac{\sqrt{b}}{\sqrt{\tilde{\gamma}}}, \quad R_A = 1 + \frac{\sqrt{\tilde{\gamma}}}{\sqrt{b}}] \end{aligned} \quad (\text{A.24})$$

where the indices enclosed between pairs of parentheses or brackets are to be symmetrized or antisymmetrized. One finds for a constant dilaton profile,

$$R_S = 1 + \frac{Pf(B)}{g_s h(\tau) V_0(\rho)} = 1 + \frac{S_\rho b_0^2(\tau)}{g_s h(\tau) V(\rho)} = 1 + 2^{4/3} |L|^4 g_s \frac{\cosh^4(\rho/2)}{I(\tau) K^2(\tau) T_3(\rho)} \left(\frac{-1 + \tau \coth \tau}{\sinh^2 \tau} \right)^2. \quad (\text{A.25})$$

The effective metric determinant in Eq. (III.17) for the χ -field action,

$$V_0^2(\rho) = -Det(\tilde{q})_{M_8} = -Det(\tilde{\gamma})_{M_8} \times Det(\tilde{q})_{\Sigma_4}, \quad [(\tilde{q})_{M_8} = diag((\tilde{\gamma})_{M_4}, (\tilde{q})_{\Sigma_4}), (\tilde{\gamma})_{\Sigma_4} = (\tilde{\gamma} + b)|_{\Sigma_4}] \quad (\text{A.26})$$

shows that the contributions from B_2^{cl} may be included by replacing $V_0(\rho) = \sqrt{-\tilde{\gamma}^{(0)}} \rightarrow \sqrt{-\tilde{q}} = R_S \sqrt{-\tilde{\gamma}^{(0)}}$ and adding the linear and quadratic order corrections in $\partial_{\mu,\rho}\chi$ from $Tr(\tilde{q}^{-1}\alpha)$ and $Tr(\tilde{q}^{-1}\alpha\tilde{q}^{-1}\alpha)$. In Eq. (III.21), for instance, the coefficient $(T_1 - T_2^2/T_3)$ of $|\partial_{\mu}\chi|^2$ is replaced by $(T_1 - T_2^2/(R_S T_3))$ and the coefficient $(T_1 - T_2^2/T_3)/T_3$ of $|\partial_{\rho}\chi|^2$ is replaced by $R_S^{-1}(T_1 - T_2^2/(T_3 R_S))/T_3$, noting that the first and second terms in these coefficients arise from linear and quadratic order corrections. The wave equations for χ modes are modified likewise by replacing in Eq. (III.21),

$$Q(\rho) \rightarrow F_1 h(\tau) (A_3 - \frac{|A_2 S_\rho|^2}{A_1 R_S}), \quad P(\rho) \rightarrow (A_3 - \frac{|A_2 S_\rho|^2}{A_1 R_S})/A_1. \quad (\text{A.27})$$

The contributions from the classical 2-form gauge fluxes are conveniently included by noting the formal similarity of F_2 with B_2 . For instance, the (1,1)-form solution derived in [41], $F_{II} = -i\lambda\epsilon_{ijk}\tilde{w}_i dw_j \wedge d\tilde{w}_k + H.c.$ is of same form as the background solution for the NSNS field B_2^{cl} , so its contribution is found by replacing $b(\tau) \rightarrow \lambda/\mu_\chi$. The comparison with Eq. (A.17) yields the explicit expression for the gauge flux 2-form in Σ_4

$$F_{II} = (f_1 d\rho \wedge \hat{h}_2 + f_2 \hat{h}_3 \wedge \hat{h}_1), \quad [f_{[1,2]} = \lambda|\mu_\chi|^2 \tilde{\mu}_\chi \cosh^2 \frac{\rho}{2} [-\sinh \frac{\rho}{2}, \cosh \frac{\rho}{2}]] \quad (\text{A.28})$$

which is represented in the antisymmetric part of the effective metric tensor by the anti-diagonal matrix, $F_{II} = \overline{diag}(f_1, f_2, -f_2, -f_1)$.

3. Reduced Dirac operator

We here examine the Dirac operator on the 4-cycle $\Sigma_4 \subset \mathcal{C}_6$ restricted to the part of the covariant derivative depending on the spin connection along the brane longitudinal directions, $\mathcal{D}|_{\Sigma_4} = (\partial + \frac{1}{4}\omega^{\hat{a}\hat{b}}\Gamma_{\hat{a}\hat{b}})$. The solution of the flat torsion condition, $0 = D_\mu e_\nu^{\hat{a}} = (D_\mu e)_\nu^{\hat{a}} - \Gamma_{\mu\nu}^\lambda e_\lambda^{\hat{a}}, [(D_\mu e)_\nu^{\hat{a}}] = \partial_\nu e_\mu^{\hat{a}} + \omega_\mu^{\hat{a}\hat{b}} e_{\nu,\hat{b}}$ provides the explicit formula for the spin connection in terms of the tangent frame vectors of Σ_4 ,

$$\omega_\mu^{\hat{a}\hat{b}} = -e^{\nu,\hat{b}}(\partial_\mu e_\nu^{\hat{a}} - \Gamma_{\mu\nu}^\lambda e_\lambda^{\hat{a}}) = \frac{1}{2}[e^{\nu\hat{a}}(\partial_\mu e_\nu^{\hat{b}} - \partial_\nu e_\mu^{\hat{b}}) - e^{\nu\hat{b}}(\partial_\mu e_\nu^{\hat{a}} - \partial_\nu e_\mu^{\hat{a}}) - e^{\lambda\hat{a}} e^{\sigma\hat{b}} e_{\mu\hat{c}}(\partial_\lambda e_\sigma^{\hat{c}} - \partial_\sigma e_\lambda^{\hat{c}})] \quad (\text{A.29})$$

where $e_\mu^{\hat{a}}(\rho, \hat{h}_k)$ carry vector indices along curved and flat space directions, $\mu = (\rho, m)$ and $\hat{a} = (\hat{\rho}, \hat{k})$. In the static gauge, $\gamma_{\alpha\beta} \rightarrow g_{\alpha\beta}$, the non vanishing components of the spin connection are evaluated from the general formulas,

$$\begin{aligned} \omega_\rho^{\hat{\rho}\hat{k}} &= 0, \quad \omega_\rho^{\hat{j}\hat{k}} = \frac{1}{2}H^{\hat{j}\hat{k}}(\eta^{\hat{j}}\eta^{\hat{k}}\partial_\rho\eta^{\hat{j}}\partial_\rho\eta^{\hat{k}}) = 2(A_{\hat{j}\hat{k}} - A_{\hat{k}\hat{j}}), \\ \omega_m^{\hat{\rho}\hat{k}} &= -\frac{1}{2}e^{\rho\hat{\rho}}(\partial_\rho e_m^{\hat{k}} + e_{m\hat{c}}\partial_\rho e_n^{\hat{c}}e_n^{\hat{k}}) = -\frac{1}{2}(e^{\rho\hat{\rho}}h_m^{\hat{k}}\partial_\rho\eta^{\hat{k}} + C_{m\hat{k}}) \\ \omega_m^{\hat{j}\hat{k}} &= e^{n\hat{j}}\partial_{[m}e_{n]}^{\hat{k}} - e^{n\hat{k}}\partial_{[m}e_{n]}^{\hat{j}} - e^{n\hat{j}}e^{p\hat{k}}e_{m\hat{c}}\partial_{[n}e_{p]}^{\hat{c}} = \frac{1}{2}X_{m,\hat{j}\hat{k}}, \\ [e^{n\hat{j}} &= \eta^{\hat{j}}(\rho)h^{n\hat{j}}(\alpha), \quad A_{\hat{j}\hat{k}} = \frac{1}{4}H^{\hat{j}\hat{k}}\eta^{\hat{j}}\partial_\rho\eta^{\hat{k}}, \quad C_{m\hat{k}} = e^{\rho\hat{\rho}}\eta^{\hat{k}}\eta_{\hat{j}}\partial_\rho\eta^{\hat{j}}h_{m\hat{i}}H^{\hat{i}\hat{k}}, \quad H^{\hat{j}\hat{k}} = h_n^{\hat{j}}h_n^{\hat{k}} = \tilde{g}^{n\hat{n}'}h_n^{\hat{j}}h_n^{\hat{k}}, \end{aligned}$$

$$X_{m,\hat{j}\hat{k}} = 2\left[\frac{\eta^{\hat{k}}}{\eta^{\hat{j}}}h_{\hat{j}}^n\partial_{[m}h_{\hat{n}]}^{\hat{k}} - \eta^{\hat{j}}\eta^{\hat{k}}h_{\hat{k}}^n\partial_{[m}h_{\hat{n}]}^{\hat{j}} - \frac{\eta^{\hat{l}2}}{\eta^{\hat{j}}\eta^{\hat{k}}}h_{\hat{j}}^n h_{\hat{k}}^p h_{\hat{m}}^{\hat{l}}\partial_{[n}h_{\hat{p}]}^{\hat{l}}\right]. \quad (\text{A.30})$$

The gamma matrices identities, $\Gamma_{\hat{\rho}}\Gamma_{\hat{l}}\Gamma_{\hat{\rho}\hat{k}} = -\Gamma_{\hat{l}}\Gamma_{\hat{k}}$, $\Gamma_{\hat{l}}\Gamma_{\hat{j}\hat{k}} = i\Gamma^{\hat{\rho}}\tilde{\gamma}_{(4)}\epsilon_{\hat{\rho}\hat{j}\hat{k}\hat{l}} + \delta_{\hat{j}\hat{l}}\Gamma_{\hat{k}} - \delta_{\hat{k}\hat{l}}\Gamma_{\hat{j}}$, yield the simplified form of the Dirac operator,

$$\begin{aligned} \tilde{D}|_{\Sigma_4} &\equiv \Gamma^{\hat{\rho}}(\partial_{\hat{\rho}} + \omega_{\hat{\rho}}) + \Gamma^m(\partial_m + \omega_m) \\ &= e_{\hat{\rho}}^{\hat{\rho}}\Gamma^{\hat{\rho}}(\partial_{\hat{\rho}} + \frac{1}{2}\eta_{\hat{k}}\partial_{\hat{\rho}}\eta^{\hat{k}} + \Gamma_{\hat{j}\hat{k}}A_{\hat{j}\hat{k}}) + e_{\hat{l}}^m\Gamma^{\hat{l}}(\partial_m - \frac{1}{4}\Gamma_{\hat{\rho}\hat{k}}C_{m\hat{k}} + \frac{1}{8}\Gamma_{\hat{j}\hat{k}}X_{m,\hat{j}\hat{k}}) \\ &= e_{\hat{\rho}}^{\hat{\rho}}\gamma_{(4)} \otimes \tilde{\gamma}^{\hat{\rho}}[\partial_{\hat{\rho}} + \frac{1}{2}\partial_{\hat{\rho}}\ln(\eta^{\hat{1}}\eta^{\hat{2}}\eta^{\hat{3}}) - i\epsilon_{\hat{\rho}\hat{j}\hat{k}\hat{l}}\gamma_{(4)} \otimes \tilde{\gamma}^{\hat{l}}\tilde{\gamma}_{(4)}A_{\hat{j}\hat{k}} \\ &\quad + \eta^{\hat{\rho}}e_{\hat{l}}^m(\tilde{\gamma}_{\hat{\rho}\hat{l}}e_{\hat{l}}^m\partial_m + \frac{1}{4}C_{m,\hat{k}}\tilde{\gamma}_{\hat{l}\hat{k}} + \frac{1}{8}X_{m,\hat{j}\hat{k}}(i\epsilon_{\hat{\rho}\hat{j}\hat{k}\hat{l}}\tilde{\gamma}_{(4)} + \delta_{\hat{j}\hat{l}}\tilde{\gamma}_{\hat{\rho}\hat{k}} - \delta_{\hat{k}\hat{l}}\tilde{\gamma}_{\hat{\rho}\hat{j}}))]. \end{aligned} \quad (\text{A.31})$$

Applying these formulas to the deformed conifold case, using Eq. (A.5) and the expressions implied by the $SO(3)$ isometry of Σ_4 ,

$$\begin{aligned} e_{\hat{m}}^{\hat{l}} &= \eta^{\hat{l}}h_{\hat{m}}^{\hat{l}}, \quad e_{\hat{l}}^m = \eta_{\hat{l}}h_{\hat{l}}^m, \quad g^{m'n'} = \sum_{\hat{l}} e_{\hat{l}}^n e_{\hat{l}}^{n'}, \quad H^{\hat{j}\hat{k}} = \delta_{\hat{j}\hat{k}}\eta_{\hat{j}}^2, \quad \sum_m h_{\hat{m}}^l h_{\hat{l}}^m = \delta_{\hat{l}\hat{l}'}, \\ g_{\hat{j}\hat{k}} &= \delta_{\hat{j}\hat{k}}\eta_{\hat{j}}^2, \quad g^{\hat{j}\hat{k}} = \delta_{\hat{j}\hat{k}}\eta_{\hat{j}}^2, \quad h^{m\hat{j}} = \eta_{\hat{j}}^2 h_{\hat{j}}^m, \quad h_{m\hat{j}} = \eta^{\hat{j}2} h_{\hat{j}}^m, \end{aligned} \quad (\text{A.32})$$

one finds that $C_{m\hat{k}}$ and $A_{\hat{j}\hat{k}} = -A_{\hat{k}\hat{j}}$ vanish identically while $e_{\hat{l}}^m X_{m,\hat{j}\hat{k}}$ is independent of angles. The resulting Dirac operator

$$\tilde{D}|_{\Sigma_4} = \frac{1}{\eta^{\hat{\rho}}}\Gamma^{\hat{\rho}}(\partial_{\hat{\rho}} + \frac{1}{2}\sum_{\hat{k}}\partial_{\hat{\rho}}(\ln\eta^{\hat{k}}) + \frac{\eta^{\hat{\rho}}}{8\eta^{\hat{l}}}h_{\hat{l}}^m X_{m,\hat{j}\hat{k}}\Gamma_{\hat{\rho}}\Gamma_{\hat{l}}\Gamma_{\hat{j}\hat{k}} + \eta^{\hat{\rho}}\Gamma_{\hat{\rho}}\Gamma_{\hat{k}}e_{\hat{k}}^m\partial_m), \quad (\text{A.33})$$

includes on side of the first radial derivative term, angle independent curvature contributions in the second and third terms (which commute with the chirality matrix Γ_{Σ_4}) and the base manifold Dirac operator $\Gamma_{\hat{k}}e_{\hat{k}}^m\partial_m = \Gamma^{\hat{k}}\nabla_{\hat{h}_{\hat{k}}} = \tilde{D}$ (S^3/Z_2) in the fourth term.

4. Undeformed conifold limit

The useful intermediate results in the undeformed conifold case are obtained from the deformed conifold results by taking the limit $\epsilon \rightarrow 0$, $\tau \rightarrow \infty$ at fixed radial conic variable r and μ , absorbing the dependence on ϵ through the change of radial variable $\tau \rightarrow r$ and of the parameter $L \rightarrow \mu = L\epsilon$. The correspondence between the 4-cycle radial variable ρ and the conifold radial variable r is given by

$$\hat{r}^3 \equiv (2/3)^{3/2}r^3 \equiv |\mu_{\chi}|^2 r_{\chi}^3 = |\mu_{\chi}|^2(1 + \cosh\rho), \quad \frac{dr_{\chi}}{d\rho} = \frac{1}{3}\left(\frac{r_{\chi}^3 - 2}{r_{\chi}}\right)^{1/2}, \quad (\text{A.34})$$

where we introduced the useful rescaled and dimensionless radial variables, \hat{r} and r_{χ} . Note that the change of variable $\rho \rightarrow r_{\chi}$ picks up the factor $\frac{d\rho}{dr_{\chi}}$ which diverges at the 4-cycle apex, $\hat{r}_{min} \equiv (2/3)^{1/2}r_{min} \equiv |\mu^2|^{1/3}(r_{\chi})_{min} = |2\mu^2|^{1/3}$. The coefficient functions $F_{1,2}$ and the warp profile have the limiting forms,

$$\begin{aligned} F_1 &\simeq (2\epsilon)^{-2/3}e^{-\tau/3} \simeq \frac{3^{1/2}2^{-3/2}}{r} = \frac{1}{2|\mu_{\chi}|^{2/3}r_{\chi}}, \quad R = \frac{F_2}{F_1} \simeq -\frac{e^{-\tau}}{3\epsilon^2} \simeq -\frac{1}{6|\mu_{\chi}|^2(1 + \cosh\rho)} = -\frac{3^{1/2}2^{-5/2}}{r^3} = -\frac{1}{6|\mu_{\chi}|^2r_{\chi}^3}, \\ \sqrt{-\gamma^{(0)}} &\simeq \frac{|\mu_{\chi}|^4}{64r^2}S_{\rho}(1 + 2C_{\rho}) = \frac{|\mu_{\chi}|^{8/3}}{96r_{\chi}^2}(r_{\chi}^3(r_{\chi}^3 - 2))^{1/2}(2r_{\chi}^3 - 1), \quad h(r) = \frac{L_{eff}^4}{r^4}, \quad [L_{eff}^4 = \frac{81}{8}(g_s M \alpha')^2 \ln \frac{r}{r_{ir}}] \end{aligned} \quad (\text{A.35})$$

where the effective curvature radius $L_{eff}(r)$ was expressed in terms of the infrared cutoff radius r_{ir} at which $L_{eff}(r_{ir}) = 0$. The conifold parameterization as a foliation by Kuperstein 4-cycle Σ_4 is described by the metric

$$\begin{aligned} d\hat{s}^2(\mathcal{C}_6) &= 2^{-5/2}3^{1/2}\frac{|\mu_{\chi}|^2}{r}\left[\frac{(1 + 2\cosh\rho)}{3}(d\rho^2 + \hat{h}_3^2) + \cosh^2(\rho/2)\hat{h}_1^2 + \sinh^2(\rho/2)\hat{h}_2^2\right. \\ &\quad \left.+ \frac{4}{3}(1 + \cosh\rho)d\chi d\bar{\chi} + \left(\frac{2}{3}\mu_{\chi}\sinh\rho(d\rho + i\hat{h}_3)d\bar{\chi} + H. c.\right)\right] \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{8\hat{r}}[\hat{r}^3\hat{h}_1^2 + (\hat{r}^3 - 2|\mu_\chi|^2)\hat{h}_2^2 + \frac{2}{3}(2\hat{r}^3 - |\mu_\chi|^2)\hat{h}_3^2 + 6\frac{\hat{r}(2\hat{r}^3 - |\mu_\chi|^2)}{(\hat{r}^3 - 2|\mu_\chi|^2)}d\hat{r}^2 \\
&+ \frac{8}{3}\hat{r}^3d\chi d\bar{\chi} + (\frac{4}{3}(\hat{r}^3(\hat{r}^3 - 2|\mu_\chi|^2))^{1/2}(3(\frac{\hat{r}}{\hat{r}^3 - 2|\mu_\chi|^2})^{1/2}d\hat{r} + i\hat{h}_3)d\bar{\chi} + H. c.].
\end{aligned} \tag{A.36}$$

We quote for convenience the metric tensor components,

$$\begin{aligned}
\tilde{g}_{\rho\rho} &= \tilde{g}_{\hat{h}_3\hat{h}_3} = 2^{-5/2}3^{1/2}\frac{|\eta_\chi|^2}{3r}(1 + 2\cosh\rho) = \frac{|\mu_\chi|^2}{12\hat{r}}(\frac{2\hat{r}^3}{|\mu_\chi|^2} - 1), \\
[\tilde{g}_{\hat{h}_1\hat{h}_1}, \tilde{g}_{\hat{h}_2\hat{h}_2}] &= \frac{2^{-7/2}3^{1/2}|\eta_\chi|^2}{r}[\pm 1 + \cosh\rho] = \frac{|\mu_\chi|^2}{8\hat{r}}[\frac{\hat{r}^3}{|\mu_\chi|^2} - 2, \frac{\hat{r}^3}{|\mu_\chi|^2}], \\
\tilde{g}_{\chi\bar{\chi}} &= \frac{\hat{r}^2}{6}, \quad \tilde{g}_{\hat{r}\bar{\chi}} = \frac{\hat{r}^2}{2}, \quad \tilde{g}_{\hat{h}_3\bar{\chi}} = -\frac{i}{6}(\hat{r}(\hat{r}^3 - 2|\mu_\chi|^2))^{1/2}.
\end{aligned} \tag{A.37}$$

The conifold metric restriction to Σ_4 is given by

$$\begin{aligned}
d\tilde{s}^2(\mathcal{C}_6)|_{\Sigma_4} &= \frac{|\mu_\chi|^{4/3}}{8r_\chi}(r_\chi^3\hat{h}_1^2 + (r_\chi^3 - 2)\hat{h}_2^2 + \frac{2}{3}(2r_\chi^3 - 1)(\hat{h}_3^2 + 9(\frac{r_\chi}{r_\chi^3 - 2})dr_\chi^2)), \\
&= \frac{3}{4}[\frac{2\hat{r}^3 - |\mu|^2}{\hat{r}^3 - 2|\mu|^2}d\hat{r}^2 + \frac{\hat{r}^2}{6}\hat{h}_1^2 + \frac{\hat{r}^3 - 2|\mu|^2}{6\hat{r}}\hat{h}_2^2 + \frac{2\hat{r}^3 - |\mu|^2}{9\hat{r}}\hat{h}_3^2].
\end{aligned} \tag{A.38}$$

The reduced $D7$ -brane action for the scalar modes descending from the open moduli field χ is given by

$$\begin{aligned}
S_B(D7) &= -\mu_7 \int d^4x d\rho \int \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 \sqrt{-\gamma^{(0)}} e^\phi [1 + \frac{A_2 S_\rho}{A_1} (\frac{\partial_\rho \bar{\chi}}{\eta_\chi} + \frac{\partial_\rho \chi}{\eta_\chi}) \\
&+ \frac{2^{1/2}}{3^{1/2}r} h(r) \frac{(1 + C_\rho)}{1 + 2C_\rho} \tilde{g}^{\mu\nu} \partial_\mu \chi \partial_\nu \bar{\chi} + \frac{4}{|\mu_\chi|^2} (\frac{1 + C_\rho}{1 + 2C_\rho} + \frac{1}{(1 + 2C_\rho)^2}) \partial_\rho \chi \partial_\rho \bar{\chi} + \dots] \\
&= -\frac{\mu_7}{g_s^2} \int d^4x \int \hat{h}_1 \wedge \hat{h}_2 \wedge \hat{h}_3 e^\phi \int d\rho \sqrt{-\gamma^{(0)}} [1 + 2\frac{(r_\chi^3(r_\chi^3 - 2))^{1/2}}{(2r_\chi^3 - 1)} (\frac{\partial_\rho \chi}{\eta_\chi} + H. c.) \\
&+ \frac{1}{3|\mu_\chi|^{2/3}} h(r) r_\chi^2 \left(1 - \frac{2(r_\chi^3 - 2)}{(2r_\chi^3 - 1)}\right) \partial_\mu \chi \partial^\mu \bar{\chi} + \frac{4r_\chi^3}{3|\mu_\chi|^2(2r_\chi^3 - 1)} \left(1 - \frac{(r_\chi^3 - 2)}{(2r_\chi^3 - 1)}\right) \partial_\rho \chi \partial_\rho \bar{\chi}].
\end{aligned} \tag{A.39}$$

The first terms (in the last line above) inside the two pairs large parentheses in front of $|\partial_{\mu,\rho}\chi|^2$, reproduce the results of [42] while the second terms arising from additional contributions to the Born-Infeld determinant are absent there. The pull-back of the NSNS classical solution on Σ_4

$$B_2^{cl} = e^{-\phi/2} h^{-1/2} |\eta_\chi|^4 b(r) \cosh^2 \frac{\rho}{2} [-\sinh \frac{\rho}{2} d\rho \wedge \hat{h}_2 + \cosh \frac{\rho}{2} \hat{h}_3 \wedge \hat{h}_1], \quad [b(r) = \frac{81}{16} \frac{g_s M \alpha'}{r^6} \ln \frac{r}{r'_{ir}}] \tag{A.40}$$

agrees with [42] and involves an infrared radius distinct from the warp profile radius in Eq. (A.35), $\ln r'_{ir} = \frac{1}{3} - \ln(2^{5/6} 3^{-1/2} \epsilon^{-2/3}) = \frac{1}{4} + \ln r_{ir}$. The contributions from the classical NSNS solution B_2^{cl} and the gauge 2-form solution [41] $F_2^{cl} = \frac{P}{r^8} \Re(i\tilde{\mu}\epsilon_{ijk} w_j dw_k)$ in Eq. (A.28),

$$B_2^{cl} + F_2^{cl} \propto \frac{2}{3r^6} (k(r) + \frac{3P}{2}), \quad [k(r) = (3/2)^4 g_s M \alpha' \ln(r/r'_{ir})] \tag{A.41}$$

(both of same dimension E^0) can be included together by shifting the above radial profile $k(r) \equiv 3r^6 b(r) \rightarrow k(r) + 3P/2$.

-
- [1] A. Giveon and D. Kutasov, [arXiv:hep-th/9802067]
[2] J. Polchinski and M. Grana, Phys. Rev. **D63**, 026001 (2000) [arXiv:hep-th/0009211]; M. Grana and J. Polchinski, Phys. Rev. **D65**, 126005 (2001) [arXiv:hep-th/0106014]
[3] A. Karch and E. Katz, [arXiv:hep-th/0205236]
[4] E.G. Gimon, L.A. Pando Zayas, J. Sonnenschein, and M.J. Strassler, [arXiv:hep-th/0212061]
[5] M. Kruczenski, D. Mateos, R.C. Myers and D.J. Winters, JHEP **C07**, 049 (2003) [arXiv:hep-th/0304032]; [arXiv:hep-th/0312071]
[6] F. Marchesano, P. McGuirk, and G. Shiu, [arXiv:0812.2247]

- [7] F. Marchesano, P. McGuirk, and G. Shiu, [arXiv:1012.2759]
- [8] B.S. Acharya, J.M. Figueroa-O'farrill, C.M. Hull, and B. Spence, [arXiv:hep-th/9808014]
- [9] J.P. Conlon, A. Maharana, and F. Quevedo, [arXiv:hep-th/0807.0789]
- [10] S. Hartnoll, [arXiv:hep-th/0201271]
- [11] B.S. Acharya, F. Benini, and R. Valandro, [arXiv:hep-th/0612192]
- [12] H. Abe, A. Oikawa, and H. Otsuka, [arXiv:hep-th/1510.03407]
- [13] S. Hong, S. Yoon, and M. Strassler, [arXiv:hep-th/0410080]
- [14] I. Kirsch, [arXiv:hep-th/0607205]
- [15] D. Mateos, R. Myers, and R.M. Thomson, [arXiv:hep-th/0701132]; D. Mateos, [arXiv:hep-ph/0709.1523]
- [16] E. Witten, Adv. Theor. Math. Phys. **2**, 505 (1998), [arXiv:hep-th/9803131]
- [17] T. Sakai and S. Sugimoto, [arXiv:hep-th/0412141]; [arXiv:hep-th/0507073]
- [18] O. Aharony, J. Sonnenschein, and S. Yankielowicz, [arXiv:hep-th/0604161]
- [19] K. Peeters and M. Zamaklar, [arXiv:hep-th/0708.1502]
- [20] J. Erdmenger, N. Evans, I. Kirsch, and E. Threlfall, [arXiv:hep-th/0711.4467]
- [21] D. Mateos, [arXiv:0709.1523]
- [22] R. Minasian and D. Tsimpis, [arXiv:hep-th/9911042]
- [23] E. Bergshoeff and P. Townsend, [arXiv:hep-th/9611173]
- [24] E. Bergshoeff, R. Kallosh, T. Ortin, and G. Papadopoulos, [arXiv:hep-th/9705040]
- [25] M. Marino, R. Minasian, G. Moore, and A. Strominger, [arXiv:hep-th/9911206]
- [26] J.P. Gauntlett, [arXiv:hep-th/0305074]
- [27] L. Martucci and J. Smyth, [arXiv:hep-th/0507099]; L. Martucci, [arXiv:hep-th/0602129]
- [28] D. Lust, F. Marchesano, L. Martucci and D. Tsimpis, JHEP **C11**, 021 (2008) [arXiv:0807.4540]
- [29] I. Klebanov and M.J. Strassler, JHEP **C08**, 052 (2000), [arXiv:hep-th/0007191]
- [30] D. Arean, D.E. Crooks and A.V. Ramallo, JHEP **C11**, 035 (2004) [arXiv:hep-th/0408210]
- [31] D. Arean Fraga, [arXiv:hep-th/0605286]
- [32] H.-Y. Chen, P. Ouyang, and G. Shiu, [arXiv:hep-th/0807.2428]
- [33] P. Ouyang, [arXiv:hep-th/0311084]
- [34] S. Kuperstein, [arXiv:hep-th/0411097]
- [35] T.S. Levi and P. Ouyang, [arXiv:hep-th/0506021]
- [36] A.L. Cotrone, A. Dymarsky, and S. Kuperstein, JHEP **C03**, 005 (2011) [arXiv:1010.1017]
- [37] T. Sakai and J. Sonnenschein, [arXiv:hep-th/0305049]
- [38] S. Kuperstein and J. Sonnenschein, [arXiv:hep-th/0807.2897]
- [39] A. Dymarsky, S. Kuperstein and J. Sonnenschein, [arXiv:hep-th/0904.0988]
- [40] A. Dymarsky, [arXiv:hep-th/0909.3083]
- [41] F. Benini and A. Dymarsky, [arXiv:hep-th/1108.4931]
- [42] F. Benini, A. Dymarsky, S. Franco, S. Kachru, D. Simic, and H. Verlinde, JHEP **C12**, 031 (2009) [arXiv:0903.0619]
- [43] F. Benini, F. Canoura, S. Cremonesi, C. Nunez and A. V. Ramallo, JHEP **C02**, 0902007 (2007) [arXiv:hep-th/0612118]; [arXiv:hep-th/076.1238]
- [44] F. Benini, [arXiv:hep-th/0710.0374]
- [45] A. Paredes, [arXiv:hep-th/0610270]
- [46] R. Casero, C. Nunez and A. Paredes, [arXiv:hep-th/0602027]
- [47] F. Bigazzi, A.L. Cotrone, and A. Paredes, [arXiv:hep-th/0807.0298]; F. Bigazzi, A.L. Cotrone, A. Paredes, and A.V. Ramallo, [arXiv:hep-th/0812.339]; [arXiv:hep-th/0903.4747];
- [48] T. Gherghetta and J. Giedt, [arXiv:hep-th/0605212]
- [49] M. Chemtob, [arXiv:hep-th/2209.15503]
- [50] S.B. Giddings, S. Kachru, and J. Polchinski, Phys. Rev. **D66**, 106006 (2002) [arXiv:hep-th/0105097]
- [51] P. Candelas and X.C. de la Ossa, Nucl. Phys. **B342**, 246 (1990)
- [52] I.R. Klebanov and E. Witten, Nucl. Phys. **B536**, 199 (1998) [arXiv:hep-th/9807080]
- [53] M. Chemtob, [arXiv:1909.07668]
- [54] A. Dymarsky, I.R. Klebanov, and N. Seiberg, [arXiv:hep-th/0511254]
- [55] R.G. Leigh and M.J. Strassler, [arXiv:hep-th/9503121] M.J. Strassler, [arXiv:hep-th/0505153]
- [56] G. Carlino, K. Konishi, and H. Murayama, [arXiv:hep-th/0005076]
- [57] M. Douglas and G. Moore, [arXiv:hep-th/9603167]
- [58] D. Sorokin, [arXiv:hep-th/9906142]
- [59] T.J. Willmore, 'An Introduction to Differential Geometry' (Dover Books on mathematics, 2012); L.P. Eisenhart, Riemannian Geometry, (Princeton Press, 1926)
- [60] L. Martucci, J. Rosseel, D. Van den Bleeken, and A. Van Proeyen, [arXiv:hep-th/0504041]
- [61] A. van Proeyen, [arXiv:hep-th/9910030]
- [62] M.K. Benna, A. Dymarsky, I.R. Klebanov, and A. Solovoyov, [arXiv:hep-th/0712.4404]
- [63] J. Minahan, [arXiv:hep-th/9811156]
- [64] J. G. Russo and K. Sfetsos, [arXiv:hep-th/9901056]
- [65] M. Krasnitz, [arXiv:hep-th/0011179], [arXiv:hep-th/0209163]
- [66] C.P. Herzog, I.R. Klebanov, and P. Ouyang, [arXiv:hep-th/0108101]
- [67] M.A. Rubin and C.R. Ordoñez, J. Math. Phys. **26**, 65 (1985)

- [68] S. Kachru, R. Kallosh, A.D. Linde, and S.P. Trivedi, [arXiv:hep-th/0301240]
- [69] D. Baumann, A. Dymarsky, I.R. Klebanov, J. Maldacena, L. McAllister, and A. Murugan, [arXiv:hep-th/0607050]
- [70] D. Baumann, A. Dymarsky, S. Kachru, I.R. Klebanov and L. McAllister, [arXiv:0706.0360]
- [71] S. Kachru, R. Kallosh, A.D. Linde, J. Maldacena, L. McAllister, and S.P. Trivedi, [arXiv:hep-th/0308055]
- [72] R. Kallosh, A. Linde, T. Wrase, and Y. Yamada, [arXiv:hep-th/2206.04210]
- [73] S. Lust and L. Randall, [arXiv:hep-th/2206.04708]
- [74] J. Babington, J. Erdmenger, N. Evans, Z. Guralnik, and I. Kirsch, [arXiv:hep-th/0306018]
- [75] J. Erdmenger, N. Evans, Y. Liu, and W. Porod, [arXiv:hep-th/2304.09190]
- [76] O. DeWolfe and S.B. Giddings, [arXiv:hep-th/0208123]
- [77] V. Kaplunovsky and J. Louis, [arXiv:hep-th/9402005]
- [78] M.R. Douglas, J. Shelton, and G. Torroba, [arXiv:hep-th/0704.4001]
- [79] O. Aharony, Y.E. Antebi, and M. Berkooz, [arXiv:hep-th/0508080]
- [80] O. DeWolfe, L. McAllister, G. Shiu, and B. Underwood, [arXiv:hep-th/0703088]
- [81] A. Krause and E. Pajer, [arXiv:hep-th/0705.4682]
- [82] S. Kachru, J. Pearson, and H.L. Verlinde, *JHEP* **C06**, 021 (2002) [arXiv:hep-th/0112197]
- [83] C.M. Brown and O. DeWolfe, *JHEP* **C01**, 039 (2009) [arXiv:hep-th/0806.4399]; *JHEP* **C05**, 018 (2009) [arXiv:hep-th/0901.4401]
- [84] O. DeWolfe, S. Kachru, and M. Mulligan, *Phys. Rev.* **D77**, 065011 (2008) [arXiv:0801.1520]
- [85] A. Karch, E. Katz, D.T. Son, and M.A. Stephanov, [arXiv:hep-th/0602229]
- [86] M. Berg, M. Haack, and W. Muck, [arXiv:hep-th/0507285]; [arXiv:hep-th/0612224]
- [87] A. Dymarsky and D. Melnikov, *JETP Lett.* **84**, 368 (2006); A. Dymarsky and D. Melnikov, *JHEP* **C05**, 035 (2008) [arXiv:hep-th/0710.4517]
- [88] O. Aharony, J. Sonnenschein, M.E. Peskin, and S. Yankielowicz, [arXiv:hep-th/9507013]
- [89] N. Evans, S.D.H Hsu, and M. Schwetz, [arXiv:hep-th/9703197], [arXiv:hep-th/9707260]
- [90] J. Erdmenger, N. Evans, Y. Liu, and W. Porod, [arXiv:hep-th/2304.09190]
- [91] A. Dymarsky, D. Melnikov, and A. Solovoyov, [arXiv:0810.5666]
- [92] S.S. Pufu, I.R. Klebanov, T. Klose, and J. Lin, [arXiv:1009.2763]
- [93] E. Witten, [arXiv:hep-th/9604030]
- [94] R. Kallosh, A.-K. Kashani-Poor, and A. Tomasiello, [arXiv:hep-th/0503138]
- [95] E. Bergshoeff, R. Kallosh, A.-K. Kashani-Poor, D. Sorokin, and A. Tomasiello, [arXiv:hep-th/0507069]
- [96] I. Bandos and D. Sorokin, [arXiv:hep-th/0607163]
- [97] M.S. Green, J.H. Schwarz, and E. Witten, ‘Superstring Theory’ Vols. I, II (Cambridge University Press, Cambridge, 1987)
- [98] S.S. Gubser, C.P. Herzog, and I.R. Klebanov, [arXiv:hep-th/0405282]
- [99] R. Argurio, G. Ferreti, and C. Petersson, [arXiv:hep-th/0601180]
- [100] S.S. Gubser, [arXiv:hep-th/9807164]; [arXiv:hep-th/0201114]
- [101] P. van Nieuwenhuizen, [arXiv:hep-th/1206.2667]
- [102] A. Higuchi, *J. Math. Phys.* **28**, 1553 (1987); Erratum *J. Math. Phys.* **43**, 6385 (2002)
- [103] O. Lechtenfeld and K. Kumar, [arXiv:hep-th/2002.01005]
- [104] R. Camporesi and A. Higuchi, [arXiv:gr-qc/9505009]
- [105] V.A. Letsios, [arXiv:hep-th/2011.07875]