

The system of the vortex-like structures: the viewpoint on a turbulence modelling

S.V. Talalov

Dept. of Applied Mathematics, Togliatti State University,
14 Belorusskaya str., Tolyatti, 445020 Russia.
svt_19@mail.ru

Abstract

In this study we suggest new approach to turbulence modeling. To develop this approach, we construct the set of the vortex-like dynamical systems evolving in the space E_3 . These systems are constructed using the AKNS hierarchy so that they generalize thin closed vortex filament described in the local induction approximation. We construct and investigate the Hamiltonian dynamics of these systems in terms of the non - standard variables. We formulate the principles of quantization these systems focusing on the quantum description of the filaments with linearized dynamics. To describe the energy of vortex-like structures, we apply a new method based on the use of the central extended Galilei group. In a special case, we calculate the energy levels of a system of quantized vortices. The proposed approach makes it possible to describe both the interaction of such vortex - like filaments and the processes of their creation and annihilation in a natural way. As a result, we have constructed a quantum turbulent flow model that demonstrates the relationship between stochastic and integrable dynamics.

keywords: vortex filament quantization; AKNS hierarchy; extended Galilei group; turbulence modelling.

1 Introduction

The turbulence, both classical and quantum, continues to be a phenomenon that is not fully understood. A common approach to describing this complex phenomenon is to consider a turbulent flow as a tangle of vortex structures. A large number of works are devoted to this issue. For the first time, the description of a turbulent medium as a tangle of vortices was proposed in Feynman's work [1]. The achievements in this matter, which followed over several decades, were summarized, for example, in the books [2, 3]. Despite many years of studying turbulence, this complex phenomenon is still of interest to researchers. Indeed, "...the understanding of turbulent flows is one of the biggest current challenges in physics" [4]. Quantum effects add complexity to the study of this problem. So, the differences between the classical and quantum turbulence were considered in the paper [5]. Along with the theoretical understanding of the problem, numerical studies were also carried out [6] (see also [7, 8] for resent results here).

As usual, the theoretical study of the phenomenon involves building some kind of model. The Gross-Pitaevskii model is the most used theory in modeling quantum turbulence. So, one of

the predictions of this theory is the values of quantum circulation $\Gamma_n \propto \hbar n$, where the numbers n are integers. The study of the statistics of the distribution of the Γ_n value in a turbulent flow was investigated in work [9] by numerical simulation. The Hall-Vinen-Bekharevich-Khalatnikov (HVBK) model is also widely used for numerically study quantum turbulence. For example, this approach was used for this purpose in the paper [11].

Currently, it is considered an established fact that turbulent flow is characterized by a wide range of length and time scales. This question was discussed, for example, in the paper [10]. Topological and geometrical aspects of a quantum turbulent flow was studied in the work [12]. More complicated problems were also considered: quantum turbulence in a channel with an counterflow of superfluid turbulent helium was studied in the paper [13].

Despite the large number of works in the field of quantum turbulence, this phenomenon cannot be considered fully investigated. In this study, the author develops new approach to the turbulent flow modeling. This approach demonstrates the relationship between the regular and stochastic dynamics of vortex-like filaments in a tangle. Moreover, it makes it possible in principle to calculate a partition function for a turbulent flow in certain cases.

2 The hierarchy of the vortex-like dynamical systems

Nevertheless, the following question would be appropriate in a framework of any approach: how to describe a single vortex in a tangle? Corresponding description will be necessary, for example, when constructing a quantum theory of such complex structures. Approaches that have been developed to describe of a single thin vortex filament $\mathbf{r}(t, s)$ are well known. One of the most common ways of such a description is to use the Local Induction Approximation. Corresponding equation (LIE) is written as follows::

$$\frac{\partial \mathbf{r}(t^*, s)}{\partial t^*} = A \frac{\partial \mathbf{r}(t^*, s)}{\partial s} \times \frac{\partial^2 \mathbf{r}(t^*, s)}{\partial s^2}, \quad (1)$$

We use notation s for the natural curve parameter and notation $t^* = t\Gamma/4\pi$ for the "time" parameter. Symbols Γ and t mean the circulation and real time correspondingly. After certain assumptions [14], a vortex filament with a nonzero flow inside the core¹ is described by the equation:

$$\begin{aligned} \frac{\partial \mathbf{r}(t^*, s)}{\partial t^*} = & A \frac{\partial \mathbf{r}(t^*, s)}{\partial s} \times \frac{\partial^2 \mathbf{r}(t^*, s)}{\partial s^2} + \\ & + B \left(\frac{\partial^3 \mathbf{r}(t^*, s)}{\partial s^3} + \frac{3}{2} \left| \frac{\partial^2 \mathbf{r}(t^*, s)}{\partial s^2} \right|^2 \frac{\partial \mathbf{r}(t^*, s)}{\partial s} \right). \end{aligned} \quad (2)$$

Of course, the application of equations (1) and (2) to describe individual vortex filaments in a tangle is impossible. Indeed, the influence of other vortices in the vortex-tangle cannot be ignored here. On the other hand, it is also impossible to take into account the real physical influence a lot of nearby random vortices on a particular filament in a turbulent flow. Most

¹We assume that the core radius is small but non-zero.

probably, this problem is no less difficult than solving the Navier-Stokes equations in a general case. In this regard, the following question arises: can we modify equations (1) and (2) so that the dynamics of the filament is described by a sufficiently large number of parameters? In this case, the influence of neighboring vortices can be taken into account indirectly, through one or another distribution of these parameters. With this approach, we save the opportunity to describe the dynamics (classical and quantum) of a separate vortex filament.

The purpose of our study is as follows. We develop a theory that can be considered as a mathematical model of a tangle of vortex-like filaments. This model will include the following items:

1. We construct the dynamical theory of the separate vortex-like structure in Hamiltonian form. This theory generalizes dynamics (1) and (2) in some way. To solve this problem, we define the Hamiltonian structure of such systems in terms of non-standard fundamental variables. So, these variables are divided into two groups: internal and external ones;
2. To quantize these systems, we linearize the dynamics of considered objects. We will also discuss the relevance of this procedure in our model;
3. We construct the quantum theory of introduced vortex-like filaments. The constructed model makes it possible to describe the interaction of the vortex-like structures at quantum level by means of the many-body theory methods;
4. In a sense, the theory combines elements of integrability and stochasticity in the behavior of the objects under consideration.

For our further studies,, we use a representation for smooth closed filament of length S in the form

$$\mathbf{r}(\tau, \xi) = \mathbf{q} + R \int_0^{2\pi} [\xi - \eta] \mathbf{j}(\tau, \eta) d\eta, \quad R = \frac{S}{2\pi}, \quad \xi = \frac{s}{R}, \quad (3)$$

where dimensionless evolution parameter $\tau = t\Gamma/4\pi R^2$ was introduced. The vector $\mathbf{q} = \mathbf{q}(\tau)$ defines the position of the filament in the coordinate system (for example, the center of the circular vortex ring). The notation $[x]$ means the integer part of the number $x/2\pi$:

$$[0] = 0, \quad [x + 2\pi] = [x] + 1, \quad \forall x. \quad (4)$$

Vector $\mathbf{j}(\tau, \eta)$ is the unit tangent (affine) vector for this filament. Let the vector $\mathbf{r}(\tau, \xi)$ satisfies equation (1). Then, the function $\mathbf{j}(\tau, \eta)$ satisfies the equation for continuous Heisenberg spin chain:

$$\partial_\tau \mathbf{j}(\tau, \xi) = \alpha \mathbf{j}(\tau, \xi) \times \partial_\xi^2 \mathbf{j}(\tau, \xi). \quad (5)$$

It is well known that this equation is gauge equivalent to the non-linear Schrödinger equation [15]. The connection between the LIE and the nonlinear Schrodinger equation was firstly established in the work [17]. This equivalence makes it possible to construct new dynamical systems in space E_3 that generalize dynamical systems (1) and (2). Despite the formal definition, such systems, according to the author, open up a new point of view on a simulation of vortex filaments behavior in a vortex tangle.

First, we will consider the Hamiltonian description of systems (1) and (2) in terms of non-standard variables. Such a description was proposed in the authors' work [18]. The main points of the suggested approach are following.

- We include the circulation Γ as an additional independent variable in our theory. Because this postulate, we take into account the movement of the surrounding fluid. Indeed, the equations (1) and (2) describe some formal dynamics of the curve $\mathbf{r}(\tau, \xi)$ only. Therefore, we describe the dynamics of a filament by the following set \mathcal{A} of the independent variables:

$$\mathcal{A} = \{ \Gamma, \mathbf{q}, \mathbf{j}(\xi) \},$$

- We replace the set \mathcal{A} by the set \mathcal{A}' , where

$$\mathcal{A}' = \{ \mathbf{p}, \mathbf{q}; \mathbf{j}(\xi) \}. \quad (6)$$

The momentum \mathbf{p} is connected with variables Γ and $\mathbf{j}(\xi)$ by means of standard hydrodynamical formula

$$\mathbf{p} = \frac{\varrho_0}{2} \int \mathbf{r} \times \mathbf{w}(\mathbf{r}) dV, \quad (7)$$

where ϱ_0 is a fluid density and vector-function $\mathbf{w}(\mathbf{r})$ means the vorticity of the vortex filament. This function is calculated as

$$\mathbf{w}(\mathbf{r}) = \Gamma \int_0^{2\pi} \hat{\delta}(\mathbf{r} - \mathbf{r}(\xi)) \partial_\xi \mathbf{r}(\xi) d\xi. \quad (8)$$

- The appearance of the momentum components p_i as independent variables makes it possible to define the Galilei boosts in a natural way. As a consequence, we assume that the central extended Galilei group² $\tilde{\mathcal{G}}_3$ is the space-time symmetry group in our model. Therefore, we apply the group-theoretical approach to the definition of the energy of thin vortex filament.
- The structure of Hamiltonian is motivated by the expressions for Casimir functions of Lie algebra of the group $\tilde{\mathcal{G}}_3$:

$$H_0(p_1, p_2, p_3; \mathbf{j}) = \frac{1}{2m_0} \sum_{i=1}^3 p_i^2 + H_{int}[\mathbf{j}; \alpha_1, \dots, \alpha_n]. \quad (9)$$

The term $H_{int}[\mathbf{j}; \alpha_1, \dots, \alpha_n]$ provides dynamics of the variables $\mathbf{j}(\xi)$ that depends on some parameters $\alpha_1, \dots, \alpha_n$.

The formulas (7) and (8) are written in a compact form:

$$\mathbf{p} = \varrho_0 R^2 \Gamma \mathbf{f}, \quad \mathbf{f} = \frac{1}{2} \int_0^{2\pi} \int_0^{2\pi} [\xi - \eta] \mathbf{j}(\eta) \times \mathbf{j}(\xi) d\xi d\eta. \quad (10)$$

²We use one parameter central extension of this group. The mass constant m_0 (central charge) is the parameter of this extension.

Please note that the replacement $\Gamma \rightarrow \mathbf{p}$ leads to the certain constraints on the set \mathcal{A}' in general. We will not discuss this issue here [26].

The theory has three natural dimensional constants that are relevant to the physical system being described. These constants are: the fluid's density ϱ_0 , the speed of sound in this fluid v_0 and the mass constant m_0 . In order to simplify some formulas, we will use the auxiliary constants³ $R_0 = \sqrt[3]{m_0/\varrho_0}$, $t_0 = R_0/v_0$ and $\mathcal{E}_0 = m_0 v_0^2$ along with constants ϱ_0 , v_0 and m_0 .

The Poisson bracket structure was postulated as following:

$$\begin{aligned} \{p_i, q_j\} &= \delta_{ij}, \quad i, j = 1, 2, 3, \\ \{j_a(\xi), j_b(\eta)\} &= -(2/R_0 m_0 v_0) \epsilon_{abc} j_c(\xi) \hat{\delta}(\xi - \eta), \quad \epsilon_{123} = 1. \end{aligned} \quad (11)$$

All other brackets vanish. In accordance with the definition (11), the function $\mathbf{j}^2(\eta)$ annulates the brackets of the fundamental variables. Thus, the constant $\mathbf{j}^2(\eta)$ numbers the symplectic sheets in the phase space of our dynamical system. Next, we consider a single symplectic sheet with $\mathbf{j}^2 = 1$.

To develop our approach it let us note that the equation (5) lead to the infinitely series of the integrals of the motion $I_n = I_n[\mathbf{j}]$, $n = 0, 1, 2, \dots$ [15]. These integrals generate Hamiltonian flows that lead to the hierarchy of the non-linear integrable differential equations, which is known as hierarhy AKNS [16]. So, we use the Hamiltonian

$$H_{int} = \mathcal{E}_0 \int_0^{2\pi} (\partial_\xi \mathbf{j}(\xi))^2 d\xi = 4\mathcal{E}_0 I_1 \quad (12)$$

to generate the dynamics in accordance with equation (5). As for the full Hamiltonian (9), it provides the dynamics of the filament according to the equation (1).

Let's now add to expression (12) the term that is proportional to the integral of motion I_2 . Then the resulting full Hamiltonian

$$\begin{aligned} H_0(p_1, p_2, p_3; \mathbf{j}) &= \frac{1}{2m_0} \sum_{i=1}^3 p_i^2 + \\ &+ \mathcal{E}_0 \left(\alpha_1 \int_0^{2\pi} (\partial_\xi \mathbf{j}(\xi))^2 d\xi + \alpha_2 \int_0^{2\pi} \partial_\xi \mathbf{j}(\xi) (\mathbf{j}(\xi) \times \partial_\xi^2 \mathbf{j}(\xi)) d\xi \right) \end{aligned} \quad (13)$$

will provide the dynamics according to the equation (2).

Considered Hamiltonians H_{int} have a natural geometrical interpretation. Indeed, taking into account Frenet-Serret equations for the curve (3), we can find that following simple formula holds:

$$H_{int}[\mathbf{j}; \alpha_1, \alpha_2] = \mathcal{E}_0 \left(\alpha_1 \int_0^{2\pi} (k(\xi))^2 d\xi + \alpha_2 \int_0^{2\pi} (k(\xi))^2 \varkappa(\xi) d\xi \right).$$

³Note that there may be other natural dimensional constants in theory: for example, the radius R_1 of the pipe in which the vortex tangle in question moves.

The function $k(\xi)$ is the curvature of the curve $\mathbf{r}(\cdot, \xi)$ here and the function $\varkappa(\xi)$ is the torsion of this curve.

The structure of the Hamiltonian (13) makes it natural to use AKNS hierarchy to generalize the dynamic system (2). Let the number $\mathbf{N} > 0$ is the natural number. To describe the hierarchy of the vortex-like filaments, we consider the Hamiltonian (9), where

$$H_{int}[\mathbf{j}; \alpha_1, \alpha_2] \rightarrow H_{int}^{\varkappa}[\mathbf{j}; \alpha_1, \dots, \alpha_{\mathbf{N}}],$$

where

$$H_{int}^{\varkappa}[\mathbf{j}; \alpha_1, \dots, \alpha_{\mathbf{N}}] = 4\mathcal{E}_0 \sum_{k=1}^{\mathbf{N}} \alpha_k I_k[\mathbf{j}], \quad \alpha_j \in \mathbb{R}. \quad (14)$$

The certain number of constants α_k may be zero. The fact that $\alpha_{k_1} = 0$ for some index k_1 means that the corresponding integral of motion $I_{k_1}[\mathbf{j}]$ does not affect the dynamics of the filament. To account for this effect, we define the real number $\varkappa \in [0, 1]$ that is written in binary form as

$$\varkappa = 0, b_{\mathbf{N}-1} b_{\mathbf{N}-2} \dots b_0 0000000 \dots \quad (15)$$

The numbers b_{k-1} are mapped to constants α_k according to the following rule: $b_{k-1} = 0$ for the case $\alpha_k = 0$ and $b_{k-1} = 1$ otherwise. The number \varkappa can take $2^{\mathbf{N}}$ values. This number determines the type of vortex-like filament evolution of concrete vortex loop in a vortex tangle.

Therefore, we have the hierarchy of the dynamical systems in framework of developed theory. These systems generalize the vortex systems (1) and (2) naturally. In general, the dynamics of the vortex-like filament (3) is described by $\mathbf{N} + 1$ -order non-linear differential equation:

$$\partial_{\tau} \mathbf{r} = \{H_{int}^{\varkappa}[\mathbf{j}; \alpha_1, \dots, \alpha_{\mathbf{N}}], \mathbf{r}\}. \quad (16)$$

In conclusion of this section, we will say a few words about the geometric interpretation of such dynamical systems. First, the functions $P_n(\xi)$ in the integrals $I_n = \int_0^{2\pi} P_n(\xi) d\xi$ are polynomials of the solution $\psi(\xi)$ of the non-linear Schrödinger equation and corresponding derivatives $\partial^n \psi(\xi)$. Second, the function $\psi(\xi)$ can be reconstructed on the curvature $k(\xi)$ and the torsion $\varkappa(\xi)$ in accordance with Hasimoto results [17]. Therefore, the Hamiltonians $H_{int}^{\varkappa}[\mathbf{j}; \alpha_1, \dots, \alpha_{\mathbf{N}}]$ will be functions of quantities $k(\xi)$ and $\varkappa(\xi)$.

3 Quantization

The problem of quantum description of vortices has a very long history. The author will not take the responsibility to give a full review of the literature on this issue here. In most of the works that have been cited above, this problem was also discussed in some way. In addition, it is necessary to mention the book [19], which also has a detailed review of the literature on the problem. The quantum Hasimoto transformation $\text{LIE} \rightarrow \text{non-linear Schrödinger equation}$ was studied in the work [20] recently.

We emphasize that the general dynamics of the vortex filaments is described by nonlinear equations, even for simplified case of the local induction approximation. The quantization of the corresponding non-linear integrable models is complex problem (see, for example, [21, 22]). According to our initial assumptions, the discussed vortex-like dynamical systems form some

”entangled structure” that models the turbulent flow. The lifetime of vortices in a turbulent flow is small; this value restricted by the period $T = 2\pi/W$, where symbol W means the vorticity [23]. Thus, it seems appropriate to consider the quantization of small perturbations of vortex loops. Such perturbations are described by the linear equations at corresponding approximation. Therefore, the quantization of these perturbations can be performed using standard methods. The main question here is follows: what fundamental variables must parametrize the phase space of the dynamical systems under consideration? It is well known that quantization of variables that are canonically equivalent in a classical theory, leads to different results at the quantum level⁴. Despite more than a century of development of quantum theory, the following Dirac’s words are still relevant: *”... methods of quantization are all of the nature of practical rules, whose application depends on consideration of simplicity”* [25].

The quantization of small oscillations of the vortex ring was suggested in the authors’ article [26] in terms of non-standard Hamiltonian variables (6). As special result, the set of acceptable values of quantized circulation $\Gamma_{[n]}$ was calculated within the framework of this approach [27]. As it turns out, this set is wider than the standard one $\Gamma_n \propto n$, where n is natural number. The developed theory allows us to interpret a quantized vortex as a certain particle with an internal structure. Consequently, we can apply the methods of the theory of many bodies to describe the processes of the interaction, creation and annihilation of the vortices [28].

Let us make the overview of our quantization scheme. First, we consider the single closed vortex-like filament that has some fixed shape. This filament is homotopically equivalent to a circle and is characterized by a certain tangent vector $\mathbf{j}_0(\eta) = \mathbf{j}_0(\tau, \eta)$:

$$\mathbf{j}_0(\eta + 2\pi) = \mathbf{j}_0(\eta), \quad \mathbf{j}_0^2(\eta) = 1.$$

We consider the small perturbation of this tangent vector:

$$\mathbf{j}_0(\tau, \xi) \rightarrow \mathbf{j}_0(\xi) + \varepsilon \mathbf{j}_p(\tau, \xi), \quad \varepsilon \ll 1. \quad (17)$$

Additionally, we suppose that these excitations be transverse:

$$\mathbf{j}_p(\tau, \xi) \mathbf{j}_0(\xi) \equiv 0 \quad (18)$$

The small perturbation of the circular vortex filament without the restriction (18) has been studied in the work [29]. Various aspects of the theory related to vortex ring oscillations have also been studied in the works [30, 31]. Besides that, the small perturbations of straight vortex filaments were studied in the work [32].

The condition (18) leads to decomposition

$$\mathbf{j}_p(\tau, \xi) = j_\rho(\tau, \xi) \mathbf{e}_\rho + j_z(\tau, \xi) \mathbf{e}_z, \quad (19)$$

where the vectors $\{\mathbf{e}_\rho, \mathbf{j}_0, \mathbf{e}_z\}$ denote the local orthonormal basis in the corresponding point of the loop. For convenience, we define the complex-valued function

$$\mathfrak{J}(\tau, \xi) = j_\rho(\tau, \xi) + i j_z(\tau, \xi).$$

⁴For example, the quantization of such a simple canonical system as harmonic oscillator in terms of action-angle variables demonstrates many unexpected and interesting results [24].

This function allows us define the complex amplitudes $j_n(\tau)$:

$$\mathfrak{J}(\tau, \xi) = \sum_n j_n(\tau) e^{i n \xi}. \quad (20)$$

For example, in the simplest case (1) amplitude $\mathfrak{J}(\tau, \xi)$ satisfies the linear equation ($A = 1$) [26]

$$\partial_\tau \mathfrak{J} = -i \partial_\xi^2 \mathfrak{J} - \frac{i}{2} (\mathfrak{J} - \overline{\mathfrak{J}}). \quad (21)$$

Consequently,

$$\mathfrak{J}(\tau, \xi) = \sum_n j_n e^{i[n\xi + n\sqrt{n^2-1}\tau]}, \quad (22)$$

where the amplitudes j_n satisfy the condition $\overline{j}_{-n} = 2[n\sqrt{n^2-1} - n^2 + \frac{1}{2}] j_n$.

Dynamics

$$j_{-n} \rightarrow j_{-n}(\tau) = j_{-n} e^{i n \sqrt{n^2-1} \tau}$$

is generated (for the conditional time $t = t_0 \tau$) by Hamiltonian

$$H_0(p_1, p_2, p_3; j) = \frac{\mathbf{p}^2}{2m_0} + \mathcal{E}_0 \sum_{n>1} |j_{-n}|^2 n \sqrt{n^2-1} \quad (23)$$

and Poisson brackets

$$\begin{aligned} \{p_i, q_j\} &= \delta_{ij}, \quad i, j = x, y, z, \\ \{j_m, \overline{j}_n\} &= (i/m_0 R_0 v_0) \delta_{mn}, \quad m, n = -1, -2, \dots \end{aligned} \quad (24)$$

In the general case (14), we have more complex but also linear equations for the amplitudes $\mathfrak{J}(\tau, \xi)$.

We assume that the non-exited filament $\mathbf{r}_0(\tau, \xi)$ (see Eq. (3), where $\mathbf{j}(\xi) \equiv \mathbf{j}_0(\xi)$), is described by the finite numbers of additional⁵ dynamical variables, except for "external" variables \mathbf{p} and \mathbf{q} . To simplify our subsequent studies, we consider the non-exited filament as a ring of radius R . In case of the Eq. (2), the following formula for the filament $\mathbf{r}_0(\tau, \xi)$ is fulfilled:

$$\mathbf{r}_0(\tau, \xi) = \mathbf{q} + R(\cos(\xi + \phi_0 + \beta\tau)\mathbf{e}_x, \sin(\xi + \phi_0 + \beta\tau)\mathbf{e}_y, A\tau\mathbf{e}_z), \quad (25)$$

where the angle $\phi_0 \in [0, 2\pi)$ and $\beta = B/R$. Therefore, the variables

$$\Delta R = \sqrt{R^2 - R_0^2}, \quad \phi(\tau) = \phi_0 + \beta\tau$$

can be considered as the additional variables. Because the inequality $R \geq R_0$ is fulfilled, the constant R_0 can characterize the average size of a molecular cluster in a flow.

To simplify our consideration, we consider here the quantization of the variables \mathbf{p} , \mathbf{q} and $\mathbf{j}_p(\tau, \xi)$ only. As regards of the quantization of the additional variables that define the filament $\mathbf{r}_0(\tau, \xi)$, this procedure was fulfilled in the articles [28, 33] in simplest case (25).

⁵to the variables $\mathbf{j}_p(\tau, \xi)$

The constructed set of the classical dynamical variables defines the quantization scheme uniquely. The quantum states of a single vortex loop are the vectors of the Hilbert space

$$\mathbf{H}_1 = \mathbf{H}_{pq} \otimes \mathbf{H}_j, \quad (26)$$

where the symbol \mathbf{H}_{pq} denotes the Hilbert space of a free structureless particle in $3D$ space \mathbb{R}_3 . As usually, we suppose that $\mathbf{H}_{pq} = L^2(\mathbb{R}_3)$. The symbol \mathbf{H}_j denotes the space for the number of the harmonic oscillators (infinite in general) which correspond to the variables j_{-n} . In order to simplify the following formulas, we assume that a finite number K of modes j_{-k} can be excited only for every vortex filament. From the author's point of view, the proposed quantization scheme is a mirror image of Lord Kelvin's old idea that particles can be considered as some kind of vortex-like structures [34]. In our approach, we describe the closed vortex-like filaments as structured particles.

As it is mentioned above, our quantization scheme make it possible to describe the interaction of vortex-like structures in terms of the many-body theory. The details can be found in the articles [28, 33] for certain special case. Here we will go the brief overview only.

- The space of a quantum states of a "vortex tangle" is constructed as the Fock space

$$\mathfrak{H} = \bigoplus_{M=0}^{\infty} \mathbf{H}_M,$$

where

$$\mathbf{H}_M = \bigotimes_{i=0}^M \mathbf{H}_1 = \underbrace{[\mathbf{H}_{pq} \otimes \mathbf{H}_j] \otimes \cdots \otimes [\mathbf{H}_{pq} \otimes \mathbf{H}_j]}_M = \mathbf{H}_{pq}^M \otimes \mathbf{H}_j^M. \quad (27)$$

Any vector $|\Phi^M\rangle \in \mathbf{H}_M$ has a form

$$\begin{aligned} |\Phi^M\rangle &= \sum_{n_1, \dots, n_M} \int \cdots \int d\mathbf{p}_1 \cdots \mathbf{p}_N \times \\ &\times f_{n_1, \dots, n_M}^M(\mathbf{p}_1, \dots, \mathbf{p}_M) |\mathbf{p}_1\rangle \cdots |\mathbf{p}_M\rangle |\mathbf{n}_1\rangle \cdots |\mathbf{n}_M\rangle, \end{aligned} \quad (28)$$

where the functions f^M are symmetrical both indexes \mathbf{n} and arguments \mathbf{p}_ℓ . The vector $|\mathbf{p}_\ell\rangle$ are eigenvector of the operator $\hat{\mathbf{p}}_\ell$ that corresponds to the filament with index " ℓ ". Here and further, we use the summation index " ℓ " to number the vortex filaments only. As regards of the "numbers" \mathbf{n}_ℓ , these symbols are reduced form for occupation numbers $n_k(\ell)$:

$$\mathbf{n}_\ell = [n_1(\ell) \dots, n_K(\ell)].$$

The occupation number $n_k(\ell)$ describes the excitation of the variable j_{-k} (see Eq. (20)) for the filament with the index " ℓ ".

- Hamiltonian

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0 + \hat{U},$$

where operator $\hat{\mathcal{H}}_0$ means the Hamiltonian for the system of the non - interacting vortex-like structures.

To write down these operators, let's first introduce notation for some operators in the Fock space \mathfrak{H} :

1. The creation and annihilation operators $\hat{a}^+(\mathbf{p}, \mathbf{n})$, $\hat{a}(\mathbf{p}, \mathbf{n})$ act in the Fock space \mathfrak{H} . They are defined in a standard way:

$$\begin{aligned} \left(\hat{a}(\mathbf{p}, \mathbf{n}) f_{\mathbf{n}_1, \dots, \mathbf{n}_M}^M \right)_{\mathbf{n}_1, \dots, \mathbf{n}_{M-1}}(\mathbf{p}_1, \dots, \mathbf{p}_{M-1}) &= \sqrt{M} f_{\mathbf{n}_1, \dots, \mathbf{n}_{M-1}, \mathbf{n}}^M(\mathbf{p}_1, \dots, \mathbf{p}_{M-1}, \mathbf{p}), \\ \left(\hat{a}^+(\mathbf{p}, \mathbf{n}) f_{\mathbf{n}_1, \dots, \mathbf{n}_M}^M \right)_{\mathbf{n}_1, \dots, \mathbf{n}_{M+1}}(\mathbf{p}_1, \dots, \mathbf{p}_{M+1}) &= \\ &= \left(1/\sqrt{M+1} \right) \sum_{j=1}^{M+1} \delta_{\mathbf{n}, \mathbf{n}_j} \delta(\mathbf{p} - \mathbf{p}_j) f_{\mathbf{n}_1, \dots, \cancel{\mathbf{n}_j}, \dots, \mathbf{n}_{M+1}}^M(\mathbf{p}_1, \dots, \cancel{\mathbf{p}_j}, \dots, \mathbf{p}_{M+1}). \end{aligned}$$

As usual, the crossed-out symbol means its absence.

2. Operators $\hat{a}_m^+(\ell)$, $\hat{a}_m(\ell)$ act in every Hilbert space \mathbf{H}_j^M . These operators create or annihilate the oscillator energy level for the mode \mathbf{j}_{-m} in the ℓ -th multiplier $[\mathbf{H}_{pq} \otimes \mathbf{H}_j]$ of decomposition (27). For example, the standard quantization rule for the value $|\mathbf{j}_{-m}|^2$ looks like this:

$$|\mathbf{j}_{-m}|^2 \rightarrow \frac{\hbar}{2} [\hat{a}_m^+(\ell) \hat{a}_m(\ell) + \hat{a}_m(\ell) \hat{a}_m^+(\ell)].$$

Thus, summarizing of our consideration, we can write down a general formula for the operator $\hat{\mathcal{H}}_0$:

$$\hat{\mathcal{H}}_0 = \bigoplus_{M=0}^{\infty} \hat{\mathcal{H}}_0^M,$$

where operators $\hat{\mathcal{H}}_0^M$ act in the spaces \mathbf{H}_M . These operators have following form:

$$\begin{aligned} \hat{\mathcal{H}}_0^M &= \frac{\hbar^2}{2m_0} \sum_{\mathbf{n}} \int \mathbf{p}^2 \hat{a}^+(\mathbf{p}, \mathbf{n}) \hat{a}(\mathbf{p}, \mathbf{n}) + \\ &+ \frac{\hbar}{2t_0} \sum_{\ell=1}^M \sum_{m=1}^K \Omega_{m\ell}(\boldsymbol{\varkappa}_\ell; \alpha_1(\ell), \dots, \alpha_{\mathbf{N}}(\ell)) [\hat{a}_m^+(\ell) \hat{a}_m(\ell) + \hat{a}_m(\ell) \hat{a}_m^+(\ell)]. \end{aligned} \quad (29)$$

The function $\Omega_{m\ell}(\boldsymbol{\varkappa}_\ell; \alpha_1, \dots, \alpha_{\mathbf{N}})$ corresponds to the vortex-like structure with number ℓ in the considered vortex tangle. Let us recall that the number \mathbf{N} determines the level of our model (see Eq. (15)). The explicit form of the functions $\Omega_{mn}(\boldsymbol{\varkappa}_n; \alpha_1, \dots, \alpha_{\mathbf{N}})$ can be deduced from the detail study of linearized dynamical equations (16). For example, let the number \mathbf{N} is equal to one and corresponding number b_0 is also equal to one: $\mathbf{N} = 1$, $b_0 = 1$. This simplest case corresponds to a vortex tangle that contains vortex loops described by the equation (1). The meaning of the constants $\alpha_i(\ell)$, $i = 1, \dots, M$ is as follows: $\alpha_1(\ell) \propto |\mathbf{v}(\ell)|$, where the vector $\mathbf{v}(\ell)$ is the velocity of the vortex loop that numbered by index " ℓ ". The operator $\hat{\mathcal{H}}_0^M$ has the

following form:

$$\begin{aligned}\hat{\mathcal{H}}_0^M &= \frac{1}{2m_0M} \sum_{\mathbf{n}} \int \mathbf{p}^2 \hat{a}^+(\mathbf{p}, \mathbf{n}) \hat{a}(\mathbf{p}, \mathbf{n}) + \\ &+ \frac{\hbar}{2t_0} \sum_{\ell=1}^M \sum_{k=1}^K \alpha_1(\ell) k \sqrt{k^2 - 1} [\hat{a}_k^+(\ell) \hat{a}_k(\ell) + \hat{a}_k(\ell) \hat{a}_k^+(\ell)].\end{aligned}\quad (30)$$

Let this considered simplest vortex tangle is located in a closed bounded domain $D \subset \mathbb{R}_3$. Then, the eigenvalue problem

$$\hat{\mathcal{H}}_0^M |\Phi_{[\sigma]}^M\rangle = E_{[\sigma]} |\Phi_{[\sigma]}^M\rangle, \quad |\Phi_{[\sigma]}^M\rangle \in \mathbf{H}_M,$$

has following solution. The eigenvectors $|\Phi_{[\sigma]}^M\rangle$ have form (28), where

$$\begin{aligned}f_{\mathbf{n}_1, \dots, \mathbf{n}_M}^M(\mathbf{p}_1, \dots, \mathbf{p}_M) &= \frac{1}{(M!)^2} \times \\ &\times \left(\sum_{[\mathbf{p}_1, \dots, \mathbf{p}_M]} \prod_{\ell=1}^M f_{w(\ell)}(\mathbf{p}_\ell) \delta(\mathbf{p}_\ell^2 - \lambda_{w(\ell)}^2) \right) \left(\sum_{[\mathbf{n}_1, \dots, \mathbf{n}_M]} \prod_{\ell=1}^M \delta_{\mathfrak{s}_\ell \mathbf{n}_\ell} \right),\end{aligned}$$

where the symbol $[c_1, \dots, c_M]$ means some rearrangement of the corresponding values c_1, \dots, c_M (summations are performed on all rearrangements). Symbols \mathfrak{s}_ℓ mean the reduced form for certain set of the occupation numbers $s_k(\ell)$:

$$\mathfrak{s}_\ell = [s_1(\ell) \dots, s_K(\ell)].$$

The function $f_i(\mathbf{p}) \delta(\mathbf{p}^2 - \lambda_i^2)$ is the momentum representation of the eigenfunction $\tilde{f}_i(\mathbf{q})$ which corresponds to certain eigenvalue λ_i^2 for the Laplace operator Δ in the domain D :

$$\Delta \tilde{f}_i(\mathbf{q}) = -\lambda_i^2 \tilde{f}_i(\mathbf{q}), \quad \mathbf{q} \in D, \quad \tilde{f}_i(\mathbf{q}) \Big|_{\mathbf{q} \in \partial D} = 0.$$

Formula (30) allows us to find eigenvalues:

$$\begin{aligned}E_{[\sigma]}(\alpha_1(1), \dots, \alpha_1(M)) &= \frac{\hbar^2}{2m_0} \sum_{\ell=1}^M \lambda_{w(\ell)}^2 + \\ &+ \frac{\hbar}{t_0} \sum_{\ell=1}^M \sum_{k=1}^K \alpha_1(\ell) k \sqrt{k^2 - 1} \left(s_k(\ell) + \frac{1}{2} \right).\end{aligned}\quad (31)$$

To simplify the writing of these formulas, we have introduced a notation for the multi-index $[\sigma]$:

$$[\sigma] = [w(1), \dots, w(M); s_1(1), \dots, s_1(M); \dots; s_K(1), \dots, s_K(M)].$$

Let's analyze the structure of formula (31). The terms, which are proportional to the value of \hbar , give the main contribution to the energy. These terms represent the energy of oscillatory

modes of the vortex rings. Terms that are proportional to the value of \hbar^2 can be interpreted as certain "fine structure" of oscillatory levels. This structure is completely determined by the geometry of the domain in which the considered vortex system is moving.

The interaction Hamiltonian \hat{U} is written as follows:

$$\hat{U} = \sum_{m,n} \varepsilon_{mn} \hat{U}_{m \leftrightarrow n}. \quad (32)$$

The constants ε_{mn} are the coupling constants which define the intensity of the vortex interaction. Operators $\hat{U}_{m \leftrightarrow n}$ define the reconnection of the considered vortex filaments in the flow. They are constructed in terms of the creation and annihilation operators $\hat{a}^+(\mathbf{p}, \mathbf{n})$ and $\hat{a}(\mathbf{p}, \mathbf{n})$. Operator $\hat{U}_{m \leftrightarrow n}$ describes the transformation of m vortex rings into n rings and vice versa, $n \rightarrow m$. For example, the theory with $U = \varepsilon \hat{U}_{1 \rightarrow 2}$ was investigated in the paper [28], where the model of the turbulent flow in it's initial stage was considered. We must emphasize that the variables \mathbf{p} and \mathbf{q} describe the virtual particles corresponding to the external degrees of freedom of the vortex-like loops. Therefore, the Hamiltonian \hat{U} describes, generally speaking, a non-local interaction. These questions, as well as corresponding examples of the operator $\hat{U}_{2 \leftrightarrow 2}$, were discussed in the article [33]. In this study, we will not dwell on this in detail.

4 Analysis and outlooks

Despite the presence of a number of uncertain parameters, the proposed theory represents certain new viewpoint to the description of turbulent flow. Some of these parameters appear to be random numbers. Apparently, this is unavoidable when describing a turbulence. Nevertheless, the approach developed in this study offers a new look at the relationship between "random" and "integrable" dynamics in the description of a fluids motion. According to the author, the proposed model can stimulate a vortex tangle even in the case of $U = 0$, especially for the large number of \mathbf{N} . Of course, the case where $U \neq 0$ is more interesting and realistic.

The proposed theory makes it possible to calculate the partition function \mathcal{Z} for any concrete domain D . In the simplest case considered above, we can write the following expression:

$$\mathcal{Z} = \prod_{\ell=1}^M \sum_n \exp \left(-\frac{\langle E_n(\ell) \rangle}{k_B T} \right),$$

where the numbers $E_n(\ell)$ are valid energy values for the vortex filament with number " ℓ ", symbol $\langle \dots \rangle$ means some averaging over the parameters $(\alpha_1(\ell), \dots, \alpha_{\mathbf{N}}(\ell))$, the value T is the temperature and the constant k_B is Boltzmann constant. This possibility is an important feature of the proposed approach from a practical viewpoint. Indeed, the function \mathcal{Z} allows us to calculate explicitly the thermodynamic characteristics of the simulated turbulent flow.

It is interesting to consider the limits $K \rightarrow \infty$ and $\mathbf{N} \rightarrow \infty$. It is clear that the proposed model, within such limits, will require the application of certain quantum field theory methods. For example, expression (31) for the energy levels would require a renormalization procedure. The limit $\mathbf{N} \rightarrow \infty$ leads, in general, to the non-local equations for the single vortex-like structures. As a cosequense, we have an additional possibility to describe the non-local effects. In this case, the number \varkappa (see Eq.(15)) takes any values in the segment $[0, 1]$. Note that non-local terms in LIE equation appear when describing the self-stretching [32]. Moreover there

is another option to introduce non-local effects here: we can consider the additional non-local integrals I_{-n} , $n = 1, 2, 3, \dots$ which can be introduced [15] for the non-linear Schrödinger equation. All these issues, as well as the physical interpretation of the proposed dynamical systems, require separate studies.

5 Novelty, limitations and possible applications

Firstly, the proposed quantization scheme leads to the richer circulation spectrum than the standard one. This issue is studied in the works of the author [26, 27, 33] in detail, so we do not discuss the relevant formulas here. From the author's point of view, this extension of a circulation spectrum provides more opportunities for practical modeling of a quantum turbulent flow.

Secondly, it is known that the standard approach to the energy definition of a thin ($a \rightarrow 0$) vortex filament leads to problems due to the divergence of the integrals. In this study, we applied a group-theoretical approach to the definition of energy.

Thirdly, as already noted above, constructed quantum systems are "inverse realization" of Lord Kelvin's old idea that any particles can be considered as some kind of vortex-like structures [34]. In our approach, we describe the closed vortex-like filament as some structured particle. In a sense, the fluid in which the vortex in question moves plays the role of "aether". Note that such "aether" is quite real here, and not hypothetical.

The proposed quantization scheme for a system of vortex-like structures takes into account the movement of the surrounding fluid in a minimal way [26]. Apparently, this fact imposes some limitations on the applicability of the constructed model. Undoubtedly, this issue requires further research.

6 Concluding remarks

The next stage in the development of the proposed model should be numerical simulation of some observable characteristics of the flow for both various types of interaction (32) and various types of domain $D \subset \mathbb{R}_3$. All these issues, as well as the physical interpretation and subsequent theoretical investigation of the constructed dynamical systems, require separate studies.

References

- [1] R. P. Feynman. *Progress in Low Temperature Physics*. ed. by C. J. Gorter. V. 1. P. 17. Amsterdam: North-Holland. (1955).
- [2] R.J. Donnelly. *Quantum Vortices in Helium II* Cambridge Univ. Press. (1991).
- [3] U. Frisch. *Turbulence. The Legacy of A.N. Kolmogorov*. Cambridge Univ. Press. (1996).
- [4] J.I. Polanco, N.P. Müller, G. Krstulovic. Nature Communication. V. 12. Pp. 1 -11 (2021) article number 7090.
- [5] W. F. Vinen. Journal of Low Temperature Physics. V. 145, pp. 7–24 (2006)

- [6] R. Aarts, *A numerical study of quantized vortices in He II*. Technische Universiteit Eindhoven. (1993)
- [7] M. Tsubota, K. Fujimoto and S. Yui. *J. Low. Temp. Phys.* 188, 119 (2017).
- [8] Koji Fukagata. *Nature Computational Science*. V. 2. pp. 68–69 (2022).
- [9] N.P. Möller, J.I. Polanco, G. Krstulovic. *Phys. Rev. X* 11, 011053 (2021).
- [10] N. Gourianov, M. Lubasch, ... D. Jaksch. *Nature Computational Science* V. 2. pp. 30 – 37 (2022).
- [11] Z. Zhang, I. Danaila, E. Lévêque, L. Danaila. *Journ. of Fluid Mechanics*, 962, A22 (2023).
- [12] Carlo F. Barenghi. Tangled vortex lines: dynamics, geometry and topology of quantum turbulence. in: *Knotted Fields*. ed. by R.L. Ricca and X. Liu, Springer (2022).
- [13] S.K. Nemirovskii. *Phys. Rev. B* 97, 134511 (2018).
- [14] S.V. Alekseenko, P.A. Kuibin, V.L. Okulov, *Theory of concentrated vortices*. Springer-Verlag, Berlin Heidelberg (2007).
- [15] L.A. Takhtajan, L.D. Faddeev, *Hamiltonian Methods in the Theory of Solitons*. Springer-Verlag, Berlin. 1987.
- [16] M.J Ablowitz, D.J. Kaup, A.C. Newell, H. Segur. The inverse scattering transform – Fourier analysis for non-linear problems. *Stud. Appl. Math.*, V.53. pp. 249 – 315 (1974)
- [17] H. Hasimoto. A soliton on a vortex filament // *J. Fluid Mech.* **23**. p. 61 - 97. 1972.
- [18] S.V. Talalov. // *Mod. Phys. Lett. B*. V. B32. No 33. 1850410. 2018. arXiv:1807.08922 [math-ph].
- [19] Sonin E.B. *Dynamics of Quantized Vortex in Superfluids*. Cambridge University Press (2016).
- [20] R. A. Van Gorder. Quantum Hasimoto transformation and nonlinear waves on a superfluid vortex filament under the quantum local induction approximation. *Phys. Rev. E* 91, 053201. (2015)
- [21] L. A. Takhtadzhian, L. D. Faddeev. The Quantum Method of the Inverse Problem and the Heizenberg XYZ Model // *Russ. Math. Surv.* V.34. 11. 1979. DOI 10.1070/RM1979v034n05ABEH003909
- [22] E.K. Sklyanin, *St. Petersburg Mathematical Journal (Formely Leningrad Mathematical Journal)*. **1**. No 2. (1989). 189 – 206.
- [23] H. Tennekes. in book: *Handbook of Turbulence*. ed. by W. Frost and T.H. Moulden. Plenum press, N.Y. (1977)
- [24] Kastrup H. A. *Annalen der Physik*. **519**. Issues 7-8. 439 - 528 (2007).

- [25] Dirac P.A.M. Generalized hamiltonian dynamics. Canadian Journal of Mathematics. V.2. No 2. pp. 129 - 148 (1950).
- [26] S.V. Talalov. Eur. Journ. Mech B/Fluids. **92**. pp. 100 - 106. (2022). arXiv: math-ph/2112.04859v1.
- [27] S.V. Talalov. Physics of Fluids. **34**. 041702 (2022)
- [28] S.V. Talalov. Physics of Fluids. **35**. 045132 (2023)
- [29] Ricca R.L. Chaos. **3**. Issue 1. 83–91 (1993)
- [30] V.F. Kop'ev, S.A. Chernyshev. *Vortex ring oscillations, the development of turbulence in vortex rings and generation of sound*. Phys. Usp. 43 663–690 (2000).
- [31] L. Kiknadze, Yu. Mamaladze. Journ. Low Temperature Physics. V. 126. Nos 1/2. Jan. (2002).
- [32] A.J. Majda and A.L. Bertozzi. *Vorticity and Incompressible Flow*. Cambridge Univ. Press. (2002).
- [33] S.V. Talalov. Phys. Rev. Fluids. **8**. 034607 (2023)
- [34] W. Thomson. On vortex atoms. Proc. R. Soc. Edinburgh 6, 94 (1869).