

INTRODUCTIVE EXERCISES TO QUANTUM FIELD THEORY

Stefano Disca, Roberto Demaria, Lorenzo Formaggio

(last revision on 24/04/2024)

Contents

Preface	3
1 Noether's Theorem and Conserved Charges	5
1.1 Complex Klein-Gordon field	5
1.2 Yukawa Lagrangian	9
2 Cross Sections and Decay Rates	11
2.1 Higgs boson decay	11
2.2 Scalar and standard QED: $\psi\bar{\psi} \rightarrow \phi\phi^*$	15
2.3 Scalar-boson interaction: $\phi\phi \rightarrow \gamma\gamma$	21
2.4 Scalar-fermion-boson interaction: $e^+e^- \rightarrow \phi\gamma$	26
2.5 Yukawa Lagrangian: $f\bar{f} \rightarrow \phi\phi$	31
2.6 Fermi Lagrangian: $\nu e^- \rightarrow \nu e^-$	36
2.7 QED with massless electron: $\gamma e^- \rightarrow \gamma e^-$	42
2.8 Chiral lagrangian: $\psi_a\psi_b \rightarrow \psi_a\psi_b$	47
3 Ward Identities	53
3.1 Radiative corrections for $\phi\phi^* \rightarrow \phi\phi^*\gamma$	53
3.2 Radiative corrections for $\psi\bar{\psi} \rightarrow \phi\phi^*\gamma$	58
A Useful formulae	63
Bibliography	69

Preface

The following is a collection of 12 exercises picked from the exam tests of the course *Elements of Quantum Field Theory*, teached by professor Mauro Moretti in the academic year 2021-22 for the Master's Degree in Physics at the University of Ferrara.

The goal of this review is to provide a general method to approach the main types of calculations in Quantum Field Theory. The topics that will be treated are: applications of Noether's theorem, checking of Ward identities, computations of cross sections and decay rates. In order to be as clear as possible in the solutions we preferred to be very explicit in the calculations, at the risk of being tedious. We must thank Thomas Lovo, for the constant help and suggestions given, and dr. Alessandro Greco, whose lecture notes of the course proved to be essential in order to write these ones.

Needless to say, this text is not free from errors; if you find errors in calculations or conceptual mistakes in general (or even more elegant solutions to the problems), please contact the authors by the following e-mail addresses: stefano.disca@edu.unife.it, roberto.demaria@edu.unife.it, lformagg@cougarnet.uh.edu.

Chapter 1

Noether's Theorem and Conserved Charges

1.1 Complex Klein-Gordon field

Text. Given a complex Klein-Gordon field, described by the Lagrangian

$$\mathcal{L} = \partial_\mu \phi \partial^\mu \phi^* - m^2 \phi \phi^*$$

determine the conserved charges of the theory.

Solution. First of all, remember that for every lagrangian theory we distinguish symmetries in *space-time* symmetries (resulting from transformations of space-time coordinates) and *internal* symmetries (resulting from field transformations which leave the coordinates untouched). The former type is common to all Lagrangians that are invariant under space-time translations, and its associated Noether currents correspond to the stress-energy tensor

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial^\nu \phi - \mathcal{L} g^{\mu\nu} \quad (1.1.1)$$

where $g^{\mu\nu}$ is the metric tensor; the following ones are the conserved charges

$$P^\nu = \int d^3x T^{0\nu} \quad (1.1.2)$$

This Lagrangian has also a *phase* symmetry

$$\begin{aligned} \phi &\rightarrow \phi' = e^{i\varepsilon} \phi \sim (1 + i\varepsilon) \phi \\ \phi^* &\rightarrow \phi'^* = e^{-i\varepsilon} \phi^* \sim (1 - i\varepsilon) \phi^* \end{aligned} \quad (1.1.3)$$

where ε is an arbitrary small parameter. The total variations of the fields are

$$\begin{aligned}\varepsilon \bar{\delta} \phi &\stackrel{def}{=} \phi' - \phi = i\varepsilon\phi \\ \varepsilon \bar{\delta} \phi^* &\stackrel{def}{=} \phi'^* - \phi^* = -i\varepsilon\phi^*\end{aligned}\quad (1.1.4)$$

so the Noether currents are

$$j^\mu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \bar{\delta} \phi + \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi^*)} \bar{\delta} \phi^* = i \left[\phi \partial_\mu \phi^* - \phi^* \partial_\mu \phi \right] \quad (1.1.5)$$

therefore its associated conserved charge is

$$Q = \int d^3x j^0 = i \int d^3x \left[\phi \partial_0 \phi^* - \phi^* \partial_0 \phi \right] =: Q_1 + Q_2 \quad (1.1.6)$$

The *real* Klein-Gordon field can be written as a combination of the ladder operators a^\dagger and a ; in the case of a *complex* scalar field we must treat ϕ and ϕ^* as two separate fields, so that we have two distinct types of ladder operators, a and b , associated to two different types of particles. Being the two fields simply related by complex conjugation, we can write them as follows

$$\begin{aligned}\phi(x) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[a_p e^{-iE_p t} e^{i\mathbf{p} \cdot \mathbf{x}} + b_p^\dagger e^{iE_p t} e^{-i\mathbf{p} \cdot \mathbf{x}} \right] \\ \phi^*(x) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[a_p^\dagger e^{iE_p t} e^{-i\mathbf{p} \cdot \mathbf{x}} + b_p e^{-iE_p t} e^{i\mathbf{p} \cdot \mathbf{x}} \right]\end{aligned}\quad (1.1.7)$$

We reobtain the case of a real Klein-Gordon field by setting $b_p = a_p$.

Let's now calculate $Q_1 = i \int d^3x \phi \partial_0 \phi^*$

$$\partial_0 \phi^*(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[iE_p a_p^\dagger e^{iE_p t} e^{-i\mathbf{p} \cdot \mathbf{x}} - iE_p b_p e^{-iE_p t} e^{i\mathbf{p} \cdot \mathbf{x}} \right] \quad (1.1.8)$$

$$\begin{aligned}\phi(x) \partial_0 \phi^*(x) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \frac{d^3q}{(2\pi)^3} \frac{1}{\sqrt{2E_q}} (iE_p) \left[a_p e^{-iE_p t} e^{i\mathbf{p} \cdot \mathbf{x}} + b_p^\dagger e^{iE_p t} e^{-i\mathbf{p} \cdot \mathbf{x}} \right] \\ &\quad \left[iE_p a_p^\dagger e^{iE_p t} e^{-i\mathbf{p} \cdot \mathbf{x}} - iE_p b_p e^{-iE_p t} e^{i\mathbf{p} \cdot \mathbf{x}} \right]\end{aligned}\quad (1.1.9)$$

so that

$$\begin{aligned}Q_1 &= i \frac{1}{(2\pi)^3} \frac{1}{2\sqrt{E_q E_p}} (iE_p) \int d^3x d^3p d^3q \left[a_q a_p^\dagger e^{i(E_q - E_p)t} e^{i(\mathbf{q} - \mathbf{p}) \cdot \mathbf{x}} + \right. \\ &\quad - a_q b_p e^{-i(E_q + E_p)t} e^{i(\mathbf{p} + \mathbf{q}) \cdot \mathbf{x}} + \\ &\quad + b_q^\dagger a_p^\dagger e^{i(E_q + E_p)t} e^{-i(\mathbf{p} + \mathbf{q}) \cdot \mathbf{x}} + \\ &\quad \left. - b_q^\dagger b_p e^{i(E_q - E_p)t} e^{i(\mathbf{p} - \mathbf{q}) \cdot \mathbf{x}} \right]\end{aligned}\quad (1.1.10a)$$

$$\begin{aligned}
Q_1 = i \frac{1}{(2\pi)^3} \frac{1}{2\sqrt{E_{\mathbf{q}} E_{\mathbf{p}}}} (iE_{\mathbf{p}}) \int d^3 p d^3 q & \left[a_{\mathbf{q}} a_{\mathbf{p}}^\dagger e^{i(E_{\mathbf{p}} - E_{\mathbf{q}})t} (2\pi)^3 \delta(\mathbf{q} - \mathbf{p}) + \right. \\
& - a_{\mathbf{q}} b_{\mathbf{p}} e^{-i(E_{\mathbf{q}} + E_{\mathbf{p}})t} (2\pi)^3 \delta(\mathbf{p} + \mathbf{q}) + \\
& + b_{\mathbf{q}}^\dagger a_{\mathbf{p}}^\dagger e^{i(E_{\mathbf{q}} + E_{\mathbf{p}})t} (2\pi)^3 \delta(\mathbf{p} + \mathbf{q}) + \\
& \left. - b_{\mathbf{q}}^\dagger b_{\mathbf{p}} e^{i(E_{\mathbf{q}} - E_{\mathbf{p}})t} (2\pi)^3 \delta(\mathbf{p} - \mathbf{q}) \right]
\end{aligned} \tag{1.1.10b}$$

$$Q_1 = i \frac{1}{(2\pi)^3} \frac{1}{2\sqrt{E_{\mathbf{q}} E_{\mathbf{p}}}} (iE_{\mathbf{p}}) (2\pi)^3 \int d^3 p \left[a_{\mathbf{p}} a_{\mathbf{p}}^\dagger - a_{-\mathbf{p}} b_{\mathbf{p}} e^{-2iE_{\mathbf{p}} t} + b_{-\mathbf{p}}^\dagger a_{\mathbf{p}}^\dagger e^{2iE_{\mathbf{p}} t} - b_{\mathbf{p}}^\dagger b_{\mathbf{p}} \right] \tag{1.1.10c}$$

By naming $\varphi = 2E_{\mathbf{p}} t$, we have

$$Q_1 = -\frac{1}{2} \frac{1}{(2\pi)^3} \int d^3 p \left[a_{\mathbf{p}} a_{\mathbf{p}}^\dagger - a_{-\mathbf{p}} b_{\mathbf{p}} e^{-i\varphi} + b_{-\mathbf{p}}^\dagger a_{\mathbf{p}}^\dagger e^{i\varphi} - b_{\mathbf{p}}^\dagger b_{\mathbf{p}} \right] \tag{1.1.11}$$

An analogous calculation holds for $Q_2 = -i \int d^3 x \phi^* \partial_0 \phi$

$$\phi^* \partial_0 \phi = (\phi \partial_0 \phi^*)^* = \dots \tag{1.1.12}$$

$$Q_2 = -\frac{1}{2} \frac{1}{(2\pi)^3} \int d^3 p \left[a_{\mathbf{p}}^\dagger a_{\mathbf{p}} - a_{-\mathbf{p}}^\dagger b_{\mathbf{p}}^\dagger e^{i\varphi} + b_{-\mathbf{p}} a_{\mathbf{p}} e^{-i\varphi} - b_{\mathbf{p}}^\dagger b_{\mathbf{p}} \right] \tag{1.1.13}$$

Finally

$$\begin{aligned}
Q = -\frac{1}{2} \frac{1}{(2\pi)^3} \int d^3 p & \left[a_{\mathbf{p}} a_{\mathbf{p}}^\dagger + a_{\mathbf{p}}^\dagger a_{\mathbf{p}} - b_{\mathbf{p}}^\dagger b_{\mathbf{p}} - b_{\mathbf{p}} b_{\mathbf{p}}^\dagger + e^{-i\varphi} \left(-a_{-\mathbf{p}} b_{\mathbf{p}} + b_{-\mathbf{p}} a_{\mathbf{p}} \right) + \right. \\
& \left. + e^{i\varphi} \left(b_{-\mathbf{p}}^\dagger a_{\mathbf{p}}^\dagger - a_{-\mathbf{p}}^\dagger b_{\mathbf{p}}^\dagger \right) \right]
\end{aligned} \tag{1.1.14}$$

The first two terms in (1.1.14) can be written as

$$a_{\mathbf{p}} a_{\mathbf{p}}^\dagger + a_{\mathbf{p}}^\dagger a_{\mathbf{p}} = 2a_{\mathbf{p}}^\dagger a_{\mathbf{p}} + [a_{\mathbf{p}}, a_{\mathbf{p}}^\dagger] = 2N_{\mathbf{p}}^a + (2\pi)^3 \delta(\mathbf{0}) \tag{1.1.15}$$

where we ignored the infinite c-number given by the Dirac delta. For what concerns the term containing the phase we have instead

$$I = \int_{-\infty}^{+\infty} d^3 p \left(-a_{-\mathbf{p}} b_{\mathbf{p}} + b_{-\mathbf{p}} a_{\mathbf{p}} \right) \tag{1.1.16}$$

$$\begin{cases} p' = -p \\ d^3 p' = d^3 p \end{cases} \tag{1.1.17}$$

$$\begin{aligned} I &= \int_{+\infty}^{-\infty} -d^3 p' \left(-a_{\mathbf{p}'} b_{-\mathbf{p}'} + b_{\mathbf{p}'} a_{-\mathbf{p}'} \right) = \\ &= \int_{-\infty}^{+\infty} d^3 p' \left(a_{-\mathbf{p}'} b_{\mathbf{p}'} - b_{-\mathbf{p}'} a_{\mathbf{p}'} \right) = -I \Rightarrow I = 0 \end{aligned} \quad (1.1.18)$$

where we used the commutation rules between ladder operators of different particles.

The conserved charge associated with the phase symmetry therefore is

$$Q = \int \frac{d^3 p}{(2\pi)^3} \left[N_{\mathbf{p}}^b - N_{\mathbf{p}}^a \right] \quad (1.1.19)$$

where the number operators $N_{\mathbf{p}}^a$ e $N_{\mathbf{p}}^b$ tell us the number of particles of type a and b . It is worth noting that $Q_1 = -Q_2$, so that the fields ϕ and ϕ^* describe particles with the same mass but opposite charge; we then identify particles of type b as the antiparticles of type a (and viceversa).

1.2 Yukawa Lagrangian

Text. Consider the following Lagrangian

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi + \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - m^2\phi^2 + y\bar{\psi}\psi\phi$$

where ψ is a Dirac spinor field and ϕ is a real Klein-Gordon field.

- Determine the conserved charges of the theory.
- For each conserved charge determine its associated operator as a function of the ladder operators and verify their action on single particle states.

Solution. Besides the stress-energy tensor, the Lagrangian produces another Noether current, associated with phase symmetries of the Dirac spinor fields

$$\begin{aligned} \psi &\rightarrow \psi' = e^{i\varepsilon}\psi \sim (1 + i\varepsilon)\psi \\ \bar{\psi} &\rightarrow \bar{\psi}' = e^{-i\varepsilon}\bar{\psi} \sim (1 - i\varepsilon)\bar{\psi} \end{aligned} \quad (1.2.1)$$

$$\begin{aligned} \varepsilon\bar{\delta}\psi &= \psi' - \psi = i\varepsilon & \bar{\delta}\psi &= i\psi \\ \varepsilon\bar{\delta}\bar{\psi} &= \bar{\psi}' - \bar{\psi} = -i\varepsilon & \bar{\delta}\bar{\psi} &= i\bar{\psi} \end{aligned} \quad (1.2.2)$$

The Noether current associated with this symmetry is

$$j^\mu = \frac{\partial\mathcal{L}}{\partial(\partial_\mu\psi)}\bar{\delta}\psi + \frac{\partial\mathcal{L}}{\partial(\partial_\mu\bar{\psi})}\bar{\delta}\bar{\psi} = -\bar{\psi}\gamma^\mu\psi \quad (1.2.3)$$

with Noether charge

$$Q = \int d^3x j^0 = - \int d^3x \bar{\psi}\gamma^0\psi \quad (1.2.4)$$

We write the charge Q as a function of the number operators

$$\begin{aligned} Q &= -\frac{1}{(2\pi)^6} \frac{1}{2\sqrt{E_{\mathbf{p}}E_{\mathbf{q}}}} \int d^3x d^3p d^3q \left[e^{-i\mathbf{q}\cdot\mathbf{x}} \left(\sum_r \bar{u}^r(\mathbf{q}) a_{\mathbf{q}}^{r\dagger} \right) + e^{i\mathbf{q}\cdot\mathbf{x}} \left(\sum_r \bar{v}^r(\mathbf{q}) b_{\mathbf{q}}^r \right) \right] \\ &\quad \gamma^0 \left[e^{i\mathbf{p}\cdot\mathbf{x}} \left(\sum_s u^s(\mathbf{p}) a_{\mathbf{p}}^s \right) + e^{-i\mathbf{p}\cdot\mathbf{x}} \left(\sum_s v^s(\mathbf{p}) b_{\mathbf{p}}^{s\dagger} \right) \right] \end{aligned} \quad (1.2.5)$$

Expanding the products we can make use of the Dirac delta in the form

$$\int d^3x e^{i(\mathbf{p}-\mathbf{q})\cdot\mathbf{x}} = (2\pi)^3 \delta(\mathbf{p} - \mathbf{q}) \quad (1.2.6)$$

so the charge Q becomes

$$\begin{aligned} Q = & -\frac{1}{(2\pi)^6} \frac{1}{2E_{\mathbf{p}}} \int d^3 p \left(\sum_r u^{r\dagger}(\mathbf{p}) a_{\mathbf{p}}^{r\dagger} \right) \left(\sum_s u^s(\mathbf{p}) a_{\mathbf{p}}^s \right) + \\ & + \left(\sum_r u^{r\dagger}(-\mathbf{p}) a_{-\mathbf{p}}^{r\dagger} \right) \left(\sum_s v^s(\mathbf{p}) b_{\mathbf{p}}^{s\dagger} \right) + \\ & + \left(\sum_r v^{r\dagger}(-\mathbf{p}) b_{-\mathbf{p}}^r \right) \left(\sum_s u^s(\mathbf{p}) a_{\mathbf{p}}^s \right) + \\ & + \left(\sum_r v^{r\dagger}(\mathbf{p}) b_{\mathbf{p}}^r \right) \left(\sum_s v^s(\mathbf{p}) b_{\mathbf{p}}^{s\dagger} \right) \end{aligned} \quad (1.2.7)$$

where we made explicit the dependence on the 3-momenta (since the time components do not get affected by the action of the Dirac delta) and exploited the following identities $\bar{u} = u^\dagger \gamma^0$, $\bar{v} = v^\dagger \gamma^0$. Thanks to (A.36) the second and third term are both equal to zero, while the first and the fourth one become

$$\begin{aligned} \left(\sum_r u^{r\dagger}(\mathbf{p}) a_{\mathbf{p}}^{r\dagger} \right) \left(\sum_s u^s(\mathbf{p}) a_{\mathbf{p}}^s \right) &= 2E_{\mathbf{p}} a_{\mathbf{p}}^{1\dagger} a_{\mathbf{p}}^1 + 2E_{\mathbf{p}} a_{\mathbf{p}}^{2\dagger} a_{\mathbf{p}}^2 \\ \left(\sum_r v^{r\dagger}(\mathbf{p}) b_{\mathbf{p}}^r \right) \left(\sum_s v^s(\mathbf{p}) b_{\mathbf{p}}^{s\dagger} \right) &= 2E_{\mathbf{p}} b_{\mathbf{p}}^1 b_{\mathbf{p}}^{1\dagger} + 2E_{\mathbf{p}} b_{\mathbf{p}}^2 b_{\mathbf{p}}^{2\dagger} \end{aligned} \quad (1.2.8)$$

so that

$$\begin{aligned} Q = & -\frac{1}{(2\pi)^3} \int d^3 p \left(a_{\mathbf{p}}^{1\dagger} a_{\mathbf{p}}^1 + a_{\mathbf{p}}^{2\dagger} a_{\mathbf{p}}^2 \right) + \left(b_{\mathbf{p}}^1 b_{\mathbf{p}}^{1\dagger} + b_{\mathbf{p}}^2 b_{\mathbf{p}}^{2\dagger} \right) = \\ = & -\frac{1}{(2\pi)^3} \int d^3 p \left(a_{\mathbf{p}}^{1\dagger} a_{\mathbf{p}}^1 + a_{\mathbf{p}}^{2\dagger} a_{\mathbf{p}}^2 \right) - \left(b_{\mathbf{p}}^{1\dagger} b_{\mathbf{p}}^1 + b_{\mathbf{p}}^{2\dagger} b_{\mathbf{p}}^2 \right) + \{b_{\mathbf{p}}^1, b_{\mathbf{p}}^{1\dagger}\} + \{b_{\mathbf{p}}^2, b_{\mathbf{p}}^{2\dagger}\} \end{aligned} \quad (1.2.9)$$

As usual we ignore the term proportional to $\delta(\mathbf{0})$ (see exercise 1.1) and the Noether charge becomes

$$Q = \int \frac{d^3 p}{(2\pi)^3} \left[N_{\mathbf{p}}^b - N_{\mathbf{p}}^a \right] \quad (1.2.10)$$

where the number operators contain both spin contributions. As expected, we have that the conserved charge is defined through the difference between the numbers of particles and antiparticles.

Chapter 2

Cross Sections and Decay Rates

2.1 Higgs boson decay

Text. The Higgs boson is a scalar particle whose interaction with the electromagnetic field is described by the following Lagrangian

$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - m^2\phi - \frac{1}{4}F_{\alpha\beta}F^{\alpha\beta} + \lambda\phi F_{\alpha\beta}F^{\alpha\beta}$$

where $F_{\alpha\beta}$ is the electromagnetic stress tensor, defined as

$$F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$$

Compute the decay rate of the Higgs boson ϕ .

Solution. We start by deriving the Feynman rules of the theory.

The complete Lagrangian is the sum of the Klein-Gordon Lagragian, the electromagnetic Lagrangian and the interaction one, which is given by

$$\mathcal{L}_{INT} = \lambda\phi F_{\alpha\beta}F^{\alpha\beta} \quad (2.1.1)$$


Explicitly

$$\begin{aligned} \mathcal{L}_{INT} &= \lambda\phi(\partial_\alpha A_\beta - \partial_\beta A_\alpha)(\partial^\alpha A^\beta - \partial^\beta A^\alpha) = \\ &= \lambda\phi(\partial_\alpha A_\beta \partial^\alpha A^\beta - \partial_\alpha A_\beta \partial^\beta A^\alpha - \partial_\beta A_\alpha \partial^\alpha A^\beta + \partial_\beta A_\alpha \partial^\beta A^\alpha) = \\ &= \lambda\phi(2\partial_\alpha A_\beta \partial^\alpha A^\beta - 2\partial_\alpha A_\beta \partial^\beta A^\alpha) = \\ &= 2\lambda\phi(\partial_\alpha A_\beta \partial^\alpha g^{\beta\gamma} A_\gamma - \partial_\alpha A_\beta \partial^\beta g^{\alpha\gamma} A_\gamma) \end{aligned} \quad (2.1.2)$$


Moving to the momentum space ($\partial_\mu \rightarrow ik_\mu$) we obtain

$$\begin{aligned} \mathcal{L}_{INT} &\rightarrow 2\lambda\phi(i k_{1\alpha} A_\beta i k_2^\alpha g^{\beta\gamma} A_\gamma - i k_{1\alpha} A_\beta i k_2^\beta g^{\alpha\gamma} A_\gamma) = \\ &= -2\lambda\phi(k_{1\alpha} k_2^\alpha g^{\beta\gamma} - k_1^\gamma k_2^\beta) A_\beta(k_1) A_\gamma(k_2) \end{aligned} \quad (2.1.3)$$

therefore the Feynman rules of the theory are




$$A_\beta \quad \text{---} \quad A_\gamma \quad \phi = -4i\lambda(k_{1\alpha}k_2^\alpha g^{\beta\gamma} - k_1^\gamma k_2^\beta) \quad (2.1.4)$$



$$A_\beta \quad \text{---} \quad A_\gamma \quad \phi = 4i\lambda(k_{1\alpha}k_2^\alpha g^{\beta\gamma} - k_1^\gamma k_2^\beta) \quad (2.1.5)$$

The factor i comes from the first order expansion, while the further factor of 2 is added in order to consider the indistinguishability of the photons.

Our goal is to compute the decay rate of the ϕ boson: $\phi \rightarrow \gamma\gamma$; at first order the process is described by the following diagram



The scattering amplitude is given by

$$\begin{aligned} \mathcal{A} &= 1 \cdot (-4i\lambda) \left[k_1 k_2 g^{\beta\gamma} - k_1^\gamma k_2^\beta \right] \left(\varepsilon_\beta^{\lambda_1}(k_1) \right)^* \left(\varepsilon_\gamma^{\lambda_2}(k_2) \right)^* = \\ &= -4i\lambda \left[k_1 k_2 g^{\beta\gamma} - k_1^\gamma k_2^\beta \right] \left(\varepsilon_\beta^{\lambda_1}(k_1) \right)^* \left(\varepsilon_\gamma^{\lambda_2}(k_2) \right)^* \end{aligned} \quad (2.1.7)$$

In order to compute the decay rate, we need the modulus square of the scattering matrix element, defined as

$$|\mathcal{M}|^2 = \sum_{\lambda_1, \lambda_2} \mathcal{A} \mathcal{A}^* \quad (2.1.8)$$

So we have

$$|\mathcal{M}|^2 = 16\lambda^2 \sum_{\lambda_1, \lambda_2} \left[k_1 k_2 g^{\beta\gamma} - k_1^\gamma k_2^\beta \right] \left[k_1 k_2 g^{\mu\nu} - k_1^\nu k_2^\mu \right] (\varepsilon_\beta^{\lambda_1})^* \varepsilon_\mu^{\lambda_1} (\varepsilon_\gamma^{\lambda_2})^* \varepsilon_\nu^{\lambda_2} \quad (2.1.9)$$

Summing over the photon polarization we can use the following relations

$$\begin{aligned} \sum_{\lambda_1} (\varepsilon_\beta^{\lambda_1})^* \varepsilon_\mu^{\lambda_1} &= -g_{\beta\mu} \\ \sum_{\lambda_2} (\varepsilon_\gamma^{\lambda_2})^* \varepsilon_\nu^{\lambda_2} &= -g_{\gamma\nu} \end{aligned} \quad (2.1.10)$$

so that

$$\begin{aligned} |\mathcal{M}|^2 &= 16\lambda^2 \left[k_1 k_2 g^{\beta\gamma} - k_1^\gamma k_2^\beta \right] \left[k_1 k_2 g^{\mu\nu} - k_1^\nu k_2^\mu \right] (-g_{\beta\mu})(-g_{\gamma\nu}) = \\ &= 16\lambda^2 \left[k_1 k_2 g_{\mu\nu} - k_{1\nu} k_{2\mu} \right] \left[k_1 k_2 g^{\mu\nu} - k_1^\nu k_2^\mu \right] = \\ &= 16\lambda^2 \left[4(k_1 k_2)^2 - (k_1 k_2)^2 - (k_1 k_2)^2 + \underbrace{k_1^2}_{=0} \underbrace{k_2^2}_{=0} \right] = \\ &= 32\lambda^2 (k_1 k_2)^2 \end{aligned} \quad (2.1.11)$$

The 2-body phase space element, after integrating on $d\varphi$, is

$$d\Phi_2 = \frac{1}{8\pi} \frac{|\mathbf{k}_1|}{E_1 + E_2} d\cos\theta \quad (2.1.12)$$

where E_1 and E_2 are the photon final energies.

The physics of the process must be independent on the reference frame chosen, so we can put ourselves into the most simple one. In this case it is useful to use the Higgs boson comoving frame, where the momenta satisfy

$$\begin{cases} k_1 = (E_1, 0, |\mathbf{k}| \cos\theta, |\mathbf{k}| \sin\theta) \\ k_2 = (E_2, 0, -|\mathbf{k}| \cos\theta, -|\mathbf{k}| \sin\theta) \end{cases} \quad (2.1.13)$$

and the following relations hold

$$\begin{aligned} E_1 + E_2 &= m \Rightarrow E_1 = E_2 = |\mathbf{k}| = \frac{m}{2} \\ k_1 k_2 &= E_1 E_2 + |\mathbf{k}|^2 = \frac{m^2}{2} \end{aligned} \quad (2.1.14)$$

We therefore obtain the differential decay rate

$$\begin{aligned} d\Gamma &= \frac{1}{2m} |\mathcal{M}|^2 d\Phi_2 = \left(\frac{1}{2m} \right) \left(32\lambda^2 \frac{m^4}{4} \right) \left(\frac{1}{8\pi} \frac{|\mathbf{k}_1|}{E_1 + E_2} d\cos\theta \right) = \\ &= \frac{\lambda^2}{4\pi} m^3 d\cos\theta \end{aligned} \quad (2.1.15)$$

and finally

$$\Gamma = \frac{\lambda^2}{4\pi} m^3 \int_{-1}^1 d\cos\theta = \frac{\lambda^2}{2\pi} m^3 \quad (2.1.16)$$

2.2 Scalar and standard QED: $\psi\bar{\psi} \rightarrow \phi\phi^*$

Text. Given the following Lagrangian

$$\mathcal{L} = i\bar{\psi}\gamma_\mu D^\mu\psi - m\bar{\psi}\psi + (D_\mu\phi)^*(D^\mu\phi) - m_\phi^2\phi\phi^* - \frac{1}{4}F_{\alpha\beta}F^{\alpha\beta} - \frac{1}{2\xi}(\partial_\alpha A^\alpha)^2$$

Compute the differential cross section for the process

$$\psi\bar{\psi} \rightarrow \phi\phi^*$$

Solution. First of all, we derive the Feynman rules in the momentum space, taking into account that the gauge-fixing term $\frac{1}{2\xi}(\partial_\alpha A^\alpha)^2$ does not modify the interactions

$$\begin{aligned} \mathcal{L}_{INT} &= ie[A_\mu\phi^*\partial^\mu\phi - A_\mu\phi\partial^\mu\phi^*] + e^2A_\mu A^\mu\phi\phi^* = \\ &= ie[A_\mu\phi^*\partial^\mu\phi - A^\mu\phi\partial_\mu\phi^*] + e^2g_{\mu\nu}A_\mu A_\nu\phi\phi^* \rightarrow \\ &\rightarrow -e[p_1^\mu + p_2^\mu]A_\mu\phi\phi^* + e^2g^{\mu\nu}A_\mu A_\nu\phi\phi^* \end{aligned} \quad (2.2.1)$$

$$= -ie[p_1^\mu - p_2^\mu] \quad (2.2.2)$$

$$= -ie[-p_1^\mu + p_2^\mu] \quad (2.2.3)$$

$$= -ie[-p_1^\mu - p_2^\mu] \quad (2.2.4)$$

$$\phi \quad p_1 \quad A_\mu \quad = -ie[p_1^\mu + p_2^\mu] \\ \phi^* \quad p_2 \quad A_\mu \quad (2.2.5)$$

$$\phi \quad A_\mu \\ \phi^* \quad A_\nu \quad = 2ie^2 g^{\mu\nu} \quad (2.2.6)$$

where we have added, as usual, a factor i due to the first order expansion of the S-matrix and a factor 2 in the last diagram to account for the exchange of the two photons; these rules will be used together with the usual rules for QED and the ones for the Klein-Gordon field (the former interaction is usually known as *scalar QED*).

The differential cross section of a 2-body interactions is defined as

$$d\sigma = \mathcal{J}|M|^2 d\Phi_2 \quad (2.2.7)$$

where

$$\begin{cases} \mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2 - m_1^2 m_2^2}} \\ |M|^2 = \sum_{s_j, \lambda_j, \dots} |A|^2 \\ d\Phi_2 = \frac{|\mathbf{q}_1|}{E_1 + E_2} \frac{d \cos \theta d\varphi}{16\pi^2} \Big|_{\mathbf{q}_2 = -\mathbf{q}_1} \end{cases} \quad (2.2.8)$$

Identifying the initial and final states as $|p_1, s_1; p_2, s_2\rangle$ and $|q_1; q_2\rangle$, the process $\psi\bar{\psi} \rightarrow \phi\phi^*$ is described, at first order by the following diagram

$$\bar{\psi} \quad p_2 \quad k \quad q_1 \quad \phi^* \\ \psi \quad p_1 \quad k \quad q_2 \quad \phi \quad (2.2.9)$$

where the four-momentum k is $k = p_1 + p_2 = q_1 + q_2$. We start calculating the scattering amplitude

$$\begin{aligned}\mathcal{A} &= \frac{ie^2 g_{\mu\nu}}{k^2} (-q_1^\nu + q_2^\nu) \bar{v}^{s_2}(p_2) \gamma^\mu u^{s_1}(p_1) = \\ &= -\frac{ie^2}{k^2} \bar{v}^{s_2}(p_2) (q_{1\mu} - q_{2\mu}) \gamma^\mu u^{s_1}(p_1) = \\ &= -\frac{ie^2}{k^2} \bar{v}^{s_2}(p_2) (\not{q}_1 - \not{q}_2) u^{s_1}(p_1)\end{aligned}\tag{2.2.10}$$

where $\not{q} := q_\mu \gamma^\mu$. We therefore compute the invariant matrix element averaging on the initial particles spins

$$\begin{aligned}|\mathcal{M}|^2 &= \sum_{s_1, s_2} |\mathcal{A}|^2 = \frac{e^4}{(k^2)^2} \sum_{s_1, s_2} \bar{v}^{s_2} (\not{q}_1 - \not{q}_2) u^{s_1} (u^{s_1})^\dagger (\not{q}_1 - \not{q}_2)^\dagger (\bar{v}^{s_2})^\dagger = \\ &= \frac{e^4}{(k^2)^2} \sum_{s_1, s_2} \bar{v}^{s_2} (\not{q}_1 - \not{q}_2) u^{s_1} \bar{u}^{s_1} \gamma^0 (\gamma^0 \not{q}_1 \gamma^0 - \gamma^0 \not{q}_2 \gamma^0) \gamma^0 v^{s_2} = \\ &= \frac{e^4}{(k^2)^2} \sum_{s_1, s_2} \bar{v}^{s_2} (\not{q}_1 - \not{q}_2) u^{s_1} \bar{u}^{s_1} (\not{q}_1 - \not{q}_2) v^{s_2} =\end{aligned}\tag{2.2.11}$$

Now we make explicit the tensorial products

$$\begin{aligned}|\mathcal{M}|^2 &= \frac{e^4}{(k^2)^2} \sum_{s_1, s_2} \sum_{a, b, c, d} (\bar{v}^{s_2})_a (\not{q}_1 - \not{q}_2)_{ab} (u^{s_1})_b (\bar{u}^{s_1})_c (\not{q}_1 - \not{q}_2)_{cd} (v^{s_2})_d = \\ &= \frac{e^4}{(k^2)^2} \sum_{a, b, c, d} (\not{q}_1 - \not{q}_2)_{ab} (\not{p}_1 + m)_{bc} (\not{q}_1 - \not{q}_2)_{cd} (\not{p}_2 - m)_{da} =\end{aligned}\tag{2.2.12}$$

This corresponds to the computation of the following trace

$$|\mathcal{M}|^2 = \frac{e^4}{(k^2)^2} Tr \left[(\not{q}_1 - \not{q}_2)(\not{p}_1 + m)(\not{q}_1 - \not{q}_2)(\not{p}_2 - m) \right] \tag{2.2.13}$$

Recall that the trace of an odd number of gamma matrices is null; therefore we can simplify the above expression in

$$|\mathcal{M}|^2 = \frac{e^4}{(k^2)^2} \left\{ Tr \left[(\not{q}_1 - \not{q}_2)\not{p}_1(\not{q}_1 - \not{q}_2)\not{p}_2 \right] - m^2 Tr \left[(\not{q}_1 - \not{q}_2)(\not{q}_1 - \not{q}_2) \right] \right\} \tag{2.2.14}$$

The explicit calculation is quite lengthy. For the first trace we have

$$Tr \left[(\not{q}_1 - \not{q}_2)\not{p}_1(\not{q}_1 - \not{q}_2)\not{p}_2 \right] = Tr \left[\not{q}_1 \not{p}_1 \not{q}_1 \not{p}_2 - \not{q}_1 \not{p}_1 \not{q}_2 \not{p}_2 - \not{q}_2 \not{p}_1 \not{q}_1 \not{p}_2 + \not{q}_2 \not{p}_1 \not{q}_2 \not{p}_2 \right] \tag{2.2.15}$$

so that

$$\begin{aligned} Tr \left[(\not{q}_1 - \not{q}_2) \not{p}_1 (\not{q}_1 - \not{q}_2) \not{p}_2 \right] &= Tr \left[(q_{1\mu} p_{1\nu} q_{1\rho} p_{2\sigma} - q_{1\mu} p_{1\nu} q_{2\rho} p_{2\sigma} - q_{2\mu} p_{1\nu} q_{1\rho} p_{2\sigma} + \right. \\ &\quad \left. + q_{2\mu} p_{1\nu} q_{2\rho} p_{2\sigma}) \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \right] \end{aligned} \quad (2.2.16)$$

For the trace of the four gamma matrices we have

$$\begin{aligned} Tr \left[(\not{q}_1 - \not{q}_2) \not{p}_1 (\not{q}_1 - \not{q}_2) \not{p}_2 \right] &= (q_{1\mu} p_{1\nu} q_{1\rho} p_{2\sigma} - q_{1\mu} p_{1\nu} q_{2\rho} p_{2\sigma} - \\ &\quad - q_{2\mu} p_{1\nu} q_{1\rho} p_{2\sigma} + q_{2\mu} p_{1\nu} q_{2\rho} p_{2\sigma}) \cdot \\ &\quad \cdot 4(g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma} + g^{\mu\sigma} g^{\nu\rho}) \end{aligned} \quad (2.2.17)$$

With some patience we can carry out the various contractions, to obtain

$$\begin{aligned} Tr \left[(\not{q}_1 - \not{q}_2) \not{p}_1 (\not{q}_1 - \not{q}_2) \not{p}_2 \right] &= 4 \left\{ (p_1 p_2) \left[-(q_1)^2 - (q_2)^2 + 2(q_1 q_2) \right] + \right. \\ &\quad + (q_1 p_2) \left[(p_1 q_1) - (p_1 q_2) \right] + \\ &\quad + (q_2 p_2) \left[(p_1 q_2) - (p_1 q_1) \right] + \\ &\quad + (q_1 p_1) \left[(q_1 p_2) - (q_2 p_2) \right] + \\ &\quad \left. + (q_2 p_1) \left[(q_2 p_2) - (q_1 p_2) \right] \right\} \end{aligned} \quad (2.2.18)$$

For the second trace instead we have

$$\begin{aligned} Tr \left[(\not{q}_1 - \not{q}_2) (\not{q}_1 - \not{q}_2) \right] &= Tr \left[\not{q}_1 \not{q}_1 - \not{q}_1 \not{q}_2 - \not{q}_2 \not{q}_1 + \not{q}_2 \not{q}_2 \right] = \\ &= Tr \left[(q_{1\mu} q_{1\nu} + q_{2\mu} q_{2\nu}) \gamma^\mu \gamma^\nu \right] - Tr \left[\{ \not{q}_1, \not{q}_2 \} \right] = \\ &= (q_{1\mu} q_{1\nu} + q_{2\mu} q_{2\nu}) \cdot 4g^{\mu\nu} - q_{1\mu} q_{2\nu} \cdot 2g^{\mu\nu} \cdot 4 = \\ &= 4 \left[(q_1)^2 + (q_2)^2 - 2(q_1 q_2) \right] \end{aligned} \quad (2.2.19)$$

Putting all together and making explicit $k^2 = (p_1)^2 + (p_2)^2 + 2(p_1 p_2) = 2m^2 + 2(p_1 p_2)$

and $(q_1)^2 = (q_2)^2 = m_\phi^2$, the final expression for $|\mathcal{M}|^2$ is

$$|\mathcal{M}|^2 = \frac{e^4}{[m^2 + (p_1 p_2)]^2} \left\{ \left[(p_1 p_2) - m^2 \right] \left[-2m_\phi^2 + 2(q_1 q_2) \right] + \right. \\ \left. + 2 \left[(p_1 q_1) - (p_1 q_2) \right] \left[(q_1 p_2) - (q_2 p_2) \right] \right\} \quad (2.2.20)$$

The flux factor is

$$\mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2 - m_1^2 m_2^2}} = \frac{1}{4\sqrt{(p_1 p_2)^2 - m^4}} \quad (2.2.21)$$

and the 2-body phase space element is

$$d\Phi_2 = \frac{|\mathbf{q}_1|}{E_1 + E_2} \frac{d \cos \theta d\varphi}{16\pi^2} = \frac{1}{8\pi} \frac{|\mathbf{q}_1|}{E_1 + E_2} d \cos \theta \quad (2.2.22)$$

where E_1 and E_2 stand for the final energies. In the centre of mass frame, setting the two final particles 3-momenta equal and opposite, and enforcing 4-momentum conservation we have

$$\begin{cases} p_1 = (E_1, 0, 0, |\mathbf{p}|) \\ p_2 = (E_2, 0, 0, -|\mathbf{p}|) \\ q_1 = (E_3, 0, |\mathbf{q}| \sin \theta, |\mathbf{q}| \cos \theta) \\ q_2 = (E_4, 0, -|\mathbf{q}| \sin \theta, -|\mathbf{q}| \cos \theta) \end{cases} \quad (2.2.23)$$

The energies are not independent from each other, since

$$\begin{aligned} (p_1)^2 = (p_2)^2 = m^2 &\implies E_1^2 - m^2 = E_2^2 - m^2 \implies E_1 = E_2 \\ (q_1)^2 = (q_2)^2 = m_\phi^2 &\implies E_3^2 - m_\phi^2 = E_4^2 - m_\phi^2 \implies E_3 = E_4 \\ E_1 + E_2 = E_3 + E_4 &\implies 2E_1 = 2E_3 \implies E_1 = E_3 \end{aligned} \quad (2.2.24)$$

It follows that $E_1 = E_2 = E_3 = E_4 = E$, so

$$\begin{cases} p_1 = (E, 0, 0, |\mathbf{p}|) \\ p_2 = (E, 0, 0, -|\mathbf{p}|) \\ q_1 = (E, 0, |\mathbf{q}| \sin \theta, |\mathbf{q}| \cos \theta) \\ q_2 = (E, 0, -|\mathbf{q}| \sin \theta, -|\mathbf{q}| \cos \theta) \end{cases} \quad (2.2.25)$$

$$\begin{aligned} p_1 p_2 &= 2E^2 - m^2 \\ q_1 q_2 &= 2E^2 - m_\phi^2 \\ p_1 q_1 = q_2 p_2 &= E^2 - \sqrt{(E^2 - m^2)(E^2 - m_\phi^2)} \cos \theta \\ p_1 q_2 = q_1 p_2 &= E^2 + \sqrt{(E^2 - m^2)(E^2 - m_\phi^2)} \cos \theta \end{aligned} \quad (2.2.26)$$

and the differential cross-section of the process, after some simplifications becomes

$$\frac{d\sigma}{d \cos \theta} = \frac{e^4}{2^7 \pi} \frac{\sqrt{(E^2 - m_\phi^2)^3 (E^2 - m^2)}}{E^6} (1 - 2 \cos^2 \theta) \quad (2.2.27)$$

2.3 Scalar-boson interaction: $\phi\phi \rightarrow \gamma\gamma$

Text. Given the Lagrangian


$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - m^2\phi - \frac{1}{4}F_{\alpha\beta}F^{\alpha\beta} - \frac{1}{2\xi}(\partial^\alpha A_\alpha)^2 + \lambda\phi F_{\alpha\beta}F^{\alpha\beta}$$

Compute the differential cross section for the process


$$\phi\phi \rightarrow \gamma\gamma$$

Solution. Since the gauge-fixing term does not affect the scattering amplitude, we can use the Feynman rules already derived in Exercise 2.1.

At first order we have the following Feynman diagram (t -channel)



where $q = k_1 - p_1 = p_2 - k_2$, and a second one with the final state photons exchanged (u -channel)



where $q = k_2 - p_1 = p_2 - k_1$.

Setting $|p_1; p_2\rangle \rightarrow |k_1, \lambda_1; k_2, \lambda_2\rangle$, the scattering amplitude of the first diagram is

$$\begin{aligned}
\mathcal{A} &= \left(\varepsilon_\beta^1\right)^* \cdot 4i\lambda \left[(k_1 q) g^{\mu\beta} - k_1^\beta q^\mu \right] \cdot \left(\frac{-ig_{\mu\nu}}{q^2} \right) \\
&\quad \cdot \left(\varepsilon_\gamma^2\right)^* \cdot (-4i\lambda) \left[(q k_2) g^{\gamma\nu} - q^\gamma k_2^\nu \right] = \\
&= \frac{-16i\lambda^2}{q^2} \left(\varepsilon_\beta^1\right)^* \left(\varepsilon_\gamma^2\right)^* g_{\mu\nu} \left[(k_1 q) g^{\mu\beta} - k_1^\beta q^\mu \right] \left[(q k_2) g^{\gamma\nu} - q^\gamma k_2^\nu \right] = \\
&= \frac{-16i\lambda^2}{q^2} \left(\varepsilon_\beta^1\right)^* \left(\varepsilon_\gamma^2\right)^* g_{\mu\nu} \left[(k_1 q) (k_2 q) g^{\mu\beta} g^{\gamma\nu} - (k_1 q) g^{\mu\beta} q^\gamma k_2^\nu - \right. \\
&\quad \left. - (q k_2) g^{\gamma\nu} k_1^\beta q^\mu + k_1^\beta q^\mu q^\gamma k_2^\nu \right] = \\
&= \frac{-16i\lambda^2}{q^2} \left(\varepsilon_\beta^1\right)^* \left(\varepsilon_\gamma^2\right)^* \left[(k_1 q) (k_2 q) g^{\beta\gamma} - (k_1 q) q^\gamma k_2^\beta - (q k_2) k_1^\beta q^\gamma + (k_2 q) k_1^\beta q^\gamma \right]
\end{aligned} \tag{2.3.3}$$

where $\varepsilon_\mu^{\lambda_j}(k_j) =: \varepsilon_\mu^j$.

We then use (A.49) to simplify the contractions of the polarization vectors, so that the amplitude reduces to

$$\begin{aligned}
\mathcal{A} &= \frac{-16i\lambda^2}{q^2} \left[(k_2 q) g^{\beta\gamma} - q^\gamma k_2^\beta \right] (k_1 q) \left(\varepsilon_\beta^1\right)^* \left(\varepsilon_\gamma^2\right)^* \\
\mathcal{A}^* &= \frac{16i\lambda^2}{q^2} \left[(k_2 q) g^{\beta\gamma} - q^\gamma k_2^\beta \right] (k_1 q) \varepsilon_\beta^1 \varepsilon_\gamma^2
\end{aligned} \tag{2.3.4}$$

where $q = k_1 - p_1 = p_2 - k_2$.

Setting $\tilde{q} = k_2 - p_1 = p_2 - k_1$, it is easy to convince ourselves that the second amplitude $\tilde{\mathcal{A}}$ can be obtained from the first one with the following substitutions

$$\begin{cases} k_1 \leftrightarrow k_2 \\ \lambda_1 \leftrightarrow \lambda_2 \\ q \rightarrow \tilde{q} \end{cases} \tag{2.3.5}$$

so that

$$\begin{aligned}
\tilde{\mathcal{A}} &= \frac{-16i\lambda^2}{\tilde{q}^2} \left[(k_1 \tilde{q}) g^{\beta\gamma} - \tilde{q}^\gamma k_1^\beta \right] (k_2 \tilde{q}) \left(\varepsilon_\beta^2\right)^* \left(\varepsilon_\gamma^1\right)^* \\
\tilde{\mathcal{A}}^* &= \frac{16i\lambda^2}{\tilde{q}^2} \left[(k_1 \tilde{q}) g^{\beta\gamma} - \tilde{q}^\gamma k_1^\beta \right] (k_2 \tilde{q}) \varepsilon_\beta^2 \varepsilon_\gamma^1
\end{aligned} \tag{2.3.6}$$

We can then sum the square modulus of \mathcal{A} over the polarization indices

$$\sum_{\lambda_1, \lambda_2} |\mathcal{A}|^2 = \frac{2^8 \lambda^4}{(q^2)^2} (k_1 q)^2 \left[(k_2 q) g^{\beta\gamma} - q^\gamma k_2^\beta \right] \left[(k_2 q) g^{\mu\nu} - q^\nu k_2^\mu \right] \sum_{\lambda_1, \lambda_2} \left(\varepsilon_\beta^1 \right)^* \left(\varepsilon_\gamma^2 \right)^* \varepsilon_\mu^1 \varepsilon_\nu^2 \quad (2.3.7)$$

Using (A.50) we get

$$\begin{aligned} \sum_{\lambda_1, \lambda_2} |\mathcal{A}|^2 &= \frac{2^8 \lambda^4}{(q^2)^2} (k_1 q)^2 \left[(k_2 q) g^{\beta\gamma} - q^\gamma k_2^\beta \right] \left[(k_2 q) g^{\mu\nu} - q^\nu k_2^\mu \right] g_{\mu\beta} g_{\nu\gamma} = \\ &= \frac{2^8 \lambda^4}{(q^2)^2} (k_1 q)^2 \left[(k_2 q) g_{\mu\nu} - q_\nu k_{2\mu} \right] \left[(k_2 q) g^{\mu\nu} - q^\nu k_2^\mu \right] = \\ &= \frac{2^8 \lambda^4}{(q^2)^2} (k_1 q)^2 \left[2(k_2 q)^2 + q^2 k_2^2 \right] \end{aligned} \quad (2.3.8)$$

Therefore, since $k_2^2 = 0$, we have

$$\begin{aligned} \sum_{\lambda_1, \lambda_2} |\mathcal{A}|^2 &= \frac{2^9 \lambda^4}{(q^2)^2} (k_1 q)^2 (k_2 q)^2 = \\ &= \frac{2^9 \lambda^4}{(q^2)^2} (k_1 p_1)^2 (k_2 p_2)^2 \end{aligned} \quad (2.3.9)$$

With the substitutions (2.3.5) we immediately obtain

$$\sum_{\lambda_1, \lambda_2} |\tilde{\mathcal{A}}|^2 = \frac{2^9 \lambda^4}{(\tilde{q}^2)^2} (k_2 \tilde{q})^2 (k_1 \tilde{q})^2 = \frac{2^9 \lambda^4}{(\tilde{q}^2)^2} (k_2 p_1)^2 (k_1 p_2)^2 \quad (2.3.10)$$

The interference terms are obtained with analogous calculations

$$\begin{aligned} \sum_{\lambda_1, \lambda_2} \mathcal{A} \tilde{\mathcal{A}}^* &= \sum_{\lambda_1, \lambda_2} \mathcal{A}^* \tilde{\mathcal{A}} = \frac{2^8 \lambda^4}{q^2 \tilde{q}^2} (k_1 q) (k_2 \tilde{q}) \left[2(k_1 \tilde{q})(k_2 q) + (q k_1)(\tilde{q} k_2) \right] = \\ &= \frac{2^8 \lambda^4}{q^2 \tilde{q}^2} (k_1 p_1) (k_2 p_1) \left[2(k_1 p_2)(k_2 p_2) + (k_1 p_1)(k_2 p_1) \right] \end{aligned} \quad (2.3.11)$$

The invariant matrix element becomes

$$\begin{aligned} \frac{|\mathcal{M}|^2}{2^9 \lambda^4} &= \frac{(k_1 p_1)^2 (k_2 p_2)^2}{(m^2 - 2k_1 p_1)^2} + \frac{(k_2 p_1)^2 (k_1 p_2)^2}{(m^2 - 2k_1 p_2)^2} + \\ &+ \frac{(k_1 p_1)(k_2 p_1)}{(m^2 - 2k_1 p_1)(m^2 - 2k_1 p_2)} \left[2(k_1 p_2)(k_2 p_2) + (k_1 p_1)(k_2 p_1) \right] \end{aligned} \quad (2.3.12)$$

where we substituted

$$\begin{aligned} q^2 &= k_1^2 + p_1^2 - 2k_1 p_1 = m^2 - 2k_1 p_1 = \\ &= p_2^2 + k_2^2 - 2p_2 k_2 = m^2 - 2k_2 p_2 \end{aligned} \quad (2.3.13)$$

and analogous relations for \tilde{q} .

The flux factor is

$$\mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2 - m^4}} \quad (2.3.14)$$

while the 2-body phase space element is

$$d\Phi_2 = \frac{1}{8\pi} \frac{|\mathbf{k}_1|}{E_1 + E_2} d\cos\theta \quad (2.3.15)$$

where E_1 and E_2 stand for the final state energies. The calculations become simpler in the centre of mass reference frame, where the two couples of particles have 3-momenta with equal modulus and opposite direction; therefore, after imposing the conservation of four-momentum, we have

$$\begin{cases} p_1 = (E, 0, 0, |\mathbf{p}|) \\ p_2 = (E, 0, 0, -|\mathbf{p}|) \\ k_1 = (E, 0, |\mathbf{k}| \sin\theta, |\mathbf{k}| \cos\theta) \\ k_2 = (E, 0, -|\mathbf{k}| \sin\theta, -|\mathbf{k}| \cos\theta) \end{cases} \quad (2.3.16)$$

$$\begin{aligned} |\mathbf{k}| &= E \\ |\mathbf{p}| &= \sqrt{E^2 - m^2} \\ p_1 p_2 &= 2E^2 - m^2 \\ k_1 p_1 &= k_2 p_2 = E^2 - E\sqrt{E^2 - m^2} \cos\theta \\ k_1 p_2 &= k_2 p_1 = E^2 + E\sqrt{E^2 - m^2} \cos\theta \end{aligned} \quad (2.3.17)$$

After some algebra we get

$$\begin{aligned} |\mathcal{M}|^2 &= 2^9 \lambda^4 \left\{ \frac{(E^2 - E\sqrt{E^2 - m^2} \cos\theta)^4}{[m^2 - 2(E^2 - E\sqrt{E^2 - m^2} \cos\theta)]^2} + \right. \\ &\quad + \frac{(E^2 + E\sqrt{E^2 - m^2} \cos\theta)^4}{[m^2 - 2(E^2 + E\sqrt{E^2 - m^2} \cos\theta)]^2} + \\ &\quad \left. + \frac{3[E^4 - E^2(E^2 - m^2) \cos^2\theta]^2}{[m^2 - 2(E^2 - E\sqrt{E^2 - m^2} \cos\theta)][m^2 - 2(E^2 + E\sqrt{E^2 - m^2} \cos\theta)]} \right\} \end{aligned} \quad (2.3.18)$$

Finally the differential cross section for this process is

$$\frac{d\sigma}{d\cos\theta} = \frac{1}{128\pi\sqrt{E^4 - E^2m^2}} |\mathcal{M}|^2 \quad (2.3.19)$$

where $|\mathcal{M}|^2$ is given by (2.3.18).

In the ultra-relativistic limit, where we can neglect the mass of the particles, the cross section simplifies considerably

$$\frac{d\sigma}{d\cos\theta} \sim \frac{\lambda^4}{4\pi} E^2 (5 - \cos^2\theta) \quad (2.3.20)$$

2.4 Scalar-fermion-boson interaction: $e^+e^- \rightarrow \phi\gamma$

Text. Let ϕ be a neutral (i.e. real) Klein-Gordon field which interacts with photons through the following interaction

$$\mathcal{L}_{INT} = \lambda\phi F_{\alpha\beta}F^{\alpha\beta}$$

Compute the differential cross section for the process


$$e^+e^- \rightarrow \phi\gamma$$

Solution. The Lagrangian is

$$\mathcal{L} = \mathcal{L}_{QED} + \mathcal{L}_{KG} + \lambda\phi F_{\alpha\beta}F^{\alpha\beta} \quad (2.4.1)$$

The Feynman rules for the theory are the usual QED ones, to which we add those derived in the exercise 2.1.

At first order, the process $e^+e^- \rightarrow \phi\gamma$ is represented by the following diagram



where $q = p_1 + p_2 = k_1 + k_2$.

Setting $|p_1, s_1; p_2, s_2\rangle \rightarrow |k_1, \lambda; k_2\rangle$ the scattering amplitude is

$$\begin{aligned}
 \mathcal{A} &= \frac{-4ie\lambda}{q^2} g_{\mu\nu} \left[qk_1g^{\nu\delta} - q^\delta k_1^\nu \right] \left(\varepsilon_\delta^\lambda(k_1) \right)^* \bar{v}^{s_2}(p_2) \gamma^\mu u^{s_1}(p_1) = \\
 &= \frac{-4ie\lambda}{q^2} \left[qk_1g_\mu^\delta - q^\delta k_{1\mu} \right] \left(\varepsilon_\delta^\lambda(k_1) \right)^* \bar{v}^{s_2}(p_2) \gamma^\mu u^{s_1}(p_1) = \\
 &= \frac{-4ie\lambda}{q^2} \left[qk_1 \left(\varepsilon_\mu^\lambda(k_1) \right)^* - q \left(\varepsilon_\lambda^\lambda(k_1) \right)^* k_{1\mu} \right] \bar{v}^{s_2}(p_2) \gamma^\mu u^{s_1}(p_1) = \\
 &= \frac{-4ie\lambda}{q^2} qk_1 \left(\varepsilon_\mu^\lambda(k_1) \right)^* \bar{v}^{s_2}(p_2) \gamma^\mu u^{s_1}(p_1) + \frac{4ie\lambda}{q^2} q \left(\varepsilon_\lambda^\lambda(k_1) \right)^* k_{1\mu} \bar{v}^{s_2}(p_2) \gamma^\mu u^{s_1}(p_1) = \\
 &=: \mathcal{A}_1 + \mathcal{A}_2
 \end{aligned} \quad (2.4.3)$$

The invariant matrix element, averaging on the initial particles spins and summing over polarizations, is

$$|\mathcal{M}|^2 = \frac{1}{4} \sum_{s_1, s_2} \sum_{\lambda} |\mathcal{A}_1|^2 + |\mathcal{A}_2|^2 + \mathcal{A}_1^* \mathcal{A}_2 + \mathcal{A}_1 \mathcal{A}_2^* \quad (2.4.4)$$

We separately compute the various terms

$$\begin{aligned} \sum_{s_1, s_2} \sum_{\lambda} |\mathcal{A}_1|^2 &= \frac{16e^2 \lambda^2}{(q^2)^2} (q k_1)^2 \sum_{s_1, s_2} \sum_{\lambda} (\varepsilon_{\mu}^{\lambda})^* \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \left[(\varepsilon_{\nu}^{\lambda})^* \bar{v}^{s_2} \gamma^{\nu} u^{s_1} \right]^* = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} (q k_1)^2 \sum_{s_1, s_2} \sum_{\lambda} (\varepsilon_{\mu}^{\lambda})^* \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \varepsilon_{\nu}^{\lambda} \bar{u}^{s_1} \gamma^0 \gamma^0 \gamma^{\nu} \gamma^0 v^{s_2} = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} (q k_1)^2 \sum_{s_1, s_2} g_{\nu \mu} \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \bar{u}^{s_1} \gamma^{\nu} v^{s_2} = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} (q k_1)^2 g_{\nu \mu} \sum_{s_1, s_2} \sum_{a, b, c, d} (\bar{v}^{s_2})_a (\gamma^{\mu})_{ab} (u^{s_1})_b (\bar{u}^{s_1})_c (\gamma^{\nu})_{cd} (v^{s_2})_d = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} (q k_1)^2 g_{\nu \mu} \sum_{a, b, c, d} (\gamma^{\mu})_{ab} (\not{p}_1 + m)_{bc} (\gamma^{\nu})_{cd} (\not{p}_2 - m)_{da} = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} (q k_1)^2 g_{\nu \mu} Tr \left[\gamma^{\mu} (\not{p}_1 + m) \gamma^{\nu} (\not{p}_2 - m) \right] \end{aligned} \quad (2.4.5)$$

$$\begin{aligned} \sum_{s_1, s_2} \sum_{\lambda} |\mathcal{A}_2|^2 &= \frac{16e^2 \lambda^2}{(q^2)^2} \sum_{s_1, s_2} \sum_{\lambda} q^{\alpha} (\varepsilon_{\alpha}^{\lambda})^* k_{1\mu} \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \left[q^{\beta} (\varepsilon_{\beta}^{\lambda})^* k_{1\nu} \bar{v}^{s_2} \gamma^{\nu} u^{s_1} \right]^* = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} q^{\alpha} q^{\beta} k_{1\mu} k_{1\nu} \sum_{s_1, s_2} \sum_{\lambda} (\varepsilon_{\alpha}^{\lambda})^* \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \varepsilon_{\beta}^{\lambda} \bar{u}^{s_1} \gamma^{\nu} v^{s_2} = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} q^{\alpha} q^{\beta} k_{1\mu} k_{1\nu} g_{\beta\alpha} \sum_{s_1, s_2} \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \bar{u}^{s_1} \gamma^{\nu} v^{s_2} = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} q^{\alpha} q_{\alpha} k_{1\mu} k_{1\nu} \sum_{s_1, s_2} \sum_{a, b, c, d} (\bar{v}^{s_2})_a (\gamma^{\mu})_{ab} (u^{s_1})_b (\bar{u}^{s_1})_c (\gamma^{\nu})_{cd} (v^{s_2})_d = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} k_{1\mu} k_{1\nu} \sum_{a, b, c, d} (\gamma^{\mu})_{ab} (\not{p}_1 + m)_{bc} (\gamma^{\nu})_{cd} (\not{p}_2 - m)_{da} = \\ &= \frac{16e^2 \lambda^2}{(q^2)^2} k_{1\mu} k_{1\nu} Tr \left[\gamma^{\mu} (\not{p}_1 + m) \gamma^{\nu} (\not{p}_2 - m) \right] \end{aligned} \quad (2.4.6)$$

$$\begin{aligned}
\sum_{s_1, s_2} \sum_{\lambda} \mathcal{A}_1^* \mathcal{A}_2 &= \frac{-16e^2 \lambda^2}{(q^2)^2} \sum_{s_1, s_2} \sum_{\lambda} (q k_1) \varepsilon_{\mu}^{\lambda} \left[\bar{v}^{s_2} \gamma^{\mu} u^{s_1} \right]^* q (\varepsilon^{\lambda})^* k_{1\nu} \bar{v}^{s_2} \gamma^{\nu} u^{s_1} = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q^{\alpha} k_{1\nu} \sum_{s_1, s_2} \sum_{\lambda} \varepsilon_{\mu}^{\lambda} (\varepsilon_{\alpha}^{\lambda})^* \left[\bar{v}^{s_2} \gamma^{\mu} u^{s_1} \right]^* \bar{v}^{s_2} \gamma^{\nu} u^{s_1} = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q^{\alpha} k_{1\nu} g_{\mu\alpha} \sum_{s_1, s_2} \bar{u}^{s_1} \gamma^{\mu} v^{s_2} \bar{v}^{s_2} \gamma^{\nu} u^{s_1} = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q_{\mu} k_{1\nu} \sum_{s_1, s_2} \sum_{a, b, c, d} (\bar{u}^{s_1})_a (\gamma^{\mu})_{ab} (v^{s_2})_b (\bar{v}^{s_2})_c (\gamma^{\nu})_{cd} (u^{s_1})_d = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q_{\mu} k_{1\nu} \sum_{a, b, c, d} (\gamma^{\mu})_{ab} (\not{p}_2 - m)_{bc} (\gamma^{\nu})_{cd} (\not{p}_1 + m)_{da} = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q_{\mu} k_{1\nu} \text{Tr} \left[\gamma^{\mu} (\not{p}_2 - m) \gamma^{\nu} (\not{p}_1 + m) \right]
\end{aligned} \tag{2.4.7}$$

$$\begin{aligned}
\sum_{s_1, s_2} \sum_{\lambda} \mathcal{A}_1 \mathcal{A}_2^* &= \frac{-16e^2 \lambda^2}{(q^2)^2} \sum_{s_1, s_2} \sum_{\lambda} (q k_1) (\varepsilon_{\mu}^{\lambda})^* \bar{v}^{s_2} \gamma^{\mu} u^{s_1} q \varepsilon^{\lambda} k_{1\nu} \left[\bar{v}^{s_2} \gamma^{\nu} u^{s_1} \right]^* = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q^{\alpha} k_{1\nu} \sum_{s_1, s_2} \sum_{\lambda} (\varepsilon_{\mu}^{\lambda})^* \varepsilon_{\alpha}^{\lambda} \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \left[\bar{v}^{s_2} \gamma^{\nu} u^{s_1} \right]^* = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q^{\alpha} k_{1\nu} g_{\alpha\mu} \sum_{s_1, s_2} \bar{v}^{s_2} \gamma^{\mu} u^{s_1} \bar{u}^{s_1} \gamma^{\nu} v^{s_2} = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q_{\mu} k_{1\nu} \sum_{s_1, s_2} \sum_{a, b, c, d} (\bar{v}^{s_2})_a (\gamma^{\mu})_{ab} (u^{s_1})_b (\bar{u}^{s_1})_c (\gamma^{\nu})_{cd} (v^{s_2})_d = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q_{\mu} k_{1\nu} \sum_{a, b, c, d} (\gamma^{\mu})_{ab} (\not{p}_1 + m)_{bc} (\gamma^{\nu})_{cd} (\not{p}_2 - m)_{da} = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (q k_1) q_{\mu} k_{1\nu} \text{Tr} \left[\gamma^{\mu} (\not{p}_1 + m) \gamma^{\nu} (\not{p}_2 - m) \right]
\end{aligned} \tag{2.4.8}$$

Thanks to the cyclic property of the trace (A.29), the two "mixed" terms turn out

to be equal

$$\begin{aligned}
\sum_{s_1, s_2} \sum_{\lambda} \mathcal{A}_1^* \mathcal{A}_2 &= \frac{-16e^2 \lambda^2}{(q^2)^2} (qk_1) q_\mu k_{1\nu} Tr \left[\gamma^\mu (\not{p}_2 - m) \gamma^\nu (\not{p}_1 + m) \right] = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (qk_1) q_\mu k_{1\nu} Tr \left[(\not{p}_2 - m) \gamma^\nu (\not{p}_1 + m) \gamma^\mu \right] = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (qk_1) q_\mu k_{1\nu} Tr \left[\gamma^\nu (\not{p}_1 + m) \gamma^\mu (\not{p}_2 - m) \right] = \\
&= \frac{-16e^2 \lambda^2}{(q^2)^2} (qk_1) q_\nu k_{1\mu} Tr \left[\gamma^\mu (\not{p}_1 + m) \gamma^\nu (\not{p}_2 - m) \right] = \\
&= \sum_{s_1, s_2} \sum_{\lambda} \mathcal{A}_1 \mathcal{A}_2^*
\end{aligned} \tag{2.4.9}$$

The computation of the scattering amplitude then reduces to

$$|\mathcal{M}|^2 = \frac{4e^2 \lambda^2}{q^2} \left[\frac{1}{q^2} (qk_1)^2 g_{\nu\mu} + k_{1\mu} k_{2\nu} - \frac{2}{q^2} (qk_1) q_\mu k_{1\nu} \right] Tr \left[\gamma^\mu (\not{p}_1 + m) \gamma^\nu (\not{p}_2 - m) \right] \tag{2.4.10}$$

that is

$$\begin{aligned}
|\mathcal{M}|^2 &= \frac{4e^2 \lambda^2}{q^2} \left[\frac{1}{q^2} (qk_1)^2 g_{\nu\mu} + k_{1\mu} k_{2\nu} - \frac{2}{q^2} (qk_1) q_\mu k_{1\nu} \right] \cdot \\
&\quad \cdot \left[p_1^\mu p_2^\nu - (p_1 p_2) g^{\mu\nu} + p_1^\nu p_2^\mu - m^2 g^{\mu\nu} \right]
\end{aligned} \tag{2.4.11}$$

Carrying out the contractions we obtain

$$\begin{aligned}
|\mathcal{M}|^2 &= \frac{4e^2 \lambda^2}{q^2} \left\{ \frac{1}{q^2} (qk_1)^2 \left[-2(p_1 p_2) - 4m^2 \right] + \right. \\
&\quad + \left[(k_1 p_1)(k_2 p_2) - (k_1 k_2)(p_1 p_2) + (k_1 p_2)(k_2 p_1) - m^2 (k_1 k_2) \right] - \\
&\quad \left. - \frac{2}{q^2} (qk_1) \left[(q p_1)(k_1 p_2) - (q k_1)(p_1 p_2) + (q p_2)(k_1 p_1) - m^2 (q k_1) \right] \right\}
\end{aligned} \tag{2.4.12}$$

Therefore we have

$$\begin{aligned}
|\mathcal{M}|^2 &= \frac{4e^2 \lambda^2}{q^4} \left\{ -2(qk_1)^2 (p_1 p_2) - 4m^2 (qk_1)^2 + q^2 (k_1 p_1)(k_2 p_2) - \right. \\
&\quad - q^2 (k_1 k_2)(p_1 p_2) + q^2 (k_1 p_2)(k_2 p_1) - m^2 q^2 (k_1 k_2) - \\
&\quad - 2(qk_1)(q p_1)(k_1 p_2) + 2(qk_1)^2 (p_1 p_2) - \\
&\quad \left. - 2(qk_1)(q p_2)(k_1 p_1) + 2m^2 (qk_1)^2 \right\}
\end{aligned} \tag{2.4.13}$$

The expressions for the flux factor, the phase space element and the 4-momenta in the centre of mass frame are the usual ones

$$\mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2 - m^4}} \quad (2.4.14)$$

$$d\Phi_2 = \frac{1}{8\pi} \frac{|\mathbf{k}_1|}{E_\gamma + E_\phi} d\cos\theta \quad (2.4.15)$$

$$\begin{cases} p_1 = (E, 0, 0, |\mathbf{p}|) \\ p_2 = (E, 0, 0, -|\mathbf{p}|) \\ k_1 = (E_\gamma, 0, |\mathbf{k}| \sin\theta, |\mathbf{k}| \cos\theta) \\ k_2 = (E_\phi, 0, -|\mathbf{k}| \sin\theta, -|\mathbf{k}| \cos\theta) \end{cases} \quad (2.4.16)$$

where $|\mathbf{p}| = \sqrt{E^2 - m^2}$ and $|\mathbf{k}| = E_\gamma = \sqrt{E_\phi^2 - m_\phi^2}$.

Recalling that $q = p_1 + p_2 = k_1 + k_2$, we can make explicit the scalar products

$$\begin{aligned} (qk_1) &= (k_1 k_2) = E_\gamma E_\phi + E_\gamma^2 \\ (qp_1) &= (qp_2) = 2E^2 \\ (p_1 p_2) &= 2E^2 - m^2 \\ (k_1 p_1) &= E_\gamma E - E_\gamma \sqrt{E^2 - m^2} \cos\theta \\ (k_2 p_2) &= E_\phi E - E_\gamma \sqrt{E^2 - m^2} \cos\theta \\ (k_1 p_2) &= E_\gamma E + E_\gamma \sqrt{E^2 - m^2} \cos\theta \\ (k_2 p_1) &= E_\phi E + E_\gamma \sqrt{E^2 - m^2} \cos\theta \\ q^2 &= 4E^2 = m_\phi^2 + 2E_\gamma E_\phi + 2E_\gamma^2 \end{aligned} \quad (2.4.17)$$

After some simplification we get

$$\begin{aligned} |\mathcal{M}|^2 &= \frac{4e^2 \lambda^2}{16E^4} \left\{ -2m^2 \left[E_\gamma^2 E_\phi^2 + E_\gamma^4 + 2E_\gamma^3 E_\phi \right] - 8E^3 E_\gamma \left[E_\gamma E_\phi + E_\gamma^2 - EE_\gamma \right] + \right. \\ &\quad \left. + 8E^2 E_\gamma^2 (E^2 - m^2) \cos^2 \theta \right\} \end{aligned} \quad (2.4.18)$$

and finally

$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \frac{e^2 \lambda^2}{256\pi} \frac{E_\gamma}{E^5 \sqrt{E^4 - E^2 m^2}} \left\{ -m^2 \left[E_\gamma^2 E_\phi^2 + E_\gamma^4 + 2E_\gamma^3 E_\phi \right] - \right. \\ &\quad - 4E^3 E_\gamma \left[E_\gamma E_\phi + E_\gamma^2 - EE_\gamma \right] + \\ &\quad \left. + 4E^2 E_\gamma^2 (E^2 - m^2) \cos^2 \theta \right\} \end{aligned} \quad (2.4.19)$$

2.5 Yukawa Lagrangian: $f\bar{f} \rightarrow \phi\phi$

Text. Consider the following lagrangian density


$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi + \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - m^2\phi^2 + y\bar{\psi}\psi\phi$$

where ψ is a Dirac spinor field and ϕ a real scalar field. Compute the differential cross section for the process


$$f\bar{f} \rightarrow \phi\phi$$

where f (\bar{f}) stands for the fermionic particle (antiparticle) and ϕ for the scalar one.


Solution. Besides the usual Feynman rules for QED, we also have the following one, due to the interaction between Dirac fields and the Klein-Gordon one (*Yukawa interaction term in the Lagrangian*).



The Feynman diagrams associated with the process in question are the following



where $q_1 = p_1 - p_3$, and



where $q_2 = p_1 - p_4$.

The two corresponding amplitudes are

$$\mathcal{A}_1 = \bar{v}^{s_2}(p_2) \left(-i\lambda \right) \frac{i(q_1 + m_f)}{q_1^2 - m_f^2} \left(-i\lambda \right) u^{s_1}(p_1) \quad (2.5.4)$$

$$\mathcal{A}_2 = \bar{v}^{s_2}(p_2) \left(-i\lambda \right) \frac{i(q_2 + m_f)}{q_2^2 - m_f^2} \left(-i\lambda \right) u^{s_1}(p_1) \quad (2.5.5)$$

We now substitute the expressions for q_1 and q_2 in the respective amplitudes. Regarding \mathcal{A}_1 , in the denominator we have

$$q_1^2 - m_f^2 = \underbrace{p_1^2}_{=m_f^2} + \underbrace{p_3^2}_{=m_\phi^2} - 2p_1 \cdot p_3 - m_f^2 = m_\phi^2 - 2p_1 \cdot p_3 \quad (2.5.6)$$

The numerator is instead

$$(q_1 + m_f)u^{s_1}(p_1) = (\not{p}_1 - \not{p}_3 + m_f)u^{s_1}(p_1) = (2m_f - \not{p}_3)u^{s_1}(p_1) \quad (2.5.7)$$

where we have exploited the Dirac equation $(\not{p} - m)u(p) = 0$.

Carrying out the same passages for \mathcal{A}_2 one obtains

$$\mathcal{A}_1 = \underbrace{\frac{-i\lambda^2}{m_\phi^2 - 2p_1 \cdot p_3}}_{=:A} \bar{v}^{s_2}(p_2) \left(2m_f - \not{p}_3 \right) u^{s_1}(p_1) \quad (2.5.8)$$

$$\mathcal{A}_2 = \underbrace{\frac{-i\lambda^2}{m_\phi^2 - 2p_1 \cdot p_4}}_{=:B} \bar{v}^{s_2}(p_2) \left(2m_f - \not{p}_4 \right) u^{s_1}(p_1) \quad (2.5.9)$$

The invariant matrix element, averaged on the initial spins, is

$$\begin{aligned}
|\mathcal{M}|^2 &= \frac{1}{4} \sum_{s_1, s_2} [\mathcal{A}_1 \mathcal{A}_1^* + \mathcal{A}_2 \mathcal{A}_2^* + \mathcal{A}_1 \mathcal{A}_2^* + \mathcal{A}_1^* \mathcal{A}_2] = \\
&= \frac{\lambda^4}{4} \sum_{s_1, s_2} \left\{ \frac{1}{A^2} \bar{v}^{s_2} \left(2m_f - p_3 \right) u^{s_1} \bar{u}^{s_1} \left(2m_f - p_3 \right) v^{s_2} + \right. \\
&\quad + \frac{1}{B^2} \bar{v}^{s_2} \left(2m_f - p_4 \right) u^{s_1} \bar{u}^{s_1} \left(2m_f - p_4 \right) v^{s_2} + \\
&\quad + \frac{1}{AB} \left[\bar{v}^{s_2} \left(2m_f - p_3 \right) u^{s_1} \bar{u}^{s_1} \left(2m_f - p_4 \right) v^{s_2} + \right. \\
&\quad \left. \left. + \bar{v}^{s_2} \left(2m_f - p_4 \right) u^{s_1} \bar{u}^{s_1} \left(2m_f - p_3 \right) v^{s_2} \right] \right\} = \\
&= \frac{\lambda^4}{4} \left\{ \frac{1}{A^2} \underbrace{Tr[(p_1 + m_f)(2m_f - p_3)(p_2 - m_f)(2m_f - p_3)]}_{(1)} + \right. \\
&\quad + \frac{1}{B^2} \underbrace{Tr[(p_1 + m_f)(2m_f - p_4)(p_2 - m_f)(2m_f - p_4)]}_{(2)} + \\
&\quad + \frac{1}{AB} \left(\underbrace{Tr[(p_1 + m_f)(2m_f - p_4)(p_2 - m_f)(2m_f - p_3)]}_{(3)} + \right. \\
&\quad \left. \left. + \underbrace{Tr[(p_1 + m_f)(2m_f - p_3)(p_2 - m_f)(2m_f - p_4)]}_{(4)} \right) \right\} \tag{2.5.10}
\end{aligned}$$

The four traces share the same structure, therefore it is sufficient to compute only one of them; the other traces will be obtained with appropriate substitutions. Let's see the computation of the trace (3).

$$\begin{aligned}
(3) &= Tr[(p_1 + m_f)(2m_f - p_4)(p_2 - m_f)(2m_f - p_3)] = \\
&= Tr[(p_1 + m_f)p_4(p_2 - m_f)p_3] + 4m_f^2 Tr[(p_1 + m_f)(p_2 - m_f)] = \\
&= Tr[p_1 p_4 p_2 p_3] - m_f^2 Tr[p_4 p_3] + 4m_f^2 Tr[p_1 p_2] - 16m_f^4 = \\
&= 4[(p_1 p_4)(p_2 p_3) - (p_1 p_2)(p_4 p_3) + (p_1 p_3)(p_4 p_2)] - 4m_f^2(p_4 p_3) + 16m_f^2(p_1 p_2) - 16m_f^4
\end{aligned} \tag{2.5.11}$$

The result of trace number (3) is symmetric under the exchange $p_3 \longleftrightarrow p_4$, so that (3) = (4).

Computing the first trace we get

$$(1) = 4[2(p_1 p_3)(p_2 p_3) - (p_1 p_2)p_3^2] - 4m_f^2 p_3^2 + 16m_f^2(p_1 p_2) - 16m_f^4 \tag{2.5.12}$$

Finally, trace (2) can be obtained from trace (1) substituting p_3 with p_4 . In the centre of mass reference frame the 4-momenta can be written as

$$\begin{cases} p_1 = (E_1, 0, 0, |\mathbf{p}|) \\ p_2 = (E_2, 0, 0, -|\mathbf{p}|) \\ p_3 = (E_3, \mathbf{q}) \\ p_4 = (E_4, -\mathbf{q}) \end{cases} \quad (2.5.13)$$

$$\begin{aligned} E_1 &= E_2 =: E \\ E_3 &= E_4 =: E' \\ p_1^2 &= p_2^2 = m_f^2 \\ p_3^2 &= p_4^2 = m_\phi^2 \end{aligned} \quad (2.5.14)$$

By introducing the Mandelstam variables

$$\begin{aligned} s &:= (p_1 + p_2)^2 \\ t &:= (p_1 - p_3)^2 \\ u &:= (p_1 - p_4)^2 \end{aligned} \quad (2.5.15)$$

the products of the 4-momenta now read

$$\begin{aligned} p_1 p_2 &= \frac{1}{2}(s - 2m_f^2) \\ p_1 p_3 = p_2 p_4 &= \frac{1}{2}(m_f^2 + m_\phi^2 - t) \\ p_1 p_4 = p_2 p_3 &= \frac{1}{2}(m_f^2 + m_\phi^2 - u) \\ p_3 p_4 &= \frac{1}{2}(s - 2m_\phi^2) \end{aligned} \quad (2.5.16)$$

where we made use of four-momentum conservation between initial and final states and also used the relation $s + t + u = \sum_i m_i^2$ (which is valid in general), where m_i are the masses of all the external particles involved in the process.

$$\begin{aligned} |\mathcal{M}|^2 &= \frac{\lambda^4}{4} \left\{ \left(\frac{1}{(t - m_f^2)^2} + \frac{1}{(u - m_f^2)^2} \right) \left[2(m_f^2 + m_\phi^2 - t)(m_f^2 + m_\phi^2 - u) - \right. \right. \\ &\quad \left. \left. - 4m_f^2 m_\phi^2 + 8m_f^2(s - 4m_f^2) \right] + \right. \\ &\quad \left. + \frac{2}{(t - m_f^2)(u - m_f^2)} \left[(m_f^2 + m_\phi^2 - t)^2 (m_f^2 + m_\phi^2 - u)^2 - \right. \right. \\ &\quad \left. \left. - (s - 2m_f^2)(s - 2m_\phi^2) + 2m_f^2 \left(4(s - 2m_f^2) - (s - 2m_\phi^2) \right) - 16m_f^4 \right] \right\} \end{aligned} \quad (2.5.17)$$

The flux factor is

$$\mathcal{J} = \frac{1}{\sqrt{4(s - 2m_f^2)^2 - 16m_f^4}} \quad (2.5.18)$$

while the two-body phase space is

$$d\Phi_2 = \frac{\sqrt{\frac{s}{4} - m_\phi^2}}{\sqrt{s}/2} \frac{d\cos\theta d\varphi}{16\pi^2} \quad (2.5.19)$$

therefore, after integrating over $d\varphi$, the differential cross section for the process is

$$\frac{d\sigma}{d\cos\theta} = \frac{1}{16\pi} \sqrt{\frac{1 - \frac{4m_\phi^2}{s}}{(s - 2m_f^2)^2 - 4m_f^4} |\mathcal{M}|^2} \quad (2.5.20)$$

where $|\mathcal{M}|^2$ is given by (2.5.17).

2.6 Fermi Lagrangian: $\nu e^- \rightarrow \nu e^-$

Text. Consider the standard QED Lagrangian for the electron; add to that a neutral and massless spinor field ψ_ν , described by the following Lagrangian

$$\mathcal{L} = i\bar{\psi}_\nu \not{\partial} \psi_\nu + G\bar{\psi}_\nu \gamma_\alpha \psi_\nu \bar{\psi}_e \gamma^\alpha \psi_e + \mu \bar{\psi}_\nu \sigma_{\alpha\beta} F^{\alpha\beta} \psi_\nu$$

Compute the differential cross section for the process

$$\nu e^- \rightarrow \nu e^-$$

Solution. The full Lagrangian of the problem is

$$\mathcal{L} = \bar{\psi}_e (i\not{\partial} - m_e) \psi_e - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i\bar{\psi}_\nu \not{\partial} \psi_\nu + G\bar{\psi}_\nu \gamma_\alpha \psi_\nu \bar{\psi}_e \gamma^\alpha \psi_e + \mu \bar{\psi}_\nu \sigma_{\alpha\beta} F^{\alpha\beta} \psi_\nu \quad (2.6.1)$$

In addition to the usual QED vertex we also have the following new ones

$$= iG\gamma_\alpha\gamma^\alpha \quad (2.6.2)$$

$$= 2i\mu(\gamma^\sigma k - k^\sigma) \quad (2.6.3)$$


This last vertex factor was computed as follows

$$\begin{aligned} \mathcal{L}_{INT} &= \mu \bar{\psi}_\nu \sigma^{\alpha\beta} F_{\alpha\beta} \psi_\nu = \mu \bar{\psi}_\nu \frac{i}{2} [\gamma^\alpha, \gamma^\beta] (\partial_\alpha A_\beta - \partial_\beta A_\alpha) \psi_\nu = \\ &= i \frac{\mu}{2} \bar{\psi}_\nu (\gamma^\alpha \gamma^\beta - \gamma^\beta \gamma^\alpha) (\partial_\alpha A_\beta - \partial_\beta A_\alpha) \psi_\nu = i \frac{\mu}{2} \bar{\psi}_\nu 2(\gamma^\alpha \gamma^\beta \partial_\alpha A_\beta - \gamma^\alpha \gamma^\beta \partial_\beta A_\alpha) = \\ &= i \mu \bar{\psi}_\nu \gamma^\alpha \gamma^\beta (g_\beta^\sigma \partial_\alpha - g_\alpha^\sigma \partial_\beta) A_\sigma = i \mu \bar{\psi}_\nu (\gamma^\alpha \gamma^\sigma \partial_\alpha - \gamma^\sigma \gamma^\beta \partial_\beta) A_\sigma \end{aligned} \quad (2.6.4)$$

In the momentum space we obtain

$$\begin{aligned}\mathcal{L}_{INT} &= i\mu\bar{\psi}_\nu \left(\gamma^\alpha \gamma^\sigma (ik_\alpha) - \gamma^\sigma \gamma^\beta (ik_\beta) \right) A_\sigma = \mu\bar{\psi}_\nu (\gamma^\sigma \mathcal{K} - \mathcal{K} \gamma^\sigma) A_\sigma \\ &= 2\mu\bar{\psi}_\nu (\gamma^\sigma \mathcal{K} - k^\sigma) A_\sigma\end{aligned}\quad (2.6.5)$$

The process $\nu e^- \rightarrow \nu e^-$ is described by the following diagrams



so that the amplitude is

$$\begin{aligned}\mathcal{A} &= \bar{u}^{s_3}(p_3) \left(-ie\gamma^\mu \right) u^{s_1}(p_1) \left(\frac{-ig_{\mu\nu}}{k^2} \right) \bar{u}^{s_4}(p_4) \left(2i\mu(\gamma^\nu \mathcal{K} - k^\nu) \right) u^{s_2} + \\ &+ iG\bar{u}^{s_4}(p_4)\gamma_\sigma u^{s_2}(p_2)\bar{u}^{s_3}(p_3)\gamma^\sigma u^{s_1}(p_1)\end{aligned}\quad (2.6.7)$$

which means

$$\mathcal{A} = -\frac{2i\mu}{k^2} \bar{u}^{s_3}\gamma_\nu u^{s_1}\bar{u}^{s_4}(\gamma^\nu \mathcal{K} - k^\nu) u^{s_2} + iG\bar{u}^{s_4}\gamma_\sigma u^{s_2}\bar{u}^{s_3}\gamma^\sigma u^{s_1} \quad (2.6.8)$$

$$\mathcal{A}^* = \frac{2i\mu}{k^2} \bar{u}^{s_2}(\mathcal{K}\gamma^\mu - k^\mu) u^{s_4}\bar{u}^{s_1}\gamma_\mu u^{s_3} - iG\bar{u}^{s_1}\gamma^\rho u^{s_3}\bar{u}^{s_2}\gamma_\rho u^{s_4} \quad (2.6.9)$$

The matrix element for an unpolarized cross section is obtained averaging over the spins of the initial particles and summing over the spins of the final ones

$$\begin{aligned}|\mathcal{M}|^2 &= \frac{1}{4} \sum_{s_1, s_2, s_3, s_4} \left\{ \frac{4\mu^2}{k^4} \bar{u}^{s_3}\gamma_\nu u^{s_1}\bar{u}^{s_4}(\gamma^\nu \mathcal{K} - k^\nu) u^{s_2}\bar{u}^{s_2}(\mathcal{K}\gamma^\mu - k^\mu) u^{s_4}\bar{u}^{s_1}\gamma_\mu u^{s_3} - \right. \\ &\quad - \frac{2\mu G}{k^2} \left[\bar{u}^{s_3}\gamma_\nu u^{s_1}\bar{u}^{s_4}(\gamma^\nu \mathcal{K} - k^\nu) u^{s_2}\bar{u}^{s_1}\gamma^\rho u^{s_3}\bar{u}^{s_2}\gamma_\rho u^{s_4} + \right. \\ &\quad \left. \left. + \bar{u}^{s_4}\gamma_\sigma u^{s_2}\bar{u}^{s_3}\gamma^\sigma u^{s_1}\bar{u}^{s_2}(\mathcal{K}\gamma^\mu - k^\mu) u^{s_4}\bar{u}^{s_1}\gamma_\mu u^{s_3} \right] + \right. \\ &\quad \left. + G^2 \bar{u}^{s_4}\gamma_\sigma u^{s_2}\bar{u}^{s_3}\gamma^\sigma u^{s_1}\bar{u}^{s_1}\gamma^\rho u^{s_3}\bar{u}^{s_2}\gamma_\rho u^{s_4} \right\}\end{aligned}\quad (2.6.10)$$

Using the Dirac spinors sum rules we obtain

$$\begin{aligned}
|\mathcal{M}|^2 = & \frac{1}{4} \left\{ \underbrace{\frac{4\mu^2}{k^4} Tr \left[(\not{p}_1 + m_e) \gamma_\mu (\not{p}_3 + m_e) \gamma_\nu \right] Tr \left[\not{p}_2 (\not{k} \gamma^\mu - k^\mu) \not{p}_4 (\gamma^\nu \not{k} - k^\nu) \right]}_{(1)(2)} + \right. \\
& - \underbrace{\frac{2\mu G}{k^2} \left[\underbrace{Tr \left[(\not{p}_1 + m_e) \gamma^\rho (\not{p}_3 + m_e) \gamma_\nu \right]}_{(3)} \underbrace{Tr \left[\not{p}_2 \gamma_\rho \not{p}_4 (\gamma^\nu \not{k} - k^\nu) \right]}_{(4)} \right.} \\
& \left. \left. + \underbrace{Tr \left[(\not{p}_1 + m_e) \gamma_\mu (\not{p}_3 + m_e) \gamma^\sigma \right] Tr \left[\not{p}_2 (\not{k} \gamma^\mu - k^\mu) \not{p}_4 \gamma_\sigma \right]}_{(5)(6)} \right] + \right. \\
& \left. + G^2 \underbrace{Tr \left[(\not{p}_1 + m_e) \gamma^\rho (\not{p}_3 + m_e) \gamma^\sigma \right]}_{(7)} \underbrace{Tr \left[\not{p}_2 \gamma_\rho \not{p}_4 \gamma_\sigma \right]}_{(8)} \right\}
\end{aligned} \tag{2.6.11}$$

Traces (4) and (6) contain an odd number of gamma matrices, so they are equal to zero; therefore the terms proportional to $-\frac{2\mu G}{k^2}$ are null.

$$\begin{aligned}
(1) = & p_{1\alpha} p_{3\gamma} Tr[\gamma^\alpha \gamma_\mu \gamma^\gamma \gamma_\nu] + m_e^2 Tr[\gamma_\mu \gamma_\nu] \\
= & 4p_{1\alpha} p_{3\gamma} (g_\mu^\alpha g_\nu^\gamma - g^{\alpha\gamma} g_{\mu\nu} + g_\nu^\alpha g_\mu^\gamma) + 4m_e^2 g_{\mu\nu} \\
= & 4[p_{1\mu} p_{3\nu} + p_{1\nu} p_{3\mu} + (m_e^2 - p_1 p_3) g_{\mu\nu}]
\end{aligned} \tag{2.6.12}$$

Trace (7) is equal to trace (1) we have just computed, so we can immediately write

$$(7) = 4[p_1^\rho p_3^\sigma + p_1^\sigma p_3^\rho + (m_e^2 - p_1 p_3) g^{\rho\sigma}] \tag{2.6.13}$$

The remaining traces are

$$\begin{aligned}
(8) = & p_{2\beta} p_{4\delta} Tr[\gamma^\beta \gamma_\rho \gamma^\delta \gamma_\sigma] \\
= & 4p_{2\beta} p_{4\delta} (g_\rho^\beta g_\sigma^\delta - g^{\beta\delta} g_{\rho\sigma} + g_\sigma^\beta g_\rho^\delta) \\
= & 4[p_{2\rho} p_{4\sigma} + p_{2\sigma} p_{4\rho} - p_2 p_4 g_{\rho\sigma}]
\end{aligned} \tag{2.6.14}$$

$$\begin{aligned}
(2) &= p_{2\beta} p_{4\delta} k_\sigma k_\rho Tr[\gamma^\beta \gamma^\sigma \gamma^\mu \gamma^\delta \gamma^\nu \gamma^\rho] + p_{2\beta} p_{4\delta} k^\mu k^\nu Tr[\gamma^\beta \gamma^\delta] \\
&= 4p_{2\beta} p_{4\delta} k_\sigma k_\rho \left[g^{\beta\sigma}(g^{\mu\delta}g^{\nu\rho} - g^{\mu\nu}g^{\delta\rho} + g^{\mu\rho}g^{\delta\nu}) - g^{\beta\mu}(g^{\sigma\delta}g^{\nu\rho} - g^{\sigma\nu}g^{\delta\rho} + g^{\sigma\rho}g^{\delta\nu}) + \right. \\
&\quad \left. + g^{\beta\delta}(g^{\sigma\mu}g^{\nu\rho} - g^{\sigma\nu}g^{\mu\rho} + g^{\sigma\rho}g^{\mu\nu}) - g^{\beta\nu}(g^{\sigma\mu}g^{\delta\rho} - g^{\sigma\delta}g^{\mu\rho} + g^{\sigma\rho}g^{\mu\delta}) + \right. \\
&\quad \left. + g^{\beta\rho}(g^{\sigma\mu}g^{\delta\nu} - g^{\sigma\delta}g^{\mu\nu} + g^{\sigma\nu}g^{\mu\delta}) \right] + 4p_{2\beta} p_{4\delta} k^\mu k^\nu g^{\beta\delta} \\
&= 4 \left[(p_2 k) \left[p_4^\mu k^\nu - (p_4 k) g^{\mu\nu} + k^\mu p_4^\nu \right] - p_2^\mu \left[(p_4 k) k^\nu - (p_4 k) k^\nu + k^2 p_4^\nu \right] + \right. \\
&\quad \left. + (p_2 p_4) \left[k^\mu k^\nu - k^\mu k^\nu + k^2 g^{\mu\nu} \right] - p_2^\nu \left[(p_4 k) k^\mu - (p_4 k) k^\mu + k^2 p_4^\mu \right] + \right. \\
&\quad \left. + (p_2 k) \left[k^\mu p_4^\nu - (p_4 k) g^{\mu\nu} + k^\nu p_4^\mu \right] \right] + 4(p_2 p_4) k^\mu k^\nu \\
&= 4 \left[2(p_2 k) \left[p_4^\mu k^\nu + p_4^\nu k^\mu - (p_4 k) g^{\mu\nu} \right] - k^2 \left[p_2^\mu p_4^\nu + p_2^\nu p_4^\mu - (p_2 p_4) g^{\mu\nu} \right] + (p_2 p_4) k^\mu k^\nu \right]
\end{aligned} \tag{2.6.15}$$

Carrying out the products (1) \times (2) and (7) \times (8), the matrix element becomes

$$\begin{aligned}
|\mathcal{M}|^2 &= \frac{16\mu^2}{k^4} \left\{ 4(p_2 k)(p_1 p_4)(p_3 k) + 4(p_2 k)(p_1 k)(p_3 p_4) + 2(p_2 p_4)(p_1 k)(p_3 k) + \right. \\
&\quad - k^2 \left[2(p_1 p_2)(p_3 p_4) + 2(p_1 p_4)(p_2 p_3) + (p_1 p_3)(p_2 p_4) \right] + \\
&\quad \left. + m_e^2 \left[3k^2(p_2 p_4) - 4(p_2 k)(p_4 k) \right] \right\} + \\
&\quad + 8G^2 \left\{ (p_1 p_2)(p_3 p_4) + (p_1 p_4)(p_2 p_3) - m_e^2(p_2 p_4) \right\}
\end{aligned} \tag{2.6.16}$$

In the centre of mass frame we can write the 4-momenta as

$$\begin{cases} p_1 = (E_1, |\mathbf{p}| \hat{z}) \\ p_2 = (E_2, -|\mathbf{p}| \hat{z}) \\ p_3 = (E_3, \mathbf{q}) \\ p_4 = (E_4, -\mathbf{q}) \\ k = p_1 - p_3 \end{cases} \tag{2.6.17}$$

Now $E_4 = E_2$, since

$$\begin{aligned}
 p_3^2 &= m_e^2 = (p_1 + p_2 - p_4)^2 = \underbrace{p_1^2}_{=m_e^2} + \underbrace{p_2^2}_{=0} + \underbrace{p_4^2}_{=0} + 2p_1p_2 - 2p_4 \cdot (p_1 + p_2) = \\
 &= m_e^2 + 2(E_1E_2 + |\mathbf{p}|^2) - 2E_4(E_1 + E_2) \\
 &\stackrel{=E_2^2}{=} \\
 \implies 0 &= (E_2 - E_4)(E_1 + E_2) \implies E_4 = E_2
 \end{aligned} \tag{2.6.18}$$

Because of conservation of energy it also follows that $E_3 = E_1$; moreover, $|\mathbf{p}| = |\mathbf{q}| = E_2$. The scalar products are

$$\begin{aligned}
 p_1p_4 &= E_2(E_1 + E_2 \cos \theta) \\
 p_3p_4 &= E_2(E_1 + E_2) \\
 p_2p_4 &= E_2^2(1 - \cos \theta) \\
 p_1p_2 &= E_2(E_1 + E_2) \\
 p_2p_3 &= E_2(E_1 + E_2 \cos \theta) \\
 p_1p_3 &= E_1^2 - E_2^2 \cos \theta \\
 p_1k &= -E_2^2(1 - \cos \theta) \\
 p_2k &= E_2^2(1 - \cos \theta) \\
 p_3k &= E_2^2(1 - \cos \theta) \\
 p_4k &= -E_2^2(1 - \cos \theta) \\
 k^2 &= 2(E_1^2 - E_2^2)(1 - \cos \theta)
 \end{aligned} \tag{2.6.19}$$

and, by defining $\chi := \frac{E_1}{E_2}$, the matrix element becomes

$$\begin{aligned}
 |\mathcal{M}|^2 &= 8\mu^2 E_2^2 \frac{\chi(2\chi^3 + 4\chi^2 + \chi - 4) + 2(\chi^4 + 2\chi^3 - 2\chi^2 - 2\chi - 2) \cos \theta + 3\chi^2 \cos^2 \theta}{(1 - \chi^2)(1 - \cos \theta)} + \\
 &+ 8G^2 E_2^4 \left[1 + (1 + \chi)^2 + [(1 + \chi)^2 - 2] \cos \theta + \cos^2 \theta \right]
 \end{aligned} \tag{2.6.20}$$

The flux factor is

$$\mathcal{J} = \frac{1}{4E_2^2(\chi + 1)} \tag{2.6.21}$$

while the 2-body phase space element is

$$d\Phi_2 = \frac{1}{\chi + 1} \frac{d \cos \theta d\varphi}{16\pi^2} \tag{2.6.22}$$

Given that, once having integrated over $d\varphi$, the differential cross section for the process becomes

$$\frac{d\sigma}{d\cos\theta} = \frac{1}{32\pi E_2^2(\chi+1)^2} |\mathcal{M}|^2 \quad (2.6.23)$$

where $|\mathcal{M}|^2$ is given by (2.6.20).

2.7 QED with massless electron: $\gamma e^- \rightarrow \gamma e^-$

Text. Consider the standard QED Lagrangian and assume the electron to be massless. Compute the differential cross section for the process

$$\gamma e^- \rightarrow \gamma e^-$$

Solution. Let's consider the initial particles 4-momenta p_j 's and the final ones q_j 's in the centre of mass reference frame.

$$p_1 = \begin{pmatrix} E_e \\ \mathbf{p} \end{pmatrix} \quad (2.7.1)$$

$$p_2 = \begin{pmatrix} E_\gamma \\ -\mathbf{p} \end{pmatrix} \quad (2.7.2)$$

$$q_1 = \begin{pmatrix} E_{e'} \\ -\mathbf{q} \end{pmatrix} \quad (2.7.3)$$

$$q_2 = \begin{pmatrix} E_{\gamma'} \\ \mathbf{q} \end{pmatrix} \quad (2.7.4)$$

The assumption of a massless electron results in the square of every 4-momentum to be null: $p_1^2 = 0$, $p_2^2 = 0$, $q_1^2 = 0$, $q_2^2 = 0$. Moreover, because of the mass-shell relation, $E_e = E_\gamma = E$, $E_{e'} = E_{\gamma'} = E'$, so that

$$p_1 = \begin{pmatrix} E \\ \mathbf{p} \end{pmatrix} \quad (2.7.5)$$



$$p_2 = \begin{pmatrix} E \\ -\mathbf{p} \end{pmatrix} \quad (2.7.6)$$

$$q_1 = \begin{pmatrix} E' \\ -\mathbf{q} \end{pmatrix} \quad (2.7.7)$$

$$q_2 = \begin{pmatrix} E' \\ \mathbf{q} \end{pmatrix} \quad (2.7.8)$$

The Feynman rules are the usual ones for QED. We proceed building the relevant

diagrams for the process



Thanks to 4-momentum conservation we have

$$p_A = p_1 - q_1 = q_2 - p_2 \quad (2.7.11)$$

$$p_B = p_1 + p_2 = q_1 + q_2 \quad (2.7.12)$$

Calling \mathcal{A} the amplitude of the first diagram and \mathcal{B} the amplitude of the second one, we have

$$\mathcal{A} = ie^2 \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_A}{p_A^2} \gamma^\mu u^{s_1} \varepsilon_\nu^{\lambda_2} \left(\varepsilon_\mu^{\lambda_1} \right)^* \quad (2.7.13)$$

$$\mathcal{B} = ie^2 \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_B}{p_B^2} \gamma^\mu u^{s_1} \varepsilon_\mu^{\lambda_2} \left(\varepsilon_\nu^{\lambda_1} \right) \quad (2.7.14)$$

The momenta p_A^2 and p_B^2 are

$$p_A^2 = p_1^2 + q_1^2 - 2p_1 q_1 = -2p_1 q_1 \quad (2.7.15)$$

$$p_B^2 = p_1^2 + p_2^2 + 2p_1 p_2 = 2p_1 p_2 \quad (2.7.16)$$

where $p_1^2 = p_2^2 = q_1^2 = 0$ since we're dealing with massless particles. Given that the amplitudes become

$$\mathcal{A} = -ie^2 \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_1 - \not{q}_1}{2p_1 q_1} \gamma^\mu u^{s_1} \varepsilon_\nu^{\lambda_2} \left(\varepsilon_\mu^{\lambda_1} \right) \quad (2.7.17)$$

$$\mathcal{B} = ie^2 \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_1 + \not{p}_2}{2p_1 p_2} \gamma^\mu u^{s_1} \varepsilon_\mu^{\lambda_2} \left(\varepsilon_\nu^{\lambda_1} \right) \quad (2.7.18)$$

For the total scattering amplitude we have

$$|\mathcal{M}|^2 = \frac{1}{4} \sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A} + \mathcal{B}|^2 = \frac{1}{4} \sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2 + |\mathcal{B}|^2 + \mathcal{A}^* \mathcal{B} + \mathcal{A} \mathcal{B}^* \quad (2.7.19)$$

where the factor $\frac{1}{4}$ comes from averaging over the initial spins.

Let's start by computing the term $\sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2$

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2 = \frac{e^4}{4(p_1 q_1)^2} g_{\nu\beta} g_{\mu\alpha} \sum_{s_1, s_2} \bar{u}^{s_2} \gamma^\nu (\not{p}_1 - \not{q}_1) \gamma^\mu u^{s_1} \bar{u}^{s_1} \gamma^\alpha (\not{p}_1 - \not{q}_1) \gamma^\beta u^{s_2} \quad (2.7.20)$$

where we have already summed over the polarizations. Contracting the metric tensors we obtain the following trace

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2 = \frac{e^4}{4(p_1 q_1)^2} Tr \left[\not{q}_2 \gamma^\nu (\not{p}_1 - \not{q}_1) \gamma^\mu \not{p}_1 \gamma_\mu (\not{p}_1 - \not{q}_1) \gamma_\nu \right] \quad (2.7.21)$$

Thanks to equation (A.20) we obtain the sum of four traces, three of which are proportional to terms of the type $\not{p}_1 \not{p}_1 = p_1^2 = 0$; we are then left with

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2 = \frac{-2e^4}{4(p_1 q_1)^2} Tr \left[\not{q}_2 \gamma^\nu \not{q}_1 \not{p}_1 \not{q}_1 \gamma_\nu \right] \quad (2.7.22)$$

Thanks to identity (A.22) we get

$$\begin{aligned} \sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2 &= \frac{4e^4}{4(p_1 q_1)^2} Tr \left[\not{q}_2 \not{q}_1 \not{p}_1 \not{q}_1 \right] = \\ &= \frac{4e^4}{(p_1 q_1)^2} q_{2\mu} q_{1\nu} p_{1\alpha} q_{1\beta} (g^{\mu\nu} g^{\alpha\beta} - g^{\mu\alpha} g^{\nu\beta} + g^{\mu\beta} g^{\nu\alpha}) \end{aligned} \quad (2.7.23)$$

Therefore

$$\begin{aligned} \sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{A}|^2 &= \frac{4e^4}{(p_1 q_1)^2} \left((q_1 q_2)(p_1 q_1) - (q_2 p_1) q_1^2 + (q_2 q_1)(q_1 p_1) \right) = \\ &= \frac{8e^4}{(p_1 q_1)^2} (q_1 q_2)(p_1 q_1) = \frac{8e^4}{(p_1 q_1)} (q_1 q_2) \end{aligned} \quad (2.7.24)$$

Note that the summation for $|\mathcal{B}|^2$ can be obtained from the above by simply exchanging $-q_1$ with $+p_2$

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} |\mathcal{B}|^2 = \frac{8e^4}{(p_1 p_2)} (p_2 q_2) \quad (2.7.25)$$

All that is left to do is to compute the terms $\mathcal{A}^*\mathcal{B}$ e $\mathcal{A}\mathcal{B}^*$

$$\mathcal{A}\mathcal{B}^* = -e^4 \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_1 - \not{q}_1}{2p_1 q_1} \gamma^\mu u^{s_1} \varepsilon_\nu^{\lambda_2} \left(\varepsilon_\mu^{\lambda_1} \right)^* \bar{u}^{s_1} \gamma^\alpha \frac{\not{p}_1 + \not{p}_2}{2p_1 p_2} \gamma^\beta u^{s_2} \left(\varepsilon_\alpha^{\lambda_2} \right)^* \varepsilon_\beta^{\lambda_1} \quad (2.7.26)$$

$$\mathcal{A}^*\mathcal{B} = -e^4 \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_1 + \not{p}_2}{2p_1 p_2} \gamma^\mu u^{s_1} \varepsilon_\mu^{\lambda_2} \left(\varepsilon_\nu^{\lambda_1} \right)^* \bar{u}^{s_1} \gamma^\alpha \frac{\not{p}_1 - \not{q}_1}{2p_1 q_1} \gamma^\beta u^{s_2} \left(\varepsilon_\beta^{\lambda_1} \right)^* \varepsilon_\alpha^{\lambda_2} \quad (2.7.27)$$

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A}\mathcal{B}^* = -e^4 \sum_{s_1, s_2, \lambda_1, \lambda_2} \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_1 - \not{q}_1}{2p_1 q_1} \gamma^\mu u^{s_1} \varepsilon_\nu^{\lambda_2} \left(\varepsilon_\mu^{\lambda_1} \right)^* \bar{u}^{s_1} \gamma^\alpha \frac{\not{p}_1 + \not{p}_2}{2p_1 p_2} \gamma^\beta u^{s_2} \left(\varepsilon_\alpha^{\lambda_2} \right)^* \varepsilon_\beta^{\lambda_1} \quad (2.7.28)$$

Making explicit the summations on the polarization indices we get two metric tensors

$$\begin{aligned} \sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A}\mathcal{B}^* &= -e^4 \sum_{s_1, s_2} \bar{u}^{s_2} \gamma^\nu \frac{\not{p}_1 - \not{q}_1}{2p_1 q_1} \gamma^\mu u^{s_1} \bar{u}^{s_1} \gamma^\alpha \frac{\not{p}_1 + \not{p}_2}{2p_1 p_2} \gamma^\beta u^{s_2} g_{\alpha\nu} g_{\beta\mu} \\ &= -\frac{e^4}{4(p_1 q_1)(p_1 p_2)} \sum_{s_1, s_2} \bar{u}^{s_2} \gamma^\nu (\not{p}_1 - \not{q}_1) \gamma^\mu u^{s_1} \bar{u}^{s_1} \gamma_\nu (\not{p}_1 + \not{p}_2) \gamma_\mu u^{s_2} \end{aligned} \quad (2.7.29)$$

and summing over the spins we arrive to the following trace

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A}\mathcal{B}^* = \frac{-e^4}{4(p_1 q_1)(p_1 p_2)} Tr \left[\not{q}_2 \gamma^\nu (\not{p}_1 - \not{q}_1) \gamma^\mu \not{p}_1 \gamma_\nu (\not{p}_1 + \not{p}_2) \gamma_\mu \right] \quad (2.7.30)$$

Using again (A.22) we obtain

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A}\mathcal{B}^* = \frac{2e^4}{4(p_1 q_1)(p_1 p_2)} Tr \left[\not{q}_2 \not{p}_1 \gamma^\mu (\not{p}_1 - \not{q}_1) (\not{p}_1 + \not{p}_2) \gamma_\mu \right] \quad (2.7.31)$$

Applying now (A.21) we have

$$\begin{aligned} \sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A}\mathcal{B}^* &= \frac{2e^4}{4(p_1 q_1)(p_1 p_2)} Tr \left[\not{q}_2 \not{p}_1 \cdot 4(p_1 - q_1)(p_1 + p_2) \right] = \\ &= \frac{8e^4}{(p_1 q_1)(p_1 p_2)} (q_2 p_1)(p_1 - q_1)(p_1 + p_2) = \\ &= \frac{8e^4}{(p_1 q_1)(p_1 p_2)} (p_1 q_2) \left[(p_1 p_2) - (q_1 p_1) - (q_1 p_2) \right] \end{aligned} \quad (2.7.32)$$

In order to compute $\sum \mathcal{A}^* \mathcal{B}$ it is sufficient to make the following substitutions: $-q_1 \rightarrow p_2$, $p_2 \rightarrow -q_1$. We then obtain

$$\sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A}^* \mathcal{B} = \sum_{s_1, s_2, \lambda_1, \lambda_2} \mathcal{A} \mathcal{B}^* = \frac{8e^4}{(p_1 q_1)(p_1 p_2)} (p_1 q_2) \left[(p_1 p_2) - (q_1 p_1) - (q_1 p_2) \right] \quad (2.7.33)$$

Putting all the pieces together, and remembering the factor $\frac{1}{4}$ in front of the summation, we have

$$|\mathcal{M}|^2 = \frac{1}{4} \left\{ \frac{8e^4}{(p_1 q_1)} (q_1 q_2) + \frac{8e^4}{(p_1 p_2)} (p_2 q_2) + \frac{16e^4}{(p_1 q_1)(p_1 p_2)} (p_1 q_2) [(p_1 p_2) - (q_1 p_1) - (q_1 p_2)] \right\} \quad (2.7.34)$$

and finally

$$|\mathcal{M}|^2 = \frac{2e^4}{(p_1 q_1)(p_1 p_2)} \left\{ (q_1 q_2)(p_1 p_2) + (p_2 q_2)(p_1 q_1) + 2(p_1 q_2)[(p_1 p_2) - (q_1 p_1) - (q_1 p_2)] \right\} \quad (2.7.35)$$

As anticipated we put ourselves in the centre of mass frame; imposing 4-momentum conservation, the moduli of the initial and final 3-momenta must be equal to each other, and so thanks to the mass shell relations we have $E' = E$. Our calculations simplify considerably, so we can write

$$|\mathcal{M}|^2 = \frac{e^4}{1 + \cos \theta} [4 + (1 + \cos \theta)^2] \quad (2.7.36)$$

The expression for the flux factor \mathcal{J} is

$$\mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2}} = \frac{1}{8E^2} \quad (2.7.37)$$

and the 2-body phase space element is

$$\begin{aligned} d\Phi_2 &= \frac{|\vec{q}_1|}{2E} \frac{d\cos\theta d\phi}{16\pi^2} = \frac{1}{2} \frac{d\cos\theta d\varphi}{16\pi^2} = \\ &= \frac{1}{2} \frac{d\cos\theta}{8\pi} \end{aligned} \quad (2.7.38)$$

Therefore, the differential cross section of the process becomes

$$\frac{d\sigma}{d\cos\theta} = \frac{1}{128} \frac{e^4}{1 + \cos\theta} [4 + (1 + \cos\theta)^2] \quad (2.7.39)$$

2.8 Chiral lagrangian: $\psi_a \psi_b \rightarrow \psi_a \psi_b$

Text. Consider the following lagrangian density

$$\mathcal{L} = \bar{\psi}_a (\partial_\mu \gamma^\mu - m_a) \psi_a + \bar{\psi}_b \partial_\mu \gamma^\mu \psi_b + \Lambda \bar{\psi}_a \gamma_\mu (g_V \mathbb{1} + g_A \gamma_5) \psi_a \bar{\psi}_b \gamma^\mu (\mathbb{1} - \gamma_5) \psi_b$$

where ψ_a and ψ_b are associated to two different fermionic species. Derive the Feynman rules of the theory and compute the cross section for the process

$$\psi_a \psi_b \rightarrow \psi_a \psi_b$$

Solution. The Lagrangian in question is made up by two Dirac Lagrangians, one for the massive fermions a and the other for the massless fermions b , plus the following interaction term

$$\mathcal{L}_{INT} = \Lambda \bar{\psi}_a \gamma^\mu (g_V \mathbb{1} + g_A \gamma^5) \psi_a \bar{\psi}_b \gamma_\mu (\mathbb{1} - \gamma^5) \psi_b \quad (2.8.1)$$

The interaction Lagrangian represents a contact term, since there's no propagator there. To the usual QED Feynman rules we add the following vertex factor

$$= i\Lambda \gamma^\mu (g_V \mathbb{1} + g_A \gamma^5) \gamma_\mu (\mathbb{1} - \gamma^5) \quad (2.8.2)$$

where, as usual, we added the factor i . At first order the process is uniquely described by the following diagram

$$(2.8.3)$$

The scattering amplitude is

$$\mathcal{A} = i\Lambda \bar{u}^{s_3}(p_3) \gamma^\mu (g_V \mathbb{1} + g_A \gamma^5) u^{s_1}(p_1) \bar{u}^{s_4}(p_4) \gamma_\mu (\mathbb{1} - \gamma^5) u^{s_2}(p_2) \quad (2.8.4)$$

with complex conjugate equal to

$$\mathcal{A}^* = -i\Lambda \bar{u}^{s_2}(p_2)(\mathbb{1} + \gamma^5)\gamma_\nu u^{s_4}(p_4) \bar{u}^{s_1}(p_1)(g_V \mathbb{1} - g_A \gamma^5)\gamma^\nu u^{s_3}(p_3) \quad (2.8.5)$$

As usual, in computing unpolarized cross sections, we have to average on the initial spins and sum over the final ones

$$|\mathcal{M}|^2 = \frac{\Lambda^2}{4} \sum_{s_1, s_2, s_3, s_4} \bar{u}^{s_3}\gamma^\mu(g_V \mathbb{1} + g_A \gamma^5)u^{s_1}\bar{u}^{s_4}\gamma_\mu(\mathbb{1} - \gamma^5)u^{s_2} \\ \bar{u}^{s_2}(\mathbb{1} + \gamma^5)\gamma_\nu u^{s_4}\bar{u}^{s_1}(g_V \mathbb{1} - g_A \gamma^5)\gamma^\nu u^{s_3} \quad (2.8.6)$$

Using the Dirac spinors sum rules

$$|\mathcal{M}|^2 = \frac{\Lambda^2}{4} \left[Tr \left[(\not{p}_1 + m_a)(g_V - g_A \gamma^5)\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu(g_V + g_A \gamma^5) \right] \right. \\ \left. Tr \left[\not{p}_2(1 + \gamma_5)\gamma_\nu \not{p}_4 \gamma_\mu(1 - \gamma_5) \right] \right] \quad (2.8.7)$$

An explicit calculation of all the above traces is excessively long and wasteful; our aim is to manipulate the former expression exploiting some properties of the Dirac matrices.

Let us consider the first trace

$$Tr \left[(\not{p}_1 + m_a)(g_V - g_A \gamma^5)\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu(g_V + g_A \gamma^5) \right] =: T_1 \quad (2.8.8)$$

Using the cyclic property of the trace (A.29), together with (A.19), we can write T_1 as follows

$$T_1 = Tr \left[(g_V + g_A \gamma^5)(\not{p}_1 + m_a)(g_V - g_A \gamma^5)\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu \right] = \\ = Tr \left[\left(m_a(g_V + g_A \gamma^5) + \not{p}_1(g_V - g_A \gamma^5) \right) (g_V - g_A \gamma^5)\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu \right] = \\ = m_a Tr \left[(g_V + g_A \gamma^5)(g_V - g_A \gamma^5)\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu \right] + \\ + Tr \left[\not{p}_1(g_V - g_A \gamma^5)^2\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu \right] = \\ = (g_V^2 - g_A^2)m_a Tr \left[\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu \right] + \\ + Tr \left[\not{p}_1(g_V^2 + g_A^2 - 2g_V g_A \gamma^5)\gamma^\nu(\not{p}_3 + m_a)\gamma^\mu \right] \quad (2.8.9)$$

Defining the following constants

$$\begin{cases} G^+ := g_V^2 + g_A^2 \\ G^- := g_V^2 - g_A^2 \\ G_{VA} := g_V g_A \end{cases} \quad (2.8.10)$$

we obtain a more compact expression

$$\begin{aligned} T_1 &= G^- m_a \text{Tr} \left[\gamma^\nu (\not{p}_3 + m_a) \gamma^\mu \right] + \text{Tr} \left[\not{p}_1 (G^+ - 2G_{VA}\gamma^5) \gamma^\nu (\not{p}_3 + m_a) \gamma^\mu \right] = \\ &= G^- m_a^2 \text{Tr} \left[\gamma^\mu \gamma^\nu \right] + G^+ \text{Tr} \left[\not{p}_1 \gamma^\nu (\not{p}_3 + m_a) \gamma^\mu \right] - \\ &\quad - 2G_{VA} \text{Tr} \left[\not{p}_1 \gamma^5 \gamma^\nu (\not{p}_3 + m_a) \gamma^\mu \right] = \\ &= 4G^- m_a^2 g^{\mu\nu} + G^+ p_{1\alpha} p_{3\gamma} \text{Tr} \left[\gamma^\alpha \gamma^\nu \gamma^\gamma \gamma^\mu \right] - 2G_{VA} p_{1\alpha} p_{3\gamma} \text{Tr} \left[\gamma^\alpha \gamma^5 \gamma^\nu \gamma^\gamma \gamma^\mu \right] \end{aligned} \quad (2.8.11)$$

The trace of an odd number of gamma matrices is null, as well as a trace containing γ^5 multiplied by an odd number of gamma matrices, so that

$$T_1 = 4G^- m_a^2 g^{\mu\nu} + 4G^+ p_{1\alpha} p_{3\gamma} (g^{\alpha\nu} g^{\gamma\mu} - g^{\alpha\gamma} g^{\nu\mu} + g^{\alpha\mu} g^{\nu\gamma}) - 2G_{VA} p_{1\alpha} p_{3\gamma} \text{Tr} \left[\gamma^5 \gamma^\nu \gamma^\gamma \gamma^\mu \gamma^\alpha \right] \quad (2.8.12)$$

and using $\text{Tr} \left[\gamma^5 \gamma^\nu \gamma^\gamma \gamma^\mu \gamma^\alpha \right] = -4i\varepsilon^{\nu\gamma\mu\alpha}$ we obtain

$$\begin{aligned} T_1 &= 4G^- m_a^2 g^{\mu\nu} + 4G^+ \left(p_1^\nu p_3^\mu - (p_1 p_3) g^{\mu\nu} + p_1^\mu p_3^\nu \right) - 8iG_{VA} p_{1\alpha} p_{3\gamma} \varepsilon^{\nu\gamma\mu\alpha} = \\ &= 4 \left[\left(G^- m_a^2 - G^+ (p_1 p_3) \right) g^{\mu\nu} + G^+ \left(p_1^\nu p_3^\mu + p_1^\mu p_3^\nu \right) \right] - 8iG_{VA} p_{1\alpha} p_{3\gamma} \varepsilon^{\nu\gamma\mu\alpha} \end{aligned} \quad (2.8.13)$$

In the expression above we can notice that the first term in square brackets is symmetric in the indices μ and ν , while the second term is antisymmetric in the same indices (due to the presence of the totally antisymmetric Levi-Civita symbol). This allows us to define the following tensors

$$\begin{cases} S^{\mu\nu} := \left[\left(G^- m_a^2 - G^+ (p_1 p_3) \right) g^{\mu\nu} + G^+ \left(p_1^\nu p_3^\mu + p_1^\mu p_3^\nu \right) \right] \\ A^{\mu\nu} := G_{VA} p_{1\alpha} p_{3\gamma} \varepsilon^{\nu\gamma\mu\alpha} \end{cases} \quad (2.8.14)$$

The trace T_1 now takes the form

$$T_1 = 4S^{\mu\nu} - 8iA^{\mu\nu} \quad (2.8.15)$$

We can handle the second trace with similar manipulations

$$\begin{aligned} T_2 &:= Tr \left[\not{p}_2 (1 + \gamma^5) \gamma_\nu \not{p}_4 \gamma_\mu (1 - \gamma^5) \right] = Tr \left[(1 - \gamma^5) \not{p}_2 (1 + \gamma^5) \gamma_\nu \not{p}_4 \gamma_\mu \right] = \\ &= Tr \left[\not{p}_2 (1 + \gamma^5) (1 + \gamma^5) \gamma_\nu \not{p}_4 \gamma_\mu \right] = 2Tr \left[\not{p}_2 \gamma_\nu \not{p}_4 \gamma_\mu \right] + 2Tr \left[\not{p}_2 \gamma^5 \gamma_\nu \not{p}_4 \gamma_\mu \right] = \\ &= 2p_2^\beta p_4^\delta Tr \left[\gamma_\beta \gamma_\nu \gamma_\delta \gamma_\mu \right] + 2p_2^\beta p_4^\delta Tr \left[\gamma^5 \gamma_\nu \gamma_\delta \gamma_\mu \gamma_\beta \right] = \\ &= 8p_2^\beta p_4^\delta \left(g_{\beta\nu} g_{\delta\mu} - g_{\beta\delta} g_{\nu\mu} + g_{\beta\mu} g_{\nu\delta} \right) - 8ip_2^\beta p_4^\delta \varepsilon_{\nu\delta\mu\beta} = \\ &= 8 \left[p_{2\nu} p_{4\mu} - (p_2 p_4) g_{\mu\nu} + p_{2\mu} p_{4\nu} \right] - 8ip_2^\beta p_4^\delta \varepsilon_{\nu\delta\mu\beta} \end{aligned} \quad (2.8.16)$$

By defining

$$\begin{cases} S'_{\mu\nu} := p_{2\nu} p_{4\mu} - (p_2 p_4) g_{\mu\nu} + p_{2\mu} p_{4\nu} \\ A'_{\mu\nu} := p_2^\beta p_4^\delta \varepsilon_{\nu\delta\mu\beta} \end{cases} \quad (2.8.17)$$

we obtain

$$T_2 = Tr \left[\not{p}_2 (1 + \gamma^5) \gamma_\nu \not{p}_4 \gamma_\mu (1 - \gamma^5) \right] = 8S'_{\mu\nu} - 8iA'_{\mu\nu} \quad (2.8.18)$$

Therefore we can make explicit the matrix element $|\mathcal{M}|^2$

$$\begin{aligned} |\mathcal{M}|^2 &= \frac{\Lambda^2}{4} \left[4S^{\mu\nu} - 8iA^{\mu\nu} \right] \left[8S'_{\mu\nu} - 8iA'_{\mu\nu} \right] = \\ &= \frac{\Lambda^2}{4} \left[32S^{\mu\nu}S'_{\mu\nu} - 32iS^{\mu\nu}A'_{\mu\nu} - 64iA^{\mu\nu}S'_{\mu\nu} - 64A^{\mu\nu}A'_{\mu\nu} \right] \end{aligned} \quad (2.8.19)$$

The second and third term are both null, since they are products of a symmetric tensor which multiplies an antisymmetric one in the same indices. It follows that

$$|\mathcal{M}|^2 = 8\Lambda^2 \left[S^{\mu\nu}S'_{\mu\nu} - 2A^{\mu\nu}A'_{\mu\nu} \right] \quad (2.8.20)$$

After some simple algebraic passages, the first term can be rewritten as

$$\begin{aligned} S^{\mu\nu}S'_{\mu\nu} &= -2 \left[G^- m_a^2 - G^+ (p_1 p_3) \right] (p_2 p_4) + \\ &\quad + 2G^+ \left[(p_1 p_2)(p_3 p_4) - (p_1 p_3)(p_2 p_4) + (p_1 p_4)(p_2 p_3) \right] \end{aligned} \quad (2.8.21)$$

For the second term we have

$$\begin{aligned} -2A^{\mu\nu} A'_{\mu\nu} &= -2 \left[G_{VA} p_{1\alpha} p_{3\gamma} \varepsilon^{\nu\gamma\mu\alpha} \right] \left[p_2^\beta p_4^\delta \varepsilon_{\nu\delta\mu\beta} \right] = \\ &= -2G_{VA} p_{1\alpha} p_2^\beta p_{3\gamma} p_4^\delta \varepsilon^{\nu\gamma\mu\alpha} \varepsilon_{\nu\delta\mu\beta} \end{aligned} \quad (2.8.22)$$

The product of two Levi-Civita symbols can be simplified in the following way

$$\begin{cases} \varepsilon^{\nu\gamma\mu\alpha} = -\varepsilon^{\nu\mu\gamma\alpha} \\ \varepsilon_{\nu\delta\mu\beta} = -\varepsilon_{\nu\mu\delta\beta} \end{cases} \Rightarrow \varepsilon^{\nu\gamma\mu\alpha} \varepsilon_{\nu\delta\mu\beta} = \varepsilon^{\nu\mu\gamma\alpha} \varepsilon_{\nu\mu\delta\beta} = -2(\delta_\delta^\gamma \delta_\beta^\alpha - \delta_\beta^\gamma \delta_\delta^\alpha) \quad (2.8.23)$$

so that

$$\begin{aligned} -2A^{\mu\nu} A'_{\mu\nu} &= 4G_{VA} p_{1\alpha} p_2^\beta p_{3\gamma} p_4^\delta (\delta_\delta^\gamma \delta_\beta^\alpha - \delta_\beta^\gamma \delta_\delta^\alpha) = \\ &= 4G_{VA} \left[(p_1 p_2)(p_3 p_4) - (p_1 p_4)(p_2 p_3) \right] \end{aligned} \quad (2.8.24)$$

Putting the previous results together we finally obtain

$$|\mathcal{M}|^2 = 16\Lambda^2 \left[-G^- m_a^2 (p_2 p_4) + (G^+ + 2G_{VA})(p_1 p_2)(p_3 p_4) + (G^+ - 2G_{VA})(p_1 p_4)(p_2 p_3) \right] \quad (2.8.25)$$

In the centre of mass frame the 4-momenta are

$$\begin{cases} p_1 = (E_1, |\mathbf{p}| \hat{z}) \\ p_2 = (E_2, -|\mathbf{p}| \hat{z}) \\ p_3 = (E_3, \mathbf{q}) \\ p_4 = (E_4, -\mathbf{q}) \end{cases} \quad (2.8.26)$$

Notice that $E_4 = E_2$, since

$$\begin{aligned} p_3^2 = m_a^2 &= (p_1 + p_2 - p_4)^2 = \underbrace{p_1^2}_{=m_a^2} + \underbrace{p_2^2}_{=0} + \underbrace{p_4^2}_{=0} + 2p_1 p_2 - 2p_4 \cdot (p_1 + p_2) = \\ &= m_a^2 + 2(E_1 E_2 + \underbrace{|\mathbf{p}|^2}_{=E_2^2}) - 2E_4(E_1 + E_2) \implies \\ &\implies 0 = (E_2 - E_4)(E_1 + E_2) \implies E_4 = E_2 \end{aligned} \quad (2.8.27)$$

Thanks to energy conservation we also have $E_3 = E_1$. The various scalar products are

$$\begin{aligned} p_2 p_4 &= E_2^2 (1 - \cos \theta) \\ p_1 p_2 &= p_3 p_4 = E_2 (E_1 + E_2) \\ p_1 p_4 &= p_2 p_3 = E_2 (E_1 + E_2 \cos \theta) \end{aligned} \quad (2.8.28)$$

The scattering amplitude then results

$$|\mathcal{M}|^2 = 16\Lambda^2 E_2^2 \left[-G^- m_a^2 (1 - \cos \theta) + (G^+ + 2G_{VA})(E_1 + E_2)^2 + (G^+ - 2G_{VA})(E_1 + E_2 \cos \theta)^2 \right] \quad (2.8.29)$$

Going back to the constants previously defined, we have

$$|\mathcal{M}|^2 = 16\Lambda^2 E_2^2 \left[(g_A^2 - g_V^2)m_a^2 (1 - \cos \theta) + (g_V + g_A)^2 (E_1 + E_2)^2 + (g_V - g_A)^2 (E_1 + E_2 \cos \theta)^2 \right] \quad (2.8.30)$$

The flux factor is

$$\mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2}} = \frac{1}{4E_2(E_1 + E_2)} \quad (2.8.31)$$

while the 2-body phase space is

$$d\Phi_2 = \frac{|\mathbf{q}|}{E_1 + E_2} \frac{d \cos \theta d\varphi}{16\pi^2} = \frac{E_2}{E_1 + E_2} \frac{d \cos \theta d\varphi}{16\pi^2} \quad (2.8.32)$$

once having integrated over $d\varphi$, the differential cross section for the process finally becomes

$$\frac{d\sigma}{d \cos \theta} = \frac{\Lambda^2}{2\pi} \frac{E_2^2}{(E_1 + E_2)^2} \left[(g_A^2 - g_V^2)m_a^2 (1 - \cos \theta) + (g_V + g_A)^2 (E_1 + E_2)^2 + (g_V - g_A)^2 (E_1 + E_2 \cos \theta)^2 \right] \quad (2.8.33)$$

Chapter 3

Ward Identities

3.1 Radiative corrections for $\phi\phi^* \rightarrow \phi\phi^*\gamma$

Text. Given the following Lagrangian



$$\mathcal{L} = (D_\mu\phi)^*(D^\mu\phi) - m^2|\phi|^2 + \bar{\psi}(i\not{D} - m_e)\psi - \frac{1}{4}F_{\alpha\beta}F^{\alpha\beta}$$

Show that the amplitude for the process

$$\phi\phi^* \rightarrow \phi\phi^*\gamma$$





vanishes after replacing $\varepsilon(k) \rightarrow k$.

Solution. Using the vertices derived in exercise 2.2 we can build the following diagrams



Notice that the amplitudes are proportional to e^3 , where e is the coupling constant of the interaction.

In order to verify the Ward identity we need to include all possible Feynman diagrams proportional to e^3 . By attaching a 3-lines vertex to external lines, which means taking into account the *radiative corrections*, we can build the following diagrams



The total scattering amplitude is the sum of these six amplitudes; the Ward identity predicts that once we make the substitution $\varepsilon_\sigma^*(q) \rightarrow q_\sigma$, the total amplitude vanishes.

Explicitely the amplitudes are

$$\mathcal{A}_1 = -ie(p_1 - p_2)^\mu \frac{-ig_{\mu\nu}}{k^2} 2ie^2 g^{\nu\sigma} \varepsilon_\sigma^*(q) \quad (3.1.7)$$

$$\mathcal{A}_2 = 2ie^2 \varepsilon_\sigma^*(q) g^{\sigma\mu} \frac{-ig_{\mu\nu}}{k^2} (-ie)(p_3 - p_4)^\nu \quad (3.1.8)$$

$$\mathcal{A}_3 = -ie(p_1 - p_2)^\mu \frac{-ig_{\mu\nu}}{k^2} (-ie)(p' - p_4)^\nu \frac{i}{p'^2 - m^2} (-ie)(p' + p_3)^\sigma \varepsilon_\sigma^*(q) \quad (3.1.9)$$

$$\mathcal{A}_4 = -ie(p_1 - p_2)^\mu \frac{-ig_{\mu\nu}}{k^2} (-ie)(p_3 - p')^\nu \frac{i}{p'^2 - m^2} (-ie)(-p' - p_4)^\sigma \varepsilon_\sigma^*(q) \quad (3.1.10)$$

$$\mathcal{A}_5 = -ie(p_1 + p')^\sigma \varepsilon_\sigma^*(q) \frac{i}{p'^2 - m^2} (-ie)(p' - p_2)^\mu \frac{-ig_{\mu\nu}}{k^2} (-ie)(p_3 - p_4)^\nu \quad (3.1.11)$$

$$\mathcal{A}_6 = -ie(-p_2 - p')^\sigma \varepsilon_\sigma^*(q) (-ie)(-p' + p_1)^\mu \frac{-ig_{\mu\nu}}{k^2} (-ie)(p_3 - p_4)^\nu \quad (3.1.12)$$

Firstly, we want to simplify each amplitude. Starting from \mathcal{A}_1 we have

$$\mathcal{A}_1 = -ie(p_1 - p_2)^\mu \frac{-ig_{\mu\nu}}{k^2} 2ie^2 g^{\nu\sigma} q_\sigma = -\frac{2ie^3(p_1 - p_2)^\mu q_\mu}{k^2} \quad (3.1.13)$$

Notice that

$$\begin{aligned} k = p_1 + p_2 \implies k^2 &= (p_1 + p_2)^2 = p_1^2 + p_2^2 + 2p_1 p_2 \\ &= m^2 + m^2 + 2p_1 p_2 \\ &= 2(m^2 + p_1 p_2) \end{aligned} \quad (3.1.14)$$

therefore

$$\mathcal{A}_1 = -\frac{ie^3(p_1 - p_2) \cdot q}{m^2 + p_1 p_2} \quad (3.1.15)$$

Elaborating in a similar fashion \mathcal{A}_2 , by using $k = p_3 + p_4$, we get

$$\mathcal{A}_2 = -\frac{ie^3(p_3 - p_4) \cdot q}{m^2 + p_3 p_4} \quad (3.1.16)$$

Exploiting once more the conservation of 4-momentum at the vertices, the amplitude \mathcal{A}_3 can be written as follows

$$\mathcal{A}_3 = -ie(p_1 - p_2)^\mu \frac{-ig_{\mu\nu}}{k^2} (-ie)(p' - p_4)^\nu \frac{i}{p'^2 - m^2} (-ie)(p' + p_3)^\sigma q_\sigma \quad (3.1.17)$$

By using $p' = q + p_3$ e $p' = p_1 + p_2 - p_4$, the products in the amplitude simplify in the following way

$$\begin{aligned} (p_1 - p_2)^\mu g_{\mu\nu} (p' - p_4)^\nu &= (p_1 - p_2) \cdot (p' - p_4) \\ &= (p_1 - p_2) \cdot (p_1 + p_2 - 2p_4) \\ &= p_1^2 - p_2^2 + 2p_1 p_2 - 2p_1 p_2 - 2p_4 \cdot (p_1 - p_2) \\ &= m^2 - m^2 - 2p_4 \cdot (p_1 - p_2) \\ &= -2p_4 \cdot (p_1 - p_2) \end{aligned} \quad (3.1.18)$$

and

$$\begin{aligned} (p' + p_3)^\sigma q_\sigma &= (q + p_3 + p_3) \cdot q \\ &= q^2 + 2p_3 q \\ &= 2p_3 q \end{aligned} \quad (3.1.19)$$

At the denominator we have

$$\begin{aligned} k^2(p'^2 - m^2) &= (p_1 + p_2)^2 \left((p_3 + q)^2 - m^2 \right) \\ &= 2(m^2 + p_1 p_2)(p_3^2 + q^2 + 2p_3 q - m^2) \\ &= 2(m^2 + p_1 p_2)(2p_3 q) \end{aligned} \quad (3.1.20)$$

where we have exploited the fact that external particles are on-shell, so that $p_1^2 = p_2^2 = p_3^2 = p_4^2 = m^2$ and $q^2 = 0$ for a real photon. In the end \mathcal{A}_3 becomes

$$\mathcal{A}_3 = \frac{ie^3}{2(m^2 + p_1 p_2)(2p_3 q)} \left(-2p_4 \cdot (p_1 - p_2) \right) (2p_3 q) = -\frac{ie^3(p_1 - p_2) \cdot p_4}{m^2 + p_1 p_2} \quad (3.1.21)$$

With similar calculations, substituting each time the more suited expressions for k and p' , for the remaining amplitudes we get the following expressions

$$\mathcal{A}_4 = -\frac{ie^3(p_1 - p_2) \cdot p_3}{m^2 + p_1 p_2} \quad (3.1.22)$$

$$\mathcal{A}_5 = \frac{ie^3(p_3 - p_4) \cdot p_2}{m^2 + p_3 p_4} \quad (3.1.23)$$

$$\mathcal{A}_6 = \frac{ie^3(p_3 - p_4) \cdot p_1}{m^2 + p_3 p_4} \quad (3.1.24)$$

We can then sum all the amplitudes:

$$\begin{aligned} \mathcal{A} &= ie^3 \left[-\frac{(p_1 - p_2) \cdot q}{m^2 + p_1 p_2} - \frac{(p_3 - p_4) \cdot q}{m^2 + p_3 p_4} - \frac{(p_1 - p_2) \cdot p_4}{m^2 + p_1 p_2} - \frac{(p_1 - p_2) \cdot p_3}{m^2 + p_1 p_2} + \right. \\ &\quad \left. + \frac{(p_3 - p_4) \cdot p_2}{m^2 + p_3 p_4} + \frac{(p_3 - p_4) \cdot p_1}{m^2 + p_3 p_4} \right] \end{aligned} \quad (3.1.25)$$

therefore

$$\mathcal{A} = ie^3 \left[\frac{(p_1 - p_2) \cdot (-q - p_4 - p_3)}{m^2 + p_1 p_2} + \frac{(p_3 - p_4) \cdot (-q + p_1 + p_2)}{m^2 + p_3 p_4} \right] \quad (3.1.26)$$

Using once again 4-momentum conservation

$$p_1 + p_2 = p_3 + p_4 + q \quad (3.1.27)$$

we can eliminate every dependence on q , finally obtaining

$$\begin{aligned} \mathcal{A} &= ie^3 \left[\frac{-(p_1 - p_2) \cdot (p_1 + p_2)}{m^2 + p_1 p_2} + \frac{(p_3 - p_4) \cdot (p_3 + p_4)}{m^2 + p_3 p_4} \right] \\ &= ie^3 \left[\frac{-(p_1^2 + p_1 p_2 - p_1 p_2 - p_2^2)}{m^2 + p_1 p_2} + \frac{p_3^2 + p_3 p_4 - p_3 p_4 - p_4^2}{m^2 + p_3 p_4} \right] = 0 \end{aligned} \quad (3.1.28)$$

since $p_1^2 = p_2^2 = p_3^2 = p_4^2 = m^2$, so the Ward identity is satisfied.

3.2 Radiative corrections for $\psi\bar{\psi} \rightarrow \phi\phi^*\gamma$

Text. Consider the interaction between a complex scalar field, a fermionic field and the electromagnetic one, described by the following Lagrangian

$$\mathcal{L} = \bar{\psi}(i\not{D} - m_\psi)\psi + (D_\mu\phi)^*(D^\mu\phi) - m_\phi^2\phi^*\phi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$$




Show that the scattering amplitude for the process



$$\psi\bar{\psi} \rightarrow \phi\phi^*\gamma$$

vanishes after the substitution $\varepsilon_\mu(q) \rightarrow q_\mu$, where ε_μ is the polarization wave vector of the photon and q_μ its 4-momentum.

Solution. The Feynman rules for this Lagrangian are given by the usual rules for QED and for scalar QED (see exercise 2.2).

For this process, at first order ($\propto e^3$), we have the following diagrams





Explicitely the amplitudes are

$$\mathcal{A}_1 = \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}2ie^2g^{\nu\sigma}\varepsilon_\sigma^*(q) \quad (3.2.6)$$

$$\mathcal{A}_2 = \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p' - p_4)^\nu\frac{i}{p'^2 - m_\phi^2}(-ie)(p' + p_3)^\sigma\varepsilon_\sigma^*(q) \quad (3.2.7)$$

$$\mathcal{A}_3 = \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p_3 - p')^\nu\frac{i}{p'^2 - m_\phi^2}(-ie)(-p' - p_4)^\sigma\varepsilon_\sigma^*(q) \quad (3.2.8)$$

$$\mathcal{A}_4 = \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)\frac{i(p' + m_\psi)}{p'^2 - m_\psi^2}(-ie\gamma^\sigma)\varepsilon_\sigma^*(q)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p_3 - p_4)^\nu \quad (3.2.9)$$

$$\mathcal{A}_5 = \bar{v}^{s_2}(p_2)(-ie\gamma^\sigma)\varepsilon_\sigma^*(q)\frac{i(-p' + m_\psi)}{p'^2 - m_\psi^2}(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p_3 - p_4)^\nu \quad (3.2.10)$$

It is worth noting that in \mathcal{A}_5 the term p' in the numerator of the fermionic propagator has negative sign, since the fermionic current flows in the opposite direction to the flow of momentum.

Substituting the photon polarization with its momentum, that is $\varepsilon_\sigma^*(q) \rightarrow q_\sigma$, we

can simplify the amplitudes as follows

$$\begin{aligned}\mathcal{A}_1 &= \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}2ie^2g^{\nu\sigma}\varepsilon_\sigma^*(q) = \\ &= \frac{-2ie^3}{(p_1 + p_2)^2}\bar{v}^{s_2}(p_2)\gamma^\mu u^{s_1}(p_1)q_\mu = \\ &= \frac{-ie^3}{m_\psi^2 + p_1 \cdot p_2}\bar{v}^{s_2}(p_2)qu^{s_1}(p_1)\end{aligned}\tag{3.2.11}$$

$$\begin{aligned}\mathcal{A}_2 &= \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p' - p_4)^\nu\frac{i}{p'^2 - m_\phi^2}(-ie)(p' + p_3)^\sigma\varepsilon_\sigma^*(q) = \\ &= \frac{ie^3}{(p_1 + p_2)^2[(p_3 + q)^2 - m_\phi^2]}\bar{v}^{s_2}(p_2)(q + p_3 - p_4)u^{s_1}(p_1)(q + 2p_3)^\sigma q_\sigma = \\ &= \frac{ie^3}{2(m_\psi^2 + p_1 \cdot p_2)(2p_3 \cdot q)}\bar{v}^{s_2}(p_2)(q + p_3 - p_4)u^{s_1}(p_1)(2p_3 \cdot q) = \\ &= \frac{ie^3}{2(m_\psi^2 + p_1 \cdot p_2)}\bar{v}^{s_2}(p_2)(q + p_3 - p_4)u^{s_1}(p_1)\end{aligned}\tag{3.2.12}$$

$$\begin{aligned}\mathcal{A}_3 &= \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p_3 - p')^\nu\frac{i}{p'^2 - m_\phi^2}(-ie)(-p' - p_4)^\sigma\varepsilon_\sigma^*(q) = \\ &= \frac{ie^3}{(p_1 + p_2)^2[(p_4 + q)^2 - m_\phi^2]}\bar{v}^{s_2}(p_2)(p_3 - q - p_4)u^{s_1}(p_1)(-q - 2p_4)^\sigma q_\sigma = \\ &= \frac{-ie^3}{2(m_\psi^2 + p_1 \cdot p_2)(2p_4 \cdot q)}\bar{v}^{s_2}(p_2)(p_3 - q - p_4)u^{s_1}(p_1)(2p_4 \cdot q) = \\ &= \frac{-ie^3}{2(m_\psi^2 + p_1 \cdot p_2)}\bar{v}^{s_2}(p_2)(p_3 - q - p_4)u^{s_1}(p_1)\end{aligned}\tag{3.2.13}$$

$$\begin{aligned}\mathcal{A}_4 &= \bar{v}^{s_2}(p_2)(-ie\gamma^\mu)\frac{i(p' + m_\psi)}{p'^2 - m_\psi^2}(-ie\gamma^\sigma)\varepsilon_\sigma^*(q)u^{s_1}(p_1)\frac{-ig_{\mu\nu}}{k^2}(-ie)(p_3 - p_4)^\nu = \\ &= \frac{ie^3}{(p_3 + p_4)^2[(p_1 - q)^2 - m_\phi^2]}\bar{v}^{s_2}(p_2)\left[(p_3 - p_4)(p_1 - q + m_\psi)q\right]u^{s_1}(p_1) = \\ &= \frac{ie^3}{2(m_\phi^2 + p_3 \cdot p_4)(-2p_1 \cdot q)}\bar{v}^{s_2}(p_2)\left[(p_3 - p_4)(p_1q - qq + m_\psi q)\right]u^{s_1}(p_1)\end{aligned}\tag{3.2.14}$$

The spinors product can be simplified as

$$\bar{v}^{s_2}(p_2) \left[(\not{p}_3 - \not{p}_4)(\not{p}_1 \not{q} - \underbrace{\not{q}\not{q}}_{=0} + m_\psi \not{q}) \right] u^{s_1}(p_1) \quad (3.2.15)$$

The term $\not{p}_1 \not{q}$ can be rewritten as $\not{p}_1 \not{q} = 2p_1 q - \not{q}\not{p}_1$, so that

$$\mathcal{A}_4 = \frac{ie^3}{2(m_\phi^2 + p_3 p_4)(-2p_1 q)} \bar{v}^{s_2}(p_2) \left[(\not{p}_3 - \not{p}_4) \left(2p_1 q - \not{q}(\not{p}_1 - m_\psi) \right) \right] u^{s_1}(p_1) \quad (3.2.16)$$

Thanks to the Dirac equation $(\not{p}_1 - m_\psi)u^{s_1}(p_1) = 0$, we have

$$\begin{aligned} \mathcal{A}_4 &= \frac{-ie^3}{2(m_\phi^2 + p_3 p_4)(2p_1 q)} \bar{v}^{s_2}(p_2) \left[(\not{p}_3 - \not{p}_4)(2p_1 q) \right] u^{s_1}(p_1) = \\ &= \frac{-ie^3}{2(m_\phi^2 + p_3 p_4)} \bar{v}^{s_2}(p_2) \left[\not{p}_3 - \not{p}_4 \right] u^{s_1}(p_1) \end{aligned} \quad (3.2.17)$$

$$\begin{aligned} \mathcal{A}_5 &= \bar{v}^{s_2}(p_2) (-ie\gamma^\sigma) \varepsilon_\sigma^*(q) \frac{i(-\not{p}' + m_\psi)}{p'^2 - m_\psi^2} (-ie\gamma^\mu) u^{s_1}(p_1) \frac{-ig_{\mu\nu}}{k^2} (-ie)(p_3 - p_4)^\nu = \\ &= \frac{ie^3}{(p_3 + p_4)^2 \left[(p_2 - q)^2 - m_\phi^2 \right]} \bar{v}^{s_2}(p_2) \left[\not{q}(-\not{p}_2 + \not{q} + m_\psi)(\not{p}_3 - \not{p}_4) \right] u^{s_1}(p_1) = \\ &= \frac{ie^3}{2(m_\phi^2 + p_3 p_4)(-2p_2 q)} \bar{v}^{s_2}(p_2) \left[(-\not{q}\not{p}_2 + \underbrace{\not{q}\not{q}}_{=0} + \not{q}m_\psi)(\not{p}_3 - \not{p}_4) \right] u^{s_1}(p_1) \end{aligned} \quad (3.2.18)$$

As did for \mathcal{A}_4 , we simplify the spinors product by moving \not{q} to the right of \not{p}_2 . We then let $(\not{p}_2 + m_\psi)\not{q}$ act on the spinor $\bar{v}^{s_2}(p_2)$ from the right, which simply gives us zero thanks again to the Dirac equation $\bar{v}^{s_2}(p_2)(\not{p}_2 + m_\psi) = 0$.

It follows that

$$\begin{aligned} \mathcal{A}_5 &= \frac{ie^3}{2(m_\phi^2 + p_3 p_4)(-2p_2 q)} \bar{v}^{s_2}(p_2) \left[(-2p_2 q)(\not{p}_3 - \not{p}_4) \right] u^{s_1}(p_1) = \\ &= \frac{ie^3}{2(m_\phi^2 + p_3 p_4)} \bar{v}^{s_2}(p_2) \left[\not{p}_3 - \not{p}_4 \right] u^{s_1}(p_1) \end{aligned} \quad (3.2.19)$$

and by summing all the amplitudes we can check the validity of the Ward identity

$$\begin{aligned} \mathcal{A} = \sum_{j=1}^5 \mathcal{A}_j &= \frac{-ie^3}{m_\psi^2 + p_1 p_2} \bar{v}^{s_2}(p_2) q u^{s_1}(p_1) + \\ &+ \frac{ie^3}{2(m_\psi^2 + p_1 p_2)} \bar{v}^{s_2}(p_2) (\not{q} + \not{p}_3 - \not{p}_4) u^{s_1}(p_1) + \\ &+ \frac{-ie^3}{2(m_\psi^2 + p_1 p_2)} \bar{v}^{s_2}(p_2) (\not{p}_3 - \not{q} - \not{p}_4) u^{s_1}(p_1) + \\ &+ \frac{-ie^3}{2(m_\phi^2 + p_3 p_4)} \bar{v}^{s_2}(p_2) [\not{p}_3 - \not{p}_4] u^{s_1}(p_1) + \\ &+ \frac{ie^3}{2(m_\phi^2 + p_3 p_4)} \bar{v}^{s_2}(p_2) [\not{p}_3 - \not{p}_4] u^{s_1}(p_1) \end{aligned} \quad (3.2.20a)$$

$$\begin{aligned} \mathcal{A} &= \frac{ie^3}{m_\psi^2 + p_1 p_2} \bar{v}^{s_2}(p_2) \left[-\not{q} + \frac{1}{2} \left(\not{q} + \not{p}_3 - \not{p}_4 \right) - \frac{1}{2} \left(\not{p}_3 - \not{q} - \not{p}_4 \right) \right] u^{s_1}(p_1) + \\ &+ \frac{ie^3}{2(m_\phi^2 + p_3 p_4)} \bar{v}^{s_2}(p_2) \left[-(\not{p}_3 - \not{p}_4) + (\not{p}_3 - \not{p}_4) \right] u^{s_1}(p_1) = 0 \end{aligned} \quad (3.2.20b)$$

Appendix A

Useful formulae

EULER-LAGRANGE EQUATIONS

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} - \frac{\partial \mathcal{L}}{\partial \phi} = 0 \quad (\text{A.1})$$

HAMILTONIAN

$$\mathcal{H} = \sum_l \frac{\partial \mathcal{L}}{\partial \dot{\phi}_l} \dot{\phi}_l - \mathcal{L} \quad (\text{A.2})$$

NOETHER CURRENTS AND CONSERVED CHARGES

Internal symmetries

$$\varepsilon \bar{\delta} \phi := \phi' - \phi \quad (\text{A.3})$$

$$\mathcal{J}^\mu = \sum \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \bar{\delta} \phi, \quad \partial_\mu \mathcal{J}^\mu = 0 \quad (\text{A.4})$$

$$Q(t) = \int d^3x \mathcal{J}^0(\bar{x}, t) \quad (\text{A.5})$$

Space-time translations symmetry

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial^\nu \phi - \mathcal{L} g^{\mu\nu}, \quad \partial_\mu T^{\mu\nu} = 0 \quad (\text{A.6})$$

$$P^\nu = \int d^3x T^{0\nu} \quad (\text{A.7})$$

KLEIN-GORDON THEORY

$$\mathcal{L}_{KG} = \partial_\mu \phi \partial^\mu \phi - m^2 \phi^2 \quad (\text{A.8})$$

$$(\square + m^2) \phi = 0 \quad (\text{A.9})$$

Field expansion

$$\phi(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left(a_p e^{-ipx} + a_p^\dagger e^{ipx} \right) \quad (\text{A.10})$$

Ladder operators

$$[a_p, a_{p'}^\dagger] = (2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{p}'), \quad [a_p, a_{p'}] = [a_p^\dagger, a_{p'}^\dagger] = 0 \quad (\text{A.11})$$

$$|\mathbf{p}\rangle = \sqrt{2E_p}a_p^\dagger |0\rangle \quad (\text{A.12})$$

PAULI MATRICES

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (\text{A.13})$$

DIRAC MATRICES (CHIRAL REPRESENTATION)

$$\begin{aligned} \gamma^0 &= \begin{pmatrix} 0 & \mathbb{1}_{2x2} \\ \mathbb{1}_{2x2} & 0 \end{pmatrix}, & \gamma^i &= \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix} \\ \gamma^5 &= i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} -\mathbb{1}_{2x2} & 0 \\ 0 & \mathbb{1}_{2x2} \end{pmatrix} \end{aligned} \quad (\text{A.14})$$

γ matrices properties

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}\mathbb{1} \quad (\text{A.15})$$

$$(\gamma^\mu)^\dagger = \gamma^0\gamma^\mu\gamma^0 \quad (\text{A.16})$$

$$\gamma^\mu\gamma_\mu = 4\mathbb{1} \quad (\text{A.17})$$

$$(\gamma^5)^\dagger = \gamma^5, \quad (\gamma^5)^2 = \mathbb{1} \quad (\text{A.18})$$

$$\{\gamma^5, \gamma^\mu\} = 0 \quad (\text{A.19})$$

$$\gamma_\mu\gamma^\nu\gamma^\mu = -2\gamma^\nu \quad (\text{A.20})$$

$$\gamma^\mu\gamma^\nu\gamma^\rho\gamma_\mu = 4g^{\nu\rho}\mathbb{1}_4 \quad (\text{A.21})$$

$$\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma\gamma_\mu = -2\gamma^\sigma\gamma^\rho\gamma^\nu \quad (\text{A.22})$$

$$P_\mu\gamma^\mu P_\nu\gamma^\nu = P^2 \quad (\text{A.23})$$

Traces of γ matrices

$$Tr[\text{odd } \# \text{ of } \gamma] = 0, \quad Tr[\gamma^5 \cdot \text{odd } \# \text{ of } \gamma] = 0 \quad (\text{A.24})$$

$$Tr[\gamma^\mu\gamma^\nu] = 4g^{\mu\nu} \quad (\text{A.25})$$

$$Tr[\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma\gamma^\alpha\gamma^\beta] = 4(g^{\mu\nu}g^{\rho\sigma} - g^{\mu\rho}g^{\nu\sigma} + g^{\mu\sigma}g^{\nu\rho}) \quad (\text{A.26})$$

$$\begin{aligned} Tr[\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma\gamma^\alpha\gamma^\beta] &= 4g^{\mu\nu}(g^{\rho\sigma}g^{\alpha\beta} - g^{\rho\alpha}g^{\sigma\beta} + g^{\rho\beta}g^{\sigma\alpha}) - \\ &\quad - 4g^{\mu\rho}(g^{\nu\sigma}g^{\alpha\beta} - g^{\nu\alpha}g^{\sigma\beta} + g^{\nu\beta}g^{\sigma\alpha}) + \\ &\quad + 4g^{\mu\sigma}(g^{\nu\rho}g^{\alpha\beta} - g^{\nu\alpha}g^{\rho\beta} + g^{\nu\beta}g^{\rho\alpha}) - \\ &\quad - 4g^{\mu\alpha}(g^{\nu\rho}g^{\sigma\beta} - g^{\nu\sigma}g^{\rho\beta} + g^{\nu\beta}g^{\rho\sigma}) + \\ &\quad + 4g^{\mu\beta}(g^{\nu\rho}g^{\sigma\alpha} - g^{\nu\sigma}g^{\rho\alpha} + g^{\nu\alpha}g^{\rho\sigma}) \end{aligned} \quad (\text{A.27})$$

$$Tr[\gamma^\mu\gamma^\nu\gamma^5] = 0 \quad (\text{A.28})$$

Cyclic property of the trace

$$Tr[ABC] = Tr[CAB] = Tr[BCA] \quad (\text{A.29})$$

DIRAC THEORY

$$\mathcal{L}_D = i\bar{\psi}\not{\partial}\psi - m\bar{\psi}\psi \quad (\text{A.30})$$

$$\begin{cases} i\partial_\mu\bar{\psi}\gamma^\mu + m\bar{\psi} = 0 \\ -i\not{\partial}\psi + m\psi = 0 \end{cases}, \quad \bar{\psi} = \psi^\dagger\gamma^0, \quad \not{\partial} = \partial_\mu\gamma^\mu \quad (\text{A.31})$$

Dirac spinors

$$\begin{cases} [i\gamma^\mu p_\mu - m]u(p) = 0 \implies u(p) = \begin{pmatrix} \sqrt{p_\mu\sigma^\mu}\xi \\ \sqrt{p_\mu\bar{\sigma}^\mu}\xi \end{pmatrix} \\ [i\gamma^\mu p_\mu + m]v(p) = 0 \implies v(p) = \begin{pmatrix} \sqrt{p_\mu\sigma^\mu}\chi \\ -\sqrt{p_\mu\bar{\sigma}^\mu}\chi \end{pmatrix} \end{cases}, \quad \sigma^\mu = (\mathbb{1}, \sigma)^T, \quad \bar{\sigma}^\mu = (\mathbb{1}, -\sigma)^T \quad (\text{A.32})$$

Dirac spinor identities

$$u^{\tau\dagger}(p)u^\rho(p) = 2E_p\delta^{\tau\rho} \quad (\text{A.33})$$

$$v^{\tau\dagger}(p)v^\rho(p) = 2E_p\delta^{\tau\rho} \quad (\text{A.34})$$

$$\bar{u}^\tau(p)v^\rho(p) = \bar{v}^\tau(p)u^\rho(p) = 0 \quad (\text{A.35})$$

$$u^{\tau\dagger}(\mathbf{p})v^\rho(-\mathbf{p}) = v^{\tau\dagger}(-\mathbf{p})u^\rho(\mathbf{p}) = 0 \quad (\text{A.36})$$

$$\sum_{s=1}^2 u^s(p)\bar{u}^s(p) = \not{p} + m \quad (\text{A.37})$$

$$\sum_{s=1}^2 v^s(p)\bar{v}^s(p) = \not{p} - m \quad (\text{A.38})$$

Fields expansions

$$\psi(x) = \sum_{s=1}^2 \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[a_p^s u^s(p) e^{-ipx} + b_p^{s\dagger} v^s(p) e^{ipx} \right] \quad (\text{A.39})$$

$$\psi^\dagger(x) = \sum_{s=1}^2 \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[a_p^{s\dagger} u^{s\dagger}(p) e^{ipx} + b_p^s v^{s\dagger}(p) e^{-ipx} \right] \quad (\text{A.40})$$

Ladder operators

$$\begin{aligned} \{a_p^s, a_q^{r\dagger}\} &= \{b_p^s, b_q^{r\dagger}\} = (2\pi)^3 \delta^{rs} \delta^{(3)}(\mathbf{p} - \mathbf{q}) \\ \{a_p^s, a_q^r\} &= \{b_p^s, b_q^r\} = 0 \\ \{a_p^{s\dagger}, a_q^{r\dagger}\} &= \{b_p^{s\dagger}, b_q^{r\dagger}\} = 0 \\ \{a_p^s, b_q^r\} &= \{a_p^{s\dagger}, b_q^{r\dagger}\} = 0 \\ \{a_p^s, b_q^{r\dagger}\} &= \{a_p^{s\dagger}, b_q^r\} = 0 \end{aligned} \quad (\text{A.41})$$

$$|\mathbf{p}, s\rangle = \sqrt{2E_p} a_p^{s\dagger} |0\rangle \quad (\text{A.42})$$

ELECTROMAGNETIC THEORY

$$\mathcal{L}_{EM} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} \quad (\text{A.43})$$

$$\partial_\mu F^{\mu\nu} = 0, \quad F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu \quad (\text{A.44})$$

Gauge transformations

$$A_\mu(x) \rightarrow A_\mu(x) + \partial_\mu \lambda(x) \quad (\text{A.45})$$

Lorentz gauge and gauge-fixing term

$$\partial_\mu A^\mu = 0 \quad (\text{A.46})$$

$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} - \frac{1}{2\xi}(\partial_\mu A^\mu)^2 \quad (\text{A.47})$$

Field expansion

$$A_\mu(x) = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2|\mathbf{p}|}} \sum_{\lambda=0}^3 \left[\varepsilon_\mu^{(\lambda)}(p) a_p^{(\lambda)} e^{-ipx} + \varepsilon_\mu^{(\lambda)*}(p) a_p^{(\lambda)\dagger} e^{ipx} \right] \quad (\text{A.48})$$

$$p^\mu \varepsilon_\mu^{(1,2)} = 0 \quad (\text{A.49})$$

$$\sum_{\lambda=1}^2 \varepsilon_{(\lambda)}^\mu [\varepsilon_{(\lambda)}^\nu]^* = g^{\mu\nu} \quad (\text{A.50})$$

Ladder operators

$$[a_p^{(\lambda)}, a_q^{(\lambda')\dagger}] = -g^{\lambda\lambda'}(2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{q}) \quad (\text{A.51})$$

$$|\mathbf{p}, \lambda\rangle = a_p^{(\lambda)\dagger} |0\rangle \quad (\text{A.52})$$

INTERACTING FIELDS

Wick's theorem: let T be the time-ordering operator and N the normal-ordering operator, then:

$$\phi(x) = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[a_p e^{-ipx} + a_p^\dagger e^{ipx} \right] := \phi^+(x) + \phi^-(x)$$

$$\langle 0 | T\{\phi(x)\phi(y)\} | 0 \rangle = \begin{cases} \langle 0 | [\phi^+(x), \phi^-(y)] | 0 \rangle, & x^0 > y^0 \\ \langle 0 | [\phi^+(y), \phi^-(x)] | 0 \rangle, & x^0 < y^0 \end{cases} := \overline{\phi(x)\phi(y)} \quad (\text{A.53})$$

$$\psi(x) = \sum_{s=1}^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \left[a_p^s u^s(p) e^{-ipx} + b_p^{s\dagger} v^s(p) e^{ipx} \right] := \psi^+(x) + \psi^-(x)$$

$$\langle 0 | T\{\psi(x)\bar{\psi}(y)\} | 0 \rangle = \begin{cases} \langle 0 | \{\psi^+(x), \psi^-(y)\} | 0 \rangle, & x^0 > y^0 \\ \langle 0 | -\{\bar{\psi}^+(y), \psi^-(x)\} | 0 \rangle, & x^0 < y^0 \end{cases} := \overline{\psi(x)\bar{\psi}(y)} \quad (\text{A.54})$$

$$T\left\{ ABC \dots Z \right\} = N \left\{ ABC \dots Z + \text{all possible contractions of } ABC \dots Z \right\} \quad (\text{A.55})$$

S-matrix and scattering amplitude

$$S = T \left\{ \exp \left[-i \int d^4 x \mathcal{H}_{INT} \right] \right\}, \quad A = \langle in | S | out \rangle \quad (\text{A.56})$$

Differential cross section $|\mathbf{p}_1, \mathbf{p}_2\rangle \rightarrow |\mathbf{q}_1, \mathbf{q}_2\rangle$

$$d\sigma = \mathcal{J}|M|^2 d\Phi_2, \quad \begin{cases} \mathcal{J} = \frac{1}{4\sqrt{(p_1 p_2)^2 - m_1^2 m_2^2}} \\ |M|^2 = \sum_{s_i, \lambda_i, \dots} |A|^2 \\ d\Phi_2 = \left. \frac{|\mathbf{q}_1|}{E_1 + E_2} \frac{d \cos \theta d\varphi}{16\pi^2} \right|_{\mathbf{q}_2 = -\mathbf{q}_1} \end{cases} \quad (\text{A.57})$$

where m_1, m_2 are the initial masses, E_1, E_2 are the final energies and $|\mathbf{p}_2| = -|\mathbf{p}_1|$
Feynman rules for ϕ^4 -theory

$$\begin{array}{c} \xrightarrow{p} \\ \text{---} \end{array} = \frac{i}{p^2 - m^2 + i\varepsilon} \quad (\text{A.58})$$

$$\begin{array}{c} \diagup \\ \diagdown \end{array} = (-i\lambda) \quad (\text{A.59})$$

$$x \begin{array}{c} \text{---} \\ \xleftarrow{p} \end{array} = 1 \quad (\text{A.60})$$

QUANTUM ELECTRODYNAMICS

$$\mathcal{L}_{QED} = i\bar{\psi}\not{D}\psi - m\bar{\psi}\psi - \frac{1}{4}F^{\mu\nu}F_{\mu\nu}, \quad D_\mu = \partial_\mu + ieA_\mu \quad (\text{A.61})$$

Feynman rules for QED

$$\begin{array}{c} \xrightarrow{p} \\ \text{---} \end{array} = u^s(p) \quad \begin{array}{c} \diagup \\ \diagdown \end{array} = \bar{u}^s(p) \quad (\text{A.62})$$

$$\begin{array}{c} \xleftarrow{p} \\ \text{---} \end{array} = \bar{v}^s(p) \quad \begin{array}{c} \diagup \\ \diagdown \end{array} = v^s(p) \quad (\text{A.63})$$

$$\mu \begin{array}{c} \xrightarrow{p} \\ \sim \sim \sim \sim \end{array} = \varepsilon_\mu(p) \quad \begin{array}{c} \xrightarrow{p} \\ \sim \sim \sim \sim \end{array} \mu = \varepsilon_\mu^*(p) \quad (\text{A.64})$$

$$\begin{array}{c} \diagup \\ \diagdown \end{array} \sim \sim \sim \sim \mu = -ie\gamma^\mu \quad (\text{A.65})$$

$$\begin{array}{ccc} \overrightarrow{\text{---}} & = & \frac{i(p+m)}{p^2-m^2+i\varepsilon} \\ \overrightarrow{\text{---}} & & \end{array} \quad (\text{A.66})$$

$$\begin{array}{ccc} \mu \sim\sim\sim\nu & = & \frac{-ig_{\mu\nu}}{p^2+i\varepsilon} \\ \overleftarrow{p} & & \end{array} \quad (\text{A.67})$$

Fundamental processes in QED ($m_e \sim 0$)

- $e^+e^- \rightarrow \mu^+\mu^-$

$$|p_1, s_1; p_2, s_2\rangle \rightarrow |k_1, s_3; k_2, s_4\rangle \quad (\text{A.68})$$

$$|M|^2 = \frac{8e^4}{(p_1 + p_2)^4} \left[(p_1 k_1)(p_2 k_2) + (p_1 k_2)(p_2 k_1) + m_\mu^2 (p_1 p_2) \right] \quad (\text{A.69})$$

$$\frac{d\sigma}{d\cos\theta} = \frac{e^4}{128\pi} \frac{1}{E^2} \sqrt{1 - \frac{m_\mu^2}{E^2}} \left[\left(1 + \frac{m_\mu^2}{E^2} \right) + \left(1 - \frac{m_\mu^2}{E^2} \right) \cos\theta \right] \quad (\text{A.70})$$

- $e^+e^- \rightarrow e^+e^-$

$$|p_1, s_1; p_2, s_2\rangle \rightarrow |k_1, s_3; k_2, s_4\rangle \quad (\text{A.71})$$

$$\begin{aligned} |M|^2 = 8e^4 & \left\{ \frac{1}{(p_1 + p_2)^4} \left[(p_1 k_1)(p_2 k_2) + (p_1 k_2)(p_2 k_1) \right] + \right. \\ & \left. + \frac{1}{(p_1 - k_1)^4} \left[(p_1 p_2)(k_1 k_2) + (p_1 k_2)(p_2 k_1) \right] + \frac{2(p_1 k_2)(p_2 k_1)}{(p_1 + p_2)^2 (p_1 - k_1)^2} \right\} \end{aligned} \quad (\text{A.72})$$

$$\frac{d\sigma}{d\cos\theta} = \frac{e^4}{64\pi} \frac{1}{E^2} \left[\frac{(1 - \cos^2\theta)^2 + (1 + \cos^2\theta)^2}{4} + \frac{4 + (1 + \cos^2\theta)^2}{(1 - \cos^2\theta)^2} - (1 + \cos^2\theta) \right] \quad (\text{A.73})$$

- $e^+e^- \rightarrow \gamma\gamma$

$$|p_1, s_1; p_2, s_2\rangle \rightarrow |k_1, \lambda_1; k_2, \lambda_2\rangle \quad (\text{A.74})$$

$$|M|^2 = e^4 \left[\frac{1}{(p_2 k_2)^2} + \frac{1}{(p_2 k_1)^2} \right] \left[(p_1 p_2)(k_1 k_2) - (p_1 k_1)(p_2 k_2) - (p_1 k_2)(p_2 k_1) \right] \quad (\text{A.75})$$

$$\frac{d\sigma}{d\cos\theta} = \frac{e^4}{16\pi} \frac{1}{E^2} \left(\frac{1 + \cos^2\theta}{\sin^2\theta} \right) \quad (\text{A.76})$$

Bibliography

- [1] M. D. Schwartz. *Quantum Field Theory and the Standard Model*. Cambridge University Press (2018).
- [2] M. E. Peskin, D. V. Schroeder. *An Introduction to Quantum Field Theory*. Westview Press (1995).
- [3] A. Greco. *Elements of Quantum Field Theory*. Lectures notes for the Master's Degree in Physics, University of Ferrara (2019).