

NON-EXISTENCE OF A HOLOMORPHIC IMBEDDING OF THE SOBOLEV LOOP SPACE INTO THE HILBERT PROJECTIVE SPACE

ANAKKAR M., IVASHKOVICH S*.

ABSTRACT. The goal of this paper is to understand the properties of meromorphic mappings with values in two model complex Hilbert manifolds: Hilbert projective space $\mathbb{P}(l^2)$ and Sobolev loop space of the Riemann sphere LP^1 . It occurs that these properties are quite different. Based on our study we obtain as a corollary that LP^1 does not admit a closed holomorphic imbedding to $\mathbb{P}(l^2)$. In other words LP^1 is *not* a Hilbert projective variety despite of the fact that it is Kähler and meromorphic functions separate points on it.

CONTENTS

1. Introduction	1
2. Meromorphic functions on Hilbert manifolds	4
3. Meromorphic mappings to Hilbert projective space	10
4. Loop space of the Riemann sphere	16
References	21

1. INTRODUCTION

1.1. On the imbedding of the loop space to the projective space. By a complex Hilbert manifold we understand a Hausdorff topological space \mathcal{X} locally homeomorphic to an open subsets in a separable Hilbert space L , i.e., \mathbb{C}^n or l^2 , such that the transition maps are holomorphic. The notation l^2 stands for the Hilbert space of sequences of complex numbers $z = \{z_k\}_{k=1}^{\infty}$ such that $\|z\|^2 := \sum_k |z_k|^2 < \infty$ with the standard Hermitian scalar product $(z, w) = \sum_k z_k \bar{w}_k$ and the standard basis $\{e_1, e_2, \dots\}$. By $B(z_0, r)$ we denote the ball of radius r centered at z_0 in L , by $B(r)$ the ball centered at origin and by B the unit ball. If we want to underline that we are speaking about unit ball in l^2 we write B^∞ . We say that meromorphic functions separate points on the complex Hilbert manifold \mathcal{X} if for every pair $p \neq q$ of points of \mathcal{X} there exists a meromorphic function f on \mathcal{X} such that f is holomorphic near p and q and $f(p) \neq f(q)$. It is not difficult to see that a compact (\Rightarrow finite dimensional) complex manifold X such that meromorphic functions separate points on it admits a *meromorphic injection* to the complex projective space \mathbb{P}^N for some N , see Proposition 4.3. By a meromorphic injection we mean a meromorphic mapping $f : X \rightarrow \mathbb{P}^N$ such that it is a holomorphic injection outside of an analytic set of codimension ≥ 2 .

Date: June 4, 2024.

2020 Mathematics Subject Classification. Primary - 32D15, Secondary - 32A20, 46G20, 46T25.

Key words and phrases. Hilbert manifold, meromorphic map, loop space.

* Partially supported by the Labex Cempi ANR-11-LABX-0007-01.

It occurs not to be the case in infinite dimensions. Namely, despite of the fact that the loop space $L\mathbb{P}^1$ of the Riemann sphere possesses the property of meromorphic separation, see Proposition 4.2, it cannot be meromorphically injected to the Hilbert projective space $\mathbb{P}(l^2)$. We prove the following statement.

Theorem 1. *$L\mathbb{P}^1$ doesn't admit a closed meromorphic injection to $\mathbb{P}(l^2)$, i.e., there doesn't exist a meromorphic map $g : L\mathbb{P}^1 \rightarrow \mathbb{P}(l^2)$ such that:*

- i) $g : L\mathbb{P}^1 \setminus I \rightarrow \mathbb{P}(l^2)$ is a holomorphic injection, where I is analytic $\text{codim } I \geq 2$;
- i) dg_z is injective and has closed range for all $z \in L\mathbb{P}^1 \setminus I$;
- ii) the full image $g(L\mathbb{P}^1)$ is closed in $\mathbb{P}(l^2)$.

The non existence of an *equivariant holomorphic imbedding* with respect to the natural group actions on $L\mathbb{P}^1$ and $\mathbb{P}(l^2)$ was proved in [Z] answering the question from [MZ]. The proof in [Z] is based on the comparison of Picard groups. Our proof is based on the comparison of extension properties of meromorphic mappings with values in these manifolds.

1.2. Hilbert projective space. Let \mathcal{X} and \mathcal{Y} be complex Hilbert manifolds. Intuitively a meromorphic mapping $\mathcal{X} \rightarrow \mathcal{Y}$ should be defined by a holomorphic map $f : \mathcal{X} \setminus I \rightarrow \mathcal{Y}$, where I is a Hilbert analytic set in \mathcal{X} of codimension ≥ 2 such that the graph Γ_f of f “extends” in some sense to $\mathcal{X} \times \mathcal{Y}$. And this “extended object” is what we call a *meromorphic graph* (or, a *meromorphic map*) and denote it still as Γ_f (or, as f). The indeterminacy set of a meromorphic mapping f is the smallest closed subset I of \mathcal{X} such that f is holomorphic on $\mathcal{X} \setminus I$. It will be denoted as I_f . We do not claim that I_f is Hilbert analytic in general, but this will be so in both model examples of Hilbert manifolds mentioned above. Furthermore we shall investigate in what sense the aforementioned extension realizes on the examples of $L\mathbb{P}^1$ and $\mathbb{P}(l^2)$. They are quite different in nature but have one common feature, namely the notion of meromorphicity for them are somehow “predefined”.

We start with the case when \mathcal{Y} is the Hilbert projective space $\mathbb{P}(l^2) := (l^2 \setminus \{0\})/w \sim \lambda w$, $\lambda \neq 0$, where $w = (w_0, w_1, \dots) \in l^2$. By $z_k^j = w_j/w_k$ we denote the affine coordinates in the affine chart $\mathcal{U}_k = \{[w] : w_k \neq 0\} = \mathbb{P}(l^2) \setminus \mathcal{H}_k$, where \mathcal{H}_k is the hyperplane

$$\mathcal{H}_k := \{[w] \in \mathbb{P}(l^2) : w_k = 0\}. \quad (1.1)$$

Let a holomorphic mapping $f : \mathcal{X} \setminus I \rightarrow \mathbb{P}(l^2)$ be given. Assuming that $f(\mathcal{X} \setminus I) \not\subset \mathcal{H}_0$ set $A_0 := f^{-1}(\mathcal{H}_0)$. Observe that $f|_{\mathcal{X} \setminus A_0}$ is a holomorphic map with values in $\mathcal{U}_0 \equiv l^2$ and therefore writes as $f = (f_1, f_2, \dots)$. Here f_k are holomorphic functions on $\mathcal{X} \setminus A_0$ and such that $\|f(z)\|_{l^2}^2 = \sum_k |f_k(z)|^2$ is locally bounded, see subsection 2.2. We prove the following

Theorem 2. *The closure \bar{A}_0 of the divisor A_0 is a divisor in \mathcal{X} and all f_k extend meromorphically to the whole of \mathcal{X} . Moreover, the orders of the poles of these extensions are locally uniformly bounded along \bar{A}_0 . In addition the following is true:*

- i) For every point $a \in \mathcal{X}$ there exists a neighborhood $\mathcal{V} \ni a$ and holomorphic in \mathcal{V} functions $\varphi_0, \varphi_1, \dots$ without common factors such that
 - a) $\|\varphi(z)\|^2 = \sum_k |\varphi_k(z)|^2$ is locally bounded and $f_k = \varphi_k/\varphi_0$ on \mathcal{V} ;
 - b) $\mathcal{V} \cap I \supset \{z \in \mathcal{V} : \varphi_k(z) = 0 \text{ for all } k\}$.
- ii) The indeterminacy set I_f of the meromorphic map f thus obtained is locally defined by the equation $\varphi(z) = 0$, i.e., $I_f \cap \mathcal{V} = \{z \in \mathcal{V} : \varphi_0(z) = \varphi_1(z) = \dots = 0\}$.

Now it is clear that by a meromorphic mapping in the case of $\mathbb{P}(l^2)$ we should understand the object obtained by the extension from $\mathcal{X} \setminus I$ to \mathcal{X} of a holomorphic map f as in this theorem. It is worth to note that on $\mathcal{V} \setminus I_f$ our map is defined as

$$f(z) = [\varphi_0(z) : \varphi_1(z) : \dots] \quad (1.2)$$

for an appropriate holomorphic map $\varphi = (\varphi_0, \varphi_1, \dots)$ with values in l^2 . Therefore it is natural to give the following

Definition 1. *A meromorphic mapping from a Hilbert manifold \mathcal{X} to the Hilbert projective space $\mathbb{P}(l^2)$ is defined by a sequence f_1, f_2, \dots of meromorphic functions on \mathcal{X} such that locally there exist holomorphic functions $\varphi_0, \varphi_1, \dots$ with $\|\varphi\|_{l^2}$ bounded and $f_k = \frac{\varphi_k}{\varphi_0}$ for all $k \in \mathbb{N}$. Mapping f is locally given by (1.2) outside of the set of common zeroes of $\varphi_0, \varphi_1, \dots$*

We prove Thullen, Levi and Hartogs type extension theorems for meromorphic mappings with values in $\mathbb{P}(l^2)$, see Theorems 3.1 and 3.2 as well as Corollary 3.1. In this Introduction we shall need the following corollary from the Hartogs type extension.

Corollary 1. *Let Γ be a \mathcal{C}^1 -smooth closed curve in \mathbb{C}^2 and let $f : \mathbb{C}^2 \setminus \Gamma \rightarrow \mathbb{P}(l^2)$ be a holomorphic map. Then f extends meromorphically to the whole of \mathbb{C}^2 .*

This corollary follows from the Hartogs type extension theorem by placing a Hartogs figure H near a given point $a \in \Gamma$ in such a way that $H \cap \Gamma = \emptyset$ but the associated to H polydisk \mathbb{D}^2 contains a .

1.3. Case of loop spaces. Our second model example when the meromorphicity is predefined is the case when \mathcal{Y} is a Sobolev loop space of the Riemann sphere \mathbb{P}^1 , i.e. $L\mathbb{P}^1 = W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$. The natural complex structure on $L\mathbb{P}^1$ will be recalled in section 4. For the moment it will be sufficient to say that a holomorphic map $f : \mathcal{X} \rightarrow L\mathbb{P}^1$ from a complex Hilbert manifold \mathcal{X} to $L\mathbb{P}^1$ can be seen as a mapping $f : \mathcal{X} \times \mathbb{S}^1 \rightarrow \mathbb{P}^1$ such that:

- for every $s \in \mathbb{S}^1$ the map $f(\cdot, s) : \mathcal{X} \rightarrow \mathbb{P}^1$ is holomorphic,
- for every $z \in \mathcal{X}$ one has $f(z, \cdot) \in W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$ and the correspondence

$$\mathcal{X} \ni z \mapsto f(z, \cdot) \in W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$$

is continuous with respect to the standard topology on \mathcal{X} and the Sobolev topology on $W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$.

Such description will be called the *representation* of f . We shall prove the following

Theorem 3. *Any holomorphic mapping $f : \mathcal{X} \setminus I \rightarrow L\mathbb{P}^1$, where \mathcal{X} is a Hilbert manifold and I a Hilbert analytic set in \mathcal{X} of codimension ≥ 2 possesses the following properties:*

- i) *for every $s \in \mathbb{S}^1$ the map $f(\cdot, s)$ is a meromorphic function which meromorphically extends to the whole of \mathcal{X} ;*
- ii) *the family of indeterminacy sets $\{I_{f(\cdot, s)} : s \in \mathbb{S}^1\}$ is locally finite in \mathcal{X} ;*
- iii) *mapping f holomorphically extends to $\mathcal{X} \setminus I_f$, and therefore $I_f = \bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}$ is the indeterminacy set of the meromorphic mapping thus obtained.*

Notice that since every $I_{f(\cdot, s)}$ is an analytic set of pure codimension two such is their union I_f . These items force us to give the following

Definition 2. *A meromorphic mapping $f : \mathcal{X} \rightarrow L\mathbb{P}^1$ from a Hilbert manifold \mathcal{X} to the Sobolev loop space $W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$ is defined by a family of meromorphic on \mathcal{X} functions $\{f(\cdot, s) : s \in \mathbb{S}^1\}$ such that:*

- i) the family of their indeterminacy sets $\{I_{f(\cdot, s)} : s \in \mathbb{S}^1\}$ is locally finite in \mathcal{X} ;
- ii) f is holomorphic when restricted to $\mathcal{X} \setminus \bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}$.

Extension properties of meromorphic mappings with values in $L\mathbb{P}^1$ occur to be not as good as for those with values in $\mathbb{P}(l^2)$. In Example 4.1 we construct a holomorphic mapping $f : \mathbb{C}^2 \setminus \Gamma \rightarrow W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$, where $\Gamma = \{\gamma(s) = (\gamma_1(s), \gamma_2(s))\}$ is a loop in \mathbb{C}^2 of class $W^{2,2} \subset \mathcal{C}^{1, \frac{1}{2}}$, such that for every $s \in \mathbb{S}^1$ the point $\gamma(s)$ is an essential singularity of f . That is f doesn't extend to any neighborhood of $\gamma(s)$ even meromorphically. This example shows that neither Thullen no Hartogs type extension theorems are valid for meromorphic mappings with values in $L\mathbb{P}^1$.

Assumption that there exists a meromorphic injection $g : L\mathbb{P}^1 \rightarrow \mathbb{P}(l^2)$ quickly leads to a contradiction between the already mentioned Example 4.1 and Corollary 1, see the end of the section 4 for more details.

2. MEROMORPHIC FUNCTIONS ON HILBERT MANIFOLDS

2.1. Hilbert manifolds and analytic sets. For standard facts from infinite dimensional complex analysis we refer to [Mu]. By a complex Hilbert manifold in this paper we shall understand a Hausdorff topological space \mathcal{X} covered by coordinate charts (U_k, h_k) , where $h_k : U_k \rightarrow V_k$ are local homeomorphisms to an open subsets of a separable complex Hilbert space L , i.e., L is \mathbb{C}^n or l^2 , such that the transition maps are holomorphic. We shall say, if needed, that \mathcal{X} is modeled over L . If $L = \mathbb{C}^n$ we say that \mathcal{X} is n -dimensional, if $L = l^2$ that \mathcal{X} is infinite dimensional.

We shall consecutively use results about Hilbert analytic sets from [Ra], where they are proved for analytic sets in Banach spaces in which every closed subspace admits a closed complement. By ${}_L\mathcal{O}_{z_0}$ we denote the ring of germs of holomorphic functions at z_0 in the separable complex Hilbert space L . For $L = \mathbb{C}^n$ we write ${}_n\mathcal{O}_{z_0}$, for $L = l^2$ we write ${}_{l^2}\mathcal{O}_{z_0}$. But more often when L is clear from the context we shall write simply \mathcal{O}_{z_0} . This is a factorial domain, see Theorem I.1.4.4 in [Ra]. Let \mathcal{X} be a Hilbert manifold modeled over L .

Definition 2.1. By a Hilbert analytic set in \mathcal{X} we understand a subset $A \subset \mathcal{X}$ such that for every point $a \in \mathcal{X}$ there exists neighborhood $\mathcal{V} \ni a$ and a holomorphic mapping $h : \mathcal{V} \rightarrow F$ to a separable complex Hilbert space F such that

$$A \cap \mathcal{V} = \{z \in \mathcal{V} : h(z) = 0\}.$$

If $F = \mathbb{C}^n$ for all $a \in A$ we say that A is an analytic set of *finite definition*. If A is locally contained in a finite dimensional locally closed submanifold M of \mathcal{X} and is an analytic set in M we say that A is *finite dimensional*.

We denote the germ of an analytic subset A at a by A_a and by $\mathcal{J}(A_a) \subset \mathcal{O}_a$ the ideal of germs of holomorphic functions that vanish identically on A_a . We say that A is principal at a if the ideal $\mathcal{J}(A_a)$ is principal and we say that A is principal if A_a is principal at every $a \in A$. Given a holomorphic map $f : \mathcal{X} \rightarrow F$ we write $V(f) = f^{-1}(0)$ and if $f = (f_1, \dots, f_n)$ is \mathbb{C}^n -valued we write $V(f_1, \dots, f_n) = V(f_1) \cap \dots \cap V(f_n)$. Given $a \in \mathcal{X}$ the zero locus of a holomorphic germ $V(f_a)$ is defined by the germ of analytic set at a of the zero locus of f , i.e $V(f_a) = (V(f))_a$. We say that a germ of an analytic subset A_a is of codimension p if the height of the ideal $\mathcal{J}(A_a)$ is equal to p . This is equivalent to saying that

$$\sup \{n \in \mathbb{N} \mid \exists E \text{ subspace of } L, \dim E = n \text{ and } (a + E) \cap A_a = \{a\}\} = p,$$

see corollary on the page 70 of [Ra]. And we say that A is of codimension p if $\text{codim}(A_a) = p$ at every $a \in A$. It is proved in [Ra], see Proposition II.3.3.3 that for any $d \in \mathbb{N}$ the germ A_a of a Hilbert analytic set A at a admits the following decomposition

$$A_a = \left(\bigcup_{k \leq d} A_a^k \right) \cup A_{a,d}, \quad (2.1)$$

where A_a^k are germs of codimension k for $k = 1, \dots, d$ and $A_{a,d}$ is the union of germs of codimension $> d$.

For a Hilbert analytic set in \mathcal{X} the regular part A^{reg} of A is the set of points $a \in A$ such that A is a submanifold of \mathcal{X} in a neighborhood of a . $A^{\text{sing}} := A \setminus A^{\text{reg}}$ is called the singular locus of A . Clearly A^{reg} is a union of arcwise connected components, which are locally closed submanifolds of \mathcal{X} . Let B^k be such a component of codimension k (suppose such exists). It is proved in [Ra], see Theorem II.4.1.2 that the closure \bar{B}^k is a Hilbert analytic subset of \mathcal{X} of codimension k which is globally irreducible in \mathcal{X} . We call \bar{B}^k an irreducible k -codimensional component of A . The union of all \bar{B}^k -s is locally finite in \mathcal{X} , see Proposition II.3.3.2 in [Ra].

If A is an analytic set in a complex Hilbert manifold \mathcal{X} of codimension ≥ 2 then A doesn't contain a principal germ at any of its points. This means in other words that in decomposition (2.1) $A_a^1 = \emptyset$ for every $a \in A$. The following statement gives a more detailed description of analytic sets of codimension ≥ 2 . It is proved in [Ra], see Lemma II.1.1.12. We shall give the proof of it as well as of Proposition 2.1 in order to make our exposition more accessible.

Lemma 2.1. *Let A be a germ at zero of an analytic set in a complex Hilbert space L . If A does not contain a principal germ at zero then there exist holomorphic germs f_1 and f_2 at zero such that $A \subset V(f_1, f_2)$. Moreover $V(f_1, f_2)$ does not contain a principal germ at zero.*

Proof. Since A is analytic there exists $h \in \mathcal{O}(L, F)$, with F a complex Hilbert space, such that $A = V(h)$. Since $h \not\equiv 0$ we find a non-zero element u_1 of the dual F^* such that $u_1 \circ h \not\equiv 0_{L^*}$. Let $u_1 \circ h = g_1^{\alpha_1} \dots g_n^{\alpha_n}$ be the decomposition to irreducible factors and set $f_1 = g_1 \dots g_m$. Choose a direction $v \in L$ such that f_1 does not vanish identically on $\langle v \rangle$ and consider the decomposition $L = \langle v \rangle^\perp \oplus \langle v \rangle$ with corresponding coordinates (z', z'') . The Weierstrass Preparation Theorem applied to f_1 gives that $f_1 = AP_1$ where $P_1(z', z'') \in \mathcal{O}(\langle v \rangle)[z'']$ is a Weierstrass polynomial and A is an invertible germ. Divide h by P_1 using the Weierstrass Division theorem and get $h = P_1Q + R$ where $Q \in \mathcal{O}(L, F)$ and $R \in \mathcal{O}(\langle v \rangle^\perp, F)[z'']$ with $\text{deg } R < \text{deg } P_1$.

Now assume that for $u \in F^*$ we have $u \circ h|_{V(f_1)} = 0$. Then there exists $q \in \mathcal{O}(L)$ such that $u \circ h = P_1q$. At the same time we have $u \circ h = P_1(u \circ Q) + u \circ R$ and we can deduce from the unicity of the Euclidean division that $u \circ R = 0$. If this holds for every $u \in F^*$ we obtain that $R = 0$ and $h = P_1Q$. Therefore A would contain the principal germ $V(P_1) = V(f_1)$. Contradiction. We conclude that exists $u_2 \in F^*$ such that $f_2 = u_2 \circ h$ satisfies $f_2|_{V(f_1)} \neq 0$. Now $A \subset V(f_1, f_2)$ and $V(f_1, f_2)$ doesn't contain a principal germ at zero as stated. □

The following proposition is a particular case of Proposition III.2.1.3 from [Ra].

Proposition 2.1. *Let \mathcal{X} be a complex Hilbert manifold and A an analytic subset of codimension ≥ 2 of \mathcal{X} . Let B be an analytic set of pure codimension one in $\mathcal{X} \setminus A$. Then \overline{B} is an analytic subset of \mathcal{X} .*

Proof. The question is local, therefore the only thing we need to do is to extend B to a neighborhood of any point $a \in \overline{B} \cap A$. In the sequel we assume that $a = 0$ and let \mathcal{U} be a neighborhood of 0 in L . By Lemma 2.1 we can suppose that $A = V(f_1, f_2)$. If for every direction $v \in L$ the analytic in $\langle v \rangle \cap (\mathcal{U} \setminus A)$ set $\langle v \rangle \cap \mathcal{U} \cap B$ is not discrete in $\langle v \rangle \cap (\mathcal{U} \setminus A)$ we have that B contains an open subset of L , which is a contradiction.

Therefore there exists a direction $v \in L$ such that $A \cap \langle v \rangle = \{0\}$ and $\langle v \rangle \cap B$ is discrete in $\mathcal{U} \setminus A$. Decompose $L = L' \oplus \langle v \rangle$ where $L' = \langle v \rangle^\perp$. Taking \mathcal{U} smaller, if necessary, we can assume that $\mathcal{U} = \mathcal{U}' \times \mathcal{U}''$ corresponding to this decomposition. Moreover, we can chose \mathcal{U}' and \mathcal{U}'' in such a way that $\overline{B} \cap (\mathcal{U}' \times \partial \mathcal{U}'') = \emptyset$. Let $\pi : \mathcal{U} \rightarrow \mathcal{U}'$ be the orthogonal projection to \mathcal{U}' and let P_1 and P_2 be Weierstrass polynomials of f_1 and f_2 respectively. Denote by $\rho(P_1, P_2) \in \mathcal{O}(\mathcal{U}')$ the resultant of P_1 and P_2 . We have that $\pi(A) = \{x \in \mathcal{U}' \mid \rho(P_1, P_2) = 0\}$ and the latter is an analytic subset of \mathcal{U}' of codimension one. Set $\mathcal{U}_0 = (\mathcal{U}' \setminus \pi(A)) \times \mathcal{U}''$. Then $(\overline{B} \cap \mathcal{U}_0, \pi|_{\overline{B} \cap \mathcal{U}_0}, \mathcal{U}' \setminus \pi(A))$ is a ramified cover. Denote by d the number of preimages of a generic $z' \in \mathcal{U}' \setminus \pi(A)$. The fiber over z' writes as

$$\pi^{-1}(\{z'\}) = \{z' + z_1''(z')v, \dots, z' + z_d''(z')v\}.$$

Consider the function $P : L' \times \mathbb{C} \rightarrow \mathbb{C}$ defined by

$$P(x', z) = \prod_{j=1}^d (z - x_j''(x')). \quad (2.2)$$

We have that $B \cap \mathcal{U}_0 = \{z' + zv \in \mathcal{U} \setminus A \mid P(z', z) = 0\}$. Elementary symmetric functions of $\{z_j''(z')\}_{j=0}^d$ are bounded and holomorphic and therefore extend holomorphically to \mathcal{U}' . This gives us the extension of B to \mathcal{U} . □

2.2. Holomorphic functions and mappings. An l^2 -valued holomorphic mapping from an open subset \mathcal{X} of l^2 is a mapping $f : \mathcal{X} \rightarrow l^2$ which is Fréchet differentiable at all points of \mathcal{X} . It is well known that $f : \mathcal{X} \rightarrow l^2$ is holomorphic if and only if it is G -holomorphic, i.e., Gâteaux differentiable, and $\|f(z)\|$ is locally bounded, see Proposition 8.6 and Theorem 8.7 in [Mu]. This implies that if f is written in coordinates as

$$f(z) = (f_1(z), f_2(z), \dots) \quad (2.3)$$

then f is holomorphic if and only if all f_k are holomorphic functions and $\|f(z)\|^2 = \sum_{k=1}^{\infty} |f_k(z)|^2$ is locally bounded. Indeed, since we need to prove the Gâteaux differentiability only we can suppose that $z \in \mathbb{C}$ and then Cauchy formula and local boundedness of $\|f(z)\|$ gave us the local boundedness of $\|f'(z)\|$ where $f'(z) = (f'_1(z), f'_2(z), \dots)$. Indeed

$$\|f'(z)\|^2 = \frac{1}{(2\pi)^2} \sum_k \left| \int_{|\zeta-z|=\varepsilon} \frac{f_k(\zeta)}{(\zeta-z)^2} d\zeta \right|^2 \leq \frac{1}{(2\pi)^2 \varepsilon^4} \sum_k \left(\int_0^{2\pi} |f_k(\zeta)| \varepsilon d\theta \right)^2 \leq \quad (2.4)$$

$$\leq \frac{1}{(2\pi\varepsilon)^2} \sum_k \int_0^{2\pi} |f_k(\zeta)|^2 d\theta \cdot 2\pi \leq \frac{1}{2\pi\varepsilon^2} \sup_{|\zeta-z|=\varepsilon} \|f(\zeta)\|^2.$$

In the same fashion one proves the local boundedness of $\|f''(z)\|$ and hence the continuity of $f'(z)$. And this in its turn implies Gâteaux differentiability:

$$\begin{aligned} \|f(z+h) - f(z) - f'(z)h\| &= \left\| \int_0^1 \frac{df(z+th)}{dt} dt - f'(z)h \right\| = \\ &= \left\| \int_0^1 (f'(z+th) - f'(z)) dt \cdot h \right\| = o(h) \end{aligned}$$

by the continuity of $f'(z)$.

Lemma 2.2. *Let A be an analytic subset of the ball $B \subset L$ that doesn't contain a principal germ at any of its points. Then every holomorphic map $h \in \mathcal{O}(B \setminus A, F)$, where F is a Hilbert space, extends holomorphically to B .*

Proof. It is sufficient to prove that h extends to a neighborhood of a given point, say a zero point. By the Lemma 2.1 there exist germs f_1 and f_2 such that the germ of A at zero is contained in $V(f_1, f_2)$. Choose a direction $a \in L$ such that f_1 and f_2 do not vanish identically on $\text{Vect}(a)$ and decompose $L = L' \oplus \text{Vect}(a)$. The Weierstrass Preparation Theorem gives the Weierstrass polynomials $P_1(z', z)$ and $P_2(z', z)$ in $\mathcal{O}(L')[z]$ such that $X \subset V(f_1, f_2) = V(P_1, P_2)$. Therefore there exists $r_1 > 0$ such that $\{0_{L'}\} \times \bar{\Delta}_{r_1} \subset B$ and $P_1(0_{L'}, z)$ and $P_2(0_{L'}, z)$ do not vanish on $\{0_{L'}\} \times \partial\Delta_{r_1}$.

The number of roots of $P_1(0_{L'}, z)$ and $P_2(0_{L'}, z)$ is finite. By continuity of P_1 and P_2 there exists r_2 such that P_1 and P_2 do not vanish on $B(0_{L'}, r_2) \times \partial\Delta_{r_1}$, i.e. $V(P_1, P_2) \subset B(0_{L'}, r_2) \times \bar{\Delta}_{r_1}$. We define \tilde{h} by

$$\tilde{h}(z', z) = \frac{1}{2\pi i} \int_{\partial\Delta_{r_1}} h(z', u) \frac{du}{u-z}. \quad (2.5)$$

This \tilde{h} a holomorphic function on $B(0_{L'}, r_2) \times \Delta_{r_1}$. Let Q_1, \dots, Q_m the irreducible factors of P_1 and consider the resultant $\rho_j = \text{Res}(Q_j, P_2)$. Q_j does not divide P_2 because in the other case $V(Q_j)$ would be a principal germ contained in $V(P_1, P_2)$. Therefore $\rho_j \neq 0$. Consider $Y = V(\rho_1 \dots \rho_m)$. Let $\pi : L \rightarrow L'$ be the canonical projection. Since $\pi(A)$ is contained in Y , \tilde{h} coincides with h on $B(0_{L'}, r_2) \setminus (Y \times \bar{\Delta}_{r_1})$. Therefore \tilde{h} is the holomorphic extension of h to a neighborhood of 0. □

Remark 2.1. Using (2.5) and the Riemann extension theorem one can prove that if A is a proper analytic subset of a Hilbert manifold \mathcal{U} and h is a bounded holomorphic function on $\mathcal{U} \setminus A$ then h extends holomorphically to \mathcal{U} .

2.3. Meromorphic functions. The field quotient of ${}_L\mathcal{O}_{z_0}$ is denoted as ${}_L\mathcal{M}_{z_0}$ (or, simply as \mathcal{M}_{z_0}) and is called the ring of germs of meromorphic functions at z_0 . A meromorphic function on a Hilbert manifold \mathcal{X} is defined by the following data: an open covering $\{\mathcal{U}_k\}$ of \mathcal{X} , holomorphic functions $h_k, g_k \in \mathcal{O}(\mathcal{U}_k)$ with $g_k \not\equiv 0$ such that $h_k g_j = h_j g_k$ on $\mathcal{U}_k \cap \mathcal{U}_j$. By $\mathcal{M}(\mathcal{X})$ we denote the field of meromorphic functions on \mathcal{X} .

Lemma 2.3. *A formal power series with coefficients in the integral domain \mathcal{O}*

$$F(\lambda) = \sum_{n=1}^{\infty} a_{-n} \lambda^n \in \mathcal{O}[[\lambda^{-1}]]$$

represents a rational function $\frac{P(\lambda)}{Q(\lambda)}$ with $P, Q \in \mathcal{O}[\lambda]$ and $\deg Q \leq N$ if and only if

$$\begin{vmatrix} a_{-n_1} & a_{-n_2} & \cdots & a_{-n_{N+1}} \\ \vdots & \vdots & \vdots & \vdots \\ a_{-n_1-N} & a_{-n_2-N} & \cdots & a_{-n_{N+1}-N} \end{vmatrix} \quad (2.6)$$

for all $(N+1)$ -tuples $n_1 < \cdots < n_{N+1}$.

Proof. Indeed we look for a non-zero polynomial $Q(\lambda) = c_0 + c_1 \lambda + \cdots + c_N \lambda^N$ with coefficients in \mathcal{O} such that $FP \in \mathcal{O}[\lambda]$. But this condition means that for every $k \geq 1$ one should have

$$a_{-k} c_0 + \cdots + a_{-k-N} c_N = 0. \quad (2.7)$$

This relation means that vectors $b_k := (a_{-k}, a_{-k-1}, \dots, a_{-k-N})$, $k \in \mathbb{N}$ belong to the hyperplane with the equation 2.7 in the \mathcal{M} -linear space \mathcal{M}^{N+1} . The latter means that every $N+1$ of them are linearly dependent, and this is precisely what tells the condition 2.6

□

Let A be a subset of B^∞ . We say that A is thick at $z_0 \in B^\infty$ if for any neighborhood $\mathcal{V} \ni z_0$ the set $A \cap \mathcal{V}$ is not contained in a proper analytic subset of \mathcal{V} . The domain R_{r_1, r_2}^∞

$$R_{r_1, r_2}^\infty = A_{r_1, r_2} \times B^\infty \quad (2.8)$$

will be called a ring domain. Here $A_{r_1, r_2} = \Delta_{r_1} \setminus \bar{\Delta}_{r_2}$ is an annulus, $0 \leq r_1 < r_2$.

Lemma 2.4. *Let f be a holomorphic function in the ring domain $R_{1-r, 1}^\infty$. Suppose that for w in some subset $A \subset B^\infty$ thick at origin restrictions $f_w := f(\cdot, w)$ meromorphically extend from $A_{1-r, 1}$ to Δ and the number of poles counted with multiplicities of these extensions is uniformly bounded. Then f extends to a meromorphic function on $\Delta \times B^\infty$.*

Proof. Write f as

$$f(z, w) = f^+(z, w) + f^-(z, w) = \sum_{n \geq 0} a_n(w) z^n + \sum_{n < 0} a_n(w) z^n \quad (2.9)$$

where $a_n \in \mathcal{O}(B^\infty)$. Notice that f^+ is already holomorphic in $\Delta \times B^\infty$. Our task therefore is to extend f^- . By Lemma 2.3 applied to the ring \mathbb{C} the extendability of $f^-(z, w)$ for $w \in B^\infty$ to the disk together with the condition on poles means that for $a_n = a_n(w)$ the determinants 2.6 vanish. Therefore they vanish identically as functions of w . And therefore, again by Lemma 2.3 but this times applied to the ring $\mathcal{O}(B^\infty)$, we have that $f^-(z, w)$ is rational over the field $\mathcal{M}(B^\infty)$. I.e., is meromorphic on $\Delta \times B^\infty$.

□

Lemma 2.5. *Let I be an analytic set with $\text{codim } I \geq 2$ in a complex Hilbert manifold \mathcal{X} and let f be a meromorphic function on $\mathcal{X} \setminus I$. Then f extends to a meromorphic function \tilde{f} on the whole of \mathcal{X} .*

Proof. Denote \mathcal{P}_f the divisor of poles of f . $P_f \subset \mathcal{X} \setminus I$ is a complex hypersurface and by Proposition 2.1 its closure of $\overline{P_f}$ is an analytic subset of \mathcal{X} . Let $a \in I$. If there is a neighborhood U of a such that $\overline{P_f} \cap U = \emptyset$ then $f \in \mathcal{O}(U \setminus I)$ extends holomorphically to U by Lemma 2.2. Let $a \in I \cap \overline{P_f}$. Choose coordinates (z, w) in a neighborhood U of $a = 0$ associated to a decomposition $L = \langle b \rangle \oplus L'$ for some $b \in L$ such that $\langle b \rangle \cap \overline{P_f}$ is discrete. Now one can place a ring domain $R_{1-r,1}^\infty \subset U \setminus \overline{P_f}$. Therefore f will be holomorphic on $R_{1-r,1}^\infty$ and for every $w_0 \in \{w \mid (z, w) \in (\Delta \times B^\infty) \setminus I\}$ $f(\cdot, w)$ extends meromorphically to Δ . By Lemma 2.4 f extends meromorphically to a neighborhood of 0 and therefore f extends meromorphically to \mathcal{X} . □

We continue with the Thullen type extension of meromorphic functions.

Theorem 2.1. *Let A be an analytic subset of a complex Hilbert manifold \mathcal{X} and let f be a meromorphic function on $\mathcal{X} \setminus A$. Suppose that for every irreducible component A_1 of A of codimension one there exists a point $p_1 \in A_1$ such that f meromorphically extends to some neighborhood V_1 of p_1 . Then f meromorphically extends onto the whole of \mathcal{X} .*

Proof. If A has codimension ≥ 2 irreducible components then by Lemma 2.5 f extends across them. Now consider A_1 a irreducible component with $\text{codim } A_1 = 1$. Let p_1 the point where f extends and take p_2 another point of A_1 . Consider the path $\gamma : [0, 1] \rightarrow A_1$ such that $\gamma(0) = p_1$ and $\gamma(1) = p_2$. Let define the set E by

$$E = \{t \in [0, 1] \mid f \text{ extends meromorphically to a neighborhood of } \gamma([0, t])\}. \quad (2.10)$$

Let $t_0 = \sup E$. We need to prove that $t_0 = 1$. Suppose not. Then one can find a complex disk Δ through $\gamma(t_0)$ transverse to A_1 which do not intersects other components of A and is not contained in the set of indeterminacy of f . We can find a local coordinates (λ, w) with center $\gamma(t_0)$ such that $\Delta \times \{0\}$ is our disk. By Lemma 2.4 f extends meromorphically to a neighborhood of $\gamma(t_0)$. Contradiction. Therefore $t_0 = 1$ and f extends meromorphically to A . □

The infinite dimensional 1-concave Hartogs figure $\mathcal{H}_1^\infty(r) \subset l^2$ was defined in the introduction.

Corollary 2.1. *Every $f \in \mathcal{M}(H_1^\infty(r))$ extends meromorphically to $\Delta \times B^\infty$.*

Proof. Notice first that the set I of such $w \in B^\infty$ that $I_f \supset A_{1-r,1} \times \{w\}$ is a proper analytic subset of B^∞ . One easily local derives equations for I from that of I_f . If $w_0 \in I$ then for $z_0 \in A_{1-r,1}$ the equation of I_f in a neighborhood of (z_0, w_0) have the form $f(z, w) = \sum_{k=0}^\infty f_k(w - w_0)(z - z_0)^k = 0$. For $w = w_0$ all $f_k \equiv 0$ and $I = \{f_1(w - w_0) = f_2(w - w_0) = \dots = 0\}$. Extension of f to $\Delta \times (B^\infty \setminus I)$ can be achieved as in the proof of Theorem 2.1 using Theorem 2.4. Now f is meromorphic on $(\Delta \times B^\infty) \setminus (\Delta \times I)$ and meromorphic on $A_{1-r,1} \times B^\infty$. So we are under assumptions of the Thullen-type Corollary 2.1 and conclude that f extends to a meromorphic functions on $\Delta \times B^\infty$. □

3. MEROMORPHIC MAPPINGS TO HILBERT PROJECTIVE SPACE

3.1. Holomorphic mappings to Hilbert projective space. In the definition of the Hilbert projective space over l^2 one requires that homogeneous coordinates $[w] = [w_0 : w_1 : \dots]$ are such that $w = (w_0, w_1, \dots) \in l^2 \setminus \{0\}$. Notice that the affine coordinate (z_1^j, z_2^j, \dots) in the chart \mathcal{U}_j belong to l^2 as well. Denote by $h_j : \mathcal{U}_j \rightarrow l^2$ the corresponding coordinate map, namely $h_j([w]) = \{\frac{w_k}{w_j}\}_{k \neq j}$. Transition maps

$$h_j \circ h_i^{-1} : l^2 \setminus \{z_j^i = 0\} \rightarrow l^2 \setminus \{z_i^j = 0\}$$

are given by

$$h_j \circ h_i^{-1} : (z_0^i, z_1^i, \dots) \mapsto (z_0^i/z_j^i, z_1^i/z_j^i, \dots) = (z_0^j, z_1^j, \dots). \quad (3.1)$$

They are obviously biholomorphic and therefore $\mathbb{P}(l^2)$ is a complex Hilbert manifold modeled over l^2 . Closed submanifolds of $\mathbb{P}(l^2)$ will be called Hilbert projective or simply projective manifolds. I.e, a Hilbert projective manifold is a closed subset \mathcal{M} of $\mathbb{P}(l^2)$ such that for every point $m \in \mathcal{M}$ there exists a neighborhood $\mathcal{U} \ni m$ and a holomorphic map $h : \mathcal{U} \rightarrow L$ to a Hilbert space L with surjective differential such that $\mathcal{M} = \{z \in \mathcal{U} : h(z) = 0\}$. Their intersections with affine coordinate charts \mathcal{U}_j will be called affine. As well as closed submanifolds of l^2 will be called Stein.

Let \mathcal{X} be a complex Hilbert manifold and let $f : \mathcal{X} \rightarrow \mathbb{P}(l^2)$ be a holomorphic mapping. Without loss of generality we assume that the image of f is not contained in any hyperplane \mathcal{H}_j , see (1.1). Otherwise we would get $f : \mathcal{X} \rightarrow \mathcal{H}_j \cong \mathbb{P}(l^2)$ and so on. Taking into account the fact that

$$\mathbb{P} := \bigcap_{\mathcal{H}_j \supset f(\mathcal{X})} \mathcal{H}_j$$

is a closed linear subspace of $\mathbb{P}(l^2)$ isomorphic either to $\mathbb{P}(l^2)$ or to \mathbb{P}^N for some $N \in \mathbb{N}$, we find ourselves under our assumption with \mathbb{P} as a target space of f . Set $A_j := f^{-1}(\mathcal{H}_j)$. These are hypersurfaces (some of them may be empty) in \mathcal{X} . For the restriction $f|_{\mathcal{X} \setminus A_0} : \mathcal{X} \setminus A_0 \rightarrow \mathcal{U}_0$ the composition $h_0 \circ f$ is a holomorphic l^2 -valued mapping. In affine coordinates $z_1^0, \dots, z_k^0, \dots$ it writes as

$$(h_0 \circ f)(z) = (f_1^0(z), f_2^0(z), \dots), \quad (3.2)$$

where f_k^0 are holomorphic and $\|(h_0 \circ f)(z)\|^2 = \sum_k |f_k^0(z)|^2$ is locally bounded in $\mathcal{X} \setminus A_0$.

Proposition 3.1. *Holomorphic on $\mathcal{X} \setminus A_0$ functions $\{f_k^0\}$ are meromorphic on the whole of \mathcal{X} . Moreover, their orders of poles along A_0 are locally bounded and consequently our mapping in a neighborhood of any point $a \in \mathcal{X}$ writes as*

$$f(z) = [\varphi_0(z) : \varphi_1(z) : \dots], \quad (3.3)$$

where φ_k are holomorphic, have no common zeros and, in addition, $\|\varphi(z)\|^2 := \sum_k |\varphi_k(z)|^2$ is locally bounded.

Proof. Take a point $a \in A_0$ such that $a \notin A_1$ i.e., $f(a) \notin \mathcal{H}_1$. Then the holomorphic map $h_1 \circ f : \mathcal{X} \setminus A_1 \rightarrow \mathcal{U}_1$ writes as $h_1 \circ f = (f_0^1, f_2^1, \dots)$ with $f_0^1 = 1/f_1^0, f_2^1 = f_2^0/f_1^0, \dots, f_k^1 = f_k^0/f_1^0, \dots$ holomorphic in a neighborhood of a and $\|(f_0^1(z), f_2^1(z), \dots)\|$ locally bounded there. This proves that f_k^0 are meromorphic in a neighborhood of a and therefore on $\mathcal{X} \setminus (A_0 \cap A_1)$. The same argument can be repeated for any A_j and since $\bigcap A_j = \emptyset$ we see that f_k^0 are meromorphic on the whole of \mathcal{X} .

To prove the second assertion fix again a point $a \in A_0$ and let $h = 0$ be the primitive equation of A_0 in a neighborhood U of a . Existence of such h follows from the fact that the ring \mathcal{O}_a is factorial. We can choose U in such a way that $U = \Delta \times B$, where B is the unit ball and $A_0 \cap (\partial\Delta \times B) = \emptyset$. Such U will be called *adapted* to A_0 .

Claim 1. For every $k \in \mathbb{N}$ there exists $N_k \in \mathbb{N}_0 := \{0\} \cup \mathbb{N}$ such that

$$f_k^0 = \frac{\psi_k}{h^{N_k}} \quad \text{with } \psi_k \text{ holomorphic and not divisible by } h \text{ in } U. \quad (3.4)$$

To prove this claim suppose first that $A_0 \cap U$ is irreducible. Let $P(z_1, z')$ be the Weierstrass polynomial for h , i.e., $h(z_1, z') = z_1^p + a_1(z')z_1^{p-1} + \dots + a_p(z')$ with $a_j \in \mathcal{O}(B), a_j(0') = 0$ and $h = P\varphi$, where φ is holomorphic in U and doesn't vanish. Since f_k^0 has poles at most on $A_0 \cap U$ we can multiply it by P^{N_k} , where $N_k \in \mathbb{N}_0$ is the order of the pole of f_k^0 along $A_0 \cap U$ to get a holomorphic function ψ_k in U which is obviously not divisible by h . If $A_0 \cap U$ is reducible represent it as a union of a finite number of irreducible components $A_0 \cap U = \bigcup_l A_0^l$ with primitive equations $h^l = 0$ and repeat the argument as above to each A_0^l . The rest is obvious.

Claim 2. Next we claim that the sequence $\{N_k\}$ in 3.4 is bounded. If not one can find a subsequence $\{k_n\}$ such that $N_{k_n} \nearrow +\infty$ and

$$f_{k_n}^0 = \frac{\psi_{k_n}}{h^{N_{k_n}}} \quad \text{with } \psi_{k_n} \text{ holomorphic and not divisible by } h \text{ in } U. \quad (3.5)$$

Since $\bigcap A_j = \emptyset$ there exists m such that $a \in A_0 \setminus A_m$. Then

$$f_j^m = \frac{f_j^0}{f_m^0} = \frac{\psi_j}{\psi_m} \cdot h^{N_m - N_j}$$

should be holomorphic in a neighborhood of a for all j . But this is certainly not true for $j = k_n$ with n big enough, contradiction.

Let $N = \max_{k \in \mathbb{N}} N_k$ and set $\varphi_0 := h^N$, $\varphi_k := f_k^0 h^N$ for $k = 1, 2, \dots$. Consider the map

$$\varphi := (\varphi_0, \varphi_1, \dots) : U \rightarrow l^2. \quad (3.6)$$

We need to prove that φ indeed takes its values in l^2 and is locally bounded on U . Remark that $\|\varphi(z)\| = |h(z)|^N \|(1, (h_0 \circ f)(z))\|$ for z in a neighborhood of $\partial\Delta \times B$ and therefore is *locally* bounded there. Using compactness of $\partial\Delta$ we deduce that $\|\varphi\|$ is bounded on $\partial\Delta \times B_\delta$, say by M , for $\delta > 0$ small enough. Each cut $\sum_{k=0}^N |\varphi_k|^2$ is plurisubharmonic on U and therefore is bounded by M on $\Delta \times B_\delta$. Tending $N \rightarrow \infty$ we get the boundedness of $\|\varphi\|$ on $\Delta \times B_\delta$.

All what left to prove is that $\varphi_0, \varphi_1, \dots$ can be taken without common zeroes. Set $I = \{z \in U : \varphi_0(z) = \varphi_1(z) = \dots = 0\}$. If φ_j -s have common divisor in ${}_L\mathcal{O}_a$ we can divide them by this divisor and this will not change our map f . Therefore $\text{codim } I \geq 2$. Suppose it is non-empty and choose some $b \in I$. Our map on $U \setminus I$ is given by

$$f(z) = [\varphi_0(z) : \varphi_1(z) : \dots].$$

Since f is holomorphic in a neighborhood of b there exists j such that $f(b) \in \mathcal{U}_j$. Therefore $\varphi_j(b) \neq 0$. Contradiction. □

Remark 3.1. 1. The affine coordinate chart \mathcal{U}_0 will be considered as distinguished and f_k^0 -s will be denoted simply as f_k -s. We proved that our holomorphic map f is represented

by a sequence of meromorphic on \mathcal{X} functions $(f_1, f_2, \dots, f_k, \dots)$ which are holomorphic on $\mathcal{X} \setminus A_0 = f^{-1}(\mathcal{U}_0)$ and $\|f\|^2 = \sum_{k=1}^{\infty} \|f_k\|^2$ is locally bounded on $\mathcal{X} \setminus A_0$.

2. Moreover, we proved that for every point $a \in \mathcal{X}$ there exists a neighborhood $\mathcal{V} \ni a$, holomorphic in \mathcal{V} functions $\varphi_0, \varphi_1, \dots$ without common zeroes such that $\varphi = (\varphi_0, \varphi_1, \dots)$ is locally bounded as a mapping $\mathcal{V} \rightarrow l^2$ and such that our f is represented on \mathcal{V} as in (3.3), i.e., $f = \pi \circ \varphi$, where $\pi : l^2 \setminus \{0\} \rightarrow \mathbb{P}(l^2)$ is the canonical projection.

3. Remark that if a holomorphic mapping $f : \mathcal{X} \rightarrow \mathbb{P}(l^2)$ is given on an open $\mathcal{V} \subset \mathcal{X}$ by a representation (3.3) then $A_j \cap \mathcal{V} = \{\varphi_j = 0\}$. Indeed, $A_j \cap \mathcal{V}$ is the preimage of \mathcal{H}_j under φ , here $\mathcal{H}_j = \{w \in l^2 : w_j = 0\}$, since it is the preimage of \mathcal{H}_j under f .

3.2. Meromorphic mappings to $\mathbb{P}(l^2)$. Recall that a meromorphic mapping between finite dimensional complex manifolds X and Y is a holomorphic mapping $f : X \setminus I \rightarrow Y$ where I is analytic set with $\text{codim } I \geq 2$, such that the closure $\bar{\Gamma}_f$ of its graph Γ_f is an analytic set in $X \times Y$ and the restriction of the natural projection $\pi_1 : X \times Y \rightarrow X$ to $\bar{\Gamma}_f$ is proper. With some ambiguity this meromorphic mapping is denoted still as f , instead of $\bar{\Gamma}_f$ one writes simply Γ_f and calls it the graph of f .

Therefore an object which pretends to be a meromorphic map in Hilbert case should be at least holomorphic outside of an analytic set of codimension two.

Proposition 3.2. *Let \mathcal{X} be a complex Hilbert manifold, I a codimension at least 2 analytic set in \mathcal{X} and let $f : \mathcal{X} \setminus I \rightarrow \mathbb{P}(l^2)$ be a holomorphic mapping. Then for every $a \in I$ there exists a neighborhood $\mathcal{U} \subset \mathcal{X}$ and holomorphic in \mathcal{U} functions $\{\varphi_k\}$ such that*

- i) $\varphi = (\varphi_0, \varphi_1, \dots)$ is a holomorphic mapping from \mathcal{U} to l^2 ;
- ii) $\mathcal{U} \cap I \supset \{z \in \mathcal{U} : \varphi_k(z) = 0 \text{ for all } k\}$;
- iii) $(\pi \circ \varphi)(z) = f(z)$ for $z \in \mathcal{U} \setminus I$, where $\pi : l^2 \setminus \{0\} \rightarrow \mathbb{P}(l^2)$ is the canonical projection.

Proof. As we saw in Proposition 3.1 mapping f is defined by a sequence $\{f_k\}_{k \in \mathbb{N}}$ of meromorphic functions on $\mathcal{X} \setminus I$ which are holomorphic on $\mathcal{X} \setminus (I \cup A_0)$ where $A_0 = f^{-1}(\mathcal{H}_0)$. We can extend f_k meromorphically to \mathcal{X} by Lemma 2.5, denote by \tilde{f}_k these extensions, and by Proposition 2.1 we can extend A_0 to an analytic set \tilde{A}_0 in \mathcal{X} . Given $a \in I$ we need to examine f in a neighborhood of a . If $a \in I \setminus \tilde{A}_0$ all f_k extend holomorphically to a neighborhood of a by Lemma 2.2, i.e., our mapping f extends holomorphically to a neighborhood of such a . Suppose now that $a \in I \cap \tilde{A}_0$. Meromorphic functions \tilde{f}_k have poles at most on \tilde{A}_0 and we need to prove that their orders stay bounded, i.e., that there exists N_k such that $h^{N_k} \tilde{f}_k$ is holomorphic on a neighborhood of a for every $k \in \mathbb{N}$ and $N = \sup_{k \in \mathbb{N}} N_k < \infty$. Here h is a minimal defining function of \tilde{A}_0 at a . We assume that that N_k are minimal that do the job. Suppose this is not true, i.e., there exists a subsequence $N_{k_j} \rightarrow \infty$ such that $h^{N_{k_j}} \tilde{f}_{k_j}$ are holomorphic but $h^{N_{k_j}-1} \tilde{f}_{k_j}$ are not. But $h = 0$ is an equation for \tilde{A}_0 in a neighborhood of a and we get in this case that $h^{N_{k_j}-1} \tilde{f}_{k_j}$ are not holomorphic in this neighborhood. This contradicts to Proposition 3.1. Now set $\varphi_0 = h^N$ and $\varphi_k = h^N \tilde{f}_k$. Then ii) and iii) are satisfied. The same consideration as in Proposition 3.1 gives us i).

□

The discussion above leads us to the following

Definition 3.1. *Let \mathcal{X} be a complex Hilbert manifold. A meromorphic map $f : \mathcal{X} \rightarrow \mathbb{P}(l^2)$ is given by a holomorphic map $f : \mathcal{X} \setminus I \rightarrow \mathbb{P}(l^2)$, where $\text{codim } I \geq 2$, such that for every $a \in I$ there exists a neighborhood \mathcal{V} of a and a holomorphic map $\varphi = \{\varphi_k\}_{k \in \mathbb{N}} : \mathcal{V} \rightarrow l^2$ such that $f = \pi \circ \varphi$, i.e. $f(z) = [\varphi_0(z) : \varphi_1(z) : \dots]$ for $z \in \mathcal{V} \setminus I$.*

The minimal such I is called the indeterminacy set of f and is denoted as I_f . We can interpret Proposition 3.2 as follows : if $f : \mathcal{X} \setminus I \rightarrow \mathbb{P}(l^2)$ is a holomorphic map and $\text{codim } I \geq 2$ then f extends meromorphically to \mathcal{X} . One more remark: every meromorphic map $f : \mathcal{X} \rightarrow \mathbb{P}(l^2)$ can be represented by globally defined meromorphic functions $\{f_k\}$ such that their denominators are bounded in the following sense. Let $A_0 = f^{-1}(\mathcal{H}_0)$ be the preimage of the hyperplane $\mathcal{H}_0 = \{z_0 = 0\}$ under the holomorphic map $f : \mathcal{X} \setminus I_f \rightarrow \mathbb{P}(l^2)$. Then A_0 extends to an analytic hypersurface in the whole of \mathcal{X} , we denote it still as A_0 . And now the poles of all f_k are on A_0 and their orders are uniformly bounded.

The indeterminacy set I of a meromorphic mapping between Hilbert manifolds might look not as an analytic set in finite dimensions. And, moreover, the graph of f might be not a direct analytic subset of $\mathcal{X} \times \mathcal{Y}$ in the sense that it cannot be locally represented as a finite analytic cover.

3.3. Example. Consider the following mapping $f : l^2 \rightarrow \mathbb{P}(l^2)$

$$f : z = \{z_k\}_{k=1}^{\infty} \mapsto [\{z_k(z_k - 1/k)\}_{k=1}^{\infty}], \quad (3.7)$$

or, in more details

$$\{z_k\}_{k=1}^{\infty} \mapsto [z_1(z_1 - 1) : \dots : z_k(z_k - 1/k) : \dots].$$

The indeterminacy set I of f (i.e., all z such that all components in (3.7) vanish) is of Cantor type, see pp. 33-34 in [Ra]. In particular it is totally disconnected and contains the sequence $\{e_1, \dots, \frac{1}{k}e_k, \dots\}$ converging to zero.

Fix $z = \{z_j\}_{j=1}^{\infty} \in l^2$ and fix $k \in \mathbb{N}^*$. Consider $y_{n,k} \in l^2$ defined for $n \in \mathbb{N}^*$ by

$$\begin{aligned} y_{n,k} &= \left(\frac{1}{k} + \frac{kz_k}{n^2} \right) e_k + \sum_{j=1, j \neq k}^n \frac{-jz_j}{n^2} e_j = \\ &= \left(\frac{-z_1}{n^2}, \frac{-2z_2}{n^2}, \dots, \frac{-(k-1)z_{k-1}}{n^2}, \frac{1}{k} + \frac{kz_k}{n^2}, \frac{-(k+1)z_{k+1}}{n^2}, \dots, \frac{-nz_n}{n^2}, 0, \dots \right). \end{aligned}$$

Notice that $y_{n,k} \xrightarrow[n \rightarrow \infty]{} \frac{1}{k}e_k$. For n big enough we have

$$f(y_{n,k}) = \left[\frac{-z_1}{n^2} \left(\frac{-z_1}{n^2} - 1 \right) : \frac{-2z_2}{n^2} \left(\frac{-2z_2}{n^2} - \frac{1}{2} \right) : \dots : \left(\frac{1}{k} + \frac{kz_k}{n^2} \right) \frac{kz_k}{n^2} : \dots \right] \quad (3.8)$$

$$= \left[-z_1 \left(\frac{-z_1}{n^2} - 1 \right) : -2z_2 \left(\frac{-2z_2}{n^2} - \frac{1}{2} \right) : \dots : \left(\frac{1}{k} + \frac{kz_k}{n^2} \right) kz_k : \dots \right]. \quad (3.9)$$

Therefore $f(y_{n,k}) \xrightarrow[n \rightarrow \infty]{} [z_1 : z_2 : \dots : z_k : \dots]$ and consequently $(y_{n,k}, f(y_{n,k})) \xrightarrow[n \rightarrow \infty]{} (\frac{1}{k}e_k, [z])$. This means that $\pi_1^{-1}(\{\frac{1}{k}e_k\}) = \mathbb{P}(l^2)$ for every $k \in \mathbb{N}^*$, i.e., the set $\pi_1^{-1}(I) \supset \{\frac{1}{k}e_k\} \times \mathbb{P}(l^2)$ is not direct.

3.4. Restrictions to complex curves. Let us prove that any meromorphic mapping $f : \mathcal{X} \rightarrow \mathbb{P}(l^2)$ restricted to a complex curve is holomorphic. Since f is meromorphic on \mathcal{X} it exists an analytic subset $I \subset \mathcal{X}$ of codimension ≥ 2 such that f is holomorphic on $\mathcal{X} \setminus I$. Let $\varphi : \Delta \rightarrow \mathcal{X}$ be a complex curve which is not contained in I . Then φ is holomorphic on $\Delta \setminus \varphi^{-1}(I)$. Moreover the subset $\varphi^{-1}(I)$ is discrete. Let $\lambda_0 \in \varphi^{<-1>}(I)$. Consider a ball $B \subset \mathcal{X}$ centered on $\varphi(\lambda_0)$ such that there exists $F \in \mathcal{O}(B, l^2)$ with $f|_{B \setminus I} = \pi \circ F$. Then there exists $0 < r < 1$ such that $\partial\Delta(\lambda_0, r) \cap I = \emptyset$ and $\varphi(\Delta(\lambda_0, r)) \subset B$.

For every $j \in \mathbb{N}$ we have $F_j \circ \varphi(\lambda_0) = 0$. So there exists $k_j \in \mathbb{N}$ such that

$$F_j \circ \varphi(\lambda) = (\lambda - \lambda_0)^{k_j} G_j(\lambda)$$

with $G_j \in \mathcal{O}(B)$ and G_j does not vanish on $\Delta(\lambda_0, r)$. Let $k = \min\{k_j \mid j \in \mathbb{N}\}$. Therefore

$$f \circ \varphi(\lambda) = [F_0 \circ \varphi(\lambda) : F_1 \circ \varphi(\lambda) : \dots] = [(\lambda - \lambda_0)^{k_0 - k} G_0(\lambda) : (\lambda - \lambda_0)^{k_1 - k} G_1(\lambda) : \dots]$$

with

$$(\lambda - \lambda_0)^{2k} \sum_{j=0}^{+\infty} |(\lambda - \lambda_0)^{k_j - k} G_j(\lambda)|^2 = \sum_{j=0}^{+\infty} |F_j \circ \varphi(\lambda)|^2 = \|F \circ \varphi(\lambda)\|^2 < \infty$$

and then $f \circ \varphi$ is holomorphic on λ_0 .

3.5. Extension properties of meromorphic maps to $\mathbb{P}(l^2)$. Now we shall prove the following generalization of the classical Levi, Thullen and Hartogs type theorems. We start with Levi-Thullen.

Theorem 3.1. *Let \mathcal{X} be a complex Hilbert manifold and let $f : \mathcal{X} \setminus A \rightarrow \mathbb{P}(l^2)$ be a meromorphic mapping where A is analytic and*

- i) *either $\text{codim } A \geq 2$ or, more generally,*
- ii) *for every irreducible component A_α of A of codimension one there exists a point $p_\alpha \in A_\alpha$ such that f meromorphically extends to some neighborhood V_α of p_α .*

Then f meromorphically extends to the whole of \mathcal{X} .

Proof. i) We start with codimension ≥ 2 and denote A as I . Our f is defined on $\mathcal{X} \setminus I$ by meromorphic functions $\{f_k\}$ as above. Let A_0 be as above, i.e., A_0 is the set of poles of all f_k and their orders along A_0 are uniformly bounded. A_0 extends to an analytic hypersurface in the whole of \mathcal{X} by Proposition 2.1 and f_k meromorphically extend to \mathcal{X} by Lemma 2.5. All we need to prove that the orders of poles of these extensions stay uniformly bounded. We proceed as in the proof of Proposition 3.1. Let $a \in \tilde{A}_0 \setminus A_0$. There exists a neighborhood U of a and $g \in \mathcal{O}(U)$ such that $U \cap \tilde{A}_0 = \{g = 0\}$. Consider the Weierstrass polynomial P of g such that $g = P\varphi$ where φ is holomorphic and does not vanish on U . Since the poles of f_k are uniformly bounded on $U \cap A_0$ we can multiply them by P^L to obtain a holomorphic map $\psi_k = f_k P^L$ on $U \cap A_0$. ψ_k extends holomorphically to U and then the poles are uniformly bounded on \tilde{A}_0 .

ii) Represent f by meromorphic functions $\{f_k\}_{k \in \mathbb{N}^*}$ on $(\mathcal{X} \setminus A) \cup_\alpha V_\alpha$. By Theorem 2.1 every f_k extends meromorphically to the whole of \mathcal{X} . Notice that at p_α the orders of poles of f_k -s are uniformly bounded by say N_α . Let $q_\alpha \in A_\alpha$ be another point. We want to prove that the orders of poles of f_k -s are bounded at q_α with the same N_α . Clearly we can assume that both p_α and q_α are smooth points of A_α . Consider a path $\gamma : [0, 1] \rightarrow A_\alpha^{\text{reg}}$ from p_α to q_α . Set

$$E = \{t \in [0, 1] \mid \text{all } f_k \text{ have orders of poles bounded by } N_\alpha \text{ at } \gamma(t)\}.$$

E is nonempty and obviously open. Let $t_0 = \sup E$ and let \mathcal{V} be a neighborhood of $\gamma(t_0)$ in \mathcal{X} . Suppose there exists k such that $h^{N_\alpha} f_k$ is not holomorphic at $\gamma(t_0)$. Take $t_1 < t_0$ such that $t_1 \in E$. Now $h^{N_\alpha} f_k$ is holomorphic in $\mathcal{V} \setminus A_\alpha \cup \{\text{neighborhood of } \gamma(t_1)\}$. By Thullen type extension for holomorphic functions it extends holomorphically to a neighborhood of $\gamma(t_0)$. Contradiction. □

The n -dimensional Hartogs figure for $n \geq 2$ is the following open subset of \mathbb{C}^n

$$\mathcal{H}_1^{n-1}(r) = (\Delta \times B^{n-1}(r)) \cup (A_{1-r,1} \times B^{n-1}). \quad (3.10)$$

Here $r \in]0, 1[$ and $A_{1-r,1} = \Delta \setminus \bar{\Delta}_{1-r}$ is an annulus. Analogously an infinite dimensional Hartogs figure is the following open subset of l^2

$$\mathcal{H}_1^\infty(r) = (\Delta \times B^\infty(r)) \cup (A_{1-r,1} \times B^\infty). \quad (3.11)$$

Denote by $H(r)$ either of these figures and by B the unit ball in \mathbb{C}^{n-1} or l^2 respectively.

Theorem 3.2. *Every $f \in \mathcal{M}(H(r), \mathbb{P}(l^2))$ extends meromorphically to $\Delta \times B$.*

Proof. We give the proof in l^2 . f is represented by functions $f_k \in \mathcal{M}(H_1^\infty(r)) \cap \mathcal{O}(H_1^\infty(r) \setminus A_0)$. By Corollary 2.1 f_k extend meromorphically to $\Delta \times B^\infty$. Let A_0 be the divisor of poles of f_1 in $\Delta \times B^\infty$. Every irreducible component of A_0 intersects $\mathcal{H}_1^\infty(r)$. To prove this choose a straight one-dimensional closed disk $D \subset B^\infty$ in such a way that $\Delta \times D$ intersects the chosen component. The question is reduced now to a two-dimensional case.

Take $a_1 \in A_0^{\text{reg}}$ and choose $a_1 \in \tilde{A}_0^{\text{reg}} \cap \mathcal{H}_1^\infty(r)$ on the same irreducible component as a_0 . Consider a path $\gamma : [0, 1] \rightarrow \tilde{A}_0^{\text{reg}}$ such that $\gamma(0) = a_0$ and $\gamma(1) = a_1$. Let E be the set

$$E = \{t \in [0, 1] \mid \text{all } f_k \text{ have bounded denominators in a neighborhood of } \gamma(t)\}. \quad (3.12)$$

Since f is meromorphic on $\mathcal{H}_1^\infty(r)$ the set $E \neq \emptyset$ and obviously open. Denote $t_0 = \sup E$. In a neighborhood \mathcal{U} of $\gamma(t_0)$ functions f_k have bounded denominators h_k^N where N is the uniform bound. In the same manner as in the proof of Theorem 3.1 one proves that at $\gamma(t_1)$ the orders of poles of f_k are bounded by N . □

Recall that a domain \mathcal{X} in a complex Hilbert manifold is pseudoconcave at a boundary point a if there exists a direction $v \in T_a^c \partial \mathcal{X}$ on which the Levi form of some defining function is negative definite. Extension of an analytic object across a pseudoconcave boundary point of a domain in a complex Hilbert manifold is equivalent to the Hilbert-Hartogs extension. This was explained in [AI] for the case of holomorphic maps. Then the previous theorem implies the following

Corollary 3.1. *If a domain \mathcal{X} in a complex Hilbert manifold \mathcal{X} is pseudoconcave at a boundary point p then every meromorphic map $f : \mathcal{X} \rightarrow \mathbb{P}(l^2)$ extends meromorphically to a neighborhood of p .*

Proof. The problem is local in \mathcal{X} and therefore we can assume that $\mathcal{X} = l^2$. The pseudoconcavity of the point $p \in \partial \mathcal{X}$ implies that there exists $U \subset l^2$ a neighborhood of p such that $U \cap \mathcal{X} = \{z \in U \mid u(z) < 0\}$ with $\nabla u \neq 0$. Take $v \in T_p \partial \mathcal{X} \subset l^2$ such that the Levi form $\mathcal{L}_{u,p}(v) < 0$. After a complex linear change of coordinates we can suppose that $p = 0$ and $v = e_1, \nabla u_p = e_2$. We define the following figure $\varphi : \mathcal{H}_1^\infty(r) \rightarrow l^2$ by

$$z \mapsto \eta z_1 e_1 + (\eta \epsilon z_2 - r') e_2 + \delta \sum_{s=3}^{\infty} z_s e_s. \quad (3.13)$$

We choose ϵ, r', δ and η such that $\varphi(\mathcal{H}_1^\infty(r)) \subset U \cap \mathcal{X}$ and $\eta \epsilon > r'$ to insure that the image of the polydisk $\varphi(\Delta \times B^\infty)$ contains the origin.

Fix h and j and consider now the map $f \circ \varphi : \mathcal{H}_1^\infty(r) \rightarrow \mathbb{P}(l^2)$. By Theorem 3.2 $f \circ \varphi$ extends meromorphically to $\tilde{f} : \Delta \times B^\infty \rightarrow \mathbb{P}(l^2)$. Then $\tilde{f} \circ \varphi^{-1}$ gives the desired extension to a neighborhood of p .

□

4. LOOP SPACE OF THE RIEMANN SPHERE

4.1. Complex structure on the loop space of a complex manifold. Let \mathbb{S}^1 be the unit circle and consider the Hilbert manifold $W^{k,2}(\mathbb{S}^1, X)$ of Sobolev $W^{k,2}$ -mappings of \mathbb{S}^1 to a complex manifold X , $\dim_{\mathbb{C}} X = n \geq 1$. Notice that by Sobolev imbedding

$$W^{1,2}(\mathbb{R}, \mathbb{R}^N) \subset \mathcal{C}^{\frac{1}{2}}(\mathbb{R}, \mathbb{R}^N), \quad N = 2n \quad \text{in our case,} \quad (4.1)$$

and this inclusion is a compact operator. In particular mappings from $W^{1,2}(\mathbb{S}^1, X)$ are Hölder $\frac{1}{2}$ -continuous. From now on we shall always assume that $k \geq 1$. Sobolev topology on $W^{k,2}(\mathbb{S}^1, X)$ can be defined as follows. Take $g \in W^{k,2}(\mathbb{S}^1, X)$ and cover $g(\mathbb{S}^1)$ by finitely many coordinate charts (X_j, α_j) , $\alpha_j : X_j \rightarrow \mathbb{C}^n$. Similarly cover \mathbb{S}^1 by charts (S_j, β_j) making sure that $g(\bar{S}_j) \subset X_j$ for all j . In Hilbert spaces $W^{k,2}(S_j, \mathbb{C}^n)$ take neighborhoods \mathcal{V}_j of $\alpha_j \circ g \circ \beta_j^{-1}$. Then define a neighborhood of g in $W^{k,2}(\mathbb{S}^1, X)$ as

$$\mathcal{V} := \{f \in W^{k,2}(\mathbb{S}^1, X) : \alpha_j \circ f \circ \beta_j^{-1} \in \mathcal{V}_j \text{ for all } j\}. \quad (4.2)$$

As for the complex structure on $W^{k,2}(\mathbb{S}^1, X)$ recall first of all that if $g, h \in W^{k,2}(\mathbb{R})$ then $gh \in W^{k,2}(\mathbb{R})$, and this operation is continuous in Sobolev norm, see for example Lemma 2.2 in [A]. This enables us to define correctly a $W^{k,2}$ -vector bundle over \mathbb{S}^1 as well as a pull-back g^*TX of a tangent bundle of X under a $W^{k,2}$ -map $g : \mathbb{S}^1 \rightarrow X$. By that we mean that the transition functions of the bundle are in $W^{k,2}$.

Take now $g \in W^{k,2}(\mathbb{S}^1, X)$. We are going to construct a complex coordinate neighborhood of g in $W^{k,2}(\mathbb{S}^1, X)$. Consider a neighborhood U_g of the graph Γ_g of g in $\mathbb{S}^1 \times X$. For every $s \in \mathbb{S}^1$ consider the following neighborhood of $g(s)$ in X

$$U_g(s) := \{x \in X : (s, x) \in U_g\}, \quad (4.3)$$

and define $\varepsilon^s : U_g(s) \rightarrow U_g$ as

$$x \rightarrow (s, x). \quad (4.4)$$

Lemma 4.1. *There exists a $W^{k,2}$ -diffeomorphism G between a neighborhood U_g of Γ_g in $\mathbb{S}^1 \times X$ and a neighborhood V_g of the zero section of the induced bundle g^*TX such that*

- i) $\{G(s, g(s)) : s \in \mathbb{S}^1\}$ is the zero section of g^*TX ;
- ii) $G^s := G \circ \varepsilon^s$ maps $U_g(s)$ biholomorphically to a neighborhood of $0 \in T_{g(s)}X$.

Mapping G is constructed roughly as follows. For a chart (X_j, α_j) in X consider a chart (S_j, β_j) in \mathbb{S}^1 with $g(S_j) \subset X_j$ as before. Consider an appropriate neighborhood U_j of the graph $g|_{S_j}$ in $S_j \times X_j$ and define $G_j : U_j \rightarrow g^*TX_j$ as follows

$$G_j : (s, x) \rightarrow (s, (d\alpha_j^{-1})_{\alpha_j(g(s))}[\alpha_j(x) - \alpha_j(g(s))]). \quad (4.5)$$

In the right hand of (4.5) we use the natural identification of $T\alpha_j(X_j)$ with $\alpha_j(X_j) \times \mathbb{C}^n$ and treat $\alpha_j(x) - \alpha_j(g(s))$ as vector in $T_{\alpha_j(g(s))}\alpha_j(X) = \{\alpha_j(g(s))\} \times \mathbb{C}$. Then glue the maps G_j together to obtain the desired $G(s, x) = \sum_j \eta_j(s)G_j(s, x)$ using a \mathcal{C}^k -partition of unity subordinate to the covering $\{S_j\}$. For the more detailed proof of the lemma we refer to [A], see Lemma 3.1 there.

Let U_g be a neighborhood of Γ_g in $\mathbb{S}^1 \times X$ as in this lemma. Set

$$\mathcal{U}_g = \{h \in W^{k,2}(\mathbb{S}^1, X) : \Gamma_h \subset U_g\}. \quad (4.6)$$

For $h \in \mathcal{U}_g$ set $\Psi_g(h) := G(\cdot, h(\cdot)) \in W^{k,2}(\mathbb{S}^1, g^*TX)$. Ψ_g is a homeomorphism between \mathcal{U}_g and a neighborhood \mathcal{V}_g of the zero section in the space $W^{k,2}(\mathbb{S}^1, g^*TX)$, which consists from sections that are contained in V_g . We take (\mathcal{U}_g, Ψ_g) as a complex l^2 -coordinate chart in $W^{k,2}(\mathbb{S}^1, X)$. Let $h \in W^{k,2}(\mathbb{S}^1, X)$ be such that $\Gamma_h \subset U_g \cap U_{g'}$, i.e., $h \in \mathcal{U}_g \cap \mathcal{U}_{g'}$. For $s \in \mathbb{S}^1$ we have

$$\begin{aligned} \Psi_{g'}(h)(s) &= (\Psi_{g'} \circ \Psi_g^{-1}) \Psi_g(h)(s) = (G'^s \circ G^{s^{-1}}) (\Psi_g(h)(s)) = \\ &= (G'(s, \cdot) \circ G(s, \cdot)^{-1}) (\Psi_g(h)(s)). \end{aligned} \quad (4.7)$$

Due to the item (ii) of Lemma 4.1 for every s mapping $G'(s, \cdot) \circ G(s, \cdot)^{-1}$ is a biholomorphism between an appropriate open subsets of $g^*T_{g(s)}X$ and $g'^*T_{g'(s)}X$. Therefore $\Psi_{g'}(h)(s)$ holomorphically depends on $\Psi_g(h)(s)$, see [F] and [L1],[L2] for more details on this construction.

From this description of the complex structure on $W^{k,2}(\mathbb{S}^1, X)$ one can deduce, see [L1] and Lemma 3.2 in [A], the following:

Proposition 4.1. *A holomorphic map from a complex Hilbert manifold \mathcal{X} to $W^{k,2}(\mathbb{S}^1, X)$ can be represented by a mapping $f : \mathcal{X} \times \mathbb{S}^1 \rightarrow X$ such that:*

- i) *for every $s \in \mathbb{S}^1$ mapping $f(\cdot, s) : \mathcal{X} \rightarrow X$ is holomorphic;*
- ii) *for every $z \in \mathcal{X}$ one has $f(z, \cdot) \in W^{k,2}(\mathbb{S}^1, X)$ and the correspondence*

$$\mathcal{X} \ni z \rightarrow f(z, \cdot) \in W^{k,2}(\mathbb{S}^1, X)$$

is continuous with respect to the Sobolev topology on $W^{k,2}(\mathbb{S}^1, X)$ and the standard topology on \mathcal{X} . Mapping $f : \mathcal{X} \rightarrow W^{k,2}(\mathbb{S}^1, X)$ is actually $z \rightarrow f(z, \cdot)$.

Remark 4.1. a) Notice that if $X = \mathbb{C}^n$ then $G(s, x) = (s, x - g(s))$ because $TX \equiv X \times \mathbb{C}^n$ in this case. The same simple form has G if the image of g is contained in one chart $X_j \sim \tilde{X}_j \subset \mathbb{C}^n$. Therefore Ψ_g has the form $\Psi_g(h)(s) = (s, h(s) - g(s))$ in these cases.

b) And one more remark, let f be a holomorphic function on X_j and let \mathcal{X}_j be the open set of such $h \in W^{k,2}(\mathbb{S}^1, X)$ that $h(s_0) \in X_j$. Then $F(h) := f(h(s_0))$ is a holomorphic function on \mathcal{X}_j . Indeed, we can assume that X_j is a coordinate neighborhood of some $g(\mathbb{S}^1)$ and consider loops h close to g . Then using the coordinate chart (\mathcal{X}_j, G) as above we see that $F(h) = f((G^{s_0})^{-1}(h(s_0) - g(s_0)))$ in the notation of Lemma 4.1. This proves the holomorphicity of F .

c) If f is meromorphic write (locally) $f = p/q$ and set $F(g) = f(h(s_0)) = p(h(s_0))/q(h(s_0))$ in the case when $h(s_0) \notin I_f$. This will be a meromorphic function on $\mathcal{X}_j \setminus \{h \in \mathcal{X}_j : h(s_0) \in I_f\}$. The analytic set $\{h \in \mathcal{X}_j : h(s_0) \in I_f\}$ is of codimension ≥ 2 in \mathcal{X}_j and therefore F extends to a meromorphic function on \mathcal{X}_j . If f was meromorphic on the whole of X we obtain a meromorphic function F on the whole of $W^{k,2}(\mathbb{S}^1, X)$, in fact a family of meromorphic functions parametrized by $s_0 \in \mathbb{S}^1$.

d) Finally let us remark that $L\mathbb{P}^1$ is homogeneous since the loop group LG , where $G = PGL(2, \mathbb{C})$ obviously acts holomorphically and transitively on $L\mathbb{P}^1$.

Proposition 4.2. *Let X be a complex manifold such that $\mathcal{M}(X)$ separates points in X . Then $\mathcal{M}(W^{k,2}(\mathbb{S}^1, X))$ separates points in $W^{k,2}(\mathbb{S}^1, X)$.*

For if $g, h \in W^{k,2}(\mathbb{S}^1, X)$ are distinct then $g(s_0) \neq h(s_0)$ for some $s_0 \in \mathbb{S}^1$. By assumption there exists a meromorphic function on X which separates $g(s_0)$ and $h(s_0)$. The corresponding F constructed as in the remark above will separate g and h .

We finish this subsection with statement mentioned in the Introduction.

Proposition 4.3. *Let X be a finite dimensional compact complex manifold such that $\mathcal{M}(X)$ separates points on X . Then X admits a meromorphic injection to the complex projective space \mathbb{P}^N for some N .*

Proof. For any point $x_1 \in X$ we can find a non-constant meromorphic function f_1 which is holomorphic in a neighborhood of x_1 . Multiplying f_1 by a constant > 1 we find another function g_1 and a neighborhood U_1 of x_1 such that $f_1(U_1) \cap g_1(U_1) = \emptyset$. By compactness of X we find a finite cover U_1, \dots, U_d of X and meromorphic on X functions $f_1, g_1, \dots, f_d, g_d$ such that f_j and g_j are holomorphic on U_j and $f_j(U_j) \cap g_j(U_j) = \emptyset$ for every j . Set $U = \bigcup_{j=1}^d U_j \times U_j$. This is an open neighborhood of the diagonal \mathbb{D} in $X \times X$.

Further on we proceed as follows. For every $(y_1, z_1) \in X \times X \setminus U$ find a neighborhood V_1 of y and W_1 of z_1 and meromorphic function h_1 on X such that h_1 is holomorphic on V_1 and on W_1 and $h_1(V_1) \cap h_1(W_1) = \emptyset$. Using compactness of $X \times X \setminus U$ find a finite cover $\{V_k \times W_k\}_{k=1, \dots, l}$ of it by such pairs. Denote by F_1, \dots, F_N all the meromorphic functions $\{f_j, g_j, h_k : j = 1, \dots, d, k = 1, \dots, l\}$ thus obtained. Consider the meromorphic mapping $F : X \rightarrow \mathbb{P}^N$ defined by F_1, \dots, F_N . We claim that F is a meromorphic injection. Denote by I the union of indeterminacy sets of all F_1, \dots, F_N . This is an analytic subset of codimension two in X and F is holomorphic on $X \setminus I$. Take two points $p \neq q$ in $X \setminus I$. If they both belong to some U_j as above functions f_j and g_j will separate them. Otherwise the pair (p, q) belong to some $V_k \times W_k$ and then h_k will separate them. □

4.2. Meromorphic maps to the loop space. From now on we assume that $X = \mathbb{P}^1$, $k = 1$ and we are going to explain that the Definition 2 from the Introduction gives the natural notion of a meromorphic mapping in the case of $L\mathbb{P}^1 = W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$ as a target manifold. Let a meromorphic (in any sense) mapping $f : \mathcal{X} \rightarrow L\mathbb{P}^1$ be given. Then the least thing we know is that there should exist an analytic of codimension ≥ 2 set I in \mathcal{X} such that the restriction of f to $\mathcal{X} \setminus I$ is holomorphic. Denote this restriction still as $f : \mathcal{X} \setminus I \rightarrow L\mathbb{P}^1$ as well as its representation $f : (\mathcal{X} \setminus I) \times \mathbb{S}^1 \rightarrow \mathbb{P}^1$. Notice that the latter satisfies items (i) and (ii) of Proposition 4.1.

Theorem 4.1. *Let a holomorphic mapping $f : \mathcal{X} \setminus I \rightarrow L\mathbb{P}^1$ be given, where I is analytic in \mathcal{X} of codimension ≥ 2 . Then*

- i) *for every $s \in \mathbb{S}^1$ mapping $f(\cdot, s)$ is a meromorphic function which extends meromorphically on the whole of \mathcal{X} and its indeterminacy set $I_{f(\cdot, s)}$ is contained in I ;*
- ii) *the family of the indeterminacy sets $\{I_{f(\cdot, s)} : s \in \mathbb{S}^1\}$ is locally finite in \mathcal{X} and $I_f = \bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}$ is an analytic set in \mathcal{X} of pure codimension two;*
- iii) *mapping f holomorphically extends to $\mathcal{X} := \mathcal{X} \setminus I_f$.*

Proof. The proof will be achieved in several steps.

Step 1. Meromorphicity of $f(\cdot, s)$. Notice that for every $s \in \mathbb{S}^1$ mapping $f(\cdot, s) : \mathcal{X} \setminus I \rightarrow \mathbb{P}^1$ is in fact a meromorphic function without indeterminacy points and it extends to a meromorphic function on the whole of \mathcal{X} by Lemma 2.5. Denote by $I_{f(\cdot, s)}$ the indeterminacy set of the extended $f(\cdot, s)$. We have that $I_{f(\cdot, s)} \subset I$ because f is holomorphic on $\mathcal{X} \setminus I$. Since $I_{f(\cdot, s)}$ is the indeterminacy set of a meromorphic function and, therefore is locally an intersection of two relatively prime divisors it is of pure codimension two. Take some point $z_0 \in \bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)} \subset I_{z_0}^2$, where $I_{z_0}^2$ is the (necessarily finite) union of codimension

two components of I at z_0 as in (2.1). Due to the description of the structure of Hilbert analytic sets of finite definition in [Ra] one can find a neighborhood $\mathcal{U} = B^\infty \times \Delta^2$ of z_0 such that $I_{z_0}^2 \cap \mathcal{U}$ is a finite cover of B^∞ in these coordinates. Therefore $(\bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}) \cap \mathcal{U}$ can have only finitely many components. We obtain as a result that the union $\bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}$ is an analytic of pure codimension two subset of \mathcal{X} .

Items (i) and (ii) are proved and we turn to item (iii). Take some $z_0 \in \mathcal{X} \setminus \bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}$ and let \mathcal{U} be a neighborhood of z_0 not intersecting $\bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)}$. Then f extends to a mapping $f : \mathcal{U} \times \mathbb{S}^1 \rightarrow \mathbb{P}^1$. We need to prove its holomorphicity. Condition (i) of Proposition 4.1 for our f is obviously satisfied. Therefore all we need to prove is the condition (ii), i.e., continuity of the mapping $z \rightarrow f(z, \cdot)$ in Sobolev topology.

Step 2. Continuity in classical topology. We shall prove first that $f : \mathcal{U} \times \mathbb{S}^1 \rightarrow \mathbb{P}^1$ is continuous in the classical sense. We need to prove this at points $(z_0, s_0) \in \mathcal{U} \times \mathbb{S}^1$ with $z_0 \in I \setminus I_f$ only. Fix such point z_0 and take an open neighborhood \mathcal{U} of z_0 biholomorphic to $B^\infty \times \Delta^2$, where bidisk $\{0\} \times \Delta^2$ intersects I at z_0 transversely, i.e., z_0 is an isolated point of $(\{0\} \times \Delta^2) \cap I$. Natural coordinates in $B^\infty \times \Delta^2$ centered at $z_0 = (0, 0)$ denote as $z = (w, t)$. Note that f is holomorphic on $B^\infty \times \partial\Delta^2$, provided B^∞ was taken sufficiently small, and that $f(w, \cdot, s_0)|_{\Delta^2}$ is holomorphic over the whole of $\{w\} \times \Delta^2$ for every fixed $w \in B^\infty$. Due to Lemma 3.1 from [AZ] to prove the continuity of f at (z_0, s_0) it is sufficient to prove that for any sequences $s_n \rightarrow s_0$ in \mathbb{S}^1 , $w_n \rightarrow 0$ in B^∞ and any neighborhood \mathcal{V} of the graph Γ_0 of $f(0, \cdot, s_0)|_{\Delta^2}$ in $\mathcal{U} \times \mathbb{P}^1$ the graphs Γ_n of $f(w_n, \cdot, s_n)|_{\Delta^2}$ are contained in \mathcal{V} for n big enough. By the main result of [AZ], i.e., Theorem 2.1 there, we can assume that \mathcal{V} is 1-complete. That is it admits a strictly plurisubharmonic exhausting function ρ . Now it is easy to conclude from Sobolev imbedding (4.1) that for w_n and s_n close enough to 0 and s_0 respectively $f(w_n, \cdot, s_n)|_{\partial\Delta^2}$ is close to $f(0, \cdot, s_0)|_{\partial\Delta^2}$. And than by maximum principle for ρ to deduce that Γ_n stays in \mathcal{V} . The step is proved.

Step 3. Continuity in Sobolev topology. Now we need to prove that a continuous in classical sense mapping $f : \mathcal{U} \times \mathbb{S}^1 \rightarrow \mathbb{P}^1$ is continuous also in Sobolev topology. Using compactness of \mathbb{S}^1 cover it by arcs $S_j(\varepsilon) =]s_j - \varepsilon, s_j + \varepsilon[$, $j = 1, \dots, N$ centered at s_j and shrink a neighborhood \mathcal{U} of z_0 so that $f(\mathcal{U} \times S_j)$ is contained in a coordinate chart $(\mathcal{V}_j, \alpha_j)$ of \mathbb{P}^1 . Make sure that $f(\overline{\mathcal{U} \times S_j(\varepsilon_1)}) \subset \mathcal{V}_j$ still holds true for some $\varepsilon_1 > \varepsilon$. Consider the mapping $f_j := \rho_j \alpha_j \circ f|_{\mathcal{U} \times S_j(\varepsilon_1)}$, where ρ_j is a cut-off function supported in $S_j(\varepsilon_1)$ and equal to 1 on $S_j(\varepsilon)$. Take \mathcal{U} this time in the form $\Delta \times B^\infty$ and find a Hartogs figure $H_1^\infty(r) = \Delta \times B^\infty(r) \cup A_{1-r, 1} \times B^\infty \subset \Delta \times B^\infty$ not intersecting I . Let (z, t) be the natural coordinates in $\Delta \times B^\infty$.

Note that:

- a) for every $(z, t) \in H_1^\infty(r)$ mapping f_j is in $W^{1,2}(S_j(\varepsilon_1), \mathbb{C})$ and has compact support.
- b) Mapping $(z, t) \rightarrow f_j(z, t, \cdot) \in W^{1,2}(S_j(\varepsilon_1), \mathbb{C})$ is continuous in Sobolev topology.
- c) Moreover, for every $s \in S_j(\varepsilon_1)$ mapping $f_j(\cdot, s)$ is holomorphic on $\hat{H}_1^\infty(r) = \Delta \times B^\infty$.

The latter is because the multiplication by a smooth cutoff function preserves the Sobolev class and continuity in Sobolev topology and since it depends only on the space variable s it doesn't spoils the holomorphicity in (z, t) . Therefore f_j is a holomorphic mapping from $H_1^\infty(r)$ to a complex Hilbert space. Consequently it extends holomorphically to $\Delta \times B^\infty$. In particular we obtain that $f|_{\mathcal{U} \times S_j(\varepsilon)}$ is continuous in Sobolev topology. By the definition of the Sobolev topology, see (4.2) we obtain the desired continuity of f . Lemma is proved.

□

Remark 4.2. As the result we reach the understanding that the Definition 2 from the Introduction is the natural definition of meromorphicity for mappings with values in $L\mathbb{P}^1$. Indeed, what we proved in Proposition 4.1 is that a holomorphic map $f : \mathcal{X} \setminus I \rightarrow L\mathbb{P}^1$, where I is Hilbert analytic of codimension ≥ 2 , extends to a meromorphic map from \mathcal{X} to $L\mathbb{P}^1$ in the sense of Definition 2.

4.3. Non-existence of an imbedding of the loop space to the projective space. We start this subsection with the following example.

Example 4.1. Let $\gamma(s) = (\gamma_1(s), \gamma_2(s))$ be loop in \mathbb{C}^2 of class $W^{1,2}$. Denote by Γ the image of γ . Consider the mapping $f : \mathbb{C}^2 \setminus \Gamma \rightarrow W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$ defined as follows

$$f(z_1, z_2, s) = [z_1 - \gamma_1(s) : z_2 - \gamma_2(s)]. \quad (4.8)$$

For every fixed $s \in \mathbb{S}^1$ mapping $f(\cdot, s)$ is holomorphic on $\mathbb{C}^2 \setminus \{\gamma(s)\}$ and moreover $\gamma(s)$ is the only indeterminacy point of $f(\cdot, s)$. Furthermore, for every $z \in \mathbb{C}^2 \setminus \Gamma$ we have that $f(z, \cdot) \in W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$ and it continuously depends on z . This means that f is a holomorphic mapping from $\mathbb{C}^2 \setminus \Gamma$ to $W^{1,2}(\mathbb{S}^1, \mathbb{P}^1)$. Notice that for any fixed s mapping $f(\cdot, s) : \mathbb{C}^2 \setminus \{\gamma(s)\} \rightarrow \mathbb{P}^1$ extends meromorphically onto the whole \mathbb{C}^2 having $\gamma(s)$ as its single indeterminacy point and therefore $\bigcup_{s \in \mathbb{S}^1} I_{f(\cdot, s)} = \Gamma$, i.e., the family $\{I_{f(\cdot, s)}\}$ is not locally finite and henceforth is not an analytic set in \mathbb{C}^2 . We see that Γ is an *essential singularity* of f in the sense that f does not extend meromorphically to a neighborhood of any point of Γ . For would f extend meromorphically to a neighborhood V of some $\gamma(s_0) \in \Gamma$ it should be holomorphic on V minus a discrete set. But f is obviously not holomorphic at any point of $\Gamma \cap V$.

Remark 4.3. This example shows that the Hartogs type extension theorem is not valid for meromorphic values in $L\mathbb{P}^1$. Taking $\Gamma \subset \mathbb{C} \subset \mathbb{C}^2$ we obtain a counterexample also for the Thullen type extension.

Proof of Theorem 1. Suppose to the contrary that $g : L\mathbb{P}^1 \rightarrow \mathbb{P}(l^2)$ is such a map and let γ be such as in Example 4.1. Take γ in addition such that it is of class $W^{2,2} \subset \mathcal{C}^{1, \frac{1}{2}}$ and that its image Γ is *not contained* in any complex curve in \mathbb{C}^2 . Consider a holomorphic map $f : \mathbb{C}^2 \setminus \Gamma \rightarrow L\mathbb{P}^1$ defined as in (4.8). Using the homogeneity of $L\mathbb{P}^1$ compose f with an automorphism of $L\mathbb{P}^1$ in order to map some point by a resulting map to the regular point of g . Denote this resulting map still as f . Now the composition $h := g \circ f$ is well defined and by Corollary 1 $h : \mathbb{C}^2 \setminus \Gamma \rightarrow \mathbb{P}(l^2)$ extends meromorphically to \mathbb{C}^2 . Denote this extension still as h . Take $a \in \Gamma \setminus I_h$ and consider a neighborhood V of a which do not contain points from I_h , i.e., the restriction $h|_V : V \rightarrow \mathbb{P}(l^2)$ is holomorphic.

Lemma 4.2. *The differential of mapping $f : \mathbb{C}^2 \setminus \Gamma \rightarrow L\mathbb{P}^1$ is very generically injective, i.e., df_z is injective for z outside of a complex curve.*

Proof. We need to prove this for the original f as in (4.8) since a composition with an automorphism doesn't affect the injectivity. Take $z^0 = (z_1^0, z_2^0) \in \mathbb{C} \setminus \Gamma$ such that $z_j^0 - \gamma_j(s)$ never vanishes. Then

$$f(z, s) = [z_1 - \gamma_1(s) : z_2 - \gamma_2(s)]$$

takes its values in $U_0 = \{[w] \in \mathbb{P}^1 : w_0 \neq 0\}$ for z in some neighborhood $U \ni z^0$. I.e.,

$$f(z, s) = \frac{z_2 - \gamma_2(s)}{z_1 - \gamma_1(s)}$$

represents a holomorphic map $f : U \rightarrow W^{1,2}(\mathbb{S}^1, \mathbb{C})$. Now for $z \in U$ and $v \in \mathbb{C}^2$ one has

$$df_z[v] = \frac{\partial f(z, s)}{\partial z_1} v_1 + \frac{\partial f(z, s)}{\partial z_2} v_2 = \left(s, \frac{z_2 - \gamma_2(s)}{[z_1 - \gamma_1(s)]^2} v_1 + \frac{v_2}{z_1 - \gamma_1(s)} \right). \quad (4.9)$$

Here the right hand side of (4.9) should be understood as a section of the pulled-back tangent bundle $f^*(z, \cdot)T\mathbb{P}^1 \cong \mathbb{S}^1 \times \mathbb{C}$. To prove the injectivity of df_z assume that $df_z[v] = df_z[w]$ for some $v \neq w$. Then

$$\frac{z_2 - \gamma_2(s)}{[z_1 - \gamma_1(s)]^2} v_1 + \frac{v_2}{z_1 - \gamma_1(s)} = \frac{z_2 - \gamma_2(s)}{[z_1 - \gamma_1(s)]^2} w_1 + \frac{w_2}{z_1 - \gamma_1(s)}$$

for all s . And this means that

$$\frac{z_2 - \gamma_2(s)}{z_1 - \gamma_1(s)} (v_1 - w_1) + v_2 - w_2 = 0.$$

Since $v \neq w$ this implies that $\frac{z_2 - \gamma_2(s)}{z_1 - \gamma_1(s)} = \text{const}$ and therefore $\gamma_2(s) - c_1 \gamma_1(s) = c_2$ for some constants c_1, c_2 and all s . I.e., Γ is contained in the complex line $C = \{z_2 c_1 - z_1 = c_2\}$. Contradiction.

We proved that df_z is injective for z in a non-empty open U and therefore it can degenerate for z in a complex curve $C \subset \mathbb{C}^2 \setminus \Gamma$ at most. □

Therefore the differential of $h = g \circ f$ is very generically injective on $V \setminus \Gamma$ and therefore on V . Since Γ is not locally contained in a complex curve we see that the locus of degeneration of dh intersects Γ by a finite number of points at most. We can move our point a together with V to ensure that not only $h|_V$ is holomorphic but also its differential is injective at all points of V .

Furthermore, $dh_z[\mathbb{C}^2]$ is closed in $T_{h(z)}\mathbb{P}(l^2)$ for all $z \in V$ since it is a two-dimensional subspace of $T_{h(z)}\mathbb{P}(l^2) \cong l^2$. We find an affine chart \mathcal{U} containing $h(V)$. For this one might need to shrink V again. Now we compose h with the orthogonal projection π of l^2 to $dh_a[\mathbb{C}^2] \cong \mathbb{C}^2$. This composition is biholomorphic in a neighborhood (again call it V) of a . Therefore $h(V)$ is a graph over $\pi(h(V))$ in $\mathbb{C}^2 \times l^2$. This proves that $h|_V : V \rightarrow h(V)$ is a biholomorphism to a local submanifold $h(V)$ of $g(L\mathbb{P}^1)$. g^{-1} is well defined on $h(V)$ and therefore $g^{-1} \circ h$ is holomorphic on V . But it coincides with f on $V \setminus \Gamma$. We got a holomorphic extension of f from $V \setminus \Gamma$ to V . Composing this extension with the inverse to the automorphism of $L\mathbb{P}^1$ taken at the beginning of the proof we get an extension of the original f , contradiction. Theorem is proved. □

REFERENCES

- [A] ANAKKAR M.: *Complex manifolds of Sobolev mappings and a Hartogs-type theorem in loop spaces*. Complex Variables and Elliptic Equations. **68**, no. 12, 2016–2033 (2023).
- [AI] ANAKKAR M., IVASHKOVICH S.: *Loop Spaces as Hilbert-Hartogs Manifolds*. Arch. Math., **115**, no. 4, 445–456 (2020).
- [AZ] ANAKKAR M., ZAGORODNYUK A.: *On Hilbert-Hartogs manifolds*. Complex Variables and Elliptic Equations, **65** no. 12, 2071–2085 (2020).
- [F] FORSTNERIC F.: *Manifolds of holomorphic mappings from strongly pseudoconvex domains*. Asian. J. Math. **11**, no. 1, 113-126 (2007).
- [L1] LEMPERT L.: *Loop spaces as complex manifolds*. J. Diff. Geom. **38**, 519-543 (1993).

- [L2] LEMPERT L.: *Analytic continuation in mapping spaces*. Pure Appl. Math. Q. **6**, no: 4, 1051-1080 (2010).
- [MZ] MILLSON J.J., ZOMBRO B. *A Kähler structure on the moduli space of isometric maps of a circle into Euclidean space*. Invent. Math., 123, 35-59.(1996)
- [Mu] MUJICA J.: *Complex Analysis in Banach Spaces*. Math. Studies **120**, North-Holland (1985).
- [Ra] RAMIS J.-P. *Sous-ensembles analytiques d'une variété banachique complexe*. Ergebnisse der Mathematik un ihrer Grenzgebiete, B. 53, Springer Verlag (1970).
- [Z] ZHANG N.: *The Picard group of the loop space of the Riemann sphere*. arXiv: math/0602667v4 (2022).

*UNIVERSITÉ DE LILLE-1, UFR DE MATHÉMATIQUES, 59655 VILLENEUVE D'ASCQ, FRANCE.

Email address: `serge.ivashkovych@univ-lille.fr`

Email address: `anakkarmohammed@yahoo.fr`