

Holographic approach to anomalous transport in a massive $U(1)$ gauge theory

Nishal Rai^{1,2} and Eugenio Megías³

Abstract In this study, we explore a massive $U(1)$ gauge holographic model with pure gauge and mixed gauge-gravitational Chern-Simons terms. By considering the full backreaction of the gauge field on the metric tensor, we explore the vortical and energy transport sectors. Our findings for the chiral vortical conductivity, σ_V , and the chiral magnetic/vortical conductivity of energy current show that $\sigma_B^\varepsilon = \sigma_V^\varepsilon$. Notably, we highlight a contribution to σ_V induced by the mixed term in the massive theory, which is absent in the massless case.

1 Introduction

AdS/CFT correspondence [1, 2] states that, in the low energy limit, the large- N_c , $\mathcal{N} = 4$ super Yang-Mills field theory in four-dimensional space is equivalent to the type IIB string theory in $AdS_5 \times S^5$ space. It has become a very powerful theoretical tool for studying strongly coupled systems. It has a wide application in different fields of physics, such as condensed matter physics, hydrodynamics and QCD [3–22] to name a few. Our study will be focused on the application of the fluid/gravity correspondence in the context of hydrodynamics, in particular the study of the anomalous transport in this kind of systems.

Quantum chiral anomalies, arising in relativistic field theories of chiral fermions beyond perturbation theory, play a crucial role in relativistic hydrodynamics [23–

Nishal Rai

¹Center for Astrophysics Gravitation and Cosmology (CAGC), SRM University Sikkim, Upper Tadong, Sikkim, India

²Department of Physics, SRM University Sikkim, Upper Tadong, Sikkim, India.
e-mail: nishalrai10@gmail.com

Eugenio Megías

³Departamento de Física Atómica, Molecular y Nuclear and Instituto Carlos I de Física Teórica y Computacional, Universidad de Granada, Avenida de Fuente Nueva s/n, E-18071 Granada, Spain

26]. Since the 1980s, anomaly-induced transport mechanisms have been widely studied [27]. Key effects include the chiral magnetic effect, where a charge current is induced parallel to an external magnetic field [28], and the chiral vortical effect, where a vortex in a charged fluid generates a current [29–32]. These effects may be detectable in heavy ion collisions at RHIC and LHC [33] and can lead to anomalous transport properties in materials like Weyl semimetals [34, 35]. In this work, we explore and investigate the chiral vortical effects in a massive $U(1)$ gauge holographic model. To achieve this, we have accounted for the full backreaction of the gauge field on the metric tensor and included a mixed gauge-gravitational Chern-Simons term as well. To evaluate the transport coefficients we will be using the Kubo formalism where the anomalous conductivities are given by the following relations

$$\begin{aligned}\sigma_V &= \lim_{k_c \rightarrow 0} \frac{i}{2k_c} \sum_{a,b} \epsilon_{abc} \langle J^a T^{0b} \rangle |_{\omega=0}, \\ \sigma_V^\varepsilon &= \lim_{k_c \rightarrow 0} \frac{i}{2k_c} \sum_{a,b} \epsilon_{abc} \langle T^{0a} T^{0b} \rangle |_{\omega=0},\end{aligned}\tag{1}$$

where σ_V is the chiral vortical conductivity, and σ_V^ε the chiral vortical conductivity of energy current, respectively. The chiral magnetic conductivities for charge, σ_B , and energy, σ_B^ε , current are given by

$$\begin{aligned}\sigma_B &= \lim_{k_c \rightarrow 0} \frac{i}{2k_c} \sum_{a,b} \epsilon_{abc} \langle J^a J^b \rangle |_{\omega=0}, \\ \sigma_B^\varepsilon &= \lim_{k_c \rightarrow 0} \frac{i}{2k_c} \sum_{a,b} \epsilon_{abc} \langle T^{0a} J^b \rangle |_{\omega=0}.\end{aligned}\tag{2}$$

These correlators can be evaluated using the AdS/CFT dictionary [36–38].

2 Holographic model

The holographic action for a massive $U(1)$ gauge boson with pure gauge and a mixed gauge-gravitational Chern-Simons term is given by [7, 36, 39]

$$\begin{aligned}S &= \frac{1}{16\pi G} \int d^5x \sqrt{-g} \left[R + 2\Lambda - \frac{1}{4} F_{MN} F^{MN} \right. \\ &\quad - \frac{m^2}{2} (A_M - \partial_M \theta) (A^M - \partial^M \theta) \\ &\quad \left. + \epsilon^{MNPQR} (A_M - \partial_N \theta) \left(\frac{\kappa}{3} F_{NP} F_{QR} + \lambda R^A{}_{BNP} R^B{}_{AQR} \right) \right] \\ &\quad + S_{\text{GH}} + S_{\text{CSK}},\end{aligned}\tag{3}$$

where S_{GH} and S_{CSK} are the Gibbons-Hawking boundary term, and a boundary term induced by the mixed gauge-gravitational anomaly, respectively. θ is a field which ensures gauge invariance (up to gauge anomalies), and thus the mass term enters in a consistent way. As it mentioned in [40–42], the Stückelberg term arises as the holographic realization of dynamical anomalies. For convenience we will define a new field $B_M \equiv A_M - \partial_M \theta$, such that θ does not appear explicitly in the equations of motion. One may check that the action still remains invariant under gauge transformations. From now onwards we will be working with the field B_μ instead of A_μ .

2.1 Background solution

The ansatz for the background metric and fields is chosen as follows in Fefferman-Graham coordinates [43, 44],

$$ds^2 = -\frac{\ell^2}{\rho} g_{\tau\tau}(\rho) d\tau^2 + \frac{\ell^2}{\rho} g_{xx}(\rho) d\vec{x}^2 + \frac{\ell^2}{4\rho^2} d\rho^2, \quad (4)$$

$$B_\mu dx^\mu = B_t(\rho) dt,$$

where the boundary lies at $\rho = 0$ and the horizon at ρ_h , with $g_{\tau\tau}(\rho_h) = 0$. Using this ansatz the equations of motion for the background are given by

$$g''_{xx}(\rho) - \frac{g'_{xx}(\rho)}{\rho} + \frac{1}{6\ell^2 \rho} \frac{g_{xx}(\rho)}{g_{\tau\tau}(\rho)} \left(\frac{m^2 \ell^2}{4} B_t(\rho)^2 + \rho^2 B'_t(\rho)^2 \right) = 0,$$

$$g'_{\tau\tau}(\rho) \left(1 - \rho \frac{g'_{xx}(\rho)}{g_{xx}(\rho)} \right) + g_{\tau\tau}(\rho) \frac{g'_{xx}(\rho)}{g_{xx}(\rho)} \left(3 - \rho \frac{g'_{xx}(\rho)}{g_{xx}(\rho)} \right) - \frac{1}{3} \frac{\rho^2}{\ell^2} B'_t(\rho)^2 + \frac{1}{12} m^2 B_t(\rho)^2 = 0,$$

$$B''_t(\rho) + \frac{1}{2} \left(3 \frac{g'_{xx}(\rho)}{g_{xx}(\rho)} - \frac{g'_{\tau\tau}(\rho)}{g_{\tau\tau}(\rho)} \right) B'_t(\rho) - \frac{\ell^2 m^2}{4\rho^2} B_t(\rho) = 0. \quad (5)$$

We will solve numerically the above equations of motion (5) subjected to the following boundary conditions

$$B_t(\rho_h) = 0, \quad \lim_{\rho \rightarrow 0} \left(\rho^{\Delta/2} B_t(\rho) \right) = \mu_5, \quad (6)$$

where μ_5 corresponds to the source of the gauge field and Δ being the anomalous dimension of the dual current [39] related to the mass of the gauge field through the following relation $m^2 \ell^2 = \Delta(\Delta + 2)$. In addition to this, we demand the solution to the metric tensor to be regular at the horizon.

It is possible to find an analytic solution of the background for the case with $\mu_5 = 0$. This is given by




Fig. 1: (color) Dependence of the background metric and gauge field with ρ . We display the results for $g_{xx}(\rho)$ (blue), $g_{\tau\tau}(\rho)$ (orange), and $\rho^{\Delta/2}B_t(\rho)$ (green). Left panel: We have considered $\mu_5 = 0$, with solid line being the analytic solution and the dots the numerical one. Right panel: We have chosen $\Delta = 0.1$ and $\mu_5 = 0.3$.

$$g_{\tau\tau}(\rho) = \frac{1}{\rho_h^2} \frac{(\rho_h^2 - \rho^2)^2}{\rho_h^2 + \rho^2}, \quad g_{xx}(\rho) = 1 + \frac{\rho^2}{\rho_h^2}, \quad B_t(\rho) = 0. \quad (7)$$

We display in Fig. 1 (left) the numerical solution of the equations of motion for $\mu_5 = 0$ (dots), and compare it with the corresponding analytical solution (solid lines). We display also in the right panel the numerical solution for $\mu_5 = 0.3$.

3 Anomalous conductivity

To compute these correlators, we will follow the general procedure of the perturbation of the fields on top of the background, and later use the AdS/CFT dictionary to compute the correlators.

3.1 Fluctuations

Within this approach, we perturb the fields on top of the numerical solution for the background displayed in Fig. 1. We will study the linear response of the fluctuations by splitting the metric and gauge field into background and linear perturbation components, i.e.

$$g_{MN} = g_{MN}^{(0)} + \epsilon h_{MN}, \quad B_M = B_M^{(0)} + \epsilon b_M, \quad (8)$$

where the terms with $^{(0)}$ correspond to the background metric/field, and the terms with ϵ the fluctuations of this metric/field. Then, we will follow the general procedure of Fourier mode decomposition [45]

$$h_{MN}(\rho, x^\mu) = \int \frac{d^d k}{(2\pi)^d} h_{MN}(\rho) e^{-i\omega t + i\vec{k} \cdot \vec{x}}, \quad (9)$$

$$b_M(\rho, x^\mu) = \int \frac{d^d k}{(2\pi)^d} b_M(\rho) e^{-i\omega t + i\vec{k} \cdot \vec{x}}. \quad (10)$$

Without loss of generality, we consider perturbations with frequency ω and momentum k in the z -direction. Since we aim to compute correlators at zero frequency, we set the frequency-dependent parts to zero in the equations and solve the system up to first order in k . In this limit, the fields h_z^i decouple from the system and become constant. Thus, we can write the system of differential equations for the shear sector as follows

$$\begin{aligned} & b_i''(\rho) + \frac{1}{2} \left(\frac{g'_{xx}(\rho)}{g_{xx}(\rho)} + \frac{g'_{\tau\tau}(\rho)}{g_{\tau\tau}(\rho)} \right) b_i'(\rho) - \frac{\Delta(\Delta+2)}{4\rho^2} b_i(\rho) \\ & + \left(\frac{4ik\epsilon_{ij}b_j(\rho)}{\sqrt{g_{xx}(\rho)g_{\tau\tau}(\rho)}} + \frac{g_{xx}(\rho)h_t^i(\rho)}{g_{\tau\tau}(\rho)} \right) B_t'(\rho) + i\lambda k\epsilon_{ij}h_t^{j'}(\rho)\Omega(\rho) = 0, \\ & h_t^{i''}(\rho) - \left(\frac{g'_{\tau\tau}(\rho)}{2g_{\tau\tau}(\rho)} - \frac{5g'_{xx}(\rho)}{2g_{xx}(\rho)} + \frac{1}{\rho} \right) h_t^{i'}(\rho) + \frac{\rho B_t'(\rho)}{g_{xx}(\rho)} b_i'(\rho) \\ & + \frac{\Delta(\Delta+2)B_t(\rho)}{4\rho g_{xx}(\rho)} b_i(\rho) + i\lambda k\epsilon_{ij}\Phi_j(\rho) = 0, \end{aligned} \quad (11)$$

$$(12)$$

where $i, j = x, y$. The explicit expressions of the functions $\Omega(\rho)$ and $\Phi_j(\rho)$ are given in Appendix A of [7].

The asymptotic behaviour of the fields up to the first subleading term are given by

$$b_i(\rho) = b_i^{(0)} \rho^{-\frac{\Delta}{2}} + b_i^{(1)} \rho^{\frac{\Delta}{2}+1} + \dots, \quad (13)$$

$$h_t^i(\rho) = h_t^{i(0)} + h_t^{i(1)} \rho^2 + \dots, \quad (14)$$

where the leading-order terms $b_i^{(0)}$ and $h_t^{i(0)}$ play the role of sources.

3.1.1 Conductivities

From the holographic description of the correlation functions, one can evaluate the one-point functions as

$$\langle J_a \rangle = \frac{\delta S_{\text{ren}}}{\delta b_a^{(0)}} = -\frac{2}{16\pi G} (\Delta+1) b_a^{(1)}, \quad \langle T_{0a} \rangle = \frac{\delta S_{\text{ren}}}{\delta h_t^a(0)} = \frac{1}{16\pi G} \left(2h_t^{a(0)} + h_t^{a(1)} \right), \quad (15)$$

with $(a = x, y)$. $S_{\text{ren}} = S + S_{\text{ct}}$ is the renormalized action, with S the action given in Eq. (3) and S_{ct} the counterterm. The two-point functions can be obtained by taking the variation of one-point function with respect to the corresponding source term and the conductivities are evaluated using the Kubo formulae given in (1) and (2),

i.e.

$$\sigma_V = \lim_{k \rightarrow 0} \frac{1}{k} \text{Im} \langle J_x T_{0y} \rangle, \quad \sigma_V^\varepsilon = \lim_{k \rightarrow 0} \frac{1}{k} \text{Im} \langle T_{0x} T_{0y} \rangle, \quad (16)$$

$$\sigma_B = \lim_{k \rightarrow 0} \frac{1}{k} \text{Im} \langle J_x J_y \rangle, \quad \sigma_B^\varepsilon = \lim_{k \rightarrow 0} \frac{1}{k} \text{Im} \langle T_{0x} J_y \rangle. \quad (17)$$

A comparison of the variation of the action under axial gauge transformations with the consistent form of the anomaly for chiral fermions [23] allows us to set the value of the anomaly coefficients to

$$\kappa = -\frac{G}{2\pi}, \quad \lambda = -\frac{G}{48\pi}. \quad (18)$$

In this regard we will be setting $G = 2\pi$, so that $\kappa = -1$ and $\lambda = -1/24$. In Fig. 2 we




Fig. 2: Upper panels: Plot of σ_V (left) and σ_V^ε (right) vs Δ . Lower panels: Plot of σ_B (left) and σ_B^ε (right) vs Δ . We have considered $\mu_5 = 0.15$ in all the panels. We have divided the conductivities by $|\kappa|$ to make them dimensionless.

have plotted the anomalous conductivities vs Δ (a parameter related to the mass of the gauge field). From this figure, we observe that the chiral vortical conductivity and the chiral magnetic conductivity for energy current are the same even at finite mass, i.e., $\sigma_V = \sigma_B^\varepsilon$, and both increase with Δ . Additionally, we see that the chiral vortical conductivity of energy current, σ_V^ε , decreases with Δ , but the rate decreases rapidly. Conversely, in the case of the chiral magnetic conductivity, σ_B , it increases with Δ as shown in Fig. 2. In the limit of vanishing mass, our results lead to $\frac{\sigma_B}{\mu_5 |\kappa|} \simeq 16/3$,

which exactly coincides with the results in [36, 39] where α has been set to μ_5 in both references ¹.

4 Discussion

In this work we have studied the mass dependence of the anomalous conductivities in a holographic model for a massive $U(1)$ gauge field, incorporating both pure gauge and mixed gauge-gravitational anomaly terms. In this model, the vortical sector was also accessible, and the conductivities corresponding to this sector were also studied. We observed that the chiral vortical conductivity, σ_V , and the chiral magnetic conductivity for energy current, σ_B^E , are equal and increase with Δ . An intriguing finding is the presence of contributions to σ_B induced by the mixed gauge-gravitational anomaly term in the massive theory, which were absent in the massless case. Furthermore, the conductivities σ_B , σ_V , and σ_B^E all increase with Δ , while the chiral vortical conductivity of energy current, σ_V^E , decreases with Δ . We have explicitly verified that all our numerical results for the conductivities at finite mass converge to the known results at zero mass as $\Delta \rightarrow 0$.

Acknowledgments

N.R. thanks the International Centre for Theoretical Sciences (ICTS) for the support through the program - Field Theory and Turbulence (ICTS/FTT2023/12). The works of N.R. and E.M. are supported by the project PID2020-114767GB-I00 and by the Ramón y Cajal Program under Grant RYC-2016-20678 funded by MCIN/AEI/10.13039/501100011033 and by “FSE Investing in your future”, as well as by the FEDER/Junta de Andalucía-Consejería de Economía y Conocimiento 2014–2020 Operational Program under Grant A-FQM-178-UGR18. The work of E.M. is also supported by Junta de Andalucía under Grant FQM-225, and by the “Prórrogas de Contratos Ramón y Cajal” Program of the University of Granada.

References

1. J. M. Maldacena, Int. J. Theor. Phys. **38**, 1113 (1999) [Adv. Theor. Math. Phys. **2**, 231 (1998)] doi:10.1023/A:1026654312961 [hep-th/9711200].
2. E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [hep-th/9802150].
3. S. Grieninger and S. Morales-Tejera, Phys. Rev. D **108** (2023) no.12, 126010 doi:10.1103/PhysRevD.108.126010 [arXiv:2308.14829 [hep-ph]].

¹ α corresponds to the asymptotic value of the gauge field A_t for $\rho \rightarrow 0$. In our case, we assume $\alpha = \mu_5$ for $\Delta = 0$.

4. M. Baggioli, Y. Bu and V. Ziogas, *JHEP* **09** (2023), 019 doi:10.1007/JHEP09(2023)019 [arXiv:2304.14173 [hep-th]].
5. S. Morales-Tejera, V. E. Ambrus and M. N. Chernodub, [arXiv:2403.19755 [hep-th]].
6. K. Landsteiner, S. Morales-Tejera and P. Saura-Bastida, *Phys. Rev. D* **107** (2023) no.12, 125003 doi:10.1103/PhysRevD.107.125003 [arXiv:2212.14088 [hep-th]].
7. N. Rai and E. Megías, *JHEP* **06** (2023), 215 doi:10.1007/JHEP06(2023)215 [arXiv:2301.00361 [hep-th]].
8. S. Mukhopadhyay and N. Rai, *JHEP* **09** (2021), 160 doi:10.1007/JHEP09(2021)160 [arXiv:2008.00432 [hep-th]].
9. N. Rai, *Eur. Phys. J. C* **79** (2019) no.11, 972 doi:10.1140/epjc/s10052-019-7469-x
10. S. Mukhopadhyay and N. Rai, *Phys. Rev. D* **96** (2017) no.6, 066001 doi:10.1103/PhysRevD.96.066001
11. S. Mukhopadhyay and N. Rai, *Phys. Rev. D* **96** (2017) no.2, 026005 doi:10.1103/PhysRevD.96.026005
12. M. Ammon, D. Arean, M. Baggioli, S. Gray and S. Griener, *JHEP* **03** (2022), 015 doi:10.1007/JHEP03(2022)015 [arXiv:2111.10305 [hep-th]].
13. S. Griener and S. Morales-Tejera, *EPJ Web Conf.* **258** (2022), 10007 doi:10.1051/epjconf/202225810007 [arXiv:2111.14116 [hep-ph]].
14. Y. Ahn, M. Baggioli, Y. Liu and X. M. Wu, *JHEP* **03** (2024), 124 doi:10.1007/JHEP03(2024)124 [arXiv:2401.07772 [hep-th]].
15. H. Chu, X. Ji and Y. W. Sun, *JHEP* **05** (2024), 166 doi:10.1007/JHEP05(2024)166 [arXiv:2403.02669 [hep-th]].
16. K. Landsteiner and Y. Liu, *Phys. Lett. B* **753** (2016), 453-457 doi:10.1016/j.physletb.2015.12.052 [arXiv:1505.04772 [hep-th]].
17. K. Landsteiner and L. Melgar, *JHEP* **10** (2012), 131 doi:10.1007/JHEP10(2012)131 [arXiv:1206.4440 [hep-th]].
18. E. Megías and F. Pena-Benitez, *JHEP* **05**, 115 (2013) doi:10.1007/JHEP05(2013)115 [arXiv:1304.5529 [hep-th]].
19. G. M. Newman and D. T. Son, *Phys. Rev. D* **73** (2006), 045006 doi:10.1103/PhysRevD.73.045006 [arXiv:hep-ph/0510049 [hep-ph]].
20. G. M. Newman, *JHEP* **01** (2006), 158 doi:10.1088/1126-6708/2006/01/158 [arXiv:hep-ph/0511236 [hep-ph]].
21. E. Megías, *EPJ Web Conf.* **164** (2017), 08001 doi:10.1051/epjconf/201716408001 [arXiv:1701.00087 [hep-th]].
22. K. Landsteiner, E. Megías and F. Pena-Benitez, *Phys. Rev. D* **90**, no.6, 065026 (2014) doi:10.1103/PhysRevD.90.065026 [arXiv:1312.1204 [hep-ph]].
23. R. A. Bertlmann, Oxford, UK: Clarendon (1996) 566 p. (International series of monographs on physics: 91).
24. F. Bastianelli and P. van Nieuwenhuizen, Cambridge, UK: Univ. Pr. (2006) 379p.
25. K. Fujikawa and H. Suzuki, Oxford, UK: Clarendon (2004) 284p.
26. K. Landsteiner, E. Megías and F. Pena-Benitez, *Lect. Notes Phys.* **871** (2013), 433-468 doi:10.1007/978-3-642-37305-3_17 [arXiv:1207.5808 [hep-th]].
27. A. Vilenkin, *Phys. Rev. D* **20** (1979) 1807. A. Vilenkin, *Phys. Rev. D* **21** (1980) 2260. A. Vilenkin, *Phys. Rev. D* **22** (1980) 3067. A. Vilenkin, *Phys. Rev. D* **22** (1980) 3080.
28. K. Fukushima, D. E. Kharzeev and H. J. Warringa, *Phys. Rev. D* **78** (2008), 074033 doi:10.1103/PhysRevD.78.074033 [arXiv:0808.3382 [hep-ph]].
29. D. Kharzeev and A. Zhitnitsky, *Nucl. Phys. A* **797** (2007), 67-79 doi:10.1016/j.nuclphysa.2007.10.001 [arXiv:0706.1026 [hep-ph]].
30. N. Banerjee, J. Bhattacharya, S. Bhattacharyya, S. Dutta, R. Loganayagam and P. Surowka, *JHEP* **01** (2011), 094 doi:10.1007/JHEP01(2011)094 [arXiv:0809.2596 [hep-th]].
31. D. T. Son and P. Surowka, *Phys. Rev. Lett.* **103** (2009), 191601 doi:10.1103/PhysRevLett.103.191601 [arXiv:0906.5044 [hep-th]].
32. K. Landsteiner, E. Megías and F. Pena-Benitez, *Phys. Rev. Lett.* **107**, 021601 (2011) doi:10.1103/PhysRevLett.107.021601 [arXiv:1103.5006 [hep-ph]].

33. B. I. Abelev *et al.* [STAR], Phys. Rev. Lett. **103** (2009), 251601 doi:10.1103/PhysRevLett.103.251601 [arXiv:0909.1739 [nucl-ex]].
34. G. Basar, D. E. Kharzeev and H. U. Yee, Phys. Rev. B **89**, no.3, 035142 (2014) doi:10.1103/PhysRevB.89.035142 [arXiv:1305.6338 [hep-th]].
35. K. Landsteiner, Phys. Rev. B **89**, no.7, 075124 (2014) doi:10.1103/PhysRevB.89.075124 [arXiv:1306.4932 [hep-th]].
36. K. Landsteiner, E. Megías, L. Melgar and F. Pena-Benitez, JHEP **09** (2011), 121 doi:10.1007/JHEP09(2011)121 [arXiv:1107.0368 [hep-th]].
37. D. T. Son and A. O. Starinets, JHEP **09** (2002), 042 doi:10.1088/1126-6708/2002/09/042 [arXiv:hep-th/0205051 [hep-th]].
38. C. P. Herzog and D. T. Son, JHEP **03** (2003), 046 doi:10.1088/1126-6708/2003/03/046 [arXiv:hep-th/0212072 [hep-th]].
39. A. Jimenez-Alba, K. Landsteiner and L. Melgar, Phys. Rev. D **90** (2014), 126004 doi:10.1103/PhysRevD.90.126004 [arXiv:1407.8162 [hep-th]].
40. I. R. Klebanov, P. Ouyang and E. Witten, Phys. Rev. D **65** (2002), 105007 doi:10.1103/PhysRevD.65.105007 [arXiv:hep-th/0202056 [hep-th]].
41. U. Gürsoy and A. Jansen, JHEP **10** (2014), 092 doi:10.1007/JHEP10(2014)092 [arXiv:1407.3282 [hep-th]].
42. R. Casero, E. Kiritsis and A. Paredes, Nucl. Phys. B **787** (2007), 98-134 doi:10.1016/j.nuclphysb.2007.07.009 [arXiv:hep-th/0702155 [hep-th]].
43. E. Megías and M. Valle, Fortsch. Phys. **66**, no.7, 1800035 (2018) doi:10.1002/prop.201800035 [arXiv:1707.04747 [hep-th]].
44. E. Megías, J. Phys. Conf. Ser. **1416**, no.1, 012022 (2019) doi:10.1088/1742-6596/1416/1/012022 [arXiv:1909.13836 [hep-th]].
45. I. Amado, K. Landsteiner and F. Pena-Benitez, JHEP **05** (2011), 081 doi:10.1007/JHEP05(2011)081 [arXiv:1102.4577 [hep-th]].
46. A. Gynther, K. Landsteiner, F. Pena-Benitez and A. Rebhan, JHEP **02** (2011), 110 doi:10.1007/JHEP02(2011)110 [arXiv:1005.2587 [hep-th]].