Applications of Sparse Hypergraph Colorings

Felix Christian Clemen

Karlsruhe Institute of Technology, 76133 Karlsruhe, Germany

June 4, 2024

Abstract

Many problems in extremal combinatorics can be reduced to determining the independence number of a specific auxiliary hypergraph. We present two such problems, one from discrete geometry and one from hypergraph Turán theory. Using results on hypergraph colorings by Cooper-Mubayi and Li-Postle, we demonstrate that for those two problems the trivial lower bound on the independence number can be improved upon:

• Erdős, Graham, Ruzsa and Taylor asked to determine the largest size, denoted by g(n), of a subset P of the grid $[n]^2$ such that every pair of points in P span a different slope. Improving on a lower bound by Zhang from 1993, we show that

$$g(n) = \Omega\left(\frac{n^{2/3}(\log\log n)^{1/3}}{\log^{1/3} n}\right).$$

• Let H_3^r denote an r-graph with r+1 vertices and 3 edges. Recently, Sidorenko proved the following lower bounds for the Turán density of this r-graph: $\pi(H_3^r) \ge r^{-2}$ for every r, and $\pi(H_3^r) \ge (1.7215 - o(1))r^{-2}$. We present an improved asymptotic bound:

$$\pi(H_3^r) = \Omega\left(r^{-2}\log^{1/2}r\right).$$

1 Introduction

A hypergraph is a pair (V, E) where V is a set whose elements are called *vertices*, and E is a family of subsets of V, called *edges*. A hypergraph is a k-graph if every edge has size k. A hypergraph has rank k if every edge contains at least 2 and at most k vertices.

A subset of vertices $I \subseteq V$ is called an *independent set* in H = (V, E) if there is no edge $e \in E$ contained in I. The *independence number* of a hypergraph H, denoted by $\alpha(H)$, is the size of a maximum independent set. Many problems in extremal combinatorics are related to the problem of determining the independence number of a specific auxiliary hypergraph, see e.g. [1,3,8,11,16]. In this paper we present two such problems, one from discrete geometry and one from hypergraph Turán theory. For both we use probabilistic tools to present an improved bound on the independence number.

1.1 Application 1: The distinct slopes in the grid problem

The unit distance problem, a classical problem in combinatorial geometry, posed by Erdős [7] in 1946, asks to determine the maximum number of unit segments in a set of n points in the plane. Since then a great variety of extremal problems in finite planar point sets have been studied. Here, we look at such a question concerning the grid $[n]^2$.

A subset $P \subseteq [n]^2$ of the grid has *distinct slopes* if there are no two pairs of points spanning lines with the same slope. Note that such a set P does not contain collinear triples, nor 4 points forming a trapezoid. Denote by g(n) the size of the largest subset P of the grid $[n]^2$ with distinct slopes. In 1992, Erdős, Graham, Ruzsa and Taylor [6] asked to determine this function and proved

$$\Omega(n^{1/2}) = g(n) = O(n^{4/5}).$$

Their lower bound comes from an algebraic construction. In 1993, Zhang [21] improved the lower bound by a deletion method argument resulting in

$$g(n) = \Omega\left(\frac{n^{2/3}}{\log^{1/3} n}\right)$$

To do so, Zhang [21] observed that the number T of trapezoids in $[n]^2$ is $O(n^6 \log n)$ and the number C of collinear triples is $O(n^4 \log n)$. Therefore, when considering a p-random subset of the grid, for $p = cn^{-4/3} \log^{-1/3} n$, where c is sufficiently small it holds that

$$pn^2 - p^4T - p^3C = \Omega\left(\frac{n^{2/3}}{\log^{1/3}n}\right).$$

Thus, there exists a subset $P \subseteq [n]^2$ of the desired size with no trapezoids and no collinear triples, and therefore P has distinct slopes. Here, we present the first improvement to this deletion method argument in 30 years.

Theorem 1.1. There exists a subset of the grid $[n]^2$ with distinct slopes and of size

$$g(n) = \Omega\left(\frac{n^{2/3}(\log\log n)^{1/3}}{\log^{1/3} n}\right)$$

1.2 Application 2: The Turán density of the *r*-graph with 3 edges on r + 1 vertices

Given an r-graph H, the Turán function ex(n, H) is the maximum number of edges in an H-free n-vertex r-graph. The Turán density of H, denote by $\pi(H)$, is

$$\pi(H) = \lim_{n \to \infty} \frac{\operatorname{ex}(n, H)}{\binom{n}{r}}.$$

It is a central question in extremal combinatorics to determine $\pi(H)$ for non r-partite r-graphs when $r \geq 3$. The Turán density of any r-graph with at most 2 edges is 0. A natural problem is to consider r-graphs with 3 edges. Denote by H_3^r an r-graph with 3 edges on r+1 vertices. Note that all such r-graphs are isomorphic. It can be observed quickly that $\pi(H_3^r) \to 0$, see e.g. [2] by Bollobás, Leader and Malvenuto. Specifically, $r!/r^r \leq \pi(H_3^r) \leq 2/(r+1)$. The lower bound holds, because H_3^r is not *r*-partite. The upper bound holds because in an H_3^r -free *r*-graph every r+1 vertices span at most 2 edges. Frankl and Füredi [9] proved $\pi(H_3^r) \geq 2^{1-r}$ using a geometric construction. Recently, this result was improved by Sidorenko [20] to $\pi(H_3^r) \geq r^{-2}$ for all r and $\pi(H_3^r) \geq (1.7155 - o(1))r^{-2}$ for $r \to \infty$. The best general upper bound is $\pi(H_3^r) \leq 1/r$ by Gunderson and Semerano [13] who used a double counting argument. See [12] for more precise bounds on $\pi(H_3^r)$ for small values of r. We present an asymptotic improvement on the lower bound.

Theorem 1.2. $\pi(H_3^r) = \Omega\left(\frac{\sqrt{\log r}}{r^2}\right).$

1.3 Sparse hypergraph coloring

In this subsection we present the tools we are using to prove Theorems 1.1 and 1.2.

A proper coloring of a hypergraph H is an assignment of colors to the vertices so that no edge is monochromatic. The smallest number of colors such that a proper coloring of H with this many colors exists, is called the *chromatic number* of H and denoted by $\chi(H)$.

Let H be a rank k hypergraph. For an integer $2 \leq l \leq k$, and a set S of vertices, $1 \leq |S| < \ell$, we define $\deg_{\ell}(S, H)$ to be the number of size ℓ edges containing the set S. When H is clear from context, we omit H and write $\deg_{\ell}(S)$ instead. The maximum l-degree of H, denoted by $\Delta_{\ell}(H)$, is the maximum of $\deg_{\ell}(\{v\}, H)$ over all vertices $v \in V(H)$. The maximum (s, ℓ) -codegree of H, denoted by $\Delta_{s,\ell}(H)$, is the maximum of $\deg_{\ell}(S, H)$ over all vertex sets S of size s in H. For a k-graph, we write $\Delta(H)$ for the maximum degree $\Delta_{1,k}(H)$. Again, when H is clear from context, we omit H and simply write Δ_{ℓ} , $\Delta_{s,\ell}$ and Δ respectively.

An elementary greedy coloring algorithm shows that any graph G has chromatic number $\chi(G) \leq \Delta_2 + 1$. In 1996, Johansson [15] proved that $\chi(G) = O(\Delta_2/\log \Delta_2)$ for any trianglefree graph G. Corresponding bounds on the independence number were established by Ajtai-Komlós-Szemerédi [1] and Shearer [19]. Frieze and Mubayi [10] generalized Johansson's theorem to all 3-uniform linear hypergraphs, and Cooper and Mubayi [5] later generalized it to all rank 3 hypergraphs. The following result by Cooper and Mubayi [4] gives an upper bound on the chromatic number $\chi(G)$ of a k-graph under a codegree condition.

Theorem 1.3 (Cooper, Mubayi [4]). Fix $k \ge 3$. Let H be a k-graph. If

$$\Delta_{\ell,k}(H) \le \Delta^{\frac{k-\ell}{k-1}} / f \qquad \text{for } \ell = 2, \dots, k-1, \tag{1}$$

then

$$\chi(H) = O\left(\left(\frac{\Delta}{\log f}\right)^{1/(k-1)}\right)$$

Recently, Li and Postle [17] generalized Theorem 1.3 to hold for rank k hypergraphs. Given two hypergraphs F_1 and F_2 , a map $\Phi: V(F_1) \to V(F_2)$ is an *isomorphism* if for all subsets $E \subseteq V(F_1)$, we have $\Phi(E) \in F_2$ if and only if $E \in F_1$. If there exists an isomorphism $\Phi: V(F_1) \to V(F_2)$, we say F_1 is isomorphic to F_2 and denoted it by $F_1 \cong_{\Phi} F_2$. For two hypergraphs F, H and a vertex $v \in V(F)$, let

$$\Delta_{F,v}(H) = \max_{u \in V(H)} |F' \subseteq H : F' \cong_{\Phi} F \text{ and } \Phi(u) = v|$$

and

$$\Delta_F(H) = \min_{v \in V(F)} \Delta_{F,v}(H).$$

Theorem 1.4 (Li, Postle [17]). Fix $k \ge 3$. Let \mathcal{H} be a rank k hypergraph. Suppose that for all $2 \le s < \ell \le k$,

$$\Delta_{s,\ell}(\mathcal{H}) \le \left(\max_{2 \le \ell \le k} \Delta_\ell(\mathcal{H})^{1/(\ell-1)}\right)^{\ell-s} / f,\tag{2}$$

and additionally for the graph triangle K_3 ,

$$\Delta_{K_3}(\mathcal{H}) \le \left(\max_{2 \le \ell \le k} \Delta_\ell(\mathcal{H})^{1/(\ell-1)}\right)^2 / f.$$
(3)

Then we have

$$\chi(\mathcal{H}) \le O\left(\max_{2 \le \ell \le k} \left\{ \left(\frac{\Delta_{\ell}(\mathcal{H})}{\log f}\right)^{\frac{1}{\ell-1}} \right\} \right)$$

We point out that Li and Postle conclude Theorem 1.4 from a more general result (see Theorem 1.7 in [17]).

This paper is organized as follows. In Section 2 we first collect several observations about the grid, and then we prove Theorem 1.1 using Theorem 1.4. Finally, in Section 3, we present the proof of Theorem 1.2 using Theorem 1.3.

2 Application 1: The distinct slopes in the grid problem

In this section we prove Theorem 1.1. To do so, we first collect several facts about the grid.

2.1 Observations about the grid

The elements from the grid $[n]^2$ are referred to as *grid points*. The slope of a line ℓ in \mathbb{R}^2 can be written in the form $\frac{u(\ell)}{v(\ell)}$, where $gcd(|u(\ell)|, |v(\ell)|) = 1$ and $v(\ell) > 0$. Define

$$H(l) := \max\{|u(\ell)|, |v(\ell)|\}.$$

The number of grid points on a line with $H(\ell) = s$ is at most O(n/s). The origin in the grid is the point o = (1, 1). Two grid points (x, y), (x', y') are mutually visible if the line segment joining them contains no further grid point. A grid point (x, y) is called visible if (x, y) and the origin are mutually visible. A classical result in the theory of visible lattice points says that the number of visible grid points is $\frac{6}{\pi^2}n^2(1 + o(1))$, see e.g. [14, 18]. We will use the following crude estimate on the number of visible lattice in a specific subset of the grid repeatedly. For a positive integer i, let V_i be the set of visible points in $[2^i]^2 \setminus [2^{i-1}]^2$.

Observation 2.1. For every *i* sufficiently large, $|V_i| \ge \frac{1}{4}2^{2i}$.

Proof. The number of visible points in $[2^i]^2$ is $\frac{6}{\pi^2}2^{2i}(1+o(1))$ and the total number of points in $[2^{i-1}]$ is $\frac{1}{4}2^{2i}$, implying this observation.

Denote by \mathcal{H} the rank 4-graph with vertex set $[n]^2$, where the 3-edges are the collinear triples and 4-edges are trapezoids. Then $g(n) = \alpha(\mathcal{H})$. We remark that Theorem 1.4 cannot be applied directly to \mathcal{H} because the codegree condition (2) on $\Delta_{2,4}$ is not satisfied. To circumvent this obstruction, we modify the hypergraph \mathcal{H} .

Definition 2.2. Let s^* be a positive integer. Denote by \mathcal{H}_{s^*} the rank 4-graph with vertex set $[n]^2$ with edges:

- 2-sets of grid points lying on a line ℓ which satisfies $H(\ell) \leq s^*$,
- 3-sets of grid points forming collinear triples,
- 4-sets of grid points which form a trapezoid with the additional property that for each choice of two parallel lines ℓ_1, ℓ_2 , going through two of the four points each, they satisfy $H(l_i) \ge s^*$. (Note that for a rhombus there are two ways of choosing two parallel lines.)

The following lemma provides several properties of the auxiliary hypergraph \mathcal{H}_{s^*} .

Lemma 2.3. Let $\varepsilon > 0$ and s^* be a positive integer, $\Omega(n^{\varepsilon}) = s^* = O(n^{1-\varepsilon})$. Then,

Proof of Lemma 2.3 a). Let i_0 be sufficiently large such that Observation 2.1 holds for $i \ge i_0$. Let $i_0 \le i \le \lfloor \log(s^*) \rfloor$. Given a visible point $p \in V_i$, let ℓ_p be the line through the origin o and p. The line ℓ_p contains $\Omega(n/2^i)$ grid points. For any grid point q, the pair qo forms a 2-edge in \mathcal{H}_{s^*} if q is on a line ℓ_p for some visible point $p \in [s^*]^2$. Thus,

$$\Delta_2(\mathcal{H}_{s^*}) \ge \deg_2(o) \ge \sum_{i=i_0}^{\lfloor \log s^* \rfloor} |V_i| \ \Omega\left(\frac{n}{2^i}\right).$$

By Observation 2.1, $|V_i| \ge 2^{2i-2}$ and therefore $\Delta_2(\mathcal{H}_{s^*}) = \Omega(ns^*)$.

For the upper bound, let $p \in [n]^2$ be an arbitrary grid point. Given $s \leq s^*$, there are at most 4s choices for slopes of lines ℓ passing through p and satisfying $H(\ell) = s$. On each such line there are at most O(n/s) grid points. Thus,

$$\deg_2(p) \le \sum_{s=1}^{s^*} 4s \ O\left(\frac{n}{s}\right) = O(ns^*).$$

Since the grid point p was chosen arbitrarily, $\Delta_2(\mathcal{H}_{s^*}) = O(ns^*)$.

Proof of Lemma 2.3 b). We will provide a lower bound on $\Delta_3(\mathcal{H}_{s^*})$ by counting the number of collinear triples containing the origin o. Let i_0 be sufficiently large such that Observation 2.1 holds for $i \geq i_0$. Let $i_0 \leq i \leq \lfloor \log(n) \rfloor$. Given a visible point $p \in V_i$, let ℓ_p be the line through the origin

o and p. The line ℓ_p contains $\Omega(n/2^i)$ grid points. For grid points q and q' a 3-set qq'o forms a 3-edge in \mathcal{H}_{s^*} iff q and q' are on a line ℓ_p for some visible point $p \in [n]^2$, we obtain

$$\Delta_3(\mathcal{H}_{s^*}) \ge \deg_3(o) \ge \sum_{i=i_0}^{\lfloor \log n \rfloor} |V_i| \ \Omega\left(\begin{pmatrix}\frac{n}{2^i}\\2\end{pmatrix}\right).$$

Using Observation 2.1, we get $\Delta_3(\mathcal{H}_{s^*}) = \Omega(n^2 \log n)$.

For the upper bound on $\Delta_3(\mathcal{H}_{s^*})$, let $p \in [n]^2$ be an arbitrary grid point. Given $s \leq s^*$, there are at most 4s choices for slopes of lines ℓ passing through p and satisfying $H(\ell) = s$. On each such line there are at most O(n/s) grid points. Thus,

$$\deg_3(p) \le \sum_{s=1}^{s^*} 4s \ O\left(\left(\frac{n}{s}\right)^2\right) = O(n^2 \log n).$$

Since p was chosen arbitrarily from $[n]^2$, $\Delta_3(\mathcal{H}_{s^*}) = O(n^2 \log n)$.

Proof of Lemma 2.3 c). For the lower bound on $\Delta_4(\mathcal{H}_{s^*})$, we count the number of 4-edges containing the origin. Let $i \in \mathbb{N}$ satisfying that $\log s^* \leq i \leq \log n$. Denote by T_i the number of trapezoids containing the origin o and exactly one pair of parallel sides (i.e. rhombi are not being counted) such that the line through the origin whose slope appears twice in the trapezoid contains a visible point $p \in V_i$. Fixing a point $p \in V_i$, there are $\Omega(n/2^i)$ points on the line ℓ_p through the origin o and p. For the third point of the trapezoid we choose an arbitrary grid point from $\{(p_1, p_2) : n/4 \leq p_i \leq 3n/4\}$ which is not on the line ℓ_p . There are $\Omega(n/2^i)$ ways to choose a forth point to complete a trapezoid, and thus in total

$$\Delta_4(\mathcal{H}_{s^*}) \ge \deg_4(o) \ge \sum_{i=\lceil \log s^* \rceil}^{\lfloor \log n \rfloor} |V_i| \ \Omega\left(\frac{n}{2^i} \cdot n^2 \cdot \frac{n}{2^i}\right).$$

Using Observation 2.1, we get $\Delta_4(\mathcal{H}_{s^*}) = \Omega(n^4 \log n)$.

Now, let $p \in [n]^2$ be an arbitrary point. We will count the number of trapezoids containing p. For a given positive integer s, let T_s be the number of trapezoids containing p and two parallel sides on lines ℓ_1, ℓ_2 satisfying $H(\ell_1) = H(\ell_2) = s$. Then, clearly

$$\deg_4(p) \le \sum_{s=1}^n T_s.$$

There are at most 4s lines ℓ through p satisfying $H(\ell) = s$. For each of them, there are at most O(n/s) grid points on it. For the third point of a trapezoid there are at most n^2 choices trivially. The last point of the trapezoid needs to be on a line ℓ' through the third point and parallel to ℓ , that are at most O(n/s) choices for such a point. Thus,

$$T_s \le O\left(4s \cdot \frac{n}{s} \cdot n^2 \cdot \frac{n}{s}\right) = O\left(\frac{n^4}{s}\right)$$

which implies

$$\deg_4(p) \le \sum_{s=1}^n T_s = O(n^4 \log n).$$

Since p was arbitrary, we conclude $\Delta_4(\mathcal{H}_{s^*}) = O(n^4 \log n)$.

Proof of Lemma 2.3 d). Since any line in \mathbb{R}^2 contains at most n grid points, we immediately obtain $\Delta_{2,3}(\mathcal{H}_{s^*}) = O(n)$.

Proof of Lemma 2.3 e). Let $A = \{p_1, p_2, p_3\}$ be a 3-element subset of the grid. If the points p_1, p_2, p_3 form a collinear triple, then they cannot be contained in a trapezoid and thus $\deg_4(A) = 0$. Otherwise, for $1 \leq i < j \leq 3$, let $\ell_{i,j}$ be the line through p_i and p_j . A grid point p forming a trapezoid with the points from A, must lie on the line thorough p_m parallel to $\ell_{i,j}$ for some choice $\{i, j, m\} = [3]$. For any such point p, the set $\{p, p_1, p_2, p_3\}$ is an edge in \mathcal{H}_{s^*} , if additionally $H(\ell_{i,j}) \geq s^*$. Thus, there are at most $O(n/s^*)$ choices for p. We conclude $\Delta_{3,4}(\mathcal{H}_{s^*}) = O(n/s^*)$. \Box

Proof of Lemma 2.3 f). By Lemma 2.3 e),

$$\Delta_{2,4}(\mathcal{H}_{s^*}) \le n^2 \Delta_{3,4}(\mathcal{H}_{s^*}) = O\left(\frac{n^3}{s^*}\right).$$

2.2 Proof of Theorem 1.1

Let $s^* = \lceil n^{1/3} \rceil$ and $f = \log^{1/2} n$. Then, by Lemma 2.3

$$\gamma := \max_{2 \le \ell \le 4} \Delta_{\ell}(\mathcal{H}_{s^*})^{1/(\ell-1)} = \Theta(\max\{n^{4/3}, n \log^{1/2} n, n^{4/3} \log^{1/3} n\}) = \Theta(n^{4/3} \log^{1/3} n).$$

Further,

$$\Delta_{2,3}(\mathcal{H}_{s^*}) = O(n), \qquad \Delta_{3,4}(\mathcal{H}_{s^*}) = O(n^{2/3}), \qquad \text{and} \qquad \Delta_{2,4}(\mathcal{H}_{s^*}) = O(n^{8/3}).$$

Therefore,

$$\Delta_{2,3}(\mathcal{H}_{s^*}) = O(n) \le \Theta\left(\frac{n^{4/3}\log^{1/3}n}{f}\right) = \frac{\gamma}{f},$$

$$\Delta_{3,4}(\mathcal{H}_{s^*}) = O(n^{2/3}) \le \Theta\left(\frac{n^{4/3}\log^{1/3}n}{f}\right) = \frac{\gamma}{f},$$

$$\Delta_{2,4}(\mathcal{H}_{s^*}) = O(n^{8/3}) \le \Theta\left(\frac{n^{8/3}\log^{2/3}n}{f}\right) = \frac{\gamma^2}{f}.$$

We conclude that the conditions (2) are satisfied. Further, for the graph triangle K_3 , by Lemma 2.3

$$\Delta_{K_3}(\mathcal{H}_{s^*}) \le \Delta_2(\mathcal{H}_{s^*})^2 = O(n^{8/3}) \le \Theta\left(\frac{n^{8/3}\log^{2/3}n}{f}\right) = \frac{\gamma^2}{f}.$$

We conclude that (3) is satisfied. Note that

$$\max\left\{\left(\frac{\Delta_{\ell}}{\log\log n}\right)^{\frac{1}{\ell-1}}\right\} = \max\left\{\left(\frac{n^{4/3}}{\log\log n}\right), \left(\frac{n^2\log n}{\log\log n}\right)^{\frac{1}{2}}, \left(\frac{n^4\log n}{\log\log n}\right)^{\frac{1}{3}}\right\} = \Theta\left(\frac{n^{4/3}\log^{1/3} n}{(\log\log n)^{1/3}}\right).$$

Therefore, by Theorem 1.4,

$$\chi(\mathcal{H}_{s^*}) = O\left(\max_{2 \le \ell \le 4} \left\{ \left(\frac{\Delta_\ell}{\log f}\right)^{\frac{1}{\ell-1}} \right\} \right) = O\left(\frac{n^{4/3} \log^{1/3} n}{(\log \log n)^{1/3}}\right).$$

Using that every independent set in \mathcal{H}_{s^*} is an independent set in \mathcal{H} , we conclude

$$g(n) = \alpha(\mathcal{H}) \ge \alpha(\mathcal{H}_{s^*}) = \Omega\left(\frac{n^{2/3}(\log\log n)^{1/3}}{\log^{1/3} n}\right).$$

This completes the proof of Theorem 1.1.

3 Application 2: The Turán density of H_3^r

In this section we prove Theorem 1.2 using Theorem 1.3.

3.1 The blow-up operation

Let G be an r-graph. The blow-up G(t) of G is the r-graph which is obtained from G by replacing each vertex $x \in V(G)$ by t vertices x^1, \ldots, x^t and each edge $x_1 \cdots x_r \in E(G)$ by t^r edges $x_1^{a_1} \cdots x_r^{a_r}$ with $1 \leq a_1, \ldots, a_r \leq t$. We say an r-graph H has complete shadow if every pair of vertices is contained in an edge. The following two observations about blow-ups are essential.

Lemma 3.1. Let $m \in \mathbb{N}$ and H be an r-graph with complete shadow. If G is an H-free r-graph, then G(m) is H-free.

Proof. Towards contradiction, assume that G(m) contains a copy of H on some vertex set $U \subseteq V(G(m))$. Then, because H has complete shadow, for any two vertices $x_1^i, x_2^j \in U$ we have $x_1 \neq x_2$. Therefore G contains a copy of H too, a contradiction.

Lemma 3.2. Let G be an r-graph with m vertices, where $m \ge r^2$. Let n be an integer divisible by m, and let t = n/m. Then

$$\frac{|E(G(t))|}{\binom{n}{r}} \ge \frac{1}{4} \frac{|E(G)|}{\binom{m}{r}}$$

Proof. We have

$$\frac{|E(G(t))|}{\binom{n}{r}} = \frac{|E(G)|t^{r}}{\binom{n}{r}} = \frac{|E(G)|t^{r}}{\binom{m}{r}\frac{n!(m-r)!}{m!(n-r)!}} = \frac{|E(G)|}{\binom{m}{r}}\frac{m!(n-r)!}{n!(m-r)!}\frac{n^{r}}{m^{r}} \ge \frac{|E(G)|}{\binom{m}{r}}\frac{m!}{(m-r)!m^{r}}$$
$$\ge \frac{|E(G)|}{\binom{m}{r}}\left(1 - \frac{r}{m}\right)^{r} \ge \frac{|E(G)|}{\binom{m}{r}}\left(1 - \frac{1}{r}\right)^{r} \ge \frac{1}{4}\frac{|E(G)|}{\binom{m}{r}}.$$

Informally, the takeaway of Lemmas 3.1 and 3.2 is that in order to understand the asymptotic behavior of Turán density of H_3^r it suffices to consider host graphs with roughly r^2 vertices.

3.2 Proof of Theorem 1.2

Let G be the auxiliary 3-graph with vertex set $V = {[r^2] \choose r}$ and edges $A \in E(G)$ iff A forms a copy of H_3^r in $K_{r^2}^r$. Note that

$$\Delta = \Delta(G) = \binom{r}{r-2}(r^2 - r),$$

because simply every edge in $K_{r^2}^r$ is contained in this many copies of H_3^r . Therefore, $r^4/3 \leq \Delta \leq r^4$ for r sufficiently large. Set $f = \Delta^{1/10}$ and let $A = \{e_1, e_2\} \subseteq V(G)$ of size 2. We will upper bound deg₃(A). If $|e_1 \cap e_2| < r - 1$, then deg₃(A) = 0. If $|e_1 \cap e_2| = r - 1$, then any $f \in V(G)$ satisfying that e_1, e_2, f form a copy of H_3^r has the property that $f \subseteq e_1 \cup e_2$. Since $f \neq e_1$ and $f \neq e_2$, we have

$$\deg_3(A) \le \binom{|e_1 \cup e_2|}{r} - 2 = \binom{r+1}{r} - 2 = r - 1.$$

We conclude $\Delta_{2,3}(G) \leq r-1$ and thus condition (1) holds for G:

$$\Delta_{2,3}(G) \le r - 1 \le \left(\frac{r^4}{3}\right)^{\frac{1}{3}} \le \Delta^{\frac{1}{3}} \le \frac{\Delta^{\frac{1}{2}}}{\Delta^{\frac{1}{10}}} = \frac{\Delta^{\frac{1}{2}}}{f}.$$

By Theorem 1.3 the 3-graph G satisfies

$$\chi(G) = O\left(\left(\frac{\Delta}{\log \Delta}\right)^{\frac{1}{2}}\right) = O\left(\frac{r^2}{\left(\log r\right)^{\frac{1}{2}}}\right).$$

Therefore G contains an independent set of size at least

$$\alpha(G) = \Omega\left(\binom{r^2}{r} \frac{(\log r)^{\frac{1}{2}}}{r^2}\right).$$

This independent set corresponds to an H_3^r -free *r*-graph *H* on r^2 vertices with $\alpha(G)$ many edges. Let $t = n/r^2$. By considering the blow-up H(t) of this *r*-graph and Lemmas 3.1 and 3.2 we obtain

$$\pi(H_3^r) \ge \limsup_{n \to \infty} \frac{|E(H(t))|}{\binom{n}{r}} \ge \frac{1}{4} \frac{|E(H)|}{\binom{r^2}{r}} = \Omega\left(\frac{(\log r)^{\frac{1}{2}}}{r^2}\right),$$

completing the proof of Theorem 1.2.

4 Acknowledgements

The author thanks Maria Axenovich, József Balogh, Alberto Espuny Díaz, Peter Kaiser, Lina Li, Dingyuan Liu, Letícia Mattos, Ethan Patrick White for valuable discussions on the topic of this paper. Special thanks are extended to Hong Liu and Zixiang Xu; Hong Liu for organizing the 1st ECOPRO combinatorial week workshop in Daejeon, and Zixiang Xu for presenting the distinct slopes problem.

References

- M. Ajtai, J. Komlós, and E. Szemerédi, A note on Ramsey numbers, Journal of Combinatorial Theory, Series A 29 (1980), no. 3, 354–360.
- [2] B. Bollobás, I. Leader, and C. Malvenuto, *Daisies and other Turán problems*, Combinatorics, Probability and Computing 20 (2011), no. 5, 743–747.
- [3] M. Campos, M. Jenssen, M. Michelen, and J. Sahasrabudhe, A new lower bound for sphere packing, arXiv preprint arXiv:2312.10026 (2023).

- [4] J. Cooper and D. Mubayi, *Coloring sparse hypergraphs*, SIAM Journal on Discrete Mathematics (2016), 1165-1180.
- [5] _____, List coloring triangle-free hypergraphs, Random Structures & Algorithms 47 (2015), no. 3, 487–519.
- [6] P. Erdős, R. Graham, I. Z Ruzsa, and H. Taylor, Bounds for arrays of dots with distinct slopes or lengths, Combinatorica 12 (1992), no. 1, 39–44.
- [7] P. Erdős, On sets of distances of n points, Amer. Math. Monthly 53 (1946), 248–250.
- [8] Irene Gil Fernández, Jaehoon Kim, Hong Liu, and Oleg Pikhurko, New lower bound on ball packing density in high-dimensional hyperbolic spaces, arXiv preprint arXiv:2405.07818 (2024).
- [9] P. Frankl and Z. Füredi, An exact result for 3-graphs, Discrete Math. 2-3 (1984), 323–328.
- [10] A. Frieze and D. Mubayi, On the chromatic number of simple triangle-free triple systems, The Electronic Journal of Combinatorics 15 (2008), no. 1.
- [11] Z. Füred, Maximal independent subsets in Steiner systems and in planar sets, SIAM Journal on Discrete Mathematics 4 (1991), no. 2, 196–199.
- [12] K. Gunderson and J. Semeraro, Turán numbers and switching, arXiv preprint arXiv:2204.10775 (2022).
- [13] _____, Tournaments, 4-uniform hypergraphs, and an exact extremal result, Journal of Combinatorial Theory, Series B 126 (2017), 114–136.
- [14] F. Herzog and B. M. Stewart, Patterns of Visible and Nonvisible Lattice Points, The American Mathematical Monthly 78 (1971), no. 5, 487–496.
- [15] A. Johansson, Asymptotic choice number for triangle free graphs (1996).
- [16] M. Krivelevich, S. Litsyn, and A. Vardy, A lower bound on the density of sphere packings via graph theory, International Mathematics Research Notices 2004 (2004), no. 43, 2271–2279.
- [17] L. Li and L. Postle, The chromatic number of triangle-free hypergraphs, arXiv preprint arXiv:2202.02839 (2022).
- [18] H. Rademacher, Lectures on Elementary Number Theory (1964).
- [19] J. B Shearer, A note on the independence number of triangle-free graphs, Discrete Mathematics 46 (1983), no. 1, 83–87.
- [20] A. Sidorenko, Turán numbers of r-graphs on r + 1 vertices, arXiv preprint arXiv:2205.02006 (2022).
- [21] Z. Zhang, A note on arrays of dots with distinct slopes, Combinatorica 13 (1993), no. 1, 127–128.