Šoltés' hypergraphs

Stijn Cambie *

June 4, 2024

Abstract

More than 30 years ago, Šoltés observed that the total distance of the graph C_{11} does not change by deleting a vertex, and wondered about the existence of other such graphs, called Šoltés graphs. We extend the definition of Šoltés' graphs to Šoltés' hypergraphs, determine all orders for which a Šoltés' hypergraph exists, observe infinitely many uniform Šoltés' hypergraphs, and find the Šoltés' hypergraph with minimum size (spoiler: it is not C_{11}).

1 Introduction

The total distance, which is proportional with the average distance, is one of the most important distance-based graph parameters. For a graph G, we denote the total distance with W(G). Here $W(G) = \sum_{u,v \in V(G)} d(u,v)$ is the sum of distances between any (unordered) pair of vertices in a graph G. We will examine one of the most easy-to-state questions related to the total distance. In 1991, Šoltés [7] observed that $W(C_{11}) = W(C_{11} \setminus u)$ for every vertex $u \in V(C_{11})$, and asked whether there are other graphs G (nowadays called Šoltés' graphs) satisfying $W(G) = W(G \setminus u)$ for all vertices u of G. It has been conjectured that C_{11} is the only Šoltés' graph ([3, conj. 51]).

One can generalize the notion of Šoltés' graphs to Šoltés' hypergraphs. A hypergraph H = (V, E), with $E \subset 2^V$, is a pair consisting of a set of vertices and a set of edges, where an edge is a set that can contain any number of vertices (we will assume at least 2, as edges of size 1 do not matter in computing the total distance). More precise terminology is given in Subsection 1.1. A Šoltés' hypergraph is a connected hypergraph H for which $W(H) = W(H \setminus v)$ (which needs to be finite) for every $v \in V(H)$. Here W(H) is the sum of distances over all pairs of vertices, where the distance d(u, v) is the minimum number of edges in a connected subhypergraph of H containing both u and v. By convention, K_1 is not considered as a Šoltés' hypergraph. In Section 2, we prove that there exists a Soltés' hypergraph of order n if and only if $n \geq 5$. The latter trivially implies that there are infinitly many Soltés' hypergraphs.

In Section 3, we consider the problem for uniform Šoltés' hypergraphs. Those are hypergraphs for which all edges have the same cardinality (graphs are 2-uniform hypergraphs). We give two infinite examples of families of uniform Šoltés' hypergraphs, indicating the abundance of solutions. By the discovery of a few examples of uniform Šoltés' hypergraphs which have larger diameter (arguably, they behave like graphs in this context), the original version of Šoltés' graph. There still remain natural subproblems, such as determining the smallest uniform Šoltés' hyper-

There still remain natural subproblems, such as determining the smallest uniform Soltes' hypergraphs. We find two uniform Šoltés' hypergraphs of order 10 (hence C_{11} is not the smallest uni-

^{*}Department of Computer Science, KU Leuven Campus Kulak-Kortrijk, 8500 Kortrijk, Belgium. Supported by a postdoctoral fellowship by the Research Foundation Flanders (FWO) with grant number 1225224N.

form Šoltés' hypergraph); the hemi-dodecahedron (see Fig. 4) and the dual of $K_5^{(3)}$ (see Fig. 5). We prove that the latter is the unique Soltés' hypergraph of minimum size.

1.1 Notation and Terminology

A hypergraph H = (V, E), where $E \subset 2^V$, consists of a set of vertices and a set of hyperedges. A hyperedge is defined as a set containing at least two vertices¹ and called an edge when no confusion can arise. The number of vertices and edges are called the order and size respectively. For a vertex $v \in V$, $H \setminus v$ denotes the hypergraph $(V \setminus v, E')$, where E' includes all hyperedges in E not containing v. In essence, $H \setminus v$ is obtained by removing v and all hyperedges containing v. The dual $H^* = (E, V^*)$ of a hypergraph H = (V, E), is the hypergraph whose vertices are the edges of H, and whose hyperedges correspond with the tuples of all edges incident to a vertex of V. A clique $K_n^{(r)}$ is the r-uniform hypergraph with a vertex set of order n, containing all possible hyperedges with r elements. So up to isomorphism, $K_n^r = ([n], {[n] \choose r})$ where $[n] = \{1, 2, \ldots, n\}$.

The degree of a vertex v, denoted by deg(v), is the number of hyperedges containing v. The minimum and maximum degrees, denoted as δ and Δ , represent the smallest and largest degrees in the hypergraph.

A path in H is represented by a sequence $(v_0, e_1, v_1, e_2, v_2, \ldots, e_r, v_r)$, where each e_i is a hyperedge containing vertices v_{i-1} and v_i for every $i, v_0 = u$, and $v_r = v$. No edge or vertex appears multiple times in the path, and r denotes the length of the path. The distance d(u, v) between two vertices u, v in a hypergraph H is defined as the length of the shortest path between them.

For a hypergraph H, the total distance or Wiener index, W(H), is the sum of distances between any pair of vertices in H, i.e., $W(H) = \sum_{u,v \in V(H)} d(u,v)$. Equivalently, W(H) equals the total distance W(G') of the underlying graph G' = (V, E'), where $E' = \bigcup \{ \binom{e}{2} \mid e \in E \}$ comprises all 2-uniform edges that are subsets of a hyperedge in E.

2 Non-uniform Šoltés' hypergraphs

Before proving the existence of non-uniform Soltés' hypergraphs of almost any order, we present a few small examples.

Example 1. Consider the hypergraphs in Fig. 1. Here edges are drawn with lines, and hyperedges with dashed ellipses. These satisfy $W(H) = W(H \setminus v) = \binom{n}{2}$ for any $v \in V(H)$.



Figure 1: Examples of non-uniform Šoltés' hypergraphs for n = 5, 6, 7, 8

We give a few remarks for the hypergraph H on 7 and 8 vertices.

Case n = 7: Since *H* has three orbits, there are three computations needed to verify that $W(H \setminus v) = W(H) = \binom{7}{2} = 21$ for every $v \in V(H)$. The hypergraph is non-regular, since

¹Edges with a single vertex would not influence the total distance.

its minimum degree equals $\delta = 4$ and the maximum degree is $\Delta = 6$. This imply that some conjectures [3, conj. 47,48] stating that a Šoltés' graph is regular or vertex-transitive, are not true for the extension towards Šoltés' hypergraphs. Also, H contains twin-vertices, which cannot happen in a Šoltés' graph.

Case n = 8: The hypergraph on 8 vertices is obtained from the circulant graph on [8] with difference set $\{1, 2\}$ together with hyperedges containing the even, odd and all vertices. Since the hypergraph is vertex-transitive, it is easy to verify that $W(H) = \binom{n}{2} = 28$ and $W(H \setminus v) = W(K_4) + W(P_3) + 3 \cdot (2 \cdot 2 + 2 \cdot 1) = 6 + 4 + 18 = 28$ for any $v \in V(H)$.

Next, we generalize the folklore result on the extrema of the Wiener index for graphs of a given order. This will be used afterwards to prove that no Šoltés' hypergraph can have order bounded by 4.

Proposition 2. For a connected hypergraph H on n vertices, we have

$$\binom{n}{2} = W(K_n) \le W(H) \le W(P_n) = \binom{n+1}{3}$$

The upper bound is strict if $H \neq P_n$, while there are many hypergraphs attaining the minimum.

Proof. The lower bound is trivial, since all distances are at least one. The upper bound is by induction. Let S be a minimum set of (twin-)vertices such that $H \setminus S$ (remaining hypergraph after removing S and edges incident with S) is connected. The existence of such a set can be concluded by starting from a vertex and adding overlapping edges one at a time. Let |S| = s. Then by induction, we have that

$$W(H) \le H(H \setminus S) + s \cdot \sum_{i=1}^{n-s} i$$
$$= \binom{n-s+1}{3} + s\binom{n-s+1}{2}$$
$$\le \binom{n-s+1}{3} + \sum_{i=n-s+1}^{n} \binom{i}{2} = \binom{n+1}{3}.$$

For equality to be possible, it is necessary that s = 1 and $H \setminus S = P_{n-1}$ (by induction). From here one easily concludes that $H = P_n$, since the distances between the vertex S and the vertices in P_{n-1} have to be exactly 1 up to n-1.

We end this section with the main result, the determination of all orders for which a Soltés' hypergraph exists.

Theorem 3. The smallest Šoltés' hypergraph has order 5 and is unique. For every $n \ge 5$, there exists a Šoltés' hypergraph of order n.

Proof. For $2 \le n \le 4$, by Proposition 2, $W(H \setminus v) \le {n \choose 3} < {n \choose 2} \le W(H)$ if both H and $H \setminus v$ are connected (which is a necessary condition). For n = 5, equality needs to be attained, so $H \setminus v \cong P_4$ for ever v and W(H) = 10, implying that H is exactly equal to a C_5 with one hyperedge containing the 5 vertices.

For $5 \le n \le 8$, examples of Soltés' hypergraphs are depicted in Fig. 1.

Finally, we show that there are Šoltés' hypergraphs of order n for every $n \ge 9$, by giving an example, which actually depends on $n \pmod{12}$.

Partition the vertex set V in $\lfloor \frac{n}{3} \rfloor$ sets A_i of size 3 or 4. Let $E_i = V \setminus A_i$ for every $1 \le i \le \lfloor \frac{n}{3} \rfloor$. We start with constructing a graph G of order n. For a set A_i of size 4, we choose $G[A_i]$ to be a *m*-factor, where $n-1 \equiv 3-m \pmod{3}$ and every vertex from A_i has $\frac{2(n-4)-m}{3}$ neighbours in E_i . Note that every two of them have at least one common neighbour since $2\frac{2(n-4)-m}{3} > n-4$.

For a set A_i of size 3, if $2 \mid n - 1$, we choose either

- $G[A_i]$ to be a triangle K_3 and each vertex in A_i has $(n-3) \frac{n-1}{2}$ neighbours in E_i
- $G[A_i]$ to be a single edge, the two vertices of the K_2 in A_i both having $(n-3) \frac{n-1}{2}$ neighbours in E_i and let the third vertex have degree $(n-3) \frac{n-1}{2} + 1$ and have a common neighbour with each of the other vertices.

For a set A_i of size 3, if $2 \mid n-2$, we choose either

- $G[A_i]$ to be an empty graph $3K_1$ and each vertex in A_i has $(n-3) \frac{n-2}{2}$ neighbours in E_i (such that every two vertices of A_i have a common neighbour)
- $G[A_i]$ to be a P_3 , where the two endvertices in A_i have $(n-3) \frac{n-2}{2}$ neighbours in E_i (at least one in common) and the third vertex of A_i has degree $(n-3) \frac{n}{2}$.

Since n > 8, there is at least one A_i of size 3 and so the choices above can be made in such a way that the sum of degrees is even. There are multiple ways to see that a valid construction is possible for every order $n \ge 8$. One can e.g. do so by giving explicit constructions depending on $n \pmod{12}$. Finally, add the hyperedges E_i for every $1 \le i \le \frac{n}{3}$ and V to G, to form a hypergraph H. Now one can check that $W(H) = W(H \setminus v) = \binom{n}{2}$ in all cases. Since $H \setminus v$ has diameter 2, it is sufficient to check that there are n - 1 pairs of vertices at distance 2. If $v \in A_i$, E_i is a clique in $H \setminus v$, thus one needs to count the non-neighbours of the remaining vertices in A_i . The n - 1 pairs are now obtained by construction. If $|A_i| = 4$, this is immediate by construction, since there are 3 - m pairs with two vertices in A_i at distance 2 and n - 4 + m = (n - 1) - (3 - m) nonadjacent pairs of vertices in $A_i \times E_i$. If $|A_i| = 3$, the same conclusion holds since $2\frac{n-1}{2}$, $\frac{n}{2} + \frac{n-2}{2}$ are all equal to n - 1.

Examples of such hypergraphs are depicted in Fig. 2 for $n \in \{9, 10\}$ (where two hyperedges for the union of two colour classes (sets A_i) are not depicted for clarity of the drawing).



Figure 2: Šoltés' hypergraphs H for n = 9, 10

We conclude that there are numerous examples of non-uniform Šoltés' hypergraphs, and additional constructions are presented in Appendix B. To narrow our focus and potentially gain deeper insights, we turn our attention to a specific subclass of hypergraphs. Considering that graphs correspond to 2-uniform hypergraphs, the most natural class to explore is that of uniform hypergraphs. Our hope is that understanding uniform Šoltés' hypergraphs will provide valuable insights in the search for Šoltés' graphs. This is exactly the scope of the next section.

3 Uniform Šoltés' hypergraphs

In this section, we prove that there are (infinitely many) uniform Šoltés' hypergraphs. We do so by constructing two infinite families; one with increasing uniformity and one with fixed uniformity.

Proposition 4. There are infinitely many values of k for which there exists a k-uniform Šoltés' hypergraph.

Proof. Let $m \ge 4$, $k = \binom{m}{2}$ and $n = \binom{m+1}{2} + 1$. Let H be the hypergraph with V = [n] and E be the edges consisting of k consecutive (modulo n) numbers in [n]. Then H is vertex-transitive and $W(H) = \binom{n}{2}$, as diam(H) = 1 due to $2k - 1 \ge n$). For any $v \in V$, $W(H \setminus v) = \binom{n-1}{2} + \sum_{i=1}^{m} i = \binom{n}{2}$. The latter since $H \setminus v$ has diameter 2 and it is equal to a path for which the first m vertices have resp. $m, m - 1 \dots, 1$ non-neighbours in $V \setminus v$.

Proposition 5. There are infinitely many 4-uniform Šoltés' hypergraphs.

Proof. Consider a Steiner 2-design H = S(2, 4, n) with block size 4, for some $n \equiv 1, 4 \pmod{12}$ and $n \geq 13$. They exist for all such n, and there are many of them for large n, see [6].

Every 2 vertices share exactly one edge, thus diam(H) = 1. If u, w are two vertices such that $\{u, v, w\}$ are part of an edge e in H, then for a vertex $z \notin e$, there are two disjoint edges (in $G \setminus v$ which contain $\{z, u\}$ and $\{z, w\}$. This implies that for every vertex $v \in V(H)$,

$$W(H \setminus v) = \binom{n-1}{2} + \frac{n-1}{3}\binom{3}{2} = \binom{n-1}{2} + (n-1) = \binom{n}{2} = W(H).$$

Thus H is indeed a Šoltés' hypergraph, which has diameter 1 and is 4-uniform.

The examples in the above families all have diameter 1, making the computations easier. We remark that there are no 3-uniform Šoltés' hypergraphs with diameter 1. In that regard, finding a 3-uniform Šoltés' hypergraph may be somewhat equally difficult as finding a Šoltés' graph. Finding one, or excluding the existence seems a hard question. From any cubic arc-transitive graph G, one can derive a 2-regular 3-uniform hypergraph H with V(H) = E(G) and E(H) being exactly equal to the triangles in the line graph of G. Using the census on cubic arc-transitive graphs [2], see https://www.math.auckland.ac.nz/~conder/AllSymmCubic-OrderUpTo10000-ByEdgeSets.txt, by taking the dual, we find many examples of 3-uniform hypergraphs H for which $W(H)-W(H \setminus v)$ can be either positive or negative, and there are examples for which it is exactly equal to 1 or -1, or near zero compared with its order. See [1, doc. arctrans10000_3unif].

Proposition 6. There are no 3-uniform Šoltés' hypergraphs with diameter 1.

Proof. Let H be a 3-uniform Šoltés' hypergraph with diameter 1. We associate with every pair of vertices (v_i, v_j) in $\binom{V}{2}$, all vertices $v' \in V \setminus \{v_i, v_j\}$ for which $\{v_i, v_j, v'\} \in E$. For every vertex v', count the number $x_{v'}$ of pairs associated with v' and no other vertex. Since there are $\binom{n}{2}$ pairs in total, and n vertices, there is a vertex v for which this number is bounded by $\frac{n-1}{2}$. Then it is easy to see that $H \setminus v$ has diameter 2 as follows. For every 2 vertices u, w in $V \setminus v$, which are not adjacent in $H \setminus v$ there are at most $\frac{n-3}{2}$ vertices which are not a neighbour of both u, w.

So they have at least $\frac{n-3}{2} \ge 1$ (we only need to consider $n \ge 5$ by Theorem 3) neighbours in common. Since $diam(H \setminus v) = 2$, we conclude that $W(H \setminus v) = \binom{n-1}{2} + x_v < \binom{n}{2}$, contradicting the choice of H.

Remark 7. Among the 18 examples for S(2, 4, 25), as mentioned in [6], there are also non-vertex-transitive examples.

With the same idea as for Proposition 6, one can prove that all 4-uniform Šoltés' hypergraphs for which diam(H) = 1 and $diam(H \setminus v) = 2 \ \forall v \in v(H)$ were determined in Proposition 5.

A first example of a Soltés' hypergraph with diameter larger than 1 is obtained by duplicating the dual of K_4 (equivalently the Fano plane with one vertex removed), with the corresponding hyperedges put together, and where the vertex set of every copy is a hyperedge as well. This one is vertex-transitive, but not (hyper)edge-transitive (there is exactly one pair of complementary hyperedges). It is presented in Fig. 3 (left), with 4 hyperedges drawn by dashed lines, and also the vertices with the same colour belonging to a hyperedge.

Remark 8. The 6-uniform hypergraph on $\{0, 1, 2, ..., 11\}$ given by its hyperedges

((0, 1, 2, 3, 4, 5), (0, 1, 6, 7, 8, 9), (2, 3, 6, 7, 10, 11), (4, 5, 8, 9, 10, 11), (0, 2, 4, 6, 8, 10), (1, 3, 5, 7, 9, 11))is a (uniform) Šoltés' hypergraph with diameter 2.



Figure 3: Uniform Šoltés' hypergraphs H with diameter 2

All the constructions we have given so far had diameter 1 or 2. The simple reason for this is that $W(H) - W(H \setminus v)$ being zero is a rare event and thinking about hypergraphs with larger diameter is hard. Using the census of https://staff.matapp.unimib.it/~spiga/census.html [4, 5] and a computer search, we managed to find 4 additional Šoltés' hypergraphs, having diameter 2, 6, 7 and 8. They are 2-regular (as is the case for C_{11}) and 4-uniform, being the dual hypergraphs of 4-regular 2-uniform hypergraphs (graphs 45705, 51044, 51045, 51046 from https://houseofgraphs.org/graphs/), whose graph6-strings are also listed in Appendix A. The smallest one (the dual of the blow-up of a C_5) being depicted in Fig. 3 (right). Note that since the initial graphs are arc-transitive, the dual hypergraph is vertex-transitive and so it is sufficient to consider $W(H) - W(H \setminus v)$ for just one vertex v. The full verification can be found at [1, doc. graphs_dual_SH].

Subjectively, they may be considered as surprising as the possible existence of Soltés' graphs different from C_{11} (since there are more uniformities to consider, there were more chances for the existence of course) and evidence that excluding the existence of Soltés' graphs different from C_{11} is likely very hard if not impossible.

The examples so far are all somewhat large (order at least 11). Next, we present a uniform Šoltés' hypergraph, which has order 10. This may well be the minimum possible order.

Example 9. Consider the Petersen graph $P_{5,2}$, as e.g. depicted in Fig. 4 (left) and choose a (6,2)-cycle-cover, that is a collection of 6 smallest cycles in $P_{5,2}$ that cover every edge twice (this is easily seen to be unique up to isomorphism). Equivalently, choose the 6 faces of the hemi-dodecahedron as presented on the right in Fig. 4 (a and a' are the same vertex).



Figure 4: The Petersen graph $P_{5,2}$ and the hemi-dodecahedron

The (6,2)-cycle-cover forms a (vertex-transitive) 5-uniform Šoltés' hypergraph of order 10 and size 6, with $V(H) = \{0, 1, 2, ..., 9\}$ and

$$E(H) = \{(0, 1, 2, 3, 4), (0, 2, 5, 6, 7), (1, 3, 6, 7, 8), (2, 4, 7, 8, 9), (0, 3, 5, 8, 9), (1, 4, 5, 6, 9)\}.$$

Indeed, it is easy to verify that W(H) = 45 and also $W(H \setminus v) = 45$ for every $v \in [10]$. The latter since the hypergraph is vertex-transitive and taking e.g. v = 5, we note that $diam(H \setminus v) = 2$ and the pairs of vertices which are at distance 2 from each other contain at least one neighbour of v. This results in $W(H \setminus v) = {9 \choose 2} + 3 \cdot 4 - {3 \choose 2}$, since each neighbour of v belongs to one C_5 and thus has 4 neighbours and 4 non-neighbours within $H \setminus v$.

Finally, we determine the Soltés' hypergraph with minimum size. It is isomorphic to a (10, 6, 3)-design (BIBD). A structure with 10 elements, 5 edges containing 6 elements each, such that every element belongs to 3 edges and every 2 elements share precisely 3 edges. See Fig. 5. Here the elements/ vertices are the rows, and an edge is presented with a colour containing a cell of the 6 corresponding rows.



Figure 5: A (10, 6, 3)-BIBD; a 6-uniform Šoltés' hypergraph with order 10 and size 5

Theorem 10. The dual of $K_5^{(3)}$ is the unique uniform Šoltés' hypergraph of minimum size.

Proof. Note that a Soltés' hypergraph has minimum degree at least 2, since if not, a vertex v of degree 1 has a neighbour whose removal disconnects v from the remaining vertices. Also the

size m and maximum degree satisfy $m \ge \Delta + 2$, as otherwise removing a vertex of maximum degree leaves either a disconnected hypergraph, or a clique (with one hyperedge).

Now $2 \le \delta \le \Delta \le m - 2$ implies that $m \ge 4$. We first exclude m = 4.

Claim 11. A Šoltés' hypergraph cannot have size 4 or less.

Proof. In this case, $\Delta = \delta = 2$ and by Theorem 3, $n \ge 6$. No two vertices can belong to the same set of hyperedges (otherwise removing one of them disconnects the other), so $n \le {m \choose 2}$. The only case where equality holds, is the 3-uniform hypergraph on 6 vertices K_4^* , which is not a Šoltés' hypergraph.

For a k-uniform hypergraph of size 5 satisfying $\delta = 2$ and $\Delta = 3$, with vertices u and v having degree 2 and 3 respectively, we have $W(H \setminus u) < W(H \setminus v)$ if both are finite. So we conclude that a k-uniform Šoltés' hypergraph of size 5 is 2- or 3-regular. Then the order n satisfies $n\Delta = 5k$ by the analogue of the handshaking lemma and thus $5 \mid n$. Together with $n \geq 6$ and $n \leq \binom{m}{2} = 10$, this implies that n = 10. The equality $n = \binom{m}{2}$ implies that we know the hypergraph exactly as a function of $\Delta \in \{2, 3\}$. We have 5 edges, and every pair resp. triple of edges intersects in exactly 1 vertex. The 4-uniform hypergraph (K_5^{\star}) is not a Šoltés' hypergraph, but the 6-uniform hypergraph $(\binom{K_5^{(3)}}{2}^{\star})$ is; $W(H) = \binom{10}{2} = \binom{9}{2} + 3^2 = W(H \setminus v)$. As we have ruled out the other possibilities, we conclude that $(K_5^{(3)})^{\star}$ is the unique uniform Šoltés' hypergraph of size 5. \Box

With a little bit more work, the above theorem can be generalized to arbitrary hypergraphs.

Remark 12. Noting that a Šoltés' hypergraph of size 5 satisfies $\delta \geq 2, \Delta \leq 3$ and the incident edges of a vertex cannot be a subset of the incident edges of another vertex, we know that the candidates correspond with the dual of an antichain in $\binom{[5]}{2,3}$. For $n \geq 9$, the antichain is uniform and hence there are only 4 candidates. A brute force check ([1, doc. NonUniformSize5]) confirms that $\left(K_5^{(3)}\right)^*$ is also the unique (not necessarily uniform) Šoltés' hypergraph of size 5.

We conclude that a Šoltés' hypergraph has order at least 5 and size at least 5, and for both quantities there is a unique Šoltés' hypergraph attaining equality.

Analogous to the question by Šoltés' [7], one can ask to characterize all the uniform Šoltés' hypergraphs. Given the two infinite families and additional examples, it may be hard to characterize all them. This intuition may be even clearer in the case of non-uniform Šoltés' hypergraphs, where we also gave various non-regular examples. A few potential subquestions one may pose to have an idea on the richness of uniform Šoltés' hypergraphs, are the following ones (the third subquestion might be as hard as finding a Šoltés' graph different from C_{11}).

Question 13. • There is no uniform Šoltés' hypergraph with order bounded by 9?

- For every n ≥ 10, there exists an uniform Šoltés' hypergraph with order n? (Note that we have e.g. found examples for every 10 ≤ n ≤ 13)
- There exist 3-uniform Šoltés' hypergraphs?
- There exist k-uniform Šoltés' hypergraphs for every k ≥ 4?
 (Note that we have e.g. found examples for every 4 ≤ k ≤ 6)
- There are uniform Šoltés' hypergraphs which are not regular? (in Remark 7 we mentioned there are non-vertex-transitive examples)

We expect that all of these subquestions may well be answered with yes. The latter would indicate that the existence of a Šoltés' graph different from C_{11} cannot easily be excluded, given the richness of the class of uniform Šoltés' hypergraphs. For that reason, any negative answer on one of the last three questions would be even more surprising or interesting.

Acknowledgments

The author thanks Dragan Stevanović, Pablo Spiga, Primož Potočnik and Marston Conder for referrals, updating on the current censuses, and generating and uploading the full census (of arc-transitive cubic graphs with order bounded by 10^4).

References

- [1] S. Cambie. Code and data for the paper "šoltés' hypergraphs". https://github.com/StijnCambie/Soltes/tree/main/SoltesHypergraphs, 2024.
- [2] M. Conder and P. Dobcsányi. Trivalent symmetric graphs on up to 768 vertices. J. Combin. Math. Combin. Comput., 40:41–63, 2002.
- [3] M. Knor, R. Škrekovski, and A. Tepeh. Selected topics on wiener index, 2023.
- [4] P. Potočnik, P. Spiga, and G. Verret. Cubic vertex-transitive graphs on up to 1280 vertices. J. Symb. Comput., 50:465–477, 2013.
- [5] P. Potočnik, P. Spiga, and G. Verret. Bounding the order of the vertex-stabiliser in 3-valent vertex-transitive and 4-valent arc-transitive graphs. J. Combin. Theory Ser. B, 111:148–180, 2015.
- [6] C. Reid and A. Rosa. Steiner systems S(2, 4, v) a survey. *Electron. J. Comb.*, DS18:34, 2010.
- [7] L. Šoltés. Transmission in graphs: a bound and vertex removing. Mathematica Slovaca, 41(1):11–16, 1991.

A Large diameter uniform Šoltés' hypergraphs

Here we give the graph6-strings of four 4-regular graphs for which its dual is a Šoltés' hypergraph.

IYIYMOre_

```
~?@Ss'AA?_C@?A?G?C?@??H??a?G??G_?C??G??@???C?G???CG???@@??GG??
@@???A????C???A?@??@????O????B????C????G????A@????OG???OC?????
_G????@?????A?C????A?W?O?E??????????????????A????O??O?????W???????
```

```
~?@os'AQ__C@_D?A?@??OOK?@O?@O?@???_??W??I??@G???o???_???O???C?O?
@ ???S???@G????W????I????A????O????B?????Q????K?????a????@ ?????
@?????????O??O?????E?????@G???????@O???????@_??????@_????????
```

B More examples of non-uniform Šoltés' hypergraphs

Proposition 14. There exist infinitely many non-uniform Šoltés' hypergraphs with diameter 2.

Proof. For every $k \ge 2$, we construct a hypergraph H_k , obtained by taking a complete k-partite graph whose partition classes have size 2k+3, minus a 2-factor and adding a hyperedge for every partition class. Examples for H_2 and H_3 (there are multiple choices for the removed 2-factor) are depicted in Fig. 6, with the hyperedges for a partition class drawn with dashed lines, and the removed edges in red. The important observation is that both H and For every 2 vertices in $H \setminus v$, which do not both belong to the same hyperedge of size 2k+3 as v in H, the distance within $H \setminus v$ is unchanged. The distance between vertices of the same partition class, changes from 1 to 2. Now, we observe that $\sum_{u \in V \setminus v} d_{H_k}(u, v) = (n-1) + 2 = k(2k+3) + 1 = \binom{2k+2}{2}$, i.e., the sum of distances using v initially equals the number of distances (between the vertices sharing the same hyperedge of size 2k + 3 with v in H_k) that increase (by one) in $H_k \setminus v$.

Example 15. Let k be even and sufficiently large. First we take a graph G with vertex set $A \cup B$. Here $B = B_1 \cup B_2$ is a set with k+2 vertices, where the k-8 vertices from B_1 have k-4 neighbours in B and the 10 vertices in B_2 have k-3 neighbours in B, i.e., $\deg_{G[B]}(v) = k-4$ if $v \in B_1$ and $\deg_{G[B]}(v) = k-3$ if $v \in B_2$. Let $A = A_1 \cup A_2$ with $|A_1| = k-2$ and $|A_2| = 2$, such that $\deg_{G[A]}(v) = k-6$ if $v \in A_1$ and $\deg_{G[A]}(v) = k-7$ if $v \in A_2$.

In the bipartite graph G[A, B], the degree of vertices of A_1 have degree $\frac{k}{2} + 7$, the ones of A_2 $\frac{k}{2} + 8$, the vertices in B_1 degree $\frac{k}{2} + 6$ and the ones in B_2 degree $\frac{k}{2} + 5$.

Finally, we add hyperedges A and B, to form a hypergraph H with order n = 2k + 2.



Figure 6: Two Šoltés' hypergraphs with diameter 2.

The degree of the vertices in A is equal to $(k-6) + \frac{k}{2} + 7 + 1 = \frac{3k}{2} + 2$ and the vertices in B have degree $k - 4 + \frac{k}{2} + 6 + 1 = \frac{3k}{2} + 3$. Hence the graph is not regular (and thus not vertex-transitive), The minimum degree of H equals $\frac{3}{4}n + \frac{1}{2}$ and as a corollary, diam(H) = 2 is also immediate. Finally, we verify that H is a Šoltés' hypergraph.

For $v \in A_1$, $\sum_{u \in V \setminus v} d_H(u, v) = 2k + 1 + \frac{k}{2} - 5 = \frac{5k}{2} - 4$. The only distances that increase (by one) correspond with non-edges in A which do not contain v, there are $\frac{5k+2}{2} - 5 = \frac{5k}{2} - 4$. This implies that $W(H) = W(H \setminus v)$. The same holds true for $v \in A_2$, B_1 and B_2 respectively.