

Spectral characterization of the constant sign derivatives of Green's function related to two point boundary value conditions

Alberto Cabada, Lucía López-Somoza and Mouhcine Yousfi

¹ CITMAga, 15782, Santiago de Compostela, Galicia, Spain

² Departamento de Estatística, Análise Matemática e Optimización

Facultade de Matemáticas, Universidade de Santiago de Compostela, Spain.

alberto.cabada@usc.es; lucia.lopez.somoza@usc.es; yousfi.mouhcine@usc.es

June 4, 2024

Abstract

In this paper we will study the set of parameters in which certain partial derivatives of the Green's function, related to a n -order linear operator $T_n[M]$, depending on a real parameter M , coupled to different two-point boundary conditions, are of constant sign. We will do it without using the explicit expression of the Green's function. The constant sign interval will be characterized by the first eigenvalue related to suitable boundary conditions of the studied operator. As a consequence of the main result, we will be able to give sufficient conditions to ensure that the derivatives of Green's function cannot be nonpositive (nonnegative). These characterizations and the obtained results can be used to deduce the existence of solutions of nonlinear problems under additional conditions on the nonlinear part. To illustrate the obtained results, some examples are given.

Keywords: Green's function, Two-point boundary conditions, Spectral theory, Comparison results, Constant sign.

1 Introduction

Many papers in the literature have studied the qualitative properties of solutions of differential equations coupled with different boundary conditions. Several techniques have been used to ensure the existence, non-existence and multiplicity of solutions. Among them, stands out the method of lower and upper solutions, coupled with monotone iterative techniques which, as it is very well-known, are equivalent to the constant sign of the Green's function of the corresponding linear problem. Moreover the method of constructing suitable cones in Banach spaces where to apply the well-known Krasnoselskii's fixed-point theorem, see for instance

[7, 9, 10, 12] is a powerful tool to deduce the existence of positive solutions of the nonlinear problem. Such cones are constructed using constant sign properties of the Green's function and some of its derivatives.

To be concise, in this paper, we consider the following n -order linear differential equation

$$T_n[M] u(t) \equiv u^{(n)}(t) + M u(t) = 0, \quad t \in I := [a, b], \quad (1.1)$$

where $M \in \mathbb{R}$ a is real parameter, coupled to the two-point boundary conditions

$$u^{(\sigma_1)}(a) = \dots = u^{(\sigma_k)}(a) = 0, \quad (1.2)$$

$$u^{(\varepsilon_1)}(b) = \dots = u^{(\varepsilon_{n-k})}(b) = 0, \quad (1.3)$$

with $k \in \{1, \dots, n-1\}$ and the sets of indices $\{\sigma_1, \dots, \sigma_k\}, \{\varepsilon_1, \dots, \varepsilon_{n-k}\} \subset \{0, \dots, n-1\}$, satisfying that

$$0 \leq \sigma_1 < \sigma_2 < \dots < \sigma_k \leq n-1, \quad 0 \leq \varepsilon_1 < \varepsilon_2 < \dots < \varepsilon_{n-k} \leq n-1.$$

This paper is devoted to study the set of parameters M in which certain partial derivatives of the Green's function g_M , related to problem (1.1)–(1.3), are of constant sign without using the explicit expression of Green's function. It should be noted that, in general, it is not possible to obtain the explicit expression of the Green's function, specially for non constant coefficients.

For the particular case of equation (1.1), together with boundary conditions (1.2)–(1.3), the interval of parameters for which the Green's function has constant sign has been characterized in [5, Theorem 8.1]. Also for boundary conditions of the type $(k, n-k)$, i.e., $\sigma_j = j-1$ for all $j = 1, \dots, k$, and $\varepsilon_j = j-1$ for all $j = 1, \dots, n-k$ which are a particular case of the previous ones, a characterization of the constant sign intervals of the Green's function is given in [3]. Moreover and the calculation of the eigenvalues that characterizes these intervals up to $n = 8$ is obtained.

The main assumption is the disconjugacy character of the operator $T_n[M]$, which, as we know, is disconjugate for the value $M = 0$ (see [3]) at any arbitrary interval $[a, b]$. In the literature, sufficient conditions have been given to ensure the disconjugacy character of linear operators, we refer to [6, 8, 11, 13] and the references therein. A characterization of this property is proved in [5], in which the optimal extremes of the intervals are given as the eigenvalues of suitable $(k, n-k)$ problems.

It is in this context that the corresponding spectral theory arises: we will prove that the constant sign interval is characterized by the first eigenvalues of the studied operator, related to suitable boundary conditions.

The study starts from conditions (1.2)–(1.3) and constructs the conditions satisfied by the derivatives of g_M , defining the necessary hypotheses that we need to obtain the main results, and deducing the relationships between the adjoint spaces of the respective related spaces. This way, we will obtain results that characterize the constant sign interval of the corresponding derivatives.

The novelty of the work is that the results generalize [5, Theorem 8.1] for the particular operator $T_n[M]$ defined in (1.1). Also, as a consequence of these results, we will deduce some

cases in which the derivatives of the Green's function cannot be of constant sign. Furthermore, we will obtain a necessary condition that allow us to ensure that the corresponding derivatives are strictly positive (strictly negative) on the interior of the square of definition.

The paper is organized as follows: in a preliminary section 2 we introduce the fundamental concepts, the results and the main hypotheses that are needed in the development of the article. The next section is devoted to prove the results that characterize the constant sign of the derivatives of Green's function by distinguishing three cases. The main results are proved through spectral theory. This section includes also a corollary that provides sufficient conditions for derivatives to be nonpositive or nonnegative, a necessary condition to ensure the negative (positive) sign of certain derivatives with respect to t of g_M , and examples of application of our main results.

2 Preliminaries, hypotheses and main assumptions

In this section, we will make a survey of several results and properties that will be used along the paper. The main part of these results are proven in [5], and many of them are given in [3].

Let us consider the n^{th} -order linear differential equation (1.1) coupled to the two-point boundary conditions (1.2)–(1.3).

Definition 2.1. Let us say that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfy property (N_a) if

$$\sum_{\sigma_j < h} 1 + \sum_{\varepsilon_j < h} 1 \geq h, \quad \forall h \in \{1, \dots, n-1\}.$$

Notation 2.2. Let us denote

$$\alpha = \min \{i \in \{0, \dots, n-1\} : i \notin \{\sigma_1, \dots, \sigma_k\}\}, \quad (2.1)$$

$$\beta = \min \{i \in \{0, \dots, n-1\} : i \notin \{\varepsilon_1, \dots, \varepsilon_{n-k}\}\}. \quad (2.2)$$

We introduce the following set of functions related to the boundary conditions (1.2)–(1.3):

$$X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} = \left\{ u \in C^n(I) \mid u^{(\sigma_1)}(a) = \dots = u^{(\sigma_k)}(a) = u^{(\varepsilon_1)}(b) = \dots = u^{(\varepsilon_{n-k})}(b) = 0 \right\}.$$

Also, we define the following sets of functions:

$$\begin{aligned} X_{\{\sigma_1, \dots, \sigma_{k-1}\}|\alpha}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} &= \left\{ u \in C^n(I) \mid u^{(\sigma_1)}(a) = \dots = u^{(\sigma_{k-1})}(a) = 0, u^{(\alpha)}(a) = 0, u^{(\varepsilon_1)}(b) = \dots = u^{(\varepsilon_{n-k})}(b) = 0 \right\}, \\ X_{\{\sigma_1, \dots, \sigma_k\}|\alpha}^{\{\varepsilon_1, \dots, \varepsilon_{n-k-1}\}} &= \left\{ u \in C^n(I) \mid u^{(\sigma_1)}(a) = \dots = u^{(\sigma_k)}(a) = 0, u^{(\alpha)}(a) = 0, u^{(\varepsilon_1)}(b) = \dots = u^{(\varepsilon_{n-k-1})}(b) = 0 \right\}, \\ X_{\{\sigma_1, \dots, \sigma_{k-1}\}|\beta}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} &= \left\{ u \in C^n(I) \mid u^{(\sigma_1)}(a) = \dots = u^{(\sigma_{k-1})}(a) = 0, u^{(\varepsilon_1)}(b) = \dots = u^{(\varepsilon_{n-k})}(b) = 0, u^{(\beta)}(b) = 0 \right\}, \\ X_{\{\sigma_1, \dots, \sigma_k\}|\beta}^{\{\varepsilon_1, \dots, \varepsilon_{n-k-1}\}} &= \left\{ u \in C^n(I) \mid u^{(\sigma_1)}(a) = \dots = u^{(\sigma_k)}(a) = 0, u^{(\varepsilon_1)}(b) = \dots = u^{(\varepsilon_{n-k-1})}(b) = 0, u^{(\beta)}(b) = 0 \right\}. \end{aligned}$$

Definition 2.3. Let us say that the operator $T_n[M]$ satisfies the property (T_d) in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ if, and only if, there exists the following decomposition:

$$T_0 u(t) = u(t), \quad T_l u(t) = \frac{d}{dt} \left(\frac{T_{l-1} u(t)}{v_l(t)} \right), \quad l = 1, \dots, n, \quad t \in I, \quad (2.3)$$

where $v_l > 0$ on I , $v_l \in C^n(I)$ such that

$$T_n[M] u(t) = v_1(t) \dots v_n(t) T_n u(t), \quad t \in I,$$

and, moreover, such a decomposition satisfies, for every $u \in X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$:

$$\begin{aligned} T_{\sigma_1} u(a) &= \dots = T_{\sigma_k} u(a) = 0, \\ T_{\varepsilon_1} u(b) &= \dots = T_{\varepsilon_{n-k}} u(b) = 0. \end{aligned}$$

Remark 2.4. Note that decomposition (2.3) is not unique, it depends on the choice of v_k for $k = 1, \dots, n$. Throughout this work, for the operator $T_n[0] u(t) = u^{(n)}(t)$ we choose the following decomposition:

$$T_0 u(t) = u(t), \quad T_k u(t) = \frac{d}{dt} \left(\frac{T_{k-1} u(t)}{v_k(t)} \right), \quad k = 1, \dots, n, \quad t \in I,$$

where $v_k \equiv 1$ on I . That is, $T_k u(t) = u^{(k)}(t)$, $t \in I$. In particular:

$$T_n[0] u(t) = v_1(t) \dots v_n(t) T_n u(t), \quad t \in I,$$

and, moreover, this decomposition satisfies, for every $u \in X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$:

$$\begin{aligned} T_{\sigma_1} u(a) &= u^{(\sigma_1)}(a) = 0, \quad \dots, \quad T_{\sigma_k} u(a) = u^{(\sigma_k)}(a) = 0, \\ T_{\varepsilon_1} u(b) &= u^{(\varepsilon_1)}(b) = 0, \quad \dots, \quad T_{\varepsilon_{n-k}} u(b) = u^{(\varepsilon_{n-k})}(b) = 0, \end{aligned}$$

or which is the same $T_n[0]$ satisfies property (T_d) in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

Lemma 2.5. [5, Lemma 3.8] Let $\bar{M} \in \mathbb{R}$ be such that $T_n[\bar{M}]$ satisfies the property (T_d) in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. Then $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfy the property (N_a) if, and only if, $M = 0$ is not an eigenvalue of $T_n[\bar{M}]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

Let us denote by G_M the Green's matrix associated with the equivalent n -dimensional first order system associated to (1.1)–(1.3), which is given by the expression

$$G_M(t, s) = \begin{pmatrix} g_1(t, s) & g_2(t, s) & \dots & g_{n-1}(t, s) & g_M(t, s) \\ \frac{\partial}{\partial t} g_1(t, s) & \frac{\partial}{\partial t} g_2(t, s) & \dots & \frac{\partial}{\partial t} g_{n-1}(t, s) & \frac{\partial}{\partial t} g_M(t, s) \\ \vdots & \vdots & \dots & \vdots & \vdots \\ \frac{\partial^{n-1}}{\partial t^{n-1}} g_1(t, s) & \frac{\partial^{n-1}}{\partial t^{n-1}} g_2(t, s) & \dots & \frac{\partial^{n-1}}{\partial t^{n-1}} g_{n-1}(t, s) & \frac{\partial^{n-1}}{\partial t^{n-1}} g_M(t, s) \end{pmatrix}, \quad (2.4)$$

where g_M is the scalar Green's function related to operator $T_n[M]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ (see [1, Section 1.4] for details). Moreover, by the axiomatic definition of G_M , we have that for all $s \in (a, b)$, the following equality holds

$$\frac{\partial}{\partial t} G_M(t, s) = A G_M(t, s), \quad \text{for all } t \in I \setminus \{s\}, \quad (2.5)$$

where $A = \left(\begin{array}{c|c} 0 & I_{n-1} \\ \hline -M & 0 \end{array} \right)$, with I_{n-1} the $(n-1)$ dimensional identity matrix, denotes the matrix that defines the n -dimensional system equivalent to our problem (1.1)–(1.3) and

$$B G_M(a, s) + C G_M(b, s) = 0, \quad \text{for all } s \in (a, b), \quad (2.6)$$

with $B, C \in \mathcal{M}_{n \times n}$, defined by $b_{j, 1+\sigma_j} = 1$ for $j = 1, \dots, k$ and $c_{j+k, 1+\varepsilon_j} = 1$ for $j = 1, \dots, n-k$; otherwise, $b_{ij} = 0$ and $c_{ij} = 0$.

In particular, it is easy to see, using the expressions (2.4) and (2.5) (see [1, Section 1.4] for details), that $g_M \in C^{n-2}(I \times I)$. Moreover, it is a C^n function on the triangles $a \leq s < t \leq b$ and $a \leq t < s \leq b$, it satisfies, as a function of t for any $s \in (a, b)$ fixed, the two-point boundary value conditions (1.2)–(1.3) and solves equation (1.1) for all $t \in I \setminus \{s\}$.

Moreover, for any $t \in (a, b)$ it satisfies

$$\frac{\partial^{n-1}}{\partial t^{n-1}} g_M(t^+, t) = \frac{\partial^{n-1}}{\partial t^{n-1}} g_M(t^-, t) + 1. \quad (2.7)$$

Studying the matrix Green's function G_M , it is proved in [3, page 13] that, for this particular case, $g_{n-j}(t, s)$ may be expressed as a function of $g_M(t, s)$ for all $j = 1, \dots, n-1$ as follows:

$$g_{n-j}(t, s) = (-1)^j \frac{\partial^j}{\partial s^j} g_M(t, s). \quad (2.8)$$

The adjoint operator of $T_n[M]$ follows the expression

$$T_n^*[M]v(t) \equiv (-1)^n v^{(n)}(t) + M v(t), \quad (2.9)$$

defined on the domain

$$D(T_n^*[M]) = \left\{ v \in C^n(I) \mid \sum_{i=0}^{n-1} (-1)^{n-1-i} v^{(n-1-i)}(b) u^{(i)}(b) = \sum_{i=0}^{n-1} (-1)^{n-1-i} v^{(n-1-i)}(a) u^{(i)}(a) \right\},$$

for all $u \in D(T_n[M])$.

We denote by $g_M^*(t, s)$ the Green's function related to operator $T_n^*[M]$. Furthermore, the following equality is satisfied (see [1, section 1.4])

$$g_M^*(t, s) = g_M(s, t), \quad (t, s) \in I \times I. \quad (2.10)$$

In such a case, if we define the following operator

$$\widehat{T}_n[(-1)^n M] := (-1)^n T_n^*[M], \quad (2.11)$$

we have, according to the above equality, that

$$\widehat{g}_{(-1)^n M}(t, s) = (-1)^n g_M^*(t, s) = (-1)^n g_M(s, t), \quad (2.12)$$

where $\widehat{g}_{(-1)^n M}(t, s)$ is the scalar Green's function related to operator $\widehat{T}_n[(-1)^n M]$ in $D(T_n^*[M])$.

As a particular case of the obtained results in [5, section 4], we can deduce that

$$D(T_n^*[M]) \equiv X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} = X_{\{\tau_1, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k\}},$$

being $\{\delta_1, \dots, \delta_k\}, \{\tau_1, \dots, \tau_{n-k}\} \subset \{0, \dots, n-1\}$, such that $\delta_i < \delta_{i+1}$ and $\tau_j < \tau_{j+1}$, for $i = 1, \dots, k-1$ and $j = 1, \dots, n-k-1$, satisfying:

$$\begin{aligned} \{\sigma_1, \dots, \sigma_k, n-1-\tau_1, \dots, n-1-\tau_{n-k}\} &= \{0, \dots, n-1\}, \\ \{\varepsilon_1, \dots, \varepsilon_{n-k}, n-1-\delta_1, \dots, n-1-\delta_k\} &= \{0, \dots, n-1\}. \end{aligned}$$

Notation 2.6. Let us define η, γ in the following way:

$$\eta = \min \{i \in \{0, \dots, n-1\} : i \notin \{\tau_1, \dots, \tau_{n-k}\}\}, \quad (2.13)$$

$$\gamma = \min \{i \in \{0, \dots, n-1\} : i \notin \{\delta_1, \dots, \delta_k\}\}. \quad (2.14)$$

Analogously, let us denote by $\widehat{G}(t, s)$ the Green's matrix associated with the equivalent n -dimensional problem of $\widehat{T}_n[(-1)^n M]v(t) = 0$, $v \in X_{\{\tau_1, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k\}}$ given by

$$\widehat{G}(t, s) = \begin{pmatrix} \widehat{g}_1(t, s) & \cdots & \widehat{g}_{n-1}(t, s) & \widehat{g}_{(-1)^n M}(t, s) \\ \frac{\partial}{\partial t} \widehat{g}_1(t, s) & \cdots & \frac{\partial}{\partial t} \widehat{g}_{n-1}(t, s) & \frac{\partial}{\partial t} \widehat{g}_{(-1)^n M}(t, s) \\ \vdots & \cdots & \vdots & \vdots \\ \frac{\partial^{n-1}}{\partial t^{n-1}} \widehat{g}_1(t, s) & \cdots & \frac{\partial^{n-1}}{\partial t^{n-1}} \widehat{g}_{n-1}(t, s) & \frac{\partial^{n-1}}{\partial t^{n-1}} \widehat{g}_{(-1)^n M}(t, s) \end{pmatrix}, \quad (2.15)$$

where $\widehat{g}_{(-1)^n M}(t, s)$ is the scalar Green's function related to operator $\widehat{T}_n[(-1)^n M]$ in $X_{\{\tau_1, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k\}}$.

Similarly to (2.8) we can deduce that

$$\widehat{g}_{n-j}(t, s) = (-1)^j \frac{\partial^j}{\partial s^j} \widehat{g}_{(-1)^n M}(t, s). \quad (2.16)$$

By definition, $\widehat{G}(t, s)$ satisfies the equality

$$\widehat{B} \widehat{G}(a, s) + \widehat{C} \widehat{G}(b, s) = 0, \quad \text{for all } s \in (a, b), \quad (2.17)$$

where $\widehat{B}, \widehat{C} \in \mathcal{M}_{n \times n}$, defined as $(\widehat{B})_{i, \tau_i+1} = 1$ for $i = 1, \dots, n-k$ and $(\widehat{C})_{j+n-k, 1+\delta_j} = 1$ for $j = 1, \dots, k$; otherwise, $(\widehat{B})_{ij} = 0$ and $(\widehat{C})_{ij} = 0$.

Remark 2.7. From the previous definitions, as it is pointed out in [5, Remarks 4.1 and 4.2] we have that $\alpha = n - 1 - \tau_{n-k}$, $\beta = n - 1 - \delta_k$, $\eta = n - 1 - \sigma_k$ and $\gamma = n - 1 - \varepsilon_{n-k}$.

Example 2.8. Let us consider operator $T_4[M]$ coupled with the boundary conditions

$$u(0) = u'(0) = u''(0) = u''(1) = 0.$$

In this case, $\{\sigma_1, \sigma_2, \sigma_3\} = \{0, 1, 2\}$, $\{\varepsilon_1\} = \{2\}$, $\{\tau_1\} = \{0\}$ and $\{\delta_1, \delta_2, \delta_3\} = \{0, 2, 3\}$. Thus, we deduce that $X_{\{0,1,2\}}^{*\{2\}} = X_{\{0\}}^{\{0,2,3\}}$. Moreover, $\alpha = 3$, $\beta = 0$, $\eta = 1$ and $\gamma = 1$.

Definition 2.9. Operator $T_n[M]$ is said to be inverse positive (negative) on $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ if every $u \in X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ such that $T_n[M]u \geq 0$ on I , satisfies $u \geq 0$ ($u \leq 0$) on I .

The above definition is equivalent to the constant sign of Green's function g_M related to $T_n[M]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ as it is shown in the following result whose proof is analogous the one made in [5, Theorem 2.9].

Theorem 2.10. Operator $T_n[M]$ is inverse positive (negative) in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ if, and only if, the Green's function related to problem (1.1)–(1.3) is non-negative (non-positive) on its square of definition.

Definition 2.11. Operator $T_n[M]$ is said to be strongly inverse positive (negative) in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ if every $u \in X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ such that $T_n[M]u \not\geq 0$ on I , satisfies that $u > 0$ (< 0) on (a, b) and, moreover, $u^{(\alpha)}(a) > 0$ (< 0) and $(-1)^\beta u^{(\beta)}(b) > 0$ (< 0), where α and β are defined in (2.1) and (2.2), respectively.

Analogously to Theorem 2.10, the following characterization can be shown:

Theorem 2.12. Operator $T_n[M]$ is strongly inverse positive (negative) in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ if, and only if, the Green's function related to problem (1.1)–(1.3) satisfies the following properties:

- $g_M(t, s) > 0$ (< 0) a.e. on (a, b) .
- $\frac{\partial^\alpha}{\partial t^\alpha} g_M(t, s) > 0$ (< 0) for a.e. $s \in (a, b)$.
- $(-1)^\beta \frac{\partial^\beta}{\partial t^\beta} g_M(t, s) > 0$ (< 0) for a.e. $s \in (a, b)$.

Let $u \in X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ be such that $T_n[M]u(t) = 0$, $t \in I$. Then, differentiating equation (1.1) with respect to t , as a direct consequence of equations (2.4) and (2.5) we deduce that, for any $s \in (a, b)$ given, the q -th derivative $v_s^q[M](t) := \frac{\partial^q}{\partial t^q} g_M(t, s)$, with $1 \leq q \leq n - 1$, satisfies the equation $T_n[M]v_s^q[M](t) = 0$, for all $t \in I \setminus \{s\}$, together with the new boundary conditions

$$(v_s^q)^{(\mu_1^q)}[M](a) = \dots = (v_s^q)^{(\mu_k^q)}[M](a) = 0, \quad (2.18)$$

$$(v_s^q)^{(\rho_1^q)}[M](b) = \dots = (v_s^q)^{(\rho_{n-k}^q)}[M](b) = 0. \quad (2.19)$$

Here, the sets of indices $\{\mu_1^q, \dots, \mu_k^q\}, \{\rho_1^q, \dots, \rho_{n-k}^q\} \subset \{0, \dots, n-1\}$, satisfy that

$$0 \leq \mu_1^q < \mu_2^q < \dots < \mu_k^q \leq n-1, \quad 0 \leq \rho_1^q < \rho_2^q < \dots < \rho_{n-k}^q \leq n-1,$$

and are defined in the following way:

- If $A_q = \{i \in \{1, \dots, k\} / \sigma_i \geq q\} \neq \emptyset$, then

$$\mu_1^q = \sigma_j - q, \mu_2^q = \sigma_{j+1} - q, \dots, \mu_{k-(j-1)}^q = \sigma_k - q, \mu_{k-(j-2)}^q = \sigma_1 - q + n, \dots, \mu_k^q = \sigma_{j-1} - q + n,$$

with

$$j = \min A_q \geq 1. \quad (2.20)$$

- If $A_q = \emptyset$, then

$$\mu_i^q = \sigma_i - q + n, \quad i = 1, \dots, k.$$

- If $B_q = \{i \in \{1, \dots, n-k\} / \varepsilon_i \geq q\} \neq \emptyset$, then

$$\rho_1^q = \varepsilon_r - q, \rho_2^q = \varepsilon_{r+1} - q, \dots, \rho_{n-k-(r-1)}^q = \varepsilon_{n-k} - q, \rho_{n-k-(r-2)}^q = \varepsilon_1 - q + n, \dots, \\ \rho_{n-k}^q = \varepsilon_{r-1} - q + n,$$

with

$$r = \min B_q \geq 1. \quad (2.21)$$

- If $B_q = \emptyset$, then

$$\rho_i^q = \varepsilon_i - q + n, \quad i = 1, \dots, n-k.$$

Remark 2.13. The above process of calculating the boundary conditions of $v_s^q[M]$ is valid for both $M \neq 0$ and $M = 0$.

Taking into account previous computations, we may define ϕ as the function which maps the space $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ into the space of the boundary conditions that satisfies the first derivative, that is, $X_{\{\mu_1^1, \dots, \mu_k^1\}}^{\{\rho_1^1, \dots, \rho_{n-k}^1\}}$. This way, we can extend this definition to the q -th derivative as

$$\phi^q = \phi \circ \dots \circ \phi : X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} \longrightarrow X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}.$$

In particular, ϕ^n is the identity function.

Example 2.14. Consider the operator $T_5[M]$ defined in $X_{\{0,2,4\}}^{\{0,2\}}$. Applying the above definition of the spaces of the derivatives we obtain the following sequence of spaces:

$$X_{\{0,2,4\}}^{\{0,2\}} \xrightarrow{\phi} X_{\{1,3,4\}}^{\{1,4\}} \xrightarrow{\phi} X_{\{0,2,3\}}^{\{0,3\}} \xrightarrow{\phi} X_{\{1,2,4\}}^{\{2,4\}} \xrightarrow{\phi} X_{\{0,1,3\}}^{\{1,3\}} \xrightarrow{\phi} X_{\{0,2,4\}}^{\{0,2\}}.$$

Analogously to the Notation 2.2, let us define α^q, β^q as follows:

$$\alpha^q = \min \{i \in \{0, \dots, n-1\} : i \notin \{\mu_1^q, \dots, \mu_k^q\}\}, \quad (2.22)$$

$$\beta^q = \min \{i \in \{0, \dots, n-1\} : i \notin \{\rho_1^q, \dots, \rho_{n-k}^q\}\}. \quad (2.23)$$

Remark 2.15. It is important to point out that for any $q \geq 1$, the function $v_s^q[M]$ is not the Green's function related to the operator $T_n[M]$ in $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$. This is due to the fact that $v_s^q[M] \in C^{n-2-q}(I \times I)$.

Remark 2.16. The operator $T_n[0]$ satisfies the property (T_d) in $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$. From Remark 2.4, we know that operator $T_n[0]u(t) = u^{(n)}(t)$ satisfies the following decomposition:

$$T_0 u(t) = u(t), \quad T_k u(t) = \frac{d}{dt} \left(\frac{T_{k-1} u(t)}{v_k(t)} \right), \quad k = 1, \dots, n,$$

where $v_k \equiv 1$ on I and

$$T_n[0] u(t) = v_1(t) \dots v_n(t) T_n u(t), \quad t \in I,$$

that is, $T_k u(t) = u^{(k)}(t)$.

Moreover, this decomposition satisfies, for every $u \in X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$:

$$\begin{aligned} T_{\mu_1^q} u(a) = u^{(\mu_1^q)}(a) = 0, \quad \dots, \quad T_{\mu_k^q} u(a) = u^{(\mu_k^q)}(a) = 0, \\ T_{\rho_1^q} u(b) = u^{(\rho_1^q)}(b) = 0, \quad \dots, \quad T_{\rho_{n-k}^q} u(b) = u^{(\rho_{n-k}^q)}(b) = 0. \end{aligned}$$

We consider the general expression of the solution of equation

$$T_n[0] u(t) + \lambda u(t) \equiv u^{(n)}(t) + \lambda u(t) = 0, \quad t \in I,$$

to obtain the different eigenvalues.

Next we present a result concerning the spectrum of the spaces $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$.

Lemma 2.17. $\bar{\lambda} \neq 0$ is an eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ if and only if $\bar{\lambda}$ is an eigenvalue of $T_n[0]$ in $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ for all $q \in \{1, \dots, n-1\}$.

Proof. Suppose that $\bar{\lambda} \neq 0$ is an eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. So, there is $u \neq 0$ in I such that $u^{(n)}(t) + \bar{\lambda} u(t) = 0$, $t \in I$ and $u \in X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. Since $u \neq 0$ in I we have that $u^{(n)} \neq 0$ in I . Therefore, $u^{(j)} \neq 0$ in I for all $j \in \{1, \dots, n-1\}$.

Taking into account that $T_n[0]u^{(j)}(t) = u^{(j+n)}(t) = -\bar{\lambda}u^{(j)}(t)$ for all $t \in I$ and $j \in \mathbb{N}$, we have that $u^{(j)}$ is an eigenvector related to the eigenvalue $\bar{\lambda}$ such that $u^{(j)} \in X_{\{\mu_1^j, \dots, \mu_k^j\}}^{\{\rho_1^j, \dots, \rho_{n-k}^j\}}$ for all $j \in \{1, \dots, n-1\}$.

For $j = n$, we have that $\mu_i^n = \sigma_i$ for all $i \in \{1, \dots, k\}$ and $\rho_l^n = \varepsilon_l$ for all $l \in \{1, \dots, n-k\}$. Therefore, the result holds. \square

Remark 2.18. As a direct consequence of Remark 2.4, we have, from Lemma 2.5, that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) if and only if $M = 0$ is not an eigenvalue of $u^{(n)}$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

Remark 2.19. Note that if $\lambda = 0$ is not an eigenvalue of $T_n[0]$ in $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$, then the spaces $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ and $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ have the same spectrum. However, it is very important to point out that $\lambda = 0$ can appear as an eigenvalue in some space $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$, as it is shown in the following example.

Example 2.20. Let us consider again the Example 2.14. As we know, the following sequence of spaces is fulfilled:

$$X_{\{0,2,4\}}^{\{0,2\}} \xrightarrow{\phi} X_{\{1,3,4\}}^{\{1,4\}} \xrightarrow{\phi} X_{\{0,2,3\}}^{\{0,3\}} \xrightarrow{\phi} X_{\{1,2,4\}}^{\{2,4\}} \xrightarrow{\phi} X_{\{0,1,3\}}^{\{1,3\}} \xrightarrow{\phi} X_{\{0,2,4\}}^{\{0,2\}}.$$

In this case, we have that $\lambda = 0$ is not an eigenvalue of $T_5[0]$ in $X_{\{0,2,4\}}^{\{0,2\}}$, $X_{\{0,2,3\}}^{\{0,3\}}$ and $X_{\{0,1,3\}}^{\{1,3\}}$, while $\lambda = 0$ is an eigenvalue of $T_5[0]$ in $X_{\{1,3,4\}}^{\{1,4\}}$ and $X_{\{1,2,4\}}^{\{2,4\}}$.

Now, we will show how to determine the adjoint space of $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ from the adjoint space of $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. We arrive to the next result.

Theorem 2.21. *The adjoint space of $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ is given by $X_{\{\tau_1^q, \dots, \tau_{n-k}^q\}}^{\{\delta_1^q, \dots, \delta_k^q\}}$, where the indices $\{\tau_1^q, \dots, \tau_{n-k}^q\}, \{\delta_1^q, \dots, \delta_k^q\} \subset \{0, \dots, n-1\}$ satisfy*

$$0 \leq \tau_1^q < \tau_2^q < \dots < \tau_{n-k}^q \leq n-1, \quad 0 \leq \delta_1^q < \delta_2^q < \dots < \delta_k^q \leq n-1,$$

and are defined as follows:

- If $C_q = \{i \in \{1, \dots, n-k\} / \tau_i + q \leq n-1\} \neq \emptyset$, then

$$\tau_1^q = \tau_{l+1} + q - n, \tau_2^q = \tau_{l+2} + q - n, \dots, \tau_{n-k-l}^q = \tau_{n-k} + q - n, \tau_{n-k-l+1}^q = \tau_1 + q, \dots, \tau_{n-k}^q = \tau_l + q,$$

with

$$l = \max C_q. \tag{2.24}$$

- If $C_q = \emptyset$, then

$$\tau_i^q = \tau_i + q - n, \quad i = 1, \dots, n-k.$$

- If $D_q = \{i \in \{1, \dots, k\} / \delta_i + q \leq n-1\} \neq \emptyset$, then

$$\delta_1^q = \delta_{p+1} + q - n, \delta_2^q = \delta_{p+2} + q - n, \dots, \delta_{k-p}^q = \delta_k + q - n, \delta_{k-p+1}^q = \delta_1 + q, \dots, \delta_k^q = \delta_p + q,$$

with

$$p = \max D_q. \tag{2.25}$$

- If $D_q = \emptyset$, then

$$\delta_i^q = \delta_i + q - n, \quad i = 1, \dots, k.$$

As a consequence, the following diagram

$$\begin{array}{ccc} X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} & \xrightarrow{\phi^q} & X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}} \\ \downarrow * & & \downarrow * \\ X_{\{\tau_1, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k\}} & \xleftarrow{\phi^q} & X_{\{\tau_1^q, \dots, \tau_{n-k}^q\}}^{\{\delta_1^q, \dots, \delta_k^q\}} \end{array} \quad (2.26)$$

is commutative.

Proof. We do the proof only for τ_i^q . For δ_i^q , the proof is done in a similar way. To prove it, we start from the definition of τ_i , we have that $\tau_i < \tau_{i+1}$ for all $i \in \{1, \dots, n-k-1\}$ and

$$\{\sigma_1, \dots, \sigma_k, n-1-\tau_{n-k}, \dots, n-1-\tau_1\} \equiv \{0, \dots, n-1\}.$$

Taking into account the definition of $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ we have that

$$\{\sigma_1, \dots, \sigma_k, n-1-\tau_{n-k}, \dots, n-1-\tau_1\} \xrightarrow{\phi^q} \{\mu_1^q, \dots, \mu_k^q, n-1-\tau_{n-k}^q, \dots, n-1-\tau_1^q\},$$

with $\{\mu_1^q, \dots, \mu_k^q, n-1-\tau_{n-k}^q, \dots, n-1-\tau_1^q\} \equiv \{0, \dots, n-1\}$ where τ_i^q has to fulfill that

$$0 \leq \tau_1^q < \tau_2^q < \dots < \tau_{n-k}^q \leq n-1.$$

To calculate them, we distinguish two cases:

- 1) If $C_q \neq \emptyset$, then:

- If $n-1-\tau_i \geq q$ (i.e, $i \leq l$), $n-1-\tau_i$ becomes $n-1-\tau_i-q$ by applying ϕ^q .
- If $n-1-\tau_i < q$ (i.e, $i > l$), $n-1-\tau_i$ becomes $n-1-\tau_i-q+n$ by applying ϕ^q .

Reordering the elements, we obtain that:

$$\begin{aligned} 0 &\leq n-1-\tau_l-q < n-1-\tau_{l-1}-q < \dots < n-1-\tau_1-q \\ &< n-1-\tau_{n-k}-q+n < \dots < n-1-\tau_{l+1}-q+n \leq n-1, \end{aligned}$$

or, equivalently,

$$0 \leq \tau_{l+1}+q-n < \tau_{l+2}+q-n < \dots < \tau_{n-k}+q-n < \tau_1+q < \dots < \tau_l+q \leq n-1.$$

Taking

$$\begin{aligned} \tau_1^q &= \tau_{l+1}+q-n, \tau_2^q = \tau_{l+2}+q-n, \dots, \tau_{n-k}^q = \tau_{n-k}+q-n, \tau_{n-k-l+1}^q = \tau_1+q, \dots, \\ \tau_{n-k}^q &= \tau_l+q, \end{aligned}$$

the first statement yields.

2) If $C_q = \emptyset$, then $n - 1 - \tau_i$ becomes $n - 1 - \tau_i - q + n$ by applying ϕ^q and we obtain that

$$0 \leq n - 1 - \tau_{n-k} - q + n < \cdots < n - 1 - \tau_2 - q + n < n - 1 - \tau_1 - q + n \leq n - 1,$$

and, reordering, we infer that

$$0 \leq \tau_1 + q - n < \tau_2 + q - n < \cdots < \tau_{n-k} + q - n \leq n - 1.$$

Taking $\tau_i^q = \tau_i + q - n$, $i = 1, \dots, n - k$, the second statement yields.

Therefore, $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{*\{\rho_1^q, \dots, \rho_{n-k}^q\}} = X_{\{\tau_1^q, \dots, \tau_{n-k}^q\}}^{\{\delta_1^q, \dots, \delta_k^q\}}$. □

Remark 2.22. Note that, unlike the direct arrows between the leading spaces, the arrows between the adjoint spaces are in the inverse direction.

Example 2.23. Consider the operator $T_4[M]$ coupled with the boundary conditions

$$u(a) = u''(a) = u'(b) = u''(b).$$

In this case, $u \in X_{\{0,2\}}^{\{1,2\}}$ and $X_{\{0,2\}}^{*\{1,2\}} = X_{\{0,2\}}^{\{0,3\}}$. Using the diagram (2.26), we get that

$$\begin{array}{ccccccc} X_{\{0,2\}}^{\{1,2\}} & \xrightarrow{\phi} & X_{\{1,3\}}^{\{0,1\}} & \xrightarrow{\phi} & X_{\{0,2\}}^{\{0,3\}} & \xrightarrow{\phi} & X_{\{1,3\}}^{\{2,3\}} \\ * \downarrow & & \downarrow * & & \downarrow * & & \downarrow * \\ X_{\{0,2\}}^{\{0,3\}} & \xleftarrow{\phi} & X_{\{1,3\}}^{\{0,1\}} & \xleftarrow{\phi} & X_{\{0,2\}}^{\{1,2\}} & \xleftarrow{\phi} & X_{\{1,3\}}^{\{2,3\}} \end{array}$$

Using the fact that the adjoint operators have the same spectrum as the original one, from Lemma 2.17, we arrive at the following result.

Lemma 2.24. $\lambda^* \neq 0$ is an eigenvalue of $T_n[0]$ in $X_{\{\tau_1, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k\}}$ if and only if λ^* is an eigenvalue of $T_n[0]$ in $X_{\{\tau_1^q, \dots, \tau_{n-k}^q\}}^{\{\delta_1^q, \dots, \delta_k^q\}}$ for all $q \in \{0, \dots, n - 1\}$. Moreover, for each fixed $q \in \{0, \dots, n - 1\}$, $\bar{\lambda} \in \mathbb{R}$ is an eigenvalue of $T_n[0]$ in $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ if and only if $\bar{\lambda}$ is an eigenvalue of $T_n[0]$ in $X_{\{\tau_1^q, \dots, \tau_{n-k}^q\}}^{\{\delta_1^q, \dots, \delta_k^q\}}$.

Following the line of Definition 2.11 and the characterization of Theorem 2.12, we introduce the following concept for $v_s^q[M]$.

Definition 2.25. Let $q \in \{0, \dots, n - 1\}$ be fixed. We said that the function $v_s^q[M]$ is strongly positive (strongly negative) on $I \times I$ if it satisfies the following properties:

- $v_s^q[M] > 0$ (< 0) a.e. on (a, b) .
- $(v_s^q)^{(\alpha^q)}[M](a) > 0$ (< 0) for a.e. $s \in (a, b)$, with α^q defined in (2.22).

- $(-1)^{\beta^q} (v_s^q)^{(\beta^q)} [M](b) > 0 (< 0)$ for a.e. $s \in (a, b)$, with β^q defined in (2.23).

Let us consider the following two conditions on $g_M(t, s)$ introduced in [1, pages 78 and 86] as follows:

- (P_g) Suppose that there is a continuous function $\phi(t) > 0$ for all $t \in (a, b)$ and $k_1, k_2 \in L^1(I)$, such that $0 < k_1(s) < k_2(s)$ for a.e. $s \in I$, satisfying

$$\phi(t) k_1(s) \leq g_M(t, s) \leq \phi(t) k_2(s), \quad \text{for a.e. } (t, s) \in I \times I.$$

- (N_g) Suppose that there is a continuous function $\phi(t) > 0$ for all $t \in (a, b)$ and $k_1, k_2 \in L^1(I)$, such that $k_1(s) < k_2(s) < 0$ for a.e. $s \in I$, satisfying

$$\phi(t) k_1(s) \leq g_M(t, s) \leq \phi(t) k_2(s), \quad \text{for a.e. } (t, s) \in I \times I.$$

As a particular case of [5, Theorem 5.1], the following result is attained.

Theorem 2.26. *Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a). Then the following properties are fulfilled:*

- *If $n - k$ is even, then $T_n[0]$ is strongly inverse positive in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ and, moreover, the related Green's function, $g_0(t, s)$, satisfies (P_g).*
- *If $n - k$ is odd, then $T_n[0]$ is strongly inverse negative in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ and, moreover, the related Green's function, $g_0(t, s)$, satisfies (N_g).*

Next, for any $q \in \{1, \dots, n-1\}$ we determine the positive (negative) sign of $v_s^q(t) := v_s^q[0](t)$ on the interval I .

Let us define $\{\sigma_1^q, \dots, \sigma_{c_q}^q\}, \{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\} \subset \{0, \dots, n-1\}$, in the following way:

$$\{\sigma_1^q, \dots, \sigma_{c_q}^q\} = \{\sigma_1 - q, \dots, \sigma_k - q\} \cap \{0, \dots, n - q - 1\}, \quad (2.27)$$

$$\{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\} = \{\varepsilon_1 - q, \dots, \varepsilon_{n-k} - q\} \cap \{0, \dots, n - q - 1\}. \quad (2.28)$$

If $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies the condition (N_a) then $c_q + d_q \leq n - q$. Moreover, if $c_q + d_q < n - q$ then $\{\sigma_1^q, \dots, \sigma_{c_q}^q\} - \{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\}$ does not satisfy (N_a), and in this case $\lambda = 0$ is an eigenvalue of $X_{\{\sigma_1^q, \dots, \sigma_{c_q}^q\}}^{\{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\}}$ (we cannot study in this case the constant sign of its corresponding derivative $v_s^q[M]$).

Remark 2.27. Taking into account that function $v_s^q(t) = \frac{\partial^q}{\partial t^q} g_0(t, s)$ is the Green's function related to operator $T_{n-q}[0]$ in the space $X_{\{\sigma_1^q, \dots, \sigma_{c_q}^q\}}^{\{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\}}$, then operator $T_{n-q}[0]$ is strongly inverse positive (negative) in the space $X_{\{\sigma_1^q, \dots, \sigma_{c_q}^q\}}^{\{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\}}$ if and only if it satisfies the Definition 2.11 with the same α^q and β^q since

$$\mu_1^q = \sigma_1^q, \mu_2^q = \sigma_2^q, \dots, \mu_{k-(j-1)}^q = \sigma_{c_q}^q,$$

and

$$\rho_1^q = \varepsilon_1^q, \rho_2^q = \varepsilon_2^q, \dots, \rho_{n-k-(r-1)}^q = \varepsilon_{d_q}^q.$$

Theorem 2.28. Let $q \in \{1, \dots, n-1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) and $c_q + d_q = n - q$. The following properties are fulfilled:

- (a) If $c_q \geq 1$ and $d_q \geq 1$, then v_s^q is strongly positive on $I \times I$ if $n - q - c_q$ is even and v_s^q is strongly negative on $I \times I$ if $n - q - c_q$ is odd.
- (b) If $c_q = n - q$ and $d_q = 0$, then $v_s^q(t) > 0, \dots, v_s^{n-1}(t) > 0$ in $(s, b]$, $v_s^q(t) = \dots = v_s^{n-2}(t) = 0$ in $[a, s]$ and $v_s^{n-1}(t) = 0$ in $[a, s)$.
- (c) If $c_q = 0$ and $d_q = n - q$, then $v_s^q(t) > 0$ in $[a, s)$ if $q = n - l$ with l even and $v_s^q(t) < 0$ in $[a, s)$ if $q = n - l$ with l odd. Moreover, $v_s^q(t) = \dots = v_s^{n-2}(t) = 0$ in $[s, b]$ and $v_s^{n-1}(t) = 0$ in $(s, b]$.

Proof. (a) In this case, it is easy to verify that the space $X_{\{\sigma_1^q, \dots, \sigma_{c_q}^q\}}^{\{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\}}$ satisfies the condition (N_a) since

$$\sum_{\sigma_j - q < h} 1 + \sum_{\varepsilon_j - q < h} 1 \geq h, \quad \forall h \in \{1, \dots, n - q - 1\}.$$

So, taking into account the fact that v_s^q is the Green's function related to operator $T_{n-q}[0]$ in the space $X_{\{\sigma_1^q, \dots, \sigma_{c_q}^q\}}^{\{\varepsilon_1^q, \dots, \varepsilon_{d_q}^q\}}$, we apply Theorem 2.26, and we have proved the first statement.

- (b) Since $v_s^{(n)}(t) = 0$ for all $t \neq 0$, we have, by (2.7) and $v_s^{(n-1)}(a) = 0$, that $\frac{\partial^{n-1}}{\partial t^{n-1}} g_0(t, s) = 0$ if $t < s$ and $\frac{\partial^{n-1}}{\partial t^{n-1}} g_0(t, s) = 1$ if $t > s$.

Since $v_s^h(a) = 0$ for all $q \leq h \leq n-2$, and $g_0 \in C^{n-2}(I \times I)$, we deduce that $\frac{\partial^h}{\partial t^h} g_0(t, s) = 0$ if $t < s$ and $\frac{\partial^h}{\partial t^h} g_0(t, s) > 0$ if $t > s$. So, we conclude the proof of second claim.

- (c) In this case arguing as in the previous case, since $v_s^{(n)}(b) = 0$, we have that $\frac{\partial^{n-1}}{\partial t^{n-1}} g_0(t, s) = 0$ if $t > s$ and $\frac{\partial^{n-1}}{\partial t^{n-1}} g_0(t, s) = -1$ if $t < s$.

As consequence, since $v_s^h(b) = 0$ for all $q \leq h \leq n-2$, and $g_0 \in C^{n-2}(I \times I)$, we deduce, by recurrence, that $\frac{\partial^h}{\partial t^h} g_0(t, s) = 0$ for all $t > s$ and $(-1)^{n-h} \frac{\partial^h}{\partial t^h} g_0(t, s) > 0$ for all $t < s$. Thus, the third assertion holds. □

Let $q \in \{0, \dots, n-1\}$ be fixed. Suppose that $(-1)^{n-q-c_q} v_s^q$ is strongly positive on $I \times I$. Then, for each $s \in (a, b)$, we obtain the following limits:

$$\begin{aligned} \ell_1^q(s) &:= \lim_{t \rightarrow a^+} \frac{(-1)^{n-q-c_q} v_s^q(t)}{(t-a)^{\alpha^q} (b-t)^{\beta^q}} = \frac{(-1)^{n-q-c_q} (v_s^q)^{(\alpha^q)}(a)}{\alpha^q! (b-a)^{\beta^q}}, \\ \ell_2^q(s) &:= \lim_{t \rightarrow b^-} \frac{(-1)^{n-q-c_q} v_s^q(t)}{(t-a)^{\alpha^q} (b-t)^{\beta^q}} = \frac{(-1)^{n-q-c_q-\beta^q} (v_s^q)^{(\beta^q)}(b)}{\beta^q! (b-a)^{\alpha^q}}. \end{aligned}$$

For each $s \in (a, b)$, let us consider the following function defined on I by

$$\tilde{u}_s^q(t) = \begin{cases} \ell_1^q(s), & t = a, \\ \frac{(-1)^{n-q-c_q} v_s^q(t)}{(t-a)^{\alpha_q} (b-t)^{\beta_q}}, & t \in (a, b), \\ \ell_2^q(s), & t = b. \end{cases}$$

It is clear that $\tilde{u}_s^q > 0$ on $[a, b]$ for all $s \in (a, b)$.

Since $g_0(t, s) \in C^{n-2}(I \times I)$, $\frac{\partial^{n-1}}{\partial t^{n-1}} g_0 \in C^\infty((I \times I) \setminus \{(t, t) / t \in I\})$ and there exists $\lim_{s \rightarrow t^\pm} \frac{\partial^{n-1}}{\partial t^{n-1}} g_0(t, s) \in \mathbb{R}$, we deduce that there exists $K^q > 0$ such that $\tilde{u}_s^q(t) \leq K^q$ for every $(t, s) \in I \times I$ and $q \in \{1, \dots, n-1\}$. Therefore, the following functions

$$\tilde{k}_1^q(s) = \min_{t \in I} \tilde{u}_s^q(t), \quad s \in [a, b],$$

$$\tilde{k}_2^q(s) = \max_{t \in I} \tilde{u}_s^q(t), \quad s \in [a, b],$$

are continuous on I and positive in (a, b) .

Taking $\phi(t) = (t-a)^{\alpha_q} (b-t)^{\beta_q} > 0$ on (a, b) , the function $v_s^q(t)$ satisfies the condition (P_g) if $n-q-c_q$ is even with $k_1(s) = \tilde{k}_1^q(s)$ and $k_2(s) = \tilde{k}_2^q(s)$ and condition (N_g) if $n-q-c_q$ is odd with $k_1(s) = -\tilde{k}_2^q(s)$ and $k_2(s) = -\tilde{k}_1^q(s)$.

As for the set of parameters of M in which the function $v_s^q[M]$, $q \in \{1, \dots, n-1\}$ maintains constant sign and monotony with respect to M , we have the following results:

Lemma 2.29. *Let $q \in \{1, \dots, n-1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) . If there exists $M \in \mathbb{R}$ be such that $\frac{\partial^q}{\partial t^q} g_M$ has constant sign, then g_M has constant sign.*

Proof. If $v_s^q[M](t) = \frac{\partial^q}{\partial t^q} g_M(t, s)$ has constant sign, then $v_s^{q-1}[M](t) = \frac{\partial^{q-1}}{\partial t^{q-1}} g_M(t, s)$ is a monotone and continuous function. Thus, $v_s^{q-1}[M]$ has at most one zero on I .

By recurrence, we have that $v_s^{q-l}[M](t) = \frac{\partial^{q-l}}{\partial t^{q-l}} g_M(t, s)$ has at most l zeros on I . In particular, $g_M(\cdot, s)$ has at most q zeros on I .

Now, since condition (N_a) implies that $g_M(\cdot, s)$ has at least q zeros of order smaller or equal to q on the boundary, we deduce that $g_M(\cdot, s)$ cannot have any zero on (a, b) . Thus, g_M has constant sign. \square

Moreover, we recall the following facts:

- If $n-k$ is even, then from Theorem 2.26 and [1, Lemma 1.8.33] we have that $g_M(t, s) > 0$ for all $(t, s) \in (a, b) \times (a, b)$ if and only if $M \in (\lambda_1, +\infty)$ or $M \in (\lambda_1, \lambda_2]$, and it is monotone decreasing with respect to M on such interval, where $\lambda_1 < 0$ is the first eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ and $\lambda_2 > 0$ (if the interval is bounded) is not an eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. The value of λ_2 is characterized in [5, Sect.7 and Sect.8].

- If $n - k$ is odd, then from Theorem 2.26 and [1, Lemma 1.8.25] we have that $g_M(t, s) < 0$ for all $(t, s) \in (a, b) \times (a, b)$ if and only if $M \in (-\infty, \lambda_1)$ or $M \in [\bar{\lambda}_2, \lambda_1)$ and it is monotone decreasing with respect to M on such interval, where $\lambda_1 > 0$ is the first eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ and $\bar{\lambda}_2 < 0$ (if the interval is bounded) is not an eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. The value of $\bar{\lambda}_2$ is characterized in [5, Sect. 7 and Sect. 8].

Therefore, as a direct consequence of Theorem 2.26, Lemma 2.29 and [2, Theorem 3.6], the following result is deduced.

Lemma 2.30. *The following properties hold:*

- If $n - k$ is even and $\frac{\partial^q}{\partial t^q} g_M(t, s) > 0$ on $(a, b) \times (a, b)$ for some $M \in \mathbb{R}$, then $\frac{\partial^q}{\partial t^q} g_M(t, s) > 0$ if and only if $M \in (\lambda_1, M_q]$, with $M_q \leq \lambda_2$. Moreover, $\frac{\partial^q}{\partial t^q} g_M$ is monotone decreasing with respect to $M \in (\lambda_1, M_q]$.
- If $n - k$ is even and $\frac{\partial^q}{\partial t^q} g_M(t, s) < 0$ on $(a, b) \times (a, b)$ for some $M \in \mathbb{R}$, then $\frac{\partial^q}{\partial t^q} g_M(t, s) < 0$ if and only if $M \in (\lambda_1, M_q]$, with $M_q \leq \lambda_2$. Moreover, $\frac{\partial^q}{\partial t^q} g_M$ is monotone increasing with respect to $M \in (\lambda_1, M_q]$.
- If $n - k$ is odd and $\frac{\partial^q}{\partial t^q} g_M(t, s) > 0$ on $(a, b) \times (a, b)$ for some $M \in \mathbb{R}$, then $\frac{\partial^q}{\partial t^q} g_M(t, s) > 0$ if and only if $M \in [M_q, \lambda_1)$, with $M_q \geq \bar{\lambda}_2$. Moreover, $\frac{\partial^q}{\partial t^q} g_M$ is monotone increasing with respect to $M \in [M_q, \lambda_1)$.
- If $n - k$ is odd and $\frac{\partial^q}{\partial t^q} g_M(t, s) < 0$ on $(a, b) \times (a, b)$ for some $M \in \mathbb{R}$, then $\frac{\partial^q}{\partial t^q} g_M(t, s) < 0$ if and only if $M \in [M_q, \lambda_1)$, with $M_q \geq \bar{\lambda}_2$. Moreover, $\frac{\partial^q}{\partial t^q} g_M$ is monotone decreasing with respect to $M \in [M_q, \lambda_1)$.

3 Study of the constant sign of the derivatives of Green's function

In this section for any $q \in \{1, \dots, n - 1\}$ be fixed, we will give a proof of the main result that characterizes the constant sign of the partial derivative with respect to t of order q of the Green's function related to operator $T_n[M]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

We distinguish three cases:

- $c_q \geq 1$ and $d_q \geq 1$,
- $c_q = n - q$ and $d_q = 0$,
- $c_q = 0$, and $d_q = n - q$,

where c_q and d_q are defined in (2.27) and (2.28) and satisfy $c_q + d_q = n - q$.

3.1 Case (a): $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$

Lemma 3.1. *Let $q \in \{1, \dots, n - 1\}$ be fixed. If $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$, then the indices $z = k - (j - 1)$ and $h = n - k - (r - 1)$ are such that*

$$\mu_z^q + q + \eta = n - 1, \quad \text{and} \quad \rho_h^q + q + \gamma = n - 1,$$

with j and r defined in (2.20) and (2.21), η and γ defined in (2.13) and (2.14).

Proof. By the definition of $\mu_z^q = \mu_{k-(j-1)}^q$ and $\rho_h^q = \rho_{n-k-(r-1)}^q$ we have that since $c_q \geq 1$, $\mu_z^q = \sigma_k - q \geq 0$ and since $d_q \geq 1$, $\rho_h^q = \varepsilon_{n-k} - q \geq 0$. Using Remark 2.7, we deduce that

$$\mu_z^q + q + \eta = \sigma_k - q + q + n - 1 - \sigma_k = n - 1,$$

and

$$\rho_h^q + q + \gamma = \varepsilon_{n-k} - q + q + n - 1 - \varepsilon_{n-k} = n - 1.$$

□

Lemma 3.2. *Let $q \in \{1, \dots, n - 1\}$ be fixed. If $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$, then $z = p$ and $h = l$ with z and h defined in Lemma 3.1, p and l defined in (2.25) and (2.24).*

Proof. First, note that $z = |A_q|$ is the cardinal of the set $A_q = \{i \in \{1, \dots, k\} / \sigma_i \geq q\}$ and $p = |D_q|$ is the cardinal of the set $D_q = \{i \in \{1, \dots, k\} / \delta_i + q \leq n - 1\}$. Moreover, by (2.27), it is clear that c_q corresponds with the number of σ 's that are equal or greater than q , that is, $z = c_q$.

On the other hand, we have that

$$\{\varepsilon_1, \dots, \varepsilon_{n-k}, n - 1 - \delta_k, \dots, n - 1 - \delta_1\} \equiv \{0, \dots, n - 1\}.$$

Among these n elements (all the numbers between 0 and $n - 1$) there are $n - q$ elements that are equal or greater than q . By (2.28), we have that the number of ε 's that are equal or greater than q is d_q . Therefore, the number of δ 's such that $n - 1 - \delta_i \geq q$ is $n - q - d_q$, that is, $p = n - q - d_q$. Since, $c_q + d_q = n - q$, we have that $z = p$. In an analogous way, we can prove that $h = l$. □

Remark 3.3. It should be noted that in the previous proof, since j and r depend on q , we have that z and h also depend on q , but we will omit such dependence in the notation for the sake of simplicity.

Remark 3.4. Indices z and h depend on the space $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ and are unique. Indeed, for the rest of the components of space $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ we have that $(\mu_i^q + q + \eta) \bmod n < n - 1$ for $i \neq z$ with $i \in \{1, \dots, k\}$ and $(\rho_l^q + q + \gamma) \bmod n < n - 1$ for $l \neq h$ with $l \in \{1, \dots, n - k\}$.

Now, we present some results that provide sufficient conditions to ensure the constant sign of solutions of (1.1) in the spaces $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$. Using Lemma 2.16, these results can be proved analogously to [5, Proposition 6.7] and [5, Proposition 6.9], respectively, and we omit them.

Let us consider the following spaces

$$X_2 := \begin{cases} X_{\{\mu_1^q, \dots, \mu_{z-1}^q, \mu_{z+1}^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q | \beta^q\}}, & \text{if } z \neq 1, \\ X_{\{\mu_2^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q | \beta^q\}}, & \text{if } z = 1, \end{cases} \quad (3.1)$$

$$X_3 := \begin{cases} X_{\{\mu_1^q, \dots, \mu_{z-1}^q, \mu_{z+1}^q, \dots, \mu_k^q | \alpha^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}, & \text{if } z \neq 1, \\ X_{\{\mu_2^q, \dots, \mu_k^q | \alpha^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}, & \text{if } z = 1, \end{cases} \quad (3.2)$$

$$X_4 := \begin{cases} X_{\{\mu_1^q, \dots, \mu_k^q | \alpha^q\}}^{\{\rho_1^q, \dots, \rho_{h-1}^q, \rho_{h+1}^q, \dots, \rho_{n-k}^q\}}, & \text{if } h \neq 1, \\ X_{\{\mu_1^q, \dots, \mu_k^q | \alpha^q\}}^{\{\rho_2^q, \dots, \rho_{n-k}^q\}}, & \text{if } h = 1, \end{cases} \quad (3.3)$$

and

$$X_5 := \begin{cases} X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{h-1}^q, \rho_{h+1}^q, \dots, \rho_{n-k}^q | \beta^q\}}, & \text{if } h \neq 1, \\ X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_2^q, \dots, \rho_{n-k}^q | \beta^q\}}, & \text{if } h = 1, \end{cases} \quad (3.4)$$

where z and h are defined in Lemma 3.1.

Proposition 3.5. *Let $q \in \{1, \dots, n-1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$. If $u \in C^n(I)$ is a solution of (1.1) on (a, b) , satisfying the boundary conditions:*

$$u^{(\mu_1^q)}(a) = \dots = u^{(\mu_{z-1}^q)}(a) = u^{(\mu_{z+1}^q)}(a) = \dots = u^{(\mu_k^q)}(a) = 0, \quad (3.5)$$

$$u^{(\rho_1^q)}(b) = \dots = u^{(\rho_{n-k}^q)}(b) = 0, \quad (3.6)$$

then it does not have any zero on (a, b) provided that one of the following assertions is satisfied:

- Let $n - k$ be even:

- If $k > 1$, $\mu_z^q \neq z - 1$ and $M \in [\lambda_3^q, \lambda_2^q]$, where:

- * $\lambda_3^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_3 .

- * $\lambda_2^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_2 .

- If $k > 1$, $\mu_z^q = z - 1$ and $M \in [\lambda_1, \lambda_2^q]$, where:
 - * $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.
 - * $\lambda_2^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_2 .
- If $k = 1$, $\mu_1^q \neq 0$ and $M \in [\lambda_3^q, +\infty)$, where:
 - * $\lambda_3^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\alpha^q\}}^{\{\rho_1^q, \dots, \rho_{n-1}^q\}}$, where $\alpha^q = 0$.
- If $k = 1$, $\mu_1^q = 0$ and $M \in [\lambda_1, +\infty)$, where:
 - * $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-1}\}}$.

• Let $n - k$ be odd:

- If $k > 1$, $\mu_z^q \neq z - 1$ and $M \in [\lambda_2^q, \lambda_3^q]$, where:
 - * $\lambda_3^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_3 .
 - * $\lambda_2^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_2 .
- If $k > 1$, $\mu_z^q = z - 1$ and $M \in [\lambda_2^q, \lambda_1]$, where:
 - * $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.
 - * $\lambda_2^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_2 .
- If $k = 1$, $\mu_1^q \neq 0$ and $M \in (-\infty, \lambda_3^q]$, where:
 - * $\lambda_3^q > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\alpha^q\}}^{\{\rho_1^q, \dots, \rho_{n-1}^q\}}$, where $\alpha^q = 0$.
- If $k = 1$, $\mu_1^q = 0$ and $M \in (-\infty, \lambda_1]$, where:
 - * $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-1}\}}$.

Proposition 3.6. Let $q \in \{1, \dots, n - 1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$. If $u \in C^n(I)$ is a solution of (1.1) on (a, b) satisfying the boundary conditions:

$$u^{(\mu_1^q)}(a) = \dots = u^{(\mu_k^q)}(a) = 0, \quad (3.7)$$

$$u^{(\rho_1^q)}(b) = \dots = u^{(\rho_{h-1}^q)}(b) = u^{(\rho_{h+1}^q)}(b) = \dots = u^{(\rho_{n-k}^q)}(b) = 0, \quad (3.8)$$

then it does not have any zero on (a, b) provided that one of the following assertions is satisfied:

• Let $n - k$ be even:

- If $\rho_h^q \neq h - 1$ and $M \in [\lambda_5^q, \lambda_4^q]$, where:
 - * $\lambda_5^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_5 .
 - * $\lambda_4^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_4 .
- If $\rho_h^q = h - 1$ and $M \in [\lambda_1, \lambda_4^q]$, where:
 - * $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

* $\lambda_4^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_4 .

• Let $n - k$ be odd:

– If $k < n - 1$, $\rho_h^q \neq h - 1$ and $M \in [\lambda_4^q, \lambda_5^q]$, where:

* $\lambda_5^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_5 .

* $\lambda_4^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_4 .

– If $k < n - 1$, $\rho_h^q = h - 1$ and $M \in [\lambda_4^q, \lambda_1]$, where:

* $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

* $\lambda_4^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_4 .

– If $k = n - 1$, $\rho_1^q \neq 0$ and $M \in (-\infty, \lambda_5^q]$, where:

* $\lambda_5^q > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\mu_1^q, \dots, \mu_{n-1}^q\}}^{\{\beta^q\}}$, where $\beta^q = 0$.

– If $k = n - 1$, $\rho_1^q = 0$ and $M \in (-\infty, \lambda_1]$, where:

* $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_{n-1}\}}^{\{\varepsilon_1\}}$.

Example 3.7. Let us consider the operator $T_4[M]$ with the boundary conditions:

$$u'(0) = u'''(0) = u(1) = u'(1) = 0.$$

The hypotheses of Propositions 3.5 and 3.6 are satisfied only for the value $q = 1$. Moreover, $z = 2$, $h = 1$, and $v_s^1[M]$ satisfies the conditions of the space $X_{\{0,2\}}^{\{0,3\}}$.

From Proposition 3.5, we can affirm that any nontrivial solution of the problem

$$T_4[M] = 0, \quad t \in [0, 1], \quad u(0) = u(1) = u'''(1) = 0, \quad (3.9)$$

does not have any zero on $(0, 1)$ for $M \in [\lambda_3^1, \lambda_2^1]$, where $\lambda_3^1 < 0$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{0,1\}}^{\{0,3\}}$ and $\lambda_2^1 > 0$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{1\}}^{\{0,1,3\}}$.

The eigenvalues of $T_4[0]$ in $X_{\{0,1\}}^{\{0,3\}}$ are given by $-\lambda^4$, where λ is a positive solution of $\sin(\lambda) = 0$. The smallest positive solution of this equation is π , so $\lambda_3^1 = -\pi^4$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{0,1\}}^{\{0,3\}}$.

Similarly, the eigenvalues of $T_4[0]$ in $X_{\{1\}}^{\{0,1,3\}}$ are given by λ^4 , where λ is a positive solution of

$$\left(-1 + e^{\sqrt{2}m}\right) \cos\left(\frac{m}{\sqrt{2}}\right) + \left(1 + e^{\sqrt{2}m}\right) \sin\left(\frac{m}{\sqrt{2}}\right) = 0.$$

Denoting by m_2 the smallest positive solution of this equation, we deduce that $\lambda_2^1 = m_2^4 \approx 3.34^4$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0\}}^{\{0,1,3\}}$.

It is easy to verify that the solutions of problem (3.9) are given as multiples of the following expression:

- If $M = -m^4 < 0$,

$$u(t) = \cosh(m - mt) \sin(m) - \sin(mt) - \cosh(m) \sin(m - mt) - \cos(m - mt) \sinh(m) + \sinh(mt) + \cos(m) \sinh(m - mt).$$

- If $M = 0$,

$$u(t) = t^2 - t.$$

- If $M = m^4 > 0$,

$$u(t) = e^{-\frac{mt}{\sqrt{2}}} \left(e^{\sqrt{2}m} (-1 + e^{\sqrt{2}mt}) \cos\left(\frac{m(-2+t)}{\sqrt{2}}\right) - e^{\sqrt{2}m} (-1 + e^{\sqrt{2}m}) \cos\left(\frac{mt}{\sqrt{2}}\right) \right) + e^{-\frac{mt}{\sqrt{2}}} (-1 + e^{\sqrt{2}m}) \left(-e^{\sqrt{2}m} + e^{\sqrt{2}mt} \right) \sin\left(\frac{mt}{\sqrt{2}}\right).$$

Analogously, from Proposition 3.6, we conclude that any solution of the problem

$$T_4[M]u(t) = 0, \quad t \in [0, 1], \quad u(0) = u''(0) = u'''(1) = 0, \quad (3.10)$$

does not have any zero on $(0, 1)$ for $M \in [\lambda_1, \lambda_4^1]$, where $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{1,3\}}^{\{0,1\}}$ and $\lambda_4^1 > 0$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1,2\}}^{\{3\}}$.

The eigenvalues of $T_4[0]$ in $X_{\{1,3\}}^{\{0,1\}}$ are given by $-\lambda^4$, where λ is a positive solution of

$$\sin(\lambda) + \cos(\lambda) \tanh(\lambda) = 0.$$

Now, if m_1 is the smallest positive solution of this equation, then $\lambda_1 = -m_1^4 \approx -2.36^4$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{0,1\}}^{\{0,3\}}$.

On the other hand, the eigenvalues of $T_4[0]$ in $X_{\{0,1,2\}}^{\{3\}}$ are given by λ^4 , where λ is a positive solution of $\cos\left(\frac{m}{\sqrt{2}}\right) = 0$. The smallest positive solution of this equation is $\frac{\sqrt{2}}{2}\pi$, so $\lambda_4^1 = \left(\frac{\sqrt{2}}{2}\pi\right)^4$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1,2\}}^{\{3\}}$.

It is immediate to show that solutions of problem (3.10) are given as multiples of:

- If $M = -m^4 < 0$,

$$u(t) = \sin(mt) + \cos(m) \operatorname{sech}(m) \sinh(mt).$$

- If $M = 0$,

$$u(t) = t.$$

- If $M = m^4 > 0$,

$$u(t) = e^{-\frac{mt}{\sqrt{2}}} \left((-1 + e^{\sqrt{2}m(1+t)}) \cos\left(\frac{m(-1+t)}{\sqrt{2}}\right) + (-e^{\sqrt{2}m} + e^{\sqrt{2}mt}) \cos\left(\frac{m(1+t)}{\sqrt{2}}\right) \right) + e^{-\frac{mt}{\sqrt{2}}} \left((1 + e^{\sqrt{2}m(1+t)}) \sin\left(\frac{m(-1+t)}{\sqrt{2}}\right) + (e^{\sqrt{2}m} + e^{\sqrt{2}mt}) \sin\left(\frac{m(1+t)}{\sqrt{2}}\right) \right).$$

Remark 3.8. It should be noted that in the above propositions we have used the fact proven in Lemma 2.17 that the first nonzero eigenvalues of the spaces $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ and $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ coincide.

Let us define the following spaces

$$X_2^* := \begin{cases} X_{\{\tau_1, \dots, \tau_{n-k}|\eta\}}^{\{\delta_1, \dots, \delta_{z-1}, \delta_{z+1}, \dots, \delta_k\}}, & \text{if } z \neq 1, \\ X_{\{\tau_1, \dots, \tau_{n-k}|\eta\}}^{\{\delta_2, \dots, \delta_k\}}, & \text{if } z = 1, \end{cases} \quad (3.11)$$

and

$$X_4^* := \begin{cases} X_{\{\tau_1, \dots, \tau_{h-1}, \tau_{h+1}, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k|\gamma\}}, & \text{if } h \neq 1, \\ X_{\{\tau_2, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k|\gamma\}}, & \text{if } h = 1. \end{cases} \quad (3.12)$$

Lemma 3.9. $\lambda \neq 0$ is an eigenvalue of $T_n[0]$ in X_2^* defined in (3.11) if and only if λ is an eigenvalue of $T_n[0]$ in X_2 . In particular, $(\lambda_2^*)^q = \lambda_2^q$ where $(\lambda_2^*)^q \neq 0$ is the least positive (biggest negative) eigenvalue of $T_n[0]$ in X_2^* and $\lambda_2^q \neq 0$ is the least positive (biggest negative) eigenvalue of $T_n[0]$ in X_2 .

Proof. Suppose that $z \neq 1$ (the case $z = 1$ is proved similarly). By the definition of adjoint space of $X_{\{\mu_1^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q\}}$ we have that

$$\{\mu_1^q, \dots, \mu_{z-1}^q, \mu_z^q, \mu_{z+1}^q, \dots, \mu_k^q, n-1-\tau_{n-k}^q, \dots, n-1-\tau_1^q\} = \{0, \dots, n-1\},$$

and

$$\{\rho_1^q, \dots, \rho_{n-k}^q, n-1-\delta_k^q, \dots, n-1-\delta_1^q\} = \{0, \dots, n-1\}.$$

Since, $\mu_z^q = \sigma_k - q$ and $\beta^q = n-1-\delta_k^q$ we deduce that

$$X_{\{\mu_1^q, \dots, \mu_{z-1}^q, \mu_{z+1}^q, \dots, \mu_k^q\}}^{\{\rho_1^q, \dots, \rho_{n-k}^q|\beta^q\}} \xrightarrow{*} X_{\{\tau_1^q, \dots, \tau_{n-k}^q|n-1-\sigma_k+q\}}^{\{\delta_1^q, \dots, \delta_{k-1}^q\}}. \quad (3.13)$$

Using that $\eta = n-1-\sigma_k$, the definition of $X_{\{\tau_1^q, \dots, \tau_{n-k}^q\}}^{\{\delta_1^q, \dots, \delta_k^q\}}$ and the diagram (2.26) we infer that

$$X_{\{\tau_1^q, \dots, \tau_{n-k}^q|n-1-\sigma_k+q\}}^{\{\delta_1^q, \dots, \delta_{k-1}^q\}} \xrightarrow{\phi^q} X_{\{\tau_1, \dots, \tau_{n-k}|\eta\}}^{\{\delta_1, \dots, \delta_{p-1}, \delta_{p+1}, \dots, \delta_k\}}.$$

From Lemma 3.2 we know that $p = z$ and therefore

$$X_{\{\tau_1^q, \dots, \tau_{n-k}^q|n-1-\sigma_k+q\}}^{\{\delta_1^q, \dots, \delta_{k-1}^q\}} \xrightarrow{\phi^q} X_{\{\tau_1, \dots, \tau_{n-k}|\eta\}}^{\{\delta_1, \dots, \delta_{z-1}, \delta_{z+1}, \dots, \delta_k\}}. \quad (3.14)$$

Considering Lemma 2.17 and the fact that the adjoint spaces have the same eigenvalues, from (3.13) and (3.14), we obtain that the spaces $X_{\{\rho_1^q, \dots, \rho_{n-k}^q | \beta^q\}}$ and $X_{\{\delta_1, \dots, \delta_{z-1}, \delta_{z+1}, \dots, \delta_k\}}_{\{\tau_1, \dots, \tau_{n-k} | \eta\}}$ have the same eigenvalues. In particular, $(\lambda_2^*)^q = \lambda_2^q$. \square

Using similar argument to the previous result, we arrive to the next one.

Lemma 3.10. $\bar{\lambda} \neq 0$ is an eigenvalue of $T_n[0]$ in X_4^* defined in (3.12) if and only if $\bar{\lambda}$ is an eigenvalue of $T_n[0]$ in X_4 . In particular, $(\lambda_4^*)^q = \lambda_4^q$ where $(\lambda_4^*)^q \neq 0$ is the least positive (biggest negative) eigenvalue of $T_n[0]$ in X_4^* and $\lambda_4^q \neq 0$ is the least positive (biggest negative) eigenvalue of $T_n[0]$ in X_4 .

Next, we state and prove the main result, which gives the characterization of the set of parameter M where $v_s^q[M]$ is of constant sign.

Theorem 3.11. Let $q \in \{1, \dots, n-1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$. The following properties are fulfilled:

- Let $2 \leq k \leq n-2$ and $n-k$ even:
 - If $n-q-c_q$ is even, then $v_s^q[M]$ is strongly positive on $I \times I$ if, and only if, $M \in (\lambda_1, \lambda^q]$, and if $n-q-c_q$ is odd, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in (\lambda_1, \lambda^q]$ where:
 - * $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}_{\{\sigma_1, \dots, \sigma_k\}}$.
 - * $\lambda^q > 0$ is the minimum between:
 - $\lambda_2^q > 0$, the least positive eigenvalue of $T_n[0]$ in X_2 defined in (3.1).
 - $\lambda_4^q > 0$, the least positive eigenvalue of $T_n[0]$ in X_4 defined in (3.3).
- Let $2 \leq k \leq n-2$ and $n-k$ odd:
 - If $n-q-c_q$ is even, then $v_s^q[M]$ is strongly positive on $I \times I$ if, and only if, $M \in [\lambda^q, \lambda_1)$, and if $n-q-c_q$ is odd, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in [\lambda^q, \lambda_1)$ where:
 - * $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}_{\{\sigma_1, \dots, \sigma_k\}}$.
 - * $\lambda^q < 0$ is the maximum between:
 - $\lambda_2^q < 0$, the biggest negative eigenvalue of $T_n[0]$ in X_2 .
 - $\lambda_4^q < 0$, the biggest negative eigenvalue of $T_n[0]$ in X_4 .
- Let $k=1$ and $n > 2$ odd:
 - If $n-q-1$ is even, then $v_s^q[M]$ is strongly positive on $I \times I$ if, and only if, $M \in (\lambda_1, \lambda_2^q]$, and if $n-q-1$ is odd, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in (\lambda_1, \lambda_2^q]$ where:
 - * $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\varepsilon_1, \dots, \varepsilon_{n-1}\}}_{\{\sigma_1\}}$.

* $\lambda_2^q > 0$ is the least positive eigenvalue of $T_n[0]$ in X_2 .

• Let $k = 1$ and $n > 2$ even:

– If $n - q - 1$ is even, then $v_s^q[M]$ is strongly positive on $I \times I$ if, and only if, $M \in [\lambda_2^q, \lambda_1)$, and if $n - q - 1$ is odd, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in [\lambda_2^q, \lambda_1)$, where:

* $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-1}\}}$.

* $\lambda_2^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_2 .

• If $k = n - 1$, $n > 2$ and $n - q - c_q = 1$, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in [\lambda_2^q, \lambda_1)$, where

* $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_{n-1}\}}^{\{\varepsilon_1\}}$.

* $\lambda_2^q < 0$ is the biggest negative eigenvalue of $T_n[0]$ in X_2 .

Proof. First, it has been shown in Theorem 2.28 that v_s^q satisfies the property (P_g) if $n - q - c_q$ is even and the property (N_g) if $n - q - c_q$ is odd. In addition, from Lemma 2.30 we know that if $v_s^q[M]$ has constant sign then the first eigenvalue λ_1 of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$ is one of the extremes of the interval where $v_s^q[M]$ holds that sign.

Again, taking into account Lemma 2.30 on the monotony of $v_s^q[M]$ with respect to M , the constant sign starts/ends at the first eigenvalue λ_1 of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$. Thus, from Theorems 2.26 and 2.28 we conclude that:

- If $n - k$ and $n - q - c_q$ are even, and $M \leq 0$, then $v_s^q[M]$ is strongly positive on $I \times I$ if, and only if, $M \in (\lambda_1, 0]$.
- If $n - k$ is even, $n - q - c_q$ is odd and $M \leq 0$, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in (\lambda_1, 0]$.
- If $n - k$ is odd, $n - q - c_q$ is even and $M \geq 0$, then $v_s^q[M]$ is strongly positive on $I \times I$ if, and only if, $M \in [0, \lambda_1)$.
- If $n - k$ and $n - q - c_q$ are odd, and $M \geq 0$, then $v_s^q[M]$ is strongly negative on $I \times I$ if, and only if, $M \in [0, \lambda_1)$.

We now need to determine the other extreme of the constant sign interval of $v_s^q[M]$ using Definition 2.25, Propositions 3.5 and 3.6. We divide the proof into five steps assuming that $n - k$ and $n - q - c_q$ are even, and $2 \leq k \leq n - 2$. For the rest of cases the proof is done in an analogous way.

Step 1. Study of $v_s^q[M]$ at $s = a$.

Step 2. Study of $v_s^q[M]$ at $s = b$.

Step 3. Study of $v_s^q[M]$ at $t = a$.

Step 4. Study of $v_s^q[M]$ at $t = b$.

Step 5. Study of $v_s^q[M]$ on $(a, b) \times (a, b)$.

Let us denote

$$g_M(t, s) = \begin{cases} g_M^1(t, s), & a \leq s \leq t \leq b, \\ g_M^2(t, s), & a < t < s < b, \end{cases}$$

as the related Green's function of $T_n[M]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

Step 1. Study of the function $v_s^q[M]$ at $s = a$.

Let us consider the function

$$w_{M,q}(t) = \frac{\partial^\eta}{\partial s^\eta} \left(\frac{\partial^q}{\partial t^q} g_M^1(t, s) \right)_{|s=a},$$

where η has been defined in (2.13).

If $\eta > 0$, since $\{0, \dots, \eta - 1\} \subset \{\tau_1, \dots, \tau_{n-k}\}$ using the equality (2.10) we obtain that

$$g_M^1(t, a) = \frac{\partial}{\partial s} g_M^1(t, a) = \dots = \frac{\partial^{\eta-1}}{\partial s^{\eta-1}} g_M^1(t, a) = 0, \quad \text{for all } t \in (a, b].$$

Therefore, for all $t \in (a, b]$ and $q \in \{1, \dots, n-1\}$ we infer that

$$\frac{\partial^q}{\partial t^q} g_M^1(t, a) = \frac{\partial^q}{\partial t^q} \left(\frac{\partial}{\partial s} g_M^1(t, s) \right)_{|s=a} = \dots = \frac{\partial^q}{\partial t^q} \left(\frac{\partial^{\eta-1}}{\partial s^{\eta-1}} g_M^1(t, s) \right)_{|s=a} = 0,$$

equivalently,

$$\frac{\partial^q}{\partial t^q} g_M^1(t, a) = \frac{\partial}{\partial s} \left(\frac{\partial^q}{\partial t^q} g_M^1(t, s) \right)_{|s=a} = \dots = \frac{\partial^{\eta-1}}{\partial s^{\eta-1}} \left(\frac{\partial^q}{\partial t^q} g_M^1(t, s) \right)_{|s=a} = 0.$$

Note that it is necessary that $w_{M,q} > 0$ to guarantee the strongly positive sign of the function $v_s^q[M]$. Indeed, if there exists $t^* \in [a, b]$, such that $w_{M,q}(t^*) < 0$, then there exists $\rho(t^*) > a$ such that $v_s^q[M](t^*) < 0$ for all $s \in (a, \rho(t^*))$, which contradicts the strongly positive sign.

Since the function $v_s^q[M]$ is a solution of $T_n[M] v_s^q[M](t) = 0$ for $t \neq s$, we have that

$$\frac{\partial^\eta}{\partial s^\eta} (T_n[M] v_s^q[M](t))_{|s=a} = T_n[M] w_{M,q}(t) = 0, \quad t \in (a, b],$$

that is, the function $w_{M,q}$ solves the equation $T_n[M] u(t) = 0$, $t \in (a, b]$.

It remains to obtain the boundary conditions satisfied by the function $w_{M,q}$.

Using the matrix of the Green's function given in (2.4), the equality (2.6), the expression of g_{n-j} , $j = 1, \dots, n-1$ given in (2.8) and the following identity

$$\frac{\partial}{\partial t} \left(\frac{\partial^q}{\partial t^q} G_M(t, s) \right) = A \frac{\partial^q}{\partial t^q} G_M(t, s), \quad \text{for all } t \in I \setminus \{s\}, q \in \{1, \dots, n-1\},$$

it follows that

$$\frac{\partial^{\mu_1^q+q}}{\partial t^{\mu_1^q+q}} g_M^2(t, s)|_{t=a} = 0, -\frac{\partial^{\mu_1^q+q+1}}{\partial t^{\mu_1^q+q} \partial s} g_M^2(t, s)|_{t=a} = 0, \dots, (-1)^\eta \frac{\partial^{\mu_1^q+q+\eta}}{\partial t^{\mu_1^q+q} \partial s^\eta} g_M^2(t, s)|_{t=a} = 0.$$

Since $(\eta + \mu_1^q + q) \bmod n < n-1$, then we are not in an element of the diagonal of $\frac{\partial^q}{\partial t^q} G_M(t, s)$, and taking in account that the above equalities are satisfied for $s = a$, by continuity we deduce that

$$\left\{ \begin{array}{l} \frac{\partial^{\mu_1^q+q}}{\partial t^{\mu_1^q+q}} g_M^1(t, s)|_{(t,s)=(a,a)} = 0, \\ -\frac{\partial^{\mu_1^q+q+1}}{\partial t^{\mu_1^q+q} \partial s} g_M^1(t, s)|_{(t,s)=(a,a)} = 0, \\ \vdots \\ (-1)^\eta \frac{\partial^{\mu_1^q+q+\eta}}{\partial t^{\mu_1^q+q} \partial s^\eta} g_M^1(t, s)|_{(t,s)=(a,a)} = 0. \end{array} \right.$$

Therefore,

$$w_{M,q}^{(\mu_1^q)}(a) = \frac{\partial^{\mu_1^q+q+\eta}}{\partial t^{\mu_1^q+q} \partial s^\eta} g_M^1(t, s)|_{(t,s)=(a,a)} = 0.$$

Making a similar reasoning we deduce that

$$w_{M,q}^{(\mu_2^q)}(a) = \dots = w_{M,q}^{(\mu_{z-1}^q)}(a) = w_{M,q}^{(\mu_{z+1}^q)}(a) = \dots = w_{M,q}^{(\mu_k^q)}(a) = 0.$$

For μ_z^q , we have the following equalities

$$\frac{\partial^{\mu_z^q+q}}{\partial t^{\mu_z^q+q}} g_M^2(t, s)|_{t=a} = 0, -\frac{\partial^{\mu_z^q+q+1}}{\partial t^{\mu_z^q+q} \partial s} g_M^2(t, s)|_{t=a} = 0, \dots, (-1)^\eta \frac{\partial^{\mu_z^q+q+\eta}}{\partial t^{\mu_z^q+q} \partial s^\eta} g_M^2(t, s)|_{t=a} = 0.$$

By assumption, we have that $\eta + \mu_z^q + q = n-1$, so we are in an element of the diagonal of $\frac{\partial^q}{\partial t^q} G_M(t, s)$, and taking into account again that the previous equalities are satisfied for $s = a$

and that $\frac{\partial^q}{\partial t^q} G_M(t, s)$ has a jump equal to 1 on the diagonal we infer that

$$\left\{ \begin{array}{l} \frac{\partial^{\mu_z^q+q}}{\partial t^{\mu_z^q+q}} g_M^1(t, s)|_{(t,s)=(a,a)} = 0, \\ -\frac{\partial^{\mu_z^q+q+1}}{\partial t^{\mu_z^q+q} \partial s} g_M^1(t, s)|_{(t,s)=(a,a)} = 0, \\ \vdots \\ (-1)^\eta \frac{\partial^{\mu_z^q+q+\eta}}{\partial t^{\mu_z^q+q} \partial s^\eta} g_M^1(t, s)|_{(t,s)=(a,a)} = 1. \end{array} \right.$$

Therefore,

$$w_{M,q}^{(\mu_z^q)}(a) = \frac{\partial^{\mu_z^q+q+\eta}}{\partial t^{\mu_z^q+q} \partial s^\eta} g_M^1(t, s)|_{(t,s)=(a,a)} = (-1)^\eta.$$

Let's see what conditions $w_{M,q}$ satisfies at $t = b$. To do this, proceeding with the previous argument we obtain that

$$\frac{\partial^{\rho_1^q+q}}{\partial t^{\rho_1^q+q}} g_M^1(t, s)|_{t=b} = 0, -\frac{\partial^{\rho_1^q+q+1}}{\partial t^{\rho_1^q+q} \partial s} g_M^1(t, s)|_{t=b} = 0, \dots, (-1)^\eta \frac{\partial^{\rho_1^q+q+\eta}}{\partial t^{\rho_1^q+q} \partial s^\eta} g_M^1(t, s)|_{t=b} = 0.$$

The above equalities are satisfied at $s = a$ and since $b \neq a$ we have that

$$w_{M,q}^{(\rho_1^q)}(b) = \frac{\partial^{\rho_1^q+q+\eta}}{\partial t^{\rho_1^q+q} \partial s^\eta} g_M^1(t, s)|_{(t,s)=(b,a)} = 0.$$

Analogously we obtain that

$$w_{M,q}^{(\rho_2^q)}(b) = \dots = w_{M,q}^{(\rho_{n-k}^q)}(b) = 0.$$

If $\eta = 0$, then it follows from the matrix argument we made earlier that the function $w_{M,q}(t) = \frac{\partial^q}{\partial t^q} g_M^1(t, a)$ satisfies the same boundary conditions (3.5)–(3.6).

Then from Proposition 3.5 and Theorem 2.28 we conclude that $w_{M,q} > 0$ on (a, b) for all $M \in [0, \lambda_2^q]$.

Now, let us see that $v_s^q[M]$ cannot be positive for $M > \lambda_2^q$.

Suppose that there exists $\widehat{M} > \lambda_2^q$ such that $v_s^q[\widehat{M}]$ is positive. Then, by monotony of $v_s^q[M]$ we have for every $M \in [\lambda_2^q, \widehat{M}]$ that $w_{\widehat{M},q} \leq w_{M,q} \leq w_{\lambda_2^q,q}$.

In particular, $w_{\widehat{M},q}^{(\beta^q)}(b) \geq w_{M,q}^{(\beta^q)}(b) \geq w_{\lambda_2^q,q}^{(\beta^q)}(b) = 0$ if β^q is even and $w_{\widehat{M},q}^{(\beta^q)}(b) \leq w_{M,q}^{(\beta^q)}(b) \leq w_{\lambda_2^q,q}^{(\beta^q)}(b) = 0$ if β^q is odd.

If $w_{\widehat{M},q}^{(\beta^q)}(b) \neq 0$, then there exists $\rho > 0$ such that $w_{\widehat{M},q}(t) < 0$ for all $t \in (b - \rho, b)$, which contradicts our assumption. Hence,

$$0 = w_{\widehat{M},q}^{(\beta^q)}(b) = w_{M,q}^{(\beta^q)}(b) = w_{\lambda_2^q,q}^{(\beta^q)}(b), \quad \forall M \in [\lambda_2^q, \widehat{M}],$$

and this fact contradicts the discrete character of the spectrum.

Thus, we conclude that, if $M \in [0, \lambda_2^q]$, then

$$\forall t \in (a, b), \quad \exists \rho^q(t) > 0 \mid v_s^q[M](t) > 0 \quad \forall s \in (a, a + \rho^q(t)).$$

Moreover, if $M > \lambda_2^q$, then $v_s^q[M]$ is not positive.

Step 2. Study of the function $v_s^q[M]$ at $s = b$.

We study the following function by applying similar reasoning to Step 1. In this case we consider

$$y_{M,q}(t) = \frac{\partial^\gamma}{\partial s^\gamma} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right) \Big|_{s=b},$$

where γ has been defined in (2.14).

In this case, using arguments analogous to Step 1 we obtain that if $\gamma > 0$, then

$$\frac{\partial^q}{\partial t^q} g_M^2(t, b) = \frac{\partial}{\partial s} g_M^2(t, s) \Big|_{s=b} = \dots = \frac{\partial^{\gamma-1}}{\partial s^{\gamma-1}} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right) \Big|_{s=b} = 0.$$

Moreover, if γ is even, then $y_{M,q} \geq 0$ and if γ is odd, then $y_{M,q} \leq 0$.

Again, we have that

$$T_n[M] y_{M,q}(t) = 0, \quad \forall t \in [a, b].$$

In the same way than above, studying $\frac{\partial^q}{\partial t^q} G_M(t, s)$ to determine the boundary conditions of $y_{M,q}$ we infer that

$$\begin{aligned} y_{M,q}^{(\mu_1^q)}(a) &= \dots = y_{M,q}^{(\mu_k^q)}(a) = 0, \\ y_{M,q}^{(\rho_1^q)}(b) &= \dots = y_{M,q}^{(\rho_{h-1}^q)}(b) = y_{M,q}^{(\rho_{h+1}^q)}(b) = \dots = y_{M,q}^{(\rho_{n-k}^q)}(b) = 0, \\ y_{M,q}^{(\rho_h^q)}(b) &= (-1)^{\gamma+1}. \end{aligned}$$

Analogously, if $\gamma = 0$, then $y_{M,q}(t) = \frac{\partial^q}{\partial t^q} g_M^2(t, b)$ satisfies the conditions (3.7)–(3.8) by an argument similar to the case $\gamma > 0$.

Then from Proposition 3.6 and Theorem 2.28 we conclude that $y_{M,q} > 0$ if γ is even and $y_{M,q} < 0$ if γ is odd on (a, b) for all $M \in [0, \lambda_4^q]$.

As in the previous step it can be seen that for $M > \lambda_4^q$, the function $v_s^q[M]$ is never positive. Therefore, we conclude from this step that, if $M \in [0, \lambda_4^q]$, then

$$\forall t \in (a, b), \quad \exists \rho^q(t) > 0 \mid v_s^q[M](t) > 0 \quad \forall s \in (b - \rho^q(t), b).$$

Moreover, if $M > \lambda_4^q$, then $v_s^q[M]$ is cannot be positive.

Step 3. Study of the function $v_s^q[M]$ at $t = a$.

Let us denote

$$\widehat{g}_{(-1)^n M}(t, s) = \begin{cases} \widehat{g}_{(-1)^n M}^1(t, s), & a \leq s \leq t \leq b, \\ \widehat{g}_{(-1)^n M}^2(t, s), & a < t < s < b, \end{cases}$$

as the related Green's function of $\widehat{T}_n[(-1)^n M]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{*\{\varepsilon_1, \dots, \varepsilon_{n-k}\}} = X_{\{\tau_1, \dots, \tau_{n-k}\}}^{\{\delta_1, \dots, \delta_k\}}$.

In this case, we study the following function by applying similar reasoning to Step 1:

$$\widehat{w}_{M,q}(t) = (-1)^n \frac{\partial^{\alpha^q+q}}{\partial s^{\alpha^q+q}} \widehat{g}_{(-1)^n M}^1(t, s)|_{s=a},$$

which is equivalent, using equality (2.12), to study the function

$$\widehat{w}_{M,q}(s) = \frac{\partial^{\alpha^q+q}}{\partial t^{\alpha^q+q}} g_M^2(t, s)|_{t=a} = \frac{\partial^{\alpha^q}}{\partial t^{\alpha^q}} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right) |_{t=a}, \quad s \in I.$$

In this case, if $\alpha^q > 0$ using that $\{0, \dots, \alpha^q - 1\} \subset \{\mu_1^q, \dots, \mu_k^q\}$ we have

$$\frac{\partial^q}{\partial t^q} g_M^2(a, s) = \frac{\partial}{\partial t} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right) |_{t=a} = \dots = \frac{\partial^{\alpha^q-1}}{\partial t^{\alpha^q-1}} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right) |_{t=a} = 0, \quad \forall s \in (a, b).$$

As in the Steps 1 and 2 we can deduce that

$$\widehat{T}_n[(-1)^n M] \widehat{w}_{M,q}(t) = 0, \quad t \in (a, b).$$

Using the arguments of Step 1, we can affirm that if there exists $t^* \in (a, b)$ such that $\widehat{w}_{M,q}(t^*) < 0$, then $v_s^q[M]$ is not positive.

By the first row of (2.17) and (2.16) we deduce that:

$$\frac{\partial^{\tau_1}}{\partial t^{\tau_1}} \widehat{g}_{(-1)^n M}^2(t, s)|_{t=a} = 0, \quad -\frac{\partial^{\tau_1+1}}{\partial t^{\tau_1} \partial s} \widehat{g}_{(-1)^n M}^2(t, s)|_{t=a} = 0, \quad \dots, \quad (-1)^{\alpha^q+q} \frac{\partial^{\tau_1+\alpha^q+q}}{\partial t^{\tau_1} \partial s^{\alpha^q+q}} \widehat{g}_{(-1)^n M}^2(t, s)|_{t=a} = 0.$$

Since $(\tau_1 + \alpha^q + q) \bmod n < n - 1$, we do not reach any diagonal element, hence the previous equalities are satisfied for $s = a$, and we obtain:

$$\left\{ \begin{array}{l} \frac{\partial^{\tau_1}}{\partial t^{\tau_1}} \widehat{g}_{(-1)^n M}^1(t, s)|_{(t,s)=(a,a)} = 0, \\ -\frac{\partial^{\tau_1+1}}{\partial t^{\tau_1} \partial s} \widehat{g}_{(-1)^n M}^1(t, s)|_{(t,s)=(a,a)} = 0, \\ \vdots \\ (-1)^{\alpha^q+q} \frac{\partial^{\tau_1+\alpha^q+q}}{\partial t^{\tau_1} \partial s^{\alpha^q+q}} \widehat{g}_{(-1)^n M}^1(t, s)|_{(t,s)=(a,a)} = 0. \end{array} \right.$$

So, we conclude that $\widehat{w}_{M,q}^{(\tau_1)}(a) = 0$.

Analogously we obtain that

$$\widehat{w}_{M,q}^{(\tau_2)}(a) = \dots = \widehat{w}_{M,q}^{(\tau_{h-1})}(a) = \widehat{w}_{M,q}^{(\tau_{h+1})}(a) = \dots = \widehat{w}_{M,q}^{(\tau_{n-k})}(a) = 0.$$

For τ_h , we have the following equalities

$$\frac{\partial^{\tau_h}}{\partial t^{\tau_h}} \widehat{g}_{(-1)^n M}^2(t, s)|_{t=a} = 0, -\frac{\partial^{\tau_h+1}}{\partial t^{\tau_h} \partial s} \widehat{g}_{(-1)^n M}^2(t, s)|_{t=a} = 0, \dots, (-1)^{\alpha^q+q} \frac{\partial^{\tau_h+\alpha^q+q}}{\partial t^{\tau_h} \partial s^{\alpha^q+q}} \widehat{g}_{(-1)^n M}^2(t, s)|_{t=a} = 0.$$

In this case, since $\tau_h + \alpha^q + q = n - 1$, we reach a diagonal element of $\widehat{G}(t, s)$ given in (2.15), and as consequence we obtain the following equalities for $s = a$:

$$\left\{ \begin{array}{l} \frac{\partial^{\tau_h}}{\partial t^{\tau_h}} \widehat{g}_{(-1)^n M}^1(t, s)|_{(t,s)=(a,a)} = 0, \\ -\frac{\partial^{\tau_h+1}}{\partial t^{\tau_h} \partial s} \widehat{g}_{(-1)^n M}^1(t, s)|_{(t,s)=(a,a)} = 0, \\ \vdots \\ (-1)^{\alpha^q+q} \frac{\partial^{\tau_h+\alpha^q+q}}{\partial t^{\tau_h} \partial s^{\alpha^q+q}} \widehat{g}_{(-1)^n M}^1(t, s)|_{(t,s)=(a,a)} = 1. \end{array} \right.$$

Therefore, $\widehat{w}_{M,q}^{(\tau_h)}(a) = (-1)^{n-\alpha^q-q}$.

Now, let us study the behavior of $\widehat{w}_{M,q}$ at $t = b$. Studying the $(n - k + 1)^{\text{th}}$ row of (2.17), we have for all $s \in (a, b)$:

$$\frac{\partial^{\delta_1}}{\partial t^{\delta_1}} \widehat{g}_{(-1)^n M}^1(t, s)|_{t=b} = 0, -\frac{\partial^{\delta_1+1}}{\partial t^{\delta_1} \partial s} \widehat{g}_{(-1)^n M}^1(t, s)|_{t=b} = 0, \dots, (-1)^{\alpha^q+q} \frac{\partial^{\delta_1+\alpha^q+q}}{\partial t^{\delta_1} \partial s^{\alpha^q+q}} \widehat{g}_{(-1)^n M}^1(t, s)|_{t=b} = 0.$$

The above equalities are satisfied at $s = a$ and since $b \neq a$ we have that $\widehat{w}_{M,q}^{(\delta_1)}(b) = 0$.

Analogously we obtain that

$$\widehat{w}_{M,q}^{(\delta_2)}(b) = \dots = \widehat{w}_{M,q}^{(\delta_k)}(b) = 0.$$

Note that if $\alpha^q = 0$, then from the same matrix argument above we deduce that the function $\widehat{w}_{M,q}(s) = \frac{\partial^q}{\partial t^q} g_M^2(a, s)$ verifies the same boundary conditions:

$$\begin{aligned} \widehat{w}_{M,q}^{(\tau_2)}(a) = \dots = \widehat{w}_{M,q}^{(\tau_{h-1})}(a) = \widehat{w}_{M,q}^{(\tau_{h+1})}(a) = \dots = \widehat{w}_{M,q}^{(\tau_{n-k})}(a) &= 0, \\ \widehat{w}_{M,q}^{(\delta_2)}(b) = \dots = \widehat{w}_{M,q}^{(\delta_k)}(b) &= 0. \end{aligned}$$

From (2.11) and (2.9) we deduce that

$$\widehat{T}_n[0] v(t) := (-1)^n T_n^*[0] v(t) = T_n[0] v(t).$$

Hence, using arguments similar to Step 1, we conclude that $\widehat{w}_{M,q} > 0$ on (a, b) for all $M \in [0, (\lambda_4^*)^q]$, where $(\lambda_4^*)^q > 0$ is the least positive eigenvalue of $\widehat{T}_n[0] \equiv T_n[0]$ in X_4^* defined in (3.12).

From Lemma 3.10, we know that $(\lambda_4^*)^q = \lambda_4^q$. Thus, from this step we have that, if $M \in [0, \lambda_4^q]$, then

$$\forall s \in (a, b), \quad \exists \rho^q(s) > 0 \mid v_s^q[M](t) > 0 \quad \forall t \in (a, a + \rho^q(s)).$$

Step 4. Study of the function $v_s^q[M]$ at $t = b$.

In this case, we study the following function by applying similar reasoning to Step 3

$$\widehat{y}_{M,q}(t) = (-1)^n \frac{\partial^{\beta^q+q}}{\partial s^{\beta^q+q}} \widehat{g}_{(-1)^n M}^2(t, s)|_{s=b},$$

which is equivalent using equality (2.12) to studying the function

$$\widehat{y}_{M,q}(s) = \frac{\partial^{\beta^q+q}}{\partial t^{\beta^q+q}} g_M^1(t, s)|_{t=b} = \frac{\partial^{\beta^q}}{\partial t^{\beta^q}} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right) |_{t=b}.$$

Since $\{0, \dots, \beta^q\} \subset \{\rho_1^q, \dots, \rho_{n-k}\}$ if $\beta^q > 0$ we obtain that

$$\frac{\partial^q}{\partial t^q} g_M(b, t) = \frac{\partial}{\partial t} \left(\frac{\partial^q}{\partial t^q} g_M(t, s) \right) |_{t=b} = \dots = \frac{\partial^{\beta^q-1}}{\partial t^{\beta^q-1}} \left(\frac{\partial^q}{\partial t^q} g_M(t, s) \right) |_{t=b} = 0.$$

Moreover, using arguments analogous to Step 1 we obtain that if β^q is even, then $\widehat{y}_{M,q} \geq 0$ and if β^q is odd, then $\widehat{y}_{M,q} \leq 0$.

Again, we have that

$$\widehat{T}_n[(-1)^n M] \widehat{y}_{M,q}(t) = 0, \quad \forall t \in [a, b].$$

A study analogous to Step 3 leads us to the fact that $\widehat{y}_{M,q}$ satisfies the boundary conditions

$$\begin{aligned} \widehat{y}_{M,q}^{(\tau_1)}(a) &= \dots = \widehat{y}_{M,q}^{(\tau_{n-k})}(a) = 0, \\ \widehat{y}_{M,q}^{(\delta_1)}(b) &= \dots = \widehat{y}_{M,q}^{(\delta_{z-1})}(b) = \widehat{y}_{M,q}^{(\delta_{z+1})}(b) = \dots = \widehat{y}_{M,q}^{(\delta_k)}(b) = 0, \\ y_{M,q}^{(\delta_z)}(b) &= (-1)^{n-\beta^q-q+1}. \end{aligned} \tag{3.15}$$

Analogously, if $\beta^q = 0$, then $\widehat{y}_{M,q}(s) = \frac{\partial^q}{\partial t^q} g_M^2(b, s)$ satisfies the conditions (3.15) by an argument similar to the case $\beta^q > 0$.

Thus, we deduce that $\widehat{y}_{M,q} > 0$ on (a, b) if β^q is even and $\widehat{y}_{M,q} < 0$ on (a, b) if β^q is odd for all $M \in [0, (\lambda_2^*)^q]$, where $(\lambda_2^*)^q > 0$ is the least positive eigenvalue of $\widehat{T}_n[0] \equiv T_n[0]$ in X_2^* defined in (3.11).

Again, from Lemma 3.9 we have that $(\lambda_2^*)^q = \lambda_2^q$. Therefore, we conclude from this step that, if $M \in [0, \lambda_2^q]$, then

$$\forall s \in (a, b), \quad \exists \rho^q(s) > 0 \mid v_s^q[M](t) > 0 \quad \forall t \in (b - \rho^q(t), b).$$

Step 5. Study of the function $v_s^q[M]$ on $(a, b) \times (a, b)$.

In this step, we verify that the function $v_s^q[M](t) > 0$ for all $(t, s) \in I \times I$ if M belongs to the given intervals. Let us denote $u_M^s(t) = g_M(t, s)$, for all $s \in (a, b)$.

The function $v_s^q[M](t)$ is in $C^{n-2-q}(I)$ and it satisfies the boundary conditions (2.18)-(2.19).

Using the definition of $v_s^q[M]$, the following equality holds for all $s \in (a, b)$:

$$T_{n-q}[0] v_s^q[M](t) = -M u_M^s(t), \quad \forall t \neq s, t \in I. \quad (3.16)$$

Let's see that for the values of the parameter M for which $v_s^q[M](t)$ has constant sign in I , it cannot have a double zero in (a, b) , and this implies that the change of the sign must be at $t = a$ or $t = b$, and the result is proven.

We do the proof for $n - k$ even (the case $n - k$ odd is similarly proved). In this case, we study the behavior of $v_s^q[M](t)$ for $M > 0$ and $u_M^s \geq 0$. From (3.16), we have that

$$\frac{d^{n-q}}{dt^{n-q}} v_s^q[M](t) = \frac{\partial^n}{\partial t^n} g_M(t, s) \leq 0.$$

Therefore, using the condition (T_d) , since $v_1 = \dots = v_n = 1$, we have that

$$\frac{d^{n-q-1}}{dt^{n-q-1}} v_s^q[M](t) = \frac{\partial^{n-1}}{\partial t^{n-1}} g_M(t, s)$$

is a decreasing function, with two continuous components and a jump $+1$ at $t = s$. Then, it has at most two zeros on I (see Figure 1).




Figure 1: $\frac{d^{n-q-1}}{dt^{n-q-1}} v_s^q[M](t)$, maximal oscillation with $t \in I = [0, 1]$.

Now, we have that

$$\frac{d^{n-q-2}}{dt^{n-q-2}} v_s^q[M](t) = \frac{\partial^{n-2}}{\partial t^{n-2}} g_M(t, s)$$




Figure 2: $\frac{d^{n-q-1}}{dt^{n-q-1}} v_s^q[M](t)$, maximal oscillation with $t \in I = [0, 1]$.

is a continuous function with at most four zeros on I (See Figure 2).

By recurrence, we have that

$$\frac{d^{n-q-l}}{dt^{n-q-l}} v_s^q[M](t) = \frac{\partial^{n-l}}{\partial t^{n-l}} g_M(t, s)$$

has at most $l + 2$ zeros on I . In particular, $v_s^q[M]$ has at most $n - q + 2$ zeros on I . Moreover, since

$$\begin{aligned} (v_s^q)^{(\sigma_1^q)}[M](a) &= \dots = (v_s^q)^{(\sigma_{c_q}^q)}[M](a) = 0, \\ (v_s^q)^{(\varepsilon_1^q)}[M](b) &= \dots = (v_s^q)^{(\varepsilon_{d_q}^q)}[M](b) = 0, \end{aligned}$$

$v_s^q[M]$ has $c_q + d_q = n - q$ zeros on the boundary. Thus, $v_s^q[M]$ has at most two zeros on (a, b) .

Now, making an argument similar to the one made in [5, Theorem 5.1], we deduce that the maximal oscillation (that is, the maximal number of zeros) is possible if and only if

$$\begin{cases} (v_s^q)^{(\alpha^q)}[M](a) \leq 0, & \text{if } n - q - c_q \text{ is even,} \\ (v_s^q)^{(\alpha^q)}[M](a) \geq 0, & \text{if } n - q - c_q \text{ is odd.} \end{cases} \quad (3.17)$$

Since we are in the case $n - q - c_q$ even, the maximal oscillation is only possible if $(v_s^q)^{(\alpha^q)}[M](a) \leq 0$. However, by definition, $(v_s^q)^{(\alpha^q)}[M](a) \neq 0$ and, since $v_s^q[M]$ is non-negative and α^q denotes the order of the smallest derivative at $t = a$ which is different from zero, necessarily $(v_s^q)^{(\alpha^q)}[M](a) > 0$. Therefore the maximal oscillation can not happen (that is, $v_s^q[M]$ cannot have two zero on (a, b)). Thus, $v_s^q[M]$ has at most one simple zero on (a, b) ,

which is not possible due to the constant sign of $v_s^q[M]$, and we conclude that $v_s^q[M]$ does not have any zero on (a, b) .

From this step we deduce that if $M > 0$ and $u_M^s \geq 0$, then $v_s^q[M] > 0$ on (a, b) .

In conclusion, from the previous steps the result is proved. \square

In the sequel, we present some examples of application of the previous result.

Example 3.12. Consider again the fourth order operator $T_4[M]$ coupled with the boundary conditions $X_{\{0,1,2\}}^{\{2\}}$ defined in Example 2.8.

The functions $\frac{\partial}{\partial t}g_M$ and $\frac{\partial^2}{\partial t^2}g_M$ satisfy the conditions of spaces $X_{\{0,1,3\}}^{\{1\}}$ and $X_{\{0,2,3\}}^{\{0\}}$, respectively.

In this case $k = 3$ and $n - k = 1$ is odd. For $q = 1$, we have that $c_q = 2$, $d_q = 1$, $z = 2 \neq 1$ and $\beta^1 = 0$. By applying Theorem 3.11 it follows that $\frac{\partial}{\partial t}g_M$ is strongly negative on $[0, 1] \times [0, 1]$ if, and only if $M \in [\lambda_2^1, \lambda_1)$, where

- * $\lambda_1 > 0$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1,2\}}^{\{2\}}$.
- * $\lambda_2^1 < 0$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{0,3\}}^{\{0,1\}}$.

The eigenvalues of $T_4[0]$ in $X_{\{0,1,2\}}^{\{2\}}$ are given by λ^4 , where λ is a positive solution of

$$\tanh(\lambda\sqrt{2}) + \tan\left(\frac{\lambda}{\sqrt{2}}\right) = 0.$$

Let us denote m_1 as the smallest positive solution of this equation. Then, $\lambda_1 = m_1^4$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1,2\}}^{\{2\}}$.

In the same way, the eigenvalues of $T_4[0]$ in $X_{\{0,3\}}^{\{0,1\}}$ are given by $-\lambda^4$, where λ is a positive solution of $\sin(\lambda) = 0$.

In this case, $\lambda_{min}^1 = \pi$ is the smallest positive solution of this equation. Then, $\lambda_2^1 = -(\lambda_{min}^1)^4 = -\pi^4$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{0,3\}}^{\{0,1\}}$.

By numerical approach, it can be seen that $\lambda_1 = m_1^4 \approx 3.34^4$.

For $q = 2$, we have that $c_q = n - q - 1 = 1$, $d_q = 1$, $z = 1$ and $\beta^2 = 1$. Again, by applying Theorem 3.11 it follows that $\frac{\partial^2}{\partial t^2}g_M$ is strongly negative on $[0, 1] \times [0, 1]$ if, and only if $M \in [\lambda_2^2, \lambda_1)$, where

- * $\lambda_1 > 0$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1,2\}}^{\{2\}}$.
- * $\lambda_2^2 < 0$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{2,3\}}^{\{0,1\}}$.

The eigenvalues of $T_4[0]$ in $X_{\{2,3\}}^{\{0,1\}}$ are given by $-\lambda^4$, where λ is a least positive solution of

$$2e^{2\lambda} + (e^{2\lambda} + 1) \cos(\lambda) = 0.$$

By denoting λ_{min}^2 as the smallest positive solution of this equation, then $\lambda_2^2 = -(\lambda_{min}^2)^4$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{2,3\}}^{\{0,1\}}$.

In this case, we have that $\lambda_2^2 = -(\lambda_{min}^2)^4 \approx -1.87^4$.

Example 3.13. We consider the fourth order operator $T_4[M]$ defined in the space of the boundary conditions $X_{\{0,1\}}^{\{1,3\}}$.

In this case $k = 2$ and $n - k$ is even. For $q = 1$ we have that $c_q = 1$, $d_q = n - q - c_q = 2$, $z = 1$, $h = 2$ and $\alpha^1 = \beta^1 = 1$. Moreover, the function $\frac{\partial}{\partial t}g_M$ satisfies the conditions of space $X_{\{0,3\}}^{\{0,2\}}$.

By applying Theorem 3.11 we obtain that $\frac{\partial}{\partial t}g_M$ is strongly positive on $[0, 1] \times [0, 1]$ if, and only if $M \in (\lambda_1, \lambda^1]$, where

- * $\lambda_1 < 0$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1\}}^{\{1,3\}}$.
- * $\lambda^1 > 0$ is the minimum between:
 - $\lambda_2^1 > 0$, the least positive eigenvalue of $T_4[0]$ in $X_{\{3\}}^{\{0,1,2\}}$.
 - $\lambda_4^1 > 0$ is the least positive eigenvalue of $T_4[0]$ in $X_{\{0,1,3\}}^{\{0\}}$.

By numerical approximation, we obtain that $\lambda_1 \approx -2.36^4$, $\lambda_2^1 \approx 2.22^4$ and $\lambda_4^1 \approx 4.44^4$. Therefore, $\frac{\partial}{\partial t}g_M$ is strongly positive on $[0, 1] \times [0, 1]$ if, and only if, $M \in (-2.36^4, 2.22^4]$.

As a consequence of the Theorem 3.11 we arrive at the following result.

Corollary 3.14. *Let $q \in \{1, \dots, n - 1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$. If either $\sigma_k = q + c_q - 1$ or $\varepsilon_{n-k} = q + d_q - 1$, we have the following properties:*

- If $n - q - c_q$ is even, then there is not any $M \in \mathbb{R}$ such that $v_s^q[M]$ is strongly negative on $I \times I$.
- If $n - q - c_q$ is odd, then there is not any $M \in \mathbb{R}$ such that $v_s^q[M]$ is strongly positive on $I \times I$.

Proof. If $\sigma_k = q + c_q - 1$, then $\eta = n - 1 - \sigma_k = n - q - c_q$.

We consider

$$w_{M,q}(t) = \frac{\partial^\eta}{\partial s^\eta} \left(\frac{\partial^q}{\partial t^q} g_M^1(t, s) \right) \Big|_{s=a},$$

defined in Step 1 of the proof of Theorem 3.11.

We know that this function satisfies the following boundary conditions:

$$\begin{aligned} w_{M,q}^{(\mu_1^q)}(a) &= \dots = w_{M,q}^{(\mu_{z-1}^q)}(a) = w_{M,q}^{(\mu_{z+1}^q)}(a) = \dots = w_{M,q}^{(\mu_k^q)}(a) = 0, \\ w_{M,q}^{(\rho_1^q)}(b) &= \dots = w_{M,q}^{(\rho_{n-k}^q)}(b) = 0, \\ w_{M,q}^{(\mu_z^q)}(a) &= (-1)^\eta = (-1)^{n-q-c_q}. \end{aligned}$$

Hence, if $n - q - c_q$ is even, then there exists $\rho > 0$, such that $w_{M,q}(t) > 0$ for all $t \in (a, a + \rho)$. So, $v_s^q[M]$ cannot be negative for any real M .

Now, if $n - q - c_q$ is odd, then there exists $\rho > 0$, such that $w_{M,q}(t) < 0$ for all $t \in (a, a + \rho)$. Thus, $v_s^q[M]$ cannot be positive for any $M \in \mathbb{R}$.

Analogously, if $\varepsilon_{n-k} = q + d_q - 1$, then $\gamma = n - \varepsilon_{n-k} - 1 = n - q - d_q = c_q$. In this case, we consider the function

$$y_{M,q}(t) = \frac{\partial^\gamma}{\partial s^\gamma} \left(\frac{\partial^q}{\partial t^q} g_M^2(t, s) \right)_{|s=b},$$

defined in Step 2 of the proof of Theorem 3.11.

We know that for all $M \in \mathbb{R}$, the function $y_{M,q}$ satisfies the following boundary conditions:

$$\begin{aligned} y_{M,q}^{(\mu_1^q)}(a) &= \dots = y_{M,q}^{(\mu_k^q)}(a) = 0, \\ y_{M,q}^{(\rho_1^q)}(b) &= \dots = y_{M,q}^{(\rho_{h-1}^q)}(b) = y_{M,q}^{(\rho_{h+1}^q)}(b) = \dots = y_{M,q}^{(\rho_{n-k}^q)}(b) = 0, \\ y_{M,q}^{(\rho_h^q)}(b) &= (-1)^{c_q+1}. \end{aligned}$$

Hence, if $n - q - c_q$ and c_q are even, then there exists $\rho > 0$, such that $y_{M,q}(t) > 0$ for all $t \in (b - \rho, b)$. So, $v_s^q[M]$ cannot be negative for any real M .

Moreover, if $n - q - c_q$ is even and c_q odd, then there exists $\rho > 0$, such that $y_{M,q}(t) < 0$ for all $t \in (b - \rho, b)$. So, $v_s^q[M]$ cannot be positive for any real M .

Now, if $n - q - c_q$ is odd and c_q even, then there exists $\rho > 0$, such that $y_{M,q}(t) < 0$ for all $t \in (b - \rho, b)$. So, $v_s^q[M]$ cannot be positive for any real M .

Finally, if $n - q - c_q$ and c_q are odd, then there exists $\rho > 0$, such that $y_{M,q}(t) > 0$ for all $t \in (b - \rho, b)$.

As consequence, $v_s^q[M]$ cannot be negative for any real M . \square

Now, we give a necessary condition of the nonpositive (nonnegative) sign of $v_s^q[M]$. This characterization will be an interval in which $v_s^q[M]$ has a nonpositive (nonnegative) sign whose infimum or supremum is given by the first eigenvalue λ_1 of $T_n[M]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

Theorem 3.15. *Let $q \in \{1, \dots, n - 1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q \geq 1$ and $d_q \geq 1$. If $\mu_z^q \neq z - 1$ and $\rho_h^q \neq h - 1$, then the following properties are fulfilled:*

- *Let $n - k$ be even:*

- *If $n - q - c_q$ is even and $v_s^q[M]$ is nonpositive on $I \times I$, then $M \in [\lambda_*^q, \lambda_1)$, and if $n - q - c_q$ is odd and $v_s^q[M]$ is nonnegative on $I \times I$, then $M \in [\lambda_*^q, \lambda_1)$, where*

- * $\lambda_1 < 0$ *is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*

- * $\lambda_*^q < 0$ *is the maximum between:*

- $\lambda_3^q < 0$ *is the biggest negative eigenvalue of $T_n[0]$ in X_3 defined in (3.2).*

- $\lambda_5^q < 0$, *the biggest negative eigenvalue of $T_n[0]$ in X_5 defined in (3.4).*

• Let $n - k$ be odd:

– If $n - q - c_q$ is even and $v_s^q[M]$ is nonpositive on $I \times I$, then $M \in (\lambda_1, \lambda_*^q]$, and if $n - q - c_q$ is odd and $v_s^q[M]$ is nonnegative on $I \times I$, then $M \in (\lambda_1, \lambda_*^q]$ where

* $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.

* $\lambda_*^q > 0$ is the minimum between:

· $\lambda_3^q > 0$, the biggest negative eigenvalue of $T_n[0]$ in X_3 .

· $\lambda_5^q > 0$, the biggest negative eigenvalue of $T_n[0]$ in X_5 .

Proof. Since $\mu_z^q \neq z - 1$ and $\rho_h^q \neq h - 1$ we have from Propositions 3.5 and 3.6 that there are eigenvalues λ_3^q and λ_5^q with corresponding related eigenfunctions of constant sign. We suppose that $n - k$ and $n - q - c_q$ are even. For the other cases the proof is done in an analogous way.

Let us assume that there exists $M^* \notin [\lambda_*^q, \lambda_1)$, such that $v_s^q[M^*]$ is nonpositive on $I \times I$. From Theorem 3.11, we can affirm that $M^* < \lambda_1$. So, $M^* < \lambda_*^q$ and for all $M \in [M^*, \lambda_1)$ the function $v_s^q[M]$ is nonpositive on $I \times I$ because the set where it takes nonpositive values is an interval. By the monotone decreasing character of $v_s^q[M]$ we infer that

$$v_s^q[\lambda_*^q](t) \leq v_s^q[M](t) \leq v_s^q[M^*](t) \leq 0.$$

So, in particular

$$w_{\lambda_*^q, q}(t) \leq w_{M, q}(t) \leq w_{M^*, q}(t) \leq 0,$$

and

$$\begin{cases} y_{\lambda_*^q, q}(t) \leq y_{M, q}(t) \leq y_{M^*, q}(t) \leq 0, & \text{if } \gamma \text{ is even,} \\ 0 \leq y_{M^*, q}(t) \leq y_{M, q}(t) \leq y_{\lambda_*^q, q}(t), & \text{if } \gamma \text{ is odd.} \end{cases}$$

If $\lambda_*^q = \lambda_3^q$, then $w_{\lambda_*^q, q}^{(\alpha^q)}(a) = 0$. So, we conclude that, for all $M \in [M^*, \lambda_*^q)$, $w_{M, q}^{(\alpha^q)}(a) = 0$, which contradicts the discrete character of the spectrum of $T_n[0]$ in X_3 .

If $\lambda_*^q = \lambda_5^q$, then $y_{\lambda_*^q, q}^{(\beta^q)}(b) = 0$. So, we conclude that, for all $M \in [M^*, \lambda_*^q)$, $y_{M, q}^{(\beta^q)}(b) = 0$, which contradicts the discrete character of the spectrum of $T_n[0]$ in X_5 .

So, we arrive to a contradiction, thus the result is proved. \square

Next, we present an example to illustrate the previous result.

Example 3.16. Consider the operator $T_5[M]$ defined in the space of the boundary conditions $X_{\{0,2,3\}}^{\{1,3\}}$.

For $q = 1$ we have that $c_q = 2$, $n - q - c_q = d_q = 2$ is even, $z = h = 2$, $\alpha^1 = 0$ and $\beta^1 = 1$. Moreover, the function $\frac{\partial}{\partial t} g_M$ satisfies the conditions of space $X_{\{1,2,4\}}^{\{0,2\}}$ that fulfills the assumptions of Theorem 3.15. Therefore, if $\frac{\partial}{\partial t} g_M$ is nonpositive on $[0, 1] \times [0, 1]$, then $M \in [\lambda_*^1, \lambda^1)$, where

* $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_5[0]$ in $X_{\{0,2,3\}}^{\{1,3\}}$.

* $\lambda_*^1 < 0$ is the maximum between:

· $\lambda_3^1 < 0$, the biggest negative eigenvalue of $T_5[0]$ in $X_{\{0,1,4\}}^{\{0,2\}}$.

· $\lambda_5^1 < 0$ is the biggest negative eigenvalue of $T_5[0]$ in $X_{\{1,2,4\}}^{\{0,1\}}$.

By numerical approximation, we obtain that $\lambda_1 \approx -2.23^5$, $\lambda_3^1 \approx -3.67^5$ and $\lambda_5^1 \approx -2.88^5$. Thus, if $\frac{\partial}{\partial} g_M$ is nonpositive on $[0, 1] \times [0, 1]$, then $M \in [-2.88^5, -2.23^5)$.

3.2 Case (b): $c_q = n - q$ and $d_q = 0$

In this subsection we deal with the constant sign of $v_s^q[M]$ for the case $c_q = n - q$ and $d_q = 0$. We arrive to the next theorem.

Theorem 3.17. *Let $q \in \{1, \dots, n - 1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q = n - q$ and $d_q = 0$. The following properties are fulfilled:*

- *If $n - k$ is even, then $v_s^q[M]$ is nonnegative on $I \times I$ if, and only if, $M \in (\lambda_1, 0]$, where $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*
- *If $n - k$ is odd, then $v_s^q[M]$ is nonnegative on $I \times I$ if, and only if, $M \in [0, \lambda_1)$, where $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*

Proof. Let us assume that $n - k$ is even. For the case $n - k$ odd the proof is done in an analogous way. We know from the Theorem 2.28 that $v_s^q \in X_{\{0, \dots, c_q\}}$ satisfies an initial problem and it is nonnegative on $[a, b]$. Moreover, $v_s^q(t) = \frac{\partial^q}{\partial t^q} g_0(t, s) > 0$ if $s < t$ and $v_s^q(t) = \frac{\partial^q}{\partial t^q} g_0(t, s) = 0$ if $s > t$.

Taking into account Lemma 2.30, we obtain that $v_s^q(t) = \frac{\partial^q}{\partial t^q} g_0(t, s) > 0$ on $(a, b) \times (a, b)$ if and only if $M \in (\lambda_1, M_q]$, with $M_q > \lambda_1$. Moreover, $\frac{\partial^q}{\partial t^q} g_M$ is monotone decreasing with respect to $M \in (\lambda_1, M_q]$.

Let's see that $M_q = 0$. Using the monotone decreasing character of $v_s^q[M]$ with respect to $M \in (\lambda_1, M_q]$ and since $v_s^q(t) = \frac{\partial^q}{\partial t^q} g_0(t, s) = 0$ if $s > t$ we have that $v_s^q[M_q](t) < v_s^q(t) = 0$ if $M_q > 0$. On the other hand, if $s < t$ then there exists $(t^*, s^*) \in (a, b) \times (a, b)$ with $s^* < t^*$ such that $0 < v_{s^*}^q[M_q](t^*) < v_{s^*}^q(t^*)$ if $M_q > 0$. Thus, $v_s^q[M]$ changes sign for $M_q > 0$ and this completes the proof. \square

Example 3.18. Consider the operator $T_4[M]$ coupled with the boundary conditions in the space $X_{\{2,3\}}^{\{0,1\}}$.

In this case $n - k$ is even. For $q = 2$ we have that $c_q = n - q = 2$ and $d_q = 0$. Then, by Theorem 3.17 we have that $\frac{\partial^2}{\partial t^2} g_M$ is nonnegative on $[0, 1] \times [0, 1]$ if, and only if, $M \in (\lambda_1, 0]$, where $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_4[0]$ in $X_{\{2,3\}}^{\{0,1\}}$.

By numerical approximation, we obtain that $\lambda_1 \approx -1.8751^4$. Thus, $\frac{\partial^2}{\partial t^2} g_M$ is nonnegative on $[0, 1] \times [0, 1]$ if, and only if, $M \in (-1.8751^4, 0]$.

For $q = 3$ we have that $c_q = n - q = 1$ and $d_q = 0$. Again, by Theorem 3.17 we have that $\frac{\partial^3}{\partial t^3} g_M$ is nonnegative on $[0, 1] \times [0, 1]$ if, and only if, $M \in (-1.8751^4, 0]$.

3.3 Case (c): $c_q = 0$ and $d_q = n - q$

In this section, arguing in a similar manner than in previous one, we can characterize the constant sign of $v_s^q[M]$ for the case $c_q = 0$ and $d_q = n - q$.

Using Theorem 2.28, Lemma 2.30, and reasoning in an analogous way to the proof of Theorem 3.17, we get to the next result.

Theorem 3.19. *Let $q \in \{1, \dots, n - 1\}$. Suppose that $\{\sigma_1, \dots, \sigma_k\} - \{\varepsilon_1, \dots, \varepsilon_{n-k}\}$ satisfies condition (N_a) , $c_q + d_q = n - q$, $c_q = 0$ and $d_q = n - q$. Then, the following properties are fulfilled:*

- *Let $n - k$ be even:*
 - *If $n - q$ is even, then $v_s^q[M]$ is nonnegative on $I \times I$ if, and only if, $M \in (\lambda_1, 0]$, where $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*
 - *If $n - q$ is odd, then $v_s^q[M]$ is nonpositive on $I \times I$ if, and only if, $M \in (\lambda_1, 0]$, where $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*
- *Let $n - k$ be odd:*
 - *If $n - q$ is even, then $v_s^q[M]$ is nonnegative on $I \times I$ if, and only if, $M \in [0, \lambda_1)$, where $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*
 - *If $n - q$ is odd, then $v_s^q[M]$ is nonpositive on $I \times I$ if, and only if, $M \in [0, \lambda_1)$, where $\lambda_1 > 0$ is the least positive eigenvalue of $T_n[0]$ in $X_{\{\sigma_1, \dots, \sigma_k\}}^{\{\varepsilon_1, \dots, \varepsilon_{n-k}\}}$.*

Example 3.20. Consider the operator $T_6[M]$ coupled with the boundary conditions in the space $X_{\{0,2\}}^{\{1,3,4,5\}}$.

In this case $n - k = 4$ is even. For $q = 3$ we have that $c_q = 0$ and $d_q = n - q = 3$ is odd. Then, by Theorem 3.19 we have that $\frac{\partial^3}{\partial t^3} g_M$ is nonpositive on $[0, 1] \times [0, 1]$ if, and only if, $M \in (\lambda_1, 0]$, where $\lambda_1 < 0$ is the biggest negative eigenvalue of $T_6[0]$ in $X_{\{0,2\}}^{\{1,3,4,5\}}$.

By numerical approximation, we obtain that $\lambda_1 \approx -1.953^6$. Thus, $\frac{\partial^3}{\partial t^3} g_M$ is nonpositive on $[0, 1] \times [0, 1]$ if, and only if, $M \in (-1.953^6, 0]$.

For $q = 4$ we have that $c_q = 0$ and $d_q = n - q = 2$ is even. Again, from Theorem 3.19 we have that $\frac{\partial^4}{\partial t^4} g_M$ is nonnegative on $[0, 1] \times [0, 1]$ if, and only if, $M \in (-1.953^6, 0]$.

Finally, for $q = 5$ we have that $c_q = 0$ and $d_q = n - q = 1$ is odd. Then, $\frac{\partial^5}{\partial t^5} g_M$ is nonpositive on $[0, 1] \times [0, 1]$ if, and only if, $M \in (-1.953^6, 0]$.

References

- [1] A. Cabada, *Green's functions in the theory of ordinary differential equations*, Springer Briefs in Mathematics, Springer, New York, NY (2014).
- [2] A. Cabada, L. López-Somoza *An explicit formula of the parameter dependence of the partial derivatives of the Green's functions related to arbitrary two-point boundary conditions*. arXiv preprint arXiv: 2405.17320.
- [3] A. Cabada, L. Saavedra, *The eigenvalue characterization for the constant sign Green's functions of $(k, n-k)$ problems*, Boundary Value Problems, SpringerOpen Journal, (2016).
- [4] A. Cabada, L. Saavedra, *Disconjugacy characterization by means of spectral $(k; n - k)$ problems*, *Appl. Math. Lett.*, 52 (2016), 21-29.
- [5] A. Cabada, L. Saavedra, *Characterization of constant sign Green's function for a two-point boundary-value problem by means of spectral theory*, *Electronic Journal of Differential Equations*, Vol. 2017 (2017), No. 146, pp. 1–95.
- [6] W.A. Coppel, *Disconjugacy*, in: Lecture Notes in Mathematics, vol. 220, Springer-Verlag, Berlin, New York, 1971.
- [7] C. De Coster, P. Habets, *Two-Point Boundary Value Problems: Lower and Upper Solutions*, Mathematics in Science and Engineering 205, Elsevier B. V., Amsterdam, 2006.
- [8] U. Elias, *Eventual disconjugacy of $y^{(n)} + \mu p(x)y = 0$ for every μ* . *Arch.Math.* 40(2), 193-200 (2004).
- [9] M. A. Krasnosel'skiĭ, *Positive Solutions of Operator Equations*, Noordhoff, Groningen, 1964.
- [10] G. S. Ladde, V. Lakshmikantham, A. S. Vatsala, *Monotone Iterative Techniques for Non-linear Differential Equations*; Pitman, Boston M. A. 1985.
- [11] W. Simons, *Some disconjugacy criteria for selfadjoint linear differential equations*. *J. Math. Anal. Appl.* 34, 445-463 (1971).
- [12] P. J. Torres, *Existence of One-Signed Periodic Solutions of Some Second-Order Differential Equations via Krasnoselskii Fixed Point Theorem*, *J. Differential Equations* 190 (2003), 2, 643-662.
- [13] A. Zettl, *A constructive characterization of disconjugacy*. *Bull. Am. Math. Soc.* 81, 145-147 (1975).