

Global Embedding of Fibre Inflation in Perturbative LVS

Swagata Bera^{†*}, Dibya Chakraborty[‡], George K. Leontaris[◇], Pramod Shukla[†]

[†]*Department of Physical Sciences, Bose Institute,*

Unified Academic Campus, EN 80, Sector V, Bidhannagar, Kolkata 700091, India

^{*}*Department of Integrated Science Education and Research Center,*

Visva Bharati University, Santiniketan 731235, India

[‡]*Centre for Strings, Gravitation and Cosmology, Department of Physics,*

Indian Institute of Technology Madras, Chennai 600036, India

[◇]*Physics Department, University of Ioannina, University Campus, Ioannina 45110, Greece*

We present a global embedding of the Fibre inflation model in perturbative large volume scenario (LVS) in which the inflationary potential is induced via a combination of string-loop corrections and higher derivative F^4 -effects. The minimal Fibre inflation model developed in the standard LVS generically faces some strong bounds on the inflaton field range arising from the Kähler cone constraints, which subsequently creates a challenge for realising the cosmological observables while consistently respecting the underlying supergravity approximations. The main attractive feature of this proposal is to naturally alleviate this issue by embedding Fibre inflation in perturbative LVS.

INTRODUCTION AND MOTIVATION

In the context of type IIB orientifold compactifications, the Large Volume Scenario (LVS) [1] is one of the most popular schemes of moduli stabilisation. The minimal LVS fixes the overall volume (\mathcal{V}) of the underlying Calabi Yau threefold (CY_3) along with another modulus τ_s controlling the volume of an exceptional del-Pezzo divisor of the CY_3 , which is needed to induce the non-perturbative superpotential contributions. Therefore a generic LVS model with 3 or more Kähler moduli, arising from CY compactifications with $h^{1,1}(CY) \geq 3$, can generically have flat directions at leading order, which can be promising inflaton candidates with a potential generated at sub-leading order. Based on the geometric nature of the inflaton field and the source of the potential, there are several popular inflationary models in LVS, e.g. Blow-up inflation [2–4], Fibre inflation [5–8], poly-instanton inflation [9–12], and loop-blowup inflation [13].

Fibre inflation [5], being a large field model, is among the most attractive class of models realised in LVS. In explicit global embedding attempts for Fibre inflation, it has been observed that the inflaton range is highly constrained due to the Kähler cone conditions (KCC) [8], which subsequently obstruct the inflationary plateau forbidding the generation of sufficient number of e-foldings. A detailed analysis of all the CY_3 with $h^{1,1} = 3$ shows that the presence of diagonal del-Pezzo divisor results in a bound on the LVS flat direction [14]. In fact the reduced moduli space after LVS minimisation turns out to be compact in such cases, and the flat direction τ_f , which is a candidate for inflaton, has the following bound [14]:

$$c_1 \langle \tau_s \rangle < \tau_f < c_2 f(\langle \mathcal{V} \rangle, \langle \tau_s \rangle), \quad (1)$$

where $f(\langle \mathcal{V} \rangle, \langle \tau_s \rangle) \simeq \langle \mathcal{V} \rangle^{2/3}$ or $\langle \mathcal{V} \rangle / \langle \tau_s \rangle$, and $c_i \simeq \mathcal{O}(1)$ and positive. One can quantify the *field-range problem* as the fact that typically one needs $\Delta\varphi \simeq \mathcal{O}(6-8)M_p$ along with $\langle \mathcal{V} \rangle \simeq \mathcal{O}(10^3-10^4)$ where $\Delta\varphi$ is the shift in the canonical field φ where $\tau_f = e^{2\varphi/\sqrt{3}}$. One may argue

to increase the field range by considering larger value for $\langle \mathcal{V} \rangle$ in order to generate the required e-foldings, however it turns out that subsequently the scalar perturbation amplitude gets accordingly diluted. Moreover, one needs to respect the following mass hierarchy to trust the supergravity approximations

$$m_{\text{inf}} < H < m_{3/2} < M_{\text{KK}} < M_s < M_p, \quad (2)$$

where m_{inf} denotes the inflaton mass, H is the Hubble parameter, $m_{3/2}$ is the gravitino mass, M_{KK} is the lightest KK-mode, and M_s is the string mass. Maintaining the mass-hierarchies on top of field range bound creates a challenge for constructing a trustworthy viable model.

In this letter we propose an alternative way of realising Fibre inflation using perturbative LVS framework [15–22] that shares most of the interesting properties of the standard LVS without the need of non-perturbative corrections. Given that such effects are not necessary in perturbative LVS, the need of exceptional 4-cycles in the CY_3 is absent and subsequently the obstruction from the KCC is also naturally avoided.

LVS SCHEMES OF MODULI STABILISATION

The F-term scalar potential governing the low energy dynamics of the four-dimensional $\mathcal{N} = 1$ effective supergravity arising from the type IIB superstring compactifications is encoded in $V = e^{\mathcal{K}}(\mathcal{K}^{A\bar{B}}(D_A W)(D_{\bar{B}} \bar{W}) - 3|W|^2)$, where the full Kähler potential (\mathcal{K}) and the holomorphic superpotential (W) depend on the various moduli, e.g. the complex-structure (CS) moduli U^i , the axio-dilaton S and the Kähler moduli T_α . For a decoupled structure of \mathcal{K} such that $\mathcal{K} \equiv K_{\text{cs}}(U^i) + K(S, T_\alpha)$, the inverse metric is block-diagonal and $V \equiv V_{\text{cs}} + V_{\text{k}}$ such that $V_{\text{cs}} = e^K K_{\text{cs}}^{i\bar{j}}(D_i W)(D_{\bar{j}} \bar{W})$ and $V_{\text{k}} = e^K (K^{A\bar{B}}(D_A W)(D_{\bar{B}} \bar{W}) - 3|W|^2)$ where $A, B \in \{S, T_\alpha\}$. Moduli stabilisation follows a two-step methodology.

First, one fixes the CS moduli U^i and the axio-dilaton S by the leading order flux superpotential W_{flux} induced by the 3-form fluxes (F_3, H_3) [23] via the supersymmetric flatness conditions $D_{U^i} W_{\text{flux}} = 0 = D_S W_{\text{flux}}$.

After achieving the supersymmetric stabilisation of the axio-dilaton and the CS moduli, one has $\langle W_{\text{flux}} \rangle = W_0$. However, the no-scale structure protects the Kähler moduli T_α which remain flat, and as a second step, they can be stabilised via including other sub-leading contributions, e.g. those arising from the whole series of α' and string-loop (g_s) corrections; see [24] and reference therein. At two-derivative order, these effects can be captured through corrections in \mathcal{K} and W ,

$$\begin{aligned} \mathcal{K} &= K_{\text{cs}} - \ln[-i(S - \bar{S})] - 2\ln\mathcal{V}, \\ W &= W_0 + W_{\text{np}}(S, T_\alpha), \end{aligned} \quad (3)$$

where K_{cs} depends on CS moduli, while \mathcal{V} generically encodes the volume/dilaton pieces such that $\mathcal{V} = \mathcal{V}$ at tree-level. Here $\mathcal{V} = \frac{1}{3!} k_{\alpha\beta\gamma} t^\alpha t^\beta t^\gamma$ where $k_{\alpha\beta\gamma}$ is the triple intersection number of the CY_3 and the 2-cycle volumes t^α 's are related to the divisor volume τ_α via $\tau_\alpha = \partial_{t^\alpha} \mathcal{V}$. **Standard LVS:** The LVS scheme of moduli stabilisation considers a combination of perturbative (α')³ corrections to \mathcal{K} [25] and a non-perturbative contribution to W [26]. The minimal LVS construction includes two Kähler moduli with a so-called Swiss-cheese like volume form: $\mathcal{V} = \lambda_b \tau_b^{3/2} - \lambda_s \tau_s^{3/2}$ where $\lambda_b = \sqrt{2}/(3\sqrt{k_{bbb}})$ and $\lambda_s = \sqrt{2}/(3\sqrt{k_{sss}})$. The Kähler potential \mathcal{K} is given by Eq. (3) with $\mathcal{V} = \mathcal{V} + \xi/(2g_s^{3/2})$ such that $\xi = -\chi(\text{CY}_3)\zeta(3)/(16\pi^3)$, and $\chi(\text{CY}_3)$ being the CY Euler characteristic and $\zeta(3) \simeq 1.202$. We note in passing that moduli stabilisation in LVS requires $\xi > 0$ or equivalently $\chi < 0$, i.e. a constraint $h^{1,1} < h^{2,1}$ on the Hodge numbers counting the Kähler- and CS-moduli.

Furthermore, non-perturbative superpotential contributions [26] can be generated by using rigid divisors, such as shrinkable del-Pezzo (dP) 4-cycles, or by rigidifying non-rigid divisors using magnetic fluxes [27–29]. In fact, in the presence of a ‘diagonal’ del-Pezzo (ddP) divisor, the so-called ‘small’ 4-cycle of the CY_3 gives: $W = W_0 + A_s e^{-i a_s T_s}$, where $T_s = c_s - i\tau_s$ for c_s being the C_4 axion, and the holomorphic pre-factor A_s is considered as a parameter after fixing CS-moduli. Subsequently, after minimising the axion c_s , the leading order pieces in the large volume expansion are collected as [1]:

$$V_{\text{LVS}} \simeq \frac{\alpha_1}{\mathcal{V}^3} - \frac{\alpha_2 \tau_s}{\mathcal{V}^2} e^{-a_s \tau_s} + \frac{\alpha_3 \sqrt{\tau_s}}{\mathcal{V}} e^{-2a_s \tau_s}, \quad (4)$$

where α_i 's are positive constants defined as $\alpha_1 = \frac{3}{4} \kappa \hat{\xi} |W_0|^2$, $\alpha_2 = 4\kappa a_s |W_0| |A_s|$, $\alpha_3 = 4\kappa a_s^2 |A_s|^2 \sqrt{2k_{sss}}$, while $\kappa = g_s e^{K_{\text{cs}}}/(8\pi)$. This potential (4) results in exponentially large $\langle \mathcal{V} \rangle$ determined by

$$\langle \mathcal{V} \rangle \simeq \frac{\alpha_2 \sqrt{\langle \tau_s \rangle}}{2\alpha_3} e^{a_s \langle \tau_s \rangle}, \quad \langle \tau_s \rangle \simeq \hat{\xi}^{2/3} \left(\frac{9k_{sss}}{8} \right)^{1/3}. \quad (5)$$

Here $k_{sss} = 9 - n$ is the degree of the dP_n divisor such that $\text{dP}_0 = \mathbb{P}^2$. Thus the LVS models based on Swiss-cheese CY with $h^{1,1}(\text{CY}) \geq 3$ have leading order flat directions which can act as promising inflaton candidate with a sub-leading potential. However, the constraints from Kähler cone conditions get crucially important, especially in large field models like Fibre inflation.

Perturbative LVS: With the inclusion of the so-called *log-loop* effects [15–20] along with the BBHL corrections [25], the Kähler potential \mathcal{K} is given by Eq. (3) with $\mathcal{V} = \mathcal{V} + \frac{\hat{\xi}}{2} + \hat{\eta}(\ln \mathcal{V} - 1)$, where $\hat{\xi} = \xi/g_s^{3/2}$, and $\hat{\eta} = -g_s^2 \hat{\xi} \zeta[2]/\zeta[3]$. Now, considering the tree-level superpotential W_0 for CS+dilaton moduli stabilisation, one gets the following scalar potential,

$$V_{\text{pertLVS}} = \frac{3\kappa \hat{\xi}}{4\mathcal{V}^3} |W_0|^2 + \frac{3\kappa \hat{\eta}(\ln \mathcal{V} - 2)}{2\mathcal{V}^3} |W_0|^2. \quad (6)$$

This subsequently results in an exponentially large VEV for the overall volume modulus determined as,

$$\langle \mathcal{V} \rangle \simeq e^{a/g_s^{2+b}}, \quad a = \frac{\zeta[3]}{2\zeta[2]} \simeq 0.365381, \quad b = \frac{7}{3}, \quad (7)$$

justifying the name– ‘perturbative LVS’, which similar to the standard LVS, corresponds to an AdS minimum with large $\langle \mathcal{V} \rangle$. For $g_s = 0.2$, Eq. (7) results in $\langle \mathcal{V} \rangle = 95593.3$ while $g_s = 0.1$ leads to $\langle \mathcal{V} \rangle = 7.61463 \cdot 10^{16}$.

FIBRE INFLATION AND KÄHLER CONE BOUNDS ON THE INFLATON

One of the simplest extensions of the minimal two-field standard LVS model is to consider the models based on K3-fibred CY_3 with $h^{1,1} = 3$ and having a diagonal dP divisor [7, 14, 30]. In a suitable divisor basis $\{D_b, D_f, D_s\}$ having the ‘big’ divisor (D_b), the ‘small’ divisor (D_s) and the K3 divisor (D_f), the typical volume-form is given as:

$$\mathcal{V} = \frac{k_{bbf}}{2} t^f (t^b)^2 + \frac{k_{sss}}{6} (t^s)^3 = \lambda_f \tau_b \sqrt{\tau_f} - \lambda_s \tau_s^{3/2}, \quad (8)$$

where $\lambda_f = 1/(2\sqrt{k_{bbf}})$ and $\lambda_s = \sqrt{2}/(3\sqrt{k_{sss}})$. It turns out that \mathcal{V} and τ_s are fixed by the minimal standard LVS scheme while τ_f remains flat and serves as the inflaton field in the presence of string-loop corrections.

With the appropriate string-loop effects to the Kähler potential, one can generate sub-leading corrections to LVS potential which depend on the third modulus τ_f and helps in driving inflation. The two typical corrections known as ‘KK’ and ‘winding’ types are [31–36]

$$\begin{aligned} V_{g_s}^{\text{KK}} &= \kappa g_s^2 \frac{|W_0|^2}{16\mathcal{V}^4} \sum_{\alpha, \beta} C_\alpha^{\text{KK}} C_\beta^{\text{KK}} (2t^\alpha t^\beta - 4\mathcal{V} k^{\alpha\beta}), \\ V_{g_s}^{\text{W}} &= -\frac{\kappa |W_0|^2}{\mathcal{V}^3} \sum_{\alpha=1}^3 \frac{C_\alpha^{\text{W}}}{t^\alpha}, \end{aligned} \quad (9)$$

where C_α^{KK} and C_α^W are some model dependent coefficients which can generically depend on the CS moduli. For the minimal Fibre inflation model, one ends up with the following effective potential for K3-fibre modulus τ_f ,

$$V(\tau_f) = V_{\text{up}} + \frac{|W_0|^2}{\mathcal{V}^2} \left(\frac{B_1}{\tau_f^2} - \frac{B_2}{\mathcal{V}\sqrt{\tau_f}} + \frac{B_3\tau_f}{\mathcal{V}^2} \right), \quad (10)$$

where V_{up} denotes some appropriate uplifting needed to obtain the dS minimum, and B_i 's are some CS moduli dependent parameters to be considered as constants for inflationary dynamics occurring at sub-leading order.

Constraints from the Kähler cone : In the minimal Fibre inflation model based on CY_3 with $h^{1,1} = 3$, two moduli, namely \mathcal{V} and τ_s , are fixed by the standard LVS while the third one corresponds to the inflaton field τ_f . The crucial thing to note here is the fact that although τ_f remains flat after fixing the overall volume \mathcal{V} , there exists a bound on the field range it can traverse during inflationary dynamics. This arises because of the fact that the Kähler cone condition (KCC): $\int_{C_i} J > 0$ holds for all the curves C_i in the Mori cone of the CY_3 which typically translates into the following set of constraints:

$$\text{KCC : } p_{\alpha\beta} t^\beta > 0, \quad \text{for some } p_{\alpha\beta} \in \mathbb{Z}, \quad (11)$$

where summation for β runs in $h^{1,1}$ while α corresponds to the number of KCC which is $h^{1,1}$ for simplicial cases but can be more for non-simplicial cases. Depending on the values of $p_{\alpha\beta}$ in a given concrete model, there will be constraints on t^α , and so on the 4-cycle volumes. For example, taking the volume-form (8) and using $\tau_b = k_{bbf} t^b t^f$, $\tau_f = \frac{k_{bbf}}{2} (t^b)^2$, $\tau_s = \frac{k_{sss}}{2} (t^s)^2$, a generic KCC of the form $p_s t^s + p_b t^b + p_f t^f > 0$ translates into,

$$p_f \left(\mathcal{V} + \lambda_s \tau_s^{3/2} \right) + 2\sqrt{2} \lambda_b p_b \tau_f^{3/2} > 3\lambda_s p_s \sqrt{\tau_s} \tau_f. \quad (12)$$

Considering the generic possibilities that p_b, p_f and p_s can have either zero or non-zero values, results in 7 possibilities for the triplet $p_T^i = \{p_b, p_f, p_s\}$ where at least one of the three takes a non-zero value. It turns out that $p_T^1 = \{p_b, 0, 0\}$, $p_T^2 = \{0, p_f, 0\}$ and $p_T^3 = \{0, 0, -1\}$ give $\{\tau_b > 0, \tau_f > 0, \tau_s > 0, \mathcal{V} > 0\}$. In the presence of exceptional divisor, there always exists a 'diagonal' basis in which one KCC becomes $t^s < 0$ or equivalently $\tau_s > 0$ corresponding to $p_T^3 = \{0, 0, -1\}$. In fact, the Swiss-cheese form appears due to this generic condition which picks up the negative sign of $t^s = \pm\sqrt{2k_{sss}\tau_s}$. Further, $p_T^4 = \{p_b, 0, p_s\}$ gives a lower bound on τ_f such that $(p_s^2 k_{sss}) / (p_b^2 k_{bbf}) \tau_s < \tau_f$. On the contrary to these, $p_T^5 = \{p_b, p_f, 0\}$ results in $\tau_f \lesssim \mathcal{V}^{2/3}$ and $p_T^6 = \{0, p_f, p_s\}$ implies that $\tau_f \lesssim \mathcal{V} / \sqrt{\tau_s}$. These cases naturally arise in explicit models [8, 14]. Thus one has an upper bound for τ_f modulus in term of $\langle \mathcal{V} \rangle$ despite the LVS potential (4) being independent of τ_f . For $h^{1,1}(\text{CY}) = 3$, KCC consists of 3 or more constraints of type (12), so the only

possibility to avoid the upper bound on τ_f is to have a CY with KCC having 3 triplets from p_T^i for $1 \leq i \leq 4$. However, a detailed analysis of the CY geometries with $h^{1,1} = 3$ arising from the Kreuzer-Skarke database [37] shows that it does not happen [14] and one always have at least one of the p_T^5, p_T^6 , and p_T^7 which give the bound.

For numerical demonstration of the problem, one can take the model proposed in [7] in which $k_{bbf} = 18, k_{sss} = 1$ corresponding to a dP_8 divisor, and we find that the three conditions from KCC in the given divisor basis are: $0 < 2\langle \tau_s \rangle < \tau_f < (3\langle \mathcal{V} \rangle + \sqrt{2}\langle \tau_s \rangle^{3/2})^{2/3}$. Using the values $\langle \mathcal{V} \rangle = 10^3, \langle \tau_s \rangle = 3, \langle \tau_f \rangle = 60$ as mentioned for the benchmark model in [7], we find that KCC conditions result in $6 < \tau_f < 208.4$. Using canonical field φ via $\tau_f = e^{2\varphi/\sqrt{3}}$ leads to $\langle \varphi \rangle = 3.55$, and $\varphi < 4.63$ corresponding to the bound $\tau_f < 208.4$, and hence $\Delta\varphi \simeq 1$ while $\Delta\varphi \simeq 7 - 10M_p$ is typically needed for a successful Fibre inflation [5]. Regarding the challenge one faces in global model building, it is worth to mention that this setup was the one among just a few models found suitable for Fibre inflation after scanning a total of 305 CY geometries with $h^{1,1} = 3$, and even this faces a problem.

In fact such an upper bound exists even beyond the minimal $h^{1,1} = 3$ models, e.g. for a K3-fibred CY with $h^{1,1} = 4$ with a dP and a toroidal-like volume-form [8, 14]

$$\mathcal{V} = k_{123} t^1 t^2 t^3 + \frac{k_{sss}}{6} (t^s)^3 = \frac{\sqrt{\tau_1 \tau_2 \tau_3}}{\sqrt{k_{123}}} - \lambda_s \tau_s^{3/2}, \quad (13)$$

the generic KCC condition $p_{\alpha\beta} t^\beta > 0$ takes the form

$$\frac{p_1 \tau_2 \tau_3}{k_{123} \mathcal{Z}} + \mathcal{Z} \left(\frac{p_2}{\tau_2} + \frac{p_3}{\tau_3} \right) > 3\lambda_s p_s \sqrt{\tau_s}, \quad (14)$$

where $\mathcal{Z} = \mathcal{V} + 3\lambda_s \tau_s^{3/2}$. An explicit example of this type has the following four Kähler cone conditions [8]:

$$-t^s > 0, \quad t^1 + t^s > 0, \quad t^2 + t^s > 0, \quad t^3 + t^s > 0, \quad (15)$$

which correspond to four set of values for $p_T^i = \{p_1, p_2, p_3, p_s\}$ given as $\{0, 0, 0, -1\}$, $\{1, 0, 0, 1\}$, $\{0, 1, 0, 1\}$, and $\{0, 0, 1, 1\}$. These four sets correspond to $\tau_s > 0, \tau_2 \tau_3 \gtrsim \mathcal{V} \sqrt{\tau_s}, \tau_2 \lesssim \mathcal{V} / \sqrt{\tau_s}$, and $\tau_3 \lesssim \mathcal{V} / \sqrt{\tau_s}$ respectively. From (15) it is evident that one of the reasons for this bound is the presence of the exceptional rigid divisor which is needed to induce the non-perturbative superpotential contribution for the standard LVS.

FIBRE INFLATION IN PERTURBATIVE LVS

The *perturbative LVS* provides the nice features of standard LVS without having any non-perturbative effects, and hence the need of an exceptional divisor can be simply bypassed, and therefore the Kähler cone conditions would not put a strong bound on the inflaton field range. We will demonstrate this feature by considering a concrete K3-fibred CY orientifold with toroidal-like volume.

Global model: We consider a CY₃ with $h^{1,1} = 3$ corresponding to the polytope Id: 249 in the CY database of [38] which has been studied earlier in [21, 22, 39]. It is described by the following toric data:

Hyp	x_1	x_2	x_3	x_4	x_5	x_6	x_7
4	0	0	1	1	0	0	2
4	0	1	0	0	1	0	2
4	1	0	0	0	0	1	2
	K3	K3	K3	K3	K3	K3	SD

The Hodge numbers are $(h^{2,1}, h^{1,1}) = (115, 3)$, the Euler number is $\chi = -224$ while the Stanley-Reisner ideal is $\text{SR} = \{x_1x_6, x_2x_5, x_3x_4x_7\}$. The analysis of the divisor topologies using *cohomCalg* [40, 41] shows that the first six toric divisors are K3 surfaces while the seventh one is described by Hodge numbers $\{h^{0,0} = 1, h^{1,0} = 0, h^{2,0} = 27, h^{1,1} = 184\}$. Considering the divisor basis $\{D_1, D_2, D_3\}$ and the Kähler form $J = t^1 D_1 + t^2 D_2 + t^3 D_3$, the second Chern class $c_2(\text{CY})$ is given as $c_2(\text{CY}) = 5D_3^2 + 12D_1D_2 + 12D_2D_3 + 12D_1D_3$, while $k_{123} = 2$ is the only non-zero intersection leading to $\mathcal{V} = 2t^1 t^2 t^3 = \frac{1}{\sqrt{2}} \sqrt{\tau_1 \tau_2 \tau_3}$ where $\tau_1 = 2t^2 t^3, \tau_2 = 2t^1 t^3, \tau_3 = 2t^1 t^2$ and $\mathcal{V} = t^1 \tau_1 = t^2 \tau_2 = t^3 \tau_3$ as in toroidal case. The Kähler cone conditions are:

$$\text{KCC: } t^1 > 0, \quad t^2 > 0, \quad t^3 > 0. \quad (16)$$

Note that unlike KCC in Eq. (15) corresponding to a CY₃ with a dP divisor [14], the KCC in Eq. (16) do not put any other bound except $\tau_\alpha > 0$ and $\mathcal{V} > 0$. Therefore, the τ_3 modulus is not heavily constrained if two of them are already fixed by some moduli stabilisation scheme.

Brane setting: For a given holomorphic involution, one needs to introduce D3/D7-branes and fluxes in order to cancel all the D3/D7 tadpoles. In fact, one can nullify the D7-tadpoles via introducing stacks of N_a D7-branes wrapped around suitable divisors (say D_a) and their images (D'_a). However, the presence of D7/O7 also contributes to the D3-tadpoles, which, in addition, receive contributions from H_3/F_3 fluxes, D3-branes and O3-planes. For nullifying the D3-tadpoles one needs [42]:

$$\text{D7: } \sum_a N_a ([D_a] + [D'_a]) = 8 [\text{O7}], \quad (17)$$

$$\text{D3: } N_3 = \frac{N_{\text{O3}}}{4} + \frac{\chi(\text{O7})}{12} + \sum_a \frac{N_a (\chi(D_a) + \chi(D'_a))}{48}.$$

where $N_3 \equiv N_{\text{D3}} + \frac{N_{\text{flux}}}{2} + N_{\text{gauge}}$ such that N_{D3} is the net number of D3-brane, $N_{\text{flux}} = (2\pi)^{-4} (\alpha')^{-2} \int_X H_3 \wedge F_3$ is the contribution from background fluxes and $N_{\text{gauge}} = -\sum_a (8\pi)^{-2} \int_{D_a} \text{tr } \mathcal{F}_a^2$ is due to D7 worldvolume fluxes. Considering the involution $x_7 \rightarrow -x_7$ results in fixed point set $\{\text{O7} = D_7\}$ without any O3-planes, and subsequently one can consider 3 stacks of D7-branes wrapping each of the three divisors $\{D_1, D_2, D_3\}$, and the D3/D7-brane tadpoles are nullified via $8 [\text{O7}] = 8 ([D_1] + [D'_1]) +$

$8 ([D_2] + [D'_2]) + 8 ([D_3] + [D'_3])$ resulting in $N_3 = 44$. Now, in order to obtain a chiral visible sector on the D7-brane stacks wrapping D_1, D_2 and D_3 , we need to consistently turn on worldvolume gauge fluxes of the form: $\mathcal{F}_i = \sum_{j=1}^{h^{1,1}} f_{ij} \hat{D}_j + \frac{1}{2} \hat{D}_i - \iota_{D_i}^* B$ with $f_{ij} \in \mathbb{Z}$. Here, the half-integer contribution is due to Freed-Witten anomaly cancellation [43, 44]. However, given that the three stacks of D7-branes are wrapping a spin divisor K3 with $c_1(\text{K3}) = 0$, and given that the intersections on this CY are even, the pullback of the B -field on any divisor D_α does not generate a half-integer flux, and therefore one can appropriately adjust fluxes such that $\mathcal{F}_\alpha \in \mathbb{Z}$ for all $\alpha = 1, 2, 3$. We shall therefore consider a non-vanishing gauge flux \mathcal{F}_3 on the worldvolume of D_3 while considering $\mathcal{F}_1 = 0 = \mathcal{F}_2$. Subsequently, similar to [8], the vanishing of FI parameter leads to

$$\tau_1 = q \tau_2, \quad \text{where } q = -q_{31}/q_{32}. \quad (18)$$

Sub-leading corrections to scalar potential : Given that there are no rigid divisors present, a priori this setup will not receive non-perturbative superpotential contributions from instanton or gaugino condensation. The divisor intersection analysis shows that all the three D7-brane stacks intersect at \mathbb{T}^2 while each of those intersect the O7-plane on a curve \mathcal{H}_9 defined by $h^{0,0} = 1$ and $h^{1,0} = 9$. Further, there are no non-intersecting D7/O7 stacks and no O3-planes, and therefore this model does not induce the KK-type string-loop corrections to the Kähler potential [31–36]. However, as the O7/D7 stacks intersect on non-shrinkable 2-cycles with size t_\cap^i , one will have string-loop effects of the winding-type [31–36]: $V_{g_s}^{\text{W}} = -\frac{\kappa |W_0|^2}{\mathcal{V}^3} \sum_i \frac{C_{wi}}{t_\cap^i}$ where C_{wi} are CS-moduli dependent coefficients. This CY has several properties similar to a toroidal case, however basis divisors being K3 implies that their second Chern numbers are non-zero, $\Pi(D_\alpha) = 24$, unlike \mathbb{T}^4 , leading to the higher derivative F^4 corrections to the scalar potential $V_{\text{F}^4} \propto \Pi_\alpha t^\alpha$ [45]. Summarising all the contributions, we have [22]:

$$\begin{aligned} V = & V_{\text{up}} + \frac{\mathcal{C}_1}{\mathcal{V}^3} \left(\hat{\xi} - 4\hat{\eta} + 2\hat{\eta} \ln \mathcal{V} \right) \\ & + \frac{\mathcal{C}_2}{\mathcal{V}^4} \left(\mathcal{C}_{w_1} \tau_1 + \mathcal{C}_{w_2} \tau_2 + \mathcal{C}_{w_3} \tau_3 + \frac{\mathcal{C}_{w_4} \tau_1 \tau_2}{2(\tau_1 + \tau_2)} + \frac{\mathcal{C}_{w_5} \tau_2 \tau_3}{2(\tau_2 + \tau_3)} \right. \\ & \left. + \frac{\mathcal{C}_{w_6} \tau_3 \tau_1}{2(\tau_3 + \tau_1)} \right) + \frac{\mathcal{C}_3}{\mathcal{V}^3} \left(\frac{1}{\tau_1} + \frac{1}{\tau_2} + \frac{1}{\tau_3} \right) + \dots, \end{aligned} \quad (19)$$

where $\mathcal{C}_1 = \frac{3}{4} \kappa |W_0|^2 = \frac{3}{4} \mathcal{C}_2$, $\mathcal{C}_3 = 24 |\lambda| \kappa^2 |W_0|^4 / g_s^{3/2}$ such that $|\lambda| = \mathcal{O}(10^{-4})$ [46, 47]. The first two pieces of (19) fixes \mathcal{V} in perturbative LVS while one of the other two moduli are fixed via (18), leaving an effective single field potential for τ_3 modulus which drives inflation.

Inflationary dynamics: Using the slow-roll parameters $\epsilon_V \equiv V'^2/(2V^2)$ and $\eta_V \equiv V''/V$, the observables such as the scalar perturbation amplitude P_s , the spectral index

n_s , and the tensor-to-scalar ratio r are defined as

$$P_s \simeq \frac{V_{\text{inf}}^*}{24\pi^2 \epsilon_H^*}, \quad n_s \simeq 1 + 2\eta_V^* - 6\epsilon_V^*, \quad r \simeq 16\epsilon_V^*, \quad (20)$$

where the observables are evaluated at the horizon exit $\phi = \phi^*$. The experimental bounds on these observables are: $P_s = 2.105 \pm 0.030 \times 10^{-9}$, $n_s = 0.9649 \pm 0.0042$, and $r < 0.036$ at a pivot scale 0.05 Mpc^{-1} [48]. The number of e-foldings N_e is given as a sum three contributions [8, 49]:

$$N_e \simeq \int_{\phi_{\text{end}}}^{\phi^*} \frac{V}{V'} d\phi \simeq 57 + \frac{1}{4} \ln(r_* V_*) - \frac{1}{3} \ln \left(\frac{10V_{\text{end}}}{m_{\text{inf}}^{3/2}} \right),$$

where m_{inf} is the inflaton mass and typically $N_e \simeq 51$ for Fibre inflation [8, 50, 51].

Using $\tau_1 = \tau_2$ for $q = 1$ in (18) one is left with two Kähler moduli, and the overall volume \mathcal{V} is fixed via perturbative LVS at leading order. Subsequently, using (19) and considering $\tau_3 \equiv \tau_f = e^{2\varphi/\sqrt{3}}$ for the canonical field φ and taking the shift $\phi = \langle \varphi \rangle + \phi$, one gets the following effective single field scalar potential for the shift ϕ ,

$$V = \mathcal{C}_0 \left(\mathcal{C}_{\text{up}} + \mathcal{R}_0 e^{-\gamma\phi} - e^{-\frac{\gamma}{2}\phi} + \mathcal{R}_1 e^{\frac{\gamma}{2}\phi} + \mathcal{R}_2 e^{\gamma\phi} \right) \quad (21)$$

where $\gamma = 2/\sqrt{3}$ and $\mathcal{C}_{\text{up}} = 1 - \mathcal{R}_0 - \mathcal{R}_1 - \mathcal{R}_2$ is the uplifting needed for realising dS minimum with small cosmological constant. The anti- $D3$ uplifting of [52–54] is not directly applicable to our case due to the absence of $O3$ -planes, however other scheme, e.g. T -brane uplifting could be used [21, 52, 55]. The other parameters are:

$$\mathcal{C}_0 = \frac{\sqrt{2}\mathcal{C}_2\tilde{\mathcal{C}}_w}{\langle \mathcal{V} \rangle^3 e^{\frac{\gamma}{2}\langle \varphi \rangle}}, \quad \mathcal{R}_0 = \frac{\mathcal{C}_3 e^{-\frac{\gamma}{2}\langle \varphi \rangle}}{\sqrt{2}\mathcal{C}_2\tilde{\mathcal{C}}_w}, \quad \mathcal{R}_1 = \frac{\sqrt{2}e^{\sqrt{3}\langle \varphi \rangle}}{\langle \mathcal{V} \rangle},$$

$$\frac{\mathcal{R}_2}{\mathcal{R}_0} = \frac{\mathcal{C}_2\mathcal{C}_{w3}e^{2\gamma\langle \varphi \rangle}}{\mathcal{C}_3\langle \mathcal{V} \rangle} \left[1 + \hat{\mathcal{C}}_w \left(1 + \frac{e^{\sqrt{3}(\phi+\langle \varphi \rangle)}}{\langle \mathcal{V} \rangle \sqrt{2}} \right)^{-1} \right], \quad (22)$$

where $\tilde{\mathcal{C}}_w = -1/4(4\mathcal{C}_{w1} + 4\mathcal{C}_{w2} + \mathcal{C}_{w4})$ and $\hat{\mathcal{C}}_w = (\mathcal{C}_{w5} + \mathcal{C}_{w6})/(2\mathcal{C}_{w3})$. The first three terms of (21) determine the minimum while the other two terms create the steepening. However $\mathcal{R}_1 \ll 1$ and $\mathcal{R}_2 \ll 1$ as they are volume suppressed, which is to be exploited in finding a sufficiently long plateau. The slow-roll parameters do not depend on \mathcal{C}_0 , hence n_s , r and N_e can be determined purely by three parameters \mathcal{R}_0 , \mathcal{R}_1 and \mathcal{R}_2 . Also, \mathcal{R}_2 depends on ϕ , but this dependence is suppressed by an extra volume factor, and vanishes for $\mathcal{C}_{w5} \simeq -\mathcal{C}_{w6}$, i.e. $\hat{\mathcal{C}}_w \simeq 0$, and so \mathcal{R}_2 can be taken as a constant. Under this assumption, $\mathcal{R}_0 = (1 + \mathcal{R}_1 + 2\mathcal{R}_2)/2 \simeq 1/2$ for setting $\langle \phi \rangle = 0$, i.e. when φ modulus reaches its minimum. So one can study the inflationary dynamics from the potential (21) and determine the range of \mathcal{R}_1 and \mathcal{R}_2 which could produce sufficient e-folding N_e and suitable n_s , r values, and then \mathcal{C}_0 is fixed by matching the scalar perturbation amplitude P_s . We find that for $\mathcal{R}_2 = 0$, one can

have a viable inflationary model with $0 \leq \mathcal{R}_1 < 0.0001$, and larger \mathcal{R}_1 kills the plateau before sufficient e-folds are gained. This is shown in Fig. 1.




FIG. 1: $V(\phi)$ and N_e are plotted for ϕ on the left side while $(n_s - r)$ plots are presented on the right side.

For $\mathcal{R}_1 = 5 \cdot 10^{-5}$, we turn-on \mathcal{R}_2 and find that inflation is achieved for $\mathcal{R}_2 < 10^{-6}$. A benchmark model is:

$$\mathcal{C}_0 = 5.78 \cdot 10^{-10}, \quad \mathcal{R}_1 = 5 \cdot 10^{-5}, \quad \mathcal{R}_2 = 10^{-7}, \quad (23)$$

with $\langle \phi \rangle = 0$, $\phi_{\text{end}} = 1.03394$ for $\epsilon_V = 1$, and the observables: $N_e^* \simeq 51$, $P_s \simeq 2.1 \cdot 10^{-9}$, $n_s^* = 0.966$ and $r = 0.0085$ for $\phi^* = 6.3735$. Such a benchmark model can be consistently realised using the stringy-parameters,

$$|W_0| = 145, \quad \tilde{\mathcal{C}}_w = 5.3, \quad \mathcal{C}_{w3} = 10^{-2}, \quad \hat{\mathcal{C}}_w = 0, \quad (24)$$

$$|\lambda| = 10^{-4}, \quad g_s = 0.3, \quad \langle \mathcal{V} \rangle \simeq 15000, \quad \langle \varphi \rangle = 0.1,$$

which results in $\mathcal{C}_0 = 5.26 \cdot 10^{-10}$, $\mathcal{R}_0 = 0.4616$, $\mathcal{R}_1 = 0.0000518$ and $\mathcal{R}_2 = 1.06 \cdot 10^{-7}$. Here, the choice of $\langle \mathcal{V} \rangle = 15000$ and $g_s = 0.3$ is made as it corresponds to a de-Sitter minimum using T -brane uplifting in perturbative LVS [21], and $\lambda = -10^{-4}$ can be considered as a natural value given the presence of a suppression factor of $(2\pi)^4$ [46, 47]. We note that $\langle \tau_f \rangle = 1.12$ and $\tau_f^* \sim 1676$ and there are no Kähler cone conditions appearing as lower/upper bounds. The effective supergravity approximations are justified by ensuring by the mass hierarchies in Eq. (2) which are: $m_{\text{inf}} \sim 3.0 \cdot 10^{-6}$, $H^* \simeq \sqrt{V^*/3} \sim 9.5 \cdot 10^{-6}$, $m_{3/2} = g_s e^{K_{\text{cs}}/2} |W_0| / (8\pi\mathcal{V}) \sim 1.28 \cdot 10^{-4}$, $M_{\text{KK}} \simeq \sqrt{\pi}/(\sqrt{\mathcal{V}}\tau_b^{1/4}) \sim 1.3 \cdot 10^{-3}$ (for the bulk modulus $\tau_b \sim \mathcal{V}$ at the minimum as $\langle \tau_f \rangle \sim 1$), and $M_s = \sqrt{\pi}g_s^{1/4}/\sqrt{\mathcal{V}} \sim 1.07 \cdot 10^{-2}$ in units of M_p .

To conclude, we have presented a concrete global embedding for one of the most attractive inflationary models, namely Fibre inflation, realised in standard LVS, where it faces a generic challenge due to a geometric Kähler cone bound on the inflaton range. By demonstrating through concrete examples, we argue that one of the main reasons for this bound is the presence of an exceptional del-Pezzo divisor which is an integral part of the standard LVS. As perturbative LVS does not need any non-perturbative superpotential contribution, such a requirement can be bypassed, and a viable inflation can occur via a combination of string-loop corrections and the higher derivative F^4 -effects. This proposal consolidates the robustness of Fibre inflation beyond the standard LVS.

Acknowledgements: We thank Xin Gao for useful discussions. PS would like to thank the *Department of Science and Technology (DST), India* for the kind support.

-
- [1] V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, *Systematics of moduli stabilisation in Calabi-Yau flux compactifications*, *JHEP* **03** (2005) 007, [[hep-th/0502058](#)].
- [2] J. P. Conlon, F. Quevedo and K. Suruliz, *Large-volume flux compactifications: Moduli spectrum and D3/D7 soft supersymmetry breaking*, *JHEP* **08** (2005) 007, [[hep-th/0505076](#)].
- [3] J. J. Blanco-Pillado, D. Buck, E. J. Copeland, M. Gomez-Reino and N. J. Nunes, *Kahler Moduli Inflation Revisited*, *JHEP* **01** (2010) 081, [[0906.3711](#)].
- [4] M. Cicoli, I. Garcia-Etxebarria, C. Mayrhofer, F. Quevedo, P. Shukla and R. Valandro, *Global Orientifolded Quivers with Inflation*, *JHEP* **11** (2017) 134, [[1706.06128](#)].
- [5] M. Cicoli, C. P. Burgess and F. Quevedo, *Fibre Inflation: Observable Gravity Waves from IIB String Compactifications*, *JCAP* **0903** (2009) 013, [[0808.0691](#)].
- [6] M. Cicoli, D. Ciupke, S. de Alwis and F. Muia, *α' Inflation: moduli stabilisation and observable tensors from higher derivatives*, *JHEP* **09** (2016) 026, [[1607.01395](#)].
- [7] M. Cicoli, F. Muia and P. Shukla, *Global Embedding of Fibre Inflation Models*, *JHEP* **11** (2016) 182, [[1611.04612](#)].
- [8] M. Cicoli, D. Ciupke, V. A. Diaz, V. Guidetti, F. Muia and P. Shukla, *Chiral Global Embedding of Fibre Inflation Models*, *JHEP* **11** (2017) 207, [[1709.01518](#)].
- [9] M. Cicoli, F. G. Pedro and G. Tasinato, *Poly-instanton Inflation*, *JCAP* **12** (2011) 022, [[1110.6182](#)].
- [10] R. Blumenhagen, X. Gao, T. Rahn and P. Shukla, *Moduli Stabilization and Inflationary Cosmology with Poly-Instantons in Type IIB Orientifolds*, *JHEP* **11** (2012) 101, [[1208.1160](#)].
- [11] X. Gao and P. Shukla, *On Non-Gaussianities in Two-Field Poly-Instanton Inflation*, *JHEP* **03** (2013) 061, [[1301.6076](#)].
- [12] X. Gao, T. Li and P. Shukla, *Cosmological observables in multi-field inflation with a non-flat field space*, *JCAP* **10** (2014) 008, [[1403.0654](#)].
- [13] S. Bansal, L. Brunelli, M. Cicoli, A. Hebecker and R. Kuespert, *Loop Blow-up Inflation*, **2403.04831**.
- [14] M. Cicoli, D. Ciupke, C. Mayrhofer and P. Shukla, *A Geometrical Upper Bound on the Inflaton Range*, *JHEP* **05** (2018) 001, [[1801.05434](#)].
- [15] I. Antoniadis, Y. Chen and G. K. Leontaris, *Perturbative moduli stabilisation in type IIB/F-theory framework*, *Eur. Phys. J. C* **78** (2018) 766, [[1803.08941](#)].
- [16] I. Antoniadis, Y. Chen and G. K. Leontaris, *Inflation from the internal volume in type IIB/F-theory compactification*, *Int. J. Mod. Phys. A* **34** (2019) 1950042, [[1810.05060](#)].
- [17] I. Antoniadis, Y. Chen and G. K. Leontaris, *Moduli stabilisation and inflation in type IIB/F-theory*, *PoS CORFU2018* (2019) 068, [[1901.05075](#)].
- [18] I. Antoniadis, Y. Chen and G. K. Leontaris, *Logarithmic loop corrections, moduli stabilisation and de Sitter vacua in string theory*, **1909.10525**.
- [19] I. Antoniadis, Y. Chen and G. K. Leontaris, *String loop corrections and de Sitter vacua*, *PoS CORFU2019* (2020) 099.
- [20] I. Antoniadis, O. Lacombe and G. K. Leontaris, *Inflation near a metastable de Sitter vacuum from moduli stabilisation*, *Eur. Phys. J. C* **80** (2020) 1014, [[2007.10362](#)].
- [21] G. K. Leontaris and P. Shukla, *Stabilising all Kähler moduli in perturbative LVS*, *JHEP* **07** (2022) 047, [[2203.03362](#)].
- [22] S. Bera, D. Chakraborty, G. K. Leontaris and P. Shukla, *Inflating in perturbative LVS: Global Embedding and Robustness*, **2405.06738**.
- [23] S. Gukov, C. Vafa and E. Witten, *CFT's from Calabi-Yau four folds*, *Nucl. Phys. B* **584** (2000) 69–108, [[hep-th/9906070](#)].
- [24] M. Cicoli, F. Quevedo, R. Savelli, A. Schachner and R. Valandro, *Systematics of the α' expansion in F-theory*, *JHEP* **08** (2021) 099, [[2106.04592](#)].
- [25] K. Becker, M. Becker, M. Haack and J. Louis, *Supersymmetry breaking and alpha-prime corrections to flux induced potentials*, *JHEP* **06** (2002) 060, [[hep-th/0204254](#)].
- [26] E. Witten, *Nonperturbative superpotentials in string theory*, *Nucl. Phys. B* **474** (1996) 343–360, [[hep-th/9604030](#)].
- [27] M. Bianchi, A. Collinucci and L. Martucci, *Magnetized E3-brane instantons in F-theory*, *JHEP* **12** (2011) 045, [[1107.3732](#)].
- [28] M. Bianchi, A. Collinucci and L. Martucci, *Freezing E3-brane instantons with fluxes*, *Fortsch. Phys.* **60** (2012) 914–920, [[1202.5045](#)].
- [29] J. Louis, M. Rummel, R. Valandro and A. Westphal, *Building an explicit de Sitter*, *JHEP* **10** (2012) 163, [[1208.3208](#)].
- [30] P. Shukla, *Classifying divisor topologies for string phenomenology*, *JHEP* **12** (2022) 055, [[2205.05215](#)].
- [31] M. Berg, M. Haack and B. Kors, *On the moduli dependence of nonperturbative superpotentials in brane inflation*, in *Themes in Unification. Proceedings: 10th International Symposium on particles, strings and cosmology, (PASCOS 2004), Pran Nath Festschrift. Part II: Boston, USA, August 18-19, 2004*, 2004, [[hep-th/0409282](#)].
- [32] G. von Gersdorff and A. Hebecker, *Kahler corrections for the volume modulus of flux compactifications*, *Phys. Lett. B* **624** (2005) 270–274, [[hep-th/0507131](#)].
- [33] M. Berg, M. Haack and B. Kors, *String loop corrections to Kahler potentials in orientifolds*, *JHEP* **11** (2005) 030, [[hep-th/0508043](#)].
- [34] M. Berg, M. Haack and E. Pajer, *Jumping Through Loops: On Soft Terms from Large Volume Compactifications*, *JHEP* **09** (2007) 031, [[0704.0737](#)].
- [35] M. Cicoli, J. P. Conlon and F. Quevedo, *Systematics of String Loop Corrections in Type IIB Calabi-Yau Flux Compactifications*, *JHEP* **01** (2008) 052, [[0708.1873](#)].
- [36] X. Gao, A. Hebecker, S. Schreyer and G. Venken, *Loops, local corrections and warping in the LVS and other type IIB models*, *JHEP* **09** (2022) 091, [[2204.06009](#)].
- [37] M. Kreuzer and H. Skarke, *Complete classification of reflexive polyhedra in four-dimensions*, *Adv. Theor. Math. Phys.* **4** (2000) 1209–1230, [[hep-th/0002240](#)].

- [38] R. Altman, J. Gray, Y.-H. He, V. Jejjala and B. D. Nelson, *A Calabi-Yau Database: Threefolds Constructed from the Kreuzer-Skarke List*, *JHEP* **02** (2015) 158, [[1411.1418](#)].
- [39] X. Gao and P. Shukla, *On Classifying the Divisor Involutions in Calabi-Yau Threefolds*, *JHEP* **1311** (2013) 170, [[1307.1139](#)].
- [40] R. Blumenhagen, B. Jurke, T. Rahn and H. Roschy, *Cohomology of Line Bundles: A Computational Algorithm*, *J. Math. Phys.* **51** (2010) 103525, [[1003.5217](#)].
- [41] R. Blumenhagen, B. Jurke and T. Rahn, *Computational Tools for Cohomology of Toric Varieties*, *Adv. High Energy Phys.* **2011** (2011) 152749, [[1104.1187](#)].
- [42] R. Blumenhagen, V. Braun, T. W. Grimm and T. Weigand, *GUTs in Type IIB Orientifold Compactifications*, *Nucl.Phys.* **B815** (2009) 1–94, [[0811.2936](#)].
- [43] R. Minasian and G. W. Moore, *K theory and Ramond-Ramond charge*, *JHEP* **11** (1997) 002, [[hep-th/9710230](#)].
- [44] D. S. Freed and E. Witten, *Anomalies in string theory with D-branes*, *Asian J. Math.* **3** (1999) 819, [[hep-th/9907189](#)].
- [45] D. Ciupke, J. Louis and A. Westphal, *Higher-Derivative Supergravity and Moduli Stabilization*, *JHEP* **10** (2015) 094, [[1505.03092](#)].
- [46] T. W. Grimm, K. Mayer and M. Weissenbacher, *One-modulus Calabi-Yau fourfold reductions with higher-derivative terms*, *JHEP* **04** (2018) 021, [[1712.07074](#)].
- [47] M. Cicoli, M. Licheri, P. Piantadosi, F. Quevedo and P. Shukla, *Higher derivative corrections to string inflation*, *JHEP* **02** (2024) 115, [[2309.11697](#)].
- [48] PLANCK collaboration, Y. Akrami et al., *Planck 2018 results. X. Constraints on inflation*, *Astron. Astrophys.* **641** (2020) A10, [[1807.06211](#)].
- [49] A. R. Liddle and S. M. Leach, *How long before the end of inflation were observable perturbations produced?*, *Phys. Rev. D* **68** (2003) 103503, [[astro-ph/0305263](#)].
- [50] S. Bhattacharya, K. Dutta, M. R. Gangopadhyay, A. Maharana and K. Singh, *Fibre Inflation and Precision CMB Data*, *Phys. Rev. D* **102** (2020) 123531, [[2003.05969](#)].
- [51] M. Cicoli and E. Di Valentino, *Fitting string inflation to real cosmological data: The fiber inflation case*, *Phys. Rev. D* **102** (2020) 043521, [[2004.01210](#)].
- [52] M. Cicoli, F. Quevedo and R. Valandro, *De Sitter from T-branes*, *JHEP* **03** (2016) 141, [[1512.04558](#)].
- [53] C. Crinò, F. Quevedo and R. Valandro, *On de Sitter String Vacua from Anti-D3-Branes in the Large Volume Scenario*, *JHEP* **03** (2021) 258, [[2010.15903](#)].
- [54] S. AbdusSalam, C. Crinò and P. Shukla, *On K3-fibred LARGE Volume Scenario with de Sitter vacua from anti-D3-branes*, *JHEP* **03** (2023) 132, [[2206.12889](#)].
- [55] M. Cicoli, I. n. G. Etxebarria, F. Quevedo, A. Schachner, P. Shukla and R. Valandro, *The Standard Model quiver in de Sitter string compactifications*, *JHEP* **08** (2021) 109, [[2106.11964](#)].