Doubly minimized Petz Rényi mutual information: Properties and operational interpretation from direct exponent

Laura Burri

Institute for Theoretical Physics, ETH Zurich, Zurich, Switzerland

The doubly minimized Petz Rényi mutual information of order α is defined as the minimization of the Petz divergence of order α of a fixed bipartite quantum state relative to any product state. In this work, we establish several properties of this type of Rényi mutual information, including its additivity for $\alpha \in [1/2, 2]$. As an application, we show that the direct exponent of certain binary quantum state discrimination problems is determined by the doubly minimized Petz Rényi mutual information of order $\alpha \in (1/2, 1)$. This provides an operational interpretation of this type of Rényi mutual information, and generalizes a previous result for classical probability distributions to the quantum setting.

1. INTRODUCTION

The relationship between operational tasks and information measures is a central topic of study in information theory. This paper is concerned with a specific type of information measure, namely, Rényi generalizations of the mutual information. We will now elucidate their relation to direct exponents of certain binary discrimination problems in both classical and quantum information theory.

Classical mutual information. The classical mutual information can be expressed in terms of the classical relative entropy (or: Kullback-Leibler divergence) in several ways [1, Proposition 8]:

$$I(X:Y)_P = D(P_{XY} || P_X P_Y) = \inf_{R_Y} D(P_{XY} || P_X R_Y) = \inf_{Q_X, R_Y} D(P_{XY} || Q_X R_Y).$$
(1.1)

Here, X and Y are random variables over finite alphabets \mathcal{X} and \mathcal{Y} , P_{XY} is a joint probability mass function (PMF) with marginals P_X and P_Y , and the minimizations are over PMFs Q_X and R_Y . Based on the Rényi divergence of order α , these expressions induce the following three types of Rényi mutual information (RMI) for $\alpha \in [0, \infty)$.

$$I_{\alpha}^{\uparrow\uparrow}(X:Y)_P \coloneqq D_{\alpha}(P_{XY} \| P_X P_Y) \tag{1.2}$$

$$I_{\alpha}^{\uparrow\downarrow}(X:Y)_P \coloneqq \inf_{R_Y} D_{\alpha}(P_{XY} \| P_X R_Y)$$
(1.3)

$$I_{\alpha}^{\downarrow\downarrow}(X:Y)_P \coloneqq \inf_{Q_X, R_Y} D_{\alpha}(P_{XY} \| Q_X R_Y)$$
(1.4)

We call them the non-minimized RMI, the singly minimized RMI, and the doubly minimized RMI, respectively. The singly minimized RMI was originally introduced in [2] and its properties were subsequently studied in [3–8]. The doubly minimized RMI was introduced in [1, 9]. Each of the three aforementioned RMIs can be related to the direct exponent of a certain binary discrimination problem, as outlined in Table I. The results in Table I provide an operational interpretation of the family of the non-minimized RMIs and the singly minimized RMIs of order $\alpha \in (0, 1)$, and the doubly minimized RMIs of order $\alpha \in (\frac{1}{2}, 1)$.

Quantum mutual information. Analogous to (1.1), the quantum mutual information can be expressed in terms of the quantum relative entropy in several ways [15, 17, 18]:

$$I(A:B)_{\rho} = D(\rho_{AB} \| \rho_A \otimes \rho_B) = \inf_{\tau_B} D(\rho_{AB} \| \rho_A \otimes \tau_B) = \inf_{\sigma_A, \tau_B} D(\rho_{AB} \| \sigma_A \otimes \tau_B).$$
(1.5)

Reference	Null hypothesis, alternative hypothesis	Direct exponent
[10–14]	$ \begin{aligned} H_0^n &= \{P_{XY}^{\times n}\} \\ H_1^n &= \{P_X^{\times n} P_Y^{\times n}\} \end{aligned} $	For any $R \in (0, \infty)$ holds $\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_n(e^{-nR}) = \sup_{s \in (0,1)} \frac{1-s}{s} (I_s^{\uparrow\uparrow}(X:Y)_P - R).$
[9]		For any $R \in (I_0^{\uparrow\downarrow}(X:Y)_P,\infty)$ holds $\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_n(e^{-nR}) = \sup_{s \in (0,1)} \frac{1-s}{s} (I_s^{\uparrow\downarrow}(X:Y)_P - R).$
[9]		For any $R \in (R_{1/2}, \infty)$ holds $\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_n(e^{-nR}) = \sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(X : Y)_P - R).$

Table I. Overview of direct exponents of certain binary discrimination problems. Let P_{XY} be a PMF. Each row pertains to a sequence of binary discrimination problems with null hypothesis H_0^n and alternative hypothesis H_1^n for $n \in \mathbb{N}_{>0}$. In all three rows, the *n*th null hypothesis is given by $P_{XY}^{\times n}$. In the first row, the nth alternative hypothesis is given by the n-fold product of the marginal PMFs of X and Y induced by P_{XY} . In the second row, the *n*th alternative hypothesis is given by PMFs of the form $P_X^{\times n} R_{Y^n}$ for permutation invariant PMFs R_{Y^n} . Another option (which leads to the same direct exponent) is to define the *n*th alternative hypothesis in the second row as $P_X^{\times n} R_Y^{\times n}$ for PMFs R_Y . In the third row, the *n*th alternative hypothesis is given by $Q_{X^n}R_{Y^n}$ for permutation invariant PMFs Q_{X^n}, R_{Y^n} . Another option is to define the *n*th alternative hypothesis in the third row as $Q_X^{\times n} R_Y^{\times n}$ for PMFs Q_X, R_Y . The papers cited in the first column derive single-letter formulas (i.e., formulas in which P_{XY} occurs only once and not $n \to \infty$ many times) for the corresponding direct exponents, which are stated in the last column. The function $\hat{\alpha}_n(\mu)$ that appears in the last column is linked to the *n*th hypothesis testing problem and is defined as the minimum type-I error when the type-II error is upper bounded by $\mu \in [0, \infty)$. This definition of $\hat{\alpha}_n(\mu)$ will be repeated in greater detail in Section 2D. In that section, we will also explain that the range of R where the formula in the second row is applicable can be extended to $R \in (0,\infty)$ due to the validity of an analogous formula for quantum states [15]. The lower bound on R in the third row is defined as $R_{1/2} \coloneqq I_{1/2}^{\downarrow\downarrow}(X:Y)_P - \frac{1}{4} \frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(X:Y)_P|_{s=1/2}$ [9]. For a negative answer to the question of potential extensions of the result in the third row, see [16].

Here, ρ_{AB} is a bipartite quantum state on finite-dimensional Hilbert spaces A and B with marginal states ρ_A and ρ_B , and the minimizations are over quantum states σ_A and τ_B . From the perspective of quantum information theory, it is desirable to extend the RMIs in (1.2)–(1.4) from the classical to the quantum domain. However, there exist several quantum generalizations of the Rényi divergence. Commonly used are, for example, the Petz quantum Rényi divergence [19] and the sandwiched quantum Rényi divergence [20, 21]. Consequently, various generalizations of the classical RMIs to the quantum setting are conceivable and discerning which of these generalizations are operationally relevant is not straightforward. This paper examines generalizations based on the Petz divergence, as this choice of divergence turns out to be suitable for generalizing the results in Table I from the classical to the quantum setting. Using the Petz divergence, we consider the following three types of Petz Rényi mutual information (PRMI) for $\alpha \in [0, \infty)$.

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \coloneqq D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B) \tag{1.6}$$

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \coloneqq \inf_{\tau_{B}} D_{\alpha}(\rho_{AB} \| \rho_{A} \otimes \tau_{B})$$
(1.7)

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \coloneqq \inf_{\sigma_{A},\tau_{B}} D_{\alpha}(\rho_{AB} \| \sigma_{A} \otimes \tau_{B})$$
(1.8)

We call them the non-minimized PRMI, the singly minimized PRMI, and the doubly minimized PRMI, respectively. The non-minimized PRMI has previously been examined in the context of quantum field theory [22, 23], and has been applied to classical-quantum channel coding [24] and quantum soft covering [25]. The singly minimized PRMI has been studied with regard to its general

properties in [15, 17]. The doubly minimized PRMI has been mentioned in previous work on binary quantum state discrimination [26] and has been investigated for the case $\alpha = 0$ in [27]. However, none of these works has studied its general properties in detail.

Main results. In this paper, we initiate the information-theoretic study of the doubly minimized PRMI. Our findings on properties of the doubly minimized PRMI are presented in Theorems 6 and 7. These theorems constitute the first main result of this work. The second main result is Theorem 8, which shows that the doubly minimized PRMI of order $\alpha \in (\frac{1}{2}, 1)$ determines the direct exponent of certain binary quantum state discrimination problems. We will now provide a summary of these results.

In Theorem 6, we show that the minimization problem in (1.8), which defines the doubly minimized PRMI, is jointly convex in σ_A and τ_B . A crucial component of the proof of this theorem is an operator inequality (Lemma 5), which follows from the subadditivity of the geometric operator mean.

In Theorem 7 (a)–(v), we enumerate several properties pertaining to the doubly minimized PRMI. Of these, the following are particularly relevant:

- (d) additivity for $\alpha \in [\frac{1}{2}, 2]$,
- (j) uniqueness of the minimizer for $\alpha \in (\frac{1}{2}, 1]$,
- (k) a fixed-point property of minimizers for $\alpha \in (\frac{1}{2}, 2]$,
- (l) asymptotic optimality of the universal permutation invariant state for $\alpha \in [0, 2]$,
- (p) continuous differentiability in α of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ on $\alpha \in (\frac{1}{2},2)$, and
- (q) convexity in α of $(\alpha 1)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ on $\alpha \in [0,2]$.

Of these properties, we will prove (j) first. According to (j), the minimizer of the optimization problem in (1.8), which defines the doubly minimized PRMI, is unique. That is, for any $\alpha \in (\frac{1}{2}, 1]$, there exists a unique pair of quantum states (σ_A, τ_B) such that $D_\alpha(\rho_{AB} || \sigma_A \otimes \tau_B) = I_\alpha^{\downarrow\downarrow}(A : B)_\rho$. Previously, the uniqueness of the minimizer of the optimization problem that defines the doubly minimized RMI has been established in [1, Lemma 20] for any $\alpha \in (\frac{1}{2}, \infty)$. Similar to their proof for the classical setting, we show that the uniqueness of the minimizer follows from the joint convexity of the optimization problem (Theorem 6).

Then, we will prove (k). The item (k) characterizes minimizers σ_A for the optimization problem (1.8) in terms of a fixed-point property. The proof of (k) is based on an extension (Lemma 15) of a lemma from previous work [15, Lemma 22] that asserts a general equivalence of optimizers and fixed-points.

Additivity on product states (d) is then a direct consequence of (k). The additivity of the doubly minimized PRMI has been previously suggested in [26, Section 3.3], but no detailed proof was given.

The central assertion in (1) is that $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \lim_{n\to\infty} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} || \omega_{A^n}^n \otimes \omega_{B^n}^n)$, where $\omega_{A^n}^n$ and $\omega_{B^n}^n$ are certain universal permutation invariant states, which will be defined in Section 2 B. According to (1), the minimization over product states on AB, which is present in the definition of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ in (1.8), can be circumvented by comparing multiple copies of ρ_{AB} to the tensor product of two universal permutation invariant states. Thus, (1) offers a qualitatively different perspective on the doubly minimized PRMI, which can be useful in applications. The proof of (1) for $\alpha \in [\frac{1}{2}, 2]$ makes use of additivity (d), and proceeds in a similar manner to an analogous proof [15] for certain types of Rényi mutual information that are based on the sandwiched divergence. In the case where $\alpha \in [0, \frac{1}{2})$, a different proof method is employed that does not require additivity.

The convexity property (q) follows directly from (l). Subsequently, we prove (p) using (j) and (q). The properties (j), (p), and (q) are valuable technical tools for applications.

We then turn to the question of whether the doubly minimized PRMI has an operational interpretation. Previous research has demonstrated that, for the family of the non-minimized PRMIs

Reference	Null hypothesis, alternative hypothesis	Direct exponent
[14, 28, 29]	$ \begin{array}{l} H_0^n = \{\rho_{AB}^{\otimes n}\} \\ H_1^n = \{\rho_A^{\otimes n} \otimes \rho_B^{\otimes n}\} \end{array} $	For any $R \in (0, \infty)$ holds $\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_n(e^{-nR}) = \sup_{s \in (0,1)} \frac{1-s}{s} (I_s^{\uparrow\uparrow}(A:B)_{\rho} - R).$
[15]		For any $R \in (0, \infty)$ holds $\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_n(e^{-nR}) = \sup_{s \in (0,1)} \frac{1-s}{s} (I_s^{\uparrow\downarrow}(A:B)_{\rho} - R).$
Theorem 8	$ \begin{aligned} H_0^n &= \{\rho_{AB}^{\otimes n}\} \\ H_1^n &= \{\sigma_{A^n} \otimes \tau_{B^n}\}_{\sigma_{A^n}, \tau_{B^n}} \end{aligned} $	For any $R \in (R_{1/2}, \infty)$ holds $\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_n(e^{-nR}) = \sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_\rho - R).$

Table II. Overview of direct exponents of certain binary quantum state discrimination problems. Let ρ_{AB} be a quantum state. Each row pertains to a sequence of binary quantum state discrimination problems with null hypothesis H_0^n and alternative hypothesis H_1^n for $n \in \mathbb{N}_{>0}$. In all three rows, the *n*th null hypothesis is given by $\rho_{AB}^{\otimes n}$. In the first row, the *n*th alternative hypothesis is given by the *n*-fold tensor product of the marginal states of ρ_{AB} on A and B, respectively. In the second row, the *n*th alternative hypothesis is given by states of the form $\rho_A^{\otimes n} \otimes \tau_{B^n}$ for permutation invariant quantum states τ_{B^n} . Another option is to define the *n*th alternative hypothesis is given by $\sigma_{A^n} \otimes \tau_{B^n}^{\otimes n}$ for quantum states σ_{A^n} . In the third row, the *n*th alternative hypothesis is given by $\sigma_{A^n} \otimes \tau_{B^n}^{\otimes n}$ for permutation invariant quantum states τ_B . In the third row, the *n*th alternative hypothesis is given by $\sigma_{A^n} \otimes \tau_{B^n}$ for permutation invariant quantum states σ_{A^n} , τ_{B^n} . Another option is to define the *n*th alternative hypothesis in the third row as $\sigma_A^{\otimes n} \otimes \sigma_A^{\otimes n} \otimes \sigma_A^{\otimes$

and the singly minimized PRMIs of order $\alpha \in (0, 1)$, an operational interpretation can be obtained from the direct exponent of certain binary quantum state discrimination problems. These results are summarized in the first and second row of Table II. As will be discussed in Section 2 D, these results generalize the corresponding results for the classical setting in the first and second row of Table I. In Theorem 8, we show that this generalization from the classical to the quantum setting is also attainable for the doubly minimized (P)RMI. The central assertion of this theorem is outlined in the third row of Table II. Thus, Theorem 8 yields an operational interpretation of the family of the doubly minimized PRMIs of order $\alpha \in (\frac{1}{2}, 1)$. Theorem 8 generalizes a previous result [9, Section IV.A.3] for classical probability distributions outlined in the third row of Table I to the quantum setting. The proof of Theorem 8 proceeds in a similar manner to corresponding proofs for the singly minimized PRMI [15] and the doubly minimized RMI [9], and makes use of several properties of the doubly minimized PRMI, including Theorem 7 (j), (l), (p), and (q).

Related work. A summary of results on single-letter formulas for the direct exponent that are determined by (P)RMIs of order $\alpha \in (0,1)$ or $\alpha \in (\frac{1}{2},1)$ was given in Tables I and II (for the classical and the quantum setting, respectively). It should be noted that the classical discrimination problems in Table I are also known to admit similar single-letter formulas for the strong converse exponent, which are again determined by the non-minimized [13, 30, 31], the singly minimized [9], and the doubly minimized RMI [9], respectively. These results on strong converse exponents can be generalized [15, 32–34] to the quantum settings in Table II by lifting the RMIs from the classical to the quantum domain based on the sandwiched divergence (instead of the Petz divergence).

Outline. In Section 2 we collect some mathematical preliminaries. We first explain our general notation (2 A). Subsequently, we state some definitions and properties related to permutation invariance (2 B), entropies and divergences (2 C), binary quantum state discrimination (2 D), and several types of Petz Rényi mutual information (2 E). In Section 3 we present our main results (Theorems 6, 7, 8).

2. PRELIMINARIES

A. Notation

We take "log" to refer to the natural logarithm. The set of natural numbers strictly smaller than $n \in \mathbb{N}$ is denoted by $[n] \coloneqq \{0, 1, \dots, n-1\}$.

Throughout this work, we restrict ourselves to finite-dimensional Hilbert spaces (over the field \mathbb{C}) for simplicity. The dimension of a Hilbert space A is denoted by $d_A := \dim(A) \in \mathbb{N}_{>0}$. The tensor product of two Hilbert spaces A and B is sometimes denoted by AB instead of $A \otimes B$, and $A^n := A^{\otimes n}$ for any $n \in \mathbb{N}_{>0}$. The set of linear maps from A to B is denoted by $\mathcal{L}(A, B)$, and the set of linear maps from A to itself is denoted by $\mathcal{L}(A) := \mathcal{L}(A, A)$. To facilitate concise notation, identities are occasionally omitted. For instance, for $X_A \in \mathcal{L}(A)$, " X_A " may be interpreted as $X_A \otimes 1_B \in \mathcal{L}(A \otimes B)$. The spectrum, kernel, and rank of $X \in \mathcal{L}(A)$ are denoted as $\operatorname{spec}(X)$, $\operatorname{ker}(X)$, and $\operatorname{rank}(X)$, respectively. The support of $X \in \mathcal{L}(A)$ is defined as the orthogonal complement of the kernel of X, and is denoted by $\sup p(X)$. For $X, Y \in \mathcal{L}(A), X \ll Y$ is true iff $\operatorname{ker}(Y) \subseteq \operatorname{ker}(X)$, and $X \notin Y$ is true iff $X \ll Y$ is false. For $X \in \mathcal{L}(A)$, $X \perp Y$ is true iff XY = YX = 0, and $X \neq Y$ is true iff $X \perp Y$ is false. For $X \in \mathcal{L}(A)$, $X \geq 0$ is true iff X is positive definite, and X > 0 is true iff X is positive definite. For two self-adjoint operators $X, Y \in \mathcal{L}(A), X \geq Y$ is true iff $X - Y \geq 0$.

The adjoint of $X \in \mathcal{L}(A)$ with respect to the inner product of A is denoted by X^{\dagger} . For a positive semidefinite $X \in \mathcal{L}(A)$, X^p is defined for $p \in \mathbb{R}$ by taking the power on the support of X. In the case where p = 1/2, the square root symbol is sometimes employed, $\sqrt{X} \coloneqq X^{1/2}$. The operator absolute value of $X \in \mathcal{L}(A)$ is defined as $|X| \coloneqq (X^{\dagger}X)^{1/2}$. The Schatten *p*-norm of $X \in \mathcal{L}(A)$ is defined as $|X| \coloneqq (X^{\dagger}X)^{1/2}$. The Schatten *p*-norm of $X \in \mathcal{L}(A)$ is defined as $|X|_p \coloneqq \operatorname{tr}[|X|^p]^{1/p}$ for $p \in [1, \infty)$, and as $||X||_{\infty} \coloneqq \sqrt{\max(\operatorname{spec}(X^{\dagger}X))}$ for $p = \infty$. The Schatten *p*-quasi-norm is defined as $||X||_p \coloneqq \operatorname{tr}[|X|^p]^{1/p}$ for $p \in (0, 1)$.

If $X, Y \in \mathcal{L}(A)$ are self-adjoint operators, then $\{X \ge Y\}$ denotes the orthogonal projection onto the subspace corresponding to the non-negative eigenvalues of X - Y. The orthogonal projection onto the subspace corresponding to the strictly negative eigenvalues of X - Y is denoted by $\{X < Y\} \coloneqq 1 - \{X \ge Y\}.$

If $X, Y \in \mathcal{L}(A)$ are positive definite, then the geometric operator mean [35–37] is defined as $X \# Y := X^{\frac{1}{2}} (X^{-\frac{1}{2}} Y X^{-\frac{1}{2}})^{\frac{1}{2}} X^{\frac{1}{2}}$. This definition is extended to the case where $X, Y \in \mathcal{L}(A)$ are positive semidefinite by setting $X \# Y := \lim_{\varepsilon \to 0} (X + \varepsilon 1) \# (Y + \varepsilon 1)$, where "lim" denotes the limit in the strong operator topology. Since we are only working with finite-dimensional Hilbert spaces, the strong operator topology coincides with the weak operator topology and with the norm topology induced by the operator norm. "lim" can therefore be interpreted as an operator limit with respect to any of these topologies. The geometric operator mean is subadditive [37], i.e., $(X \# Y) + (X' \# Y') \leq (X + X') \# (Y + Y')$ for all positive semidefinite operators $X, X', Y, Y' \in \mathcal{L}(A)$.

The set of unitary operators on A is denoted by $\mathcal{U}(A) \subseteq \mathcal{L}(A)$. The set of quantum states on A is $\mathcal{S}(A) := \{\rho \in \mathcal{L}(A) : \rho \ge 0, \operatorname{tr}[\rho] = 1\}$. The set of positive definite quantum states on A is denoted by $\mathcal{S}_{>0}(A) := \{\rho \in \mathcal{S}(A) : \rho > 0\}$. Moreover, the following constrained versions of $\mathcal{S}(A)$ are defined with respect to a self-adjoint $X \in \mathcal{L}(A)$.

$$\mathcal{S}_{\not\perp X}(A) \coloneqq \{ \rho \in \mathcal{S}(A) : \rho \not\perp X \}$$

$$(2.1)$$

$$\mathcal{S}_{\ll X}(A) \coloneqq \{\rho \in \mathcal{S}(A) : \rho \ll X\}$$
(2.2)

$$\mathcal{S}_{X\ll}(A) \coloneqq \{\rho \in \mathcal{S}(A) : X \ll \rho\}$$
(2.3)

$$\mathcal{S}_{\ll X \ll}(A) \coloneqq \{ \rho \in \mathcal{S}(A) : \rho \ll X, X \ll \rho \}$$
(2.4)

The set of completely positive, trace-preserving linear maps from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ is denoted by $\operatorname{CPTP}(A, B)$.

B. Permutation invariance

The symmetric group of degree $n \in \mathbb{N}_{>0}$ is denoted by S_n . The unitary operator $U(\pi)_{A^n} \in \mathcal{L}(A^{\otimes n})$ associated with $\pi \in S_n$ is defined by the requirement that

$$U(\pi)_{A^n}|\psi_1\rangle \otimes \cdots \otimes |\psi_n\rangle = |\psi_{\pi^{-1}(1)}\rangle \otimes \cdots \otimes |\psi_{\pi^{-1}(n)}\rangle \qquad \forall |\psi_1\rangle, \dots, |\psi_n\rangle \in A.$$
(2.5)

The symmetric subspace of $A^{\otimes n}$ is denoted by

$$\operatorname{Sym}^{n}(A) \coloneqq \{ |\psi\rangle_{A^{n}} \in A^{\otimes n} : U(\pi)_{A^{n}} |\psi\rangle_{A^{n}} = |\psi\rangle_{A^{n}} \,\forall \pi \in S_{n} \},$$
(2.6)

and the orthogonal projection onto $\operatorname{Sym}^n(A)$ is denoted by $P_{\operatorname{sym}}^n \in \mathcal{L}(A^n)$. The set of *permutation* invariant operators is

$$\mathcal{L}_{\text{sym}}(A^{\otimes n}) \coloneqq \{ X_{A^n} \in \mathcal{L}(A^{\otimes n}) : U(\pi)_{A^n} X_{A^n} U(\pi)_{A^n}^{\dagger} = X_{A^n} \, \forall \pi \in S_n \},$$
(2.7)

and the set of *permutation invariant states* is $\mathcal{S}_{sym}(A^{\otimes n}) \coloneqq \mathcal{S}(A^{\otimes n}) \cap \mathcal{L}_{sym}(A^{\otimes n})$.

We will make use of the following construction of a permutation invariant state [15, 38, 39]. Let A and A' be isomorphic Hilbert spaces of dimension $d_A \in \mathbb{N}_{>0}$. The universal permutation invariant state is defined for any $n \in \mathbb{N}_{>0}$ as

$$\omega_{A^n}^n \coloneqq \frac{1}{g_{n,d_A}} \operatorname{tr}_{A'^n}[(P_{\operatorname{sym}}^n)_{A^n A'^n}], \quad \text{where} \quad g_{n,d_A} \coloneqq \dim(\operatorname{Sym}^n(AA')) = \binom{n + d_A^2 - 1}{n}.$$
(2.8)

The following properties of $\omega_{A^n}^n$ have been established in previous work.

Proposition 1 (Universal permutation invariant state). [15, 38, 39] Let $n \in \mathbb{N}_{>0}$. Then all of the following hold.

- (a) $\omega_{A^n}^n \in \mathcal{S}_{sym}(A^{\otimes n}).$
- (b) $\sigma_{A^n} \leq g_{n,d_A} \omega_{A^n}^n$ for all $\sigma_{A^n} \in \mathcal{S}_{sym}(A^{\otimes n})$, and $1 \leq g_{n,d_A} \leq (n+1)^{d_A^2 1}$. As a consequence, $\lim_{n \to \infty} \frac{1}{n} \log g_{n,d_A} = 0.$
- (c) $\sigma_{A^n} \omega_{A^n}^n = \omega_{A^n}^n \sigma_{A^n}$ for all $\sigma_{A^n} \in \mathcal{S}_{sym}(A^{\otimes n})$.
- (d) Let $|0\rangle_{AA'} \in AA'$ be an arbitrary but fixed unit vector. Then

$$(P_{\text{sym}}^n)_{A^n A'^n} = g_{n,d_A} \int_{\mathcal{U}(AA')} \mathrm{d}\mu_H(U) \left(|\sigma(U)\rangle\!\langle \sigma(U)|_{AA'} \right)^{\otimes n}, \tag{2.9}$$

where $d\mu_H$ denotes the Haar measure on the unitary group $\mathcal{U}(AA')$ normalized so that $\int_{\mathcal{U}(AA')} d\mu_H(U) = 1$, and $|\sigma(U)\rangle_{AA'} \coloneqq U|0\rangle_{AA'}$. As a consequence,

$$\omega_{A^n}^n = \int_{\mathcal{U}(AA')} \mathrm{d}\mu_H(U) \,\sigma(U)_A^{\otimes n},\tag{2.10}$$

where $\sigma(U)_A \coloneqq \operatorname{tr}_{A'}[|\sigma(U)\rangle\!\langle\sigma(U)|_{AA'}].$

C. Entropies and divergences

The von Neumann entropy of $\rho \in \mathcal{S}(A)$ is defined as $H(A)_{\rho} \coloneqq -\operatorname{tr}[\rho \log \rho]$. For $\rho \in \mathcal{S}(AB)$, the conditional entropy of A given B is $H(A|B)_{\rho} \coloneqq H(AB)_{\rho} - H(B)_{\rho}$ and the mutual information between A and B is $I(A:B)_{\rho} \coloneqq H(A)_{\rho} + H(B)_{\rho} - H(AB)_{\rho}$. The Rényi entropy (of order α) of $\rho \in \mathcal{S}(A)$ is defined as $H_{\alpha}(A)_{\rho} \coloneqq \frac{1}{1-\alpha} \log \operatorname{tr}[\rho^{\alpha}]$ for $\alpha \in (-\infty, 1) \cup (1, \infty)$, and for $\alpha \in \{1, \infty\}$ as the corresponding limits.

The quantum relative entropy is defined for $\rho \in \mathcal{S}(A)$ and a positive semidefinite $\sigma \in \mathcal{L}(A)$ as

$$D(\rho \| \sigma) \coloneqq \operatorname{tr}[\rho(\log \rho - \log \sigma)]$$
(2.11)

if $\rho \ll \sigma$ and $D(\rho \| \sigma) \coloneqq \infty$ else. The mutual information of $\rho_{AB} \in \mathcal{S}(AB)$ can be expressed in terms of the quantum relative entropy in the following ways.

$$I(A:B)_{\rho} = D(\rho_{AB} \| \rho_A \otimes \rho_B) = \inf_{\tau_B \in \mathcal{S}(B)} D(\rho_{AB} \| \rho_A \otimes \tau_B) = \inf_{\substack{\sigma_A \in \mathcal{S}(A), \\ \tau_B \in \mathcal{S}(B)}} D(\rho_{AB} \| \sigma_A \otimes \tau_B)$$
(2.12)

The latter two equalities can be derived via Klein's inequality, as elucidated in [18, Eq. (35)]. The argument based on Klein's inequality implies (due to the positive definiteness of the quantum relative entropy) that the corresponding minimizers are uniquely given by ρ_A and ρ_B , i.e.,

$$\underset{\tau_B \in \mathcal{S}(B)}{\arg\min} D(\rho_{AB} \| \rho_A \otimes \tau_B) = \{\rho_B\}, \qquad (2.13)$$

$$\underset{(\sigma_A,\tau_B)\in\mathcal{S}(A)\times\mathcal{S}(B)}{\arg\min} D(\rho_{AB} \| \sigma_A \otimes \tau_B) = \{(\rho_A,\rho_B)\}.$$
(2.14)

More generally, the argument based on Klein's inequality implies that for any $\sigma_A \in \mathcal{S}_{\rho_A \ll}(A)$

$$\underset{\tau_B \in \mathcal{S}(B)}{\arg\min} D(\rho_{AB} \| \sigma_A \otimes \tau_B) = \{\rho_B\}.$$
(2.15)

The quantum information variance is defined as $V(\rho \| \sigma) \coloneqq \operatorname{tr}[\rho(\log \rho - \log \sigma - D(\rho \| \sigma))^2]$ for $\rho, \sigma \in \mathcal{S}(A)$ [40, 41]. The mutual information variance of $\rho_{AB} \in \mathcal{S}(AB)$ is defined as [15]

$$V(A:B)_{\rho} \coloneqq V(\rho_{AB} \| \rho_A \otimes \rho_B) = \operatorname{tr}[\rho_{AB}(\log \rho_{AB} - \log(\rho_A \otimes \rho_B) - I(A:B)_{\rho})^2].$$
(2.16)

The Petz (quantum Rényi) divergence (of order α) is defined for $\alpha \in (0,1) \cup (1,\infty), \rho \in \mathcal{S}(A)$ and any positive semidefinite $\sigma \in \mathcal{L}(A)$ as [19]

$$D_{\alpha}(\rho \| \sigma) \coloneqq \frac{1}{\alpha - 1} \log \operatorname{tr}[\rho^{\alpha} \sigma^{1 - \alpha}]$$
(2.17)

if $(\alpha < 1 \land \rho \not\perp \sigma) \lor \rho \ll \sigma$ and $D_{\alpha}(\rho \| \sigma) \coloneqq \infty$ else. Moreover, D_0 and D_1 are defined as the limits of D_{α} for $\alpha \to \{0,1\}$. For $\alpha \in (-\infty, \infty)$, we define $Q_{\alpha}(\rho \| \sigma) \coloneqq \operatorname{tr}[\rho^{\alpha} \sigma^{1-\alpha}]$ for all positive semidefinite $\rho, \sigma \in \mathcal{L}(A)$. We will make use of the following properties of the Petz divergence.

Proposition 2 (Petz divergence). Let $\rho \in S(A)$ and let $\sigma \in \mathcal{L}(A)$ be positive semidefinite. Then all of the following hold. [19, 42–45]

- (a) Data-processing inequality: $D_{\alpha}(\rho \| \sigma) \geq D_{\alpha}(\mathcal{M}(\rho) \| \mathcal{M}(\sigma))$ for any $\mathcal{M} \in \text{CPTP}(A, A')$ and all $\alpha \in [0, 2]$.
- (b) Invariance under isometries: $D_{\alpha}(V\rho V^{\dagger} || V\sigma V^{\dagger}) = D_{\alpha}(\rho || \sigma)$ for any isometry $V \in \mathcal{L}(A, A')$ and all $\alpha \in [0, \infty)$.
- (c) Additivity: Let $\rho'_B \in \mathcal{S}(B)$ and let $\sigma'_B \in \mathcal{L}(B)$ be positive semidefinite. Then $D_{\alpha}(\rho_A \otimes \rho'_B \| \sigma_A \otimes \sigma'_B) = D_{\alpha}(\rho_A \| \sigma_A) + D_{\alpha}(\rho'_B \| \sigma'_B)$ for all $\alpha \in [0, \infty)$.
- (d) Normalization: $D_{\alpha}(\rho \| c\sigma) = D_{\alpha}(\rho \| \sigma) \log c \text{ for all } \alpha \in [0, \infty), c \in (0, \infty).$
- (e) Dominance: If $\sigma' \in \mathcal{L}(A)$ is positive semidefinite and such that $\sigma \leq \sigma'$, then $D_{\alpha}(\rho \| \sigma) \geq D_{\alpha}(\rho \| \sigma')$ for all $\alpha \in [0, 2]$.
- (f) Non-negativity: If $\sigma \in \mathcal{S}(A)$, then $D_{\alpha}(\rho \| \sigma) \in [0, \infty]$ for all $\alpha \in [0, \infty)$.

- (g) Positive definiteness: Let $\alpha \in (0, \infty)$. If $\sigma \in \mathcal{S}(A)$, then $D_{\alpha}(\rho \| \sigma) = 0$ iff $\rho = \sigma$. Furthermore, $D_0(\rho \| \rho) = 0$.
- (h) Rényi order $\alpha = 1$: $D_1(\rho \| \sigma) = D(\rho \| \sigma)$.
- (i) Monotonicity in α : If $\alpha, \beta \in [0, \infty)$ are such that $\alpha \leq \beta$, then $D_{\alpha}(\rho \| \sigma) \leq D_{\beta}(\rho \| \sigma)$.
- (j) Continuity in α : If $\rho \not\perp \sigma$, then the function $[0,1) \to \mathbb{R}, \alpha \mapsto D_{\alpha}(\rho \| \sigma)$ is continuous. If $\rho \ll \sigma$, then the function $[0,\infty) \to \mathbb{R}, \alpha \mapsto D_{\alpha}(\rho \| \sigma)$ is continuous.
- (k) Differentiability in α : If $\rho \not\perp \sigma$, then the function $(0,1) \to \mathbb{R}, \alpha \mapsto D_{\alpha}(\rho \| \sigma)$ is continuously differentiable. If $\rho \ll \sigma$, then the function $(0,\infty) \to \mathbb{R}, \alpha \mapsto D_{\alpha}(\rho \| \sigma)$ is continuously differentiable. Furthermore, if $\rho \ll \sigma$ and $\sigma \in \mathcal{S}(A)$, then $\frac{d}{d\alpha} D_{\alpha}(\rho \| \sigma)|_{\alpha=1} = \frac{1}{2} V(\rho \| \sigma)$.
- (l) Convexity in α : If $\rho \not\perp \sigma$, then the function $[0,1) \to \mathbb{R}, \alpha \mapsto (\alpha 1)D_{\alpha}(\rho \| \sigma)$ is convex. If $\rho \ll \sigma$, then the function $[0,\infty) \to \mathbb{R}, \alpha \mapsto (\alpha 1)D_{\alpha}(\rho \| \sigma)$ is convex.

The sandwiched (quantum Rényi) divergence (of order α) is defined for $\alpha \in (0,1) \cup (1,\infty), \rho \in \mathcal{S}(A)$ and any positive semidefinite $\sigma \in \mathcal{L}(A)$ as [20, 21]

$$\widetilde{D}_{\alpha}(\rho \| \sigma) \coloneqq \frac{1}{\alpha - 1} \log \operatorname{tr}[(\sigma^{\frac{1 - \alpha}{2\alpha}} \rho \sigma^{\frac{1 - \alpha}{2\alpha}})^{\alpha}]$$
(2.18)

if $(\alpha < 1 \land \rho \not\perp \sigma) \lor \rho \ll \sigma$ and $\widetilde{D}_{\alpha}(\rho \| \sigma) \coloneqq \infty$ else. Moreover, \widetilde{D}_1 and \widetilde{D}_{∞} are defined as the limits of \widetilde{D}_{α} for $\alpha \to \{1, \infty\}$. The limit $\alpha \to 1$ is identical to the quantum relative entropy, $\lim_{\alpha \to 1} \widetilde{D}_{\alpha}(\rho \| \sigma) = D(\rho \| \sigma)$. For $\alpha \in (0, \infty)$, we define $\widetilde{Q}_{\alpha}(\rho \| \sigma) \coloneqq \operatorname{tr}[(\sigma^{\frac{1-\alpha}{2\alpha}}\rho\sigma^{\frac{1-\alpha}{2\alpha}})^{\alpha}]$ for all positive semidefinite $\rho, \sigma \in \mathcal{L}(A)$.

The Araki-Lieb-Thirring inequality [46–48] implies that $\widetilde{Q}_{\alpha}(\rho \| \sigma) \geq Q_{\alpha}(\rho \| \sigma)$ if $\alpha \in (0, 1]$ and $\widetilde{Q}_{\alpha}(\rho \| \sigma) \leq Q_{\alpha}(\rho \| \sigma)$ if $\alpha \in [1, \infty)$ for all positive semidefinite $\rho, \sigma \in \mathcal{L}(A)$.

D. Binary quantum state discrimination

In this section, we will explain the notation related to binary discrimination. First, we will describe the classical setting, then the quantum setting, and then we will explain how the classical setting can be regarded as a special case of the quantum setting under suitable conditions.

Classical setting. Classical binary discrimination pertains to a scenario where one is given a PMF S_X over \mathcal{X} that is an element of H_0 or H_1 , both of which are non-empty subsets of the set of PMFs over \mathcal{X} . The task is to decide which is true: $S_X \in H_0$ (the null hypothesis) or $S_X \in H_1$ (the alternative hypothesis). The test is a function $T : \mathcal{X} \to [0, 1]$, and the decision is determined by the binary test (T, 1 - T) applied to S_X . If the event corresponding to T occurs, then the decision is made that the null hypothesis is true. Conversely, if the event corresponding to 1 - T occurs, then the decision is made that the alternative hypothesis is true. This paper solely addresses discrimination problems with a simple null hypothesis, i.e., $H_0 = \{P_X\}$ for some PMF P_X .

We are mainly concerned with sequences of binary discrimination problems of the following form, for a fixed PMF P_{XY} over $\mathcal{X} \times \mathcal{Y}$. For each $n \in \mathbb{N}_{>0}$, the null hypothesis is $H_0^n = \{P_{XY}^{\times n}\}$, the alternative hypothesis H_1^n is a non-empty subset of the set of PMFs over $(\mathcal{X} \times \mathcal{Y})^{\times n}$, and the test is a function $T^n : (\mathcal{X} \times \mathcal{Y})^{\times n} \to [0, 1]$. The type-I error and the (worst case) type-II error are then, respectively,

$$\alpha_n(T^n) \coloneqq \sum_{\substack{x_1, \dots, x_n \in \mathcal{X}, \\ y_1, \dots, y_n \in \mathcal{Y}}} P_{XY}(x_1, y_1) \cdot \dots \cdot P_{XY}(x_n, y_n) (1 - T^n(x_1, y_1, \dots, x_n, y_n)),$$
(2.19)

$$\beta_n(T^n) \coloneqq \sup_{\substack{Q_{X^nY^n} \in H_1^n \\ y_1, \dots, y_n \in \mathcal{Y}}} \sum_{\substack{x_1, \dots, x_n \in \mathcal{X}, \\ y_1, \dots, y_n \in \mathcal{Y}}} Q_{X^nY^n}(x_1, y_1, \dots, x_n, y_n) T^n(x_1, y_1, \dots, x_n, y_n).$$
(2.20)

$$\hat{\alpha}_n(\mu) \coloneqq \inf_{T^n} \{ \alpha_n(T^n) : \beta_n(T^n) \le \mu \},$$
(2.21)

where the minimization is over all functions $T^n : (\mathcal{X} \times \mathcal{Y})^{\times n} \to [0, 1].$

Quantum setting. Binary quantum state discrimination pertains to a scenario where one is given a quantum state $\xi \in S(A)$ that is an element of H_0 or H_1 , both of which are non-empty subsets of S(A). The task is to decide which is true: $\xi \in H_0$ (the null hypothesis) or $\xi \in H_1$ (the alternative hypothesis). The test is some $T \in \mathcal{L}(A)$ that satisfies $0 \leq T \leq 1$. The decision is determined by the binary measurement (T, 1 - T) applied to ξ . If the measurement outcome is the one associated with T, then the decision is made that the null hypothesis is true. Conversely, if the measurement outcome is the one associated with 1 - T, then the decision is made that the alternative hypothesis is true. This paper solely addresses quantum state discrimination problems with a simple null hypothesis, i.e., $H_0 = \{\rho\}$ for some $\rho \in S(A)$.

We are mainly concerned with sequences of binary quantum state discrimination problems of the following form, for a fixed $\rho_{AB} \in \mathcal{S}(AB)$. For each $n \in \mathbb{N}_{>0}$, the null hypothesis is $H_0^n = \{\rho_{AB}^{\otimes n}\}$, the alternative hypothesis H_1^n is a non-empty subset of $\mathcal{S}(A^nB^n)$, and the test is some $T_{A^nB^n}^n \in \mathcal{L}(A^nB^n)$ that satisfies $0 \leq T_{A^nB^n}^n \leq 1$. The type-I error and the (worst case) type-II error are then, respectively,

$$\alpha_n(T_{A^nB^n}^n) \coloneqq \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^nB^n}^n)], \qquad (2.22)$$

$$\beta_n(T_{A^nB^n}^n) \coloneqq \sup_{\sigma_{A^nB^n} \in H_1^n} \operatorname{tr}[\sigma_{A^nB^n} T_{A^nB^n}^n].$$
(2.23)

For the *n*th hypothesis testing problem, the *minimum* type-I error when the type-II error is upper bounded by $\mu \in [0, \infty)$ is denoted by

$$\hat{\alpha}_{n}(\mu) \coloneqq \inf_{\substack{T_{A^{n}B^{n}} \in \mathcal{L}(A^{n}B^{n}):\\ 0 \le T_{A^{n}B^{n}} \le 1}} \{ \alpha_{n}(T_{A^{n}B^{n}}) : \beta_{n}(T_{A^{n}B^{n}}) \le \mu \}.$$
(2.24)

The trade-off between type-I and type-II errors can be quantified in the asymptotics $(n \to \infty)$ by various error exponents [9, 49]. This paper solely addresses one of them: the direct exponent. The *direct exponent* with respect to $R \in [0, \infty)$ is defined as $\liminf_{n\to\infty} -\frac{1}{n}\log \hat{\alpha}_n(e^{-nR})$ if this limit exists, and as $+\infty$ else [9, 49]. The parameter R is called the *type-II rate*.

CC states. Binary quantum state discrimination can be compared to classical binary discrimination by restricting the former to the special case of CC states, as we will elaborate in the following. Let $\mathcal{X} := [d_A], \mathcal{Y} := [d_B]$, let P_{XY} be a PMF over $\mathcal{X} \times \mathcal{Y}$, and let $\rho_{AB} :=$ $\sum_{x \in \mathcal{X}, y \in \mathcal{Y}} P_{XY}(x, y) |a_x, b_y\rangle \langle a_x, b_y|_{AB}$, where $\{|a_x\rangle_A\}_{x \in [d_A]}$ and $\{|b_y\rangle_B\}_{y \in [d_B]}$ are orthonormal bases for A and B, respectively.

Consider a sequence of binary quantum state discrimination problems as above with null hypothesis $H_0^{q,n} = \{\rho_{AB}^{\otimes n}\}$ and alternative hypothesis $H_1^{q,n}$, and let the function in (2.24) be denoted by $\hat{\alpha}_n^q$. For notational convenience, let us define for any $\sigma_{A^nB^n} \in H_1^{q,n}$:

$$Q_{X^nY^n}^{\sigma}(x_1, y_1, \dots, x_n, y_n) \coloneqq \langle a_{x_1}, b_{y_1}, \dots, a_{x_n}, b_{y_n} | \sigma_{A^nB^n} | a_{x_1}, b_{y_1}, \dots, a_{x_n}, b_{y_n} \rangle$$
(2.25)

for all $x_1, \ldots, x_n \in \mathcal{X}, y_1, \ldots, y_n \in \mathcal{Y}$. To facilitate subsequent comparison with the classical setting, suppose that the alternative hypothesis is closed under pinching with respect to the eigenbasis of the CC state ρ_{AB} , i.e., for all $\sigma_{A^nB^n} \in H_1^{q,n}$:

$$\sum_{\substack{x_1,\dots,x_n \in \mathcal{X}, \\ y_1,\dots,y_n \in \mathcal{Y}}} Q_{X^n Y^n}^{\sigma}(x_1, y_1,\dots,x_n, y_n) | a_{x_1}, b_{y_1},\dots, a_{x_n}, b_{y_n} \rangle \langle a_{x_1}, b_{y_1},\dots, a_{x_n}, b_{y_n} | \in H_1^{q,n}.$$
(2.26)

Consider now a sequence of *classical* binary discrimination problems as above with null hypothesis $H_0^{c,n} = \{P_{XY}^{\times n}\}$ and alternative hypothesis $H_1^{c,n}$, and let the function in (2.21) be denoted by $\hat{\alpha}_n^c$. Suppose that the alternative hypotheses for the quantum and the classical setting are *analogous* in the sense that

$$\bigcup_{\sigma_{A^n B^n} \in H_1^{q,n}} \{ Q_{X^n Y^n}^{\sigma} \} = H_1^{c,n}.$$
(2.27)

Assuming that the conditions in (2.26) and (2.27) are satisfied, it is straightforward to derive that $\hat{\alpha}_n^{\rm q}(\mu) = \hat{\alpha}_n^{\rm c}(\mu)$ for all $\mu \in [0, \infty)$, as shown in Appendix A. Thus, certain classical binary discrimination problems can be regarded as special cases of binary quantum state discrimination problems.

In particular, the conditions in (2.26) and (2.27) are satisfied for the examples in Table I and Table II. Therefore, the classical discrimination problems in Table I can be regarded as special cases of the corresponding binary quantum state discrimination problems in Table II, so propositions in Table II imply corresponding propositions in Table I. In particular, this implies that the proposition in the second row of Table I can be extended to $R \in (0, \infty)$.

E. Petz Rényi mutual information

This section reiterates the definitions of the three types of Petz Rényi mutual information (PRMI) that were mentioned in the introduction. Prior to that, two generalized PRMIs will be introduced.

We define for $\alpha \in [0, \infty)$, $\rho_{AB} \in \mathcal{S}(AB)$ and any positive semidefinite $\sigma_A \in \mathcal{L}(A)$

$$I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A) \coloneqq D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \rho_B), \qquad (2.28)$$

$$I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) \coloneqq \inf_{\tau_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B).$$
(2.29)

We call them the non-minimized generalized PRMI (of order α) and the minimized generalized PRMI (of order α), respectively. The minimized generalized PRMI was introduced in [15]. For an overview, we include a list of properties of these two generalized PRMIs in Appendix B. Some of these properties have been established in previous work [15], and the remaining properties follow immediately from the definitions in (2.28) and (2.29), as explained in Appendix C.

As in the introduction, we define the following PRMIs for $\alpha \in [0, \infty)$ and $\rho_{AB} \in \mathcal{S}(AB)$.

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \coloneqq D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B) = I_{\alpha}^{\uparrow}(\rho_{AB} \| \rho_A)$$
(2.30)

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \coloneqq \inf_{\tau_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \tau_B) = \inf_{\tau_B \in \mathcal{S}(B)} I_{\alpha}^{\uparrow}(\rho_{AB} \| \tau_B) = I_{\alpha}^{\downarrow}(\rho_{AB} \| \rho_A)$$
(2.31)

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \coloneqq \inf_{\substack{\sigma_A \in \mathcal{S}(A), \\ \tau_B \in \mathcal{S}(B)}} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) = \inf_{\sigma_A \in \mathcal{S}(A)} I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A)$$
(2.32)

We call them the non-minimized PRMI (of order α), the singly minimized PRMI (of order α), and the doubly minimized PRMI (of order α), respectively. It should be noted that the singly minimized PRMI is not generally invariant under a swap of A and B for $\alpha \neq 1$. To provide an overview, we present a list of properties of the non-minimized PRMI and the singly minimized PRMI below. The majority of these properties can be derived directly from the definitions in (2.30) and (2.31). The remaining properties have been established in previous work or can be derived from it immediately, as elucidated in Appendix D. **Proposition 3** (Non-minimized PRMI). Let $\rho_{AB} \in \mathcal{S}(AB)$. Then all of the following hold.

- (a) Symmetry: $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\uparrow}(B:A)_{\rho}$ for all $\alpha \in [0,\infty)$.
- (b) Non-increase under local operations: $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \geq I_{\alpha}^{\uparrow\uparrow}(A':B')_{\mathcal{M}\otimes\mathcal{N}(\rho)}$ for any $\mathcal{M} \in \operatorname{CPTP}(A,A'), \mathcal{N} \in \operatorname{CPTP}(B,B')$ and all $\alpha \in [0,2]$.
- (c) Invariance under local isometries: $I_{\alpha}^{\uparrow\uparrow}(A':B')_{V\otimes W\rho V^{\dagger}\otimes W^{\dagger}} = I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho}$ for any isometries $V \in \mathcal{L}(A,A'), W \in \mathcal{L}(B,B')$ and all $\alpha \in [0,\infty)$.
- (d) Additivity: Let $\alpha \in [0, \infty)$ and $\rho'_{DE} \in \mathcal{S}(DE)$. Then

$$I_{\alpha}^{\uparrow\uparrow}(AD:BE)_{\rho_{AB}\otimes\rho'_{DE}} = I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho_{AB}} + I_{\alpha}^{\uparrow\uparrow}(D:E)_{\rho'_{DE}}.$$
(2.33)

- (e) Duality: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in [0,1) \cup (1,\infty)$ and $\beta \coloneqq 2 \alpha$. Then $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = -\frac{1}{\beta-1}\log Q_{\beta}(\rho_{AC} \| \rho_A^{-1} \otimes \rho_C)$.
- (f) Non-negativity: $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \geq 0$ for all $\alpha \in [0,\infty)$.
- (g) Upper bound: Let $\alpha \in [0, \frac{3}{2}]$ and $r_A \coloneqq \operatorname{rank}(\rho_A)$. Then $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \leq 2\log r_A$. Furthermore, if $\alpha \in [0, \frac{3}{2})$, then $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = 2\log r_A$ iff $\operatorname{spec}(\rho_A) \subseteq \{0, 1/r_A\}$ and $H(A|B)_{\rho} = -\log r_A$.
- (h) Rényi order $\alpha = 1$: $I_1^{\uparrow\uparrow}(A:B)_{\rho} = I(A:B)_{\rho}$.
- (i) Monotonicity in α : If $\alpha, \beta \in [0, \infty)$ are such that $\alpha \leq \beta$, then $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \leq I_{\beta}^{\uparrow\uparrow}(A:B)_{\rho}$.
- (j) Continuity in α : The function $[0,\infty) \to [0,\infty), \alpha \mapsto I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho}$ is continuous.
- (k) Differentiability in α : The function $(0, \infty) \to [0, \infty), \alpha \mapsto I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho}$ is continuously differentiable, and the derivative at $\alpha \in (0, \infty)$ is

$$\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B).$$
(2.34)

In particular, $\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho}|_{\alpha=1} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B)|_{\alpha=1} = \frac{1}{2} V(A:B)_{\rho}.$

- (l) Convexity in α : The function $[0,\infty) \to \mathbb{R}, \alpha \mapsto (\alpha-1)I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho}$ is convex.
- (m) Product states: If $\rho_{AB} = \rho_A \otimes \rho_B$, then $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = 0$ for all $\alpha \in [0,\infty)$. Conversely, for any $\alpha \in (0,\infty)$, if $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = 0$, then $\rho_{AB} = \rho_A \otimes \rho_B$.
- (n) AC-independent states: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. If $\rho_{AC} = \rho_A \otimes \rho_C$, then $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = 2H_{3-2\alpha}(A)_{\rho}$ for all $\alpha \in [0,\infty)$.
- (o) Pure states: [50] If there exists $|\rho\rangle_{AB} \in AB$ such that $\rho_{AB} = |\rho\rangle\langle\rho|_{AB}$, then for all $\alpha \in [0, \infty)$

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{|\rho\rangle\langle\rho|} = 2H_{3-2\alpha}(A)_{\rho}.$$
(2.35)

(p) CC states: Let P_{XY} be the joint PMF of two random variables X, Y over $\mathcal{X} \coloneqq [d_A], \mathcal{Y} \coloneqq [d_B]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x\in[d_A]}, \{|b_y\rangle_B\}_{y\in[d_B]}$ for A, B such that $\rho_{AB} = \sum_{x\in\mathcal{X}}\sum_{y\in\mathcal{Y}}P_{XY}(x,y)|a_x,b_y\rangle\langle a_x,b_y|_{AB}$, then for all $\alpha \in [0,\infty)$

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\uparrow}(X:Y)_{P}.$$
(2.36)

(q) Copy-CC states: Let P_X be the PMF of a random variable X over $\mathcal{X} := [\min(d_A, d_B)]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x\in[d_A]}, \{|b_y\rangle_B\}_{y\in[d_B]}$ for A, B such that $\rho_{AB} = \sum_{x\in\mathcal{X}} P_X(x)|a_x, b_x\rangle\langle a_x, b_x|_{AB}$, then for all $\alpha \in [0, \infty)$

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\uparrow}(X:Y)_{P} = H_{2-\alpha}(A)_{\rho}.$$
(2.37)

Remark 1 (Order in the list). The properties in Proposition 3 are organized into three groups. The first group (a)–(g) concerns general properties for a fixed Rényi order α . The second group (h)–(l) deals with special values of α and the behavior as α is varied. The third group (m)–(q) addresses special states. This qualitative organization into three groups is also retained in subsequent lists, with varying numbers of items within the groups.

Proposition 4 (Singly minimized PRMI). Let $\rho_{AB} \in \mathcal{S}(AB)$. Then all of the following hold.

- (a) Non-increase under local operations: $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \geq I_{\alpha}^{\uparrow\downarrow}(A':B')_{\mathcal{M}\otimes\mathcal{N}(\rho)}$ for any $\mathcal{M} \in \operatorname{CPTP}(A,A'), \mathcal{N} \in \operatorname{CPTP}(B,B')$ and all $\alpha \in [0,2]$.
- (b) Invariance under local isometries: $I_{\alpha}^{\uparrow\downarrow}(A':B')_{V\otimes W\rho V^{\dagger}\otimes W^{\dagger}} = I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho}$ for any isometries $V \in \mathcal{L}(A,A'), W \in \mathcal{L}(B,B')$ and all $\alpha \in [0,\infty)$.
- (c) Additivity: [15] Let $\alpha \in [0, \infty)$ and $\rho'_{DE} \in \mathcal{S}(DE)$. Then

$$I_{\alpha}^{\uparrow\downarrow}(AD:BE)_{\rho_{AB}\otimes\rho'_{DE}} = I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho_{AB}} + I_{\alpha}^{\uparrow\downarrow}(D:E)_{\rho'_{DE}}.$$
(2.38)

- (d) Duality: [15] Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in [0,\infty)$ and $\beta := \frac{1}{\alpha} \in (0,\infty]$. Then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = -\widetilde{D}_{\beta}(\rho_{AC} \| \rho_A^{-1} \otimes \rho_C)$.
- (e) Non-negativity: $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \ge 0$ for all $\alpha \in [0,\infty)$.
- (f) Upper bound: Let $\alpha \in [0,\infty)$ and $r_A \coloneqq \operatorname{rank}(\rho_A)$. Then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \leq 2H_{-1}(A)_{\rho}$, and if $\alpha \in [0,2]$, then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \leq 2\log r_A$. Furthermore, if $\alpha \in [0,2)$, then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 2\log r_A$ iff $\operatorname{spec}(\rho_A) \subseteq \{0,1/r_A\}$ and $H(A|B)_{\rho} = -\log r_A$.
- (g) Existence and uniqueness of minimizer: [15, 17] Let $\alpha \in [0, \infty)$. If $\alpha \in (0, \infty)$, then $\hat{\tau}_B := (\operatorname{tr}_A[\rho_{AB}^{\alpha}\rho_A^{1-\alpha}])^{\frac{1}{\alpha}}/\operatorname{tr}[(\operatorname{tr}_A[\rho_{AB}^{\alpha}\rho_A^{1-\alpha}])^{\frac{1}{\alpha}}] \in \mathcal{S}_{\ll \rho_B \ll}(B)$ and

$$\underset{\tau_B \in \mathcal{S}(B)}{\arg\min} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \tau_B) = \{ \hat{\tau}_B \}.$$
(2.39)

If $\alpha = 0$, then $\emptyset \neq \arg \min_{\tau_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \tau_B) \subseteq \mathcal{S}_{\ll \rho_B}(B)$. (h) Closed-form expression: [15, 17] Let $\alpha \in (0, 1) \cup (1, \infty)$. Then

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = \frac{1}{\alpha - 1} \log \| \operatorname{tr}_{A}[\rho_{AB}^{\alpha} \rho_{A}^{1 - \alpha}] \|_{\frac{1}{\alpha}}.$$
 (2.40)

(i) Asymptotic optimality of universal permutation invariant state: Let $\alpha \in [0, 2]$. Then

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = \lim_{n \to \infty} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \rho_{A}^{\otimes n} \otimes \omega_{B^{n}}^{n})$$
(2.41)

and for any $n \in \mathbb{N}_{>0}$

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = \inf_{\tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \rho_A^{\otimes n} \otimes \tau_{B^n}) = \inf_{\tau_{B^n} \in \mathcal{S}(B^n)} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \rho_A^{\otimes n} \otimes \tau_{B^n}).$$

$$(2.42)$$

(j) Rényi order $\alpha \in \{0,1\} {:}~ I_1^{\uparrow \downarrow}(A:B)_\rho = I(A:B)_\rho$ and

$$I_0^{\uparrow\downarrow}(A:B)_{\rho} = -\log \|\operatorname{tr}_A[\rho_{AB}^0 \rho_A]\|_{\infty} = \min_{\substack{|\tau\rangle_B \in \operatorname{supp}(\rho_B):\\\langle\tau|\tau\rangle_B = 1}} D_0(\rho_{AB} \|\rho_A \otimes |\tau\rangle\langle\tau|_B).$$
(2.43)

- (k) Monotonicity in α : If $\alpha, \beta \in [0, \infty)$ are such that $\alpha \leq \beta$, then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \leq I_{\beta}^{\uparrow\downarrow}(A:B)_{\rho}$.
- (l) Continuity in α : [17] The function $[0,\infty) \to [0,\infty), \alpha \mapsto I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho}$ is continuous.
- (m) Differentiability in α : The function $(0,2) \to [0,\infty), \alpha \mapsto I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho}$ is continuously differentiable. For any $\alpha \in (0,2)$ and any fixed $\tau_B \in \arg \min_{\tau'_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \tau'_B)$, the derivative at α is

$$\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = \frac{\partial}{\partial\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \tau_B).$$
(2.44)

In particular, $\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho}|_{\alpha=1} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B)|_{\alpha=1} = \frac{1}{2} V(A:B)_{\rho}.$

- (n) Convexity in α : The function $[0,2] \to \mathbb{R}, \alpha \mapsto (\alpha-1)I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho}$ is convex.
- (o) Product states: If $\rho_{AB} = \rho_A \otimes \rho_B$, then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 0$ for all $\alpha \in [0,\infty)$. Conversely, for any $\alpha \in (0,\infty)$, if $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 0$, then $\rho_{AB} = \rho_A \otimes \rho_B$.
- (p) AC-independent states: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. If $\rho_{AC} = \rho_A \otimes \rho_C$, then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 2H_{\frac{2-\alpha}{\alpha}}(A)_{\rho}$ for all $\alpha \in (0,\infty)$.
- (q) Pure states: If there exists $|\rho\rangle_{AB} \stackrel{\alpha}{\in} AB$ such that $\rho_{AB} = |\rho\rangle\langle\rho|_{AB}$, then for all $\alpha \in (0,\infty)$

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{|\rho\rangle\langle\rho|} = D_{\alpha}(|\rho\rangle\langle\rho|_{AB} \|\rho_A \otimes \tau_B) = 2H_{\frac{2-\alpha}{\alpha}}(A)_{\rho}, \qquad (2.45)$$

where $\tau_B \coloneqq \rho_B^{\frac{2-\alpha}{\alpha}} / \operatorname{tr}[\rho_B^{\frac{2-\alpha}{\alpha}}].$

(r) CC states: Let P_{XY} be the joint PMF of two random variables X, Y over $\mathcal{X} \coloneqq [d_A], \mathcal{Y} \coloneqq [d_B]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x\in[d_A]}, \{|b_y\rangle_B\}_{y\in[d_B]}$ for A, B such that $\rho_{AB} = \sum_{x\in\mathcal{X}}\sum_{y\in\mathcal{Y}}P_{XY}(x,y)|a_x,b_y\rangle\langle a_x,b_y|_{AB}$, then for all $\alpha \in [0,\infty)$

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\downarrow}(X:Y)_{P}.$$
(2.46)

(s) Copy-CC states: Let P_X be the PMF of a random variable X over $\mathcal{X} := [\min(d_A, d_B)]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x\in[d_A]}, \{|b_y\rangle_B\}_{y\in[d_B]}$ for A, B such that $\rho_{AB} = \sum_{x\in\mathcal{X}} P_X(x)|a_x, b_x\rangle\langle a_x, b_x|_{AB}$, then for all $\alpha \in (0, \infty)$

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\downarrow}(X:Y)_{P} = D_{\alpha}(\rho_{AB} \| \rho_{A} \otimes \tau_{B}) = H_{\frac{1}{\alpha}}(A)_{\rho}, \qquad (2.47)$$

where $\tau_B := \rho_B^{\frac{1}{\alpha}} / \operatorname{tr}[\rho_B^{\frac{1}{\alpha}}].$

3. MAIN RESULTS

A. Properties of the doubly minimized Petz Rényi mutual information

In this section, we present our findings on properties of the doubly minimized PRMI of order α . We thereby focus on $\alpha \in (\frac{1}{2}, 1)$ because this will turn out to be the relevant range for the subsequent application of the doubly minimized PRMI in binary quantum state discrimination (Theorem 8). Several assertions are extended to larger ranges of α whenever the proof technique permits a straightforward extension.

In Theorem 6, we show that the minimization problem underlying the doubly minimized PRMI of order α is jointly convex for any $\alpha \in [\frac{1}{2}, 1)$, see (3.4). In [1], joint convexity has been proved for the classical case. More precisely, in [1, Lemma 15], the joint convexity of $D_{\alpha}(P_{XY}||Q_XR_Y)$ in Q_X and R_Y has been proved for any $\alpha \in [\frac{1}{2}, \infty)$. Our proof for the quantum case follows the same scheme as the aforementioned proof for the classical case. The main difference lies in one particular proof step where the proof for the classical case employs the Cauchy-Schwarz inequality for \mathbb{R}^2 with the standard inner product [1, Eq. (227)] in the form

$$\sqrt{x}\sqrt{y} + \sqrt{x'}\sqrt{y'} \le \sqrt{x+x'}\sqrt{y+y'} \qquad \forall x, y, x', y' \in [0,\infty).$$
(3.1)

In order to prove joint convexity for the quantum case, we employ an operator version of this inequality instead (Lemma 5), which follows from the subadditivity of the geometric operator mean. Remarkably, Lemma 5 shows that the Cauchy-Schwarz inequality (3.1) can be lifted from the positive real axis to positive semidefinite operators. The proof of Lemma 5 is given in Appendix E 1 and the proof of Theorem 6 is given in Appendix E 2.

Lemma 5 (Operator inequality from subadditivity of geometric operator mean). Let $X_A, X'_A \in$ $\mathcal{L}(A), Y_B, Y'_B \in \mathcal{L}(B)$ be positive semidefinite. Then

$$\sqrt{X_A} \otimes \sqrt{Y_B} + \sqrt{X_A'} \otimes \sqrt{Y_B'} \le \sqrt{X_A + X_A'} \otimes \sqrt{Y_B + Y_B'}.$$
(3.2)

Theorem 6 (Joint concavity/convexity). Let $\alpha \in [\frac{1}{2}, 1)$. Let $\rho_{AB} \in \mathcal{S}(AB), \sigma_A, \sigma'_A \in \mathcal{S}(A), \tau_B, \tau'_B \in \mathcal{S}(A)$ $\mathcal{S}(B)$, and let $\lambda, \lambda' \in (0,1)$ be such that $\lambda + \lambda' = 1$. Then

$$Q_{\alpha}(\rho_{AB} \| (\lambda \sigma_A + \lambda' \sigma'_A) \otimes (\lambda \tau_B + \lambda' \tau'_B)) \ge \lambda Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) + \lambda' Q_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tau'_B), \quad (3.3)$$

$$D_{\alpha}(\rho_{AB} \| (\lambda \sigma_A + \lambda' \sigma'_A) \otimes (\lambda \tau_B + \lambda' \tau'_B)) \leq \lambda D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) + \lambda' D_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tau'_B), \quad (3.4)$$

$$I^{\downarrow}_{\alpha}(\rho_{AB} \| \lambda \sigma_A + \lambda' \sigma'_A) \le \lambda I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) + \lambda' I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma'_A).$$
(3.5)

Furthermore, if $\alpha \in (\frac{1}{2}, 1), \sigma_A, \sigma'_A \in \mathcal{S}_{\ll \rho_A \ll}(A), \tau_B, \tau'_B \in \mathcal{S}_{\ll \rho_B \ll}(B)$, and at least one of the inequalities in (3.3) and (3.4) holds with equality, then $\sigma_A = \sigma'_A$ and $\tau_B = \tau'_B$. If $\alpha \in (\frac{1}{2}, 1), \sigma_A, \sigma'_A \in \mathcal{S}_{\ll \rho_A \ll}(A)$ $\mathcal{S}_{\ll \rho_A \ll}(A)$, and (3.5) holds with equality, then $\sigma_A = \sigma'_A$.

In Theorem 7, we enumerate several properties of the doubly minimized PRMI. The proof of Theorem 7 is given in Appendix E4. For the proof of Theorem 7 (k), we will use a lemma that asserts a general equivalence of optimizers and fixed-points. This lemma is stated and proved beforehand in Appendix $\mathbf{E3}$.

Theorem 7 (Doubly minimized PRMI). Let $\rho_{AB} \in \mathcal{S}(AB)$. Then all of the following hold.

- (a) Symmetry: $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\downarrow\downarrow}(B:A)_{\rho}$ for all $\alpha \in [0,\infty)$.
- (b) Non-increase under local operations: $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \geq I_{\alpha}^{\downarrow\downarrow}(A':B')_{\mathcal{M}\otimes\mathcal{N}(\rho)}$ for any $\mathcal{M} \in \mathcal{M}$ $\operatorname{CPTP}(A, A'), \mathcal{N} \in \operatorname{CPTP}(B, B') \text{ and all } \alpha \in [0, 2].$
- (c) Invariance under local isometries: $I_{\alpha}^{\downarrow\downarrow}(A':B')_{V\otimes W\rho V^{\dagger}\otimes W^{\dagger}} = I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ for any isometries $V \in \mathcal{L}(A, A'), W \in \mathcal{L}(B, B') \text{ and all } \alpha \in [0, \infty).$
- (d) Additivity: Let $\alpha \in [\frac{1}{2}, 2]$ and $\rho'_{DE} \in \mathcal{S}(DE)$. Then

$$I_{\alpha}^{\downarrow\downarrow}(AD:BE)_{\rho_{AB}\otimes\rho'_{DE}} = I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho_{AB}} + I_{\alpha}^{\downarrow\downarrow}(D:E)_{\rho'_{DE}}.$$
(3.6)

(e) Duality: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in (0,\infty)$ and $\beta :=$ $\frac{1}{\alpha} \in (0,\infty)$. Then

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \begin{cases} \inf_{\substack{\sigma_{A} \in \mathcal{S}(A):\\\rho_{A} \neq \sigma_{A}}} -\frac{1}{\beta-1} \log \widetilde{Q}_{\beta}(\rho_{AC} \| \sigma_{A}^{-1} \otimes \rho_{C}) & \text{if } \alpha \in (0,1) \\ \inf_{\substack{\rho_{A} \neq \sigma_{A}}} -\widetilde{D}_{\beta}(\rho_{AC} \| \sigma_{A}^{-1} \otimes \rho_{C}) & \text{if } \alpha \in [1,\infty). \end{cases}$$
(3.7)

- (f) Non-negativity: $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \geq 0$ for all $\alpha \in [0,\infty)$.
- (g) Upper bound: Let $\alpha \in [0,\infty)$ and $r_A \coloneqq \operatorname{rank}(\rho_A)$. Then $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \leq 2\log r_A$. Furthermore, if $\alpha \in [\frac{1}{2}, 2]$, then $I_{\alpha}^{\downarrow\downarrow}(A : B)_{\rho} = 2 \log r_A$ iff $\operatorname{spec}(\rho_A) \subseteq \{0, 1/r_A\}$ and $H(A|B)_{\rho} = -\log r_A.$ If $\alpha \in [0, \frac{1}{2})$ instead, then $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \leq \frac{1}{1-\alpha}H_{\infty}(A)_{\rho} \leq \frac{1}{1-\alpha}\log r_A < 2\log r_A$.
- (h) Existence of minimizers: Let $\alpha \in [0, \infty)$. Then

$$\emptyset \neq \operatorname*{arg\,min}_{(\sigma_A,\tau_B)\in\mathcal{S}(A)\times\mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) \subseteq \mathcal{S}_{\ll\rho_A}(A) \times \mathcal{S}_{\ll\rho_B}(B).$$
(3.8)

(i) Fixed-point property of partial minimizers: Let $\alpha \in (0, \infty)$. Let us define the following maps.

$$\mathcal{N}_{A \to B}: \quad \mathcal{S}_{\not{\perp}\rho_A}(A) \to \mathcal{S}_{\ll \rho_B}(B), \quad \sigma_A \mapsto \frac{(\mathrm{tr}_A[\rho_{AB}^{\alpha}\sigma_A^{1-\alpha}])^{\frac{1}{\alpha}}}{\mathrm{tr}[(\mathrm{tr}_A[\rho_{AB}^{\alpha}\sigma_A^{1-\alpha}])^{\frac{1}{\alpha}}]} \tag{3.9}$$

$$\mathcal{N}_{B\to A}: \quad \mathcal{S}_{\not{\perp}\rho_B}(B) \to \mathcal{S}_{\ll\rho_A}(A), \quad \tau_B \mapsto \frac{(\mathrm{tr}_B[\rho_{AB}^{\alpha}\tau_B^{1-\alpha}])^{\frac{1}{\alpha}}}{\mathrm{tr}[(\mathrm{tr}_B[\rho_{AB}^{\alpha}\tau_B^{1-\alpha}])^{\frac{1}{\alpha}}]} \tag{3.10}$$

$$\mathcal{N}_{A \to A}: \quad \mathcal{S}_{\not\perp \rho_A}(A) \to \mathcal{S}_{\ll \rho_A}(A), \quad \sigma_A \mapsto \mathcal{N}_{B \to A} \circ \mathcal{N}_{A \to B}(\sigma_A)$$
(3.11)

Furthermore, let

$$\mathcal{M}_{\alpha} \coloneqq \operatorname*{arg\,min}_{\sigma_{A} \in \mathcal{S}(A)} I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_{A}), \tag{3.12}$$

$$\mathcal{F}_{\alpha} \coloneqq \{ \sigma_A \in \mathcal{S}_{\ll \rho_A}(A) : \mathcal{N}_{A \to A}(\sigma_A) = \sigma_A \}.$$
(3.13)

Then $\emptyset \neq \mathcal{M}_{\alpha} \subseteq \mathcal{F}_{\alpha}$.

(j) Uniqueness of minimizer: Let $\alpha \in (\frac{1}{2}, 1]$. Then there exists $(\hat{\sigma}_A, \hat{\tau}_B) \in \mathcal{S}_{\ll \rho_A \ll}(A) \times \mathcal{S}_{\ll \rho_B \ll}(B)$ such that

$$\underset{(\sigma_A,\tau_B)\in\mathcal{S}(A)\times\mathcal{S}(B)}{\arg\min} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) = \{ (\hat{\sigma}_A, \hat{\tau}_B) \}.$$
(3.14)

(k) Fixed-point property of minimizers: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in (\frac{1}{2}, 2]$,

$$\mathcal{M}_{\alpha} \coloneqq \underset{\sigma_{A} \in \mathcal{S}(A)}{\arg\min} I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_{A}), \tag{3.15}$$

$$\mathcal{F}_{\alpha} \coloneqq \left\{ \sigma_{A} \in \mathcal{S}_{\ll \rho_{A} \ll}(A) : \sigma_{A} = \frac{\operatorname{tr}_{C}[\left(\sigma_{A}^{\frac{1-\alpha}{2}} \otimes \rho_{C}^{\frac{\alpha-1}{2}} \rho_{AC} \sigma_{A}^{\frac{1-\alpha}{2}} \otimes \rho_{C}^{\frac{\alpha-1}{2}}\right)^{\frac{1}{\alpha}}]}{\operatorname{tr}[\left(\sigma_{A}^{\frac{1-\alpha}{2}} \otimes \rho_{C}^{\frac{\alpha-1}{2}} \rho_{AC} \sigma_{A}^{\frac{1-\alpha}{2}} \otimes \rho_{C}^{\frac{\alpha-1}{2}}\right)^{\frac{1}{\alpha}}]} \right\}.$$
(3.16)

Then $\mathcal{M}_{\alpha} = \mathcal{F}_{\alpha}$.

(l) Asymptotic optimality of universal permutation invariant state: Let $\alpha \in [0, 2]$. Then

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \lim_{n \to \infty} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \omega_{A^{n}}^{n} \otimes \omega_{B^{n}}^{n}) = \lim_{n \to \infty} \inf_{\substack{\sigma_{A^{n}} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^{n}} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^{n}} \otimes \tau_{B^{n}}).$$
(3.17)

Moreover, if $\alpha \in [\frac{1}{2}, 2]$, then for any $n \in \mathbb{N}_{>0}$

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \inf_{\substack{\sigma_{A^{n}} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^{n}} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^{n}} \otimes \tau_{B^{n}}) = \inf_{\substack{\sigma_{A^{n}} \in \mathcal{S}(A^{n}), \\ \tau_{B^{n}} \in \mathcal{S}(B^{n})}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^{n}} \otimes \tau_{B^{n}}).$$
(3.18)

(m) Rényi order $\alpha \in \{0,1\}$: $I_1^{\downarrow\downarrow}(A:B)_{\rho} = I(A:B)_{\rho}$ and

$$I_0^{\downarrow\downarrow}(A:B)_{\rho} = \min_{\substack{|\sigma\rangle_A \in \operatorname{supp}(\rho_A), |\tau\rangle_B \in \operatorname{supp}(\rho_B):\\\langle\sigma|\sigma\rangle_A = 1, \langle\tau|\tau\rangle_B = 1}} D_0(\rho_{AB} |||\sigma\rangle\langle\sigma|_A \otimes |\tau\rangle\langle\tau|_B).$$
(3.19)

- (n) Monotonicity in α : If $\alpha, \beta \in [0, \infty)$ are such that $\alpha \leq \beta$, then $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \leq I_{\beta}^{\downarrow\downarrow}(A:B)_{\rho}$. (o) Continuity in α : The function $[0, \infty) \to [0, \infty), \alpha \mapsto I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ is continuous.

(p) Differentiability in α : The function $(\frac{1}{2}, 2) \to [0, \infty), \alpha \mapsto I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ is continuously differentiable. For any $\alpha \in (\frac{1}{2}, 2)$ and any fixed $(\sigma_A, \tau_B) \in \arg\min_{(\sigma'_A, \tau'_B) \in \mathcal{S}(A) \times \mathcal{S}(B)} D_\alpha(\rho_{AB} \| \sigma'_A \otimes \mathcal{S}(B))$ τ'_B), the derivative at α is

$$\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \frac{\partial}{\partial\alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B).$$
(3.20)

In particular, $\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}|_{\alpha=1} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B)|_{\alpha=1} = \frac{1}{2} V(A:B)_{\rho}.$ Moreover, $\frac{\partial}{\partial \alpha^+} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}|_{\alpha=1/2} = \lim_{\beta \to 1/2^+} \frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}|_{\alpha=\beta} \in [0,\infty).$ (q) Convexity in α : The function $[0,2] \to \mathbb{R}, \alpha \mapsto (\alpha-1)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ is convex.

- (r) Product states: If $\rho_{AB} = \rho_A \otimes \rho_B$, then $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = 0$ for all $\alpha \in [0,\infty)$. Conversely, for any $\alpha \in (0,\infty)$, if $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = 0$, then $\rho_{AB} = \rho_A \otimes \rho_B$.
- (s) AC-independent states: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. If $\rho_{AC} =$ $\rho_A \otimes \rho_C$, then for all $\alpha \in [0,\infty)$

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \begin{cases} \frac{1}{1-\alpha}H_{\infty}(A)_{\rho} & \text{if } \alpha \in [0,\frac{1}{2}]\\ 2H_{\frac{1}{2\alpha-1}}(A)_{\rho} & \text{if } \alpha \in (\frac{1}{2},\infty). \end{cases}$$
(3.21)

(t) Pure states: If there exists $|\rho\rangle_{AB} \in AB$ such that $\rho_{AB} = |\rho\rangle\langle\rho|_{AB}$, then for all $\alpha \in [0,\infty)$

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{|\rho\rangle\!\langle\rho|} = D_{\alpha}(|\rho\rangle\!\langle\rho|_{AB} \| \sigma_A \otimes \tau_B) = \begin{cases} \frac{1}{1-\alpha} H_{\infty}(A)_{\rho} & \text{if } \alpha \in [0, \frac{1}{2}]\\ 2H_{\frac{1}{2\alpha-1}}(A)_{\rho} & \text{if } \alpha \in (\frac{1}{2}, \infty), \end{cases}$$
(3.22)

where $\sigma_A \coloneqq \rho_A^{\frac{1}{2\alpha-1}} / \operatorname{tr}[\rho_A^{\frac{1}{2\alpha-1}}], \tau_B \coloneqq \rho_B^{\frac{1}{2\alpha-1}} / \operatorname{tr}[\rho_B^{\frac{1}{2\alpha-1}}]$ if $\alpha \in (\frac{1}{2}, \infty)$, and if $\alpha \in [0, \frac{1}{2}]$, then $|\sigma\rangle_A \in A$ is defined as a unit eigenvector of ρ_A corresponding to the largest eigenvalue of ρ_A , $\sigma_A \coloneqq |\sigma\rangle\!\langle\sigma|_A, \ |\tau\rangle_B \coloneqq \langle\sigma|_A |\rho\rangle_{AB} / \sqrt{\langle\sigma|_A \rho_A |\sigma\rangle_A}, \ and \ \tau_B \coloneqq |\tau\rangle\!\langle\tau|_B.$

(u) CC states: Let P_{XY} be the joint PMF of two random variables X, Y over $\mathcal{X} \coloneqq [d_A], \mathcal{Y} \coloneqq$ $[d_B]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x\in[d_A]}, \{|b_y\rangle_B\}_{y\in[d_B]}$ for A, B such that $\rho_{AB} =$ $\sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} P_{XY}(x, y) |a_x, b_y\rangle \langle a_x, b_y|_{AB}$, then for all $\alpha \in [0, \infty)$

$$I^{\downarrow\downarrow}_{\alpha}(A:B)_{\rho} = I^{\downarrow\downarrow}_{\alpha}(X:Y)_{P}.$$
(3.23)

(v) Copy-CC states: Let P_X be the PMF of a random variable X over $\mathcal{X} \coloneqq [\min(d_A, d_B)]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x\in[d_A]}, \{|b_y\rangle_B\}_{y\in[d_B]}$ for A, B such that $\rho_{AB} =$ $\sum_{x \in \mathcal{X}} P_X(x) |a_x, b_x\rangle \langle a_x, b_x |_{AB}$, then for all $\alpha \in [0, \infty)$

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\downarrow\downarrow}(X:Y)_{P} = D_{\alpha}(\rho_{AB} \| \sigma_{A} \otimes \tau_{B}) = \begin{cases} \frac{\alpha}{1-\alpha} H_{\infty}(A)_{\rho} & \text{if } \alpha \in [0, \frac{1}{2}] \\ H_{\frac{\alpha}{2\alpha-1}}(A)_{\rho} & \text{if } \alpha \in (\frac{1}{2}, \infty), \end{cases}$$

$$(3.24)$$

where
$$\sigma_A \coloneqq \rho_A^{\frac{\alpha}{2\alpha-1}} / \operatorname{tr}[\rho_A^{\frac{\alpha}{2\alpha-1}}], \tau_B \coloneqq \rho_B^{\frac{\alpha}{2\alpha-1}} / \operatorname{tr}[\rho_B^{\frac{\alpha}{2\alpha-1}}]$$
 if $\alpha \in (\frac{1}{2}, \infty)$, and if $\alpha \in [0, \frac{1}{2}]$, then we let $\hat{x} \in \operatorname{arg\,max}_{x \in \mathcal{X}} P_X(x)$ be arbitrary but fixed, and $\sigma_A \coloneqq |a_{\hat{x}}\rangle\langle a_{\hat{x}}|_A, \tau_B \coloneqq |b_{\hat{x}}\rangle\langle b_{\hat{x}}|_B$.

Remark 2 (Inequivalence of PRMIs). The non-minimized, the singly minimized, and the doubly minimized PRMI of a fixed quantum state ρ_{AB} are not necessarily the same for $\alpha \neq 1$. For illustrative purposes, two examples are provided in Figure 1 and Figure 2.

Remark 3 (Classical case). According to Theorem 7 (u), the doubly minimized PRMI of a CC state ρ_{AB} reduces to the doubly minimized RMI. The doubly minimized RMI can therefore be regarded as a specific instance of the doubly minimized PRMI. For more properties of the doubly minimized RMI, we direct the reader to [1, Theorem 1].



Figure 1. Comparison of PRMIs for a pure state. Suppose $d_A = 2$, $d_B = 2$, and let $\{|i\rangle_A\}_{i=0}^1, \{|i\rangle_B\}_{i=0}^1$ be orthonormal vectors in A, B. Let $\rho_{AB} \coloneqq |\rho\rangle\langle\rho|_{AB}$, where $|\rho\rangle_{AB} \coloneqq \sqrt{p}|0,0\rangle_{AB} + \sqrt{1-p}|1,1\rangle_{AB}$ and $p \coloneqq 0.2$. The solid lines depict the behavior of three PRMIs for ρ_{AB} , computed according to Proposition 3 (o), Proposition 4 (q), and Theorem 7 (t), respectively. For comparison, the values of certain Rényi entropies of $\rho_A = p|0\rangle\langle 0|_A + (1-p)|1\rangle\langle 1|_A$ are indicated by dashed lines. The plot shows that the three PRMIs differ from each other for all $\alpha \in [0, 1) \cup (1, \infty)$.



Figure 2. Comparison of PRMIs for a copy-CC state. Suppose $d_A = 2$, $d_B = 2$, and let $\{|i\rangle_A\}_{i=0}^1, \{|i\rangle_B\}_{i=0}^1$ be orthonormal vectors in A, B. Let $\rho_{AB} := p|0, 0\rangle\langle 0, 0|_{AB} + (1-p)|1, 1\rangle\langle 1, 1|_{AB}$, where p := 0.2. The solid lines depict the behavior of three PRMIs for ρ_{AB} , computed according to Proposition 3 (q), Proposition 4 (s), and Theorem 7 (v), respectively. For comparison, the values of certain Rényi entropies of $\rho_A = p|0\rangle\langle 0|_A + (1-p)|1\rangle\langle 1|_A$ are indicated by dashed lines. The plot shows that the three PRMIs differ from each other for all $\alpha \in [0, 1) \cup (1, \infty)$.

B. Operational interpretation from direct exponent

Correlation detection. Let ρ_{AB} be a bipartite quantum state that is correlated, i.e., $\rho_{AB} \neq \rho_A \otimes \rho_B$. How well can this quantum state be distinguished from any uncorrelated quantum state $\sigma_A \otimes \tau_B$? To make this question more precise, we will use the terminology for binary quantum state discrimination, as outlined above in Section 2 D. Consider the null hypothesis $H_0 \coloneqq \{\rho_{AB}\}$ and the alternative hypothesis $H_1 \coloneqq \{\sigma_A \otimes \tau_B : \sigma_A \in \mathcal{S}(A), \tau_B \in \mathcal{S}(B)\}$. Then the minimum type-I error when the type-II error is upper bounded by $\mu \in [0, \infty)$ is

$$\min_{\substack{T_{AB}\in\mathcal{L}(AB):\\0\leq T_{AB}\leq 1}} \{ \operatorname{tr}[\rho_{AB}(1-T_{AB})] : \max_{\substack{\sigma_{A}\in\mathcal{S}(A),\\\tau_{B}\in\mathcal{S}(B)}} \operatorname{tr}[\sigma_{A}\otimes\tau_{B}T_{AB}] \leq \mu \}.$$
(3.25)

(3.25) is the minimum probability with which one erroneously decides that the given quantum state is uncorrelated, under the constraint that the probability with which one erroneously decides that the given quantum state is ρ_{AB} is upper bounded by μ . The goal of this section is to show that the single-letter formula for the direct exponent in the third row of Table I, where the doubly minimized RMI occurs, can be generalized from the classical to the quantum setting by means of the doubly minimized PRMI. Accordingly, we are interested in any one of the following variants of correlation detection as described in (3.25). It should be noted that several variants of the problem are introduced, as all of them will be found to have the same direct exponent, and this exponent is determined by the doubly minimized PRMI analogously to the classical setting.

First, one may consider the i.i.d. (independent and identically distributed) version of (3.25), where $H_0^n \coloneqq \{\rho_{AB}^{\otimes n}\}$ and $H_1^n \coloneqq \{\sigma_A^{\otimes n} \otimes \tau_B^{\otimes n} : \sigma_A \in \mathcal{S}(A), \tau_B \in \mathcal{S}(B)\}$. The minimum type-I error is then

$$\hat{\alpha}_{n,\rho}^{\text{iid}}(\mu) \coloneqq \min_{\substack{T_{A^n B^n} \in \mathcal{L}(A^n B^n):\\ 0 \le T_{A^n B^n} \le 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \max_{\substack{\sigma_A \in \mathcal{S}(A),\\ \tau_B \in \mathcal{S}(B)}} \operatorname{tr}[\sigma_A^{\otimes n} \otimes \tau_B^{\otimes n} T_{A^n B^n}^n] \le \mu \},$$
(3.26)

which shall be defined for any $n \in \mathbb{N}_{>0}, \rho_{AB} \in \mathcal{S}(AB), \mu \in [0, \infty).$

Second, one may impose the i.i.d. assumption on the null hypothesis only, and take the alternative hypothesis to be given by all states that are uncorrelated between A^n and B^n , and permutation invariant on both A^n and B^n , i.e., $H_0^n \coloneqq \{\rho_{AB}^{\otimes n}\}$ and $H_1^n \coloneqq \{\sigma_{A^n} \otimes \tau_{B^n} : \sigma_{A^n} \in \mathcal{S}_{sym}(A^{\otimes n}), \tau_{B^n} \in \mathcal{S}_{sym}(B^{\otimes n})\}$. The minimum type-I error is then

$$\hat{\alpha}_{n,\rho}(\mu) \coloneqq \min_{\substack{T_{A^n B^n} \in \mathcal{L}(A^n B^n):\\ 0 \le T_{A^n B^n} \le 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \max_{\substack{\sigma_A n \in \mathcal{S}_{\operatorname{sym}}(A^{\otimes n}),\\ \tau_{B^n} \in \mathcal{S}_{\operatorname{sym}}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T_{A^n B^n}^n] \le \mu \}.$$
(3.27)

Third, the second option may be modified by imposing the permutation invariance constraint on A^n only, i.e., $H_0^n \coloneqq \{\rho_{AB}^{\otimes n}\}$ and $H_1^n \coloneqq \{\sigma_{A^n} \otimes \tau_{B^n} : \sigma_{A^n} \in \mathcal{S}_{sym}(A^{\otimes n}), \tau_{B^n} \in \mathcal{S}(B^n)\}$. The minimum type-I error then matches that of the second option,

$$\hat{\alpha}_{n,\rho}(\mu) = \min_{\substack{T_{A^n B^n} \in \mathcal{L}(A^n B^n):\\ 0 \le T_{A^n B^n} \le 1}} \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \max_{\substack{\sigma_{A^n} \in \mathcal{S}_{\operatorname{sym}}(A^{\otimes n}),\\ \tau_{B^n} \in \mathcal{S}(B^n)}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T_{A^n B^n}^n] \le \mu\}.$$
(3.28)

The proof of the equality in (3.28) is deferred to the appendix, see Lemma 16 (c). Given (3.28), it is natural to inquire whether the permutation invariance constraint on the alternative hypothesis may be omitted entirely, without compromising the desired connection between the direct exponent and the doubly minimized PRMI. This question will be answered in the negative in Remark 4.

Problem formulation. The problem we are interested in is to find a single-letter formula for the direct exponent of the quantum state discrimination problems associated with $\hat{\alpha}_{n,\rho}^{\text{iid}}$ and $\hat{\alpha}_{n,\rho}$ as defined in (3.26) and (3.27). This is accomplished in the following theorem. It shows that if the type-II rate R is sufficiently large but below the threshold given by $I(A:B)_{\rho}$, then the minimum type-I error decreases to 0 exponentially fast in n, and the corresponding exponent is determined by the family of the doubly minimized PRMIs of order $s \in (\frac{1}{2}, 1)$.

Theorem 8 (Direct exponent). Let $\rho_{AB} \in \mathcal{S}(AB)$ and let

$$R_{1/2} \coloneqq I_{1/2}^{\downarrow\downarrow}(A:B)_{\rho} - \frac{1}{4} \frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_{\rho} \big|_{s=1/2} \in [I_0^{\downarrow\downarrow}(A:B)_{\rho}, I_{1/2}^{\downarrow\downarrow}(A:B)_{\rho}].$$
(3.29)

Then, for any $R \in (R_{1/2}, \infty)$

$$\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}(e^{-nR}) = \sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R),$$
(3.30)

and the same is true if $\hat{\alpha}_{n,\rho}$ in (3.30) is replaced by $\hat{\alpha}_{n,\rho}^{\text{iid}}$.

Furthermore, for any $R \in (0, \infty)$, the right-hand side of (3.30) lies in $[0, \max(0, I(A : B)_{\rho} - R)]$, and it is strictly positive iff $R < I(A : B)_{\rho}$. The proof of Theorem 8 is given in Appendix F 2 and uses two lemmas that are established in advance in Appendix F 1. The proof of Theorem 8 is divided into two parts: a proof of achievability and a proof of optimality. The proof of achievability uses a quantum Neyman-Pearson test that compares $\rho_{AB}^{\otimes n}$ with $\omega_{A^n}^n \otimes \omega_{B^n}^n$, and leverages the asymptotic optimality of the universal permutation invariant state, which has been established in Theorem 7 (l). This proof method is an adapted version of an analogous proof of achievability for the minimized generalized PRMI [15, Section V.A]. The proof of optimality employs techniques for classical binary hypothesis testing from [9], and makes use of several properties of the doubly minimized PRMI, including Theorem 7 (j), (p), and (q).

The proof of Theorem 8 implies the following corollary, which can be viewed as an alternative formulation of Theorem 8.

Corollary 9 (Direct exponent). Let $\rho_{AB} \in \mathcal{S}(AB)$ and let

$$R: (1/2, 1] \to [0, I(A:B)_{\rho}], \quad s \mapsto I_s^{\downarrow\downarrow}(A:B)_{\rho} - s(1-s)\frac{\mathrm{d}}{\mathrm{d}s}I_s^{\downarrow\downarrow}(A:B)_{\rho}. \tag{3.31}$$

Then R is continuous and monotonically increasing. Let $R_{1/2} := \lim_{s \to 1/2^+} R(s)$, $R(1/2) := R_{1/2}$, $s_{1/2} := \max\{s \in [\frac{1}{2}, 1] : R(s) = R_{1/2}\}$, and $s_1 := \min\{s \in [\frac{1}{2}, 1] : R(s) = I(A : B)_{\rho}\}$. Then for any $s \in (s_{1/2}, s_1)$

$$\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}(e^{-nR(s)}) = \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_\rho - R(s)) = (1-s)^2 \frac{\mathrm{d}}{\mathrm{d}s} I_s^{\downarrow\downarrow}(A:B)_\rho.$$
(3.32)

Moreover, the same is true if $\hat{\alpha}_{n,\rho}$ in (3.32) is replaced by $\hat{\alpha}_{n,\rho}^{\text{iid}}$.

The proof of achievability for Theorem 8 immediately leads to the following corollary. The proof of Corollary 10 is given in Appendix F 3.

Corollary 10 (Asymptotic minimum type-I error). Let $\rho_{AB} \in \mathcal{S}(AB)$ and let $R \in (-\infty, I(A : B)_{\rho})$. Then $\lim_{n\to\infty} \hat{\alpha}_{n,\rho}(e^{-nR}) = 0$. Moreover, the same is true if $\hat{\alpha}_{n,\rho}$ is replaced by $\hat{\alpha}_{n,\rho}^{\text{iid}}$.

Remark 4 (Necessity of permutation invariance of alternative hypothesis). Consider the following variant of $\hat{\alpha}_{n,\rho}$ where the alternative hypothesis is constrained only by the independence of A^n and B^n .

$$\hat{\alpha}_{n,\rho}^{\text{ind}}(\mu) \coloneqq \min_{\substack{T_{A^n B^n} \in \mathcal{L}(A^n B^n):\\ 0 \le T_{A^n B^n} \le 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \max_{\substack{\sigma_{A^n} \in \mathcal{S}(A^n),\\ \tau_{B^n} \in \mathcal{S}(B^n)}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T_{A^n B^n}^n] \le \mu \}$$
(3.33)

In light of the equality in (3.28), it is natural to inquire whether Theorem 8 remains valid when $\hat{\alpha}_{n,\rho}$ is replaced by $\hat{\alpha}_{n,\rho}^{\text{ind}}$. This is not the case; an explicit counterexample where ρ_{AB} is a correlated CC state is provided in Appendix F 4.

ACKNOWLEDGMENTS

We thank Renato Renner for fruitful discussions and many comments that helped improve the presentation of our results. This work was supported by the Swiss National Science Foundation via grant No. 200021_188541 and the National Centre of Competence in Research SwissMAP, and the Quantum Center at ETH Zurich.

Appendix A: Proof for Section 2D

Proof. Let $n \in \mathbb{N}_{>0}, \mu \in [0, \infty)$. Consider the optimization problem that defines $\hat{\alpha}_n^{\mathbf{q}}(\mu)$. Let $T_{A^nB^n}^n \in \mathcal{L}(A^nB^n)$ be in the feasible set of this optimization problem, i.e., $0 \leq T_{A^nB^n}^n \leq 1$ and $\sup_{\sigma_{A^nB^n} \in H_1^{\mathbf{q},n}} \operatorname{tr}[\sigma_{A^nB^n}T_{A^nB^n}^n] \leq \mu$. Let

$$\hat{T}^{n}(x_{1}, y_{1}, \dots, x_{n}, y_{n}) \coloneqq \langle a_{x_{1}}, b_{y_{1}}, \dots, a_{x_{n}}, b_{y_{n}} | T^{n}_{A^{n}B^{n}} | a_{x_{1}}, b_{y_{1}}, \dots, a_{x_{n}}, b_{y_{n}} \rangle$$
(A.1)

for all $x_1, \ldots, x_n \in \mathcal{X}, y_1, \ldots, y_n \in \mathcal{Y}$, and let

$$\hat{T}^{n}_{A^{n}B^{n}} \coloneqq \sum_{\substack{x_{1},\dots,x_{n}\in\mathcal{X},\\y_{1},\dots,y_{n}\in\mathcal{Y}}} \hat{T}^{n}(x_{1},y_{1},\dots,x_{n},y_{n})|a_{x_{1}},b_{y_{1}},\dots,a_{x_{n}},b_{y_{n}}\rangle\langle a_{x_{1}},b_{y_{1}},\dots,a_{x_{n}},b_{y_{n}}|.$$
(A.2)

Since ρ_{AB} is a CC state, $\operatorname{tr}[\rho_{AB}^{\otimes n}(1-T_{A^{n}B^{n}}^{n})] = \operatorname{tr}[\rho_{AB}^{\otimes n}(1-\hat{T}_{A^{n}B^{n}}^{n})]$. By (2.26), we have

$$\sup_{\sigma_{A^n B^n} \in H_1^{\mathbf{q}, n}} \operatorname{tr}[\sigma_{A^n B^n} T^n_{A^n B^n}] \le \sup_{\sigma_{A^n B^n} \in H_1^{\mathbf{q}, n}} \operatorname{tr}[\sigma_{A^n B^n} T^n_{A^n B^n}] \le \mu.$$
(A.3)

Therefore,

$$\hat{\alpha}_{n}^{q}(\mu) = \inf_{\hat{T}^{n}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - \hat{T}_{A^{n}B^{n}}^{n})] : \sup_{\sigma_{A^{n}B^{n}} \in H_{1}^{q,n}} \operatorname{tr}[\sigma_{A^{n}B^{n}} \hat{T}_{A^{n}B^{n}}^{n}] \le \mu \},$$
(A.4)

where the minimization is over all functions $\hat{T}^n : (\mathcal{X} \times \mathcal{Y})^{\times n} \to [0, 1]$, and the expressions inside the brackets are evaluated for $\hat{T}^n_{A^n B^n}$ as in (A.2). Then,

$$\hat{\alpha}_{n}^{q}(\mu) = \inf_{\hat{T}^{n}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - \hat{T}_{A^{n}B^{n}}^{n})] : \\ \sup_{\substack{\sigma_{A^{n}B^{n}} \in H_{1}^{q,n} \\ y_{1}, \dots, y_{n} \in \mathcal{Y}}} \sum_{\substack{\chi_{1}, \dots, \chi_{n} \in \mathcal{X}, \\ y_{1}, \dots, y_{n} \in \mathcal{Y}}} Q_{X^{n}Y^{n}}^{\sigma}(x_{1}, y_{1}, \dots, x_{n}, y_{n}) \hat{T}^{n}(x_{1}, y_{1}, \dots, x_{n}, y_{n}) \leq \mu \}$$
(A.5)

$$=\hat{\alpha}_{n}^{c}(\mu). \tag{A.6}$$

(A.5) follows from (A.4). (A.6) follows from (2.27).

Appendix B: Properties of the generalized PRMIs

Proposition 11 (Non-minimized generalized PRMI). Let $\rho_{AB} \in \mathcal{S}(AB)$ and let $\sigma_A \in \mathcal{S}(A)$. Then all of the following hold.

- (a) Non-increase under local operations: $I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A) \geq I^{\uparrow}_{\alpha}(\mathcal{M} \otimes \mathcal{N}(\rho_{AB}) \| \mathcal{M}(\sigma_A))$ for any $\mathcal{M} \in \operatorname{CPTP}(A, A'), \mathcal{N} \in \operatorname{CPTP}(B, B')$ and all $\alpha \in [0, 2]$.
- (b) Invariance under local isometries: $I^{\uparrow}_{\alpha}(V \otimes W \rho_{AB} V^{\dagger} \otimes W^{\dagger} || V \sigma_A V^{\dagger}) = I^{\uparrow}_{\alpha}(\rho_{AB} || \sigma_A)$ for any isometries $V \in \mathcal{L}(A, A'), W \in \mathcal{L}(B, B')$ and all $\alpha \in [0, \infty)$.
- (c) Additivity: Let $\alpha \in [0,\infty)$ and $\rho'_{DE} \in \mathcal{S}(DE), \sigma'_D \in \mathcal{S}(D)$. If $(\alpha \in [0,1) \land \rho_A \not\perp \sigma_A \land \rho'_D \not\perp \sigma'_D) \lor (\rho_A \ll \sigma_A \land \rho'_D \ll \sigma'_D)$, then

$$I_{\alpha}^{\uparrow}(\rho_{AB} \otimes \rho_{DE}' \| \sigma_A \otimes \sigma_D') = I_{\alpha}^{\uparrow}(\rho_{AB} \| \sigma_A) + I_{\alpha}^{\uparrow}(\rho_{DE}' \| \sigma_D').$$
(B.1)

(d) Duality: [15] Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in [0,1) \cup (1,\infty)$ and $\beta \coloneqq 2 - \alpha$. If $(\alpha \in [0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then

$$I_{\alpha}^{\uparrow}(\rho_{AB} \| \sigma_A) = -\frac{1}{\beta - 1} \log Q_{\beta}(\rho_{AC} \| \sigma_A^{-1} \otimes \rho_C).$$
(B.2)

- (e) Non-negativity: Let $\alpha \in [0, \infty)$. Then $I_{\alpha}^{\uparrow}(\rho_{AB} \| \sigma_A) \in [0, \infty]$. Furthermore, $I_{\alpha}^{\uparrow}(\rho_{AB} \| \sigma_A)$ is finite iff $(\alpha \in [0, 1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$.
- (f) Rényi order $\alpha = 1$: $I_1^{\uparrow}(\rho_{AB} \| \sigma_A) = D(\rho_{AB} \| \sigma_A \otimes \rho_B).$
- (g) Monotonicity in α : If $\alpha, \beta \in [0, \infty)$ are such that $\alpha \leq \beta$, then $I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A) \leq I^{\uparrow}_{\beta}(\rho_{AB} \| \sigma_A)$.
- (h) Continuity in α : If $\rho_A \not\perp \sigma_A$, then the function $[0,1) \to [0,\infty), \alpha \mapsto I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuous. ous. If $\rho_A \ll \sigma_A$, then the function $[0,\infty) \to [0,\infty), \alpha \mapsto I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuous.
- (i) Differentiability in α : If $\rho_A \ll \sigma_A$, then all of the following hold. The function $(0,\infty) \to [0,\infty), \alpha \mapsto I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuously differentiable and the derivative at $\alpha \in (0,\infty)$ is

$$\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\uparrow}(\rho_{AB} \| \sigma_A) = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \rho_B).$$
(B.3)

In particular, $\frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\uparrow}(\rho_{AB} \| \sigma_A)|_{\alpha=1} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \rho_B)|_{\alpha=1} = \frac{1}{2} V(\rho_{AB} \| \sigma_A \otimes \rho_B).$

- (j) Convexity in α : If $\rho_A \not\perp \sigma_A$, then the function $[0,1) \to \mathbb{R}, \alpha \mapsto (\alpha-1)I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is convex. If $\rho_A \ll \sigma_A$, then the function $[0,\infty) \to \mathbb{R}, \alpha \mapsto (\alpha-1)I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is convex.
- (k) Product states: If $\rho_{AB} = \rho_A \otimes \rho_B$, then $I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A) = D_{\alpha}(\rho_A \| \sigma_A)$ and $I^{\uparrow}_{\alpha}(\rho_{AB} \| \rho_A) = 0$ for all $\alpha \in [0, \infty)$. Conversely, for any $\alpha \in (0, \infty)$, if $I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A) = 0$, then $\rho_{AB} = \rho_A \otimes \rho_B$ and $\sigma_A = \rho_A$.
- (l) AC-independent states: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in [0,1) \cup (1,\infty)$ and $\beta \coloneqq 2 \alpha$. If $\rho_{AC} = \rho_A \otimes \rho_C$ and $(\alpha \in [0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then $I^{\uparrow}_{\alpha}(\rho_{AB} \| \sigma_A) = -\frac{1}{\beta 1} \log Q_{\beta}(\rho_A \| \sigma_A^{-1})$. (m) Pure states: Let $\alpha \in [0,1) \cup (1,\infty)$ and $\beta \coloneqq 2 - \alpha$. If there exists $|\rho\rangle_{AB} \in AB$ such that $\rho_{AB} = -\frac{1}{\beta - 1} \log Q_{\beta}(\rho_A \| \sigma_A^{-1})$.
- (m) Pure states: Let $\alpha \in [0,1) \cup (1,\infty)$ and $\beta \coloneqq 2-\alpha$. If there exists $|\rho\rangle_{AB} \in AB$ such that $\rho_{AB} = |\rho\rangle\langle\rho|_{AB}$ and $(\alpha \in [0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then $I_{\alpha}^{\uparrow}(|\rho\rangle\langle\rho|_{AB} \|\sigma_A) = -\frac{1}{\beta-1} \log Q_{\beta}(\rho_A \|\sigma_A^{-1})$.

Proposition 12 (Minimized generalized PRMI). Let $\rho_{AB} \in \mathcal{S}(AB)$ and let $\sigma_A \in \mathcal{S}(A)$. Then all of the following hold.

- (a) Non-increase under local operations: $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) \geq I^{\downarrow}_{\alpha}(\mathcal{M} \otimes \mathcal{N}(\rho_{AB}) \| \mathcal{M}(\sigma_A))$ for any $\mathcal{M} \in \operatorname{CPTP}(A, A'), \mathcal{N} \in \operatorname{CPTP}(B, B')$ and all $\alpha \in [0, 2]$.
- (b) Invariance under local isometries: $I^{\downarrow}_{\alpha}(V \otimes W \rho_{AB} V^{\dagger} \otimes W^{\dagger} || V \sigma_A V^{\dagger}) = I^{\downarrow}_{\alpha}(\rho_{AB} || \sigma_A)$ for any isometries $V \in \mathcal{L}(A, A'), W \in \mathcal{L}(B, B')$ and all $\alpha \in [0, \infty)$.
- (c) Additivity: [15] Let $\alpha \in [0,\infty)$ and $\rho'_{DE} \in \mathcal{S}(DE), \sigma'_D \in \mathcal{S}(D)$. If $(\alpha \in [0,1) \land \rho_A \not\perp \sigma_A \land \rho'_D \not\perp \sigma'_D) \lor (\rho_A \ll \sigma_A \land \rho'_D \ll \sigma'_D)$, then

$$I_{\alpha}^{\downarrow}(\rho_{AB} \otimes \rho'_{DE} \| \sigma_A \otimes \sigma'_D) = I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) + I_{\alpha}^{\downarrow}(\rho'_{DE} \| \sigma'_D).$$
(B.4)

(d) Duality: [15] Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in (0,1) \cup (1,\infty)$ and $\beta \coloneqq \frac{1}{\alpha}$. If $(\alpha \in (0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = -\frac{1}{\beta - 1} \log \widetilde{Q}_{\beta}(\rho_{AC} \| \sigma_A^{-1} \otimes \rho_C).$$
(B.5)

- (e) Non-negativity: Let $\alpha \in [0, \infty)$. Then $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) \in [0, \infty]$. Furthermore, $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is finite iff $(\alpha \in [0, 1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$.
- (f) Existence and uniqueness of minimizer: [15] Let $\alpha \in [0, \infty)$. If $\alpha \neq 0 \land ((\alpha \in (0, 1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A)$, then $\hat{\tau}_B \coloneqq (\operatorname{tr}_A[\rho_{AB}^{\alpha}\sigma_A^{1-\alpha}])^{\frac{1}{\alpha}} / \operatorname{tr}[(\operatorname{tr}_A[\rho_{AB}^{\alpha}\sigma_A^{1-\alpha}])^{\frac{1}{\alpha}}] \in \mathcal{S}_{\ll \rho_B}(B)$ and

$$\underset{\tau_B \in \mathcal{S}(B)}{\arg\min} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) = \{ \hat{\tau}_B \}.$$
(B.6)

If $\alpha = 0 \land \rho_A \not\perp \sigma_A$, then $\emptyset \neq \operatorname{arg\,min}_{\tau_B \in \mathcal{S}(B)} D_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B) \subseteq \mathcal{S}_{\ll \rho_B}(B)$.

(g) Closed-form expression: [15] Let $\alpha \in (0,1) \cup (1,\infty)$. If $(\alpha \in (0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = \frac{1}{\alpha - 1} \log \| \operatorname{tr}_A[\rho_{AB}^{\alpha} \sigma_A^{1 - \alpha}] \|_{\frac{1}{\alpha}}.$$
 (B.7)

(h) Asymptotic optimality of universal permutation invariant state: Let $\alpha \in [0,2]$. If $(\alpha \in [0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = \lim_{n \to \infty} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \omega_{B^n}^n)$$
(B.8)

and for any $n \in \mathbb{N}_{>0}$

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = \inf_{\tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_{B^n}) = \inf_{\tau_{B^n} \in \mathcal{S}(B^n)} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_{B^n}).$$
(B.9)

(i) Rényi order $\alpha \in \{0,1\}$: $I_1^{\downarrow}(\rho_{AB} \| \sigma_A) = D(\rho_{AB} \| \sigma_A \otimes \rho_B)$. Furthermore, if $\rho_A \not\perp \sigma_A$, then

$$I_0^{\downarrow}(\rho_{AB} \| \sigma_A) = -\log \| \operatorname{tr}_A[\rho_{AB}^0 \sigma_A] \|_{\infty} = \min_{\substack{|\tau\rangle_B \in \operatorname{supp}(\rho_B):\\\langle \tau | \tau \rangle_B = 1}} D_0(\rho_{AB} \| \sigma_A \otimes |\tau\rangle \langle \tau |_B).$$
(B.10)

- (j) Monotonicity in α : If $\alpha, \beta \in [0, \infty)$ are such that $\alpha \leq \beta$, then $I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) \leq I_{\beta}^{\downarrow}(\rho_{AB} \| \sigma_A)$.
- (k) Continuity in α : If $\rho_A \not\perp \sigma_A$, then the function $[0,1) \rightarrow [0,\infty), \alpha \mapsto I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuous. ous. If $\rho_A \ll \sigma_A$, then the function $[0,\infty) \rightarrow [0,\infty), \alpha \mapsto I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuous.
- (l) Differentiability in α : If $\rho_A \ll \sigma_A$, then all of the following hold. The function $(0,2) \to [0,\infty), \alpha \mapsto I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuously differentiable. For any $\alpha \in (0,2)$ and any fixed $\tau_B \in \arg\min_{\tau'_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau'_B)$, the derivative at α is

$$\frac{\mathrm{d}}{\mathrm{d}\alpha}I^{\downarrow}_{\alpha}(\rho_{AB}\|\sigma_A) = \frac{\partial}{\partial\alpha}D_{\alpha}(\rho_{AB}\|\sigma_A\otimes\tau_B). \tag{B.11}$$

In particular, $\frac{\mathrm{d}}{\mathrm{d}\alpha}I_{\alpha}^{\downarrow}(\rho_{AB}\|\sigma_A)|_{\alpha=1} = \frac{\mathrm{d}}{\mathrm{d}\alpha}D_{\alpha}(\rho_{AB}\|\sigma_A\otimes\rho_B)|_{\alpha=1} = \frac{1}{2}V(\rho_{AB}\|\sigma_A\otimes\rho_B).$

- (m) Convexity in α : If $\rho_A \not\perp \sigma_A$, then the function $[0,1) \to \mathbb{R}, \alpha \mapsto (\alpha 1)I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is convex. If $\rho_A \ll \sigma_A$, then the function $[0,2] \to \mathbb{R}, \alpha \mapsto (\alpha - 1)I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is convex.
- (n) Product states: If $\rho_{AB} = \rho_A \otimes \rho_B$, then $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) = D_{\alpha}(\rho_A \| \sigma_A)$ and $I^{\downarrow}_{\alpha}(\rho_{AB} \| \rho_A) = 0$ for all $\alpha \in [0, \infty)$. Conversely, for any $\alpha \in (0, \infty)$, if $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) = 0$, then $\rho_{AB} = \rho_A \otimes \rho_B$ and $\sigma_A = \rho_A$.
- (o) AC-independent states: Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in (0,1) \cup (1,\infty)$ and $\beta := \frac{1}{\alpha}$. If $\rho_{AC} = \rho_A \otimes \rho_C$ and $(\alpha \in (0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) = -\frac{1}{\beta-1} \log \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1})$.
- (p) Pure states: Let $\alpha \in (0,1) \cup (1,\infty)$ and $\beta \coloneqq \frac{1}{\alpha}$. If there exists $|\rho\rangle_{AB} \in AB$ such that $\rho_{AB} = |\rho\rangle\langle\rho|_{AB}$ and $(\alpha \in (0,1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then $I^{\downarrow}_{\alpha}(|\rho\rangle\langle\rho|_{AB} \|\sigma_A) = -\frac{1}{\beta-1}\log \widetilde{Q}_{\beta}(\rho_A \|\sigma_A^{-1})$. (q) CC states: Let $\alpha \in [0,\infty)$. Let P_{XY} be the joint PMF of two random variables X, Y over
- (q) CC states: Let $\alpha \in [0, \infty)$. Let P_{XY} be the joint PMF of two random variables X, Y over $\mathcal{X} := [d_A], \mathcal{Y} := [d_B]$. If there exist orthonormal bases $\{|a_x\rangle_A\}_{x \in [d_A]}, \{|b_y\rangle_B\}_{y \in [d_B]}$ for A, B such that $\rho_{AB} = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} P_{XY}(x, y) |a_x, b_y\rangle\langle a_x, b_y|_{AB}$ and $(\alpha \in [0, 1) \land \rho_A \not\perp \sigma_A) \lor \rho_A \ll \sigma_A$, then

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = \min_{\substack{\tau_B \in \mathcal{S}(B): \\ \exists (t_y)_{y \in \mathcal{Y}} \in [0,1]^{\times |\mathcal{Y}|: \\ \tau_B = \sum_{y \in \mathcal{Y}} t_y | b_y \rangle \langle b_y |_B}} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B).$$
(B.12)

Appendix C: Proofs for Appendix B

1. Proof of Proposition 11

We prove the listed items not in alphabetical order, but in a different order.

Proof of (a), (b), (c), (e), (f), (g), (h), (i), (j), (k). These properties follow from the corresponding properties of the Petz divergence, see Proposition 2. In particular, (e) follows from the nonnegativity of the Petz divergence because $\rho_{AB} \not\perp \sigma_A \otimes \rho_B$ iff $\rho_A \not\perp \sigma_A$, and $\rho_{AB} \ll \sigma_A \otimes \rho_B$ iff $\rho_A \ll \sigma_A$. (k) follows from the additivity and positive definiteness of the Petz divergence.

Proof of (d). Duality has been proved in [15, Lemma 6] under the assumption that $(\alpha \in [0, 1) \cup (1, 2]) \land \rho_A \ll \sigma_A$. However, as can be easily verified, their proof still works under the slightly weaker conditions specified in (d).

Proof of (l), (m). The assertion in (l) follows from duality (d), and (m) follows from (l). \Box

2. Proof of Proposition 12

Proof of (a), (e), (j), (n). These properties follow from the corresponding properties of the Petz divergence, see Proposition 2. In particular, (n) follows from the additivity and positive definiteness of the Petz divergence.

Proof of (i).
$$I_1^{\downarrow}(\rho_{AB} \| \sigma_A) = D(\rho_{AB} \| \sigma_A \otimes \rho_B)$$
 follows from (2.15).
Now, suppose $\rho_A \not\perp \sigma_A$. Then

$$\exp(-I_0^{\downarrow}(\rho_{AB} \| \sigma_A)) = \max_{\tau_B \in \mathcal{S}(B)} \operatorname{tr}[\rho_{AB}^0 \sigma_A \otimes \tau_B]$$
(C.1)

$$= \max_{\substack{|\tau\rangle_B \in \operatorname{supp}(\rho_B):\\\langle\tau|\tau\rangle_B = 1}} \operatorname{tr}[\rho_{AB}^0 \sigma_A \otimes |\tau\rangle\langle\tau|_B] = \|\operatorname{tr}_A[\rho_{AB}^0 \sigma_A]\|_{\infty}.$$
(C.2)

Proof of (f). Let
$$\alpha \in [0, \infty)$$
.

Case 1: $\alpha \neq 0$. For this case, the assertion has been proved in [15, Eq. (3.10)] under the assumption $\alpha \in (0, \infty) \land \rho_A \ll \sigma_A$ by means of a quantum Sibson identity [15, Eq. (B10)] (see also [51, Proposition 2]). However, as can be easily verified, their proof still works under the slightly weaker conditions specified in (f) because the quantum Sibson identity still applies.

Case 2: $\alpha = 0$. For this case, the assertion follows from (i).

Proof of (g). The closed-form expression follows from the explicit expression for the minimizer in (f). \Box

Proof of (b). Let $\alpha \in [0,\infty)$. Let $\hat{\tau}_B \in \mathcal{S}(B)$ be such that $I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B)$. Then

$$I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) = D_{\alpha}(V \otimes W \rho_{AB} V^{\dagger} \otimes W^{\dagger} \| V \sigma_A V^{\dagger} \otimes W \hat{\tau}_B W^{\dagger})$$
(C.3)

$$\geq I^{\downarrow}_{\alpha}(V \otimes W \rho_{AB} V^{\dagger} \otimes W^{\dagger} \| V \sigma_A V^{\dagger}) \tag{C.4}$$

$$= I_{\alpha}^{\downarrow}(W\rho_{AB}W^{\dagger} \| \sigma_A) \tag{C.5}$$

$$= \inf_{\substack{\tau_{B'} \in \mathcal{S}(B'):\\\tau_{B'} \ll W \rho_B W^{\dagger}}} D_{\alpha}(W \rho_{AB} W^{\dagger} \| \sigma_A \otimes \tau_{B'})$$
(C.6)

$$\geq \inf_{\tau_B \in \mathcal{S}(B)} D_{\alpha} (W \rho_{AB} W^{\dagger} \| \sigma_A \otimes W \tau_B W^{\dagger})$$
(C.7)

$$= I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A). \tag{C.8}$$

(C.3), (C.5) and (C.8) follow from the isometric invariance of the Petz divergence. (C.6) follows from (f). $\hfill \Box$

Proof of (c). Let $\alpha \in [0, \infty)$. Case 1: $\alpha \in (0, 1) \cup (1, \infty)$. Then additivity follows from the closed-form expression (g). Case 2: $\alpha \in \{0, 1\}$. Then additivity follows from (i).

Proof of (d). Duality has been proved in [15, Lemma 6] under the assumption that $\rho_A \ll \sigma_A$. However, as can be easily verified, their proof still works under the slightly weaker conditions specified in (d) due to (g).

Proof of (h). Let $\alpha \in [0, 2]$. Then

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = \inf_{\tau_B \in \mathcal{S}(B)} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_B^{\otimes n})$$
(C.9a)

$$\geq \inf_{\tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})} \frac{1}{n} D_\alpha(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_{B^n})$$
(C.9b)

$$\geq \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \omega_{B^n}^n) - \frac{\log g_{n,d_B}}{n}$$
(C.9c)

$$\geq \frac{1}{n} I_{\alpha}^{\downarrow}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n}) - \frac{\log g_{n,d_B}}{n} = I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) - \frac{\log g_{n,d_B}}{n}.$$
 (C.9d)

(C.9a) follows from the additivity of the Petz divergence. (C.9c) follows from Proposition 1 (b). (C.9d) follows from additivity (c). The assertion in (B.8) follows from (C.9) by taking the limit $n \to \infty$ and using Proposition 1 (b).

It remains to prove the assertion in (B.9). For any $n \in \mathbb{N}_{>0}$

$$I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) \ge \inf_{\tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_{B^n})$$
(C.10)

$$\geq \inf_{\tau_{B^n} \in \mathcal{S}(B^n)} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_{B^n}) = \frac{1}{n} I_{\alpha}^{\downarrow}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n}) = I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A).$$
(C.11)

(C.10) follows from (C.9b). (C.11) follows from additivity (c).

Proof of (k). If $\rho_A \not\perp \sigma_A$, then continuity on [0, 1) follows from the continuity in α of the Petz divergence.

Now, suppose $\rho_A \ll \sigma_A$. Then continuity on [0, 1) and on $[1, \infty)$ follows from the continuity in α of the Petz divergence. It remains to prove left-continuity at $\alpha = 1$. For any $n \in \mathbb{N}_{>0}$

$$\frac{1}{n}D_1(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \omega_{B^n}^n) - \frac{\log g_{n,d_B}}{n} \le \lim_{\alpha \to 1^-} I_\alpha^{\downarrow}(\rho_{AB} \| \sigma_A) \le I_1^{\downarrow}(\rho_{AB} \| \sigma_A).$$
(C.12)

The first inequality in (C.12) follows from (C.9). The second inequality in (C.12) follows from monotonicity (j). By Proposition 1 (b), the second term on the left-hand side of (C.12) vanishes in the limit $n \to \infty$. Thus,

$$I_{1}^{\downarrow}(\rho_{AB}\|\sigma_{A}) = \lim_{n \to \infty} \frac{1}{n} D_{1}(\rho_{AB}^{\otimes n} \| \sigma_{A}^{\otimes n} \otimes \omega_{B^{n}}^{n}) \le \lim_{\alpha \to 1^{-}} I_{\alpha}^{\downarrow}(\rho_{AB}\|\sigma_{A}) \le I_{1}^{\downarrow}(\rho_{AB}\|\sigma_{A}),$$
(C.13)

where the first equality in (C.13) follows from (h). Hence, $\lim_{\alpha \to 1^{-}} I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) = I_{1}^{\downarrow}(\rho_{AB} \| \sigma_A)$. \Box

Proof of (m). Convexity is inherited from the Petz divergence because, according to (B.8) in (h), $(\alpha - 1)I^{\downarrow}_{\alpha}(\rho_{AB} || \sigma_A)$ is the pointwise limit of a sequence of functions that are convex in α .

Proof of (l). We will first prove differentiability on $\alpha \in (0,1) \cup (1,2)$. By the closed-form expression (g), $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is differentiable in α on $\alpha \in (0,1) \cup (1,2)$. In particular, the right and left derivatives of this function at $\alpha \in (0,1) \cup (1,2)$ coincide,

$$\frac{\mathrm{d}}{\mathrm{d}\alpha}I^{\downarrow}_{\alpha}(\rho_{AB}\|\sigma_A) = \frac{\partial}{\partial\alpha^+}I^{\downarrow}_{\alpha}(\rho_{AB}\|\sigma_A) = \frac{\partial}{\partial\alpha^-}I^{\downarrow}_{\alpha}(\rho_{AB}\|\sigma_A).$$
(C.14)

Let $\alpha \in (0,1) \cup (1,2)$ and let $\tau_B \in \arg \min_{\tau'_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau'_B)$ be fixed. Then the right derivative of $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ at α is upper bounded by its left derivative at α because

$$\frac{\partial}{\partial \alpha^{+}} I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_{A}) = \lim_{\varepsilon \to 0^{+}} \frac{1}{\varepsilon} (I^{\downarrow}_{\alpha + \varepsilon}(\rho_{AB} \| \sigma_{A}) - I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_{A}))$$
(C.15)

$$\leq \lim_{\varepsilon \to 0^+} \frac{1}{\varepsilon} (D_{\alpha+\varepsilon}(\rho_{AB} \| \sigma_A \otimes \tau_B) - D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B))$$
(C.16)

$$= \frac{\partial}{\partial \alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) \tag{C.17}$$

$$= \lim_{\varepsilon \to 0^{-}} \frac{1}{\varepsilon} (D_{\alpha + \varepsilon}(\rho_{AB} \| \sigma_A \otimes \tau_B) - D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B))$$
(C.18)

$$\leq \lim_{\varepsilon \to 0^{-}} \frac{1}{\varepsilon} (I_{\alpha+\varepsilon}^{\downarrow}(\rho_{AB} \| \sigma_A) - I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A)) = \frac{\partial}{\partial \alpha^{-}} I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A).$$
(C.19)

(C.17) and (C.18) follow from the differentiability in α of the Petz divergence, see Proposition 2. By (C.14), all inequalities must be saturated. This proves (B.11) for all $\alpha \in (0, 1) \cup (1, 2)$.

We will now prove that $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is *continuously* differentiable on $\alpha \in (0,1) \cup (1,2)$. Let $g(\alpha) := (\alpha - 1)I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ for all $\alpha \in (0,1) \cup (1,2)$. By the product rule, g is differentiable on $\alpha \in (0,1) \cup (1,2)$. By (m), g is convex, so the differentiability of g implies its continuous differentiability. By the product rule, this implies that also $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ is continuously differentiable on $\alpha \in (0,1) \cup (1,2)$.

Next, we will prove differentiability of $I^{\downarrow}_{\alpha}(\rho_{AB} || \sigma_A)$ at $\alpha = 1$. The combination of (2.15) and a quantum Sibson identity [15, Eq. (B10)] implies that the limits

$$\lim_{\beta \to 1^{-}} \frac{\mathrm{d}}{\mathrm{d}\alpha} I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) \big|_{\alpha = \beta} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \rho_B) \big|_{\alpha = 1},$$
(C.20a)

$$\lim_{\beta \to 1^+} \frac{\mathrm{d}}{\mathrm{d}\alpha} I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) \big|_{\alpha = \beta} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \rho_B) \big|_{\alpha = 1}$$
(C.20b)

exist. Therefore, they are identical to the left and right derivative of $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ at $\alpha = 1$, respectively. By (C.20), the left and right derivative of $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ at $\alpha = 1$ coincide. This proves differentiability at $\alpha = 1$. The *continuous* differentiability of $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A)$ at $\alpha = 1$ follows from the quantum Sibson identity [15, Eq. (B10)].

Proof of (o), (p). The assertion in (o) follows from duality (d), and (p) follows from (o). \Box

Proof of (q). Let $\alpha \in [0, \infty)$.

Case 1: $\alpha \in (0, \infty)$. By (f), the unique minimizer is then

$$\hat{\tau}_B = \text{const.} \ (\text{tr}_A[\rho_{AB}^{\alpha}\sigma_A^{1-\alpha}])^{\frac{1}{\alpha}} = \text{const.} \ \sum_{y \in \mathcal{Y}} \left(\sum_{x \in \mathcal{X}} P_{XY}(x,y)^{\alpha} \langle a_x |_A \sigma_A^{1-\alpha} | a_x \rangle_A\right)^{\frac{1}{\alpha}} |b_y\rangle \langle b_y |_B, \ (C.21)$$

so $\hat{\tau}_B$ has the desired form.

Case 2: $\alpha = 0$. For all $y \in \mathcal{Y}$, let $c_y \coloneqq \sum_{x \in \mathcal{X}: P_{XY}(x,y) \neq 0} \langle a_x |_A \sigma_A | a_x \rangle_A$. Let $\hat{y} \in \arg \max_{y \in \mathcal{Y}} c_y$ and let $|\hat{\tau}\rangle_B \coloneqq |b_{\hat{y}}\rangle_B$. By (i),

$$\exp(-I_0^{\downarrow}(\rho_{AB} \| \sigma_A)) = \left\| \sum_{y \in \mathcal{Y}} c_y | b_y \rangle \langle b_y |_B \right\|_{\infty} = \max_{y \in \mathcal{Y}} c_y = c_{\hat{y}} = \exp(-D_0(\rho_{AB} \| \sigma_A \otimes |\hat{\tau}\rangle \langle \hat{\tau} |_B)).$$
(C.22)

Therefore, $\hat{\tau}_B \coloneqq |\hat{\tau}\rangle\langle\hat{\tau}|_B = |b_{\hat{y}}\rangle\langle b_{\hat{y}}|_B$ is a minimizer that has the desired form.

Appendix D: Proofs for Section 2 E

1. Proof of Proposition 3

Proof of (a). This assertion follows from the symmetry of the definition of the non-minimized PRMI in (2.30) with respect to A and B.

Proof of (b), (c), (d), (e), (f), (h), (i), (j), (k), (l), (m). Since $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow}(\rho_{AB} \| \rho_A)$, these properties follow from the corresponding properties of the non-minimized generalized PRMI, see Proposition 11.

Proof of (n). Let $\alpha \in [0, \infty)$.

Case 1: $\alpha \in [0,1) \cup (1,\infty)$. Let $\beta \coloneqq 2 - \alpha$. By Proposition 11 (l),

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow}(\rho_{AB} \| \rho_A) = -\frac{1}{\beta - 1} \log Q_{\beta}(\rho_A \| \rho_A^{-1}) = 2H_{2\beta - 1}(A)_{\rho} = 2H_{3-2\alpha}(A)_{\rho}.$$
(D.1)

Case 2: $\alpha = 1$. Then the assertion follows from case 1 by continuity in α (j).

Proof of (o). This assertion follows from (n).

Proof of (g). Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$. Let $\alpha \in [0, \frac{3}{2}]$. By (b) and the expression for pure states (o),

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} \le I_{\alpha}^{\uparrow\uparrow}(A:BC)_{|\rho\rangle\langle\rho|} = 2H_{3-2\alpha}(A)_{\rho} \le 2H_0(A)_{\rho} = 2\log r_A.$$
(D.2)

Let now $\alpha \in [0, \frac{3}{2})$ instead.

First, suppose spec $(\rho_A) \subseteq \{0, 1/r_A\}$ and $H(A|B)_{\rho} = -\log r_A$. Then $\rho_A = \rho_A^0/r_A$ and $\rho_{AC} = \rho_A \otimes \rho_C$. By (n), $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = 2H_{3-2\alpha}(A)_{\rho} = 2\log r_A$.

Now, suppose $I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = 2 \log r_A$ instead. Then the inequalities in (D.2) must be saturated. Hence, $H_{3-2\alpha}(A)_{\rho} = \log r_A$. Since $3-2\alpha > 0$, it follows that $\operatorname{spec}(\rho_A) \subseteq \{0, 1/r_A\}$. By duality (e),

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = -D_{2-\alpha}(\rho_{AC} \| \rho_A^{-1} \otimes \rho_C) = 2\log r_A - D_{2-\alpha}(\rho_{AC} \| \rho_A \otimes \rho_C) \le 2\log r_A, \quad (D.3)$$

where the last inequality follows from the non-negativity of the Petz divergence. Since $I_{\alpha}^{\uparrow\uparrow}(A : B)_{\rho} = 2 \log r_A$, the inequality in (D.3) must be saturated, so $D_{2-\alpha}(\rho_{AC} || \rho_A \otimes \rho_C) = 0$. By the positive definiteness of the Petz divergence, $\rho_{AC} = \rho_A \otimes \rho_C$. Therefore, $H(A|B)_{\rho} = -H(A|C)_{\rho} = -H(A)_{\rho} = -\log r_A$.

Proof of (p). Let $\alpha \in [0, \infty)$.

Case 1: $\alpha \in (0, 1) \cup (1, \infty)$. Then

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = \frac{1}{\alpha - 1} \log \sum_{\substack{x \in \mathcal{X}, \\ y \in \mathcal{Y}}} P_{XY}(x, y)^{\alpha} P_X(x)^{1 - \alpha} P_Y(y)^{1 - \alpha} = I_{\alpha}^{\uparrow\uparrow}(X:Y)_P.$$
(D.4)

Case 2: $\alpha \in \{0, 1\}$. Then the assertion follows from case 1 by continuity in α (j).

Proof of (q). Let $\alpha \in [0, \infty)$.

Case 1: $\alpha \in (0,1) \cup (1,\infty)$. Let Y be a random variable over $\mathcal{Y} := \mathcal{X}$ and let $P_{XY}(x,y) :=$ $P_X(x)\delta_{x,y}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$. Then $\rho_{AB} = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} P_{XY}(x,y) |a_x, b_y|_{AB}$. By the expression for CC states (p),

$$I_{\alpha}^{\uparrow\uparrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\uparrow}(X:Y)_{P} = \frac{1}{\alpha - 1} \log \sum_{\substack{x \in \mathcal{X}, \\ y \in \mathcal{Y}}} P_{XY}(x,y)^{\alpha} P_{X}(x)^{1 - \alpha} P_{Y}(y)^{1 - \alpha}$$
(D.5)

$$= \frac{1}{\alpha - 1} \log \sum_{x \in \mathcal{X}} P_X(x)^{\alpha} P_X(x)^{1 - \alpha} P_X(x)^{1 - \alpha} = \frac{1}{\alpha - 1} \log \operatorname{tr}[\rho_A^{2 - \alpha}] = H_{2 - \alpha}(A)_{\rho}.$$
 (D.6)

Case 2: $\alpha \in \{0,1\}$. Then the assertion follows from case 1 by continuity in α (j).

2. Proof of Proposition 4

Proof of (a), (b), (c), (d), (e), (g), (h), (i), (j), (k), (l), (m), (n), (o), (r). Since $I_{\alpha}^{\uparrow\downarrow}(A : B)_{\rho} =$ $I^{\downarrow}_{\alpha}(\rho_{AB} \| \rho_A)$, these properties follow from the corresponding properties of the minimized generalized PRMI, see Proposition 12.

Proof of (p). Let $\alpha \in (0, \infty)$.

Case 1: $\alpha \in (0,1) \cup (1,\infty)$. Let $\beta := \frac{1}{\alpha}$. By Proposition 12 (o),

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\downarrow}(\rho_{AB} \| \rho_A) = -\frac{1}{\beta - 1} \log \widetilde{Q}_{\beta}(\rho_A \| \rho_A^{-1}) = 2H_{2\beta - 1}(A)_{\rho} = 2H_{\frac{2-\alpha}{\alpha}}(A)_{\rho}.$$
(D.7)

Case 2: $\alpha = 1$. Then the assertion follows from case 1 by continuity in α (l).

Proof of (q). $I_{\alpha}^{\uparrow\downarrow}(A:B)_{|\rho\rangle\langle\rho|} = 2H_{\frac{2-\alpha}{\alpha}}(A)_{\rho}$ follows from (p). The assertion regarding τ_B can be verified by inserting τ_B into (2.45).

Proof of (f). Let $|\rho\rangle_{ABC} \in ABC$ be such that $\rho_{AB} = \text{tr}_C[|\rho\rangle\langle\rho|_{ABC}]$. Let $\alpha \in [0,\infty)$. Then $\frac{1}{\alpha} \in (0, \infty]$. By duality (d),

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = -\widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_{A}^{-1} \otimes \rho_{C}) = -\widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_{A}^{-1} / \operatorname{tr}[\rho_{A}^{-1}] \otimes \rho_{C}) + \log \operatorname{tr}[\rho_{A}^{-1}] \leq 2H_{-1}(A)_{\rho},$$
(D.8)

where the last inequality follows from the non-negativity of the sandwiched divergence.

Let now $\alpha \in [0,2]$. Let $\gamma := \frac{2-\alpha}{\alpha} \in [0,\infty]$. By (a) and the expression for pure states (q),

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} \le I_{\alpha}^{\uparrow\downarrow}(A:BC)_{|\rho\rangle\!\langle\rho|} = 2H_{\gamma}(A)_{\rho} \le 2H_0(A)_{\rho} = 2\log r_A.$$
(D.9)

Let now $\alpha \in [0,2)$. Let $\gamma \coloneqq \frac{2-\alpha}{\alpha} \in (0,\infty]$. First, suppose $\operatorname{spec}(\rho_A) \subseteq \{0,1/r_A\}$ and $H(A|B)_{\rho} = -\log r_A$. Then $\rho_A = \rho_A^0/r_A$ and $\rho_{AC} = \frac{1}{2} \log r_A$. $\rho_A \otimes \rho_C. \text{ By (p)}, I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 2H_{\gamma}(A)_{\rho} = 2\log r_A.$

Now, suppose $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 2\log r_A$ instead. Then the inequalities in (D.9) must be saturated. Thus, $H_{\gamma}(A)_{\rho} = \log r_A$. Since $\gamma > 0$, it follows that $\operatorname{spec}(\rho_A) \subseteq \{0, 1/r_A\}$. By duality (d),

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = -\widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_A^{-1} \otimes \rho_C) = 2\log r_A - \widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_A \otimes \rho_C) \le 2\log r_A,$$
(D.10)

where the last inequality follows from the non-negativity of the sandwiched divergence. Since $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = 2\log r_A$, the inequality in (D.10) must be saturated, so $\widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_A \otimes \rho_C) = 0$. By the positive definiteness of the sandwiched divergence, $\rho_{AC} = \rho_A \otimes \rho_C^{\alpha}$. Therefore, $H(A|B)_{\rho} =$ $-H(A|C)_{\rho} = -H(A)_{\rho} = -\log r_A.$ Proof of (s). Let $\alpha \in (0, \infty)$.

Case 1: $\alpha \in (0,1) \cup (1,\infty)$. Let Y be a random variable over $\mathcal{Y} := \mathcal{X}$ and let $P_{XY}(x,y) := P_X(x)\delta_{x,y}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$. Then $\rho_{AB} = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} P_{XY}(x,y) |a_x, b_y\rangle \langle a_x, b_y|_{AB}$. By the expression for CC states (r) and the closed-form expression (h),

$$I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\uparrow\downarrow}(X:Y)_{P} = \frac{1}{\alpha - 1} \log \left\| \sum_{\substack{x \in \mathcal{X}, \\ y \in \mathcal{Y}}} P_{XY}(x,y)^{\alpha} P_{X}(x)^{1 - \alpha} |b_{y}\rangle \langle b_{y}|_{B} \right\|_{\frac{1}{\alpha}}$$
(D.11)

$$= \frac{\alpha}{\alpha - 1} \log \sum_{y \in \mathcal{Y}} (P_X(y)^{\alpha} P_X(y)^{1 - \alpha})^{\frac{1}{\alpha}} = \frac{\alpha}{\alpha - 1} \log \operatorname{tr}[\rho_A^{\frac{1}{\alpha}}] = H_{\frac{1}{\alpha}}(A)_{\rho}.$$
 (D.12)

The assertion regarding τ_B can be verified by inserting τ_B into (2.47).

Case 2: $\alpha = 1$. Then $I_{\alpha}^{\uparrow\downarrow}(A:B)_{\rho} = H_{\frac{1}{\alpha}}(A)_{\rho}$ follows from case 1 by continuity in α (l). The assertion regarding τ_B can be verified by inserting τ_B into (2.47).

Appendix E: Proofs for Section 3 A

1. Proof of Lemma 5

Proof.

$$\sqrt{X_A} \otimes \sqrt{Y_B} + \sqrt{X'_A} \otimes \sqrt{Y'_B} = (X_A \otimes 1_B) \# (1_A \otimes Y_B) + (X'_A \otimes 1_B) \# (1_A \otimes Y'_B)$$
(E.1)

$$\leq ((X_A \otimes 1_B) + (X'_A \otimes 1_B)) \# ((1_A \otimes Y_B) + (1_A \otimes Y'_B))$$
 (E.2)

$$=\sqrt{X_A + X'_A \otimes \sqrt{Y_B + Y'_B}} \tag{E.3}$$

The inequality follows from the subadditivity of the geometric operator mean. \Box

2. Proof of Theorem 6

Lemma 13 (Saturation of operator inequality from saturation of trace inequality). Let $X, Y, Z \in \mathcal{L}(A)$ be positive semidefinite and such that $X \leq Y$. Then all of the following hold.

(a) $\operatorname{tr}[XZ] \leq \operatorname{tr}[YZ].$ (b) If $\operatorname{tr}[XZ] = \operatorname{tr}[YZ]$ and $Y \ll Z$, then X = Y.

Proof. By spectral decomposition, $Z = \sum_{\lambda \in \operatorname{spec}(Z)} \lambda P_{\lambda}$, where $P_{\lambda} \in \mathcal{L}(A)$ denotes the orthogonal projection onto the eigenspace corresponding to λ .

We will now prove (a).

$$\operatorname{tr}[YZ] - \operatorname{tr}[XZ] = \operatorname{tr}[(Y-X)Z] = \sum_{\lambda \in \operatorname{spec}(Z)} \lambda \operatorname{tr}[(Y-X)P_{\lambda}] = \sum_{\substack{\lambda \in \operatorname{spec}(Z):\\\lambda \neq 0}} \lambda \operatorname{tr}[(Y-X)P_{\lambda}] \ge 0$$
(E.4)

We will now prove (b). Suppose $\operatorname{tr}[XZ] = \operatorname{tr}[YZ]$. Then, (E.4) implies that $\operatorname{tr}[(Y - X)P_{\lambda}] = 0$ for all $\lambda \in \operatorname{spec}(Z)$ such that $\lambda \neq 0$. Let us denote the pinching map with respect to Z as $\mathcal{P}_Z : \mathcal{L}(A) \to \mathcal{L}(A), K \mapsto \sum_{\lambda \in \operatorname{spec}(Z)} P_{\lambda}KP_{\lambda}$. Since the pinching map is trace-preserving,

$$\operatorname{tr}[Y-X] = \operatorname{tr}[\mathcal{P}_Z(Y-X)] = \sum_{\lambda \in \operatorname{spec}(Z)} \operatorname{tr}[(Y-X)P_{\lambda}] = \sum_{\substack{\lambda \in \operatorname{spec}(Z):\\\lambda \neq 0}} \operatorname{tr}[(Y-X)P_{\lambda}] = 0, \quad (E.5)$$

where we used that $X \leq Y$ and $Y \ll Z$ implies $(Y - X) \ll Z$. By (E.5), tr[Y - X] = 0. Because $(Y - X) \geq 0$, the trace of (Y - X) vanishes iff the operator itself vanishes. Therefore, X = Y.

Lemma 14 (Strict concavity for states). Let $\sigma, \sigma' \in \mathcal{S}(A), p \in (1, \infty)$, and let $\lambda, \lambda' \in (0, 1)$ be such that $\lambda + \lambda' = 1$. Then

$$\lambda \sigma^{\frac{1}{p}} + \lambda' \sigma'^{\frac{1}{p}} \le (\lambda \sigma + \lambda' \sigma')^{\frac{1}{p}}.$$
(E.6)

Moreover, if (E.6) holds with equality, then $\sigma = \sigma'$.

Proof. The inequality in (E.6) follows from the operator concavity of $X \mapsto X^{\frac{1}{p}}$ since $\frac{1}{p} \in (0, 1)$.

Now, suppose (E.6) holds with equality. Let $X \coloneqq \sigma^{\frac{1}{p}}$ and $X' \coloneqq {\sigma'}^{\frac{1}{p}}$. By the saturation of the inequality (E.6), $\lambda X + \lambda' X' = (\lambda X^p + \lambda' X'^p)^{\frac{1}{p}}$. Hence,

$$\|\lambda X + \lambda' X'\|_{p} = \|(\lambda X^{p} + \lambda' X'^{p})^{\frac{1}{p}}\|_{p} = (\operatorname{tr}[\lambda X^{p} + \lambda' X'^{p}])^{\frac{1}{p}} = (\lambda + \lambda')^{\frac{1}{p}} = 1.$$
(E.7)

By the subadditivity of norms,

$$1 = \|\lambda X + \lambda' X'\|_{p} \le \|\lambda X\|_{p} + \|\lambda' X'\|_{p} = \lambda + \lambda' = 1.$$
 (E.8)

Hence, the inequality in (E.8) must be saturated. The saturation of the inequality in (E.8) implies that $(\lambda X)^p$ is proportional to $(\lambda' X')^p$ for some strictly positive proportionality constant (due to the variational characterization of the Schatten norms [45, Lemma 3.2]). Since $(\lambda X)^p = \lambda^p \sigma$ and $(\lambda' X')^p = (\lambda')^p \sigma'$, we conclude that $\sigma = \sigma'$.

Proof of Theorem 6. Let $\alpha \in [\frac{1}{2}, 1)$. We will now prove (3.3).

=

$$\lambda Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) + \lambda' Q_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tau'_B)$$
(E.9)

$$= \lambda \operatorname{tr}[\rho_{AB}^{\alpha}(\sigma_A \otimes \tau_B)^{1-\alpha}] + \lambda' \operatorname{tr}[\rho_{AB}^{\alpha}(\sigma_A' \otimes \tau_B')^{1-\alpha}]$$
(E.10)

$$= \operatorname{tr}[\rho_{AB}^{\alpha}(\lambda(\sigma_A \otimes \tau_B)^{1-\alpha} + \lambda'(\sigma_A' \otimes \tau_B')^{1-\alpha})]$$
(E.11)

$$= \operatorname{tr}[\rho_{AB}^{\alpha}(\sqrt{\lambda\sigma_{A}^{2(1-\alpha)}} \otimes \sqrt{\lambda\tau_{B}^{2(1-\alpha)}} + \sqrt{\lambda'\sigma_{A}'^{2(1-\alpha)}} \otimes \sqrt{\lambda'\tau_{B}'^{2(1-\alpha)}})]$$
(E.12)

$$\leq \operatorname{tr}[\rho_{AB}^{\alpha}\sqrt{\lambda\sigma_{A}^{2(1-\alpha)} + \lambda'\sigma_{A}'^{2(1-\alpha)}} \otimes \sqrt{\lambda\tau_{B}^{2(1-\alpha)} + \lambda'\tau_{B}'^{2(1-\alpha)}}]$$
(E.13)

$$\leq \operatorname{tr}[\rho_{AB}^{\alpha}\sqrt{(\lambda\sigma_{A}+\lambda'\sigma_{A}')^{2(1-\alpha)}} \otimes \sqrt{\lambda\tau_{B}^{2(1-\alpha)}+\lambda'\tau_{B}'^{2(1-\alpha)}}]$$
(E.14)

$$\leq \operatorname{tr}[\rho_{AB}^{\alpha}\sqrt{(\lambda\sigma_A + \lambda'\sigma_A')^{2(1-\alpha)}} \otimes \sqrt{(\lambda\tau_B + \lambda'\tau_B')^{2(1-\alpha)}}]$$
(E.15)

$$= \operatorname{tr}[\rho_{AB}^{\alpha}(\lambda\sigma_{A} + \lambda'\sigma_{A}')^{1-\alpha} \otimes (\lambda\tau_{B} + \lambda'\tau_{B}')^{1-\alpha}]$$
(E.16)

$$= Q_{\alpha}(\rho_{AB} \| (\lambda \sigma_A + \lambda' \sigma'_A) \otimes (\lambda \tau_B + \lambda' \tau'_B))$$
(E.17)

(E.13) follows from Lemma 5. (E.14) and (E.15) hold because $X \mapsto X^{2(1-\alpha)}$ is operator concave since $2(1-\alpha) \in (0,1]$, and $X \mapsto \sqrt{X}$ is operator monotone. This completes the proof of (3.3).

We will now prove (3.4). If $\rho_{AB} \perp \sigma_A \otimes \tau_B$, then $D_{\alpha}(\rho_{AB} || \sigma_A \otimes \tau_B) = \infty$, so (3.4) is trivially true. If $\rho_{AB} \perp \sigma'_A \otimes \tau'_B$, then (3.4) is trivially true for the same reason. It remains to prove (3.4) for the case where both $\rho_{AB} \not\perp \sigma_A \otimes \tau_B$ and $\rho_{AB} \not\perp \sigma'_A \otimes \tau'_B$ hold. Then

$$\exp\left((\alpha-1)(\lambda D_{\alpha}(\rho_{AB}\|\sigma_A\otimes\tau_B)+\lambda' D_{\alpha}(\rho_{AB}\|\sigma'_A\otimes\tau'_B))\right)$$
(E.18)

$$= \operatorname{tr}[\rho_{AB}^{\alpha}(\sigma_A \otimes \tau_B)^{1-\alpha}]^{\lambda} \operatorname{tr}[\rho_{AB}^{\alpha}(\sigma_A' \otimes \tau_B')^{1-\alpha}]^{\lambda'}$$
(E.19)

$$\leq \lambda \operatorname{tr}[\rho_{AB}^{\alpha}(\sigma_A \otimes \tau_B)^{1-\alpha}] + \lambda' \operatorname{tr}[\rho_{AB}^{\alpha}(\sigma_A' \otimes \tau_B')^{1-\alpha}]$$
(E.20)

$$= \lambda Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) + \lambda' Q_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tau'_B)$$
(E.21)

$$\leq Q_{\alpha}(\rho_{AB} \| (\lambda \sigma_A + \lambda' \sigma'_A) \otimes (\lambda \tau_B + \lambda' \tau'_B)) \tag{E.22}$$

$$= \exp\left((\alpha - 1)D_{\alpha}(\rho_{AB} \| (\lambda \sigma_A + \lambda' \sigma'_A) \otimes (\lambda \tau_B + \lambda' \tau'_B))\right).$$
(E.23)

(E.20) follows from the weighted arithmetic-geometric mean inequality. (E.22) follows from (3.3). This completes the proof of (3.4).

We will now prove (3.5). If $\rho_A \perp \sigma_A$, then $I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) = \infty$, so (3.5) is trivially true. For the same reason, (3.5) is trivially true if $\rho_A \perp \sigma'_A$. It remains to prove (3.5) for the case where both $\rho_A \not\perp \sigma_A$ and $\rho_A \not\perp \sigma'_A$ hold. Let $\hat{\tau}_B \in \arg\min_{\tilde{\tau}_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tilde{\tau}_B)$ and let $\hat{\tau}'_B \in$ $\arg\min_{\tilde{\tau}_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tilde{\tau}_B)$. Then

$$\lambda I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) + \lambda' I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma'_A) = \lambda D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B) + \lambda' D_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \hat{\tau}'_B)$$
(E.24)

$$\geq D_{\alpha}(\rho_{AB} \| (\lambda \sigma_A + \lambda' \sigma'_A) \otimes (\lambda \hat{\tau}_B + \lambda' \hat{\tau}'_B))$$
(E.25)

$$\geq I^{\downarrow}_{\alpha}(\rho_{AB} \| \lambda \sigma_A + \lambda' \sigma'_A). \tag{E.26}$$

(E.25) follows from (3.4). This completes the proof of (3.5).

We will now prove the assertions below (3.5). Suppose $\alpha \in (\frac{1}{2}, 1)$.

First, suppose in addition that $\sigma_A, \sigma'_A \in S_{\ll \rho_A \ll}(A), \tau_B, \tau'_B \in S_{\ll \rho_B \ll}(B)$, and that (3.3) or (3.4) holds with equality. The proof above then implies that (3.3) holds with equality. In particular, the inequalities in (E.13), (E.14), and (E.15) must then hold with equality. Let us define the following positive semidefinite operators.

$$X_B \coloneqq \sqrt{\lambda \tau_B^{2(1-\alpha)} + \lambda' {\tau_B'}^{2(1-\alpha)}} \tag{E.27}$$

$$Y_B \coloneqq \sqrt{(\lambda \tau_B + \lambda' \tau'_B)^{2(1-\alpha)}}$$
(E.28)

$$Z_B := \operatorname{tr}_A[\rho_{AB}^{\alpha} \sqrt{(\lambda \sigma_A + \lambda' \sigma'_A)^{2(1-\alpha)}}]$$
(E.29)

The saturation of the inequality in (E.15) can then be expressed as $tr[X_BZ_B] = tr[Y_BZ_B]$. Since $\tilde{X} \mapsto \tilde{X}^{2(1-\alpha)}$ is operator concave, $X_B^2 \leq Y_B^2$. Since $\tilde{X} \mapsto \tilde{X}^{1/2}$ is operator monotone, $X_B \leq Y_B$. Furthermore, $Y_B \ll \rho_B$ and $\rho_B \ll Z_B$, so $Y_B \ll Z_B$. By applying Lemma 13 (b), it follows that $X_B = Y_B$, i.e.,

$$\lambda \tau_B^{2(1-\alpha)} + \lambda' \tau_B'^{2(1-\alpha)} = \left(\lambda \tau_B + \lambda' \tau_B'\right)^{2(1-\alpha)}.$$
(E.30)

An analogous argument can be made to conclude that

$$\lambda \sigma_A^{2(1-\alpha)} + \lambda' \sigma_A'^{2(1-\alpha)} = \left(\lambda \sigma_A + \lambda' \sigma_A'\right)^{2(1-\alpha)}.$$
(E.31)

By applying Lemma 14, we deduce from (E.30) and (E.31) that $\sigma_A = \sigma'_A$ and $\tau_B = \tau'_B$.

Now, suppose $\sigma_A, \sigma'_A \in \mathcal{S}_{\ll \rho_A \ll}(A)$, and that (3.5) holds with equality instead. Let $\hat{\tau}_B \in \arg\min_{\tilde{\tau}_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tilde{\tau}_B)$ and let $\hat{\tau}'_B \in \arg\min_{\tilde{\tau}_B \in \mathcal{S}(B)} D_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tilde{\tau}_B)$. By Proposition 12 (f), $\hat{\tau}_B, \hat{\tau}'_B \in \mathcal{S}_{\ll \rho_B \ll}(B)$. Since the inequality in (E.25) must be saturated, it follows from the saturation of (3.4) that $\sigma_A = \sigma'_A$.

3. Lemma for Theorem 7 (k)

In order to prove the following lemma, we will use Fréchet derivatives. We will now elucidate the notation for these derivatives. Consider $\mathcal{B}_A := \{X_A \in \mathcal{L}(A) : X_A \text{ is self-adjoint}\}$ with $\|\cdot\|_{\infty}$ as

a Banach space over \mathbb{R} . Similarly, consider $\mathcal{B}_B := \{Y_B \in \mathcal{L}(B) : Y_B \text{ is self-adjoint}\}$ with $\|\cdot\|_{\infty}$ as a Banach space over \mathbb{R} . Let $U \subseteq \mathcal{B}_A$ be an open set, and let $f : U \to \mathcal{B}_B$ be a Fréchet differentiable function. Then we denote the Fréchet derivative of f at $X \in U$ by $Df(X) \in \mathcal{L}(\mathcal{B}_A, \mathcal{B}_B)$. (Since we are working exclusively with finite-dimensional Hilbert spaces A and B, the Fréchet derivative of f remains unchanged if the norms of \mathcal{B}_A and \mathcal{B}_B are given by some other Schatten p-norm for $p \in [1, \infty)$ due to norm equivalence.) For a Fréchet differentiable map $f : \mathcal{S}_{>0}(A) \to \mathcal{B}_B$, we define the directional derivative of f at $\sigma \in \mathcal{S}_{>0}(A)$ in the direction of $\omega \in \mathcal{S}(A)$ as $\partial_{\omega} f(\sigma) \coloneqq Df(\sigma)(\omega - \sigma)$. For some basic properties of the directional derivative ∂_{ω} , we refer the reader to [15, Appendix C1].

The following lemma is an extension of [15, Lemma 22].

Lemma 15 (Equivalence of optimizers and fixed-points). Let $\alpha \in (0, \infty), \beta \in \mathbb{R}, \gamma \in [-1, 0) \cup (0, 1)$. Let $\rho_{AB} \in \mathcal{S}(AB)$ and $\tau_B \in \mathcal{S}_{\not{\perp}\rho_B}(B)$. Let $X_{AB} \coloneqq \tau_B^\beta \rho_{AB}^{\frac{1}{2}}$ and $X_A \coloneqq \operatorname{tr}_B[X_{AB}]$. Let us define the following functions and sets.

$$\chi_{\alpha,\gamma}: \quad \mathcal{S}(A) \to [0,\infty), \ \sigma_A \mapsto \chi_{\alpha,\gamma}(\sigma_A) \coloneqq \operatorname{tr}[(X_{AB}^{\dagger} \sigma_A^{\gamma} X_{AB})^{\alpha}]$$
(E.32)

$$\mathcal{X}_{\alpha,\gamma}: \quad \mathcal{S}_{\ll X_A \ll}(A) \to \mathcal{S}_{\ll X_A \ll}(A), \ \sigma_A \mapsto \mathcal{X}_{\alpha,\gamma}(\sigma_A) \coloneqq \frac{\operatorname{tr}_B[(\sigma_A^{\frac{\gamma}{2}} X_{AB} X_{AB}^{\dagger} \sigma_A^{\frac{\gamma}{2}})^{\alpha}]}{\operatorname{tr}[(\sigma_A^{\frac{\gamma}{2}} X_{AB} X_{AB}^{\dagger} \sigma_A^{\frac{\gamma}{2}})^{\alpha}]} \tag{E.33}$$

$$\mathcal{F}_{\alpha,\gamma} \coloneqq \{ \sigma_A \in \mathcal{S}_{\ll X_A \ll}(A) : \mathcal{X}_{\alpha,\gamma}(\sigma_A) = \sigma_A \}$$
(E.34)

$$\mathcal{M}_{\alpha,\gamma} \coloneqq \begin{cases} \underset{\sigma_A \in \mathcal{S}_{\ll X_A \ll}(A)}{\arg \min} \chi_{\alpha,\gamma}(\sigma_A) & \text{if } \gamma \in (0,1) \\ \underset{\sigma_A \in \mathcal{S}_{\ll X_A \ll}(A)}{\arg \min} \chi_{\alpha,\gamma}(\sigma_A) & \text{if } \gamma \in [-1,0) \end{cases}$$
(E.35)

Then all of the following hold.

(a) If $\gamma \in (0,1)$ and $\alpha \in (0,\frac{1}{\gamma}]$, then $\mathcal{F}_{\alpha,\gamma} = \mathcal{M}_{\alpha,\gamma} \neq \emptyset$. (b) If $\gamma \in [-1,0)$, then $\mathcal{F}_{\alpha,\gamma} = \mathcal{M}_{\alpha,\gamma} \neq \emptyset$.

Proof of (a). X_A is positive semidefinite. Without loss of generality, suppose X_A is positive definite. (Otherwise, the same proof works if A is restricted to $\text{supp}(X_A)$.) Then the sets defined in (E.34) and (E.35) can be expressed as follows.

$$\mathcal{F}_{\alpha,\gamma} = \{ \sigma_A \in \mathcal{S}_{>0}(A) : \mathcal{X}_{\alpha,\gamma}(\sigma_A) = \sigma_A \}$$
(E.36)

$$\mathcal{M}_{\alpha,\gamma} = \operatorname*{arg\,max}_{\sigma_A \in \mathcal{S}_{>0}(A)} \chi_{\alpha,\gamma}(\sigma_A) \tag{E.37}$$

We will now prove that there exists a maximizer for

$$\max_{\sigma_A \in \mathcal{S}_{>0}(A)} \chi_{\alpha,\gamma}(\sigma_A).$$
(E.38)

To this end, we will show that the directional derivative of $\chi_{\alpha,\gamma}(\sigma_A)$ at $\sigma_A \in S_{>0}(A)$ in the direction of the maximally mixed state ω_A is strictly positive if (at least) one of the eigenvalues of σ_A becomes sufficiently small. Let $\sigma_A \in S_{>0}(A)$, let $\omega_A := 1_A/d_A \in S(A)$ and let us define the map $f: S_{>0}(A) \to \mathcal{L}(A), \sigma_A \mapsto \sigma_A^{\gamma}$. Its Fréchet derivative at σ_A is $Df(\sigma_A)(X_A) = f^{[1]}(\sigma_A) \odot X_A$ for all self-adjoint $X_A \in \mathcal{L}(A)$, where \odot denotes the Hadamard product taken in an eigenbasis of σ_A and $f^{[1]}$ is the first-order divided difference of f [52]. Then,

$$\partial_{\omega_A} \chi_{\alpha,\gamma}(\sigma_A) = \alpha \operatorname{tr}[X_{AB} (X_{AB}^{\dagger} \sigma_A^{\gamma} X_{AB})^{\alpha - 1} X_{AB}^{\dagger} \partial_{\omega_A} f(\sigma_A)]$$
(E.39)

$$= \alpha \operatorname{tr}[\operatorname{tr}_B[X_{AB}(X_{AB}^{\dagger}\sigma_A^{\gamma}X_{AB})^{\alpha-1}X_{AB}^{\dagger}]\partial_{\omega_A}f(\sigma_A)]$$
(E.40)

$$= \alpha \gamma \operatorname{tr}[\operatorname{tr}_{B}[X_{AB}(X_{AB}^{\dagger}\sigma_{A}^{\gamma}X_{AB})^{\alpha-1}X_{AB}^{\dagger}]\left(\frac{1}{d_{A}}\sigma_{A}^{\gamma-1}-\sigma_{A}^{\gamma}\right)].$$
(E.41)

(E.41) holds because

$$\partial_{\omega_A} f(\sigma_A) = Df(\sigma_A)(\omega_A - \sigma_A) = f^{[1]}(\sigma_A) \odot (\omega_A - \sigma_A)$$
(E.42)

$$=\gamma\sigma_A^{\gamma-1}\odot(\omega_A-\sigma_A)=\gamma\sigma_A^{\gamma-1}(\omega_A-\sigma_A)=\gamma\left(\frac{1}{d_A}\sigma_A^{\gamma-1}-\sigma_A^{\gamma}\right),\qquad(E.43)$$

where the first two equalities in (E.43) hold because $\omega_A - \sigma_A$ is diagonal in any eigenbasis of σ_A . Since $\alpha > 0$ and $\gamma > 0$, the prefactor in (E.41) is strictly positive. The trace in (E.41) is the trace of the product of two self-adjoint operators on A, and the first operator is positive definite. Now, consider the case where (at least) one of the eigenvalues of σ_A is arbitrarily small but non-zero. Since $0 < \gamma < 1$, the corresponding eigenvalue of $\sigma_A^{\gamma-1}$ becomes arbitrarily large, whereas none of the eigenvalues of σ_A^{γ} diverges in this limit. Consequently, if one of the eigenvalues of σ_A is sufficiently small, then $\partial_{\omega_A} \chi_{\alpha,\gamma}(\sigma_A)$ is strictly positive. Hence, there exists a maximizer for the optimization problem in (E.38), so $\mathcal{M}_{\alpha,\gamma} \neq \emptyset$.

It remains to prove that $\mathcal{M}_{\alpha,\gamma} = \mathcal{F}_{\alpha,\gamma}$. The function $\chi_{\alpha,\gamma}$ is concave [53, Theorem 2.1(a)] (see also [54–56]). Hence, for any $\sigma_A \in \mathcal{S}_{>0}(A)$: $\sigma_A \in \mathcal{M}_{\alpha,\gamma}$ iff $\partial_{\omega_A}\chi_{\alpha,\gamma}(\sigma_A) = 0 \forall \omega_A \in \mathcal{S}(A)$. The argument in [15, Proof of Lemma 22] shows that the second condition is equivalent to the condition that

$$\sigma_A^{-\frac{1}{2}} \operatorname{tr}_B[(\sigma_A^{\frac{\gamma}{2}} X_{AB} X_{AB}^{\dagger} \sigma_A^{\frac{\gamma}{2}})^{\alpha}] \sigma_A^{-\frac{1}{2}}$$
(E.44)

is proportional to the identity. This is equivalent to $\sigma_A \in \mathcal{F}_{\alpha,\gamma}$. Therefore, $\mathcal{M}_{\alpha,\gamma} = \mathcal{F}_{\alpha,\gamma}$.

Proof of (b). X_A is positive semidefinite. Without loss of generality, suppose X_A is positive definite. Then the sets defined in (E.34) and (E.35) can be expressed as follows.

$$\mathcal{F}_{\alpha,\gamma} = \{ \sigma_A \in \mathcal{S}_{>0}(A) : \mathcal{X}_{\alpha,\gamma}(\sigma_A) = \sigma_A \}$$
(E.45)

$$\mathcal{A}_{\alpha,\gamma} = \operatorname*{arg\,min}_{\sigma_A \in \mathcal{S}_{>0}(A)} \chi_{\alpha,\gamma}(\sigma_A) \tag{E.46}$$

Since $\alpha > 0, \gamma < 0$, and $\chi_{\alpha,\gamma}(\sigma_A) = \operatorname{tr}[(X_{AB}^{\dagger}\sigma_A^{\gamma}X_{AB})^{\alpha}]$, it is clear that $\chi_{\alpha,\gamma}(\sigma_A)$ diverges to $+\infty$ if (at least) one of the eigenvalues of σ_A becomes arbitrarily small. Hence, there exists a minimizer for $\min_{\sigma_A \in \mathcal{S}_{>0}(A)} \chi_{\alpha,\gamma}(\sigma_A)$, so $\mathcal{M}_{\alpha,\gamma} \neq \emptyset$.

It remains to prove that $\mathcal{M}_{\alpha,\gamma} = \mathcal{F}_{\alpha,\gamma}$. The function $\chi_{\alpha,\gamma}$ is convex [53, Theorem 2.1(b)] (see also [56]). Hence, for any $\sigma_A \in \mathcal{S}_{>0}(A)$: $\sigma_A \in \mathcal{M}_{\alpha,\gamma}$ iff $\partial_{\omega_A}\chi_{\alpha,\gamma}(\sigma_A) = 0 \forall \omega_A \in \mathcal{S}(A)$. The argument in [15, Proof of Lemma 22] shows that the second condition is equivalent to the condition that the operator in (E.44) is proportional to the identity. Therefore, $\mathcal{M}_{\alpha,\gamma} = \mathcal{F}_{\alpha,\gamma}$.

4. Proof of Theorem 7

Proof of (a). This assertion follows from the symmetry of the definition of the doubly minimized PRMI in (2.32) with respect to A and B. \Box

Proof of (b), (e), (f), (h), (i), (n), (r). Since $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \inf_{\sigma_A \in \mathcal{S}(A)} I_{\alpha}^{\downarrow}(\rho_{AB} || \sigma_A)$, these properties follow from corresponding properties of the minimized generalized PRMI, see Proposition 12. In particular, (h) and (i) follow from Proposition 12 (f).

Proof of (c).

$$I_{\alpha}^{\downarrow\downarrow}(A':B')_{V\otimes W\rho_{AB}V^{\dagger}\otimes W^{\dagger}} = \inf_{\sigma_{A'}\in\mathcal{S}(A')} I_{\alpha}^{\downarrow}(V\otimes W\rho_{AB}V^{\dagger}\otimes W^{\dagger}\|\sigma_{A'})$$
(E.47)

$$= \inf_{\sigma_{A'} \in \mathcal{S}(A')} I^{\downarrow}_{\alpha}(V \rho_{AB} V^{\dagger} \| \sigma_{A'}) = I^{\downarrow\downarrow}_{\alpha}(A':B)_{V \rho_{AB} V^{\dagger}}$$
(E.48)

$$= \inf_{\tau_B \in \mathcal{S}(B)} I^{\downarrow}_{\alpha} (V \rho_{AB} V^{\dagger} \| \tau_B)$$
(E.49)

$$= \inf_{\tau_B \in \mathcal{S}(B)} I^{\downarrow}_{\alpha}(\rho_{AB} \| \tau_B) = I^{\downarrow\downarrow}_{\alpha}(A:B)_{\rho_{AB}}$$
(E.50)

Above, we have used the invariance of the minimized generalized PRMI under local isometries, see Proposition 12 (b), twice: for the first equality in (E.48), and for the first equality in (E.50). \Box

Proof of (j). If $\alpha = 1$, then the assertion is true due to (2.14). It remains to prove (j) for $\alpha \in (\frac{1}{2}, 1)$. Let $\rho_{AB} \in \mathcal{S}(AB), \alpha \in (\frac{1}{2}, 1)$ be fixed. Let $\mathcal{M}_{\alpha} \coloneqq \arg\min_{\sigma_A \in \mathcal{S}(A)} I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A)$. We will now prove that $\mathcal{M}_{\alpha} \subseteq \mathcal{S}_{\ll \rho_A \ll}(A)$ by cases.

Case 1: $\rho_{AB} = \rho_A \otimes \rho_B$. By the positive definiteness of the Petz divergence, $\mathcal{M}_{\alpha} = \{\rho_A\}$. Therefore, $\mathcal{M}_{\alpha} \subseteq \mathcal{S}_{\ll \rho_A \ll}(A)$.

Case 2: $\rho_{AB} \neq \rho_A \otimes \rho_B$. For this case, we will prove that

=

$$\underset{(\sigma_A,\tau_B)\in\mathcal{S}(A)\times\mathcal{S}(B)}{\arg\min} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) = \underset{(\sigma_A,\tau_B)\in\mathcal{S}_{\ll\rho_A}\ll(A)\times\mathcal{S}_{\ll\rho_B}\ll(B)}{\arg\min} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B).$$
(E.51)

Clearly, the assertion that $\mathcal{M}_{\alpha} \subseteq \mathcal{S}_{\ll \rho_A \ll}(A)$ then follows from (E.51). In order to prove (E.51), let

$$(\hat{\sigma}_A, \hat{\tau}_B) \in \underset{(\sigma_A, \tau_B) \in \mathcal{S}(A) \times \mathcal{S}(B)}{\operatorname{arg\,min}} D_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B).$$
(E.52)

By Proposition 12 (f), $\hat{\sigma}_A \ll \rho_A$ and $\hat{\tau}_B \ll \rho_B$. It remains to show that $\rho_A \ll \hat{\sigma}_A$ and $\rho_B \ll \hat{\tau}_B$. Note that if $\rho_A \ll \hat{\sigma}_A$, then $\rho_B \ll \hat{\tau}_B$ by Proposition 12 (f). So it suffices to show that $\rho_A \ll \hat{\sigma}_A$. We will prove this by contradiction.

Suppose it is false that $\rho_A \ll \hat{\sigma}_A$. Let $\sigma_A \in \mathcal{S}(A)$ be the quantum state that is proportional to the orthogonal projection onto ker $(\hat{\sigma}_A)$. Let $\tau_B \in \mathcal{S}(B)$, to be specified later. Let

$$\hat{f} \coloneqq Q_{\alpha}(\rho_{AB} \| \hat{\sigma}_A \otimes \hat{\tau}_B) = \max_{(\sigma'_A, \tau'_B) \in \mathcal{S}(A) \times \mathcal{S}(B)} Q_{\alpha}(\rho_{AB} \| \sigma'_A \otimes \tau'_B) > 0.$$
(E.53)

Let us define the following two functions of $\lambda \in [0, 1]$.

$$f_1(\lambda) \coloneqq Q_\alpha(\rho_{AB} \| ((1-\lambda)\hat{\sigma}_A + \lambda\sigma_A) \otimes ((1-\lambda)\hat{\tau}_B + \lambda\tau_B))$$
(E.54)

$$f_2(\lambda) \coloneqq \hat{f} - (2 - \alpha)\lambda \hat{f} + \lambda^{1 - \alpha} Q_\alpha(\rho_{AB} \| \sigma_A \otimes ((1 - \lambda)\hat{\tau}_B + \lambda \tau_B))$$
(E.55)

Then, for all $\lambda \in [0, 1]$

$$\hat{f} \ge f_1(\lambda)$$

$$= (1-\lambda)^{1-\alpha} Q_\alpha(\rho_{AB} \| \hat{\sigma}_A \otimes ((1-\lambda)\hat{\tau}_B + \lambda\tau_B)) + \lambda^{1-\alpha} Q_\alpha(\rho_{AB} \| \sigma_A \otimes ((1-\lambda)\hat{\tau}_B + \lambda\tau_B))$$
(E.56b)
(E.56b)

$$\geq (1-\lambda)^{1-\alpha}((1-\lambda)\hat{f} + \lambda Q_{\alpha}(\rho_{AB} \| \hat{\sigma}_A \otimes \tau_B)) + \lambda^{1-\alpha} Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes ((1-\lambda)\hat{\tau}_B + \lambda \tau_B))$$
(E.56c)

$$\geq (1-\lambda)^{2-\alpha} \hat{f} + \lambda^{1-\alpha} Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes ((1-\lambda)\hat{\tau}_B + \lambda\tau_B))$$
(E.56d)

$$\geq (1 - (2 - \alpha)\lambda)\hat{f} + \lambda^{1 - \alpha}Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes ((1 - \lambda)\hat{\tau}_B + \lambda\tau_B)) = f_2(\lambda).$$
(E.56e)

(E.56a) follows from (E.52). (E.56b) holds because $\hat{\sigma}_A \perp \sigma_A$ implies that $((1 - \lambda)\hat{\sigma}_A + \lambda\sigma_A)^{1-\alpha} = ((1 - \lambda)\hat{\sigma}_A)^{1-\alpha} + (\lambda\sigma_A)^{1-\alpha}$. (E.56c) follows from the concavity of $Q_\alpha(\rho_{AB} \parallel \cdot)$ [45]. (E.56e) holds because $(1 - \lambda)^{2-\alpha} \geq 1 - (2 - \alpha)\lambda$ for all $\lambda \in [0, 1]$.

Case 2.1: $Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B) > 0$. Then we define $\tau_B \coloneqq \hat{\tau}_B$. By (E.55),

$$f_2(\lambda) = \hat{f} + \lambda(-(2-\alpha)\hat{f} + \lambda^{-\alpha}Q_\alpha(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B))$$
(E.57)

for all $\lambda \in [0,1]$. Let $\lambda_0 \coloneqq \left(\frac{Q_\alpha(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B)}{(2-\alpha)\hat{f}}\right)^{\frac{1}{\alpha}} > 0$. Then $f_2(\lambda_0) = \hat{f}$. Since $-\alpha < 0$, it follows from (E.57) that $f_2(\lambda) > \hat{f}$ for all $\lambda \in (0, \lambda_0)$. This contradicts (E.56).

Case 2.2: $Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B) = 0$. Then we define τ_B as the quantum state which is proportional to the orthogonal projection onto ker $(\hat{\tau}_B)$. Then, for all $\lambda \in [0, 1]$

$$Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes ((1-\lambda)\hat{\tau}_B + \lambda\tau_B)) = \lambda^{1-\alpha} Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) + (1-\lambda)^{1-\alpha} Q_{\alpha}(\rho_{AB} \| \sigma_A \otimes \hat{\tau}_B)$$
(E.58a)

$$=\lambda^{1-\alpha}Q_{\alpha}(\rho_{AB}\|\sigma_A\otimes\tau_B).$$
(E.58b)

The left-hand side of (E.58a) is non-zero for any $\lambda \in (0,1)$ because $\sigma_A \ll \rho_A$ and $(1-\lambda)\hat{\tau}_B + \lambda \tau_B$ has full rank. Hence, the expression in (E.58b) must be non-zero for any $\lambda \in (0,1)$, which implies that $Q_{\alpha}(\rho_{AB} || \sigma_A \otimes \tau_B) > 0$. The combination of (E.58) and (E.55) implies that

$$f_2(\lambda) = \hat{f} + \lambda(-(2-\alpha)\hat{f} + \lambda^{1-2\alpha}Q_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B)) \qquad \forall \lambda \in [0,1].$$
(E.59)

Let $\lambda_0 \coloneqq (\frac{Q_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B)}{(2-\alpha)\hat{f}})^{\frac{1}{2\alpha-1}} > 0$. Then $f_2(\lambda_0) = \hat{f}$. Since $1 - 2\alpha < 0$, it follows from (E.59) that $f_2(\lambda) > \hat{f}$ for all $\lambda \in (0, \lambda_0)$. This contradicts (E.56). This completes the proof of the inclusion $\mathcal{M}_\alpha \subseteq \mathcal{S}_{\ll \rho_A \ll}(A)$.

We will now prove that \mathcal{M}_{α} cannot contain more than one element. Let $\sigma_A, \sigma'_A \in \mathcal{M}_{\alpha}$. Let $\lambda := \frac{1}{2}, \lambda' := \frac{1}{2}$. Then, (3.5) in Theorem 6 implies that also $\lambda \sigma_A + \lambda' \sigma'_A \in \mathcal{M}_{\alpha}$. We have

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\downarrow}(\rho_{AB} \| \lambda \sigma_A + \lambda' \sigma'_A) = \lambda I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A) + \lambda' I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma'_A).$$
(E.60)

Hence, the inequality in (3.5) is saturated. By Theorem 6, $\sigma_A = \sigma'_A$. This proves that \mathcal{M}_{α} contains at most one element. We conclude that there exists $\hat{\sigma}_A \in \mathcal{S}_{\ll \rho_A \ll}(A)$ such that $\mathcal{M}_{\alpha} = \{\hat{\sigma}_A\}$. The assertion now follows from Proposition 12 (f).

Proof of (k). Let $\alpha \in (\frac{1}{2}, 2]$.

Case 1: $\alpha = 1$. Then $\mathcal{M}_{\alpha} = \{\rho_A\}$ due to (2.15). For any $\sigma_A \in \mathcal{S}_{\ll \rho_A \ll}(A)$: $\sigma_A \in \mathcal{F}_{\alpha}$ iff $\sigma_A = \operatorname{tr}_C[\rho_A C]$. Hence, $\mathcal{F}_{\alpha} = \{\rho_A\}$. Therefore, $\mathcal{M}_{\alpha} = \mathcal{F}_{\alpha}$.

Case 2: $\alpha \in (\frac{1}{2}, 1)$. Let $\beta := \frac{1}{\alpha} \in (1, 2)$ and $\gamma := 1 - \alpha = \frac{\beta - 1}{\beta} \in (0, \frac{1}{2})$. Let $X_{AC} := \rho_C^{\frac{1 - \beta}{2\beta}} \rho_{AC}^{\frac{1}{2}}$ and $X_A := \operatorname{tr}_C[X_{AC}]$. Then $\operatorname{supp}(X_A) = \operatorname{supp}(\rho_A)$. By (j) and the duality of the minimized generalized PRMI, see Proposition 12 (d),

$$\mathcal{M}_{\alpha} = \underset{\sigma_{A} \in \mathcal{S}_{\ll \rho_{A} \ll}(A)}{\operatorname{arg\,max}} \widetilde{Q}_{\beta}(\rho_{AC} \| \sigma_{A}^{-1} \otimes \rho_{C}) = \underset{\sigma_{A} \in \mathcal{S}_{\ll X_{A} \ll}(A)}{\operatorname{arg\,max}} \operatorname{tr}[(X_{AC}^{\dagger} \sigma_{A}^{\gamma} X_{AC})^{\beta}].$$
(E.61)

Lemma 15 (a) implies that $\mathcal{M}_{\alpha} = \mathcal{F}_{\alpha}$ since $\gamma \in (0, 1)$ and $0 < \beta \leq 2 \leq \frac{1}{\gamma}$.

Case 3: $\alpha \in (1,2]$. Let $\beta := \frac{1}{\alpha} \in [\frac{1}{2}, 1)$ and $\gamma := 1 - \alpha = \frac{\beta - 1}{\beta} \in [-1, 0)$. Let $X_{AC} := \rho_C^{\frac{1 - \beta}{2\beta}} \rho_{AC}^{\frac{1}{2}}$ and $X_A := \operatorname{tr}_C[X_{AC}]$. Then $\operatorname{supp}(X_A) = \operatorname{supp}(\rho_A)$. By (h) and the duality of the minimized generalized PRMI, see Proposition 12 (d),

$$\mathcal{M}_{\alpha} = \operatorname*{arg\,min}_{\sigma_A \in \mathcal{S}_{\ll \rho_A \ll}(A)} \widetilde{Q}_{\beta}(\rho_{AC} \| \sigma_A^{-1} \otimes \rho_C) = \operatorname*{arg\,min}_{\sigma_A \in \mathcal{S}_{\ll X_A \ll}(A)} \operatorname{tr}[(X_{AC}^{\dagger} \sigma_A^{\gamma} X_{AC})^{\beta}]. \tag{E.62}$$

Lemma 15 (b) implies that $\mathcal{M}_{\alpha} = \mathcal{F}_{\alpha}$ since $\gamma \in [-1, 0)$ and $0 < \beta$.

Proof of (d). Let $\alpha \in [\frac{1}{2}, 2]$.

Case 1: $\alpha \in (\frac{1}{2}, 2]$. Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$, and let $|\rho'\rangle_{DEF} \in \mathcal{O}(\mathbb{R})$ DEF be such that $\operatorname{tr}_F[|\rho'\rangle\langle\rho'|_{DEF}] = \rho'_{DE}$.

Let $\sigma_A \in \arg\min_{\tilde{\sigma}_A \in \mathcal{S}(A)} I^{\downarrow}_{\alpha}(\rho_{AB} \| \tilde{\sigma}_A)$. According to (k), $\sigma_A \in \mathcal{S}_{\ll \rho_A \ll}(A)$ and

$$\sigma_A = \frac{\operatorname{tr}_C\left[\left(\sigma_A^{\frac{1-\alpha}{2}} \otimes \rho_C^{\frac{\alpha-1}{2}} \rho_{AC} \sigma_A^{\frac{1-\alpha}{2}} \otimes \rho_C^{\frac{\alpha-1}{2}}\right)^{\frac{1}{\alpha}}\right]}{\operatorname{tr}\left[\left(\sigma_A^{\frac{1-\alpha}{2}} \otimes \rho_C^{\frac{\alpha-1}{2}} \rho_{AC} \sigma_A^{\frac{1-\alpha}{2}} \otimes \rho_C^{\frac{\alpha-1}{2}}\right)^{\frac{1}{\alpha}}\right]}.$$
(E.63)

Let $\sigma'_D \in \arg\min_{\tilde{\sigma}'_D \in \mathcal{S}(D)} I^{\downarrow}_{\alpha}(\rho'_{DE} \| \tilde{\sigma}'_D)$. According to (k), $\sigma'_D \in \mathcal{S}_{\ll \rho'_D \ll}(D)$ and

$$\sigma'_D = \frac{\operatorname{tr}_F[(\sigma'_D^{\frac{1-\alpha}{2}} \otimes \rho'_F^{\frac{\alpha-1}{2}} \rho'_{DF} \sigma'_D^{\frac{1-\alpha}{2}} \otimes \rho'_F^{\frac{\alpha-1}{2}})^{\frac{1}{\alpha}}]}{\operatorname{tr}[(\sigma'_D^{\frac{1-\alpha}{2}} \otimes \rho'_F^{\frac{\alpha-1}{2}} \rho'_{DF} \sigma'_D^{\frac{1-\alpha}{2}} \otimes \rho'_F^{\frac{\alpha-1}{2}})^{\frac{1}{\alpha}}]}.$$
(E.64)

Clearly, we have $\sigma_A \otimes \sigma'_D \in \mathcal{S}_{\ll \rho_A \otimes \rho'_D \ll}(AD)$, and the combination of (E.63) and (E.64) implies that

$$\sigma_A \otimes \sigma'_D = \frac{\operatorname{tr}_{CF}[((\sigma_A \otimes \sigma'_D)^{\frac{1-\alpha}{2}} \otimes (\rho_C \otimes \rho'_F)^{\frac{\alpha-1}{2}} \rho_{AC} \otimes \rho'_{DF} (\sigma_A \otimes \sigma'_D)^{\frac{1-\alpha}{2}} \otimes (\rho_C \otimes \rho'_F)^{\frac{\alpha-1}{2}})^{\frac{1}{\alpha}}]}{\operatorname{tr}[((\sigma_A \otimes \sigma'_D)^{\frac{1-\alpha}{2}} \otimes (\rho_C \otimes \rho'_F)^{\frac{\alpha-1}{2}} \rho_{AC} \otimes \rho'_{DF} (\sigma_A \otimes \sigma'_D)^{\frac{1-\alpha}{2}} \otimes (\rho_C \otimes \rho'_F)^{\frac{\alpha-1}{2}})^{\frac{1}{\alpha}}]}.$$
(E.65)

We can deduce from (E.65) that $\sigma_A \otimes \sigma'_D \in \arg\min_{\tilde{\sigma}_{AD} \in \mathcal{S}(AD)} I^{\downarrow}_{\alpha}(\rho_{AB} \otimes \rho'_{DE} \| \tilde{\sigma}_{AD})$ due to (k). Hence,

$$I_{\alpha}^{\downarrow\downarrow}(AD:BE)_{\rho_{AB}\otimes\rho_{DE}'} = I_{\alpha}^{\downarrow}(\rho_{AB}\otimes\rho_{DE}' \| \sigma_A\otimes\sigma_D')$$
(E.66)

$$= I^{\downarrow}_{\alpha}(\rho_{AB} \| \sigma_A) + I^{\downarrow}_{\alpha}(\rho'_{DE} \| \sigma'_D)$$
(E.67)

$$= I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho_{AB}} + I_{\alpha}^{\downarrow\downarrow}(D:E)_{\rho'_{DE}}.$$
(E.68)

(E.67) follows from the additivity of the minimized generalized PRMI, see Proposition 12 (c).

Case 2: $\alpha = \frac{1}{2}$. Then the assertion follows from case 1 by taking the limit $\alpha \to 1/2^+$ due to the continuity in α of the Petz divergence.

Proof of (l). Let $\alpha \in [0,2]$. Then, for any $n \in \mathbb{N}_{>0}$

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \inf_{\substack{\sigma_A \in \mathcal{S}(A), \\ \tau_B \in \mathcal{S}(B)}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_A^{\otimes n} \otimes \tau_B^{\otimes n})$$
(E.69a)

$$\geq \inf_{\substack{\sigma_{A^n} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^n} \otimes \tau_{B^n})$$
(E.69b)

$$\geq \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) - \frac{\log g_{n,d_A}}{n} - \frac{\log g_{n,d_B}}{n}$$
(E.69c)

$$\geq \inf_{\substack{\sigma_{A^n} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^n} \otimes \tau_{B^n}) - \frac{\log g_{n,d_A}}{n} - \frac{\log g_{n,d_B}}{n}.$$
(E.69d)

(E.69a) follows from the additivity of the Petz divergence. (E.69c) follows from Proposition 1 (b). (E.69d) follows from Proposition 1 (a). In the limit $n \to \infty$, the two terms on the right-hand side of (E.69c) vanish due to Proposition 1 (b). Hence, $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \geq \limsup_{n\to\infty} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} || \omega_{A^n}^n \otimes \omega_{B^n}^n).$

Case 1: $\alpha \in [\frac{1}{2}, 2]$. Then, for any $n \in \mathbb{N}_{>0}$

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \ge \inf_{\substack{\sigma_{A^{n}} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^{n}} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^{n}} \otimes \tau_{B^{n}})$$
(E.70)

$$\geq \inf_{\substack{\sigma_{A^n} \in \mathcal{S}(A^n), \\ \tau_{B^n} \in \mathcal{S}(B^n)}} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \sigma_{A^n} \otimes \tau_{B^n}) = \frac{1}{n} I_{\alpha}^{\downarrow\downarrow}(A^n : B^n)_{\rho^{\otimes n}} = I_{\alpha}^{\downarrow\downarrow}(A : B)_{\rho}.$$
(E.71)

(E.70) follows from (E.69b). (E.71) follows from additivity (d). This proves the assertion in (3.18). Combining this result with (E.69) and taking the limit $n \to \infty$ proves the assertion in (3.17) for $\alpha \in [\frac{1}{2}, 2]$.

Case 2: $\alpha \in (0, \frac{1}{2})$. Let A' and B' be Hilbert spaces isomorphic to A and B, respectively. Then, for any $n \in \mathbb{N}_{>0}$

$$\exp((\alpha - 1)D_{\alpha}(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n)) \tag{E.72}$$

$$= \operatorname{tr}[(\rho_{AB}^{\otimes n})^{\alpha}(\omega_{A^{n}}^{n} \otimes \omega_{B^{n}}^{n})^{1-\alpha}]$$
(E.73)

$$= \operatorname{tr}[(\rho_{AB}^{\otimes n})^{\alpha}\omega_{A^{n}}^{n} \otimes \omega_{B^{n}}^{n}(\omega_{A^{n}}^{n} \otimes \omega_{B^{n}}^{n})^{-\alpha}]$$
(E.74)

$$= \int_{\mathcal{U}(AA')} d\mu_H(U) \int_{\mathcal{U}(BB')} d\mu_H(V) \operatorname{tr}[(\rho_{AB}^{\otimes n})^{\alpha} \sigma(U)_A^{\otimes n} \otimes \tau(V)_B^{\otimes n} (\omega_{A^n}^n \otimes \omega_{B^n}^n)^{-\alpha}]$$
(E.75)

$$= \int_{\mathcal{U}(AA')} d\mu_H(U) \int_{\mathcal{U}(BB')} d\mu_H(V) \operatorname{tr}[(\sigma(U)_A^{\otimes n} \otimes \tau(V)_B^{\otimes n})^{\frac{1}{2}} (\rho_{AB}^{\otimes n})^{\alpha} (\sigma(U)_A^{\otimes n} \otimes \tau(V)_B^{\otimes n})^{\frac{1}{2}} (\omega_{A^n}^n \otimes \omega_{B^n}^n)^{-\alpha}]$$
(E.76)

$$\leq g_{n,d_A}^{\alpha} g_{n,d_B}^{\alpha} \int_{\mathcal{U}(AA')} \mathrm{d}\mu_H(U) \int_{\mathcal{U}(BB')} \mathrm{d}\mu_H(V) \operatorname{tr} \left[(\sigma(U)_A^{\otimes n} \otimes \tau(V)_B^{\otimes n})^{\frac{1}{2}} (\rho_{AB}^{\otimes n})^{\alpha} (\sigma(U)_A^{\otimes n} \otimes \tau(V)_B^{\otimes n})^{\frac{1}{2} - \alpha} \right]$$
(E.77)

$$=g_{n,d_A}^{\alpha}g_{n,d_B}^{\alpha}\int_{\mathcal{U}(AA')}\mathrm{d}\mu_H(U)\int_{\mathcal{U}(BB')}\mathrm{d}\mu_H(V)(\mathrm{tr}[(\rho_{AB})^{\alpha}(\sigma(U)_A\otimes\tau(V)_B)^{1-\alpha}])^n\tag{E.78}$$

$$\leq g_{n,d_A}^{\alpha} g_{n,d_B}^{\alpha} \sup_{\sigma_A \in \mathcal{S}(A)} \sup_{\tau_B \in \mathcal{S}(B)} (\operatorname{tr}[(\rho_{AB})^{\alpha} (\sigma_A \otimes \tau_B)^{1-\alpha}])^n \tag{E.79}$$

$$=g_{n,d_A}^{\alpha}g_{n,d_B}^{\alpha}\exp((\alpha-1)nI_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}).$$
(E.80)

(E.75) follows from Proposition 1 (d). (E.76) follows from Proposition 1 (c). (E.77) follows from Proposition 1 (b) and the operator anti-monotonicity of $X \mapsto X^{-\alpha}$ for $\alpha \in (0, 1)$. Hence, for any $n \in \mathbb{N}_{>0}$

$$\frac{1}{n}D_{\alpha}(\rho_{AB}^{\otimes n}\|\omega_{A^{n}}^{n}\otimes\omega_{B^{n}}^{n})\geq I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}-\frac{\alpha}{1-\alpha}\left(\frac{\log g_{n,d_{A}}}{n}+\frac{\log g_{n,d_{B}}}{n}\right).$$
(E.81)

By Proposition 1 (b), the second term on the right-hand side vanishes in the limit $n \to \infty$. Therefore,

$$\liminf_{n \to \infty} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) \ge I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}.$$
(E.82)

Case 3: $\alpha = 0$. By monotonicity in α (n), we have for any $n \in \mathbb{N}_{>0}$

$$I_0^{\downarrow\downarrow}(A:B)_{\rho} \le \liminf_{\alpha \to 0^+} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \le \liminf_{\alpha \to 0^+} \frac{1}{n} D_{\alpha}(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) = \frac{1}{n} D_0(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n),$$
(E.83)

where we have used (E.81) and the continuity in α of the Petz divergence, see Proposition 2. Therefore, (E.82) also holds for $\alpha = 0$.

Proof of (m). $I_1^{\downarrow\downarrow}(A:B)_{\rho} = I(A:B)_{\rho}$ follows from (2.12). Furthermore, we have

$$I_0^{\downarrow\downarrow}(A:B)_\rho = \inf_{\sigma_A \in \mathcal{S}(A)} I_0^{\downarrow}(\rho_{AB} \| \sigma_A) = \inf_{\substack{\sigma_A \in \mathcal{S}(A) \mid \tau \rangle_B \in \text{supp}(\rho_B):\\\langle \tau \mid \tau \rangle_B = 1}} D_0(\rho_{AB} \| \sigma_A \otimes |\tau\rangle \langle \tau |_B)$$
(E.84)

$$= \inf_{\substack{|\tau\rangle_B \in \operatorname{supp}(\rho_B):\\\langle\tau|\tau\rangle_B = 1}} I_0^{\downarrow}(\rho_{AB} |||\tau\rangle\!\langle\tau|_B)$$
(E.85)

$$= \inf_{\substack{|\tau\rangle_B \in \operatorname{supp}(\rho_B): |\sigma\rangle_A \in \operatorname{supp}(\rho_A):\\ \langle \tau | \tau \rangle_B = 1 \quad \langle \sigma | \sigma \rangle_A = 1}} \inf_{\substack{D_0(\rho_{AB} \| |\sigma\rangle \langle \sigma|_A \otimes |\tau\rangle \langle \tau|_B).}} (E.86)$$

Above, we have used Proposition 12 (i) twice: for the second equality in (E.84), and for (E.86). \Box

Proof of (o). The continuity of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ on $\alpha \in [0,1)$ and on $\alpha \in [1,\infty)$ follows from the continuity in α of the Petz divergence. It remains to prove left-continuity at $\alpha = 1$. By (E.69), we have for any $n \in \mathbb{N}_{>0}$

$$\frac{1}{n}D_1(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) - \frac{\log g_{n,d_A}}{n} - \frac{\log g_{n,d_B}}{n} \le \lim_{\alpha \to 1^-} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \le I_1^{\downarrow\downarrow}(A:B)_{\rho}, \qquad (E.87)$$

where the last inequality follows from the monotonicity in α (n). By Proposition 1 (b), the second term and the third term on the left-hand side of (E.87) vanish in the limit $n \to \infty$. Therefore,

$$I_1^{\downarrow\downarrow}(A:B)_\rho = \lim_{n \to \infty} \frac{1}{n} D_1(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) \le \lim_{\alpha \to 1^-} I_\alpha^{\downarrow\downarrow}(A:B)_\rho \le I_1^{\downarrow\downarrow}(A:B)_\rho,$$
(E.88)

where the first equality in (E.88) follows from (l). Hence, $\lim_{\alpha \to 1^{-}} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = I_{1}^{\downarrow\downarrow}(A:B)_{\rho}$. \Box

Proof of (q). Convexity is inherited from the Petz divergence because, according to the first equality in (3.17) in (l), $(\alpha - 1)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ is the pointwise limit of a sequence of functions that are convex in α .

Proof of (p). Let us define the following two functions.

$$f: \quad (1/2,2) \to \mathbb{R}, \quad \alpha \mapsto I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \tag{E.89}$$

$$g: (1/2,2) \to \mathbb{R}, \quad \alpha \mapsto (\alpha - 1)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$$
 (E.90)

By (o), f is continuous. By (o) and (q), g is convex and continuous. Due to the convexity of g, the left and right derivative of g exist at all points within its domain. Since $f(\alpha) = \frac{1}{\alpha-1}g(\alpha)$, we have for any $\alpha \in (\frac{1}{2}, 1) \cup (1, 2)$

$$\frac{\partial}{\partial \alpha^{-}} f(\alpha) = -\frac{1}{(\alpha - 1)^{2}} g(\alpha) + \frac{1}{\alpha - 1} \frac{\partial}{\partial \alpha^{-}} g(\alpha) = -\frac{1}{\alpha - 1} f(\alpha) + \frac{1}{\alpha - 1} \frac{\partial}{\partial \alpha^{-}} g(\alpha), \quad (E.91a)$$

$$\frac{\partial}{\partial \alpha^{+}} f(\alpha) = -\frac{1}{(\alpha - 1)^{2}} g(\alpha) + \frac{1}{\alpha - 1} \frac{\partial}{\partial \alpha^{+}} g(\alpha) = -\frac{1}{\alpha - 1} f(\alpha) + \frac{1}{\alpha - 1} \frac{\partial}{\partial \alpha^{+}} g(\alpha).$$
(E.91b)

For any $\alpha \in (\frac{1}{2}, 1) \cup (1, 2)$ and any fixed $(\sigma_A, \tau_B) \in \arg\min_{(\sigma'_A, \tau'_B) \in \mathcal{S}(A) \times \mathcal{S}(B)} D_\alpha(\rho_{AB} \| \sigma'_A \otimes \tau'_B)$

$$\frac{\partial}{\partial \alpha^+} f(\alpha) = \lim_{\varepsilon \to 0^+} \frac{1}{\varepsilon} (I_{\alpha+\varepsilon}^{\downarrow\downarrow}(A:B)_{\rho} - I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho})$$
(E.92a)

$$\leq \lim_{\varepsilon \to 0^+} \frac{1}{\varepsilon} (D_{\alpha+\varepsilon}(\rho_{AB} \| \sigma_A \otimes \tau_B) - D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B))$$
(E.92b)

$$= \frac{\partial}{\partial \alpha} D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B) \tag{E.92c}$$

$$= \lim_{\varepsilon \to 0^{-}} \frac{1}{\varepsilon} (D_{\alpha+\varepsilon}(\rho_{AB} \| \sigma_A \otimes \tau_B) - D_{\alpha}(\rho_{AB} \| \sigma_A \otimes \tau_B))$$
(E.92d)

$$\leq \lim_{\varepsilon \to 0^{-}} \frac{1}{\varepsilon} (I_{\alpha+\varepsilon}^{\downarrow\downarrow}(A:B)_{\rho} - I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}) = \frac{\partial}{\partial \alpha^{-}} f(\alpha).$$
(E.92e)

(E.92c) and (E.92d) follow from the differentiability in α of the Petz divergence, see Proposition 2.

We will now prove the continuous differentiability of f on $\alpha \in (1,2)$. Since g is convex, $\frac{\partial}{\partial \alpha^-}g(\alpha) \leq \frac{\partial}{\partial \alpha^+}g(\alpha)$ for all $\alpha \in (1,2)$. Hence, $\frac{\partial}{\partial \alpha^-}f(\alpha) \leq \frac{\partial}{\partial \alpha^+}f(\alpha)$ for all $\alpha \in (1,2)$ due to (E.91). By (E.92), it follows that the left and right derivative of f coincide, so f is differentiable on $\alpha \in (1,2)$ and (3.20) holds. Since f is differentiable and $g(\alpha) = (\alpha - 1)f(\alpha)$ for all $\alpha \in (1,2)$, also g is differentiable on $\alpha \in (1,2)$. Since g is convex, its differentiability on $\alpha \in (1,2)$ implies its continuous differentiability on $\alpha \in (1,2)$. By the product rule, this implies that also f is continuously differentiable on $\alpha \in (1,2)$.

We will now prove the continuous differentiability of f on $\alpha \in (\frac{1}{2}, 1)$. For any $\alpha \in (\frac{1}{2}, 1)$, let

$$(\sigma_A^{(\alpha)}, \tau_B^{(\alpha)}) \in \operatorname*{arg\,min}_{(\sigma_A, \tau_B) \in \mathcal{S}(A) \times \mathcal{S}(B)} D_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B)$$
(E.93)

denote the unique minimizer, see (j). Let us define the function

$$(1/2,1) \to \mathbb{R}, \quad \alpha \mapsto h(\alpha) \coloneqq f(\alpha) + (\alpha - 1) \frac{\partial}{\partial \alpha} D_{\alpha}(\rho_{AB} \| \sigma_A^{(\alpha)} \otimes \tau_B^{(\alpha)}),$$
 (E.94)

where $\sigma_A^{(\alpha)}$ and $\tau_B^{(\alpha)}$ are held fixed. The map $\alpha \mapsto (\sigma_A^{(\alpha)}, \tau_B^{(\alpha)})$ is continuous on $\alpha \in (\frac{1}{2}, 1)$ due to the uniqueness of $(\sigma_A^{(\alpha)}, \tau_B^{(\alpha)})$. By the continuous differentiability of the Petz divergence, see Proposition 2, it follows that h is continuous. For any $\alpha \in (\frac{1}{2}, 1)$,

$$\frac{\partial}{\partial \alpha^{+}} f(\alpha) \leq \frac{\partial}{\partial \alpha} D_{\alpha}(\rho_{AB} \| \sigma_{A}^{(\alpha)} \otimes \tau_{B}^{(\alpha)}) \leq \frac{\partial}{\partial \alpha^{-}} f(\alpha), \tag{E.95}$$

$$\frac{\partial}{\partial \alpha^{-}} g(\alpha) \le h(\alpha) \le \frac{\partial}{\partial \alpha^{+}} g(\alpha).$$
(E.96)

(E.95) follows from (E.92), and it is understood that $\sigma_A^{(\alpha)}$ and $\tau_B^{(\alpha)}$ are held fixed in (E.95). (E.96) follows from (E.91) and (E.95). Therefore, for any $\alpha \in (\frac{1}{2}, 1)$,

$$h(\alpha) = \lim_{\varepsilon \to 0^+} h(\alpha - \varepsilon) \le \lim_{\varepsilon \to 0^+} \frac{\partial}{\partial \alpha^+} g(\alpha - \varepsilon)$$
(E.97a)

$$\leq \frac{\partial}{\partial \alpha^{+}} g(\alpha) \leq \lim_{\varepsilon \to 0^{+}} \frac{\partial}{\partial \alpha^{-}} g(\alpha + \varepsilon) \leq \lim_{\varepsilon \to 0^{+}} h(\alpha + \varepsilon) = h(\alpha).$$
(E.97b)

The first two inequalities in (E.97b) follow from the convexity of g. It follows that all inequalities in (E.97) must be saturated, so $\frac{\partial}{\partial \alpha^+}g(\alpha) = h(\alpha)$ for all $\alpha \in (\frac{1}{2}, 1)$. Since h is continuous, also $\frac{\partial}{\partial \alpha^+}g(\alpha)$ is continuous on $\alpha \in (\frac{1}{2}, 1)$. Since g is convex, the continuity of the right derivative of gimplies that g is differentiable and $g'(\alpha) = \frac{\partial}{\partial \alpha^+}g(\alpha) = h(\alpha)$ for all $\alpha \in (\frac{1}{2}, 1)$. Since h is continuous, this proves that g is continuously differentiable on $\alpha \in (\frac{1}{2}, 1)$. By the product rule, this implies that also f is continuously differentiable on $\alpha \in (\frac{1}{2}, 1)$.

Next, we will prove the continuous differentiability of f at $\alpha = 1$. The combination of (2.14) and a quantum Sibson identity [15, Eq. (B10)] implies that $\lim_{\alpha \to 1^-} (\sigma_A^{(\alpha)}, \tau_B^{(\alpha)}) = (\rho_A, \rho_B)$ and that the limits

$$\lim_{\beta \to 1^{-}} \frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \big|_{\alpha=\beta} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B) \big|_{\alpha=1},$$
(E.98a)

$$\lim_{\beta \to 1^+} \frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \big|_{\alpha=\beta} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \rho_A \otimes \rho_B) \big|_{\alpha=1}$$
(E.98b)

exist. Therefore, they are identical to the left and right derivative of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ at $\alpha = 1$, respectively. By (E.98), the left and right derivative of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ at $\alpha = 1$ coincide. This proves differentiability at $\alpha = 1$. The *continuous* differentiability of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ at $\alpha = 1$ follows from the quantum Sibson identity [15, Eq. (B10)].

It remains to prove that $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ is right-differentiable at $\alpha = 1/2$. Let $(\sigma_A^{(1/2)}, \tau_B^{(1/2)}) := \lim_{\alpha \to 1/2^+} (\sigma_A^{(\alpha)}, \tau_B^{(\alpha)})$. Then, $\rho_{AB} \neq \sigma_A^{(1/2)} \otimes \tau_B^{(1/2)}$, and the limit

$$\lim_{\beta \to 1/2^+} \frac{\mathrm{d}}{\mathrm{d}\alpha} I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \big|_{\alpha=\beta} = \frac{\mathrm{d}}{\mathrm{d}\alpha} D_{\alpha}(\rho_{AB} \| \sigma_A^{(1/2)} \otimes \tau_B^{(1/2)}) \big|_{\alpha=1/2}$$
(E.99)

exists. Therefore, this limit is identical to the right derivative of $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}$ at $\alpha = 1/2$. Note that (E.99) lies in $[0,\infty)$ due to the differentiability and monotonicity in α of the Petz divergence, see Proposition 2.

Proof of (s). Let $\alpha \in [0, \infty)$.

Case 1: $\alpha \in (0, \frac{1}{2}]$. Let $\beta \coloneqq \frac{1}{\alpha} \in [2, \infty)$. Let $|\hat{\sigma}\rangle_A \in A$ be a unit eigenvector of ρ_A corresponding to its largest eigenvalue. By duality (e),

$$\exp((1-\beta)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}) = \sup_{\sigma_A \in \mathcal{S}(A)} \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1}) \le \sup_{\sigma_A \in \mathcal{S}(A)} Q_{\beta}(\rho_A \| \sigma_A^{-1})$$
(E.100)

$$= \sup_{\sigma_A \in \mathcal{S}(A)} \operatorname{tr}[\rho_A^\beta \sigma_A^{\beta-1}] = \|\rho_A\|_{\infty}^{\beta}$$
(E.101)

$$= \widetilde{Q}_{\beta}(\rho_A \| |\hat{\sigma}\rangle\!\langle \hat{\sigma} |_A^{-1}) \le \sup_{\sigma_A \in \mathcal{S}(A)} \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1}),$$
(E.102)

so all inequalities must be saturated. Therefore, $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \frac{\beta}{\beta-1}H_{\infty}(A)_{\rho} = \frac{1}{1-\alpha}H_{\infty}(A)_{\rho}$.

Case 2: $\alpha \in (\frac{1}{2}, 1)$. Let $\beta \coloneqq \frac{1}{\alpha} \in (1, 2)$. Let $\hat{\sigma}_A \coloneqq \rho_A^{\frac{1}{2\alpha-1}} / \operatorname{tr}[\rho_A^{\frac{1}{2\alpha-1}}] = \rho_A^{\frac{\beta}{2-\beta}} / \operatorname{tr}[\rho_A^{\frac{\beta}{2-\beta}}]$. By duality (e),

$$\exp((1-\beta)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}) = \sup_{\sigma_A \in \mathcal{S}(A)} \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1}) \le \sup_{\sigma_A \in \mathcal{S}(A)} Q_{\beta}(\rho_A \| \sigma_A^{-1})$$
(E.103)

$$= \sup_{\sigma_A \in \mathcal{S}(A)} \operatorname{tr}[\rho_A^\beta \sigma_A^{\beta-1}] = \|\rho_A^\beta\|_{\frac{1}{2-\beta}} = \operatorname{tr}[\rho_A^{\frac{\beta}{2-\beta}}]^{2-\beta}$$
(E.104)

$$= \widetilde{Q}_{\beta}(\rho_A \| \hat{\sigma}_A^{-1}) \le \sup_{\sigma_A \in \mathcal{S}(A)} \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1}),$$
(E.105)

so all inequalities must be saturated. In (E.104), we have used the variational characterization of the Schatten norms [45, Lemma 3.2]. Therefore, $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = 2H_{\frac{\beta}{2-\beta}}(A)_{\rho} = 2H_{\frac{1}{2\alpha-1}}(A)_{\rho}$.

Case 3: $\alpha \in (1,\infty)$. Let $\beta \coloneqq \frac{1}{\alpha} \in (0,1)$. Let $\hat{\sigma}_A \coloneqq \rho_A^{\frac{1}{2\alpha-1}} / \operatorname{tr}[\rho_A^{\frac{1}{2\alpha-1}}] = \rho_A^{\frac{\beta}{2-\beta}} / \operatorname{tr}[\rho_A^{\frac{\beta}{2-\beta}}]$. By duality (e),

$$\exp((1-\beta)I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho}) = \inf_{\substack{\sigma_A \in \mathcal{S}(A):\\\rho_A \ll \sigma_A}} \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1}) \ge \inf_{\substack{\sigma_A \in \mathcal{S}(A):\\\rho_A \ll \sigma_A}} Q_{\beta}(\rho_A \| \sigma_A^{-1})$$
(E.106)

$$= \inf_{\substack{\sigma_A \in \mathcal{S}(A):\\\rho_A \ll \sigma_A}} \operatorname{tr}[\rho_A^\beta \sigma_A^{\beta-1}] = \|\rho_A^\beta\|_{\frac{1}{2-\beta}} = \operatorname{tr}[\rho_A^{\frac{\beta}{2-\beta}}]^{2-\beta}$$
(E.107)

$$= \widetilde{Q}_{\beta}(\rho_A \| \hat{\sigma}_A^{-1}) \ge \inf_{\substack{\sigma_A \in \mathcal{S}(A):\\\rho_A \ll \sigma_A}} \widetilde{Q}_{\beta}(\rho_A \| \sigma_A^{-1}),$$
(E.108)

so all inequalities must be saturated. In (E.107), we have used the variational characterization of the Schatten quasi-norms [45, Lemma 3.2]. Therefore, $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = 2H_{\frac{\beta}{2-\beta}}(A)_{\rho} = 2H_{\frac{1}{2\alpha-1}}(A)_{\rho}$.

Case 4: $\alpha \in \{0, 1\}$. Then the assertion follows from the previous cases due to the continuity in α (o).

Proof of (t). The assertion regarding σ_A and τ_B can be verified by inserting σ_A and τ_B into (3.22). The rest of the assertion follows from (s).

Proof of (g). Let $|\rho\rangle_{ABC} \in ABC$ be such that $\operatorname{tr}_C[|\rho\rangle\langle\rho|_{ABC}] = \rho_{AB}$.

Let $\alpha \in [0, \infty)$. Then $\frac{1}{\alpha} \in (0, \infty]$. By duality of the minimized generalized PRMI, see Proposition 12 (d),

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \leq I_{\alpha}^{\downarrow}(\rho_{AB} \| \rho_A^0 / r_A) = -\widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| (\rho_A^0 / r_A)^{-1} \otimes \rho_C)$$
(E.109a)

$$= 2\log r_A - \widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_A^0 / r_A \otimes \rho_C) \le 2\log r_A.$$
(E.109b)

The last inequality follows from the non-negativity of the sandwiched divergence.

Let now $\alpha \in [\frac{1}{2}, 2]$. Let $\gamma \coloneqq \frac{1}{2\alpha - 1} \in [\frac{1}{3}, \infty]$.

First, suppose spec $(\rho_A) \subseteq \{0, 1/r_A\}$ and $H(A|B)_{\rho} = -\log r_A$. Then $\rho_A = \rho_A^0/r_A$ and $\rho_{AC} = \rho_A \otimes \rho_C$. By (s), $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = 2H_{\gamma}(A)_{\rho} = 2\log r_A$.

Now, suppose $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = 2\log r_A$ instead. Then the inequalities in (E.109) must be saturated, so $\widetilde{D}_{\frac{1}{\alpha}}(\rho_{AC} \| \rho_A^0/r_A \otimes \rho_C) = 0$. Since $\frac{1}{\alpha} \geq \frac{1}{2}$, it follows from the positive definiteness of the sandwiched divergence that $\rho_{AC} = \rho_A^0/r_A \otimes \rho_C$. Hence, $\rho_A = \rho_A^0/r_A$, which implies that $\operatorname{spec}(\rho_A) \subseteq \{0, 1/r_A\}$. Therefore, $H(A|B)_{\rho} = -H(A|C)_{\rho} = -H(A)_{\rho} = -\log r_A$.

Let now $\alpha \in [0, \frac{1}{2})$. By (b) and the expression for pure states in (t), we have

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} \le I_{\alpha}^{\downarrow\downarrow}(A:BC)_{|\rho\rangle\langle\rho|} = \frac{1}{1-\alpha} H_{\infty}(A)_{\rho} \le \frac{1}{1-\alpha} \log r_A < 2\log r_A.$$
(E.110)

Proof of (u). Let $\alpha \in [0, \infty)$.

$$I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = \inf_{\sigma_A \in \mathcal{S}(A)} I_{\alpha}^{\downarrow}(\rho_{AB} \| \sigma_A)$$
(E.111)

$$= \inf_{\substack{\sigma_A \in \mathcal{S}(A) \\ \exists (t_y)_{y \in \mathcal{Y}} \in [0,1]^{\times |\mathcal{Y}|:} \\ \tau_B = \sum_{y \in \mathcal{Y}} t_y |b_y\rangle \langle b_y|_B}} D_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B)$$
(E.112)

$$= \inf_{\substack{\tau_B \in \mathcal{S}(B):\\ \exists (t_y)_{y \in \mathcal{Y}} \in [0,1]^{\times |\mathcal{Y}|:}\\ \tau_B = \sum_t t_y |b_y \rangle \langle b_y |_B}} I_{\alpha}^{\downarrow}(\rho_{AB} \| \tau_B)$$
(E.113)

$$= \inf_{\substack{\tau_B \in \mathcal{S}(B): \\ \exists (t_y)_{y \in \mathcal{Y}} \in [0,1]^{\times |\mathcal{Y}|: } \exists (s_x)_{x \in \mathcal{X}} \in [0,1]^{\times |\mathcal{X}|: } \\ \tau_B = \sum_{y \in \mathcal{Y}} t_y |b_y\rangle \langle b_y|_B \sigma_A = \sum_{x \in \mathcal{X}} s_x |a_x\rangle \langle a_x|_A} D_\alpha(\rho_{AB} \| \sigma_A \otimes \tau_B) = I_\alpha^{\downarrow\downarrow}(X:Y)_P \quad (E.114)$$

Above, we have used Proposition 12 (q) twice: for (E.112), and for the first equality in (E.114). \Box

Proof of (v). Let $\alpha \in [0,\infty)$. Let Y be a random variable over $\mathcal{Y} := \mathcal{X}$ and let $P_{XY}(x,y) :=$ $P_X(x)\delta_{x,y}$ for all $x \in \mathcal{X}, y \in \mathcal{Y}$. Then $\rho_{AB} = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} P_{XY}(x,y) |a_x, b_y| \langle a_x, b_y | AB$. By (u), $I_{\alpha}^{\downarrow\downarrow}(A:B)_{\rho} = I_{\alpha}^{\downarrow\downarrow}(X:Y)_{P}$. As shown in [1, Lemma 11], $I_{\alpha}^{\downarrow\downarrow}(X:Y)_{P} = \frac{\alpha}{1-\alpha}H_{\infty}(A)_{\rho}$ if $\alpha \in [0, \frac{1}{2}]$, and $I^{\downarrow\downarrow}_{\alpha}(X:Y)_P = H_{\frac{\alpha}{2\alpha-1}}(A)_{\rho}$ if $\alpha \in (\frac{1}{2},\infty)$. The assertion regarding σ_A and τ_B can be verified by inserting σ_A and τ_B into (3.24).

Appendix F: Proofs for Section 3 B

Lemmas for Theorem 8 1.

Let us define the following function of $\mu \in [0, \infty)$ for any $\rho_{AB} \in \mathcal{S}(AB), n \in \mathbb{N}_{>0}$.

$$\hat{\alpha}_{n,\rho}'(\mu) = \min_{\substack{T_{A^n B^n} \in \mathcal{L}(A^n B^n):\\ 0 \le T_{A^n B^n} \le 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \max_{\substack{\sigma_{A^n} \in \mathcal{S}_{\operatorname{sym}}(A^{\otimes n}),\\ \tau_{B^n} \in \mathcal{S}(B^n)}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T_{A^n B^n}^n] \le \mu \}$$
(F.1)

The following lemma describes some basic properties of the functions $\hat{\alpha}_{n,\rho}^{\text{iid}}, \hat{\alpha}_{n,\rho}, \hat{\alpha}_{n,\rho}^{\text{ind}}, \hat{\alpha}_{n,\rho}'$, as defined in (3.26), (3.27), (3.33), (F.1).

Lemma 16 (Minimum type-I errors). Let $\rho_{AB} \in \mathcal{S}(AB), n \in \mathbb{N}_{>0}$. Then all of the following hold.

- (a) The function $[0,\infty) \to \mathbb{R}, \mu \mapsto \hat{\alpha}_{n,\rho}(\mu)$ is monotonically decreasing, and the same is true for
- $\hat{\alpha}_{n,\rho}^{\text{iid}}, \hat{\alpha}_{n,\rho}^{\text{ind}}, \text{ and } \hat{\alpha}_{n,\rho}'.$ $(b) \quad 0 \leq \hat{\alpha}_{n,\rho}^{\text{iid}}(\mu) \leq \hat{\alpha}_{n,\rho}(\mu) \leq \hat{\alpha}_{n,\rho}'(\mu) \leq \hat{\alpha}_{n,\rho}^{\text{ind}}(\mu) \leq \max(0, 1-\mu) \leq 1 \text{ for all } \mu \in [0,\infty).$
- (c) $\hat{\alpha}_{n,\rho}(\mu) = \hat{\alpha}'_{n,\rho}(\mu)$ for all $\mu \in [0,\infty)$.

Proof of (a). This assertion follows directly from the definitions of the functions.

Proof of (b). Let $\mu \in [0,\infty)$. $\mathcal{S}_{sym}(A^{\otimes n}) \otimes \mathcal{S}(B^{\otimes n}) \subseteq \mathcal{S}(A^{\otimes n}) \otimes \mathcal{S}(B^{\otimes n})$ implies that $\hat{\alpha}'_{n,\rho}(\mu) \leq \mathcal{S}(A^{\otimes n})$ $\hat{\alpha}_{n,\rho}^{\mathrm{ind}}(\mu)$. $\mathcal{S}_{\mathrm{sym}}(A^{\otimes n}) \otimes \mathcal{S}_{\mathrm{sym}}(B^{\otimes n}) \subseteq \mathcal{S}_{\mathrm{sym}}(A^{\otimes n}) \otimes \mathcal{S}(B^n)$ implies that $\hat{\alpha}_{n,\rho}(\mu) \leq \hat{\alpha}_{n,\rho}'(\mu)$. $\mathcal{S}(A)^{\otimes n} \otimes \mathcal{S}(B^n)$ $\mathcal{S}(B)^{\otimes n} \subseteq \mathcal{S}_{\text{sym}}(A^{\otimes n}) \otimes \mathcal{S}_{\text{sym}}(B^{\otimes n}) \text{ implies that } \hat{\alpha}_{n,\rho}^{\text{iid}}(\mu) \leq \hat{\alpha}_{n,\rho}(\mu).$

Let $T_{A^nB^n}^n \in \mathcal{L}(A^nB^n)$ be in the feasible set of the optimization problem that defines $\hat{\alpha}_{n,\rho}^{\text{iid}}$. Then, $0 \leq T_{A^n B^n}^n \leq 1$. Hence, $1 - T_{A^n B^n}^n \geq 0$, which implies that $\hat{\alpha}_{n,\rho}^{\text{iid}}(\mu) \geq 0$.

If $\mu \in [0,1]$, then choosing the test $T^n_{A^nB^n} \coloneqq \mu 1$ implies that $\hat{\alpha}^{\text{ind}}_{n,\rho}(\mu) \leq 1-\mu$. If $\mu \in (1,\infty)$, then choosing the test $T^n_{A^nB^n} \coloneqq 1$ implies that $\hat{\alpha}^{\text{ind}}_{n,\rho}(\mu) \leq 0$. Hence, $\hat{\alpha}^{\text{ind}}_{n,\rho}(\mu) \leq \max(0, 1-\mu)$.

Proof of (c). Let $\mu \in [0, \infty)$. By (b), it suffices to prove that $\hat{\alpha}'_{n,\rho}(\mu) \leq \hat{\alpha}_{n,\rho}(\mu)$.

Let $T_{A^nB^n}^n \in \mathcal{L}(A^nB^n)$ be in the feasible set of the optimization problem that defines $\hat{\alpha}_{n,\rho}(\mu)$. Let $\hat{T}^n_{A^nB^n} \coloneqq \frac{1}{|S_n|} \sum_{\pi \in S_n} U(\pi)_{A^n} \otimes U(\pi)_{B^n} T^n_{A^nB^n} U(\pi)^{\dagger}_{A^n} \otimes U(\pi)^{\dagger}_{B^n}$. Then, for any $\sigma_{A^n} \in \mathcal{T}^n_{A^nB^n}$ $\mathcal{S}_{\mathrm{sym}}(A^{\otimes n}), \tau_{B^n} \in \mathcal{S}_{\mathrm{sym}}(B^{\otimes n})$

$$\operatorname{tr}[\sigma_{A^{n}} \otimes \tau_{B^{n}} T^{n}_{A^{n}B^{n}}] = \frac{1}{|S_{n}|} \sum_{\pi \in S_{n}} \operatorname{tr}[(U(\pi)^{\dagger}_{A^{n}} \sigma_{A^{n}} U(\pi)_{A^{n}}) \otimes (U(\pi)^{\dagger}_{B^{n}} \tau_{B^{n}} U(\pi)_{B^{n}}) T^{n}_{A^{n}B^{n}}]$$
(F.2)

$$=\frac{1}{|S_n|}\sum_{\pi\in S_n}\operatorname{tr}[\sigma_{A^n}\otimes\tau_{B^n}U(\pi)_{A^n}\otimes U(\pi)_{B^n}T^n_{A^nB^n}U(\pi)^{\dagger}_{A^n}\otimes U(\pi)^{\dagger}_{B^n}] \quad (F.3)$$

$$= \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} \hat{T}^n_{A^n B^n}]. \tag{F.4}$$

Hence,

$$\max_{\substack{\sigma_{A^n} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T^n_{A^n B^n}] = \max_{\substack{\sigma_{A^n} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \\ \tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T^n_{A^n B^n}].$$
(F.5)

Since $\rho_{AB}^{\otimes n} \in S_{\text{sym}}((AB)^{\otimes n})$, $\text{tr}[\rho_{AB}^{\otimes n}(1-T_{A^nB^n}^n)] = \text{tr}[\rho_{AB}^{\otimes n}(1-\hat{T}_{A^nB^n}^n)]$. Since $\hat{T}_{A^nB^n}^n$ is permutation invariant, it follows that

$$\hat{\alpha}_{n,\rho}(\mu) = \min_{\substack{T_{A^n B^n} \in \mathcal{L}_{\mathrm{sym}}((AB)^{\otimes n}):\\ 0 \le T_{A^n B^n} \le 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \max_{\substack{\sigma_{A^n} \in \mathcal{S}_{\mathrm{sym}}(A^{\otimes n}),\\ \tau_{B^n} \in \mathcal{S}_{\mathrm{sym}}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T_{A^n B^n}^n] \le \mu \}.$$
(F.6)

Let $\widetilde{T}^n_{A^nB^n} \in \mathcal{L}_{\text{sym}}((AB)^{\otimes n})$ be positive semidefinite. Then, for all $\sigma_{A^n} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}), \tau_{B^n} \in \mathcal{S}(B^n)$

$$\operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} \widetilde{T}^n_{A^n B^n}] = \frac{1}{|S_n|} \sum_{\pi \in S_n} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} (U(\pi)_{A^n} \otimes U(\pi)_{B^n} \widetilde{T}^n_{A^n B^n} U(\pi)^{\dagger}_{A^n} \otimes U(\pi)^{\dagger}_{B^n})] \quad (F.7)$$

$$=\frac{1}{|S_n|}\sum_{\pi\in S_n} \operatorname{tr}[\underbrace{(U(\pi)_{A^n}^{\dagger}\sigma_{A^n}U(\pi)_{A^n})}_{=\sigma_{A^n}} \otimes (U(\pi)_{B^n}^{\dagger}\tau_{B^n}U(\pi)_{B^n})\widetilde{T}_{A^nB^n}^n]$$
(F.8)

$$= \operatorname{tr}[\sigma_{A^{n}} \otimes \underbrace{\frac{1}{|S_{n}|} \sum_{\pi \in S_{n}} (U(\pi)_{B^{n}}^{\dagger} \tau_{B^{n}} U(\pi)_{B^{n}}) \widetilde{T}_{A^{n}B^{n}}^{n}]}_{= \mathfrak{s}_{A^{n}}}.$$
(F.9)

$$\in \mathcal{S}_{\mathrm{sym}}(B^{\otimes n})$$

Hence,

$$\max_{\substack{\sigma_{A^n} \in \mathcal{S}_{sym}(A^{\otimes n}), \\ \tau_{B^n} \in \mathcal{S}(B^n)}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} \widetilde{T}^n_{A^n B^n}] = \max_{\substack{\sigma_{A^n} \in \mathcal{S}_{sym}(A^{\otimes n}), \\ \tau_{B^n} \in \mathcal{S}_{sym}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} \widetilde{T}^n_{A^n B^n}].$$
(F.10)

We conclude that

$$\begin{aligned} \hat{\alpha}_{n,\rho}^{\prime}(\mu) &\leq \min_{\substack{T_{A^{n}B^{n}} \in \mathcal{L}_{\text{sym}}((AB)^{\otimes n}):\\ 0 \leq T_{A^{n}B^{n}} \leq 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^{n}B^{n}}^{n})] : \max_{\substack{\sigma_{A^{n}} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}),\\ \tau_{B^{n}} \in \mathcal{S}(B^{n})}} \operatorname{tr}[\sigma_{A^{n}} \otimes \tau_{B^{n}} T_{A^{n}B^{n}}^{n}] \leq \mu \} \end{aligned}$$

$$= \min_{\substack{T_{A^{n}B^{n}} \in \mathcal{L}_{\text{sym}}((AB)^{\otimes n}):\\ 0 \leq T_{A^{n}B^{n}}^{n} \leq 1}} \{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^{n}B^{n}}^{n})] : \max_{\substack{\sigma_{A^{n}} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}),\\ \tau_{B^{n}} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^{n}} \otimes \tau_{B^{n}} T_{A^{n}B^{n}}^{n}] \leq \mu \} \end{aligned}$$

$$= \hat{\alpha}_{n,\rho}(\mu). \tag{F.12}$$

(F.12) follows from (F.10). (F.13) follows from (F.6).

The following lemma is a variant of [9, Lemma 20] that does not require ϕ to be differentiable.

Lemma 17 (A property of convex functions). Let $a, b \in \mathbb{R}$ be such that $0 \leq a < b$ and let $\phi : (a, b) \to \mathbb{R}$ be a convex function. Let $\psi(s) \coloneqq s \frac{\partial}{\partial s^+} \phi(s) - \phi(s)$ for all $s \in (a, b)$. Then, ψ is monotonically increasing.

Proof. Let $s_0, s_1 \in \mathbb{R}$ be such that $a < s_0 < s_1 < b$. Then,

$$\psi(s_1) - \psi(s_0) = s_1 \frac{\partial}{\partial s^+} \phi(s_1) - s_0 \frac{\partial}{\partial s^+} \phi(s_0) - (\phi(s_1) - \phi(s_0)) \tag{F.14}$$

$$\geq s_1 \frac{\partial}{\partial s^+} \phi(s_1) - s_0 \frac{\partial}{\partial s^+} \phi(s_0) - (s_1 - s_0) \frac{\partial}{\partial s^+} \phi(s_1) \tag{F.15}$$

$$= s_0 \left(\frac{\partial}{\partial s^+} \phi(s_1) - \frac{\partial}{\partial s^+} \phi(s_0) \right) \ge 0, \tag{F.16}$$

where the inequalities follow from the convexity of ϕ .

2. Proof of Theorem 8

First, we prove the bounds on $R_{1/2}$. It follows from Theorem 7 (q) that $I_s^{\downarrow\downarrow}(A:B)_{\rho}$ is rightdifferentiable on $s \in (0,1)$. By the monotonicity of the doubly minimized PRMI in the Rényi order, see Theorem 7 (n),

$$\frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_\rho \ge 0 \qquad \forall s \in (0,1).$$
(F.17)

For any $s \in (0,1)$ and any fixed $(\sigma_A, \tau_B) \in \arg\min_{(\sigma'_A, \tau'_B) \in \mathcal{S}(A) \times \mathcal{S}(B)} D_s(\rho_{AB} \| \sigma'_A \otimes \tau'_B)$, the right derivative at s is upper bounded as $\frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_{\rho} \leq \frac{\partial}{\partial s^+} D_s(\rho_{AB} \| \sigma_A \otimes \tau_B)$, which implies that $0 \leq \liminf_{s \to 0^+} \frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_{\rho} \leq \limsup_{s \to 0^+} \frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_{\rho} < \infty$. Hence,

$$I_0^{\downarrow\downarrow}(A:B)_\rho = \lim_{s \to 0^+} (I_s^{\downarrow\downarrow}(A:B)_\rho - s(1-s)\frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_\rho)$$
(F.18)

$$\leq \left(I_s^{\downarrow\downarrow}(A:B)_\rho - s(1-s)\frac{\partial}{\partial s^+}I_s^{\downarrow\downarrow}(A:B)_\rho\right)\Big|_{s=1/2} = R_{1/2} \tag{F.19}$$

$$\leq I_{1/2}^{\downarrow\downarrow}(A:B)_{\rho}.\tag{F.20}$$

(F.19) follows from Lemma 17 due to the convexity of $(s-1)I_s^{\downarrow\downarrow}(A:B)_{\rho}$ on $s \in (0,1)$, see Theorem 7 (q). (F.20) follows from (F.17).

Next, we derive the bounds on the right-hand side of (3.30). For any $R \in (0, \infty)$

$$0 = \lim_{s \to 1^{-}} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R) \le \sup_{s \in (\frac{1}{2},1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R)$$
(F.21a)
$$\le \sup_{s \in (\frac{1}{2},1)} \frac{1-s}{s} (I_1^{\downarrow\downarrow}(A:B)_{\rho} - R) = \max(0, I(A:B)_{\rho} - R),$$
(F.21b)

where we have used the monotonicity and continuity of the doubly minimized PRMI in the Rényi order and $I_1^{\downarrow\downarrow}(A:B)_{\rho} = I(A:B)_{\rho}$, see Theorem 7 (m), (n), (o). On the one hand, the bounds in (F.21) imply that for any $R \in [I(A:B)_{\rho}, \infty)$, the right-hand side of (3.30) vanishes. On the other hand, if $R \in (0, I(A:B)_{\rho})$, then the right-hand side of (3.30) is strictly positive due to Theorem 7 (m), (n), (o).

We will now prove the equality in (3.30). The proof of (3.30) is divided into two parts: a proof of achievability for $\hat{\alpha}_{n,\rho}$ and a proof of optimality for $\hat{\alpha}_{n,\rho}^{\text{iid}}$. The assertion in Theorem 8 follows from these two parts because $\hat{\alpha}_{n,\rho}^{\text{iid}}(\mu) \leq \hat{\alpha}_{n,\rho}(\mu)$ for all $\mu \in [0, \infty)$, see Lemma 16 (b).

a. Proof of achievability

Below, we will prove that for any $R \in (0, \infty)$

$$\liminf_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}(e^{-nR}) \ge \sup_{s \in (0,1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R)$$
(F.22)

$$\geq \sup_{s \in (\frac{1}{2}, 1)} \frac{1 - s}{s} (I_s^{\downarrow\downarrow}(A : B)_{\rho} - R).$$
 (F.23)

The inequality in (F.23) is trivially true, so it suffices to prove (F.22).

Proof. Let $R \in (0, \infty)$ and $s \in (0, 1)$ be arbitrary but fixed. For all $n \in \mathbb{N}_{>0}$, we define

$$\lambda_n \coloneqq \frac{1}{s} \left(\log g_{n,d_A} + \log g_{n,d_B} + nR - (1-s)D_s(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) \right)$$
(F.24)

and the test $T_{A^nB^n}^n \coloneqq \{\rho_{AB}^{\otimes n} \ge e^{\lambda_n} \omega_{A^n}^n \otimes \omega_{B^n}^n\}$. For this test holds

$$\max_{\substack{\sigma_{A^n} \in \mathcal{S}_{\text{sym}}(A^{\otimes n}),\\ \tau_{B^n} \in \mathcal{S}_{\text{sym}}(B^{\otimes n})}} \operatorname{tr}[\sigma_{A^n} \otimes \tau_{B^n} T^n_{A^n B^n}] \le g_{n, d_A} g_{n, d_B} \operatorname{tr}[\omega^n_{A^n} \otimes \omega^n_{B^n} \{ e^{-\lambda_n} \rho^{\otimes n}_{A^B} \ge \omega^n_{A^n} \otimes \omega^n_{B^n} \}] \quad (F.25a)$$

$$\leq g_{n,d_A}g_{n,d_B}\operatorname{tr}[(e^{-\lambda_n}\rho_{AB}^{\otimes n})^s(\omega_{A^n}^n\otimes\omega_{B^n}^n)^{1-s}]$$
(F.25b)

$$= g_{n,d_A}g_{n,d_B}e^{-s\lambda_n}\exp\left(-(1-s)D_s(\rho_{AB}^{\otimes n}\|\omega_{A^n}^n\otimes\omega_{B^n}^n)\right) \quad (F.25c)$$

$$= e^{-nR}.$$
 (F.25d)

(F.25a) follows from Proposition 1 (b). (F.25b) follows from [15, Eq. (2.2)]. (F.25d) follows from (F.24). Furthermore,

$$\operatorname{tr}[\rho_{AB}^{\otimes n}(1-T_{A^{n}B^{n}}^{n})] = \operatorname{tr}[\rho_{AB}^{\otimes n}\{\rho_{AB}^{\otimes n} < e^{\lambda_{n}}\omega_{A^{n}}^{n} \otimes \omega_{B^{n}}^{n}\}]$$
(F.26a)

$$\leq \operatorname{tr}[(\rho_{AB}^{\otimes n})^{s}(e^{\lambda_{n}}\omega_{A^{n}}^{n}\otimes\omega_{B^{n}}^{n})^{1-s}]$$
(F.26b)

$$= e^{\lambda_n (1-s)} \exp\left(-(1-s)D_s(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n)\right)$$
(F.26c)

$$= \exp\left(\frac{1-s}{s}\left(\log g_{n,d_A} + \log g_{n,d_B} - \left(D_s(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) - nR\right)\right)\right).$$
(F.26d)

(F.26b) follows from [15, Eq. (2.2)]. (F.26d) follows from (F.24). We conclude that

$$\liminf_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}(e^{-nR}) \ge \liminf_{n \to \infty} -\frac{1}{n} \log \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)]$$
(F.27a)

$$\geq \frac{1-s}{s} \left(\liminf_{n \to \infty} \frac{1}{n} D_s(\rho_{AB}^{\otimes n} \| \omega_{A^n}^n \otimes \omega_{B^n}^n) - R \right)$$
(F.27b)

$$=\frac{1-s}{s}(I_s^{\downarrow\downarrow}(A:B)_{\rho}-R).$$
(F.27c)

(F.27a) follows from (F.25). (F.27b) follows from (F.26) and Proposition 1 (b). (F.27c) follows from Theorem 7 (l). Since $s \in (0,1)$ can be chosen arbitrarily, the assertion in (F.22) follows from (F.27).

b. Proof of optimality

Below, we will prove that for any $R \in (R_{1/2}, \infty)$

$$\limsup_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}^{\text{iid}}(e^{-nR}) \le \sup_{s \in (\frac{1}{2},1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R).$$
(F.28)

Proof. Let $R \in (R_{1/2}, \infty)$ be arbitrary but fixed. Since $R > R_{1/2} \ge 0$, the converse quantum Hoeffding bound [14, 29] implies that for any $(\sigma_A, \tau_B) \in \mathcal{S}(A) \times \mathcal{S}(B)$

$$\limsup_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}^{\text{iid}}(e^{-nR}) \le \sup_{s \in (0,1)} \frac{1-s}{s} (D_s(\rho_{AB} \| \sigma_A \otimes \tau_B) - R).$$
(F.29)

Case 1: $R \in [I(A:B)_{\rho}, \infty)$. Then,

$$\limsup_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}^{\text{iid}}(e^{-nR}) \le \sup_{s \in (0,1)} \frac{1-s}{s} (I_s^{\uparrow\uparrow}(A:B)_{\rho} - R)$$
(F.30)

$$= 0$$
 (F.31)

$$= \sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R).$$
 (F.32)

(F.30) follows from the evaluation of (F.29) for $(\sigma_A, \tau_B) = (\rho_A, \rho_B)$. (F.31) follows from $I_1^{\uparrow\uparrow}(A : B)_{\rho} = I(A : B)_{\rho} \leq R$ and the monotonicity in the Rényi order of the non-minimized PRMI, see Proposition 3 (h), (i). (F.32) follows from (F.21).

Case 2: $R \in (R_{1/2}, I(A : B)_{\rho})$. Let us define the following functions of $s \in (\frac{1}{2}, 1)$.

$$\phi(s) \coloneqq (s-1)I_s^{\downarrow\downarrow}(A:B)_\rho \tag{F.33}$$

$$\psi(s) \coloneqq s\phi'(s) - \phi(s) = I_s^{\downarrow\downarrow}(A:B)_\rho - s(1-s)\frac{\mathrm{d}}{\mathrm{d}s}I_s^{\downarrow\downarrow}(A:B)_\rho \tag{F.34}$$

$$g(s) := \frac{1}{s}((s-1)R - \phi(s)) = \frac{1-s}{s}(I_s^{\downarrow\downarrow}(A:B)_{\rho} - R)$$
(F.35)

Note that $\lim_{s\to 1/2^+} \psi(s) = R_{1/2}$ and $\lim_{s\to 1^-} \psi(s) = I_1^{\downarrow\downarrow}(A:B)_{\rho} = I(A:B)_{\rho}$ due to Theorem 7 (m). ϕ is convex and continuously differentiable due to Theorem 7 (p), (q). Since ϕ is continuously differentiable, ψ is continuous. The convexity of ϕ implies that ψ is monotonically increasing, see [9, Lemma 20] or Lemma 17. Since ϕ is continuously differentiable, also g is continuously differentiable, so $g'(s) = \frac{1}{s^2}(R - \psi(s))$ is continuous. Since ψ is monotonically increasing, g' is monotonically decreasing. Furthermore, $\lim_{s\to 1/2^+} g'(s) = 4(R - R_{1/2}) > 0$ and $\lim_{s\to 1^-} g'(s) = R - I(A:B)_{\rho} < 0$. Thus, g'(s) = 0 iff g is maximal at s.

Let $\hat{s} \in (\frac{1}{2}, 1)$ be such that $g'(\hat{s}) = 0$ and

s

$$\sup_{\in (\frac{1}{2},1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R) = \frac{1-\hat{s}}{\hat{s}} (I_{\hat{s}}^{\downarrow\downarrow}(A:B)_{\rho} - R).$$
(F.36)

Such an \hat{s} exists because $R \in (R_{1/2}, I(A : B)_{\rho})$, see also [9, Lemma 21].

Let $(\sigma_A^{(\hat{s})}, \tau_B^{(\hat{s})}) \in \arg\min_{(\sigma_A, \tau_B) \in \mathcal{S}(A) \times \mathcal{S}(B)} D_{\hat{s}}(\rho_{AB} \| \sigma_A \otimes \tau_B)$ be the unique minimizer, see Theorem 7 (j). Let us define the following functions of $s \in (0, 1)$.

$$\bar{\phi}(s) \coloneqq (s-1)D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) \tag{F.37}$$

$$\bar{\psi}(s) \coloneqq s\bar{\phi}'(s) - \bar{\phi}(s) = D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) - s(1-s)\frac{\mathrm{d}}{\mathrm{d}s} D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) \tag{F.38}$$

$$\bar{g}(s) \coloneqq \frac{1}{s}((s-1)R - \bar{\phi}(s)) = \frac{1-s}{s}(D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) - R)$$
(F.39)

 $\bar{\phi}$ is convex and continuously differentiable due to Proposition 2. The convexity of $\bar{\phi}$ implies that $\bar{\psi}$ is monotonically increasing, see [9, Lemma 20] or Lemma 17. Since $\bar{\phi}$ is continuously differentiable, also \bar{g} is continuously differentiable. Hence, $\bar{g}'(s) = \frac{1}{s^2}(R - \bar{\psi}(s))$ is continuous and monotonically decreasing. Therefore, if $\bar{g}'(s) = 0$ for some $s \in (0, 1)$, then \bar{g} is maximal at s.

decreasing. Therefore, if $\bar{g}'(s) = 0$ for some $s \in (0, 1)$, then \bar{g} is maximal at s. By the definition of $(\sigma_A^{(\hat{s})}, \tau_B^{(\hat{s})})$, it is clear that $D_{\hat{s}}(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) = I_{\hat{s}}^{\downarrow\downarrow}(A:B)_{\rho}$. According to Theorem 7 (p), $\frac{d}{ds} D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})})|_{s=\hat{s}} = \frac{d}{ds} I_s^{\downarrow\downarrow}(A:B)_{\rho}|_{s=\hat{s}}$. Importantly, this implies that ϕ and $\bar{\phi}$ are the same up to first order at \hat{s} , i.e.,

$$\bar{\phi}(\hat{s}) = (\hat{s} - 1)I_{\hat{s}}^{\downarrow\downarrow}(A:B)_{\rho} = \phi(\hat{s}) \quad \text{and} \quad \bar{\phi}'(\hat{s}) = \phi'(\hat{s}).$$
 (F.40)

(F.40) implies that $\bar{g}'(\hat{s}) = g'(\hat{s}) = 0$, so \bar{g} achieves its maximum at \hat{s} . Therefore,

$$\sup_{s \in (0,1)} \frac{1-s}{s} (D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) - R) = \frac{1-\hat{s}}{\hat{s}} (D_{\hat{s}}(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) - R)$$
(F.41a)

$$=\frac{1-\hat{s}}{\hat{s}}(I_{\hat{s}}^{\downarrow\downarrow}(A:B)_{\rho}-R).$$
 (F.41b)

By evaluating (F.29) for $(\sigma_A, \tau_B) = (\sigma_A^{(\hat{s})}, \tau_B^{(\hat{s})})$, it follows that

$$\limsup_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}^{\text{iid}}(e^{-nR}) \le \sup_{s \in (0,1)} \frac{1-s}{s} (D_s(\rho_{AB} \| \sigma_A^{(\hat{s})} \otimes \tau_B^{(\hat{s})}) - R).$$
(F.42)

The assertion then follows from the combination of (F.36), (F.41), and (F.42).

3. Proof of Corollary 10

Proof. Let $\rho_{AB} \in \mathcal{S}(AB)$ and let $R \in (-\infty, I(A:B)_{\rho})$. By Lemma 16 (b),

$$0 \le \liminf_{n \to \infty} \hat{\alpha}_{n,\rho}(e^{-nR}) \le \limsup_{n \to \infty} \hat{\alpha}_{n,\rho}(e^{-nR}).$$
(F.43)

We will now prove that $\lim_{n\to\infty} \hat{\alpha}_{n,\rho}(e^{-nR}) = 0$ by cases.

Case 1: $R \in (0, I(A : B)_{\rho})$. By the proof of achievability for Theorem 8, see (F.23),

$$\liminf_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}(e^{-nR}) \ge \sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R) > 0,$$
(F.44)

where the strict inequality follows from Theorem 8. (F.44) implies that $\limsup_{n\to\infty} \hat{\alpha}_{n,\rho}(e^{-nR}) = 0$. By (F.43), it follows that $\lim_{n\to\infty} \hat{\alpha}_{n,\rho}(e^{-nR}) = 0$.

Case 2: $R \in (-\infty, 0]$. By Lemma 16 (a), (b),

$$\limsup_{n \to \infty} \hat{\alpha}_{n,\rho}(e^{-nR}) \le \limsup_{n \to \infty} \hat{\alpha}_{n,\rho}(e^0) = 0.$$
 (F.45)

By (F.43), it follows that $\lim_{n\to\infty} \hat{\alpha}_{n,\rho}(e^{-nR}) = 0.$

This completes the proof of $\lim_{n\to\infty} \hat{\alpha}_{n,\rho}(e^{-nR}) = 0$. The assertion regarding $\hat{\alpha}_{n,\rho}^{iid}$ follows from this due to Lemma 16 (b).

4. Example for Remark 4

Before we begin constructing a counterexample, let us note that for any $\rho_{AB} \in \mathcal{S}(AB), n \in \mathbb{N}_{>0}, \mu \in [0, \infty),$

$$\hat{\alpha}_{n,\rho}^{\text{ind}}(\mu) = \min_{\substack{T_{A^n B^n} \in \mathcal{L}(A^n B^n):\\ 0 \le T_{A^n B^n} \le 1}} \left\{ \operatorname{tr}[\rho_{AB}^{\otimes n}(1 - T_{A^n B^n}^n)] : \right.$$

$$\max_{\substack{m \in \mathbb{N}_{>0} \\ (p_i)_{i \in [m]} \in [0,1]^{\times m}: \\ i \in [m]}} \max_{\substack{p_i \operatorname{tr}[\sigma_{A^n}^{(i)} \otimes \tau_{B^n}^{(i)} T_{A^n B^n}^n] \le \mu }} \sum_{i \in [m]} p_i \operatorname{tr}[\sigma_{A^n}^{(i)} \otimes \tau_{B^n}^{(i)} T_{A^n B^n}^n] \le \mu } \right\}. \quad (F.46)$$

We will now construct a counterexample for the consideration in Remark 4. Suppose $d_A \geq 2, d_B \geq 2$, and let $\{|i\rangle_A\}_{i=0}^1, \{|i\rangle_B\}_{i=0}^1$ be orthonormal vectors in A, B. Let $p \in (\frac{1}{2}, 1)$ and let $\rho_{AB} \coloneqq p|0,0\rangle\langle 0,0|_{AB} + (1-p)|1,1\rangle\langle 1,1|_{AB}$. By Theorem 7 (v),

$$R_{1/2} \coloneqq I_{1/2}^{\downarrow\downarrow}(A:B)_{\rho} - \frac{1}{4} \frac{\partial}{\partial s^+} I_s^{\downarrow\downarrow}(A:B)_{\rho} \big|_{s=\frac{1}{2}} = -\log p - \frac{1}{4} (-4\log\max(p,1-p)) = 0.$$
(F.47)

Consider now the left-hand side of (3.30) with $\hat{\alpha}_{n,\rho}$ replaced by $\hat{\alpha}_{n,\rho}^{\text{ind}}$. Since ρ_{AB} is separable between A and B, also $\rho_{AB}^{\otimes n}$ is separable between A^n and B^n for any $n \in \mathbb{N}_{>0}$. Hence, (F.46) implies that $\hat{\alpha}_{n,\rho}^{\text{ind}}(\mu) \geq 1 - \mu$ for all $\mu \in [0, 1]$. On the other hand, due to Lemma 16 (b), we have $\hat{\alpha}_{n,\rho}^{\text{ind}}(\mu) \leq 1 - \mu$ for all $\mu \in [0, 1]$. Therefore, $\hat{\alpha}_{n,\rho}^{\text{ind}}(\mu) = 1 - \mu$ for all $\mu \in [0, 1]$. This implies that $\lim_{n\to\infty} \hat{\alpha}_{n,\rho}^{\text{ind}}(e^{-nR}) = 1$ for any $R \in (0, \infty)$. Hence, for any $R \in (0, \infty)$

$$\lim_{n \to \infty} -\frac{1}{n} \log \hat{\alpha}_{n,\rho}^{\text{ind}}(e^{-nR}) = 0.$$
 (F.48)

Consider now the right-hand side of (3.30). For any $R \in (0, H(A)_{\rho})$

$$\sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (I_s^{\downarrow\downarrow}(A:B)_{\rho} - R) = \sup_{s \in (\frac{1}{2}, 1)} \frac{1-s}{s} (H_{\frac{s}{2s-1}}(A)_{\rho} - R) > 0.$$
(F.49)

The equality in (F.49) follows from Theorem 7 (v). The strict inequality in (F.49) follows from the continuity of the Rényi entropy in the Rényi order and $R < H(A)_{\rho} = H_1(A)_{\rho}$. A comparison of (F.48) and (F.49) reveals that the equality in (3.30) is violated if $\hat{\alpha}_{n,\rho}$ is replaced by $\hat{\alpha}_{n,\rho}^{\text{ind}}$. Therefore, Theorem 8 does not hold if $\hat{\alpha}_{n,\rho}$ is replaced by $\hat{\alpha}_{n,\rho}^{\text{ind}}$.

- Amos Lapidoth and Christoph Pfister. Two Measures of Dependence. Entropy, 21(778), 2019. DOI: 10.3390/e21080778.
- [2] Robin Sibson. Information radius. Zeitschrift f
 ür Wahrscheinlichkeitstheorie und Verwandte Gebiete, 14:149–160, 1969. DOI: 10.1007/BF00537520.
- [3] Imre Csiszár. Generalized Cutoff Rates and Rényi's Information Measures. IEEE Transactions on Information Theory, 41(1):26-34, 1995. DOI: 10.1109/18.370121.
- [4] Siu-Wai Ho and Sergio Verdú. Convexity/Concavity of Renyi Entropy and α-Mutual Information. In 2015 IEEE International Symposium on Information Theory (ISIT), pages 745–749, 2015.
 DOI: 10.1109/ISIT.2015.7282554.
- [5] Sergio Verdú. α-Mutual Information. In 2015 Information Theory and Applications Workshop (ITA), 2015. DOI: 10.1109/ITA.2015.7308959.
- [6] Sergio Verdú. Error Exponents and α-Mutual Information. Entropy, 23(2):199, 2021.
 DOI: https://doi.org/10.3390/e23020199.
- [7] Amedeo Roberto Esposito, Adrien Vandenbroucque, and Michael Gastpar. On Sibson's α-Mutual Information. In 2022 IEEE International Symposium on Information Theory (ISIT), pages 2904–2909, 2022. DOI: 10.1109/ISIT50566.2022.9834428.
- [8] Amedeo Roberto Esposito, Michael Gastpar, and Ibrahim Issa. Sibson's α-Mutual Information and its Variational Representations, 2024. DOI: 10.48550/arXiv.2405.08352.
- [9] Marco Tomamichel and Masahito Hayashi. Operational Interpretation of Rényi Information Measures via Composite Hypothesis Testing Against Product and Markov Distributions. *IEEE Transactions on Information Theory*, 64(2):1064–1082, 2018. DOI: 10.1109/TIT.2017.2776900.
- [10] Wassily Hoeffding. Asymptotically optimal tests for multinomial distributions. The Annals of Mathematical Statistics, 36(2):369–401, 1965. DOI: 10.1214/aoms/1177700150.
- [11] Wassily Hoeffding. On probabilities of large deviations. Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability, 5(1):203–219, 1967.

- [12] Imre Csiszár and Giuseppe Longo. On the error exponent for source coding and for testing simple statistical hypotheses. *Studia Scientiarum Mathematicarum Hungarica*, 6:181–191, 1971.
- [13] Richard Blahut. Hypothesis Testing and Information Theory. IEEE Transactions on Information Theory, 20(4):405-417, 1974. DOI: 10.1109/TIT.1974.1055254.
- [14] Koenraad M. R. Audenaert, Michael Nussbaum, Arleta Szkoła, and Frank Verstraete. Asymptotic Error Rates in Quantum Hypothesis Testing. *Communications in Mathematical Physics*, 279(1):251– 283, 2008. DOI: 10.1007/s00220-008-0417-5.
- [15] Masahito Hayashi and Marco Tomamichel. Correlation detection and an operational interpretation of the Rényi mutual information. *Journal of Mathematical Physics*, 57(102201), 2016. DOI: 10.1063/1.4964755.
- [16] Amos Lapidoth and Christoph Pfister. Testing Against Independence and a Rényi Information Measure. In 2018 IEEE Information Theory Workshop (ITW), 2018. DOI: 10.1109/ITW.2018.8613520.
- [17] Manish K. Gupta and Mark M. Wilde. Multiplicativity of Completely Bounded p-Norms Implies a Strong Converse for Entanglement-Assisted Capacity. *Communications in Mathematical Physics*, 334(2):867–887, 2014. DOI: 10.1007/s00220-014-2212-9.
- [18] Alexander McKinlay and Marco Tomamichel. Decomposition rules for quantum Rényi mutual information with an application to information exclusion relations. *Journal of Mathematical Physics*, 61(7), 2020. DOI: 10.1063/1.5143862.
- [19] Dénes Petz. Quasi-entropies for finite quantum systems. Reports on Mathematical Physics, 23(1):57–65, 1986. DOI: 10.1016/0034-4877(86)90067-4.
- [20] Mark M. Wilde, Andreas Winter, and Dong Yang. Strong Converse for the Classical Capacity of Entanglement-Breaking and Hadamard Channels via a Sandwiched Rényi Relative Entropy. Communications in Mathematical Physics, 331(2):593–622, 2014. DOI: 10.1007/s00220-014-2122-x.
- [21] Martin Müller-Lennert, Frédéric Dupuis, Oleg Szehr, Serge Fehr, and Marco Tomamichel. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*, 54(12):122203, 2013. DOI: 10.1063/1.4838856.
- [22] Jonah Kudler-Flam, Laimei Nie, and Akash Vijay. Rényi mutual information in quantum field theory, tensor networks, and gravity, 2023. DOI: 10.48550/arXiv.2308.08600.
- [23] Jonah Kudler-Flam. Rényi Mutual Information in Quantum Field Theory. Physical Review Letters, 130(021603), 2023. DOI: 10.1103/PhysRevLett.130.021603.
- [24] Hao-Chung Cheng. Simple and Tighter Derivation of Achievability for Classical Communication Over Quantum Channels. PRX Quantum, 4:040330, 2023. DOI: 10.1103/PRXQuantum.4.040330.
- [25] Hao-Chung Cheng and Li Gao. Error Exponent and Strong Converse for Quantum Soft Covering. IEEE Transactions on Information Theory, 70(5):3499–3511, 2024. DOI: 10.1109/TIT.2023.3307437.
- [26] Mario Berta, Fernando G. S. L. Brandão, and Christoph Hirche. On Composite Quantum Hypothesis Testing. Communications in Mathematical Physics, 385(1):55–77, 2021. DOI: 10.1007/s00220-021-04133-8.
- [27] Yuan Zhai, Bo Yang, and Zhengjun Xi. Chain rules for a mutual information based on Rényi zerorelative entropy. *Physical Review A*, 108(1):012413, 2023. DOI: 10.1103/PhysRevA.108.012413.
- [28] Masahito Hayashi. Error exponent in asymmetric quantum hypothesis testing and its application to classical-quantum channel coding. *Physical Review A*, 76(6), 2007. DOI: 10.1103/physreva.76.062301.
- [29] Hiroshi Nagaoka. The Converse Part of The Theorem for Quantum Hoeffding Bound, 2006. DOI: 10.48550/arXiv.quant-ph/0611289.
- [30] Te Sun Han and Kingo Kobayashi. The strong converse theorem for hypothesis testing. IEEE Transactions on Information Theory, 35(1):178–180, 1989. DOI: 10.1109/18.42188.
- [31] Kenji Nakagawa and Fumio Kanaya. On the Converse Theorem in Statistical Hypothesis Testing. IEEE Transactions on Information Theory, 39(2):623–628, 1993. DOI: 10.1109/18.212293.
- [32] Milán Mosonyi and Tomohiro Ogawa. Quantum Hypothesis Testing and the Operational Interpretation of the Quantum Rényi Relative Entropies. *Communications in Mathematical Physics*, 334(3):1617–1648, 2015. DOI: 10.1007/s00220-014-2248-x.
- [33] Milán Mosonyi and Tomohiro Ogawa. Two Approaches to Obtain the Strong Converse Exponent of Quantum Hypothesis Testing for General Sequences of Quantum States. *IEEE Transactions on Information Theory*, 61(12):6975–6994, 2015. DOI: 10.1109/TIT.2015.2489259.
- [34] Laura Burri. Doubly minimized sandwiched Rényi mutual information: Properties and operational

interpretation from strong converse exponent (to appear).

- [35] Wiesław Pusz and Stanisław L. Woronowicz. Functional calculus for sesquilinear forms and the purification map. Reports on Mathematical Physics, 8(2):159–170, 1975.
 DOI: 10.1016/0034-4877(75)90061-0.
- [36] Tsuyoshi Ando. On some operator inequalities. Mathematische Annalen, 279(1):157–159, 1987. DOI: BF01456197.
- [37] Fumio Kubo and Tsuyoshi Ando. Means of positive linear operators. *Mathematische Annalen*, 246(3):205–224, 1980. DOI: 10.1007/BF01371042.
- [38] Renato Renner. Security of Quantum Key Distribution, 2006. DOI: 10.48550/arXiv.quant-ph/0512258.
- [39] Matthias Christandl, Robert König, and Renato Renner. Postselection Technique for Quantum Channels with Applications to Quantum Cryptography. *Physical Review Letters*, 102(2), 2009. DOI: 10.1103/physrevlett.102.020504.
- [40] Marco Tomamichel and Masahito Hayashi. A Hierarchy of Information Quantities for Finite Block Length Analysis of Quantum Tasks. *IEEE Transactions on Information Theory*, 59(11):7693–7710, 2013. DOI: 10.1109/TIT.2013.2276628.
- [41] Ke Li. Second-order asymptotics for quantum hypothesis testing. The Annals of Statistics, 42(1):171– 189, 2014. DOI: 10.1214/13-aos1185.
- [42] Masanori Ohya and Dénes Petz. Quantum Entropy and Its Use. Springer, Berlin, 1993.
- [43] Michael Nussbaum and Arleta Szkoła. The Chernoff lower bound for symmetric quantum hypothesis testing. The Annals of Statistics, 37(2):1040–1057, 2009. DOI: 10.1214/08-A0S593.
- [44] Simon M. Lin and Marco Tomamichel. Investigating properties of a family of quantum Rényi divergences. Quantum Information Processing, 14(4):1501-1512, 2015. DOI: 10.1007/s11128-015-0935-y.
- [45] Marco Tomamichel. Quantum Information Processing with Finite Resources. Springer International Publishing, 2016. DOI: 10.1007/978-3-319-21891-5.
- [46] Huzihiro Araki. On an inequality of Lieb and Thirring. Letters in Mathematical Physics, 19:167—170, 1990. DOI: 10.1007/BF01045887.
- [47] Elliott H. Lieb and Walter E. Thirring. Inequalities for the Moments of the Eigenvalues of the Schrödinger Hamiltonian and Their Relation to Sobolev Inequalities, pages 135–169. Springer Berlin Heidelberg, Berlin, Heidelberg, 1991. DOI: 10.1007/978-3-662-02725-7.
- [48] Rajendra Bhatia. Matrix Analysis. Springer. DOI: 10.1007/978-1-4612-0653-8.
- [49] Milán Mosonyi, Zsombor Szilágyi, and Mihály Weiner. On the Error Exponents of Binary State Discrimination With Composite Hypotheses. *IEEE Transactions on Information Theory*, 68(2):1032–1067, 2022. DOI: 10.1109/TIT.2021.3125683.
- [50] Samuel O. Scalet, Álvaro M. Alhambra, Georgios Styliaris, and J. Ignacio Cirac. Computable Rényi mutual information: Area laws and correlations. *Quantum*, 5:541, 2021. DOI: 10.22331/q-2021-09-14-541.
- [51] Hao-Chung Cheng, Li Gao, and Min-Hsiu Hsieh. Properties of Noncommutative Rényi and Augustin Information. *Communications in Mathematical Physics*, 390:501–544, 2022. DOI: 10.1007/s00220-022-04319-8.
- [52] Rajendra Bhatia. Positive Definite Matrices. Princeton University Press, 2007.
- [53] Eric Evert, Scott McCullough, Tea Strekelj, and Anna Vershynina. Convexity of a certain operator trace functional. *Linear Algebra and its Applications*, 643:218–234, 2022. DOI: 10.1016/j.laa.2022.02.033.
- [54] Henri Epstein. Remarks on Two Theorems of E. Lieb. Communications in Mathematical Physics, 31:317–325, 1973. DOI: 10.1007/BF01646492.
- [55] Eric A. Carlen and Elliott H. Lieb. A Minkowski Type Trace Inequality and Strong Subadditivity of Quantum Entropy II: Convexity and Concavity. *Letters in Mathematical Physics*, 83(2):107–126, 2008. DOI: 10.1007/s11005-008-0223-1.
- [56] Fumio Hiai. Concavity of certain matrix trace and norm functions. Linear Algebra and its Applications, 439(5):1568–1589, 2013. DOI: 10.1016/j.laa.2013.04.020.