

LANDAU TYPE THEOREM FOR α -HARMONIC MAPPINGS

VASUDEVARAO ALLU AND ROHIT KUMAR

ABSTRACT. In this paper, we first obtain an estimate of the coefficients for α -harmonic mappings. By applying these coefficient estimates, we prove the Landau type theorem for α -harmonic mappings defined on the unit disc \mathbb{D} .

1. INTRODUCTION

Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ represents the unit disc in the complex plane \mathbb{C} . We denote the complex differential operators

$$\frac{\partial}{\partial z} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \quad \text{and} \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$$

for $z = x + iy \in \mathbb{C}$ where x and y are real. For a continuously differentiable function f , denote

$$\Lambda_f = \max_{0 \leq \theta \leq 2\pi} |f_z + e^{-2i\theta} f_{\bar{z}}| = |f_z| + |f_{\bar{z}}|$$

and

$$\lambda_f = \min_{0 \leq \theta \leq 2\pi} |f_z + e^{-2i\theta} f_{\bar{z}}| = ||f_z| - |f_{\bar{z}}||,$$

where $f_z = \partial f / \partial z$ and $f_{\bar{z}} = \partial f / \partial \bar{z}$. Throughout this paper, we denote by $C^n(\mathbb{D})$ the set of all n -times continuously differentiable complex-valued functions in \mathbb{D} , where $n \in \{1, 2, \dots\}$.

For $\alpha \in \mathbb{R}$ and $z \in \mathbb{D}$, let

$$T_\alpha = -\frac{\alpha^2}{4} (1 - |z|^2)^{-\alpha-1} + \frac{\alpha}{2} (1 - |z|^2)^{-\alpha-1} \left(z \frac{\partial}{\partial z} + \bar{z} \frac{\partial}{\partial \bar{z}} \right) + \frac{1}{4} (1 - |z|^2)^{-\alpha} \Delta$$

be the second order elliptic partial differential operator, where Δ is the usual complex Laplacian operator

$$\Delta := 4 \frac{\partial^2}{\partial z \partial \bar{z}} = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}.$$

The corresponding homogeneous differential equation is

$$T_\alpha(f) = 0 \quad \text{in } \mathbb{D}. \quad (1.1)$$

File: Rohit_Allu-03-06-2024-arxiv.tex, printed: 2024-6-7, 0.47

2020 *Mathematics Subject Classification.* Primary 31A30, 30C99; Secondary 31A05, 31A35.

Key words and phrases. α -harmonic mapping, Landau type theorem, Coefficient estimate, Gauss hypergeometric function.

and its associated Dirichlet boundary value problem is as follows

$$\begin{cases} T_\alpha(f) = 0 & \text{in } \mathbb{D}, \\ f = f^* & \text{on } \partial\mathbb{D}. \end{cases} \quad (1.2)$$

Here, the boundary data $f^* \in \mathfrak{D}'(\partial\mathbb{D})$ is a distribution on the boundary $\partial\mathbb{D}$ of \mathbb{D} , and the boundary condition in (1.2) is interpreted in the distributional sense that $f_r \rightarrow f^*$ in $\mathfrak{D}'(\partial\mathbb{D})$ as $r \rightarrow 1^-$, where

$$f_r(e^{i\theta}) = f(re^{i\theta}), e^{i\theta} \in \partial\mathbb{D},$$

for $r \in [0, 1)$. In 2014, Olofsson [26] proved that, for parameter values $\alpha > -1$, if a function $f \in C^2(\mathbb{D})$ satisfies (1.1) with $\lim_{r \rightarrow 1^-} f_r = f^* \in \mathfrak{D}'(\partial\mathbb{D})$, then it has the form of a Poisson type integral

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} K_\alpha(ze^{-i\tau}) f^*(e^{i\tau}) d\tau \quad \text{for } z \in \mathbb{D},$$

where

$$K_\alpha(z) = c_\alpha \frac{(1 - |z|^2)^{\alpha+1}}{|1 - z|^{\alpha+2}},$$

$c_\alpha = (\Gamma(\alpha/2 + 1))^2/\Gamma(1 + \alpha)$ and $\Gamma(s) = \int_0^\infty t^{s-1}e^{-t}dt$ for $s > 0$ is the standard Gamma function.

If we take $\alpha = 2(n - 1)$, then f is polyharmonic (or n -harmonic), where $n \in \{1, 2, 3, \dots\}$. For a detailed study on polyharmonic mappings, we refer to [5, 11, 16, 17, 23, 24, 25, 26]. In particular, if $\alpha = 0$, then f is harmonic. Thus, f is a kind of generalization of classical harmonic mappings.

The Gauss hypergeometric function is defined by the following series

$$F(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n x^n}{(c)_n n!}$$

for $|x| < 1$, where $(a)_0 = 1$ and $(a)_n = a(a+1)\cdots(a+n-1)$ for $n = 1, 2, \dots$ are the Pochhammer symbols. Obviously, for $n = 0, 1, 2, \dots$, $(a)_n = \Gamma(a+n)/\Gamma(a)$. It is easy to see that

$$\frac{d}{dx} F(a, b; c; x) = \frac{ab}{c} F(a+1, b+1; c+1; x).$$

Furthermore, for $\text{Re}(c - a - b) > 0$, we have (see [4, Theorem 2.2.2])

$$F(a, b; c; 1) = \lim_{x \rightarrow 1} F(a, b; c; x) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}. \quad (1.3)$$

The following result concerns the solution to the equation (1.1).

Theorem A. [26] *Let $\alpha \in \mathbb{R}$ and $f \in C^2(\mathbb{D})$. Then f satisfies (1.1) if, and only if, it has a series expansion of the form*

$$f(z) = \sum_{k=0}^{\infty} c_k F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2\right) z^k + \sum_{k=1}^{\infty} c_{-k} F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2\right) \bar{z}^k, \quad (1.4)$$

for some sequence $\{c_k\}_{-\infty}^{\infty}$ of complex number satisfying

$$\limsup_{|k| \rightarrow \infty} |c_k|^{\frac{1}{|k|}} \leq 1. \quad (1.5)$$

In particular, the expansion (1.4), subject to the condition (1.5), converges in $C^\infty(\mathbb{D})$, and every solution f of (1.1) is $C^\infty(\mathbb{D})$ in the unit disc \mathbb{D} .

In 2014, Olofsson [26] pointed out that if $\alpha \leq -1$, $f \in C^2(\mathbb{D})$ satisfies (1.1), and the boundary limit $f^* = \lim_{r \rightarrow 1^-} f_r$ exists in $\mathfrak{D}'(\partial\mathbb{D})$, then $f(z) = 0$ for all $z \in \mathbb{D}$. Therefore, in this paper, we always assume that $\alpha > -1$.

The classical Landau theorem (see [15]) asserts that if f is a holomorphic mapping with $f(0) = 0$, $f'(0) = 1$ and $|f(z)| < M$ for $z \in \mathbb{D}$, then f is univalent in \mathbb{D}_{r_0} , and $f(\mathbb{D}_{r_0})$ contains a disc \mathbb{D}_{σ_0} , where

$$r_0 = \frac{1}{M + \sqrt{M^2 - 1}} \quad \text{and} \quad \sigma_0 = Mr_0^2.$$

The quantities r_0 and σ_0 can't be improved as the function $f_0(z) = Mz \left(\frac{1-Mz}{M-z} \right)$ shows the sharpness of the quantities r_0 and σ_0 .

In 2000, Chen *et al.* [6] obtained the Landau type theorems for harmonic mappings in the unit disc \mathbb{D} .

Theorem B. [6] *Let f be a harmonic mapping of the unit disc \mathbb{D} such that $f(0) = 0$, $f_{\bar{z}}(0) = 0$, $f_z(0) = 1$, and $|f(z)| < M$ for $z \in \mathbb{D}$. Then, f is univalent on a disc \mathbb{D}_{ρ_0} with*

$$\rho_0 = \frac{\pi^2}{16mM},$$

and $f(\mathbb{D}_{\rho_0})$ contains a schlicht disc \mathbb{D}_{R_0} with

$$R_0 = \rho_0/2 = \frac{\pi^2}{32mM},$$

where $m \approx 6.85$ is the minimum of the function $(3 - r^2)/(r(1 - r^2))$ for $0 < r < 1$.

Theorem C. [6] *Let f be a harmonic mapping of the unit disc \mathbb{D} such that $f(0) = 0$, $\lambda_f(0) = 1$ and $\Lambda_f(z) \leq \Lambda$ for $z \in \mathbb{D}$. Then, f is univalent on a disc \mathbb{D}_{ρ_0} with*

$$\rho_0 = \frac{\pi}{4(1 + \Lambda)},$$

and $f(\mathbb{D}_{\rho_0})$ contains a schlicht disc \mathbb{D}_{R_0} with

$$R_0 = \frac{1}{2}\rho_0 = \frac{\pi}{8(1 + \Lambda)}.$$

The radii ρ_0 and R_0 in Theorem B and Theorem C are not sharp. These results have been improved by Chen *et al.* [9], Dorff and Nowak [12], Grigoryan [13], Huang [14], Liu [19, 20], Liu and Chen [21] and Zhu [28]. In particular, Liu [20] has proved that under the hypothesis of Theorem C, $\Lambda \geq 1$, and when $\Lambda = 1$, f is univalent in the disc \mathbb{D}_ρ and $f(\mathbb{D}_\rho)$ contains a schlicht disc \mathbb{D}_R with $\rho = R = 1$ being sharp. In 2018, Liu and Chen [21] established the sharp version of Theorem C for $\Lambda > 1$ by applying geometric method. The Landau type theorem has also been studied

for elliptic harmonic mappings [3], logharmonic mappings [2] and log- p -harmonic mappings [22].

In 2008, Abdulhadi and Muhanna [1] established two versions of Landau-type theorems for certain bounded biharmonic mappings. After that, the Landau type theorem for functions f satisfying (1.1) with $\alpha = 2(n - 1)$, where $n \in \{1, 2, 3, \dots\}$, has been extensively studied by many authors (see [1, 7, 10, 18, 27]).

2. COEFFICIENT ESTIMATES

There have been many discussions in the literature regarding the coefficient estimates of the functions f satisfying (1.1) with $\alpha = 0$ and $\alpha = 2$. In 2015, Chen and Vuorinen [11] proved the coefficient estimates for bounded functions satisfying (1.1) with $\alpha > -1$. In this section, we prove a different coefficient estimates for α -harmonic mappings when $\alpha > 0$.

Theorem 2.1. *For $\alpha > 0$, let $f \in C^2(\mathbb{D})$ be the solution of (1.1) with series expansion of the form (1.4) and $\Lambda_f(z) \leq \Lambda$ for each $z \in \mathbb{D}$, where Λ is a positive number. Then, for $k \in \{1, 2, 3, \dots\}$,*

$$\frac{\alpha|2k - \alpha|}{2(k + 1)} \left(\left| c_k F \left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; 1 \right) \right| + \left| c_{-k} F \left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; 1 \right) \right| \right) \leq \Lambda.$$

Proof. We know that

$$F \left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2 \right) = \sum_{n=0}^{\infty} \frac{\left(-\frac{\alpha}{2}\right)_n \left(k - \frac{\alpha}{2}\right)_n |z|^{2n}}{(k + 1)_n n!}. \quad (2.1)$$

Differentiating (2.1) partially with respect to z , we obtain

$$\begin{aligned} \frac{\partial}{\partial z} F \left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2 \right) &= \bar{z} \sum_{n=1}^{\infty} \frac{\left(-\frac{\alpha}{2}\right)_n \left(k - \frac{\alpha}{2}\right)_n |z|^{2(n-1)}}{(k + 1)_n (n - 1)!} \\ &= \bar{z} \frac{-\frac{\alpha}{2} \left(k - \frac{\alpha}{2}\right)}{k + 1} F \left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; |z|^2 \right). \end{aligned}$$

Similarly, we have

$$\frac{\partial}{\partial \bar{z}} F \left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2 \right) = z \frac{-\frac{\alpha}{2} \left(k - \frac{\alpha}{2}\right)}{k + 1} F \left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; |z|^2 \right).$$

For $r \in [0, 1)$, let

$$G_k(r, \alpha) = \frac{-\frac{\alpha}{2} \left(k - \frac{\alpha}{2}\right)}{k + 1} F \left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; r^2 \right).$$

A simple computation shows that

$$f_z(re^{i\theta}) = \sum_{k=1}^{\infty} c_k r^{k+1} G_k(r, \alpha) e^{i(k-1)\theta} + \sum_{k=1}^{\infty} k c_k F \left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; r^2 \right) r^{k-1} e^{i(k-1)\theta}$$

$$+ \sum_{k=1}^{\infty} c_{-k} r^{k+1} G_k(r, \alpha) e^{-i(k+1)\theta} \quad (2.2)$$

and

$$\begin{aligned} f_{\bar{z}}(r e^{i\theta}) &= \sum_{k=1}^{\infty} c_k r^{k+1} G_k(r, \alpha) e^{i(k+1)\theta} + \sum_{k=1}^{\infty} k c_{-k} F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; r^2\right) r^{k-1} e^{-i(k-1)\theta} \\ &\quad + \sum_{k=1}^{\infty} c_{-k} r^{k+1} G_k(r, \alpha) e^{-i(k-1)\theta}, \end{aligned} \quad (2.3)$$

where $z = r e^{i\theta}$. By multiplying (2.2) by $e^{i(k+1)\theta}$ and then integrating from 0 to 2π , we obtain

$$c_{-k} r^{k+1} G_k(r, \alpha) = \frac{1}{2\pi} \int_0^{2\pi} f_z(r e^{i\theta}) e^{i(k+1)\theta} d\theta. \quad (2.4)$$

Similarly, multiplying (2.3) by $e^{-i(k+1)\theta}$ and then integrating from 0 to 2π , we obtain

$$c_k r^{k+1} G_k(r, \alpha) = \frac{1}{2\pi} \int_0^{2\pi} f_{\bar{z}}(r e^{i\theta}) e^{-i(k+1)\theta} d\theta. \quad (2.5)$$

Since $\Lambda_f(z) \leq \Lambda$, we may conclude from equations (2.4) and (2.5) that

$$\begin{aligned} (|c_k G_k(r, \alpha)| + |c_{-k} G_k(r, \alpha)|) r^{k+1} &\leq \frac{1}{2\pi} \int_0^{2\pi} (|f_z(r e^{i\theta})| + |f_{\bar{z}}(r e^{i\theta})|) d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \Lambda_f(r e^{i\theta}) d\theta \leq \Lambda. \end{aligned} \quad (2.6)$$

By letting $r \rightarrow 1^-$ and using (1.3), we obtain

$$|c_k G_k(1, \alpha)| + |c_{-k} G_k(1, \alpha)| \leq \Lambda.$$

This completes the proof. \square

We obtain the following corollary from Theorem 2.1.

Corollary 2.1. *Let $\alpha > 0$ and not an even integer. Let $f \in C^2(\mathbb{D})$ be the solution of (1.1) with series expansion of the form (1.4) and $\Lambda_f(z) \leq \Lambda$ for each $z \in \mathbb{D}$. Then, for $k \in \{1, 2, 3, \dots\}$,*

$$|c_k| + |c_{-k}| \leq \frac{2\Lambda\Gamma(1 + \frac{\alpha}{2})\Gamma(k + 1 + \frac{\alpha}{2})}{k!\Gamma(\alpha + 1)|2k - \alpha|}. \quad (2.7)$$

Proof. Since $\alpha > 0$ and α is not a positive even integer, by using Theorem 2.1 and (1.3), we obtain the following inequality

$$\frac{\Gamma(k + 2)\Gamma(\alpha)}{\Gamma(k + 1 + \frac{\alpha}{2})\Gamma(1 + \frac{\alpha}{2})} (|c_k| + |c_{-k}|) \leq \frac{2\Lambda(k + 1)}{\alpha|2k - \alpha|}.$$

This implies that

$$|c_k| + |c_{-k}| \leq \frac{2\Lambda\Gamma(1 + \frac{\alpha}{2})\Gamma(k + 1 + \frac{\alpha}{2})}{k!\Gamma(\alpha + 1)|2k - \alpha|},$$

which completes the proof. \square

3. THE LANDAU TYPE THEOREM

In 2015, Chen and Vuorinen[11] obtained the Landau type theorem for α -harmonic mappings when $\alpha \in (-1, 0]$ and in 2021, Long and Wang [23] obtained it when $\alpha \in (0, 2)$. In this section, we derive another Landau type theorem for α -harmonic mappings when $\alpha \in (0, 2)$.

We first prove the following lemma, which is vital to the proof of the main result.

Lemma 3.1. *For $x \in [0, 1)$, let*

$$\varphi(x) = \beta - \frac{2\Lambda}{2-\alpha} \left((2-\alpha) \frac{x^2}{1-x} + \frac{2a}{(1-x)^3} - 2a + (2a-1) \frac{x^2}{1-x^2} \right),$$

where $\alpha \in (0, 2)$, $\beta > 0$ and $\Lambda > 0$ are constants and $a = \frac{\Gamma(1+\frac{\alpha}{2})}{\Gamma(1+\alpha)}$. Then φ is strictly decreasing and there is a unique $x_0 \in (0, 1)$ such that $\varphi(x_0) = 0$.

Proof. It is easy to see that $1/2 < a < 1$ for $\alpha \in (0, 2)$ (see [23, Lemma 3.1]). Let

$$f_1(x) = (2-\alpha) \frac{x^2}{1-x},$$

$$f_2(x) = \frac{2a}{(1-x)^3} - 2a,$$

and

$$f_3(x) = (2a-1) \frac{x^2}{1-x^2}.$$

Clearly,

$$\varphi(x) = \beta - \frac{2\Lambda}{2-\alpha} \left(f_1(x) + f_2(x) + f_3(x) \right).$$

A simple computation shows that $f_1'(x) > 0$, $f_2'(x) > 0$ and $f_3'(x) > 0$ for $x \in (0, 1)$. This implies that $\varphi(x)$ is strictly decreasing in $[0, 1)$.

Furthermore, we can observe that

$$\lim_{x \rightarrow 0} \varphi(x) = \beta > 0 \quad \text{and} \quad \lim_{x \rightarrow 1^-} \varphi(x) = -\infty.$$

Therefore, there exists a unique $x_0 \in (0, 1)$ such that $\varphi(x_0) = 0$. This completes the proof of this lemma. \square

Theorem 3.1. *For $\alpha \in (0, 2)$, let $a = \Gamma(\alpha/2 + 1)/\Gamma(1 + \alpha)$, $f \in C^2(\mathbb{D})$ be the solution of (1.1) satisfying $f(0) = 0$, $\lambda_f(0) \geq \beta$ and $\Lambda_f(z) \leq \Lambda$ for each $z \in \mathbb{D}$, where Λ, β are positive constants. Then f is univalent on a disc \mathbb{D}_{ρ_0} , where ρ_0 satisfies the following equation*

$$\beta - \frac{2\Lambda}{2-\alpha} \left((2-\alpha) \frac{\rho_0^2}{1-\rho_0} + \frac{2a}{(1-\rho_0)^3} - 2a + (2a-1) \frac{\rho_0^2}{1-\rho_0^2} \right) = 0.$$

Moreover, $f(\mathbb{D}_{\rho_0})$ contains a schlicht disc of radius

$$R_0 \geq \rho_0 \left(\beta - \frac{2\Lambda}{2-\alpha} \left(\frac{2-\alpha}{3} \frac{\rho_0^2}{1-\rho_0} + \frac{2a}{(1-\rho_0)^3} - 2a + \frac{2a-1}{3} \frac{\rho_0^2}{1-\rho_0^2} \right) \right).$$

Proof. By Theorem A, we may assume that

$$f(z) = \sum_{k=0}^{\infty} c_k F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2\right) z^k \\ + \sum_{k=1}^{\infty} c_{-k} F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; |z|^2\right) \bar{z}^k, \quad z \in \mathbb{D},$$

for some sequence $\{c_k\}_{k=-\infty}^{\infty}$ of complex numbers satisfying

$$\limsup_{|k| \rightarrow \infty} |c_k|^{\frac{1}{|k|}} \leq 1.$$

For $\alpha \in (0, 2)$ and $k \in \{1, 2, 3, \dots\}$, we observe that

$$\frac{d}{dt} F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; t\right) = \frac{\left(-\frac{\alpha}{2}\right)\left(k - \frac{\alpha}{2}\right)}{k + 1} F\left(1 - \frac{\alpha}{2}, k + 1 - \frac{\alpha}{2}; k + 2; t\right) < 0.$$

This shows that $F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; t\right)$ is decreasing on $t \in [0, 1)$. Therefore, we have

$$F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; t\right) \leq F\left(-\frac{\alpha}{2}, k - \frac{\alpha}{2}; k + 1; 0\right) = 1. \quad (3.1)$$

Further, for $\alpha \in (0, 2)$ and $k \in \{1, 2, 3, \dots\}$, we have

$$\frac{d}{dt} F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; t\right) \\ = \frac{\left(-\frac{\alpha}{2} + 1\right)\left(k - \frac{\alpha}{2} + 1\right)}{k + 2} F\left(-\frac{\alpha}{2} + 2, k - \frac{\alpha}{2} + 2; k + 3; t\right) > 0$$

which implies that $F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; t\right)$ is increasing function on $[0, 1)$.

Note that the function $F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; t\right)$ is nonnegative as its parameters are nonnegative. Therefore, we have

$$\left|F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; t\right)\right| \leq \left|F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; 1\right)\right|.$$

From Theorem 2.1, we have

$$\frac{\alpha(2k - \alpha)}{2(k + 1)} (|c_k| + |c_{-k}|) \left|F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; t\right)\right| \\ \leq \frac{\alpha(2k - \alpha)}{2(k + 1)} (|c_k| + |c_{-k}|) \left|F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; 1\right)\right| \leq \Lambda. \quad (3.2)$$

For $k \in \{1, 2, 3, \dots\}$ and $\alpha \in (0, 2)$, Long and Wang [23] have proved the following inequality.

$$\left|\frac{\left(-\frac{\alpha}{2}\right)_n \left(k - \frac{\alpha}{2}\right)_n}{(k + 1)_n} \frac{1}{n!}\right| \leq 1 - \frac{\Gamma(k + 1)\Gamma(1 + \alpha)}{\Gamma(k + 1 + \frac{\alpha}{2})\Gamma(1 + \frac{\alpha}{2})}. \quad (3.3)$$

Using (3.3) and Corollary 2.1, we obtain

$$\left(|c_1| + |c_{-1}|\right) \left|\frac{\left(-\frac{\alpha}{2}\right)_n \left(k - \frac{\alpha}{2}\right)_n}{(k + 1)_n} \frac{1}{n!}\right| \leq \frac{2\Lambda}{2 - \alpha} \left(\frac{\Gamma(1 + \frac{\alpha}{2})\Gamma(2 + \frac{\alpha}{2})}{\Gamma(2)\Gamma(\alpha + 1)} - 1\right) \\ < \frac{2\Lambda}{2 - \alpha} (2\alpha - 1), \quad (3.4)$$

where $a = \Gamma(\alpha/2 + 1)/\Gamma(1 + \alpha)$ and $\alpha \in (0, 2)$. Since $f(0) = c_0 = 0$, a simple computation shows that

$$\begin{aligned}
f_z(z) - f_z(0) &= \sum_{k=1}^{\infty} c_k \frac{-\frac{\alpha}{2}(k - \frac{\alpha}{2})}{k+1} F\left(-\frac{\alpha}{2} + 1; k - \frac{\alpha}{2} + 1; k + 2; |z|^2\right) \bar{z} z^k \\
&\quad + \sum_{k=1}^{\infty} c_{-k} \frac{-\frac{\alpha}{2}(k - \frac{\alpha}{2})}{k+1} F\left(-\frac{\alpha}{2} + 1; k - \frac{\alpha}{2} + 1; k + 2; |z|^2\right) \bar{z}^{k+1} \\
&\quad + \sum_{k=2}^{\infty} k c_k F\left(-\frac{\alpha}{2}; k - \frac{\alpha}{2}; k + 1; |z|^2\right) z^{k-1} \\
&\quad + c_1 \left(F\left(-\frac{\alpha}{2}; 1 - \frac{\alpha}{2}; 2; |z|^2\right) - 1\right)
\end{aligned} \tag{3.5}$$

and

$$\begin{aligned}
f_{\bar{z}}(z) - f_{\bar{z}}(0) &= \sum_{k=1}^{\infty} c_k \frac{-\frac{\alpha}{2}(k - \frac{\alpha}{2})}{k+1} F\left(-\frac{\alpha}{2} + 1; k - \frac{\alpha}{2} + 1; k + 2; |z|^2\right) z^{k+1} \\
&\quad + \sum_{k=1}^{\infty} c_{-k} \frac{-\frac{\alpha}{2}(k - \frac{\alpha}{2})}{k+1} F\left(-\frac{\alpha}{2} + 1; k - \frac{\alpha}{2} + 1; k + 2; |z|^2\right) z \bar{z}^k \\
&\quad + \sum_{k=2}^{\infty} k c_{-k} F\left(-\frac{\alpha}{2}; k - \frac{\alpha}{2}; k + 1; |z|^2\right) \bar{z}^{k-1} \\
&\quad + c_{-1} \left(F\left(-\frac{\alpha}{2}; 1 - \frac{\alpha}{2}; 2; |z|^2\right) - 1\right).
\end{aligned} \tag{3.6}$$

For $|z| = r$, in view of (3.5) and (3.6), we obtain

$$\begin{aligned}
&|f_z(z) - f_z(0)| + |f_{\bar{z}}(z) - f_{\bar{z}}(0)| \\
&\leq 2 \sum_{k=1}^{\infty} (|c_k| + |c_{-k}|) \frac{\alpha(2k - \alpha)}{2(k+1)} \left| F\left(-\frac{\alpha}{2} + 1, k - \frac{\alpha}{2} + 1; k + 2; r^2\right) \right| r^{k+1} \\
&\quad + \sum_{k=2}^{\infty} k(|c_k| + |c_{-k}|) \left| F\left(-\frac{\alpha}{2}; k - \frac{\alpha}{2}; k + 1; r^2\right) \right| r^{k-1} \\
&\quad + (|c_1| + |c_{-1}|) \left| F\left(-\frac{\alpha}{2}; 1 - \frac{\alpha}{2}; 2; r^2\right) - 1 \right|.
\end{aligned}$$

Applying (3.1), (3.2), (3.4) and Corollary 2.1, we obtain

$$\begin{aligned}
&|f_z(z) - f_z(0)| + |f_{\bar{z}}(z) - f_{\bar{z}}(0)| \\
&\leq 2\Lambda \sum_{k=1}^{\infty} r^{k+1} + \sum_{k=2}^{\infty} k(|c_k| + |c_{-k}|) r^{k-1} + (|c_1| + |c_{-1}|) \sum_{n=1}^{\infty} \left| \frac{(-\frac{\alpha}{2})_n (k - \frac{\alpha}{2})_n r^{2n}}{(k+1)_n n!} \right| \\
&\leq 2\Lambda \sum_{k=1}^{\infty} r^{k+1} + \sum_{k=2}^{\infty} \frac{2\Lambda a \Gamma(k+1 + \frac{\alpha}{2})}{\Gamma(k+1)(2k - \alpha)} r^{k-1} + \frac{2\Lambda}{2 - \alpha} (2a - 1) \sum_{n=1}^{\infty} r^{2n} \\
&< 2\Lambda \sum_{k=1}^{\infty} r^{k+1} + \frac{2\Lambda}{2 - \alpha} a \sum_{k=2}^{\infty} k(k+1) r^{k-1} + \frac{2\Lambda}{2 - \alpha} (2a - 1) \sum_{n=1}^{\infty} r^{2n}
\end{aligned}$$

$$\begin{aligned}
&= 2\Lambda \frac{r^2}{1-r} + \frac{2\Lambda}{2-\alpha} \left(\frac{2a}{(1-r)^3} - 2a \right) + \frac{2\Lambda}{2-\alpha} (2a-1) \frac{r^2}{1-r^2} \\
&= \frac{2\Lambda}{2-\alpha} \left((2-\alpha) \frac{r^2}{1-r} + \frac{2a}{(1-r)^3} - 2a + (2a-1) \frac{r^2}{1-r^2} \right). \tag{3.7}
\end{aligned}$$

To prove the univalence of f in \mathbb{D}_{ρ_0} , we choose two distinct points $z_1, z_2 \in \mathbb{D}_{\rho_0}$ and let $[z_1, z_2]$ denote the line segment from z_1 to z_2 with ends points z_1 and z_2 , where ρ_0 satisfies the following equation

$$\beta - \frac{2\Lambda}{2-\alpha} \left((2-\alpha) \frac{\rho_0^2}{1-\rho_0} + \frac{2a}{(1-\rho_0)^3} - 2a + (2a-1) \frac{\rho_0^2}{1-\rho_0^2} \right) = 0.$$

By using (3.7) and Lemma 3.1, we have

$$\begin{aligned}
|f(z_2) - f(z_1)| &= \left| \int_{[z_1, z_2]} f_z(z) dz + f_{\bar{z}}(z) d\bar{z} \right| \\
&= \left| \int_{[z_1, z_2]} f_z(0) dz + f_{\bar{z}}(0) d\bar{z} \right| \\
&\quad - \left| \int_{[z_1, z_2]} (f_z(z) - f_z(0)) dz + (f_{\bar{z}}(z) - f_{\bar{z}}(0)) d\bar{z} \right| \\
&\geq \lambda_f(0) |z_2 - z_1| \\
&\quad - \int_{[z_1, z_2]} (|f_z(z) - f_z(0)| + |f_{\bar{z}}(z) - f_{\bar{z}}(0)|) |dz| \\
&> |z_1 - z_2| \left\{ \beta - \frac{2\Lambda}{2-\alpha} \left((2-\alpha) \frac{\rho_0^2}{1-\rho_0} + \frac{2a}{(1-\rho_0)^3} - 2a \right. \right. \\
&\quad \left. \left. + (2a-1) \frac{\rho_0^2}{1-\rho_0^2} \right) \right\} \\
&= 0.
\end{aligned}$$

Therefore, $f(z_1) \neq f(z_2)$. This shows that f is univalent in the ball \mathbb{D}_{ρ_0} .

To prove the second part of the theorem, let $\zeta = \rho_0 e^{i\theta} \in \partial\mathbb{D}_{\rho_0}$, then we have

$$\begin{aligned}
|f(\zeta) - f(0)| &= \left| \int_{[0, \zeta]} f_z(z) dz + f_{\bar{z}}(z) d\bar{z} \right| \\
&= \left| \int_{[0, \zeta]} f_z(0) dz + f_{\bar{z}}(0) d\bar{z} \right| \\
&\quad - \left| \int_{[0, \zeta]} (f_z(z) - f_z(0)) dz + (f_{\bar{z}}(z) - f_{\bar{z}}(0)) d\bar{z} \right| \\
&\geq \lambda_f(0) |\zeta| - \int_{[0, \zeta]} (|f_z(z) - f_z(0)| + |f_{\bar{z}}(z) - f_{\bar{z}}(0)|) |dz| \\
&\geq \beta \rho_0 - \frac{2\Lambda}{2-\alpha} \int_0^{\rho_0} \left((2-\alpha) \frac{r^2}{1-r} + \frac{2a}{(1-r)^3} - 2a \right.
\end{aligned}$$

$$\begin{aligned}
& + (2a - 1) \frac{r^2}{1 - r^2} \Big) dr \\
& \geq \beta \rho_0 - \frac{2\Lambda}{2 - \alpha} \left(\frac{2 - \alpha}{1 - \rho_0} \int_0^{\rho_0} r^2 dr + \left(\frac{2a}{(1 - \rho_0)^3} - 2a \right) \int_0^{\rho_0} dr \right. \\
& \quad \left. + \frac{2a - 1}{1 - \rho_0^2} \int_0^{\rho_0} r^2 dr \right) \\
& = \beta \rho_0 - \frac{2\Lambda}{2 - \alpha} \left(\frac{2 - \alpha}{1 - \rho_0} \frac{\rho_0^3}{3} + \left(\frac{2a}{(1 - \rho_0)^3} - 2a \right) \rho_0 + \frac{2a - 1}{1 - \rho_0^2} \frac{\rho_0^3}{3} \right) \\
& = \rho_0 \left(\beta - \frac{2\Lambda}{2 - \alpha} \left(\frac{2 - \alpha}{1 - \rho_0} \frac{\rho_0^2}{3} + \frac{2a}{(1 - \rho_0)^3} - 2a + \frac{2a - 1}{1 - \rho_0^2} \frac{\rho_0^2}{3} \right) \right).
\end{aligned}$$

Hence $f(\mathbb{D}_{\rho_0})$ contains a Schlicht disc of radius R_0 with

$$R_0 \geq \rho_0 \left(\beta - \frac{2\Lambda}{2 - \alpha} \left(\frac{2 - \alpha}{1 - \rho_0} \frac{\rho_0^2}{3} + \frac{2a}{(1 - \rho_0)^3} - 2a + \frac{2a - 1}{1 - \rho_0^2} \frac{\rho_0^2}{3} \right) \right).$$

This completes the proof. \square

Corollary 3.1. *For $\alpha \in (0, 2)$, let $a = \Gamma(\alpha/2 + 1)/\Gamma(1 + \alpha)$, $f \in C^2(\mathbb{D})$ be the solution of (1.1) satisfying $f(0) = |J_f(0)| - \beta = 0$ and $\Lambda_f(z) \leq \Lambda$ for each $z \in \mathbb{D}$, where Λ, β are positive constants and J_f is the Jacobian of f . Then f is univalent on a disc \mathbb{D}_{ρ_1} , where ρ_1 satisfies the following equation*

$$\frac{\beta}{\Lambda} - \frac{2\Lambda}{2 - \alpha} \left((2 - \alpha) \frac{\rho_1^2}{1 - \rho_1} + \frac{2a}{(1 - \rho_1)^3} - 2a + (2a - 1) \frac{\rho_1^2}{1 - \rho_1^2} \right) = 0.$$

Moreover, $f(\mathbb{D}_{\rho_1})$ contains a schlicht disc of radius

$$R_1 \geq \rho_1 \left(\frac{\beta}{\Lambda} - \frac{2\Lambda}{2 - \alpha} \left(\frac{2 - \alpha}{1 - \rho_1} \frac{\rho_1^2}{3} + \frac{2a}{(1 - \rho_1)^3} - 2a + \frac{2a - 1}{1 - \rho_1^2} \frac{\rho_1^2}{3} \right) \right).$$

Proof. Since, $|J_f(0)| = \beta$, we have

$$\beta = |J_f(0)| = |\det D_f(0)| = \Lambda_f(0) \lambda_f(0) \leq \Lambda \lambda_f(0),$$

which gives

$$\lambda_f(0) = ||f_z(0)| - |f_{\bar{z}}(0)|| \geq \frac{\beta}{\Lambda}.$$

Let $z_1, z_2 \in \mathbb{D}_{\rho_1}$, then by using the proof of Theorem 3.1 and Lemma 3.1, we have

$$\begin{aligned}
|f(z_1) - f(z_2)| & > |z_1 - z_2| \left\{ \frac{\beta}{\Lambda} - \frac{2\Lambda}{2 - \alpha} \left((2 - \alpha) \frac{\rho_1^2}{1 - \rho_1} + \frac{2a}{(1 - \rho_1)^3} - 2a \right. \right. \\
& \quad \left. \left. + (2a - 1) \frac{\rho_1^2}{1 - \rho_1^2} \right) \right\} = 0.
\end{aligned}$$

Therefore, $f(z_1) \neq f(z_2)$. This shows that f is univalent in the ball \mathbb{D}_{ρ_1} .

To prove the second part of the corollary, let $\zeta = \rho_1 e^{i\theta} \in \partial\mathbb{D}_{\rho_1}$. By using the proof of Theorem 3.1, we obtain

$$|f(\zeta)| \geq \rho_1 \left(\frac{\beta}{\Lambda} - \frac{2\Lambda}{2-\alpha} \left(\frac{2-\alpha}{1-\rho_1} \frac{\rho_1^2}{3} + \frac{2a}{(1-\rho_1)^3} - 2a + \frac{2a-1}{1-\rho_1^2} \frac{\rho_1^2}{3} \right) \right).$$

Hence $f(\mathbb{D}_{\rho_1})$ contains a Schlicht disc of radius R_1 with

$$R_1 \geq \rho_1 \left(\frac{\beta}{\Lambda} - \frac{2\Lambda}{2-\alpha} \left(\frac{2-\alpha}{1-\rho_1} \frac{\rho_1^2}{3} + \frac{2a}{(1-\rho_1)^3} - 2a + \frac{2a-1}{1-\rho_1^2} \frac{\rho_1^2}{3} \right) \right).$$

This completes the proof. □

Acknowledgement: The second named author thank CSIR for their support.

REFERENCES

- [1] Z. ABDULHADI and Y. ABU MUHANNA, Landau's theorem for biharmonic mappings, *J. Math. Anal. Appl.* **338** (2008), 705–709.
- [2] Z. ABDULHADI, Y. ABU MUHANNA and R.M. ALI, Landau's theorem for functions with logharmonic Laplacian, *Appl. Math. Comput.* **218** (2012), 6798–6802.
- [3] V. ALLU and R. KUMAR, Landau-Bloch type theorem for elliptic and quasiregular harmonic mappings, *J. Math. Anal. Appl.* **535** (2024), 128215.
- [4] G.E. ANDREWS, R. ASKEY and R. ROY, Volume 71 of Encyclopedia of Mathematics and its Applications, *Cambridge University Press*, Cambridge, 1999.
- [5] A. BORICHEV and H. HEDENMALM, Weighted integrability of polyharmonic functions, *Adv. Math.* **264** (2014), 464–505.
- [6] S. CHEN, P.M. GAUTHIER and W. HENGARTNER, Bloch constants for planer harmonic mappings, *Proc. Amer. Math. Soc.* **128** (2000), 3231–3240.
- [7] S. CHEN, S. PONNUSAMY and X. WANG, Landau's theorem for certain biharmonic mappings, *Appl. Math. Comput.* **208** (2009), 427–433.
- [8] S. CHEN, S. PONNUSAMY and X. WANG, Bloch constant and Landau's theorem for planar p-harmonic mappings, *J. Math. Anal. Appl.* **373** (2011), 102–110.
- [9] S. CHEN, S. PONNUSAMY and X. WANG, Coefficient estimates and Landau–Bloch's theorem for planar harmonic mappings, *Bull. Malays. Math. Sci. Soc.* **34**(2) (2011), 255–265.
- [10] J. CHEN, A. RASILA and X. WANG, Landau's theorem for polyharmonic mappings, *J. Math. Anal. Appl.* **409** (2014), 934–945.
- [11] S. CHEN and M. VUORINEN, Some properties of a class of elliptic partial differential operators, *J. Math. Anal. Appl.* **431**(2) (2015), 1124–1137.
- [12] M. DORFF and M. NOWAK, Landau's theorem for planar harmonic mappings, *Comput. Methods Funct. Theory* **4** (2004), 151–158.
- [13] A. GRIGOYAN, Landau and Bloch theorems for harmonic mappings, *Complex Var. Elliptic Equ.* **51**(1) (2006), 81–87.
- [14] X. Z. HUANG, Estimates on Bloch constants for planar harmonic mappings, *J. Math. Anal. Appl.* **337** (2008), 880–887.
- [15] E. LANDAU, Der Picard-Schottkysche Satz und die Blochsche Konstanten, *Sitzungsber. Preuss. Akad. Wiss. Berlin Phys.-Math. Kl.* (1926), 467–474.
- [16] P.J. LI and S. PONNUSAMY, Lipschitz continuity of quasiconformal mappings and of the solutions to second order elliptic PDE with respect to the distance ratio metric, *Complex Anal. Oper. Theory* **12** (2018), 1991–2001.
- [17] P. J. LI, X. T. WANG and Q. H. XIAO, Several properties of α -harmonic functions in the unit disk, *Monatsh. Math.* **184** (2017), 627–640.

- [18] M.S. LIU, Landau's theorems for biharmonic mappings, *Complex Var. Elliptic Equ.* **53**(9) (2008), 843–855.
- [19] M. S. LIU, Landau's theorems for planer harmonic mappings, *Comput. Math. Appl.* **57**(7) (2009), 1142–1146.
- [20] M.S. LIU, Estimates on Bloch constants for planar harmonic mappings, *Sci. China Ser. A-Math.* **52**(1) (2009), 87–92.
- [21] M.S. LIU and H.H. CHEN, The Landau–Bloch type theorems for planar harmonic mappings with bounded dilation, *J. Math. Anal. Appl.* **468**(2) (2018), 1066–1081.
- [22] P.J. LI and X.T. WANG, Landau's theorem for log-p-harmonic mappings, *Appl. Math. Comput.* **218**(2012), 4806–4812.
- [23] B.-Y. LONG and Q.-H. WANG, Starlikeness, convexity and Landau type theorem of the real kernel α -harmonic mappings, *Filomat* **35**(8) (2021), 2629–2644.
- [24] B.-Y. LONG and Q.-H. WANG, Some coefficient estimates on real kernel α -harmonic mappings, *Proc. Amer. Math. Soc.* **150**(4) (2022), 1529–1540.
- [25] A. OLOFSSON, A representation formula for radially weighted biharmonic functions in the unit disc, *Publ. Math.* **49** (2005), 393–415.
- [26] A. OLOFSSON, Differential operators for a scale of Poisson type kernels in the unit disc, *J. Anal. Math.* **123** (2014), 227–249.
- [27] Y.C. ZHU and M.S. LIU, Landau-type theorems for certain planar harmonic mappings or biharmonic mappings, *Complex Var. Elliptic Equ.* **58**(12) (2013), 1667–1676.
- [28] J.F. ZHU, Landau theorem for planar harmonic mappings, *Complex Anal. Oper. Theory* **9** (2015), 1819–1826.

VASUDEVARAO ALLU, SCHOOL OF BASIC SCIENCE, INDIAN INSTITUTE OF TECHNOLOGY BHUBANESWAR, BHUBANESWAR-752050, ODISHA, INDIA.

Email address: avrao@iitbbs.ac.in

ROHIT KUMAR, SCHOOL OF BASIC SCIENCE, INDIAN INSTITUTE OF TECHNOLOGY BHUBANESWAR, BHUBANESWAR-752050, ODISHA, INDIA.

Email address: rohitk12798@gmail.com