Global existence of solutions to the fully parabolic chemotaxis system with logistic source under nonlinear Neumann boundary condition

Minh Le

June 5, 2024

Abstract

We study the existence of global boundedness solutions to the fully parabolic chemotaxis systems with logistic sources, $ru - \mu u^2$, under nonlinear Neumann boundary conditions, $\frac{\partial u}{\partial \nu} = |u|^p$ where p > 1 in smooth bounded domain $\Omega \subset \mathbb{R}^n$ with $n \ge 2$. A recent study by Le (2023) has shown that the logistic sources can ensure that solutions are global and bounded when n = 2 with $p < \frac{3}{2}$ and n = 3 with $p < \frac{7}{5}$. In this paper, we extend the previous findings by demonstrating the existence of global bounded solutions when $p < \frac{3}{2}$ in any spatial dimension $n \ge 2$.

1 Introduction

In this paper, we consider the following PDEs arising from chemotaxis

$$\begin{cases} u_t = \Delta u - \nabla \cdot (u \nabla v) + au - \mu u^2 & x \in \Omega, \ t \in (0, T_{\max}), \\ \tau v_t = \Delta v + u - v & x \in \Omega, \ t \in (0, T_{\max}), \end{cases}$$
(1.1)

in a smooth, convex, bounded domain $\Omega \subset \mathbb{R}^n$ with $n \geq 2$, where $a, \mu > 0$ and $\tau \in \{0, 1\}$. The system (1.1) is complemented with the nonnegative initial conditions in $C^2(\overline{\Omega})$, not identically zero:

$$u(x,0) = u_0(x), \qquad v(x,0) = v_0(x), \qquad x \in \Omega,$$
(1.2)

and the nonlinear Neumann boundary conditions

$$\frac{\partial u}{\partial \nu} = |u|^p, \qquad \frac{\partial v}{\partial \nu} = 0, \qquad x \in \partial\Omega, \ t \in (0, T_{\max}), \tag{1.3}$$

where p > 1 and ν is the outward normal vector.

Under the homogeneous Neumann boundary conditions, the system (1.1) when $a = \mu = 0$ used to describe the movement of cells toward chemical concentration known as chemotaxis has been first introduced in [4]. This system in 2d has an intriguing property known as critical mass, which states that if the total mass is less than a certain value solutions are global and bounded (see [2, 1, 11, 14, 15]) while it is greater than that value, solutions blow up in finite time (see [12, 13, 18, 19]). In higher dimensional case $n \ge 3$, this property has been proven in [23] to fail to hold, precisely it was proven that finite-time blow-up solutions can be constructed for any given initial mass. In attempt to avoid blow-up phenomenon, the logistic sources, $au-\mu u^2$, introduced in [21] can guarantee that solutions are globally bounded when $\mu > \frac{n-2}{n}$ with $\tau = 0$. The similar result for fully parabolic chemotaxis system when $\tau > 0$, was later established in [25]. Additionally, those interested in further exploration of qualitative and quantitative aspects of chemotaxis systems with logistic sources can consult the following references: [3, 5, 7, 8, 9, 20, 24, 26, 27].

In 1990s, nonlinear parabolic equations under the nonlinear Neumann boundary conditions has been studied in [10, 17, 16]. Particularly, the authors study the global existence and finite-time blow-up solutions of the following equation:

$$\begin{cases} U_t = \Delta U - \mu U^Q & x \in \Omega, \ t \in (0, T_{\max}), \\ \frac{\partial U}{\partial \nu} = U^P & x \in \Omega, \ t \in (0, T_{\max}), \\ U(x, 0) = U_0(x) & x \in \overline{\Omega}, \end{cases}$$
(NBC)

where Ω is a smooth bounded domain in \mathbb{R}^n , Q, P > 1, $\mu > 0$ and $U_0 \in W^{1,\infty}(\Omega)$ is a nonnegative function. It was shown that $P = \frac{Q+1}{2}$ is critical for the blow up in the following sense:

- 1. if $P < \frac{Q+1}{2}$ then all solutions of (NBC) exist globally and are globally bounded,
- 2. If $P > \frac{Q+1}{2}$ (or $P = \frac{Q+1}{2}$ and μ is sufficiently small) then there exist initial functions U_0 such that the corresponding solutions of (NBC) blow-up in L^{∞} -norm.

Motivated by those works, [6] introduces and studies the chemotaxis systems (1.1) under the nonlinear Neumann boundary conditions (1.3). The main objective of that work is to give an answer to the question: "What is the largest value p so that logistic damping still avoids blow-up?"

In [6], it was shown that solutions of the parabolic-elliptic system (1.1) when $\tau = 0$ exist globally and remain bounded in time when $p < \frac{3}{2}$. However, in case $\tau = 1$, the existence of globally bounded solutions was only shown for n = 2 when $p < \frac{3}{2}$ and n = 3 when $p < \frac{7}{5}$ with additional largeness assumption on μ . In this paper, we will show that largest value pso that logistic damping still excludes blow-up is $\frac{3}{2}$ for the fully parabolic system (1.1) when $\tau = 1$ in any spacial dimension. This result is consistent with the results of (NBC) and [6] with $\tau = 0$, where critical power is shown to be $P = \frac{Q+1}{2}$ with Q = 2 in our problem. To be more precise, our main result reads as follows:

Theorem 1.1. Let Ω be a bounded, convex domain with smooth boundary $\partial \Omega \in C^{\infty}$ in \mathbb{R}^n with $n \geq 2$. Assume that $\tau = 1$, $1 and initial data <math>u_0, v_0$ in Holder space $C^{2+\frac{1}{2}}(\overline{\Omega})$ are nonnegative, not identically zero such that

$$\frac{\partial u_0}{\partial \nu} = |u_0|^p.$$

The system (1.1) under boundary condition (1.3) possesses a unique nonnegative classical solution $(u, v) \in (C^{2+p-1}(\bar{\Omega} \times [0, \infty)))^2$. Moreover, u, v > 0 in $\Omega \times (0, \infty)$ and remain

bounded in the sense that

$$\sup_{t>0} \left\{ \|u(\cdot,t)\|_{L^\infty(\Omega)} + \|v(\cdot,t)\|_{W^{1,\infty}(\Omega)} \right\} < \infty.$$

The key idea of the proof for $\tau = 1$ when n = 3 in [6] motivated by [25] is to investigate the energy functional

$$y(t) = \int_{\Omega} u^2 + \frac{1}{3} \int_{\Omega} u |\nabla v|^2 + \int_{\Omega} |\nabla v|^4.$$
 (1.4)

This allows us to take advantages of the term $-\mu u^2$, but also raises difficulties on the boundary terms when doing integral by parts. This method works well in 3D but requires $p < \frac{7}{5}$, which is far from the critical power $p = \frac{3}{2}$ in the parabolic-elliptic case. However, it cannot be utilized for higher dimensional cases when $n \ge 4$. Instead of using this idea, there is another way to deal with fully parabolic chemotaxis problems [28] by only considering the energy functional $\int_{\Omega} u^p$. This eases the computational significantly since we do not need the deal with the chemical concentration function v directly. This can be done thanks to the help of Lemma 2.2, which enables us to bound Δv by u directly.

In the following sections, we will briefly recall the local-wellposedness results for system (1.1) as well as some important estimates in Section 2, and prove the main theorem in Section 3.

2 Preliminaries

Let us commence this section by stating the local existence result established in [6][Theorem 2.2].

Lemma 2.1. Assume that Ω is a open convex bounded domain in \mathbb{R}^n , where $n \geq 1$ with smooth boundary $\partial \Omega \in C^{\infty}$ and nonnegative functions u_0, v_0 are in $C^{2+p-1}(\overline{\Omega})$ such that

$$\frac{\partial u_0}{\partial \nu} = |u_0|^p \qquad on \ \partial\Omega, \tag{2.1}$$

where $p \in (1, 2)$. Then there exists $T_{\max} \in (0, \infty]$ such that problem (1.1) under the boundary condition

$$\frac{\partial u}{\partial \nu} = |u|^p, \qquad \frac{\partial v}{\partial \nu} = 0, \qquad on \; \partial \Omega \times (0, T_{\max}),$$

admits a unique nonnegative solution u, v in $C^{2+p-1,1+\frac{p-1}{2}}(\overline{\Omega} \times [0, T_{\max}))$. Moreover, if u_0, v_0 are not identically zero in Ω then u, v are strictly positive in $\Omega \times (0, T_{\max})$. If $T_{\max} < \infty$, then

$$\lim_{t \to T_{\max}} \sup_{t \to T_{\max}} \left\{ \| u(\cdot, t) \|_{L^{\infty}(\Omega)} + \| v(\cdot, t) \|_{W^{1,\infty}(\Omega)} \right\} = \infty.$$
(2.2)

The forthcoming lemma concerned with the maximal Sobolev regularity for heat equation serves as a key estimate in the proof of the main result. Interested readers are referred to [22][Lemma 2.3] for details of the proof.

Lemma 2.2. Assuming that $\Omega \subset \mathbb{R}^n$ with $n \geq 1$, $p \in (n, \infty)$, $f \in L^p((0, T) \times \Omega)$, $g_0 \in W^{2,p}(\Omega)$ and g is a classical solution to the following system

$$\begin{cases} g_t = \Delta g - g + f & in \ \Omega \times (0, T), \\ \frac{\partial g}{\partial \nu} = 0 & on \ \partial \Omega \times (0, T), \\ g(\cdot, 0) = g_0 & in \ \Omega \end{cases}$$
(2.3)

for some $T \in (0, \infty]$. There exists $C = C(n, p, \Omega) > 0$ such that

$$\int_{t_0}^t e^{\frac{ps}{2}} \int_{\Omega} |\Delta g|^p \, dx \, ds \le C \left(\int_{t_0}^t e^{\frac{ps}{2}} \int_{\Omega} |f|^p \, dx \, ds + e^{\frac{pt_0}{2}} \, \|\Delta g(\cdot, t_0)\|_{L^p(\Omega)}^p \right) \tag{2.4}$$

for all $t \in (t_0, T)$ with $t_0 := \max\{1, \frac{T}{2}\}.$

The following lemma established in [6][Lemma 3.6] allows us to control boundary integrals by using the diffusion and the quadratic degradation.

Lemma 2.3. If $r \geq \frac{1}{2}$, $p \in (1, \frac{3}{2})$, and $g \in C^1(\overline{\Omega})$, then for every $\epsilon > 0$, there exists a positive constant $C = C(\epsilon, \Omega, p, r)$ such that

$$\int_{\partial\Omega} |g|^{p+2r-1} \le \epsilon \int_{\Omega} |g|^{2r+1} + \epsilon \int_{\Omega} |\nabla g^r|^2 + C.$$
(2.5)

3 Proof of the main result

We are now in position to prove our main result.

Proof of Theorem 1.1. First, we will show that $u \in L^{\infty}((0, T_{\max}); L^{r}(\Omega))$ for some fixed r > n. A direct calculation shows

$$\frac{1}{r}\frac{d}{dt}\int_{\Omega}u^{r} = \int_{\Omega}u^{r-1}u_{t}$$

$$= \int_{\Omega}u^{r-1}\left[\Delta u - \nabla(u\nabla v) + au - \mu u^{2}\right]$$

$$= -\frac{4(r-1)}{r^{2}}\int_{\Omega}|\nabla u^{\frac{r}{2}}|^{2} - \frac{r-1}{r}\int_{\Omega}u^{r}\Delta v + a\int_{\Omega}u^{r}$$

$$-\mu\int_{\Omega}u^{r+1} + \int_{\partial\Omega}u^{r+p-1}dS.$$
(3.1)

Applying Lemma 2.3 with $\epsilon = \frac{r-1}{r^2}$ yields

$$r \int_{\partial \Omega} u^{r+p-1} dS \le \frac{r-1}{r} \int_{\Omega} |\nabla u^{\frac{r}{2}}|^2 + \frac{r-1}{r} \int_{\Omega} u^{r+1} + c_2, \qquad (3.2)$$

where $c_2 = C(r) > 0$. In light of Young's inequality, it follows that

$$(ra + \frac{r+1}{2})\int_{\Omega} u^r \le \frac{r\mu}{4}\int_{\Omega} u^{r+1} + c_3,$$
 (3.3)

where $c_3 = C(r, a, \Omega) > 0$. By Young's inequality, we have

$$-(r-1)\int_{\Omega}u^{r}\Delta v \leq \frac{r\mu}{4}\int_{\Omega}u^{r+1} + c_{4}\int_{\Omega}|\Delta v|^{r+1},$$
(3.4)

where $c_4 = (r-1)^{r+1} \left(\frac{2}{\mu(r+1)}\right)^r$. Collecting from (3.1) to (3.4) leads to

$$\frac{d}{dt} \int_{\Omega} u^r + \frac{r+1}{2} \int_{\Omega} u^r \le c_4 \int_{\Omega} |\Delta v|^{r+1} - \left(\frac{r\mu}{2} - \frac{r-1}{r}\right) \int_{\Omega} u^{r+1} + c_5, \tag{3.5}$$

where $c_5 = c_2 + c_3$. Choosing μ sufficiently large such that $\frac{r\mu}{4} > \frac{r-1}{r}$ and plugging into (3.5) implies

$$\frac{d}{dt} \int_{\Omega} u^r + \frac{r+1}{2} \int_{\Omega} u^r \le c_4 \int_{\Omega} |\Delta v|^{r+1} - \frac{r\mu}{2} \int_{\Omega} u^{r+1} + c_5.$$
(3.6)

Multiplying both sides of (3.6) by $e^{(r+1)t}$ and integrating from t_0 to t where $t_0 = \min\left\{1, \frac{T_{\max}}{2}\right\}$, entails

$$\int_{\Omega} u^{r}(\cdot, t) \leq c_{4} \int_{t_{0}}^{t} e^{-\frac{(r+1)(t-s)}{2}} \int_{\Omega} |\Delta v(\cdot, s)|^{r+1} - \frac{r\mu}{4} \int_{t_{0}}^{t} e^{-\frac{(r+1)(t-s)}{2}} \int_{\Omega} u^{r+1}(\cdot, s) + \int_{\Omega} u^{r}(\cdot, t_{0}) + \frac{2c_{5}}{r+1}.$$
(3.7)

Applying Lemma 2.2 to the first term of the right hand side of (3.7) yields

$$c_{4} \int_{t_{0}}^{t} e^{-\frac{(r+1)(t-s)}{2}} \int_{\Omega} |\Delta v(\cdot,s)|^{r+1} \leq c_{6} \int_{t_{0}}^{t} e^{-\frac{(r+1)(t-s)}{2}} \int_{\Omega} u^{r+1}(\cdot,s) + c_{6} e^{-\frac{(r+1)(t-t_{0})}{2}} \int_{\Omega} |\Delta v(\cdot,t_{0})|^{r+1},$$
(3.8)

where $c_6 = C(n, r, \Omega) > 0$. From (3.7) and (3.8), we obtain

$$\int_{\Omega} u^{r}(\cdot, t) \le \left(c_{6} - \frac{r\mu}{4}\right) \int_{t_{0}}^{t} e^{-\frac{(r+1)(t-s)}{2}} \int_{\Omega} u^{r+1}(\cdot, s) + c_{7},$$
(3.9)

where $c_7 = c_6 e^{-\frac{(r+1)(t-t_0)}{2}} \int_{\Omega} |\Delta v(\cdot, t_0)|^{r+1} + \int_{\Omega} u^r(\cdot, t_0) + \frac{2c_5}{r+1}$. Choosing $\mu > \frac{4c_6}{r}$ yields that $u \in L^{\infty}((0, T_{\max}); L^r(\Omega))$ for some r > n. Applying Moser-iteration procedure established in [6][Lemma 5.2] for nonlinear Neumann boundary conditions implies that $L^{\infty}((0, T_{\max}); L^{\infty}(\Omega))$. This, together with Lemma 2.1 implies that $T_{\max} = \infty$ and u, v are in $C^{2+p-1}(\bar{\Omega} \times [0, \infty))$. The proof is now complete.

Acknowledgments

The author acknowledges support from the Mathematics Graduate Research Award Fellowship at Michigan State University.

References

- A. Blanchet, J. Dolbeault, and B. Perthame. "Two-dimensional Keller-Segel model: Optimal critical mass and qualitative properties of the solutions". In: *Electronic Jour*nal of Differential Equations, (May 2006).
- [2] J. Dolbeault and B. Perthame. "A Optimal critical mass in the two dimensional Keller–Segel model in R2". In: C. R. Acad. Sci. Paris, Ser. I 339 (2004), pp. 611– 616.
- [3] H. Jin and T. Xiang. "Chemotaxis effect vs. logistic damping on boundedness in the2-D minimal Keller–Segel model". In: C. R. Acad. Sci. Paris, Ser.I 356 (2018), pp. 875– 885.
- [4] F. Keller and A. Segel. "Initiation of Slime Mold Aggregation Viewed as an Instability". In: Journal of Theoretical Biology 26 (1970), pp. 399–415.
- [5] J. Lankeit. "Chemotaxis can prevent thresholds on population density". In: *Discrete* and Continuous Dynamical Systems - B 20 (2015), pp. 1499–1527.
- [6] M. Le. "Global existence of solutions to the chemotaxis system with logistic source under nonlinear Neumann boundary conditions". In: *Journal of Differential Equations* 377 (Dec. 2023), pp. 1–37.
- [7] D. Li et al. "Large time behavior of solution to an attraction-repulsion chemotaxis system with logistic source in three dimensions". In: *Journal of Mathematical Analysis and Applications* 448 (2017), pp. 914–936.
- [8] X. Li and Y. Wang. "On a fully parabolic chemotaxis system with Lotka–Volterra competitive kinetics". In: *Journal of Mathematical Analysis and Applications* 471 (Mar. 2019), pp. 584–598.
- [9] K. Lin, C. Mu, and H. Zhong. "A blow-up result for a quasilinear chemotaxis system with logistic source in higher dimensions". In: *Journal of Mathematical Analysis and Applications* 464 (Aug. 2018), pp. 435–455.
- [10] P. Quittner M. Chipot M. Fila. "Stationary solutions, blow up and convergence to stationary solutions for semilinear parabolic equations with nonlinear boundary conditions". In: Acta Math. Univ. Comenianae 60 (1991), pp. 35–103.
- [11] N. Mizoguchi. "Global existence for the Cauchy problem of the parabolic-parabolic Keller-Segel system on the plane". In: *Calculus of Variations and Partial Differential Equations* 48 (2013), pp. 491–505.
- [12] T. Nagai. "Blow-up of radially symmetric solutions to a chemotaxis system". In: Adv. Math. Sci. Appl. 5 (1995), pp. 581–601.
- [13] T. Nagai. "Global existence and blowup of solutions to a chemotaxis system". In: Nonlinear Analysis 47 (2001), pp. 777–787.
- [14] T. Nagai, T. Senba, and K. Yoshida. "Application of the Trudinger-Moser inequality to a Parabolic System of Chemotaxis," in: *Funkcilaj Ekvacioj* 40 (1997), pp. 411–433.
- [15] K. Osaki and A. Yagi. "Global existence for a chemotaxis-growth system in \mathbb{R}^{2} ". In: Adv. Math. Sci. Appl. 2 (Dec. 2002), pp. 587–606.

- [16] P. Quittner. "Global existence of solutions of parabolic problems with nonlinear boundary conditions". In: *Banach Center Publ.* 33 (1996), pp. 309–314.
- [17] P. Quittner. "On global existence and stationary solutions for two classes of semilinear parabolic problems". In: *Comment. Math. Univ. Carolin.* 34 (1993), pp. 105–124.
- [18] T. Senba T. Nagai and T. Suzuki. "Blowup of nonradial solutions to parabolic-elliptic systems modeling chemotaxis in two-dimensional domains". In: J. Inequal. Appl. 6 (2001), pp. 37–55.
- [19] T. Senba T. Nagai and T. Suzuki. "Chemotaxis collapse in a parabolic system of mathematical biology". In: *HiroshimaMath. J.* 20 (2000), pp. 463–497.
- [20] Y. Tao and M. Winkler. "Persistence of mass in a chemotaxis system with logistic source". In: *Journal of Differential Equations* 259 (Dec. 2015), pp. 6142–6161.
- [21] J. Tello and M. Winkler. "A Chemotaxis System with Logistic Source". In: Communications in Partial Differential Equations 32 (June 2007), pp. 849–877.
- [22] S. Zheng W. Wang M. Zhuang. "Positive effects of repulsion in boundedness in a fully parabolic attraction-repulsion chemotaxis system with logistic source". In: *Journal Differential Equations* 264 (2018), pp. 2011–2027.
- [23] M. Winkler. "Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model". In: Journal Differential Equations 248(12) (2010), pp. 2889–2905.
- [24] M. Winkler. "Blow-up in a higher-dimensional chemotaxis system despite logistic growth restriction". In: Journal of Mathematical Analysis and Applications 384 (Dec. 2011), pp. 261–272.
- [25] M. Winkler. "Boundedness in the Higher-Dimensional Parabolic-Parabolic Chemotaxis System with Logistic Source". In: Communications in Partial Differential Equations 35 (2010), pp. 1516–1537.
- [26] C. Yang et al. "Boundedness in a quasilinear fully parabolic Keller–Segel system of higher dimension with logistic source". In: Journal of Mathematical Analysis and Applications 430 (Oct. 2015), pp. 585–591.
- [27] J. Zheng et al. "A new result for global existence and boundedness of solutions to a parabolic–parabolic Keller–Segel system with logistic source". In: Journal of Mathematical Analysis and Applications 462 (June 2018), pp. 1–25.
- [28] M. Zhuang, W. Wang, and S. Zheng. "Boundedness in a fully parabolic chemotaxis system with logistic-type source and nonlinear production". In: Nonlinear Analysis: Real World Applications 47 (June 2019), pp. 473–483.