# A Ramsey-type theorem on deficiency

Jin Sun<sup>a</sup>, Xinmin Hou<sup>a,b,c\*</sup>

<sup>a</sup> School of Mathematical Sciences

University of Science and Technology of China, Hefei, Anhui 230026, China.

 $^{b}$  CAS Key Laboratory of Wu Wen-Tsun Mathematics

University of Science and Technology of China, Hefei, Anhui 230026, China.

 $^{c}$ Hefei National Laboratory,

University of Science and Technology of China, Hefei 230088, Anhui, China

#### Abstract

Ramsey's Theorem states that a graph G has bounded order if and only if G contains no complete graph  $K_n$  or empty graph  $E_n$  as its induced subgraph. The Gyárfás-Sumner conjecture says that a graph G has bounded chromatic number if and only if it contains no induced subgraph isomorphic to  $K_n$  or a tree T. The deficiency of a graph is the number of vertices that cannot be covered by a maximum matching. In this paper, we prove a Ramsey type theorem for deficiency, i.e., we characterize all the forbidden induced subgraphs for graphs G with bounded deficiency. As an application, we answer a question proposed by Fujita, Kawarabayashi, Lucchesi, Ota, Plummer and Saito (JCTB, 2006).

Keywords: Ramsey-type problem, deficiency, forbidden subgraph.

## 1 Introduction

In this paper, all the graphs we consider are finite and simple. For positive integer n, write [n] for  $\{1, 2, ..., n\}$ . Let G be a graph with vertex set V(G) and edge set E(G). Write |G| = |E(G)| for the size of G and call |V(G)| the order of G. For a vertex  $x \in V(G)$ , let  $N_G(x) = \{y \in V(G) : xy \in E(G)\}$  be the neighborhood of x. Write  $d_G(x) = |N_G(x)|$  for the degree of G. A leaf is a vertex of degree one in a graph. If there is no confusion from the context, we omit the index G. Write  $\delta(G)$  and  $\Delta(G)$  for the minimum degree and maximum degree of G, respectively. A graph G is called k-regular if  $\delta(G) = \Delta(G) = k$ .

<sup>\*</sup>e-mail: jinsun@mail.ustc.edu.cn (J. Sun), xmhou@ustc.edu.cn (X. Hou)

For two vertices x, y in a connected graph G, the *distance* between x and y is  $dist_G(x, y) = min\{|P| : P \text{ is an } x-y \text{ path in } G\}$ , and the *diameter* of G is

$$\operatorname{diam}(G) = \max_{x,y \in V(G)} \operatorname{dist}(x,y).$$

For a subset  $X \subset V(G)$ , let G[X] denote the subgraph of G induced by X, and set  $N_G(X) = \bigcup_{x \in X} N_G(x) - X$ . As usual, let  $P_n$ ,  $C_n$ ,  $K_n$  and  $E_n$  denote a path, a cycle, a complete graph, and an empty graph of order n, respectively. Let  $K_{s,t}$  be the complete bipartite graph with partitions of orders s and t. When s = 1, we call  $K_{1,t}$  a star. The center of a graph G is a vertex  $c \in V(G)$  with  $d(c, x) \leq \lceil \frac{\operatorname{diam}(G)}{2} \rceil$  for any vertex  $x \in V(G)$ . Clearly, a star  $K_{1,t}$  has a unique center, and a path  $P_n$  has a unique center if n is odd and two centers if n is even. One can refer to [4] for any other notation not introduced here.

For graphs  $H_1$  and  $H_2$ , we say  $H_1 \prec H_2$  if  $H_2$  contains an induced subgraph isomorphic to  $H_1$ , otherwise, write  $H_1 \not\prec H_2$ . For a family  $\mathcal{H}$  of graphs, we say G is  $\mathcal{H}$ -free if  $H \not\prec G$ for any  $H \in \mathcal{H}$ . For two families  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , we say  $\mathcal{H}_1 \leq \mathcal{H}_2$  if for every  $H_2 \in \mathcal{H}_2$ , there exists  $H_1 \in \mathcal{H}_1$  such that  $H_1 \prec H_2$ . The following result is a simple observation; however, it is important.

**Proposition 1.1.** The relation ' $\leq$ ' is transitive but not antisymmetric. Therefore, if  $\mathcal{H}_1 \leq \mathcal{H}_2$ , then every  $\mathcal{H}_1$ -free graph is also  $\mathcal{H}_2$ -free.

Using this language, the classical Ramsey's Theorem [14] and its connected version can be rephrased as follows.

**Theorem 1.2.** (A) (Ramsey's Theorem [14], 1929) For a family  $\mathcal{H}$  of graphs, there is a constant  $c = c(\mathcal{H})$  such that |V(G)| < c for every  $\mathcal{H}$ -free graph G if and only if  $\mathcal{H} \leq \{K_n, E_n\}$  for some positive integer n.

(B) (Proposition 9.4.1 in [4]) For a family  $\mathcal{H}$  of graphs, there is a constant  $c = c(\mathcal{H})$ such that |V(G)| < c for every connected  $\mathcal{H}$ -free graph G if and only if  $\mathcal{H} \leq \{K_n, K_{1,n}, P_n\}$ for some positive integer n.

The least integer  $c = c(\mathcal{H})$  when  $\mathcal{H} = \{K_m, E_n\}$  in (A) of Theorem 1.2 is denoted by R(m, n), called the *Ramsey number*.

In this article, we consider a more general Ramsey-type problem as follows. Let  $\mathcal{G}_c$  be the family of connected graphs. Given a graph parameter  $\mu$  and a constant b > 0, define

$$b\text{-B-}\mu = \{\mathcal{H} \subset \mathcal{G}_c : \mu(G) < b \text{ for any } \mathcal{H}\text{-free graph } G \in \mathcal{G}_c\},\$$

and

$$B-\mu = \{\mathcal{H} \subset \mathcal{G}_c : \text{there is a constant } b > 0 \text{ such that } \mathcal{H} \in b-B-\mu\}$$

### **Problem 1.3.** Given a graph parameter $\mu$ , determine $\mathcal{H} \in B$ - $\mu$ .

Let  $\mathcal{F}$  be a family of graphs. An *(induced)*  $\mathcal{F}$ -cover (or *(induced)*  $\mathcal{F}$ -partition) number of a graph G is the minimum n such that there exist  $G_1, G_2, \cdots, G_n$  satisfying every vertex of G is covered by at least (or exactly) one  $G_i$ , where each  $G_i$  is a (induced) subgraph of Gisomorphic to a member of  $\mathcal{F}$ . Write  $\mathcal{F}$ -c(G) (or  $\mathcal{F}$ -p(G)) and ind- $\mathcal{F}$ -c(G) (or ind- $\mathcal{F}$ -p(G)) for the  $\mathcal{F}$ -cover (or  $\mathcal{F}$ -partition) number and induced  $\mathcal{F}$ -cover (or induced  $\mathcal{F}$ -partition) number of a graph G, respectively. Different graph parameters can be formulated using this language. For example, the chromatic number  $\chi(G) = \text{ind-}\mathcal{F}$ -p(G), where  $\mathcal{F}$  consists of empty graphs; the order  $\operatorname{ord}(G) = \mathcal{F}$ -p(G), where  $\mathcal{F}$  consists of  $K_1$ ; and the domination number  $\gamma(G) = \mathcal{F}$ -p(G), where  $\mathcal{F}$  consists of stars. The Gyárfás-Sumner conjecture [7] can be restated as follows. This conjecture is still open in general although it has been widely considered in literatures. One can refer to a survey [16] about the progress of this conjecture.

**Conjecture 1.4** (Gyárfás-Sumner conjecture [7]). For a finite family  $\mathcal{H}$  of connected graphs,  $\mathcal{H} \in B$ - $\chi$  if and only if  $\mathcal{H} \leq \{K_n, T\}$ , where T is a tree and n is a positive integer.

In the following, we list more results on different parameters of Problem 1.3.

- (1) The connected version of the Ramsey theorem ((B) in Theorem 1.2) can be rephrased as  $\mathcal{H} \in B$ -ord if and only if  $\mathcal{H} \leq \{K_n, K_{1,n}, P_n\}$ .
- (2) Chiba and Furuya determined *H* ∈ B-*μ* when *μ* is the *F*-cover/partition number for *F* being the set of paths in [1], and when *μ* is the induced *F*-cover/partition number for *F* being the set of paths and stars in [2].
- (3) Choi, Furuya, Kim and Park [3] determined H ∈ B-µ when µ is the induced matching number (the maximum size of an induced 1-regular subgraph of a graph) or matching number (the maximum size of a 1-regular subgraph of a graph).
- (4) Furuya [5] determined  $\mathcal{H} \in B-\gamma$ , where  $\gamma$  is the domination number of a graph.
- (5) Lozin [11] determined *H* ∈ B-*µ*, where *µ* is the neighborhood diversity or VC-dimension (two parameters related to the property of the closed neighborhood of vertices, one can refer to [11] for details).
- (5) Lozin and Razgon [12] determined *H* ∈ B-μ, where μ(G) is the tree-width of graph G, which is the smallest integer such that graph G has a tree-decomposition into parts with at most k + 1 vertices.

(6) Kierstead and Penrice [9], and Scott, Seymour, and Spirkl [17] determined  $\mathcal{H} \in B-\partial$ , where  $\partial(G)$ , the degeneracy of graph G, is the smallest integer d such that every nonnull subgraph of G has a vertex of degree at most d.

In this paper, we focus on the deficiency of graphs. Let M be a matching of a graph G. For a vertex  $v \in V(G)$ , we call M saturates v if  $v \in V(M)$ ; otherwise, M misses v. The *matching number*  $\nu(G)$  of graph G is the cardinality of a maximum matching of G. The *deficiency* of graph G, denoted by def(G), is defined as

$$\operatorname{def}(G) = |V(G)| - 2\nu(G).$$

According to the definition,  $def(G) \equiv |V(G)| \pmod{2}$ . Note that  $def(G) = \mathcal{F} \cdot p(G) - 1$ where  $\mathcal{F} = \{K_1, mK_2 : m \geq 1\}$ . A vertex set X in a graph G is called an *independent* set, or stable set if G[X] is an empty graph. For a graph G, let  $\alpha(G)$  be the independence number of G. We call a vertex  $v \in V(G)$  the *claw center* if  $\alpha(G[N(v)]) \geq 3$ . We call G *claw-free* if G contains no claw center. The classical results of (near-)perfect matching on claw-free graph due to Las Vergnas [10], Sumner [15], Jünger, Pulleyblank and Reinelt [8] can be described as follows.

**Theorem 1.5** ([8, 10, 15]). Every connected claw-free graph has deficiency at most one.

Theorem 1.5 can be restated as  $\mathcal{H} \in 2$ -B- def if and only if  $\mathcal{H} \leq \{K_{1,3}\}$ . The main result of this article characterizes the finite family of graphs  $\mathcal{H} \in B$ - def.

In the following constructions, attaching a path  $P_n$  at a vertex v of a graph G means that we identify an end of  $P_n$  with v. We define additional families of graphs that will be used in this article. (See Figures 1 and 2.)

- $F_n^1$ : the graph obtained by attaching  $P_2$  at the center of path  $P_{2n+1}$ .
- $F_n^2$ : the graph obtained by attaching a copy of  $P_n$  at each vertex of triangle  $K_3$ .
- $F_n^3$ : the graph obtained by attaching a copy of  $P_n$  at each vertex of two fixed nonadjacent vertices of cycle  $C_4$ .
- $F_n^4$ : the graph obtained by adding an edge to connect the only two vertices of degree 3 in  $F_n^3$ .
- $T_n$ : the graph obtained by attaching two copies of  $P_2$  at one end of a path  $P_n$ . We call this end of  $P_n$  the branch vertex of  $T_n$ , and the other end of  $P_n$  is the end of  $T_n$ .

Note that  $T_1$  is actually  $P_3$ . The center of  $P_3$  is not only the branch vertex but also the end vertex of  $T_1$ . We continue to define graphs by attaching  $T_p$  at a vertex v of a graph G, i.e., we identify the end of  $T_p$  with v.




Figure 1: Graphs  $F_n^1$ ,  $F_n^2$ ,  $F_n^3$ ,  $F_n^4$  and  $\check{F}_n^p$ .

- $B_i (i \ge 0)$ : the graph obtained by attaching two copies of  $P_2$  at the end of  $T_{i+2}$ .
- $\check{K}_n^p$ : the graph obtained by attaching a copy of  $T_p$  at each vertex of  $K_n$ .
- $\check{F}_n^p$ : the graph obtained by attaching a  $T_p$  at one vertex of  $K_3$ , and attaching a copy of  $P_n$  at each of the remaining vertices of  $K_3$ .

Now, we state our main theorem.

**Theorem 1.6.** Let  $\mathcal{H}$  be a finite family of connected graphs. Then  $\mathcal{H} \in B$ -def if and only if

$$\mathcal{H} \le \{K_{1,n}, T_n, \check{K}_n^p : 1 \le p \le \frac{n}{2} - 1\}$$

or

$$\mathcal{H} \le \{K_{1,n}, F_n^1, F_n^2, F_n^3, F_n^4, \check{F}_n^p, \check{K}_n^p : 1 \le p \le n-2\}$$

for some even integer  $n \geq 4$ .

For any integer  $d \ge 2$ , let d-B'-def consist of inclusion-minimal elements of d-B-def, i.e. forbidden sets  $\mathcal{H} \in d$ -B-def while  $\mathcal{H}' \notin d$ -B- def for any proper subset  $\mathcal{H}' \subset \mathcal{H}$ . According to Theorem 1.5,  $\mathcal{H} \in 2$ -B'-def if and only if  $\mathcal{H} = \{K_{1,3}\}$ . Plummer and Saito [13] further showed that, given a graph H, for any integer  $d \ge 2$ ,  $\{H\} \in d$ -B'-def if and only if  $H \prec K_{1,3}$ . Fujita, Kawarabayashi, Lucchesi, Ota, Plummer, and Saito [6] asked the following question.

**Question 1.7.** For any integer  $d \ge 3$ , determine  $\mathcal{H}$  with  $|\mathcal{H}| = 2$  and  $\mathcal{H} \in d$ -B'-def.




Figure 2: Graphs  $\check{K}_n^p$ ,  $T_n$  and  $B_i$ .

Furthermore, they proved the following extension of Theorem 1.5.

**Theorem 1.8** ([6]). Let G be a connected  $\{K_{1,s}, B_i : i \ge 0\}$ -free graph where  $s \ge 4$ . Then  $def(G) \le s - 2$ .

As an application of Theorem 1.6, we answer Question 1.7.

**Theorem 1.9.** Let  $\mathcal{H} = \{H_1, H_2\}$  and  $d \geq 3$  be an integer. Then,  $\mathcal{H} \in d$ -B'-def if and only if  $\mathcal{H} = \{K_{1,s}, T_3\}$  or  $\{K_{1,s}, P_4\}$  for some s with  $4 \leq s \leq d + 1$ .

*Proof.* Note that, for  $\mathcal{H} = \{K_{1,s}, T_3\}$  or  $\{K_{1,s}, P_4\}$ ,  $\mathcal{H} \leq \{K_{1,s}, B_i : i \geq 0\}$ . Thus, by Theorem 1.8, we have  $def(G) \leq s - 2 < d$  since  $4 \leq s \leq d + 1$ , i.e.  $\mathcal{H} \in d$ -B-def. Clearly,  $\mathcal{H}$  is inclusion-minimal. Thus  $\mathcal{H} \in d$ -B'-def.

Now suppose  $\mathcal{H} \in d$ -B'- def. Let  $\mathcal{H}_1 = \{K_{1,n}, T_n, \check{K}_n^p : 1 \leq p \leq \frac{n}{2} - 1\}$  and  $\mathcal{H}_2 = \{K_{1,n}, F_n^1, F_n^2, F_n^3, F_n^4, \check{K}_n^p, \check{F}_n^p : 1 \leq p \leq n - 2\}$ . Thus,  $\mathcal{H}_1, \mathcal{H}_2 \leq \{K_{1,n}, F_n^1, \check{K}_n^1\}$ . Since  $\mathcal{H} \in d$ -B'- def  $\subset$  B- def, we have  $\mathcal{H} \leq \{K_{1,n}, F_n^1, \check{K}_n^1\}$ . Without loss of generality, assume  $H_1 \prec K_{1,n}$ . By Theorem 1.5 and the minimality of  $\mathcal{H}$ , we obtain  $H_1 = K_{1,s}$  for some  $s \geq 4$ . Thus,  $H_1$  is not an induced subgraph of  $F_n^1$  or  $\check{K}_n^1$ . Hence  $\{H_2\} \leq \{F_n^1, \check{K}_n^1\}$ . However,  $H_2 \prec F_n^1$  implies that  $H_2$  is triangle-free. Hence,  $H_2$  contains at most two vertices of the largest clique of  $\check{K}_n^1$ . Therefore,  $H_2 \prec B_0$ . Since  $B_0 \not\prec F_n^1$ , we have  $H_2 \prec T_3$ . By the minimality of  $\mathcal{H}$ , we have  $H_2 = T_3$  or  $P_4$ . Therefore, we have  $\mathcal{H} = \{K_{1,s}, T_3\}$  or  $\{K_{1,s}, P_4\}$ .  $s \leq d+1$  as def $(K_{1,s-1}) = s - 2 < d$  and  $K_{1,s-1}$  is  $\mathcal{H}$ -free.

The rest of this article is arranged as follows. We present some preliminaries in Section 2 and prove Theorem 1.6 in Section 3. We provide some discussion and remarks in the last section.

# 2 Preliminaries

We first construct several graphs with large deficiencies. Let s, t be positive integers, and let  $Q_i = u_i^1 u_i^2 \dots u_i^t$ ,  $i \in [s+1]$  be s+1 pairwise vertex disjoint paths. Define

- $H_{s,t}^1$ : the graph obtained from the union of paths  $Q_i, i \in [s+1]$  by adding 2s vertices  $\{v_i, w_i : i \in [s]\}$  and 3s edges  $\{v_i w_i, v_i u_i^t, v_i u_{i+1}^1 : i \in [s]\}$ .
- $H^3_{s,t}$ : the graph obtained from the union of paths  $Q_i, i \in [s+1]$  by adding 2s+2 vertices  $\{x, y\} \cup \{v_i, w_i : i \in [s]\}$  and 4s+2 edges  $\{xu_1^1, u_{s+1}^ty\} \cup \{v_iu_i^t, v_iu_{i+1}^1, w_iu_i^t, w_iu_{i+1}^1 : i \in [s]\}.$
- $H_{s,t}^4$ : the graph obtained from  $H_{s,t}^3$  by adding s edges  $\{u_i^t u_{i+1}^1 : i \in [s]\}$ .
- $\check{H}_{s,t}^p$ : the graph obtained from the union of paths  $Q_i, i \in [s+1]$  and  $R_i \cong T_p, i \in [s]$  with end  $v_i$  by adding 3s edges  $\{u_i^t u_{i+1}^1, u_i^t v_i, u_{i+1}^1 v_i : i \in [s]\}$ .

For convience, let  $H_{0,t}^1 = P_t$  and  $H_{0,t}^3 = H_{0,t}^4 = P_{t+2}$ .

Lemma 2.1. (1)

$$\operatorname{def}(\breve{H}_{s,t}^p) = \begin{cases} s & \text{if } t(s+1) + s(p+2) \text{ is even,} \\ s+1 & \text{otherwise.} \end{cases}$$

(2)

$$\operatorname{def}(\check{K}_{n}^{p}) = \begin{cases} n & \text{if } n(p+1) \text{ is even,} \\ n+1 & \text{otherwise.} \end{cases}$$

Proof. Recall the definition of  $\check{H}_{s,t}^p$  (resp.  $\check{K}_n^p$ ), for any  $i \in [s]$  (resp.  $i \in [n]$ ), each  $R_i (\cong T_p)$  contains a branch vertex that is adjacent to two leaves. Given a maximum matching M of  $\check{H}_{s,t}^p$  (or  $\check{K}_n^p$ ), M misses at least one of these two leaves. Thus M misses at least s (resp. n) vertices of  $\check{H}_{s,t}^p$  (resp.  $\check{K}_n^p$ ). Therefore,  $\operatorname{def}(\check{H}_{s,t}^p) \geq s$  (resp.  $\operatorname{def}(\check{K}_n^p) \geq n$ ).

Let H be the graph obtained by deleting a leaf adjacent to every branch vertex from  $\check{H}^p_{s,t}$  (resp.  $\check{K}^p_n$ ). Then H is connected and claw-free. By Theorem 1.5,  $\operatorname{def}(H) \leq 1$ . Hence  $\operatorname{def}(\check{H}^p_{s,t}) \leq s + 1$  (resp.  $\operatorname{def}(\check{K}^p_n) \leq n + 1$ ).

Since  $def(G) \equiv |V(G)| \pmod{2}$  for any graph G, the proof is complete.

Next we handle  $H_{s,t}^j$ :  $j \in \{1,3,4\}$ . When t is even, it can be easily checked that  $def(H_{s,t}^1) = def(H_{s,t}^3) = def(H_{s,t}^4) = 0$ , which is not what we want. When t is odd, we show that  $H_{s,t}^j$  has large deficiency for  $j \in \{1,3,4\}$ .

**Lemma 2.2.** If t is odd, then  $def(H_{s,t}^1) = s + 1$ .

*Proof.* Let  $H = H_{s,t}^1 - \{w_1, w_2, \cdots, w_s\}$ . Then H is a path, thus  $def(H) \leq 1$ . As a maximum matching of H is also a matching of  $H_{s,t}^1$ , we have  $def(H_{s,t}^1) \leq s + 1$ .

Now we prove that  $def(H_{s,t}^1) \geq s + 1$  by induction on s. For the base case s = 0,  $H_{0,t}^1 = P_t$  has odd vertices. Thus  $def(H_{0,t}^1) \geq 1$ . Assume that  $s \geq 1$ . Let M be a maximum matching of  $H_{s,t}^1$ . If  $v_1 \notin V(M)$ , then  $w_1$  must not be in V(M). Thus  $M \cup \{v_1 w_1\}$  is a larger matching, a contradiction. Therefore,  $v_1 \in V(M)$ . If  $\{v_1 u_1^t\} \in M$  (or  $\{v_1 u_2^1\} \in M$ ), then  $w_1$  is missed by M. Hence  $M' = M \cup \{v_1 w_1\} \setminus \{v_1 u_1^t\}$  (or  $M' = M \cup \{v_1 w_1\} \setminus \{v_1 u_2^1\}$ ) is still a maximum matching. Thus we may assume that  $\{v_1 w_1\} \in M$ . Then  $H' = H_{s,t}^1 - \{v_1, w_1\}$ consists of two components,  $H_{0,t}^1$  and  $H_{s-1,t}^1$ . By induction hypothesis,  $def(H_{0,t}^1) \geq 1$  and  $def(H_{s-1,t}^1) \geq s$ . Hence M misses at least 1 vertex in  $H_{0,t}^1$  and s vertices in  $H_{s-1,t}^1$ . Therefore,  $def(H_{s,t}^1) \geq s + 1$ .

**Lemma 2.3.** If t is odd, then  $def(H_{s,t}^3) = def(H_{s,t}^4) = s + 1$ .

*Proof.* Since  $H_{s,t}^3$  is a spanning subgraph of  $H_{s,t}^4$ , we have  $def(H_{s,t}^4) \leq def(H_{s,t}^3)$ . Let  $H = H_{s,t}^3 - \{w_1, w_2, \cdots, w_s\}$ . Then H is a path too. Thus  $def(H) \leq 1$ . As a maximum matching of H is also a matching of  $H_{s,t}^3$ , we have  $def(H_{s,t}^3) \leq s + 1$  too.

Now we prove that  $def(H_{s,t}^4) \geq s + 1$  by induction on s. For the base case s = 0,  $H_{0,t}^4 = P_{t+2}$ . Thus  $def(H_{0,t}^4) \geq 1$  as t is odd. Assume that  $s \geq 1$ . Let M be a maximum matching of  $H_{s,t}^4$ . If  $u_1^t \notin V(M)$ , then at least one of  $v_1$  and  $w_1$  cannot be saturated by M since both can be matched with only  $u_2^1$ . By symmetry, we may assume  $v_1 \notin V(M)$ . Then  $M \cup \{u_1^t v_1\}$  is a larger matching of  $H_{s,t}^4$ , a contradiction. Now assume  $u_1^t \in V(M)$ . If  $u_1^t u_2^1 \in M$ , then both  $v_1$  and  $w_1$  are missed by M. Therefore,  $(M \setminus \{u_1^t u_2^1\}) \cup \{u_1^t v_1, u_2^1 w_1\}$  is a larger matching, a contradiction too. If  $u_1^t u_1^{t-1} \in M$  and  $v_1$  is not saturated by M, then  $M' = M \setminus \{u_1^t u_1^{t-1}\} \cup \{u_1^t v_1\}$  is still a maximum matching.

Thus we may always assume  $u_1^t v_1 \in M$  by symmetry of  $v_1$  and  $w_1$  in  $H_{s,t}^4$ . Let  $H' = H_{s,t}^4 - \{u_1^t, v_1\}$ , then H' consists of two components,  $H_{0,t-2}^4$  and  $H_{s-1,t}^4$ . By induction hypothesis, def $(H_{0,t-2}^4) \geq 1$  and def $(H_{s-1,t}^4) \geq s$ . Hence M misses at least one vertex in  $H_{0,t}^4$  and at least s vertices in  $H_{s-1,t}^4$ . Therefore, def $(H_{s,t}^4) \geq s+1$ . The proof is complete.  $\Box$ 

Let X be a set of vertices in a graph G. Denote by  $\delta_G(X) = \min\{d_G(x) : x \in X\}$ and  $\Delta_G(X) = \max\{d_G(x) : x \in X\}$ . For two vertex sets X and Y in G, we say that X dominates Y if  $Y \subseteq N_G(X)$ . Recall that an induced matching M of G is an induced 1-regular subgraph of G. The *induced matching number* of G is defined as

 $\nu'(G) = \max\{|M| : M \text{ is an induced matching of } G\}.$ 

Next we present a technical lemma that has also been used implicitly in [1, 5].

**Lemma 2.4.** Let G be a bipartite graph with vertex partition sets X and Y. Suppose  $\delta(X) \ge 1$  and  $\Delta(Y) \le n$ . If  $|X| \ge n(p-1)+1$ , then  $\nu'(G) \ge p$ . Moreover, the lower bound of |X| is tight.

Proof. Since  $\delta(X) \geq 1$ , Y dominates X. Take a minimal subset  $Y' \subset Y$  dominating X, i.e.  $N_G(Y') = X$  and for any proper subset Y'' of Y',  $N_G(Y'') \neq X$ . Therefore, Y' is irredundant, i.e., every  $y \in Y'$  has a *private* neighbor  $x_y \in X$  (i.e.,  $N_G(x_y) \cap Y' = \{y\}$ ). Since  $\Delta(Y) \leq n$ , every  $y \in Y'$  dominates at most n vertices in X. Since  $|X| \geq n(p-1)+1$ , we have  $|Y'| \geq p$ . Hence, we can choose p vertices in Y' and a private neighbor for every one; they together form an induced matching  $M_p$ . Therefore,  $\nu'(G) \geq p$ .

Let  $G = (X \cup Y, E)$  be a bipartite graph with  $X = \{x_{ij} : i \in [p-1], j \in [n]\},$  $Y = \{y_i : i \in [p-1]\}, \text{ and } E = \{(x_{ij}, y_i) : i \in [p-1]\}.$  Then we have  $\delta(X) = 1, \Delta(Y) = n,$ but  $\nu'(G) = p - 1.$ 

For two disjoint vertex sets X and Y of a graph G, write G(X, Y) for the bipartite subgraph induced by the edges between X and Y in G. The following result is important in this article and has its own meaning on this subject.

**Theorem 2.5.** Let n and N be two integers greater than 3. Let G be a connected  $\{K_{1,n}, \check{K}_n^p : p \ge 1\}$ -free graph. If diam $(G) \le N$ , then there is an integer f(n, N) such that def $(G) \le f(n, N)$ .

*Proof.* Fix a vertex  $x_0 \in V(G)$ , for  $i \ge 0$ , define

$$X_i = \{ v \in V(G) : \operatorname{dist}(x_0, v) = i \}.$$

Since diam $(G) \leq N$ , we may assume that  $X_N$  is the last level. Then  $V(G) = \bigcup_{i=0}^N X_i$ . Now we construct a matching level by level. First, find a maximum matching  $M_N$  in  $G[X_N]$ and let  $Y_N = X_N \setminus V(M_N)$ . Then  $Y_N$  is a stable set. Next, find a maximum matching  $M'_N$  in  $G(X_{N-1}, Y_N)$  such that  $G_{N-1} = G - V(M_N) - V(M'_N)$  is still connected. Let  $Z_N = Y_N - V(M'_N)$  and  $B_{N-1} = N_{G_{N-1}}(Z_N)$ . Let  $H_N = G(Z_N, B_{N-1})$ .

**Claim 1.** For every vertex  $u \in B_{N-1}$ , there exist x and y in  $Z_N$  such that  $N_{G_{N-1}}(x) = N_{G_{N-1}}(y) = \{u\}$ , i.e., u has at least two private neighbors in  $Z_N$ .

Proof. Since  $u \in B_{N-1} = N_{G_{N-1}}(Z_N)$ , there is a vertex  $z \in Z_N$  adjacent to u. Then  $G_{N-1} - \{u, z\}$  is disconnected; otherwise,  $M'_N \cup \{uz\}$  will be a larger matching in  $G(X_{N-1}, Y_N)$  such that  $G - V(M_N) - V(M' \cup \{uz\})$  is still connected. Hence there is a vertex  $x \in Z_N$  such that  $N_{G_{N-1}}(x) = \{u\}$ . For the same reason,  $G_{N-1} - \{x, u\}$  is disconnected. Thus we can find another vertex  $y \in Z_N$  (possibly z) such that  $N_{G_{N-1}}(y) = \{u\}$ .

Claim 2. If  $N \ge 2$ , then  $\Delta_{H_N}(B_{N-1}) \le n-2$ .

Proof. Suppose to the contrary that  $\Delta_{H_N}(B_{N-1}) \ge n-1$ . Then the neighbors of a vertex  $u \in B_{N-1}$  with  $d_{H_N}(u) = n-1$  together with a neighbor of u in  $X_{N-2}$  induces a star  $K_{1,n}$ , a contradiction.

Set  $\alpha_1 = n$  and  $\beta_1 = R(n, n)$ . For  $i \ge 1$ , recursively define

$$\alpha_{i+1} = (n-2)(\beta_i - 1) + 1, \quad \beta_{i+1} = R(n, \alpha_{i+1}).$$

Claim 3.  $|Z_N| \le (\beta_{N-1} - 1)(n-2).$ 

Proof. Suppose to the contrary that  $|Z_N| > (\beta_{N-1} - 1)(n-2)$ . Then By Claim 2, we have  $|B_{N-1}| \ge |Z_N|/(n-2) > \beta_{N-1} - 1$ , i.e.,  $|B_{N-1}| \ge \beta_{N-1}$ . By Claim 1, for each vertex  $u \in B_{N-1}$ , u has at least two private neighbors x and y in  $Z_N$ . Recall that  $|B_{N-1}| \ge \beta_{N-1} = R(n, \alpha_{N-1})$ . By the Ramsey Theorem,  $G[B_{N-1}]$  contains a clique  $K_n$  or a stable set of order  $\alpha_{N-1}$ . If  $G[B_{N-1}]$  contains a clique  $K_n$ , then  $G[V(K_n) \cup N_{H_N}(V(K_n))]$  must contain an induced  $\check{K}_n^1$ , a contradiction.

Now, assume that there is a stable set  $A_{N-1}$  in  $G[B_{N-1}]$  with  $|A_{N-1}| \ge \alpha_{N-1}$ . Set  $H_{N-1} = G(A_{N-1}, X_{N-2})$ . Clearly,  $\delta_{H_{N-1}}(A_{N-1}) \ge 1$ . For the same reason as in Claim 2, we have  $\Delta_{H_{N-1}}(X_{N-2}) \le n-2$ . By Lemma 2.4,  $H_{N-1}$  has an induced matching  $M_{N-1}$  with  $|M_{N-1}| \ge \beta_{N-2}$ . Let  $B_{N-2} = V(M_{N-1}) \cap X_{N-2}$ . Then  $|B_{N-2}| = |M_{N-1}| \ge \beta_{N-2} = R(n, \alpha_{N-2})$ . Note that each vertex of  $V(M_{N-1}) \cap A_{N-1}$  has at least two private neighbors in  $Z_N$ . Hence, if  $G[B_{N-2}]$  contains a clique  $K_n$ , then G must contain an induced  $\check{K}_n^2$ , a contradiction. Thus there is a stable set  $A_{N-2}$  in  $G[B_{N-2}]$  with  $|A_{N-2}| \ge \alpha_{N-2}$ . Set  $H_{N-2} = G(A_{N-2}, X_{N-3})$ . With the same discussion as for  $H_{N-1}$ ,  $H_{N-2}$  has an induced matching  $M_{N-2}$  with  $|M_{N-2}| = |B_{N-3}| \ge \beta_{N-3}$  and a stable set  $A_{N-3} \subseteq B_{N-3}$  with  $|A_{N-3}| \ge \alpha_{N-3}$ , where  $B_{N-3} = V(M_{N-2}) \cap X_{N-3}$ . The above process is continued until we find a stable set  $A_1 \subseteq B_1 \subseteq X_1$  with  $|A_1| \ge \alpha_1 = n$ . However,  $\{x_0\} \cup A_1$  induces a copy of  $K_{1,n}$ , a contradiction. This claim holds.

Recall that every vertex of  $B_{N-1}$  has a private neighbor in  $Z_N$ . Hence  $H_N$  has an induced matching  $M''_N$  saturating all vertices of  $B_{N-1}$ . Let  $Z'_N = Z_N \setminus V(M''_N)$ . Then

$$|Z'_N| = |Z_N| - |B_{N-1}| \le \left(1 - \frac{1}{n-2}\right) |Z_N| \le (n-3)(\beta_{N-1} - 1).$$

Therefore, the matching  $M_N \cup M'_N \cup M''_N$  saturates the vertices in  $X_N$  except for  $|Z'_N| \leq (n-3)(\beta_{N-1}-1)$  vertices. Let  $G'_{N-1}$  be the graph obtained from G by deleting all vertices in  $V(M_N) \cup V(M'_N) \cup V(M''_N) \cup Z'_N$  and update  $X_{N-1}$  by  $X_{N-1} - (V(M'_N) \cup V(M''_N))$ . Then  $V(G'_{N-1}) = \bigcup_{i=0}^{N-1} X_i$  and  $G'_{N-1}$  is still connected.

We can continue the above process to  $G'_i$  for each i = N - 1, N - 2, ..., 2 such that we can construct the matching  $M_i \cup M'_i \cup M''_i$  in  $G'_i$  that saturates vertices in  $X_i$  except for  $|Z'_i| \leq (n-3)(\beta_{i-1}-1)$  vertices. Finally, let  $G'_1 = G'_2 - (V(M_2) \cup V(M'_2) \cup V(M''_2) \cup Z'_2)$  and update  $X_1$  by  $X_1 - (V(M'_2) \cup V(M''_2))$ .

Find a maximum matching  $M_1$  in  $G[X_1]$  and let  $Y_1 = X_1 - V(M_1)$ . Thus  $Y_1$  is a stable set dominated by  $x_0$ . Since G is  $K_{1,n}$ -free, we have  $|Y_1| \leq n-1$ . Let  $M'_1$  be a maximum matching of  $G[\{x_0\} \cup Y_1]$ . Set  $M''_1 = \emptyset$ .

Finally, we have a matching  $\bigcup_{i=1}^{N} (M_i \cup M'_i \cup M''_i)$  of G missing at most

$$n - 2 + (n - 3) \sum_{i=1}^{N-1} (\beta_i - 1) = f(n, N)$$

vertices. Thus  $def(G) \leq f(n, N)$ .

## 3 Proof of Theorem 1.6

The "only if" part: Suppose that  $\mathcal{H} \in B$ - def, i.e., there is a constant  $c = c(\mathcal{H})$  such that every connected  $\mathcal{H}$ -free graph G satisfies def(G) < c. Since  $\mathcal{H}$  is a finite family, the value  $h = \max\{|V(H)| : H \in \mathcal{H}\}$  is well-defined.

Choose s > c, and p > h as integers and t > h as an odd integer. By Lemma 2.1, we have def $(\check{H}_{s,t}^p) \ge s > c$ . Hence  $\check{H}_{s,t}^p$  is not  $\mathcal{H}$ -free, i.e., there is an  $H \in \mathcal{H}$  such that  $H \prec \check{H}_{s,t}^p$ . Since  $t > h \ge |V(H)|$  and  $p > h \ge |V(H)|$ , H contains at most one of the end vertex and the branch vertex of each  $R_i : i \in [s]$ . Thus there is some even integer  $n > h \ge |V(H)|$ (we choose even n just for convenience) such that  $H \prec T_n$  or  $H \prec F_n^2$ . We may assume n > c + 1 further.

If  $H \prec T_n$ , then  $\mathcal{H} \leq \{T_n\}$ . Since

$$\operatorname{def}(K_{1,n}) = n - 1 > c, \text{ and } \operatorname{def}(\check{K}_n^p) \ge n > c, \text{ for any } p \in \left[\frac{n}{2} - 1\right],$$

 $K_{1,n}$  and  $\check{K}_n^p: 1 \le p \le \frac{n}{2} - 1$  are not  $\mathcal{H}$ -free either. Therefore,  $\mathcal{H} \le \{K_{1,n}, T_n, \check{K}_n^p: 1 \le p \le \frac{n}{2} - 1\}$ .

For the other case  $H \prec F_n^2$ , we have  $\mathcal{H} \leq \{F_n^2\}$ . By Lemma 2.2 and 2.3,  $\operatorname{def}(H_{s,t}^j) = s+1 > c$  for  $j \in \{1,3,4\}$ . Thus  $H_{s,t}^j$  is not  $\mathcal{H}$ -free for  $j \in \{1,3,4\}$ . Hence there exists  $H_j \in \mathcal{H}$  such that  $H_j \prec H_{s,t}^j$ . Since  $|V(H_j)| \leq h$ , we have  $|\{i \in [s] : V(H_j) \cap \{v_i, w_i\} \neq \emptyset\}| \leq 1$ . This implies that  $H_j \leq F_n^j$ . By Lemma 2.1,  $\operatorname{def}(\check{H}_{s,t}^p) > s > c$  for  $1 \leq p \leq n-2$ . Thus  $\check{H}_{s,t}^p$  contains an induced subgraph  $H_p \in \mathcal{H}$  too. Since  $|H_p| \leq h < t$ ,  $|\{i \in [s] : v_i \in V(H_p)\}| \leq 1$  too. This implies that  $H_p \prec \check{F}_n^p$ . Note that both  $K_{1,n}$  and  $\check{K}_n^p : 1 \leq p \leq n-2$  are not  $\mathcal{H}$ -free. In summary, we have

$$\mathcal{H} \leq \{K_{1,n}, F_n^1, F_n^2, F_n^3, F_n^4, \check{K}_n^p, \check{F}_n^p : 1 \leq p \leq n-2\}.$$




Figure 3: The decomposition of G.

The proof of the "only if" part is completed.

The "if" part: Either

$$\mathcal{H} \le \{K_{1,n}, T_n, \check{K}_n^p : 1 \le p \le \frac{n}{2} - 1\}$$

or

$$\mathcal{H} \le \{K_{1,n}, F_n^1, F_n^2, F_n^3, F_n^4, \check{F}_n^p, \check{K}_n^p : 1 \le p \le n-2\}$$

implies that every  $\mathcal{H}$ -free graph G must be  $\{K_{1,n}, F_n^1, F_n^3, F_n^4, \check{F}_n^p, \check{K}_n^p : p \ge 1\}$ -free. Take a longest induced path  $P = u_1 u_2 \dots u_m$  of G. We may assume  $m \ge n^2 - n - 1$ , otherwise, by Theorem 2.5 the proof is completed since  $diam(G) \le m-1 < n^2$ . We consider the following decomposition of G. (See Figure 3.)

Let

$$X_0 = \{u_i : 1 \le i \le n \text{ or } m - n + 1 \le i \le m\}$$

Let  $Y_0 = \{u_i : n+1 \le i \le m-n\}$ . Set

$$X_1 = N_G(X_0) - V(P)$$
, and  $Y_1 = N_G(Y_0) - (X_0 \cup X_1)$ .

For  $i \geq 2$ , recursively define

$$X_i = N_G(X_{i-1}) - (Y_0 \cup Y_1) - \bigcup_{j=1}^{i-2} X_j.$$

Claim 4.  $X_{n+1} = \emptyset$ .

Proof. Suppose to the contrary that there exists a vertex  $x_{n+1}$  in  $X_{n+1}$ . Then we can recursively trace back to its ancestors  $x_{n+1-i} \in N_G(x_{n+2-i}) \cap X_{n+1-i}$  for each  $i \in [n+1]$ . Assume  $x_0 = u_k$  for some k with  $1 \le k \le n$  or  $m - n + 1 \le k \le m$ . By symmetry, we may assume  $k \in [n]$  and that such a k is as large as possible in [n]. Consider the vertices in  $N_G(x_1) \cap V(P)$ . Let  $j_1 = \min\{j \in [m] : x_1u_j \in E(G)\}$ . For  $p \ge 2$ , we recursively define

$$j_p = \min\{j : j_{p-1} + 2 \le j \le m, x_1 u_j \in E(G)\},\$$

and stop when

$$\{j: j_p + 2 \le j \le m, \ x_1 u_j \in E(G)\} = \emptyset$$

Let  $S = \{u_{j_p} : p \ge 1\}$ , and set s = |S|. Since  $j_p \ge j_{p-1} + 2$  and  $P = u_1 u_2 \dots u_m$  is an induced path, we have S is a stable set of G. Thus  $\{x_1, x_2\} \cup S$  induces a copy of  $K_{1,s+1}$  with center  $x_1$  in G. Since G is  $K_{1,n}$ -free, we have  $s + 1 \le n - 1$ .

We claim that  $s \ge 2$ . If s = 1, then  $\alpha(N(x_1) \cap V(P)) = 1$ . Since  $x_1u_k \in E(G)$ ,  $x_1u_i \notin E(G)$  for any  $i \ge k+2$ . If  $x_1u_{k+1} \in E(G)$ , then  $x_{n+1}x_n \dots x_1u_{k+1}u_{k+2}\dots u_m$  is an induced path with n + m + 1 - k > m vertices, a contradiction to the longest P. On the other hand, if  $x_1u_{k+1} \notin E(G)$ , then  $x_{n+1}x_n \dots x_1u_ku_{k+1}u_{k+2}\dots u_m$  is an induced path with n + m + 2 - k > m vertices, which is also a contradiction.

Let  $j^* = \max\{j \in [m] : x_1u_j \in E(G)\}$ . Note that  $j^* \in \{j_s, j_s+1\}$ . Let  $Q_1 = u_1u_2 \cdots u_{j_1}$ ,  $Q_p = u_{j_{p-1}+2}u_{j_{p-1}+3} \cdots u_{j_p}$  for  $p \in [2, s]$ , and  $Q_{s+1} = u_{j^*}u_{j^*+1} \cdots u_m$ . Then

$$V(P) - \bigcup_{p=1}^{s+1} V(Q_p) = \{u_{j_p+1} : p \in [s-1]\},\$$

and hence

$$m = |V(P)| = \left| V(P) - \bigcup_{p=1}^{s+1} V(Q_p) \right| + \left| \bigcup_{p=1}^{s+1} V(Q_p) \right|$$
$$\leq (s-1) + \sum_{p=1}^{s+1} |V(Q_p)|$$
$$= \sum_{p=1}^{s+1} (|V(Q_p)| + 1) - 2.$$

This implies that

$$\sum_{p=1}^{s+1} (|V(Q_p)| + 1) \ge m + 2 \ge n^2 - n + 1.$$

If  $|V(Q_p)| \le n-1$  for all  $p \in [s+1]$ , then

$$n^{2} - n + 1 \le \sum_{p=1}^{s+1} (|V(Q_{p})| + 1) \le (s+1)n \le (n-1)n,$$

a contradiction. Thus there is some  $q \in [s+1]$  such that  $|V(Q_q)| \ge n$ . Note that  $|N_G(x_1) \cap V(Q_q)| = 1$ . Write  $N_G(x_1) \cap V(Q_q) = \{u\}$ , i.e.,  $u = u_{j_q}$  if  $q \ne s+1$ , and  $u = u_{j^*} \in \{u_{j_s}, u_{j_s+1}\}$  if q = s+1. Since  $|S| \ge 2$ , we can choose anther vertex  $v \in S$  such that  $N_G(v) \cap V(Q_q) = \emptyset$  (e.g., choose  $v \in S - \{u\}$  if  $q \ne s+1$ , and, otherwise, let  $v = u_{j_1}$ ). Note that u is an end vertex of  $Q_q$ ,  $|V(Q_q)| \ge n$ . We have a copy of  $F_n^1 \prec G[\{v, x_1, x_2, \ldots, x_{n+1}\} \cup V(Q_q)]$ , a contradiction.

Now set

$$X = \bigcup_{i=0}^{n} X_i.$$

Recall that  $Y_0 = \{u_i : n+1 \le i \le m-n\}$ , and  $Y_1 = N_G(Y_0) - X$ . For  $i \ge 2$ , recursively define

$$Y_i = N_G(Y_{i-1}) - X - \bigcup_{j=0}^{i-2} Y_j.$$

Suppose N is the largest index with  $Y_N \neq \emptyset$  and  $Y_{N+1} = \emptyset$ . Set

$$Y = \bigcup_{i=0}^{N} Y_i.$$

Then  $V(G) = X \cup Y$ . Take vertices  $v, w \in Y_1$  (if any) such that

$$i = \min\{j : vu_j \in E(G)\} = \min\{j : wu_j \in E(G)\}.$$

Let

$$j = \max\{k : vu_k \in E(G)\}, \text{ and } j' = \max\{k : wu_k \in E(G)\}$$

By symmetry, we may assume  $j \ge j'$ . Note that  $Y_1 \cap N_G(X_0) = \emptyset$ . Hence  $n + 1 \le i \le j' \le j \le m - n$ .

Claim 5.  $vu_{i+1} \in E(G)$ .

*Proof.* Suppose to the contrary that  $vu_{i+1} \notin E(G)$ . Thus  $j \neq i+1$ . If j = i, then

$$\{u_{i-n}, u_{i-n+1}, \dots, u_{i-1}, u_i, v, u_{i+1}, u_{i+2}, \dots, u_{i+n}\}$$

induces a copy of  $F_n^1$  in G, a contradiction. If j = i + 2, then

$$\{u_{i-n+1}, u_{i-n+2}, \dots, u_i, v, u_{i+1}, u_{i+2}, u_{i+3}, \dots, u_{i+n+1}\}$$

induces a copy of  $F_n^3$  in G, a contradiction again. Now assume j > i + 2. Then

$$\{u_{i-n}, u_{i-n+1}, \dots, u_{i-1}, u_i, u_{i+1}, v, u_j, u_{j+1}, \dots, u_{j+n-2}\}$$

induces a copy of  $F_n^1$ , a contradiction. This completes the proof of the claim.

Claim 6. If there exists  $r \in N(v) \cap Y_2$ , then  $N_G(v) \cap Y_0 = \{u_i, u_{i+1}\}$ .

*Proof.* By Claim 5,  $j \ge i+1$ . It is sufficient to show that j = i+1. Suppose to the contrary that  $u_i$  and  $u_j$  are not adjacent in P. Then

$$\{u_{i-n+1}, u_{i-n}, \dots, u_{i-1}, u_i, v, r, u_j, u_{j+1}, \dots, u_{j+n-1}\}$$

induces a copy of  $F_n^1$  in G, a contradiction.

Claim 7.  $vw \in E(G)$ .

*Proof.* Suppose to the contrary that  $vw \notin E(G)$ . If j = j', then

$$\{u_{i-n+1}, u_{i-n+2}, \dots, u_i, v, w, u_j, u_{j+1}, \dots, u_{j+n-1}\}$$

induces a copy of  $F_n^3$  or  $F_n^4$  in G according to whether j = i + 1 or not, which is a contradiction. Thus j > j'. Therefore,  $wu_j \notin E(G)$ . Then

$$\{u_{i-n}, u_{i-n+1}, \ldots, u_{i-1}, u_i, w, v, u_j, u_{j+1}, \ldots, u_{j+n-2}\}$$

induces a copy of  $F_n^1$  in G, a contradiction too.

Since G is  $\check{F}_n^p$ -free for any  $p \ge 1$ , we know that for any  $y \in Y_i$ ,  $i \ge 1$ , the children of y in  $Y_{i+1}$  are pairwise adjacent. Otherwise, we can construct a copy of  $\check{F}_n^p$  for some  $p \ge 1$  as an induced subgraph in G according to Claim 6. In other words,  $N_G(y) \cap Y_{i+1}$  induces a clique in G for any  $y \in Y_i$ ,  $i \ge 1$ .

Now we construct a matching step by step from the last level  $Y_N$ . First, we find a maximum matching  $M_N$  in  $G[Y_N]$ . Then  $Z_N := Y_N \setminus V(M_N)$  is a stable set. Hence for every vertex  $z \in Z_N$ , there exists a  $p_z \in Y_{N-1}$  such that  $N(p_z) \cap Z_N = \{z\}$  as long as  $N \geq 2$ . Therefore,  $M'_N = \{zp_z : z \in Z_N \text{ and } p_z \in Y_{N-1}\}$  is a matching in G. Set  $Y'_{N-1} = Y_{N-1} - V(M'_N)$  and repeat the above process to vertices in  $Y'_{N-1}$ , we obtain matchings  $M_{N-1}$  and  $M'_{N-1}$ . Set  $Y'_{N-2} = Y_{N-2} - V(M'_{N-1})$ . This process is continued until we obtain  $Y'_1 = Y_1 - V(M'_2)$ . We can still find a maximum matching  $M_1$  in  $G[Y'_1]$  and a stable set  $Z_1 = Y'_1 - V(M_1)$ . By Claim 7, for every  $u_i \in Y_0$ , there is at most one vertex in  $Z_1$  satisfying  $i = \min\{j : v \in Z_1 \text{ and } vu_j \in E(G)\}$ , denote such a vertex by  $v_i$  (if any). By Claim 5,  $v_i u_{i+1} \in E(G)$ . We retain the two edges  $v_i u_i$  and  $v_i u_{i+1}$  and delete all the other edges incident to  $v_i$  for every possible  $v_i$ ; let L be the resulting subgraph of  $G[Y_0 \cup Z_1]$ .

Claim 8. L is connected and claw-free.

*Proof.* Note that  $V(L) = Y_0 \cup Z_1$  as  $Z_1 \subseteq N(Y_0)$ . By the construction of L, we have L is connected and  $d_L(v_i) = 2$  for each  $v_i \in Z_1$ . Thus no vertex in  $Z_1$  can be a claw center. For

each vertex  $u_i \in Y_0$ , we have  $d_L(u_i) \leq 4$  as the possible neighbors of  $u_i$  are  $u_{i-1}, u_{i+1} \in Y_0$ and  $v_{i-1}, v_i \in Z_1$ . As  $N_L(v_{i-1}) = \{u_{i-1}, u_i\}$  and  $N_L(v_i) = \{u_i, u_{i+1}\}, u_i$  cannot be a claw center either.

Claim 9.  $def(G[Y]) \leq 1$ .

*Proof.* By Claim 8 and Theorem 1.5, we have  $def(L) \leq 1$ . Since all vertices in  $Y \setminus (Z_1 \cup Y_0)$  has been saturated by the matching  $\bigcup_{i=1}^{N} (M_i \cup M'_i)$ , where we set  $M'_1 = \emptyset$  for convenience. Therefore, we have  $def(G[Y]) \leq 1$ .

Since  $G[X_0]$  consists of two disjoint paths, G[X] is composed of at most two components. According to Claim 4, each component has diameter at most 3n. By Theorem 2.5, def(G[X]) is bounded by a function b(n) = f(3n, n). Together with  $def(G[Y]) \leq 1$ , we know that def(G) is bounded. The proof of Theorem 1.6 is complete.

# 4 Remarks and Discussions

According to the proofs of Theorems 1.6 and 2.5, every  $\mathcal{H}$ -free graph G with  $\mathcal{H} \in B$ -def has deficiency bounded by a function dependent on the Ramsey number  $R(n, \alpha_i)$  and its diameter, where  $\alpha_i$  is defined recursively dependent on  $R(n, \alpha_{i-1})$ . Seemingly, it is difficult to determine the exact value of def(G) for an  $\mathcal{H}$ -free graph G with  $\mathcal{H} \in B$ - def. However, it will be very interesting to determine the family  $\mathcal{H} \in d$ -B- def for a fixed integer d as in Theorem 1.8.

With a similar discussion as in Theorem 1.9, we have further determined  $\mathcal{H} \in B$ -def with  $|\mathcal{H}| = 3$ . However, it is interminable to write down.

### Acknowledgements

This work was supported by the National Key Research and Development Program of China (2023YFA1010200), the National Natural Science Foundation of China (No. 12071453), and the Innovation Program for Quantum Science and Technology (2021ZD0302902).

### References

- S. Chiba and M. Furuya. Ramsey-type results for path covers and path partitions. Electron. J. Combin. 29(4) (2022), P4.8.
- [2] S. Chiba and M. Furuya. Ramsey-type problems on induced covers and induced partitions toward Gyárfás-Sumner conjecture. arXiv: 2205.14466.

- [3] I. Choi, M. Furuya, R. Kim and B. Park. A Ramsey-type theorem for the matching number regarding connected graphs. Discrete Math. 343 (2020), 111648.
- [4] R. Diestel. Graph theory, volume 173 of Graduate Texts in Mathematics. Springer, Berlin, fifth edition, 2017.
- [5] M. Furuya. Forbidden subgraphs for constant domination number. Discrete Math. Theor. 20(1) (2018), #19.
- [6] S. Fujita, K. Kawarabayashi, C. L. Lucchesi, K. Ota, M. Plummer and A. Saito. A pair of forbidden subgraphs and perfect matchings. J. Combin. Theory Ser. B 96 (2006), 315-324.
- [7] A. Gyárfás. Problems from the world surrounding perfect graphs. In Proc. Int. Conf. on Comb. Analysis and Applications (Pokrzywna, 1985). Zastos. Mat. 19, (1987) 413-441.
- [8] M. Jünger, W. R. Pulleyblank and G. Reinelt. On partitioning the edges of graphs into connected subgraphs, J. Graph Theory 9 (1985) 539-549.
- [9] H. A. Kierstead and S. G. Penrice. Radius two trees specify  $\chi$ -bounded classes. J. Graph Theory 18 (1994) 119-129.
- [10] M. Las Vergnas. A note on matchings in graphs, Colloque sur la Théorie des Graphes (Paris 1974), Cahiers Centre Études Rech. Opér. 17 (1975) 257-260.
- [11] V. V. Lozin. Graph parameters and Ramsey theory. Lecture Notes in Computer Science, 10765 (2018) 185-194.
- [12] V. Lozin and I. Razgon. Tree-width dichotomy. European J. Combin., 103 (2022), 103517.
- [13] M. D. Plummer and A. Saito. Forbidden subgraphs and bounds on the size of a maximum matching, J. Graph Theory 50 (2005) 1-12.
- [14] F. P. Ramsey. On a problem of formal logic. Proc. London Math. Soc. (2), 30(4), (1929), 264-286.
- [15] D. P. Sumner. 1-factors and antifactor sets, J. London Math. Soc. 13 (1976) 351-359.
- [16] A. Scott and P. Seymour. A survey of  $\chi$ -boundedness. J. Graph Theory 95 (2020) 473-504.
- [17] A. Scott, P. Seymour and S. Spirkl. Polynomial bounds for chromatic number. I. Excluding a biclique and an induced tree, J. Graph Theory 102 (2023) 458-471.