

Computation of weighted Bergman inner products on bounded symmetric domains and restriction to subgroups II

Ryosuke Nakahama^{*†}

*NTT Institute for Fundamental Mathematics,
NTT Communication Science Laboratories,
Nippon Telegraph and Telephone Corporation,
3-9-11 Midori-cho, Musashino-shi, Tokyo 180-8585, Japan*

October 14, 2024

Abstract

Let (G, G_1) be a symmetric pair of holomorphic type, and we consider a pair of Hermitian symmetric spaces $D_1 = G_1/K_1 \subset D = G/K$, realized as bounded symmetric domains in complex vector spaces $\mathfrak{p}_1^+ \subset \mathfrak{p}^+$ respectively. Then the universal covering group \tilde{G} of G acts unitarily on the weighted Bergman space $\mathcal{H}_\lambda(D) \subset \mathcal{O}(D)$ on D . Its restriction to the subgroup \tilde{G}_1 decomposes discretely and multiplicity-freely, and its branching law is given explicitly by Hua–Kostant–Schmid–Kobayashi’s formula in terms of the K_1 -decomposition of the space $\mathcal{P}(\mathfrak{p}_2^+)$ of polynomials on the orthogonal complement \mathfrak{p}_2^+ of \mathfrak{p}_1^+ in \mathfrak{p}^+ . The object of this article is to compute explicitly the inner product $\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \rangle_\lambda$ for $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+) \subset \mathcal{P}(\mathfrak{p}_2^+)$, $x = (x_1, x_2), z \in \mathfrak{p}^+ = \mathfrak{p}_1^+ \oplus \mathfrak{p}_2^+$, and to construct explicitly \tilde{G}_1 -intertwining operators (symmetry breaking operators) $\mathcal{H}_\lambda(D)|_{\tilde{G}_1} \rightarrow \mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+))$ from holomorphic discrete series representations of \tilde{G} to those of \tilde{G}_1 , which are unique up to constant multiple for sufficiently large λ . In this article, we treat the case $\mathfrak{p}^+, \mathfrak{p}_2^+$ are both simple of tube type and $\text{rank } \mathfrak{p}^+ = \text{rank } \mathfrak{p}_2^+$. When $\text{rank } \mathfrak{p}^+ = 3$, we treat all partitions \mathbf{k} , and when $\text{rank } \mathfrak{p}^+$ is general, we treat partitions of the form $\mathbf{k} = (k, \dots, k, k-l)$. This article is a continuation of the author’s previous article [34].

Keywords: weighted Bergman spaces; holomorphic discrete series representations; branching laws; intertwining operators; symmetry breaking operators; highest weight modules.

2020 Mathematics Subject Classification: 22E45; 43A85; 17C30; 33C67.

Contents

1	Introduction	2
2	Preliminaries	6
2.1	Hermitian positive Jordan triple systems	6
2.2	Jordan algebras	7
2.3	Structure groups and the Kantor–Koecher–Tits construction	9
2.4	Simultaneous Peirce decomposition	9

^{*}This work was supported by JST CREST Grant Number JPMJCR2113, Japan.

[†]Email: ryosuke.nakahama@ntt.com

2.5	Space of polynomials on Jordan triple systems	10
2.6	Projection of some polynomials	11
2.7	Holomorphic discrete series representations	19
2.8	Classification	22
2.9	Restriction to symmetric subgroups	23
3	Rank 3 case	27
3.1	Jordan algebra structure on $\text{Herm}(3, \mathbb{F})^{\mathbb{C}}$	27
3.2	Definition of some functions	29
3.3	Results on weighted Bergman inner products	35
3.4	Results on restriction of $\mathcal{H}_{\lambda}(D)$ to subgroups	46
4	General rank case	48
4.1	Definition of some functions	49
4.2	Results on weighted Bergman inner products	58
4.3	Results on restriction of $\mathcal{H}_{\lambda}(D)$ to subgroups	72

1 Introduction

The purpose of this article is to compute explicitly the weighted Bergman inner product of polynomials on some subspace of a bounded symmetric domain, and study the decomposition of the restriction of a holomorphic discrete series representation to some subgroup in detail. This article is a direct continuation of the author's previous article [34], and also related to the works [33, 35]. In this article, we construct symmetry breaking operators for pairs of representations that were missing in [34].

We consider a Hermitian symmetric space $D \simeq G/K$, realized as a bounded symmetric domain $D \subset \mathfrak{p}^+$ in a complex vector space \mathfrak{p}^+ centered at the origin. We assume that G is connected, and K is the isotropy subgroup at $0 \in \mathfrak{p}^+$. Let \tilde{G}, \tilde{K} denote the universal covering groups of G, K , let (τ, V) be a finite-dimensional representation of \tilde{K} , and we consider the homogeneous vector bundle $\tilde{G} \times_{\tilde{K}} V \rightarrow \tilde{G}/\tilde{K} \simeq D$. Then we can trivialize this bundle, and the space of holomorphic sections is identified with the space $\mathcal{O}(D, V) = \mathcal{O}_{\tau}(D, V)$ of V -valued holomorphic functions on D , on which \tilde{G} acts by

$$(\hat{\tau}(g)f)(x) := \tau(\kappa(g^{-1}, x))^{-1} f(g^{-1}x) \quad (g \in \tilde{G}, x \in D, f \in \mathcal{O}(D, V)),$$

by using some function $\tau: \tilde{G} \times D \rightarrow \tilde{K}^{\mathbb{C}}$. According to the choice of the representation (τ, V) of \tilde{K} , there may or may not exist a Hilbert subspace $\mathcal{H}_{\tau}(D, V) \subset \mathcal{O}_{\tau}(D, V)$ on which \tilde{G} acts unitarily. The classification of (τ, V) such that $\mathcal{H}_{\tau}(D, V)$ exists is given by [6, 14]. Especially, if the \tilde{G} -invariant inner product of $\mathcal{H}_{\tau}(D, V)$ is given by an integral on D , then this is called a *weighted Bergman inner product* and $(\hat{\tau}, \mathcal{H}_{\tau}(D, V))$ is called a *holomorphic discrete series representation*. When (τ, V) is of the form $(\tau, V) = (\chi^{-\lambda} \otimes \tau_0, V)$ with a suitably normalized character χ of \tilde{K} and a fixed representation (τ_0, V) of K , we write $\mathcal{H}_{\tau}(D, V) =: \mathcal{H}_{\lambda}(D, V) \subset \mathcal{O}_{\tau}(D, V) =: \mathcal{O}_{\lambda}(D, V)$. In addition, if $(\tau, V) = (\chi^{-\lambda}, \mathbb{C})$, then we write $\mathcal{H}_{\tau}(D, V) =: \mathcal{H}_{\lambda}(D) \subset \mathcal{O}_{\tau}(D, V) =: \mathcal{O}_{\lambda}(D)$.

Let us give a concrete example. We consider

$$G = SO^*(2s) := \left\{ g \in GL(2s, \mathbb{C}) \mid {}^t g \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} g = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, g \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} \bar{g} \right\}.$$

Then G acts transitively on

$$D := \{x \in \text{Alt}(s, \mathbb{C}) \mid I - xx^* \text{ is positive definite}\} \subset \mathfrak{p}^+ := \text{Alt}(s, \mathbb{C})$$

by the linear fractional transform

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot x := (ax + b)(cx + d)^{-1}.$$

Next let $\lambda \in \mathbb{C}$, and let (τ_0, V) be an irreducible representation of $K^\mathbb{C} = GL(s, \mathbb{C})$. Then the universal covering group \tilde{G} acts on $\mathcal{O}(D)$ by

$$\left(\tau_\lambda \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} \right) f \right) (x) := \det(cx + d)^{-\lambda/2} \tau_0(t(cx + d)) f((ax + b)(cx + d)^{-1}).$$

We note that $\det(cx + d)^{-\lambda/2}$ is not well-defined on $G \times D$ unless $\lambda \in 2\mathbb{Z}$, but is well-defined on the universal covering space $\tilde{G} \times D$. If $\lambda \in \mathbb{R}$ is sufficiently large, then this preserves the weighted Bergman inner product

$$\langle f, g \rangle_\lambda := C_\lambda \int_D (f(x), g(x))_{\tau_0} \det(I - xx^*)^{(\lambda - 2(s-1))/2} dx,$$

and the corresponding Hilbert space $\mathcal{H}_\lambda(D, V) \subset \mathcal{O}(D, V)$ gives a holomorphic discrete series representation, with the lowest \tilde{K} -type $\chi^{-\lambda} \otimes V = \det^{-\lambda/2} \otimes V$.

Next we consider an involution σ on G , and let $G_1 := (G^\sigma)_0$ be the identity component of the group of fixed points by σ . Without loss of generality we may assume σ stabilizes K , and let $K_1 := G_1 \cap K$. Then $D_1 := G_1 \cdot 0 \simeq G_1/K_1$ is either a complex submanifold or a totally real submanifold of $D \simeq G/K$. The pair (G, G_1) is called a symmetric pair of *holomorphic type* for the former case, and of *anti-holomorphic type* for the latter case (see [18, Section 3.4]). In the following we assume (G, G_1) is of holomorphic type. Then σ induces a holomorphic action on $\mathfrak{p}^+ \simeq T_0(G/K)$, and let $(\mathfrak{p}^+)^\sigma =: \mathfrak{p}_1^+$, $(\mathfrak{p}^+)^{-\sigma} =: \mathfrak{p}_2^+$, so that $\mathfrak{p}^+ = \mathfrak{p}_1^+ \oplus \mathfrak{p}_2^+$ holds. Now we consider the restriction of a holomorphic discrete series representation $\mathcal{H}_\tau(D, V)$ of \tilde{G} to the subgroup \tilde{G}_1 . Then the restriction $\mathcal{H}_\tau(D, V)|_{\tilde{G}_1}$ decomposes into a Hilbert direct sum of irreducible representations of \tilde{G}_1 , and this decomposition is given in terms of the decomposition of $\mathcal{P}(\mathfrak{p}_2^+) \otimes (V|_{\tilde{K}_1})$ under \tilde{K}_1 , where $\mathcal{P}(\mathfrak{p}_2^+)$ denotes the space of polynomials on \mathfrak{p}_2^+ . That is, if $\mathcal{P}(\mathfrak{p}_2^+) \otimes (V|_{\tilde{K}_1})$ is decomposed under \tilde{K}_1 as

$$\mathcal{P}(\mathfrak{p}_2^+) \otimes (V|_{\tilde{K}_1}) \simeq \bigoplus_j m(\tau, \rho_j)(\rho_j, W_j) \quad (m(\tau, \rho_j) \in \mathbb{Z}_{\geq 0}),$$

then $\mathcal{H}_\tau(D, V)|_{\tilde{G}_1}$ is decomposed abstractly into the Hilbert direct sum

$$\mathcal{H}_\tau(D, V)|_{\tilde{G}_1} \simeq \sum_j^{\oplus} m(\tau, \rho_j) \mathcal{H}_{\rho_j}(D_1, W_j)$$

(see Kobayashi [18, Lemma 8.8], [17, Section 8]. For earlier results, see also [15], [31]). We note that if the unitary subrepresentation $\mathcal{H}_\tau(D, V)$ is not holomorphic discrete, then its \tilde{K} -finite part $\mathcal{H}_\tau(D, V)_{\tilde{K}}$ may be strictly smaller than $\mathcal{O}_\tau(D, V)_{\tilde{K}} = \mathcal{P}(\mathfrak{p}^+) \otimes V$, and the above decomposition may not hold in general.

One of main problems on the restriction of representations is to determine the \tilde{G}_1 -intertwining operators

$$\begin{aligned} \mathcal{F}_{\tau\rho_j}^\downarrow : \mathcal{H}_\tau(D, V)|_{\tilde{G}_1} &\longrightarrow \mathcal{H}_{\rho_j}(D_1, W_j) \quad \text{or} \quad \mathcal{O}_\tau(D, V)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\rho_j}(D_1, W_j), \\ \mathcal{F}_{\tau\rho_j}^\uparrow : \mathcal{H}_{\rho_j}(D_1, W_j) &\longrightarrow \mathcal{H}_\tau(D, V)|_{\tilde{G}_1} \quad \text{or} \quad \mathcal{O}_{\rho_j}(D_1, W_j) \longrightarrow \mathcal{O}_\tau(D, V)|_{\tilde{G}_1}. \end{aligned}$$

$\mathcal{F}_{\tau\rho_j}^\downarrow$ is called a symmetry breaking operator (SBO) and $\mathcal{F}_{\tau\rho_j}^\uparrow$ is called a holographic operator, according to the terminology introduced in [24, 25] and [26] respectively. Such a problem is proposed by Kobayashi from the viewpoint of the representation theory (see [20]), and studied from various viewpoints for holomorphic discrete series, principal series, and complementary series representations. For example, differential SBOs for tensor products of holomorphic discrete series representations of $SL(2, \mathbb{R})$ are called Rankin–Cohen brackets. The study of such operators is initiated by [4, 39] from the viewpoint of automorphic forms, and their generalizations for larger groups are studied by e.g., [2, 3, 36, 37, 38]. Similarly, the study of differential SBOs for spherical principal series representations of $(G, G_1) = (SO_0(1, n+1), SO_0(1, n))$ is initiated by Juhl [16] from the viewpoint of conformal geometry on the sphere S^n , and generalized to the indefinite case [23]. Also, differential SBOs for symmetric pairs (G, G_1) of higher split rank are studied by e.g., Ibukiyama–Kuzumaki–Ochiai [13] and the author [33, 34, 35]. In addition, the complete classification of general (possibly non-differential) SBOs for spherical principal series representations of $(G, G_1) = (O(1, n+1), O(1, n))$ is given by Kobayashi–Speh [27], and general SBOs for various symmetric pairs are being studied by e.g., [10, 11, 12, 22, 28, 32]. On the other hand, the study of holographic operators is recently initiated by Kobayashi–Pevzner [26] and is studied by, e.g., [29, 33].

Especially, it is proved by [19, 24] that in the holomorphic setting, symmetry breaking operators $\mathcal{F}_{\tau\rho_j}^\downarrow : \mathcal{O}_\tau(D, V)|_{\tilde{G}_1} \rightarrow \mathcal{O}_{\rho_j}(D_1, W_j)$ are always given by differential operators, that is, $\mathcal{F}_{\tau\rho_j}^\downarrow$ are given by the form

$$\mathcal{F}_{\tau\rho_j}^\downarrow : \mathcal{O}_\tau(D, V)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\rho_j}(D_1, W_j), \quad f(x) = f(x_1, x_2) \longmapsto F_{\tau\rho_j}^\downarrow \left(\frac{\partial}{\partial x} \right) f(x) \Big|_{x_2=0}$$

for some operator-valued polynomials $F_{\tau\rho_j}^\downarrow(z) \in \mathcal{P}(\mathfrak{p}^-) \otimes \text{Hom}_{\mathbb{C}}(V, W_j)$ on the dual space \mathfrak{p}^- of \mathfrak{p}^+ (localness theorem), and $F_{\tau\rho_j}^\downarrow(z)$ are characterized as polynomial solutions of certain systems of differential equations (F-method). This F-method gives a new insight on the study of differential symmetry breaking operators, which unifies Rankin–Cohen brackets and Juhl’s operators, and gives new results [21, 23, 25]. Also, in [33], the author proved that when $\mathcal{H}_\tau(D, V) \subset \mathcal{O}_\tau(D, V)$ is holomorphic discrete, $F_{\tau\rho_j}^\downarrow(z)$ is given by the inner product

$$F_{\tau\rho_j}^\downarrow(z) = \left\langle e^{(x|z)_{\mathfrak{p}^+}}, K(x_2) \right\rangle_{\mathcal{H}_\tau(D, V), x}, \quad K(x_2) \in (\mathcal{P}(\mathfrak{p}_2^+) \otimes \text{Hom}_{\mathbb{C}}(\overline{V}, \overline{W_j}))^{\tilde{K}_1}$$

up to constant multiple. Here $(\cdot|\cdot)_{\mathfrak{p}^+}$ is a suitable non-degenerate pairing on $\mathfrak{p}^+ \times \mathfrak{p}^-$. Especially, if $\mathcal{H}_\tau(D, V) = \mathcal{H}_\lambda(D)$ is a unitary highest weight representation of scalar type, then since its restriction $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ decomposes multiplicity-freely by [18, Theorem A], symmetry breaking operators from $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ are unique up to constant multiple. On the other hand, if λ is not in the holomorphic discrete range, we do not know a priori whether symmetry breaking operators from $\mathcal{O}_\lambda(D)|_{\tilde{G}_1}$ are unique or not. Indeed, for tensor product case, the space of symmetry breaking operators from $\mathcal{O}_\lambda(D) \hat{\otimes} \mathcal{O}_\mu(D)$ may become greater than 1-dimensional for some $(\lambda, \mu) \in \mathbb{C}$ (see [25, Section 9], [35, Section 8]).

In the following, we assume G is simple, and consider the branching of the holomorphic discrete series representations of scalar type $\mathcal{H}_\tau(D, V) = \mathcal{H}_\lambda(D)$. We recall that $G_1 = (G^\sigma)_0$, and set $G_2 := (G^{\sigma\vartheta})_0$, where ϑ is the Cartan involution of G corresponding to K . We additionally assume that both G and the non-compact part of G_2 are of tube type, or equivalently, both \mathfrak{p}^+ and \mathfrak{p}_2^+ have unital Jordan algebra structures. Then one of the following holds.

- (1) \mathfrak{p}_2^+ is a direct sum of two simple Jordan algebras, and $\text{rank } \mathfrak{p}_2^+ = \text{rank } \mathfrak{p}^+$.
- (2) \mathfrak{p}_2^+ is simple, and $\text{rank } \mathfrak{p}_2^+ = \text{rank } \mathfrak{p}^+/2$.
- (3) \mathfrak{p}_2^+ is simple, and $\text{rank } \mathfrak{p}_2^+ = \text{rank } \mathfrak{p}^+$.

For Case (1), i.e., when (G, G_1, G_2) is one of

$$\begin{aligned} & (SO_0(2, d+2), \quad SO_0(2, d) \times SO(2), \quad SO_0(2, 2) \times SO(d) \quad), \\ & (Sp(r, \mathbb{R}), \quad U(r', r''), \quad Sp(r', \mathbb{R}) \times Sp(r'', \mathbb{R}) \quad), \\ & (U(r, r), \quad U(r', r'') \times U(r'', r'), \quad U(r', r') \times U(r'', r'') \quad), \\ & (SO^*(4r), \quad U(2r', 2r''), \quad SO^*(4r') \times SO^*(4r'') \quad), \\ & (E_{7(-25)}, \quad U(1) \times E_{6(-14)}, \quad SL(2, \mathbb{R}) \times Spin_0(2, 10) \quad) \end{aligned}$$

$(r = r' + r'')$, let $\mathfrak{p}_2^+ = \mathfrak{p}_{11}^+ \oplus \mathfrak{p}_{22}^+$. Then the decomposition of $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ is given by the form

$$\mathcal{H}_\lambda(D)|_{\tilde{G}_1} \simeq \sum_{(\mathbf{k}, \mathbf{l}) \in \mathbb{Z}_{++}^{r'} \times \mathbb{Z}_{++}^{r''}}^\oplus \mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_\mathbf{l}(\mathfrak{p}_{22}^+)),$$

where $\mathbb{Z}_{++}^r := \{\mathbf{k} = (k_1, \dots, k_r) \in \mathbb{Z}^r \mid k_1 \geq \dots \geq k_r \geq 0\}$, and $\varepsilon_1 \in \{1, 2\}$. If $\mathbf{k} = (k, \dots, k)$ and $\mathbf{l} = (l, \dots, l)$, then $\mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_{(k, \dots, k)}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_{(l, \dots, l)}(\mathfrak{p}_{22}^+))$ is of scalar type. In [33, 34], the author constructed the intertwining operators

$$\begin{aligned} \mathcal{F}_{\lambda, \mathbf{k}, \mathbf{l}}^\uparrow: \mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_\mathbf{l}(\mathfrak{p}_{22}^+))_{\tilde{K}_1} &\longrightarrow \mathcal{H}_\lambda(D)_{\tilde{K}}|_{\mathfrak{g}_1}, \\ \mathcal{F}_{\lambda, \mathbf{k}, \mathbf{l}}^\downarrow: \mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\longrightarrow \mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_\mathbf{l}(\mathfrak{p}_{22}^+)), \end{aligned}$$

when \mathbf{k} is of the form $\mathbf{k} = (k, \dots, k)$ ($k \in \mathbb{Z}_{\geq 0}$) and $\mathbf{l} \in \mathbb{Z}_{++}^{r''}$ is arbitrary. Next, for Case (2), i.e., when (G, G_1, G_2) is one of

$$\begin{aligned} & (SO_0(2, n), \quad SO_0(2, n-1), \quad SO_0(2, 1) \times SO(n-1) \quad), \\ & (Sp(2r_2, \mathbb{R}), \quad Sp(r_2, \mathbb{R}) \times Sp(r_2, \mathbb{R}), \quad U(r_2, r_2) \quad), \\ & (SU(2r_2, 2r_2), \quad Sp(2r_2, \mathbb{R}), \quad SO^*(4r_2) \quad), \end{aligned}$$

the decomposition of $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ is given by the form

$$\mathcal{H}_\lambda(D)|_{\tilde{G}_1} \simeq \sum_{\mathbf{k} \in \mathbb{Z}_{++}^{r_2}}^\oplus \mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+)).$$

If $\mathbf{k} = (k, \dots, k)$, then $\mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_{(k, \dots, k)}(\mathfrak{p}_2^+)) \simeq \mathcal{H}_{\varepsilon_1(\lambda+k)}(D_1)$ is of scalar type. In [33, 34], the author constructed the intertwining operators $\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow, \mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ when \mathbf{k} is of the form $\mathbf{k} = (k+1, \dots, k+1, k, \dots, k)$, so that $\mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+))$ is multiplicity-free under the maximal compact subgroup $\tilde{K}_1 \subset \tilde{G}_1$. Next, for Case (3), i.e., when (G, G_1, G_2) is one of

$$\begin{aligned} & (SO_0(2, n), \quad SO_0(2, n') \times SO(n''), \quad SO_0(2, n'') \times SO(n') \quad), \\ & (SU(r, r), \quad SO^*(2r), \quad Sp(r, \mathbb{R}) \quad), \\ & (SO^*(4r), \quad SO^*(2r) \times SO^*(2r), \quad U(r, r) \quad), \\ & (E_{7(-25)}, \quad SU(2, 6), \quad SU(2) \times SO^*(12) \quad) \end{aligned}$$

$(n'' \geq 3)$, the decomposition of $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ is given by the form

$$\mathcal{H}_\lambda(D)|_{\tilde{G}_1} \simeq \sum_{\mathbf{k} \in \mathbb{Z}_{++}^r}^\oplus \mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+)).$$

If $\mathbf{k} = (k, \dots, k)$, then $\mathcal{H}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_{(k, \dots, k)}(\mathfrak{p}_2^+)) \simeq \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1)$ is of scalar type. In [33], the author constructed the holographic operators $\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow$ when \mathbf{k} is of the form one of

- (3-a) $\mathbf{k} = (k+l, k, \dots, k)$ ($k, l \in \mathbb{Z}_{\geq 0}$),
- (3-b) $\mathbf{k} = (k, \dots, k, k-l)$ ($k, l \in \mathbb{Z}_{\geq 0}$, $k \geq l$),
- (3-c) $\mathbf{k} = (k_1, k_2, k_3) \in \mathbb{Z}_{++}^3$ (when $r = \text{rank } \mathfrak{p}^+ = 3$),

so that $\mathcal{H}_{\varepsilon_1\lambda}(D_1, \mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+))$ is multiplicity-free under $\tilde{K}_1 \subset \tilde{G}_1$ for classical G , and in [34], the author constructed the symmetry breaking operators $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ when \mathbf{k} is of the form (3-a). In any case, if $\mathbf{k} = (k, \dots, k)$, $\mathbf{l} = (l, \dots, l)$, then $\mathcal{F}_{\lambda, \mathbf{k}, \mathbf{l}}^\uparrow$, $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ are given by infinite-order differential operators whose symbols are given by multivariate ${}_0F_1$ functions, and $\mathcal{F}_{\lambda, \mathbf{k}, \mathbf{l}}^\downarrow$, $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ are given by finite-order differential operators whose symbols are given by multivariate ${}_2F_1$ polynomials (or equivalently, Heckman–Opdam’s hypergeometric polynomials of type BC , see [1]). In this article, we consider (G, G_1, G_2) of Case (3), and we construct the symmetry breaking operators $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ when \mathbf{k} is of the form (3-b) and (3-c). The result for (3-c) is conjectured in [35, Section 7], and this is verified in this paper.

In addition, in [35], the author determined the operator norms of $\mathcal{F}_{\lambda, \mathbf{k}, \mathbf{l}}^\downarrow$, $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ for all (\mathbf{k}, \mathbf{l}) , \mathbf{k} under a suitable normalization, and determined the Parseval–Plancherel type formula. Also, in the same paper, the author found polynomials $b_{\mathbf{k}, \mathbf{l}}(\lambda), b_{\mathbf{k}}(\lambda) \in \mathbb{C}[\lambda]$ such that

$$\begin{aligned} b_{\mathbf{k}, \mathbf{l}}(\lambda) \left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} & \quad (f(x_2) \in \mathcal{P}_\mathbf{k}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_\mathbf{l}(\mathfrak{p}_{22}^+)), \\ b_{\mathbf{k}}(\lambda) \left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} & \quad (f(x_2) \in \mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+)) \end{aligned}$$

are holomorphically continued for all $\lambda \in \mathbb{C}$ for all (\mathbf{k}, \mathbf{l}) , \mathbf{k} . This result gives some information on the branching of $\mathcal{O}_\lambda(D)|_{\tilde{K}}$ for λ such that $\mathcal{O}_\lambda(D)|_{\tilde{K}}$ is reducible. By [34], when $f(x_2) \neq 0$, the above give non-zero polynomials for all $\lambda \in \mathbb{C}$ if, for example, (G, G_1, G_2) is of Case (1) and $\mathbf{k} = (k, \dots, k)$, $\mathbf{l} \in \mathbb{Z}_{++}^{\min\{r', r''\}} \subset \mathbb{Z}_{++}^{r''}$, (G, G_1, G_2) is of Case (2) and $\mathbf{k} = (k+1, \dots, k+1, k, \dots, k)$, or (G, G_1, G_2) is of Case (3) with r even and $\mathbf{k} = (k+l, k, \dots, k)$ ((3-a)). However, for many other cases, the non-zeroness of the above formula is not proved. In this paper, we prove the non-zeroness for (G, G_1, G_2) of Case (3) (with r both even and odd) and \mathbf{k} is of the form either (3-a), (3-b) or (3-c).

This paper is organized as follows. In Section 2, we review Jordan triple systems, Jordan algebras and holomorphic discrete series representations. In Section 3, we consider the case $\text{rank } \mathfrak{p}^+ = 3$ and general $\mathbf{k} \in \mathbb{Z}_{++}^3$, i.e., consider (3-c) case. In Section 4, we consider \mathfrak{p}^+ of general rank, and \mathbf{k} of the form (3-b). For the convenience of the readers, we also rewrite the results for \mathbf{k} of the form (3-a) given in [34].

2 Preliminaries

In this section we review Jordan triple systems, Jordan algebras and holomorphic discrete series representations. We use almost the same notations as in [34, 35], and readers who read [34, 35] can skip this section, except for Section 2.6. For detail see, e.g., [7, Parts III, V], [9], [30], [40].

2.1 Hermitian positive Jordan triple systems

We consider a Hermitian positive Jordan triple system $(\mathfrak{p}^+, \mathfrak{p}^-, \{\cdot, \cdot, \cdot\}, \bar{\cdot})$, where \mathfrak{p}^\pm are finite-dimensional vector spaces over \mathbb{C} , with a non-degenerate bilinear form $(\cdot|\cdot)_{\mathfrak{p}^\pm} : \mathfrak{p}^\pm \times$

$\mathfrak{p}^\mp \rightarrow \mathbb{C}$, $\{\cdot, \cdot, \cdot\}: \mathfrak{p}^\pm \times \mathfrak{p}^\mp \times \mathfrak{p}^\pm \rightarrow \mathfrak{p}^\pm$ is a \mathbb{C} -trilinear map satisfying

$$\begin{aligned}\{x, y, z\} &= \{z, y, x\}, \\ \{u, v, \{x, y, z\}\} &= \{\{u, v, x\}, y, z\} - \{x, \{v, u, y\}, z\} + \{x, y, \{u, v, z\}\}, \\ (\{u, v, x\}|y)_{\mathfrak{p}^\pm} &= (x|\{v, u, y\})_{\mathfrak{p}^\pm}\end{aligned}$$

for any $u, x, z \in \mathfrak{p}^\pm$, $v, y \in \mathfrak{p}^\mp$, and $\bar{\cdot}: \mathfrak{p}^\pm \rightarrow \mathfrak{p}^\mp$ is a \mathbb{C} -antilinear involutive isomorphism satisfying $(x|\bar{x})_{\mathfrak{p}^\pm} \geq 0$ for any $x \in \mathfrak{p}^\pm$. Let $D, B: \mathfrak{p}^\pm \times \mathfrak{p}^\mp \rightarrow \text{End}_{\mathbb{C}}(\mathfrak{p}^\pm)$, $Q: \mathfrak{p}^\pm \times \mathfrak{p}^\pm \rightarrow \text{Hom}_{\mathbb{C}}(\mathfrak{p}^\mp, \mathfrak{p}^\pm)$, $Q: \mathfrak{p}^\pm \rightarrow \text{Hom}_{\mathbb{C}}(\mathfrak{p}^\mp, \mathfrak{p}^\pm)$ be the maps given by

$$\begin{aligned}D(x, y)z &= Q(x, z)y := \{x, y, z\}, \\ Q(x) &:= \frac{1}{2}Q(x, x), \\ B(x, y) &:= I_{\mathfrak{p}^\pm} - D(x, y) + Q(x)Q(y)\end{aligned}$$

for $x, z \in \mathfrak{p}^\pm$, $y \in \mathfrak{p}^\mp$, let $h = h_{\mathfrak{p}^\pm}: \mathfrak{p}^\pm \times \mathfrak{p}^\mp \rightarrow \mathbb{C}$ be the *generic norm*, which is a polynomial on $\mathfrak{p}^+ \times \mathfrak{p}^-$ irreducible on each simple Jordan triple subsystem, and write $B(x) := B(x, \bar{x})$, $h(x) := h(x, \bar{x})$. If \mathfrak{p}^+ is simple, then $B(x, y)$ and $h(x, y)$ are related as

$$h(x, y)^p = \text{Det}_{\mathfrak{p}^+} B(x, y) \quad (x \in \mathfrak{p}^+, y \in \mathfrak{p}^-) \quad (2.1)$$

for some $p \in \mathbb{Z}_{>0}$.

If $e \in \mathfrak{p}^+$ is a tripotent, i.e., $\{e, \bar{e}, e\} = 2e$, then $D(e, \bar{e}) \in \text{End}_{\mathbb{C}}(\mathfrak{p}^+)$ has the eigenvalues 0, 1, 2. For $j = 0, 1, 2$ let

$$\begin{aligned}\mathfrak{p}^+(e)_j &= \mathfrak{p}^+(\bar{e})_j := \{x \in \mathfrak{p}^+ \mid D(e, \bar{e})x = jx\} \subset \mathfrak{p}^+, \\ \mathfrak{p}^-(\bar{e})_j &= \mathfrak{p}^-(e)_j := \{x \in \mathfrak{p}^- \mid D(\bar{e}, e)x = jx\} \subset \mathfrak{p}^-, \end{aligned} \quad (2.2)$$

so that $\mathfrak{p}^\pm = \mathfrak{p}^\pm(e)_2 \oplus \mathfrak{p}^\pm(e)_1 \oplus \mathfrak{p}^\pm(e)_0$ holds (*Peirce decomposition*). A non-zero tripotent $e \in \mathfrak{p}^+$ is called *primitive* if $\mathfrak{p}^+(e)_2 = \mathbb{C}e$, and *maximal* if $\mathfrak{p}^+(e)_0 = \{0\}$. If $\mathfrak{p}^+(e)_2 = \mathfrak{p}^+$ holds for some (or equivalently any) maximal tripotent $e \in \mathfrak{p}^+$, then we say that \mathfrak{p}^+ is of *tube type*. Throughout the paper we assume that the bilinear form $(\cdot|\cdot)_{\mathfrak{p}^\pm}: \mathfrak{p}^\pm \times \mathfrak{p}^\mp \rightarrow \mathbb{C}$ is normalized such that $(e|\bar{e})_{\mathfrak{p}^+} = (\bar{e}|e)_{\mathfrak{p}^-} = 1$ holds for any primitive tripotent $e \in \mathfrak{p}^+$. If \mathfrak{p}^+ is simple, then under this normalization $(x|y)_{\mathfrak{p}^+}$ and $D(x, y)$ are related as

$$p(x|y)_{\mathfrak{p}^+} = \text{Tr}_{\mathfrak{p}^+} D(x, y) \quad (x \in \mathfrak{p}^+, y \in \mathfrak{p}^-) \quad (2.3)$$

with the same $p \in \mathbb{Z}_{>0}$ as in (2.1).

2.2 Jordan algebras

Next we consider a complex Jordan algebra $\mathfrak{n}^{+\mathbb{C}}$ with the Euclidean real form $\mathfrak{n}^+ \subset \mathfrak{n}^{+\mathbb{C}}$ and the unit element $e \in \mathfrak{n}^+$, that is, the product on $\mathfrak{n}^{+\mathbb{C}}$ satisfies

$$x \circ y = y \circ x, \quad x^2 \circ (x \circ y) = x \circ (x^2 \circ y) \quad (x, y \in \mathfrak{n}^{+\mathbb{C}}),$$

where $x^2 := x \circ x$, and there exists a symmetric bilinear form $(\cdot|\cdot)_{\mathfrak{n}^+}: \mathfrak{n}^{+\mathbb{C}} \times \mathfrak{n}^{+\mathbb{C}} \rightarrow \mathbb{C}$, which is positive definite on \mathfrak{n}^+ , satisfying

$$(x \circ y|z)_{\mathfrak{n}^+} = (x|y \circ z)_{\mathfrak{n}^+} \quad (x, y, z \in \mathfrak{n}^{+\mathbb{C}}).$$

Let $L: \mathfrak{n}^{+\mathbb{C}} \rightarrow \text{End}_{\mathbb{C}}(\mathfrak{n}^{+\mathbb{C}})$, $D_{\mathfrak{n}^+}, P: \mathfrak{n}^{+\mathbb{C}} \times \mathfrak{n}^{+\mathbb{C}} \rightarrow \text{End}_{\mathbb{C}}(\mathfrak{n}^{+\mathbb{C}})$, $P: \mathfrak{n}^{+\mathbb{C}} \rightarrow \text{End}_{\mathbb{C}}(\mathfrak{n}^{+\mathbb{C}})$ be the maps given by

$$\begin{aligned} L(x)y &:= x \circ y, \\ D_{\mathfrak{n}^+}(x, y)z &= P(x, z)y := 2(x \circ (y \circ z) + z \circ (y \circ x) - (x \circ z) \circ y), \\ P(x)y &:= 2x \circ (x \circ y) - x^2 \circ y = \frac{1}{2}P(x, x)y, \end{aligned}$$

let $\text{tr}_{\mathfrak{n}^+}$, $\det_{\mathfrak{n}^+}$ be the trace form and the determinant polynomial of $\mathfrak{n}^{+\mathbb{C}}$, and we normalize $(\cdot | \cdot)_{\mathfrak{n}^+}$ by $(x|e)_{\mathfrak{n}^+} = \text{tr}_{\mathfrak{n}^+}(x)$. If $\mathfrak{n}^{+\mathbb{C}}$ is simple of rank r , dimension n , then we have

$$(x|y)_{\mathfrak{n}^+} = \text{tr}_{\mathfrak{n}^+}(x \circ y) = \frac{r}{n} \text{Tr}(L(x \circ y)) = \frac{r}{2n} \text{Tr}(D_{\mathfrak{n}^+}(x, y)), \quad \det_{\mathfrak{n}^+}(x)^{2n/r} = \text{Det}(P(x)).$$

For $x \in \mathfrak{n}^{+\mathbb{C}}$, if $\det_{\mathfrak{n}^+}(x) \neq 0$, then the inverse x^{-1} and the adjugate element x^\sharp are given by

$$x^{-1} := P(x)^{-1}x, \quad x^\sharp := \det_{\mathfrak{n}^+}(x)x^{-1}.$$

Then the directional derivatives of x^{-1} and $\det_{\mathfrak{n}^+}(x)$ are given by

$$\left. \frac{d}{dt} \right|_{t=0} (x + ty)^{-1} = -P(x)^{-1}y, \quad \left. \frac{d}{dt} \right|_{t=0} \det_{\mathfrak{n}^+}(x + ty) = \det_{\mathfrak{n}^+}(x)(x^{-1}|y)_{\mathfrak{n}^+} = (x^\sharp|y)_{\mathfrak{n}^+}.$$

In addition, let

$$\begin{aligned} \Omega &:= (\text{connected component of } \{x \in \mathfrak{n}^+ \mid P(x) \text{ is positive definite}\} \text{ which contains } e) \\ &= (\text{connected component of } \{x \in \mathfrak{n}^+ \mid \det_{\mathfrak{n}^+}(x) > 0\} \text{ which contains } e) \end{aligned}$$

be the *symmetric cone*.

For a given Hermitian positive Jordan triple system \mathfrak{p}^\pm , if it is of tube type, then for a fixed maximal tripotent $e \in \mathfrak{p}^+$,

$$Q(\bar{e}): \mathfrak{p}^+ \longrightarrow \mathfrak{p}^-, \quad Q(e): \mathfrak{p}^- \longrightarrow \mathfrak{p}^+$$

are mutually inverse, and \mathfrak{p}^+ becomes a Jordan algebra by the product

$$x \circ y := \frac{1}{2}\{x, \bar{e}, y\} \quad (x, y \in \mathfrak{p}^+),$$

with the unit element $e \in \mathfrak{p}^+$, and the Euclidean real form

$$\mathfrak{n}^+ = \{x \in \mathfrak{p}^+ \mid Q(e)\bar{x} = x\}.$$

Then the corresponding operations and the generic norm are given by

$$\begin{aligned} D_{\mathfrak{n}^+}(x, y) &= D(x, Q(\bar{e})y), \quad P(x, y) = Q(x, y)Q(\bar{e}), \quad P(x) = Q(x)Q(\bar{e}), \\ (x|y)_{\mathfrak{n}^+} &= (x|Q(\bar{e})y)_{\mathfrak{p}^+} = (y|Q(\bar{e})x)_{\mathfrak{p}^+}, \quad h_{\mathfrak{n}^+}(x, y) = h_{\mathfrak{p}^+}(x, Q(\bar{e})y). \end{aligned}$$

Throughout the paper, for a Hermitian positive Jordan triple system \mathfrak{p}^\pm of tube type, when a maximal tripotent $e \in \mathfrak{p}^+$ is fixed, we omit $Q(e), Q(\bar{e})$, and identify $\mathfrak{p}^+ = \mathfrak{p}^-$ by the identity map, so that $\bar{\cdot}: \mathfrak{p}^\pm \rightarrow \mathfrak{p}^\mp = \mathfrak{p}^\pm$ is the complex conjugate with respect to the Euclidean real form $\mathfrak{n}^+ \subset \mathfrak{p}^+$, and the corresponding operations are

$$\begin{aligned} D_{\mathfrak{n}^+}(x, y) &= D(x, y), \quad P(x, y) = Q(x, y), \quad P(x) = Q(x), \\ (x|y)_{\mathfrak{n}^+} &= (x|y)_{\mathfrak{p}^+}, \quad h_{\mathfrak{n}^+}(x, y) = h_{\mathfrak{p}^+}(x, y). \end{aligned}$$

Conversely, a complex unital Jordan algebra $\mathfrak{n}^{+\mathbb{C}}$ with the Euclidean real form $\mathfrak{n}^+ \subset \mathfrak{n}^{+\mathbb{C}}$ becomes a Hermitian positive Jordan triple system by $\{x, y, z\} = D_{\mathfrak{n}^+}(x, y)z$ and by the complex conjugate $\bar{\cdot}: \mathfrak{n}^{+\mathbb{C}} \rightarrow \mathfrak{n}^{+\mathbb{C}}$ with respect to \mathfrak{n}^+ .

2.3 Structure groups and the Kantor–Koecher–Tits construction

In this subsection we consider some Lie algebras corresponding to Jordan triple systems \mathfrak{p}^\pm . For $l \in \text{End}_{\mathbb{C}}(\mathfrak{p}^+)$, let $\bar{l}, {}^t l \in \text{End}_{\mathbb{C}}(\mathfrak{p}^-)$, $l^* \in \text{End}_{\mathbb{C}}(\mathfrak{p}^+)$ be the elements given by $\bar{l}\bar{x} = \bar{l}x$, $(lx|\bar{y})_{\mathfrak{p}^+} = (x|{}^t l\bar{y})_{\mathfrak{p}^+} = (x|\bar{l}^*y)_{\mathfrak{p}^+}$ for $x, y \in \mathfrak{p}^+$. Then the *Structure group* $\text{Str}(\mathfrak{p}^+)$ and the *automorphism group* $\text{Aut}(\mathfrak{p}^+)$ are defined as

$$\begin{aligned}\text{Str}(\mathfrak{p}^+) &:= \{l \in GL_{\mathbb{C}}(\mathfrak{p}^+) \mid \{lx, {}^t l^{-1}y, lz\} = l\{x, y, z\} \quad (x, z \in \mathfrak{p}^+, y \in \mathfrak{p}^-)\}, \\ \text{Aut}(\mathfrak{p}^+) &:= \{k \in \text{Str}(\mathfrak{p}^+) \mid k^{-1} = k^*\}.\end{aligned}$$

Let $\mathfrak{str}(\mathfrak{p}^+) = \mathfrak{k}^{\mathbb{C}}$ and $\mathfrak{der}(\mathfrak{p}^+) = \mathfrak{k}$ denote the Lie algebras of $\text{Str}(\mathfrak{p}^+)$ and $\text{Aut}(\mathfrak{p}^+)$ respectively. Then $D(\mathfrak{p}^+, \mathfrak{p}^-) = \mathfrak{str}(\mathfrak{p}^+)$ holds.

Next we recall the *Kantor–Koecher–Tits construction*. As vector spaces let

$$\begin{aligned}\mathfrak{g}^{\mathbb{C}} &:= \mathfrak{p}^+ \oplus \mathfrak{k}^{\mathbb{C}} \oplus \mathfrak{p}^-, \\ \mathfrak{g} &:= \{(x, k, \bar{x}) \mid x \in \mathfrak{p}^+, k \in \mathfrak{k}\} \subset \mathfrak{g}^{\mathbb{C}},\end{aligned}$$

and give the Lie algebra structure on $\mathfrak{g}^{\mathbb{C}}$ by

$$[(x, k, y), (z, l, w)] := (kz - lx, [k, l] + D(x, w) - D(z, y), -{}^t kw + {}^t ly).$$

Then $\mathfrak{g}^{\mathbb{C}}$ becomes a Lie algebra and \mathfrak{g} becomes a real form of $\mathfrak{g}^{\mathbb{C}}$. Let $(\cdot|\cdot)_{\mathfrak{g}^{\mathbb{C}}} : \mathfrak{g}^{\mathbb{C}} \times \mathfrak{g}^{\mathbb{C}} \rightarrow \mathbb{C}$ be the $\mathfrak{g}^{\mathbb{C}}$ -invariant bilinear form normalized such that $(x|y)_{\mathfrak{g}^{\mathbb{C}}} = (x|y)_{\mathfrak{p}^+}$ holds for any $x \in \mathfrak{p}^+$, $y \in \mathfrak{p}^-$. We fix a connected complex Lie group $G^{\mathbb{C}}$ with the Lie algebra $\mathfrak{g}^{\mathbb{C}}$, and let $G, K^{\mathbb{C}}, K, P^+, P^- \subset G^{\mathbb{C}}$ be the connected closed subgroups corresponding to the Lie algebras $\mathfrak{g}, \mathfrak{k}^{\mathbb{C}}, \mathfrak{k}, \mathfrak{p}^+, \mathfrak{p}^- \subset \mathfrak{g}^{\mathbb{C}}$ respectively. Then $Ad|_{\mathfrak{p}^+} : K^{\mathbb{C}} \rightarrow \text{Str}(\mathfrak{p}^+)_0$ gives a covering map, and for $l \in K^{\mathbb{C}}, x \in \mathfrak{p}^+$, we abbreviate $Ad(l)x =: lx$.

Next suppose \mathfrak{p}^+ is of tube type. We fix a tripotent $e \in \mathfrak{p}^+$, regard \mathfrak{p}^+ as a Jordan algebra, and let $\mathfrak{n}^+ \subset \mathfrak{p}^+$ be the Euclidean real form. Also, let

$$\begin{aligned}\mathfrak{n}^- &:= \overline{\mathfrak{n}^+} = Q(\bar{e})\mathfrak{n}^+ \subset \mathfrak{p}^-, \\ \mathfrak{l} &:= D(\mathfrak{n}^+, \mathfrak{n}^-) = [\mathfrak{n}^+, \mathfrak{n}^-] \subset \mathfrak{k}^{\mathbb{C}}, \\ {}^c \mathfrak{g} &:= \mathfrak{n}^+ \oplus \mathfrak{l} \oplus \mathfrak{n}^- \subset \mathfrak{g}^{\mathbb{C}}.\end{aligned}$$

These become real forms of the right hand sides. Let ${}^c G \subset G^{\mathbb{C}}$ be the connected closed subgroup corresponding to the Lie algebra ${}^c \mathfrak{g} \subset \mathfrak{g}^{\mathbb{C}}$, and let $L := K^{\mathbb{C}} \cap {}^c G$, $K_L := L \cap K$, $\mathfrak{k}_L := \mathfrak{l} \cap \mathfrak{k}$. Then G and ${}^c G$ are isomorphic via the Cayley transform in $G^{\mathbb{C}}$, L acts transitively on the symmetric cone $\Omega \subset \mathfrak{n}^+$, and K_L acts on \mathfrak{n}^+ as Jordan algebra automorphisms. Also, for $l \in \mathfrak{k}^{\mathbb{C}} \subset \text{End}_{\mathbb{C}}(\mathfrak{p}^+) = \text{End}_{\mathbb{C}}(\mathfrak{n}^{+\mathbb{C}})$, we abbreviate $Q(e){}^t l Q(\bar{e}) =: {}^t l \in \mathfrak{k}^{\mathbb{C}}$, and extend it to the anti-automorphism on $K^{\mathbb{C}}$.

2.4 Simultaneous Peirce decomposition

In this subsection we assume \mathfrak{p}^+ is simple, or equivalently, the corresponding Lie algebra \mathfrak{g} is simple. We fix a Jordan frame $\{e_1, \dots, e_r\} \subset \mathfrak{p}^+$, i.e., a maximal set of primitive tripotents in \mathfrak{p}^+ satisfying $D(e_i, \bar{e_j}) = 0$ for $i \neq j$, where $r = \text{rank } \mathfrak{p}^+ = \text{rank}_{\mathbb{R}} \mathfrak{g}$. Using this, let

$$\begin{aligned}\mathfrak{p}_{ij}^+ &:= \{x \in \mathfrak{p}^+ \mid D(e_l, \bar{e_l})x = (\delta_{il} + \delta_{jl})x \quad (l = 1, \dots, r)\}, \quad \mathfrak{p}_{ij}^- := \overline{\mathfrak{p}_{ij}^+} \quad (1 \leq i \leq j \leq r), \\ \mathfrak{p}_{0j}^+ &:= \{x \in \mathfrak{p}^+ \mid D(e_l, \bar{e_l})x = \delta_{jl}x \quad (l = 1, \dots, r)\}, \quad \mathfrak{p}_{0j}^- := \overline{\mathfrak{p}_{0j}^+} \quad (1 \leq i \leq r),\end{aligned}$$

so that

$$\mathfrak{p}^\pm = \bigoplus_{\substack{0 \leq i \leq j \leq r \\ (i,j) \neq (0,0)}} \mathfrak{p}_{ij}^\pm$$

holds. This is called the *simultaneous Peirce decomposition*. Using this decomposition, we define integers (d, b, p, n) by

$$\begin{aligned} d &:= \dim \mathfrak{p}_{ij}^+ \quad (1 \leq i < j \leq r), & b &:= \dim \mathfrak{p}_{0j}^+ \quad (1 \leq j \leq r), \\ p &:= 2 + d(r-1) + b, & n &:= \dim \mathfrak{p}^+ = r + \frac{d}{2}r(r-1) + br. \end{aligned} \quad (2.4)$$

We note that if $r = 1$, then d is not determined uniquely, and any number is allowed. Then p coincides with the one in (2.1), (2.3).

2.5 Space of polynomials on Jordan triple systems

In this subsection we consider the space $\mathcal{P}(\mathfrak{p}^\pm)$ of polynomials on the Jordan triple system \mathfrak{p}^\pm , on which $K^\mathbb{C}$ acts by

$$\begin{aligned} (Ad|_{\mathfrak{p}^+}(l))^\vee f(x) &= f(l^{-1}x) & (l \in K^\mathbb{C}, f \in \mathcal{P}(\mathfrak{p}^+), x \in \mathfrak{p}^+), \\ (Ad|_{\mathfrak{p}^-}(l))^\vee f(y) &= f({}^tly) & (l \in K^\mathbb{C}, f \in \mathcal{P}(\mathfrak{p}^-), y \in \mathfrak{p}^-). \end{aligned}$$

We assume \mathfrak{p}^+ is simple, fix a Jordan frame $\{e_1, \dots, e_r\} \subset \mathfrak{p}^+$, and consider the tripotents $e^k := \sum_{j=1}^k e_j$. Then the subalgebras

$$\mathfrak{p}^+(e^k)_2 = \bigoplus_{1 \leq i \leq j \leq k} \mathfrak{p}_{ij}^+$$

are of tube type, and have Jordan algebra structures. Let $\mathfrak{n}^+(e^k) \subset \mathfrak{p}^+(e^k)_2$ be the Euclidean real forms, and we extend the determinant polynomials $\det_{\mathfrak{n}^+(e^k)}$ on $\mathfrak{p}^+(e^k)_2$ to polynomials on \mathfrak{p}^+ . We also write $e^r =: e$, $\mathfrak{n}^+(e^r) =: \mathfrak{n}^+$. Using these, for

$$\mathbf{m} \in \mathbb{Z}_{++}^r := \{\mathbf{m} = (m_1, \dots, m_r) \in \mathbb{Z}^r \mid m_1 \geq m_2 \geq \dots \geq m_r \geq 0\},$$

we define the polynomial $\Delta_{\mathbf{m}}^{\mathfrak{n}^+}(x)$ on \mathfrak{p}^+ by

$$\Delta_{\mathbf{m}}^{\mathfrak{n}^+}(x) := \prod_{k=1}^r \det_{\mathfrak{n}^+(e^k)}(x)^{m_k - m_{k+1}},$$

where we set $m_{r+1} := 0$, and let

$$\begin{aligned} \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+) &:= \text{span}_{\mathbb{C}}\{\Delta_{\mathbf{m}}^{\mathfrak{n}^+}(l^{-1}x) \mid l \in K^\mathbb{C}\} \subset \mathcal{P}(\mathfrak{p}^+), \\ \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^-) &:= \{\overline{f(\bar{y})} \mid f(x) \in \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)\} \subset \mathcal{P}(\mathfrak{p}^-). \end{aligned}$$

Then the following holds.

Theorem 2.1 (Hua–Kostant–Schmid, [7, Part III, Theorem V.2.1]). *Under the $K^\mathbb{C}$ -action, $\mathcal{P}(\mathfrak{p}^\pm)$ is decomposed into the sum of irreducible submodules as*

$$\mathcal{P}(\mathfrak{p}^\pm) = \bigoplus_{\mathbf{m} \in \mathbb{Z}_{++}^r} \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^\pm).$$

Next suppose \mathfrak{p}^+ is of tube type. For $\mu \in \mathbb{C}$, let

$$\underline{\mu}_r := (\underbrace{\mu, \dots, \mu}_r) \in \mathbb{C}^r,$$

and for $\mathbf{m} \in \mathbb{Z}_{++}^r + \underline{\mu}_r$, let

$$\mathcal{P}_{\mathbf{m}}(\mathfrak{p}^\pm) := \mathcal{P}_{\mathbf{m} - \underline{\mu}_r}(\mathfrak{p}^\pm) \det_{\mathfrak{n}^\pm}(x)^\mu \subset \mathcal{P}(\mathfrak{p}^\pm) \det_{\mathfrak{n}^\pm}(x)^\mu,$$

where $\det_{\mathfrak{n}^-}(x)$ is the determinant polynomial on $\mathfrak{p}^- = \mathfrak{n}^{-\mathbb{C}}$. Then $\mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+) \simeq \mathcal{P}_{-\mathbf{m}^\vee}(\mathfrak{p}^-)$ holds as a $\widetilde{K}^{\mathbb{C}}$ -module, where $\mathbf{m}^\vee := (m_r, \dots, m_1)$ and $\widetilde{K}^{\mathbb{C}}$ is the universal covering group of $K^{\mathbb{C}}$. In addition, for $\lambda \in \mathbb{C}$, $\mathbf{s} \in \mathbb{C}^r$, $\mathbf{m} \in (\mathbb{Z}_{\geq 0})^r$, let

$$\Gamma_r^d(\lambda + \mathbf{s}) := (2\pi)^{dr(r-1)/4} \prod_{j=1}^r \Gamma\left(\lambda + s_j - \frac{d}{2}(j-1)\right), \quad (2.5)$$

$$(\lambda + \mathbf{s})_{\mathbf{m}, d} := \prod_{j=1}^r \left(\lambda + s_j - \frac{d}{2}(j-1)\right)_{m_j}, \quad (2.6)$$

where $(\lambda)_m := \lambda(\lambda+1)(\lambda+2)\cdots(\lambda+m-1)$, and we abbreviate $\Gamma_r^d(\lambda + \underline{0}_r) =: \Gamma_r^d(\lambda)$, $\Gamma_r^d(0 + \mathbf{s}) =: \Gamma_r^d(\mathbf{s})$, $(\lambda + \underline{0}_r)_{\mathbf{m}, d} =: (\lambda)_{\mathbf{m}, d}$, $(0 + \mathbf{s})_{\mathbf{m}, d} =: (\mathbf{s})_{\mathbf{m}, d}$. Then the following hold.

Lemma 2.2 (Gindikin, [9, Lemma XI.2.3, Section IX.3]). *Let $\mathbf{m} \in \mathbb{Z}_{++}^r$, $f \in \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$.*

(1) *For $\operatorname{Re} \lambda > \frac{n}{r} - 1$, $w \in \Omega + \sqrt{-1}\mathfrak{n}^+$, we have*

$$\int_{\Omega} e^{-(z|w)_{\mathfrak{n}^+}} f(z) \det_{\mathfrak{n}^+}(z)^{\lambda - \frac{n}{r}} dz = \Gamma_r^d(\lambda + \mathbf{m}) f(w^{-1}) \det_{\mathfrak{n}^+}(w)^{-\lambda}.$$

(2) *For $\operatorname{Re} \lambda > \frac{2n}{r} - 1$, $z, a \in \Omega$, we have*

$$\frac{\Gamma_r^d(\lambda + \mathbf{m})}{(2\pi\sqrt{-1})^n} \int_{a + \sqrt{-1}\mathfrak{n}^+} e^{(z|w)_{\mathfrak{n}^+}} f(w^{-1}) \det_{\mathfrak{n}^+}(w)^{-\lambda} dw = f(z) \det_{\mathfrak{n}^+}(z)^{\lambda - \frac{n}{r}},$$

and this does not depend on the choice of $a \in \Omega$.

Lemma 2.3 ([9, Proposition VII.1.6]). *For $k \in \mathbb{Z}_{\geq 0}$, $\mathbf{m} \in \mathbb{Z}_{++}^r + \mathbb{C}\underline{1}_r := \bigcup_{\mu \in \mathbb{C}} (\mathbb{Z}_{++}^r + \underline{\mu}_r)$, $f(x) \in \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$, we have*

$$\begin{aligned} \det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^k f(x) &= \left(\frac{d}{2}(r-1) + 1 - k + \mathbf{m} \right)_{\underline{k}_r, d} f(x) \det_{\mathfrak{n}^+}(x)^{-k} \\ &= (-1)^{kr} (-\mathbf{m}^\vee)_{\underline{k}_r, d} f(x) \det_{\mathfrak{n}^+}(x)^{-k}. \end{aligned}$$

2.6 Projection of some polynomials

In this subsection, we assume $\mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}}$ is of tube type. For $\mathbf{m} \in \mathbb{Z}_{++}^r$, let

$$\operatorname{Proj}_{\mathbf{m}}^{\mathfrak{p}^+} : \mathcal{P}(\mathfrak{p}^+) \longrightarrow \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$$

be the orthogonal projection. Also, let $y \in \mathfrak{p}^+$ be an element of rank 1. Then for $\mathbf{k} \in \mathbb{Z}_{++}^r$, $l \in \mathbb{Z}_{\geq 0}$, $f(x) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}^+)$, as a function of x we have

$$(x|y)_{\mathfrak{n}^+}^l f(x) = \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r, |\mathbf{m}| = |\mathbf{k}| + l \\ k_{j-1} \geq m_j \geq k_j}} \operatorname{Proj}_{\mathbf{m}}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l f(x)), \quad (2.7)$$

where we set $k_0 := +\infty$. More generally, for $\mu \in \mathbb{C}$, $\mathbf{m} \in \mathbb{Z}_{++}^r + \underline{\mu}_r$, let

$$\text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+} : \mathcal{P}(\mathfrak{p}^+) \det_{\mathfrak{n}^+}(x)^\mu \longrightarrow \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+) = \mathcal{P}_{\mathbf{m} - \underline{\mu}_r}(\mathfrak{p}^+) \det_{\mathfrak{n}^+}(x)^\mu$$

be the orthogonal projection, with the same symbol.

Lemma 2.4. Suppose that $\mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}}$ is of tube type. We fix a primitive idempotent $e' \in \mathfrak{n}^+$, let $\mathfrak{p}^+(e')_2 = \mathbb{C}e' =: \mathfrak{p}'$, $\mathfrak{p}^+(e')_0 =: \mathfrak{p}''$ be as in (2.2), and for $x \in \mathfrak{p}^+$, let $x' \in \mathfrak{p}'$, $x'' \in \mathfrak{p}''$ be the orthogonal projections. Also, let $y \in \mathfrak{p}^+$ be an element of rank 1.

(1) For $\mu \in \mathbb{C}$, $\mathbf{m} \in \mathbb{Z}_{++}^r + \underline{\mu}_r$, $k, l \in \mathbb{Z}_{\geq 0}$, $f(x) \in \mathcal{P}(\mathfrak{p}^+) \det_{\mathfrak{n}^+}(x)^\mu$, we have

$$\text{Proj}_{\mathbf{m} + \underline{k}_r}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \det_{\mathfrak{n}^+}(x)^k f(x)) = \det_{\mathfrak{n}^+}(x)^k \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l f(x)).$$

(2) For $\mathbf{k} \in \mathbb{Z}_{++}^{r-1}$, $\mathbf{m} \in \mathbb{Z}_{++}^r$, $k, l \in \mathbb{Z}_{\geq 0}$ with $k \leq \min\{l, m_r\}$ and for $f(x'') \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}'')$, we have

$$\begin{aligned} & \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l f(x'')) \\ &= \frac{(-l)_k (-\mathbf{k}^\vee)_{\underline{k}_{r-1}, d}}{(-\mathbf{m}^\vee)_{\underline{k}_r, d}} \det_{\mathfrak{n}^+}(x)^k (e'|y)_{\mathfrak{n}^+}^k \text{Proj}_{\mathbf{m} - \underline{k}_r}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^{l-k} \det_{\mathfrak{n}^{++}}(x'')^{-k} f(x'')). \end{aligned}$$

(3) For $\mathbf{k} \in \mathbb{Z}_{++}^{r-1}$, $l \in \mathbb{Z}_{\geq 0}$ with $l \leq k_{r-1}$ and for $f(x'') \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}'')$, we have

$$\text{Proj}_{(\mathbf{k}, l)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l f(x'')) = \frac{(-\mathbf{k}^\vee)_{l_{r-1}, d}}{(-\frac{d}{2} - \mathbf{k}^\vee)_{l_{r-1}, d}} \det_{\mathfrak{n}^+}(x)^l (e'|y)_{\mathfrak{n}^+}^l \det_{\mathfrak{n}^{++}}(x'')^{-l} f(x'').$$

(4) For $\mathbf{k} \in \mathbb{Z}_{++}^{r-1}$, $l \in \mathbb{Z}_{\geq 0}$ with $l \geq k_1$ and for $f(x'') \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}'')$, we have

$$\text{Proj}_{(l, \mathbf{k})}^{\mathfrak{p}^+}((x|e')_{\mathfrak{n}^+}^l f(x'')) \Big|_{x=x'+x'' \in \mathfrak{p}^{+'} \oplus \mathfrak{p}^{++}} = \frac{(-l)_{\mathbf{k}, d}}{(-l - \frac{d}{2})_{\mathbf{k}, d}} (x'|e')_{\mathfrak{n}^+}^l f(x'').$$

(5) There exists an element $y \in \mathfrak{p}^+$ of rank 1 such that for $\mu \in \mathbb{C}$, $\mathbf{k}, \mathbf{m} \in \mathbb{Z}_{++}^r + \underline{\mu}_r$, $l \in \mathbb{Z}_{\geq 0}$ with $k_{j-1} \geq m_j \geq k_j$, $|\mathbf{m}| = |\mathbf{k}| + l$ we have

$$\text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^+}(x)) \neq 0.$$

Proof. (1) Clear from $\mathcal{P}_{\mathbf{m} + \underline{k}_r}(\mathfrak{p}^+) = \det_{\mathfrak{n}^+}(x)^k \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$.

(2) Suppose $k_r = 0$, $f(x) = f(x'') \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}'')$, and we apply $\det_{\mathfrak{n}^+}(\frac{\partial}{\partial x})^k$ to the both sides of (2.7). Then the right hand side is computed as

$$\begin{aligned} & \det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^k \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r, |\mathbf{m}| = |\mathbf{k}| + l \\ k_{j-1} \geq m_j \geq k_j}} \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l f(x'')) \\ &= \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r, |\mathbf{m}| = |\mathbf{k}| + l \\ k_{j-1} \geq m_j \geq k_j \\ k_{r-1} \geq m_r \geq k}} (-1)^{kr} (-\mathbf{m}^\vee)_{\underline{k}_r, d} \det_{\mathfrak{n}^+}(x)^{-k} \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l f(x'')). \end{aligned}$$

Similarly, for the left hand side, since y is of rank 1, the operator $\det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^k e^{(x|y)_{\mathfrak{n}^+}}$ is characterized by

$$\begin{aligned} & \det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^k e^{(x|y)_{\mathfrak{n}^+}} e^{(x|z)_{\mathfrak{n}^+}} = \det_{\mathfrak{n}^+} (y + z)^k e^{(x|y)_{\mathfrak{n}^+}} e^{(x|z)_{\mathfrak{n}^+}} \\ &= (\det_{\mathfrak{n}^+}(z) + (z^\sharp | y)_{\mathfrak{n}^+})^k e^{(x|y)_{\mathfrak{n}^+}} e^{(x|z)_{\mathfrak{n}^+}} \\ &= \sum_{l'=0}^k \sum_{l''=0}^{\infty} \binom{k}{l'} \frac{1}{l''!} \det_{\mathfrak{n}^+}(z)^{k-l'} (z^\sharp | y)_{\mathfrak{n}^+}^{l'} (x|y)_{\mathfrak{n}^+}^{l''} e^{(x|z)_{\mathfrak{n}^+}} \quad (z \in \mathfrak{p}^+), \end{aligned}$$

and by taking the homogeneous terms with respect to y , we have

$$\det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^k (x|y)_{\mathfrak{n}^+}^l = l! \sum_{l'+l''=l} \binom{k}{l'} \frac{1}{l''!} (x|y)_{\mathfrak{n}^+}^{l''} \det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^{k-l'} \left(\left(\frac{\partial}{\partial x} \right)^\sharp \Big| y \right)_{\mathfrak{n}^+}^{l'}.$$

Then since $\det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^{k-l'} f(x'')$ vanishes for $l' < k$, we get

$$\begin{aligned} & \det_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x} \right)^k (x|y)_{\mathfrak{n}^+}^l f(x'') = \frac{l!}{(l-k)!} (x|y)_{\mathfrak{n}^+}^{l-k} \left(\left(\frac{\partial}{\partial x} \right)^\sharp \Big| y \right)_{\mathfrak{n}^+}^k f(x'') \\ &= (-1)^k (-l)_k (x|y)_{\mathfrak{n}^+}^{l-k} (e'|y)_{\mathfrak{n}^+}^k \det_{\mathfrak{n}^{++}} \left(\frac{\partial}{\partial x''} \right)^k f(x'') \\ &= (-1)^{kr} (-l)_k (-\mathbf{k}^\vee)_{\underline{k}_{r-1}, d} (e'|y)_{\mathfrak{n}^+}^k (x|y)_{\mathfrak{n}^+}^{l-k} \det_{\mathfrak{n}^{++}} (x'')^{-k} f(x'') \\ &= (-1)^{kr} (-l)_k (-\mathbf{k}^\vee)_{\underline{k}_{r-1}, d} (e'|y)_{\mathfrak{n}^+}^k \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r, |\mathbf{m}|=|\mathbf{k}|+l \\ k_{j-1} \geq m_j \geq k_j \\ k_{r-1} \geq m_r \geq k}} \text{Proj}_{\mathbf{m}-\underline{k}_r}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^{l-k} \det_{\mathfrak{n}^{++}} (x'')^{-k} f(x'')). \end{aligned}$$

Comparing the both sides, for $m_r \geq k$ we have

$$\begin{aligned} & (-\mathbf{m}^\vee)_{\underline{k}_r, d} \det_{\mathfrak{n}^+} (x)^{-k} \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l f(x'')) \\ &= (-l)_k (-\mathbf{k}^\vee)_{\underline{k}_{r-1}, d} (e'|y)_{\mathfrak{n}^+}^k \text{Proj}_{\mathbf{m}-\underline{k}_r}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^{l-k} \det_{\mathfrak{n}^{++}} (x'')^{-k} f(x'')), \end{aligned}$$

and we get the desired formula.

(3) In (2), we set $\mathbf{m} = (\mathbf{k}, l)$, $k = l$. Then we have

$$\begin{aligned} & \text{Proj}_{(\mathbf{k}, l)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l f(x'')) \\ &= \frac{(-l)_l (-\mathbf{k}^\vee)_{\underline{l}_{r-1}, d}}{(-(l, \mathbf{k}^\vee))_{\underline{l}_r, d}} \det_{\mathfrak{n}^+} (x)^l (e'|y)_{\mathfrak{n}^+}^l \text{Proj}_{(\mathbf{k}, l)-\underline{l}_r}^{\mathfrak{p}^+} (\det_{\mathfrak{n}^{++}} (x'')^{-l} f(x'')) \\ &= \frac{(-l)_l (-\mathbf{k}^\vee)_{\underline{l}_{r-1}, d}}{(-l)_l (-\frac{d}{2} - \mathbf{k}^\vee)_{\underline{l}_{r-1}, d}} \det_{\mathfrak{n}^+} (x)^l (e'|y)_{\mathfrak{n}^+}^l \det_{\mathfrak{n}^{++}} (x'')^{-l} f(x'') \\ &= \frac{(-\mathbf{k}^\vee)_{\underline{l}_{r-1}, d}}{(-\frac{d}{2} - \mathbf{k}^\vee)_{\underline{l}_{r-1}, d}} \det_{\mathfrak{n}^+} (x)^l (e'|y)_{\mathfrak{n}^+}^l \det_{\mathfrak{n}^{++}} (x'')^{-l} f(x''). \end{aligned}$$

(4) Since the map

$$\mathcal{P}_{\mathbf{k}}(\mathfrak{p}^{++}) \longrightarrow \mathcal{P}(\mathfrak{p}^{+}) \otimes \mathcal{P}(\mathfrak{p}^{++}), \quad f(x'') \mapsto \text{Proj}_{(l, \mathbf{k})}^{\mathfrak{p}^+} ((x|e')_{\mathfrak{n}^+}^l f(x'')) \Big|_{x=x'+x'' \in \mathfrak{p}^{+} \oplus \mathfrak{p}^{++}}$$

is $\mathfrak{k}^{\mathbb{C}''} := [\mathfrak{p}^{+''}, \mathfrak{p}^{+''}]$ -equivariant, it is enough to prove when $f(x'') \in \mathcal{P}(\mathfrak{p}^{+''})$ is a lowest weight vector. We fix a Jordan frame $\{e_2, \dots, e_r\}$ of $\mathfrak{n}^{+''}$ so that $\{e_1 = e', e_2, \dots, e_r\}$ is a Jordan frame of \mathfrak{n}^+ , and define $\Delta_{\mathbf{k}}^{\mathfrak{n}^{+''}}(x'') \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}^{+''})$, $\Delta_{\mathbf{m}}^{\mathfrak{n}^+}(x) \in \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$ by using these Jordan frames. We prove

$$\text{Proj}_{(l,\mathbf{k})}((x|e')_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{+''}}(x'')) = \frac{(-l)_{\mathbf{k},d}}{(-l - \frac{d}{2})_{\mathbf{k},d}} \Delta_{(l,\mathbf{k})}^{\mathfrak{n}^+}(x) \quad (2.8)$$

by induction on $r = \text{rank } \mathfrak{p}^+$. When $\mathbf{k} = \underline{0}_{r-1}$ (“ $r = 1$ case”), this is trivial. Next we assume (2.8) for $r - 1$, and prove it for r . Then by (2) with $k = k_{r-1}$, $y = e'$, we have

$$\begin{aligned} & \text{Proj}_{(l,\mathbf{k})}((x|e')_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{+''}}(x'')) \\ &= \frac{(-l)_{k_{r-1}} (-\mathbf{k}^\vee)_{\underline{k_{r-1}}_{r-1},d}}{(-(\mathbf{k}^\vee, l))_{\underline{k_{r-1}}_r,d}} \\ &\quad \times \det_{\mathfrak{n}^+}(x)^{k_{r-1}} \text{Proj}_{(l,\mathbf{k})-\underline{k_{r-1}}_r}^{\mathfrak{p}^+}((x|e')_{\mathfrak{n}^+}^{l-k_{r-1}} \det_{\mathfrak{n}^{+''}}(x'')^{-k_{r-1}} \Delta_{\mathbf{k}}^{\mathfrak{n}^{+''}}(x'')) \\ &= \frac{(-l)_{k_{r-1}} (-\mathbf{k}^\vee)_{\underline{k_{r-1}}_{r-1},d}}{(-\mathbf{k}^\vee)_{\underline{k_{r-1}}_{r-1},d} (-l - \frac{d}{2}(r-1))_{k_{r-1},d}} \\ &\quad \times \det_{\mathfrak{n}^+}(x)^{k_{r-1}} \text{Proj}_{(l,\mathbf{k})-\underline{k_{r-1}}_r}^{\mathfrak{p}^+}((x|e')_{\mathfrak{n}^+}^{l-k_{r-1}} \Delta_{\mathbf{k}-\underline{k_{r-1}}_{r-1}}^{\mathfrak{n}^{+''}}(x'')) \\ &= \frac{(-l)_{k_{r-1}}}{(-l - \frac{d}{2}(r-1))_{k_{r-1}}} \det_{\mathfrak{n}^+}(x)^{k_{r-1}} \frac{(-l + k_{r-1})_{\mathbf{k}-\underline{k_{r-1}}_{r-1},d}}{(-l + k_{r-1} - \frac{d}{2})_{\mathbf{k}-\underline{k_{r-1}}_{r-1},d}} \Delta_{(l,\mathbf{k})-\underline{k_{r-1}}_r}^{\mathfrak{n}^+}(x) \\ &= \frac{(-l)_{k_{r-1}}}{(-l - \frac{d}{2}(r-1))_{k_{r-1}}} \frac{(-l)_{\mathbf{k},d}}{(-l - \frac{d}{2})_{\mathbf{k},d}} \frac{(-l - \frac{d}{2})_{\underline{k_{r-1}}_{r-1},d}}{(-l)_{\underline{k_{r-1}}_{r-1},d}} \Delta_{(l,\mathbf{k})}^{\mathfrak{n}^+}(x) \\ &= \frac{(-l)_{\mathbf{k},d}}{(-l - \frac{d}{2})_{\mathbf{k},d}} \frac{(-l)_{\underline{k_{r-1}}_r,d}}{(-l)_{\underline{k_{r-1}}_r,d}} \Delta_{(l,\mathbf{k})}^{\mathfrak{n}^+}(x) = \frac{(-l)_{\mathbf{k},d}}{(-l - \frac{d}{2})_{\mathbf{k},d}} \Delta_{(l,\mathbf{k})}^{\mathfrak{n}^+}(x), \end{aligned}$$

where we have applied the induction hypothesis (2.8) for $\mathfrak{p}^+(e_r)_0 = \mathfrak{p}^+(\sum_{j=1}^{r-1} e_j)_2 \subset \mathfrak{p}^+$ at the 3rd equality. This proves (2.8) for all r . Then since

$$\Delta_{(l,\mathbf{k})}^{\mathfrak{n}^+}(x)|_{x=x'+x'' \in \mathfrak{p}^{+'} \oplus \mathfrak{p}^{+''}} = (x'|e')_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{+''}}(x'')$$

holds, we get the desired formula.

(5) Let $\{e_1, \dots, e_r\} \subset \mathfrak{n}^+$ be the Jordan frame defining $\Delta_{\mathbf{k}}^{\mathfrak{n}^+}(x)$, and we take an element $y \in \mathfrak{p}^+$ of rank 1 such that $(e_j|y)_{\mathfrak{n}^+} \neq 0$ holds for all $j = 1, \dots, r$. We prove

$$\text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^+}(x)) \neq 0 \quad (2.9)$$

by induction on r . When $r = 1$ this is clear. Next we assume (2.9) for $r - 1$, and prove it for r . Then by (1) we have

$$\text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^+}(x)) = \det_{\mathfrak{n}^+}(x)^{k_r} \text{Proj}_{\mathbf{m}-\underline{k_r}_r}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}-\underline{k_r}}^{\mathfrak{n}^+}(x)),$$

and it suffices to prove (2.9) when $k_r = 0$. Then by applying (2) for $e' = e_r$, $k = m_r$ so that $\mathfrak{p}^{+''} = \mathfrak{p}^+(e_r)_0 = \mathfrak{p}^+(\sum_{j=1}^{r-1} e_j)_2$, we have

$$\begin{aligned} & \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^+}(x)) = \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{+''}}(x'')) \\ &= \frac{(-l)_{m_r} (-\mathbf{k}^\vee)_{\underline{m_r}_{r-1},d}}{(-\mathbf{m}^\vee)_{\underline{m_r}_r,d}} \det_{\mathfrak{n}^+}(x)^{m_r} (e_r|y)_{\mathfrak{n}^+}^{m_r} \text{Proj}_{\mathbf{m}-\underline{m_r}_r}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^{l-m_r} \Delta_{\mathbf{k}-\underline{m_r}_{r-1}}^{\mathfrak{n}^{+''}}(x'')), \end{aligned}$$

and it suffices to prove (2.9) when $k_r = m_r = 0$. Then for $k_r = 0$, by restricting

$$(x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'') = \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r, |\mathbf{m}|=|\mathbf{k}|+l \\ k_{j-1} \geq m_j \geq k_j}} \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'')),$$

to the subalgebra $\mathfrak{p}^{++} = \mathfrak{p}^+(e_r)_0 \subset \mathfrak{p}^+$, we get

$$\begin{aligned} & \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r, |\mathbf{m}|=|\mathbf{k}|+l \\ k_{j-1} \geq m_j \geq k_j \\ m_r=0}} \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'')) \Big|_{\mathfrak{p}^{++}} = (x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'') \Big|_{\mathfrak{p}^{++}} \\ & = (x''|y'')_{\mathfrak{n}^{++}}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'') = \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^{r-1}, |\mathbf{m}|=|\mathbf{k}|+l \\ k_{j-1} \geq m_j \geq k_j}} \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^{++}}((x''|y'')_{\mathfrak{n}^{++}}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'')), \end{aligned}$$

and hence for $k_r = m_r = 0$ we have

$$\text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'')) \Big|_{\mathfrak{p}^{++}} = \text{Proj}_{\mathbf{m}}^{\mathfrak{p}^{++}}((x''|y'')_{\mathfrak{n}^{++}}^l \Delta_{\mathbf{k}}^{\mathfrak{n}^{++}}(x'')).$$

By the induction hypothesis, this is non-zero. Hence (2.9) holds for all r . \square

Also, the following holds.

Proposition 2.5. Suppose that \mathfrak{p}^+ is of tube type and $\text{rank } \mathfrak{p}^+ \geq 2$. Let $k, l, m \in \mathbb{Z}_{\geq 0}$, $m \leq k, l$, and let $y, z \in \mathfrak{p}^+$ be elements of rank 1. Then we have

$$\begin{aligned} & \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\ & = \frac{(-1)^m}{(-k-l+m-\frac{d}{2}(r-1))_m m!} \sum_{j=m}^{\min\{k,l\}} \frac{(-k)_j (-l)_j}{(-k-l+2m-\frac{d}{2}(r-1)+1)_{j-m} (j-m)!} \\ & \quad \times (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j, \\ & \text{Proj}_{(k+l-m, m, \underline{0}_{r-2})}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x|z)_{\mathfrak{n}^+}^k) \\ & = \frac{(-1)^m}{(-k-l+m-\frac{d}{2})_m m!} \sum_{j=m}^{\min\{k,l\}} \frac{(-k)_j (-l)_j}{(-k-l+2m-\frac{d}{2}+1)_{j-m} (j-m)!} \\ & \quad \times (x|y)_{\mathfrak{n}^+}^{l-j} (x|z)_{\mathfrak{n}^+}^{k-j} ((x|y)_{\mathfrak{n}^+} (x|z)_{\mathfrak{n}^+} - (P(x)y|z)_{\mathfrak{n}^+})^j. \end{aligned}$$

Proof. (1) First we prove

$$\bigoplus_{j=0}^{\min\{k,l\}} \mathbb{R} (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j = \bigoplus_{m=0}^{\min\{k,l\}} \mathbb{R} \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k). \quad (2.10)$$

Indeed, by the definition of $\text{Proj}_{\mathbf{m}}^{\mathfrak{p}^+}$ we have

$$(x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k = \sum_{m=0}^{\min\{k,l\}} \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k).$$

We fix a primitive idempotent $e' \in \mathfrak{n}^+$, and let $\mathfrak{n}^+(e')_0 =: \mathfrak{n}^{++}$. Then since z is of rank 1, $z = ge'$ holds for some $g \in K^\mathbb{C}$. Then as in the proof of Lemma 2.4(2), for $j = 0, 1, \dots, \min\{k, l\}$ we have

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(x)^{j-k} \det_{\mathfrak{n}^+}\left(\frac{\partial}{\partial x}\right)^j \det_{\mathfrak{n}^+}(x)^k (x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k \\
&= \det_{\mathfrak{n}^+}(g^{-1}x)^{j-k} \det_{\mathfrak{n}^+}\left(t g \frac{\partial}{\partial x}\right)^j \det_{\mathfrak{n}^+}(g^{-1}x)^k (g^{-1}x|tgy)_{\mathfrak{n}^+}^l ((g^{-1}x)^{-1}|e')_{\mathfrak{n}^+}^k \\
&= \det_{\mathfrak{n}^+}(w)^{j-k} \det_{\mathfrak{n}^+}\left(\frac{\partial}{\partial w}\right)^j (w|tgy)_{\mathfrak{n}^+}^l \det_{\mathfrak{n}^{++}}(w'')^k \Big|_{w=g^{-1}x} \\
&= (-1)^{jr} (-l)_j (-k)_{\underline{j}_{r-1}, d} \det_{\mathfrak{n}^+}(w)^{j-k} (w|tgy)_{\mathfrak{n}^+}^{l-j} \det_{\mathfrak{n}^{++}}(w'')^{k-j} (e'|tgy)_{\mathfrak{n}^+}^j \Big|_{w=g^{-1}x} \\
&= (-1)^{jr} (-l)_j (-k)_{\underline{j}_{r-1}, d} (g^{-1}x|tgy)_{\mathfrak{n}^+}^{l-j} ((g^{-1}x)^{-1}|e')_{\mathfrak{n}^+}^{k-j} (tgy|e')_{\mathfrak{n}^+}^j \\
&= (-1)^{jr} (-l)_j (-k)_{\underline{j}_{r-1}, d} (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j. \tag{2.11}
\end{aligned}$$

On the other hand, for $j, m = 0, 1, \dots, \min\{k, l\}$ we have

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(x)^{j-k} \det_{\mathfrak{n}^+}\left(\frac{\partial}{\partial x}\right)^j \det_{\mathfrak{n}^+}(x)^k \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\
&= (-1)^{jr} (-(m, \underline{k}_{r-2}, k+l-m))_{\underline{j}_r, d} \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k), \tag{2.12}
\end{aligned}$$

and hence

$$\begin{aligned}
& (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j \in \mathbb{R} \det_{\mathfrak{n}^+}(x)^{j-k} \det_{\mathfrak{n}^+}\left(\frac{\partial}{\partial x}\right)^j \det_{\mathfrak{n}^+}(x)^k (x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k \\
&= \mathbb{R} \det_{\mathfrak{n}^+}(x)^{j-k} \det_{\mathfrak{n}^+}\left(\frac{\partial}{\partial x}\right)^j \det_{\mathfrak{n}^+}(x)^k \sum_{m=0}^{\min\{k, l\}} \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\
&\subset \bigoplus_{m=j}^{\min\{k, l\}} \mathbb{R} \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k).
\end{aligned}$$

Therefore, the left hand side of (2.10) is contained in the right hand side. Since both sides have the dimension $\min\{k, l\} + 1$, we get (2.10).

Next we determine $\text{Proj}_{(l, \underline{0}_{r-2}, -k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k)$. By (2.10) this is of the form

$$\text{Proj}_{(l, \underline{0}_{r-2}, -k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) = \sum_{j=0}^{\min\{k, l\}} a_j (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j$$

for some $a_j \in \mathbb{C}$. Then since

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(x)^k \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \in \mathcal{P}_{(k+l-m, \underline{k}_{r-2}, m)}(\mathfrak{p}^+), \\
& \det_{\mathfrak{n}^+}(x)^k (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} \in \mathcal{P}_{(\underline{k}_{r-1}, j)}(\mathfrak{p}^+)
\end{aligned}$$

vanish for $m, j \geq 1$ if x is not invertible, we have

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(x)^k \text{Proj}_{(l, \underline{0}_{r-2}, -k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) = (x|y)_{\mathfrak{n}^+}^l (x^\sharp|z)_{\mathfrak{n}^+}^k, \\
& \det_{\mathfrak{n}^+}(x)^k \sum_{j=0}^{\min\{k, l\}} a_j (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j = a_0 (x|y)_{\mathfrak{n}^+}^l (x^\sharp|z)_{\mathfrak{n}^+}^k
\end{aligned}$$

for non-invertible x , and hence we get $a_0 = 1$. To determine other a_j , we consider the differential operator

$$\mathcal{L} := \sum_{\alpha\beta} (P(x)e_\alpha^\vee | e_\beta^\vee)_{\mathfrak{n}^+} \frac{\partial^2}{\partial x_\alpha \partial x_\beta},$$

where $\{e_\alpha\}, \{e_\alpha^\vee\} \subset \mathfrak{n}^+$ are mutually dual bases of \mathfrak{n}^+ and $\frac{\partial}{\partial x_\alpha}$ is the directional derivative along e_α . Then by [9, Proposition VI.4.4], $\text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k)$ is the eigenfunction of \mathcal{L} with the eigenvalue

$$\begin{aligned} & (l-m)^2 - (l-m) + (m-k)^2 - (m-k)(1+d(r-1)) \\ &= 2m^2 - m(2k+2l+d(r-1)) + k^2 + l^2 + k(1+d(r-1)) - l, \end{aligned}$$

which are distinct for all $m = 0, 1, \dots, \min\{k, l\}$. Also, we have

$$\begin{aligned} & \mathcal{L}(x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j \\ &= (\mathcal{L}(x|y)_{\mathfrak{n}^+}^{l-j}) (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j + (x|y)_{\mathfrak{n}^+}^{l-j} (\mathcal{L}(x^{-1}|z)_{\mathfrak{n}^+}^{k-j}) (y|z)_{\mathfrak{n}^+}^j \\ &\quad + 2 \sum_{\alpha\beta} (P(x)e_\alpha^\vee | e_\beta^\vee)_{\mathfrak{n}^+} \left(\frac{\partial}{\partial x_\alpha} (x|y)_{\mathfrak{n}^+}^{l-j} \right) \left(\frac{\partial}{\partial x_\beta} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} \right) (y|z)_{\mathfrak{n}^+}^j \\ &= ((l-j)^2 - (l-j)) (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j \\ &\quad + ((j-k)^2 - (j-k)(1+d(r-1))) (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j \\ &\quad - 2(l-j)(k-j)(P(x)y|P(x)^{-1}z)_{\mathfrak{n}^+} (x|y)_{\mathfrak{n}^+}^{l-j-1} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j-1} (y|z)_{\mathfrak{n}^+}^j \\ &= (2j^2 - j(2k+2l+d(r-1)) + k^2 + l^2 + k(1+d(r-1)) - l) (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j \\ &\quad - 2(l-j)(k-j)(x|y)_{\mathfrak{n}^+}^{l-j-1} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j-1} (y|z)_{\mathfrak{n}^+}^{j+1}. \end{aligned}$$

Hence we have

$$\begin{aligned} 0 &= (\mathcal{L} - (k^2 + l^2 + k(1+d(r-1)) - l)) \text{Proj}_{(l, \underline{0}_{r-2}, -k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\ &= \sum_{j=0}^{\min\{k,l\}} a_j ((2j^2 - j(2k+2l+d(r-1))) (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j \\ &\quad - 2(l-j)(k-j)(x|y)_{\mathfrak{n}^+}^{l-j-1} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j-1} (y|z)_{\mathfrak{n}^+}^{j+1}) \\ &= 2 \sum_{j=0}^{\min\{k,l\}-1} \left(a_{j+1} \left(j - k - l - \frac{d}{2}(r-1) + 1 \right) (j+1) - a_j(j-k)(j-l) \right) \\ &\quad \times (x|y)_{\mathfrak{n}^+}^{l-j-1} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j-1} (y|z)_{\mathfrak{n}^+}^{j+1}, \end{aligned}$$

and since $a_0 = 1$, we have

$$a_j = \frac{(-k)_j (-l)_j}{(-k - l - \frac{d}{2}(r-1) + 1)_j j!},$$

that is, we get

$$\begin{aligned} & \text{Proj}_{(l, \underline{0}_{r-2}, -k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\ &= \sum_{j=0}^{\min\{k,l\}} \frac{(-k)_j (-l)_j}{(-k - l - \frac{d}{2}(r-1) + 1)_j j!} (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j. \end{aligned}$$

Finally we determine $\text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k)$ for $m \geq 1$. If $z = e' \in \mathfrak{n}^+$ is a primitive idempotent, then by Lemma 2.4 (2) we have

$$\begin{aligned}
& \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\
&= \det_{\mathfrak{n}^+}(x)^{-k} \text{Proj}_{(k+l-m, \underline{k}_{r-2}, m)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l \det_{\mathfrak{n}^{++}}(x'')^k) \\
&= \frac{(-l)_m (-k)_{\underline{m}_{r-1}, d}}{(-(m, \underline{k}_{r-2}, k+l-m))_{\underline{m}_r, d}} \\
&\quad \times \det_{\mathfrak{n}^+}(x)^{m-k} \text{Proj}_{(k+l-2m, \underline{k}-\underline{m}_{r-2}, 0)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^{l-m} \det_{\mathfrak{n}^{++}}(x'')^{k-m}) (e'|y)_{\mathfrak{n}^+}^m \\
&= \frac{(-l)_m (-k)_m (-k - \frac{d}{2})_{\underline{m}_{r-2}, d}}{(-m)_m (-k - \frac{d}{2})_{\underline{m}_{r-2}, d} (-k - l + m - \frac{d}{2}(r-1))_m} \\
&\quad \times \text{Proj}_{(l-m, \underline{0}_{r-2}, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^{l-m} (x^{-1}|z)_{\mathfrak{n}^+}^{k-m}) (y|z)_{\mathfrak{n}^+}^m \\
&= \frac{(-1)^m (-l)_m (-k)_m}{(-k - l + m - \frac{d}{2}(r-1))_m m!} \\
&\quad \times \sum_{j=0}^{\min\{k-m, l-m\}} \frac{(-k+m)_j (-l+m)_j}{(-k-l+2m-\frac{d}{2}(r-1)+1)_j j!} (x|y)_{\mathfrak{n}^+}^{l-m-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-m-j} (y|z)_{\mathfrak{n}^+}^{j+m} \\
&= \frac{(-1)^m}{(-k - l + m - \frac{d}{2}(r-1))_m m!} \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j}{(-k - l + 2m - \frac{d}{2}(r-1) + 1)_{j-m} (j-m)!}.
\end{aligned}$$

For general $z \in \mathfrak{p}^+$ of rank 1, when $z = ge'$ with $g \in K^\mathbb{C}$, by replacing x, y with $g^{-1}x, {}^tgy$ and applying the above, we get the desired formula.

(2) First we consider the case $\text{rank } \mathfrak{p}^+ = 2$. Then for $x, z \in \mathfrak{p}^+$ we have

$$(x|z)_{\mathfrak{n}^+} = (x^\sharp|z^\sharp)_{\mathfrak{n}^+} = \det_{\mathfrak{n}^+}(x)(x^{-1}|z^\sharp)_{\mathfrak{n}^+},$$

and z^\sharp is of rank 1 if z is of rank 1. Hence by (1) we have

$$\begin{aligned}
& \text{Proj}_{(k+l-m, m)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x|z)_{\mathfrak{n}^+}^k) = \det_{\mathfrak{n}^+}(x)^k \text{Proj}_{(l-m, m-k)}^{\mathfrak{p}^+}((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z^\sharp)_{\mathfrak{n}^+}^k) \\
&= \frac{(-1)^m}{(-k - l + m - \frac{d}{2})_m m!} \det_{\mathfrak{n}^+}(x)^k \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z^\sharp)_{\mathfrak{n}^+}^{k-j} (y|z^\sharp)_{\mathfrak{n}^+}^j}{(-k - l + 2m - \frac{d}{2} + 1)_{j-m} (j-m)!} \\
&= \frac{(-1)^m}{(-k - l + m - \frac{d}{2})_m m!} \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j (x|y)_{\mathfrak{n}^+}^{l-j} (x|z)_{\mathfrak{n}^+}^{k-j} (y|z^\sharp)_{\mathfrak{n}^+}^j \det_{\mathfrak{n}^+}(x)^j}{(-k - l + 2m - \frac{d}{2} + 1)_{j-m} (j-m)!}.
\end{aligned}$$

Hence it suffices to show

$$(x|y)_{\mathfrak{n}^+} (x|z)_{\mathfrak{n}^+} - (P(x)y|z)_{\mathfrak{n}^+} = (y|z^\sharp)_{\mathfrak{n}^+} \det_{\mathfrak{n}^+}(x). \quad (2.13)$$

Indeed, when $y = z$ we have

$$\begin{aligned}
(x|y)_{\mathfrak{n}^+}^2 - (P(x)y|y)_{\mathfrak{n}^+} &= (e|P(x^{1/2})y)_{\mathfrak{n}^+}^2 - (P(x^{1/2})y|P(x^{1/2})y)_{\mathfrak{n}^+} \\
&= 2 \det_{\mathfrak{n}^+}(P(x^{1/2})y) = 2 \det_{\mathfrak{n}^+}(x) \det_{\mathfrak{n}^+}(y) = \det_{\mathfrak{n}^+}(x)(y|y^\sharp)_{\mathfrak{n}^+},
\end{aligned}$$

and by polarization we get (2.13). Thus we get the desired formula when \mathfrak{p}^+ is of rank 2. For \mathfrak{p}^+ of general rank, since the both sides of the desired formula depend only on the rank 2 subalgebra containing y and z , this is reduced to the rank 2 case. \square

Remark 2.6. By (2.11), (2.12) and Proposition 2.5, for $i = 0, 1, \dots, \min\{k, l\}$ we have

$$\begin{aligned}
& (-l)_i (-k)_i (x|y)_{\mathfrak{n}^+}^{l-i} (x^{-1}|z)_{\mathfrak{n}^+}^{k-i} (y|z)_{\mathfrak{n}^+}^i \\
&= \sum_{m=i}^{\min\{k,l\}} (-m)_i \left(-k - l + m - \frac{d}{2}(r-1) \right)_i \text{Proj}_{(l-m, 0_{r-2}, m-k)}^{\mathfrak{p}^+} ((x|y)_{\mathfrak{n}^+}^l (x^{-1}|z)_{\mathfrak{n}^+}^k) \\
&= \sum_{m=i}^{\min\{k,l\}} \frac{(-1)^{m-i}}{(-k - l + i + m - \frac{d}{2}(r-1))_{m-i} (m-i)!} \\
&\quad \times \sum_{j=m}^{\min\{k,l\}} \frac{(-k)_j (-l)_j}{(-k - l + 2m - \frac{d}{2}(r-1) + 1)_{j-m} (j-m)!} (x|y)_{\mathfrak{n}^+}^{l-j} (x^{-1}|z)_{\mathfrak{n}^+}^{k-j} (y|z)_{\mathfrak{n}^+}^j,
\end{aligned}$$

that is, for $j = i+1, i+2, \dots, \min\{k, l\}$ we have

$$\begin{aligned}
& \sum_{m=i}^j \frac{(-1)^{m-i}}{(-k - l + i + m - \frac{d}{2}(r-1))_{m-i} (m-i)!} \frac{(-k)_j (-l)_j}{(-k - l + 2m - \frac{d}{2}(r-1) + 1)_{j-m} (j-m)!} \\
&= \sum_{m=0}^{j-i} \frac{(-1)^m (-k)_j (-l)_j}{(-k - l + 2i + m - \frac{d}{2}(r-1))_m m! (-k - l + 2i + 2m - \frac{d}{2}(r-1) + 1)_{j-i-m} (j-i-m)!} \\
&= 0.
\end{aligned}$$

In general, for $\mu \in \mathbb{C}$, $j \in \mathbb{Z}_{>0}$, we can directly verify

$$\begin{aligned}
& \sum_{m=0}^j \frac{(-1)^m}{(\mu+m)_m m! (\mu+2m+1)_{j-m} (j-m)!} = \sum_{m=0}^j \frac{(-1)^m (\mu)_m (\mu+2m) (\mu+j+m+1)_{j-m}}{(\mu)_{2j+1} m! (j-m)!} \\
&= \frac{(-1)^j}{(\mu)_{2j+1}} \sum_{m=0}^j \frac{(\mu)_m (\mu+2m)}{m!} \frac{(-\mu-2j)_{j-m}}{(j-m)!} = 0,
\end{aligned}$$

by comparing the coefficients for t^j of

$$\begin{aligned}
& \sum_{q=0}^{\infty} \sum_{m=0}^q \frac{(\mu)_m (\mu+2m)}{m!} \frac{(-\mu-2j)_{q-m}}{(q-m)!} t^q = \sum_{m=0}^{\infty} \frac{(\mu)_m (\mu+2m)}{m!} t^m \sum_{n=0}^{\infty} \frac{(-\mu-2j)_n}{n!} t^n \\
&= \mu(1+t)(1-t)^{-\mu-1} (1-t)^{\mu+2j} = \mu(1+t)(1-t)^{2j-1} = -\frac{\mu}{j} \sum_{q=0}^{\infty} \frac{(-2j)_q (-j+q)}{q!} t^q.
\end{aligned}$$

2.7 Holomorphic discrete series representations

In this subsection we review holomorphic discrete series representations. First we recall the bounded symmetric domain realization (Harish-Chandra realization) of the Hermitian symmetric space G/K via the Borel embedding,

$$\begin{array}{ccc}
G/K & \longrightarrow & G^{\mathbb{C}}/K^{\mathbb{C}} P^- \\
\downarrow \lambda & & \uparrow \exp \\
D^{\mathbb{C}} & \longrightarrow & \mathfrak{p}^+
\end{array}$$

where

$$\begin{aligned} D &= (\text{connected component of } \{x \in \mathfrak{p}^+ \mid B(x) \text{ is positive definite}\} \text{ which contains } 0) \\ &= (\text{connected component of } \{x \in \mathfrak{p}^+ \mid h(x) > 0\} \text{ which contains } 0). \end{aligned}$$

Here we write $B(x) := B(x, \bar{x})$, $h(x) := h(x, \bar{x})$. Then its Bergman–Shilov boundary $\Sigma \subset \partial D$ consists of all maximal tripotents in \mathfrak{p}^+ . For $g \in G^\mathbb{C}$, $x \in D$, if $g \exp(x) \in P^+ K^\mathbb{C} P^-$ holds, then we write

$$g \exp(x) = \exp(\pi^+(g, x)) \kappa(g, x) \exp(\pi^-(g, x)),$$

where $\pi^\pm(g, x) \in \mathfrak{p}^\pm$ and $\kappa(g, x) \in K^\mathbb{C}$. Then the map $\pi^+: G \times D \rightarrow D$ gives an action of G on D , and we abbreviate $\pi^+(g, x) =: gx$.

Next let (τ, V) be an irreducible holomorphic representation of the universal covering group $\tilde{K}^\mathbb{C}$ of $K^\mathbb{C}$, with the \tilde{K} -invariant inner product $(\cdot, \cdot)_\tau$. Then the universal covering group \tilde{G} of G acts on the space $\mathcal{O}(D, V) = \mathcal{O}_\tau(D, V)$ of V -valued holomorphic functions on D by

$$(\hat{\tau}(g)f)(x) := \tau(\kappa(g^{-1}, x))^{-1} f(g^{-1}x) \quad (g \in \tilde{G}, x \in D, f \in \mathcal{O}(D, V)),$$

where we lift the map $\kappa: G \times D \rightarrow K^\mathbb{C}$ to the universal covering spaces, and represent by the same symbol $\kappa: \tilde{G} \times D \rightarrow \tilde{K}^\mathbb{C}$. Its differential action is given by

$$(d\hat{\tau}(z, k, w)f)(x) = d\tau(k - D(x, w))f(x) + \frac{d}{dt} \Big|_{t=0} f(x - t(z + kx - Q(x)w)) \quad (2.14)$$

for $z \in \mathfrak{p}^+, k \in \mathfrak{k}^\mathbb{C}, w \in \mathfrak{p}^-$. This becomes a highest weight representation with the minimal \tilde{K} -type (τ, V) . If this contains a unitary subrepresentation $\mathcal{H}_\tau(D, V) \subset \mathcal{O}_\tau(D, V)$, then its reproducing kernel is proportional to $\tau(B(x, \bar{y}))$, and such unitary subrepresentation is unique. Here we lift the map $B: D \times \overline{D} \rightarrow \text{Str}(\mathfrak{p}^+)_0 \subset \text{End}_{\mathbb{C}}(\mathfrak{p}^+)$ to the universal covering space, and represent by the same symbol $B: D \times \overline{D} \rightarrow \tilde{K}^\mathbb{C}$. Especially, if the \tilde{G} -invariant inner product is given by the converging integral

$$\langle f, g \rangle_{\hat{\tau}} := C_\tau \int_D (\tau(B(x)^{-1})f(x), g(x))_\tau \text{Det}_{\mathfrak{p}^+}(B(x))^{-1} dx$$

(a *weighted Bergman inner product*), then $(\hat{\tau}, \mathcal{H}_\tau(D, V))$ is called a *holomorphic discrete series representation*. Here we normalize the Lebesgue measure dx on $D \subset \mathfrak{p}^+$ with respect to the inner product $(\cdot, \cdot)_{\mathfrak{p}^+}$, and determine the constant C_τ such that $\|v\|_{\hat{\tau}} = |v|_\tau$ holds for all constant functions $v \in V$.

Next, let $\chi: \tilde{K}^\mathbb{C} \rightarrow \mathbb{C}^\times$ be the character of $\tilde{K}^\mathbb{C}$ normalized such that

$$d\chi([x, y]) = (x|y)_{\mathfrak{p}^+} \quad (x \in \mathfrak{p}^+, y \in \mathfrak{p}^-), \quad (2.15)$$

so that $h(x, y) = \chi(B(x, y))$ holds, and we fix a representation (τ_0, V) of $K^\mathbb{C}$. When (τ, V) is of the form $(\tau, V) = (\chi^{-\lambda} \otimes \tau_0, V)$, we write $\mathcal{H}_\tau(D, V) = \mathcal{H}_\lambda(D, V) \subset \mathcal{O}_\tau(D, V) = \mathcal{O}_\lambda(D, V)$. In addition, if $(\tau, V) = (\chi^{-\lambda}, \mathbb{C})$, then we write $\mathcal{H}_\tau(D, V) = \mathcal{H}_\lambda(D) \subset \mathcal{O}_\tau(D, V) = \mathcal{O}_\lambda(D)$ and $\hat{\tau} = \tau_\lambda$.

In the rest of this subsection we assume \mathfrak{p}^+ is simple and $(\tau, V) = (\chi^{-\lambda}, \mathbb{C})$. Then $\mathcal{H}_\lambda(D)$ is holomorphic discrete if $\lambda > p - 1$, and then the inner product is given by

$$\begin{aligned} \langle f, g \rangle_\lambda &= \langle f, g \rangle_{\lambda, \mathfrak{p}^+} := C_\lambda \int_D f(x) \overline{g(x)} h(x)^{\lambda-p} dx, \\ C_\lambda &:= \frac{1}{\pi^n} \frac{\Gamma_r^d(\lambda)}{\Gamma_r^d(\lambda - \frac{n}{r})}, \end{aligned} \quad (2.16)$$

where $\Gamma_r^d(\lambda)$ is as in (2.5). We consider another inner product on $\mathcal{P}(\mathfrak{p}^+)$, called the *Fischer inner product* (see e.g. [9, Section XI.1]), defined by

$$\langle f, g \rangle_F = \langle f, g \rangle_{F, \mathfrak{p}^+} := \frac{1}{\pi^n} \int_{\mathfrak{p}^+} f(x) \overline{g(x)} e^{-(x|\bar{x})_{\mathfrak{p}^+}} dx = \overline{g\left(\frac{\partial}{\partial x}\right)} f(x) \Big|_{x=0}. \quad (2.17)$$

Here $\overline{g(\cdot)}$ is a holomorphic polynomial on \mathfrak{p}^- , and we normalize $\frac{\partial}{\partial x}$ with respect to the bilinear form $(\cdot|\cdot)_{\mathfrak{p}^+} : \mathfrak{p}^+ \times \mathfrak{p}^- \rightarrow \mathbb{C}$. Then the following holds.

Theorem 2.7 (Faraut–Korányi, [8], [7, Part III, Corollary V.3.9]). *For $\lambda > p - 1$, $\mathbf{m} \in \mathbb{Z}_{++}^r$, $f \in \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$, $g \in \mathcal{P}(\mathfrak{p}^+)$, we have*

$$\langle f, g \rangle_\lambda = \frac{1}{(\lambda)_{\mathbf{m}, d}} \langle f, g \rangle_F,$$

where $(\lambda)_{\mathbf{m}, d}$ is as in (2.6).

Since the reproducing kernel on $\mathcal{P}(\mathfrak{p}^+)$ with respect to $\langle \cdot, \cdot \rangle_F$ is given by $e^{(x|\bar{y})_{\mathfrak{p}^+}}$, the following holds.

Corollary 2.8. *For $\lambda > p - 1$, $\mathbf{m} \in \mathbb{Z}_{++}^r$, $f(x) \in \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$, we have*

$$\left\langle f(x), e^{(x|\bar{y})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} = \frac{1}{(\lambda)_{\mathbf{m}, d}} f(y).$$

Here the subscript x stands for the variable of integration. By this theorem, $\langle \cdot, \cdot \rangle_\lambda$ is meromorphically continued for all $\lambda \in \mathbb{C}$, and the \tilde{K} -finite part $\mathcal{O}_\lambda(D)_{\tilde{K}} = \chi^{-\lambda} \otimes \mathcal{P}(\mathfrak{p}^+)$ is reducible as a $(\mathfrak{g}, \tilde{K})$ -module if and only if λ is a pole of $\langle \cdot, \cdot \rangle_\lambda$. Especially, for $j = 1, 2, \dots, r$, $\lambda \in \frac{d}{2}(j-1) - \mathbb{Z}_{\geq 0}$,

$$M_j(\lambda) = M_j^{\mathfrak{g}}(\lambda) := \bigoplus_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r \\ m_j \leq \frac{d}{2}(j-1)-\lambda}} \chi^{-\lambda} \otimes \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+) \subset \chi^{-\lambda} \otimes \mathcal{P}(\mathfrak{p}^+) = \mathcal{O}_\lambda(D)_{\tilde{K}} \quad (2.18)$$

gives a $(\mathfrak{g}, \tilde{K})$ -submodule. Moreover, $\langle \cdot, \cdot \rangle_\lambda$ is positive definite on $\mathcal{P}(\mathfrak{p}^+)$ for $\lambda > \frac{d}{2}(r-1)$, and on $M_j^{\mathfrak{g}}(\frac{d}{2}(j-1))$ for $\lambda = \frac{d}{2}(j-1)$, $j = 1, \dots, r$, that is, $\mathcal{O}_\lambda(D)_{\tilde{K}}$ contains a unitary submodule $\mathcal{H}_\lambda(D)$ if λ sits in the *Wallach set*

$$\lambda \in \left\{ 0, \frac{d}{2}, d, \dots, \frac{d}{2}(r-1) \right\} \cup \left(\frac{d}{2}(r-1), \infty \right), \quad (2.19)$$

and its \tilde{K} -finite part is given by

$$\mathcal{H}_\lambda(D)_{\tilde{K}} = \begin{cases} \chi^{-\lambda} \otimes \mathcal{P}(\mathfrak{p}^+) & (\lambda > \frac{d}{2}(r-1)), \\ M_j^{\mathfrak{g}}(\frac{d}{2}(j-1)) & (\lambda = \frac{d}{2}(j-1), j = 1, 2, \dots, r). \end{cases}$$

In addition, the quotient module $\mathcal{O}_\lambda(D)_{\tilde{K}}/M_r(\lambda)$ ($\lambda \in \frac{d}{2}(r-1) - \mathbb{Z}_{\geq 0}$) also gives an infinitesimally unitary module (see [8]). Also by the corollary, for $\mathbf{k} \in \mathbb{Z}_{++}^r$, $f(x) \in \mathcal{P}(\mathfrak{p}^+)$, $(\lambda)_{\mathbf{k}, d} \langle f(x), e^{(x|\bar{y})_{\mathfrak{p}^+}} \rangle_{\lambda, x}$ is holomorphically continued for all $\lambda \in \mathbb{C}$ if and only if

$$f(x) \in \bigoplus_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^r \\ m_j \leq k_j \ (j=1,\dots,r)}} \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$$

holds, and if this is satisfied, then for $j = 1, \dots, r$,

$$f(x) \in M_j(\lambda) \quad \text{holds if } \lambda \in \frac{d}{2}(j-1) - k_j - \mathbb{Z}_{\geq 0}.$$

2.8 Classification

A simple Hermitian positive Jordan triple system \mathfrak{p}^\pm is isomorphic to one of the following.

$$\begin{array}{lll} \mathfrak{p}^\pm = \mathbb{C}^n & (n \geq 3), & \text{Sym}(r, \mathbb{C}), \quad M(q, s; \mathbb{C}), \\ & \text{Alt}(s, \mathbb{C}), & \text{Herm}(3, \mathbb{O})^\mathbb{C}, \quad M(1, 2; \mathbb{O})^\mathbb{C}. \end{array}$$

Here $\text{Sym}(r, \mathbb{C})$ and $\text{Alt}(s, \mathbb{C})$ denote the spaces of symmetric and alternating matrices over \mathbb{C} respectively, and $\text{Herm}(3, \mathbb{O})^\mathbb{C}$ denotes the complexification of the space of 3×3 Hermitian matrices over the octonions \mathbb{O} . Then the corresponding Lie groups G and their maximal compact subgroups K are given by

$$(G, K) = \begin{cases} (SO_0(2, n), SO(2) \times SO(n)) & (\mathfrak{p}^\pm = \mathbb{C}^n), \\ (Sp(r, \mathbb{R}), U(r)) & (\mathfrak{p}^\pm = \text{Sym}(r, \mathbb{C})), \\ (SU(q, s), S(U(q) \times U(s))) & (\mathfrak{p}^\pm = M(q, s; \mathbb{C})), \\ (SO^*(2s), U(s)) & (\mathfrak{p}^\pm = \text{Alt}(s, \mathbb{C})), \\ (E_{7(-25)}, U(1) \times E_6) & (\mathfrak{p}^\pm = \text{Herm}(3, \mathbb{O})^\mathbb{C}), \\ (E_{6(-14)}, U(1) \times \text{Spin}(10)) & (\mathfrak{p}^\pm = M(1, 2; \mathbb{O})^\mathbb{C}) \end{cases}$$

(up to covering), and the numbers (n, r, d, b, p) (see (2.4)) are given by

$$(n, r, d, b, p) = \begin{cases} (n, 2, n - 2, 0, n) & (\mathfrak{p}^\pm = \mathbb{C}^n, n \geq 3), \\ \left(\frac{1}{2}r(r+1), r, 1, 0, r+1\right) & (\mathfrak{p}^\pm = \text{Sym}(r, \mathbb{C})), \\ (qs, \min\{q, s\}, 2, |q-s|, q+s) & (\mathfrak{p}^\pm = M(q, s; \mathbb{C})), \\ \left(\frac{1}{2}s(s-1), \frac{s}{2}, 4, 0, 2(s-1)\right) & (\mathfrak{p}^\pm = \text{Alt}(s, \mathbb{C}), s: \text{even}), \\ \left(\frac{1}{2}s(s-1), \lfloor \frac{s}{2} \rfloor, 4, 2, 2(s-1)\right) & (\mathfrak{p}^\pm = \text{Alt}(s, \mathbb{C}), s: \text{odd}), \\ (27, 3, 8, 0, 18) & (\mathfrak{p}^\pm = \text{Herm}(3, \mathbb{O})^\mathbb{C}), \\ (16, 2, 6, 4, 12) & (\mathfrak{p}^\pm = M(1, 2; \mathbb{O})^\mathbb{C}). \end{cases}$$

Here, if $r = 1$, then d is not determined uniquely, and any number is allowed. When $b = 0$, \mathfrak{p}^\pm is of tube type, and has a Jordan algebra structure. That is, $\mathfrak{p}^\pm = \mathbb{C}^n$, $\text{Sym}(r, \mathbb{C})$, $M(r, \mathbb{C})$, $\text{Alt}(2r, \mathbb{C})$ and $\text{Herm}(3, \mathbb{O})^\mathbb{C}$ are of tube type. Their corresponding Euclidean real forms $\mathfrak{n}^+ \subset \mathfrak{p}^+$ are given by

$$\mathfrak{n}^+ = \begin{cases} \mathbb{R}^{1, n-1} \\ \text{Sym}(r, \mathbb{R}) \\ \text{Herm}(r, \mathbb{C}) \\ \text{Herm}(r, \mathbb{H}) \\ \text{Herm}(3, \mathbb{O}) \end{cases} \subset \mathfrak{p}^+ = \begin{cases} \mathbb{C}^n \\ \text{Sym}(r, \mathbb{C}) \\ M(r, \mathbb{C}) \\ \text{Alt}(2r, \mathbb{C}) \\ \text{Herm}(3, \mathbb{O})^\mathbb{C}, \end{cases}$$

where $\mathbb{R}^{1, n-1} \subset \mathbb{C}^n$ is a real form on which the quadratic form $\det_{\mathfrak{n}^+}(x)$ has the signature $(1, n-1)$. For these cases $p = \frac{2n}{r} = d(r-1) + 2$ holds.

Next we fix the parametrization of finite-dimensional irreducible representations of $\widetilde{GL}(s, \mathbb{C})$ and $\text{Spin}(n, \mathbb{C})$ ($n \geq 3$). We take a basis $\{\epsilon_j\}_{j=1}^s \subset \mathfrak{h}^{\mathbb{C}\vee}$ of the dual space of a Cartan subalgebra $\mathfrak{h}^\mathbb{C} \subset \mathfrak{gl}(s, \mathbb{C})$ such that the positive root system is given by $\{\epsilon_i - \epsilon_j \mid 1 \leq i < j \leq s\}$. For $\mathbf{m} \in \mathbb{C}^s$ with $m_j - m_{j+1} \in \mathbb{Z}_{\geq 0}$, let $V_{\mathbf{m}}^{(s)}$ be the irreducible representation of $\widetilde{GL}(s, \mathbb{C})$ with the highest weight $\sum_j m_j \epsilon_j$, and let $V_{\mathbf{m}}^{(s)\vee}$ be that with the

lowest weight $-\sum_j m_j \epsilon_j$. If $\mathbf{m} \in \mathbb{Z}^r$, then $V_{\mathbf{m}}^{(s)}, V_{\mathbf{m}}^{(s)\vee}$ are reduced to the representations of $GL(s, \mathbb{C})$. Similarly, we take a basis $\{\epsilon_j\}_{j=1}^{\lfloor n/2 \rfloor} \subset \mathfrak{h}^{\mathbb{C}\vee}$ of the dual space of a Cartan subalgebra $\mathfrak{h}^{\mathbb{C}} \subset \mathfrak{so}(n, \mathbb{C})$ such that the positive root system is given by

$$\begin{aligned} \{\epsilon_i \pm \epsilon_j \mid 1 \leq i < j \leq n/2\} & \quad (n: \text{even}), \\ \{\epsilon_i \pm \epsilon_j \mid 1 \leq i < j \leq \lfloor n/2 \rfloor\} \cup \{\epsilon_j \mid 1 \leq j \leq \lfloor n/2 \rfloor\} & \quad (n: \text{odd}). \end{aligned}$$

For $\mathbf{m} \in \mathbb{Z}^{\lfloor n/2 \rfloor} \cup (\mathbb{Z} + \frac{1}{2})^{\lfloor n/2 \rfloor}$ with $m_1 \geq \dots \geq m_{n/2-1} \geq |m_{n/2}|$ for even n , $m_1 \geq \dots \geq m_{\lfloor n/2 \rfloor} \geq 0$ for odd n , let $V_{\mathbf{m}}^{[n]}$ be the irreducible representation of $Spin(n, \mathbb{C})$ with the highest weight $\sum_j m_j \epsilon_j$, and let $V_{\mathbf{m}}^{[n]\vee}$ be that with the lowest weight $-\sum_j m_j \epsilon_j$. If $\mathbf{m} \in \mathbb{Z}^{\lfloor n/2 \rfloor}$, then $V_{\mathbf{m}}^{[n]}, V_{\mathbf{m}}^{[n]\vee}$ are reduced to the representations of $SO(n, \mathbb{C})$. Under this notation, the \tilde{K} -type decompositions of the holomorphic discrete series representations of scalar type

$$\mathcal{O}_\lambda(D)_{\tilde{K}} = \chi^{-\lambda} \otimes \mathcal{P}(\mathfrak{p}^+) = \chi^{-\lambda} \otimes \bigoplus_{\mathbf{m} \in \mathbb{Z}_{++}^r} \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+)$$

are given as

$$\begin{aligned} \chi^{-\lambda} &\simeq \begin{cases} \mathbb{C}_{-\lambda} \boxtimes V_{(0, \dots, 0)}^{[n]\vee} & (\mathfrak{p}^+ = \mathbb{C}^n, n \geq 3), \\ V_{(\lambda, \dots, \lambda)}^{(r)\vee} & (\mathfrak{p}^+ = \text{Sym}(r, \mathbb{C})), \\ V_{\left(\frac{s\lambda}{q+s}, \dots, \frac{s\lambda}{q+s}\right)}^{(q)\vee} \boxtimes V_{\left(\frac{q\lambda}{q+s}, \dots, \frac{q\lambda}{q+s}\right)}^{(s)} & (\mathfrak{p}^+ = M(q, s; \mathbb{C})), \\ V_{\left(\frac{\lambda}{2}, \dots, \frac{\lambda}{2}\right)}^{(s)\vee} & (\mathfrak{p}^+ = \text{Alt}(s, \mathbb{C})), \\ \mathbb{C}_{-\lambda} \boxtimes V_{(0, \dots, 0)}^{[10]\vee} & (\mathfrak{p}^+ = M(1, 2; \mathbb{O})^{\mathbb{O}}), \end{cases} \\ \mathcal{P}_{\mathbf{m}}(\mathfrak{p}^+) &\simeq \begin{cases} \mathbb{C}_{-m_1-m_2} \boxtimes V_{(m_1-m_2, 0, \dots, 0)}^{[n]\vee} & (\mathfrak{p}^+ = \mathbb{C}^n, n \geq 3), \\ V_{(2m_1, \dots, 2m_r)}^{(r)\vee} =: V_{2\mathbf{m}}^{(r)\vee} & (\mathfrak{p}^+ = \text{Sym}(r, \mathbb{C})), \\ V_{\mathbf{m}}^{(q)\vee} \boxtimes V_{(m_1, \dots, m_q, 0, \dots, 0)}^{(s)} =: V_{\mathbf{m}}^{(q)\vee} \boxtimes V_{\mathbf{m}}^{(s)} & (\mathfrak{p}^+ = M(q, s; \mathbb{C}), q \leq s), \\ V_{(m_1, \dots, m_s, 0, \dots, 0)}^{(q)\vee} \boxtimes V_{\mathbf{m}}^{(s)} =: V_{\mathbf{m}}^{(q)\vee} \boxtimes V_{\mathbf{m}}^{(s)} & (\mathfrak{p}^+ = M(q, s; \mathbb{C}), q \geq s), \\ V_{(m_1, m_1, m_2, m_2, \dots, m_{\lfloor s/2 \rfloor}, m_{\lfloor s/2 \rfloor}, (0))}^{(s)\vee} =: V_{\mathbf{m}^2}^{(s)\vee} & (\mathfrak{p}^+ = \text{Alt}(s, \mathbb{C})), \\ \mathbb{C}_{-\frac{3}{4}|\mathbf{m}|} \boxtimes V_{\left(\frac{m_1+m_2}{2}, \frac{m_1-m_2}{2}, \frac{m_1-m_2}{2}, \frac{m_1-m_2}{2}, \frac{m_1-m_2}{2}\right)}^{[10]\vee} & (\mathfrak{p}^+ = M(1, 2; \mathbb{O})^{\mathbb{C}}) \end{cases} \end{aligned} \tag{2.20}$$

when $K^{\mathbb{C}}$ is classical, if we normalize the representations $\mathbb{C}_{-\lambda}$ of $\widetilde{SO}(2) \simeq \widetilde{U}(1)$ for the first and the last cases suitably.

2.9 Restriction to symmetric subgroups

In this subsection we consider a \mathbb{C} -linear involution σ on a Hermitian positive Jordan triple system \mathfrak{p}^{\pm} , i.e., a Jordan triple system automorphism $\sigma: \mathfrak{p}^{\pm} \rightarrow \mathfrak{p}^{\mp}$ of order 2 which commutes with the \mathbb{C} -antilinear map $\bar{\cdot}: \mathfrak{p}^{\pm} \rightarrow \mathfrak{p}^{\mp}$, and extend to the involution of the Lie algebra $\mathfrak{g}^{\mathbb{C}} = \mathfrak{p}^+ \oplus \mathfrak{k}^{\mathbb{C}} \oplus \mathfrak{p}^-$ by letting σ act on $\mathfrak{k}^{\mathbb{C}} = \mathfrak{str}(\mathfrak{p}^+) \subset \text{End}_{\mathbb{C}}(\mathfrak{p}^+)$ by $\sigma(l) := \sigma l \sigma$.

Also let $\vartheta := -I_{\mathfrak{p}^+} + I_{\mathfrak{k}^\mathbb{C}} - I_{\mathfrak{p}^-}$. Using these, we set

$$\begin{aligned}\mathfrak{p}_1^\pm &:= (\mathfrak{p}^\pm)^\sigma = \{x \in \mathfrak{p}^\pm \mid \sigma(x) = x\}, \\ \mathfrak{p}_2^\pm &:= (\mathfrak{p}^\pm)^{-\sigma} = \{x \in \mathfrak{p}^\pm \mid \sigma(x) = -x\}, \\ \mathfrak{k}_1^\mathbb{C} &:= (\mathfrak{k}^\mathbb{C})^\sigma = \{l \in \mathfrak{k}^\mathbb{C} \mid \sigma(l) = l\}, \\ \mathfrak{k}_1 &:= \mathfrak{k}^\sigma = \mathfrak{k}_1^\mathbb{C} \cap \mathfrak{k}, \\ \mathfrak{g}_1^\mathbb{C} &:= (\mathfrak{g}^\mathbb{C})^\sigma = \mathfrak{p}_1^+ \oplus \mathfrak{k}_1^\mathbb{C} \oplus \mathfrak{p}_1^-, \\ \mathfrak{g}_2^\mathbb{C} &:= (\mathfrak{g}^\mathbb{C})^{\sigma\vartheta} = \mathfrak{p}_2^+ \oplus \mathfrak{k}_1^\mathbb{C} \oplus \mathfrak{p}_2^-, \\ \mathfrak{g}_1 &:= \mathfrak{g}^\sigma = \mathfrak{g}_1^\mathbb{C} \cap \mathfrak{g}, \\ \mathfrak{g}_2 &:= \mathfrak{g}^{\sigma\vartheta} = \mathfrak{g}_2^\mathbb{C} \cap \mathfrak{g},\end{aligned}$$

and let $G_1, G_1^\mathbb{C}, G_2, G_2^\mathbb{C}, K_1, K_1^\mathbb{C} \subset G^\mathbb{C}$ be the connected closed subgroups corresponding to $\mathfrak{g}_1, \mathfrak{g}_1^\mathbb{C}, \mathfrak{g}_2, \mathfrak{g}_2^\mathbb{C}, \mathfrak{k}_1, \mathfrak{k}_1^\mathbb{C}$ respectively. Such (G, G_1) is called a symmetric pair of holomorphic type (see [18, Section 3.4]). Also let $\tilde{G}_1 \subset \tilde{G}$ and $\tilde{K}_1^\mathbb{C} \subset \tilde{K}^\mathbb{C}$ be the connected closed subgroups of the universal covering groups of G and $K^\mathbb{C}$ corresponding to \mathfrak{g}_1 and $\mathfrak{k}_1^\mathbb{C}$ respectively, and let $D \subset \mathfrak{p}^+$, $D_1 \subset \mathfrak{p}_1^+$ be the corresponding bounded symmetric domains, so that $D \simeq G/K$, $D_1 \simeq G_1/K_1$ hold. Let χ be the character of $K^\mathbb{C}$ given in (2.15). Similarly, we normalize the bilinear form $(\cdot|\cdot)_{\mathfrak{p}_j^+}: \mathfrak{p}_j^+ \times \mathfrak{p}_j^- \rightarrow \mathbb{C}$ ($j = 1, 2$) such that $(e|\bar{e})_{\mathfrak{p}_j^+} = 1$ holds for any primitive tripotent $e \in \mathfrak{p}_j^+$, and define the characters χ_j ($j = 1, 2$) of $K_1^\mathbb{C}$ by

$$\begin{aligned}d\chi_j([x, y]) &= (x|y)_{\mathfrak{p}_j^+} & (x \in \mathfrak{p}_j^+, y \in \mathfrak{p}_j^-), \\ d\chi_j(l) &= 0 & (l \in [\mathfrak{p}_j^+, \mathfrak{p}_j^-]^\perp \subset \mathfrak{k}_1^\mathbb{C}).\end{aligned}$$

As in Section 2.7, for $\lambda \in \mathbb{R}$ and for an irreducible representation V of $K_1^\mathbb{C}$, let $\mathcal{H}_\lambda(D) \subset \mathcal{O}_\lambda(D)$ and $\mathcal{H}_\lambda(D_1, V) \subset \mathcal{O}_\lambda(D_1, V)$ be the unitary representations of \tilde{G} and \tilde{G}_1 with the minimal \tilde{K} -type $\chi^{-\lambda}$ and with the minimal \tilde{K}_1 -type $\chi_1^{-\lambda} \otimes V$ respectively, if they exist.

In the following we assume G is simple. Then \mathfrak{p}_2^+ is a direct sum of at most two simple Jordan triple systems. Let $\text{rank } \mathfrak{p}^+ =: r$, $\text{rank } \mathfrak{p}_2^+ =: r_2$, and define $\varepsilon_1, \varepsilon_2 \in \{1, 2\}$ by $d\chi|_{\mathfrak{k}_1^\mathbb{C}} = \varepsilon_j d\chi_j$, or equivalently, by

$$(x|y)_{\mathfrak{p}^+} = \varepsilon_j (x|y)_{\mathfrak{p}_j^+} \quad (j = 1, 2, x \in \mathfrak{p}_j^+, y \in \mathfrak{p}_j^-). \quad (2.21)$$

When \mathfrak{p}_2^+ is not simple, we write $\mathfrak{p}_2^+ =: \mathfrak{p}_{11}^+ \oplus \mathfrak{p}_{22}^+$, $\mathfrak{p}_1^+ =: \mathfrak{p}_{12}^+$, and let $\text{rank } \mathfrak{p}_{11}^+ =: r'$, $\text{rank } \mathfrak{p}_{22}^+ =: r''$. Now we consider the restriction of the representation $\mathcal{H}_\lambda(D)$ of \tilde{G} to the subgroup \tilde{G}_1 . If λ satisfies (2.19), then the unitary representation $\mathcal{H}_\lambda(D)$ exists, $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ is discretely decomposable, and every \tilde{G}_1 -submodule in $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ contains a \mathfrak{p}_1^+ -null vector, that is, has an intersection with $(\mathcal{H}_\lambda(D)_{\tilde{K}})^{\mathfrak{p}_1^+} = (\chi_1^{-\varepsilon_1\lambda} \otimes \mathcal{P}(\mathfrak{p}_2^+)) \cap \mathcal{H}_\lambda(D)_{\tilde{K}}$. Especially, $\mathcal{H}_\lambda(D)_{\tilde{K}} = \chi^{-\lambda} \otimes \mathcal{P}(\mathfrak{p}^+)$ holds if $\lambda > \frac{d}{2}(r-1)$, and $\mathcal{H}_\lambda(D)$ is holomorphic discrete if $\lambda > p-1$. For such λ , according to the decomposition

$$\begin{aligned}\mathcal{P}(\mathfrak{p}_2^+) &= \bigoplus_{\mathbf{k} \in \mathbb{Z}_{++}^{r_2}} \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+) & (\mathfrak{p}_2^+ : \text{simple}), \\ \mathcal{P}(\mathfrak{p}_2^+) &= \bigoplus_{\mathbf{k} \in \mathbb{Z}_{++}^{r'}} \bigoplus_{\mathbf{l} \in \mathbb{Z}_{++}^{r''}} \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_{\mathbf{l}}(\mathfrak{p}_{22}^+) & (\mathfrak{p}_2^+ : \text{non-simple})\end{aligned}$$

(Theorem 2.1), the following holds.

Theorem 2.9 (Kobayashi, [18, Theorems 8.3]). *For $\lambda > \frac{d}{2}(r-1)$, the restriction of $\mathcal{H}_\lambda(D)$ to the subgroup \tilde{G}_1 is decomposed into the Hilbert direct sum of irreducible representations as*

$$\begin{aligned}\mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\simeq \sum_{\mathbf{k} \in \mathbb{Z}_{++}^{r_2}}^{\oplus} \mathcal{H}_{\varepsilon_1\lambda}(D_1, \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)) && (\mathfrak{p}_2^+ : \text{simple}), \\ \mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\simeq \sum_{\mathbf{k} \in \mathbb{Z}_{++}^{r'}}^{\oplus} \sum_{\mathbf{l} \in \mathbb{Z}_{++}^{r''}}^{\oplus} \mathcal{H}_{\varepsilon_1\lambda}(D_1, \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_{\mathbf{l}}(\mathfrak{p}_{22}^+)) && (\mathfrak{p}_2^+ : \text{non-simple}).\end{aligned}$$

In the following, when \mathfrak{p}_2^+ is not simple, we write $\mathcal{P}_{(\mathbf{k}, \mathbf{l})}(\mathfrak{p}_2^+) := \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_{11}^+) \boxtimes \mathcal{P}_{\mathbf{l}}(\mathfrak{p}_{22}^+)$. Then by the above decomposition, for each $\lambda > \frac{d}{2}(r-1)$ and $\mathbf{k} \in \mathbb{Z}_{++}^{r_2}$ or $\mathbf{k} \in \mathbb{Z}_{++}^{r'} \times \mathbb{Z}_{++}^{r''}$, there exist \tilde{G}_1 -intertwining operators

$$\begin{aligned}\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow: \mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\longrightarrow \mathcal{H}_{\varepsilon_1\lambda}(D_1, \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)), \\ \mathcal{F}_{\lambda, \mathbf{k}}^\uparrow: \mathcal{H}_{\varepsilon_1\lambda}(D_1, \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)) &\longrightarrow \mathcal{H}_\lambda(D)|_{\tilde{G}_1},\end{aligned}$$

which are unique up to constant multiple. $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ is called a *symmetry breaking operator*, and $\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow$ is called a *holographic operator*, according to the terminology introduced in [24, 25] and [26] respectively. Then the construction of $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ is reduced to the computation of some inner product. In the following, let $V_{\mathbf{k}}$ be an abstract K_1 -module isomorphic to $\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, let $\|\cdot\|_{\varepsilon_1\lambda, \mathbf{k}, \mathfrak{p}_1^+}$ be the \tilde{G}_1 -invariant norm on $\mathcal{H}_{\varepsilon_1\lambda}(D_1, V_{\mathbf{k}})$ normalized such that $\|v\|_{\varepsilon_1\lambda, \mathbf{k}, \mathfrak{p}_1^+} = |v|_{V_{\mathbf{k}}}$ holds for all constant functions $v \in V_{\mathbf{k}}$, and let $\|\cdot\|_{F, \mathfrak{p}^+}$ be the Fischer norm on \mathfrak{p}^+ given in (2.17).

Theorem 2.10 ([33, Theorem 3.10(1)], [35, Proposition 2.9]). *For $\lambda > p-1$ and for $\mathbf{k} \in \mathbb{Z}_{++}^{r_2}$ (when \mathfrak{p}_2^+ is simple) or $\mathbf{k} \in \mathbb{Z}_{++}^{r'} \times \mathbb{Z}_{++}^{r''}$ (when \mathfrak{p}_2^+ is not simple), let $C_{\mathfrak{p}^+, \mathfrak{p}_2^+}(\lambda, \mathbf{k}) \in \mathbb{C}$ be the constant satisfying*

$$\left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \Big|_{z_1=0} = C_{\mathfrak{p}^+, \mathfrak{p}_2^+}(\lambda, \mathbf{k}) f(z_2) \quad (f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)).$$

We take a vector-valued polynomial $K_{\mathbf{k}}(x_2) \in \mathcal{P}(\mathfrak{p}_2^+, \overline{V_{\mathbf{k}}})^{K_1}$ normalized such that

$$|\langle f(x_2), K_{\mathbf{k}}(x_2) \rangle_{F, \mathfrak{p}^+}|_{V_{\mathbf{k}}}^2 = \|f\|_{F, \mathfrak{p}^+}^2 \quad (f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)),$$

and define the vector-valued polynomial $F_{\lambda, \mathbf{k}}^\downarrow(z) \in \mathcal{P}(\mathfrak{p}^-, V_{\mathbf{k}})$ by

$$F_{\lambda, \mathbf{k}}^\downarrow(z) := \frac{1}{C_{\mathfrak{p}^+, \mathfrak{p}_2^+}(\lambda, \mathbf{k})} \left\langle e^{(x|z)_{\mathfrak{p}^+}}, K_{\mathbf{k}}(x_2) \right\rangle_{\lambda, x}. \quad (2.22)$$

Then the differential operator

$$\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow: \mathcal{H}_\lambda(D)|_{\tilde{G}_1} \longrightarrow \mathcal{H}_{\varepsilon_1\lambda}(D_1, V_{\mathbf{k}}), \quad (\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow f)(x_1) := F_{\lambda, \mathbf{k}}^\downarrow \left(\frac{\partial}{\partial x} \right) f(x) \Big|_{x_2=0} \quad (2.23)$$

becomes a symmetry breaking operator satisfying

$$\|\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow f\|_{\varepsilon_1\lambda, \mathbf{k}, \mathfrak{p}_1^+}^2 = \|f\|_{F, \mathfrak{p}^+}^2 \quad (f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)).$$

Its operator norm is given by

$$\|\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow\|_{\text{op}}^2 = C_{\mathfrak{p}^+, \mathfrak{p}_2^+}(\lambda, \mathbf{k})^{-1}.$$

The notion of symmetry breaking operator is extended to the map

$$\mathcal{F}_{\lambda, \mathbf{k}}^{\downarrow}: \mathcal{O}_{\lambda}(D)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\varepsilon_1 \lambda}(D_1, \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+))$$

for all $\lambda \in \mathbb{C}$. Especially, the operators constructed in the above theorem are meromorphically continued for all $\lambda \in \mathbb{C}$, and give symmetry breaking operators on $\mathcal{O}_{\lambda}(D)$ for all $\lambda \in \mathbb{C}$ except for poles. Moreover, symmetry breaking operators for holomorphic settings are always given by differential operators (localness theorem, [24, Theorem 5.3]). However, the uniqueness of $\mathcal{F}_{\lambda, \mathbf{k}}^{\downarrow}$ for non-unitary range is not trivial. Indeed, the uniqueness of symmetry breaking operators for tensor product case fails for non-unitary range. See [25, Section 9], [35, Section 8]. The symbols of differential symmetry breaking operators are characterized as polynomial solutions of some differential equations as follows. For $\lambda \in \mathbb{C}$, let $\mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}}$ be the vector-valued differential operator

$$\begin{aligned} \mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}}: \mathcal{P}(\mathfrak{p}^{\pm}) &\longrightarrow \mathcal{P}(\mathfrak{p}^{\pm}) \otimes \mathfrak{p}^{\mp}, \\ \mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}} f(z) &:= \sum_{\alpha, \beta} \frac{1}{2} Q(e_{\alpha}^{\mp}, e_{\beta}^{\mp}) z \frac{\partial^2 f}{\partial z_{\alpha}^{\pm} \partial z_{\beta}^{\pm}}(z) + \lambda \sum_{\alpha} e_{\alpha}^{\mp} \frac{\partial f}{\partial z_{\alpha}^{\pm}}(z), \end{aligned} \quad (2.24)$$

where $\{e_{\alpha}^{\mp}\} \subset \mathfrak{p}^{\mp}$ is a basis of \mathfrak{p}^{\mp} , and $\{z_{\alpha}^{\pm}\}$ is the coordinate of \mathfrak{p}^{\pm} dual to $\{e_{\alpha}^{\mp}\}$ (*Bessel operator* [5], [9, Section XV.2]), let

$$(\mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}})_1: \mathcal{P}(\mathfrak{p}^{\pm}) \longrightarrow \mathcal{P}(\mathfrak{p}^{\pm}) \otimes \mathfrak{p}_1^{\mp}$$

be the orthogonal projection of $\mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}}$ onto \mathfrak{p}_1^{\mp} , and let

$$\text{Sol}_{\mathcal{P}(\mathfrak{p}^{\pm})}((\mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}})_1) := \{f(z) \in \mathcal{P}(\mathfrak{p}^{\pm}) \mid (\mathcal{B}_{\lambda}^{\mathfrak{p}^{\pm}})_1 f = 0\}.$$

Theorem 2.11 (F-method, [25, Theorem 3.1]). *Let V be a $K_1^{\mathbb{C}}$ -module, and V^{\vee} be its contragredient. Then we have*

$$\begin{aligned} \text{Hom}_{\tilde{G}_1}(\mathcal{O}_{\lambda}(D), \mathcal{O}_{\varepsilon_1 \lambda}(D_1, V)) &\simeq \text{Hom}_{\mathfrak{k}_1^{\mathbb{C}} \oplus \mathfrak{p}_1^-} \left(\chi_1^{\varepsilon_1 \lambda} \otimes V^{\vee}, \text{ind}_{\mathfrak{k}^{\mathbb{C}} \oplus \mathfrak{p}^-}^{\mathfrak{g}^{\mathbb{C}}}(\chi^{\lambda}) \right) \\ &\simeq \left(\text{Sol}_{\mathcal{P}(\mathfrak{p}^{\pm})}((\mathcal{B}_{\lambda}^{\mathfrak{p}^-})_1) \otimes V \right)^{K_1}. \end{aligned}$$

Especially, if $\lambda > \frac{d}{2}(r-1)$ then since $\mathcal{H}_{\lambda}(D)|_{\tilde{G}_1}$ decomposes multiplicity-freely, the space of symmetry breaking operators is 1-dimensional for each $V \simeq \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, and combining the above two theorems, for $\lambda > \frac{d}{2}(r-1)$ we get

$$\text{Hom}_{K_1} \left(\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+), \text{Sol}_{\mathcal{P}(\mathfrak{p}^+)}((\mathcal{B}_{\lambda}^{\mathfrak{p}^+})_1) \right) = \mathbb{C} \left(f(x_2) \mapsto \left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \right).$$

In this article, we consider the symmetry breaking operators for the case $\mathfrak{p}^+, \mathfrak{p}_2^+$ are simple of tube type and $\varepsilon_2 = 1$, so that $r = r_2$ holds. In [34, Section 6], we treated $\mathbf{k} = (k+l, k, \dots, k)$ case. In this article, we treat general $\mathbf{k} \in \mathbb{Z}_{++}^3$ for $r = 3$ case in Section 3, and treat $\mathbf{k} = (k, \dots, k, k-l)$ case for general r in Section 4.

3 Rank 3 case

In this section we treat $\mathfrak{p}^\pm = (\mathfrak{p}^\pm)^\sigma \oplus (\mathfrak{p}^\pm)^{-\sigma} = \mathfrak{p}_1^\pm \oplus \mathfrak{p}_2^\pm$ such that $\mathfrak{p}^\pm, \mathfrak{p}_2^\pm$ are both simple, of tube type, and of rank 3, that is, we treat

$$(\mathfrak{p}^\pm, \mathfrak{p}_1^\pm, \mathfrak{p}_2^\pm) = (\text{Herm}(3, \mathbb{F})^\mathbb{C}, \text{Alt}(3, \mathbb{F}')^\mathbb{C}, \text{Herm}(3, \mathbb{F}')^\mathbb{C}) \quad ((\mathbb{F}, \mathbb{F}') = (\mathbb{C}, \mathbb{R}), (\mathbb{H}, \mathbb{C}), (\mathbb{O}, \mathbb{H})) \\ = \begin{cases} (M(3, \mathbb{C}), \text{Alt}(3, \mathbb{C}), \text{Sym}(3, \mathbb{C})) & ((\mathbb{F}, \mathbb{F}') = (\mathbb{C}, \mathbb{R})), \\ (\text{Alt}(6, \mathbb{C}), \text{Alt}(3, \mathbb{C}) \oplus \text{Alt}(3, \mathbb{C}), M(3, \mathbb{C})) & ((\mathbb{F}, \mathbb{F}') = (\mathbb{H}, \mathbb{C})), \\ (\text{Herm}(3, \mathbb{O})^\mathbb{C}, M(2, 6; \mathbb{C}), \text{Alt}(6, \mathbb{C})) & ((\mathbb{F}, \mathbb{F}') = (\mathbb{O}, \mathbb{H})). \end{cases}$$

Then the corresponding symmetric pairs $(G, (G^\sigma)_0, (G^{\sigma\vartheta})_0) = (G, G_1, G_2)$ are given by

$$(G, G_1, G_2) = \begin{cases} (SU(3, 3), SO^*(6), Sp(3, \mathbb{R})) & ((\mathbb{F}, \mathbb{F}') = (\mathbb{C}, \mathbb{R})), \\ (SO^*(12), SO^*(6) \times SO^*(6), U(3, 3)) & ((\mathbb{F}, \mathbb{F}') = (\mathbb{H}, \mathbb{C})), \\ (E_{7(-25)}, SU(2, 6), SU(2) \times SO^*(12)) & ((\mathbb{F}, \mathbb{F}') = (\mathbb{O}, \mathbb{H})) \end{cases}$$

(up to covering). We have $(d, d_2) = (\dim_{\mathbb{R}} \mathbb{F}, \dim_{\mathbb{R}} \mathbb{F}') \in \{(2, 1), (4, 2), (8, 4)\}$, and $\varepsilon_1 = 2$ for $(\mathbb{F}, \mathbb{F}') = (\mathbb{C}, \mathbb{R})$, $\varepsilon_1 = 1$ for $(\mathbb{F}, \mathbb{F}') = (\mathbb{H}, \mathbb{C}), (\mathbb{O}, \mathbb{H})$. \mathfrak{p}^+ and \mathfrak{p}_2^+ have Jordan algebra structures with the Euclidean real forms $\mathfrak{n}^+ = \text{Herm}(3, \mathbb{F})$ and $\mathfrak{n}_2^+ = \text{Herm}(3, \mathbb{F}')$ respectively.

3.1 Jordan algebra structure on $\text{Herm}(3, \mathbb{F})^\mathbb{C}$

First we look at the Jordan algebra structure on $\mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}} = \text{Herm}(3, \mathbb{F})^\mathbb{C}$ for $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$ given by

$$x \circ y := \frac{1}{2}(xy + yx),$$

with the unit element $e = I_3$, the Euclidean real form $\mathfrak{n}^+ = \text{Herm}(3, \mathbb{F})$, the symmetric cone $\Omega = \text{Herm}_+(3, \mathbb{F}) \subset \mathfrak{n}^+$ (the set of positive definite matrices), and the associative bilinear form $(\cdot|\cdot) = (\cdot|\cdot)_{\mathfrak{n}^+} : \mathfrak{n}^{+\mathbb{C}} \times \mathfrak{n}^{+\mathbb{C}} \rightarrow \mathbb{C}$,

$$(x|y) = (x|y)_{\mathfrak{n}^+} := \text{Re}_{\mathbb{F}} \text{tr}(xy) = \text{Re}_{\mathbb{F}} \text{tr}(yx),$$

where $\text{Re}_{\mathbb{F}} : \mathbb{F}^\mathbb{C} = \mathbb{F} \otimes_{\mathbb{R}} \mathbb{C} \rightarrow \mathbb{C}$ is the \mathbb{C} -linear extension of the \mathbb{F} -real part. Next, for $x \in \mathfrak{n}^{+\mathbb{C}}$, let x^\sharp be the adjugate matrix given by

$$\begin{pmatrix} a_1 & x_3 & \hat{x}_2 \\ \hat{x}_3 & a_2 & x_1 \\ x_2 & \hat{x}_1 & a_3 \end{pmatrix} := \begin{pmatrix} a_2a_3 - x_1\hat{x}_1 & \hat{x}_2\hat{x}_1 - a_3x_3 & x_3x_1 - a_2\hat{x}_2 \\ x_1x_2 - a_3\hat{x}_3 & a_3a_1 - x_2\hat{x}_2 & \hat{x}_3\hat{x}_2 - a_1x_1 \\ \hat{x}_1\hat{x}_3 - a_2x_2 & x_2x_3 - a_1\hat{x}_1 & a_1a_2 - x_3\hat{x}_3 \end{pmatrix} \quad (a_i \in \mathbb{C}, x_i \in \mathbb{F}^\mathbb{C}),$$

where $x_i \mapsto \hat{x}_i$ is the \mathbb{C} -linear extension of the \mathbb{F} -conjugate, and for $x, y \in \text{Herm}(3, \mathbb{F})^\mathbb{C}$, let $x \times y$ be the *Freudenthal product* given by

$$x \times y := (x + y)^\sharp - x^\sharp - y^\sharp,$$

so that $x \times x = 2x^\sharp$. Then the determinant polynomial, the inverse element, the generic norm, and the map $P(x) \in \text{End}_{\mathbb{C}}(\mathfrak{n}^{+\mathbb{C}})$ are given by

$$\det(x) = \det_{\mathfrak{n}^+}(x) := \frac{1}{3}(x|x)^\sharp, \quad x^{-1} := \frac{1}{\det(x)}x^\sharp,$$

$$h(x, y) := 1 - (x|y) + (x^\sharp|y^\sharp) - \det(x)\det(y), \quad P(x)y := (x|y)x - x^\sharp \times y.$$

$xx^\sharp = x^\sharp x = \det(x)I$ holds for $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$, and $\frac{1}{2}(xx^\sharp + x^\sharp x) = \det(x)I$ holds for $\mathbb{F} = \mathbb{O}$. Also, for $x, y, z \in \mathfrak{n}^{+\mathbb{C}}$ we have

$$(x \times y|z) = (y \times z|x) = (z \times x|y), \quad (x^\sharp)^\sharp = \det(x)x.$$

If $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$, then by taking the directional derivative of $zz^\sharp = \det(z)I$ along y , we get

$$yz^\sharp + z(z \times y) = (z^\sharp|y)I, \quad (3.1)$$

and by substituting $z = x^\sharp$ and multiplying $\det(x)^{-1}x$ from the left, we get

$$P(x)y = (x|y)x - x^\sharp \times y = xyx.$$

We note that these do not hold for $\mathbb{F} = \mathbb{O}$. Let $\bar{\cdot}: \mathfrak{n}^{+\mathbb{C}} \rightarrow \mathfrak{n}^{+\mathbb{C}}$ be the complex conjugate with respect to the real form $\mathfrak{n}^+ \subset \mathfrak{n}^{+\mathbb{C}}$. Then the bounded symmetric domain $D \subset \mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}}$ is the connected component of $\{x \in \mathfrak{n}^{+\mathbb{C}} \mid h(x, \bar{x}) > 0\}$ which contains the origin.

Next we consider the Cayley–Dickson extension $\mathbb{F} = \mathbb{F}' \oplus \mathbb{F}'j$ for $(\mathbb{F}, \mathbb{F}') = (\mathbb{C}, \mathbb{R}), (\mathbb{H}, \mathbb{C}), (\mathbb{O}, \mathbb{H})$, and as in [41, 42] we consider the involution σ on $\mathfrak{p}^\pm = \text{Herm}(3, \mathbb{F})$,

$$\sigma \begin{pmatrix} a_1 & x_3 + y_3j & \hat{x}_2 - y_2j \\ \hat{x}_3 - y_3j & a_2 & x_1 + y_1j \\ x_2 + y_2j & \hat{x}_1 - y_1j & a_3 \end{pmatrix} := - \begin{pmatrix} a_1 & x_3 - y_3j & \hat{x}_2 + y_2j \\ \hat{x}_3 + y_3j & a_2 & x_1 - y_1j \\ x_2 - y_2j & \hat{x}_1 + y_1j & a_3 \end{pmatrix}$$

$(a_i \in \mathbb{C}, x_i, y_i \in \mathbb{F}'^\mathbb{C})$, so that we have

$$\begin{aligned} (\mathfrak{p}_1^\pm, \mathfrak{p}_2^\pm) &:= ((\mathfrak{p}^\pm)^\sigma, (\mathfrak{p}^\pm)^{-\sigma}) = (\text{Alt}(3, \mathbb{F}')^\mathbb{C}j, \text{Herm}(3, \mathbb{F}')^\mathbb{C}), \\ (\mathfrak{n}_1^+, \mathfrak{n}_2^+) &:= ((\mathfrak{n}^+)^\sigma, (\mathfrak{n}^+)^{-\sigma}) = (\text{Alt}(3, \mathbb{F}')j, \text{Herm}(3, \mathbb{F}')). \end{aligned}$$

$\mathfrak{p}_2^+ = \mathfrak{n}_2^{+\mathbb{C}}$ becomes a Jordan subalgebra with the Euclidean real form \mathfrak{n}_2^+ and the symmetric cone $\Omega_2 := \Omega \cap \mathfrak{n}_2^+ \subset \mathfrak{n}_2^+$, but $\mathfrak{p}_1^+ = \mathfrak{n}_1^{+\mathbb{C}}$ is not. We identify $\mathfrak{p}_1^+ = \mathfrak{n}_1^{+\mathbb{C}} = \text{Alt}(3, \mathbb{F}')^\mathbb{C}j$ with $(\mathbb{F}'^\mathbb{C})^3$ (the space of row vectors) by

$$(\mathbb{F}'^\mathbb{C})^3 \xrightarrow{\sim} \text{Alt}(3, \mathbb{F}')^\mathbb{C}j, \quad (y_1, y_2, y_3) \mapsto \begin{pmatrix} 0 & y_3j & -y_2j \\ -y_3j & 0 & y_1j \\ y_2j & -y_1j & 0 \end{pmatrix}.$$

In the following, we use lowercase Greek alphabets and capital Latin alphabets to express elements in $\mathfrak{n}_1^{+\mathbb{C}} \simeq (\mathbb{F}'^\mathbb{C})^3$ and $\mathfrak{n}_2^{+\mathbb{C}} = \text{Herm}(3, \mathbb{F}')^\mathbb{C}$ respectively. For $\xi, \eta \in (\mathbb{F}'^\mathbb{C})^3$, let

$$(\xi|\eta) := \text{Re}_{\mathbb{F}'}(\xi^t \hat{\eta}).$$

Then for $x = (x_1, x_2) = (\xi, X)$, $y = (y_1, y_2) = (\eta, Y) \in \mathfrak{n}^{+\mathbb{C}} = (\mathbb{F}'^\mathbb{C})^3 \oplus \text{Herm}(3, \mathbb{F}')^\mathbb{C}$ we have

$$\begin{aligned} (x|y)_{\mathfrak{n}^+} &= (x_1|y_1)_{\mathfrak{n}^+} + (x_2|y_2)_{\mathfrak{n}^+} = 2(\xi|\eta) + (X|Y), \\ (\xi, X)^\sharp &= (-\xi X, X^\sharp - {}^t \hat{\xi} \xi), \\ \det((\xi, X)) &= \det(X) - \xi X^t \hat{\xi}, \\ P((\xi, X))(\eta, Y) &= (\xi^t \hat{\eta} \xi + \eta X^\sharp + (X|Y)\xi - \xi XY, \\ &\quad XYX + ({}^t \hat{\xi} \xi) \times Y + 2(\xi|\eta)X - X^t \hat{\xi} \eta - {}^t \hat{\eta} \xi X). \end{aligned}$$

$\mathfrak{p}_1^\pm, \mathfrak{p}_2^\pm$ are preserved by the action of $K_1 := G_1 \cap G_2$. Especially, the action of the real form $L_2 = K_1^\mathbb{C} \cap L \subset K_1^\mathbb{C}$,

$$L_2 := \begin{cases} \{1\} \times GL(3, \mathbb{R}) & \{g \in GL(3, \mathbb{C}) \mid \det(g) \in \mathbb{R}\} \quad ((\mathbb{F}, \mathbb{F}') = (\mathbb{C}, \mathbb{R})), \\ \{1\} \times GL(3, \mathbb{C}) & GL(3, \mathbb{H}) \quad ((\mathbb{F}, \mathbb{F}') = (\mathbb{H}, \mathbb{C})), \\ Sp(1) \times GL(3, \mathbb{H}) & \mathbb{R}_{>0} \times E_{6(-26)} \quad ((\mathbb{F}, \mathbb{F}') = (\mathbb{O}, \mathbb{H})), \end{cases}$$

on $\mathfrak{n}_1^+ \oplus \mathfrak{n}_2^+ = \mathbb{F}'^3 \oplus \text{Herm}(3, \mathbb{F}')$ is given by, for $g = (g_1, g_2) \in L_2$,

$$gx = (gx_1, gx_2) = (\chi(g_2)g_1\xi g_2^{-1}, g_2 X^t \hat{g}_2),$$

and this preserves the symmetric cone $\Omega_2 \subset \mathfrak{n}_2^+$.

3.2 Definition of some functions

In this subsection we define some functions on $\mathfrak{p}_1^\pm \oplus \mathfrak{p}_2^\pm$. First we note that, for $m \in \mathbb{Z}_{\geq 0}$, $(y_1, x_2) = (\eta, X) \in \mathfrak{p}_1^\mp \oplus \mathfrak{p}_2^\pm$, $(x_1, x_2) = (\xi, X) \in \mathfrak{p}_1^\pm \oplus \mathfrak{p}_2^\pm$, we have

$$\begin{aligned} ((y_1)^\sharp | (x_2)^\sharp)^m &= (\eta X^{\sharp t} \hat{\eta})^m \in \mathcal{P}_{2m}(\mathfrak{p}_1^\mp)_{y_1} \otimes \mathcal{P}_{(m, m, 0)}(\mathfrak{p}_2^\pm)_{x_2}, \\ \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^\pm}(x_2)} \right)^m &= \left(\frac{\xi X^t \hat{\xi}}{\det(X)} \right)^m \in \mathcal{P}_{2m}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{(0, -m, -m)}(\mathfrak{p}_2^\pm)_{x_2}, \end{aligned}$$

where $\mathcal{P}_m(\mathfrak{p}_1^\mp)$ is the space of homogeneous polynomials on \mathfrak{p}_1^\mp of degree m . Next let $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3 = \mathbb{Z}_{++}^3 + \mathbb{C}(1, 1, 1)$, and take $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^\pm) = \mathcal{P}_{(k_1-k_3, k_2-k_3, 0)}(\mathfrak{p}_2^\pm) \det_{\mathfrak{n}_2^\pm}(x_2)^{k_3}$. Then we have

$$\begin{aligned} ((y_1)^\sharp | (x_2)^\sharp)^m f(x_2) &\in \bigoplus_{\substack{\mathbf{l} \in \mathbb{Z}^3, |\mathbf{l}|=m \\ 0 \leq l_1 \leq k_1 - k_2 \\ 0 \leq l_2 \leq k_2 - k_3 \\ 0 \leq l_3}} \mathcal{P}_{2m}(\mathfrak{p}_1^\mp)_{y_1} \otimes \mathcal{P}_{(k_1+l_2+l_3, k_2+l_1+l_3, k_3+l_1+l_2)}(\mathfrak{p}_2^\pm)_{x_2}, \\ \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^\pm}(x_2)} \right)^m f(x_2) &\in \bigoplus_{\substack{\mathbf{l} \in \mathbb{Z}^3, |\mathbf{l}|=m \\ 0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2 \\ 0 \leq l_3 \leq k_2 - k_3}} \mathcal{P}_{2m}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{(k_1-l_2-l_3, k_2-l_1-l_3, k_3-l_1-l_2)}(\mathfrak{p}_2^\pm)_{x_2}. \end{aligned}$$

According to these decompositions, for $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^3$ we define the polynomials $F_{\mathbf{k}, \mathbf{l}}^\uparrow[f](y_1, x_2) = F_{\mathbf{k}, \mathbf{l}}^\uparrow[f](\eta, X)$ and $F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) = F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](\xi, X)$ by

$$\begin{aligned} F_{\mathbf{k}, \mathbf{l}}^\uparrow[f](y_1, x_2) &:= \text{Proj}_{(k_1+l_2+l_3, k_2+l_1+l_3, k_3+l_1+l_2), x_2}^{\mathfrak{p}_2^\pm} \left(\frac{1}{|\mathbf{l}|!} ((y_1)^\sharp | (x_2)^\sharp)^{|\mathbf{l}|} f(x_2) \right) \\ &\in \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^\mp)_{y_1} \otimes \mathcal{P}_{(k_1+l_2+l_3, k_2+l_1+l_3, k_3+l_1+l_2)}(\mathfrak{p}_2^\pm)_{x_2}, \\ F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) &:= \text{Proj}_{(k_1-l_2-l_3, k_2-l_1-l_3, k_3-l_1-l_2), x_2}^{\mathfrak{p}_2^\pm} \left(\frac{1}{|\mathbf{l}|!} \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^\pm}(x_2)} \right)^{|\mathbf{l}|} f(x_2) \right) \\ &\in \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{(k_1-l_2-l_3, k_2-l_1-l_3, k_3-l_1-l_2)}(\mathfrak{p}_2^\pm)_{x_2}. \end{aligned}$$

Under the identification $\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^\pm) \simeq \mathcal{P}_{-\mathbf{k}^\vee}(\mathfrak{p}_2^\mp)$, $f \mapsto f((\cdot)^{-1})$, where $\mathbf{k}^\vee := (k_3, k_2, k_1)$, these are related as

$$F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) = F_{-\mathbf{k}^\vee, \mathbf{l}^\vee}^\uparrow[f((\cdot)^{-1})](x_1, x_2^{-1}).$$

We also set

$$\begin{aligned}\tilde{F}_{\mathbf{k},\mathbf{l}}[f](x_1, x_2) &:= F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2) \det_{\mathfrak{n}_2^\pm}(x_2)^{|\mathbf{l}|} = \text{Proj}_{\mathbf{k}+\mathbf{l}, x_2}^{\mathfrak{p}_2^\pm} \left(\frac{1}{|\mathbf{l}|!} ((x_1)^\sharp |x_2|)^{|\mathbf{l}|} f(x_2) \right) \\ &\in \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{\mathbf{k}+\mathbf{l}}(\mathfrak{p}_2^\pm)_{x_2}.\end{aligned}$$

Since $F_{\mathbf{k},\mathbf{l}}^\uparrow[f](y_1, x_2)$ is recovered from $F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2)$, in the following we only consider $F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2)$. For $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$, let $d_{\mathbf{k}}^{\mathfrak{p}_2^+} := \dim \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, and let $\Sigma_2 \subset \mathfrak{p}_2^+$ be the Bergman–Shilov boundary of $D_2 := D \cap \mathfrak{p}_2^+$.

Proposition 3.1. *Let $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$, $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^3$ with $0 \leq l_2 \leq k_1 - k_2$, $0 \leq l_3 \leq k_2 - k_3$.*

(1) *For $x_1 \in \mathfrak{p}_1^+$, $x_2 = ge \in \mathfrak{p}_2^+$ with $g \in \tilde{K}_1^\mathbb{C}$, we have*

$$|F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2)| \leq \frac{\sqrt{d_{\mathbf{k}}^{\mathfrak{p}_2^+} d_{(|\mathbf{l}|, 0, 0)}^{\mathfrak{p}_2^+}}}{|\mathbf{l}|!} \frac{(g^{-1} x_1 |g^{-1} x_1|_{\mathfrak{p}_2^+})^{|\mathbf{l}|}}{2^{|\mathbf{l}|}} \|f(g(\cdot))\|_{L^2(\Sigma_2)}.$$

(2) *If $f(x_2) \neq 0$, then $F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2) \neq 0$.*

Proof. (1) Since $f(x_2) \mapsto F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2)$ is $\tilde{K}_1^\mathbb{C}$ -equivariant, we have

$$F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, x_2) = F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, ge) = F_{\mathbf{k},\mathbf{l}}^\downarrow[f(g(\cdot))](g^{-1} x_1, e),$$

and hence it suffices to prove when $x_2 = e$, so that $F_{\mathbf{k},\mathbf{l}}^\downarrow[f](x_1, e) = \tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, e)$. Next, since the reproducing kernel of $\mathcal{P}_{\mathbf{k}+\mathbf{l}}(\mathfrak{p}_2^+)$ with respect to the $L^2(\Sigma_2)$ -inner product is given by $K'_{\mathbf{k}+\mathbf{l}}(X, Z) := d_{\mathbf{k}+\mathbf{l}}^{\mathfrak{p}_2^+} \Phi_{\mathbf{k}+\mathbf{l}}^{\mathfrak{n}_2^+}(P(\overline{Z}^{1/2})X) = d_{\mathbf{k}+\mathbf{l}}^{\mathfrak{p}_2^+} \Phi_{\mathbf{k}+\mathbf{l}}^{\mathfrak{n}_2^+}(P(X^{1/2})\overline{Z})$, where $\Phi_{\mathbf{k}}^{\mathfrak{n}_2^+}(X) := \int_{K_1 \cap K_L} \Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}(kX) dk$ (see [9, Proposition XII.2.4]), we have

$$\begin{aligned}|\tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, e)| &= |\langle \tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, \cdot), K'_{\mathbf{k}+\mathbf{l}}(\cdot, e) \rangle_{L^2(\Sigma_2)}| \\ &\leq \|\tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, \cdot)\|_{L^2(\Sigma_2)} \|K'_{\mathbf{k}+\mathbf{l}}(\cdot, e)\|_{L^2(\Sigma_2)} \\ &= \|\tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, \cdot)\|_{L^2(\Sigma_2)} \sqrt{K'_{\mathbf{k}+\mathbf{l}}(e, e)} = \sqrt{d_{\mathbf{k}+\mathbf{l}}^{\mathfrak{p}_2^+}} \|\tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, \cdot)\|_{L^2(\Sigma_2)}.\end{aligned}$$

Now since $\tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, X)$ is defined by the orthogonal projection of $(\xi X^t \hat{\xi})^{|\mathbf{l}|} f(X)/|\mathbf{l}|!$, we have

$$\|\tilde{F}_{\mathbf{k},\mathbf{l}}[f](\xi, \cdot)\|_{L^2(\Sigma_2)} \leq \left\| \frac{1}{|\mathbf{l}|!} (\xi X^t \hat{\xi})^{|\mathbf{l}|} f(X) \right\|_{L^2(\Sigma_2), X} \leq \frac{1}{|\mathbf{l}|!} \|\xi X^t \hat{\xi}\|_{L^\infty(\Sigma_2), X}^{|\mathbf{l}|} \|f\|_{L^2(\Sigma_2)},$$

and for $X \in \Sigma_2$ we have

$$|\xi X^t \hat{\xi}| = |(\xi | \bar{\xi})(\xi X | \bar{\xi} X)| \leq \sqrt{(\xi | \bar{\xi})(\xi X | \bar{\xi} X)} = (\xi | \bar{\xi}) = \frac{1}{2} (x_1 | \bar{x}_1)_{\mathfrak{p}^+}.$$

We also have

$$d_{\mathbf{k}+\mathbf{l}}^{\mathfrak{p}_2^+} = \dim \mathcal{P}_{\mathbf{k}+\mathbf{l}}(\mathfrak{p}_2^+) \leq \dim (\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+) \otimes \mathcal{P}_{(|\mathbf{l}|, 0, 0)}(\mathfrak{p}_2^+)) = d_{\mathbf{k}}^{\mathfrak{p}_2^+} d_{(|\mathbf{l}|, 0, 0)}^{\mathfrak{p}_2^+}.$$

Hence the proposition follows.

(2) Again since $f(x_2) \mapsto F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2)$ is $\tilde{K}_1^{\mathbb{C}}$ -equivariant, it is enough to prove when $f(x_2) = \Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}(x_2) = \Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}(X)$. Then since ${}^t\hat{\xi}\xi \in \mathfrak{p}_2^+$ is at most of rank 1 for $x_1 = \xi \in \mathfrak{p}_1^+$ and every rank 1 element is written as this form, by Lemma 2.4(5) we have

$$F_{\mathbf{k}, \mathbf{l}}^\downarrow[\Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}](x_1, x_2) = \frac{1}{|\mathbf{l}|!} \det(X)^{-|\mathbf{l}|} \text{Proj}_{\mathbf{k}+\mathbf{l}, X}^{\mathfrak{p}_2^+}((X| {}^t\hat{\xi}\xi)^{|\mathbf{l}|} \Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}(X)) \neq 0$$

for some $x_1 = \xi \in \mathfrak{p}_1^+$. Hence we get the desired formula. \square

The following proposition gives some examples of $F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2)$.

Proposition 3.2. *Let $x = (x_1, x_2) = (\xi, X) \in \mathfrak{p}_1^+ \oplus \mathfrak{p}_2^+$.*

(1) *For $k \in \mathbb{C}$, $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^3$, $\mathbf{k} \in \mathbb{Z}_{++}^3$, $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, we have*

$$F_{\mathbf{k}+(k, k, k), \mathbf{l}}^\downarrow[f \det_{\mathfrak{n}_2^+}^k](x_1, x_2) = \det_{\mathfrak{n}_2^+}(x_2)^k F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2).$$

(2) *For $k \in \mathbb{C}$, $l \in \mathbb{Z}_{\geq 0}$, we have*

$$\begin{aligned} F_{(k, k, k), (l, 0, 0)}^\downarrow[\det_{\mathfrak{n}_2^+}^k](x_1, x_2) &= \frac{1}{l!} \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k = \frac{1}{l!} \left(\frac{\xi X^t \hat{\xi}}{\det(X)} \right)^l \det(X)^k, \\ F_{(k, k, k), (0, 0, l)}^\uparrow[\det_{\mathfrak{n}_2^+}^k](x_1, x_2) &= \frac{1}{l!} \left(-((x_1)^\sharp | (x_2)^\sharp) \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k = \frac{1}{l!} (\xi X^\sharp t \hat{\xi})^l \det(X)^k. \end{aligned}$$

(3) *Let $e' \in \mathfrak{n}_2^+$ be a primitive idempotent, and let $\mathfrak{p}^+(e')_0 =: \mathfrak{p}^{+''}$, $\mathfrak{p}_2^+(e')_0 =: \mathfrak{p}_2^{+''}$ be as in (2.2). Then for $f(x_2'') \in \mathcal{P}_{(k_1, k_2)}(\mathfrak{p}_2^{+''})$, $l = 0, 1, \dots, k_2$, we have*

$$\begin{aligned} F_{(k_1, k_2, 0), (0, 0, l)}^\downarrow[f](x_1, x_2) &= \frac{(-k_2)_l (-k_1 - \frac{d_2}{2})_l}{(-k_2 - \frac{d_2}{2})_l (-k_1 - d_2)_l l!} \left(-\frac{\det_{\mathfrak{n}_2^{+''}}(x_2'')}{\det_{\mathfrak{n}_2^{+''}}(x_1'')} \right)^l f(x_2'') \\ &= \frac{(-k_2)_l (-k_1 - \frac{d_2}{2})_l}{(-k_2 - \frac{d_2}{2})_l (-k_1 - d_2)_l l!} \left(\frac{({}^t\hat{\xi}\xi | e')}{\det_{\mathfrak{n}_2^{+''}}(X'')} \right)^l f(X''). \end{aligned}$$

(4) *Let $w_2 = W \in \mathfrak{p}_2^+$ be of rank 1. Then for $k, l_1, l_2 \in \mathbb{Z}_{\geq 0}$ with $l_2 \leq k$ we have*

$$\begin{aligned} F_{(k, 0, 0), (l_1, l_2, 0)}^\downarrow[(|w_2|_{\mathfrak{n}_2^+}^k)](x_1, x_2) &= F_{(0, 0, -k), (0, l_2, l_1)}^\uparrow[((\cdot)^{-1} | w_2|_{\mathfrak{n}_2^+}^k)](x_1, x_2^{-1}) \\ &= \frac{(-1)^{l_1} \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_2}}{(-k - l_1 - \frac{d_2}{2})_{l_2} l_2! (l_1 + l_2)!} \sum_{j=l_2}^{\min\{k, l_1 + l_2\}} \frac{(-k)_j (-l_1 - l_2)_j}{(-k - l_1 + l_2 - \frac{d_2}{2} + 1)_{j-l_2} (j - l_2)!} \\ &\quad \times (x_2 | (x_1)^\sharp)^{l_1 + l_2 - j} (x_2 | w_2)^{k-j} ((x_2 | (x_1)^\sharp)(x_2 | w_2) - (x_2 | (x_1)^\sharp x_2 | w_2))^j \\ &= \frac{(-1)^{l_2} \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_2}}{(-k - l_1 - \frac{d_2}{2})_{l_2} l_2! (l_1 + l_2)!} \sum_{j=l_2}^{\min\{k, l_1 + l_2\}} \frac{(-k)_j (-l_1 - l_2)_j}{(-k - l_1 + l_2 - \frac{d_2}{2} + 1)_{j-l_2} (j - l_2)!} \\ &\quad \times (X | {}^t\hat{\xi}\xi)^{l_1 + l_2 - j} (X | W)^{k-j} ((X | {}^t\hat{\xi}\xi)(X | W) - (X | {}^t\hat{\xi}\xi X | W))^j \det(X)^{-l_1 - l_2}. \end{aligned}$$

(5) Let $w_2 = W \in \mathfrak{p}_2^+$ be of rank 1. Then for $k, l_1, l_3 \in \mathbb{Z}_{\geq 0}$ with $l_3 \leq k$ we have

$$\begin{aligned} & F_{(0,0,-k),(l_1,0,l_3)}^\downarrow \left[((\cdot)^{-1}|w_2)_{\mathfrak{n}_2^+}^k \right] (x_1, x_2) = F_{(k,0,0),(l_3,0,l_1)}^\uparrow \left[(\cdot|w_2)_{\mathfrak{n}_2^+}^k \right] (x_1, x_2^{-1}) \\ &= \frac{(-1)^{l_1}}{(-k - l_1 - d_2)_{l_3} l_3! (l_1 + l_3)!} \sum_{j=l_3}^{\min\{k, l_1 + l_3\}} \frac{(-k)_j (-l_1 - l_3)_j}{(-k - l_1 + l_3 - d_2 + 1)_{j-l_3} (j - l_3)!} \\ & \quad \times (x_2 | (x_1)^\sharp)^{l_1 + l_3 - j} (x_2^{-1} | w_2)^{k-j} ((x_1)^\sharp | w_2)^j \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_3} \\ &= \frac{(-1)^{l_3}}{(-k - l_1 - d_2)_{l_3} l_3! (l_1 + l_3)!} \sum_{j=l_3}^{\min\{k, l_1 + l_3\}} \frac{(-k)_j (-l_1 - l_3)_j}{(-k - l_1 + l_3 - d_2 + 1)_{j-l_3} (j - l_3)!} \\ & \quad \times (X | {}^t \hat{\xi} \xi)^{l_1 + l_3 - j} (X^{-1} | W)^{k-j} ({}^t \hat{\xi} \xi | W)^j \det(X)^{-l_1 - l_3}. \end{aligned}$$

Proof. (1) Clear from $\mathcal{P}_{\mathbf{k}+(k,k,k)}(\mathfrak{p}_2^+) = \det_{\mathfrak{n}_2^+}(x_2)^k \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$.

(2) By the definition of $F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2)$, $F_{\mathbf{k}, \mathbf{l}}^\uparrow[f](x_1, x_2)$,

$$\begin{aligned} F_{(k,k,k), (l,0,0)}^\downarrow \left[\det_{\mathfrak{n}_2^+}^k \right] (x_1, x_2) &= \text{Proj}_{(k,k-l,k-l), x_2}^{\mathfrak{p}_2^+} \left(\frac{1}{l!} \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k \right) \\ &= \frac{1}{l!} \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k = \frac{1}{l!} \left(\frac{\xi X {}^t \hat{\xi}}{\det(X)} \right)^l \det(X)^k, \\ F_{(k,k,k), (0,0,l)}^\uparrow \left[\det_{\mathfrak{n}_2^+}^k \right] (x_1, x_2) &= \text{Proj}_{(k+l, k+l, k), x_2}^{\mathfrak{p}_2^+} \left(\frac{1}{l!} \left(-((x_1)^\sharp | (x_2)^\sharp) \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k \right) \\ &= \frac{1}{l!} \left(-((x_1)^\sharp | (x_2)^\sharp) \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k = \frac{1}{l!} (\xi X^\sharp {}^t \hat{\xi})^l \det(X)^k. \end{aligned}$$

(3) By Lemma 2.4 (3) we have

$$\begin{aligned} F_{(k_1, k_2, 0), (0,0,l)}^\downarrow[f](x_1, x_2) &= \tilde{F}_{(k_1, k_2, 0), (0,0,l)}[f](x_1, x_2) \det_{\mathfrak{n}_2^+}(x_2)^{-l} \\ &= \frac{(-1)^l}{l!} \text{Proj}_{(k_1, k_2, l), x_2}^{\mathfrak{p}_2^+} ((x_2 | (x_1)^\sharp)^l f(x_2'')) \det_{\mathfrak{n}_2^+}(x_2)^{-l} \\ &= \frac{(-1)^l}{l!} \frac{(-k_2)_l (-k_1 - \frac{d_2}{2})_l}{(-k_2 - \frac{d_2}{2})_l (-k_1 - d_2)_l} \frac{((x_1)^\sharp | e')^l}{\det_{\mathfrak{n}_2^{++}}(x_2'')^l} f(x_2''), \end{aligned}$$

and we have

$$((x_1)^\sharp | e') = \det_{\mathfrak{n}^{++}}(x_2''), \quad ((x_1)^\sharp | e') = -({}^t \hat{\xi} \xi | e').$$

(4) By Proposition 2.5 (2) we have

$$\begin{aligned} F_{(k,0,0), (l_1, l_2, 0)}^\downarrow \left[(\cdot | w_2)_{\mathfrak{n}_2^+}^k \right] (x_1, x_2) &= \tilde{F}_{(k,0,0), (l_1, l_2, 0)} \left[(\cdot | w_2)_{\mathfrak{n}_2^+}^k \right] (x_1, x_2) \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_2} \\ &= \frac{(-1)^{l_1 + l_2}}{(l_1 + l_2)!} \text{Proj}_{(k+l_1, l_2, 0), x_2}^{\mathfrak{p}_2^+} ((x_2 | w_2)^k (x_2 | (x_1)^\sharp)^{l_1 + l_2}) \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_2} \\ &= \frac{(-1)^{l_1 + l_2}}{(l_1 + l_2)!} \frac{(-1)^{l_2}}{(-k - l_1 - \frac{d_2}{2})_{l_2} l_2!} \sum_{j=l_2}^{\min\{k, l_1 + l_2\}} \frac{(-k)_j (-l_1 - l_2)_j}{(-k - l_1 + l_2 - \frac{d_2}{2} + 1)_{j-l_2} (j - l_2)!} \\ & \quad \times (x_2 | (x_1)^\sharp)^{l_1 + l_2 - j} (x_2 | w_2)^{k-j} ((x_2 | (x_1)^\sharp)(x_2 | w_2) - (x_2 | (x_1)^\sharp x_2 | w_2))^j \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_2} \end{aligned}$$

$$= \frac{1}{(l_1 + l_2)!} \frac{(-1)^{l_2}}{\left(-k - l_1 - \frac{d_2}{2}\right)_{l_2} l_2!} \sum_{j=l_2}^{\min\{k, l_1 + l_2\}} \frac{(-k)_j (-l_1 - l_2)_j}{\left(-k - l_1 + l_2 - \frac{d_2}{2} + 1\right)_{j-l_2} (j - l_2)!} \\ \times (X|^{t\hat{\xi}\xi})^{l_1 + l_2 - j} (X|W)^{k-j} ((X|^{t\hat{\xi}\xi})(X|W) - (X^{t\hat{\xi}\xi} X|W))^j \det(X)^{-l_1 - l_2}.$$

(5) By Proposition 2.5 (1) we have

$$\begin{aligned} F_{(0,0,-k), (l_1,0,l_3)}^\downarrow [((\cdot)^{-1}|w_2)_{\mathfrak{n}_2^+}^k](x_1, x_2) &= \tilde{F}_{(0,0,-k), (l_1,0,l_3)}^\downarrow [((\cdot)^{-1}|w_2)_{\mathfrak{n}_2^+}^k](x_1, x_2) \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_3} \\ &= \frac{(-1)^{l_1 + l_3}}{(l_1 + l_3)!} \text{Proj}_{(l_1,0,l_3-k), x_2}^{\mathfrak{p}_2^+} ((x_2^{-1}|w_2)^k (x_2|(x_1)^\sharp)^{l_1 + l_3}) \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_3} \\ &= \frac{(-1)^{l_1 + l_3}}{(l_1 + l_3)!} \frac{(-1)^{l_3}}{(-k - l_1 - d_2)_{l_3} l_3!} \sum_{j=l_3}^{\min\{k, l_1 + l_3\}} \frac{(-k)_j (-l_1 - l_3)_j}{(-k - l_1 + l_3 - d_2 + 1)_{j-l_3} (j - l_3)!} \\ &\quad \times (x_2|(x_1)^\sharp)^{l_1 + l_3 - j} (x_2^{-1}|w_2)^{k-j} ((x_1)^\sharp|w_2)^j \det_{\mathfrak{n}_2^+}(x_2)^{-l_1 - l_3} \\ &= \frac{1}{(l_1 + l_3)!} \frac{(-1)^{l_3}}{(-k - l_1 - d_2)_{l_3} l_3!} \sum_{j=l_3}^{\min\{k, l_1 + l_3\}} \frac{(-k)_j (-l_1 - l_3)_j}{(-k - l_1 + l_3 - d_2 + 1)_{j-l_3} (j - l_3)!} \\ &\quad \times (X|^{t\hat{\xi}\xi})^{l_1 + l_3 - j} (X^{-1}|W)^{k-j} (t\hat{\xi}\xi|W)^j \det(X)^{-l_1 - l_3}. \end{aligned}$$

□

Using $F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2)$, for $\lambda \in \mathbb{C}$, $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$ and for $f(x) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^\pm)$, we define a function of $x = (x_1, x_2) \in \Omega$ by

$$\begin{aligned} F^\downarrow(\lambda, \mathbf{k}; f; x) &:= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2 \\ 0 \leq l_3 \leq k_2 - k_3}} \frac{((-k_3, -k_2, -k_1))_{(l_1 + l_2, l_1 + l_3, l_2 + l_3), d_2}}{(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}} F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) \\ &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2 \\ 0 \leq l_3 \leq k_2 - k_3}} \frac{(-k_1 - \frac{d}{2})_{l_2 + l_3} (-k_2 - \frac{d}{4})_{l_1 + l_3} (-k_3)_{l_1 + l_2} F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2)}{(-\lambda - k_2 - k_3 + \frac{3}{4}d + 1)_{l_1} (-\lambda - k_1 - k_3 + \frac{1}{2}d + 1)_{l_2} (-\lambda - k_1 - k_2 + \frac{1}{4}d + 1)_{l_3}}, \end{aligned} \tag{3.2}$$

where $(\lambda + \mathbf{s})_{\mathbf{m}, d_2}$ is as in (2.6), and recall that $d_2 = d/2$. This has continuous parameters $(\lambda, k_3) \in \mathbb{C}^2$ and discrete parameters $(k'_1, k'_2) := (k_1 - k_3, k_2 - k_3) \in \mathbb{Z}_{++}^2$. When we put $f'(x_2) := f(x_2) \det_{\mathfrak{n}_2^\pm}(x_2)^{-k_3} \in \mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^\pm)$, we have

$$\begin{aligned} F^\downarrow(\lambda, \mathbf{k}; f; x) &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k'_1 - k'_2 \\ 0 \leq l_3 \leq k'_2}} \frac{(-k_3 - (0, k'_2, k'_1))_{(l_1 + l_2, l_1 + l_3, l_2 + l_3), d_2}}{(-\lambda - 2k_3 + \frac{3}{2}d_2 + 1 - (k'_2, k'_1, k'_1 + k'_2))_{(l_1, l_2, l_3), d_2}} \\ &\quad \times \det_{\mathfrak{n}_2^\pm}(x_2)^{k_3} F_{(k'_1, k'_2, 0), \mathbf{l}}^\downarrow[f'](x_1, x_2), \end{aligned}$$

and when we fix $(k'_1, k'_2) \in \mathbb{Z}_{++}^2$, as a function of $(\lambda, k_3) \in \mathbb{C}^2$ this has poles at

$$\begin{aligned} \lambda + 2k_3 \in & \left\{ \frac{3}{2}d_2 + 1 - k'_2 + j \mid j \in \mathbb{Z}_{\geq 0} \right\} \cup \{d_2 + 1 - k'_1 + j \mid j \in \mathbb{Z}, 0 \leq j < k'_1 - k'_2\} \\ & \cup \left\{ \frac{1}{2}d_2 + 1 - (k'_1 + k'_2) + j \mid j \in \mathbb{Z}, 0 \leq j < k'_2 \right\}. \end{aligned}$$

If we additionally fix $k_3 \in \mathbb{Z}_{\geq 0}$, then $F^\downarrow(\lambda, \mathbf{k}; f; x)$ becomes a polynomial in x , and the singularities $\lambda \in \{\frac{3}{2}d_2 + 1 - k'_2 - 2k_3 + j \mid j \in \mathbb{Z}_{\geq k_3}\}$ are removable. We use the same notation $F^\downarrow(\lambda, \mathbf{k}; f; x)$ to express this polynomial also for these λ .

Proposition 3.3. *Suppose $\lambda \in \mathbb{C}$ is not a pole of $F^\downarrow(\lambda, \mathbf{k}; f; x)$. Then $F^\downarrow(\lambda, \mathbf{k}; f; x)$ converges if $x_2 = ge$ for some $g \in K_1^{\mathbb{C}}$ and $(g^{-1}x_1|\overline{g^{-1}x_1})_{\mathfrak{p}^+} < 2$. Especially, this converges for $x = (x_1, x_2) \in \Omega$.*

Proof. By Proposition 3.1, for $x_1 \in \mathfrak{p}_1^+$, $x_2 \in ge \in \mathfrak{p}_2^+$ with $g \in \tilde{K}_1^{\mathbb{C}}$, we have

$$|F^\downarrow(\lambda, \mathbf{k}; f; x)| \leq \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2 \\ 0 \leq l_3 \leq k_2 - k_3}} \frac{|((-k_3, -k_2, -k_1))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2}| \sqrt{d_{(\mathbf{l}, 0, 0)}^{\mathfrak{p}_2^+} d_{(\mathbf{l}, 0, 0)}^{\mathfrak{p}_2^+}}}{|(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}| |\mathbf{l}|!} \times \frac{(g^{-1}x_1|\overline{g^{-1}x_1})_{\mathfrak{p}^+}^{|\mathbf{l}|}}{2^{|\mathbf{l}|}} \|f(g(\cdot))\|_{L^2(\Sigma_2)}.$$

Then

$$d_{(\mathbf{l}, 0, 0)}^{\mathfrak{p}_2^+} = \dim \mathcal{P}_{(\mathbf{l}, 0, 0)}(\mathfrak{p}_2^+) \leq \dim \mathcal{P}_{|\mathbf{l}|}(\mathfrak{p}_2^+) = \binom{|\mathbf{l}| + n_2 - 1}{n_2 - 1}$$

is of polynomial growth, where $n_2 := \dim \mathfrak{p}_2^+$. Also, since l_2, l_3 run over the finite sets, by Stirling's formula, there exist constants $C_{\lambda, \mathbf{k}}, C'_{\lambda, \mathbf{k}} \in \mathbb{R}_{>0}$ such that

$$\begin{aligned} & \frac{|((-k_3, -k_2, -k_1))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2}|}{|(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}| |\mathbf{l}|!} \\ & \leq C_{\lambda, \mathbf{k}} \left| \frac{(l_1 + l_2)^{-k_3 + l_1 + l_2 - \frac{1}{2}} e^{-l_1 - l_2} (l_1 + l_3)^{-k_2 - \frac{1}{2}d_2 + l_1 + l_3 - \frac{1}{2}} e^{-l_1 - l_3}}{l_1^{-\lambda + \frac{3}{2}d_2 - k_2 - k_3 + l_1 + \frac{1}{2}} e^{-l_1} (l_1 + l_2 + l_3)^{l_1 + l_2 + l_3 + \frac{1}{2}} e^{-l_1 - l_2 - l_3}} \right| \leq C'_{\lambda, \mathbf{k}} |\mathbf{l}|^{\operatorname{Re} \lambda - 2d_2 - 2} \end{aligned}$$

holds. Hence the above converges if $(g^{-1}x_1|\overline{g^{-1}x_1})_{\mathfrak{p}^+} < 2$. Especially, if $x = (x_1, x_2) = (x_1, ge) \in \Omega$ with $g \in L_2 = K_1^{\mathbb{C}} \cap L$, then by

$$\begin{aligned} 0 < \det_{\mathfrak{n}^+}(x) &= \det_{\mathfrak{n}^+}(x_2) \det_{\mathfrak{n}^+}(e + g^{-1}x_1) = \det_{\mathfrak{n}_2^+}(x_2) \left(1 - \frac{(g^{-1}x_1|\overline{g^{-1}x_1})_{\mathfrak{n}^+}}{2} \right) \\ &= \det_{\mathfrak{n}_2^+}(x_2) \left(1 - \frac{(g^{-1}x_1|\overline{g^{-1}x_1})_{\mathfrak{p}^+}}{2} \right), \end{aligned}$$

we have $(g^{-1}x_1|\overline{g^{-1}x_1})_{\mathfrak{p}^+} < 2$. Hence the above converges if $x = (x_1, x_2) \in \Omega$. \square

This $F^\downarrow(\lambda, \mathbf{k}; f; x)$ has the following property. Recall that $\underline{k}_3 := (k, k, k)$.

Proposition 3.4. *For $\lambda \in \mathbb{C}$, $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$, $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, $k \in \mathbb{Z}_{\geq 0}$, we have*

$$\begin{aligned} & \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial x_2} \right)^k F^\downarrow(\lambda, \mathbf{k}; f; x) \\ &= (-1)^k ((-k_3, -k_2, -k_1))_{\underline{k}_3, d_2} F^\downarrow(\lambda + 2k, \mathbf{k} - \underline{k}_3; f \det_{\mathfrak{n}_2^+}^{-k}; x). \end{aligned}$$

Proof. Since $F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) \in \mathcal{P}_{2|\mathbf{k}|}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{(k_1-l_2-l_3, k_2-l_1-l_3, k_3-l_1-l_2)}(\mathfrak{p}_2^+)$ holds, by Lemma 2.3 we have

$$\begin{aligned} & \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial x_2} \right)^k ((-k_3, -k_2, -k_1))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2} F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) \\ &= (-1)^{3k} ((-k_3 + l_1 + l_2, -k_2 + l_1 + l_3, -k_1 + l_2 + l_3))_{(k, k, k), d_2} \\ &\quad \times ((-k_3, -k_2, -k_1))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2} F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](x_1, x_2) \det_{\mathfrak{n}_2^+}(x_2)^{-k} \\ &= (-1)^k ((-k_3, -k_2, -k_1))_{(k+l_1+l_2, k+l_1+l_3, k+l_2+l_3), d_2} F_{\mathbf{k}-\underline{k}_3, \mathbf{l}}^\downarrow[f \det_{\mathfrak{n}_2^+}^{-k}](x_1, x_2) \\ &= (-1)^k ((-k_3, -k_2, -k_1))_{(k, k, k), d_2} \\ &\quad \times ((-k_3 + k, -k_2 + k, -k_1 + k))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2} F_{\mathbf{k}-\underline{k}_3, \mathbf{l}}^\downarrow[f \det_{\mathfrak{n}_2^+}^{-k}](x_1, x_2), \end{aligned}$$

and easily we have

$$\begin{aligned} & \left(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2) \right)_{1, d_2} \\ &= \left(-(\lambda + 2k) + \frac{3}{2}d_2 + 1 - ((k_2 - k) + (k_3 - k), (k_1 - k) + (k_3 - k), (k_1 - k) + (k_2 - k)) \right)_{1, d_2}. \end{aligned}$$

Hence the desired formula follows from the definition of $F^\downarrow(\lambda, \mathbf{k}; f; x)$. \square

3.3 Results on weighted Bergman inner products

For $\lambda \in \mathbb{C}$, $\mathbf{k} \in \mathbb{Z}_{++}^3$, let

$$\begin{aligned} C^{d, d_2}(\lambda, \mathbf{k}) &:= \frac{\prod_{1 \leq i < j \leq 3} (\lambda - \frac{d}{4}(i+j-2))_{k_i+k_j}}{\prod_{1 \leq i < j \leq 4} (\lambda - \frac{d}{4}(i+j-3))_{k_i+k_j}} \\ &= \frac{(\lambda + k_1 + k_3 - \frac{d}{4})_{k_2-k_3} (\lambda + \max\{k_1, k_2 + k_3\} - \frac{d}{2})_{\min\{k_3, k_1-k_2\}} (\lambda + k_2 - \frac{3}{4}d)_{k_3}}{(\lambda)_{k_1+k_2} (\lambda - \frac{d}{2})_{\min\{k_1, k_2+k_3\}} (\lambda - d)_{k_3}} \\ &= \frac{(\lambda - \frac{d_2}{2} + (k_1 + k_3, \max\{k_1, k_2 + k_3\}, k_2))_{(k_2-k_3, \min\{k_3, k_1-k_2\}, k_3), d_2}}{(\lambda)_{(k_1+k_2, \min\{k_1, k_2+k_3\}, k_3), d}}, \end{aligned} \tag{3.3}$$

where $k_4 := 0$, $d_2 = d/2$. Then the following holds.

Theorem 3.5. Suppose $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3 = \mathbb{Z}_{++}^3 + \mathbb{C}(1, 1, 1)$, $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$.

(1) When $\mathbf{k} \in \mathbb{Z}_{++}^3$, for $\operatorname{Re} \lambda > 2d + 1$ we have

$$\left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} = C^{d, d_2}(\lambda, \mathbf{k}) F^\downarrow(\lambda, \mathbf{k}; f; z).$$

(2) When $\mathbf{k} \in \mathbb{Z}_{++}^3$, as a function of λ ,

$$(\lambda)_{(k_1+k_2, \min\{k_1, k_2+k_3\}, k_3), d} \left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x}$$

is holomorphically continued for all \mathbb{C} , and give non-zero polynomials in z for all $\lambda \in \mathbb{C}$ if $f(x_2) \neq 0$.

(3) For $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$, $\lambda \in \mathbb{C}$, we have

$$\begin{aligned} & F^\downarrow(\lambda, \mathbf{k}; f; z) \\ &= \det_{\mathfrak{n}^+}(z)^{-\lambda+d+1} F^\downarrow\left(-\lambda + 2(d+1), \mathbf{k} + \underline{\lambda - d - 1}_3; f \det_{\mathfrak{n}_2^+}^{\lambda-d-1}; z\right). \end{aligned}$$

Especially if $\mathbf{k} \in \mathbb{Z}_{++}^3$ and $\lambda = d+1-a$ with $a = 1, 2, \dots, k_3$, then $F^\downarrow(\lambda, \mathbf{k}; f; z)$ is factorized as

$$F^\downarrow(d+1-a, \mathbf{k}; f; z) = \det_{\mathfrak{n}^+}(z)^a F^\downarrow\left(d+1+a, \mathbf{k} - \underline{a}_3; f \det_{\mathfrak{n}_2^+}^{-a}; z\right).$$

Theorem 3.5 (1), (3) solves [35, Conjecture 7.2], and coincides with [34, Theorem 6.3 (1)] for $k_2 = k_3$ case. Also, in (2) the holomorphy is already proved in [35, Theorem 6.2], and the non-zeroness is new except for $k_1 = k_2 = k_3$ or $k_3 = 0$ ([34, Corollary 6.6 (2)]).

Remark 3.6. By [35, Theorem 3.3 (1)], for $\mathbf{k} \in \mathbb{Z}_{++}^3$, $f(x_2) \in \mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+)$, $\operatorname{Re} \lambda > 2d+1$ we have

$$\begin{aligned} \left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} &= \frac{1}{(\lambda)_{2k_3, d}} \det_{\mathfrak{n}^+}(z)^{-\lambda+d+1} \det_{\mathfrak{n}_2^+}\left(\frac{\partial}{\partial z_2}\right)^{k_3} \det_{\mathfrak{n}^+}(z)^{\lambda+2k_3-d-1} \\ &\quad \times \left\langle \det_{\mathfrak{n}_2^+}(x_2)^{-k_3} f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda+2k_3, x}. \end{aligned} \quad (3.4)$$

Combining this with Proposition 3.4 and Theorem 3.5 (1), (3), we have

$$\begin{aligned} & \left\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \frac{C^{d, d_2}(\lambda + 2k_3, \mathbf{k} - \underline{k}_3)}{(\lambda)_{2k_3, d}} \det_{\mathfrak{n}^+}(z)^{-\lambda+d+1} \det_{\mathfrak{n}_2^+}\left(\frac{\partial}{\partial z_2}\right)^{k_3} \det_{\mathfrak{n}^+}(z)^{\lambda+2k_3-d-1} \\ &\quad \times F^\downarrow\left(\lambda + 2k_3, \mathbf{k} - \underline{k}_3; f \det_{\mathfrak{n}_2^+}^{-k_3}; z\right) \\ &= \frac{(\lambda + k_1 + k_3 - \frac{d_2}{2})_{k_2-k_3}}{(\lambda)_{(k_1+k_2, k_2+k_3, 2k_3), d}} \det_{\mathfrak{n}^+}(z)^{-\lambda+d+1} \det_{\mathfrak{n}_2^+}\left(\frac{\partial}{\partial z_2}\right)^{k_3} \\ &\quad \times F^\downarrow\left(-\lambda - 2k_3 + 2(d+1), \mathbf{k} + \underline{\lambda + k_3 - d - 1}_3; f \det_{\mathfrak{n}_2^+}^{\lambda+k_3-d-1}; z\right) \\ &= C^{d, d_2}(\lambda, \mathbf{k}) \det_{\mathfrak{n}^+}(z)^{-\lambda+d+1} F^\downarrow\left(-\lambda + 2(d+1), \mathbf{k} + \underline{\lambda - d - 1}_3; f \det_{\mathfrak{n}_2^+}^{\lambda-d-1}; z\right) \\ &= C^{d, d_2}(\lambda, \mathbf{k}) F^\downarrow(\lambda, \mathbf{k}; f; z). \end{aligned}$$

First we prove (1). By [35, Theorem 6.1], we have

$$\frac{\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \rangle_{\lambda, x}|_{z_1=0}}{f(z_2)} = \frac{\prod_{1 \leq i < j \leq 3} (\lambda - \frac{d}{4}(i+j-2))_{k_i+k_j}}{\prod_{1 \leq i < j \leq 4} (\lambda - \frac{d}{4}(i+j-3))_{k_i+k_j}} = C^{d, d_2}(\lambda, \mathbf{k}).$$

Therefore to prove (1), it is enough to prove that $\langle f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \rangle_{\lambda, x}$ is proportional to $F^\downarrow(\lambda, \mathbf{k}; f; z)$, and by the F-method, enough to prove

$$(f(x_2) \mapsto F^\downarrow(\lambda, \mathbf{k}; f; z)) \in \operatorname{Hom}_{K_1}(\mathcal{P}_\mathbf{k}(\mathfrak{p}_2^+), \operatorname{Sol}_{\mathcal{P}(\mathfrak{p}^+)}((\mathcal{B}_\lambda)_1)), \quad (3.5)$$

where $(\mathcal{B}_\lambda)_1$ is the orthogonal projection of $\mathcal{B}_\lambda = \mathcal{B}_\lambda^{\mathfrak{p}^+} : \mathcal{P}(\mathfrak{p}^+) \rightarrow \mathcal{P}(\mathfrak{p}^+) \otimes \mathfrak{p}^+$ given in (2.24) (with the identification $\mathfrak{p}^- \simeq \mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}}$) onto \mathfrak{p}_1^+ . This follows from the next proposition. Recall that we write $z = (z_1, z_2) = (\zeta, Z) \in \mathfrak{p}_1^+ \oplus \mathfrak{p}_2^+ \simeq (\mathbb{F}^{\mathbb{C}})^3 \oplus \text{Herm}(3, \mathbb{F}')^{\mathbb{C}}$, and use lowercase Greek alphabets and capital Latin alphabets to express elements in $(\mathbb{F}'^{\mathbb{C}})^3$ and in $\text{Herm}(3, \mathbb{F}')^{\mathbb{C}}$ respectively. Also we abbreviate $\det_{\mathfrak{n}_2^+}(X) =: \det(X)$, and let $(X|Y) := \text{Re}_{\mathbb{F}'} \text{tr}(XY)$, $(\xi|\eta) := \text{Re}_{\mathbb{F}'}(\xi^t \hat{\eta})$, so that $((\xi, X)|(\eta, Y))_{\mathfrak{n}^+} = 2(\xi|\eta) + (X|Y)$ holds. Especially we have $-((z_1)^\sharp|z_2)_{\mathfrak{n}_2^+} = \zeta Z^t \hat{\zeta}$. Let $\Gamma_3^{d_2}(\lambda + \mathbf{s})$ be as in (2.5).

Proposition 3.7. *Let $\lambda \in \mathbb{C}$, $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$, $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$.*

(1) *If $\text{Re } k_1 < -d_2$, $\text{Re}(\lambda + k_1 + k_2) < -\frac{1}{2}d_2$, then for $z = (\zeta, Z) \in \Omega$, $A \in \Omega_2$, we have*

$$\begin{aligned} F^\downarrow(\lambda, \mathbf{k}; f; \zeta, Z) \\ = \frac{\Gamma_3^{d_2}(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))} \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda + |\mathbf{k}| - 2d_2 - 1} \\ \times \left(\frac{1}{(2\pi\sqrt{-1})^{n_2}} \int_{A + \sqrt{-1}\mathfrak{n}_2^+} e^{(X|Y^\sharp)} e^{\zeta X^{-1} t \hat{\zeta}} \det(X)^{\lambda + |\mathbf{k}| - \frac{3}{2}d_2 - 1} f(X^{-1}) dX \right) dY. \end{aligned}$$

(2) *Suppose $\lambda \in \mathbb{C}$ is not a pole of $F^\downarrow(\lambda, \mathbf{k}; f; \zeta, Z)$. Then for $g \in \tilde{K}_1^{\mathbb{C}}$ we have*

$$F^\downarrow(\lambda, \mathbf{k}; f(g(\cdot)); \zeta, Z) = F^\downarrow(\lambda, \mathbf{k}; f; g(\zeta, Z)).$$

(3) *$F^\downarrow(\lambda, \mathbf{k}; f; \zeta, Z)$ satisfies the differential equation*

$$(\mathcal{B}_\lambda)_1 F^\downarrow(\lambda, \mathbf{k}; f; \zeta, Z) = 0.$$

Proof. (1) By the definition of $\tilde{F}_{\mathbf{k}, \mathbf{l}}[f]$ and Lemma 2.2, we have

$$\begin{aligned} & \frac{\Gamma_3^{d_2}(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))} \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda + |\mathbf{k}| - 2d_2 - 1} \\ & \times \left(\frac{1}{(2\pi\sqrt{-1})^{n_2}} \int_{A + \sqrt{-1}\mathfrak{n}_2^+} e^{(X|Y^\sharp)} e^{\zeta X^{-1} t \hat{\zeta}} \det(X)^{\lambda + |\mathbf{k}| - \frac{3}{2}d_2 - 1} f(X^{-1}) dX \right) dY \\ &= \frac{\Gamma_3^{d_2}(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))} \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda + |\mathbf{k}| - 2d_2 - 1} \\ & \times \left(\frac{1}{(2\pi\sqrt{-1})^{n_2}} \int_{A + \sqrt{-1}\mathfrak{n}_2^+} e^{(X|Y^\sharp)} \det(X)^{\lambda + |\mathbf{k}| - \frac{3}{2}d_2 - 1} \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2 \\ 0 \leq l_3 \leq k_2 - k_3}} \tilde{F}_{\mathbf{k}, \mathbf{l}}[f](\zeta, X^{-1}) dX \right) dY \\ &= \frac{1}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))} \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda + |\mathbf{k}| - 2d_2 - 1} \det(Y^\sharp)^{-\lambda - |\mathbf{k}| + \frac{1}{2}d_2} \\ & \times \sum_{\mathbf{l}} \frac{\Gamma_3^{d_2}(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))}{\Gamma_3^{d_2}(-\lambda - |\mathbf{k}| + \frac{3}{2}d_2 + 1 + (k_1 + l_1, k_2 + l_2, k_3 + l_3))} \tilde{F}_{\mathbf{k}, \mathbf{l}}[f](\zeta, Y^\sharp) dY \\ &= \frac{1}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))} \int_{\Omega_2} \sum_{\mathbf{l}} \frac{e^{-(Y|Z)} \det(Y)^{-|\mathbf{k}| - d_2 - 1} \tilde{F}_{\mathbf{k}, \mathbf{l}}[f](\zeta, \det(Y)Y^{-1})}{(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}} dY \\ &= \frac{1}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))} \int_{\Omega_2} \sum_{\mathbf{l}} \frac{e^{-(Y|Z)} \det(Y)^{|\mathbf{l}| - d_2 - 1} \tilde{F}_{\mathbf{k}, \mathbf{l}}[f](\zeta, Y^{-1})}{(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}} dY \end{aligned}$$

$$\begin{aligned}
&= \sum_{\mathbf{l}} \frac{\Gamma_3^{d_2}(|\mathbf{l}| + (-k_3 - l_3, -k_2 - l_2, -k_1 - l_1)) \det(Z)^{-|\mathbf{l}|} \tilde{F}_{\mathbf{k}, \mathbf{l}}[f](\zeta, Z)}{\Gamma_3^{d_2}((-k_3, -k_2, -k_1))(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}} \\
&= \sum_{\mathbf{l}} \frac{((-k_3, -k_2, -k_1))_{(l_1 + l_2, l_1 + l_3, l_2 + l_3), d_2} F_{\mathbf{k}, \mathbf{l}}^{\downarrow}[f](\zeta, Z)}{(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_1 + k_3, k_1 + k_2))_{(l_1, l_2, l_3), d_2}} = F^{\downarrow}(\lambda, \mathbf{k}; f; \zeta, Z).
\end{aligned}$$

Hence we get the desired formula.

(2) Clear from the $\tilde{K}_1^{\mathbb{C}}$ -equivariance of $f(X) \mapsto F_{\mathbf{k}, \mathbf{l}}^{\downarrow}[f](\zeta, Z)$.

(3) We take bases $\{E_{\alpha}\} \subset \mathfrak{n}_2^+$, $\{\epsilon_{\alpha}\} \subset \mathfrak{n}_1^+$, let $\{E_{\alpha}^{\vee}\} \subset \mathfrak{n}_2^+$ be the dual basis of $\{E_{\alpha}\}$ with respect to $(\cdot | \cdot)_{\mathfrak{n}_2^+}$, and let $\{\epsilon_{\alpha}^{\vee}\} \subset \mathfrak{n}_1^+$ be the dual basis of $\{\epsilon_{\alpha}\}$ with respect to the bilinear form $(\xi | \eta) = \text{Re}_{\mathbb{F}'}(\xi^t \hat{\eta})$, so that $\{\epsilon_{\alpha}, E_{\alpha}\}$ and $\{\frac{1}{2}\epsilon_{\alpha}^{\vee}, E_{\alpha}^{\vee}\}$ are mutually dual with respect to $(\cdot | \cdot)_{\mathfrak{n}^+}$. Also let $\partial/\partial Z_{\alpha}$ and $\partial/\partial \zeta_{\alpha}$ denote the directional derivatives along E_{α} and ϵ_{α} respectively. Then $(\mathcal{B}_{\lambda})_1$ is given by

$$\begin{aligned}
(\mathcal{B}_{\lambda})_1 &= \frac{1}{8} \sum_{\alpha \beta} P(\epsilon_{\alpha}^{\vee}, \epsilon_{\beta}^{\vee}) \zeta \frac{\partial^2}{\partial \zeta_{\alpha} \partial \zeta_{\beta}} + \frac{1}{2} \sum_{\alpha \beta} P(E_{\alpha}^{\vee}, E_{\beta}^{\vee}) \zeta \frac{\partial^2}{\partial Z_{\alpha} \partial Z_{\beta}} \\
&\quad + \frac{1}{2} \sum_{\alpha \beta} P(\epsilon_{\alpha}^{\vee}, E_{\beta}^{\vee}) Z \frac{\partial^2}{\partial \zeta_{\alpha} \partial Z_{\beta}} + \frac{\lambda}{2} \sum_{\alpha} \epsilon_{\alpha}^{\vee} \frac{\partial}{\partial \zeta_{\alpha}}.
\end{aligned}$$

We put $\det(X)^{\lambda + |\mathbf{k}| - \frac{3}{2}d_2 - 1} f(X^{-1}) =: F(X)$. By (1) it is enough to prove, for $\text{Re } k_1 < -d_2$, $\text{Re } (\lambda + k_1 + k_2) < -\frac{1}{2}d_2$,

$$(\mathcal{B}_{\lambda})_1 \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda + |\mathbf{k}| - 2d_2 - 1} \left(\int_{A + \sqrt{-1}\mathfrak{n}_2^+} e^{(X|Y^{\sharp})} e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY = 0.$$

First we consider $(\mathcal{B}_{\lambda})_1 e^{-(Y|Z)} e^{\zeta X^{-1} t \hat{\zeta}}$. In the following we use the notation $([\frac{\partial}{\partial w} f] g) h := \frac{\partial f}{\partial w} gh + f \frac{\partial g}{\partial w} h$. Then we have

$$\begin{aligned}
&(\mathcal{B}_{\lambda})_1 e^{-(Y|Z)} e^{\zeta X^{-1} t \hat{\zeta}} \\
&= \left[\frac{1}{8} \sum_{\alpha \beta} P(\epsilon_{\alpha}^{\vee}, \epsilon_{\beta}^{\vee}) \zeta \frac{\partial^2}{\partial \zeta_{\alpha} \partial \zeta_{\beta}} + \frac{1}{2} \sum_{\alpha \beta} P(E_{\alpha}^{\vee}, E_{\beta}^{\vee}) \zeta \frac{\partial^2}{\partial Z_{\alpha} \partial Z_{\beta}} \right. \\
&\quad \left. + \frac{1}{2} \sum_{\alpha \beta} P(\epsilon_{\alpha}^{\vee}, E_{\beta}^{\vee}) Z \frac{\partial^2}{\partial \zeta_{\alpha} \partial Z_{\beta}} + \frac{\lambda}{2} \sum_{\alpha} \epsilon_{\alpha}^{\vee} \frac{\partial}{\partial \zeta_{\alpha}} \right] e^{-(Y|Z)} e^{\zeta X^{-1} t \hat{\zeta}} \\
&= \left(\frac{1}{8} \sum_{\alpha \beta} P(\epsilon_{\alpha}^{\vee}, \epsilon_{\beta}^{\vee}) \zeta (4(\zeta X^{-1} | \epsilon_{\alpha})(\zeta X^{-1} | \epsilon_{\beta}) + 2(\epsilon_{\alpha} X^{-1} | \epsilon_{\beta})) + \frac{1}{2} P(Y, Y) \zeta \right. \\
&\quad \left. - \sum_{\alpha} P(\epsilon_{\alpha}^{\vee}, Y) Z (\zeta X^{-1} | \epsilon_{\alpha}) + \lambda \sum_{\alpha} \epsilon_{\alpha}^{\vee} (\zeta X^{-1} | \epsilon_{\alpha}) \right) e^{-(Y|Z)} e^{\zeta X^{-1} t \hat{\zeta}} \\
&= \left(P(\zeta X^{-1}) \zeta + \frac{1}{4} \sum_{\alpha} P(\epsilon_{\alpha}^{\vee}, \epsilon_{\alpha} X^{-1}) \zeta + P(Y) \zeta - P(\zeta X^{-1}, Y) Z + \lambda \zeta X^{-1} \right) e^{-(Y|Z)} e^{\zeta X^{-1} t \hat{\zeta}} \\
&= \left[P(\zeta X^{-1}) \zeta + \zeta Y^{\sharp} + \left(\lambda - \frac{d_2}{2} + 1 \right) \zeta X^{-1} + \sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} \frac{\partial}{\partial Y_{\alpha}} \right] e^{-(Y|Z)} e^{\zeta X^{-1} t \hat{\zeta}},
\end{aligned}$$

where at the last equality we have used

$$\sum_{\alpha} P(\epsilon_{\alpha}^{\vee}, \epsilon_{\alpha} X^{-1}) \zeta = \sum_{\alpha} (\epsilon_{\alpha}^{\vee} t \hat{\zeta} \epsilon_{\alpha} X^{-1} + \epsilon_{\alpha} X^{-1} t \hat{\zeta} \epsilon_{\alpha}^{\vee}) = 2(2 - d_2) \zeta X^{-1},$$

which follows from $\sum_{\alpha} \epsilon_{\alpha}^{\vee t} \hat{\xi} \epsilon_{\alpha} = \sum_{\alpha} \epsilon_{\alpha}^t \hat{\xi} \epsilon_{\alpha}^{\vee} = (2 - d_2) \xi$. Thus we get

$$\begin{aligned}
& (\mathcal{B}_{\lambda})_1 \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{(X|Y^\sharp)} e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY \\
&= \int_{\Omega_2} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} \left(\left[P(\zeta X^{-1}) \zeta + \zeta Y^\sharp + \left(\lambda - \frac{d_2}{2} + 1 \right) \zeta X^{-1} \right. \right. \right. \right. \\
&\quad \left. \left. \left. \left. + \sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} \frac{\partial}{\partial Y_{\alpha}} \right] e^{-(Y|Z)} \right) \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY \\
&= \int_{\Omega_2} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{-(Y|Z)} \left(\left[P(\zeta X^{-1}) \zeta + \zeta Y^\sharp + \left(\lambda - \frac{d_2}{2} + 1 \right) \zeta X^{-1} \right. \right. \right. \\
&\quad \left. \left. \left. - \sum_{\alpha} \frac{\partial}{\partial Y_{\alpha}} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} \right] \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \right) e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY.
\end{aligned}$$

Next we consider

$$\begin{aligned}
& \sum_{\alpha} \frac{\partial}{\partial Y_{\alpha}} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \\
&= \sum_{\alpha} P(\zeta X^{-1}, E_{\alpha}) E_{\alpha}^{\vee} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \\
&\quad + (2\lambda + |\mathbf{k}| - 2d_2 - 1) \sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} (Y^{-1}|E_{\alpha}) \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \\
&\quad + \sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} (X|Y \times E_{\alpha}) \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)}.
\end{aligned}$$

Then for the 1st term we have

$$\begin{aligned}
\sum_{\alpha} P(\zeta X^{-1}, E_{\alpha}) E_{\alpha}^{\vee} &= \sum_{\alpha} ((E_{\alpha}|E_{\alpha}^{\vee}) \zeta X^{-1} - \zeta X^{-1} E_{\alpha} E_{\alpha}^{\vee}) \\
&= 3(1 + d_2) \zeta X^{-1} - (1 + d_2) \zeta X^{-1} = 2(1 + d_2) \zeta X^{-1},
\end{aligned}$$

for the 2nd term we have

$$\begin{aligned}
\sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} (Y^{-1}|E_{\alpha}) &= P(\zeta X^{-1}, Y) Y^{-1} = (Y|Y^{-1}) \zeta X^{-1} - \zeta X^{-1} Y Y^{-1} \\
&= 3\zeta X^{-1} - \zeta X^{-1} = 2\zeta X^{-1},
\end{aligned}$$

and for the 3rd term we have

$$\begin{aligned}
\sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} (X|Y \times E_{\alpha}) &= \sum_{\alpha} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} (X \times Y|E_{\alpha}) \\
&= P(\zeta X^{-1}, Y) (X \times Y) = (Y|X \times Y) \zeta X^{-1} - \zeta X^{-1} Y (X \times Y) \\
&= \zeta X^{-1} (2(X|Y^\sharp) I - Y(X \times Y)) = \zeta X^{-1} ((X|Y^\sharp) I + XY^\sharp) = (X|Y^\sharp) \zeta X^{-1} + \zeta Y^\sharp,
\end{aligned}$$

where we have used (3.1) at the 5th equality. Hence we have

$$\begin{aligned}
& \left[P(\zeta X^{-1}) \zeta + \zeta Y^\sharp + \left(\lambda - \frac{d_2}{2} + 1 \right) \zeta X^{-1} - \sum_{\alpha} \frac{\partial}{\partial Y_{\alpha}} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} \right] \\
& \quad \times \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)}
\end{aligned}$$

$$\begin{aligned}
&= \left(P(\zeta X^{-1})\zeta + \zeta Y^\sharp + \left(\lambda - \frac{d_2}{2} + 1 \right) \zeta X^{-1} - (X|Y^\sharp)\zeta X^{-1} - \zeta Y^\sharp \right. \\
&\quad \left. - 2(1+d_2)\zeta X^{-1} - 2(2\lambda + |\mathbf{k}| - 2d_2 - 1)\zeta X^{-1} \right) \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \\
&= \left[P(\zeta X^{-1})\zeta + \left(-3\lambda - 2|\mathbf{k}| + \frac{3}{2}d_2 + 1 \right) \zeta X^{-1} - \zeta X^{-1} \sum_{\alpha} (X|E_{\alpha}^{\vee}) \frac{\partial}{\partial X_{\alpha}} \right] \\
&\quad \times \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)},
\end{aligned}$$

and we get

$$\begin{aligned}
&(\mathcal{B}_{\lambda})_1 \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{(X|Y^\sharp)} e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY \\
&= \int_{\Omega_2} e^{-(Y|Z)} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} \left(\left[P(\zeta X^{-1})\zeta + \zeta Y^\sharp + \left(\lambda - \frac{d_2}{2} + 1 \right) \zeta X^{-1} \right. \right. \right. \\
&\quad \left. \left. \left. - \sum_{\alpha} \frac{\partial}{\partial X_{\alpha}} P(\zeta X^{-1}, Y) E_{\alpha}^{\vee} \right] \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \right) e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY \\
&= \int_{\Omega_2} e^{-(Y|Z)} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} \left(\left[P(\zeta X^{-1})\zeta + \left(-3\lambda - 2|\mathbf{k}| + \frac{3}{2}d_2 + 1 \right) \zeta X^{-1} \right. \right. \right. \\
&\quad \left. \left. \left. - \zeta X^{-1} \sum_{\alpha} (X|E_{\alpha}^{\vee}) \frac{\partial}{\partial X_{\alpha}} \right] \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} e^{(X|Y^\sharp)} \right) e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY \\
&= \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{(X|Y^\sharp)} \left(\left[P(\zeta X^{-1})\zeta \right. \right. \right. \\
&\quad \left. \left. \left. + \left(-3\lambda - 2|\mathbf{k}| + \frac{3}{2}d_2 + 1 \right) \zeta X^{-1} + \sum_{\alpha} \frac{\partial}{\partial X_{\alpha}} \zeta X^{-1} (X|E_{\alpha}^{\vee}) \right] e^{\zeta X^{-1} t \hat{\zeta}} F(X) \right) dX \right) dY.
\end{aligned}$$

Then since we have

$$\begin{aligned}
&\sum_{\alpha} \frac{\partial}{\partial X_{\alpha}} \zeta X^{-1} (X|E_{\alpha}^{\vee}) e^{\zeta X^{-1} t \hat{\zeta}} F(X) \\
&= - \sum_{\alpha} \zeta X^{-1} E_{\alpha} X^{-1} (X|E_{\alpha}^{\vee}) e^{\zeta X^{-1} t \hat{\zeta}} F(X) + \sum_{\alpha} \zeta X^{-1} (E_{\alpha}|E_{\alpha}^{\vee}) e^{\zeta X^{-1} t \hat{\zeta}} F(X) \\
&\quad - \sum_{\alpha} \zeta X^{-1} (X|E_{\alpha}^{\vee}) (\zeta X^{-1} E_{\alpha} X^{-1} t \hat{\zeta}) e^{\zeta X^{-1} t \hat{\zeta}} F(X) + \sum_{\alpha} \zeta X^{-1} (X|E_{\alpha}^{\vee}) e^{\zeta X^{-1} t \hat{\zeta}} \frac{\partial F}{\partial X_{\alpha}}(X) \\
&= -\zeta X^{-1} XX^{-1} e^{\zeta X^{-1} t \hat{\zeta}} F(X) + 3(1+d_2)\zeta X^{-1} e^{\zeta X^{-1} t \hat{\zeta}} F(X) \\
&\quad - (\zeta X^{-1} XX^{-1} t \hat{\zeta}) \zeta X^{-1} e^{\zeta X^{-1} t \hat{\zeta}} F(X) + \zeta X^{-1} e^{\zeta X^{-1} t \hat{\zeta}} \sum_{\alpha} (X|E_{\alpha}^{\vee}) \frac{\partial F}{\partial X_{\alpha}}(X) \\
&= e^{\zeta X^{-1} t \hat{\zeta}} \left[(2+3d_2)\zeta X^{-1} - P(\zeta X^{-1})\zeta + \zeta X^{-1} \sum_{\alpha} (X|E_{\alpha}^{\vee}) \frac{\partial}{\partial X_{\alpha}} \right] F(X),
\end{aligned}$$

we get

$$\begin{aligned}
&(\mathcal{B}_{\lambda})_1 \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{(X|Y^\sharp)} e^{\zeta X^{-1} t \hat{\zeta}} F(X) dX \right) dY \\
&= \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{(X|Y^\sharp)} \left(\left[P(\zeta X^{-1})\zeta \right. \right. \right. \\
&\quad \left. \left. \left. + \left(-3\lambda - 2|\mathbf{k}| + \frac{3}{2}d_2 + 1 \right) \zeta X^{-1} + \sum_{\alpha} \frac{\partial}{\partial X_{\alpha}} \zeta X^{-1} (X|E_{\alpha}^{\vee}) \right] e^{\zeta X^{-1} t \hat{\zeta}} F(X) \right) dX \right) dY
\end{aligned}$$

$$\begin{aligned}
&= \int_{\Omega_2} e^{-(Y|Z)} \det(Y)^{2\lambda+|\mathbf{k}|-2d_2-1} \left(\int_{A+\sqrt{-1}\mathbf{n}_2^+} e^{(X|Y^\sharp)} e^{\zeta X^{-1}t\hat{\zeta}} \right. \\
&\quad \times \left. \left[\left(-3\lambda - 2|\mathbf{k}| + \frac{9}{2}d_2 + 3 \right) \zeta X^{-1} + \zeta X^{-1} \sum_{\alpha} (X|E_{\alpha}^\vee) \frac{\partial}{\partial X_{\alpha}} \right] F(X) dX \right) = 0,
\end{aligned}$$

where the last equality holds since $F(X) = \det(X)^{\lambda+|\mathbf{k}|-2d_2-1} f(X^{-1})$ is homogeneous of degree $3\lambda + 2|\mathbf{k}| - \frac{9}{2}d_2 - 3$. Hence we get the desired formula. \square

Especially, when $\mathbf{k} \in \mathbb{Z}_{++}^3$ we have (3.5), and this proves Theorem 3.5 (1).

Proof of Theorem 3.5 (2). By Theorem 3.5 (1) we have

$$\begin{aligned}
&(\lambda)_{(k_1+k_2, \min\{k_1, k_2+k_3\}, k_3), d} \langle f(x_2), e^{(x|\bar{z})_{\mathbf{p}^+}} \rangle_{\lambda, x} \\
&= \left(\lambda - \frac{d_2}{2} + (k_1 + k_3, \max\{k_1, k_2 + k_3\}, k_2) \right)_{(k_2-k_3, \min\{k_3, k_1-k_2\}, k_3), d_2} F^\downarrow(\lambda, \mathbf{k}; f; z) \\
&= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2 \\ 0 \leq l_3 \leq k_2 - k_3 \\ l_1 + l_2 \leq k_3}} \left(\lambda - \frac{d_2}{2} + (k_1 + k_3, \max\{k_1, k_2 + k_3\}, k_2) \right)_{(k_2-k_3-l_3, \min\{k_3, k_1-k_2\}-l_2, k_3-l_1), d_2} \\
&\quad \times (-1)^{|l|} ((-k_3, -k_2, -k_1))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2} F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](z_1, z_2),
\end{aligned}$$

and this is analytically continued for all $\lambda \in \mathbb{C}$. Moreover, all $F_{\mathbf{k}, \mathbf{l}}^\downarrow[f](z_1, z_2)$ are non-zero by Proposition 3.1 (2) and linearly independent, and the coefficients of $F_{\mathbf{k}, (k_3, 0, k_2-k_3)}^\downarrow[f](z_1, z_2)$ and $F_{\mathbf{k}, (\max\{0, k_3-k_1+k_2\}, \min\{k_3, k_1-k_2\}, k_2-k_3)}^\downarrow[f](z_1, z_2)$,

$$\begin{aligned}
&\left(\lambda - \frac{d_2}{2} + (k_1 + k_3, \max\{k_1, k_2 + k_3\}, k_2) \right)_{(0, \min\{k_3, k_1-k_2\}, 0), d_2} \\
&= (\lambda - d_2 + \max\{k_1, k_2 + k_3\})_{\min\{k_3, k_1-k_2\}}, \\
&\left(\lambda - \frac{d_2}{2} + (k_1 + k_3, \max\{k_1, k_2 + k_3\}, k_2) \right)_{(0, 0, \min\{k_3, k_1-k_2\}), d_2} \\
&= \left(\lambda - \frac{3}{2}d_2 + k_2 \right)_{\min\{k_3, k_1-k_2\}}
\end{aligned}$$

do not vanish simultaneously. Hence $(\lambda)_{(k_1+k_2, \min\{k_1, k_2+k_3\}, k_3), d} \langle f(x_2), e^{(x|\bar{z})_{\mathbf{p}^+}} \rangle_{\lambda, x}$ is non-zero. \square

Proof of Theorem 3.5 (3). By combining Theorem 3.5 (1) and (3.4), for $\operatorname{Re} \lambda > 2d + 1$, $\mathbf{k} \in \mathbb{Z}_{++}^3$ we have

$$\begin{aligned}
C^{d, d_2}(\lambda, \mathbf{k}) F^\downarrow(\lambda, \mathbf{k}; f; z) &= \frac{C^{d, d_2}(\lambda + 2k_3, \mathbf{k} - \underline{k}_{33})}{(\lambda)_{2\underline{k}_{33}, d}} \det_{\mathbf{n}^+}(z)^{-\lambda+d+1} \det_{\mathbf{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k_3} \\
&\quad \times \det_{\mathbf{n}^+}(z)^{\lambda+2k_3-d-1} F^\downarrow \left(\lambda + 2k_3, \mathbf{k} - \underline{k}_{33}; f \det_{\mathbf{n}_2^+}^{-k_3}; z \right),
\end{aligned}$$

that is,

$$\begin{aligned}
F^\downarrow(\lambda, \mathbf{k}; f; z) &= \frac{1}{(\lambda - d_2 + \mathbf{k})_{\underline{k}_{33}, d_2}} \det_{\mathbf{n}^+}(z)^{-\lambda+d+1} \det_{\mathbf{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k_3} \\
&\quad \times \det_{\mathbf{n}^+}(z)^{\lambda+2k_3-d-1} F^\downarrow \left(\lambda + 2k_3, \mathbf{k} - \underline{k}_{33}; f \det_{\mathbf{n}_2^+}^{-k_3}; z \right).
\end{aligned}$$

Then this is meromorphically continued for all $\lambda \in \mathbb{C}$, and if $\lambda = d + 1 - a$, $a = 1, 2, \dots, k_3$, then the right hand side of

$$\begin{aligned} \det_{\mathfrak{n}^+}(z)^{-a} F^\downarrow(d + 1 - a, \mathbf{k}; f; z) &= \frac{1}{(d_2 + 1 - a + \mathbf{k})_{k_{33}, d_2}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k_3} \det_{\mathfrak{n}^+}(z)^{2k_3 - a} \\ &\times F^\downarrow(d + 1 - a + 2k_3, \mathbf{k} - \underline{k}_{33}; f \det_{\mathfrak{n}_2^+}^{-k_3}; z) \end{aligned}$$

becomes a polynomial. Hence $F^\downarrow(d + 1 - a, \mathbf{k}; f; z)$ is divisible by $\det_{\mathfrak{n}^+}(z)^a$. Also, by $\mathcal{B}_{\frac{n}{r}-a} \det_{\mathfrak{n}^+}(z)^a = \det_{\mathfrak{n}^+}(z)^a \mathcal{B}_{\frac{n}{r}+a}$ (see the proof of [9, Proposition XV.2.4]), we have

$$\begin{aligned} &(\mathcal{B}_{d+1+a})_1 \det_{\mathfrak{n}^+}(z)^{-a} F^\downarrow(d + 1 - a, \mathbf{k}; f; z) \\ &= \det_{\mathfrak{n}^+}(z)^{-a} (\mathcal{B}_{d+1-a})_1 F^\downarrow(d + 1 - a, \mathbf{k}; f; z) = 0, \end{aligned}$$

and hence

$$\begin{aligned} &\left(f \det_{\mathfrak{n}_2^+}^{-a} \mapsto \det_{\mathfrak{n}^+}(z)^{-a} F^\downarrow(d + 1 - a, \mathbf{k}; f; z) \right) \\ &\in \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{\mathbf{k} - \underline{a}_3}(\mathfrak{p}_2^+), \text{Sol}_{\mathcal{P}(\mathfrak{p}^+)}((\mathcal{B}_{d+1+a})_1)) \\ &\simeq \mathbb{C} \left(f \det_{\mathfrak{n}_2^+}^{-a} \mapsto F^\downarrow(d + 1 + a, \mathbf{k} - \underline{a}_3; f \det_{\mathfrak{n}_2^+}^{-a}; z) \right) \end{aligned}$$

holds. Then comparing the values at $z_1 = 0$, for $a = 1, 2, \dots, k_3$ we get

$$F^\downarrow(d + 1 - a, \mathbf{k}; f; z) = \det_{\mathfrak{n}^+}(z)^a F^\downarrow(d + 1 + a, \mathbf{k} - \underline{a}_3; f \det_{\mathfrak{n}_2^+}^{-a}; z)$$

(see [35, Theorem 6.6] for more general case). We put $(k'_1, k'_2) := (k_1 - k_3, k_2 - k_3) \in \mathbb{Z}_{++}^2$, let $f'(x_2) := \det_{\mathfrak{n}_2^+}(x_2)^{-k_3} f(x_2) \in \mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^+)$, and rewrite the above as

$$\begin{aligned} &F^\downarrow(d + 1 - a, (k'_1, k'_2, 0) + \underline{k}_{33}; f' \det_{\mathfrak{n}_2^+}^{k_3}; z) \\ &= \det_{\mathfrak{n}^+}(z)^a F^\downarrow(d + 1 + a, (k'_1, k'_2, 0) + \underline{k}_3 - \underline{a}_3; f' \det_{\mathfrak{n}_2^+}^{k_3 - a}; z) \quad (3.6) \\ &\quad ((a, k_3) \in (\mathbb{Z}_{\geq 0})^2, a \leq k_3). \end{aligned}$$

Now we want to prove this for general $(a, k_3) \in \mathbb{C}^2$. We recall that the left hand side of (3.6) is given by

$$\begin{aligned} &F^\downarrow(d + 1 - a, (k'_1, k'_2, 0) + \underline{k}_{33}; f' \det_{\mathfrak{n}_2^+}^{k_3}; z) \\ &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k'_1 - k'_2 \\ 0 \leq l_3 \leq k'_2}} \frac{(-k_3 - (0, k'_2, k'_1))_{(l_1 + l_2, l_1 + l_3, l_2 + l_3), d_2}}{(a - 2k_3 - \frac{d_2}{2} - (k'_2, k'_1, k'_1 + k'_2))_{(l_1, l_2, l_3), d_2}} \det_{\mathfrak{n}_2^+}(z_2)^{k_3} F^\downarrow_{(k'_1, k'_2, 0), 1}[f'](z_1, z_2). \quad (3.7) \end{aligned}$$

Similarly, the right hand side is computed as

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(z)^a F^\downarrow(d + 1 + a, (k'_1, k'_2, 0) + \underline{k_3 - a}; f' \det_{\mathfrak{n}_2^+}^{k_3-a}; z) \\
&= \sum_{\substack{0 \leq l'_1 \\ 0 \leq l'_2 \leq k'_1 - k'_2 \\ 0 \leq l'_3 \leq k'_2}} \frac{(a - k_3 - (0, k'_2, k'_1))_{(l'_1 + l'_2, l'_1 + l'_3, l'_2 + l'_3), d_2}}{(a - 2k_3 - \frac{d_2}{2} - (k'_2, k'_1, k'_1 + k'_2))_{(l'_1, l'_2, l'_3), d_2}} \det_{\mathfrak{n}_2^+}(z_2)^{k_3-a} F^\downarrow_{(k'_1, k'_2, 0), \mathbf{l}'}[f'](z_1, z_2) \\
&\quad \times \det_{\mathfrak{n}_2^+}(z_2)^a \left(1 + \frac{((z_1)^\sharp | z_2)}{\det_{\mathfrak{n}_2^+}(z_2)}\right)^a \\
&= \sum_{\substack{0 \leq l'_1 \\ 0 \leq l'_2 \leq k'_1 - k'_2 \\ 0 \leq l'_3 \leq k'_2}} \frac{(a - k_3 - (0, k'_2, k'_1))_{(l'_1 + l'_2, l'_1 + l'_3, l'_2 + l'_3), d_2}}{(a - 2k_3 - \frac{d_2}{2} - (k'_2, k'_1, k'_1 + k'_2))_{(l'_1, l'_2, l'_3), d_2}} \det_{\mathfrak{n}_2^+}(z_2)^{k_3} F^\downarrow_{(k'_1, k'_2, 0), \mathbf{l}'}[f'](z_1, z_2) \\
&\quad \times \sum_{m=0}^{\infty} \frac{(-a)_m}{m!} \left(-\frac{((z_1)^\sharp | z_2)}{\det_{\mathfrak{n}_2^+}(z_2)}\right)^m.
\end{aligned}$$

Then we have

$$\begin{aligned}
& \left(f' \mapsto \frac{1}{m!} F^\downarrow_{(k'_1, k'_2, 0), \mathbf{l}'}[f'](z_1, z_2) \left(-\frac{((z_1)^\sharp | z_2)}{\det_{\mathfrak{n}_2^+}(z_2)}\right)^m \right) \\
& \in \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^+), \mathcal{P}_{2|\mathbf{l}'|+2m}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{(k'_1 - l'_2 - l'_3, k'_2 - l'_1 - l'_3, -l'_2 - l'_3)}(\mathfrak{p}_2^+) \otimes \mathcal{P}_{(0, -m, -m)}(\mathfrak{p}_2^+)) \\
& \simeq \bigoplus_{\substack{\mathbf{l}'' \in (\mathbb{Z}_{\geq 0})^3, \\ |\mathbf{l}''|=m}} \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^+), \mathcal{P}_{2|\mathbf{l}'|+2m}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{(k'_1 - l'_2 - l'_3 - l''_2 - l''_3, k'_2 - l'_1 - l'_3 - l''_1 - l''_3, -l'_2 - l'_3 - l''_2 - l''_3)}(\mathfrak{p}_2^+)) \\
& = \bigoplus_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^3, |\mathbf{l}|=|\mathbf{l}'|+m \\ l_j \geq l'_j}} \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^+), \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{(k'_1 - l_2 - l_3, k'_2 - l_1 - l_3, -l_2 - l_3)}(\mathfrak{p}_2^+)),
\end{aligned}$$

and by the Pieri rule,

$$\begin{aligned}
& \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^+), \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{(k'_1 - l_2 - l_3, k'_2 - l_1 - l_3, -l_2 - l_3)}(\mathfrak{p}_2^+)) \\
& \simeq \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(k'_1, k'_2, 0)}(\mathfrak{p}_2^+) \otimes \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^+)^{\vee}, \mathcal{P}_{(k'_1 - l_2 - l_3, k'_2 - l_1 - l_3, -l_2 - l_3)}(\mathfrak{p}_2^+)) \neq \{0\}
\end{aligned}$$

holds only if $0 \leq l_2 \leq k'_1 - k'_2$, $0 \leq l_3 \leq k'_2$. Therefore, for $|\mathbf{l}| = |\mathbf{l}'| + m$ we set

$$\begin{aligned}
& F_{(k'_1, k'_2, 0), \mathbf{l}', \mathbf{l}}[f'](z_1, z_2) \\
&:= \text{Proj}_{(k'_1 - l_2 - l_3, k'_2 - l_1 - l_3, -l_1 - l_2), z_2}^{\mathfrak{p}_2^+} \left(\frac{1}{m!} F^\downarrow_{(k'_1, k'_2, 0), \mathbf{l}'}[f'](z_1, z_2) \left(-\frac{((z_1)^\sharp | z_2)}{\det_{\mathfrak{n}_2^+}(z_2)}\right)^m \right) \\
& \in \mathcal{P}_{2|\mathbf{l}|}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{(k'_1 - l_2 - l_3, k'_2 - l_1 - l_3, -l_1 - l_2)}(\mathfrak{p}_2^+),
\end{aligned}$$

so that

$$\frac{1}{m!} F^\downarrow_{(k'_1, k'_2, 0), \mathbf{l}'}[f'](z_1, z_2) \left(-\frac{((z_1)^\sharp | z_2)}{\det_{\mathfrak{n}_2^+}(z_2)}\right)^m = \sum_{\substack{|\mathbf{l}|=|\mathbf{l}'|+m \\ l'_1 \leq l_1 \\ l'_2 \leq l_2 \leq k'_1 - k'_2 \\ l'_3 \leq l_3 \leq k'_2}} F_{(k'_1, k'_2, 0), \mathbf{l}', \mathbf{l}}[f'](z_1, z_2).$$

Then the right hand side of (3.6) becomes

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(z)^a F^\downarrow \left(d + 1 + a, (k'_1, k'_2, 0) + \underline{k_3 - a}; f' \det_{\mathfrak{n}_2^+}^{k_3-a}; z \right) \\
&= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k'_1 - k'_2 \\ 0 \leq l_3 \leq k'_2}} \sum_{\substack{0 \leq l'_1 \leq l_1 \\ 0 \leq l'_2 \leq l_2 \\ 0 \leq l'_3 \leq l_3}} \frac{(a - k_3 - (0, k'_2, k'_1))_{(l'_1+l'_2, l'_1+l'_3, l'_2+l'_3), d_2} (-a)_{|\mathbf{l}|-|\mathbf{l}'|}}{(a - 2k_3 - \frac{d_2}{2} - (k'_2, k'_1, k'_1 + k'_2))_{(l'_1, l'_2, l'_3), d_2}} \\
&\quad \times \det_{\mathfrak{n}_2^+}(z_2)^{k_3} F_{(k'_1, k'_2, 0), \mathbf{l}', \mathbf{l}}[f'](z_1, z_2). \tag{3.8}
\end{aligned}$$

If $a, k_3 \in \mathbb{Z}_{\geq 0}$, $a \leq k_3$, then in (3.7) and (3.8) only $l_1 + l_2 \leq k_3$ terms remain, and all denominators in the remaining terms are non-zero. Hence for $l_1 + l_2 \leq k_3$, by comparing both formulas we get

$$\begin{aligned}
& \frac{(-k_3 - (0, k'_2, k'_1))_{(l_1+l_2, l_1+l_3, l_2+l_3), d_2}}{(a - 2k_3 - \frac{d_2}{2} - (k'_2, k'_1, k'_1 + k'_2))_{(l_1, l_2, l_3), d_2}} F_{(k'_1, k'_2, 0), \mathbf{l}}[f'](z_1, z_2) \\
&= \sum_{0 \leq l'_j \leq l_j} \frac{(a - k_3 - (0, k'_2, k'_1))_{(l'_1+l'_2, l'_1+l'_3, l'_2+l'_3), d_2} (-a)_{|\mathbf{l}|-|\mathbf{l}'|}}{(a - 2k_3 - \frac{d_2}{2} - (k'_2, k'_1, k'_1 + k'_2))_{(l'_1, l'_2, l'_3), d_2}} F_{(k'_1, k'_2, 0), \mathbf{l}', \mathbf{l}}[f'](z_1, z_2).
\end{aligned}$$

Then since $\{(a, k_3) \in (\mathbb{Z}_{\geq 0})^2 \mid k_3 \geq l_1 + l_2, k_3 \geq a\} \subset \mathbb{C}^2$ is Zariski dense, this holds for all $(a, k_3) \in \mathbb{C}^2$ without the assumption $l_1 + l_2 \leq k_3$, and hence (3.6) holds for all $(a, k_3) \in \mathbb{C}^2$. This completes the proof of Theorem 3.5 (3). \square

Example 3.8. Let $\mathbf{k} = (k, k, k)$ with $k \in \mathbb{C}$ and $f(x_2) = \det_{\mathfrak{n}_2^+}(x_2)^k$. Then by Proposition 3.2 (2), for $x = (x_1, x_2) = (\xi, X) \in \mathfrak{p}^+$ we have

$$\begin{aligned}
F^\downarrow \left(\lambda, \underline{k_3}; \det_{\mathfrak{n}_2^+}^k; x \right) &= \sum_{l=0}^{\infty} \frac{(-k)_{(l, l, 0), d_2}}{(-\lambda - 2k + \frac{3}{2}d_2 + 1)_{(l, 0, 0), d_2}} F_{(k, k, k), (l, 0, 0)}[\det_{\mathfrak{n}_2^+}^k](x_1, x_2) \\
&= \sum_{l=0}^{\infty} \frac{(-k)_l (-k - \frac{d_2}{2})_l}{(-\lambda - 2k + \frac{3}{2}d_2 + 1)_l l!} \left(-\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right)^l \det_{\mathfrak{n}_2^+}(x_2)^k \\
&= {}_2F_1 \left(\begin{matrix} -k, -k - \frac{d_2}{2} \\ -\lambda - 2k + \frac{3}{2}d_2 + 1 \end{matrix}; \frac{\xi X^t \hat{\xi}}{\det(X)} \right) \det(X)^k.
\end{aligned}$$

Also, for $k \in \mathbb{Z}_{\geq 0}$ we have

$$C^{d, d_2}(\lambda, \underline{k_3}) = \frac{(\lambda - \frac{d_2}{2} + (2k, 2k, k))_{(0, 0, k), d_2}}{(\lambda)_{(2k, k, k), d}} = \frac{(\lambda + k - \frac{3}{2}d_2)_k}{(\lambda)_{(2k, k, k), d}}.$$

Combining these we get

$$\begin{aligned}
& \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k, e^{(x|\bar{x})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\
&= \frac{(\lambda + k - \frac{3}{2}d_2)_k}{(\lambda)_{(2k, k, k), d}} {}_2F_1 \left(\begin{matrix} -k, -k - \frac{d_2}{2} \\ -\lambda - 2k + \frac{3}{2}d_2 + 1 \end{matrix}; -\frac{((z_1)^\sharp | z_2)}{\det_{\mathfrak{n}_2^+}(z_2)} \right) \det_{\mathfrak{n}_2^+}(z_2)^k.
\end{aligned}$$

In addition, Theorem 3.5 (3) is equivalent to

$$\begin{aligned}
& {}_2F_1 \left(\begin{matrix} -k, -k - \frac{1}{4}d \\ -\lambda - 2k + \frac{3}{4}d + 1 \end{matrix}; -\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right) \\
&= \left(1 + \frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right)^{-\lambda+d+1} {}_2F_1 \left(\begin{matrix} -\lambda - k + d + 1, -\lambda - k + \frac{3}{4}d + 1 \\ -\lambda - 2k + \frac{3}{4}d + 1 \end{matrix}; -\frac{((x_1)^\sharp | x_2)}{\det_{\mathfrak{n}_2^+}(x_2)} \right).
\end{aligned}$$

Example 3.9. Let $\mathbf{k} = (k_1, k_2, 0) \in \mathbb{Z}_{++}^3$, let $e' \in \mathfrak{n}_2^+$ be a primitive idempotent, and let $\mathfrak{p}^+(e')_0 =: \mathfrak{p}^{+''}$, $\mathfrak{p}_2^+(e')_0 =: \mathfrak{p}_2^{+''}$ be as in (2.2). We take $f(x_2) = f(x_2'') \in \mathcal{P}_{(k_1, k_2)}(\mathfrak{p}_2^{+''})$. Then by Proposition 3.2 (3), for $\xi = (\xi, X) \in \mathfrak{p}^+$ we have

$$\begin{aligned} & F^\downarrow(\lambda, (k_1, k_2, 0); f; x) \\ &= \sum_{l=0}^{k_2} \frac{((0, -k_2, -k_1))_{(0, l, l), d_2}}{(-\lambda + \frac{3}{2}d_2 + 1 - (k_2, k_1, k_1 + k_2))_{(0, 0, l), d_2}} F^\downarrow_{(k_1, k_2, 0), (0, 0, l)}[f](x_1, x_2) \\ &= \sum_{l=0}^{k_2} \frac{(-k_2 - \frac{d_2}{2})_l (-k_1 - d_2)_l}{(-\lambda - k_1 - k_2 + \frac{d_2}{2} + 1)_l} \frac{(-k_2)_l (-k_1 - \frac{d_2}{2})_l}{(-k_2 - \frac{d_2}{2})_l (-k_1 - d_2)_l l!} \left(-\frac{\det_{\mathfrak{n}^{+''}}(x_1'')}{\det_{\mathfrak{n}^{+''}}(x_2'')} \right)^l f(x_2'') \\ &= \sum_{l=0}^{k_2} \frac{(-k_2)_l (-k_1 - \frac{d_2}{2})_l}{(-\lambda - k_1 - k_2 + \frac{d_2}{2} + 1)_l l!} \left(-\frac{\det_{\mathfrak{n}^{+''}}(x_1'')}{\det_{\mathfrak{n}^{+''}}(x_2'')} \right)^l f(x_2'') \\ &= {}_2F_1 \left(\begin{matrix} -k_2, -k_1 - \frac{d_2}{2} \\ -\lambda - k_1 - k_2 + \frac{d_2}{2} + 1 \end{matrix}; \frac{(\hat{t}\xi\xi|e')}{\det_{\mathfrak{n}_2^{+''}}(X'')} \right)^l f(X''). \end{aligned}$$

We also have

$$C^{d, d_2}(\lambda, (k_1, k_2, 0)) = \frac{(\lambda - \frac{d_2}{2} + (k_1, k_1, k_2))_{(k_2, 0, 0), d_2}}{(\lambda)_{(k_1+k_2, k_2, 0), d}} = \frac{(\lambda + k_1 - \frac{d_2}{2})_{k_2}}{(\lambda)_{(k_1+k_2, k_2), d}}.$$

Combining these we get

$$\begin{aligned} & \left\langle f(x_2''), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \frac{(\lambda + k_1 - \frac{d_2}{2})_{k_2}}{(\lambda)_{(k_1+k_2, k_2), d}} {}_2F_1 \left(\begin{matrix} -k_2, -k_1 - \frac{d_2}{2} \\ -\lambda - k_1 - k_2 + \frac{d_2}{2} + 1 \end{matrix}; -\frac{\det_{\mathfrak{n}^{+''}}(z_1'')}{\det_{\mathfrak{n}^{+''}}(z_2'')} \right)^l f(z_2''). \end{aligned}$$

Example 3.10. Let $\mathbf{k} = (k_1, k_2, k_2)$ with $k_1 - k_2 \in \mathbb{Z}_{\geq 0}$, $k_2 \in \mathbb{C}$, let $w_2 \in \mathfrak{p}_2^+$ be of rank 1, and let $f(x_2) := (x_2|w_2)_{\mathfrak{n}_2^+}^{k_1-k_2} \det_{\mathfrak{n}_2^+}(x_2)^{k_2}$. Then we have

$$\begin{aligned} & F^\downarrow(\lambda, (k_1, k_2, k_2); f; x) \\ &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2}} \frac{((-k_2, -k_2, -k_1))_{(l_1+l_2, l_1, l_2), d_2}}{(-\lambda + \frac{3}{2}d_2 + 1 - (2k_2, k_1 + k_2, k_1 + k_2))_{(l_1, l_2, 0), d_2}} F^\downarrow_{(k_1, k_2, k_2), (l_1, l_2, 0)}[f](x_1, x_2) \\ &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_2 \leq k_1 - k_2}} \frac{(-k_2)_{l_1+l_2} (-k_2 - \frac{d_2}{2})_{l_1} (-k_1 - d_2)_{l_2}}{(-\lambda - 2k_2 + \frac{3}{2}d_2 + 1)_{l_1} (-\lambda - k_1 - k_2 + d_2 + 1)_{l_2}} \\ &\quad \times F^\downarrow_{(k_1 - k_2, 0, 0), (l_1, l_2, 0)}[(\cdot|w_2)_{\mathfrak{n}_2^+}^{k_1-k_2}](x_1, x_2) \det_{\mathfrak{n}_2^+}(x_2)^{k_2}, \end{aligned}$$

and $F^\downarrow_{(k_1 - k_2, 0, 0), (l_1, l_2, 0)}[(\cdot|w_2)_{\mathfrak{n}_2^+}^{k_1-k_2}](x_1, x_2)$ is given in Proposition 3.2 (4). Also, when $k_2 \in \mathbb{Z}_{\geq 0}$ we have

$$\begin{aligned} C^{d, d_2}(\lambda, (k_1, k_2, k_2)) &= \frac{(\lambda - \frac{d_2}{2} + (k_1 + k_2, \max\{k_1, 2k_2\}, k_2))_{(0, \min\{k_2, k_1 - k_2\}, k_2), d_2}}{(\lambda)_{(k_1+k_2, \min\{k_1, 2k_2\}, k_2), d}} \\ &= \frac{(\lambda + \max\{k_1, 2k_2\} - d_2)_{\min\{k_2, k_1 - k_2\}} (\lambda + k_2 - \frac{3}{2}d_2)_{k_2}}{(\lambda)_{(k_1+k_2, \min\{k_1, 2k_2\}, k_2), d}}, \end{aligned}$$

and $\langle (x_2|w_2)_{\mathfrak{n}_2^+}^{k_1-k_2} \det_{\mathfrak{n}_2^+}(x_2)^{k_2}, e^{(x|\bar{x})_{\mathfrak{p}^+}} \rangle_{\lambda,x}$ is given by the product of the above two formulas.

Example 3.11. Let $\mathbf{k} = (k_2, k_2, k_3)$ with $k_2 - k_3 \in \mathbb{Z}_{\geq 0}$, $k_2 \in \mathbb{C}$, let $w_2 \in \mathfrak{p}_2^+$ be of rank 1, and let $f(x_2) := (x_2^{-1}|w_2)_{\mathfrak{n}_2^+}^{k_2-k_3} \det_{\mathfrak{n}_2^+}(x_2)^{k_2}$. Then we have

$$\begin{aligned} & F^\downarrow(\lambda, (k_2, k_2, k_3); f; x) \\ &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_3 \leq k_2 - k_3}} \frac{((-k_3, -k_2, -k_2))_{(l_1, l_1 + l_3, l_3), d_2}}{(-\lambda + \frac{3}{2}d_2 + 1 - (k_2 + k_3, k_2 + k_3, 2k_2))_{(l_1, 0, l_3), d_2}} F^\downarrow_{(k_2, k_2, k_3), (l_1, 0, l_3)}[f](x_1, x_2) \\ &= \sum_{\substack{0 \leq l_1 \\ 0 \leq l_3 \leq k_2 - k_3}} \frac{(-k_3)_{l_1} (-k_2 - \frac{d_2}{2})_{l_1 + l_3} (-k_2 - d_2)_{l_3}}{(-\lambda - k_2 - k_3 + \frac{3}{2}d_2 + 1)_{l_1} (-\lambda - 2k_2 + \frac{d_2}{2} + 1)_{l_3}} \\ &\quad \times F^\downarrow_{(0, 0, k_3 - k_2), (l_1, 0, l_3)}[((\cdot)^{-1}|w_2)_{\mathfrak{n}_2^+}^{k_2 - k_3}](x_1, x_2) \det_{\mathfrak{n}_2^+}(x_2)^{k_2}, \end{aligned}$$

and $F^\downarrow_{(0, 0, k_3 - k_2), (l_1, 0, l_3)}[((\cdot)^{-1}|w_2)_{\mathfrak{n}_2^+}^{k_2 - k_3}](x_1, x_2)$ is given in Proposition 3.2(5). Also, when $k_3 \in \mathbb{Z}_{\geq 0}$ we have

$$\begin{aligned} C^{d, d_2}(\lambda, (k_2, k_2, k_3)) &= \frac{(\lambda - \frac{d_2}{2} + (k_2 + k_3, k_2 + k_3, k_2))_{(k_2 - k_3, 0, k_3), d_2}}{(\lambda)_{(2k_2, k_2, k_3), d}} \\ &= \frac{(\lambda + k_2 + k_3 - \frac{d_2}{2})_{k_2 - k_3} (\lambda + k_2 - \frac{3}{2}d_2)_{k_3}}{(\lambda)_{(2k_2, k_2, k_3), d}}, \end{aligned}$$

and $\langle (x_2^{-1}|w_2)_{\mathfrak{n}_2^+}^{k_2 - k_3} \det_{\mathfrak{n}_2^+}(x_2)^{k_2}, e^{(x|\bar{x})_{\mathfrak{p}^+}} \rangle_{\lambda,x}$ is given by the product of the above two formulas.

3.4 Results on restriction of $\mathcal{H}_\lambda(D)$ to subgroups

In this subsection we consider the decomposition of $\mathcal{H}_\lambda(D)$ under \tilde{G}_1 . Let $V_{\mathbf{k}}$ be an irreducible K_1 -module isomorphic to $\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$. Then $\mathcal{H}_\lambda(D)$ is decomposed under \tilde{G}_1 as

$$\mathcal{H}_\lambda(D)|_{\tilde{G}_1} \simeq \sum_{\mathbf{k} \in \mathbb{Z}_{++}^3}^{\oplus} \mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}}).$$

For each case, $\mathcal{H}_\lambda(D)$ has the minimal \tilde{K} -type

$$\chi^{-\lambda} \simeq \begin{cases} V_{\frac{\lambda}{2}, 3}^{(3)\vee} \boxtimes V_{\underline{\lambda}, 3}^{(3)} \simeq V_{\underline{\lambda}, 3}^{(3)\vee} \boxtimes V_{\underline{0}, 3}^{(3)} \simeq V_{\underline{0}, 3}^{(3)\vee} \boxtimes V_{\underline{\lambda}, 3}^{(3)} & (G = SU(3, 3)), \\ V_{\frac{\lambda}{2}, 6}^{(6)\vee} & (G = SO^*(12)), \\ \chi^{-\lambda} & (G = E_{7(-25)}), \end{cases}$$

and $\mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}})$ has the minimal \tilde{K}_1 -type

$$\chi_1^{-\varepsilon_1 \lambda} \otimes V_{\mathbf{k}} \simeq \begin{cases} V_{\underline{\lambda}_3 + (2k_1, 2k_2, 2k_3)}^{(3)\vee} & (G_1 = SO^*(6)), \\ V_{\underline{\lambda}, 3}^{(3)\vee} \boxtimes V_{\underline{\lambda}/2, 3}^{(3)\vee} & (G_1 = SO^*(6) \times SO^*(6)), \\ V_{\frac{3}{4}\lambda + \frac{1}{2}|\mathbf{k}|, 2}^{(2)\vee} \boxtimes V_{\frac{1}{4}\lambda + \frac{1}{2}|\mathbf{k}|, 6}^{(6)} & (G_1 = SU(2, 6)). \\ \simeq V_{\underline{0}, 2}^{(2)\vee} \boxtimes V_{\underline{\Delta}_6 + (k_1 + k_2, k_1 + k_2, k_1 + k_3, k_1 + k_3, k_2 + k_3, k_2 + k_3)}^{(6)} & \end{cases}$$

To describe the intertwining operators, for $\lambda \in \mathbb{C}$, $\mathbf{k} \in \mathbb{Z}_{++}^3 + \mathbb{C}\underline{1}_3$, $f(x_2) \in \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, we define a function $F^\uparrow(\lambda, \mathbf{k}; f; y_1, x_2)$ on $\mathfrak{p}_1^- \oplus \mathfrak{p}_2^+$ by

$$\begin{aligned}
& F^\uparrow(\lambda, \mathbf{k}; f; y_1, x_2) \\
&:= \sum_{\substack{\mathbf{l} \in \mathbb{Z}^3 \\ 0 \leq l_1 \leq k_1 - k_2 \\ 0 \leq l_2 \leq k_2 - k_3 \\ 0 \leq l_3}} \frac{1}{(\lambda - \frac{d_2-2}{2} + (k_1 + k_2, k_1 + k_3, k_2 + k_3))_{(l_3, l_2, l_1), d_2}} F_{\mathbf{k}, \mathbf{l}}^\uparrow[f](y_1, x_2) \\
&= \sum_{\substack{\mathbf{l} \in \mathbb{Z}^3 \\ 0 \leq l_1 \leq k_1 - k_2 \\ 0 \leq l_2 \leq k_2 - k_3 \\ 0 \leq l_3}} \frac{F_{\mathbf{k}, \mathbf{l}}^\uparrow[f](y_1, x_2)}{(\lambda + k_2 + k_3 - \frac{3}{4}d + 1)_{l_1} (\lambda + k_1 + k_3 - \frac{1}{2}d + 1)_{l_2} (\lambda + k_1 + k_2 - \frac{1}{4}d + 1)_{l_3}}. \tag{3.9}
\end{aligned}$$

Theorem 3.12. Let $\operatorname{Re} \lambda > 1 + 2d$, $\mathbf{k} \in \mathbb{Z}_{++}^3$.

(1) ([33, Theorems 5.7(5), 5.12]). Let $K_{\mathbf{k}}(x_2) \in (\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+) \otimes \operatorname{Hom}(V_{\mathbf{k}}, \mathbb{C}))^{K_1}$. Then the linear map

$$\begin{aligned}
\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow: \mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}})|_{\tilde{K}_1} &\longrightarrow \mathcal{H}_\lambda(D)|_{\tilde{K}}|_{(\mathfrak{g}_1, \tilde{K}_1)}, \\
(\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow f)(x) &:= F^\uparrow\left(\lambda, \mathbf{k}; K_{\mathbf{k}}; \frac{1}{\varepsilon_1} \frac{\partial}{\partial x_1}, x_2\right) f(x_1)
\end{aligned}$$

intertwines the $(\mathfrak{g}_1, \tilde{K}_1)$ -action, where F^\uparrow is as in (3.9).

(2) Let $K_{\mathbf{k}}^\vee(y_2) \in (\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^-) \otimes V_{\mathbf{k}})^{K_1}$. Then the linear map

$$\begin{aligned}
\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow: \mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\longrightarrow \mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}}), \\
(\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow f)(x_1) &:= F^\downarrow\left(\lambda, \mathbf{k}; K_{\mathbf{k}}^\vee; \frac{\partial}{\partial x}\right) f(x)\Big|_{x_2=0}
\end{aligned}$$

intertwines the \tilde{G}_1 -action, where F^\downarrow is as in (3.2).

(3) ([35, Corollary 6.7]). Suppose $c_{\mathbf{k}}, c_{\mathbf{k}}^\vee \in \mathbb{R}_{>0}$ satisfy, for $v \in V_{\mathbf{k}}$,

$$\|K_{\mathbf{k}}(x_2)v\|_{F, x_2}^2 = c_{\mathbf{k}}|v|_{V_{\mathbf{k}}}^2, \quad \|(v, K_{\mathbf{k}}^\vee(\overline{x_2}))\|_{F, x_2}^2 = c_{\mathbf{k}}^\vee|v|_{V_{\mathbf{k}}}^2.$$

Then the operator norms of $\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow$, $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ are given by

$$c_{\mathbf{k}}^{-1} \|\mathcal{F}_{\lambda, \mathbf{k}}^\uparrow\|_{\text{op}}^2 = c_{\mathbf{k}}^\vee \|\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow\|_{\text{op}}^{-2} = C^{d, d_2}(\lambda, \mathbf{k}),$$

where $C^{d, d_2}(\lambda, \mathbf{k})$ is as in (3.3).

Remark 3.13. In [33, Theorem 5.12], the subscripts $(m_3, m_2, m_1), 2$ are wrong, and $(m_3, m_2, m_1), 1$ (in (1)), $(m_3, m_2, m_1), 4$ (in (2)) are correct.

The differential operators $\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow$ are meromorphically continued for all $\lambda \in \mathbb{C}$, and give \tilde{G}_1 -intertwining operators

$$\mathcal{F}_{\lambda, \mathbf{k}}^\downarrow: \mathcal{O}_\lambda(D)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}})$$

for all $\lambda \in \mathbb{C}$ except for poles. Then by Theorem 3.5 (3), for special λ the following holds.

Theorem 3.14 ([35, Corollary 6.8(4)]). *Let $\mathbf{k} \in \mathbb{Z}_{++}^3$. For $a = 1, 2, \dots, k_3$, we fix a $[K_1, K_1]$ -isomorphism $V_{\mathbf{k}} \simeq V_{\mathbf{k}-\underline{a}_3}$, and assume that*

$$K_{\mathbf{k}}^\vee(y_2) = \det_{\mathfrak{n}^-}(y_2)^a K_{\mathbf{k}-\underline{a}_3}^\vee(y_2) \in (\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^-) \otimes V_{\mathbf{k}})^{K_1}$$

holds. Then we have

$$\begin{aligned} \mathcal{F}_{d+1-a, \mathbf{k}}^\downarrow &= \mathcal{F}_{d+1+a, \mathbf{k}-\underline{a}_3}^\downarrow \circ \det_{\mathfrak{n}^-} \left(\frac{\partial}{\partial x} \right)^a : \\ \mathcal{O}_{d+1-a}(D)|_{\tilde{G}_1} &\longrightarrow \mathcal{O}_{\varepsilon_1(d+1-a)}(D_1, V_{\mathbf{k}}) \simeq \mathcal{O}_{\varepsilon_1(d+1+a)}(D_1, V_{\mathbf{k}-\underline{a}_3}). \end{aligned}$$

For $\lambda > d$ we have

$$\mathcal{H}_{\varepsilon_1\lambda}(D_1, V_{\mathbf{k}})_{\tilde{K}_1} \simeq d\tau_\lambda(\mathcal{U}(\mathfrak{g}_1))\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+) \subset \mathcal{H}_\lambda(D)_{\tilde{K}},$$

but this does not hold for smaller λ in general. For such λ , by Theorem 3.5(2) the following holds. For $i = 1, 2, 3$, $\lambda \in \frac{d}{2}(i-1) - \mathbb{Z}_{\geq 0}$, let $M_i^{\mathfrak{g}}(\lambda) \subset \mathcal{O}_\lambda(D)_{\tilde{K}}$ be the $(\mathfrak{g}, \tilde{K})$ -submodule given in (2.18).

Theorem 3.15. *Let $\mathbf{k} \in \mathbb{Z}_{++}^3$. For $i = 1, 2, 3$,*

$$d\tau_\lambda(\mathcal{U}(\mathfrak{g}_1))\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+) \subset M_i^{\mathfrak{g}}(\lambda)$$

holds if and only if

$$\lambda \in \begin{cases} -(k_1 + k_2) - \mathbb{Z}_{\geq 0} & (i = 1), \\ \frac{d}{2} - \min\{k_1, k_2 + k_3\} - \mathbb{Z}_{\geq 0} & (i = 2), \\ d - k_3 - \mathbb{Z}_{\geq 0} & (i = 3). \end{cases}$$

Here, “if” follows from the holomorphy in Theorem 3.5(2) and this is already given in [35, Corollary 6.8(1)]. On the other hand, “only if” follows from the non-zeroness in Theorem 3.5(2) and this is new except for $k_1 = k_2 = k_3$ or $k_3 = 0$.

4 General rank case

In this section we treat $\mathfrak{p}^\pm = (\mathfrak{p}^\pm)^\sigma \oplus (\mathfrak{p}^\pm)^{-\sigma} = \mathfrak{p}_1^\pm \oplus \mathfrak{p}_2^\pm$ such that $\mathfrak{p}^\pm, \mathfrak{p}_2^\pm$ are both simple, classical, of tube type, and $\text{rank } \mathfrak{p}^\pm = \text{rank } \mathfrak{p}_2^\pm$, that is, we treat

$$(\mathfrak{p}^\pm, \mathfrak{p}_1^\pm, \mathfrak{p}_2^\pm) = \begin{cases} (\mathbb{C}^n, \mathbb{C}^{n'}, \mathbb{C}^{n''}) & (n = n' + n'', n'' \geq 3) \quad (\text{Case 1}), \\ (M(r, \mathbb{C}), \text{Alt}(r, \mathbb{C}), \text{Sym}(r, \mathbb{C})) & (\text{Case 2}), \\ (\text{Alt}(2r, \mathbb{C}), \text{Alt}(r, \mathbb{C}) \oplus \text{Alt}(r, \mathbb{C}), M(r, \mathbb{C})) & (\text{Case 3}). \end{cases}$$

Then the corresponding symmetric pairs $(G, (G^\sigma)_0, (G^{\sigma\vartheta})_0) = (G, G_1, G_2)$ are given by

$$(G, G_1, G_2) = \begin{cases} (SO_0(2, n), SO_0(2, n') \times SO(n''), SO_0(2, n'') \times SO(n')) & (\text{Case 1}), \\ (SU(r, r), SO^*(2r), Sp(r, \mathbb{R})) & (\text{Case 2}), \\ (SO^*(4r), SO^*(2r) \times SO^*(2r), U(r, r)) & (\text{Case 3}) \end{cases}$$

(up to covering). $(r, d, d_2, \varepsilon_1)$ are given as follows, and we define δ by the following.

$$(r, d, d_2, \varepsilon_1, \delta) = \begin{cases} (2, n - 2, n'' - 2, 1, n' - 2) & (\text{Case 1, } n' \geq 2), \\ (2, n - 2, n'' - 2, 2, -1) & (\text{Case 1, } n' = 1), \\ (r, 2, 1, 2, -1) & (\text{Case 2}), \\ (r, 4, 2, 1, 0) & (\text{Case 3}). \end{cases} \quad (4.1)$$

Especially $d = 2d_2$ holds if $r \geq 3$. We fix a maximal tripotent $e \in \mathfrak{p}_2^+ \subset \mathfrak{p}^+$, regard $\mathfrak{p}_2^+ \subset \mathfrak{p}^+$ as Jordan algebras with the unit element e , the Euclidean real forms $\mathfrak{n}_2^+ \subset \mathfrak{n}^+$, the symmetric cones $\Omega_2 \subset \Omega$, and identify $\mathfrak{p}^+ \simeq \mathfrak{p}^-$, $\mathfrak{p}_2^+ \simeq \mathfrak{p}_2^-$ via $Q(e), Q(\bar{e})$.

4.1 Definition of some functions

In this subsection we define some functions on $\mathfrak{p}_1^\pm \oplus \mathfrak{p}_2^\pm$. First, for $\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}$, let $\mathbf{m}^2 := (m_1, m_1, m_2, m_2, \dots, m_{[r/2]}, m_{[r/2]}(, 0)) \in \mathbb{Z}_{++}^r$, let

$$\mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^\pm) := \begin{cases} \mathcal{P}_{(m, m)}(\mathbb{C}^{n'}) & (\text{Case 1, } n' \geq 3), \\ \mathcal{P}_m(\mathbb{C}) \otimes \mathcal{P}_m(\mathbb{C}) & (\text{Case 1, } n' = 2), \\ \mathcal{P}_{2m}(\mathbb{C}) & (\text{Case 1, } n' = 1), \\ \mathcal{P}_{2\mathbf{m}}(\text{Alt}(r, \mathbb{C})) & (\text{Case 2}), \\ \mathcal{P}_{\mathbf{m}}(\text{Alt}(r, \mathbb{C})) \otimes \mathcal{P}_{\mathbf{m}}(\text{Alt}(r, \mathbb{C})) & (\text{Case 3}) \end{cases}$$

(where $\mathbb{C}^2 = \mathbb{C} \oplus \mathbb{C}$ corresponds to the local isomorphism $SO_0(2, 2) \simeq SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$), and $\Phi_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2), \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) \in \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{\mathbf{m}^2}(\mathfrak{p}_2^\mp)_{y_2}$ be the polynomials satisfying

$$e^{\frac{1}{2}(x_1|P(y_2)x_1)_{\mathfrak{n}^\pm}} = \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}} \frac{d_{\mathbf{m}}^{\tilde{\mathfrak{p}}}}{(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1)_{\mathbf{m}, d}} \Phi_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) = \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2),$$

where

$$\tilde{\mathfrak{p}} := \begin{cases} \mathbb{C} & (\text{Case 1}), \\ M(\lfloor \frac{r}{2} \rfloor, \mathbb{C}) & (\text{Case 2}), \\ \text{Alt}(2\lfloor \frac{r}{2} \rfloor, \mathbb{C}) & (\text{Case 3}), \end{cases} \quad d_{\mathbf{m}}^{\tilde{\mathfrak{p}}} := \dim \mathcal{P}_{\mathbf{m}}(\tilde{\mathfrak{p}}) = \begin{cases} 1 & (\text{Case 1}), \\ \dim(V_{\mathbf{m}}^{(\lfloor r/2 \rfloor)^\vee})^{\otimes 2} & (\text{Case 2}), \\ \dim V_{\mathbf{m}^2}^{(2\lfloor r/2 \rfloor)^\vee} & (\text{Case 3}). \end{cases}$$

Then for $\mu \in \mathbb{C}$ we have

$$\begin{aligned} h_{\mathfrak{n}^\pm}(x_1, P(y_2)x_1)^{-\mu/2} &= \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}} \frac{(\mu)_{\mathbf{m}, d} d_{\mathbf{m}}^{\tilde{\mathfrak{p}}}}{(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1)_{\mathbf{m}, d}} \Phi_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) \\ &= \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}} (\mu)_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2), \\ \det_{\mathfrak{n}^\pm}(x_1 + x_2)^{-\mu} &= \det_{\mathfrak{n}^\pm}(x_2)^{-\mu} h_{\mathfrak{n}^\pm}(x_1, P(x_2^{-1})x_1)^{-\mu/2} \\ &= \det_{\mathfrak{n}^\pm}(x_2)^{-\mu} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}} \frac{(\mu)_{\mathbf{m}, d} d_{\mathbf{m}}^{\tilde{\mathfrak{p}}}}{(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1)_{\mathbf{m}, d}} \Phi_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, x_2^{-1}) \\ &= \det_{\mathfrak{n}^\pm}(x_2)^{-\mu} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]}} (\mu)_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, x_2^{-1}) \end{aligned} \quad (4.2)$$

(for details see [34, Propositions 2.4, 6.2]).

Example 4.1. Let $\mu = -1$. Then for $x = x_1 + x_2 \in \mathfrak{p}^+$ we have

$$\begin{aligned}\det_{\mathfrak{n}^+}(x_1 + x_2) &= \det_{\mathfrak{n}_2^+}(x_2) h_{\mathfrak{n}^+}(x_1, P(x_2^{-1})x_1)^{1/2} \\ &= \det_{\mathfrak{n}_2^+}(x_2) \sum_{j=0}^{\lfloor r/2 \rfloor} (-1)^j \binom{\lfloor r/2 \rfloor}{j} \Phi_{(\underline{1}_j, \underline{0}_{\lfloor r/2 \rfloor-j})}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, x_2^{-1}),\end{aligned}$$

where $\underline{k}_j := (\overbrace{k, \dots, k}^j)$, and comparing the homogeneous terms of degree $2\lfloor \frac{r}{2} \rfloor$ with respect to x_1 , we have

$$(-1)^{\lfloor r/2 \rfloor} \det_{\mathfrak{n}_2^+}(x_2) \Phi_{\underline{1}_{\lfloor r/2 \rfloor}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, x_2^{-1}) = \begin{cases} \det_{\mathfrak{n}^+}(x_1) & (r: \text{even}), \\ ((x_1)^\sharp | x_2)_{\mathfrak{n}^+} & (r: \text{odd}), \end{cases}$$

where $x \mapsto x^\sharp$ is the adjugate in $\mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}}$. That is, for $x_1 \in \mathfrak{p}_1^+$, $y_2 \in \mathfrak{p}_2^-$, we have

$$\Phi_{\underline{1}_{\lfloor r/2 \rfloor}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) = \begin{cases} (-1)^{r/2} \det_{\mathfrak{n}^+}(x_1) \det_{\mathfrak{n}_2^-}(y_2) & (r: \text{even}), \\ (-1)^{\lfloor r/2 \rfloor} ((x_1)^\sharp | (y_2)^\sharp)_{\mathfrak{n}^+} & (r: \text{odd}). \end{cases}$$

More generally, for $k \in \mathbb{Z}_{\geq 0}$ we have

$$\Phi_{\underline{k}_{\lfloor r/2 \rfloor}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) = \Phi_{\underline{1}_{\lfloor r/2 \rfloor}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2)^k = \begin{cases} (-1)^{kr/2} \det_{\mathfrak{n}^+}(x_1)^k \det_{\mathfrak{n}_2^-}(y_2)^k & (r: \text{even}), \\ (-1)^{k\lfloor r/2 \rfloor} ((x_1)^\sharp | (y_2)^\sharp)_{\mathfrak{n}^+}^k & (r: \text{odd}). \end{cases}$$

Next, for $\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}$, $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil}$, we define $\phi_\pm(\mathbf{m}, \mathbf{l}) \in \mathbb{Z}^r$ by

$$\begin{aligned}\phi_+(\mathbf{m}, \mathbf{l}) &:= \begin{cases} (m_1 + l_1, m_1, m_2 + l_2, m_2, \dots, m_{r/2} + l_{r/2}, m_{r/2}) & (r: \text{even}), \\ (m_1 + l_1, m_1, m_2 + l_2, m_2, \dots, m_{\lfloor r/2 \rfloor} + l_{\lfloor r/2 \rfloor}, m_{\lfloor r/2 \rfloor}, l_{\lceil r/2 \rceil}) & (r: \text{odd}), \end{cases} \\ \phi_-(\mathbf{m}, \mathbf{l}) &:= \begin{cases} (m_1, m_1 - l_1, m_2, m_2 - l_2, \dots, m_{r/2}, m_{r/2} - l_{r/2}) & (r: \text{even}), \\ (m_1, m_1 - l_1, m_2, m_2 - l_2, \dots, m_{\lfloor r/2 \rfloor}, m_{\lfloor r/2 \rfloor} - l_{\lfloor r/2 \rfloor}, -l_{\lceil r/2 \rceil}) & (r: \text{odd}). \end{cases}\end{aligned}$$

Then for $l \in \mathbb{Z}_{\geq 0}$, $f(y_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^\mp)$, we have

$$\begin{aligned}\tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) f(y_2) &\in \bigoplus_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{\phi_+(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^\mp)_{y_2}, \\ \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) f(y_2^{-1}) &\in \bigoplus_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{\phi_-(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^\mp)_{y_2},\end{aligned}$$

where $\mathcal{P}_{\phi_+(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^\mp)$, $\mathcal{P}_{\phi_-(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^\mp) = \{0\}$ if $\phi_+(\mathbf{m}, \mathbf{l})$, $\phi_-(\mathbf{m}, \mathbf{l}) \notin \mathbb{Z}_+^r := \{\mathbf{k} \in \mathbb{Z}^r \mid k_1 \geq \dots \geq k_r\}$. According to these decompositions, for $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil}$ with $|\mathbf{l}| = l$, we define the polynomials $F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2)$ and $F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, y_2)$ by

$$\begin{aligned}F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2) &:= \text{Proj}_{\phi_+(\mathbf{m}, \mathbf{l}), y_2}^{\mathfrak{p}_2^\mp} \left(\tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) f(y_2) \right) \in \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{\phi_+(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^\mp)_{y_2}, \\ F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, y_2) &:= \text{Proj}_{\phi_-(\mathbf{m}, \mathbf{l}), y_2}^{\mathfrak{p}_2^\mp} \left(\tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^\pm, \mathfrak{p}_2^\mp}(x_1, y_2) f(y_2^{-1}) \right) \in \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{\phi_-(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^\mp)_{y_2}.\end{aligned}$$

Then $F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2) \neq 0$ holds only if

$$\begin{aligned} 0 \leq l_1, \quad 0 \leq l_j \leq m_{j-1} - m_j \quad (1 < j \leq \frac{r}{2}) & \quad (r: \text{even}), \\ 0 \leq l_1, \quad 0 \leq l_j \leq m_{j-1} - m_j \quad (1 < j \leq \lfloor \frac{r}{2} \rfloor), \quad 0 \leq l_{\lceil r/2 \rceil} \leq m_{\lfloor r/2 \rfloor}, & \quad (r: \text{odd}), \end{aligned} \quad (4.3)$$

and $F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, y_2) \neq 0$ holds only if

$$\begin{aligned} 0 \leq l_j \leq m_j - m_{j+1} \quad (1 \leq j < \frac{r}{2}), \quad 0 \leq l_{r/2} & \quad (r: \text{even}), \\ 0 \leq l_j \leq m_j - m_{j+1} \quad (1 \leq j < \lfloor \frac{r}{2} \rfloor), \quad 0 \leq l_{\lfloor r/2 \rfloor} \leq m_{\lfloor r/2 \rfloor}, \quad 0 \leq l_{\lceil r/2 \rceil} & \quad (r: \text{odd}). \end{aligned}$$

If $r = 3$, then these polynomials and $F_{\mathbf{k}, \mathbf{l}}^\uparrow[f], F_{\mathbf{k}, \mathbf{l}}^\downarrow[f]$ in the previous section are related as

$$\begin{aligned} F_{m, (l_1, l_2)}^+[f](x_1, y_2) &= F_{(l_1 + l_2, 0, 0), (l_2, 0, m - l_2)}^\uparrow [f](x_1, y_2) = F_{(0, 0, -l_1 - l_2), (m - l_2, 0, l_2)}^\downarrow [f((\cdot)^{-1})](x_1, y_2^{-1}) \\ &\in \mathcal{P}_{\langle m \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{(m + l_1, m, l_2)}(\mathfrak{p}_2^\mp)_{y_2}, \\ F_{m, (l_1, l_2)}^-[f](x_1, y_2) &= F_{(0, 0, -l_1 - l_2), (0, l_1, m - l_1)}^\uparrow [f((\cdot)^{-1})](x_1, y_2) = F_{(l_1 + l_2, 0, 0), (m - l_1, l_1, 0)}^\downarrow [f](x_1, y_2^{-1}) \\ &\in \mathcal{P}_{\langle m \rangle}(\mathfrak{p}_1^\pm)_{x_1} \otimes \mathcal{P}_{(m, m - l_1, -l_2)}(\mathfrak{p}_2^\mp)_{y_2}. \end{aligned}$$

For general r , $F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, y_2)$ satisfies the following. For $\mathbf{k} \in \mathbb{Z}_+^r$, let $d_{\mathbf{k}}^{\mathfrak{p}_2^+} := \dim \mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$, and let $\Sigma_2 \subset \mathfrak{p}_2^+$ be the Bergman–Shilov boundary of $D_2 := D \cap \mathfrak{p}_2^+$. For $x \in \mathfrak{p}^+$, let $|x|_\infty := \inf\{t > 0 \mid t^{-1}x \in D\}$ be the spectral norm of x .

Proposition 4.2. *For $\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}$, $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil}$, $|\mathbf{l}| = l$, $f(y_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^-)$ and for $x_1 \in \mathfrak{p}_1^+$, $y_2 = ge \in \mathfrak{p}_2^-$ with $g \in K_1^{\mathbb{C}}$, we have*

$$\begin{aligned} |F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2)| &\leq \frac{\sqrt{d_{\mathbf{m}}^{\mathfrak{p}_2^+} d_{(l, \underline{0}_{r-1})}^{\mathfrak{p}_2^+}} d_{\mathbf{m}}^{\tilde{\mathfrak{p}}}}{\left(\frac{d}{2}\left(\lfloor \frac{r}{2} \rfloor - 1\right) + 1\right)_{\mathbf{m}, d}} (|{}^t g x_1|_\infty)^{2|\mathbf{m}|} \|f(g(\cdot))\|_{L^2(\Sigma_2)}, \\ |F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, y_2)| &\leq \frac{\sqrt{d_{\mathbf{m}}^{\mathfrak{p}_2^+} d_{(l, \underline{0}_{r-1})}^{\mathfrak{p}_2^+}} d_{\mathbf{m}}^{\tilde{\mathfrak{p}}}}{\left(\frac{d}{2}\left(\lfloor \frac{r}{2} \rfloor - 1\right) + 1\right)_{\mathbf{m}, d}} (|{}^t g x_1|_\infty)^{2|\mathbf{m}|} \|f({}^t g^{-1}(\cdot))\|_{L^2(\Sigma_2)}. \end{aligned}$$

Proof. Since $f(x_2) \mapsto F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2)$, $f(x_2^{-1}) \mapsto F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, y_2)$ are $K_1^{\mathbb{C}}$ -equivariant, we have

$$\begin{aligned} F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2) &= F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, ge) = F_{\mathbf{m}, \mathbf{l}}^+[f(g(\cdot))]({}^t g x_1, e), \\ F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, y_2) &= F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, ge) = F_{\mathbf{m}, \mathbf{l}}^-[f({}^t g^{-1}(\cdot))]({}^t g x_1, e), \end{aligned}$$

and hence it suffices to prove when $y_2 = e$. Next, since the reproducing kernel of $\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^-)$ with respect to the $L^2(\Sigma_2)$ -inner product is given by $K'_{\mathbf{k}}(y_2, z_2) := d_{\mathbf{k}}^{\mathfrak{p}_2^+} \Phi_{\mathbf{k}}^{\mathfrak{n}_2^-}(P(\overline{z_2}^{1/2})y_2) = d_{\mathbf{k}}^{\mathfrak{p}_2^+} \Phi_{\mathbf{k}}^{\mathfrak{n}_2^-}(P(y_2^{1/2})\overline{z_2})$, where $\Phi_{\mathbf{k}}^{\mathfrak{n}_2^-}(x_2) := \int_{K_1 \cap K_L} \Delta_{\mathbf{k}}^{\mathfrak{n}_2^-}(kx_2) dk$ (see [9, Proposition XII.2.4]), we have

$$\begin{aligned} |F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, e)| &= |\langle F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, \cdot), K'_{\phi_\pm(\mathbf{m}, \mathbf{l})}(\cdot, e) \rangle_{L^2(\Sigma_2)}| \\ &\leq \|F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, \cdot)\|_{L^2(\Sigma_2)} \|K'_{\phi_\pm(\mathbf{m}, \mathbf{l})}(\cdot, e)\|_{L^2(\Sigma_2)} \\ &= \|F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, \cdot)\|_{L^2(\Sigma_2)} \sqrt{K'_{\phi_\pm(\mathbf{m}, \mathbf{l})}(e, e)} = \sqrt{d_{\phi_\pm(\mathbf{m}, \mathbf{l})}^{\mathfrak{p}_2^+}} \|F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, \cdot)\|_{L^2(\Sigma_2)}. \end{aligned}$$

Now since $F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, y_2)$ is defined by the orthogonal projection of $\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x_1, y_2)f(y_2)$ or $\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^-, \mathbf{p}_2^+}(x_1, y_2)f(y_2^{-1})$, we have

$$\begin{aligned} \|F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, \cdot)\|_{L^2(\Sigma_2)} &\leq \|\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x_1, y_2)f(y_2)\|_{L^2(\Sigma_2), y_2} \leq \|\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x_1, \cdot)\|_{L^\infty(\Sigma_2)} \|f\|_{L^2(\Sigma_2)}, \\ \|F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, \cdot)\|_{L^2(\Sigma_2)} &\leq \|\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^-, \mathbf{p}_2^+}(x_1, y_2)f(y_2^{-1})\|_{L^2(\Sigma_2), x_2} \leq \|\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^-, \mathbf{p}_2^+}(x_1, \cdot)\|_{L^\infty(\Sigma_2)} \|f\|_{L^2(\Sigma_2)}. \end{aligned}$$

In addition, let $\Sigma_1 \subset \mathbf{p}_1^+$ be the Bergman–Shilov boundary of $D_1 := D \cap \mathbf{p}_1^+$. Then we have

$$\|\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x_1, \cdot)\|_{L^\infty(\Sigma_2)} \leq \sup_{\substack{|x'_1|_\infty \leq |x_1|_\infty \\ y_2 \in \Sigma_2}} |\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x'_1, y_2)| = (|x_1|_\infty)^{2|\mathbf{m}|} \sup_{\substack{x'_1 \in \Sigma_1 \\ y_2 \in \Sigma_2}} |\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x'_1, y_2)|.$$

Also, since $\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x'_1, y_2)$ is defined by using the roots of $t^{\lfloor r/2 \rfloor} h_{\mathbf{n}^+}(t^{-1}x'_1, P(y_2)x'_1)^{1/2}$, and these roots have absolute values 1 when $x'_1 \in \Sigma_1$, $y_2 \in \Sigma_2$, by [9, Theorem XII.1.1] we have

$$|\tilde{\Phi}_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x'_1, y_2)| = \frac{d_{\mathbf{m}}^{\tilde{\mathbf{m}}} |\Phi_{\mathbf{m}}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(x'_1, y_2)|}{\left(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1\right)_{\mathbf{m}, d}} \leq \frac{d_{\mathbf{m}}^{\tilde{\mathbf{m}}}}{\left(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1\right)_{\mathbf{m}, d}} \quad (x'_1 \in \Sigma_1, y_2 \in \Sigma_2).$$

We also have

$$d_{\phi_\pm(\mathbf{m}, \mathbf{l})}^{\mathbf{p}_2^\pm} = \dim \mathcal{P}_{\phi_\pm(\mathbf{m}, \mathbf{l})}(\mathbf{p}_2^\pm) \leq \dim (\mathcal{P}_{\mathbf{m}^2}(\mathbf{p}_2^+) \otimes \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathbf{p}_2^\pm)) = d_{\mathbf{m}^2}^{\mathbf{p}_2^+} d_{(l, \underline{0}_{r-1})}^{\mathbf{p}_2^+}.$$

Combining these, we get the desired formulas. \square

The following proposition gives some examples of $F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, y_2)$.

Proposition 4.3. *Let $\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}$, $\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil}$, $|\mathbf{l}| = l$.*

(1) Suppose r is even. For $k \in \mathbb{Z}_{\geq 0}$, $f(y_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathbf{p}_2^-)$, we have

$$F_{\mathbf{m} + \underline{k}_{r/2}, \mathbf{l}}^\pm[f](x_1, y_2) = \frac{1}{(-k - \mathbf{m}^\vee)_{\underline{k}_{r/2}, d}} F_{\mathbf{m}, \mathbf{l}}^\pm[f](x_1, y_2) \det_{\mathbf{n}^+}(x_1)^k \det_{\mathbf{n}_2^-}(y_2)^k,$$

where $\mathbf{m}^\vee = (m_{\lfloor r/2 \rfloor}, \dots, m_1)$.

(2) Suppose r is odd, and let $w_2 \in \mathbf{p}_2^+$ be of rank 1. Then for $k, l, m \in \mathbb{Z}_{\geq 0}$ with $m \leq k, l$ we have

$$\begin{aligned} &F_{\underline{k}_{\lfloor r/2 \rfloor}, (l-m, \underline{0}_{\lfloor r/2 \rfloor-1}, m)}^+[(\cdot | w_2)_{\mathbf{n}_2^+}^l](x_1, y_2) \\ &= \frac{(-1)^m \det_{\mathbf{n}_2^+}(y_2)^k}{(-k)_{\underline{k}_{\lfloor r/2 \rfloor}, d} (-k - l + m - \frac{d_2}{2}(r-1))_m m!} \\ &\times \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j (y_2^{-1} | (x_1)^\sharp)_{\mathbf{n}_2^+}^{k-j} (y_2 | w_2)_{\mathbf{n}_2^+}^{l-j} ((x_1)^\sharp | w_2)_{\mathbf{n}_2^+}^j}{(-k - l + 2m - \frac{d_2}{2}(r-1) + 1)_{j-m} (j-m)!}. \end{aligned}$$

(3) Suppose r is odd, and let $w_2 \in \mathfrak{p}_2^+$ be of rank 1. Then for $k, l, m \in \mathbb{Z}_{\geq 0}$ with $m \leq k, l$, we have

$$\begin{aligned} & F_{\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor - 1}, m, l-m)}^- [(\cdot | w_2)_m^l] (x_1, y_2) \\ &= \frac{(-1)^m \det_{\mathfrak{n}_2^+}(y_2)^k}{(-k)_{\underline{k}_{\lfloor r/2 \rfloor}, d} (-k-l+m-\frac{d_2}{2})_m m!} \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j}{(-k-l+2m-\frac{d_2}{2}+1)_{j-m} (j-m)!} \\ & \quad \times (y_2^{-1} | (x_1)^\sharp)_m^{k-j} (y_2^{-1} | w_2)_m^{l-j} ((y_2^{-1} | (x_1)^\sharp)_m^{k-j} (y_2^{-1} | w_2)_m^{l-j} - (P(y_2)^{-1} (x_1)^\sharp | w_2)_m^{k-j}). \end{aligned}$$

(4) Suppose $r \geq 3$. Let $\mathbf{m} \in \mathbb{Z}_{++}^{[r/2]-1}$, $l \in \mathbb{Z}_{\geq 0}$, $m_1 \leq l$, and put $m_{r/2} := 0$ for even r . We define $\mathbf{l}(\mathbf{m}) \in (\mathbb{Z}_{\geq 0})^{[r/2]}$ by

$$\mathbf{l}(\mathbf{m}) := (l - m_1, m_1 - m_2, m_2 - m_3, \dots, m_{\lceil r/2 \rceil - 2} - m_{\lceil r/2 \rceil - 1}, m_{\lceil r/2 \rceil - 1}),$$

so that

$$\begin{aligned} \phi_+(\mathbf{m}, \mathbf{l}(\mathbf{m})) &= \begin{cases} (l, m_1, m_1, m_2, m_2, \dots, m_{r/2-1}, m_{r/2-1}, 0) & (r: \text{even}), \\ (l, m_1, m_1, m_2, m_2, \dots, m_{\lceil r/2 \rceil - 1}, m_{\lceil r/2 \rceil - 1}) & (r: \text{odd}) \end{cases} \\ &= (l, \mathbf{m}^2) \in \mathbb{Z}_{++}^r. \end{aligned}$$

Let $e' \in \mathfrak{n}_2^+$ be a primitive idempotent, let $\mathfrak{p}^+(e')_0 =: \mathfrak{p}^{+''}$ be as in (2.2), let $\mathfrak{p}_1^+ \cap \mathfrak{p}^{+''} =: \mathfrak{p}_1^{+''}$, $\mathfrak{p}_2^+ \cap \mathfrak{p}^{+''} =: \mathfrak{p}_2^{+''}$, and for $x \in \mathfrak{p}^+$ let $x'' \in \mathfrak{p}^{+''}$ be the orthogonal projection. Then we have

$$\det_{\mathfrak{n}_2^+}(x_2)^l F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+ [(\cdot | e')_m^l] (x_1, x_2^{-1}) = \frac{(-l)_{\mathbf{m}, d}}{(-l - \frac{d}{2})_{\mathbf{m}, d}} \det_{\mathfrak{n}_2^{+''}}(x_2'')^l \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^{+''}, \mathfrak{p}_2^{-''}}(x_1'', (x_2'')^{-1}).$$

Proof. (1) Follows from

$$\begin{aligned} \tilde{\Phi}_{\mathbf{m} + \underline{k}_{r/2}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) &= \frac{d^{\tilde{\mathbf{p}}}_{\mathbf{m} + \underline{k}_{r/2}}}{(\frac{d}{2}(\frac{r}{2} - 1) + 1)_{\mathbf{m} + \underline{k}_{r/2}, d}} \Phi_{\mathbf{m} + \underline{k}_{r/2}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) \\ &= \frac{d^{\tilde{\mathbf{p}}}_{\mathbf{m}}}{(\frac{d}{2}(\frac{r}{2} - 1) + 1)_{\mathbf{m} + \underline{k}_{r/2}, d}} \Phi_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) \Phi_{\underline{k}_{r/2}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2)^k \\ &= \frac{(\frac{d}{2}(\frac{r}{2} - 1) + 1)_{\mathbf{m}, d}}{(\frac{d}{2}(\frac{r}{2} - 1) + 1)_{\mathbf{m} + \underline{k}_{r/2}, d}} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) \Phi_{\underline{k}_{r/2}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2)^k \\ &= \frac{(-1)^{kr/2}}{(-k - \mathbf{m}^\vee)_{\underline{k}_{r/2}, d}} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) \Phi_{\underline{k}_{r/2}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2)^k \\ &= \frac{1}{(-k - \mathbf{m}^\vee)_{\underline{k}_{r/2}, d}} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1, y_2) \det_{\mathfrak{n}_2^+}(x_1)^k \det_{\mathfrak{n}_2^-}(y_2)^k. \end{aligned}$$

(2) Since r is odd, $x_1 \in \mathfrak{p}_1^+$ is not invertible, and $(x_1)^\sharp \in \mathfrak{p}_2^+$ is at most of rank 1. By

Example 4.1 and Proposition 2.5 (1) we have

$$\begin{aligned}
& F_{k_{\lfloor r/2 \rfloor}, (l-m, 0_{\lfloor r/2 \rfloor-1}, m)}^+ [(\cdot | w_2)_n^l] (x_1, y_2) = \text{Proj}_{(k+l-m, k_{r-2}, m), y_2}^{\mathfrak{p}_2^-} (\tilde{\Phi}_{k_{\lfloor r/2 \rfloor}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-} (x_1, y_2) (y_2 | w_2)_n^l) \\
&= \frac{(-1)^{k \lfloor r/2 \rfloor}}{\left(\frac{d}{2} \left(\lfloor \frac{r}{2} \rfloor - 1\right) + 1\right)_{k_{\lfloor r/2 \rfloor}, d}} \det_{n_2^+}(y_2)^k \text{Proj}_{(l-m, 0_{r-2}, m-k), y_2}^{\mathfrak{p}_2^-} ((y_2^{-1} | (x_1)^\sharp)_n^k (y_2 | w_2)_n^l) \\
&= \frac{\det_{n_2^+}(y_2)^k}{(-k)_{k_{\lfloor r/2 \rfloor}, d}} \frac{(-1)^m}{(-k-l+m-\frac{d_2}{2}(r-1))_m m!} \\
&\times \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j (y_2^{-1} | (x_1)^\sharp)_n^{k-j} (y_2 | w_2)_n^{l-j} ((x_1)^\sharp | w_2)_n^j}{(-k-l+2m-\frac{d_2}{2}(r-1)+1)_{j-m} (j-m)!}.
\end{aligned}$$

(3) By Example 4.1 and Proposition 2.5 (2) we have

$$\begin{aligned}
& F_{k_{\lfloor r/2 \rfloor}, (0_{\lfloor r/2 \rfloor-1}, m, l-m)}^- [(\cdot | w_2)_n^l] (x_1, y_2) = \text{Proj}_{(k_{r-2}, k-m, m-l), y_2}^{\mathfrak{p}_2^-} (\tilde{\Phi}_{k_{\lfloor r/2 \rfloor}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-} (x_1, y_2) (y_2^{-1} | w_2)_n^l) \\
&= \frac{(-1)^{k \lfloor r/2 \rfloor}}{\left(\frac{d}{2} \left(\lfloor \frac{r}{2} \rfloor - 1\right) + 1\right)_{k_{\lfloor r/2 \rfloor}, d}} \det_{n_2^+}(y_2)^k \text{Proj}_{(0_{r-2}, -m, m-k-l), y_2}^{\mathfrak{p}_2^-} ((y_2^{-1} | (x_1)^\sharp)_n^k (y_2^{-1} | w_2)_n^l) \\
&= \frac{\det_{n_2^+}(y_2)^k}{(-k)_{k_{\lfloor r/2 \rfloor}, d}} \frac{(-1)^m}{(-k-l+m-\frac{d_2}{2})_m m!} \sum_{j=m}^{\min\{k, l\}} \frac{(-k)_j (-l)_j}{(-k-l+2m-\frac{d_2}{2}+1)_{j-m} (j-m)!} \\
&\times (y_2^{-1} | (x_1)^\sharp)_n^{k-j} (y_2^{-1} | w_2)_n^{l-j} ((y_2^{-1} | (x_1)^\sharp)_n^k (y_2^{-1} | w_2)_n^l - (P(y_2)^{-1} (x_1)^\sharp | w_2)_n^l)^j.
\end{aligned}$$

(4) Let $\mathfrak{p}_2^+(e')_2 = \mathbb{C}e' =: \mathfrak{p}_2^{+'}$, and let $P_{K_1} \subset K_1^\mathbb{C}$ be a maximal parabolic subgroup stabilizing $\mathcal{P}(\mathfrak{p}_1^{++}) \subset \mathcal{P}(\mathfrak{p}_1^+)$ and $\mathcal{P}(\mathfrak{p}_2^{++}) \subset \mathcal{P}(\mathfrak{p}_2^+)$. First, we have

$$\begin{aligned}
& (\det_{n_2^+}(x_2)^l f(x_2^{-1})) \mapsto \det_{n_2^+}(x_2)^l F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+ [f](x_1, x_2^{-1}) \\
& \in \text{Hom}_{K_1}(\mathcal{P}_{(l_{r-1}, 0)}(\mathfrak{p}_2^+), \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{l_r - \phi_+(\mathbf{m}, \mathbf{l}(\mathbf{m}))^\vee}(\mathfrak{p}_2^+)) \\
& \simeq \text{Hom}_{K_1}(\mathcal{P}_{(l, 0_{r-1})}(\mathfrak{p}_2^-) \otimes \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^-), \mathcal{P}_{(l, \mathbf{m}^2)}(\mathfrak{p}_2^-)),
\end{aligned}$$

and this space is 1-dimensional by the Pieri rule for Cases 2, 3. Especially, since $\det_{n_2^+}(x_2)^l (x_2^{-1} | e')_n^l = \det_{n_2^{++}}(x_2'')^l \in \mathcal{P}_{(l_{r-1}, 0)}(\mathfrak{p}_2^+)$ is relatively invariant under the parabolic subgroup P_{K_1} , $\det_{n_2^+}(x_2)^l F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+ [(\cdot | e')_n^l] (x_1, x_2^{-1}) \in \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{l_r - (l, \mathbf{m}^2)^\vee}(\mathfrak{p}_2^+)$ is also relatively invariant under P_{K_1} , with the weight same with $\det_{n_2^+}(x_2)^l (x_2^{-1} | e')_n^l = \det_{n_2^{++}}(x_2'')^l$, and such function is unique up to constant multiple. Hence

$$\det_{n_2^+}(x_2)^l F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+ [(\cdot | e')_n^l] (x_1, x_2^{-1}) = c \det_{n_2^{++}}(x_2'')^l \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^{++}, \mathfrak{p}_2^{--}}(x_1'', (x_2'')^{-1})$$

holds for some $c \in \mathbb{C}$. To determine c , it is enough to consider the case $x_1 = x_1'' \in \mathfrak{p}^{++}$. Then since $\tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(x_1'', y_2) = \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^{++}, \mathfrak{p}_2^{--}}(x_1'', y_2'')$ holds, by Lemma 2.4 (4) we have

$$\begin{aligned}
F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+ [(\cdot | e')_n^l] (x_1'', x_2^{-1})|_{x_2=x_2'+x_2''} &= \text{Proj}_{(l, \mathbf{m}^2)}^{\mathfrak{p}_2^-} ((y_2 | e')_n^l \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^{++}, \mathfrak{p}_2^{--}}(x_1'', y_2''))|_{y_2^{-1}=x_2'+x_2''} \\
&= \frac{(-l)_{\mathbf{m}^2, d_2}}{(-l - \frac{d_2}{2})_{\mathbf{m}^2, d_2}} ((x_2')^{-1} | e')_n^l \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^{++}, \mathfrak{p}_2^{--}}(x_1'', (x_2'')^{-1}),
\end{aligned}$$

and hence we get

$$c = \frac{(-l)_{\mathbf{m}^2, d_2}}{(-l - \frac{d_2}{2})_{\mathbf{m}^2, d_2}} = \frac{(-l)_{\mathbf{m}, d} (-l - \frac{d_2}{2})_{\mathbf{m}, d}}{(-l - \frac{d_2}{2})_{\mathbf{m}, d} (-l - \frac{d}{2})_{\mathbf{m}, d}} = \frac{(-l)_{\mathbf{m}, d}}{(-l - \frac{d}{2})_{\mathbf{m}, d}}. \quad \square$$

Using $F_{\mathbf{m}, 1}^\pm[f](x_1, y_2)$, for $l \in \mathbb{Z}_{\geq 0}$, $\mu, \nu \in \mathbb{C}$ and for $f(x_2) \in \mathcal{P}_{(l, 0, \dots, 0)}(\mathfrak{p}_2^\pm)$, we define a function $F_{l,+}^\downarrow(\frac{\nu}{\mu}; f; x)$ by

$$F_{l,+}^\downarrow\left(\frac{\nu}{\mu}; f; x\right) := \det_{\mathfrak{n}_2^\pm}(x_2)^{-\nu} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} F_{\mathbf{m}, 1}^-[f](x_1, x_2^{-1}) \\ \times \begin{cases} \frac{(\nu)_{\mathbf{m}, d} (\nu - \frac{d_2}{2} - (\underline{0}_{r/2-1}, l))_{\mathbf{m}-\mathbf{l}+(\underline{0}_{r/2-1}, l), d}}{(\mu - \frac{d_2}{2} - (\underline{0}_{r/2-1}, l))_{\mathbf{m}-\mathbf{l}+(\underline{0}_{r/2-1}, l), d}} & (r: \text{even}), \\ \frac{(\nu)_{\mathbf{m}, d} (\nu - \frac{d_2}{2})_{\mathbf{m}-\mathbf{l}', d} (\nu - \frac{d_2}{2}(r-1) - l)_{l-l_{\lceil r/2 \rceil}}}{(\mu - \frac{d_2}{2})_{\mathbf{m}-\mathbf{l}', d} (\mu - \frac{d_2}{2}(r-1) - l)_{l-l_{\lceil r/2 \rceil}}} & (r: \text{odd}), \end{cases} \quad (4.4)$$

where, when r is odd, let $\mathbf{l}' := (l_1, \dots, l_{\lfloor r/2 \rfloor})$ so that $\mathbf{l} = (\mathbf{l}', l_{\lceil r/2 \rceil})$ holds, and we define a function $F_{l,-}^\downarrow(\frac{\nu}{\mu}; f; x)$ by

$$F_{l,-}^\downarrow\left(\frac{\nu}{\mu}; f; x\right) := \det_{\mathfrak{n}_2^\pm}(x_2)^{-\nu} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} F_{\mathbf{m}, 1}^+[f](x_1, x_2^{-1}) \\ \times \frac{(\nu + (l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{0}_{\lceil r/2 \rceil-1}), d} (\nu - \frac{d_2}{2})_{\mathbf{m}, d}}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{0}_{\lceil r/2 \rceil-1}), d}}, \quad (4.5)$$

where we identify $\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}$ and $(\mathbf{m}, 0) \in \mathbb{Z}_{++}^{\lceil r/2 \rceil}$ when r is odd. Then as a function of μ , $F_{l,+}^\downarrow(\frac{\nu}{\mu}; f; x)$ has poles at

$$\mu \in \bigcup_{i=1}^{r/2-1} \left\{ \frac{d_2}{2} + \frac{d}{2}(i-1) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq 0 \end{array} \right\} \cup \left\{ \frac{d_2}{2} + \frac{d}{2}\left(\frac{r}{2} - 1\right) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq -l \end{array} \right\} \quad (r: \text{even}), \\ \mu \in \bigcup_{i=1}^{\lfloor r/2 \rfloor} \left\{ \frac{d_2}{2} + \frac{d}{2}(i-1) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq 0 \end{array} \right\} \cup \left\{ \frac{d_2}{2} + \frac{d}{2}\left(\frac{r}{2} - 1\right) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ -l \leq j < 0 \end{array} \right\} \quad (r: \text{odd})$$

(we note that $d = 2d_2$ holds if r is odd), and $F_{l,-}^\downarrow(\frac{\nu}{\mu}; f; x)$ has poles at

$$\mu \in \left\{ \frac{d_2}{2} - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq l \end{array} \right\} \cup \bigcup_{i=2}^{r/2} \left\{ \frac{d_2}{2} + \frac{d}{2}(i-1) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq 0 \end{array} \right\} \quad (r: \text{even}), \\ \mu \in \left\{ \frac{d_2}{2} - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq l \end{array} \right\} \cup \bigcup_{i=2}^{\lfloor r/2 \rfloor} \left\{ \frac{d_2}{2} + \frac{d}{2}(i-1) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ j \geq 0 \end{array} \right\} \\ \cup \left\{ \frac{d_2}{2} + \frac{d}{2}\left(\left\lceil \frac{r}{2} \right\rceil - 1\right) - j \mid \begin{array}{l} j \in \mathbb{Z}, \\ 0 \leq j < l \end{array} \right\} \quad (r: \text{odd})$$

for generic $\nu \in \mathbb{C}$. On the other hand, if $\nu = -k \in -\mathbb{Z}_{\geq 0}$ then $F_{l,+}^\downarrow(\underline{\mu}; f; x)$ becomes a finite sum and is a polynomial in x , and if $\nu = -k \in -\mathbb{Z}_{\geq l}$ then $F_{l,-}^\downarrow(\underline{\mu}; f; x)$ also becomes a polynomial. For these cases, all poles satisfying $\mu \leq \frac{d_2}{2} - k$ become removable, and we use the same notations $F_{l,+}^\downarrow(\underline{\mu}; f; x)$, $F_{l,-}^\downarrow(\underline{\mu}; f; x)$ to express these polynomials also for μ at such removable singularities. If $r = 3$, then since $d = 2d_2$ holds, $F_{l,\pm}^\downarrow(\underline{\mu}; f; x)$ and $F^\downarrow(\lambda, \mathbf{k}; f; x)$ in the previous section are related as

$$F^\downarrow\left(\lambda, (k_1, k_2, k_2); f \det_{\mathfrak{n}_2^+}^{k_2}; x\right) = F_{k_1-k_2,+}^\downarrow\left(\begin{matrix} -k_2 \\ -\lambda - 2k_2 + d + 1 \end{matrix}; f; x\right),$$

$$F^\downarrow\left(\lambda, (k_2, k_2, k_3); f((\cdot)^{-1}) \det_{\mathfrak{n}_2^+}^{k_2}; x\right) = F_{k_2-k_3,-}^\downarrow\left(\begin{matrix} -k_2 \\ -\lambda - 2k_2 + d + 1 \end{matrix}; f; x\right).$$

Proposition 4.4. Suppose $\mu \in \mathbb{C}$ is not a pole of $F_{l,\pm}^\downarrow(\underline{\mu}; f; x)$. Then $F_{l,\pm}^\downarrow(\underline{\mu}; f; x)$ converges if $x_2 = ge$ for some $g \in K_1^{\mathbb{C}}$ and $|g^{-1}x_1|_\infty < 1$ holds. Especially, this converges for $x = (x_1, x_2) \in \Omega$.

Proof. By Proposition 4.2 we have

$$|F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, x_2^{-1})| \leq \frac{\sqrt{d_{\mathbf{m}^2}^{\mathfrak{p}_2^+} d_{(l, \underline{0}_{r-1})}^{\mathfrak{p}_2^+}} d_{\tilde{\mathbf{m}}}^{\tilde{\mathfrak{p}}}}{\left(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1\right)_{\mathbf{m}, d}} (|g^{-1}x_1|_\infty)^{2|\mathbf{m}|} \|f(g(\cdot))\|_{L^2(\Sigma_2)},$$

$$|F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, x_2^{-1})| \leq \frac{\sqrt{d_{\mathbf{m}^2}^{\mathfrak{p}_2^+} d_{(l, \underline{0}_{r-1})}^{\mathfrak{p}_2^+}} d_{\tilde{\mathbf{m}}}^{\tilde{\mathfrak{p}}}}{\left(\frac{d}{2}(\lfloor \frac{r}{2} \rfloor - 1) + 1\right)_{\mathbf{m}, d}} (|g^{-1}x_1|_\infty)^{2|\mathbf{m}|} \|f(tg^{-1}(\cdot))\|_{L^2(\Sigma_2)},$$

and we have

$$d_{\mathbf{m}^2}^{\mathfrak{p}_2^+} = \dim \mathcal{P}_{\mathbf{m}^2}(\mathfrak{p}_2^+) \leq \dim \mathcal{P}_{2|\mathbf{m}|}(\mathfrak{p}_2^+) = \binom{2|\mathbf{m}| + n_2 - 1}{n_2 - 1},$$

$$d_{\tilde{\mathbf{m}}}^{\tilde{\mathfrak{p}}} = \dim \mathcal{P}_{\mathbf{m}}(\tilde{\mathfrak{p}}) \leq \dim \mathcal{P}_{|\mathbf{m}|}(\tilde{\mathfrak{p}}) = \binom{|\mathbf{m}| + \tilde{n} - 1}{\tilde{n} - 1},$$

where $n_2 := \dim \mathfrak{p}_2^+$, $\tilde{n} := \dim \tilde{\mathfrak{p}}$. Then the convergence for $|g^{-1}x_1|_\infty < 1$ follows from standard argument. Especially when $x = (x_1, x_2) \in \Omega$, we have

$$0 < \det_{\mathfrak{n}^+}(x) = \det_{\mathfrak{n}_2^+}(x_2) h_{\mathfrak{n}^+}(x_1, P(x_2)^{-1}x_1) = \det_{\mathfrak{n}_2^+}(x_2) h_{\mathfrak{n}^+}(P(x_2^{-1/2})x_1, P(x_2^{-1/2})x_1),$$

and hence $P(x_2^{-1/2})x_1 \in D \cap \Omega$, that is, $|P(x_2^{-1/2})x_1|_\infty < 1$ holds. Thus $F_{l,\pm}^\downarrow(\underline{\mu}; f; x)$ converges for $x \in \Omega$. \square

These $F_{l,\pm}^\downarrow(\underline{\mu}; f; x)$ has the following property.

Proposition 4.5. For $\mu, \nu \in \mathbb{C}$, $k, l \in \mathbb{Z}_{\geq 0}$, $f(x_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^\pm)$, we have

$$\begin{aligned} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial x_2} \right)^k F_{l,+}^\downarrow \left(\frac{\nu}{\mu}; f; x \right) &= (-1)^{kr} (\nu - (\underline{0}_{r-1}, l))_{\underline{k}_r, d_2} F_{l,+}^\downarrow \left(\frac{\nu + k}{\mu}; f; x \right) \\ &= (-1)^{kr} (\nu)_{\underline{k}_{\lfloor r/2 \rfloor}, d} \left(\nu - \frac{d_2}{2} \right)_{\underline{k}_{\lceil r/2 \rceil - 1}, d} \left(\nu - \frac{d_2}{2}(r-1) - l \right)_k F_{l,+}^\downarrow \left(\frac{\nu + k}{\mu}; f; x \right), \\ \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial x_2} \right)^k F_{l,-}^\downarrow \left(\frac{\nu}{\mu}; f; x \right) &= (-1)^{kr} (\nu + (l, \underline{0}_{r-1}))_{\underline{k}_r, d_2} F_{l,-}^\downarrow \left(\frac{\nu + k}{\mu}; f; x \right) \\ &= (-1)^{kr} (\nu + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\underline{k}_{\lceil r/2 \rceil}, d} \left(\nu - \frac{d_2}{2} \right)_{\underline{k}_{\lfloor r/2 \rfloor}, d} F_{l,-}^\downarrow \left(\frac{\nu + k}{\mu}; f; x \right). \end{aligned}$$

Proof. First, the numerator of each term of $F_{l,\pm}^\downarrow \left(\frac{\nu}{\mu}; f; x \right)$ satisfies

$$\begin{aligned} &(\nu - (\underline{0}_{r-1}, l))_{\phi_-(\mathbf{m}, \mathbf{l}) + (\underline{0}_{r-1}, l), d_2} \\ &= \begin{cases} (\nu)_{\mathbf{m}, d} \left(\nu - \frac{d_2}{2} - (\underline{0}_{r/2-1}, l) \right)_{\mathbf{m}-\mathbf{l} + (\underline{0}_{r/2-1}, l), d} & (r: \text{even}), \\ (\nu)_{\mathbf{m}, d} \left(\nu - \frac{d_2}{2} \right)_{\mathbf{m}-\mathbf{l}', d} \left(\nu - \frac{d_2}{2}(r-1) - l \right)_{l - l_{\lceil r/2 \rceil}} & (r: \text{odd}), \end{cases} \\ &(\nu + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) - (l, \underline{0}_{r-1}), d_2} = (\nu + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m}+\mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d} \left(\nu - \frac{d_2}{2} \right)_{\mathbf{m}, d}. \end{aligned}$$

Then since $F_{\mathbf{m}, \mathbf{l}}^\mp[f](x_1, y_2) \in \mathcal{P}(\mathfrak{p}_1^+) \otimes \mathcal{P}_{\phi_\mp(\mathbf{m}, \mathbf{l})}(\mathfrak{p}_2^-)$ holds, by Lemma 2.3 we have

$$\begin{aligned} &\det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial x_2} \right)^k (\nu - (\underline{0}_{r-1}, l))_{\phi_-(\mathbf{m}, \mathbf{l}) + (\underline{0}_{r-1}, l), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu} F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, x_2^{-1}) \\ &= (-1)^{kr} (\nu + \phi_-(\mathbf{m}, \mathbf{l}))_{\underline{k}_r, d_2} (\nu - (\underline{0}_{r-1}, l))_{\phi_-(\mathbf{m}, \mathbf{l}) + (\underline{0}_{r-1}, l), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu-k} F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, x_2^{-1}) \\ &= (-1)^{kr} (\nu - (\underline{0}_{r-1}, l))_{\phi_-(\mathbf{m}, \mathbf{l}) + \underline{k}_r + (\underline{0}_{r-1}, l), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu-k} F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, x_2^{-1}) \\ &= (-1)^{kr} (\nu - (\underline{0}_{r-1}, l))_{\underline{k}_r, d_2} \\ &\quad \times (\nu + k - (\underline{0}_{r-1}, l))_{\phi_-(\mathbf{m}, \mathbf{l}) + (\underline{0}_{r-1}, l), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu-k} F_{\mathbf{m}, \mathbf{l}}^-[f](x_1, x_2^{-1}), \\ &\det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial x_2} \right)^k (\nu + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) - (l, \underline{0}_{r-1}), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu} F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, x_2^{-1}) \\ &= (-1)^{kr} (\nu + \phi_+(\mathbf{m}, \mathbf{l}))_{\underline{k}_r, d_2} (\nu + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) - (l, \underline{0}_{r-1}), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu-k} F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, x_2^{-1}) \\ &= (-1)^{kr} (\nu + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) + \underline{k}_r - (l, \underline{0}_{r-1}), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu-k} F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, x_2^{-1}) \\ &= (-1)^{kr} (\nu + (l, \underline{0}_{r-1}))_{\underline{k}_r, d_2} \\ &\quad \times (\nu + k + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) - (l, \underline{0}_{r-1}), d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\nu-k} F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, x_2^{-1}). \end{aligned}$$

Hence the 1st and 3rd equalities follow from the definition of $F_{l,\pm}^\downarrow \left(\frac{\nu}{\mu}; f; x \right)$. The 2nd and 4th equalities are easy. \square

4.2 Results on weighted Bergman inner products

For $\lambda \in \mathbb{C}$, $k, l \in \mathbb{Z}_{\geq 0}$, we set

$$C_{r,+}^{d,d_2}(\lambda, k, l) := \begin{cases} \frac{(\lambda + k - \frac{d}{4}r + \frac{d_2}{2} + (l, \underline{0}_{r/2-1}))_{\underline{k}_{r/2}, d}}{(\lambda)_{(2k+l, \underline{2k}_{r/2-1}, \underline{k}_{r/2}), d}} & (r: \text{even}), \\ \frac{(\lambda - \frac{d}{2}\lfloor \frac{r}{2} \rfloor + \max\{2k, k+l\})_{\min\{k, l\}} (\lambda + k - \frac{d}{2}\lceil \frac{r}{2} \rceil + \frac{d_2}{2})_{\underline{k}_{\lfloor r/2 \rfloor}, d}}{(\lambda)_{(2k+l, \underline{2k}_{\lfloor r/2 \rfloor-1}, \min\{2k, k+l\}, \underline{k}_{\lfloor r/2 \rfloor}), d}} & (r: \text{odd}), \end{cases} \quad (4.6)$$

and when $k \geq l$ we set

$$C_{r,-}^{d,d_2}(\lambda, k, l) := \begin{cases} \frac{(\lambda + k - \frac{d}{4}r + \frac{d_2}{2})_{(\underline{k}_{r/2-1}, k-l), d}}{(\lambda)_{(\underline{2k}_{r/2-1}, 2k-l, \underline{k}_{r/2-1}, k-l), d}} & (r: \text{even}), \\ \frac{(\lambda + k - \frac{d}{2}\lfloor \frac{r}{2} \rfloor + \frac{d_2}{2} + (k-l, \underline{0}_{\lfloor r/2 \rfloor}))_{(l, \underline{k}_{\lfloor r/2 \rfloor-1}, k-l), d}}{(\lambda)_{(\underline{2k}_{\lfloor r/2 \rfloor}, \underline{k}_{\lfloor r/2 \rfloor}, k-l), d}} & (r: \text{odd}). \end{cases} \quad (4.7)$$

If $r = 3$, then since $d = 2d_2$ holds, $C_{3,\pm}^{d,d_2}(\lambda, k, l)$ and $C^{d,d_2}(\lambda, \mathbf{k})$ in (3.3) are related as

$$\begin{aligned} C^{d,d_2}(\lambda, (k_1, k_2, k_3)) &= C_{3,+}^{d,d_2}(\lambda, k_2, k_1 - k_2), \\ C^{d,d_2}(\lambda, (k_2, k_2, k_3)) &= C_{3,-}^{d,d_2}(\lambda, k_2, k_2 - k_3). \end{aligned}$$

Also, for general r we have

$$C_{r,+}^{d,d_2}(\lambda, k, 0) = C_{r,-}^{d,d_2}(\lambda, k, 0) = C_{r+1,-}^{d,d_2}(\lambda, k, k) = \frac{(\lambda + k - \frac{d}{2}\lceil \frac{r}{2} \rceil + \frac{d_2}{2})_{\underline{k}_{\lfloor r/2 \rfloor}, d}}{(\lambda)_{(\underline{2k}_{\lfloor r/2 \rfloor}, \underline{k}_{\lceil r/2 \rceil}), d}}.$$

Now, for $\det_{\mathfrak{n}_2^+}(x_2)^k f(x_2) \in \mathcal{P}_{(k+l, \underline{k}_{r-1})}(\mathfrak{p}_2^+)$, the following holds.

Theorem 4.6. Let $k, l \in \mathbb{Z}_{\geq 0}$, $f(x_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$.

(1) For $\operatorname{Re} \lambda > \frac{2n}{r} - 1$ we have

$$\left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} = C_{r,+}^{d,d_2}(\lambda, k, l) F_{l,+}^{\downarrow} \left(-\lambda - 2k + \frac{n}{r}; f; z \right).$$

(2) As functions of λ ,

$$\begin{aligned} (\lambda)_{(2k+l, \underline{2k}_{r/2-1}, \underline{k}_{r/2}), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} &\quad (r: \text{even}), \\ (\lambda)_{(2k+l, \underline{2k}_{\lfloor r/2 \rfloor-1}, \min\{2k, k+l\}, \underline{k}_{\lfloor r/2 \rfloor}), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} &\quad (r: \text{odd}) \end{aligned}$$

are holomorphically continued for all $\lambda \in \mathbb{C}$, and give non-zero polynomials in $z \in \mathfrak{p}^+$ for all $\lambda \in \mathbb{C}$ if $f(x_2) \neq 0$.

(3) For $\mu, \nu \in \mathbb{C}$ we have

$$F_{l,+}^{\downarrow} \left(\frac{\nu}{\mu}; f; z \right) = \det_{\mathfrak{n}^+}(z)^{\mu-2\nu} F_{l,+}^{\downarrow} \left(\frac{\mu-\nu}{\mu}; f; z \right).$$

Especially if $\mu = a - 2k$ and $\nu = -k$ with $a, k \in \mathbb{Z}_{\geq 0}$, $a \leq k$, then $F_{l,+}^{\downarrow}(\frac{\nu}{\mu}; f; z)$ is factorized as

$$F_{l,+}^{\downarrow}\left(\begin{matrix} -k \\ a - 2k \end{matrix}; f; z\right) = \det_{\mathfrak{n}^+}(z)^a F_{l,+}^{\downarrow}\left(\begin{matrix} a - k \\ a - 2k \end{matrix}; f; z\right).$$

Theorem 4.6 (1) and (3) for $\nu = -k \in \mathbb{Z}_{\geq 0}$ case are already proved in [34, Theorem 6.3 (1)], and we can prove (3) for general $\nu \in \mathbb{C}$ by using the Zariski denseness of $-\mathbb{Z}_{\geq 0} \subset \mathbb{C}$. For Theorem 4.6 (2), their holomorphy for general r, k, l and non-zeroness for r even or $kl = 0$ are given in [34, Corollary 6.6 (1)], and the only new result is the non-zeroness for odd r with $k, l \neq 0$. Next, for $\det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}) \in \mathcal{P}_{(\underline{k}_{r-1}, k-l)}(\mathfrak{p}_2^+)$, the following holds.

Theorem 4.7. *Let $k, l \in \mathbb{Z}_{\geq 0}$ with $k \geq l$, and $f(x_2) \in \mathcal{P}_{(l, \underline{k}_{r-1})}(\mathfrak{p}_2^+)$.*

(1) *For $\operatorname{Re} \lambda > \frac{2n}{r} - 1$ we have*

$$\left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} = C_{r,-}^{d,d_2}(\lambda, k, l) F_{l,-}^{\downarrow}\left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z\right).$$

(2) *As functions of λ ,*

$$\begin{aligned} & (\lambda)_{(2\underline{k}_{r/2-1}, 2k-l, \underline{k}_{r/2-1}, k-l), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} && (r: \text{even}), \\ & (\lambda)_{(2k_{\lfloor r/2 \rfloor}, k_{\lfloor r/2 \rfloor}, k-l), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} && (r: \text{odd}) \end{aligned}$$

are holomorphically continued for all $\lambda \in \mathbb{C}$, and give non-zero polynomials in $z \in \mathfrak{p}^+$ for all $\lambda \in \mathbb{C}$ if $f(z_2) \neq 0$.

(3) *For $\mu, \nu \in \mathbb{C}$ we have*

$$F_{l,-}^{\downarrow}\left(\begin{matrix} \nu \\ \mu \end{matrix}; f; z\right) = \det_{\mathfrak{n}^+}(z)^{\mu-2\nu} F_{l,-}^{\downarrow}\left(\begin{matrix} \mu - \nu \\ \mu \end{matrix}; f; z\right).$$

Especially if $\mu = a - 2k$ and $\nu = -k$ with $a, k \in \mathbb{Z}_{\geq 0}$, $a \leq k - l$, then $F_{l,-}^{\downarrow}(\frac{\nu}{\mu}; f; z)$ is factorized as

$$F_{l,-}^{\downarrow}\left(\begin{matrix} -k \\ a - 2k \end{matrix}; f; z\right) = \det_{\mathfrak{n}^+}(z)^a F_{l,-}^{\downarrow}\left(\begin{matrix} a - k \\ a - 2k \end{matrix}; f; z\right).$$

The holomorphy in Theorem 4.7 (2) is already proved in [35, Theorem 6.2], but the other results are new except for $l = 0$ (Example 4.10) or $r = 2$ (Example 4.11).

Remark 4.8. *For + case, in [34, Theorem 6.3 (1)] we proved*

$$\begin{aligned} & \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \frac{1}{(\lambda)_{(2k+l, \underline{2k}_{r-1}), d}} \det_{\mathfrak{n}^+}(z)^{-\lambda + \frac{n}{r}} \det_{\mathfrak{n}_2^+}\left(\frac{\partial}{\partial z_2}\right)^k \det_{\mathfrak{n}^+}(z)^{\lambda + 2k - \frac{n}{r}} f(z_2) \\ &= \frac{1}{(\lambda)_{(2k+l, \underline{2k}_{r-1}), d}} \det_{\mathfrak{n}^+}(z)^{-\lambda + \frac{n}{r}} \det_{\mathfrak{n}_2^+}\left(\frac{\partial}{\partial z_2}\right)^k F_{l,+}^{\downarrow}\left(\begin{matrix} -\lambda - 2k + \frac{n}{r} \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z\right) \\ &= C_{r,+}^{d,d_2}(\lambda, k, l) \det_{\mathfrak{n}^+}(z)^{-\lambda + \frac{n}{r}} F_{l,+}^{\downarrow}\left(\begin{matrix} -\lambda - k + \frac{n}{r} \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z\right) \\ &= C_{r,+}^{d,d_2}(\lambda, k, l) F_{l,+}^{\downarrow}\left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z\right). \end{aligned}$$

Similarly, for $-$ case, by [35, Theorem 3.3 (1)] we have

$$\begin{aligned} & \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} \\ &= \frac{\det_{\mathfrak{n}^+}(z)^{-\lambda+\frac{n}{r}}}{(\lambda)_{\underline{2(k-l)}_r, d}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k-l} \det_{\mathfrak{n}^+}(x)^{\lambda+2(k-l)-\frac{n}{r}} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^l f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda+2(k-l),x}, \end{aligned} \quad (4.8)$$

and combining with Proposition 4.5 and Theorem 4.7(1), (3), we have

$$\begin{aligned} & \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} \\ &= \frac{C_{r,-}^{d,d_2}(\lambda+2(k-l), l, l)}{(\lambda)_{\underline{2(k-l)}_r, d}} \det_{\mathfrak{n}^+}(z)^{-\lambda+\frac{n}{r}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k-l} \det_{\mathfrak{n}^+}(z)^{\lambda+2(k-l)-\frac{n}{r}} \\ & \quad \times F_{l,-}^{\downarrow} \left(\begin{matrix} -l \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right) \\ &= \frac{(\lambda+2k-l-\frac{d}{2}\lfloor\frac{r}{2}\rfloor+\frac{d_2}{2})_{\underline{l}_{\lceil r/2 \rceil-1}, d}}{(\lambda)_{(2k_{\lceil r/2 \rceil-1}, 2k-l_{\lfloor r/2 \rfloor}, 2k-2l), d}} \det_{\mathfrak{n}^+}(z)^{-\lambda+\frac{n}{r}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k-l} \\ & \quad \times F_{l,-}^{\downarrow} \left(\begin{matrix} -\lambda - 2k + l + \frac{n}{r} \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right) \\ &= C_{r,-}^{d,d_2}(\lambda, k, l) \det_{\mathfrak{n}^+}(z)^{-\lambda+\frac{n}{r}} F_{l,-}^{\downarrow} \left(\begin{matrix} -\lambda - k + \frac{n}{r} \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right) \\ &= C_{r,-}^{d,d_2}(\lambda, k, l) F_{l,-}^{\downarrow} \left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right). \end{aligned}$$

First we prove Theorem 4.7(1). By [35, Theorem 6.1] with $\mathbf{k} = (k_{r-1}, k-l)$, $k_{r+1} = 0$, when $d = 2d_2$ we have

$$\begin{aligned} & \frac{\left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x}}{\det_{\mathfrak{n}_2^+}(z_2)^k f(z_2^{-1})} = \frac{\prod_{a=2}^{2r-2} \prod_{i=\max\{1, a+1-r\}}^{\lfloor a/2 \rfloor} (\lambda - \frac{d}{4}(a-1))_{k_i+k_{a+1-i}}}{\prod_{a=1}^{2r-1} \prod_{i=\max\{1, a+1-r\}}^{\lceil a/2 \rceil} (\lambda - \frac{d}{4}(a-1))_{k_i+k_{a+2-i}}} \\ &= \prod_{a=1}^{r-2} \frac{\prod_{i=1}^{\lfloor a/2 \rfloor} (\lambda - \frac{d}{4}(a-1))_{2k}}{\prod_{i=1}^{\lceil a/2 \rceil} (\lambda - \frac{d}{4}(a-1))_{2k}} \prod_{a=r-1}^{r-1} \frac{\prod_{i=1}^{\lfloor a/2 \rfloor} (\lambda - \frac{d}{4}(a-1))_{2k}}{(\lambda - \frac{d}{4}(a-1))_{2k-l} \prod_{i=2}^{\lceil a/2 \rceil} (\lambda - \frac{d}{4}(a-1))_{2k}} \\ & \quad \times \prod_{a=r}^{2r-3} \frac{(\lambda - \frac{d}{4}(a-1))_{2k-l} \prod_{i=a+2-r}^{\lfloor a/2 \rfloor} (\lambda - \frac{d}{4}(a-1))_{2k}}{(\lambda - \frac{d}{4}(a-1))_k (\lambda - \frac{d}{4}(a-1))_{2k-l} \prod_{i=a+3-r}^{\lceil a/2 \rceil} (\lambda - \frac{d}{4}(a-1))_{2k}} \\ & \quad \times \prod_{a=2r-2}^{2r-2} \frac{(\lambda - \frac{d}{4}(a-1))_{2k-l}}{(\lambda - \frac{d}{4}(a-1))_k} \prod_{a=2r-1}^{2r-1} \frac{1}{(\lambda - \frac{d}{4}(a-1))_{k-l}} \\ &= \frac{1}{\prod_{a=1, \text{odd}}^{r-2} (\lambda - \frac{d}{4}(a-1))_{2k}} \frac{\prod_{a=r-1, \text{even}}^{r-1} (\lambda - \frac{d}{4}(a-1) + 2k-l)_l}{\prod_{a=r-1, \text{odd}}^{r-1} (\lambda - \frac{d}{4}(a-1))_{2k-l}} \\ & \quad \times \frac{\prod_{a=r, \text{even}}^{2r-3} (\lambda - \frac{d}{4}(a-1) + k)_k \prod_{a=2r-2}^{2r-2} (\lambda - \frac{d}{4}(a-1) + k)_{k-l}}{\prod_{a=r, \text{odd}}^{2r-3} (\lambda - \frac{d}{4}(a-1))_k \prod_{a=2r-1}^{2r-1} (\lambda - \frac{d}{4}(a-1))_{k-l}} \end{aligned}$$

$$= \begin{cases} \frac{(\lambda - \frac{d}{4}(r-1) + k)_{(\underline{k}_{r/2-1}, k-l), d}}{(\lambda)_{(2\underline{k}_{r/2-1}, 2k-l, \underline{k}_{r/2-1}, k-l), d}} = \frac{(\lambda + k - \frac{d}{4}r + \frac{d_2}{2})_{(\underline{k}_{r/2-1}, k-l), d}}{(\lambda)_{(2\underline{k}_{r/2-1}, 2k-l, \underline{k}_{r/2-1}, k-l), d}} & (r: \text{even}), \\ \frac{(\lambda - \frac{d}{4}(r-2) + (2k-l, \underline{k}_{\lfloor r/2 \rfloor}))_{(l, \underline{k}_{\lfloor r/2 \rfloor}-1, k-l), d}}{(\lambda)_{(2\underline{k}_{\lfloor r/2 \rfloor}, \underline{k}_{\lfloor r/2 \rfloor}, k-l), d}} \\ = \frac{(\lambda + k - \frac{d}{2}\lfloor \frac{r}{2} \rfloor + \frac{d_2}{2} + (k-l, \underline{0}_{\lfloor r/2 \rfloor}))_{(l, \underline{k}_{\lfloor r/2 \rfloor}-1, k-l), d}}{(\lambda)_{(2\underline{k}_{\lfloor r/2 \rfloor}, \underline{k}_{\lfloor r/2 \rfloor}, k-l), d}} & (r: \text{odd}), \end{cases}$$

and again by [35, Theorem 6.1] this also holds for $r = 2$, $d \neq 2d_2$ case. Therefore, to prove (1), it is enough to prove that $\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \rangle_{\lambda, x}$ is proportional to $F_{l,-}^\downarrow(-\lambda - \frac{-k}{2k+\frac{n}{r}}; f; z)$, and by the F-method, enough to prove

$$\left(\det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}) \mapsto F_{l,-}^\downarrow(-\lambda - \frac{-k}{2k+\frac{n}{r}}; f; z) \right) \in \text{Hom}_{K_1}(\mathcal{P}_{(\underline{k}_{r-1}, k-l)}(\mathfrak{p}_2^+), \text{Sol}_{\mathcal{P}(\mathfrak{p}^+)}((\mathcal{B}_\lambda)_1)), \quad (4.9)$$

where $(\mathcal{B}_\lambda)_1$ is the orthogonal projection of $\mathcal{B}_\lambda = \mathcal{B}_\lambda^{\mathfrak{p}^+}: \mathcal{P}(\mathfrak{p}^+) \rightarrow \mathcal{P}(\mathfrak{p}^+) \otimes \mathfrak{p}^+$ given in (2.24) (with the identification $\mathfrak{p}^- \simeq \mathfrak{p}^+ = \mathfrak{n}^{+\mathbb{C}}$) onto \mathfrak{p}_1^+ .

Proposition 4.9. *Let $\mu, \nu \in \mathbb{C}$, $l \in \mathbb{Z}_{\geq 0}$, $f(x_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$.*

(1) *For $z = z_1 + z_2 \in \Omega$, $a_2 \in \Omega_2$ with $z_1 + a_2 \in \Omega$ and for $\text{Re } \mu > l + \frac{2n_2}{r} - 1$, $\text{Re } \nu > l + \frac{n_2}{r} - 1$, we have*

$$F_{l,+}^\downarrow\left(\frac{\nu}{\mu}; f; z\right) = \frac{1}{(2\pi\sqrt{-1})^{n_2}} \frac{\Gamma_r^{d_2}(\mu - (\underline{0}_{r-1}, l))}{\Gamma_r^{d_2}(\nu - (\underline{0}_{r-1}, l))} \int_{\Omega_2} e^{-(y_2|z_2)_{\mathfrak{n}_2^+}} \det_{\mathfrak{n}_2^+}(y_2)^{\nu-\mu} \\ \times \left(\int_{a_2 + \sqrt{-1}\mathfrak{n}_2^+} e^{(x_2|y_2)_{\mathfrak{n}_2^+}} \det_{\mathfrak{n}^+}(z_1 + x_2)^{-\mu} f(x_2) dx_2 \right) dy_2.$$

Similarly, for $\text{Re } \mu > \frac{d_2}{2} + \frac{2n_2}{r} - 1$, $\text{Re } \nu > \frac{n_2}{r} - 1$, we have

$$F_{l,-}^\downarrow\left(\frac{\nu}{\mu}; f; z\right) = \frac{1}{(2\pi\sqrt{-1})^{n_2}} \frac{\Gamma_r^{d_2}(\mu - \frac{d_2}{2} + (l, \underline{0}_{r-1}))}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \int_{\Omega_2} e^{-(y_2|z_2)_{\mathfrak{n}_2^+}} \det_{\mathfrak{n}_2^+}(y_2)^{\nu-\mu+\frac{d_2}{2}} \\ \times \left(\int_{a_2 + \sqrt{-1}\mathfrak{n}_2^+} e^{(x_2|y_2)_{\mathfrak{n}_2^+}} \det_{\mathfrak{n}_2^+}(x_2)^{-\frac{d_2}{2}} \det_{\mathfrak{n}^+}(z_1 + x_2)^{-\mu+d_2} f(x_2^{-1}) dx_2 \right) dy_2.$$

(2) Suppose $\mu \in \mathbb{C}$ is not a pole of $F_{l,\pm}^\downarrow(\frac{\nu}{\mu}; f; z)$. Then for $g \in \widetilde{K}_1^{\mathbb{C}}$ we have

$$F_{l,+}^\downarrow\left(\frac{\nu}{\mu}; f(g(\cdot)); z\right) = \det_{\mathfrak{n}_2^+}(ge)^\nu F_{l,+}^\downarrow\left(\frac{\nu}{\mu}; f; gz\right), \\ F_{l,-}^\downarrow\left(\frac{\nu}{\mu}; f(t g^{-1}(\cdot)); z\right) = \det_{\mathfrak{n}_2^+}(ge)^\nu F_{l,-}^\downarrow\left(\frac{\nu}{\mu}; f; gz\right).$$

(3) $F_{l,\pm}^\downarrow(\frac{\nu}{\mu}; f; z)$ satisfies the differential equation

$$(\mathcal{B}_{-\mu+2\nu+\frac{n}{r}})_1 F_{l,\pm}^\downarrow\left(\frac{\nu}{\mu}; f; z\right) = 0.$$

For $F_{l,+}^\downarrow(\frac{\nu}{\mu}; f; z)$, this proposition is a repost of [34, Proposition 6.4], and we only prove this for $F_{l,-}^\downarrow(\frac{\nu}{\mu}; f; z)$.

Proof. (1) By (4.2) and the definition of $F_{\mathbf{m}, \mathbf{l}}^+[f](x_1, y_2)$, we have

$$\begin{aligned} & \det_{\mathfrak{n}^+}(z_1 + x_2)^{-\mu+d_2} f(x_2^{-1}) = \det_{\mathfrak{n}_2^+}(x_2)^{-\mu+d_2} h_{\mathfrak{n}^+}(z_1, P(x_2^{-1})z_1)^{(-\mu+d_2)/2} f(x_2^{-1}) \\ &= \det_{\mathfrak{n}_2^+}(x_2)^{-\mu+d_2} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} (\mu - d_2)_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, x_2^{-1}) f(x_2^{-1}) \\ &= \det_{\mathfrak{n}_2^+}(x_2)^{-\mu+d_2} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} (\mu - d_2)_{\mathbf{m}, d} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, x_2^{-1}), \end{aligned}$$

and hence by Lemma 2.2 we get

$$\begin{aligned} & \frac{1}{(2\pi\sqrt{-1})^{n_2}} \frac{\Gamma_r^{d_2}(\mu - \frac{d_2}{2} + (l, \underline{0}_{r-1}))}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \int_{\Omega_2} e^{-(y_2|z_2)}_{\mathfrak{n}_2^+} \det_{\mathfrak{n}_2^+}(y_2)^{\nu - \mu + \frac{d_2}{2}} \\ & \quad \times \left(\int_{a_2 + \sqrt{-1}\mathfrak{n}_2^+} e^{(x_2|y_2)}_{\mathfrak{n}_2^+} \det_{\mathfrak{n}_2^+}(x_2)^{-\frac{d_2}{2}} \det_{\mathfrak{n}^+}(z_1 + x_2)^{-\mu+d_2} f(x_2^{-1}) dx_2 \right) dy_2 \\ &= \frac{1}{(2\pi\sqrt{-1})^n} \frac{\Gamma_r^{d_2}(\mu - \frac{d_2}{2} + (l, \underline{0}_{r-1}))}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \int_{\Omega_2} e^{-(y_2|z_2)}_{\mathfrak{n}_2^+} \det_{\mathfrak{n}_2^+}(y_2)^{\nu - \mu + \frac{d_2}{2}} \\ & \quad \times \left(\int_{a_2 + \sqrt{-1}\mathfrak{n}_2^+} e^{(x_2|y_2)}_{\mathfrak{n}_2^+} \det_{\mathfrak{n}_2^+}(x_2)^{-\mu + \frac{d_2}{2}} \sum_{\mathbf{m}, \mathbf{l}} (\mu - d_2)_{\mathbf{m}, d} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, x_2^{-1}) dx_2 \right) dy_2 \\ &= \frac{\Gamma_r^{d_2}(\mu - \frac{d_2}{2} + (l, \underline{0}_{r-1}))}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \int_{\Omega_2} e^{-(y_2|z_2)}_{\mathfrak{n}_2^+} \det_{\mathfrak{n}_2^+}(y_2)^{\nu - \frac{n_2}{r}} \sum_{\mathbf{m}, \mathbf{l}} \frac{(\mu - d_2)_{\mathbf{m}, d} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, y_2)}{\Gamma_r^{d_2}(\mu - \frac{d_2}{2} + \phi_+(\mathbf{m}, \mathbf{l}))} dy_2 \\ &= \frac{1}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \int_{\Omega_2} e^{-(y_2|z_2)}_{\mathfrak{n}_2^+} \sum_{\mathbf{m}, \mathbf{l}} \frac{(\mu - d_2)_{\mathbf{m}, d} \det_{\mathfrak{n}_2^+}(y_2)^{\nu - \frac{n_2}{r}} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, y_2)}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) - (l, \underline{0}_{r-1}), d}} dy_2 \\ &= \frac{1}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \int_{\Omega_2} e^{-(y_2|z_2)}_{\mathfrak{n}_2^+} \sum_{\mathbf{m}, \mathbf{l}} \frac{\det_{\mathfrak{n}_2^+}(y_2)^{\nu - \frac{n_2}{r}} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, y_2)}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} dy_2 \\ &= \sum_{\mathbf{m}, \mathbf{l}} \frac{\Gamma_r^{d_2}(\nu + \phi_+(\mathbf{m}, \mathbf{l}))}{\Gamma_r^{d_2}(\nu + (l, \underline{0}_{r-1}))} \frac{\det_{\mathfrak{n}_2^+}(z_2)^{-\nu} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1})}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} \\ &= \sum_{\mathbf{m}, \mathbf{l}} \frac{(\nu + (l, \underline{0}_{r-1}))_{\phi_+(\mathbf{m}, \mathbf{l}) - (l, \underline{0}_{r-1}), d}}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} \det_{\mathfrak{n}_2^+}(z_2)^{-\nu} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}) \\ &= \sum_{\mathbf{m}, \mathbf{l}} \frac{(\nu + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d} (\nu - \frac{d_2}{2})_{\mathbf{m}, d}}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} \det_{\mathfrak{n}_2^+}(z_2)^{-\nu} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}) \\ &= F_{l,-}^\downarrow\left(\frac{\nu}{\mu}; f; z\right). \end{aligned}$$

(2) Since $F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1})$ satisfies

$$F_{\mathbf{m}, \mathbf{l}}^+[f(tg^{-1}(\cdot))](z_1, z_2^{-1}) = F_{\mathbf{m}, \mathbf{l}}^+[f](gz_1, (gz_2)^{-1}),$$

we get the desired formula.

(3) As in the proof of [34, Proposition 6.4], we can show that

$$\begin{aligned}
& (\mathcal{B}_{-\mu+2\nu+\frac{n}{r}})_1 \int_{\Omega_2} e^{-(y_2|z_2)_{\mathfrak{n}_2^+}} \det_{\mathfrak{n}_2^+}(y_2)^{\nu-\mu+\frac{d_2}{2}} \left(\int_{a_2+\sqrt{-1}\mathfrak{n}_2^+} e^{(x_2|y_2)_{\mathfrak{n}_2^+}} \right. \\
& \quad \times \det_{\mathfrak{n}^+}(z_1+x_2)^{-\mu+d_2} \det_{\mathfrak{n}_2^+}(x_2)^{-\frac{d_2}{2}} f(x_2^{-1}) dx_2 \Big) dy_2 \\
& = \int_{\Omega_2} e^{-(y_2|z_2)_{\mathfrak{n}_2^+}} \det_{\mathfrak{n}_2^+}(y_2)^{\nu-\mu+\frac{d_2}{2}} \left(\int_{a_2+\sqrt{-1}\mathfrak{n}_2^+} e^{(x_2|y_2)_{\mathfrak{n}_2^+}} \right. \\
& \quad \times \det_{\mathfrak{n}^+}(z_1+x_2)^{-\mu+d_2} (\mathcal{B}_{-\mu+d_2+\frac{n}{r}})_1 \det_{\mathfrak{n}_2^+}(x_2)^{-\frac{d_2}{2}} f(x_2^{-1}) dx_2 \Big) dy_2.
\end{aligned}$$

Hence it is enough to prove $(\mathcal{B}_\lambda)_1 \det_{\mathfrak{n}_2^+}(x_2)^{-d_2/2} f(x_2^{-1}) = 0$. In the following we write $(z_1, x_2) = (\zeta, X)$, let $\{E_\alpha\} \subset \mathfrak{p}_2^+$ be a basis with the dual basis $\{E_\alpha^\vee\} \subset \mathfrak{p}_2^+$, and let $\frac{\partial}{\partial X_\alpha}$, $\frac{\partial}{\partial Y_\alpha^\vee}$ denote the directional derivatives along E_α and E_α^\vee . Then we have

$$\begin{aligned}
& (\mathcal{B}_\lambda)_1 \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} f(X^{-1}) = \frac{1}{2} \sum_{\alpha\beta} P(E_\alpha^\vee, E_\beta^\vee) \zeta \frac{\partial^2}{\partial X_\alpha \partial X_\beta} \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} f(X^{-1}) \\
& = \frac{1}{2} \sum_{\alpha\beta} P(E_\alpha^\vee, E_\beta^\vee) \zeta \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} \left(\left(\frac{d_2^2}{4} (E_\alpha|X^{-1})_{\mathfrak{n}_2^+} (E_\beta|X^{-1})_{\mathfrak{n}_2^+} \right. \right. \\
& \quad + \frac{d_2}{2} (E_\beta|P(X)^{-1}E_\alpha)_{\mathfrak{n}_2^+} \Big) f(X^{-1}) - d_2 (E_\beta|X^{-1})_{\mathfrak{n}_2^+} \frac{\partial}{\partial X_\alpha} f(X^{-1}) + \frac{\partial^2}{\partial X_\alpha \partial X_\beta} f(X^{-1}) \Big) \\
& = \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} \left(\left(\frac{d_2^2}{4} P(X^{-1}) \zeta + \frac{d_2}{4} \sum_\alpha P(E_\alpha^\vee, P(X)^{-1}E_\alpha) \zeta \right) f(X^{-1}) \right. \\
& \quad + \frac{d_2}{2} \sum_\alpha P(E_\alpha^\vee, X^{-1}) \zeta \sum_\gamma (E_\gamma|P(X)^{-1}E_\alpha)_{\mathfrak{n}_2^+} \frac{\partial f}{\partial Y_\gamma^\vee}(X^{-1}) \\
& \quad + \frac{1}{2} \sum_{\alpha\beta} P(E_\alpha^\vee, E_\beta^\vee) \zeta \left(\sum_{\gamma\delta} (E_\gamma|P(X^{-1})E_\alpha)_{\mathfrak{n}_2^+} (E_\delta|P(X^{-1})E_\beta)_{\mathfrak{n}_2^+} \frac{\partial^2 f}{\partial Y_\gamma^\vee \partial Y_\delta^\vee}(X^{-1}) \right. \\
& \quad \left. \left. + \sum_\gamma (E_\gamma|P(X^{-1}, P(X)^{-1}E_\alpha)E_\beta)_{\mathfrak{n}_2^+} \frac{\partial f}{\partial Y_\gamma^\vee}(X^{-1}) \right) \right) \\
& = \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} \left(\left(\frac{d_2^2}{4} P(X)^{-1} \zeta + \frac{d_2}{4} \sum_\alpha P(E_\alpha^\vee, P(X)^{-1}E_\alpha) \zeta \right) f(X^{-1}) \right. \\
& \quad + \frac{d_2}{2} \sum_\gamma P(P(X)^{-1}E_\gamma, X^{-1}) \zeta \frac{\partial f}{\partial Y_\gamma^\vee}(X^{-1}) \\
& \quad + \frac{1}{2} \sum_{\gamma\delta} P(P(X)^{-1}E_\gamma, P(X)^{-1}E_\delta) \zeta \frac{\partial^2 f}{\partial Y_\gamma^\vee \partial Y_\delta^\vee}(X^{-1}) \\
& \quad \left. + \frac{1}{2} \sum_{\alpha\gamma} P(E_\alpha^\vee, P(X^{-1}, P(X)^{-1}E_\alpha)E_\gamma) \zeta \frac{\partial f}{\partial Y_\gamma^\vee}(X^{-1}) \right).
\end{aligned}$$

Then at the 5th term, since $\sum_\alpha P(E_\alpha^\vee, P(X)^{-1}, P(X)^{-1}E_\alpha)E_\gamma) \zeta$ does not depend on the choice of $\{E_\alpha\}$ and $\{E_\alpha^\vee\}$, we may replace these with $\{P(X^{1/2})E'_\alpha\}$ and $\{P(X^{-1/2})E'_\alpha\}$ by using an orthonormal basis $\{E'_\alpha\} \subset \mathfrak{n}_2^+$. Then by using the equality $X^{-1} = P(X^{-1/2})E$,

where $E \in \mathfrak{n}_2^+$ is the unit element, and $P(P(X)Y, P(X)Z) = P(X)P(Y, Z)P(X)$, we have

$$\begin{aligned}
& \sum_{\alpha} P(E_{\alpha}^{\vee}, P(X^{-1}, P(X)^{-1}E_{\alpha})E_{\gamma})\zeta \\
&= \sum_{\alpha} P(P(X^{-1/2})E'_{\alpha}, P(P(X^{-1/2})E, P(X^{-1/2})E'_{\alpha})E_{\gamma})\zeta \\
&= \sum_{\alpha} P(X^{-1/2})P(E'_{\alpha}, P(E, E'_{\alpha})P(X^{-1/2})E_{\gamma})P(X^{-1/2})\zeta \\
&= P(X^{-1/2}) \sum_{\alpha} (P(E'_{\alpha})D_{\mathfrak{n}^+}(P(X^{-1/2})E_{\gamma}, E) + D_{\mathfrak{n}^+}(E, P(X^{-1/2})E_{\gamma})P(E'_{\alpha}))P(X^{-1/2})\zeta \\
&= P(X^{-1/2}) \left(-\frac{d_2}{2} D_{\mathfrak{n}^+}(P(X^{-1/2})E_{\gamma}, E) - \frac{d_2}{2} D_{\mathfrak{n}^+}(E, P(X^{-1/2})E_{\gamma}) \right) P(X^{-1/2})\zeta \\
&= -d_2 P(X^{-1/2})P(P(X^{-1/2})E_{\gamma}, E)P(X^{-1/2})\zeta = -d_2 P(P(X)^{-1}E_{\gamma}, X^{-1})\zeta,
\end{aligned}$$

where at the 3nd equality we have used [7, Part V, Proposition I.2.1 (J2.3)], at the 4th equality $\sum_{\alpha} P(E'_{\alpha})\eta = -\frac{d_2}{2}\eta$ follows from direct computation with an explicit basis, and at the 5th equality we have used $D_{\mathfrak{n}^+}(Y, E) = D_{\mathfrak{n}^+}(E, Y) = P(Y, E)$. Similarly, at the 2nd term we have

$$\begin{aligned}
& \sum_{\alpha} P(E_{\alpha}^{\vee}, P(X)^{-1}E_{\alpha})\zeta = \sum_{\alpha} P(P(X^{-1/2})E_{\alpha}^{\vee}, P(X^{-1/2})E_{\alpha})\zeta \\
&= P(X^{-1/2}) \sum_{\alpha} P(E'_{\alpha}, E'_{\alpha})P(X^{-1/2})\zeta = -d_2 P(X^{-1/2})^2\zeta = -d_2 P(X)^{-1}\zeta.
\end{aligned}$$

Therefore we get

$$\begin{aligned}
& (\mathcal{B}_{\lambda})_1 \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} f(X^{-1}) \\
&= \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} \left(\left(\frac{d_2^2}{4} P(X)^{-1}\zeta - \frac{d_2^2}{4} P(X)^{-1}\zeta \right) f(X^{-1}) \right. \\
&\quad + \frac{d_2}{2} \sum_{\gamma} P(P(X)^{-1}E_{\gamma}, X^{-1})\zeta \frac{\partial f}{\partial Y_{\gamma}^{\vee}}(X^{-1}) \\
&\quad + \frac{1}{2} \sum_{\gamma\delta} P(P(X)^{-1}E_{\gamma}, P(X)^{-1}E_{\delta})\zeta \frac{\partial^2 f}{\partial Y_{\gamma}^{\vee} \partial Y_{\delta}^{\vee}}(X^{-1}) \\
&\quad \left. - \frac{d_2}{2} \sum_{\gamma} P(P(X)^{-1}E_{\gamma}, X^{-1})\zeta \frac{\partial f}{\partial Y_{\gamma}^{\vee}}(X^{-1}) \right) \\
&= \frac{1}{2} \det_{\mathfrak{n}_2^+}(X)^{-d_2/2} P(X)^{-1} \sum_{\gamma\delta} P(E_{\gamma}, E_{\delta})P(X)^{-1}\zeta \frac{\partial^2 f}{\partial Y_{\gamma}^{\vee} \partial Y_{\delta}^{\vee}}(X^{-1}).
\end{aligned}$$

Now, let $\mathfrak{p}^+ = \bigoplus_{1 \leq i < j \leq r} \mathfrak{p}_{ij}^+$ be the simultaneous Peirce decomposition given in Section 2.4 with respect to a Jordan frame $\{e_1, \dots, e_r\} \subset \mathfrak{n}_2^+ \subset \mathfrak{n}^+$. If $f(Y) = (Y_{11})^l \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$, then since $\mathfrak{p}_{11}^+ \cap \mathfrak{p}_1^+ = \{0\}$, for $\eta \in \mathfrak{p}_1^+$ we have

$$\sum_{\gamma\delta} P(E_{\gamma}, E_{\delta})\eta \frac{\partial^2 f}{\partial Y_{\gamma}^{\vee} \partial Y_{\delta}^{\vee}}(Y) = l(l-1)P(E_{11}, E_{11})\eta(Y_{11})^{l-2} = 0,$$

and by the K_1 -equivariance, $\sum_{\gamma\delta} P(E_{\gamma}, E_{\delta})\eta \frac{\partial^2 f}{\partial Y_{\gamma}^{\vee} \partial Y_{\delta}^{\vee}}(Y) = 0$ holds for all $f \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$. Hence we get $(\mathcal{B}_{\lambda})_1 \det_{\mathfrak{n}_2^+}(x_2)^{-d_2/2} f(x_2^{-1}) = 0$ for all $f \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$, and also get

$(\mathcal{B}_{-\mu+2\nu+\frac{n}{r}})_1 F_{l,-}^{\downarrow}(\nu; f; z) = 0$ for $\operatorname{Re} \mu > \frac{d_2}{2} + \frac{2n_2}{r} - 1$, $\operatorname{Re} \nu > \frac{n_2}{r} - 1$. Then by analytic continuation, this holds for all $\mu, \nu \in \mathbb{C}$ except for poles. \square

Especially, when $\mu = -\lambda - 2k + \frac{n}{r}$, $\nu = -k$ with $k \in \mathbb{Z}_{\geq 0}$, $k \geq l$, we have (4.9), and this proves Theorem 4.7 (1). Next we prove Theorems 4.6 (2) and 4.7 (2).

Proof of Theorem 4.6 (2). Let $\mu := -\lambda - 2k + \frac{n}{r} = -\lambda - 2k + 1 + \frac{d}{2}(r-1)$. We recall from the proof of [34, Corollary 6.6 (1)] that, by Theorem 4.6 (1), for $f(x_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$, for even r we have

$$\begin{aligned} & (\lambda)_{(2k+l, \underline{2k}_{r/2-1}, k_{r/2}), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \left(\lambda + k - \frac{d}{4}r + \frac{d_2}{2} + (l, \underline{0}_{r/2-1}) \right)_{\underline{k}_{r/2}, d} F_{l,+}^{\downarrow} \left(\begin{matrix} -k \\ \mu \end{matrix}; f; z \right) \\ &= (-1)^{kr/2} \left(\mu - \frac{d_2}{2} - (\underline{0}_{r/2-1}, l) \right)_{\underline{k}_{r/2}, d} \det_{\mathfrak{n}_2^+}(z_2)^k \\ &\quad \times \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \frac{(-k)_{\mathbf{m}, d} \left(-k - \frac{d_2}{2} - (\underline{0}_{r/2-1}, l) \right)_{\mathbf{m}-\mathbf{l}+(\underline{0}_{r/2-1}, l), d}}{(\mu - \frac{d_2}{2} - (\underline{0}_{r/2-1}, l))_{\mathbf{m}-\mathbf{l}+(\underline{0}_{r/2-1}, l), d}} F_{\mathbf{m}, \mathbf{l}}^-[f](z_1, z_2^{-1}), \end{aligned} \tag{4.10}$$

and for odd r we have

$$\begin{aligned} & (\lambda)_{(2k+l, \underline{2k}_{\lfloor r/2 \rfloor-1}, \min\{2k, k+l\}, \underline{k}_{\lfloor r/2 \rfloor}), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \left(\lambda - \frac{d}{2} \left\lfloor \frac{r}{2} \right\rfloor + \max\{2k, k+l\} \right)_{\min\{k, l\}} \left(\lambda + k - \frac{d}{2} \left\lceil \frac{r}{2} \right\rceil + \frac{d_2}{2} \right)_{\underline{k}_{\lfloor r/2 \rfloor}, d} F_{l,+}^{\downarrow} \left(\begin{matrix} -k \\ \mu \end{matrix}; f; z \right) \\ &= (-1)^{k\lfloor r/2 \rfloor + \min\{k, l\}} \left(\mu - \frac{d_2}{2} \right)_{\underline{k}_{\lfloor r/2 \rfloor}, d} \left(\mu - \frac{d_2}{2}(r-1) - l \right)_{\min\{k, l\}} \det_{\mathfrak{n}_2^+}(z_2)^k \\ &\quad \times \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \frac{(-k)_{\mathbf{m}, d} \left(-k - \frac{d_2}{2} \right)_{\mathbf{m}-\mathbf{l}', d} \left(-k - \frac{d_2}{2}(r-1) - l \right)_{l-l^{\lceil r/2 \rceil}}}{(\mu - \frac{d_2}{2})_{\mathbf{m}-\mathbf{l}', d} \left(\mu - \frac{d_2}{2}(r-1) - l \right)_{l-l^{\lceil r/2 \rceil}}} F_{\mathbf{m}, \mathbf{l}}^-[f](z_1, z_2^{-1}). \end{aligned} \tag{4.11}$$

These are holomorphically continued for all $\mu \in \mathbb{C}$. Moreover, all non-zero $F_{\mathbf{m}, \mathbf{l}}^-[f](z_1, z_2^{-1})$ are linearly independent, and when r is even, the coefficient of (4.10) for

$$F_{\underline{k}_{r/2}, (\underline{0}_{r/2-1}, l)}^-[f](z_1, z_2^{-1}) = \tilde{\Phi}_{\underline{k}_{r/2}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1}) f(z_2)$$

is independent of μ and non-zero. Hence (4.10) is non-zero for all $\mu \in \mathbb{C}$ when $f(x_2) \neq 0$. Similarly, when r is odd, the coefficients of (4.11) for $(\mathbf{m}, \mathbf{l}) = (\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor}, l))$ and

$$(\mathbf{m}, \mathbf{l}) = (\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor - 1}, \min\{k, l\}, \max\{l - k, 0\})),$$

$$\begin{aligned} & \frac{(\mu - \frac{d_2}{2})_{\underline{k}_{\lfloor r/2 \rfloor}, d} (\mu - \frac{d_2}{2}(r-1) - l)_{\min\{k, l\}}}{(\mu - \frac{d_2}{2})_{\underline{k}_{\lfloor r/2 \rfloor} - \underline{0}_{\lfloor r/2 \rfloor}, d} (\mu - \frac{d_2}{2}(r-1) - l)_{l-l}} = \left(\mu - \frac{d_2}{2}(r-1) - l \right)_{\min\{k, l\}}, \\ & \frac{(\mu - \frac{d_2}{2})_{\underline{k}_{\lfloor r/2 \rfloor}, d} (\mu - \frac{d_2}{2}(r-1) - l)_{\min\{k, l\}}}{(\mu - \frac{d_2}{2})_{\underline{k}_{\lfloor r/2 \rfloor} - (\underline{0}_{\lfloor r/2 \rfloor - 1}, \min\{k, l\}), d} (\mu - \frac{d_2}{2}(r-1) - l)_{l-\max\{l-k, 0\}}} \\ &= \left(\mu - \frac{d_2}{2}(r-2) + \max\{0, k-l\} \right)_{\min\{k, l\}} \end{aligned}$$

do not vanish simultaneously, and if $f(x_2) = (x_2|w_2)_{\mathfrak{n}_2^+}^l$ for some $w_2 \in \mathfrak{p}_2^+$ of rank 1, then by Proposition 4.3 (3) we have

$$F_{\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor}, l)}^-[f](z_1, z_2^{-1}) \neq 0, \quad F_{\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor - 1}, \min\{k, l\}, \max\{l-k, 0\})}^+[f](z_1, z_2^{-1}) \neq 0.$$

By the $K_1^{\mathbb{C}}$ -equivariance these hold for all non-zero $f(x_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$. Hence (4.11) is non-zero for all $\mu \in \mathbb{C}$ if $f(x_2) \neq 0$. \square

Proof of Theorem 4.7(2). Again let $\mu := -\lambda - 2k + \frac{n}{r} = -\lambda - 2k + 1 + \frac{d}{2}(r-1)$. By Theorem 4.7(1), for even r we have

$$\begin{aligned} & (\lambda)_{(2\underline{k}_{r/2-1}, 2k-l, \underline{k}_{r/2-1}, k-l), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \left(\lambda + k - \frac{d}{4}r + \frac{d_2}{2} \right)_{(k_{r/2-1}, k-l), d} F_{l, -}^{\downarrow} \left(\begin{matrix} -k \\ \mu \end{matrix}; f; z \right) \\ &= (-1)^{kr/2-l} \left(\mu - \frac{d_2}{2} + (l, \underline{0}_{r/2-1}) \right)_{(k-l, \underline{k}_{r/2-1}), d} \det_{\mathfrak{n}_2^+}(z_2)^k \\ &\quad \times \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \frac{(-k + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{0}_{\lceil r/2 \rceil - 1}), d} (-k - \frac{d_2}{2})_{\mathbf{m}, d}}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}), \end{aligned} \tag{4.12}$$

and for odd r we have

$$\begin{aligned} & (\lambda)_{(2\underline{k}_{\lfloor r/2 \rfloor}, \underline{k}_{\lfloor r/2 \rfloor}, k-l), d} \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} \\ &= \left(\lambda + k - \frac{d}{2} \left\lfloor \frac{r}{2} \right\rfloor + \frac{d_2}{2} + (k-l, \underline{0}_{\lfloor r/2 \rfloor}) \right)_{(l, \underline{k}_{\lfloor r/2 \rfloor - 1}, k-l), d} F_{l, -}^{\downarrow} \left(\begin{matrix} -k \\ \mu \end{matrix}; f; z \right) \\ &= (-1)^{k\lfloor r/2 \rfloor} \left(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lfloor r/2 \rfloor}) \right)_{(k-l, \underline{k}_{\lfloor r/2 \rfloor - 1}, l), d} \det_{\mathfrak{n}_2^+}(z_2)^k \\ &\quad \times \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \frac{(-k + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{0}_{\lceil r/2 \rceil - 1}), d} (-k - \frac{d_2}{2})_{\mathbf{m}, d}}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}). \end{aligned} \tag{4.13}$$

These are holomorphically continued for all $\mu \in \mathbb{C}$. Moreover, all non-zero $F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1})$ are linearly independent. When r is even, the coefficient of (4.12) for $(\mathbf{m}, \mathbf{l}) = ((\underline{k}_{r/2-1},$

$(k-l), (\underline{0}_{r/2-1}, l))$ is independent of μ and non-zero, and when r is odd, the coefficient of (4.13) for $(\mathbf{m}, \mathbf{l}) = (\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor}, l))$ is independent of μ and non-zero. Moreover, if $f(x_2) = (x_2|e')^l_{\mathfrak{n}_2^+}$ for some primitive idempotent $e' \in \mathfrak{n}_2^+$, then for even r , by Proposition 4.3 (1), (4) and Example 4.1 we have

$$\begin{aligned} & \det_{\mathfrak{n}_2^+}(z_2)^k F_{(\underline{k}_{r/2-1}, k-l), (\underline{0}_{r/2-1}, l)}^+ [(\cdot|e')^l_{\mathfrak{n}_2^+}] (z_1, z_2^{-1}) \\ &= \frac{1}{(-(k-l, \underline{k}_{r/2-1}))_{\underline{k}-l_{r/2}, d}} \det_{\mathfrak{n}^+}(z_1)^{k-l} \det_{\mathfrak{n}_2^+}(z_2)^l F_{(\underline{l}_{r/2-1}, 0), (\underline{0}_{r/2-1}, l)}^+ [(\cdot|e')^l_{\mathfrak{n}_2^+}] (z_1, z_2^{-1}) \\ &= \frac{(-k)_l}{(-k)_{(k, \underline{k}-l_{r/2-1}), d}} \frac{(-l)_{\underline{l}_{r/2-1}, d}}{(-l - \frac{d}{2})_{\underline{l}_{r/2-1}, d}} \det_{\mathfrak{n}^+}(z_1)^{k-l} \det_{\mathfrak{n}_2^{++}}(z_2^{\prime\prime})^l \tilde{\Phi}_{\underline{l}_{r/2-1}}^{\mathfrak{p}_1^{++}, \mathfrak{p}_2^{--}} (z_1^{\prime\prime}, (z_2^{\prime\prime})^{-1}) \\ &= \frac{(-k)_l}{(-k)_{\underline{k}_{r/2}, d}} \det_{\mathfrak{n}^+}(z_1)^{k-l} ((z_1^{\prime\prime})^\sharp | z_2^{\prime\prime})^l_{\mathfrak{n}_2^{++}} \neq 0, \end{aligned}$$

and for odd r , by Proposition 4.3 (2) we have

$$\begin{aligned} & \det_{\mathfrak{n}_2^+}(z_2)^k F_{(\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor}, l))}^+ [(\cdot|e')^l_{\mathfrak{n}_2^+}] (z_1, z_2^{-1}) \\ &= \frac{(-1)^l (-k)_l (-l)_l}{(-k)_{\underline{k}_{\lfloor r/2 \rfloor}, d} (-k - \frac{d_2}{2}(r-1))_l l!} (z_2 | (z_1)^\sharp)^{k-l}_{\mathfrak{n}_2^+} ((z_1)^\sharp | e')^l_{\mathfrak{n}_2^+} \\ &= \frac{(-k)_l}{(-k)_{(\underline{k}_{\lfloor r/2 \rfloor}, l), d}} (z_2 | (z_1)^\sharp)^{k-l}_{\mathfrak{n}_2^+} ((z_1)^\sharp | e')^l_{\mathfrak{n}_2^+} \neq 0, \end{aligned}$$

and by the $K_1^{\mathbb{C}}$ -equivariance,

$$\begin{aligned} F_{(\underline{k}_{r/2-1}, k-l), (\underline{0}_{r/2-1}, l)}^+ [f] (z_1, z_2^{-1}) &\neq 0 & (r: \text{even}), \\ F_{(\underline{k}_{\lfloor r/2 \rfloor}, (\underline{0}_{\lfloor r/2 \rfloor}, l))}^+ [f] (z_1, z_2^{-1}) &\neq 0 & (r: \text{odd}) \end{aligned}$$

hold for all non-zero $f(x_2) \in \mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+)$. Hence (4.13) is non-zero for all $\mu \in \mathbb{C}$ if $f(x_2) \neq 0$. \square

Proof of Theorem 4.7(3). Combining Theorem 4.7 (1) and (4.8), for $\operatorname{Re} \lambda > \frac{2n}{r} - 1$, $k, l \in \mathbb{Z}_{\geq 0}$, $k \geq l$ we have

$$\begin{aligned} C_{r,-}^{d,d_2}(\lambda, k, l) F_{l,-}^{\downarrow} \left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right) &= \frac{C_{r,-}^{d,d_2}(\lambda + 2(k-l), l, l)}{(\lambda)_{2(k-l), d}} \det_{\mathfrak{n}^+}(z)^{-\lambda + \frac{n}{r}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k-l} \\ &\quad \times \det_{\mathfrak{n}^+}(z)^{\lambda + 2(k-l) - \frac{n}{r}} F_{l,-}^{\downarrow} \left(\begin{matrix} -l \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right), \end{aligned}$$

that is,

$$\begin{aligned} F_{l,-}^{\downarrow} \left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right) &= \frac{\det_{\mathfrak{n}^+}(z)^{-\lambda + \frac{n}{r}}}{(\lambda - \frac{d}{4}r + \frac{d_2}{2} + (\underline{k}_{r-1}, k-l))_{\underline{k}-l_r, d_2}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k-l} \\ &\quad \times \det_{\mathfrak{n}^+}(z)^{\lambda + 2(k-l) - \frac{n}{r}} F_{l,-}^{\downarrow} \left(\begin{matrix} -l \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; f; z \right). \end{aligned}$$

Then this is meromorphically continued for all $\lambda \in \mathbb{C}$, and if $\lambda = \frac{n}{r} - a = \frac{d}{2}(r-1) + 1 - a$, $a = 1, 2, \dots, k-l$, then the left hand side of

$$\begin{aligned} \det_{\mathfrak{n}^+}(z)^{-a} F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right) &= \frac{1}{(\frac{d_2}{2}(r-1) + 1 - a + (\underline{k}_{r-1}, k-l))_{\underline{k}-l_r, d_2}} \det_{\mathfrak{n}_2^+} \left(\frac{\partial}{\partial z_2} \right)^{k-l} \\ &\times \det_{\mathfrak{n}^+}(z)^{2(k-l)-a} F_{l,-}^\downarrow \left(\begin{matrix} -l \\ a-2k \end{matrix}; f; z \right) \end{aligned}$$

becomes a polynomial (note that $d_2 = d/2$ or $r = 2$ hold). Hence $F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right)$ is divisible by $\det_{\mathfrak{n}^+}(z)^a$. Also, by $\mathcal{B}_{\frac{n}{r}-a} \det_{\mathfrak{n}^+}(z)^a = \det_{\mathfrak{n}^+}(z)^a \mathcal{B}_{\frac{n}{r}+a}$ (see the proof of [9, Proposition XV.2.4]), we have

$$(\mathcal{B}_{\frac{n}{r}+a})_1 \det_{\mathfrak{n}^+}(z)^{-a} F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right) = \det_{\mathfrak{n}^+}(z)^{-a} (\mathcal{B}_{\frac{n}{r}-a})_1 F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right) = 0,$$

and hence

$$\begin{aligned} &\left(\det_{\mathfrak{n}_2^+}(x_2)^{k-a} f(x_2^{-1}) \mapsto \det_{\mathfrak{n}_2^+}(z)^{-a} F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right) \right) \\ &\in \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(k-\underline{a}_{r-1}, k-l-a)}(\mathfrak{p}_2^+), \text{Sol}_{\mathcal{P}(\mathfrak{p}^+)}((\mathcal{B}_{\frac{n}{r}+a})_1)) \\ &\simeq \mathbb{C} \left(\det_{\mathfrak{n}_2^+}(x_2)^{k-a} f(x_2^{-1}) \mapsto F_{l,-}^\downarrow \left(\begin{matrix} a-k \\ a-2k \end{matrix}; f; z \right) \right) \end{aligned}$$

holds. Then comparing the values at $z_1 = 0$, for $a, k \in \mathbb{Z}_{\geq 0}$ with $a \leq k-l$, we get

$$F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right) = \det_{\mathfrak{n}^+}(z)^a F_{l,-}^\downarrow \left(\begin{matrix} a-k \\ a-2k \end{matrix}; f; z \right) \quad (4.14)$$

(see [35, Theorem 6.6] for more general case). Now we want to prove this for general $(a, k) \in \mathbb{C}^2$. We recall that the left hand side of (4.14) is given by

$$\begin{aligned} F_{l,-}^\downarrow \left(\begin{matrix} -k \\ a-2k \end{matrix}; f; z \right) &= \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l}} \frac{(-k + (l, 0_{\lceil r/2 \rceil-1}))_{\mathbf{m}+\mathbf{l}-(l, 0_{\lceil r/2 \rceil-1}), d} (-k - \frac{d_2}{2})_{\mathbf{m}, d}}{(a-2k - \frac{d_2}{2} + (l, 0_{\lceil r/2 \rceil-1}))_{\mathbf{m}+\mathbf{l}-(l, 0_{\lceil r/2 \rceil-1}), d}} \\ &\times \det_{\mathfrak{n}_2^+}(z_2)^k F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}). \end{aligned} \quad (4.15)$$

Similarly, by (4.2), the right hand side is computed as

$$\begin{aligned} &\det_{\mathfrak{n}^+}(z)^a F_{l,-}^\downarrow \left(\begin{matrix} a-k \\ a-2k \end{matrix}; f; z \right) \\ &= \sum_{\mathbf{m}' \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l}' \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}'|=l}} \frac{(a-k + (l, 0_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, 0_{\lceil r/2 \rceil-1}), d} (a-k - \frac{d_2}{2})_{\mathbf{m}', d}}{(a-2k - \frac{d_2}{2} + (l, 0_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, 0_{\lceil r/2 \rceil-1}), d}} \\ &\times \det_{\mathfrak{n}_2^+}(z_2)^{k-a} F_{\mathbf{m}', \mathbf{l}'}^+[f](z_1, z_2^{-1}) \det_{\mathfrak{n}_2^+}(z_2)^a h_{\mathfrak{n}^+}(z_1, P(z_2^{-1}) z_1)^{a/2} \\ &= \sum_{\mathbf{m}' \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l}' \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}'|=l}} \frac{(a-k + (l, 0_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, 0_{\lceil r/2 \rceil-1}), d} (a-k - \frac{d_2}{2})_{\mathbf{m}', d}}{(a-2k - \frac{d_2}{2} + (l, 0_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, 0_{\lceil r/2 \rceil-1}), d}} \\ &\times \det_{\mathfrak{n}_2^+}(z_2)^k F_{\mathbf{m}', \mathbf{l}'}^+[f](z_1, z_2^{-1}) \sum_{\mathbf{m}'' \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} (-a)_{\mathbf{m}'', d} \tilde{\Phi}_{\mathbf{m}''}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1}). \end{aligned}$$

Then we have

$$\begin{aligned}
& (f \mapsto F_{\mathbf{m}', \mathbf{l}'}^+[f](z_1, y_2) \tilde{\Phi}_{\mathbf{m}''}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(z_1, y_2)) \\
& \in \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(l, \underline{0}_{r-1})}(\mathbf{p}_2^-), \mathcal{P}_{\langle \mathbf{m}' \rangle}(\mathbf{p}_1^+) \otimes \mathcal{P}_{\phi_+(\mathbf{m}', \mathbf{l}')}(\mathbf{p}_2^-) \otimes \mathcal{P}_{\langle \mathbf{m}'' \rangle}(\mathbf{p}_1^+) \otimes \mathcal{P}_{\mathbf{m}''^2}(\mathbf{p}_2^-)) \\
& \subset \bigoplus_{\substack{|\mathbf{k}|=|\langle \mathbf{m}' \rangle|+|\langle \mathbf{m}'' \rangle| \\ k_j \geq \max\{\langle \mathbf{m}' \rangle_j, \langle \mathbf{m}'' \rangle_j\}}} \bigoplus_{\substack{\mathbf{k}' \in \mathbb{Z}_{++}^{r-1} \\ |\mathbf{k}'|=|\phi_+(\mathbf{m}', \mathbf{l}')|+|\mathbf{m}''^2| \\ k'_j \geq \max\{\phi_+(\mathbf{m}', \mathbf{l}')_j, (\mathbf{m}''^2)_j\}}} \text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(l, \underline{0}_{r-1})}(\mathbf{p}_2^-), \mathcal{P}_{\mathbf{k}}(\mathbf{p}_1^+) \otimes \mathcal{P}_{\mathbf{k}'}(\mathbf{p}_2^-)),
\end{aligned}$$

where $\mathbf{k} \in \mathbb{Z}_{++}^{\lfloor r/\varepsilon_1 \rfloor}$ (Case 1 ($n' \neq 2$), Case 2) or $\mathbf{k} \in (\mathbb{Z}_{++}^{\lfloor r/2 \rfloor})^2$ (Case 1 ($n' = 2$), Case 3). Now since $\text{Hom}_{K_1^{\mathbb{C}}}(\mathcal{P}_{(l, \underline{0}_{r-1})}(\mathbf{p}_2^-), \mathcal{P}_{\mathbf{k}}(\mathbf{p}_1^+) \otimes \mathcal{P}_{\mathbf{k}'}(\mathbf{p}_2^-)) \neq \{0\}$ implies $(\mathbf{k}, \mathbf{k}') = (\langle \mathbf{m} \rangle, \phi_+(\mathbf{m}, \mathbf{l}))$ for some $(\mathbf{m}, \mathbf{l}) \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor} \times (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil}$ satisfying $|\mathbf{l}| = l$ and (4.3), we set

$$\begin{aligned}
F_{\mathbf{m}', \mathbf{l}', \mathbf{m}'', \mathbf{m}, \mathbf{l}}[f](z_1, y_2) &:= \text{Proj}_{\phi_+(\mathbf{m}, \mathbf{l}), y_2}^{\mathbf{p}_2^-}(F_{\mathbf{m}', \mathbf{l}'}^+[f](z_1, y_2) \tilde{\Phi}_{\mathbf{m}''}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(z_1, y_2)) \\
&\in \mathcal{P}_{\langle \mathbf{m} \rangle}(\mathbf{p}_1^+)_{z_1} \otimes \mathcal{P}_{\phi_+(\mathbf{m}, \mathbf{l})}(\mathbf{p}_2^-)_{y_2},
\end{aligned}$$

so that

$$F_{\mathbf{m}', \mathbf{l}'}^+[f](z_1, y_2) \tilde{\Phi}_{\mathbf{m}''}^{\mathbf{p}_1^+, \mathbf{p}_2^-}(z_1, y_2) = \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor} \\ |\mathbf{m}|=|\mathbf{m}'|+|\mathbf{m}''| \\ m_j \geq \max\{m'_j, m''_j\}}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}|=l \\ m_j + l_j \geq m'_j + l'_j}} F_{\mathbf{m}', \mathbf{l}', \mathbf{m}'', \mathbf{m}, \mathbf{l}}[f](z_1, y_2).$$

Then the right hand side of (4.14) becomes

$$\begin{aligned}
& \det_{\mathfrak{n}^+}(z)^a F_{l, -}^{\downarrow} \left(\begin{matrix} a-k \\ a-2k \end{matrix}; f; z \right) \\
& = \det_{\mathfrak{n}_2^+}(z_2)^k \sum_{\substack{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor} \\ |\mathbf{l}'|=l}} \sum_{\substack{\mathbf{m}', \mathbf{m}'' \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor} \\ |\mathbf{m}'|+|\mathbf{m}''|=|\mathbf{m}| \\ m'_j, m''_j \leq m_j}} \sum_{\substack{\mathbf{l}' \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}'|=l \\ m'_j + l'_j \leq m_j + l_j}} \sum_{\substack{(-a)_{\mathbf{m}'', d} \\ |\mathbf{l}'|=l \\ m'_j + l'_j \leq m_j + l_j}} (-a)_{\mathbf{m}'', d} \\
& \quad \times \frac{(a-k+(l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, \underline{0}_{\lceil r/2 \rceil-1}), d} (a-k-\frac{d_2}{2})_{\mathbf{m}', d}}{(a-2k-\frac{d_2}{2}+(l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, \underline{0}_{\lceil r/2 \rceil-1}), d}} F_{\mathbf{m}', \mathbf{l}', \mathbf{m}'', \mathbf{m}, \mathbf{l}}[f](z_1, z_2^{-1}). \tag{4.16}
\end{aligned}$$

If $a, k \in \mathbb{Z}_{\geq 0}$, $a \leq k-l$, then in (4.15) only $m_1 + l_1 \leq k$ terms remain, and both in (4.15) and (4.16), all denominators in the terms with $m_1 + l_1 \leq k$ are non-zero. Hence for $m_1 + l_1 \leq k$, by comparing the both formulas we get

$$\begin{aligned}
& \frac{(-k+(l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}+1-(l, \underline{0}_{\lceil r/2 \rceil-1}), d} (-k-\frac{d_2}{2})_{\mathbf{m}, d}}{(a-2k-\frac{d_2}{2}+(l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}+1-(l, \underline{0}_{\lceil r/2 \rceil-1}), d}} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}) \\
& = \sum_{\substack{\mathbf{m}', \mathbf{m}'' \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor} \\ |\mathbf{m}'|+|\mathbf{m}''|=|\mathbf{m}| \\ m'_j, m''_j \leq m_j}} \sum_{\substack{\mathbf{l}' \in (\mathbb{Z}_{\geq 0})^{\lceil r/2 \rceil} \\ |\mathbf{l}'|=l \\ m'_j + l'_j \leq m_j + l_j}} (-a)_{\mathbf{m}'', d} \\
& \quad \times \frac{(a-k+(l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, \underline{0}_{\lceil r/2 \rceil-1}), d} (a-k-\frac{d_2}{2})_{\mathbf{m}', d}}{(a-2k-\frac{d_2}{2}+(l, \underline{0}_{\lceil r/2 \rceil-1}))_{\mathbf{m}'+\mathbf{l}'-(l, \underline{0}_{\lceil r/2 \rceil-1}), d}} F_{\mathbf{m}', \mathbf{l}', \mathbf{m}'', \mathbf{m}, \mathbf{l}}[f](z_1, z_2^{-1}).
\end{aligned}$$

Then since $\{(a, k) \in (\mathbb{Z}_{\geq 0})^2 \mid k \geq a + l, k \geq m_1 + l_1\} \subset \mathbb{C}^2$ is Zariski dense, this holds for all $(a, k) \in \mathbb{C}^2$ without the assumption $m_1 + l_1 \leq k$, and hence (4.14) holds for all $(a, k) \in \mathbb{C}^2$. This completes the proof of Theorem 4.7(3). \square

Example 4.10. Suppose $l = 0$, $f(x_2) = 1 \in \mathcal{P}_{(0, \dots, 0)}(\mathfrak{p}_2^+)$. Then we have

$$F_{\mathbf{m}, \underline{\Omega}_{\lceil r/2 \rceil}}^-[1](z_1, z_2^{-1}) = F_{\mathbf{m}, \underline{\Omega}_{\lceil r/2 \rceil}}^+[1](z_1, z_2^{-1}) = \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1}),$$

$$F_{0, \pm}^\downarrow \left(\frac{\nu}{\mu}; 1; z \right) = \det_{\mathfrak{n}_2^+}(z_2)^{-\nu} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \frac{(\nu)_{\mathbf{m}, d} (\nu - \frac{d_2}{2})_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1})}{(\mu - \frac{d_2}{2})_{\mathbf{m}, d}}.$$

Especially, for $k \in \mathbb{Z}_{\geq 0}$, $\operatorname{Re} \lambda > \frac{2n}{r} - 1$, we have

$$\begin{aligned} & \left\langle \det_{\mathfrak{n}_2^+}(x_2)^k, e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda, x} = C_{r, \pm}^{d, d_2}(\lambda, k, 0) F_{0, \pm}^\downarrow \left(-k, -\lambda - 2k + \frac{n}{r}; 1; z \right) \\ &= \frac{(\lambda + k - \frac{d}{2} \lceil \frac{r}{2} \rceil + \frac{d_2}{2})_{\underline{k}_{\lceil r/2 \rceil}, d}}{(\lambda)_{(2k_{\lceil r/2 \rceil}, \underline{k}_{\lceil r/2 \rceil}), d}} \det_{\mathfrak{n}_2^+}(z_2)^k \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \frac{(-k)_{\mathbf{m}, d} (-k - \frac{d_2}{2})_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1})}{(-\lambda - 2k + \frac{n}{r} - \frac{d_2}{2})_{\mathbf{m}, d}}. \end{aligned}$$

In addition, Theorems 4.6(3), 4.7(3) are equivalent to

$$\begin{aligned} & \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \frac{(\nu)_{\mathbf{m}, d} (\nu - \frac{d_2}{2})_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1})}{(\mu - \frac{d_2}{2})_{\mathbf{m}, d}} \\ &= h_{\mathfrak{n}^+}(z_1, P(z_2)^{-1} z_1)^{(\mu - 2\nu)/2} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \frac{(\mu - \nu)_{\mathbf{m}, d} (\mu - \nu - \frac{d_2}{2})_{\mathbf{m}, d} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^+, \mathfrak{p}_2^-}(z_1, z_2^{-1})}{(\mu - \frac{d_2}{2})_{\mathbf{m}, d}}. \end{aligned}$$

Example 4.11. Suppose $r = 2$. We fix a Jordan frame $\{e', e''\} \subset \mathfrak{n}_2^+$, and consider $f(x_2) = (x_2|e')_{\mathfrak{n}_2^+}^l$ or $f(y_2) = (y_2|e'')_{\mathfrak{n}_2^+}^l$. Then we have

$$\begin{aligned} F_{m, l}^-[(\cdot|e')_{\mathfrak{n}_2^+}^l](z_1, z_2^{-1}) &= \frac{1}{m!} \left(-\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right)^m (z_2|e')_{\mathfrak{n}_2^+}^l, \\ F_{m, l}^+[(\cdot|e'')_{\mathfrak{n}_2^+}^l](z_1, z_2^{-1}) &= \frac{1}{m!} \left(-\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right)^m (z_2^{-1}|e'')_{\mathfrak{n}_2^+}^l, \\ F_{l, +}^\downarrow \left(\frac{\nu}{\mu}; (\cdot|e')_{\mathfrak{n}_2^+}^l; z \right) &= \det_{\mathfrak{n}^+}(z_2)^{-\nu} \sum_{m=0}^{\infty} \frac{(\nu)_m (\nu - l - \frac{d_2}{2})_m}{(\mu - l - \frac{d_2}{2})_m m!} \left(-\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right)^m (z_2|e')_{\mathfrak{n}_2^+}^l \\ &= \det_{\mathfrak{n}^+}(z_2)^{-\nu} {}_2F_1 \left(\nu, \nu - l - \frac{d_2}{2}; \mu - l - \frac{d_2}{2}; -\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right) (z_2|e')_{\mathfrak{n}_2^+}^l, \\ F_{l, -}^\downarrow \left(\frac{\nu}{\mu}; (\cdot|e'')_{\mathfrak{n}_2^+}^l; z \right) &= \det_{\mathfrak{n}^+}(z_2)^{-\nu} \sum_{m=0}^{\infty} \frac{(\nu + l)_m (\nu - \frac{d_2}{2})_m}{(\mu + l - \frac{d_2}{2})_m m!} \left(-\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right)^m (z_2^{-1}|e'')_{\mathfrak{n}_2^+}^l \\ &= \det_{\mathfrak{n}^+}(z_2)^{-\nu - l} {}_2F_1 \left(\nu + l, \nu - \frac{d_2}{2}; \mu + l - \frac{d_2}{2}; -\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right) (z_2|e')_{\mathfrak{n}_2^+}^l. \end{aligned}$$

Especially, for $\mathbf{k} \in \mathbb{Z}_{++}^2$, $\operatorname{Re} \lambda > \frac{2n}{r} - 1$, we have

$$\begin{aligned}
& \left\langle \Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}(x_2), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} \\
&= \left\langle (x_2|e')_{\mathfrak{n}_2^+}^{k_1-k_2} \det_{\mathfrak{n}_2^+}(x_2)^{k_2}, e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} = \left\langle (x_2^{-1}|e'')_{\mathfrak{n}_2^+}^{k_1-k_2} \det_{\mathfrak{n}_2^+}(x_2)^{k_1}, e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} \\
&= C_{2,+}^{d,d_2}(\lambda, k_2, k_1 - k_2) F_{k_1-k_2,+}^{\downarrow} \left(\begin{matrix} -k_2 \\ -\lambda - 2k_2 + \frac{n}{2}; (\cdot|e')_{\mathfrak{n}_2^+}^{k_1-k_2}; z \end{matrix} \right) \\
&= C_{2,-}^{d,d_2}(\lambda, k_1, k_1 - k_2) F_{k_1-k_2,-}^{\downarrow} \left(\begin{matrix} -k_1 \\ -\lambda - 2k_1 + \frac{n}{2}; (\cdot|e'')_{\mathfrak{n}_2^+}^{k_1-k_2}; z \end{matrix} \right) \\
&= \frac{(\lambda + k_1 - \frac{d-d_2}{2})_{k_2}}{(\lambda)_{(k_1+k_2,k_2),d}} {}_2F_1 \left(\begin{matrix} -k_2, -k_1 - \frac{d_2}{2} \\ -\lambda - k_1 - k_2 + \frac{d-d_2}{2} + 1 \end{matrix}; -\frac{\det_{\mathfrak{n}^+}(z_1)}{\det_{\mathfrak{n}^+}(z_2)} \right) \Delta_{\mathbf{k}}^{\mathfrak{n}_2^+}(z_2).
\end{aligned}$$

Example 4.12. Suppose $r \geq 3$, $-\nu = k = l \in \mathbb{Z}_{\geq 0}$, $f(y_2) \in \mathcal{P}_{(l,\underline{0}_{r-1})}(\mathfrak{p}_2^+)$. Then since

$$\begin{aligned}
& (-l + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}) \\
&= ((0, \underline{-l}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l} - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d} F_{\mathbf{m}, \mathbf{l}}^+[f](z_1, z_2^{-1}) \neq 0
\end{aligned}$$

implies $m_1 + l_1 = l$ and $\phi_+(\mathbf{m}, \mathbf{l}) \in \mathbb{Z}_+^r$, that is,

$$\begin{aligned}
m_1 &\leq l, \quad m_{\lceil r/2 \rceil} = 0 \quad (\text{for even } r), \\
\mathbf{l} = \mathbf{l}(\mathbf{m}) &:= (l - m_1, m_1 - m_2, m_2 - m_3, \dots, m_{\lceil r/2 \rceil - 2} - m_{\lceil r/2 \rceil - 1}, m_{\lceil r/2 \rceil - 1}),
\end{aligned}$$

so that $\phi_+(\mathbf{m}, \mathbf{l}(\mathbf{m})) = (l, \mathbf{m}^2)$, $\mathbf{m} + \mathbf{l}(\mathbf{m}) = (l, \mathbf{m})$, we have

$$\begin{aligned}
& F_{l,-}^{\downarrow} \left(\begin{matrix} -l \\ \mu \end{matrix}; f; z \right) \\
&= \det_{\mathfrak{n}_2^+}(z_2)^l \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lceil r/2 \rceil - 1}} \frac{((0, \underline{-l}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l}(\mathbf{m}) - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d} (-l - \frac{d_2}{2})_{\mathbf{m}, d} F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+[f](z_1, z_2^{-1})}{(\mu - \frac{d_2}{2} + (l, \underline{0}_{\lceil r/2 \rceil - 1}))_{\mathbf{m} + \mathbf{l}(\mathbf{m}) - (l, \underline{0}_{\lceil r/2 \rceil - 1}), d}} \\
&= \det_{\mathfrak{n}_2^+}(z_2)^l \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lceil r/2 \rceil - 1}} \frac{(-l - \frac{d}{2})_{\mathbf{m}, d} (-l - \frac{d_2}{2})_{\mathbf{m}, d}}{(\mu - \frac{d+d_2}{2})_{\mathbf{m}, d}} F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+[f](z_1, z_2^{-1}),
\end{aligned}$$

and hence for $\operatorname{Re} \lambda > \frac{2n}{r} - 1$ we have

$$\begin{aligned}
& \left\langle \det_{\mathfrak{n}_2^+}(x_2)^l f(x_2^{-1}), e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} = C_{r,-}^{d,d_2}(\lambda, l, l) F_{l,-}^{\downarrow} \left(\begin{matrix} -l \\ -\lambda - 2l + \frac{n}{r}; f; z \end{matrix} \right) \\
&= \frac{(\lambda + l - \frac{d}{2} \lfloor \frac{r}{2} \rfloor + \frac{d_2}{2})_{l_{\lceil r/2 \rceil - 1}, d}}{(\lambda)_{(2l_{\lceil r/2 \rceil - 1}, l_{\lceil r/2 \rceil}), d}} \\
&\quad \times \det_{\mathfrak{n}_2^+}(z_2)^l \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lceil r/2 \rceil - 1}} \frac{(-l - \frac{d}{2})_{\mathbf{m}, d} (-l - \frac{d_2}{2})_{\mathbf{m}, d}}{(-\lambda - 2l + \frac{d}{2}(r-2) + 1 - \frac{d_2}{2})_{\mathbf{m}, d}} F_{\mathbf{m}, \mathbf{l}(\mathbf{m})}^+[f](z_1, z_2^{-1}).
\end{aligned}$$

Especially, we fix a primitive idempotent $e' \in \mathfrak{n}_2^+$ and let $\mathfrak{p}_2^+(e')_0 =: \mathfrak{p}_2^{+''}$, $\mathfrak{p}^+(e')_0 \cap \mathfrak{p}_1^+ =: \mathfrak{p}_1^{+''}$ be as in (2.2). If $f(y_2) = (y_2|e')_{\mathfrak{n}_2^+}^l$ so that $\det_{\mathfrak{n}_2^+}(x_2)^l f(x_2^{-1}) = \det_{\mathfrak{n}_2^{+''}}(x_2^{''})^l$, then by

Proposition 4.3(4) we have

$$\begin{aligned} & \left\langle \det_{\mathfrak{n}_2^{+''}}(x_2'')^l, e^{(x|\bar{z})_{\mathfrak{p}^+}} \right\rangle_{\lambda,x} = C_{r,-}^{d,d_2}(\lambda, l, l) F_{l,-}^{\downarrow} \left(-l; -\lambda - 2l + \frac{n}{r}; (\cdot|e')_{\mathfrak{n}_2^+}^l; z \right) \\ &= \frac{(\lambda + l - \frac{d}{2}\lfloor \frac{r}{2} \rfloor + \frac{d_2}{2})_{\lfloor r/2 \rfloor - 1, d}}{(\lambda)_{(2\lfloor r/2 \rfloor - 1, l_{\lfloor r/2 \rfloor}), d}} \\ & \quad \times \det_{\mathfrak{n}_2^{+''}}(z_2'')^l \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor - 1}} \frac{(-l)_{\mathbf{m}, d} (-l - \frac{d_2}{2})_{\mathbf{m}, d}}{(-\lambda - 2l + \frac{d}{2}(r-2) + 1 - \frac{d_2}{2})_{\mathbf{m}, d}} \tilde{\Phi}_{\mathbf{m}}^{\mathfrak{p}_1^{+''}, \mathfrak{p}_2^{-''}}(z_1'', (z_2'')^{-1}). \end{aligned}$$

This fits with Example 4.10 applied for $\mathfrak{p}_2^{+''}$.

4.3 Results on restriction of $\mathcal{H}_\lambda(D)$ to subgroups

In this subsection we consider the decomposition of $\mathcal{H}_\lambda(D)$ under \tilde{G}_1 . Let $V_{\mathbf{k}}$ be an irreducible K_1 -module isomorphic to $\mathcal{P}_{\mathbf{k}}(\mathfrak{p}_2^+)$. Then $\mathcal{H}_\lambda(D)$ is decomposed under \tilde{G}_1 as

$$\mathcal{H}_\lambda(D)|_{\tilde{G}_1} \simeq \sum_{\mathbf{k} \in \mathbb{Z}_{++}^r}^{\oplus} \mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}}).$$

For each case, $\mathcal{H}_\lambda(D)$ has the minimal \tilde{K} -type

$$\chi^{-\lambda} \simeq \begin{cases} \mathbb{C}_{-\lambda} \boxtimes V_{\underline{\Omega}_{\lfloor n/2 \rfloor}}^{[n]^\vee} & (G = SO_0(2, n)), \\ V_{\lambda/2}^{(r)\vee} \boxtimes V_{\underline{\lambda/2}_r}^{(r)} \simeq V_{\underline{\lambda}_r}^{(r)\vee} \boxtimes V_{\underline{\Omega}_r}^{(r)} \simeq V_{\underline{\Omega}_r}^{(r)\vee} \boxtimes V_{\underline{\lambda}_r}^{(r)} & (G = SU(r, r)), \\ V_{\underline{\lambda/2}_{2r}}^{(2r)\vee} & (G = SO^*(4r)), \end{cases}$$

and $\mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{\mathbf{k}})$ has the minimal \tilde{K}_1 -type

$$\begin{aligned} & \chi_1^{-\varepsilon_1 \lambda} \otimes V_{\mathbf{k}} \\ & \simeq \begin{cases} \mathbb{C}_{-(\lambda+k_1+k_2)} \boxtimes V_{\underline{\Omega}_{\lfloor n'/2 \rfloor}}^{[n']^\vee} \boxtimes V_{(k_1-k_2, \underline{\Omega}_{\lfloor n''/2 \rfloor - 1})}^{[n'']^\vee} & (G_1 = SO_0(2, n') \times SO(n''), n' \geq 3), \\ \mathbb{C}_{-(\lambda+k_1+k_2)} \boxtimes \mathbb{C}_{-(\lambda+k_1+k_2)} \boxtimes V_{(k_1-k_2, \underline{\Omega}_{\lfloor n''/2 \rfloor - 1})}^{[n'']^\vee} & (G_1 = SO_0(2, n') \times SO(n''), n' = 2), \\ \mathbb{C}_{-2(\lambda+k_1+k_2)} \boxtimes V_{(k_1-k_2, \underline{\Omega}_{\lfloor n''/2 \rfloor - 1})}^{[n'']^\vee} & (G_1 = SO_0(2, n') \times SO(n''), n' = 1), \\ V_{\underline{\lambda}_r + 2\mathbf{k}}^{(r)\vee} & (G_1 = SO^*(2r)), \\ V_{\underline{\lambda/2}_r + \mathbf{k}}^{(r)\vee} \boxtimes V_{\underline{\lambda/2}_r + \mathbf{k}}^{(r)\vee} & (G_1 = SO^*(2r) \times SO^*(2r)) \end{cases} \end{aligned}$$

(note that $SO_0(2, 2) \simeq SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$, $SO_0(2, 1) \simeq SL(2, \mathbb{R})$ hold up to covering). In this section we only deal with the cases $\mathbf{k} = (k+l, \underline{\Omega}_{r-1})$ or $\mathbf{k} = (\underline{\Omega}_{r-1}, k-l)$. Let $V_l := V_{(l, \underline{\Omega}_{r-1})}$, and let V_l^\vee be its contragredient. Then we have

$$\mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{(k+l, \underline{\Omega}_{r-1})}) \simeq \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l), \quad \mathcal{H}_{\varepsilon_1 \lambda}(D_1, V_{(\underline{\Omega}_{r-1}, k-l)}) \simeq \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l^\vee).$$

To describe the intertwining operators, for $\mu \in \mathbb{C}$, $l \in \mathbb{Z}_{\geq 0}$ and for $f(x_2) \in \mathcal{P}_{(l, \underline{\Omega}_{r-1})}(\mathfrak{p}_2^+)$, we define a function $F_{l,+}^\uparrow(\bar{\mu}; f; y_1, x_2)$ on $\mathfrak{p}_1^- \oplus \mathfrak{p}_2^+$ by

$$F_{l,+}^\uparrow(\bar{\mu}; f; y_1, x_2) := \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lfloor r/2 \rfloor} \\ |\mathbf{l}|=l}} \frac{F_{\mathbf{m}, \mathbf{l}}^+[f](y_1, x_2)}{(\mu - \frac{\delta}{2} + (l, \underline{\Omega}_{\lfloor r/2 \rfloor - 1}))_{\mathbf{m}+\mathbf{l}-(l, \underline{\Omega}_{\lfloor r/2 \rfloor - 1}), d}}, \quad (4.17)$$

where δ is given in (4.1), and define a function $F_{l,-}^\uparrow(\underline{\mu}; f; y_1, x_2)$ on $\mathfrak{p}_1^- \oplus \mathfrak{p}_2^+$ by

$$F_{l,-}^\uparrow\left(\underline{\mu}; f; y_1, x_2\right) := \begin{cases} \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lfloor r/2 \rfloor} \\ |\mathbf{l}|=l}} \frac{F_{\mathbf{m}, \mathbf{l}}^-[f](y_1, x_2)}{(\mu - \frac{\delta}{2} - (0_{\lfloor r/2 \rfloor-1}, l))_{\mathbf{m}-\mathbf{l}+(0_{\lfloor r/2 \rfloor-1}, l), d}} & (r: \text{even}), \\ \sum_{\mathbf{m} \in \mathbb{Z}_{++}^{\lfloor r/2 \rfloor}} \sum_{\substack{\mathbf{l} \in (\mathbb{Z}_{\geq 0})^{\lfloor r/2 \rfloor} \\ |\mathbf{l}|=l}} \frac{F_{\mathbf{m}, \mathbf{l}}^-[f](y_1, x_2)}{(\mu - \frac{\delta}{2})_{\mathbf{m}-\mathbf{l}', d}(\mu - l - \frac{\delta}{2} - \frac{d}{2}(r-1))_{l-l_{\lfloor r/2 \rfloor}}} & (r: \text{odd}), \end{cases} \quad (4.18)$$

where for odd r let $\mathbf{l}' := (l_1, \dots, l_{\lfloor r/2 \rfloor})$.

Theorem 4.13. *Let $\operatorname{Re} \lambda > \frac{2n}{r} - 1$, $k, l \in \mathbb{Z}_{\geq 0}$.*

- (1) ([33, Theorems 5.7(3), 5.17(2), 5.19(1)]). *Let $K_l(x_2) \in (\mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^+) \otimes \operatorname{Hom}(V_l, \mathbb{C}))^{K_1}$. Then the linear map*

$$\begin{aligned} \mathcal{F}_{\lambda, k, l, +}^\uparrow: \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l)_{\tilde{K}_1} &\longrightarrow \mathcal{H}_\lambda(D)_{\tilde{K}}|_{(\mathfrak{g}_1, \tilde{K}_1)}, \\ (\mathcal{F}_{\lambda, k, l, +}^\uparrow f)(x) &:= \det_{\mathfrak{n}^+}(x_2)^k F_{l,+}^\uparrow\left(\begin{matrix} - \\ \lambda + 2k \end{matrix}; K_l; \frac{1}{\varepsilon_1} \frac{\partial}{\partial x_1}, x_2\right) f(x_1) \end{aligned}$$

intertwines the $(\mathfrak{g}_1, \tilde{K}_1)$ -action, where $F_{l,+}^\uparrow$ is as in (4.17).

- (2) ([34, Theorem 8.5]). *Let $K_l^\vee(y_2) \in (\mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^-) \otimes V_l)^{K_1}$. Then the linear map*

$$\begin{aligned} \mathcal{F}_{\lambda, k, l, +}^\downarrow: \mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\longrightarrow \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l), \\ (\mathcal{F}_{\lambda, k, l, +}^\downarrow f)(x_1) &:= F_{l,+}^\downarrow\left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r} \end{matrix}; K_l^\vee; \frac{\partial}{\partial x}\right) f(x)\Big|_{x_2=0} \end{aligned}$$

intertwines the \tilde{G}_1 -action, where $F_{l,+}^\downarrow$ is as in (4.4).

- (3) ([34, Theorem 8.5]). *Suppose $c_{k,l,+}$, $c_{k,l,+}^\vee \in \mathbb{C}$ satisfy, for $v \in V_l$,*

$$\begin{aligned} \|\det_{\mathfrak{n}_2^+}(x_2)^k K_l(x_2)v\|_{F, x_2}^2 &= c_{k,l,+}|v|_{V_l}^2, \\ \|\det_{\mathfrak{n}_2^+}(x_2)^k(v, K_l^\vee(\overline{x_2}))\|_{F, x_2}^2 &= c_{k,l,+}^\vee|v|_{V_l}^2. \end{aligned}$$

Then the operator norms of $\mathcal{F}_{\lambda, k, l, +}^\uparrow$, $\mathcal{F}_{\lambda, k, l, +}^\downarrow$ are given by

$$c_{k,l,+}^{-1} \|\mathcal{F}_{\lambda, k, l, +}^\uparrow\|_{\operatorname{op}}^2 = c_{k,l,+}^\vee \|\mathcal{F}_{\lambda, k, l, +}^\downarrow\|_{\operatorname{op}}^{-2} = C_{r,+}^{d,d_2}(\lambda, k, l),$$

where $C_{r,+}^{d,d_2}(\lambda, k, l)$ is as in (4.6).

Theorem 4.14. *Let $\operatorname{Re} \lambda > \frac{2n}{r} - 1$, $k, l \in \mathbb{Z}_{\geq 0}$ with $k \geq l$.*

- (1) ([33, Theorems 5.7(4), 5.19(1), 7.2]). *Let $K_l^\vee(y_2) \in (\mathcal{P}_{(l, \underline{0}_{r-1})}(\mathfrak{p}_2^-) \otimes \operatorname{Hom}(V_l^\vee, \mathbb{C}))^{K_1}$. Then the linear map*

$$\begin{aligned} \mathcal{F}_{\lambda, k, l, -}^\uparrow: \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l^\vee)_{\tilde{K}_1} &\longrightarrow \mathcal{H}_\lambda(D)_{\tilde{K}}|_{(\mathfrak{g}_1, \tilde{K}_1)}, \\ (\mathcal{F}_{\lambda, k, l, -}^\uparrow f)(x) &:= \det_{\mathfrak{n}^+}(x_2)^k F_{l,-}^\uparrow\left(\begin{matrix} - \\ \lambda + 2k \end{matrix}; K_l^\vee; \frac{1}{\varepsilon_1} \frac{\partial}{\partial x_1}, x_2\right) f(x_1) \end{aligned}$$

intertwines the $(\mathfrak{g}_1, \tilde{K}_1)$ -action, where $F_{l,-}^\uparrow$ is as in (4.18).

(2) Let $K_l(x_2) \in (\mathcal{P}_{(l, \underline{k}_{r-1})}(\mathfrak{p}_2^+) \otimes V_l^\vee)^{K_1}$. Then the linear map

$$\begin{aligned}\mathcal{F}_{\lambda, k, l, -}^\downarrow : \mathcal{H}_\lambda(D)|_{\tilde{G}_1} &\longrightarrow \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l^\vee), \\ (\mathcal{F}_{\lambda, k, l, -}^\downarrow f)(x_1) &:= F_{l, -}^\downarrow \left(\begin{matrix} -k \\ -\lambda - 2k + \frac{n}{r}; K_l^\vee; \frac{\partial}{\partial x} \end{matrix} \right) f(x) \Big|_{x_2=0}\end{aligned}$$

intertwines the \tilde{G}_1 -action, where $F_{l, -}^\downarrow$ is as in (4.5).

(3) ([35, Corollary 6.7]). Suppose $c_{k, l, -}, c_{k, l, -}^\vee \in \mathbb{R}_{>0}$ satisfy, for $v \in V_l^\vee$,

$$\begin{aligned}\|\det_{\mathfrak{n}_2^+}(x_2)^k K_l^\vee(x_2^{-1})v\|_{F, x_2}^2 &= c_{k, l, -}|v|_{V_l^\vee}^2, \\ \|\det_{\mathfrak{n}_2^+}(x_2)^k(v, K_l(\overline{x_2^{-1}}))\|_{F, x_2}^2 &= c_{k, l, -}^\vee|v|_{V_l^\vee}^2.\end{aligned}$$

Then the operator norms of $\mathcal{F}_{\lambda, k, l, -}^\uparrow, \mathcal{F}_{\lambda, k, l, -}^\downarrow$ are given by

$$c_{k, l, -}^{-1} \|\mathcal{F}_{\lambda, k, l, -}^\uparrow\|_{\text{op}}^2 = c_{k, l, -}^\vee \|\mathcal{F}_{\lambda, k, l, -}^\downarrow\|_{\text{op}}^{-2} = C_{r, -}^{d, d_2}(\lambda, k, l),$$

where $C_{r, -}^{d, d_2}(\lambda, k, l)$ is as in (4.7).

The differential operators $\mathcal{F}_{\lambda, k, l, \pm}^\downarrow$ are meromorphically continued for all $\lambda \in \mathbb{C}$, and give \tilde{G}_1 -intertwining operators

$$\mathcal{F}_{\lambda, k, l, +}^\downarrow : \mathcal{O}_\lambda(D)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\varepsilon_1(\lambda+2k)}(D_1, V_l), \quad \mathcal{F}_{\lambda, k, l, -}^\downarrow : \mathcal{O}_\lambda(D)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\varepsilon_1(\lambda+2k)}(D_1, V_l^\vee)$$

for all $\lambda \in \mathbb{C}$ except for poles. Then by Theorems 4.6 (3), 4.7 (3), for special λ the following holds.

Theorem 4.15 ([34, Theorem 8.9 (1)], [35, Corollary 6.8 (4)]). *Let $k, l \in \mathbb{Z}_{\geq 0}$. For $a = 1, 2, \dots, k$ we have*

$$\mathcal{F}_{\frac{n}{r}-a, k, l, +}^\downarrow = \mathcal{F}_{\frac{n}{r}+a, k-a, l, +}^\downarrow \circ \det_{\mathfrak{n}^-} \left(\frac{\partial}{\partial x} \right)^a : \mathcal{O}_{\frac{n}{r}-a}(D)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\varepsilon_1(\frac{n}{r}-a+2k)}(D_1, V_l).$$

Similarly, when $k \geq l$, for $a = 1, 2, \dots, k-l$ we have

$$\mathcal{F}_{\frac{n}{r}-a, k, l, -}^\downarrow = \mathcal{F}_{\frac{n}{r}+a, k-a, l, -}^\downarrow \circ \det_{\mathfrak{n}^-} \left(\frac{\partial}{\partial x} \right)^a : \mathcal{O}_{\frac{n}{r}-a}(D)|_{\tilde{G}_1} \longrightarrow \mathcal{O}_{\varepsilon_1(\frac{n}{r}-a+2k)}(D_1, V_l^\vee).$$

For $\lambda > \frac{n}{r} - 1$ we have

$$\begin{aligned}\mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l)|_{\tilde{K}_1} &\simeq d\tau_\lambda(\mathcal{U}(\mathfrak{g}_1)) \mathcal{P}_{(k+l, \underline{k}_{r-1})}(\mathfrak{p}_2^+) \subset \mathcal{H}_\lambda(D)|_{\tilde{K}}, \\ \mathcal{H}_{\varepsilon_1(\lambda+2k)}(D_1, V_l^\vee)|_{\tilde{K}_1} &\simeq d\tau_\lambda(\mathcal{U}(\mathfrak{g}_1)) \mathcal{P}_{(\underline{k}_{r-1}, k-l)}(\mathfrak{p}_2^+) \subset \mathcal{H}_\lambda(D)|_{\tilde{K}},\end{aligned}$$

but this does not hold for smaller λ in general. For such λ , by Theorems 4.6 (2), 4.7 (2), the following holds. For $i = 1, 2, \dots, r$, $\lambda \in \frac{d}{2}(i-1) - \mathbb{Z}_{\geq 0}$, let $M_i^\mathfrak{g}(\lambda)$ be the $(\mathfrak{g}, \tilde{K})$ -submodule given in (2.18).

Theorem 4.16. (1) Let $k, l \in \mathbb{Z}_{\geq 0}$. For $i = 1, 2, \dots, r$,

$$d\tau_\lambda(\mathcal{U}(\mathfrak{g}_1)) \mathcal{P}_{(k+l, \underline{k}_{r-1})}(\mathfrak{p}_2^+) \subset M_i^\mathfrak{g}(\lambda)$$

holds if and only if

$$\lambda \in \begin{cases} \frac{d}{2}(i-1) - (2k+l) - \mathbb{Z}_{\geq 0} & (i=1), \\ \frac{d}{2}(i-1) - 2k - \mathbb{Z}_{\geq 0} & (2 \leq i \leq \lfloor r/2 \rfloor), \\ \frac{d}{2}(i-1) - \min\{2k, k+l\} - \mathbb{Z}_{\geq 0} & (r: \text{odd}, i = \lceil r/2 \rceil), \\ \frac{d}{2}(i-1) - k - \mathbb{Z}_{\geq 0} & (\lceil r/2 \rceil + 1 \leq i \leq r). \end{cases}$$

(2) Let $k, l \in \mathbb{Z}_{\geq 0}$ with $k \geq l$. For $i = 1, 2, \dots, r$,

$$d\tau_\lambda(\mathcal{U}(\mathfrak{g}_1))\mathcal{P}_{(k_{r-1}, k-l)}(\mathfrak{p}_2^+) \subset M_i^\mathfrak{g}(\lambda)$$

holds if and only if

$$\lambda \in \begin{cases} \frac{d}{2}(i-1) - 2k - \mathbb{Z}_{\geq 0} & (1 \leq i \leq \lceil r/2 \rceil - 1), \\ \frac{d}{2}(i-1) - (2k-l) - \mathbb{Z}_{\geq 0} & (r: even, i = r/2), \\ \frac{d}{2}(i-1) - k - \mathbb{Z}_{\geq 0} & (\lceil r/2 \rceil + 1 \leq i \leq r-1), \\ \frac{d}{2}(i-1) - (k-l) - \mathbb{Z}_{\geq 0} & (i = r). \end{cases}$$

Here, “if” follows from the holomorphy in Theorems 4.6 (2), 4.7 (2), and these are already given in [34, Section 7], [35, Corollary 6.8 (1)]. On the other hand, “only if” follows from the non-zeroness in Theorems 4.6 (2), 4.7 (2), and these are new except for r even or $kl = 0$ for + case.

References

- [1] R.J. Beerends and E.M. Opdam, *Certain hypergeometric series related to the root system BC*. Trans. Amer. Math. Soc. **339** (1993), no. 2, 581–609.
- [2] S. Ben Saïd, J.L. Clerc and K. Koufany, *Conformally covariant bi-differential operators on a simple real Jordan algebra*. Int. Math. Res. Not. IMRN 2020, no. 8, 2287–2351.
- [3] J.L. Clerc, *Symmetry breaking differential operators, the source operator and Rodrigues formulae*. Pacific J. Math. **307** (2020), no. 1, 79–107.
- [4] H. Cohen, *Sums involving the values at negative integers of L-functions of quadratic characters*. Math. Ann. **217** (1975), no. 3, 271–285.
- [5] H. Dib, *Fonctions de Bessel sur une algèbre de Jordan*. J. Math. Pures Appl. (9) **69** (1990), no. 4, 403–448.
- [6] T. Enright, R. Howe and N. Wallach, *A classification of unitary highest weight modules*. Representation theory of reductive groups (Park City, Utah, 1982), 97–143, Progr. Math., 40, Birkhäuser Boston, Boston, MA, 1983.
- [7] J. Faraut, S. Kaneyuki, A. Korányi, Q.k. Lu and G. Roos, *Analysis and geometry on complex homogeneous domains*. Progress in Mathematics, 185. Birkhäuser Boston, Inc., Boston, MA, 2000.
- [8] J. Faraut and A. Korányi, *Function spaces and reproducing kernels on bounded symmetric domains*. J. Funct. Anal. **88** (1990), no. 1, 64–89.
- [9] J. Faraut and A. Korányi, *Analysis on symmetric cones*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1994.
- [10] J. Frahm and B. Ørsted, *Knapp-Stein type intertwining operators for symmetric pairs II. —The translation principle and intertwining operators for spinors*. SIGMA Symmetry Integrability Geom. Methods Appl. **15** (2019), Paper No. 084, 50 pp.

- [11] J. Frahm and C. Weiske, *Symmetry breaking operators for line bundles over real projective spaces*. J. Lie Theory **29** (2019), no. 2, 511–558.
- [12] J. Frahm and C. Weiske, *Symmetry breaking operators for real reductive groups of rank one*. J. Funct. Anal. **279** (2020), no. 5, 108568, 70 pp.
- [13] T. Ibukiyama, T. Kuzumaki and H. Ochiai, *Holonomic systems of Gegenbauer type polynomials of matrix arguments related with Siegel modular forms*. J. Math. Soc. Japan **64** (2012), no. 1, 273–316.
- [14] H.P. Jakobsen, *Hermitian symmetric spaces and their unitary highest weight modules*. J. Funct. Anal. **52** (1983), no. 3, 385–412.
- [15] H.P. Jakobsen and M. Vergne, *Restrictions and expansions of holomorphic representations*. J. Funct. Anal. **34** (1979), no. 1, 29–53.
- [16] A. Juhl, *Families of conformally covariant differential operators, Q-curvature and holography*. Progress in Mathematics, 275. Birkhäuser Verlag, Basel, 2009.
- [17] T. Kobayashi, *Multiplicity-free representations and visible actions on complex manifolds*. Publ. Res. Inst. Math. Sci. **41** (2005), no. 3, 497–549.
- [18] T. Kobayashi, *Multiplicity-free theorems of the restrictions of unitary highest weight modules with respect to reductive symmetric pairs*. Representation theory and automorphic forms, 45–109, Progr. Math., 255, Birkhäuser Boston, Boston, MA, 2008.
- [19] T. Kobayashi, *F-method for constructing equivariant differential operators*. Geometric analysis and integral geometry, 139–146, Contemp. Math., 598, Amer. Math. Soc., Providence, RI, 2013.
- [20] T. Kobayashi, *A program for branching problems in the representation theory of real reductive groups*. Representations of reductive groups, 277–322, Progr. Math., 312, Birkhäuser/Springer, Cham, 2015.
- [21] T. Kobayashi, T. Kubo and M. Pevzner, *Conformal symmetry breaking operators for differential forms on spheres*. Lecture Notes in Math. 2170, Springer, Singapore, 2016, ix+192 pp.
- [22] T. Kobayashi and A. Leontiev, *Image of conformally covariant, symmetry breaking operators for $\mathbb{R}^{p,q}$* . Quantum theory and symmetries with Lie theory and its applications in physics. Vol. 1, 3–35. Springer Proc. Math. Stat., 263, Springer, Singapore, 2018.
- [23] T. Kobayashi, B. Ørsted, P. Somberg and V. Souček, *Branching laws for Verma modules and applications in parabolic geometry. I*. Adv. Math. **285** (2015), 1796–1852.
- [24] T. Kobayashi and M. Pevzner, *Differential symmetry breaking operators: I. General theory and F-method*. Selecta Math. (N.S.) **22** (2016), no. 2, 801–845.
- [25] T. Kobayashi and M. Pevzner, *Differential symmetry breaking operators: II. Rankin–Cohen Operators for Symmetric Pairs*. Selecta Math. (N.S.) **22** (2016), no. 2, 847–911.
- [26] T. Kobayashi and M. Pevzner, *Inversion of Rankin–Cohen operators via Holographic Transform*. Ann. Inst. Fourier (Grenoble) **70** (2020), no. 5, 2131–2190.

- [27] T. Kobayashi and B. Speh, *Symmetry breaking for representations of rank one orthogonal groups*. Mem. Amer. Math. Soc. 238 (2015), no. 1126.
- [28] T. Kobayashi and B. Speh, *Symmetry breaking for representations of rank one orthogonal groups II*. Lecture Notes in Mathematics, 2234. Springer, Singapore, 2018. xv+342 pp.
- [29] Q. Labriet, *A geometrical point of view for branching problems for holomorphic discrete series of conformal Lie groups*. Internat. J. Math. **33** (2022), no. 10–11, Paper No. 2250069, 70 pp.
- [30] O. Loos, *Bounded symmetric domains and Jordan pairs*. Math. Lectures, Univ. of California, Irvine, 1977.
- [31] S. Martens, *The characters of the holomorphic discrete series*. Proc. Nat. Acad. Sci. U.S.A. **72** (1975), no. 9, 3275–3276.
- [32] J. Möllers and Y. Oshima, *Restriction of most degenerate representations of $O(1, N)$ with respect to symmetric pairs*. J. Math. Sci. Univ. Tokyo **22** (2015), no. 1, 279–338.
- [33] R. Nakahama, *Construction of intertwining operators between holomorphic discrete series representations*. SIGMA Symmetry Integrability Geom. Methods Appl. **15** (2019), Paper No. 036, 101 pp.
- [34] R. Nakahama, *Computation of weighted Bergman inner products on bounded symmetric domains and restriction to subgroups*. SIGMA Symmetry Integrability Geom. Methods Appl. **18** (2022), Paper No. 033, 105 pp.
- [35] R. Nakahama, *Computation of weighted Bergman inner products on bounded symmetric domains and Parseval–Plancherel-type formulas under subgroups*. SIGMA Symmetry Integrability Geom. Methods Appl. **19** (2023), Paper No. 049, 74 pp.
- [36] V. Ovsienko and P. Redou, *Generalized transvectants–Rankin–Cohen brackets*. Lett. Math. Phys. **63** (2003), no. 1, 19–28.
- [37] J. Peetre, *Hankel forms of arbitrary weight over a symmetric domain via the transvectant*. Rocky Mountain J. Math. **24** (1994), no. 3, 1065–1085.
- [38] L. Peng and G. Zhang, *Tensor products of holomorphic representations and bilinear differential operators*. J. Funct. Anal. **210** (2004), no. 1, 171–192.
- [39] R.A. Rankin, *The construction of automorphic forms from the derivatives of a given form*. J. Indian Math. Soc. (N.S.) **20** (1956), 103–116.
- [40] I. Satake, *Algebraic structures of symmetric domains*. Kanô Memorial Lectures, 4. Iwanami Shoten, Tokyo; Princeton University Press, Princeton, N.J., 1980.
- [41] I. Yokota, *Realizations of involutive automorphisms σ and G^σ of exceptional linear Lie groups G . I. $G = G_2, F_4$ and E_6* . Tsukuba J. Math. **14** (1990), no. 1, 185–223.
- [42] I. Yokota, *Exceptional Lie groups*. arXiv:0902.0431 (2009), 204 pages.