Extended Massive Ambitwistor String

Christian Kunz

E-mail: kunz.christian.321@gmail.com

June 5, 2024

Contents

1	Introduction	2
2	Extended Massive Ambitwistor String 2.1 Consistency Checks: Einstein-Yang-Mills Amplitudes and Massless Limit . 2.2 Factorization of Tree Amplitudes	3 8 10
3	Loop Amplitudes and Factorization3.1 Nonseparating Degeneration3.2 Separating Degeneration3.3 Higher Loops and Cosmological Constant	11 13 15 16
4	Compton Scattering Using Massive Twistor Strings	17
5	Summary and Outlook	21

Abstract

This work considers a variation of the massive ambitwistor string model presented in arXiv:2301.11227 that describes supergravity and super-Yang-Mills on the Coulomb branch simultaneously with a single Lagrangian. All-multiplicity tree and one-loop amplitudes are evaluated and shown to have proper unitary factorization. The massless limit provides Einstein Yang-Mills Amplitudes including multiple gluon traces. It is argued that the cosmological constant vanishes at all orders of perturbation theory. As application results for Compton scattering are obtained and compared with ones in the literature.

1 Introduction

There has been a lot of progress in calculating on-shell scattering amplitudes for massless and massive particles using the spinor-helicity formalism and generalized unitarity [1, 2, 3, 4, 5]. There also has been some progress in deriving such amplitudes directly from a massive ambitwistor string theory centered on polarized scattering equations [6, 7, 8, 9, 10]. What has been still missing, was a consolidation of supergravity and super-Yang-Mills (SYM) on the Coulomb branch into a unified theory that exhibits proper unitary factorization justifying the use of generalized unitarity. This article is a step forward in this direction.

The paper is organized as follows:

In section 2 the supergravity version of the massive ambitwistor string presented in [9, 10] is extended to include additional gauge symmetries. It has some similarity with the supergravity model of Skinner [11] and is shown to be anomaly-free in the same manner as the original massive ambitwistor string. It also has the same $\mathcal{N}=8$ supergravity spectrum. The all-multiplicity tree amplitude is calculated.

In subsection 2.1 some consistency checks are performed. First a massive Einstein-Yang-Mills (EYM) amplitude is derived which includes SYM on the Coulomb branch as special case and has the proper massless limit [12]. Also the massless limit of the general tree amplitude is shown to encompass known supergravity amplitudes [13, 7].

In subsection 2.2 proper factorization of tree amplitudes is established.

In section 3 one-loop amplitudes are examined. They vanish on even spin structures. Their form on the odd spin structure is evaluated.

In subsection 3.1 unitary factorization of the amplitude is demonstrated for nonseparating degeneration of genus 1 Riemann surfaces.

In subsection 3.2 the same is done for separating degeneration.

In subsection 3.3 some remarks concerning higher loop amplitudes are made. It is argued that the cosmological constant is zero at all orders of perturbation theory, and that all higher loop amplitudes vanish on even spin structures. Further, not even all odd spin structures can contribute to loop amplitudes, only a particular subset. It is also mentioned that the original ambitwistor model shares many of the features of the extended model, in particular with regards to even/odd spin structures and the cosmological constant.

Section 4 serves as another consistency check by applying the model to the Compton scattering of a massless particle hitting a massive target. A difference to the original massive ambitwistor string for supergravity is that the same-helicity amplitude does not vanish for 'gravitinos', and the result for 'gluons' is consistent with the one arising from SYM on the Coulomb branch.

Section 5 contains summary and discussion.

2 Extended Massive Ambitwistor String

In this section the massive ambitwistor string of [10] for maximal supergravity is extended with an enlarged supersymmetric gauge symmetry. It can also be viewed as generalization of a massless ambitwistor string model in [14] to include massive particles.

The twistor action used here is, in similar notation to [14],

$$S = \int_{\Sigma} \mathcal{Z}^{a} \cdot \bar{\partial} \mathcal{Z}_{a} + A_{ab} \mathcal{Z}^{a} \cdot \mathcal{Z}^{b} + a \langle \lambda_{A} \lambda_{B} \rangle \Omega^{AB} + \tilde{a} \langle \eta_{\mathcal{I}} \eta_{\mathcal{J}} \rangle \Omega^{\mathcal{I}\mathcal{J}} + S_{\rho_{1}} + S_{(\rho_{2},\tau)},$$

$$S_{\rho} = \int_{\Sigma} \frac{1}{2} \langle \rho_{A} \bar{\partial} \rho_{B} \rangle \Omega^{AB} + B_{ab} \lambda_{A}^{a} \rho_{B}^{b} \Omega^{AB} + b \langle \lambda_{A} \rho_{B} \rangle \Omega^{AB},$$

$$S_{(\rho,\tau)} = \int_{\Sigma} \frac{1}{2} (\langle \rho_{A} \bar{\partial} \rho_{B} \rangle \Omega^{AB} + \langle \tau_{\mathcal{I}} \bar{\partial} \tau_{\mathcal{J}} \rangle \Omega^{\mathcal{I}\mathcal{J}}) + D_{ab} (\lambda_{A}^{a} \rho_{B}^{b} \Omega^{AB} + \eta_{\mathcal{I}}^{a} \tau_{\mathcal{J}}^{b} \Omega^{\mathcal{I}\mathcal{J}})$$

$$+ d (\langle \lambda_{A} \rho_{B} \rangle \Omega^{AB} + \langle \eta_{\mathcal{I}} \tau_{\mathcal{J}} \rangle \Omega^{\mathcal{I}\mathcal{J}}),$$

$$(2.1)$$

where a = 1, 2 is the little group index with contraction notation $\xi_a \zeta_b \varepsilon^{ab} = \xi_a \zeta^a \equiv \langle \xi \zeta \rangle$ and \mathcal{Z}_a is a pair of supertwistor fields that are worldsheet spinors, repackaged into *Dirac* supertwistors of the form

$$\mathcal{Z} = (\lambda_A, \mu^A, \eta_\iota) : \lambda_A = (\lambda_\alpha, \tilde{\lambda}_{\dot{\alpha}}), \mu^A = (\mu^{\dot{\alpha}}, \tilde{\mu}^\alpha), \ \eta_\iota = (\eta_\mathcal{I}, \tilde{\eta}^\mathcal{I}), \eta_\mathcal{I} = (\eta_I, \tilde{\eta}_{\dot{I}}), \tilde{\eta}^\mathcal{I} = (\tilde{\eta}^{\prime I}, {\eta^{\prime \dot{I}}}),$$

$$I, \dot{I} = 1 \dots \frac{\mathcal{N}}{2}, \mathcal{I} = 1, \dots, \mathcal{N}, \ \iota = 1, \dots, 2\mathcal{N}, \ \mathcal{N} = 8,$$

where λ_A and μ^A are Dirac spinors made up of the homogeneous chiral and antichiral components of the twistor $Z = (\lambda_{\alpha}, \mu^{\dot{\alpha}})$ and dual twistor $\tilde{Z} = (\tilde{\lambda}_{\dot{\alpha}}, \tilde{\mu}^{\alpha})$, and $\eta_{\iota} = (\eta_{\mathcal{I}}, \tilde{\eta}^{\mathcal{I}})$ are fermionic components with $\mathcal{I} = 1, \ldots, \mathcal{N}$ as the R-symmetry index, with $\mathcal{N} = 8$ for maximal supergravity. Indices are raised and lowered with skew 4×4 forms $\Omega^{EF}, \Omega_{EF}, \Omega^{EG}\Omega_{GF} = \delta_F^E$. The $\mathcal{N} \times \mathcal{N}$ form $\Omega_{\mathcal{I}\mathcal{J}}$ decomposes into two $\frac{\mathcal{N}}{2} \times \frac{\mathcal{N}}{2}$ forms: $\Omega_{\mathcal{I}\mathcal{J}} = \Omega_{IJ} \otimes \Omega_{\dot{I}\dot{J}}$. The inner product is defined as

 $\mathcal{Z}_a \cdot \mathcal{Z}_b = \frac{1}{2} (\tilde{Z}_a \cdot \tilde{Z}_b + \tilde{Z}_b \cdot Z_a + \tilde{\eta}_{a\mathcal{I}} \eta_b^{\mathcal{I}} + \tilde{\eta}_{b\mathcal{I}} \eta_a^{\mathcal{I}}), \quad \tilde{Z}_a \cdot Z_b = \tilde{\mu}_a^{\alpha} \lambda_{b\alpha} + \tilde{\lambda}_{a\dot{\alpha}} \mu_b^{\dot{\alpha}},$ with special treatment of $\bar{\partial} Z$ when taking the dual: $\widetilde{\partial} Z = -\bar{\partial} \tilde{Z}$.

 $\rho_{rA}^a(r=1,2)$ and $\tau_{\mathcal{I}}^a$ are auxiliary fermionic and bosonic worldsheet spinors, respectively, and (B_{ab},b,D_{ab},d) are fermionic Lagrange multipliers for the constraints $\lambda_A^{(a}\rho_{1B}^{b)}\Omega^{AB} = 0 = \langle \lambda_A\rho_{1B} \rangle , \lambda_A^{(a}\rho_{2B}^{b)}\Omega^{AB} + \eta_{\mathcal{I}}^{(a}\tau_{\mathcal{J}}^{b)}\Omega^{\mathcal{I}\mathcal{J}} = 0 = \langle \lambda_A\rho_{2B} \rangle \Omega^{AB} + \langle \eta_{\mathcal{I}}\tau_{\mathcal{J}} \rangle \Omega^{\mathcal{I}\mathcal{J}}$ on the supersymmetric gauge currents.

Little group transformations for the twistors are gauged by the fields $A_{ab} = A_{(ab)}$, and \tilde{a} are worldsheet (0,1)-forms that also act as Lagrange multipliers, required in order to have a closed current algebra.

It can be noted that the action is somewhat similar to the $\mathcal{N}=8$ supergravity model of Skinner [11] which has the same kind of auxiliary pair of supertwistors with reverse statistics as in $S_{(\rho,\tau)}$, but only a single main supertwistor. The action is even more similar to the massive ambitwistor model of [9, 10] and can be viewed as an extension of it. The difference consists in doubling the fermionic components in the main supertwistor, gauging them with auxiliary τ fields using the same worldsheet supersymmetry as for ρ_2 , and replacing the constraints on mass operators with constraints on currents that close the current algebra.

BRST quantization can be performed similarly to [8]. In addition to the familiar fermionic (b, c) ghost system related to worldsheet gravity (the action (2.1) is already written in conformal gauge), the Lagrange multipliers in (2.1) are associated with corresponding (antighost, ghost) pairs:

the bosonic fields $\{A_{ab}, a, \tilde{a}\}$ with fermionic ghosts $\{(M^{ab}, N_{ab}), (m, n), (\tilde{m}, \tilde{n})\}$ and the fermionic fields $\{B_{ab}, b, D_{ab}, d\}$ with bosonic ghosts $\{(\beta^{ab}, \gamma_{ab}), (\beta, \gamma), (\tilde{\beta}^{ab}, \tilde{\gamma}_{ab}), (\tilde{\beta}, \tilde{\gamma})\}$. All the Lagrange multipliers can be gauged to zero and the BRST operator becomes $Q = \oint dz J_{BRST}(z)$, where the BRST current $J_{BRST}(z)$ is (based on the formula $Q = c^i K_i - \frac{1}{2} f_k^{ij} c^i c^j b_k$ for the current algebra K_i with structure constants f_k^{ij} , ghosts c^i , and antighosts b_k):

$$\begin{split} J_{BRST} &= c\,T + N_{ab}(J^{ab} + M^a_c N^{bc}) + n\langle\lambda_A\lambda^A\rangle + \tilde{n}\langle\eta_\mathcal{I}\eta^\mathcal{I}\rangle + \gamma_{ab}\lambda^{aA}\rho^b_{1A} + \gamma\langle\lambda^A\rho_{1A}\rangle + m\gamma\gamma \\ &\quad + \tilde{\gamma}_{ab}(\lambda^{aA}\rho^b_{2A} + \eta^{a\mathcal{I}}\tau^b_\mathcal{I}) + \tilde{\gamma}(\langle\lambda^A\rho_{2A}\rangle + \langle\eta^\mathcal{I}\tau_\mathcal{I}\rangle) + (m+\tilde{m})\tilde{\gamma}\tilde{\gamma}\;,\\ T &= \mathcal{Z}^a \cdot \partial\mathcal{Z}_a + \frac{1}{2}\sum_{r=1,2}\langle\rho_{rA}\partial\rho^A_r\rangle + \beta^{ab}\partial\gamma_{ab} + \beta\partial\gamma + \frac{1}{2}\langle\tau_\mathcal{I}\partial\tau^\mathcal{I}\rangle + \tilde{\beta}^{ab}\partial\tilde{\gamma}_{ab} + \tilde{\beta}\partial\tilde{\gamma}\\ &\quad + M^{ab}\partial N_{ab} + m\partial n + \tilde{m}\partial\tilde{n} + \frac{1}{2}(b\partial c + \partial(bc))\;,\\ J^{ab} &= \mathcal{Z}^a \cdot \mathcal{Z}^b + \frac{1}{2}\sum_{r=1,2}\rho^{(a}_{rA}\rho^{b)A}_r + \beta^{c(a}\gamma^b_c + \frac{1}{2}\tau^{(a}_\mathcal{I}\tau^{b)\mathcal{I}} + \tilde{\beta}^{c(a}\tilde{\gamma}^b_c)\;, \end{split}$$

where T is the energy-momentum current and J^{ab} the sl(2, \mathbb{C}) current. Obstructions to the nilpotency of the BRST charge Q can only come from the sl(2, \mathbb{C}) algebra and the Virasoro

algebra. The sl(2, \mathbb{C}) anomaly coefficient is, using $\operatorname{tr}_{\operatorname{adj}}(t^kt^k) = 6$, $\operatorname{trf}_{\operatorname{F}}(t^kt^k) = 3/2$,

$$a_{sl2} = \frac{1}{2} \text{trf}_{F}(t^{k}t^{k})((8-2\mathcal{N})_{\mathcal{Z}} - 4_{\rho_{1}} - 4_{\rho_{2}} + \mathcal{N}_{\tau}) + \text{tr}_{adj}(t^{k}t^{k})(-1_{MN} + 1_{\beta\gamma} + 1_{\tilde{\beta}\tilde{\gamma}}) = \frac{3}{4}(8-\mathcal{N}) = 0.$$

For the central charge it is assumed that there is a contribution of 12 from compactifying 6 dimensions like for the massive ambitwistor model of [10], e.g. with help of 6 bosonic scalars as in appendix A of [10]. Then it can be seen to vanish for $\mathcal{N} = 8$:

$$c = (-8 + 2\mathcal{N})_{\mathcal{Z}} - 26_{bc} - 6_{MN} - 2_{mn} - 2_{\tilde{m}\tilde{n}} + 4_{\rho_1} + 8_{\beta\gamma} + 4_{\rho_2} - \mathcal{N}_{\tau} + 8_{\tilde{\beta}\tilde{\gamma}} + 12_{\text{comp}} = -8 + \mathcal{N} = 0.$$

Therefore, the model can be viewed as conditionally anomaly-free. The additional contribution to the central charge can also be obtained by gauging 6 additional worldsheet supersymmetries between 3 or 4 additional auxiliary supertwistors $(\rho_i^A, \tau_i^A), i = 3, 4, 5, 6$ that do not affect the spectrum and vertex operators. In this manner the model can become anomaly-free without resorting to extra spacetime dimensions. Of course, one could argue that the auxiliary spinors are remnants from compactifying extra dimensions in twistor space, and there might be many more ways to come up with an auxiliary action that achieves the same outcome.

To find the spectrum one needs to determine the Virasoro L_0 constant a [14]. In the NS sector a is calculated by changing in the calculation of the central charge the bc contribution from -26_{bc} to -2_{bc} and dividing the result by 24, such that a=1 in the NS sector. In the R sector a is:

$$24a = (16 - 4\mathcal{N})_{\mathcal{Z}} - 2_{bc} - 6_{MN} - 2_{mn} - 2_{\tilde{m}\tilde{n}} - 8_{\rho_1} + 8_{\beta\gamma} - (8_{\rho_2} - 2\mathcal{N}_{\tau}) + 8_{\tilde{\beta}\tilde{\gamma}} + 12_{\text{comp}} = 2(8 - \mathcal{N}) = 0.$$

i.e. a = 0 in the R sector.

The situation is parallel to the massive ambitwistor model of [10], and based on BRST cohomology the spectrum is again the one of $\mathcal{N}=8$ supergravity with SU(4) × SU(4) R-symmetry. Although the fermionic modes in the supertwistors have been doubled, similarly to the exclusion of μ_n^{aA} modes from the spectrum because of BRST cohomology, now the $\tilde{\eta}_n^{a\mathcal{I}}$ cannot contribute either to the spectrum because of the requirement that the non-negative modes of the $\eta^{a\mathcal{I}}\tau_{\mathcal{I}}^{b}$ and $\langle \eta^{\mathcal{I}}\tau_{\mathcal{I}} \rangle$ currents annihilate physical states. One other difference is that the action in [10] contains mass operators that ensure that the supersymmetric multiplet has a particular mass whereas the current model leaves the value of the mass open.

Moving on to vertex operators, following [8, 9, 10] vertex operators are sought in the form

$$\begin{pmatrix} \mathcal{V} \\ V \end{pmatrix} = \int \mathrm{d}^2 u \, \mathrm{d}^2 v \begin{pmatrix} \mathcal{W}(u) \\ w(u) \end{pmatrix} \bar{\delta}(\langle v \epsilon \rangle - 1) \bar{\delta}^4(\langle u \lambda_A \rangle - \langle v \kappa_A \rangle) \bar{\delta}^{\mathcal{N}}(\langle u \eta_{\mathcal{I}} \rangle - \langle v \zeta_{\mathcal{I}} \rangle) e^{\langle u \mu^A \rangle \epsilon_A + \langle u \tilde{\eta}^{\mathcal{I}} \rangle q_{\mathcal{I}}}.$$
 (2.2)

Here ϵ_a are little group spinors, $\kappa_a^A = (\kappa_a^\alpha, \tilde{\kappa}_a^{\dot{\alpha}})$ are momentum spinors with associated polarization spinors $\epsilon^A = \langle \epsilon \kappa^A \rangle$, $\zeta_a^{\mathcal{I}}$ are supermomenta with superpolarization spinors $q^{\mathcal{I}} = \langle \epsilon \zeta^{\mathcal{I}} \rangle$, and \mathcal{V} and V stand for fixed and integrated vertex operators, respectively, which differ only by an operator \mathcal{W} or w with \mathcal{W} typically being just a product of fermionic ghost fields and delta functions of bosonic ghost fields. As the handling of fixed and integrated vertex operators concerning the (b,c) ghosts is standard, the difference being just an integration over the worldsheet, the (b,c) ghosts and worldsheet integrations will be omitted in the formulae for \mathcal{W} and w, but then added later in the correlators. Also the effect of (m,n) is just to fix $\langle \lambda_\alpha \lambda^\alpha \rangle = \langle \lambda_{\dot{\alpha}} \lambda^{\dot{\alpha}} \rangle$ which can be viewed as a little group transformation outside of sl(2, \mathbb{C}) [2, 15] and will be skipped as well, and similarly for (\tilde{m}, \tilde{n}) . Finally it is assumed that the bosonic fields from compactifying extra dimensions or additional supersymmetric gauge symmetries do not affect the spectrum or vertex operators¹. The remainder can be simplified even further because the form (2.2) is such that it is sufficient that the delta functions of γ_{ab} and $\tilde{\gamma}_{ab}$ in \mathcal{W} force only the ghost components orthogonal to u to vanish [8]. Consequently

$$\begin{split} \mathcal{W}(u) &= \delta(\gamma) \, \delta(u^a u^b \gamma_{ab}) \, \delta(\tilde{\gamma}) \, \delta(u^a u^b \tilde{\gamma}_{ab}), \\ w(u) &= \left(\frac{\langle \hat{u} \lambda_A \rangle \, \epsilon^A}{\langle u \hat{u} \rangle} + \frac{\langle \rho_{1A} \rho_{1B} \rangle}{2} \epsilon^A \epsilon^B \right) \left(\frac{\langle \hat{u} \lambda_A \rangle \epsilon^A}{\langle u \hat{u} \rangle} + \frac{\langle \rho_{2A} \rho_{2B} \rangle}{2} \epsilon^A \epsilon^B + \frac{\langle \hat{u} \eta_\mathcal{I} \rangle q^\mathcal{I}}{\langle u \hat{u} \rangle} + \frac{\langle \tau_\mathcal{I} \tau_\mathcal{J} \rangle}{2} q^\mathcal{I} q^\mathcal{J} \right). \end{split}$$

Here \hat{u} is a reference little group spinor that forms a basis with u such that $\langle u\hat{u}\rangle \neq 0$. In amplitudes it will fall out based on solutions of the scattering equations [8].

The all-multiplicity tree amplitude of this model can now be evaluated:

$$\mathcal{A}_{n}^{\text{tree}} = \left\langle c \mathcal{V}_{1}^{(-1,-1)} c \mathcal{V}_{2}^{(-1,0)} c \mathcal{V}_{3}^{(-1,0)} \prod_{j=4}^{n} \int_{\Sigma} V_{j} \right\rangle,\,$$

where $\mathcal{V}^{(-1,-1)}$ denotes a fixed vertex operator with regards to all $(\beta^{ab}, \gamma_{ab}), (\tilde{\beta}^{ab}, \tilde{\gamma}_{ab}), (\beta, \gamma), (\tilde{\beta}, \tilde{\gamma})$ ghosts and $\mathcal{V}^{(-1,0)}$ only fixed with regards to $(\beta^{ab}, \gamma_{ab}), (\tilde{\beta}^{ab}, \tilde{\gamma}_{ab})$.

Integrating out the μ^A and $\tilde{\eta}^{\mathcal{I}}$ leads, on the support of the delta functions, to the polarized scattering equations (using the notation $\sigma_{ij} = \sigma_i - \sigma_j$):

$$\langle v_r \kappa_r^A \rangle = \langle u_r \lambda^A(\sigma_r) \rangle , \quad \lambda_A^a(\sigma_r) = \sum_{l \neq r} \frac{u_l^a \epsilon_{lA}}{\sigma_{rl}},$$

$$\langle v_r \zeta_r^{\mathcal{I}} \rangle = \langle u_r \eta^{\mathcal{I}}(\sigma_r) \rangle , \quad \eta_{\mathcal{I}}^a(\sigma_r) = \sum_{l \neq r} \frac{u_l^a q_{l\mathcal{I}}}{\sigma_{rl}}.$$
(2.3)

¹They will be crucial for loop amplitudes in the next section.

Based on these scattering equations and because of the delta functions enforcing $\langle v_r \epsilon_r \rangle = 1$ it follows that $\langle u_r \lambda^A(\sigma_r) \rangle \epsilon_{rA} = \langle v_r \kappa_r^A \rangle \epsilon_{rA} = 0$ and $\langle u_r \eta^{\mathcal{I}}(\sigma_r) \rangle q_{r\mathcal{I}} = \langle v_r \zeta_r^{\mathcal{I}} \rangle q_{r\mathcal{I}} = 0$ such that the dependence on \hat{u} falls out of the correlator.

Then the amplitude is evaluated to

$$\mathcal{A}_n^{\text{tree}} = \int d\mu_n^{\text{pol}|\mathcal{N}} \, \mathcal{I}_n \,, \tag{2.4}$$

where $\mathrm{d}\mu_n^{\mathrm{pol}|\mathcal{N}}$ is the measure which localizes on the scattering equations (2.3) and also accounts for the quotient by the volume of $\mathrm{SL}(2,\mathbb{C})_{\sigma} \times \mathrm{SL}(2,\mathbb{C})_{u}$ from the path integral over the (b,c) and (M^{ab},N_{ab}) ghost systems:

$$d\mu_n^{\text{pol}|\mathcal{N}} := \frac{\prod_l d\sigma_l d^2 u_l d^2 v_l}{\text{vol}(\text{SL}(2, \mathbb{C})_{\sigma} \times \text{SL}(2, \mathbb{C})_u)} \prod_{r=1}^n \bar{\delta}(\langle v_r \epsilon_r \rangle - 1) \bar{\delta}^4(\langle u_r \lambda^A \rangle - \langle v_r \kappa_r^A \rangle) \bar{\delta}^{\mathcal{N}}(\langle u_r \eta^{\mathcal{I}} \rangle - \langle v_r \zeta_r^{\mathcal{I}} \rangle)$$
(2.5)

and where the integrand \mathcal{I}_n is the product of two reduced determinants,

$$\mathcal{I}_{n} = \det' H \det' G = \frac{\det H_{[kl]}^{[ij]}}{\langle u_{i}u_{j}\rangle \langle u_{k}u_{l}\rangle} \frac{\det G_{[st]}^{[pr]}}{\langle u_{p}u_{r}\rangle \langle u_{s}u_{t}\rangle},$$

$$H_{ij} = \frac{\epsilon_{iA}\epsilon_{j}^{A}}{\sigma_{ij}}, \qquad u_{i}^{a}H_{ii} = -\lambda_{A}^{a}\epsilon_{i}^{A},$$

$$G_{ij} = \frac{\epsilon_{iA}\epsilon_{j}^{A} + q_{i\mathcal{I}}q_{j}^{\mathcal{I}}}{\sigma_{ij}}, \qquad u_{i}^{a}G_{ii} = -\lambda_{A}^{a}\epsilon_{i}^{A} - \eta_{\mathcal{I}}^{a}q_{i}^{\mathcal{I}},$$
(2.6)

where H and G are $n \times n$ matrices with co-rank 2 and $H_{[kl]}^{[ij]}$ means that rows i and j and columns k and l have been removed. The reduced determinants are independent of the choice of rows and columns to remove.

The amplitude (2.4) has been worked out in the so-called little group (LG) representation [7], mainly because the derivation required 'scattering equations' (2.3) for the η fields, but in the following it is more convenient to go over to the R-symmetry preserving representation, by Fourier transforming both halves of the $q_{\mathcal{I}} = (q_I, \tilde{q}_{\dot{I}}) = (q_I, \langle \epsilon \zeta_I \rangle) = (\langle \epsilon \tilde{\zeta}_{\dot{I}} \rangle, \tilde{q}_{\dot{I}})$ variables, using $N = \tilde{N} = \frac{N}{2}$ and $v_a = \xi_a + \langle \xi v \rangle \epsilon_a$ where (ϵ, ξ) form a spinor basis with $\langle \xi \epsilon \rangle = 1$ [7]:

$$2^{-N} \int \prod_{i=1}^{n} d^{N} q_{i}^{I} \prod_{j=1}^{n} e^{-\frac{1}{2} q_{i}^{I} \langle \xi_{i} \zeta_{iI} \rangle} e^{F_{N}} = \prod_{i=1}^{n} \delta^{N} \left(\sum_{j=1}^{n} \frac{\langle u_{i} u_{j} \rangle}{\sigma_{ij}} \langle \epsilon_{j} \zeta_{jI} \rangle - \langle v_{i} \zeta_{jI} \rangle \right),$$

$$F_{N} = \sum_{i < j} \frac{\langle u_{i} u_{j} \rangle}{\sigma_{ij}} q_{iI} q_{j}^{I} - \frac{1}{2} \sum_{i=1}^{n} \langle \xi_{i} v_{i} \rangle q_{i}^{2},$$

$$2^{-\tilde{N}} \int \prod_{i=1}^{n} d^{\tilde{N}} \tilde{q}_{i}^{\dot{I}} \prod_{j=1}^{n} e^{-\frac{1}{2} \tilde{q}_{i}^{\dot{I}} \langle \xi_{i} \tilde{\zeta}_{i\dot{I}} \rangle} e^{F_{\tilde{N}}} = \prod_{i=1}^{n} \delta^{\tilde{N}} \left(\sum_{j=1}^{n} \frac{\langle u_{i} u_{j} \rangle}{\sigma_{ij}} \langle \epsilon_{j} \tilde{\zeta}_{j\dot{I}} \rangle - \langle v_{i} \tilde{\zeta}_{j\dot{I}} \rangle \right),$$
$$\tilde{F}_{\tilde{N}} = \sum_{i < j} \frac{\langle u_{i} u_{j} \rangle}{\sigma_{ij}} \tilde{q}_{i\dot{I}} \tilde{q}_{j}^{\dot{I}} - \frac{1}{2} \sum_{i=1}^{n} \langle \xi_{i} v_{i} \rangle \tilde{q}_{i}^{2}.$$

When doing this, the q variables in the matrix G and in $F_{\mathcal{N}} = F_N + \tilde{F}_{\tilde{N}}$ will have transposed halves, but because the inner products are the same one can just transpose one set of these q variables, e.g. the ones in $F_{\mathcal{N}}$. One gets

$$\mathcal{A}_{n}^{\text{tree}} = \int d\mu_{n}^{\text{pol}} \,\mathcal{I}_{n} e^{F_{\mathcal{N}}}, \qquad F_{\mathcal{N}} = \sum_{i < j} \frac{\langle u_{i} u_{j} \rangle}{\sigma_{ij}} q_{i\mathcal{I}} q_{j}^{\mathcal{I}} - \frac{1}{2} \sum_{i=1}^{n} \langle \xi_{i} v_{i} \rangle \, q_{i\mathcal{I}} q_{i}^{\mathcal{I}}, \qquad (2.7)$$
$$d\mu_{n}^{\text{pol}} := \frac{\prod_{l} d\sigma_{l} d^{2} u_{l} d^{2} v_{l}}{\text{vol}(\text{SL}(2, \mathbb{C})_{\sigma} \times \text{SL}(2, \mathbb{C})_{u})} \prod_{r=1}^{n} \bar{\delta} (\langle v_{r} \epsilon_{r} \rangle - 1) \bar{\delta}^{4} (\langle u_{r} \lambda^{A} \rangle - \langle v_{r} \kappa_{r}^{A} \rangle),$$

where compared to (2.5) the measure has lost the delta function in the fermionic fields, replaced by the exponential factor $\exp(F_{\mathcal{N}})$.

Now the only difference in the tree amplitude to the original one [7, 5] is that a second occurrence of $\det' H$ is replaced by $\det' G$. The elements in G might look surprising as the terms containing ϵ^A and the ones containing q^T seem to refer to incompatible quantities, but one has to keep in mind that the amplitude in (2.7) is a sum of component amplitudes describing different physical situations, and the sum of quantities in elements from the matrix G are always split up among component amplitudes. In other words, (2.7) should be viewed as a power series in the q variables with the coefficients being component amplitudes describing different physics, although some of them are just permutations of each other. In terms of natural units the elements in G containing ϵ^A should be viewed as multiplied with the gravitational coupling constant $\kappa = \sqrt{G_N}$, i.e. the q^T -terms in G effectively lower the power in gravitational couplings, in contrast to the terms in $\exp(F_N)$. One important benefit of (2.7) is that component amplitudes exist that contain a term $q_i^T q_{jT}$ even when $\langle u_i u_j \rangle = 0$, allowing for more realistic situations, particularly between massless particles of the same helicity when the exponential factor $\exp(F_N)$ cannot contribute because of $\langle u_i u_j \rangle = 0$.

2.1 Consistency Checks: Einstein-Yang-Mills Amplitudes and Massless Limit

The purpose of this subsection is to check whether some component amplitudes are compatible with known amplitudes that describe interesting physics. One example are Einstein-Yang-Mills (EYM) amplitudes.

First it is to be noted that all off-diagonal elements in the G matrix come from the auxiliary spinor fields and only the diagonal elements reflect contractions of the main supertwistors that are relevant for physical interpretation. Collect all diagonal elements with non-zero q-dependence and consider all component amplitudes that contain the product of these diagonal elements with the contributions from λ_A removed and all other component amplitudes with the same (maximum) power in q variables from $\det' G$. Consider the case where all q-products are connected to each other to form a single trace (loop) of length m. Then the component amplitude arises from the determinant of the product of two matrices \hat{H} and \hat{G} , with $(n-m)\times(n-m)$ matrix \hat{H} having all rows and columns containing a q dependence removed and $m\times m$ matrix \hat{G} containing only the q dependent elements that have been removed from \hat{H} . Obviously \hat{G} inherits co-rank 2 from the original matrix G, and \hat{H} is no longer singular. Therefore, for a single trace amplitude:

$$\mathcal{A}_{n,1\text{tr}}^{\text{tree}} = \int d\mu_n^{\text{pol}} \frac{\det H_{[kl]}^{[ij]}}{\langle u_i u_j \rangle \langle u_k u_l \rangle} \det \hat{H} \frac{\det \hat{G}_{[st]}^{[pr]}}{\langle u_p u_r \rangle \langle u_s u_t \rangle} e^{F_{\mathcal{N}}}.$$
 (2.8)

By pulling down $\frac{\langle u_p u_r \rangle}{\sigma_{pr}} q_{p\mathcal{I}} q_r^{\mathcal{I}}$ and $\frac{\langle u_s u_t \rangle}{\sigma_{st}} q_{s\mathcal{I}} q_t^{\mathcal{I}}$ from the exponential factor $F_{\mathcal{N}}$ to close the trace (also recognizing that the term $\langle \xi_i v_i \rangle q_{i\mathcal{I}} q_i^{\mathcal{I}}$ in $F_{\mathcal{N}}$ vanishes for a particle referred to by a singly indexed q and does not contribute) one gets a Parke-Taylor(PT) factor with q-dependent numerator N(q):

$$\mathcal{A}_{n,1\text{tr}}^{\text{tree}} = \int d\mu_n^{\text{pol}} \det' H \, \det \hat{H} \, \text{PT} \, e^{F_{\mathcal{N}}}, \quad \text{PT} = \sum_{\text{Perm}} \frac{N(q)}{\sigma_{m1} \prod_{i=1}^{m-1} \sigma_{i\,i+1}}. \tag{2.9}$$

This is the single trace 'massive EYM' amplitude with no restrictions on the individual masses. When m=n, \hat{H} is empty and (2.9) is then a SYM amplitude on the Coulomb branch [7, 10], i.e. the current model covers both supergravity and SYM. But it should be noted that by being embedded in $\mathcal{N}=8$ supergravity this SYM has more supersymmetry than the conventional SYM.

Concerning multiple traces the same procedure of removing all non-zero q-dependent elements in the matrix G still would result in a degeneration of G into a product of two matrices \hat{H} and \hat{G} , but \hat{G} would have a co-rank of twice the number of traces. Therefore, instead of removing all non-zero q-dependent elements in G one should keep 2 rows and 2 columns for all traces except one and sum over all ways of doing this. In each summand \hat{H} gets all q-dependence removed and is then a non-singular $t \times t$ matrix with $t = n - 2 - \sum_k (l_k - 2)$, where l_k is the length of trace k, whereas \hat{G} again has co-rank 2. To get a PT factor for each trace one needs to pull down two elements from the exponential factor F_N for each trace corresponding either to the two rows and columns kept in \hat{H} or to the two rows and columns removed from \hat{G} like for the single trace.

Another consistency check is to take the massless limit. This is a trivial generalization of section 6 in [7]. The amplitude for k negative helicity and n-k positive helicity massless particles becomes, using the standard notation $\langle \epsilon_i \epsilon_j \rangle = \varepsilon^{\alpha\beta} \epsilon_{i\alpha} \epsilon_{j\beta} = -\langle \epsilon_j \epsilon_i \rangle$, $[\epsilon_i \epsilon_j] = \varepsilon^{\dot{\alpha}\dot{\beta}} \epsilon_{i\dot{\alpha}} \epsilon_{\dot{j}\dot{\beta}} = -[\epsilon_j \epsilon_i]$ (not to be confused with similar usage of $\langle \rangle$ for little group contractions)

$$\mathcal{A}_{n,k}^{4d,\text{tree}} = \int d\mu_{n,k}^{4d} \det' G_{-} \det' G_{+} e^{F_{N}^{k}}, \quad \det' G_{\pm} = \frac{\det G_{\pm}^{|st|}}{u_{s} u_{t}}, \qquad (2.10)$$

$$d\mu_{n,k}^{4d} := \frac{\prod_{l=1}^{n} d\sigma_{l} du_{l} / u_{l}}{\text{vol}(GL(2, \mathbb{C}))} \prod_{i=1}^{k} \bar{\delta}^{2} (u_{i} \tilde{\lambda}_{\dot{\alpha}}(\sigma_{i}) - \frac{\tilde{\kappa}_{i\dot{\alpha}}}{\epsilon_{i}}) \prod_{p=k+1}^{n} \bar{\delta}^{2} (u_{p} \lambda_{\alpha}(\sigma_{p}) - \frac{\kappa_{p\alpha}}{\tilde{\epsilon}_{p}}),$$

$$F_{N}^{k} = \sum_{i=1}^{k} \sum_{j=k+1}^{n} \frac{u_{i} u_{j}}{\sigma_{ij}} q_{i\mathcal{I}} q_{j}^{\mathcal{I}}, \quad \lambda_{\alpha}(\sigma) = \sum_{i=1}^{k} \frac{u_{i} \epsilon_{i\alpha}}{\sigma - \sigma_{i}}, \quad \tilde{\lambda}_{\dot{\alpha}}(\sigma) = \sum_{i=k+1}^{n} \frac{u_{i} \epsilon_{i\dot{\alpha}}}{\sigma - \sigma_{i}},$$

$$\eta_{I}(\sigma) = \sum_{i=1}^{k} \frac{u_{i} q_{iI}}{\sigma - \sigma_{i}}, \quad \tilde{\eta}_{\dot{I}}(\sigma) = \sum_{i=k+1}^{n} \frac{u_{i} q_{i\dot{\alpha}}}{\sigma - \sigma_{i}},$$

$$G_{-}^{ij} = \frac{\langle \epsilon_{i} \epsilon_{j} \rangle + q_{iI} q_{j}^{I}}{\sigma_{ij}}, \quad G_{+}^{ii} = -\frac{\langle \epsilon_{i} \lambda(\sigma_{i}) \rangle + q_{i}^{I} \eta_{I}(\sigma_{i})}{u_{i}}, \quad 1 \leq i, j \leq k,$$

$$G_{+}^{ij} = \frac{[\tilde{\epsilon}_{i} \tilde{\epsilon}_{j}] + q_{iI} q_{j}^{I}}{\sigma_{ij}}, \quad G_{+}^{ii} = -\frac{[\tilde{\epsilon}_{i} \tilde{\lambda}(\sigma_{i})] + q_{i}^{I} \tilde{\eta}_{I}(\sigma_{i})}{u_{i}}, \quad k+1 \leq i, j \leq n,$$

where $G^{[st]}$ denotes a matrix G with row s and column t removed. Up to the additional entries in the matrices coming from fermionic fields, this is the tree amplitude of [13, 7]. The direct massless limit of the single trace EYM amplitude (2.8) is obtained similarly:

$$\mathcal{A}_{n,k,1\text{tr}}^{4d,\text{tree}} = \int d\mu_{n,k}^{4d} \det \hat{H}_{-} \det \hat{H}_{+} \, \text{PT} \, e^{F_{\mathcal{N}}^{k}},$$

where \hat{H}_{\pm} is the matrix obtained from G_{\pm} by removing all rows and columns containing non-vanishing fermionic entries. This is exactly the single trace EYM amplitude of [12] if there the reduced determinants are calculated by removing the last row and column representing 'squeezed' gluons. It is interesting to notice that trying to get EYM amplitudes as component amplitudes directly from (2.10) runs into the limitation that all positive helicity gluons must be adjacent to each other. The same obstruction occurred in the massless models of [14].

2.2 Factorization of Tree Amplitudes

Again this becomes a straightforward exercise by building upon work done in section 7 of [7]. There it has been shown that the measure $d\mu_n^{\text{pol}}$, the exponential factor e^{F_N} , and the reduced determinants $\det' H$ without fermionic elements factorize correctly. Therefore,

what remains to be shown is proper factorization of the reduced determinant of G with fermionic contributions included. One observation is that the matrix elements of G can be rewritten as, when using the 'super' notation $\epsilon_j^A = (\epsilon_j^A, q_i^{\mathcal{I}}), \zeta_A^a = (\lambda_A^a, \eta_{\mathcal{I}}^a)$:

$$G_{ij} = \frac{\epsilon_{i\mathcal{A}}\epsilon_{j}^{\mathcal{A}}}{\sigma_{ij}}, \quad u_{i}^{a}G_{ii} = -\zeta_{\mathcal{A}}^{a}\epsilon_{i}^{\mathcal{A}},$$

with the 'super' spinor fulfilling the polarized scattering equations (2.3) in unified notation

$$\langle v_i \kappa_i^{\mathcal{A}} \rangle = \langle u_i \zeta^{\mathcal{A}}(\sigma_i) \rangle, \quad \zeta_{\mathcal{A}}^a(\sigma_i) = \sum_{j \neq i} \frac{u_j^a \epsilon_{j\mathcal{A}}}{\sigma_{ij}},$$

where $\kappa_i^{\mathcal{A}} = (\kappa_i^A, \zeta_i^{\mathcal{I}})$. Then the proof of factorization of the reduced determinants in [7] can be repeated basically word by word by replacing $(\epsilon^A, \lambda_A, \kappa^A)$ with $(\epsilon^A, \zeta_A, \kappa^A)$.

More effort will be spent on the proof of factorization at the one-loop level in the next section.

3 Loop Amplitudes and Factorization

When calculating multi-loop amplitudes in QFT, this is typically done assuming generalized unitarity [4]. On the other hand, in this work using a model with a Lagrangian, consistency requires that this model exhibits proper factorization of scattering amplitudes to justify using generalized unitarity.

Before evaluating one-loop amplitudes it is interesting to look at partition functions. The model has two groups of 16 bosonic and 24 fermionic spinors each, one representing twistors and one representing dual twistors, 8 bosonic and 6 fermionic ghost-antighost pairs, and additionally 6 bosonic scalars or ghost-antighost pairs, so in balance the partition function is the same as for the RNS ambitwistor string [16] when the spin structures of the two particle groups are taken independently, but is modular invariant even when they are selected to be the same, because of the modularity of $\eta(\tau)^{-24}(\theta_2(\tau)^8 + \theta_3(\tau)^8 + \theta_4(\tau)^8)$ of weight -8. On the other hand, because of Jacobi's abstruse identity $\theta_2(\tau)^4 - \theta_3(\tau)^4 + \theta_4(\tau)^4 = 0$ in the former case one obtains a zero cosmological constant at the one-loop level like for RNS, which is preferable.

Moving on to the one-loop scattering amplitudes themselves, the scattering equations

(2.3) change to

$$\langle v_r \kappa_r^A \rangle = \langle u_r \lambda^A(\sigma_r) \rangle , \quad \lambda_A^a(\sigma_r) = \lambda_{0A}^a + \sum_{l \neq r} u_l^a \epsilon_{lA} \frac{S_\alpha(\sigma_r, \sigma_l; \tau)}{\sqrt{\mathrm{d}\sigma_r} \sqrt{\mathrm{d}\sigma_l}},$$

$$\langle v_r \zeta_r^{\mathcal{I}} \rangle = \langle u_r \eta^{\mathcal{I}}(\sigma_r) \rangle , \quad \eta_{\mathcal{I}}^a(\sigma_r) = \eta_{0\mathcal{I}}^a + \sum_{l \neq r} u_l^a q_{l\mathcal{I}} \frac{S_\alpha(\sigma_r, \sigma_l; \tau)}{\sqrt{\mathrm{d}\sigma_r} \sqrt{\mathrm{d}\sigma_l}}.$$
(3.1)

where λ_{0A}^a and $\eta_{0\mathcal{I}}^a$ are zero modes on an odd spin structure $\alpha = 1$ but generally do not exist on even spin structures $\alpha = 2, 3, 4$, and $S_{\alpha}(\sigma, \tau)$ is the Szegö kernel for spin structure α . Zero modes of the auxiliary fields $\rho_{1,2}^a$ and τ^a are not allowed as these fields do not contribute to the spectrum and, therefore, are only allowed as virtual fields, so their zero modes are omitted, but all propagators in the amplitudes, including the ones of the auxiliary fields, are changed to Szegö kernels.

One important change for the one-loop level is that there is only one zero mode of the c ghost left and none for the ghosts of the supersymmetric gauge symmetries, i.e. only one fixed vertex operator with regard to the conformal symmetry.

It follows that for even spin structures the amplitude looks like (2.4) with $SL(2, \mathbb{C})_{\sigma} \times SL(2, \mathbb{C})_u$ replaced with $GL(1)_{\sigma}$, all propagators $1/\sigma_{rl}$ with Szegö kernels $S_{\alpha}(\sigma_r, \sigma_l; \tau)/(\sqrt{d\sigma_r}\sqrt{d\sigma_l})$, reduced determinants with their full versions, and with an additional integration over the torus modulus τ . Because the partition function and the Szegö kernels are modular invariant so is the amplitude, like for the RNS ambitwistor string. On the other hand, it becomes immediately clear that all these determinants still have co-rank 2 and vanish, i.e. the even spin structures do not contribute to the one-loop scattering amplitude. Although this seems surprising, one should keep in mind, that in contrast to the conventional RNS string the spectrum of the current model is the same for the NS and R sector [14], so it is understandable that only the odd spin structure contributes a non-zero amplitude, saved from vanishing by the zero modes, as will be now shown explicitly.

The one-loop amplitude for odd spin structure becomes

$$\mathcal{A}_{n}^{1-\text{loop}} = \delta^{4} \left(\sum_{i=1}^{n} \langle \kappa_{i}^{\alpha} \tilde{\kappa}_{i}^{\dot{\alpha}} \rangle \right) \int d\mu_{n|1-\text{loop}}^{\text{pol}} \det H \det G \ e^{\frac{1}{2} \langle \eta_{0\mathcal{I}} \eta_{0}^{\mathcal{I}} \rangle + F_{\mathcal{N}}^{1-\text{loop}}}$$

$$d\mu_{n|1-\text{loop}}^{\text{pol}} := \frac{\prod_{l=1}^{n} d\sigma_{l} d^{2} u_{l} d^{2} v_{l} \ d\tau}{\text{vol}(GL(1)_{\sigma})} d^{8} \lambda_{0A}^{a} d^{2\mathcal{N}} \eta_{0\mathcal{I}}^{a} \prod_{r=1}^{n} \bar{\delta}(\langle v_{r} \epsilon_{r} \rangle - 1) \bar{\delta}^{4} (\langle u_{r} \lambda^{A} \rangle - \langle v_{r} \kappa_{r}^{A} \rangle),$$

$$F_{\mathcal{N}} = \sum_{i < j} \langle u_{i} u_{j} \rangle \ q_{i\mathcal{I}} q_{j}^{\mathcal{I}} \frac{S_{\alpha}(\sigma_{i}, \sigma_{j}; \tau)}{\sqrt{d\sigma_{i}} \sqrt{d\sigma_{j}}} + \sum_{i=1}^{n} \langle u_{i} \eta_{0\mathcal{I}} \rangle \ q_{i}^{\mathcal{I}} - \frac{1}{2} \sum_{i=1}^{n} \langle \xi_{i} v_{i} \rangle \ q_{i\mathcal{I}} q_{i}^{\mathcal{I}},$$

$$H_{ij} = \epsilon_{iA} \epsilon_{j}^{A} \frac{S_{\alpha}(\sigma_{i}, \sigma_{j}; \tau)}{\sqrt{d\sigma_{i}} \sqrt{d\sigma_{j}}}, \qquad u_{i}^{a} H_{ii} = -\lambda_{A}^{a} \epsilon_{i}^{A},$$

$$(3.2)$$

$$G_{ij} = (\epsilon_{iA}\epsilon_j^A + q_{i\mathcal{I}}q_j^{\mathcal{I}}) \frac{S_{\alpha}(\sigma_i, \sigma_j; \tau)}{\sqrt{\mathrm{d}\sigma_i}\sqrt{\mathrm{d}\sigma_j}}, \quad u_i^a G_{ii} = -\lambda_A^a \epsilon_i^A - \eta_{\mathcal{I}}^a q_i^{\mathcal{I}},$$

where λ_A^a and η_I^a are to be inserted from (3.1) and momentum conservation is expressed explicitly with a delta function, considered to come from the integration over the zero modes μ_0^{aA} . An additional factor $\exp(\frac{1}{2}\langle\eta_{0\mathcal{I}}\eta_0^{\mathcal{I}}\rangle)$ has been added to $\exp(F_{\mathcal{N}})$ to bring the integration over $\eta_{0\mathcal{I}}^a$ into normal form [17]². It could also be regarded as substitute for the same effect as a delta function arising from the integration over the zero modes $\tilde{\eta}_0^{a\mathcal{I}}$ when working in the LG representation, cf. (2.4).

Including the zero modes of μ_0^{aA} and $\tilde{\eta}_0^{a\mathcal{I}}$ which already have been integrated out, there are 16 eight bosonic and 32 fermionic spinor zero modes resulting in a modular weight of -8, making the amplitude modular invariant and UV-finite, when taking the 8 bosonic contributions from the momentum integrations over the loop and disregarded zero modes of the bc ghosts into account, like for the partition function.

3.1 Nonseparating Degeneration

This section checks whether the one-loop amplitude (3.2) factorizes properly for a nonseparating degeneration of the worldsheet. Pinching a nonseparating cycle occurs in the limit $q = e^{2\pi i \tau} \to 0$. One obstruction to dealing with this limit by simply taking the residue at 0 of the integrand as a function in q is that the integrand is not an analytic function in q because the torus Szegö kernel $S_1(\sigma_1, \sigma_2; \tau)$ contains an $Im(\tau)$ -dependent term:

$$S_1(\sigma_1, \sigma_2; \tau) = \left(\frac{\theta'_1(\sigma_{12}, \tau)}{\theta_1(\sigma_{12}, \tau)} + 4\pi \frac{\operatorname{Im}(\sigma_{12})}{\operatorname{Im}(\tau)}\right).$$

The behavior of S_1 can be examined by modeling the torus as a Riemann sphere with a handle far away from all punctures, with ends located at two reference points σ_+ and σ_- , i.e. these two points are added as additional punctures, modeled in terms of local coordinates by

$$(\sigma - \sigma_+)(\sigma - \sigma_-) = q.$$

Then, because according to [18] all Szegö kernels are holomorphic in q for small enough q, the integrand in (3.2) becomes a manifestly meromorphic function in q.

The asymptotic behavior of the Szegö kernel is [19, 20]

$$S_1(\sigma_{ij};\tau)\sqrt{\mathrm{d}\sigma_i\mathrm{d}\sigma_j} \to \frac{\sqrt{\mathrm{d}z_i\mathrm{d}z_j}}{2z_{ij}}\left(\sqrt{\frac{z_{i-}z_{j+}}{z_{i+}z_{j-}}} + \sqrt{\frac{z_{i+}z_{j-}}{z_{i-}z_{j+}}}\right) + O(q), \tag{3.3}$$

²A similar exponential factor appears in [12, 10].

in terms of the coordinates $z=e^{2\pi i(\sigma-\frac{\tau}{2})}$ on the Riemann sphere. As $q\to 0$, the handle containing z_+ and z_- stretches and pinches, such that the values of the square roots in (3.3) can be replaced by 1 up to O(q), i.e. S_1 becomes just the genus 0 propagator in this limit.

On the Riemann sphere the torus zero modes λ_0 and η_0 are lifted up to fields λ_0/z and η_0/z . To proceed, the points z_+ and z_- are chosen temporarily as 0 and ∞ . Then the right hand side of the scattering equations (3.1) for λ can be rewritten as, in abuse of notation changing back from z to σ ,

$$\lambda^{aA}(\sigma) = \frac{\lambda_0^{aA}}{\sigma} + \sum_l \frac{u_l^a \epsilon_l^A}{\sigma - \sigma_l} + O(q) = \frac{\langle u_+^a \epsilon_+^A + u_-^a \epsilon_-^A \rangle}{\sigma} + \sum_l \frac{u_l^a \epsilon_l^A}{\sigma - \sigma_l} + O(q),$$

where λ_0^{aA} has been expanded into a basis of little spinors u_{\pm}^a with $\langle u_+ u_- \rangle \neq 0$. Now reverting to general σ_+ and σ_- , designating $u_{\pm}^a \epsilon_{\pm}^A$ to belong to σ_{\pm} by requiring that $\langle u_{\pm} \lambda^A(\sigma_{\pm}) \rangle < \infty$, defining a spinor basis ϵ_{\pm}^a with $\langle \epsilon_+ \epsilon_- \rangle = 1$ and momenta $\kappa_{\pm A}^a$ with $\kappa_{+A}^1 = \kappa_{-A}^1, \kappa_{+A}^2 = -\kappa_{-A}^2$ such that $\epsilon_{\pm}^A = \langle \epsilon_{\pm} \kappa_{\pm}^A \rangle$, and finally viewing σ_{\pm} as additional punctures σ_{n+1} and σ_{n+2} one gets

$$\lambda^{aA}(\sigma) = \sum_{l=1}^{n+2} \frac{u_l^a \epsilon_l^A}{\sigma - \sigma_l} + O(q).$$

A similar procedure for η_0 gives

$$\eta^{a\mathcal{I}}(\sigma) = \sum_{l=1}^{n+2} \frac{u_l^a q_l^{\mathcal{I}}}{\sigma - \sigma_l} + O(q) \text{ with } q_{n+\frac{1}{2}}^{\mathcal{I}} = \langle \epsilon_{n+\frac{1}{2}} \zeta_{n+\frac{1}{2}}^{\mathcal{I}} \rangle, \zeta_{n+1\mathcal{I}}^1 = \zeta_{n+2\mathcal{I}}^1, \zeta_{n+1\mathcal{I}}^2 = -\zeta_{n+2\mathcal{I}}^2.$$
 (3.4)

This way scattering equations for n+2 particles have been obtained, so far fulfilled by the first n particles. Now the measure needs to be modified. This is done by setting

$$\begin{split} \delta^4\!\!\left(\sum_{i=1}^n \left\langle \kappa_i^\alpha \tilde{\kappa}_i^{\dot{\alpha}} \right\rangle \right) & \frac{\mathrm{d}^8 \lambda_{0A}^a \, \mathrm{d}^{2\mathcal{N}} \eta_{0\mathcal{I}}^a}{\mathrm{vol}(\mathrm{GL}(1)_\sigma)} \frac{\mathrm{d}q}{q} = \int \!\! \frac{\prod_{l=n+1}^{n+2} \mathrm{d}\sigma_l \, \mathrm{d}^2 u_l \, \mathrm{d}^2 v_l}{\mathrm{vol}(\mathrm{SL}(2,\mathbb{C})_\sigma \! \times \! \mathrm{SL}(2,\mathbb{C})_u)} \\ & \frac{\prod_{l=n+1}^{n+2} \mathrm{d}^4 \epsilon_l^A \, \mathrm{d}^{\mathcal{N}} q_l^{\mathcal{I}}}{\left\langle u_{n+1} u_{n+2} \right\rangle^4} \prod_{r=n+1}^{n+2} \bar{\delta}(\!\langle v_r \epsilon_r \rangle \! - \! 1) \bar{\delta}^4 (\!\langle u_r \lambda^A \rangle \! - \! \langle v_r \kappa_r^A \rangle). \end{split}$$

That the left hand side is equal to the right hand side can be seen as follows: on the tree level two more σ_l values can be fixed (as was done with σ_{n+1} and σ_{n+2}) and three of the (u_l^a, v_l^a) variables (e.g. by choosing $u_{n+1}^a = (1,0), u_{n+2}^a = (u_{n+2},1)$), increasing the denominator to $\text{vol}(\text{SL}(2,\mathbb{C})_{\sigma}\times \text{SL}(2,\mathbb{C})_u)$ and fixing 5 of the 10 delta functions on the right. On the left $\mathrm{d}^8\lambda_{0A}^a\mathrm{d}q/q$ is equivalent to $\mathrm{d}^4l_\mu\delta(l^2-m^2)\mathrm{d}q/q\sim -(\mathrm{d}^4l_\mu/(l^2-m^2))\mathrm{d}q\bar{\delta}(q)$, and because momentum conservation follows from the scattering equations for the n+2 particles the

remaining 5 delta functions on both sides balance each other out. Finally $\langle u_{n+1}u_{n+2}\rangle^{4-\mathcal{N}}$ = $\langle u_{n+1}u_{n+2}\rangle^{-4}$ is the Jacobian going from $d^8\lambda_0 d^{2\mathcal{N}}\eta_0$ to $\prod_l d^4\epsilon_l d^{\mathcal{N}}q_l$. $\mathcal{N}=8$ is crucial here.

It remains to check $F_{\mathcal{N}}$. Inserting

$$\eta_0(\sigma) = u_{n+1}q_{n+1}/(\sigma - \sigma_{n+1}) + u_{n+2}q_{n+2}/(\sigma - \sigma_{n+2})$$
(3.5)

from (3.4) into F_N in (3.2) gives F_N of the tree amplitude (2.7) up to a missing piece:

$$F_{\mathcal{N}} = \sum_{1 \leq i < j \leq n+2} \frac{\langle u_i u_j \rangle}{\sigma_{ij}} q_{i\mathcal{I}} q_j^{\mathcal{I}} - \frac{1}{2} \sum_{i=1}^{n+2} \langle \xi_i v_i \rangle q_{i\mathcal{I}} q_i^{\mathcal{I}} - C$$

$$C = \frac{\langle u_{n+1} u_{n+2} \rangle}{\sigma_{n+1n+2}} q_{n+1\mathcal{I}} q_{n+2}^{\mathcal{I}} - \frac{1}{2} \sum_{i=n+1}^{n+2} \langle \xi_i v_i \rangle q_{i\mathcal{I}} q_i^{\mathcal{I}}.$$

The second term in C vanishes because of $\langle \xi v \rangle \epsilon^a = v^a - \xi^a$ and the special relationship between values for ζ_{n+1} and ζ_{n+2} in (3.4). The first term can be recognized as arising from $\frac{1}{2}\langle \eta_{0\mathcal{I}}\eta_0^{\mathcal{I}}\rangle$ when going from the torus to the Riemann sphere:

$$\frac{1}{2}\langle \eta_{0\mathcal{I}}\eta_{0}^{\mathcal{I}}\rangle \xrightarrow{(3.5)} \frac{1}{2}(q_{n+1}^{\mathcal{I}}\langle u_{n+1}\eta_{0\mathcal{I}}(\sigma_{n+1})\rangle + q_{n+2}^{\mathcal{I}}\langle u_{n+2}\eta_{0\mathcal{I}}(\sigma_{n+2})\rangle) = \frac{\langle u_{n+1}u_{n+2}\rangle}{\sigma_{n+1}n+2}q_{n+1\mathcal{I}}q_{n+2}^{\mathcal{I}}.$$

This proves that the one-loop amplitude (3.2) factorizes correctly into a tree amplitude (2.7) where two additional punctures are connected through a complete set of intermediate physical states:

$$\begin{split} \mathcal{A}_{n}^{\text{1-loop}} &= \int \prod_{l=n+1}^{n+2} \mathrm{d}^{4} \epsilon_{l}^{A} \, \mathrm{d}^{\mathcal{N}} q_{l}^{\mathcal{I}} \, \, \mathcal{A}_{n+2}^{\text{tree}}, \qquad \left\langle \epsilon_{n+1} \epsilon_{n+2} \right\rangle = 1, \\ \epsilon_{n+\frac{1}{2}}^{A} &= \left\langle \epsilon_{n+\frac{1}{2}} \kappa_{n+\frac{1}{2}}^{A} \right\rangle, \, \kappa_{n+1A}^{1} = \kappa_{n+2A}^{1}, \, \kappa_{n+1A}^{2} = -\kappa_{n+2A}^{2}, \\ q_{n+\frac{1}{2}}^{\mathcal{I}} &= \left\langle \epsilon_{n+\frac{1}{2}} \zeta_{n+\frac{1}{2}}^{\mathcal{I}} \right\rangle, \, \zeta_{n+1\mathcal{I}}^{1} = \zeta_{n+2\mathcal{I}}^{1}, \, \zeta_{n+1\mathcal{I}}^{2} = -\zeta_{n+2\mathcal{I}}^{2}. \end{split}$$

3.2 Separating Degeneration

Pinching a separating cycle involves the degeneration of the Riemann surface Σ of genus $g = g_L + g_R$ into a surface Σ_L of genus g_L and a surface Σ_R of genus g_R and can again be modeled in terms of local coordinates by

$$(\sigma - \sigma_L)(\sigma - \sigma_R) = q,$$

where σ_L is an extra puncture on Σ_L and σ_R one on Σ_R with their neighborhoods sewed together and both points identified in the pinching limit $q \to 0$.

Proper factorization requires to show that during such a separating degeneration an n-particle amplitude on Σ can be represented as an n_L+1 -particle amplitude on Σ_L and an n_R+1 -particle amplitude on Σ_R with $n=n_L+n_R$ and additional punctures σ_L and σ_R which are connected through insertion of a complete set of physical states.

First it needs to be recognized that when Σ has an odd spin structure, then the spin structure of Σ_L can only be odd if Σ_R has even spin structure or genus $g_R = 0$ and vice versa. Therefore, because the beginning of this section has shown that one-loop amplitudes vanish on even spin structures and the next subsection will argue that the same is generally true for all loop amplitudes, a separating degeneration can only occur when one side is a tree amplitude.

Now proper factorization for separate degeneration of a one-loop amplitude into a tree and a one-loop amplitude can be simply demonstrated by first factorizing the one-loop amplitude into a tree amplitude using nonseparating factorization, applying factorization of tree amplitudes from subsection 2.2 with the two additional punctures from the loop amplitude necessarily ending up on the same side, and then reversing the nonseparating degeneration for that side, possibly with a reduced number of punctures.

3.3 Higher Loops and Cosmological Constant

Looking first at partition functions at two or more loops, there is a strictly positive number of zero modes of anti-ghosts and, therefore, at the minimum one picture-changing operator for every supersymmetric gauge symmetry resulting either in zero because of no contractions possible or in a matrix with a determinant that vanishes on even spin structures like at the one-loop level. Because the partition function on odd spin structures is always zero due to the presence of a Dirac zero mode, the cosmological constant vanishes at every order of perturbation theory (this assumes that at the one-loop level the Jacobi abstruse identity can be applied on even spin structures). Of course, non-perturbative effects like IR renormalization for massless particles cannot be excluded and might give the cosmological constant a non-vanishing value.

Multi-loop amplitudes with a non-zero number of vertex operators will again generally vanish on even spin structures and develop a single zero mode for the λ_A^a and η_N^a fields on odd spin structures. But assuming the validity of unitary factorization it can be seen that not even all odd spin structures contribute. The total number of odd spin structures can be calculated by gluing Riemann surfaces of genus 1 to an existing surface of genus g:

$$\#_{\text{odd}} = \sum_{m \text{ odd}} \binom{g}{m} 1^m 3^{g-m} = 2^{g-1} (2^g - 1)$$
 (3.6)

When m is larger than 1 in a term of (3.6) there is no way to use generalized unitarity and factorize down to a tree amplitude using nonseparating degeneration without going through an even spin structure as intermediate step, i.e. such a spin structure cannot contribute to multi-loop amplitudes. It follows that only $g \, 3^{g-1}$ odd spin structures have non-vanishing multi-loop amplitudes.

It is interesting to note that many of the results in this article also apply to the original massive ambitwistor model of [7, 9, 10]. This includes zero cosmological constant, non-vanishing amplitudes only on the same selected subset of odd spin structures, and proper factorization of one-loop amplitudes.

4 Compton Scattering Using Massive Twistor Strings

In this section it is important to note that in the following the usual convention is used that in amplitudes all legs are taken to be outgoing, i.e. incoming particles will have their momenta and helicities reversed.

The 4-point amplitude for the original massive ambitwistor model of [7, 9, 10] is calculated to be [7]

$$M_{4} = \frac{\langle 1234 \rangle^{4}}{(2p_{1} \cdot p_{2})(2p_{2} \cdot p_{3})(2p_{2} \cdot p_{4})} e^{F_{\mathcal{N}}},$$

$$\langle 1234 \rangle = \varepsilon^{ABCD} e_{1A} e_{2B} e_{3C} e_{4D},$$

$$F_{\mathcal{N}} = \frac{1}{2} \sum_{i,j=1}^{4} \frac{\langle u_{i} u_{j} \rangle}{\sigma_{ij}} q_{i\mathcal{I}} q_{j}^{\mathcal{I}} - \frac{1}{2} \sum_{i=1}^{4} \langle \xi_{i} v_{i} \rangle q_{i\mathcal{I}} q_{i}^{\mathcal{I}}.$$

$$(4.1)$$

More details can be found in $[7]^3$.

The amplitude (4.1) is valid for 4 massive particles up to spin 2. It is now specialized to the case of Compton scattering of a massless particle with momenta $p_2 \to k_2, p_3 \to k_3$ against a heavy massive particle with mass M much larger than the energy $\hbar\omega$ of the massless particle. Switching from the polarized formalism to a standard little-group based amplitude, it can readily be seen that the amplitude becomes either

$$M_4^{+-} = \frac{(\langle 43 \rangle [12] - \langle 13 \rangle [42])^4}{(2p_1 \cdot k_2)(2k_2 \cdot k_3)(2k_2 \cdot p_4)} e^{F_{\mathcal{N}}}$$
(4.2)

³In [7] the amplitude is given for 6 dimensions, with $\langle 1234 \rangle^4$ replaced by $\langle 1234 \rangle^2 [1234]^2$, and F_N with $F_N + \tilde{F}_N$, but the contributions become equal via dimensional reduction.

or

$$M_4^{++} = \frac{\langle \mathbf{14} \rangle^4 [23]^4}{(2p_1 \cdot k_2)(2k_2 \cdot k_3)(2k_2 \cdot p_4)} e^{F_N}, \tag{4.3}$$

depending on whether the incoming massless particle with negative helicity preserves or flips the helicity, respectively⁴. In order to simplify formulas, the same notation as in [2] is used by bolding the massive spinor helicity variables and suppressing the little group indices. (4.2) matches the amplitude (5.35) in [2], up to the supersymmetric exponential factor.

The supersymmetric factor is useful when trying to switch from spin 2 to lower spin particles, by taking derivatives with respect to the supermomenta $q_{\mathcal{I}}$ and setting them to zero. It was shown in [7] that in order to lower the spin of the massless particle it is required to multiply the amplitude with a factor of

$$\frac{\langle u_2 u_3 \rangle}{\sigma_{23}} = -\frac{\langle k_2 14 \rangle}{\langle 1234 \rangle}$$

for every spin $\frac{1}{2}$ downward change and for the massive particle with a factor of

$$\frac{\langle u_1 u_4 \rangle}{\sigma_{14}} = -\frac{\langle p_1 23 \rangle}{\langle 1234 \rangle}.$$

Because $\langle k_2 14 \rangle = 0$ when the massless particle flips its helicity if follows that the amplitude (4.3) vanishes for massless particles with spin<2. Therefore, in this model only a graviton can scatter with reversing its helicity [21].

Concerning the amplitude that conserves helicity, when disregarding the difference between $p_1 \cdot k_2$ and $p_1 \cdot k_3$ (valid in the classical limit $\hbar \omega / M \to 0$) in equation (5.26) of [7] when applied to 4 dimensions:

$$\langle k_2 14 \rangle \langle p_1 23 \rangle = -2p_1 \cdot k_3 \ e_{1A} e_2^A e_{4B} e_3^B - 2p_1 \cdot k_2 \ e_{1A} e_3^A e_{4B} e_2^B \to -2p_1 \cdot k_3 \langle 1234 \rangle, \tag{4.4}$$

and switching to the little-group helicity-based formalism the amplitude (4.2) becomes

$$M_4^{+-} = \frac{\langle 3|p_1|2|^{2h} (2p_1 \cdot k_3)^{2(2-h)}}{(2p_1 \cdot k_2)(2k_2 \cdot k_3)(2k_2 \cdot p_4)} \left(\frac{\langle 43 \rangle [12] - \langle 13 \rangle [42]}{\langle 3|p_1|2]}\right)^{2s},\tag{4.5}$$

where s is the spin of the massive particle and h the spin of the massless particle.

What is different with the model of this article? Assume first that the fermionic contribution in $\det' G$ in (2.6) is for a particle of spin 3/2 ('gravitino') or spin 1 ('gluon') still

⁴Remember from the note at the beginning of the section that because of the convention to regard all particles outgoing, conserving/flipping the helicity means that the incoming and outgoing massless particles have the opposite/same helicity.

preserving helicity. This represents a massive EYM amplitude as mentioned in subsection 2.1 with a 2-particle PT factor for the 'gluon' and an open trace for the 'gravitino':

$$\det' G \to \begin{cases} \left(\frac{q_{2\mathcal{I}}q_{3}^{\mathcal{I}}}{\sigma_{23}}\right)^{2} \det H_{[32]}^{[23]} = \det' H\left(\frac{\langle u_{2}u_{3}\rangle q_{2\mathcal{I}}q_{3}^{\mathcal{I}}}{\sigma_{23}}\right)^{2}, & \text{'gluon'}, \\ \frac{q_{2\mathcal{I}}q_{3}^{\mathcal{I}}}{\sigma_{23}} \frac{\det H_{[32]}^{[23]}}{\langle u_{3}u_{2}\rangle} = \det' H\frac{\langle u_{2}u_{3}\rangle q_{2\mathcal{I}}q_{3}^{\mathcal{I}}}{\sigma_{23}}, & \text{'gravitino'}. \end{cases}$$
(4.6)

It turns out that the amplitude is the same as the helicity-conserving case (4.5), which is good for consistency. Because a PT factor for massless particles needs to have at least one particle of each helicity, the 'gluon' case cannot flip the helicity, but it is possible for a 'gravitino':

$$\det G_{[12]}^{[34]} = G_{13}G_{24} - G_{14}G_{23} = \frac{\epsilon_1^A \epsilon_{4A} q_{2\mathcal{I}} q_3^{\mathcal{I}}}{\sigma_{14}\sigma_{32}} = \frac{q_{2\mathcal{I}} q_3^{\mathcal{I}}}{\epsilon_2^A \epsilon_{3A}} \det H_{[12]}^{[34]}.$$

Therefore, going from spin 2 to 3/2 just lowers the power of [23] in (4.3), but it also lowers the power in the gravitational coupling, in contrast to the helicity-preserving case. The same is true for the massive particle:

$$M_4^{++} = \frac{\langle \mathbf{14} \rangle^{2s} [23]^{2h}}{(2p_1 \cdot k_2)(2k_2 \cdot k_3)(2k_2 \cdot p_4)}, \quad 3/2 \le h, s \le 2.$$

Finally one can consider both, the massless and the massive, particles as 'gluons' leading to replace $\det'G$ with a PT factor for two incoming and two outgoing particles, i.e. the situation is now described by the Coulomb branch of SYM for the scattering between 'gluons' and 'W bosons' (see 2.1):

$$< Wg^{+}g^{+}\bar{W}> = \frac{\langle \mathbf{14} \rangle^{2} [23]}{(2k_{2} \cdot p_{4}) \langle 23 \rangle}, \quad < Wg^{+}g^{-}\bar{W}> = \frac{(\langle \mathbf{43} \rangle [12] - \langle \mathbf{13} \rangle [42])^{2}}{(2k_{2} \cdot p_{4})(2k_{2} \cdot k_{3})}$$

which is consistent with corresponding results in [1, 3, 10].

For gravitational Compton scattering on a Kerr black hole, to make contact with the results in [22, 23, 21, 24], the kinematic variables are now taken to be the same as in [21, 24]. The black hole is treated as a massive spin particle with mass M and a spin vector $a^{\mu} = s^{\mu}/M = (0, a_x, a_y, a_z)$. The incoming/outgoing momenta of the scattered particle are still labeled as k_2/k_3 and of the black hole as p_1/p_4 . In the chosen reference frame the black hole is taken to be initially at rest and the scattering process to happen in the x-z plane. This leads to the following explicit representation of the kinematic variables:

$$\begin{split} p_1^{\mu} &= (M,0,0,0), \\ k_2^{\mu} &= \hbar \omega (1,0,0,1), \\ k_3^{\mu} &= \gamma \, \hbar \omega (1,\sin\theta,0,\cos\theta), \\ \gamma &= 1/(1+2\frac{\hbar \omega}{M}\sin^2\frac{\theta}{2}) \\ p_4^{\mu} &= p_1^{\mu} + k_2^{\mu} - k_3^{\mu} \, . \end{split}$$

Here θ is the scattering angle and the value of γ is determined by the on-shell condition $p_4^2 = M^2$ for the outgoing massive momentum. It should be kept in mind that this setup violates the standard notation mentioned at the beginning of the section, by reversing the incoming momenta p_1 and k_2 .

It will be convenient to introduce a basis of spinor helicity variables in terms of θ to represent the momentum and the polarization matrices of the outgoing massless particle:

$$[3] = \sqrt{2\gamma\hbar\omega} \left(\cos\frac{\theta}{2}, \sin\frac{\theta}{2}\right), \qquad [\mu] = \left(-\sin\frac{\theta}{2}, \cos\frac{\theta}{2}\right),$$

$$|3\rangle = \sqrt{2\gamma\hbar\omega} \left(\cos\frac{\theta}{2}, \sin\frac{\theta}{2}\right), \qquad |\mu\rangle = \left(-\sin\frac{\theta}{2}, \cos\frac{\theta}{2}\right),$$

$$k_{3\alpha\dot{\alpha}} = |3\rangle_{\alpha}[3|_{\dot{\alpha}},$$

$$\epsilon_{3\alpha\dot{\alpha}}^{+} = \sqrt{2}\frac{|\mu\rangle_{\alpha}[3|_{\dot{\alpha}}}{\langle\mu3\rangle}, \qquad \epsilon_{3\alpha\dot{\alpha}}^{-} = \sqrt{2}\frac{|3\rangle_{\alpha}[\mu|_{\dot{\alpha}}}{[3\mu]},$$

where $|\mu\rangle$ and $[\mu|$ serve as reference spinors. The incoming massless states associated to k_2 are recovered by setting $\theta=0$. Other choices for the reference spinors are allowed. Actually, the reference spinor $|\mu\rangle$ in ϵ_3^+ and ϵ_2^+ could be replaced by $|2\rangle$ and $|3\rangle$, respectively, and $[\mu|$ in ϵ_3^- and ϵ_2^- by [2] and [3], respectively, a choice which will be made presently.

In this kinematics the classical limit $(s \to \infty, \gamma \to 1)$ of M_4^{+-} in (4.5) can be written as [22, 23, 24]

$$M_4^{+-} = \frac{M^2(\cos\frac{\theta}{2})^{2h}}{\sin^2\frac{\theta}{2}} \exp[(k_3 - k_2 + 2w) \cdot a]$$

$$= \frac{M^2(\cos\frac{\theta}{2})^{2h}}{\sin^2\frac{\theta}{2}} \exp\left[-\frac{\hbar\omega}{M} \left(2a_z \sin^2\frac{\theta}{2} - a_x \sin\theta + 2(a_x - ia_y) \tan\frac{\theta}{2}\right)\right],$$
(4.7)

where the vector w is defined as $w = -p_1 \cdot k_3 \frac{|3\rangle[2]}{\langle 3|p_1|2|} = -\frac{p_1 \cdot k_3}{p_1 \cdot \epsilon_3} \epsilon_3^-$.

Amplitudes (4.7) and (4.5) for s > h exhibit a spurious pole for $\langle 3|p_1|2 \rangle = 2M\hbar\omega \cos \frac{\theta}{2}$ that stems from the vector w and needs to be remedied with help of contact terms [24]. However, the issue of contact terms will not be considered here. But it is intriguing to note that the spurious pole is missing in the reverse-helicity scattering amplitude M_4^{++} and that indeed according to (4.4) w can be rewritten as $w = \frac{\langle k_2 14 \rangle}{2\langle 1234 \rangle} |3\rangle [2| = -\langle u_2 u_3 \rangle/(2\sigma_{23}) |3\rangle [2|$ which vanishes for massless particles when flipping the helicity, that is, from the point of view of the ambitwistor string model the appearance or absence of the spurious pole is directly related to solutions of the polarized scattering equations.

In summary, although this section has not provided any new results (besides maybe the gravitino amplitudes), it was interesting to show generally and in the context of a black hole how, starting from a Lagrangian, the ambitwistor string models with a single amplitude (4.1) can describe multiple aspects of the Compton scattering of a massless particle of various spin hitting a massive target particle with higher spin.

5 Summary and Outlook

This work presented an anomaly-free extension to the massive ambitwistor string found in [8, 9, 10] as a unified model for supergravity and the Coulomb branch of SYM. The all-multiplicity tree amplitudes were shown to have the expected massless limit, to include EYM amplitudes for multiple gluon traces, and to have proper unitary factorization. At every order of perturbation theory the cosmological constant is zero and even spin structures of genus $g \ge 1$ Riemann surfaces do not contribute to loop amplitudes. Even odd spin structures are restricted to contain a single odd cycle. The all-multiplicity one-loop amplitude on an odd spin structure was calculated and established to have unitary factorization for separating and nonseparating degeneration. Then Compton scattering of a massless particle against a massive target was used to obtain scattering amplitudes for both cases of conserved and reversed helicity. They were compatible with results in the literature. One surprise score might be counted for the non-vanishing reverse-helicity amplitude of gravitino scattering.

Future work would include the calculation of higher loop amplitudes together with the demonstration of proper factorization. Because the model will be IR-divergent when massless particles like the graviton and photon are included, IR-renormalization will be required. In order to do that correctly, a string field theory might need to be developed. It would be interesting to see how model features would be affected by renormalization and whether for instance the cosmological constant could pick up a non-vanishing value. Further, the $\mathcal{N}=8$ supergravity spectrum with $\mathrm{SU}(4)\times\mathrm{SU}(4)$ R-symmetry should be studied for various symmetry breakdown scenarios to get more realistic spectra. Regarding the spectrum, one issue that has come up in [14] is that in the NS sector the spectrum is truncated because of the absence of oscillator zero modes for the twistors, and it was argued that the issue might be solved through spectral flow. The current work showed that because of the vanishing of amplitudes on even spin structures the spectrum in the R sector is sufficient up to at least the one-loop level, but is not clear whether the NS sector obstruction would not present itself again at higher loop level. This is another item to look into in more detail.

⁵Note that the $\mathcal{N}=8$ supergravity spectrum contains exactly the spin $\frac{1}{2}$ content for all the fermions of the standard model and also for 8 massive gravitinos, a fact that is well known [25, 26].

And, of course, the origin of all the auxiliary spinors in the action (2.1) is still a mystery. They are excluded from the physical spectrum and the question arises whether they materialize as a ghost system used for mathematical convenience or whether they emanate from compactifying extra twistor space dimensions (in which case there still might not be a mandatory necessity to relate them to extra spacetime dimensions). More knowledge in this regard might also reveal the exact nature of the unknown six bosonic degrees of freedom that make the model anomaly-free.

References

- [1] N. Craig, H. Elvang, M. Kiermaier, and T. Slatyer, "Massive amplitudes on the Coulomb branch of N=4 SYM," JHEP 12 (2011) p.097, [1104.2050 [hep-th]].
- [2] N. Arkani-Hamed, T.-C. Huang, and Y.-t. Huang, "Scattering amplitudes for all masses and spins," JHEP 11 (2021) p.70, [1709.04891 [hep-th]].
- [3] F. Cachazo, A. Guevara, M. Heydeman, S. Mizera, J. H. Schwarz, and C. Wen, "The S Matrix of 6D Super Yang-Mills and Maximal Supergravity from Rational Maps," JHEP 09 (2018) p.125, [1805.11111 [hep-th]].
- [4] Z. Bern and Y.-t. Huang, "Basics of Generalized Unitarity," J. Phys. A 44 (2011) p.454003, [1103.1869 [hep-th]].
- [5] C. Wen and S.-Q. Zhang, "D3-Brane Loop Amplitudes from M5-Brane Tree Amplitudes," JHEP 07 (2020) p.098, [2004.02735 [hep-th]].
- [6] Y. Geyer and L. Mason, "Polarized Scattering Equations for 6D Superamplitudes," Phys. Rev. Lett. 122 (2019) p.101601, no. 10, [1812.05548 [hep-th]].
- [7] G. Albonico, Y. Geyer, and L. Mason, "Recursion and worldsheet formulae for 6d superamplitudes," JHEP 08 (2020) p.66, [2001.05928 [hep-th]].
- [8] Y. Geyer, L. Mason, and D. Skinner, "Ambitwistor strings in six and five dimensions," JHEP 08 (2021) p.153, [2012.15172 [hep-th]].
- [9] G. Albonico, Y. Geyer, and L. Mason, "From Twistor-Particle Models to Massive Amplitudes," SIGMA 18 (2022) p.45, [2203.08087 [hep-th]].
- [10] G. Albonico, Y. Geyer, and L. Mason, "Massive ambitwistor-strings; twistorial models," JHEP 01 (2024) p.127, [2301.11227 [hep-th]].
- [11] D. Skinner, "Twistor strings for $\mathcal{N}=8$ supergravity," JHEP **04** (2020) p.147, [1301.0868 [hep-th]].

- [12] T. Adamo, E. Casali, K. A. Roehrig, and D. Skinner, "On tree amplitudes of supersymmetric Einstein-Yang-Mills theory," JHEP 12 (2015) p.177, [1507.02207 [hep-th]].
- [13] Y. Geyer, A. E. Lipstein, and L. J. Mason, "Ambitwistor Strings in Four Dimensions," Phys. Rev. Lett. 113 (2014) p.81602, no. 8, [1404.6219 [hep-th]].
- [14] C. Kunz, "Spectrum of Massive and Massless Ambitwistor Strings," arXiv:2305.01734 [hep-th], 2023.
- [15] R. Penrose and W. Rindler, SPINORS AND SPACE-TIME. VOL. 2: SPINOR AND TWISTOR METHODS IN SPACE-TIME GEOMETRY. Cambridge Monographs on Mathematical Physics, Cambridge University Press, 4 1988.
- [16] T. Adamo, E. Casali, and D. Skinner, "Ambitwistor strings and the scattering equations at one loop," JHEP 04 (2014) p.104, [1312.3828 [hep-th]].
- [17] F. A. Berezin, "The method of second quantization," Pure Appl. Phys. 24 (1966) p.1–228.
- [18] M. P. Tuite and A. Zuevsky, "The Szegö Kernel on a Sewn Riemann Surface," Commun. Math. Phys. 306 (2011) p.617-645, [1002.4114 [math.QA]].
- [19] E. Casali and P. Tourkine, "Infrared behaviour of the one-loop scattering equations and supergravity integrands," JHEP 04 (2015) p.13, [1412.3787 [hep-th]].
- [20] Y. Geyer, Ambitwistor Strings: Worldsheet Approaches to perturbative Quantum Field Theories. PhD thesis, Oxford U., Inst. Math., 2016, [1610.04525 [hep-th]].
- [21] Y. F. Bautista, A. Guevara, C. Kavanagh, and J. Vines, "Scattering in black hole backgrounds and higher-spin amplitudes. Part I," JHEP 03 (2023) p.136, [2107.10179 [hep-th]].
- [22] A. Guevara, A. Ochirov, and J. Vines, "Scattering of Spinning Black Holes from Exponentiated Soft Factors," JHEP 09 (2019) p.56, [1812.06895 [hep-th]].
- [23] R. Aoude, K. Haddad, and A. Helset, "On-shell heavy particle effective theories," JHEP 05 (2020) p.51, [2001.09164 [hep-th]].
- [24] Y. F. Bautista, A. Guevara, C. Kavanagh, and J. Vines, "Scattering in black hole backgrounds and higher-spin amplitudes. Part II," JHEP 05 (2023) p.211, [2212.07965 [hep-th]].
- [25] K. A. Meissner and H. Nicolai, "Standard Model Fermions and N=8 supergravity," Phys. Rev. D 91 (2015) p.65029, [1412.1715 [hep-th]].
- [26] K. A. Meissner and H. Nicolai, "Standard Model Fermions and Infinite-Dimensional R-Symmetries," Phys. Rev. Lett. 121 (2018) p.91601, no. 9, [1804.09606 [hep-th]].