# Some remarks on almost projective envelopes

Xiaolei Zhang

School of Mathematics and Statistics, Shandong University of Technology Zibo 255049, China

Corresponding author, E-mail: zxlrghj@163.com

Wei Qi

School of Mathematics and Statistics, Shandong University of Technology Zibo 255049, China

E-mail: qwrghj@126.com

Dechuan Zhou Southwest University of Science and Technology Mianyang 621010, China E-mail: dechuan11119@sina.com

## Abstract

Let R be a commutative ring. An R-module M is said to be almost projective if  $\operatorname{Ext}_{R}^{1}(M, N) = 0$  for any  $R_{\mathfrak{m}}$ -module N and any maximal ideal  $\mathfrak{m}$  of R. In this paper, we investigate rings R over which every R-module has an almost projective envelope. We also characterize rings R over which every R-module has an almost projective envelope with some special properties.

*Key Words:* almost projective module, envelope, coherent ring, locally Artinian ring.

2020 Mathematics Subject Classification: 13C10; 13C11.

Throughout this paper, R is a commutative ring with identity and all R-modules are unitary. We always denote by Max(R) the spectrum of all maximal ideals of R, K. dim(R)the Krull dimension of R and w.gld(R) the weak global dimension of R.

The notion of locally free modules, in the sense that R-modules M satisfying that  $M_{\mathfrak{m}}$  is a free  $R_{\mathfrak{m}}$ -module for any maximal ideal  $\mathfrak{m}$  of R, is a key terminology to the areas of both commutative algebra and algebra geometry. Recently, Zhou et al. [14] found

that an *R*-module *M* is locally free if and only if  $\operatorname{Ext}_{R}^{1}(M, N) = 0$  for any  $R_{\mathfrak{m}}$ -module *N* and any maximal ideal  $\mathfrak{m}$  of *R*, and the later is called almost projective modules in [14]. The authors also defined the almost global dimension  $A.\operatorname{gld}(R)$  of a ring *R* in terms of almost projective modules, and obtained that rings *R* with  $A.\operatorname{gld}(R) = 0$  are exactly von Neumann regular rings, and domains with  $A.\operatorname{gld}(R) \leq 1$  are exactly almost Dedekind domains. Subsequently, Zhou et al. [15] studied the existence of almost projective covers, and showed every *R*-module has an almost projective cover if and only if *R* is a locally perfect ring, i.e., rings *R* satisfying  $R_{\mathfrak{m}}$  is a perfect ring for any maximal ideal  $\mathfrak{m}$  of *R*.

It is well known that every R-module has a projective envelope if and only if R is an Artinian ring (see [1]). If R is a domain or has w.gld $(R) < \infty$ , then every R-module has a flat envelope if and only if w.gld $(R) \le 2$  (see [1, 5, 10]). However, the characterizations of rings with existences of flat envelopes for all R-modules remain open in general case. The main motivation is to investigate the existence of almost projective envelopes. We show that every R-module has an almost projective envelope implies that R is a coherent locally Artinian ring. The converse is also true in some cases (see Theorem 4). We also characterize rings over which every R-module has a monoic, an epimorphic almost projective envelope or an almost projective envelope with unique mapping (see Theorem 6, Theorem 7, Theorem 8).

### 1. Main results

Let M be an R-module and  $\mathscr{P}$  be a class of R-modules. Recall from [6] that an R-homomorphism  $f : M \to P$  with  $P \in \mathscr{P}$  is said to be a  $\mathscr{P}$ -preenvelope provided that for any  $P' \in \mathscr{P}$ , the natural homomorphism  $\operatorname{Hom}_R(P, P') \to \operatorname{Hom}_R(M, P')$  is an epimorphism. If, moreover, every endomorphism h such that f = hf is an automorphism, then  $f : M \to P$  is said to be a  $\mathscr{P}$ -envelope.

**Lemma 1.** [13, Lemma 2.1.2] Let  $f : M \to P$  be a  $\mathscr{P}$ -envelope, and assume  $\mathscr{P}$  is closed extensions. Set  $A = \operatorname{Coker}(f)$ . Then  $\operatorname{Ext}^{1}_{R}(A, P') = 0$  for all  $P' \in \mathscr{P}$ .

The proof of [2, Proposition 3.4] can be applied to a wider range of situations. We give its proof for completeness.

**Proposition 2.** Let R be a ring and  $\mathscr{P}$  be a class of R-modules closed under isomorphisms and extensions. Let M be an R-module and  $f: M \to P$  be a  $\mathscr{P}$ -envelope. Suppose  $\alpha : M \to M$  is an R-isomorphism and  $\beta : P \to P$  is an R-homomorphism such that  $\beta f = f \alpha$ . Then  $\beta$  is also an R-isomorphism.

*Proof.* Note that there is an exact sequence of commutative diagram

$$\begin{array}{cccc} M & \stackrel{f}{\longrightarrow} P & \stackrel{\pi}{\longrightarrow} A & \longrightarrow 0 \\ \downarrow^{\alpha} & \downarrow^{\beta} \\ M & \stackrel{f}{\longrightarrow} P & \stackrel{\pi}{\longrightarrow} A & \longrightarrow 0 \end{array}$$

with  $A = \operatorname{coker}(f)$ .

We will first show  $\beta$  is a monomorphism. Since  $\alpha$  is an isomorphism and  $\beta f = f\alpha$ ,  $M \xrightarrow{\beta f} B$  is also a  $\mathscr{P}$ -envelope. So, there exists an *R*-homomorphism  $g: P \to P$  such that the following diagram is commutative



So  $g\beta f = gf\alpha = f$ . Since f is a  $\mathscr{P}$ -envelope, we have  $g\beta$  is an automorphism. Hence  $\beta$  is a monomorphism.

Next, we will show  $\beta$  is an epimorphism. Since  $\mathscr{P}$  is closed under isomorphisms,  $\beta(P) \in \mathscr{P}$ . Since  $M = \alpha(M)$ ,  $f(M) = f(\alpha(M)) = \beta(f(M))$ . Hence  $\operatorname{Im}(f) \subseteq \operatorname{Im}(\beta)$ . And so there is a commutative diagram of epimorphisms



Consider the exact sequence



Since  $\mathscr{P}$  is closed under extensions, we have  $\operatorname{Ext}^{1}_{R}(A, P) = 0$  by Lemma 1. So there is  $h: A \to P$  such that the above diagram is commutative. Consider the *R*-homomorphism  $\gamma$  :  $\operatorname{Id}_{P} - h\pi : P \to P$ . Then  $\gamma f = f - h\pi f = f$ . So  $\gamma$  is an automorphism as f

is a  $\mathscr{P}$ -envelope. Hence  $\gamma(P) = P$ . Note that  $\rho\gamma = \rho - \rho h\pi = \rho - \tau\pi = 0$ . So  $P = \gamma(P) \subseteq \text{Ker}(\rho) = \beta(P)$ . Hence  $\beta$  is an epimorphism.

**Corollary 3.** Let R be a ring and  $\mathfrak{p}$  be a prime ideal of R. Let M be an  $R_{\mathfrak{p}}$ -module and  $f: M \to P$  be an almost projective envelope. Then f is a projective envelope of M as an  $R_{\mathfrak{p}}$ -module.

Proof. Let  $s \in R - \mathfrak{p}$ . Then the multiplication by s is an isomorphism of  $R_{\mathfrak{p}}$ -modules. Note that the class of almost projective modules is closed under isomorphisms and extensions. It follows by Proposition 2 that P is also an  $R_{\mathfrak{p}}$ -module. Since  $\operatorname{Hom}_{R}(M, N) \cong$  $\operatorname{Hom}_{R_{\mathfrak{p}}}(M, N)$  for any  $R_{\mathfrak{p}}$ -module N, f is also an almost projective (thus a projective) envelope as  $R_{\mathfrak{p}}$ -modules.

It is well-known that the every R-module has a projective envelope if and only if R is an Artinian ring (see [1, Corollary 3.6]). However, the existence of flat envelope for every R-module is still open for general rings. The next Theorem investigates rings R over which every R-module has an almost projective envelope.

**Theorem 4.** Let R be a ring. Suppose every R-module has an almost projective envelope. Then R is a coherent locally Artinian ring.

Moreover, suppose R satisfies one of the following conditions:

- (1) R is a reduced ring;
- (2)  $w.\operatorname{gld}(R) < \infty;$
- (3) R is a Noetherian ring;
- (4)  $R_{\mathfrak{m}}$  is a finitely generated projective *R*-module for each  $\mathfrak{m} \in \operatorname{Max}(R)$ .

Then the converse is also true.

Proof. Suppose every *R*-module has an almost projective envelope. We will first show *R* is a coherent ring. Let  $\Lambda$  be any index set and  $\prod_{i \in \Lambda} R^{(i)} \to P$  be an almost projective preenvelope with  $R^{(i)} \cong R$  for each  $i \in \Lambda$ . Since *R* is almost projective, Then is a decomposition  $\prod_{i \in \Lambda} R^{(i)} \to P \to R^{(i)}$  for each  $i \in \Lambda$ . Hence the composition  $\prod_{i \in \Lambda} R^{(i)} \to$  $P \to \prod_{i \in \Lambda} R^{(i)}$  is the identity of  $\prod_{i \in \Lambda} R^{(i)}$ . So  $\prod_{i \in \Lambda} R^{(i)}$  is a direct summand of *P*, and so is also an almost projective (and thus a flat) *R*-module. Consequently, *R* is a coherent ring. Next, we will show *R* is a locally Artinian ring. Let **p** be any prime ideal of *R*, *M* be any  $R_{p}$ -module, and  $f: M \to P$  be an almost projective envelope. Then *f* is a projective envelope of M by Corollary 3. Hence  $R_{\mathfrak{p}}$  is an Artinian ring by [1, Corollary 3.6]. In conclusion, R is a coherent locally Artinian ring.

On the other hand, we first suppose R is a coherent locally Artinian ring satisfying (1) or (2). Then R is a reduced ring with K.dim(R) = 0 (see [7, Corollary 4.24] for (2)). That is R is a von Neumann regular ring (see [8, Page 5]). So every R-module is almost projective by [14, Theorem 4.8]. Hence every R-module has an almost projective envelope. Next we suppose R is a coherent locally Artinian ring satisfying (3). Then R is an Artinian ring. So every almost projective module is projective. Consequently, the result follows by [1, Corollary 3.6].

Finally we suppose R is a coherent locally Artinian ring satisfying (4). Let M be an R-module and  $f: M \to P$  be an almost projective envelope. For each  $\mathfrak{m} \in \operatorname{Max}(R)$ , we claim that  $f_{\mathfrak{m}}: M_{\mathfrak{m}} \to P_{\mathfrak{m}}$  is a projective preenvelope of  $M_{\mathfrak{m}}$ . Indeed, let  $g: M_{\mathfrak{m}} \to X$  be an  $R_{\mathfrak{m}}$ -homomorphism with X a projective  $R_{\mathfrak{m}}$ -module, so it is a flat R-module, and hence an almost projective R-module module by assumption. Since f is an almost projective preenvelope, there is an R-homomorphism  $h: P \to X$  such that the following diagram is commutative



where  $\phi$  and  $\psi$  are natural homomorphisms. By localizing at  $\mathfrak{m}$ , we have an  $R_{\mathfrak{m}}$ -homomorphism  $h_{\mathfrak{m}}: P_{\mathfrak{m}} \to X$  such that  $g = h_{\mathfrak{m}} f_{\mathfrak{m}}$ . Hence  $f_{\mathfrak{m}}$  is a projective preenvelope as  $R_{\mathfrak{m}}$ -modules. It follows by [7, Lemma 5.3] that there is a direct summand  $Q^{\mathfrak{m}}$  of  $P_{\mathfrak{m}}$  such that  $\operatorname{Im}(f_{\mathfrak{m}}) \subseteq Q^{\mathfrak{m}}$  and  $f'_{\mathfrak{m}}: M_{\mathfrak{m}} \to Q^{\mathfrak{m}}$ , having the same correspondence rule as  $f_{\mathfrak{m}}$ , is a projective envelope of  $M_{\mathfrak{m}}$  as  $R_{\mathfrak{m}}$ -modules. Consider the following commutative diagram



Set  $Q = \bigcap_{\mathfrak{m} \in \operatorname{Max}(R)} \psi^{-1}(Q^{\mathfrak{m}})$ . Then  $\operatorname{Im}(f) \subseteq Q$ . Set  $f' : M \to Q$  to have the same correspondence rule as f. We claim that f' is the almost projective envelope of M. First we will show Q is an almost projective R-module. Indeed, since  $R_{\mathfrak{m}'}$  is a finitely generated projective R-module for each  $\mathfrak{m}' \in \operatorname{Max}(R)$ , then

$$Q_{\mathfrak{m}'} = (\bigcap_{\mathfrak{m}\in\operatorname{Max}(R)} \psi^{-1}(Q^{\mathfrak{m}}))_{\mathfrak{m}'}$$
$$= (\bigcap_{\mathfrak{m}\in\operatorname{Max}(R)} \psi^{-1}(Q^{\mathfrak{m}})) \otimes_R R_{\mathfrak{m}'}$$
$$= \bigcap_{\mathfrak{m}\in\operatorname{Max}(R)} (\psi^{-1}(Q^{\mathfrak{m}}) \otimes_R R_{\mathfrak{m}'})$$
$$= \bigcap_{\mathfrak{m}\in\operatorname{Max}(R)} \psi^{-1}(Q^{\mathfrak{m}})_{\mathfrak{m}'}$$
$$= Q^{\mathfrak{m}'}$$

The last equality follows by  $(\psi^{-1}(Q^{\mathfrak{m}}))_{\mathfrak{m}'} = P_{\mathfrak{m}'}$  and  $(\psi^{-1}(Q^{\mathfrak{m}'}))_{\mathfrak{m}'} = Q^{\mathfrak{m}'}$  for any distinct two maximal ideals  $\mathfrak{m}, \mathfrak{m}'$ . Indeed, we trivially have  $(\psi^{-1}(Q^{\mathfrak{m}'}))_{\mathfrak{m}'} = Q^{\mathfrak{m}'}$ . Now let  $\mathfrak{m}, \mathfrak{m}'$  be two distinct two maximal ideals of R. since  $R_{\mathfrak{m}}$  is Artinian, we have  $(P_{\mathfrak{m}})_{\mathfrak{m}'} \cong (F_{\mathfrak{m}})_{\mathfrak{m}'} = 0$ for some free R-module F. Hence  $Q_{\mathfrak{m}'}^{\mathfrak{m}}$ , which can be seen as a submodule of  $(P_{\mathfrak{m}})_{\mathfrak{m}'}$ , is also equal to 0. Localizing the last commutative diagram at  $\mathfrak{m}'$ , we have  $(\psi^{-1}(Q^{\mathfrak{m}}))_{\mathfrak{m}'} =$  $(\psi^{-1}(0))_{\mathfrak{m}'} = P_{\mathfrak{m}'}$ . So Q is an almost projective R-module. And consequently, f' is an almost projective preenvelope. Now let h be an endomorphism of Q such that hf' = f'. Then  $h_{\mathfrak{m}}f'_{\mathfrak{m}} = f'_{\mathfrak{m}}$  for each  $\mathfrak{m} \in \operatorname{Max}(R)$ . So  $h_{\mathfrak{m}}$  is an automorphism since  $f'_{\mathfrak{m}}$  is a projective envelope as  $R_{\mathfrak{m}}$ -modules. Hence f is also an automorphism. In conclusion, f' is an almost projective envelope.

Remark 5. Note that the conditions (1), (2) and (3) in Theorem 4 are not essential. Indeed, let R be a non-field Artinian ring. Then (1), (2) do not hold. However, since every almost projective R-module is projective over Artinian rings, every R-module has an almost projective envelope by [1, Corollary 3.6]. Let R be a non-semisimple von Neumann regular ring. Then (3) does not hold. Since every R-module is almost projective over von Neumann regular rings (see [14, Theorem 4.8]), every R-module trivially has an almost projective envelope. However, we do not known whether the condition (4) is essential. For further study, we propose the following question: Question: How to characterize rings R over which every R-module has an almost projective envelope?

Since over locally Artinian rings, the class of flat modules is equal to that of almost projective modules by [14, Theorem 2.5]. So to solve the above question, it is only need to characterize rings R over which every R-module has a flat envelope.

In the rest of this paper, we will mainly characterize rings R over which every R-module has a monoic, an epimorphic almost projective envelope or an almost projective envelope with the unique mapping property.

Let M be an R-module and  $\mathscr{P}$  be a class of R-modules. Let  $f: M \to P$  be a  $\mathscr{P}$ envelope of M. If for any  $f': M \to P'$  with  $P' \in \mathscr{P}$ , there exists a unique  $g: P \to P'$ such that f' = gf. Then f is said to a  $\mathscr{P}$ -envelope with the unique mapping property. If f is a monomorphism (resp., an epimorphism), then f is said to be a monoic (resp., an
epimorphic)  $\mathscr{P}$ -envelope.

**Theorem 6.** Let R be a ring. Then the following statements are equivalent.

- (1) R is a von Neumann regular ring.
- (2) Every R-module has an almost projective envelope with the unique mapping property.

*Proof.* (1)  $\Rightarrow$  (2) Suppose *R* is a von Neumann regular ring. Then every *R*-module is almost projective by [14, Theorem 4.8]. So it is trivial that every *R*-module has an almost projective envelope with the unique mapping property.

 $(2) \Rightarrow (1)$  Suppose every *R*-module has an almost projective envelope with the unique mapping property. We first have *R* is a coherent locally Artinian ring by Theorem 4. Let  $0 \to K \xrightarrow{\alpha} P_0 \xrightarrow{\beta} P_1$  be an exact sequence with  $P_0$  and  $P_1$  almost projective. Let  $f: K \to H$  be an almost projective envelope of *K*. Then there is a unique homomorphism  $h: H \to P_0$  such that the following diagram commutative



So  $\beta hf = 0$ , and hence  $\beta h = 0$  since f has unique mapping property. It follows that we have a unique homomorphism  $g: H \to K$  with  $\alpha g = h$  and so  $\alpha gf = \alpha$ . Hence  $gf = \mathrm{Id}_K$ .

So K is a direct summand of H. It follows that K is an almost projective module. Hence  $w.\operatorname{gld}(R) \leq 2$ . It follows by Theorem 4 and [7, Corollary 4.24] that R is a reduced ring with with  $\operatorname{K.dim}(R) = 0$ , that is, R is a von Neumann regular ring.

**Theorem 7.** Let R be a ring. Then the following statements are equivalent.

- (1) R is a von Neumann regular ring.
- (2) Every R-module has an almost projective envelope which is an epimorphism.

*Proof.*  $(1) \Rightarrow (2)$  Trivially if R is a von Neumann regular ring, then every R-module has an almost projective envelope which is an epimorphism.

 $(2) \Rightarrow (1)$  On the other hand, let P be a projective module, K a submodule of P and f:  $K \twoheadrightarrow P'$  an epimorphic almost projective envelope. Then there exists an homomorphism  $g: P' \to P$  such that the following diagram commutative



Thus f is also a monomorphism. Cosequently,  $K \cong P'$  is almost projective, and so is flat. It follows that  $w.\operatorname{gld}(R) \leq 1$ . Consequently, R is a von Neumann regular ring by Theorem 4 and [7, Corollary 4.24] again.

The authors in [1, Theorem 2.2] and [11, Theorem 2.1] characterized rings over which every R-module has a monoic flat envelope. The following result shows that the related property also characterizes rings over which every R-module has a monoic almost projective envelope.

**Theorem 8.** Let R be a ring. Then the following statements are equivalent.

- (1) every *R*-module has an almost projective envelope which is a monomorphism.
- (2) For each m ∈ Max(R), R<sub>m</sub> is either a field or a QF-ring which is projective as an R-module.
- (3) There exists a family of primitive orthogonal idempotents {e<sub>α</sub> | α ∈ Γ} such that:
  (a) Re<sub>α</sub> is a local non-field QF-ring for each α ∈ Γ;
  - (b)  $R / \bigoplus_{\alpha \in \Gamma} Re_{\alpha}$  is a von Neuman regular ring.
- (4) every *R*-module has a flat envelope which is a monomorphism.

- (5) R is a subdirect product of a locally QF-ring with monoic flat (almost projective) envelope  $R_1$  and a von Neuman regular ring  $R_2$  that is flat as an R-module by restriction of scalars.
- (6) R is an essential subdirect product of rings  $R_1$  and  $R_2$  as in (5).

*Proof.* (1) ⇒ (3) We first show  $R_{\mathfrak{m}}$  is a QF-ring for any  $\mathfrak{m} \in \operatorname{Max}(R)$ . It follows by the proof of Theorem 4 that every  $R_{\mathfrak{m}}$ -module has a monoic projective envelope in the category of all  $R_{\mathfrak{m}}$ -modules. So every injective  $R_{\mathfrak{m}}$ -module is a submodule (hence a direct summand) of a projective  $R_{\mathfrak{m}}$ -module. Hence every injective  $R_{\mathfrak{m}}$ -module is projective, that is,  $R_{\mathfrak{m}}$  is a QF-ring. So every flat module is almost projective. It follows by the proof of [1, Theorem 2.2] that (3) holds.

 $(2) \Rightarrow (1)$  Note that R is a locally perfect ring, so every flat module is almost projective. Hence the result follows by [1, Theorem 2.2].

- $(2) \Leftrightarrow (4)$  See [1, Theorem 2.2].
- $(3) \Leftrightarrow (4) \Leftrightarrow (5) \Leftrightarrow (6)$  See [11, Theorem 2.1].

## Acknowledgement.

The second author was supported by the National Natural Science Foundation of China (No.12201361), the third author was supported by the National Natural Science Foundation of China (No. 12101515).

#### References

- Asensio Mayor J. and Martíez Hernádez J. (1993). On flat and projective envelopes. J. Algebra 160(2): 434-440.
- Bazzoni S. (2010). Divisible envelopes, *P*<sub>1</sub>-covers and weak-injective modules. J. Algebra Appl. 9(4): 531-542.
- [3] Chase S. U. (1960). Direct products of modules. Trans. Amer. Math. Soc. 97: 457-473.
- [4] Ding Q. N. (1996). On envelopes with the unique mapping property. Commun. Algebra 24(4): 1459-1470.
- [5] Enochs E. E. (1981). Injective and flat covers, envelopes and resolvents. Israel J. Math. 39: 189-209.
- [6] Enochs E. E. and Jenda O. M. (2011). *Relative homological algebra*, Volume I, Expositions in Mathematics 30, Walter de Gruyter.
- [7] Glaz S. (1989). Commutative Coherent Rings, Lecture Notes in Mathematics, vol. 1371, Spring-Verlag, Berlin.
- [8] Huckaba J. A. (1988). Commutative rings with zero divisors, Monographs and Textbooks in Pure and Applied Mathematics, vol. 117, Marcel Dekker, Inc., New York.

- [9] Matsumura H. (1989). Commutative ring theory, volume 8 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, second edition.
- [10] Martinez Hernandez J. (1986). Relatively flat envelopes. Commun. Algebra 14: 867-884.
- [11] Saorin M. (1995). The structure of commutative rings with monomorphic flat envelopes. Commun. Algebra 23(14): 5383-5394.
- [12] Wang F. G. and Kim H. (2016). Foundations of Commutative Rings and Their Modules, Singapore: Springer.
- [13] Xu J. Z. (1996). Flat covers of modules, Lecture Notes in Mathematics, 1634, Springer-Verlag, Berlin.
- [14] Zhou D. C., Kim H., Wang F. G. and Hu K. (2020). Homological characterizations of almost Dedekind domains. J. Algebra Appl. 19(7): 2050128, 12 pages.
- [15] Zhou D. C., Kim H., Zhang X. L. and Hu K. (2021). Some Remarks on Locally Perfect Rings. J. Algebra Appl. 20(2): 2150009, 13 pages.