

On the structure of Kauffman bracket skein algebra of a surface

Haimiao Chen

Abstract

Suppose R is a commutative ring with identity and a fixed invertible element $q^{\frac{1}{2}}$ such that $q + q^{-1}$ is invertible. For an oriented surface Σ , let $\mathcal{S}(\Sigma; R)$ denote the Kauffman bracket skein algebra of Σ over R . It is shown that to each embedded graph $G \subset \Sigma$ satisfying that $\Sigma \setminus G$ is homeomorphic to a disk and some other mild conditions, one can associate a generating set for $\mathcal{S}(\Sigma; R)$, and the ideal of defining relations is generated by relations of degree at most 6 supported by certain small subsurfaces.

Keywords: Kauffman bracket; skein algebra; quantization; character variety; oriented surface

MSC2020: 57K16, 57K31

1 Introduction

Let R be a commutative ring with identity and a fixed invertible element $q^{\frac{1}{2}}$. For an oriented surface Σ , the *Kauffman bracket skein algebra* of Σ over R , denoted by $\mathcal{S}(\Sigma; R)$, is defined as the quotient of the free R -module generated by isotopy classes of framed links embedded in $\Sigma \times (0, 1)$ by the submodule generated by the skein relations

$$\begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = q^{\frac{1}{2}} \begin{array}{c} \diagdown \\ \diagup \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} + q^{-\frac{1}{2}} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \diagdown \\ \diagup \end{array}, \quad \bigcirc = -(q + q^{-1}) \emptyset.$$

As a convention, R is identified with $R\emptyset \subset \mathcal{S}(M; R)$ via $\eta \mapsto \eta\emptyset$. Each link is equipped with the blackboard framing.

Given $L_1, L_2 \subset \Sigma \times (0, 1)$, their product $L_1 L_2$ is defined by stacking L_1 over L_2 in the $[0, 1]$ direction. Using the local relations, each element of $\mathcal{S}(\Sigma; R)$ can be written as a R -linear combination of multi-curves in Σ , where a multicurve is considered as a link in $\Sigma \times (0, 1)$ by identifying Σ with $\Sigma \times \{1/2\}$.

The description of the structure of $\mathcal{S}(\Sigma; \mathbb{Z}[q^{\pm\frac{1}{2}}])$ was raised as [9] Problem 1.92 (J) (proposed by Bullock and Przytycki), and also [10] Problem 4.5. A finite set of generators was given by Bullock [2]. So the problem is really to determine the defining relations.

Let $\Sigma_{g,h}$ denote the oriented surface of genus g with h boundary circles; let $\Sigma_g = \Sigma_{g,0}$. In the previous work [4], we found a generating set of $\mathcal{S}(\Sigma_{0,n+1}; R)$ consisting of $n + \binom{n}{2} + \binom{n}{3}$ elements, and showed that relations among them can be “localized”, in the sense that the ideal of defining relations is generated by certain relations of degree at most 6. In this paper, we extend the result to $\Sigma_{g,h}$

for all g, h . Setting $q^{\frac{1}{2}} = -1$ will lead to a presentation for $\mathbb{C}[\mathcal{X}_{\text{SL}(2, \mathbb{C})}(\Sigma_g)]$, which was regarded as a difficult problem and no result is seen in the literature.

When $q + q^{-1}$ is not assumed to be invertible, in principle a presentation for $\mathcal{S}(\Sigma_{g,h}; \mathbb{Z}[q^{\pm \frac{1}{2}}])$ can still be obtained without essential difficulty. Through this approach, the structure of $\mathcal{S}(\Sigma_{0,5}; \mathbb{Z}[q^{\pm \frac{1}{2}}])$ had been determined in [5].

The content is organized as follows. In Section 2 we introduce some necessary notations and conventions (most of which have been given in [4]), and propose the notion ‘‘cutting system’’; we define admissible expressions for all connected 1-submanifolds in $\Sigma \times [0, 1]$ for all surface Σ . In Section 3 we formulate Theorem 3.1 which (roughly) asserts that relations can be ‘‘localized’’, and outline the process of proving it; the core of the proof is a ‘‘tortuous induction’’ which was implemented in [4]. In Section 4, several examples of cutting systems on surfaces are given. Section 5 is an appendix, containing detailed formulas supporting Lemma 2.3.

2 Generating set arising from a cutting system

2.1 Preparation

Throughout, denote q^{-1} as \bar{q} (and also denote $q^{-\frac{1}{2}}$ as $\bar{q}^{\frac{1}{2}}$, etc), to make expressions as compact as possible. Let $\Sigma = \Sigma_{g,h}$. Let $\pi : \Sigma \times [0, 1] \rightarrow \Sigma$ denote the projection. We usually abbreviate $\mathcal{S}(\Sigma_{g,h}; R)$ to \mathcal{S} .

For a finite set Y , let $\#Y$ denote its cardinality.

We always use ‘isotopy’ to mean ordinary isotopy, without further explanations. A *fine isotopy* is a family of homeomorphisms $\varphi_t : \Sigma \times [0, 1] \rightarrow \Sigma \times [0, 1]$, $0 \leq t \leq 1$, such that $\varphi_0 = \text{id}$, $\pi(\varphi_t(\mathbf{a}, z)) = \pi(\varphi_t(\mathbf{a}, 0))$ for all $(\mathbf{a}, z) \in \Sigma \times [0, 1]$, and $\varphi_t(\Gamma_k) = \Gamma_k$ for all t, k . This notion is introduced for keeping track of the information on crossings. Given 1-submanifolds $X, X' \subset \Sigma \times [0, 1]$, call them *finely isotopic* and denote $X \approx X'$, if there exists a fine isotopy φ_t with $\varphi_1(X) = X'$; in this case, $\lambda(X) = \lambda(X')$.

Call a (most likely non-connected) embedded graph $G \subset \Sigma$ a *cutting system* for Σ if the following conditions hold: (i) $\Sigma \setminus G$ is homeomorphic to a disk; (ii) each point in $G \cap \partial\Sigma$ is a univalent vertex; (iii) the arcs incident to each interior vertex of G (i.e. one lying in the interior of Σ) belong to 2, 3, or 4 edges; (iv) each edge of G is oriented.

Fix a cutting system G . Numerate the edges of G as $\gamma_1, \dots, \gamma_n$. Let $\Gamma_i = \gamma_i \times [0, 1] \subset \Sigma \times [0, 1]$; let $\Gamma = \bigcup_{i=1}^n \Gamma_i$.

Given $\{i_1, \dots, i_r\} \subset \{1, \dots, n\}$, let $\Sigma(i_1, \dots, i_r)$ denote the surface obtained from cutting Σ along the γ_j 's for $j \in \{1, \dots, n\} \setminus \{i_1, \dots, i_r\}$.

A 1-submanifold $X \subset \Sigma \times [0, 1]$ is always assumed to be *in generic position*, meaning that $\pi(X)$ is stable under small perturbations of X . In particular, $\pi(X) \cap G = \emptyset$. Let $\text{Cr}(X)$ denote the set of crossings of X . Let $|X| = \#(X \cap \Gamma)$, called the *degree* of X , and let $\text{cn}(X) = \#\text{Cr}(X)$. Call X *simple* if $\text{cn}(X) = 0$. Define $\lambda(X) = (|X|, \text{cn}(X))$.

Each arc is assumed to be oriented, although sometimes the orientation is irrelevant. For an arc F , write $\partial F = \{\partial_- F, \partial_+ F\}$ so that F is oriented from $\partial_- F$ to $\partial_+ F$. Let \overline{F} denote the arc obtained from F by reversing its orientation.

Given $\Omega = \sum_i a_i X_i$ with $0 \neq a_i \in R$ and X_i being a 1-submanifold, let $\text{md}_\Omega(v) = \max_i \text{md}_{X_i}(v)$, and let $|\Omega| = \sum_{v=1}^n \text{md}_\Omega(v)$. For linear combinations Ω, Ω' , write $\text{md}_\Omega \leq \text{md}_{\Omega'}$ if $\text{md}_\Omega(v) \leq \text{md}_{\Omega'}(v)$ for all v .

Introduce a linear order \preceq on \mathbb{N}^2 , by declaring $(m', c') \preceq (m, c)$ if either $m' < m$, or $m' = m$, $c' \leq c$. Denote $(m', c') \prec (m, c)$ if $(m', c') \preceq (m, c)$ and $(m', c') \neq (m, c)$. If $\lambda(X) \prec \lambda(X')$, then we say that X is *simpler* than X' .

Let \mathcal{F} denote the free R -algebra generated by fine isotopy classes of links, with multiplication defined via stacking. Let $\theta' : \mathcal{F} \twoheadrightarrow \mathcal{S}$ denote the canonical quotient. For $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{F}$, we say $\mathbf{u}_1 = \mathbf{u}_2$ in \mathcal{S} if $\theta'(\mathbf{u}_1) = \theta'_n(\mathbf{u}_2)$.

Let \mathcal{V} denote the free R -module generated by isotopy classes of multi-curves. For a link L , let $\Theta(L) \in \mathcal{V}$ denote the linear combination of isotopy classes of multi-curves resulting from resolving all crossings of L . Extend this via linearity to $\Theta : \mathcal{S} \rightarrow \mathcal{V}$; by definition, it descends to $\Theta : \mathcal{S} \rightarrow \mathcal{V}$. By [7, 14], $\Theta : \mathcal{S} \cong \mathcal{V}$ is an isomorphism of R -modules.

When A, B are arcs with $\partial_+ A = \partial_- B$ and $\partial_- A \neq \partial_+ B$, let AB denote the arc obtained by identifying $\partial_+ A$ with $\partial_- B$, i.e. $AB = A \cup B$; when $\partial_+ A = \partial_- B$ and $\partial_- A = \partial_+ B$, let $\text{tr}(AB) = A \cup B$, which is a knot. In the construction of such kind, we may perturb A or B if necessary, to ensure $A \cup B$ to be generic. This convention will be always adopted.

Let $\mathbf{w}(F)$ denote the word in $\mathbf{x}_1^{\pm 1}, \dots, \mathbf{x}_n^{\pm 1}$ determined as follows: starting from $\partial_- F$, walk along F guided by the orientation of F , record \mathbf{x}_i (resp. \mathbf{x}_i^{-1}) whenever passing through Γ_i from left to right (resp. from right to left), and multiply all the recorded $\mathbf{x}_i^{\pm 1}$ together when reaching $\partial_+ F$. Call F *reduced* if $\mathbf{w}(F)$ is a reduced word; otherwise, call F *reducible*.

Given simple curves S, S' , if there exist simple arcs A, F, F' such that $|F|, |F'| \leq 4$, $S = \text{tr}(AF)$, $S' = \text{tr}(AF')$, and $\text{tr}(FF')$ is a short simple curve enclosing some interior vertex of G , then we say that S' is obtained from S by sliding F' to F across the vertex.

Lemma 2.1. *Two simple curves are isotopic if and only if they can be related by fine isotopy, shrinking degree 2 arcs, and sliding across interior vertices.*

Proof. The “if” part is trivial.

For the “only if” part, suppose S, S' are isotopic simple curves. Remove a small disk around each interior vertex. Then Σ becomes a surface with boundary Σ° , and G becomes a cutting system G° for Σ° which is a disjoint union of arcs. We may assume $S, S' \subset \Sigma^\circ$.

By sliding across interior vertices and shrinking degree 2 arcs whenever necessary, we may assume that S, S' are reduced and isotopic in Σ° .

Then the proof of [4] Lemma 2.2 can be easily extended to Σ° here, to show that S, S' are finely isotopic. \square

2.2 Admissible expressions

Let Δ denote the closure of $\Sigma \setminus G$, which by hyperthesis is a closed disk. Let $\text{gl} : \Delta \rightarrow \Sigma$ stand for the gluing map.

Each simple arc $F \subset \Sigma$ of degree 3 can be presented in the following way. Suppose F intersects $\gamma_{i_1}, \gamma_{i_2}, \gamma_{i_3}$ consecutively in the order given by the orientation, with i_1, i_2, i_3 not necessarily distinct. Take a homeomorphism $f : \Delta \rightarrow D^2$ (the unit disk) such that the 6 points constituting $f(\text{gl}^{-1}(F \cap G))$ is equidistributed on ∂D^2 . Let $\tilde{F} = \text{gl}^{-1}(F)$. Call $f(\tilde{F})$ the *symbol* of F .

Two examples of arcs are shown in Figure 1, where $\Sigma = \Sigma_2$.

Up to orientation-reversion of F and self-homeomorphism of D^2 , all possible symbols are shown in Figure 2.

Choose a point $x_i \in \gamma_i$ for each i . Suppose $\text{gl}^{-1}(x_i) = \{\tilde{x}_i^+, \tilde{x}_i^-\}$.

For each i , take an arc $\tilde{t}_i \subset \Delta$ connecting \tilde{x}_i^+ to \tilde{x}_i^- , and let $t_i = \text{gl}(\tilde{t}_i)$. For $i < j$, choose once for all a disjoint union \tilde{t}_{ij} of two arcs connecting \tilde{x}_i^\pm to $\tilde{x}_j^{\pm\epsilon}$, with $\epsilon \in \{\pm\}$, and let $t_{ij} = \text{gl}(\tilde{t}_{ij})$. For $i < j < k$, choose once for all a disjoint union \tilde{t}_{ijk} of three arcs connecting \tilde{x}_i^+ to \tilde{x}_j^ϵ , connecting $\tilde{x}_j^{-\epsilon}$ to \tilde{x}_k^\emptyset , and connecting $\tilde{x}_k^{-\emptyset}$ to \tilde{x}_i^- , where $\epsilon, \emptyset \in \{\pm\}$, and let $t_{ijk} = \text{gl}(\tilde{t}_{ijk})$.

Remark 2.2. When $\tilde{x}_i^\pm, \tilde{x}_j^\pm$ are not interlaced, \tilde{t}_{ij} is unique. Otherwise, there are two choices for \tilde{t}_{ij} ; however, denoting the other choice by \tilde{t}'_{ij} , we have that $\text{gl}(\tilde{t}'_{ij})$ equals a linear combination of $t_i t_j$ and t_{ij} .

The situation for $\tilde{x}_i^\pm, \tilde{x}_j^\pm, \tilde{x}_k^\pm$ is similar, although more complicated. It is not difficult to see that the $\text{gl}(\tilde{t}'_{ijk})$ for any other choice \tilde{t}'_{ijk} for \tilde{t}_{ijk} can be written as a linear combination of $t_i t_j t_k, t_i t_{jk}, t_j t_{ik}, t_k t_{ij}$ and t_{ijk} .

Let \mathcal{T} be the free R -algebra generated by \mathfrak{T} , where

$$\mathfrak{T} = \{t_{i_1 \dots i_r} : 1 \leq i_1 < \dots < i_r \leq n, 1 \leq r \leq 3\}.$$

Let $\theta : \mathcal{T} \rightarrow \mathcal{S}$ denote the canonical map.




Figure 1: Upper row (from left to right): a cutting system of Σ_2 ; the 4-holed sphere obtained from cutting two curves; the disk obtained from cutting along the curves. Middle and lower rows: degree 3 simple arcs, with orientations indicated by arrows.




Figure 2: Various symbols for degree 3 simple arcs. Denote the ones in the first two rows by I_1 to I_{10} ; denote the ones in the third and fourth rows by II_1 to II_8 ; denote the ones in the last two rows by III_1 to III_7 .

Given $x, y \in \Sigma \times \{z\}$ with $z \in \{0, 1\}$, let $\mathcal{H}_z(x, y)$ denote the set of 1-

submanifolds $X = C \cup L \subset \Sigma \times [0, 1]$, where L is a link and C is an arc with $\partial_- C = x$, $\partial_+ C = y$. Let $\mathcal{S}(x, y)$ denote the R -module generated by relative isotopy classes of elements of $\mathcal{H}_z(x, y)$, modulo skein relations. Here a relative isotopy means an isotopy φ_t of $\Sigma \times [0, 1]$ with $\varphi_t(x) = x$ and $\varphi_t(y) = y$ for all t . Let $[X] \in \mathcal{S}(x, y)$ denote the element represented by X . When $z = 0$ (resp. $z = 1$), $\mathcal{S}(x, y)$ is a left (resp. right) \mathcal{S} -module. Let $\mathcal{M}(x, y)$ denote the subset of $\mathcal{H}_z(x, y)$ consisting of simple reduced arcs C with $|C| \leq 2$.

Lemma 2.3. *Let $F \in \mathcal{H}_0(x, y)$ be a degree 3 arc.*

- (i) *If $z = 0$, then there exist $\mathbf{a}_s \in \mathcal{T}$, $C_s \in \mathcal{M}(x, y)$ such that $[F] = \mathfrak{s}_u(F) := \sum_s \mathbf{a}_s [C_s]$ in $\mathcal{S}(x, y)$, and $\text{md}_{\mathfrak{s}_u(F)} \leq \text{md}_F$.*
- (ii) *If $z = 1$, then there exist $\mathbf{b}_t \in \mathcal{T}$, $D_t \in \mathcal{M}(x, y)$ such that $[F] = \mathfrak{s}_d(F) := \sum_t [D_t] \mathbf{b}_t$ in $\mathcal{S}(x, y)$, and $\text{md}_{\mathfrak{s}_d(F)} \leq \text{md}_F$.*

Proof. The result is an extension of [4] Lemma 3.1.

Say that I_1 can be chopped if an arc of type I_1 satisfies the assertion, and so forth. As a key ingredient of the proof of [4] Lemma 3.1, it was shown that I_1, II_1, III_1 can be chopped.




Figure 3: Copied from [4] Figure 3. The first two are of type II_1 , the third is of type III_1 , and the last is of type I_1 .

All the other symbols can also be chopped, as ensured by the identities given in the figures in Section 5. \square

Let $\mathcal{S}_u(F)$ denote the set of formal sums $\sum_s \mathbf{a}_s C_s$ satisfying Lemma 2.3 (i); let $\mathcal{S}_d(F)$ denote the set of formal sums $\sum_t D_t \mathbf{b}_t$ satisfying Lemma 2.3 (ii). Unlike in [4], now it is no longer likely to find a preferred element of $\mathcal{S}_u(F)$ or $\mathcal{S}_d(F)$, so we need to deal with all possible choices.

For a generic $\mathfrak{s} = \sum_s \mathbf{a}_s C_s \in \mathcal{S}_u(F^\sharp)$ in the sense that each $(K^F | F^\sharp | C_s)$ is in generic position, put

$$\mathfrak{s}^\sharp(K, F, \mathfrak{s}) = U(K, F) + q^{\hat{\epsilon}(K, F)} \sum_s \mathbf{a}_s (K^F | F^\sharp | C_s) \in \mathcal{F}; \quad (1)$$

similarly, for a generic $\mathfrak{t} = \sum_t D_t \mathbf{b}_t \in \mathcal{S}_d(F^\flat)$ in the sense that each $(K_F | F^\flat | D_t)$ is in generic position, put

$$\mathfrak{s}_\flat(K, F, \mathfrak{t}) = D(K, F) + q^{\hat{\epsilon}(K, F)} \sum_t (K_F | F^\flat | D_t) \mathbf{b}_t \in \mathcal{F}. \quad (2)$$

For a 1-submanifold $X \subset \Sigma \times [0, 1]$, let $\mathfrak{A}_3(X)$ denote the set of degree 3 arcs of X .

For each knot K , recursively define the set $\mathcal{A}(K) \subset \mathcal{T}$ of *admissible expressions*. When $|K| \leq 3$, put $\mathcal{A}(K) = \{\Theta(K)\}$. Suppose $|K| > 3$ and that $\mathcal{A}(J)$ has been defined for each knot J simpler than K . Take $F \in \mathfrak{A}_3(K)$, take generic $\mathfrak{s} \in \mathfrak{S}_u(F)$, $\mathfrak{t} \in \mathfrak{S}_d(F)$, and write $\mathfrak{s}^\sharp(K, F, \mathfrak{s}) = \sum_i \mathfrak{c}_i M_i$, $\mathfrak{s}_b(K, F, \mathfrak{t}) = \sum_j N_j \mathfrak{d}_j$, with $\mathfrak{c}_i, \mathfrak{d}_j \in \mathcal{T}$ and M_i, N_j being knots subsequent to K ; put

$$\mathcal{A}_u(K, F, \mathfrak{s}) = \left\{ \sum_i \mathfrak{c}_i \mathfrak{g}_i : \mathfrak{g}_i \in \mathcal{A}(M_i) \right\}, \quad (3)$$

$$\mathcal{A}_d(K, F, \mathfrak{t}) = \left\{ \sum_j \mathfrak{h}_j \mathfrak{d}_j : \mathfrak{h}_j \in \mathcal{A}(N_j) \right\}. \quad (4)$$

Set

$$\mathcal{A}(K) = \bigcup_{F \in \mathfrak{A}_3(K)} \left(\left(\bigcup_s \mathcal{A}_u(K, F, \mathfrak{s}) \right) \cup \left(\bigcup_t \mathcal{A}_d(K, F, \mathfrak{t}) \right) \right).$$

For each arc $A \in \mathcal{H}_0(\mathbf{x}, \mathbf{y})$ with $\mathbf{x}, \mathbf{y} \in \Sigma \times \{0\}$, recursively define the set $\mathcal{A}_{\text{pu}}(A) \subset \mathcal{TS}(\mathbf{x}, \mathbf{y})$ of *purely upward admissible expressions*. Put $\mathcal{A}_{\text{pu}}(A) = \{\Theta(A)\}$ if $|A| \leq 2$; put $\mathcal{A}_{\text{pu}}(A) = \{\mathfrak{s}_u(A)\}$ if $|A| = 3$. Suppose $|A| > 3$ and that $\mathcal{A}_{\text{pu}}(B)$ has been defined for each arc B simpler than A . Take $F \in \mathfrak{A}_3(A)$, and define A^F , $U(A, F)$, “arcs subsequent to A ”, and so forth, similarly as above. Take a generic $\mathfrak{s} = \sum_s \mathfrak{a}_s C_s \in \mathfrak{S}_u(F^\sharp)$. Writing

$$s^\sharp(A, F, \mathfrak{s}) := U(A, F) + q^{\varepsilon(A, F)} \sum_s \mathfrak{a}_s (A^F | F^\sharp | C_s)$$

as $\sum_i \mathfrak{c}_i G_i$, where the G_i 's are arcs subsequent to A , put

$$\mathcal{A}_u(A, F, \mathfrak{s}) = \left\{ \sum_i \mathfrak{c}_i \mathfrak{g}_i : \mathfrak{g}_i \in \mathcal{A}_{\text{pu}}(G_i) \right\}.$$

Set

$$\mathcal{A}_{\text{pu}}(A) = \bigcup_{F \in \mathfrak{A}_3(A)} \left(\bigcup_s \mathcal{A}_u(A, F, \mathfrak{s}) \right).$$

For each arc $A \in \mathcal{H}_1(\mathbf{x}, \mathbf{y})$ with $\mathbf{x}, \mathbf{y} \in \Sigma \times \{1\}$, define the set $\mathcal{A}_{\text{pd}}(A) \subset \mathcal{S}(\mathbf{x}, \mathbf{y})\mathcal{T}$ of *purely downward admissible expressions* in a parallel way.

Lemma 2.4. *Suppose $K = \text{tr}(AB)$, where A, B are arcs with $\text{Cr}(B/A) = \emptyset$. If $\sum_i \mathfrak{a}_i [C_i] \in \mathcal{A}_{\text{pu}}(A)$, $\sum_j [D_j] \mathfrak{b}_j \in \mathcal{A}_{\text{pd}}(B)$, and $\mathfrak{f}_{i,j} \in \mathcal{A}(\text{tr}(C_i D_j))$ for all i, j , then $\sum_{i,j} \mathfrak{a}_i \mathfrak{f}_{i,j} \mathfrak{b}_j \in \mathcal{A}(K)$.*

For a stacked link $L = K_1 \cdots K_r$, put

$$\mathcal{A}(L) = \left\{ \mathfrak{e}_1 \cdots \mathfrak{e}_r : \mathfrak{e}_1 \in \mathcal{A}(K_1), \dots, \mathfrak{e}_r \in \mathcal{A}(K_r) \right\}.$$

Suppose $M = S_1 \sqcup \cdots \sqcup S_r$ is a multicurve. Each $\sigma \in \text{Sym}(r)$ (the permutation group of $\{1, \dots, r\}$) gives rise to a stacked link $S_{\sigma(1)} \cdots S_{\sigma(r)}$, which is not equivalent to $S_1 \cdots S_r$ unless σ is the identity. Put

$$\mathcal{A}(M) = \bigcup_{\sigma \in \text{Sym}(r)} \mathcal{A}(S_{\sigma(1)} \cdots S_{\sigma(r)}). \quad (5)$$

The construction based on Lemma 2.3 has ensured $\mathcal{A}(K) \neq \emptyset$ for any knot K . Hence $\mathcal{A}(M) \neq \emptyset$ for any multicurve M , establishing the surjectivity of θ .

3 Relations can be localized

For a 1-submanifold X , let $\Sigma(X) = \Sigma(v_1, \dots, v_r)$ if $X \cap \Gamma_v \neq \emptyset$ exactly for $v = v_1, \dots, v_r$.

Let Λ be the set of (v_1, \dots, v_k) with $1 \leq v_1 < \dots < v_k \leq n$ and $2 \leq k \leq 6$. For each $\vec{v} = (v_1, \dots, v_k) \in \Lambda$, let $\Sigma(\vec{v}) = \Sigma(v_1, \dots, v_k)$, and let $\mathcal{Z}(\vec{v}) \subset \mathcal{T}$ denote the R -module of polynomials \mathbf{u} in $t_{i_1 \dots i_r}$'s with $r \leq 3$ and $i_1, \dots, i_r \in \{v_1, \dots, v_k\}$ such that $|\mathbf{u}| \leq 6$ and $\mathbf{u} = 0$ in $\mathcal{S}(\Sigma(\vec{v}))$. For $\mathbf{u} \in \mathcal{Z}(\vec{v})$, we say that the relation $\mathbf{u} = 0$ is *supported* by $\Sigma(v_1, \dots, v_k)$.

Let \mathcal{I} denote the two-sided ideal of \mathcal{T} generated by $\bigcup_{\vec{v} \in \Lambda} \mathcal{Z}(\vec{v})$.

Theorem 3.1. *The ideal of defining relations of \mathcal{S} is \mathcal{I} , i.e., $\mathcal{I} = \ker \theta$.*

Let $\zeta : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{I}$ denote the quotient map. For $\mathbf{e}, \mathbf{e}' \in \mathcal{T}$, we say that \mathbf{e} is *congruent* to \mathbf{e}' and denote $\mathbf{e} \equiv \mathbf{e}'$, if $\mathbf{e} - \mathbf{e}' \in \mathcal{I}$, i.e. $\zeta(\mathbf{e}) = \zeta(\mathbf{e}')$.

For a knot K , if elements of $\mathcal{A}(K)$ are congruent to each other, then we denote $\check{\mathbf{a}}(K) \in \mathcal{T}/\mathcal{I}$ for the unique element of $\zeta(\mathcal{A}(K))$ and say that $\check{\mathbf{a}}(K)$ is *well-defined*. Note that $\check{\mathbf{a}}(K)$ is well-defined whenever $|K| \leq 6$.

Given $\Omega = \sum_i a_i N_i$, with $a_i \in R$ and N_i being a stacked link (probably a knot), let $\mathcal{A}(\Omega) = \{\sum_i a_i \mathbf{e}_i : \mathbf{e}_i \in \mathcal{A}(N_i)\}$; say that $\check{\mathbf{a}}(\Omega)$ is well-defined if $\#\zeta(\mathcal{A}(\Omega)) = 1$. For two linear combinations Ω_1, Ω_2 , denote $\Omega_1 \equiv \Omega_2$ if $\check{\mathbf{a}}(\Omega_1), \check{\mathbf{a}}(\Omega_2)$ are well-defined and equal, i.e., $\mathbf{e}_1 \equiv \mathbf{e}_2$ for any $\mathbf{e}_1 \in \mathcal{A}(\Omega_1), \mathbf{e}_2 \in \mathcal{A}(\Omega_2)$.




Figure 4: First row: the EP (A, B) ; $L_\times = \text{tr}(A)\text{tr}(B)$, $L_\infty = \text{tr}(AB)$, $L_0 = \text{tr}(A\bar{B})$. Second row: the EP (A, B) ; $L_\times = \text{tr}(AB)$, $L_\infty = \text{tr}(A)\text{tr}(B)$, $L_0 = \text{tr}(A\bar{B})$. Third row: the EP (A, B) ; $L_\times = \text{tr}(AB)$, $L_\infty = \text{tr}(A\bar{B})$, $L_0 = \text{tr}(A)\text{tr}(B)$.

An *elementary skein triple* (EST) is a triple of the form $(L_\times, L_\infty, L_0)$ in one of the three cases shown in Figure 4.

Here is the main procedure of establishing Theorem 3.1. It will be shown that $\check{\mathbf{a}}(K)$ is well-defined for each knot K , so that $\check{\mathbf{a}}(L)$ is well-defined for each stacked link L . The transformation from L to $\Theta(L)$ can be implemented via crossing-resolving of specific kind in the following sense: there exists a series of linear-combinations of stacked links $L = \Omega_0, \Omega_1, \dots, \Omega_m = \Theta(L)$ such that

$$\Omega_{i+1} - \Omega_i = \mathbf{a}(L_\times - q^{\frac{1}{2}}L_\infty - \bar{q}^{\frac{1}{2}}L_0)\mathbf{b}$$

for some EST $(L_\times, L_\infty, L_0)$ and $\mathbf{a}, \mathbf{b} \in \mathcal{T}$. We will show that $L_\times \equiv q^{\frac{1}{2}}L_\infty + \bar{q}^{\frac{1}{2}}L_0$ always. Thus, $L \equiv \Theta(L)$. In particular, $\mathbf{m} \equiv \Theta(\mathbf{m})$ for each monomial \mathbf{m} . As a consequence, whenever $\mathbf{f} = 0$ in \mathcal{S} , we have $\mathbf{f} \equiv \Theta(\mathbf{f}) = 0$, i.e. $\mathbf{f} \in \mathcal{I}$.

Technically, we need to prove several intermediate lemmas by ‘‘tortuous’’ inductions.

For $m \geq 6$ and $c \geq 0$, let $\phi_{m,c}$ stand for the statement that $\check{\mathbf{a}}(K)$ is well-defined for each knot K with $\lambda(K) = (m, c)$; let $\Phi_{m,c}$ stand for the statement that $\phi_{m',c'}$ holds for all $(m', c') \preceq (m, c)$.

Lemma 3.2. *Suppose $\Phi_{m,c}$ holds, and K is a knot with $\lambda(K) = (m, c)$.*

- (a) *If $F \in \mathfrak{A}_3(K)$, then $K \equiv \mathfrak{s}^\sharp(K, F, \mathfrak{s}) \equiv \mathfrak{s}_b(K, F, \mathfrak{t})$ for all generic $\mathfrak{s} \in \mathfrak{S}_u(F)$ and $\mathfrak{t} \in \mathfrak{S}_d(F)$.*
- (b) *For any arc $F \subset K$,*

$$K \equiv q^{\check{z}(K,F)}K^F + U(K, F) \equiv q^{\check{z}(K,F)}K_F + D(K, F).$$

Lemma 3.3. *Suppose $\Phi_{m,c}$ holds. Then*

- (a) *$L_\times \equiv q^{\frac{1}{2}}L_\infty + \bar{q}^{\frac{1}{2}}L_0$ for each EST $(L_\times, L_\infty, L_0)$ with $\lambda(L_\times) \preceq (m, c)$;*
- (b) *$K_1K_2 \equiv K_2K_1$ for any disjoint knots K_1, K_2 such that $\lambda(K_1K_2) \prec (m, c)$.*

Lemma 3.4. *Suppose $\Phi_{m,0}$ holds.*

- (a) *Let S, S' be simple curves such that $|S| \leq m$ and S' results from shrinking a degree 2 arc of S . Then $S \equiv S'$.*
- (b) *Let S, S' be simple curves such that $|S|, |S'| \leq m$ and S' results from sliding S across an interior vertex. Then $S \equiv S'$.*

Proof for (b). Assume $S = \text{tr}(AF)$, $S' = \text{tr}(AF')$, for some arcs A, F, F' such that $|F|, |F'| \leq 4$ and $\text{tr}(F'\bar{F})$ is a simple curve encircling an interior vertex.

Take $\sum_i \mathbf{a}_i[C_i] \in \mathcal{A}_{\text{pu}}(A)$. For each i , we have $\text{tr}(C_iF') \equiv \text{tr}(C_iF)$, since $\text{tr}(C_iF') = \text{tr}(C_iF)$ is a relation of degree at most 6 supported by $\Sigma(C_i \cup F \cup F')$. Hence

$$\check{\mathbf{a}}(S) - \check{\mathbf{a}}(S') = \sum_i \mathbf{a}_i(\check{\mathbf{a}}(C_iF) - \check{\mathbf{a}}(C_iF')) = 0.$$

□

Lemma 3.5. *Suppose $\Phi_{m,1}$ holds.*

- (a) *If S_1, \dots, S_r are disjoint simple curves with $|S_1| + \dots + |S_r| \leq m$, then $S_{\sigma(1)} \cdots S_{\sigma(r)} \equiv S_1 \cdots S_r$ for each $\sigma \in \text{Sym}(r)$. Consequently, elements of $\mathcal{A}(S_1 \sqcup \dots \sqcup S_r)$ are congruent to each other.*
- (b) *If M, M' are isotopic multicurves of degree at most m , then $M \equiv M'$.*

Proof for (b). This is based on Lemma 2.1. □

Lemma 3.6. Suppose $\Phi_{m,c}$ holds. Then the following statements are true.

- (a) $K \equiv \Theta(K)$ for any knot K with $\lambda(K) = (m, c)$.
- (b) $K_1 K_2 \equiv \Theta(K_1 K_2)$ for any knots K_1, K_2 with $\lambda(K_1 K_2) \preceq (m, c)$.
- (c) If $\sum_j a_j K_j = 0$ in \mathcal{S} , where each K_j is a knot with $\lambda(K_j) \preceq (m, c)$, then $\sum_j a_j \Theta(K_j) \equiv 0$.

Lemma 3.7. $\phi_{m,c}$ holds for all $m \geq 6$ and all $c \geq 0$.

Lemma 3.8. $K_1 \cdots K_r \equiv \Theta(K_1 \cdots K_r)$, for any knots K_1, \dots, K_r .

Proof of Theorem 3.1. Suppose $\mathfrak{f} = \sum_i a_i \mathfrak{g}_i = 0$ in \mathcal{S} , where $a_i \in R$ and \mathfrak{g}_i is a monomial. Since each \mathfrak{g}_i itself is a stacked link, by Lemma 3.8, $\mathfrak{g}_i \equiv \Theta(\mathfrak{g}_i)$. Hence $\mathfrak{f} \equiv \sum_i a_i \Theta(\mathfrak{g}_i) = \Theta(\mathfrak{f}) = 0$. \square

4 Examples of cutting systems

4.1 $\Sigma_{0,4}$ revised

Figure 5 shows a cutting system, together with an associated generating set.




Figure 5: From left to right: a cutting system; t_1, t_2, t_3 ; t_{12}, t_{13}, t_{23} ; t_{123} .

4.2 $\Sigma_{1,3}$




Figure 6: Left: $\Sigma_{1,3}$, with a cutting system G drawn in red. Middle and right: the closure of $\Sigma_{1,3} \setminus G$.




Figure 7: The t_i 's, t_{ij} 's, t_{ijk} 's are respectively drawn in purple, green, orange.

4.3 $\Sigma_{g,k+1}$

Display $\Sigma_{g,k+1}$ as in Figure 8. A preferred cutting system is $\bigcup_{i=1}^{2g+k} \gamma_i$, where $\gamma_1, \dots, \gamma_{2g+k}$ are indicated by the dotted lines.




Figure 8: $\Sigma_{g,k+1}$; the dotted lines (from left to right) are $\gamma_1, \dots, \gamma_{2g+k}$.

In particular, see Figure 9 for $\Sigma_{2,1}$.




Figure 9: Left: $\Sigma_{2,1}$. Right: the closure of $\Sigma_{2,1} \setminus (\gamma_1 \cup \dots \cup \gamma_4)$.




Figure 10: The t_i 's, t_{ij} 's, t_{ijk} 's are respectively drawn in red, green, orange.

A generating set for $\mathcal{S}(\Sigma_{2,1})$ is shown in Figure 10.

A presentation for $\mathcal{S}(\Sigma_2)$ will be obtained by adding relations given by capping off the boundary.

5 Appendix: pictorial proofs for basic identities

An apology: at first sight these formulas seem to be cluttered arbitrarily; but actually they are obtained based on various attempts, and have been arranged to be nearly optimal.

$$\begin{aligned}
 & \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \\ I_3 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \quad \diagdown \\ 1 \quad 3 \\ 2 \\ I_8 \end{array} \\
 & \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \quad \diagdown \\ 1 \quad 3 \\ 2 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \\ III_6 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \quad \diagdown \\ 1 \quad 3 \\ 2 \\ I_3 \end{array} \\
 & \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \\ I_8 \end{array} + q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \quad \diagdown \\ 1 \quad 3 \\ 2 \\ III_6 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \quad \diagdown \\ 1 \quad 3 \\ 2 \end{array} \\
 & \quad \quad \quad + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \end{array} + q \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \quad \diagdown \\ 1 \quad 3 \\ 2 \end{array} + \bar{q} \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \quad \diagup \\ 1 \quad 3 \\ 2 \end{array}
 \end{aligned}$$

Figure 11: Taking charge of I_3 , I_8 , III_6 .

$$\begin{aligned}
& \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \\ 2 \quad 1 \\ 3 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{I}_6 \\ 2 \quad 1 \\ 3 \end{array} \\
& \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \\ 2 \quad 1 \\ 3 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{I}_6 \\ 2 \quad 1 \\ 3 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} \\
& \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{---} \\ 2 \quad 1 \\ 3 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} \\
& \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagup \diagdown \\ 2 \quad 1 \\ 3 \end{array} = q^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} + \bar{q}^{\frac{1}{2}} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} \\
& \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \\ 2 \quad 1 \\ 3 \end{array} = q \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} + \bar{q} \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{II}_7 \\ 2 \quad 1 \\ 3 \end{array} \\
& \quad \quad \quad + \begin{array}{c} 3 \\ 1 \quad 2 \\ \text{---} \\ 2 \quad 1 \\ 3 \end{array} + \begin{array}{c} 3 \\ 1 \quad 2 \\ \diagdown \\ 2 \quad 1 \\ 3 \end{array}
\end{aligned}$$

Figure 14: Taking charge of I_6 and II_7 .

$$\begin{aligned}
& \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 2 = q^{\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 3 + q^{-\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 2 \text{ and } 3 \\
& \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 2 = q^{\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 3 + q^{-\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 2 \text{ and } 3 \\
& \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 2 = q^{\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 3 + q^{-\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 2 \text{ and } 3 \\
& \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 2 = q^{\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 3 + q^{-\frac{1}{2}} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 2 \text{ and } 3 \\
& \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 2 = q \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 3 + \bar{q} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 2 \text{ and } 3 \\
& \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 2 = q \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 1 \text{ and } 3 + \bar{q} \text{Circle with strands } 1, 2, 3 \text{ and a crossing between } 2 \text{ and } 3
\end{aligned}$$

Figure 15: Taking charge of I_4 , I_9 and III_7 .

$$\begin{array}{ccc}
\begin{array}{c} 1 \\ \circlearrowleft \\ \text{II}_8 \\ \circlearrowright \\ 1 \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{II}_8 \\ \circlearrowright \\ 1 \end{array} - \bar{q} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{I}_4 \\ \circlearrowright \\ 1 \end{array} \\
\begin{array}{c} 1 \\ \circlearrowleft \\ \text{I}_2 \\ \circlearrowright \\ 1 \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{I}_2 \\ \circlearrowright \\ 1 \end{array} - q & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_7 \\ \circlearrowright \\ 1 \end{array} \\
\begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_3 \\ \circlearrowright \\ 1 \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_3 \\ \circlearrowright \\ 1 \end{array} - \bar{q} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_7 \\ \circlearrowright \\ 1 \end{array} \\
\begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_2 \\ \circlearrowright \\ 1 \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_2 \\ \circlearrowright \\ 1 \end{array} - q & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{III}_5 \\ \circlearrowright \\ 1 \end{array} \\
\begin{array}{c} 1 \\ \circlearrowleft \\ \text{I}_{10} \\ \circlearrowright \\ 1 \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{I}_{10} \\ \circlearrowright \\ 1 \end{array} - q & \begin{array}{c} 1 \\ \circlearrowleft \\ \text{I}_3 \\ \circlearrowright \\ 1 \end{array}
\end{array}$$

Figure 16: Taking charge of II_8 , I_2 , III_3 , III_2 , I_{10} .

$$\begin{array}{c}
\begin{array}{ccc}
\begin{array}{c} \textcircled{\text{I}_5} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} \textcircled{\text{I}_5} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} - q \begin{array}{c} \textcircled{\text{III}_6} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array}
\end{array} \\
\\
\begin{array}{ccc}
\begin{array}{c} \textcircled{\text{II}_2} \\ \text{3} \\ \text{1} \quad \text{2} \\ \text{1} \quad \text{3} \\ \text{2} \end{array} & = \bar{q}^{\frac{1}{2}} & \begin{array}{c} \textcircled{\text{II}_2} \\ \text{3} \\ \text{1} \quad \text{2} \\ \text{1} \quad \text{3} \\ \text{2} \end{array} - \bar{q} \begin{array}{c} \textcircled{\text{I}_3} \\ \text{3} \\ \text{1} \quad \text{2} \\ \text{1} \quad \text{3} \\ \text{2} \end{array}
\end{array} \\
\\
\begin{array}{ccc}
\begin{array}{c} \textcircled{\text{II}_5} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} & = \bar{q}^{\frac{1}{2}} & \begin{array}{c} \textcircled{\text{II}_5} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} - \bar{q} \begin{array}{c} \textcircled{\text{II}_2} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array}
\end{array} \\
\\
\begin{array}{ccc}
\begin{array}{c} \textcircled{\text{II}_3} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} & = q^{\frac{1}{2}} & \begin{array}{c} \textcircled{\text{II}_3} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} - q \begin{array}{c} \textcircled{\text{II}_2} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array}
\end{array} \\
\\
\begin{array}{ccc}
\begin{array}{c} \textcircled{\text{III}_4} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} & = \bar{q}^{\frac{1}{2}} & \begin{array}{c} \textcircled{\text{III}_4} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array} - \bar{q} \begin{array}{c} \textcircled{\text{II}_5} \\ \text{3} \\ \text{1} \quad \text{1} \\ \text{2} \quad \text{3} \\ \text{2} \end{array}
\end{array}
\end{array}$$

Figure 17: Taking charge of I_5 , II_2 , II_5 , II_3 , III_4 .

References

- [1] D. Bullock, Rings of $\text{SL}_2(\mathbb{C})$ -characters and the Kauffman bracket skein module, *Comment. Math. Helv.* 72 (1997), 521–542. Doi: 10.1007/s000140050032.
- [2] D. Bullock, A finite set of generators for the Kauffman bracket skein algebra, *Math. Z.* 231 (1999), 91–101. Doi: 10.1007/PL00004727.

- [3] D. Bullock, and J.H. Przytycki, Multiplicative structure of Kauffman bracket skein module quantizations, *Proc. Amer. Math. Soc.* 128 (2000), no. 3, 923–931. Doi: 10.1090/s0002-9939-99-05043-1.
- [4] H.-M. Chen, On skein algebras of planar surfaces, arXiv: 2206.07856.
- [5] H.-M. Chen, Integral structure of the skein algebra of the 5-punctured sphere, arXiv: 2304.06605.
- [6] J. Cooke and A. Lacabanne, Higher rank Askey-Wilson algebras as skein algebras, arXiv: 2205.04414.
- [7] L. Charles and J. Marché, Multicurves and regular functions on the representation variety of a surface in $SU(2)$, *Comment. Math. Helv.* 87 (2012), 409–431. DOI: 10.4171/cmh/258.
- [8] J. Cooke and P. Samuelson, On the genus two skein algebra, *J. Lond. Math. Soc.* 104 (2021), no. 5, 2260–2298. Doi: 10.1112/jlms.12497.
- [9] R. Kirby, *Problems in low-dimensional topology*, AMS/IP Stud. Adv. Math., 2.2, Geometric topology (Athens, GA, 1993), 35–473, Amer. Math. Soc., Providence, RI, 1997.
- [10] T. Ohtsuki, *Problems on invariants of knots and 3-manifolds*, with an introduction by J. Roberts. *Geom. Topol. Monogr.*, 4, Invariants of knots and 3-manifolds (Kyoto, 2001), i–iv, 377–572, *Geom. Topol. Publ.*, Coventry, 2002.
- [11] J. Przytycki and A.S. Sikora, On skein algebras and $Sl_2(\mathbb{C})$ -character varieties, *Topology* 39 (2000), no. 1, 115–148. Doi: 10.1016/S0040-9383(98)00062-7.
- [12] J. Przytycki and A.S. Sikora, Skein algebras of surfaces, *Trans. Amer. Math. Soc.* 371 (2019), no. 2, 1309–1332. Doi: 10.1090/tran/7298.
- [13] R. Santharoubane, Algebraic generators of the skein algebra of a surface, To appear in *Algebr. Geom. Topol.*
- [14] A.S. Sikora and B.W. Westbury, Confluence theory for graphs, *Algebr. Geom. Topol.* 7 (2007), 439–478. Doi:10.2140/agt.2007.7.439.

Haimiao Chen (orcid: 0000-0001-8194-1264) *chenhm@math.pku.edu.cn*
 Department of Mathematics, Beijing Technology and Business University,
 Liangxiang Higher Education Park, Fangshan District, Beijing, China.