

Identifying Sample Size and Accuracy and Precision of the Estimators in Case-Crossover Designs with Distributed Lags of Heteroskedastic Spatiotemporally Varying Continuous Exposures Measured with Simple or Complex Error

Honghyok Kim

Division of Environmental and Occupational Health Sciences, School of Public Health,
University of Illinois Chicago, Chicago, IL, 60661, USA

Research and Management Center for Health Risk of Particulate Matter, Seoul, 02841, Republic
of Korea

Corresponding author: Honghyok Kim, honghyok@uic.edu

Abstract

Understanding of sample size and the accuracy and precision of the estimator may facilitate power and bias analyses and help in designing reproducible research and interpreting findings. However, such understanding can become complicated in practice for several reasons. First, exposures varying spatiotemporally may be heteroskedastic. Second, distributed lags of exposures may be used to identify critical exposure time-windows. Third, exposure measurement error may exist, impacting the accuracy and/or precision of the estimator. Fourth, different study designs may affect sample size and power differently. For example, case-crossover designs as matched case-control designs, are used to estimate health effects of short-term exposures. Therefore, this article develops approximation equations for sample size, estimates of the estimators and standard errors, including polynomials for non-linear effect estimation in the presence of exposure measurement error. With real-world air pollution estimates, simulation experiments were conducted to examine novel approximation equations. Overall, sample size, the accuracy and precision of the estimators in these complex settings can be accurately approximated. For distributed lags, approximations may perform well if residual confounding due to covariate measurement errors is not severe. This condition may be difficult to identify without exposure validation data, so validation research is recommended in identifying critical exposure time-windows.

Keywords

Measurement error, Differential error, Autocorrelated error, Bias, Standard error

1. Introduction

Understanding of sample size and the accuracy and precision of the estimator may facilitate power and bias analyses. This helps in designing robust and reproducible research and ascertaining whether findings may imply causal effects or be attributable to random or systematic errors.

Case-crossover designs are matched case-control designs that utilize information from the comparison between the time of each incident (i.e., index time) and self-control times selected from pre-specified time points to estimate causal effects (Maclure 1991; Lee and Schwartz 1999; Mittleman and Mostofsky 2014). These designs are used to estimate various health effects of short-term exposures or acute health events using conditional logistic regression. Maclure (1991) proposed case-crossover designs for identifications of acute effects on myocardial infarction (MI). Mittleman and Mostofsky (2014) discussed exchangeability in case-crossover designs with examples of the relationship between sexual intercourse and MI, between illicit drug use and MI, between asthma exacerbation and death. Lee and Schwartz (1999) exemplified the use of case-crossover designs for environmental exposures such as air pollution. Numerous studies for other environmental exposures such as temperature and extreme weather events have been conducted (Bhaskaran et al. 2012; Tong, Wang, and Guo 2012; Wade et al. 2014). Associations between air pollution and COVID-19 have been also analyzed (Kim, Samet, and Bell 2022).

Despite its widespread use, understanding of sample size and the accuracy and precision of estimators remains limited. An existing sample size calculation method for conditional logistic regression (Lachin 2008) may not be directly applicable in research settings for several reasons.

First, the variance of exposures that may spatiotemporally vary may be heteroskedastic. Second, distributed (non-linear) lag models may be used to identify critical exposure time-windows or isolate exposure effects at a single time (Gasparrini, Armstrong, and Kenward 2010; Kim and Lee 2019; Mork et al. 2024). Third, exposure variables may be measured or estimated with error. The error may vary spatiotemporally. Various exposure measurement error structures may differently affect the accuracy and/or the precision of the estimator, consequently affecting sample size. Furthermore, in identifying critical exposure time-windows or isolating the effect of the exposure at a single time, residual confounding due to confounder/covariate measurement error (Greenland and Robins 1985; Fewell, Davey Smith, and Sterne 2007) may be concerning, because other times of the exposure may act as confounders, which may result in under/overestimation of the exposure time-window or the effect of exposure at a single time (Kim and Lee 2019). Fourth, in designing research, exposure validation data may not be directly available, but investigators may have information about summary statistics regarding the accuracy of exposure measurement, such as the coefficient of determination and regression coefficients. Whether this information can be used to predict the accuracy and precision of the estimators and consequently identify the sample size remains unclear.

Therefore, this article develops approximation equations for sample size, estimates of the estimators and standard errors for case-crossover designs with heteroskedastic distributed lag variables in the absence and presence of exposure measurement errors. For the accuracy of the estimators, this article develops several approximation equations including the most accurate one which can be used when exposure validation data is available and the least accurate one which

can be used when exposure validation data is not available. Simulation experiments based on real-world air pollution data were conducted to examine these equations.

2. Notations

Let $x_{i,t}$ denote the unmeasured true exposure of interest, x , for a case or self-control i at day t .

Let $x_{i,t-l}$ denote the temporally l -lagged variable of $x_{i,t}$, called *Lag* exposure. Let $x_{g,t}$ denote the exposure variable measured or estimated at spatial grid/area g (e.g., a 1km grid cell or a geographical delineation such as city or census tract) at day t . If not noted, I suppose that g is spatially less granular than case/self-control, i 's location (e.g., street address). $\mathbf{x}_{i \text{ or } g} =$

$(x_{i \text{ or } g,t}, \dots, x_{i \text{ or } g,t-L})$ and $\mathbf{x}_{i \text{ or } g}^{-l} = (x_{i \text{ or } g,t}, \dots, x_{i \text{ or } g,t-(l-1)}, x_{i \text{ or } g,t-(l+1)}, \dots, x_{i \text{ or } g,t-L})$ may be used. Let x_{t-l} denote $x_{i \text{ or } g,t-l}$ for notational convenience.

Let Y_i denote the outcome of a self-control ($Y_i = 0$) or a case ($Y_i = 1$). Let \mathbf{z} denote a set of covariates other than the main exposure variable(s) such as potential confounders, including dummy variables indicating each case-self-controls matched set. Assume n cases independently originate from a source population given x and \mathbf{z} , each matched to their respective self-controls, resulting in n matched sets. Each set has one case and $m_h - 1$ self-controls, resulting in a m_h -sized matched set, where h indicates h -th case. $x_{h,j,l}$ indicates x_{t-l} for j -th observation in h -th matched set. Let 1-st observation in each set be the case and the others be self-controls. Let $\sigma_h^2(v)$ denote the variance of a variable, v , in h -th matched set without “-1” correction. $V_{0h}(v)$ is the variance of a variable v in h -th matched set without “-1” correction (e.g., $\sigma_h^2(x_{t-l}) \equiv V_{0h}(x_{t-l})$). Superscript ep such as x^{ep} is used to denote a linear or classical (-like) error-prone variable. Other notations may be introduced when needed.

3. Development of Calculation Methods for Sample Size and Accuracy and Precision of Estimators

3.1. In the Absence of Exposure Measurement Error

3.1.1. Single Exposure Variable with Heteroskedastic Variance

By refining Lachin (2008)'s method, development begins with a single exposure variable x_{t-l} with its heteroskedastic variance as the only covariate. Under the alternative hypothesis, H_1 , the score equation, $U(\theta_l)$ for the coefficient for x_{t-l} in conditional logistic regression, θ_l , is

$$U(\theta_l) = \sum_{h=1}^n x_{h,1,l} - \frac{\sum_{j=1}^{m_h} x_{h,j,l} \exp(\theta_l x_{h,j,l})}{\sum_{j=1}^{m_h} \exp(\theta_l x_{h,j,l})}$$

The Fisher information is

$$I(\theta_l) = \sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} x_{h,j,l}^2 \exp(\theta_l x_{h,j,l})}{\sum_{j=1}^{m_h} \exp(\theta_l x_{h,j,l})} - \left(\frac{\sum_{j=1}^{m_h} x_{h,j,l} \exp(\theta_l x_{h,j,l})}{\sum_{j=1}^{m_h} \exp(\theta_l x_{h,j,l})} \right)^2 \right] = \sum_{h=1}^n V_{1h}(x_{t-l})$$

Under the null hypothesis, H_0 ,

$$\begin{aligned} U(\theta_l = 0) &= \sum_{h=1}^n \left(x_{h,1,l} - \frac{\sum_{j=1}^{m_h} x_{h,j,l}}{m_h} \right) = \sum_{h=1}^n (x_{h,1,l} - \bar{x}_{h,l}) \\ I(\theta_l = 0) &= \sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} x_{h,j,l}^2}{m_h} - \left(\frac{\sum_{j=1}^{m_h} x_{h,j,l}}{m_h} \right)^2 \right] \\ &= \sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} (x_{h,j,l} - \bar{x}_{h,l})^2}{m_h} \right] = \sum_{h=1}^n V_{0h}(x_{t-l}) \end{aligned}$$

where $\bar{x}_{h,l} = \frac{\sum_{j=1}^{m_h} x_{h,j,l}}{m_h}$. Recall $V_{0h}(x_{t-l}) \equiv \sigma_h^2(x_{t-l})$. Using a Taylor's expansion, the equation

for power analysis for a value of θ_l under H_1 becomes

$$|\theta_l| \sqrt{\sum_{h=1}^n V_{0h}(x_{t-l})} = Z_{1-\alpha} + Z_{1-\beta} \sqrt{\frac{\sum_{h=1}^n V_{1h}(x_{t-l})}{\sum_{h=1}^n V_{0h}(x_{t-l})}}$$

$$Z_{1-\beta} = \frac{|\theta_l| \sqrt{\sum_{h=1}^n V_{0h}(x_{t-l})} - Z_{1-\alpha}}{\sqrt{\frac{\sum_{h=1}^n V_{1h}(x_{t-l})}{\sum_{h=1}^n V_{0h}(x_{t-l})}}} \approx |\theta_l| \sqrt{\sum_{h=1}^n \sigma_h^2(x_{t-l})} - Z_{1-\alpha}$$

where $\sum_{h=1}^n V_{1h}(x_{t-l}) \approx \sum_{h=1}^n V_{0h}(x_{t-l})$ under a local alternative, $Z_{1-\alpha}$ is the critical value for the test (for a two-sided test, $Z_{1-\alpha/2}$). The standard error (SE) is

$$SE(\hat{\theta}_l) = \sqrt{I(\theta_l)}^{-1} = \sqrt{\sum_{h=1}^n V_{1h}(x_{t-l})}^{-1} \approx \sqrt{\sum_{h=1}^n \sigma_h^2(x_{t-l})}^{-1} \quad (\text{Eq.1})$$

Thus, $\sigma_h^2(x_{t-l})$ is a key factor that determines the precision of the estimator and power. Suppose heteroskedastic $\sigma_h^2(x_{t-l})$ over h because the variance of exposures can vary by space and time.

Let $\sum_{h=1}^n \sigma_h^2(x_{t-l}) = n\bar{\sigma}^2$ where $\bar{\sigma}^2 = \frac{\sum_{h=1}^n \sigma_h^2(x_{t-l})}{n}$. The variance of each temporally lagged variable is almost identical to that of other: $V(x_{t-1}) \approx V(x_{t-l})$, so that $\sigma_h^2(x_t) \approx \sigma_h^2(x_{t-l})$. To provide a desired level of power $1 - \beta$ when using a test at level α , n is required as

$$n \cong \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\theta_l^2 \bar{\sigma}^2} \quad (\text{Eq.2})$$

Although this is not a closed form because of $\bar{\sigma}^2$, an iterative process may be used to identify n . If investigators find that $\bar{\sigma}^2$ reaches a certain value over n , Eq.2 would be a closed solution. In practice, $\bar{\sigma}^2$ may be calculated using external data (e.g., population statistics). See Appendix S1.

3.1.2. Covariate Adjustment

With multiple covariates, the score test would be a C test and sample size calculation methods could be derived using a covariance matrix, which may be practically tedious. A deflation factor may be useful (Lachin 2008) as shown in exponential regression (Bernardo et al. 2000) and as covariate adjustment would reduce the precision of the estimator in logistic regression in observational studies (Robinson and Jewell 1991), so that

$$SE(\hat{\theta}_l|\mathbf{z}) \cong \sqrt{I(\theta_l)(1 - R_{x_{t-l}|\mathbf{z}}^2)^{-1}} \quad (\text{Eq.3})$$

For sample size calculation, n approximates

$$n \cong \frac{(z_{1-\alpha} + z_{1-\beta})^2}{\theta_l^2 \bar{\sigma}^2 (1 - R_{x_{t-l}|\mathbf{z}}^2)} \quad (\text{Eq.4})$$

where $R_{x_{t-l}|\mathbf{z}}^2$ is the coefficient of determination in the regression of x_{t-l} on \mathbf{z} , or squared Pearson correlation between x_{t-l} and the predicted x_{t-l} from the regression.

3.1.3. Distributed Lag Models

When the effect of x_{t-l} on Y is of interest, the exposure at other times, \mathbf{x}^{-l} can be seen as confounders when their effects exist (Kim and Lee 2019). Eqs.3–4 may be refined by adding \mathbf{x}^{-l} to \mathbf{z} . On the other hand, investigators may wish to identify the effect of cumulative exposure, from day 0 to day L . Then, the estimand for the effect of cumulative exposure is

$$\theta_0 x_t + \theta_1 x_{t-1} + \cdots + \theta_L x_{t-L} = \sum_{l=0}^L \theta_l x_{t-l}$$

For high L , constrained distributed lag variables may be used (Gasparrini, Armstrong, and Kenward 2010; Mork et al. 2024). For sample size calculation, it may be convenient to consider a weighted average as $\bar{x}_t = \sum_{l=0}^L \frac{\theta_l}{\bar{\theta}} x_{t-l}$ where $\bar{\theta} = \sum_{l=0}^L \theta_l$. For critical time-window identifications, the following may be alternatively used (Langholz et al. 1999),

$$\bar{x}_{t-l_s} = \sum_{l=l_s}^L w_l x_{t-l}$$

where w_l is a weight and $l_s > 0$. The weight may be pre-specified or estimated. For illustration simplicity, I adhere to \bar{x}_t . Then,

$$n \approx \frac{(z_{1-\alpha} + z_{1-\beta})^2}{\bar{\theta}^2 \bar{\sigma}_{\bar{x}_t}^2 (1 - R_{\bar{x}_t|\mathbf{z}}^2)} \quad (\text{Eq.5})$$

$$SE(\hat{\theta}|\mathbf{z}) \cong \sqrt{I(\bar{\theta})(1 - R_{\bar{x}_t|\mathbf{z}}^2)^{-1}} \quad (\text{Eq.6})$$

where $\bar{\sigma}_{\bar{x}_t}^2 = \frac{\sum_{h=1}^n \sigma_h^2(\bar{x}_t)}{n}$ and $I(\bar{\theta}) = \sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} \bar{x}_{h,j}^2 \exp(\bar{\theta} \bar{x}_{h,j})}{\sum_{j=1}^{m_h} \exp(\bar{\theta} \bar{x}_{h,j})} - \left(\frac{\sum_{j=1}^{m_h} \bar{x}_{h,j} \exp(\bar{\theta} \bar{x}_{h,j})}{\sum_{j=1}^{m_h} \exp(\bar{\theta} \bar{x}_{h,j})} \right)^2 \right] \approx n \bar{\sigma}_{\bar{x}_t}^2$.

When the weight is unknown, investigators may use different pre-specified weights and examine how sample size and precision may vary over weights.

3.2. Non-Differential Additive Errors

Non-differential exposure error means that the error is independent of Y .

Exposure variables such as air pollution and temperature variables may be provided at locations (e.g., measurement at environmental monitors), spatial grids, or a geographical delineation (e.g., city, census boundary). When $x_{g,t}$ is used in the place of $x_{i,t}$, the resulting measurement error is Berkson error as

$$x_{i,t} = x_{g,t} + u_{i,t}^B$$

$$u_{i,t}^B \sim N(0, \varphi_B^2)$$

where $u_{i,t}^B$ is independent and identically distributed with constant φ_B^2 . The Berkson error does not introduce bias in effect estimates in linear regression. In non-linear models such as logistic regression, Berkson error introduces negligible bias, particularly if the effect size is small (Keogh et al. 2020).

Suppose a distributed lag model with \mathbf{x}_g and \mathbf{z} as covariates are fit to estimate θ_l . Then,

$$n \cong \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\theta_l^2 \bar{\sigma}_{x_{g,t-l}}^2 \left(1 - R_{x_{g,t-l}|\mathbf{z}}^2 \right)} \quad (\text{Eq.7})$$

$$SE(\hat{\theta}_l | \mathbf{x}_g^{-l}, \mathbf{z}) \cong \sqrt{\sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} x_{g,h,j,l}^2 \exp(\theta_l x_{g,h,j,l})}{\sum_{j=1}^{m_h} \exp(\theta_l x_{g,h,j,l})} - \left(\frac{\sum_{j=1}^{m_h} x_{g,h,j,l} \exp(\theta_l x_{g,h,j,l})}{\sum_{j=1}^{m_h} \exp(\theta_l x_{g,h,j,l})} \right)^2 \right]} \left(1 - R_{x_{g,t-l} | \mathbf{x}_g^{-l}, \mathbf{z}}^2\right)^{-1} \approx \sqrt{n \bar{\sigma}_{x_{g,t-l}}^2 \left(1 - R_{x_{g,t-l} | \mathbf{x}_g^{-l}, \mathbf{z}}^2\right)^{-1}} \quad (\text{Eq.8})$$

where $\bar{\sigma}_{x_{g,t-l}}^2 = \frac{\sum_{h=1}^n \sigma_h^2(x_{g,t-l})}{n}$. The variability of $x_{g,t-l}$ is smaller than the variability of $x_{i,t-l}$ by φ_B^2 , which explains the precision reduction. Covariate measurement errors in \mathbf{x}_g^{-l} would not introduce residual confounding if the error is non-differential Berkson because $u_{i,t-l}^B$ is not correlated with $x_{g,t-l}$.

Other error may arise. Exposure measurements and estimation methods may be error-prone.

Suppose a linear measurement error model,

$$x_{g,t}^{ep} = \gamma_0 + \gamma_1 x_{g,t} + u_{g,t}^L$$

$$u_{g,t}^L \sim N(0, \varphi_L^2)$$

where $u_{g,t}^L$ is independent and identically distributed with constant φ_L^2 . For illustration, suppose the Berkson error is absent (i.e., $x_g = x_i$). In linear regression, this error results in either underestimation or overestimation of effects by a factor of

$$\lambda_l = \frac{\gamma_1 V(x_{g,t-l} | \mathbf{z})}{\gamma_1^2 V(x_{g,t-l} | \mathbf{z}) + V(u_{g,t-l}^L | \mathbf{z})} = \frac{\gamma_1 V(x_{g,t-l} | \mathbf{z})}{V(x_{g,t-l}^{ep} | \mathbf{z})}$$

which is called multiplicative factor (Carroll et al. 2006) or bias factor (Innes et al. 2021).

$$\theta_l^L = \lambda_l \theta_l$$

where θ_l^L denotes the quantity biased by the error, assuming no residual confounding by \mathbf{x}_g^{-l} due to the use of \mathbf{x}_g^{ep-l} (i.e., covariate measurement errors). This assumption will be lifted later. $V(\cdot|\mathbf{z})$ denotes the residual variance of a variable conditional on \mathbf{z} . For temporally lagged variables, it is likely $V(x_{g,t}|\mathbf{z}) \approx V(x_{g,t-l}|\mathbf{z})$. $\lambda\theta_l$ is a good approximation in logistic regression if the measurement error is small, the effect size is small (Carroll et al. 2006), or logistic regression in case-crossover analyses may be equal or similar to aggregated time-series analysis with log-linear regression (Lu and Zeger 2007). When $\gamma_1 = 1$, the resulting error is classical error, which results in underestimation: $0 < \lambda = \frac{V(x_{g,t-l}|\mathbf{z})}{V(x_{g,t-l}^{ep}|\mathbf{z})} < 1$.

Sample size may be calculated as

$$n \cong \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\lambda_l^2 \theta_l^2 \bar{\sigma}_{x_{g,t-l}^{ep}}^2 \left(1 - R_{x_{g,t-l}^{ep}|\mathbf{x}_g^{ep-l}, \mathbf{z}}^2\right)} \quad (\text{Eq.9})$$

$$SE\left(\hat{\theta}_l^L|\mathbf{x}_g^{ep-l}, \mathbf{z}\right) \cong$$

$$\sqrt{\sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} x_{g,h,j,l}^{ep2} \exp(\theta_l^L x_{g,h,j,l}^{ep})}{\sum_{j=1}^{m_h} \exp(\theta_l^L x_{g,h,j,l}^{ep})} - \left(\frac{\sum_{j=1}^{m_h} x_{g,h,j,l}^{ep} \exp(\theta_l^L x_{g,h,j,l}^{ep})}{\sum_{j=1}^{m_h} \exp(\theta_l^L x_{g,h,j,l}^{ep})} \right)^2 \right] \left(1 - R_{x_{g,t-l}^{ep}|\mathbf{x}_g^{ep-l}, \mathbf{z}}^2\right)}^{-1} \cong$$

$$\sqrt{n \bar{\sigma}_{x_{g,t-l}^{ep}}^2 \left(1 - R_{x_{g,t-l}^{ep}|\mathbf{x}_g^{ep-l}, \mathbf{z}}^2\right)}^{-1} \quad (\text{Eq.10})$$

When both Berkson and linear (or classical) errors arise (i.e., $x_g \neq x_i$), the error becomes compound error (CE). Eqs. 9 and 10 subsume CE.

When residual confounding by x_g^{-l} is not negligible, θ_t^l in Eq.9 needs to account for the residual confounding. Appendix S2 develops an approximation equation for θ_t^l using a re-written error model that can be seen in the context of regression calibration (Boe et al. 2023; Kim 2024). As will be shown, this may be the most accurate approximation equation in this paper, which is only available in practice when exposure validation data is available. Other approximation equations that can be used when exposure validation data is not available will be introduced later (Section 3.4).

3.3. Complex Additive Errors and Motivating Examples

3.3.1. Autocorrelated Errors

Autocorrelated errors may exist in environmental exposure estimates from prediction models. For example, let \mathbf{Q} denote a set of causal factors of air pollution or temperature exposures, which may exhibit spatiotemporal patterns. \mathbf{Q} may include emission activities, atmospheric process, solar radiation, and interaction with built-environment for air pollution; or atmospheric process, sunlight, and built-environment (e.g., urban heat island effect) for high temperature.

Spatiotemporally autocorrelated error may stem from unmeasured/overlooked causal factors and/or \mathbf{Q} measured with error, collectively, termed \mathbf{Q}^e . Let $f(x|\mathbf{Q})$ denote a true deterministic function of x . Let $f^e(x|\mathbf{Q}_s^e)$ denote an estimated or exposure prediction simulation function. $x_{g,t}^{ep}$ from $f^e(x|\mathbf{Q}_s^e)$ may have autocorrelated error, which may be described as

$$x_{g,t}^{ep} = \gamma_0 + \sum_{j=0}^J \eta_j x_{g,t-j} + u_{g,t}^{LL}$$

$\sum_{j=0}^J \eta_j x_{g,t-j}$ describes spatiotemporal autocorrelation process of $x_{g,t}^{ep}$ to some extent, analogous to autoregression. $u_{g,t}^{LL}$ may still be autocorrelated and not be independent of $x_{g,t}$. Let $u_{g,t}^{LL} = \delta(\mathbf{Q}^e) + u_{g,t}^{NI} + u_{g,t}^L$; $u_{g,t}^{NI} \sim g$. $\mathcal{E}(\mathbf{Q}^e)$ indicates association between \mathbf{Q}^e and $x_{g,t}^{ep}$ that explains

spatiotemporal autocorrelation process to some extent. $u_{g,t}^{NI}$ is an error term indicating the remaining spatiotemporal autocorrelation, which follows an unknown multivariate distribution \mathcal{g} . The superscript, NI , indicates non-independence. Let this error model be referred to as a linear-like (LL) error model because this error model subsumes linear error (and classical error). For illustrational simplicity, suppose

$$x_{g,t}^{ep} = \gamma_0 + \gamma_1 x_{g,t} + u_{g,t}^{LL}$$

by letting $u_{g,t}^{LL} = \sum_{j=1}^J \eta_j x_{g,t-j} + \delta(\mathbf{Q}^e) + u_{g,t}^{NI} + u_{g,t}^L$ and $\gamma_1 \equiv \eta_0$.

The approximation equation developed in Appendix S2 and other approximations that will be introduced in Section 3.4 can be applied to θ_t^{LL} , the quantity biased by LL error. Finally,

$$n \cong \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\theta_t^{LL^2} \bar{\sigma}_{x_{g,t-l}^{ep}}^2 \left(1 - R_{x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z}}^2 \right)} \quad (\text{Eq.11})$$

$$SE \left(\hat{\theta}_t^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z} \right) \cong$$

$$\sqrt{\sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} x_{g,h,j,l}^{ep^2} \exp(\theta_t^{LL} x_{g,h,j,l}^{ep})}{\sum_{j=1}^{m_h} \exp(\theta_t^{LL} x_{g,h,j,l}^{ep})} - \left(\frac{\sum_{j=1}^{m_h} x_{g,h,j,l}^{ep} \exp(\theta_t^{LL} x_{g,h,j,l}^{ep})}{\sum_{j=1}^{m_h} \exp(\theta_t^{LL} x_{g,h,j,l}^{ep})} \right)^2 \right] \left(1 - R_{x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z}}^2 \right)^{-1}} \cong$$

$$\sqrt{n \bar{\sigma}_{x_{g,t-l}^{ep}}^2 \left(1 - R_{x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z}}^2 \right)^{-1}} \quad (\text{Eq.12})$$

And for the estimator for cumulative exposure,

$$n = \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\bar{\theta}^{LL^2} \bar{\sigma}_{x_{g,t}}^2 \left(1 - R_{x_{g,t}^{ep} | \mathbf{z}}^2 \right)} \quad (\text{Eq.13})$$

$$SE(\hat{\theta}^{LL}|\mathbf{z}) = \sqrt{\sum_{h=1}^n \left[\frac{\sum_{j=1}^{m_h} \bar{x}_{g,h,j}^{ep2} \exp(\bar{\theta}^{LL} \bar{x}_{g,h,j}^{ep})}{\sum_{j=1}^{m_h} \exp(\bar{\theta}^{LL} \bar{x}_{g,h,j}^{ep})} - \left(\frac{\sum_{j=1}^{m_h} \bar{x}_{g,h,j} \exp(\bar{\theta}^{LL} \bar{x}_{g,h,j}^{ep})}{\sum_{j=1}^{m_h} \exp(\bar{\theta}^{LL} \bar{x}_{g,h,j}^{ep})} \right)^2 \right]} \left(1 - R_{\bar{x}_{g,t}^{ep}|\mathbf{z}}^2\right)^{-1} \approx \sqrt{n \bar{\sigma}_{\bar{x}_{g,t}^{ep}}^2 \left(1 - R_{\bar{x}_{g,t}^{ep}|\mathbf{z}}^2\right)^{-1}} \quad (\text{Eq.14})$$

where $\bar{x}_{g,t}^{ep} = \sum_{l=0}^L \frac{\theta_l^{LL}}{\bar{\theta}^{LL}} x_{g,t-l}^{ep} \approx \sum_{l=0}^L \frac{\theta_l}{\bar{\theta}} x_{g,t-l}^{ep}$.

Berkson-like (BL) error may arise ($x_g \neq x_i$). For illustration, suppose the error model,

$$x_{i,t} = x_{g,t} + u_{i,t}^{BL}$$

where $u_{i,t}^{BL} = u_{i,t}^B + u_{i,t}^{NI}$ and $u_{i,t}^{NI}$ denotes *NI* error. Unlike Berkson error, BL error may introduce bias in effect estimates because $u_{i,t}^{BL}$ may be correlated with $x_{g,t}$. Let θ_l^{BL} denote the quantity biased by the error,

$$\theta_l^{BL} \cong \theta_l + \sum_{k=0}^L \theta_k \varrho_{u_{i,t-k}^{BL}, x_{g,t-l} | x_g^{-l}, \mathbf{z}}$$

where $\varrho_{u_{i,t-k}^{BL}, x_{g,t-l} | x_g^{-l}, \mathbf{z}}$ is the regression coefficient for $x_{g,t-l}$ when $u_{i,t-k}^{BL}$ is regressed on $x_{g,t}$ and \mathbf{z} . For sample size, Eqs.7–10 can be used by replacing θ_l^B and $\bar{\theta}^B$ with θ_l^{BL} and $\bar{\theta}^{BL}$. When both BL and LL errors exist, Eqs.11–14 can be used by replacing θ_l^{LL} and $\bar{\theta}^{LL}$ with $\theta_l^{CE} \cong \theta_l^{LL} + \sum_{k=0}^L \theta_k \varrho_{u_{i,t-k}^{BL}, x_{g,t-l} | x_g^{-l}, \mathbf{z}}$ and $\bar{\theta}^{CE} = \sum_{l=0}^L \theta_l^{CE}$.

3.3.2. Differential Errors

Differential exposure measurement error means that the error is not independent of Y . I find at least five mechanisms through which the error can become differential: 1) the correlation between $x_{g,t}$ and $u_{g,t}^{LL}$ through which $u_{g,t}^{LL}$ is correlated with Y when $x_{g,t}$ has effects on Y but $x_{g,t}^{ep}$

is used (Figure 1A); 2) the correlation between $x_{g,t}$ and $u_{i,t}^{BL}$ through which $u_{i,t}^{BL}$ is correlated with Y when $x_{i,t}$ has effects on Y but $x_{g,t}$ is used (Figure 1B); 3) $u_{g,t}^{LL}$ or $u_{g,t}^{BL}$ as a risk factor of Y or related to non-confounder risk factors of Y (Figures 1C–D); 4) the correlation of $u_{g,t}^{LL}$, or $u_{g,t}^{BL}$ with unmeasured confounders; and 5) the correlation of $u_{g,t}^{LL}$, or $u_{g,t}^{BL}$ with confounder measurement error when a confounder is measured with error. Thus, BL and LL errors would be differential if there exist exposure effects.




Figure 1. Graphical descriptions of linear-like (LL) exposure measurement error (A, C) and Berkson-like (BL) exposure measurement error (B, D)

Note. Dashed lines denote potential correlation.

Potential bias to the estimators needs to be considered to accurately calculate sample size. But the bias may be smaller than what the face value of differential error may imply. For example, differential errors in distribute lag models may mitigate residual confounding due to the use of x_g^{ep-l} (Appendix S3). Differential errors may be (coincidentally) adjusted for (to some degree) by using self-control matching or adjustment for covariates that makes the error differential, so that the resulting error may become conditionally non-differential (Kim 2024). Variables or statistical terms for adjustment for differential errors could be subsumed into z (Appendix S4). For illustration, suppose both the error and the risk of Y vary over seasons or years. The error is

differential by the definition. Self-control selections may adjust for this differential error when for effect estimation, the comparison of $x_{g,t-l}^{ep}$ of a case to that of self-controls within a short-time period was used, in which the error may be locally non-differential. The matched strata can be described as matching stratum indicator variables(s) in \mathbf{z} .

3.3.3. Multiplicative Errors

Exposure measurement error may be multiplicative. Multiplicative errors may be transformed to additive errors. For example, multiplicative Berkson (-like) (MB(L)) error may be described as

$$\log(x_{i,t}) = \log(x_{g,t}) + u_{i,t}^{B(L)}$$

$$x_{i,t} = x_{g,t} \exp(u_{i,t}^{B(L)})$$

If $\log(x_{g,t})$ is used as the exposure variable in an outcome model, the consequence would be identical to that for the additive error, assuming that $\log(x_{g,t})$ can correctly represent $x - Y$ relationship. If $x_{g,t}$ is used and the error is not small, the consequence may differ. If residual confounding due to the use of x_g is negligible, $\theta_l^{MB(L)}$, the quantity biased by the error would be

$$\theta_l^{MB(L)} \approx \theta_l E \left[\exp(u_{i,t}^{B(L)}) \right]$$

Note $E \left[\exp(u_{i,t}^{B(L)}) \right] \neq \exp \left(E[u_{i,t}^{B(L)}] \right) = 1$.

Multiplicative linear (-like) (L(L)) error may be described as

$$\log(x_{g,t}^{ep}) = \gamma_{m0} + \gamma_{m1} \log(x_{g,t}) + u_{g,t}^{L(L)}$$

$$x_{g,t}^{ep} = (x_{g,t})^{\gamma_{m1}} \exp(\gamma_{m0}) \exp(u_{g,t}^{L(L)})$$

If $x_{g,t}^{ep}$ is used as the exposure variable and the error is not small, the consequence would differ.

For example, for multiplicative classical (MC) error (i.e., $\gamma_{m1} = 1$ and $\gamma_{m0} = 0$), assuming that the residual confounding due to the use of x_g^{ep-l} is negligible,

$$\theta_l^{MC} \cong \theta_l \frac{V(x_{g,t-l}|\mathbf{z})}{V(x_{g,t-l}|\mathbf{z}) \left(1 + V(u_{g,t}^L|\mathbf{z})\right) + E[x_{g,t-l}|\mathbf{z}]^2 V(u_{g,t}^L|\mathbf{z})}$$

demonstrating attenuation.

To approximate $\theta_l^{ML(L)}$, the quantity biased by the multiplicative L(L) (ML(L)) error, the approximation equation in Appendix S2 can be used for many different error models (Appendix S4) but requires the use of exposure validation data. Without exposure validation data, approximation equations that will be introduced in Section 3.4 may also be used.

Approximation of the variance of the estimator can also be obtained. The standard error of $\theta_l^{ML(L)}$ may be approximated as

$$SE\left(\hat{\theta}_l^{ML(L)}|\mathbf{z}, x_g^{ep-l}\right) \cong \sqrt{\left(n\bar{\sigma}_{x_{g,t-l}}^{ML(L)}\left(1 - R_{x_{g,t-l}^{ep}|x_g^{ep-l},\mathbf{z}}^2\right)\right)^{-1} + \theta_l^{ML(L)2}\left(n\bar{\sigma}_{\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})}^2\left(1 - R_{x_{g,t-l}^{ep}|x_g^{ep-l},\mathbf{z}}^2\right)\right)^{-1}} \quad (\text{Eq.15})$$

where let $x_{g,t-l}^{ML(L)} = (x_{g,t-l})^{\gamma_{m1}}$ (Appendix S5). The term $(x_{g,t-l})^{\gamma_{m1}}$ demonstrates the standard error can be either attenuated or inflated, depending on γ_{m1} . The inflation may arise if $\gamma_{m1} < 1$ when $x_{g,t-l} > 1$. The second component in the square root demonstrates that the standard error may be inflated, but this component could be relatively small compared to the first component, especially for small effect size, so that this may be negligible.

Similarly, the standard error of $\bar{\theta}^{ML(L)}$, which is the quantity regarding the cumulative exposure effect, may be approximated as

$$SE\left(\hat{\theta}^{ML(L)}|\mathbf{z}\right) \cong \sqrt{\left(n\bar{\sigma}_{\bar{x}_{g,t}}^{ML(L)}\left(1-R_{\bar{x}_{g,t}^{ep}|\mathbf{z}}^2\right)\right)^{-1} + \bar{\theta}^{ML(L)2}\left(n\bar{\sigma}_{\exp(\gamma_{m0})\exp(\bar{u}_{g,t}^{L(L)})}^2\left(1-R_{\bar{x}_{g,t}^{ep}|\mathbf{z}}^2\right)\right)^{-1}} \quad (\text{Eq.16})$$

where $\bar{x}_{g,t}^{ML(L)} = \exp(\gamma_{m0})E\left[\exp(u_{g,t}^{L(L)})\right]\left(\frac{\theta_0}{\bar{\theta}}(x_{g,t})^{\gamma_{m1}} + \dots + \frac{\theta_L}{\bar{\theta}}(x_{g,t-L})^{\gamma_{m1}}\right)$ and $\bar{u}_{g,t}^{L(L)} = \frac{\theta_0}{\bar{\theta}}u_{g,t}^{L(L)} + \dots + \frac{\theta_L}{\bar{\theta}}u_{g,t}^{L(L)}$.

Sample size may be calculated as

$$n \cong \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\theta_l^{ML(L)2} \bar{\sigma}_{x_{g,t-l}}^{ML(L)2} \left(1 - R_{x_{g,t-l}^{ep}|\mathbf{x}_g^{ep-l}, \mathbf{z}}^2\right) VCF} \quad (\text{Eq.17})$$

$$n \cong \frac{(Z_{1-\alpha} + Z_{1-\beta})^2}{\bar{\theta}^{ML(L)2} \bar{\sigma}_{\bar{x}_{g,t}}^{ML(L)2} \left(1 - R_{\bar{x}_{g,t}^{ep}|\mathbf{z}}^2\right) VCF} \quad (\text{Eq.18})$$

where VCF denotes a variance correction factor,

$$VCF = \frac{\left(n\bar{\sigma}_{x_{g,t-l}}^{ML(L)}\right)^{-1}}{\left(n\bar{\sigma}_{x_{g,t-l}}^{ML(L)}\right)^{-1} + \theta_l^{ML(L)2} \left(n\bar{\sigma}_{\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})}^2\right)^{-1}}$$

or

$$VCF = \frac{\left(n\bar{\sigma}_{\bar{x}_{g,t}}^{ML(L)}\right)^{-1}}{\left(n\bar{\sigma}_{\bar{x}_{g,t}}^{ML(L)}\right)^{-1} + \bar{\theta}^{ML(L)2} \left(n\bar{\sigma}_{\exp(\gamma_{m0})\exp(\bar{u}_{g,t}^{L(L)})}^2\right)^{-1}}$$

for small $\theta_l^{ML(L)}$ or $\bar{\theta}^{ML(L)}$, $VCF \approx 1$

For multiplicative CE (i.e., $x_{g,t} \neq x_{i,t}$), Eqs.15–18 can be used by considering additional bias due to BL error to the estimator. For only MBL error, Eqs. 7–10 can be used.

3.4. Without Exposure Validation Data in Hand

Investigators may not have exposure validation data in practice (i.e., having only $x_{g,t}$ or $x_{g,t}^{ep}$, but not $x_{i,t}$).

3.4.1. $\bar{\sigma}^2$ Calculation

For additive errors, the exposure variable measured/estimated with error that investigators have, which would be $x_{g,t}$ for B(L) error or $x_{g,t}^{ep}$ for non-Berkson error or CE, can be directly used to calculate $\bar{\sigma}^2$ (Note Eqs.7–14 that are based on $x_{g,t}$ or $x_{g,t}^{ep}$). For ML(L), what investigators have would be $x_{g,t}^{ep}$. $x_{g,t}^{ep}$ needs to be divided by the square root of $\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})$, which is unknown without validation data. Since the variance decomposition for $(x_{g,t})^{\gamma_{m1}}\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})$ may vary by the distribution of $x_{g,t}$ and $u_{i,t}^{L(L)}$, the generalizable decomposition may not be obtainable if $u_{i,t}^{L(L)}$ is unknown. Alternatively, it may be reasonable to consider some factor c as

$$\bar{\sigma}_{x_{g,t-l}^{ML(L)}}^2 \cong \bar{\sigma}_{x_{g,t-l}^{ep}}^2 c$$

and conduct numerical analyses about how sample size numbers could vary by several c values. However, investigators may, sometimes, have information about the error from external sources such as published documents. Suppose that the coefficient of determination in the regression of the logarithm of $x_{g,t}^{ep}$ against the logarithm of $x_{g,t}$ ($R_{\log(x_{g,t}^{ep}),\log(x_{g,t})}^2$) and γ_{m1}^{crude} as the regression

coefficient are known and the coefficient in the regression of $x_{g,t}$ against $x_{g,t}^{ep}$ are known. Recall the following ML(L) model,

$$\begin{aligned}\log(x_{g,t}^{ep}) &= \gamma_{m0} + \gamma_{m1} \log(x_{g,t}) + u_{g,t}^{L(L)} \\ \log(x_{g,t-l}^{ep}) &= \gamma_{m0} + \gamma_{m1} \log(x_{g,t-l}) + u_{g,t-l}^{L(L)}\end{aligned}$$

and,

$$R_{\log(x_{g,t}^{ep}), \log(x_{g,t})}^2 \cong R_{\log(x_{g,t-l}^{ep}), \log(x_{g,t-l})}^2 = \frac{V(\gamma_{m0} + \gamma_{m1} \log(x_{g,t-l}))}{V(\log(x_{g,t-l}^{ep}))} = \frac{V(\gamma_{m1} \log(x_{g,t-l}))}{V(\log(x_{g,t-l}^{ep}))}$$

for $\gamma_{m1} > 0$,

$$\bar{\sigma}_{x_{g,t-l}^{ML(L)}}^2 \approx \bar{\sigma}_{\log(x_{g,t-l}^{ep})}^2 E[x_{g,t-l}]^{2\gamma_{m1}} R_{\log(x_{g,t-l}^{ep}), \log(x_{g,t-l})}^2$$

(Appendix S6). Thus, by assuming $\gamma_{m1}^{crude} \cong \gamma_{m1}$, which may be reasonable in practice when

investigators have domain knowledge about exposure measurements or estimations, and

calculating $E[x_{g,t-l}]$ based on the information of the regression of $x_{g,t}$ against $x_{g,t}^{ep}$, they can

calculate $\bar{\sigma}_{\log(x_{g,t-l}^{ep})}^2$ with their data and then approximate $\bar{\sigma}_{x_{g,t-l}^{ML(L)}}^2$. $\bar{\sigma}_{x_g^{ML(L)}}^2$ can also be

approximated similarly. Sometimes, $R_{\log(x_{g,t}^{ep}), \log(x_{g,t})}^2$ is not known, but $R_{x_{g,t-l}^{ep}, x_{g,t-l}}^2$, which may

be used instead.

3.4.2. Bias Consideration

For non-Berkson error, the estimator may be biased so that investigators should consider the size

of bias when calculating sample size. Recall the additive L(L) error model,

$$x_{g,t}^{ep} = \gamma_0 + \gamma_1 x_{g,t} + u_{g,t}^{LL}$$

If γ_1^{crude} and $R_{x_{g,t}^{ep}, x_{g,t}}^2$ are known, by assuming $\gamma_1^{crude} = \gamma_1$, in distributed lag models,

$$\theta_l^{L(L)} \approx R_{x_{g,t}, x_{g,t}}^{2ep} \left(\frac{\theta_l \pm (1 - R_{x_{g,t}, x_{g,t}}^{2ep}) \sum_{j \neq l}^L \theta_j}{\gamma_1} \right) \quad (\text{Eq.19})$$

and this approximation may perform reasonably if residual confounding due to the use of x_g^{ep-l} is not severe (Appendix S7). If “+” is used for “±” in the numerator, Eq.19 may indicate the upper bound of $\theta_l^{L(L)}$ to identify whether $\theta_l^{L(L)} \leq \theta_l$ is plausible but may result in underapproximation of sample size. If $x_{g,t}$ is highly autocorrelated and $x_{g,t}^{ep}$ is a good measure of $x_{g,t}$, “-” may be more accurate than “+” (Appendix S7). If residual confounding is negligible, or $\bar{\theta}$ is of interest,

$$\theta_l^{L(L)} \approx \frac{R_{x_{g,t}, x_{g,t}}^{2ep} \theta_l}{\gamma_1} \text{ or } \bar{\theta}^{L(L)} \approx \frac{R_{x_{g,t}, x_{g,t}}^{2ep} \bar{\theta}}{\gamma_1} \quad (\text{Eq.20})$$

Multiplicative error may be treated as additive error. The ML(L) error model may be re-written as

$$x_{g,t}^{ep} = \gamma_0^{ML(L)} + \gamma_1^{crude} x_{g,t} + u_{g,t}^{ML(L)}$$

where $u_{g,t}^{ML(L)}$ mimicks $u_{g,t}^{L(L)}$ in which an interaction between $x_{g,t}$ and another error component(s) exists but is unknown. Validation studies may report γ_1^{crude} and $R_{x_t^{ep}, x_t}^2$ when whether the error is additive or multiplicative remains uncertain.

Eq.20 relies on the assumption $R_{x_{g,t}, x_g}^{2ep} \approx R_{x_{g,t-l}, x_{g,t-l} | x_g^{ep-l}, z}^2$. It is not unrealistic that $R_{x_{g,t}, x_g}^{2ep} \gg$

$R_{x_{g,t-l}, x_{g,t-l} | x_g^{ep-l}, z}^2$ in practice (Appendix S7). If $R_{x_{g,t-l}, x_{g,t-l} | x_g^{ep-l}, z}^2$ is not known, it may be wise

to apply a multiplication factor to $R_{x_{g,t}, x_g}^{2ep}$ and examine how sample size can vary accordingly by

changing that factor.

3.5. Polynomials for Non-Linear Effects

Non-linear terms such as splines and polynomials may be used. Considering exact variable transformation may be too tedious. It would be practical to focus on a second-order polynomial term, $x_{t-l}^{ep\ 2}$, because testing this term may demonstrate whether the linear term is inadequate to explain Y . If the exposure effect is believed to be non-linear only within the certain range of x_{t-l}^{ep} (e.g., high temperature effect manifests only after a certain threshold), the variable could be transformed to consider this.

Covariate adjustment should be considered as $R^2_{(x_{t-l}^{ep})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z}} \cdot x_{t-l}^{ep} \times \mathbf{x}^{ep-l}$ denotes interaction pairs of x_{t-l}^{ep} and each element in \mathbf{x}^{ep-l} . \mathbf{z}^2 denote the squared variables in \mathbf{z} . $\mathbf{x}^{ep} \times \mathbf{z}$ denotes interaction pairs of x_{t-l}^{ep} and each element in \mathbf{z} . The bias factor of the estimator for $(x_{t-l}^{ep})^2$ may be approximated as by assuming that residual confounding due to the use of \mathbf{x}^{ep-l} is negligible and the residual of $(x_{t-l}^{ep})^2$ conditional on $\mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z}$ and the residual of x_{t-l}^{ep} conditional on $\mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2$ follow a normal distribution,

$$\lambda_l^2 \approx \frac{\left(R^2_{x_{t-l}^{ep}, x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2} \right)^2}{\gamma_1^2} \approx \frac{\left(R^2_{x_t^{ep}, x_t} \right)^2}{\gamma_1^2}$$

(Appendix S8). Again, $R^2_{x_t^{ep}, x_t}$ should be a good approximate. Approximations would become tedious if the assumption is not correct.

4. Simulation Experiments

Simulation experiments were motivated by air pollution modelled estimates. Figure 2 presents relationships of ground-based measurements for daily O_3 and $PM_{2.5}$ with estimated levels in Rio de Janeiro, Brazil (2010–2017) (A, B, E, and F) obtained from elsewhere (Kim et al. 2024) and estimated levels in the contiguous United States (2020 summer) (C, D, G, and H) from a frequently cited model in air pollution epidemiology (Reff 2023). These relationships were identified using linear regression (red lines) from the BL error perspective (A–D) and from the LL error perspective. Notably, a regression coefficient of nearly one from one perspective does not necessarily imply that from the other perspective, which is statistically legitimate. Depending on the prediction modeling strategy, the correct perspective in health effect estimation may differ, which is beyond the scope of this paper. Table 1 show findings of linear regressions for data in the United States suggesting the error of these estimates may be BL or LL error in health effect estimation. Figure S1 in Appendix S9 show residual plots and residuals are autocorrelated.

Simulation experiments were conducted. These mimicked identifications of the association between short-term exposure to $PM_{2.5}$ and a health outcome, using $PM_{2.5}$ variables measured with errors. Detailed methods are provided in Appendix S9. Confounders to be considered include short-term exposure to O_3 and temperature, as well as unmeasured time-varying confounders that manifest as seasonality, time-trend, and day-of-the-week effects. Several additive and multiplicative BL, and LL errors in $PM_{2.5}$ variables were considered, using ground-based measurements at EPA monitors and modelled estimates. Figure 3 present scatter plots for gold standard variables and variables measured with error in simulations.

Table 1. Linear regressions of measured O₃/PM_{2.5} at EPA monitors against distributed temporal lags of estimated O₃/PM_{2.5} (Berkson-like error perspective) by U.S. EPA’s FAQSD models (Jun–Aug. 2020) and of estimated O₃/PM_{2.5} by U.S. EPA’s FAQSD models (Jun–Aug. 2020) against distributed temporal lags of measured O₃/PM_{2.5} at EPA monitors (Linear-like error perspective)

Note. Figure S1 presents residual plots including partial autocorrelation plots that demonstrate that residuals are not random.

Parameter	Regression coefficient estimates (95% confidence intervals)			
	O ₃		PM _{2.5}	
	Berkson-Like error perspective	Linear-Like error perspective	Berkson-Like error perspective	Linear-Like error perspective
Intercept	-0.931 (-1.011, -0.851)	2.225 (2.152, 2.299)	-0.362 (-0.404, -0.321)	0.901 (0.866, 0.936)
Lag0	1.029 (1.027, 1.032)	0.929 (0.927, 0.931)	1.065 (1.061, 1.070)	0.840 (0.837, 0.844)
Lag1	-0.009 (-0.011, -0.006)	0.023 (0.021, 0.025)	-0.026 (-0.032, -0.020)	0.059 (0.055, 0.063)
Lag2	-0.001 (-0.004, 0.001)	0.002 (0, 0.004)	0 (-0.006, 0.005)	0 (-0.005, 0.004)
Lag3	0 (-0.002, 0.003)	0.001 (-0.001, 0.003)	-0.006 (-0.012, -0.001)	0.008 (0.003, 0.012)
Lag4	0 (-0.002, 0.003)	-0.002 (-0.004, 0)	0.002 (-0.004, 0.008)	-0.003 (-0.007, 0.001)
Lag5	0.001 (-0.001, 0.004)	-0.003 (-0.005, -0.001)	0 (-0.006, 0.006)	0 (-0.004, 0.004)
Lag6	0 (-0.002, 0.003)	-0.002 (-0.004, 0)	0.001 (-0.005, 0.006)	-0.002 (-0.006, 0.002)
Lag7	0 (-0.002, 0.002)	-0.002 (-0.003, 0)	-0.005 (-0.01, -0.001)	0.003 (-0.001, 0.006)
R-sq.	96.6%	96.6%	92.0%	92.2%




Figure 2. Relationship between daily estimated O₃/PM_{2.5} and daily measured O₃/PM_{2.5} at monitors in Rio de Janeiro, Brazil for the years 2012–2017 (A, B, E, and F) and in the contiguous United States (C, D, G, and H) for Jun.–Aug. 2020

Note. The difference between the top four panels (Berkson-like error perspective, A–D) and the bottom four panels (linear-like error perspective, E–H) is that the same data were analyzed differently by switching the x and y axes. Dashed diagonal lines indicates 100% agreement. Solid lines indicate linear regression fit. Points indicate the number of data points.




Figure 3. Scatter plots for several sets of gold standard variables and variables measured with error created in simulation experiments

Note. Eight measurement error scenarios were created. MLL=Multiplicative Linear-Like error; ALL=Additive Linear-Like error; ABL=Additive Berkson-Like error; and MBL=Multiplicative Berkson-Like error. Only randomly selected 10,000 data points from each of the first 100 simulation samples (for MLL and ALL) and randomly selected 10,000 data points from each of the first 250 simulation samples (for ABL and MBL) are presented to enhance visualizations. Solid lines represent the linear regression fit for each sample that consists of 10,000 data points and represent the average of these linear regression fits. For MLL data generation, $\gamma_1=0.9$ (MLL1 and MLL2) or $\gamma_1=1.1$ (MLL3 and MLL4) were used.

Sample sizes, $E[\hat{\theta}_l]$, $E[\hat{\theta}]$, $E[\widehat{SE}(\hat{\theta}_l)]$, and $E[\widehat{SE}(\hat{\theta})]$ were calculated using developed approximation equations earlier. They were compared to corresponding empirical quantities estimated from time-stratified case-crossover analyses for 2,500 simulation runs for each error scenario. Findings demonstrate that the calculations perform well. Tables 2 and 3 demonstrate empirical effect estimates and standard error estimates are consistent with the approximates. As expected, Eqs.19 and 20 may sometimes provide accurate approximates for effect estimates, suggesting that it would not be easy to accurately consider bias in the estimator without validation data when residual confounding due to distributed lags measured with error is not small. Figure 4 shows the consistency between empirical statistical powers (i.e., the percentage of rejecting the null hypothesis when the null hypothesis is false) and the calculated power curves. The inconsistencies were found in only very low statistical power settings for θ_l estimation. These may have been attributed to a failure of the convergence due to correlation between distributed lags in such settings because the accuracy (See “Approx. 1” in Table 2) and precision of the estimators were still accurately calculated (Table 3).

Table 2. Empirical estimates of θ_l and $\bar{\theta}$ and various approximations for θ_l and $\bar{\theta}$ in simulation experiments

Note. MLL=Multiplicative Linear-Like error; ALL=Additive Linear-Like error; ABL=Additive Berkson-Like error; MBL=Multiplicative Berkson-Like error. See Figure 3 for distributions of MLL1–4, ALL1–2, ABL, and MBL.

Lag0= θ_0 ; Lag1= θ_1 ; Lag2= θ_2 ; and Lag02= $\bar{\theta} = \theta_0 + \theta_1 + \theta_2$. $\theta_0 = 0.001$, $\theta_1 = 0.0024$, $\theta_3 = 0.0006$, and $\bar{\theta} = 0.004$ were used to generate simulation samples. Except for ABL, most of the effect estimates were biased due to the error. See “a. Empirical”, which shows $E[\hat{\theta}_l] \times 1000$.

Error	Quantities	Lag0	Lag1	Lag2	Lag02
No error	θ used to generate data $\times 1000$	1.00	2.40	0.60	4.00
MLL1	a. Empirical ($\times 1000$)	1.62	3.38	1.02	6.01
	b. Approx.1 ($\times 1000$) [Appendix S2]	1.56	3.42	1.01	5.99
	c. Difference (%) (i.e., (b-a)/a)	-3.4	1.2	-0.7	-0.4
	d. Approx.2 ($\times 1000$) [Eq. 19] “+”	1.71	3.73	1.13	6.57

	e. Difference (%) (i.e., (d-a)/a)	5.8	10.5	11.3	9.4
	f. Approx.3($\times 1000$) [Eq. 19] “-”	1.31	3.52	0.68	5.52
	g. Difference (%) (i.e., (f-a)/a)	-18.7	4.3	-32.8	-8.2
	h. Approx.4 ($\times 1000$) [Eq. 20]	1.51	3.63	0.91	6.05
	i. Difference (%) (i.e., (h-a)/a)	-6.4	7.4	-10.7	0.6
MLL2	a. Empirical ($\times 1000$)	1.08	3.08	0.64	4.81
	b. Approx.1 ($\times 1000$) [Appendix S2]	1.09	3.14	0.64	4.87
	c. Difference (%) (i.e., (b-a)/a)	1.09	3.14	0.64	4.87
	d. Approx.2 ($\times 1000$) [Eq. 19] “+”	1.8	3.65	1.27	6.73
	e. Difference (%) (i.e., (d-a)/a)	67.2	18.5	98.4	40.1
	f. Approx.3($\times 1000$) [Eq. 19] “-”	1.07	3.26	0.45	4.78
	g. Difference (%) (i.e., (f-a)/a)	-0.7	5.8	-30.2	-0.5
	h. Approx.4 ($\times 1000$) [Eq. 20]	1.44	3.46	0.87	5.77
	i. Difference (%) (i.e., (h-a)/a)	33.7	12.3	34.9	20.1
MLL3	a. Empirical ($\times 1000$)	0.79	1.76	0.50	3.05
	b. Approx.1 ($\times 1000$) [Appendix S2]	0.77	1.80	0.48	3.05
	c. Difference (%) (i.e., (b-a)/a)	-2.2	2.2	-3.0	0.2
	d. Approx.2 ($\times 1000$) [Eq. 19] “+”	0.86	1.94	0.56	3.36
	e. Difference (%) (i.e., (d-a)/a)	9.5	10.1	11.8	10.2
	f. Approx.3($\times 1000$) [Eq. 19] “-”	0.72	1.86	0.39	2.96
	g. Difference (%) (i.e., (f-a)/a)	-9.3	5.6	-21.8	-2.7
	h. Approx.4 ($\times 1000$) [Eq. 20]	0.79	1.90	0.47	3.16
	i. Difference (%) (i.e., (h-a)/a)	0.1	7.9	-5.0	3.7
MLL4	a. Empirical ($\times 1000$)	0.58	1.67	0.36	2.61
	b. Approx.1 ($\times 1000$) [Appendix S2]	0.60	1.70	0.35	2.64
	c. Difference (%) (i.e., (b-a)/a)	2.4	1.4	-3.1	1.0
	d. Approx.2 ($\times 1000$) [Eq. 19] “+”	0.90	1.91	0.62	3.43
	e. Difference (%) (i.e., (d-a)/a)	55.6	13.9	73.3	31.3
	f. Approx.3($\times 1000$) [Eq. 19] “-”	0.62	1.75	0.29	2.67
	g. Difference (%) (i.e., (g-a)/a)	6.4	4.9	-17.4	2.2
	h. Approx.4 ($\times 1000$) [Eq. 20]	0.76	1.83	0.46	3.05
	i. Difference (%) (i.e., (i-a)/a)	31.2	9.5	28.4	16.9
ALL1	a. Empirical ($\times 1000$)	0.13	2.19	-0.15	2.16
	b. Approx.1 ($\times 1000$) [Appendix S2]	0.11	2.17	-0.14	2.15
	c. Difference (%) (i.e., (b-a)/a)	-10.9	-0.8	-9.8	-0.7
	d. Approx.2 ($\times 1000$) [Eq. 19] “+”	1.35	2.48	1.04	4.88
	e. Difference (%) (i.e., (d-a)/a)	957.8	13.4	-780.4	125.5
	f. Approx.3($\times 1000$) [Eq. 19] “-”	0.49	2.00	0.06	2.55
	g. Difference (%) (i.e., (f-a)/a)	282.0	-8.7	-140.5	17.9
	h. Approx.4 ($\times 1000$) [Eq. 20]	0.93	2.24	0.56	3.74
	i. Difference (%) (i.e., (h-a)/a)	629.7	2.5	-466.0	72.8
ALL2	a. Empirical ($\times 1000$)	-0.44	1.90	-0.57	0.9
	b. Approx.1 ($\times 1000$) [Appendix S2]	-0.45	1.87	-0.60	0.82

	c. Difference (%) (i.e., (b-a)/a)	3.4	-1.8	4.5	-8.5
	d. Approx.2 ($\times 1000$) [Eq. 19] “+”	1.36	1.92	1.21	4.49
	e. Difference (%) (i.e., (d-a)/a)	-413.3	0.8	-310.6	400.4
	f. Approx.3($\times 1000$) [Eq. 19] “-”	-0.16	1.07	-0.5	0.42
	g. Difference (%) (i.e., (f-a)/a)	-63.4	-43.6	-13.3	-53.4
	h. Approx.4 ($\times 1000$) [Eq. 20]	0.63	1.5	0.38	2.5
	i. Difference (%) (i.e., (h-a)/a)	-243.7	-21.2	-165.4	178.7
ABL	a. Empirical ($\times 1000$)	1.07	2.38	0.61	4.05
	b. Approx.1 ($\times 1000$) [Appendix S2]	1.02	2.45	0.61	4.08
	c. Difference (%) (i.e., (b-a)/a)	-4.7	3.0	1.5	0.7
	d. Approx.3 ($\times 1000$) [Eq. 20]	1.02	2.44	0.61	4.07
	e. Difference (%) (i.e., (d-a)/a)	-4.6	2.7	1.0	0.5
MBL	a. Empirical ($\times 1000$)	1.08	3.02	0.61	4.71
	b. Approx.1 ($\times 1000$) [Appendix S2]	1.14	2.84	0.66	4.64
	c. Difference (%) (i.e., (b-a)/a)	5.1	-5.9	9.0	-1.4
	d. Approx.3 ($\times 1000$) [Eq. 20]	1.13	2.71	0.68	4.52
	e. Difference (%) (i.e., (d-a)/a)	4.6	-10.1	11.5	-3.9

Table 3. Empirical standard error estimates and approximations in simulation experiments

Note. Lag0= θ_0 ; Lag1= θ_1 ; Lag2= θ_2 ; and Lag02= $\bar{\theta} = \theta_0 + \theta_1 + \theta_2$

MLL=Multiplicative Linear-Like error; ALL=Additive Linear-Like error; ABL=Additive Berkson-Like error; MBL=Multiplicative Berkson-Like error. See Figure 3 for distributions of MLL1–4, ALL1–2, ABL, and MBL.

Error	Quantities	Lag0	Lag1	Lag2	Lag02
MLL1	a. Empirical ($\times 1000$)	1.535	1.648	1.489	2.007
	b. Approx. ($\times 1000$) (Eq. 15)	1.491	1.611	1.449	1.993
	c. Difference (%) (i.e., (b-a)/a)	-2.8	-2.3	-2.7	-0.7
MLL2	a. Empirical ($\times 1000$)	1.478	1.626	1.447	1.813
	b. Approx. ($\times 1000$) (Eqs. 15 and 16)	1.473	1.617	1.441	1.783
	c. Difference (%) (i.e., (b-a)/a)	-0.3	-0.5	-0.4	-1.7
MLL3	a. Empirical ($\times 1000$)	0.797	0.869	0.778	1.016
	b. Approx. ($\times 1000$) (Eqs. 15 and 16)	0.768	0.843	0.753	0.999
	c. Difference (%) (i.e., (b-a)/a)	-3.7	-2.9	-3.2	-1.6
MLL4	a. Empirical ($\times 1000$)	0.774	0.859	0.76	0.942
	b. Approx. ($\times 1000$) (Eqs. 15 and 16)	0.761	0.844	0.749	0.917
	c. Difference (%) (i.e., (b-a)/a)	-1.7	-1.7	-1.4	-2.7
ALL1	a. Empirical ($\times 1000$)	1.041	1.247	1.033	1.037
	b. Approx. ($\times 1000$) (Eqs. 12 and 14)	1.119	1.332	1.109	1.006
	c. Difference (%) (i.e., (b-a)/a)	7.5	6.8	7.3	-3.0
ALL2	a. Empirical ($\times 1000$)	0.917	1.234	0.928	0.656
	b. Approx. ($\times 1000$) (Eqs. 12 and 14)	0.957	1.292	0.969	0.644

	c. Difference (%) (i.e., (b-a)/a)	4.4	4.7	4.4	-1.9
ABL	a. Empirical ($\times 1000$)	4.386	6.794	4.547	2.267
	b. Approx. ($\times 1000$) (Eqs. 8 and 10)	4.418	6.774	4.575	2.239
	c. Difference (%) (i.e., (b-a)/a)	0.7	-0.3	0.6	-1.2
MBL	a. Empirical ($\times 1000$)	5.568	8.828	5.777	2.657
	b. Approx. ($\times 1000$) (Eqs. 8 and 10)	5.773	9.074	5.956	2.673
	c. Difference (%) (i.e., (b-a)/a)	3.7	2.8	3.1	0.6




Figure 4. Power curves obtained from sample size calculation equations presented in the main text (dashed curves) and empirical statistical powers (points) from simulation experiments

Note. Each data point was obtained from 2,500 simulation samples. Different exposure measurement error scenarios were considered: MLL (Multiplicative Linear Like Error); ALL (Additive Linear Like Error); ABL (Additive Berkson Like Error); MBL (Multiplicative Berkson Like Error). For ALL1, ALL2, ABL, and MBL, lag0 and lag2 were not presented here because empirical statistical powers were extremely low.

5. Discussion

Understanding of the determinants of sample size and the accuracy and precision of estimators is helpful in designing robust and reproducible research and interpreting findings. Several key lessons are summarized here. First, if exposure levels vary spatiotemporally, the variance may be heteroskedastic. This needs to be appropriately considered in understanding sample size. Second, even without exposure validation data, investigators may calculate sample size for a desired power and the size of the estimators based on external information about exposure measurement error. Third, for distributed lag models, it may sometimes be difficult to consider the accuracy of the estimators of lag variables due to potentially severe residual confounding from covariate measurement errors. For example, theoretically, the absence of associations between certain lag times and Y may be attributed to residual confounding (e.g., ALL1 in Table 2). In identifying critical exposure time-windows, designing validation research is desirable. Fourth, in cases where validation data is unavailable, conducting numerical analyses to examine how sample size and the accuracy of estimators can vary across different bias scenarios would be beneficial for designing research and interpreting findings. The main limitation of this paper is that it focuses on case-crossover design. Nevertheless, as the development was based on an existing sample size calculation method for conditional logistic regression (Lachin 2008), developments for other matched case-control designs and survival analyses with Cox proportional hazard models merit future research.

Funding

This work was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2021R1A6A3A14039711), by University of Illinois Chicago School of Public Health Seed Funding Program, and by the National Institute of Environmental Research (NIER) funded by the Ministry of Environment (MOE) of the Republic of Korea (NIER-2021-03-03-007)

Disclosure Statement

There is no conflict of interest

References

- Bernardo, M. P., Lipsitz, S. R., Harrington, D. P., and Catalano, P. J. (2000), "Sample size calculations for failure time random variables in non-randomized studies," *Journal of the Royal Statistical Society Series D: The Statistician*, 49 (1), 31-40.
- Bhaskaran, K., Armstrong, B., Hajat, S., Haines, A., Wilkinson, P., and Smeeth, L. (2012), "Heat and risk of myocardial infarction: hourly level case-crossover analysis of MINAP database," *BMJ*, 345.
- Boe, L. A., Shaw, P. A., Midthune, D., Gustafson, P., Kipnis, V., Park, E., Sotres-Alvarez, D., Freedman, L., of the STRATOS Initiative, o. b. o. t. M. E., and Group, M. T. (2023), "Issues in Implementing Regression Calibration Analyses," *American Journal of Epidemiology*, 192 (8), 1406-1414.
- Carroll, R. J., Ruppert, D., Stefanski, L. A., and Crainiceanu, C. M. (2006), *Measurement error in nonlinear models: a modern perspective*: Chapman and Hall/CRC.
- Fewell, Z., Davey Smith, G., and Sterne, J. A. (2007), "The impact of residual and unmeasured confounding in epidemiologic studies: a simulation study," *American Journal of Epidemiology*, 166 (6), 646-655.
- Gasparini, A., Armstrong, B., and Kenward, M. G. (2010), "Distributed lag non-linear models," *Statistics in Medicine*, 29 (21), 2224-2234.
- Greenland, S., and Robins, J. M. (1985), "Confounding and misclassification," *American Journal of Epidemiology*, 122 (3), 495-506.
- Innes, G. K., Bhondoekhan, F., Lau, B., Gross, A. L., Ng, D. K., and Abraham, A. G. (2021), "The measurement error elephant in the room: challenges and solutions to measurement error in epidemiology," *Epidemiologic Reviews*, 43 (1), 94-105.
- Keogh, R. H., Shaw, P. A., Gustafson, P., Carroll, R. J., Deffner, V., Dodd, K. W., Küchenhoff, H., Tooze, J. A., Wallace, M. P., and Kipnis, V. (2020), "STRATOS guidance document on measurement error and misclassification of variables in observational epidemiology: part 1—basic theory and simple methods of adjustment," *Statistics in Medicine*, 39 (16), 2197-2231.
- Kim, H. (2024), "A Unification of Exchangeability and Continuous Exposure and Confounder Measurement Errors: Probabilistic Exchangeability," *arXiv preprint arXiv:2405.07910*.
- Kim, H., and Lee, J.-T. (2019), "On inferences about lag effects using lag models in air pollution time-series studies," *Environmental research*, 171, 134-144.
- Kim, H., Samet, J. M., and Bell, M. L. (2022), "Association between short-term exposure to air pollution and COVID-19 mortality: a population-based case-crossover study using individual-level mortality registry confirmed by medical examiners," *Environmental Health Perspectives*, 130 (11), 117006.
- Kim, H., Son, J.-Y., Junger, W., and Bell, M. L. (2024), "Exposure to particulate matter and ozone, locations of regulatory monitors, and sociodemographic disparities in the city of Rio de Janeiro: Based on local air pollution estimates generated from machine learning models," *Atmospheric Environment*, 322, 120374.
- Lachin, J. M. (2008), "Sample size evaluation for a multiply matched case-control study using the score test from a conditional logistic (discrete Cox PH) regression model," *Statistics in Medicine*, 27 (14), 2509-2523.

- Langholz, B., Thomas, D., Xiang, A., and Stram, D. (1999), "Latency analysis in epidemiologic studies of occupational exposures: application to the Colorado Plateau uranium miners cohort," *American Journal of Industrial Medicine*, 35 (3), 246-256.
- Lee, J.-T., and Schwartz, J. (1999), "Reanalysis of the effects of air pollution on daily mortality in Seoul, Korea: A case-crossover design," *Environmental Health Perspectives*, 107 (8), 633-636.
- Lu, Y., and Zeger, S. L. (2007), "On the equivalence of case-crossover and time series methods in environmental epidemiology," *Biostatistics*, 8 (2), 337-344.
- Maclure, M. (1991), "The case-crossover design: a method for studying transient effects on the risk of acute events," *American Journal of Epidemiology*, 133 (2), 144-153.
- Mittleman, M. A., and Mostofsky, E. (2014), "Exchangeability in the case-crossover design," *International Journal of Epidemiology*, 43 (5), 1645-1655.
- Mork, D., Kioumourtzoglou, M.-A., Weisskopf, M., Coull, B. A., and Wilson, A. (2024), "Heterogeneous distributed lag models to estimate personalized effects of maternal exposures to air pollution," *Journal of the American Statistical Association*, 119 (545), 14-26.
- Reff, A. P., Sharon; Eyth, Alison; Godfrey, Janice; Vukovich, Jeff; Mintz, Mintz (2023), "Bayesian Space-time Downscaling Fusion Model (Downscaler) - Derived Estimates of Air Quality for 2020," *Environmental Protection Agency*, EPA-454/R-23-004.
- Robinson, L. D., and Jewell, N. P. (1991), "Some surprising results about covariate adjustment in logistic regression models," *International Statistical Review/Revue Internationale de Statistique*, 227-240.
- Tong, S., Wang, X. Y., and Guo, Y. (2012), "Assessing the short-term effects of heatwaves on mortality and morbidity in Brisbane, Australia: comparison of case-crossover and time series analyses," *PloS one*, 7 (5), e37500.
- Wade, T. J., Lin, C. J., Jagai, J. S., and Hilborn, E. D. (2014), "Flooding and emergency room visits for gastrointestinal illness in Massachusetts: a case-crossover study," *PloS one*, 9 (10), e110474.

Online Supplementary Materials

Identifying Sample Size and Accuracy and Precision of the Estimators in Case-Crossover Designs with Distributed Lags of Heteroskedastic Spatiotemporally Varying Continuous Exposures Measured with Simple or Complex Error

Appendix S1. Heteroscedastic variance calculations.

Appendix S2. Approximation for the estimators in the presence of additive linear (-like) error.

Appendix S3. Residual confounding in distributed lag models when the error is differential.

Appendix S4. Adjustment for differential errors.

Appendix S5. Standard error decomposition in the presence of multiplicative error.

Appendix S6. Heteroscedastic exposure variance approximation for multiplicative error.

Appendix S7. Approximation for the estimators without validation data in hand.

Appendix S8. Approximation for the estimator of a second-order polynomial.

Appendix S9. Simulation experiments.

Figure S1. Residual plots including partial autocorrelation plots for linear regressions in Table 1 of the main manuscript

Appendix S1. Heteroscedastic variance calculations.

If health data are not available before the initiation of a study (e.g., investigators need to recruit patients, but their residential history is unknown before recruitment), investigators would not be able to exactly calculate σ_h^2 . Instead, they may be able to have external information about a source population. They could approximately calculate σ_h^2 with the information about k -th sub-source populations.

Suppose that investigators have exposure data for x at spatial grid cells or census boundaries (k -th sub-source populations) and exact case-self-control locations (e.g., street-level address) are unknown. $\bar{\sigma}^2$ can be approximated to the expected case number-weighted σ^2 as

$$\bar{\sigma}^2 \cong \frac{\sum_{k=1}^K E[Cases_k] \sigma_k^2}{\sum_{k=1}^K E[Cases_k]}$$

where $E[Cases_k]$ is the expected number of cases in k -th sub-source population and K is the total number of sub-source populations. Investigators may use external data such as population statistics or summary of electronic health records to identify the expected number of cases. One way to estimate $\bar{\sigma}^2$ is

$$\bar{\sigma}^2 \cong \frac{\sum_{fl=1}^{N_{fl}} Pop_{fl} \sum_{s=1}^S R_{fl,s} \sigma_{fl,s}^2}{\sum_{fl=1}^{N_{fl}} Pop_{fl} \sum_{s=1}^S R_{fl,s}}$$

where Pop_{fl} denotes a sub-source population count for a fixed location fl . (e.g., the number of residents in a census tract fl). N_{fl} is the total number of the sub-source populations for a study (e.g., the number of census tracts in a city). $\sigma_{fl,s}^2$ is the variance for fl and matched stratum s (i.e., matching each case to their self-control(s)). S is the total number of the strata. For example, if investigators use time-stratification using the calendar year, month, and day of the week, s

indicates each time-stratified set (e.g., January 2020–Monday; January 2020–Tuesday; ..., December 2020–Sunday). $R_{f,l,s}$ is the baseline hazard (rate) for f and s .

Appendix S2. Approximation for the estimators in the presence of additive linear (-like) error.

Suppose the correct linear outcome model, $E[Y_{i,t}] = \vartheta + \sum_{l=0}^L \theta_l x_{i,t-l} + \boldsymbol{\vartheta} \mathbf{z}$ and $x_g = x_i$. I use this linear model instead of conditional logistic regression, because for small θ_l or small exposure measurement error, the consequence from linear regression may be applicable.

In practice, investigators may use $x_{g,t}^{ep}$ in place of $x_{i,t}$. Suppose a linear (-like) (L(L)) error model, $x_{g,t}^{ep} = \gamma_0 + \eta_0 x_{g,t} + u_{g,t}^{L(L)}$. (See Section 3.3). To consider the impact of this error model on effect estimators, it is convenient to consider following re-written error models based on regression calibration,

$$\begin{aligned} x_{g,t} &= \gamma_{0,0}^* + \sum_{l=0}^L \eta_{0,l}^* x_{g,t-l}^{ep} + \boldsymbol{\eta}_{0,z} \mathbf{z} + u_{g,t}^* \\ x_{g,t-1} &= \gamma_{1,0}^* + \sum_{l=0}^L \eta_{1,l}^* x_{g,t-l}^{ep} + \boldsymbol{\eta}_{1,z} \mathbf{z} + u_{g,t-1}^* \\ &\dots \\ x_{g,t-L} &= \gamma_{L,0}^* + \sum_{l=0}^L \eta_{L,l}^* x_{g,t-l}^{ep} + \boldsymbol{\eta}_{L,z} \mathbf{z} + u_{g,t-L}^* \end{aligned}$$

For $x_{g,t-l}^{ep}$, the following parameters are important:

$$\eta_{0,l}^* = \frac{\text{Cov}(x_{g,t}, x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \cong \frac{\eta_0 \text{Cov}(x_{g,t}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \text{Cov}(x_{g,t}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \eta_0 \text{Cov}(x_{g,t-l}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

$$\eta_{1,l}^* = \frac{\text{Cov}(x_{g,t-1}, x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \cong \frac{\eta_0 \text{Cov}(x_{g,t-1}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \text{Cov}(x_{g,t-1}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \eta_0 \text{Cov}(x_{g,t-l}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

...

$$\eta_{L,l}^* = \frac{\text{Cov}(x_{g,t-L}, x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \cong \frac{\eta_0 \text{Cov}(x_{g,t-L}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \text{Cov}(x_{g,t-L}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \eta_0 \text{Cov}(x_{g,t-l}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}.$$

Thus, generally,

$$\eta_{j,l}^* = \frac{\eta_0 \text{Cov}(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \text{Cov}(x_{g,t-j}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \eta_0 \text{Cov}(x_{g,t-l}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

If the error is independent and identically distributed (independent of $x_{g,t}$) (i.e., non-differential

error), then $\text{Cov}(x_{g,t-j}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}) = 0$ so that

$$\eta_{j,l}^* = \frac{\eta_0 \text{Cov}(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^* | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

and

$$\eta_{l,l}^* = \frac{\eta_0 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^* | \mathbf{x}_g^{ep-l}, \mathbf{z})} \equiv \lambda_l$$

where λ_l is a bias factor.

The correct linear outcome model can be re-expressed as

$$E[Y_{i,t}] = \vartheta + \sum_{l=0}^L \theta_l (\gamma_{l,0}^* + \sum_{j=0}^L \eta_{l,j}^* x_{g,t-j}^{ep} + \boldsymbol{\eta}_{l,z} \mathbf{z} + u_{g,t-l}^*) + \boldsymbol{\vartheta} \mathbf{z}$$

But, investigators use

$$E[Y_{i,t-l}] = \vartheta^* + \sum_{l=0}^L \theta_l^{L(L)} x_{g,t-l}^{ep} + \boldsymbol{\vartheta}^* \mathbf{z}$$

If $\eta_{l,j}^* = 0$ for $j \neq l$, which occurs when $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) = 0$ (i.e., no autocorrelation conditional on $\mathbf{x}_g^{ep-l}, \mathbf{z}$)

$$\theta_l^{L(L)} \cong \theta_l \eta_{l,l}^* + \sum_{k=0}^L \theta_k \rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$$

If u_{g-k}^* is independent and identically distributed (and independent of $x_{g,t-l}^{ep}$) given \mathbf{x}_g^{ep-l} and \mathbf{z} , which may occur when the error is non-differential, $\rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}} = 0$. Then,

$$\theta_l^L \cong \theta_l \eta_{l,l}^* = \theta_l \lambda_l$$

If $\eta_{l,j}^* \neq 0$ (e.g., non-zero correlations between lagged variables of $x_{g,t}$) and/or u_{g-k}^* is not independent of $x_{g,t-l}^{ep}$ even conditional on \mathbf{x}_g^{ep-l} and \mathbf{z} ($\rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}} \neq 0$), then the

estimator would vary. Recall the following components in the re-written correct outcome model

$$\sum_{j=0}^L \theta_0 \eta_{0,j}^* x_{g,t-j}^{ep} = \theta_0 \eta_{0,0}^* x_{g,t}^{ep} + \theta_0 \eta_{0,1}^* x_{g,t-1}^{ep} + \dots + \theta_0 \eta_{0,L}^* x_{g,t-L}^{ep} \text{ (for } l = 0 \text{)}$$

$$\sum_{j=0}^L \theta_1 \eta_{1,j}^* x_{g,t-j}^{ep} = \theta_1 \eta_{1,0}^* x_{g,t}^{ep} + \theta_1 \eta_{1,1}^* x_{g,t-1}^{ep} + \dots + \theta_1 \eta_{1,L}^* x_{g,t-L}^{ep} \text{ (for } l = 1 \text{)}$$

...

$$\sum_{j=0}^L \theta_L \eta_{L,j}^* x_{g,t-j}^{ep} = \theta_L \eta_{L,0}^* x_{g,t}^{ep} + \theta_L \eta_{L,1}^* x_{g,t-1}^{ep} + \dots + \theta_L \eta_{L,L}^* x_{g,t-L}^{ep} \text{ (for } l = L \text{)}$$

so that $\theta_l^{L(L)}$ can be approximated as

$$\theta_l^{L(L)} \cong \sum_{j=0}^L \theta_j \eta_{j,l}^* + \sum_{k=0}^L \theta_k \rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$$

This completes the development of an approximation equation for $\theta_l^{L(L)}$. Below is an investigation of the behavior of $\theta_l^{L(L)}$.

The approximation equation above shows that the following two merit investigations:

- 1) How $\theta_i^{L(L)}$ for $j \neq l$ depends on $\eta_{j,l}^*$ and $\rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$
- 2) The covariance component in the numerator of $\eta_{j,l}^*$, which is

$$Cov\left(x_{g,t-j}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}\right)$$

For 1), when $x_{g,t}$ is autocorrelated (e.g., $x_{g,t} = \zeta_1 x_{g,t-1} + \zeta_2 x_{g,t-2} + \dots$) and $\mathbf{x}_g^{ep-l}, \mathbf{z}$ are not sufficient to explain this autocorrelation, then, for some j ($j \neq l$), $Cov\left(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}\right) \neq 0$ so that $\eta_{l,j}^* \neq 0$. Note that by the definition, if $x_{g,t-l}$ has an effect on Y and is correlated with $x_{g,t-j}$, this would be seen as a confounder. Residual confounding would likely occur if the error in $x_{g,t}^{ep}$ is not small because if this is not small, $\mathbf{x}_g^{ep-l}, \mathbf{z}$ would be unlikely to be sufficient to make $Cov\left(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}\right) = 0$ so that $\eta_{l,j}^* \neq 0$. $\rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$ may be negligible, depending on the structure of $x_{g,t}, x_{g,t}^{ep}, \mathbf{z}$.

For 2), $Cov\left(x_{g,t-j}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}\right)$ may be zero if $u_{g,t}^L$ is completely random. Otherwise, this may not be zero.

Appendix S3. Residual confounding in distributed lag models when the error is differential.

Recall

$$\theta_l^{L(L)} \cong \sum_{j=0}^L \theta_j \eta_{j,l}^* + \sum_{k=0}^L \theta_k \rho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$$

and

$$Cov(x_{g,t-j}, u_{g,t-l}^{L(L)} | x_g^{ep-l}, \mathbf{z}) \text{ in } \eta_{j,l}^*$$

in Appendix S2. These may be applicable for non-linear models such as logistic regression when the effect size is small, or the measurement error is small.

If the error component is $u_{g,t-l}^{LL}$, it is convenient to distinguish between

$$Cov(x_{g,t-l}, u_{g,t-l}^{LL} | x_g^{ep-l}, \mathbf{z}) \text{ (i.e., } j = l) \text{ and } Cov(x_{g,t-j}, u_{g,t-l}^{LL} | x_g^{ep-l}, \mathbf{z}) \text{ for } j \neq l.$$

Let me re-investigate 2) in Appendix S2 to examine the behavior of $\theta_l^{L(L)}$ in more detail.

2-1) If $Cov(x_{g,t-l}, u_{g,t-l}^{LL} | x_g^{ep-l}, \mathbf{z}) \neq 0$, then $\eta_{l,l}^* \neq \lambda$.

This is a consequence when adjustment for x_g^{ep-l}, \mathbf{z} is not adequate to eliminate the correlation between $x_{g,t-l}$ and $u_{g,t-l}^{LL}$. This means that the exposure measurement error would be still differential even after conditioning on x_g^{ep-l}, \mathbf{z} because the correlation between $u_{g,t-l}^{LL}$ and Y would be non-zero through a non-zero correlation between $x_{g,t-l}$ and $u_{g,t-l}^{LL}$ given x_g^{ep-l}, \mathbf{z} if X has an effect on Y . Recall that the definition of differential exposure measurement error is that the error is not independent of Y . To understand the behavior of $\theta_l^{L(L)}$, suppose $\eta_0 > 0$, which would be realistic in epidemiological investigations because we would otherwise not use x^{ep} at all. Then,

$$\begin{aligned}
\eta_{l,l}^* &= \frac{\eta_0 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \text{Cov}(x_{g,t-l}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \\
&= \frac{\eta_0 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \frac{\text{Cov}(x_{g,t-l}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z}) V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \\
&= \frac{\eta_0 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \varrho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}} V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \\
&= \frac{(\eta_0 + \varrho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}}) V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \\
&= \frac{(\eta_0 + \varrho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}}) V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \eta_0 \text{Cov}(x_{g,t-l}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \\
&= \frac{(\eta_0 + \varrho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}})}{\eta_0 + \frac{V(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} + \frac{V(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}} \\
&= \frac{1}{\eta_0 + \frac{V(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} (\eta_0 + \varrho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}})}
\end{aligned}$$

If the error is small so that $\frac{v(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{v(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})} \approx 0$, then $\frac{v(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{v(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) (\eta_0 + \varrho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}})} \approx 0$,

then finally,

$$\eta_{l,l}^* \approx \frac{1}{\eta_0}$$

Or, if the error is not small, let $a = \frac{v(u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{v(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) (\eta_0 + \rho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}})}$, then

$$\eta_{l,l}^* \approx \frac{1}{\eta_0 + a}$$

If the error is large so that $a \geq 1$, then $0 < \eta_{l,l}^* < 1$.

If $\eta_0 \geq 1$, then $0 < \eta_{l,l}^* \leq 1$, which is similar to the consequence by λ_l for non-differential linear error. To compare this with λ_l ,

$$\lambda_l = \frac{\eta_0 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})} = \frac{1}{\eta_0 + \frac{V(u_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) (\eta_0)}}$$

If $x_{g,t-l}$ is autocorrelated, then $x_{g,t-l}$ would also be correlated with \mathbf{x}_g^{ep-l} unless the error is not very large. This implies that $\rho_{u_{g,t-l}^{LL}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}}$ may not be high because \mathbf{x}_g^{ep-l} is also correlated with u_{t-l} so that the correlation between $u_{g,t-l}^{LL}$ and $x_{g,t-l}$ may be adjusted for to some degree by conditioning on $\mathbf{x}_g^{ep-l}, \mathbf{z}$. Thus, it is reasonable to conclude that $\eta_{l,l}^*$ may act like λ_l of non-differential linear error in distributed lag models if the error is not large.

2-2) Regarding $Cov(x_{g,t-j}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z}) \neq 0$ for $j \neq l$.

This term subsumes the situation that the exposure measurement error is correlated with $x_{g,t-j}$ after adjustment for $\mathbf{x}_g^{ep-l}, \mathbf{z}$ was performed. There are two things to consider:

Q1. Can $x_{g,t-j}$ is correlated with $u_{g,t-l}^{LL}$?

A1. If $x_{g,t}$ is autocorrelated and $u_{g,t-l}^{LL}$ is autocorrelated (e.g., because both have patterns over space, time, or both). Recall that $u_{g,t-l}^{LL}$ may include $\sum_{j=1}^J \eta_j x_{g,t-j}$ (See the main text) where η_j is non-zero. So, it would be difficult to say that these two are not correlated.

Q2. If $x_{g,t-j}$ is correlated with $u_{g,t-l}^{LL}$, can this correlation be adjusted for using $x_{g,t-j}^{ep}$?

Adjusting for $x_{g,t-j}^{ep}$ implies that adjusting for $x_{g,t-j}$ and $u_{g,t-j}^{LL}$ simultaneously using one variable, but not separately (Recall a linear-like error model: $x_{g,t-j}^{ep} = \gamma_0 + \gamma_1 x_{g,t-j} + u_{g,t-j}^{LL}$). If the spatiotemporal pattern of $x_{g,t-j}$ and $u_{g,t-j}^{LL}$ would be similar, they would be highly correlated, then the performance of adjustment for $x_{g,t-j}^{ep}$ in terms of adjustment for $x_{g,t-j}$ would be substantial. Furthermore, the performance of adjustment for $x_{g,t-j}^{ep}$ in terms of adjustment for $u_{g,t-l}^{LL}$ would also be substantial. If the pattern of $x_{g,t-j}$ and $u_{g,t-j}^{LL}$ would not be similar, then these two would not be highly correlated, then the performance of adjustment for $x_{g,t-j}^{ep}$ would not be substantial. The extreme case is a non-differential error model (e.g., $x_{g,t-j}^{ep} = \gamma_0 + \gamma_1 x_{g,t-j} + u_{g,t-j}^L$ where $u_{g,t-j}^L$ is completely random).

These suggest that if the exposure measurement error is differential in a way that $u_{g,t-j}^{LL}$ is highly correlated with $x_{g,t-j}$, it would not be a stretch to say that $Cov(x_{g,t-j}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})$ may be negligible due to conditioning on $\mathbf{x}_g^{ep-l}, \mathbf{z}$.

If $Cov(x_{g,t-j}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z})$ for $j \neq l$ is negligible (i.e., ≈ 0) due to the high correlation between $x_{g,t-j}$ and $u_{g,t-l}^{LL}$, the term, $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})$ in $\eta_{j,l}^*$ may also be negligible due to conditioning on $\mathbf{x}_g^{ep-l}, \mathbf{z}$.

Recall

$$\eta_{j,l}^* = \frac{\eta_0 Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + Cov(x_{g,t-j}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{\eta_0^2 V(x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + \eta_0 Cov(x_{g,t-l}, u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z}) + V(u_{g,t-l}^{L(L)} | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

in Appendix S2. If $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) \approx 0$ and $Cov(x_{g,t-j}, u_{g,t-l}^{LL} | \mathbf{x}_g^{ep-l}, \mathbf{z}) \approx 0$ for $j \neq l$, then,

$$\eta_{j,l}^* \approx 0$$

Recall,

$$\theta_l^{L(L)} \cong \sum_{j=0}^L \theta_j \eta_{j,l}^* + \sum_{k=0}^L \theta_k \varrho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$$

and $\eta_{j,l}^* \approx 0$ for $j \neq l$ implies

$$\theta_l^{L(L)} \approx \theta_l \eta_{l,l}^* + \sum_{k=0}^L \theta_k \varrho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$$

and $\varrho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$ may also become negligible if $\mathbf{x}_{g,t}^{ep-l}$ is correlated with $x_{g,t-l}^{ep}$, then,

$$\theta_l^{L(L)} \approx \theta_l \eta_{l,l}^*$$

In 2-1), it was revealed that $\eta_{l,l}^*$ may act similar to λ_l .

In conclusion, if the additive error is autocorrelated such as additive LL error, the risk of the residual confounding due to the use of \mathbf{x}_g^{ep-l} may be smaller than what the risk would have been if the error was non-differential additive linear error.

Appendix S4. Adjustment for differential errors.

Before discussing adjustment for differential errors, recall the approximation proposed in Appendix S2,

$$\theta_l^{L(L)} \cong \sum_{j=0}^L \theta_j \eta_{j,l}^* + \sum_{k=0}^L \theta_k \varrho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$$

this may be applicable for non-linear models such as logistic regression when the effect size is small, or the measurement error is small. This approximation may be applicable for many error models in the form of $x_{g,t}^{ep} = g(x_{g,t}, v, u^{(A/M)L(L)})$ because this approximation does not rely on error models, but on re-written error models using regression calibration (Appendix S2), where $u^{L(L)}$ is non-differential error term, (including the autoregressive error model and multiplicative errors in the main manuscript) and g is a transformation function, v is an additional component describing the exposure measurement error (e.g., $x_{g,t-j}$ or $\delta(\mathbf{Q}^e)$). Simulation analyses in the Section 4 (Simulation Experiments) demonstrate that this approximation equation performs well.

Regarding adjustment for differential errors, v could be subsumed into \mathbf{z} . As shown above and discussed in Appendix S2 and S3, the magnitude of the components in the approximation varies by $x_{g,t}^{ep-l}, \mathbf{z}$. Thus, if \mathbf{z} includes some variables that are related to the error, the contribution of the error to $\eta_{j,l}^*$ and $\varrho_{u_{g,t-k}^*, x_{g,t-l}^{ep} | x_{g,t}^{ep-l}, \mathbf{z}}$ may decrease. If all components that make error differential are controlled for, the remaining conditional error would become non-differential, which is referred to as conditional non-differential error.

Appendix S5. Standard error decomposition in the presence of multiplicative error.

Consider the variance decomposition,

$$\begin{aligned}
 & SE\left(\hat{\theta}_i^{ML(L)}|\mathbf{z}, \mathbf{x}_g^{ep-l}\right)^2 \\
 &= E\left[SE\left(\hat{\theta}_i^{ML(L)}|\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)}), \mathbf{x}_g^{ep-l}, \mathbf{z}\right)^2\right] \\
 &+ V\left(E\left[\hat{\theta}_i^{ML(L)}|\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)}), \mathbf{x}_g^{ep-l}, \mathbf{z}\right]\right) \\
 &E\left[SE\left(\hat{\theta}_i^{ML(L)}|\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)}), \mathbf{x}_g^{ep-l}, \mathbf{z}\right)^2\right] \cong \left(\sum_{h=1}^n V_{0h}((x_{g,t-l})^{\gamma_{m1}})\left(1 - R_{x_{g,t-l}^{ep}|\mathbf{x}_g^{ep-l}, \mathbf{z}}^2\right)\right)^{-1} \\
 &V\left(E\left[\hat{\theta}_i^{ML(L)}|\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)}), \mathbf{x}_g^{ep-l}, \mathbf{z}\right]\right) \\
 &\cong \theta_i^{ML(L)2} \left(\sum_{h=1}^n V_{0h}\left(\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})\right)\left(1 - R_{x_{g,t-l}^{ep}|\mathbf{x}_g^{ep-l}, \mathbf{z}}^2\right)\right)^{-1}
 \end{aligned}$$

and,

$$\begin{aligned}
 n\bar{\sigma}_{x_{g,t-l}^{ML(L)}}^2 &= \sum_{h=1}^n V_{0h}((x_{g,t-l})^{\gamma_{m1}}) \\
 n\bar{\sigma}_{\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})}^2 &= \sum_{h=1}^n V_{0h}\left(\exp(\gamma_{m0})\exp(u_{g,t}^{L(L)})\right)
 \end{aligned}$$

Appendix S6. Heteroscedastic exposure variance approximation for multiplicative error.

Using the Delta method,

$$V(\gamma_{m1} \log(x_{g,t-l})) \approx \frac{\gamma_{m1}^2 V(x_{g,t-l})}{E[x_{g,t-l}]^2}$$

$$V((x_{g,t-l})^{\gamma_{m1}}) \approx \gamma_{m1}^2 E[x_{g,t-l}]^{2(\gamma_{m1}-1)} V(x_{g,t-l})$$

Recall $R_{\log(x_{g,t-l}^{ep}), \log(x_{g,t-l})}^2 = \frac{V(\gamma_{m0} + \gamma_{m1} \log(x_{g,t-l}))}{V(\log(x_{g,t-l}^{ep}))} = \frac{V(\gamma_{m1} \log(x_{g,t-l}))}{V(\log(x_{g,t-l}^{ep}))}$. So,

$$\begin{aligned} V((x_{g,t-l})^{\gamma_{m1}}) &\approx E[x_{g,t-l}]^{2\gamma_{m1}} V(\gamma_{m1} \log(x_{g,t-l})) \\ &= E[x_{g,t-l}]^{2\gamma_{m1}} R_{\log(x_{g,t-l}^{ep}), \log(x_{g,t-l})}^2 V(\log(x_{g,t-l}^{ep})) \end{aligned}$$

Since $\bar{\sigma}_{x_{g,t-l}}^2 = \frac{\sum_{h=1}^n \sigma_h^2(x_{g,t-l}^{ML(L)})}{n}$, $\bar{\sigma}_{\log(x_{g,t-l}^{ep})}^2 = \frac{\sum_{h=1}^n \sigma_h^2(\log(x_{g,t-l}^{ep}))}{n}$,

$\sigma_h^2(x_{g,t-l}^{ML(L)}) \approx E[x_{h,g,l}]^{2\gamma_{m1}} R_{\log(x_{g,t-l}^{ep}), \log(x_{g,t-l})}^2 \sigma_h^2(\log(x_{g,t-l}^{ep}))$, and

$\frac{\sum_{h=1}^n E[x_{h,g,l}]^{2\gamma_{m1}}}{n} \approx E[x_{g,t-l}]^{2\gamma_{m1}}$ for $\gamma_{m1} > 0$,

$$\bar{\sigma}_{x_{g,t-l}}^2 \approx \bar{\sigma}_{\log(x_{g,t-l}^{ep})}^2 E[x_{g,t-l}]^{2\gamma_{m1}} R_{\log(x_{g,t-l}^{ep}), \log(x_{g,t-l})}^2$$

Appendix S7. Approximation for the estimators without validation data in hand.

Recall

$$\begin{aligned}\eta_{j,l}^* &= \frac{\text{Cov}\left(x_{g,t-j}, x_{g,t-l}^{ep} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right)}{V\left(X_{g,t-l}^{ep} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right)} \\ &= \frac{\eta_0 \text{Cov}\left(x_{g,t-j}, x_{g,t-l} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right) + \text{Cov}\left(x_{g,t-j}, u_{g,t-l}^{L(L)} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right)}{\eta_0^2 V\left(x_{g,t-l} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right) + \eta_0 \text{Cov}\left(x_{g,t-l}, u_{g,t-l}^{L(L)} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right) + V\left(u_{g,t-l}^{L(L)} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right)}\end{aligned}$$

in Appendices S2 and S3. This may be applicable for non-linear models such as logistic regression when the effect size is small, or the measurement error is small.

For $j \neq l$, without knowing $x_{g,t-j}$, we may use $x_{g,t-j}^{ep}$ to approximate this quantity. For illustration, consider $V(x_{g,t-j} \mid \mathbf{x}_g^{ep-l}, \mathbf{z})$. Using the notation in Appendix S2, recall

$$x_{g,t-j} = \gamma_{j,0}^* + \sum_{l=0}^L \eta_{j,l}^* x_{g,t-l}^{ep} + \boldsymbol{\eta}_{j,z} \mathbf{z} + u_{g,t-j}^*$$

Assume $V(x_{g,t-j} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}) \approx V(x_{g,t-j} \mid \mathbf{x}_g^{ep}, \mathbf{z}) = V(u_{g,t-j}^*)$ so that

$\text{Cov}\left(x_{g,t-j}, x_{g,t-l} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right) \approx V(u_{g,t-j}^*)$, which will be discussed below. Then,

$$\eta_{j,l}^* \approx \frac{\eta_0 V(u_{g,t-j}^*) + \text{Cov}\left(x_{g,t-j}, u_{g,t-l}^{L(L)} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right)}{V\left(X_{g,t-l}^{ep} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}\right)}$$

Note

$$R_{x_{g,t-l}^{ep}, x_{g,t-l} \mid \mathbf{x}_g^{ep-l}, \mathbf{z}}^2 = \frac{V(\gamma_0 + \eta_0 x_{g,t-l} \mid \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} \mid \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

In the main manuscript, the notation γ_1 is used in place of η_0 while η_0 is used in Appendix S2 because this notation is used to highlight LL error models introduced in Section 3.3.

This can be further extended to

$$\begin{aligned}
R^2_{x_{g,t-l}^{ep}, x_{g,t-l} | x_g^{ep-l}, \mathbf{z}} &\approx \frac{V(\gamma_0 + \eta_0 x_{g,t-l} | x_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})} = \frac{\eta_0^2 V(x_{g,t-l} | x_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})} \\
&\approx \frac{\eta_0^2 (V(u_{g,t}^*) + V(\eta_{i,l}^* x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z}))}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})} = \frac{\eta_0^2 (V(u_{g,t}^*) + \eta_{i,l}^{*2} V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z}))}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})} \\
&= \frac{\eta_0^2 V(u_{g,t}^*)}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})} + \eta_0^2 \eta_{i,l}^{*2}
\end{aligned}$$

Now compare

$$\begin{aligned}
R^2_{x_{g,t-l}^{ep}, x_{g,t-l} | x_g^{ep-l}, \mathbf{z}} &\approx \frac{\eta_0^2 V(u_{g,t}^*)}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})} + \eta_0^2 \eta_{i,l}^{*2} \\
\eta_{j,l}^* &\approx \frac{\eta_0 V(u_{g,t-j}^*) + \text{Cov}(x_{g,t-j}, u_{g,t-l}^{L(L)} | x_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})}
\end{aligned}$$

Appendices S2–S4 discusses when $\text{Cov}(x_{g,t-j}, u_{g,t-l}^{L(L)} | x_g^{ep-l}, \mathbf{z}) \approx 0$. So, it is possible

$$\eta_{j,l}^* \approx \frac{\eta_0 V(u_{g,t-j}^*)}{V(x_{g,t-l}^{ep} | x_g^{ep-l}, \mathbf{z})}$$

If $R^2_{x_{g,t}, x_g} \approx R^2_{x_{g,t-l}, x_{g,t-l} | x_g^{ep-l}, \mathbf{z}}$, which is likely when \mathbf{z} is not a strong determinant of x_g ,

$$R^2_{x_{g,t}, x_g} \approx \eta_0 \eta_{j,l}^* + \eta_0^2 \eta_{i,l}^{*2}$$

For $\eta_{i,l}^*$, see Appendix S2 or recall 2-1) in Appendix S3. If $\eta_{i,l}^* \approx \lambda \approx \lambda_l$

where $\lambda = \frac{\eta_0 V(x_{g,t} | \mathbf{x}_g^{ep-0}, \mathbf{z})}{\eta_0^2 V(x_{g,t} | \mathbf{x}_g^{ep-0}, \mathbf{z}) + V(u_{g,t} | \mathbf{x}_g^{ep-0}, \mathbf{z})}$,

$$R_{x_{g,t}, x_g}^{2ep} - \eta_0^2 \lambda^2 \approx \eta_0 \eta_{j,l}^*$$

and,

$$\eta_0 \lambda = \frac{V(\gamma_0 + \eta_0 x_{g,t} | \mathbf{x}_g^{ep-0}, \mathbf{z})}{\eta_0^2 V(x_{g,t} | \mathbf{x}_g^{ep-0}, \mathbf{z}) + V(u_{g,t} | \mathbf{x}_g^{ep-0}, \mathbf{z})} = R_{x_{g,t}, x_{g,t} | \mathbf{x}_g^{ep0}, \mathbf{z}}^2$$

Recall

$$\theta_l^{L(L)} \cong \sum_{j=0}^L \theta_j \eta_{j,l}^* + \sum_{k=0}^L \theta_k \varrho_{u_{g,t-k}^* | x_{g,t-l}^{ep} | \mathbf{x}_{g,t}^{ep-l}, \mathbf{z}}$$

Assuming $\varrho_{u_{g,t-k}^* | x_{g,t-l}^{ep} | \mathbf{x}_{g,t}^{ep-l}, \mathbf{z}} = 0$,

$$\theta_l^L \cong \sum_{j=0}^L \theta_j \eta_{j,l}^*$$

Finally,

$$\theta_l^{L(L)} \approx R_{x_{g,t}, x_g}^{2ep} \left(\frac{\theta_l + \left(1 - R_{x_{g,t}, x_g}^{2ep}\right) \sum_{j \neq l}^L \theta_j}{\eta_0} \right)$$

This approximation would perform reasonably if the assumption $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) \approx V(u_{g,t-j}^*)$ is reasonable. Importantly, $V(u_{g,t-j}^*)$ is only positive so that that approximation would perform reasonably if there were no severe negative residual confounding. In practice,

$Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})$ can be negative so that

$$\eta_{j,l}^* \approx \frac{\eta_0 Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})}{V(x_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})} < 0$$

which is not considered in the approximation.

To consider that, why and when could $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})$ be negative so that $\eta_{j,l}^*$ be negative?

To answer this question, I provide an analogy here. It is known that in time-series analyses, the temporal autocorrelation of the residual of $\bar{Y}_t = \sum Y_{i,t}$ conditional on variables including variables that adjust for seasonality and long-term time-trend as unmeasured time-varying confounders can diminish and eventually randomly fluctuate around zero over time lag, which implies $Cov(\bar{Y}_t, \bar{Y}_{t-j} | \mathbf{z}) \approx 0$ for $j > 0$. This is a desirable consequence of the adequate adjustment because the non-negligible crude autocorrelation, $Cov(\bar{Y}_t, \bar{Y}_{t-j}) \neq 0$ may be a sign of the existence of unmeasured time-varying confounding. It is also known that redundant adjustment for seasonality and long-term time-trend can make $Cov(\bar{Y}_t, \bar{Y}_{t-j} | \mathbf{z}) < 0$. To apply analogy to our context, replace $Cov(\bar{Y}_t, \bar{Y}_{t-j} | \mathbf{z})$ with $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z})$. Recall that \mathbf{x}_g^{ep-l} may have time-trend (because \mathbf{x}_g^{-l} may have time-trend), case-self-control matching scheme that can be seen as matching strata variables in \mathbf{z} is inherently related to time-trend (See Lu and Zeger 2007 for detail) and other covariates in \mathbf{z} that may also have time-trend. Thus, it would not be surprising that $Cov(x_{g,t-j}, x_{g,t-l} | \mathbf{x}_g^{ep-l}, \mathbf{z}) < 0$ can occur for some j s and its magnitude may be non-negligible. Finally, it is reasonable to consider not only

$$\eta_{j,l}^* \approx \frac{\eta_0 V(u_{g,t-j}^*)}{V(X_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

but also

$$\eta_{j,l}^* \approx \frac{-\eta_0 V(u_{g,t-j}^*)}{V(X_{g,t-l}^{ep} | \mathbf{x}_g^{ep-l}, \mathbf{z})}$$

so that, by adopting \pm ,

$$\theta_l^{L(L)} \approx R_{x_{g,t},x_g}^{2,ep} \left(\frac{\theta_l \pm \left(1 - R_{x_{g,t},x_g}^{2,ep}\right) \sum_{j \neq l}^L \theta_j}{\eta_0} \right)$$

This approximation may perform well if residual confounding is not severe. This may perform in many types of errors because $u_{g,t-j}^*$ is designed from regression calibration (Appendix S2). For multiplicative L(L) errors, this approximation may be usable if treating this error as additive is reasonable and $R_{x_{g,t},x_g}^{2,ep}$ and γ_1^{crude} is known.

Note. This approximation relies on the assumption $R_{x_{g,t},x_g}^{2,ep} \approx R_{x_{g,t-l},x_{g,t-l}|x_g^{ep-l},z}^{2,ep}$. This assumption may not necessarily hold depending on, particularly \mathbf{z} , which includes how to match case and self-controls. If \mathbf{z} may strongly predict x_g (here “strong” is relative compared to the magnitude of the exposure measurement error) it may be likely $R_{x_{g,t},x_g}^{2,ep} \gg R_{x_{g,t-l},x_{g,t-l}|x_g^{ep-l},z}^{2,ep}$ because the correlation between the residual of $x_{g,t-l}^{ep}$ and the residual of $x_{g,t-l}$ would be mainly determined by the exposure measurement error while the crude correlation between $x_{g,t-l}^{ep}$ and $x_{g,t-l}$ is determined by not just the error but also \mathbf{z} .

Appendix S8. Approximation for the estimator of a second-order polynomial.

Let θ'_i be the coefficient for $(x_{t-l})^2$. If the residual of $(x_{t-l}^{ep})^2$ conditional on $\mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z}$ and the residual of x_{t-l}^{ep} conditional on $\mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2$ approximately follow a normal distribution and residual confounding due to the use of \mathbf{x}^{ep-l} is negligible, $\theta_i'^L$ in linear regression,

$$\begin{aligned} \theta_i'^L &= \frac{\text{Cov}(Y, (x_{t-l}^{ep})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})}{V((x_{t-l}^{ep})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})} \\ &= \frac{\text{Cov}(Y, \gamma_0^2 + 2\gamma_0\gamma_1 x_{t-l} + 2u^L \gamma_1 x_{t-l} + 2\gamma_0 u_t^L + \gamma_1^2 (x_{t-l})^2 + u_t^{L2} | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})}{V((x_{t-l}^{ep})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})} \\ &\approx \frac{(2\gamma_0\gamma_1 \text{Cov}(Y, x_{t-l} | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z}) + \gamma_1^2 \text{Cov}(Y, (x_{t-l})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z}))}{2V(x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2} \\ &\approx \frac{(\gamma_1^2 \text{Cov}(Y, (x_{t-l})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z}))}{V((x_{t-l})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})} \frac{V((x_{t-l})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})}{2V(x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2} \\ &= \frac{\gamma_1^2 \theta'_i V((x_{t-l})^2 | \mathbf{z}, \mathbf{z}^2, x_{t-l}^{ep}, x_{t-l}^{ep} \times \mathbf{x}^{ep-l}, \mathbf{x}^{ep} \times \mathbf{z})}{2V(x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2} \approx \frac{2V(x_{t-l} | \mathbf{x}_t^{ep-l}, \mathbf{x}_t^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2}{2V(x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2} \gamma_1^2 \theta'_i \\ &= \frac{\gamma_1^2 V(x_{t-l} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2}{V(x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2} \theta'_i = \lambda_i^2 \theta'_i \end{aligned}$$

$$\text{where } \lambda_i^2 = \left(\frac{\gamma_1^2 V(x_{t-l} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2}{V(x_{t-l}^{ep} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2)^2} \right) / \gamma_1^2 = \frac{\left(R_{x_{t-l}^{ep}, x_{t-l} | \mathbf{x}^{ep-l}, \mathbf{x}^{ep-l} \times \mathbf{z}, \mathbf{z}, \mathbf{z}^2}^2 \right)^2}{\gamma_1^2} \approx \frac{\left(R_{x_t^{ep}, x_t}^2 \right)^2}{\gamma_1^2}$$

This may be applicable for non-linear models such as logistic regression when the effect size is small, or the measurement error is small.

Appendix S9. Simulation experiments.

Modelled estimates of PM_{2.5}, O₃, (United States Environmental Protection Agency (EPA)'s Fused Air Quality Surface Using Downscaling (FAQSD)) and temperature (Daymet) in the United States were used.

For all census tracts in the United States for June to August 2020 from the United States, a generalized additive model was fit to regress the difference between the daily FAQSD PM_{2.5} estimate (Lag0) and daily PM_{2.5} measurement (Lag0) at EPA monitor against census tract-level FAQSD PM_{2.5} estimates from (Lag0, Lag1, and Lag2) and a three-way spline (Longitude, Latitude, Time). Generalized cross-validation was used to determine degrees of freedom of the spline. All terms were statistically significant, but the model explained the dependent variable $\leq 10\%$ according to the deviance explained. This low deviance explained implies the existence of unexplained variability of the difference to a great degree. The residuals were still not randomly distributed.

Using this model, the predicted variable was obtained for all census tracts in Chicago and was normalized. Then, the covariance of this predicted variable (i.e., spatiotemporal covariance) was non-parametrically estimated using the method developed by Yang and Qiu (2019).

Several exposure measurement error scenarios for linear-like errors were considered.

$X_{i,t}^{ep} = \gamma_1 X_{i,t} + \kappa_{additive} \times error_{1,i,t} + \gamma_0$ for additive linear-like errors

$X_{i,t}^{ep} = \exp(\gamma_1 \log(X_{i,t})) + \kappa_{multiplicative} \times error_{1,i,t} - error_{2,i,t}/2$ for multiplicative linear-like errors

where $error_{1,i,t}$ follows a multivariate normal distribution with the covariance estimated and $error_{2,i,t}$ follows a gamma distribution. $X_{i,t}$ is a FAQSD estimate, which is spatiotemporally autocorrelated. $error_{1,i,t}$ is highly spatiotemporally autocorrelated. $error_{2,i,t}$ is completely random. Several κ values were used. A large κ is reasonable to mimick real-world scenarios because the generalized additive model used above was able to explain the difference to a very limited degree.

For Berkson-like errors,

$$X_{i,t}^{new} = \exp(1.2X_{i,t} + 7error_{1,i,t} - error_{2,i,t})$$

and then,

$X_{g,t}$ was obtained using a population-weighted average, where g represents ZIP Code in Chicago

$E_g[X_{i,t}^{new}]$ for additive Berkson-like error

$\exp(E_g[\log(X_{i,t}^{new})])$ for multiplicative Berkson-like error

The US Department of Housing and Urban Development's USPS ZIP Code-Census Tract Crosswalk file was used to create population-weighted averages at each ZIP Code. A total of 784 census tracts and 89 ZIP Codes in Chicago were considered.

The outcome generating model was designed to draw samples using $Y_{i,t} \sim Binom(p_{i,t})$ where $p_{i,t} = Kp_{0,i,t} \exp(\sum_{l=0}^2 \theta_l PM2.5_{i,t-l} + \sum_{l=0}^2 \vartheta_l O3_{i,t-l} + \sum_{l=0}^1 \varsigma_l HighTemp_{i,t-l})$. $p_{0,i,t}$, time-varying baseline hazard, was determined using statistics on underlying cause of death and population size from the Centers for Disease Control and Prevention's WONDER and 2020 decennial Census from the Census Bureau.

$\theta_0 = 0.001$, $\theta_1 = 0.0024$, and $\theta_3 = 0.0006$

$\vartheta_0 = 0.0001$, $\vartheta_1 = 0.0003$, and $\vartheta_2 = 0.0001$ (the absolute level of O_3 is approximately 10 times bigger than $PM_{2.5}$, so that the effects of O_3 and $PM_{2.5}$ are comparable)
 $\zeta_0 = 0.01$ and $\zeta_1 = 0.005$ (temperature level, °C).

For each simulation sample, time-stratified case-crossover analysis with conditional logistic regression was conducted to estimate θ_l . The percentage of rejecting the null hypothesis (empirical statistical power), $E[\hat{\theta}_l]$ and $E[\widehat{SE}[\hat{\theta}_l]]$ were obtained.




Figure S1. Residual plots including partial autocorrelation plots for linear regressions in Table 1 of the main manuscript

References in Online Supplementary Materials

Yang K, Qiu P. Nonparametric estimation of the spatio-temporal covariance structure. *Statistics in Medicine* 2019;**38**(23):4555-4565.