

A LAZARD CORRESPONDENCE FOR POST-LIE RINGS AND SKEW BRACES

S. TRAPPENIERS

ABSTRACT. We study how the Lazard correspondence behaves on semi-direct sums of Lie rings and use this to develop a Lazard correspondence between post-Lie rings and skew braces that satisfy a natural completeness condition. In particular, for a prime p and $k < p$, this yields a correspondence between skew braces of order p^k and left nilpotent post-Lie rings of order p^k on a nilpotent Lie ring.

1. INTRODUCTION

In 1954, Lazard proved a correspondence between Lie rings and groups satisfying certain completeness conditions [22]. This built upon earlier work by Malcev [23]. The nowadays better known version of the Lazard correspondence, although more restrictive than the original version, states that for p^k a prime power, there is a bijective correspondence between nilpotent Lie rings of order p^k and nilpotency class at most $p - 1$, and nilpotent groups of order p^k of nilpotency class at most $p - 1$. In particular, one finds a correspondence between groups of order p^k and nilpotent Lie rings of order p^k for $k < p$. This has proved to be very useful in the classification of finite p -groups, see for example [26].

Pre-Lie algebras, also called left-symmetric algebras, arose as early as in 1881 in the work of Cayley [12], only to resurface again 80 years later in the works of Vinberg [40] and Koszul [21]. See [8] for a survey on pre-Lie algebras. One of the motivations to study pre-Lie algebras comes from geometry, as they appear naturally in the study of regular affine actions of Lie groups [20]. More recently, the notion of a post-Lie algebra was introduced in the works of Vallette [39], see also [16]. For R a commutative ring, a *post-Lie R -algebra* is a Lie R -algebra \mathfrak{a} , with Lie bracket $[-, -]$, together with an R -bilinear map $\triangleright : \mathfrak{a} \times \mathfrak{a} \rightarrow \mathfrak{a}$ satisfying:

$$(1.1) \quad x \triangleright [y, z] = [x \triangleright y, z] + [y, x \triangleright z],$$

$$(1.2) \quad [x, y] \triangleright z = (x, y, z)_{\triangleright} - (y, x, z)_{\triangleright},$$

with $(x, y, z)_{\triangleright} := x \triangleright (y \triangleright z) - (x \triangleright y) \triangleright z$. If \mathfrak{a} is abelian, then $(\mathfrak{a}, \triangleright)$ is a *pre-Lie R -algebra*. A post-Lie \mathbb{Z} -algebra, respectively pre-Lie \mathbb{Z} -algebra, is called a *post-Lie ring*, respectively a *pre-Lie ring*.

Braces were introduced in 2006 by Rump [29, 30] in the study of the set-theoretical solutions to the Yang–Baxter equation. They were subsequently generalized by Guarnieri and Vendramin to skew braces [18]. A *skew brace* is a triple (A, \cdot, \circ) with A a set and (A, \cdot) and (A, \circ) group structures such that

$$a \circ (b \cdot c) = (a \circ b) \cdot a^{-1} \cdot (a \circ b),$$

holds for all $a, b, c \in A$. Here a^{-1} denotes the inverse of a in (A, \cdot) . When (A, \cdot) is abelian, then (A, \cdot, \circ) is called a *brace*. For braces, the notation $(A, +, \circ)$ is also

2020 *Mathematics Subject Classification*. 20N99, 17D25, 20F40.

Key words and phrases. post-Lie ring, pre-Lie ring, brace, skew brace, left nilpotent, Lazard correspondence.

common (in fact, this is also the case for skew braces, but to avoid any possible confusion we will abstain from using $+$ for a non-abelian group operation). Skew braces are essentially the same as bijective 1-cocycles of groups or regular subgroups of the holomorph of a group. In this sense, they were already being studied in a geometric framework in the form of regular affine actions of Lie groups on vector spaces [3, 25] long before they were formally introduced. Since pre-Lie algebras played an important role there, some connection between skew braces and pre-Lie rings is to be expected. This was indeed first stated by Rump [31] for \mathbb{R} -braces. Bachiller then used the Lazard correspondence in order to give a counterexample of a conjecture on braces [4], starting from a counterexample by Burde [7] to a question by Milnor [25] (note that the first counterexample to Milnor's question was given earlier by Benoist [6]).

Smoktunowicz proved that the construction of the group of flows, as defined for pre-Lie algebras in [2], could be adapted to left nilpotent pre-Lie rings of prime power order p^k , with $k+1 < p$, in order to obtain braces [36] with the same additive structure. Here, a post-Lie ring $(\mathfrak{a}, \triangleright)$ is *left nilpotent* if there exists some n such that $\mathfrak{a}^n = 0$ where $\mathfrak{a}^1 = \mathfrak{a}$ and \mathfrak{a}^{i+1} is the subgroup of $(\mathfrak{a}, +)$ generated by $a \triangleright b$ for $a \in \mathfrak{a}$, $b \in \mathfrak{a}^i$, *right nilpotency* is defined analogously. Similarly, a skew brace (A, \cdot, \circ) is *left nilpotent* if there exists some n such that $A^n = 1$, where $A^1 = A$ and A^{i+1} is the subgroup of (A, \cdot) generated by the elements $a * b = a^{-1} \cdot (a \circ b) \cdot b^{-1}$ for all $a \in A$, $b \in A^i$, *right nilpotency* is defined analogously. Recall that a skew brace of prime power order is always left nilpotent [13]. It is not immediately clear whether Smoktunowicz's construction is invertible, an inverse construction is given only in the case of strongly nilpotent braces of order p^k , with $k+1 < p$, and with strong nilpotency index at most $p-1$. Here strong nilpotency is equivalent to the structures being both left and right nilpotent [33]. This correspondence was used to classify all strongly nilpotent braces of order p^4 , for $p > 5^5$, in [27] while the non-strongly nilpotent ones of this order were classified earlier in [28]. In [35], a construction is given starting from a brace $(A, +, \circ)$ of order p^k , with $k+1 < p$, in order to obtain a pre-Lie ring structure with additive group $(A, +)/\{a \in A \mid p^2 a = 0\}$. Although the strong nilpotency condition is no longer present, one can not hope that this yields a correspondence as one quotients by a non-trivial subgroup.

Recently, Bai, Guo, Sheng and Tang showed how from a regular affine action of a Lie group onto another Lie group, one obtains a post-Lie algebra through differentiation [5] (note that, although not explicitly stated in this way, this construction also appears in [9]). This extended the earlier techniques to obtain a pre-Lie algebra in the case that the latter group is \mathbb{R}^n . Conversely, also in [5] a construction by formal integration was given to construct a skew brace starting from a post-Lie algebra of characteristic 0 and with some completeness conditions.

In Section 2 we recall the necessary information on the Lazard correspondence, and some basic notions on post-Lie rings and skew braces. In Section 3 and Section 4 we introduce the notion of filtered post-Lie rings and filtered skew braces and for both structures we define the property of being Lazard in a natural way. We construct a natural filtration on any post-Lie ring or skew brace. As a consequence, we prove that if (A, \cdot, \circ) is a left nilpotent skew brace and (A, \cdot) is a nilpotent group, then also (A, \circ) is a nilpotent group. This result was previously only known for finite skew braces [13, Theorem 4.8]. In fact we have a stronger result, for which we first have to introduce a new notion of nilpotency.

Let (A, \cdot, \circ) be a skew brace, define inductively $L^1(A) = A$ and $L^{i+1}(A)$ the subgroup of (A, \cdot) generated by all elements of the form $a * b := a^{-1} \cdot (a \circ b) \cdot b^{-1}$ and $a \cdot b \cdot a^{-1} \cdot b^{-1}$ for $a \in A$, $b \in L^i(A)$. Note that $L^i(A)$ is the left hand version of the series $R_n(A, A)$ as defined in [1]. By induction, one sees that the $L^i(A)$ are

a descending chain of normal subgroups of (A, \cdot) . We now define L -nilpotency of skew braces, which for braces clearly coincides with left nilpotency.

Definition 1.1. A skew brace (A, \cdot, \circ) is L -nilpotent if there $L^{k+1}(A) = 1$ for some $k \geq 0$. The minimal such k , if it exists, is the L -nilpotency class of A .

Recall that the annihilator $\text{Ann}(A)$ of a skew brace (A, \cdot, \circ) is the set of the elements $a \in A$ such that $a + b = a \circ b = b \circ a = b + a$ for all $b \in A$. We can now give a more precise formulation of our results which imply the earlier claim.

Lemma 1.2. Let (A, \cdot, \circ) be a skew brace. Then A is L -nilpotent if and only if A is left nilpotent and (A, \cdot) is a nilpotent group.

Theorem 1.3. Let (A, \cdot, \circ) be an L -nilpotent skew brace of class k . Then (A, \circ) is nilpotent of class at most k and the k -th term in the lower central series of (A, \circ) is contained in $\text{Ann}(A)$.

In Section 5 we first prove that the exponential and logarithmic map give a correspondence between certain derivations and automorphisms of Lazard Lie rings, mimicking the geometric correspondence given by integration and differentiation. We then prove the final correspondence between Lazard post-Lie rings and Lazard skew braces, which extends both Smoktunowicz's correspondence on pre-Lie rings and braces of prime power order and also comprises the formal integration by Bai, Guo, Sheng and Tang. Before formulating (a restricted version of) this result, we introduce the notion of L -nilpotency for post-Lie rings.

Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring. Inductively define $L^1(\mathfrak{a}) = \mathfrak{a}$ and $L^{i+1}(\mathfrak{a})$ the subgroup of $(\mathfrak{a}, +)$ generated by all elements of the form $a \triangleright b$ and $[a, b]$ where $a \in \mathfrak{a}$, $b \in L^i(\mathfrak{a})$. One easily verifies that $L^i(\mathfrak{a})$ gives a descending filtration of ideals of the Lie ring \mathfrak{a} . We now are ready to define L -nilpotency, which for pre-Lie rings is the same as left nilpotency.

Definition 1.4. A post-Lie ring $(\mathfrak{a}, \triangleright)$ is L -nilpotent if $L^{k+1}(\mathfrak{a}) = 0$ for some $k \geq 0$. The minimal such k , if it exists, is the L -nilpotency class of $(\mathfrak{a}, \triangleright)$.

We are now ready to formulate a restrictive version of the correspondence in Theorem 5.9.

Theorem 1.5. Let p be a prime and $k < p$. There is a bijective correspondence between post-Lie rings $(\mathfrak{a}, \triangleright)$ of L -nilpotency class k such that $(\mathfrak{a}, +)$ is a p -group and skew braces (A, \cdot, \circ) of L -nilpotency class k such that (A, \cdot) and (A, \circ) are p -groups.

In particular, there is a bijective correspondence between skew braces of size p^k and L -nilpotent post-Lie rings of size p^k .

This correspondence is explicitly described in Proposition 5.7 and Proposition 5.8, which in particular implies that for brace one direction of this correspondence is precisely the group of flows construction as in [36]. The proof of Theorem 1.5 is given at the end of Section 5.

In [5] the authors propose to find a suitable differentiation and formal integration theory for finite skew braces and post-Lie rings. One such approach is precisely the one using the exponential and the logarithmic map as in Section 5. On the other hand, some techniques used by Smoktunowicz in [34, 36] could also be seen as an discrete version of differentiation. Therefore, in Section 6 we motivate why this is the case and we give an application to skew braces which extends an application by Smoktunowicz. We end by emphasizing a possible advantage of this technique and ask whether this could possibly lead to a relaxation on the nilpotency condition imposed by the exponent of the group in [36, Theorem 13].

2. PRELIMINARIES

Let us recall the precise statement of the Lazard correspondence together with some useful related results. Although we use slightly different terminology, all of the statements regarding the Lazard correspondence follow from the treatment in [19, 22].

For a group G and P a set of prime numbers, G is P -divisible if for every element $g \in G$ and every n whose prime divisors are contained in P , there exists a unique $h \in G$ such that $h^n = g$. This unique element is denoted $g^{\frac{1}{n}}$. If G is an abelian group and we denote its group operation by $+$, then we use the notation $\frac{1}{n}g$.

A *filtered group* is a group G with a filtration of normal subgroups

$$G = G_0 \supseteq G_1 \supseteq G_2 \supseteq \dots$$

such that $[G_i, G_j] \subseteq G_{i+j}$. Any subgroup H of a filtered group G has a natural filtration given by $H_i = H \cap G_i$. We endow a filtered group with the topology with base $\{gG_i \mid g \in G, i \geq 1\}$. A homomorphism of filtered groups $f : G \rightarrow H$ is a group homomorphism such that $f(G_i) \subseteq H_i$. A filtered group G is *complete* if it is Hausdorff and complete as topological space. A *Lazard group* is a complete filtered group G such that $G_0 = G_1$ and G_i is P_i -divisible for every $i \geq 1$, where P_i is the set of all primes less than or equal to i .

Example 2.1. On any group G one obtains a natural filtration by setting $G_0 = G_1 = G$ and $G_{i+1} = [G, G_i]$ for $i \geq 1$. If there exists some k such that $G_{k+1} = \{1\}$ (meaning that G is nilpotent) then the induced topology is discrete hence complete. If we moreover assume that G is a p -group for some prime $k < p$, then G is easily seen to be a Lazard group since every p -group is P_i -divisible for $i < p$. However, note that any P_p -divisible p -group is necessarily trivial.

Similarly, a *filtered Lie ring* is a Lie ring \mathfrak{a} with a filtration of ideals

$$\mathfrak{a} = \mathfrak{a}_0 \supseteq \mathfrak{a}_1 \supseteq \mathfrak{a}_2 \supseteq \dots$$

such that $[\mathfrak{a}_i, \mathfrak{a}_j] \subseteq \mathfrak{a}_{i+j}$ for all $i, j \geq 1$. A homomorphism of filtered Lie rings $f : \mathfrak{a} \rightarrow \mathfrak{b}$ is a Lie ring homomorphism such that $f(\mathfrak{a}_i) \subseteq \mathfrak{b}_i$. We endow a filtered Lie ring with the topology with base $\{x + \mathfrak{a}_i \mid x \in \mathfrak{a}, i \geq 1\}$. A filtered Lie ring is *complete* if its underlying additive group (with the same filtration) is complete. A *Lazard Lie ring* is a Lie ring whose underlying additive group is Lazard.

Example 2.2. On any Lie ring \mathfrak{a} one obtains a natural filtration by setting $\mathfrak{a}_0 = \mathfrak{a}_1 = \mathfrak{a}$ and $\mathfrak{a}_{i+1} = [\mathfrak{a}, \mathfrak{a}_i]$ for $i \geq 1$. If there exists some k such that $\mathfrak{a}_{k+1} = \{1\}$ (meaning that \mathfrak{a} is nilpotent) then the induced topology is discrete hence complete. If $(\mathfrak{a}, +)$ is a p -group for $k < p$ then \mathfrak{a} is Lazard.

The benefit of working in a complete Lie ring \mathfrak{a} is that the Baker-Campbell-Hausdorff (BCH) formula can be evaluated in any two elements. Recall that this formula originates in Lie theory and its first terms are given by

$$(2.1) \quad \text{BCH}(x, y) = x + y + \frac{1}{2}[x, y] + \frac{1}{12}([x, [x, y]] + [y, [y, x]]) - \frac{1}{24}[y, [x, [x, y]]] + \dots,$$

where higher terms are of order at least 5. The prime divisors of the denominator of coefficients of order n only involve primes $\leq n$, which indeed ensures that these are well-defined in \mathfrak{a} . The element $\text{BCH}(x, y)$, for $x, y \in \mathfrak{a}$, is then interpreted as $\lim_{k \rightarrow \infty} \text{BCH}_k(x, y)$ where BCH_k is obtained by truncating the formula at order k . Note in particular that if $x, y \in \mathfrak{a}$ commute, then $\text{BCH}(x, y) = x + y$. In particular, $\text{BCH}(x, -x) = 0$ and $\text{BCH}(x, x) = 2x$.

Similarly, in a Lazard group A , one can evaluate formal infinite products of commutators as long as their order goes to infinity and as long as the fractional

powers involve only primes $\leq n$ for commutators of order n . Two examples of such formulae are the first and second inverse BCH formula (see also [15], [19]), whose first terms are

$$(2.2) \quad P(g, h) = gh[g, h]^{-\frac{1}{2}}[g, [g, h]]^{\frac{1}{12}}[g, [g, [g, h]]]^{-\frac{1}{24}}[h, [h, [g, h]]]^{\frac{1}{24}} \cdots,$$

and

$$(2.3) \quad Q(g, h) = [g, h][g, [g, h]]^{\frac{1}{2}}[h, [g, h]]^{\frac{1}{2}}[g, [g, [g, h]]]^{\frac{1}{3}}[h, [g, [g, h]]]^{\frac{1}{4}}[h, [h, [g, h]]]^{\frac{1}{3}} \cdots,$$

where in this case $[g, h]$ is the group theoretic commutator $ghg^{-1}h^{-1}$ and further factors involve iterated brackets of order at least 5.

Theorem 2.3. *Let \mathfrak{a} be a Lazard Lie ring, then $\mathbf{Laz}(\mathfrak{a}) := (\mathfrak{a}, \text{BCH})$ is a Lazard group with respect to the same filtration. If $f : \mathfrak{a} \rightarrow \mathfrak{b}$ is a homomorphism of Lazard Lie rings, then $f : \mathbf{Laz}(\mathfrak{a}) \rightarrow \mathbf{Laz}(\mathfrak{b})$ is a homomorphism of Lazard groups.*

Conversely, let A be a Lazard group. Then $\mathbf{Laz}^{-1}(A) := (A, +, [-, -])$, where $a + b := P(a, b)$ and $[a, b] := Q(a, b)$, is a Lazard Lie ring with respect to the same filtration. This construction is also functorial. \mathbf{Laz} and \mathbf{Laz}^{-1} are inverse constructions.

Corollary 2.4. *Let \mathfrak{a} be a Lazard Lie ring, then the Lazard sub Lie rings of \mathfrak{a} and Lazard subgroups of $\mathbf{Laz}(\mathfrak{a})$ coincide. Also, Lazard ideals of \mathfrak{a} and Lazard normal subgroups of $\mathbf{Laz}(\mathfrak{a})$ coincide.*

A filtered ring R is a ring together with a descending filtration of ideals

$$A_0 \supseteq A_1 \supseteq A_2 \supseteq \dots$$

such that $A_i A_j \subseteq A_{i+j}$. A filtered ring R is *Lazard* if its underlying additive group is Lazard. If on a Lazard ring we consider the commutator bracket, then we obtain a Lazard Lie ring. Note that a Lazard ring is necessarily non-unital, but one can adjoin a unit to obtain the Dorroh extension $R' = \mathbb{Z} \oplus R$ with filtration $R'_0 = R'$, $R'_i = \{0\} \oplus R_i$ for $i \geq 1$. Then the exponential

$$\exp(a) = \sum_{k=0}^{\infty} \frac{1}{k!} a^k = \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{1}{k!} a^k \in 1 + R$$

is well-defined for $a \in R$. Similarly, for any $a \in 1 + R$ the logarithm

$$\log(a) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} (a - 1) = \lim_{n \rightarrow \infty} \sum_{k=1}^n (-1)^{k+1} \frac{1}{k} (a - 1) \in R$$

is a well-defined element and \exp and \log are mutually inverse. Then the Baker-Campbell-Hausdorff formula is obtained as $\text{BCH}(a, b) = \log(\exp(a) \exp(b))$. Similarly, $P(g, h) = \exp(\log(g) + \log(h))$ and $Q(g, h) = \exp(\log(g) \log(h) - \log(h) \log(g))$ where we evaluate P and Q in the Lazard group $1 + R$. Explicitly we have the following result.

Proposition 2.5. *Let \mathfrak{a} be a Lazard Lie ring embedding in a Lazard ring R . Then $\exp(\mathfrak{a})$ is a Lazard group for the filtration $\exp(\mathfrak{a})_i = \exp(\mathfrak{a}_i)$ and $\exp : \mathbf{Laz}(\mathfrak{a}) \rightarrow \exp(\mathfrak{a})$ is an isomorphism of Lazard groups.*

Let \mathfrak{a} be a filtered Lie ring. We define $\text{End}(\mathfrak{a})$ as the ring of endomorphisms of the underlying additive group $(\mathfrak{a}, +)$. We have a natural filtration on $\text{End}(\mathfrak{a})$ given by

$$\text{End}(\mathfrak{a})_j = \{f \mid f(\mathfrak{a}_i) \subseteq \mathfrak{a}_{i+j}\},$$

which makes $\text{End}(\mathfrak{a})$ into a filtered ring. If \mathfrak{a} is complete, then also $\text{End}(\mathfrak{a})$ is easily seen to be complete. Let $\text{End}(\mathfrak{a})^+$ denote the filtered subring of $\text{End}(\mathfrak{a})$ consisting

of all elements in $\text{End}(\mathfrak{a})_1$. If \mathfrak{a} is Lazard, then it follows directly that $\text{End}(\mathfrak{a})^+$ is Lazard.

Remark 2.6. Note that although the difference between $\text{End}(\mathfrak{a})^+$ and $\text{End}(\mathfrak{a})_1$ is subtle, we consider $\text{End}(\mathfrak{a})_1$ simply as an ideal of $\text{End}(\mathfrak{a})$ while $\text{End}(\mathfrak{a})^+$ is seen as a filtered ring on its own. Throughout this paper we use a similar notation for any filtered ring, group or Lie ring to indicate the filtered substructure consisting of all elements contained in the second term in the filtration.

For \mathfrak{a} a filtered Lie ring and $a \in \mathfrak{a}$, we denote by ad_a the adjoint map $b \mapsto [a, b]$. Clearly, $\text{ad} : \mathfrak{a} \rightarrow \text{End}(\mathfrak{a}) : a \mapsto \text{ad}_a$ is a homomorphism of filtered Lie rings, where on $\text{End}(\mathfrak{a})$ we consider the commutator bracket.

Lemma 2.7. *Let \mathfrak{a} be a Lazard Lie ring, then $\text{BCH}(a, \text{BCH}(b, -a)) = \exp(\text{ad}_a)(b)$ for all $a, b \in \mathfrak{g}$.*

At last, we fix some notation and recall some important substructures of post-Lie rings and skew braces.

Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring and $a \in \mathfrak{a}$. We obtain a new Lie bracket on the abelian group \mathfrak{a} by defining $\{a, b\} := [a, b] + a \triangleright b - b \triangleright a$. This Lie ring is denoted by \mathfrak{a}° . We will denote left multiplication by a as \mathcal{L}_a , so $\mathcal{L}_a(b) = a \triangleright b$. Note that (1.2) states precisely that $\mathfrak{a}^\circ \rightarrow \text{End}(\mathfrak{a}) : a \mapsto \mathcal{L}_a$ is a Lie ring homomorphism.

Definition 2.8. Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring. A sub Lie ring I of $(\mathfrak{a}, \triangleright)$ is

- (1) a *sub post-Lie ring* if I is closed under the operation \triangleright ,
- (2) a *left ideal* if $x \triangleright y \in I$ for all $x \in \mathfrak{a}, y \in I$,
- (3) a *strong left ideal* if it is a left ideal and it is an ideal of \mathfrak{a} ,
- (4) an *ideal* if it is a strong left ideal and moreover I is an ideal of \mathfrak{a}° .

Example 2.9. Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring, then

$$\text{Fix}(\mathfrak{a}) := \{a \in \mathfrak{a} \mid b \triangleright a = 0 \text{ for all } b \in \mathfrak{a}\},$$

is a left ideal and the *socle*

$$\text{Soc}(\mathfrak{a}) := \{a \in \mathfrak{a} \mid a \triangleright b = [a, b] = 0 \text{ for all } b \in \mathfrak{a}\},$$

and *annihilator*

$$\text{Ann}(\mathfrak{a}) := \{a \in \mathfrak{a} \mid a \triangleright b = b \triangleright a = [a, b] = 0 \text{ for all } b \in \mathfrak{a}\},$$

are ideals. For the latter statement, note that for $a \in \text{Soc}(\mathfrak{a})$ and $b, c \in \mathfrak{a}$, it follows from (1.1) that $0 = b \triangleright [a, c] = [b \triangleright a, c]$ and from (1.2) that $0 = [a, b] \triangleright c = -(b \triangleright a) \triangleright c$ hence $b \triangleright a \in \text{Soc}(\mathfrak{a})$. As $\{b, a\} = b \triangleright a$, $\text{Soc}(\mathfrak{a})$ and $\text{Ann}(\mathfrak{a})$ are ideals.

Let (A, \cdot, \circ) be a skew brace. Then we define $\lambda_a(b) := a^{-1} \cdot (a \circ b)$ and in this way we obtain an action of (A, \circ) on (A, \cdot) by automorphisms

$$\lambda : (A, \circ) \rightarrow \text{Aut}(A, \cdot) : a \mapsto \lambda_a.$$

Note that $a * b$ can also be expressed as $\lambda_a(b) \cdot b^{-1}$.

Definition 2.10. Let (A, \cdot, \circ) be a skew brace. A subgroup I of (A, \cdot) is

- (1) a *sub skew brace* if I is a subgroup of (A, \circ) ,
- (2) a *left ideal* if $\lambda_a(b) \in I$ for all $a \in A, b \in I$,
- (3) a *strong left ideal* if I is a left ideal and also a normal subgroup of (A, \cdot) ,
- (4) an *ideal* if I is a strong left ideal and also a normal subgroup of (A, \circ) .

Example 2.11. Let (A, \cdot, \circ) be a skew brace, then

$$\text{Fix}(A) := \{a \in A \mid b \circ a = b + a \text{ for all } b \in A\},$$

is a left ideal and the *socle*

$$\text{Soc}(A) := \{a \in A \mid a \circ b = a + b = b + a \text{ for all } b \in A\},$$

is an ideal.

3. FILTERED POST-LIE RINGS

Definition 3.1. A *filtered post-Lie ring* is a post-Lie ring $(\mathfrak{a}, \triangleright)$ together with a descending chain of strong left ideals \mathfrak{a}_i such that it is a filtration for \mathfrak{a} .

Let \mathfrak{a} be a filtered Lie ring. We define $\text{Der}(\mathfrak{a})$ as the set of all derivations of \mathfrak{a} contained in $\text{End}(\mathfrak{a})$, these form a sub Lie ring of $\text{End}(\mathfrak{a})$ under the usual commutator Lie bracket. Thus they have a natural filtration $\text{Der}(\mathfrak{a})_i = \text{End}(\mathfrak{a})_i \cap \text{Der}(\mathfrak{a})$. Let $\delta : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{a})$ be a homomorphism of filtered Lie rings. We define the semi-direct sum $\mathfrak{a} \oplus_\delta \mathfrak{g}$ as the direct sum of the additive groups together with the bracket

$$[(a, x), (b, y)] = ([a, b] + \delta_x(b) - \delta_y(a), [x, y]).$$

It is well-known that this makes $\mathfrak{a} \oplus_\delta \mathfrak{g}$ into a Lie ring. The proof of the following statement is straightforward; the fact that this is indeed a filtration follows directly from the Jacobi identity.

Lemma 3.2. *Let $\mathfrak{a}, \mathfrak{g}$ be filtered Lie rings and $\delta : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{a})$ a filtered Lie ring homomorphism. Then $\mathfrak{a} \oplus_\delta \mathfrak{g}$ is a filtered Lie ring for the filtration $(\mathfrak{a} \oplus_\delta \mathfrak{g})_i = \mathfrak{a}_i \oplus_\delta \mathfrak{g}_i$. If moreover \mathfrak{a} and \mathfrak{g} are Lazard rings then $\mathfrak{a} \oplus_\delta \mathfrak{g}$ is a Lazard ring.*

In particular, we will be interested in the semidirect sum $\text{aff}(\mathfrak{a}) := \mathfrak{a} \oplus_\delta \text{Der}(\mathfrak{a})$ where $\delta = \text{id}_{\text{Der}(\mathfrak{a})}$ and its sub Lie ring $\text{aff}(\mathfrak{a})^+ = \mathfrak{a}^+ \oplus \text{Der}(\mathfrak{a})^+$. If \mathfrak{a} is Lazard, then so is $\text{Der}(\mathfrak{a})^+$ since it is closed in $\text{End}(\mathfrak{a})$, and therefore also $\text{aff}(\mathfrak{a})^+$ is Lazard.

Definition 3.3. Let \mathfrak{a} be a Lie ring, a sub Lie ring \mathfrak{g} of $\text{aff}(\mathfrak{a})$ is *t-surjective*, respectively *t-bijective* if the projection $\text{pr}_\mathfrak{a} : \text{aff}(\mathfrak{a}) \rightarrow \mathfrak{a}$ to $\text{pr}_\mathfrak{a} : \mathfrak{g} \rightarrow \mathfrak{a}$ restricted to \mathfrak{g} yields a surjective, respectively bijective, map from \mathfrak{g} to \mathfrak{a} .

Proposition 3.4. *Let \mathfrak{a} be a filtered Lie ring. Then there is a bijective correspondence between operations $\triangleright : \mathfrak{a} \times \mathfrak{a} \rightarrow \mathfrak{a}$ making $(\mathfrak{a}, \triangleright)$ into a filtered post-Lie ring and t-bijective sub Lie rings of $\text{aff}(\mathfrak{a})$.*

Proof. This proof is a slight variation of the proof of [10, Proposition 2.12]. We therefore only give a sketch of the correspondence.

If $(\mathfrak{a}, \triangleright)$ is a filtered post-Lie ring then $\{(a, \mathcal{L}_a) \mid a \in \mathfrak{a}\}$ is its associated t-bijective sub Lie ring of $\text{aff}(\mathfrak{a})$, where $\mathcal{L}_a(b) = a \triangleright b$. Since the filtration on \mathfrak{a} consists of left ideals we find indeed that the maps \mathcal{L}_a are contained in $\text{End}(\mathfrak{a})$.

Conversely, if we are given a t-bijective sub Lie ring \mathfrak{g} of $\text{aff}(\mathfrak{a})$ then we define $a \triangleright b$ as $\mathcal{L}_a(b)$ where \mathcal{L}_a is the unique endomorphism of \mathfrak{a} satisfying $(a, \mathcal{L}_a) \in \mathfrak{g}$. In this way $(\mathfrak{a}, \triangleright)$ is a filtered post-Lie ring. Since the maps \mathcal{L}_a respect the filtration on \mathfrak{a} we find that \mathfrak{a}_i is a left ideal for each $i \geq 0$. \square

Corollary 3.5. *Let $(\mathfrak{a}, \triangleright)$ be a filtered post-Lie ring. We obtain a Lie ring structure \mathfrak{a}° on \mathfrak{a} with filtration $\mathfrak{a}_i^\circ = \{a \in \mathfrak{a}_i \mid \mathcal{L}_a \in \text{Der}(\mathfrak{a})_i\}$ and Lie bracket*

$$\{a, b\} := [a, b] + a \triangleright b - b \triangleright a.$$

Proof. Let \mathfrak{g} be the associated t-bijective sub Lie ring of $\text{aff}(\mathfrak{a})$ as in Proposition 3.4. The statement follows by transferring the Lie ring structure of \mathfrak{g} onto \mathfrak{a} through the bijection $\text{pr}_\mathfrak{a} : \mathfrak{g} \rightarrow \mathfrak{a}$. \square

Definition 3.6. Let $(\mathfrak{a}, \triangleright)$ be a filtered post-Lie ring, then $(\mathfrak{a}, \triangleright)$ is *Lazard* if both \mathfrak{a} and \mathfrak{a}° are Lazard Lie rings.

Lemma 3.7. *Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring, then setting $\mathfrak{a}_0 = \mathfrak{a}$ and $\mathfrak{a}_i = L^i(\mathfrak{a})$ for $i \geq 1$ makes $(\mathfrak{a}_i, \triangleright)$ into a filtered post-Lie ring.*

Proof. It is straightforward to see that the \mathfrak{a}_i form a descending chain of strong left ideals. It remains to show that $[L^i(\mathfrak{a}), L^j(\mathfrak{a})] \subseteq L^{i+j}(\mathfrak{a})$ for all $i, j \geq 1$. Let $\mathfrak{h} = \{\mathcal{L}_a \mid a \in \mathfrak{a}\} \leq \text{Der}(\mathfrak{a})$ and consider the semi-direct sum $\mathfrak{t} = \mathfrak{a} \oplus_{\delta} \mathfrak{h}$ where $\delta : \mathfrak{h} \rightarrow \text{Der}(\mathfrak{a})$ is the inclusion map. Then we find $[(a, \mathcal{L}_b), (c, 0)] = ([a, c] + b \triangleright c)$. It follows that $[\mathfrak{t}, \mathfrak{t}] = L^2(\mathfrak{a}) \times [\mathfrak{h}, \mathfrak{h}]$. For a Lie ring \mathfrak{g} we introduce the following notation for iterated commutators: $\gamma^1(\mathfrak{g}) = \mathfrak{g}$ and $[\gamma^{i+1}(\mathfrak{g})] = [\mathfrak{g}, \gamma^i(\mathfrak{g})]$. Then the previous equality can be rewritten as $\gamma^2(\mathfrak{t}) = L^2(\mathfrak{a}) \times \gamma^2(\mathfrak{h})$ and by induction one finds $\gamma^i(\mathfrak{t}) = L^i(\mathfrak{a}) \times \gamma^i(\mathfrak{h})$. It is well-known that $[\gamma^i(\mathfrak{t}), \gamma^j(\mathfrak{t})] \subseteq \gamma^{i+j}(\mathfrak{t})$, so in particular $([L^i(\mathfrak{a}), L^j(\mathfrak{a})]) \times \{0\} = [L^i(\mathfrak{a}) \times \{0\}, L^j(\mathfrak{a}) \times \{0\}] \subseteq L^{i+j}(\mathfrak{a}) \times \gamma^{i+j}(\mathfrak{h})$ thus $[L^i(\mathfrak{a}), L^j(\mathfrak{a})] \subseteq L^{i+j}(\mathfrak{a})$. \square

Let \mathbb{F} be a field of characteristic 0 and $(\mathfrak{a}, \triangleright)$ be a finite dimensional pre-Lie \mathbb{F} -algebra. Then $(\mathfrak{a}, \triangleright)$ is complete [20] if for every $a \in \mathfrak{a}$, the map $r_a : \mathfrak{a} \rightarrow \mathfrak{a} : b \mapsto b + a \triangleright b$ is a bijection. However, if \mathfrak{a}° is nilpotent, then this is equivalent to the left multiplication maps \mathcal{L}_a being nilpotent for all $a \in \mathfrak{a}$ [32, Proposition 2]. By Engel's theorem, this is equivalent with the existence of some n such that $\mathcal{L}_{a_1} \cdots \mathcal{L}_{a_n} = 0$ for all $a_1, \dots, a_n \in \mathfrak{a}$. It follows that this notion of completeness coincides with L -nilpotency, or equivalently with the fact that its associated filtered pre-Lie ring is Lazard. Indeed, the divisibility condition is trivially satisfied, as $\mathfrak{a}_k = 0$ for some $k \geq 0$, also $\mathfrak{a}_k^\circ = 0$ and thus also the completeness follows for both \mathfrak{a} and \mathfrak{a}° . Note that the condition that all \mathcal{L}_a are nilpotent also appears in [9, Theorem 3.1].

Lemma 3.8. *Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring. Then $(\mathfrak{a}, \triangleright)$ is L -nilpotent if and only if $(\mathfrak{a}, \triangleright)$ is left nilpotent and \mathfrak{a} is a nilpotent Lie ring.*

Proof. From the definition of $L^i(\mathfrak{a})$ it is clear that if $(\mathfrak{a}, \triangleright)$ is L -nilpotent then it is left nilpotent and \mathfrak{a} is a nilpotent Lie ring.

Conversely, assume that $(\mathfrak{a}, \triangleright)$ is left nilpotent and \mathfrak{a} is a nilpotent Lie ring. Let $\mathfrak{a} = \gamma^1(\mathfrak{a}) \supseteq \dots \supseteq \gamma^d(\mathfrak{a}) = \{0\}$ denote the lower central series of \mathfrak{a} and let c be such that $\mathfrak{a}^{c+1} = \{0\}$ (recall that this is the left derived series defined in the introduction). Assume $L^i(\mathfrak{a}) \subseteq \gamma^k(\mathfrak{a})$ for some $i, k \geq 1$, then $[L^i(\mathfrak{a}), \mathfrak{a}] \subseteq \gamma^{k+1}(\mathfrak{a})$ hence $L^{i+1}(\mathfrak{a}) \subseteq (\mathfrak{a} \triangleright L^i(\mathfrak{a})) + \gamma^{k+1}(\mathfrak{a})$. In particular, $L^{i+c}(\mathfrak{a}) \subseteq \mathfrak{a}^{c+1} + \gamma^{k+1}(\mathfrak{a}) = \gamma^{k+1}(\mathfrak{a})$. We now conclude by induction that $L^{kc+1}(\mathfrak{a}) \subseteq \gamma^{k+1}(\mathfrak{a}) = 1$. \square

The proof of Theorem 1.3 is analogous to the proof of Theorem 3.9.

Theorem 3.9. *Let $(\mathfrak{a}, \triangleright)$ be an L -nilpotent post-Lie ring of class k . Then \mathfrak{a}° is nilpotent of class at most k and the k -th term in the lower central series of \mathfrak{a}° is contained in $\text{Ann}(\mathfrak{a})$.*

Proof. Let $(\mathfrak{a}, \triangleright)$ be L -nilpotent of class k and consider it as a filtered post-Lie ring with its canonical filtration. Then $\mathfrak{a}_{k+1}^\circ \subseteq \mathfrak{a}_{k+1} = 0$, so in particular \mathfrak{a}° is nilpotent of class at most k . Also, since $\mathfrak{a}_k^\circ = \mathfrak{a}_k \cap \{a \in \mathfrak{a} \mid \mathcal{L}_a \in \text{End}(\mathfrak{a})_k\}$ but $\text{End}(\mathfrak{a})_k = 0$ and clearly $\mathfrak{a}_k \in Z(\mathfrak{a}) \cap \text{Fix}(\mathfrak{a})$, the statement follows. \square

4. FILTERED SKEW BRACES

Definition 4.1. A *filtered skew brace* is a skew brace (A, \cdot, \circ) together with a descending chain of strong left ideals A_i such that it is a filtration for (A, \cdot)

Definition 4.2. Let G be a filtered group. By $\text{Aut}(G)$ we denote the filtered group automorphisms of G and we define a filtration by

$$\text{Aut}(G)_i = \{f \mid f(g)g^{-1} \in G_{i+j}, \text{ for all } g \in G_j\}.$$

Lemma 4.3. *Let G be a filtered group, then $\text{Aut}(G)$ is a filtered group.*

Proof. Consider the semidirect product $T = G \rtimes \text{Aut}(G)$. We identify G and $\text{Aut}(G)$ with the subgroups $G \times \{1\}$ and $\{1\} \times \text{Aut}(G)$. Note that the subgroups G_i are normal in T . By assumption, $[\text{Aut}(G)_i, [\text{Aut}(G)_j, G_k]] \subseteq G_{i+j+k}$, so by the Hall-Witt identity we find that $[G_k, [\text{Aut}(G)_i, \text{Aut}(G)_j]] \subseteq G_{i+j+k}$ and thus we conclude that $[\text{Aut}(G)_i, \text{Aut}(G)_j] \subseteq \text{Aut}(G)_{i+j}$. \square

Lemma 4.4. *Let A, G be filtered groups and $\lambda : G \rightarrow \text{Aut}(A)$ a homomorphism of filtered groups. Then the semidirect product $A \rtimes_\lambda G$ is a filtered group for the filtration $(A \rtimes_\lambda G)_i = A_i \rtimes_\lambda G_i$ is a filtered group.*

Proof. Since we are working in a semi-direct product we find that $[A_i \rtimes_\lambda G_i, A_j \rtimes_\lambda G_j] = [A_i, A_j][A_i, G_j][A_j, G_i][G_i, G_j]$ where we identify A and G with the corresponding subgroup of $A \rtimes G$. Clearly $[A_i, A_j] \subseteq A_{i+j}$ and $[G_i, G_j] \subseteq G_{i+j}$. Since λ is a homomorphism of filtered group, also $[A_i, G_j] \subseteq A_{i+j}$ and $[A_j, G_i] \subseteq A_{i+j}$ which concludes the proof. \square

Let A be a filtered group. We will particularly be interested in the *holomorph* of A , which by definition is the semi-direct product $\text{Hol}(A) := A \rtimes \text{Aut}(A)$.

Definition 4.5. Let A be a filtered group. A subgroup $G \leq \text{Hol}(A)$ is *transitive*, respectively *regular*, if $\text{pr}_G : H \rightarrow G$ is surjective, respectively bijective.

Note that if we consider the natural action of $\text{Hol}(G)$ on G , i.e. $(g, \lambda) \cdot h = g\lambda(h)$, then transitive subgroups of $\text{Hol}(G)$ are precisely the ones such that the associated action is transitive. Regular subgroups are precisely the ones that induces a regular (or also called simply transitive) action.

Proposition 4.6. *Let (A, \cdot) be a filtered group. There is a bijective correspondence between group structures (A, \circ) making (A, \cdot, \circ) into a filtered skew brace and regular subgroups of $\text{Hol}(A, \cdot)$.*

Proof. The proof is only a slight variation of the proof of [18, Theorem 4.2], therefore we only give a sketch. Let (A, \circ) be a group operation such that (A, \cdot, \circ) is a filtered skew brace. Define $G = \{(a, \lambda_a) \mid a \in A\} \leq \text{Hol}(A)$ then G is a regular subgroup of $\text{Hol}(A)$. Note that A_i are strong left ideals so indeed λ_a respects this filtration for all $a \in A$.

Conversely, if G is a regular subgroup of $\text{Hol}(A)$, we transfer the group structure of G onto A with the help of the bijection $\text{pr}_A : G \rightarrow A$ (or equivalently $a \circ b = a + \lambda_a(b)$ with λ_a the unique map such that $(a, \lambda_a) \in G$) and we denote this new group operation by \circ . Then (A, \cdot, \circ) becomes a skew brace and by assumption the maps λ_a preserve the filtration A_i , hence the A_i are strong left ideals. \square

Corollary 4.7. *Let (A, \cdot, \circ) be a filtered skew brace. Then (A, \circ) with the filtration $(A, \circ)_i = A_i \cap \{a \in A \mid \lambda_a \in \text{Aut}(A)_i\}$ is a filtered group.*

Proof. Let (A, \cdot, \circ) be a filtered skew brace and let G be the associated regular subgroup of $\text{Hol}(A, \cdot)$ as in Proposition 4.6. Then $G_i = G \cap \text{Hol}(A, \cdot)_i$ makes G into a filtered group. Transferring this filtration of G onto (A, \circ) through the isomorphism $(A, \circ) \rightarrow G : a \mapsto (a, \lambda_a)$ now yields the statement. \square

The notion of a Lazard skew brace is now the natural one.

Definition 4.8. Let (A, \cdot, \circ) be a filtered skew brace, then (A, \cdot, \circ) is *Lazard* if both (A, \cdot) and (A, \circ) are Lazard.

Lemma 4.9. *Let (A, \cdot, \circ) be a skew brace. Then setting $A_0 = A$ and $A_i = L^i(A)$ for $i \geq 1$ makes (A, \cdot, \circ) into a filtered skew brace.*

Proof. The proof is almost word for word the same as Lemma 3.7. The only non-trivial part of the statement is that $[L^i(A), L^j(A)] \subseteq L^{i+j}(A)$ for all $i, j \geq 1$. Let $H = \{\lambda_a \mid a \in A\} \leq \text{Aut}(A, \cdot)$ and consider the semi-direct product $T = (A, \cdot) \rtimes_{\beta} H$ where $\beta : H \rightarrow \text{Aut}(A, \cdot)$ is the inclusion map. Then we find $[(a, \lambda_b), (c, 1)] = ([a, c] + b * c, 1)$. It follows that $[T, T] = L^2(A) \times [H, H]$. For a group G we define its lower central series as $\gamma^1(G) = G$ and $[\gamma^{i+1}(G) = [\mathfrak{g}, \gamma^i(G)]$. Then the previous equality can be rewritten as $\gamma^2(T) = L^2(A) \times \gamma^2(H)$ and by induction one finds $\gamma^i(T) = L^i(A) \times \gamma^i(H)$. It is well-known that $[\gamma^i(T), \gamma^j(T)] \subseteq \gamma^{i+j}(T)$, so in particular $([L^i(A), L^j(A)] \times \{0\}) = [L^i(\mathfrak{a}) \times \{0\}, L^j(\mathfrak{a}) \times \{0\}] \subseteq L^{i+j}(\mathfrak{a}) \times \gamma^{i+j}(\mathfrak{h})$ thus $[L^i(\mathfrak{a}), L^j(\mathfrak{a})] \subseteq L^{i+j}(\mathfrak{a})$. \square

5. A CORRESPONDENCE BETWEEN LAZARD POST-LIE RINGS AND LAZARD SKEW BRACES

Let \mathfrak{a} be a filtered Lie ring. We define $\text{Aut}(\mathfrak{a})$ as all filtered Lie ring automorphisms of \mathfrak{a} and $\text{Aut}(\mathfrak{a})_i = \text{Aut}(\mathfrak{a}) \cap \text{End}(\mathfrak{a})_i$ for $i \geq 0$.

Lemma 5.1. *Let \mathfrak{a} be a Lazard Lie ring. Then $\text{Aut}(\mathfrak{a})_i = \exp(\text{Der}(\mathfrak{a})_i)$ for all $i \geq 1$. In particular, $\exp : \mathbf{Laz}(\text{Der}(\mathfrak{a})^+) \rightarrow \text{Aut}(\mathfrak{a})^+$ is a group isomorphism and thus $\text{Aut}(\mathfrak{a})^+$ is a Lazard group.*

Proof. Clearly $\exp(\text{End}(\mathfrak{a})_i) = 1 + \text{End}(\mathfrak{a})_i$ for $i \geq 1$. Let $\delta \in \text{Der}(\mathfrak{a})_1$ and $x, y \in \mathfrak{a}$. By induction one finds that $\delta^n([a, b]) = \sum_{j=0}^n \binom{n}{j} [\delta^j(a), \delta^{n-j}(b)]$ for $n \geq 0$, hence

$$\begin{aligned} \exp(\delta)([a, b]) &= \sum_{n=0}^{\infty} \sum_{j=0}^i \frac{1}{n!} \binom{n}{j} [\delta^j(a), \delta^{n-j}(b)] = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{1}{i!j!} [\delta^j(a), \delta^i(b)] \\ &= [\exp(\delta)(a), \exp(\delta)(b)]. \end{aligned}$$

Conversely, let $\delta \in \text{End}(\mathfrak{a})_1$ such that $\exp(\delta) \in \text{Aut}(\mathfrak{a})$. We will prove by induction on k that

$$\delta([a, b]) \in [\delta(a), b] + [a, \delta(b)] + \mathfrak{a}_{i+j+k},$$

for all $a \in \mathfrak{a}_i, b \in \mathfrak{a}_j$ which implies that $\delta([a, b]) - [\delta(a), b] - [a, \delta(b)] \in \bigcap_{k=0}^{\infty} \mathfrak{a}_k = \{0\}$ and thus $\delta \in \text{Der}(\mathfrak{a})$. For $k = 1$, the statement is trivial. Assume that it holds for some $k \geq 1$. We prove by induction on n that

$$(5.1) \quad \delta^n([a, b]) \in \sum_{j=0}^n \binom{n}{j} [\delta^j(a), \delta^{n-j}(b)] + \mathfrak{a}_{i+j+k+1}$$

for $a \in \mathfrak{a}_i, b \in \mathfrak{a}_j$ and $n \geq 2$. By assumption,

$$\delta([a, b]) \in [\delta(a), b] + [a, \delta(b)] + \mathfrak{a}_{i+j+k},$$

so by applying δ on both sides we find

$$\delta^2([a, b]) \in \delta([\delta(a), b]) + \delta([a, \delta(b)]) + \mathfrak{a}_{i+j+k+1}.$$

By assumption,

$$\delta([\delta(a), b]) \in [\delta^2(a), b] + [\delta(a), \delta(b)] + \mathfrak{a}_{i+j+k+1},$$

$$\delta([a, \delta(b)]) \in [\delta(a), \delta^2(b)] + [a, \delta^2(b)] + \mathfrak{a}_{i+j+k+1},$$

so indeed (5.1) holds for $n = 2$. Similarly it follows that (5.1) holds for $n \geq 2$. We then use (5.1) to obtain

$$\exp(\delta)([a, b]) \in [a, b] + \delta([a, b]) + \sum_{n=2}^{\infty} \sum_{j=0}^n \frac{1}{j!(n-j)!} [\delta^j(a), \delta^{n-j}(b)] + \mathfrak{a}_{i+j+k+1}.$$

while also

$$[\exp(\delta)(a), \exp(\delta)(b)] = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{1}{i!j!} [\delta^j(a), \delta^i(b)]$$

Since $\exp(\delta)([a, b]) = [\exp(\delta)(a), \exp(\delta)(b)]$ we find $\delta([a, b]) \in [\delta(a), b] + [a, \delta(b)] + \mathfrak{a}_{i+j+k+1}$. We conclude that indeed $\exp(\text{Der}(\mathfrak{a})_i) = \text{Aut}(\mathfrak{a})_i$. The last part of the statement now follows from Proposition 2.5. \square

Lemma 5.2. *Let \mathfrak{a} be a Lazard Lie ring, then $\text{Aut}(\mathfrak{a}) = \text{Aut}(\mathbf{Laz}(\mathfrak{a}))$ as filtered groups.*

Proof. We know by Theorem 2.3 that automorphisms of \mathfrak{a} and $\mathbf{Laz}(\mathfrak{a})$ coincide. It remains to verify that the filtrations coincide. Let $x, y \in \mathfrak{a}$ such that $x+y \in \mathfrak{a}_i \setminus \mathfrak{a}_{i+1}$, then $[x, y] + \mathfrak{a}_{i+1} = [x, y] - [x+y, y] + \mathfrak{a}_{i+1} = \mathfrak{a}_{i+1}$, hence $x+y + \mathfrak{a}_{i+1} = \text{BCH}(x, y) + \mathfrak{a}_{i+1}$ and thus $\text{BCH}(x, y) \in \mathfrak{a}_i \setminus \mathfrak{a}_{i+1}$. We conclude that $x+y \in \mathfrak{a}_i$ if and only if $\text{BCH}(x, y) \in \mathfrak{a}_i$. In particular, for $\text{Aut}(\mathfrak{a})$ and $x \in \mathfrak{a}_i$ we find $\text{BCH}(\phi(x), -x) \in \mathfrak{a}_{i+j}$ if and only if $\phi(x) - x \in \mathfrak{a}_{i+j}$. We conclude that indeed $\text{Aut}(\mathfrak{a})_i = \text{Aut}(\mathbf{Laz}(\mathfrak{a}))_i$. \square

Proposition 5.3. *Let $\mathfrak{a}, \mathfrak{g}$ be Lazard lie rings and $\delta : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{a})$ a homomorphism of filtered Lie rings. Then there is a unique filtered group isomorphism*

$$\gamma : \mathbf{Laz}(\mathfrak{a} \oplus_{\delta} \mathfrak{g}) \rightarrow \mathbf{Laz}(\mathfrak{a}) \rtimes_{\lambda} \mathbf{Laz}(\mathfrak{g}),$$

such that $\gamma(a, 0) = (a, 0)$ and $\gamma(0, x) = (0, x)$, where $\lambda : \mathbf{Laz}(\mathfrak{g}) \rightarrow \text{Aut}(\mathbf{Laz}(\mathfrak{a}))$ is given by $\lambda_x = \exp(\delta_x)$.

Proof. Consider the split short exact sequence of Lazard Lie rings

$$0 \longrightarrow \mathfrak{a} \longrightarrow \mathfrak{a} \oplus_{\delta} \mathfrak{g} \xleftarrow{\quad} \mathfrak{g} \longrightarrow 0.$$

The functoriality of \mathbf{Laz} then yields a short exact sequence of Lazard groups

$$1 \longrightarrow \mathbf{Laz}(\mathfrak{a}) \longrightarrow \mathbf{Laz}(\mathfrak{a} \oplus_{\delta} \mathfrak{g}) \xleftarrow{\quad} \mathbf{Laz}(\mathfrak{g}) \longrightarrow 1.$$

Together with Lemma 2.7 this proves the statement. \square

We continue with the same setting as Proposition 5.3. Since $\mathfrak{a} \times \{0\}$ is a normal subgroup of both $\mathbf{Laz}(\mathfrak{a} \oplus_{\delta} \mathfrak{g})$ and $\mathbf{Laz}(\mathfrak{a}) \rtimes_{\lambda} \mathbf{Laz}(\mathfrak{g})$ and $\gamma(\mathfrak{a} \times \{0\}) = \mathfrak{a} \times \{0\}$, we find that γ does not affect the second component. Define $V : \mathfrak{a} \times \mathfrak{g} \rightarrow \mathfrak{a}$ as $\text{pr}_{\mathfrak{a}} \gamma$, thus $\gamma(a, x) = (V(a, x), x)$. Then in $\mathbf{Laz}(\mathfrak{a}) \rtimes_{\lambda} \mathbf{Laz}(\mathfrak{g})$ we find

$$(V(a, x), 0) = (V(a, x), x)(0, x)^{-1} = \gamma(a, x)\gamma(0, x)^{-1} = \gamma(\text{BCH}((a, x), (0, -x))),$$

from which we conclude that $V(a, x) = \text{pr}_{\mathfrak{a}} \text{BCH}((a, x), (0, -x))$. From (2.1) we find

$$(5.2) \quad V(a, x) = a + \frac{1}{2}\delta_x(a) + \frac{1}{6}\delta_x^2(a) + \frac{1}{12}[a, \delta_x(a)] + \frac{1}{24}(\delta_x([a, \delta_x(a)]) + \delta_x^3(a)) + \dots$$

where further terms are of order at least 5.

Similarly, we define $U : \mathfrak{a} \times \mathfrak{g} \rightarrow \mathfrak{a}$ as $U = \text{pr}_{\mathfrak{a}} \gamma^{-1}$ hence $\gamma^{-1}(a, x) = (U(a, x), x)$. We find $U(a, x) = \text{pr}_{\mathfrak{a}}(P((a, \lambda), (0, \lambda^{-1})))$ with P as in (2.2). Explicitly the terms of order less than 5 are given by

$$(5.3) \quad \begin{aligned} U(a, \lambda) &= a(a\lambda^{-1}(a)^{-1})^{-\frac{1}{2}}(a\lambda(a)a^{-2}\lambda^{-1}(a)a^{-1})^{\frac{1}{12}} \\ &\quad (a\lambda(a)\lambda^2(a)\lambda(a)^{-2}a\lambda(a)^{-1}\lambda^{-1}(a)^{-1}a^2\lambda(a^{-1})a^{-1})^{-\frac{1}{24}} \\ &\quad (\lambda^2(a)\lambda(a)^{-1}a\lambda(a)^{-1}a\lambda^{-1}(a)^{-1}a\lambda(a)^{-1})^{\frac{1}{24}} \dots \end{aligned}$$

In the case that \mathfrak{a} is abelian we have more explicit formulae.

Lemma 5.4. *Let $\mathfrak{a}, \mathfrak{g}$ be Lazard lie rings and $\delta : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{a})$ a homomorphism of filtered Lie rings. Let $V, U : \mathfrak{a} \times \mathfrak{g} \rightarrow \mathfrak{a}$ be as defined above. If \mathfrak{a} is abelian, then*

$$(5.4) \quad V(a, x) = \sum_{k=1}^{\infty} \frac{1}{k!} \delta_x^{k-1}(a),$$

$$(5.5) \quad U(a, x) = \sum_{k=2}^{\infty} \frac{1}{k} (-1)^{k+1} (\lambda_x - \text{id})^{k-1}(a).$$

Proof. Let $R = \text{End}(\mathfrak{a})^+$. Define $\rho : \mathfrak{a} \oplus_{\delta} \mathfrak{g} \rightarrow R$ as $\rho_{(a,x)}(n, b) = (0, na + \delta_x(b))$. By Proposition 2.5 we find that $\kappa : \mathbf{Laz}(\mathfrak{aff}(\mathfrak{a})^+) \rightarrow 1 + R$ with $\kappa = \exp \rho$ is a group homomorphism. Since $(V(a, x), 0) = \text{BCH}((a, x), (0, -x))$ it follows that $\exp(\rho_{\text{BCH}((a,x),(0,-x))}) = \exp(\rho_{(a,x)}) \exp(\rho_{(0,-x)})$. We now obtain (5.2) from

$$\begin{aligned} (1, V(a, \delta)) &= \exp(\rho_{\text{BCH}((a,x),(0,-x))})(1, 0) = \exp(\rho_{(a,x)}) \exp(\rho_{(0,-x)})(1, 0) \\ &= \exp(\rho_{(a,x)})(1, 0) = (1, \sum_{k=1}^{\infty} \frac{1}{k!} \delta_x^{k-1}(a)). \end{aligned}$$

Next we prove (5.3). Since κ is a group homomorphism, so is $f = \kappa \gamma^{-1}$. Thus $\kappa_{(U(a,x),x)} = f(a, x) = f(a, 0)f(0, \lambda) = \kappa_{(a,0)} \kappa_{(0,x)}$ and therefore

$$(0, U(a, x)) = \rho_{(U(a,x),x)}(1, 0) = \log(\kappa_{(a,0)} \kappa_{(0,x)})(1, 0).$$

We obtain (5.3) since $(\kappa_{(a,0)} \kappa_{(0,x)} - \text{id})^k(1, 0) = (0, (\lambda_x - \text{id})^{k-1}(a))$ for $k \geq 1$. \square

We now consider a particular case of the above. Let \mathfrak{a} be a Lazard lie ring, then by Proposition 5.3 we have a Lazard group isomorphism $\gamma : \mathbf{Laz}(\mathfrak{aff}(\mathfrak{a})^+) \rightarrow \mathbf{Laz}(\mathfrak{a}) \times \mathbf{Laz}(\text{Der}(\mathfrak{a})^+)$. Identifying $\mathbf{Laz}(\text{Der}(\mathfrak{a})^+)$ with $\text{Aut}(\mathfrak{a})^+$ as in Lemma 5.1 and $\text{Aut}(\mathfrak{a})^+$ with $\text{Aut}(\mathbf{Laz}(\mathfrak{a})^+)$ we find an isomorphism

$$\gamma : \mathbf{Laz}(\mathfrak{aff}(\mathfrak{a})^+) \rightarrow \text{Hol}(\mathbf{Laz}(\mathfrak{a})^+),$$

such that $\gamma(a, 0) = (a, 0)$ and $\gamma(0, \delta) = (0, \exp(\delta))$.

Proposition 5.5. *Let \mathfrak{a} be a Lazard Lie ring. Then γ yields a bijective correspondence between Lazard sub Lie rings \mathfrak{g} of $\mathfrak{aff}(\mathfrak{a})^+$ and Lazard subgroups G of $\text{Hol}(\mathbf{Laz}(\mathfrak{a})^+)$. Moreover, $\text{pr}_{\mathfrak{a}} : \mathfrak{g} \rightarrow \mathfrak{a}$ is injective, respectively surjective, if and only if $\text{pr}_{\mathfrak{a}} : \gamma(\mathfrak{g}) \rightarrow \mathfrak{a}$ is injective, respectively surjective.*

Proof. The first part follows directly from Theorem 2.3 and Proposition 5.3.

Assume that \mathfrak{g} is a Lazard sub Lie ring of $\mathfrak{aff}(\mathfrak{a})^+$ such that $\text{pr}_{\mathfrak{a}} : \mathfrak{g} \rightarrow \mathfrak{a}$ is not injective. Then there exists some non-zero element $(0, \delta) \in \mathfrak{g}$. But then clearly $\gamma(0, \delta) = (0, \exp(\delta)) \in \gamma(\mathfrak{g})$ and thus $\text{pr}_{\mathfrak{a}} : \gamma(\mathfrak{g}) \rightarrow \mathfrak{a}$ is not injective.

Conversely, assume that there exist $(a, \lambda), (a, \lambda') \in \gamma(\mathfrak{g})$ with $\lambda \neq \lambda'$. Then $(a, \lambda)^{-1}(a, \lambda') = (0, \lambda^{-1}\lambda') \in \gamma(\mathfrak{g})$. Hence $\gamma^{-1}(0, \lambda^{-1}\lambda') = (0, \log(\lambda^{-1}\lambda')) \in \mathfrak{g}$ and thus $\text{pr}_{\mathfrak{a}} : \mathfrak{g} \rightarrow \mathfrak{a}$ is not injective.

Assume that $\text{pr}_{\mathfrak{a}} : \mathfrak{g} \rightarrow \mathfrak{a}$ is surjective. Since pr_A and γ are continuous, the set $\text{pr}_{\mathfrak{a}} \gamma(\mathfrak{g}) = V(\mathfrak{g})$ is closed in \mathfrak{a} . Assume that there exists some $a \in \mathfrak{a} \setminus V(\mathfrak{g})$, then there exists a d such that $V(\mathfrak{g}) \cap (a + \mathfrak{a}_d) \neq \emptyset$ but $X \cap (a + \mathfrak{a}_{d+1}) = \emptyset$. Let $(b, \delta) \in \mathfrak{g}$ such that $V(b, \delta) \in a + \mathfrak{a}_d$. By assumption there exists some $\delta' \in \text{Der}(\mathfrak{a})^+$ such that $(-V(b, \delta) + a, \delta') \in \mathfrak{g}$. Since $-V(b, \delta) + a \in \mathfrak{a}_d$, we have

$$(5.6) \quad -V(b, \delta) + a + \mathfrak{a}_{d+1} = V(-V(b, \delta) + a, \delta') + \mathfrak{a}_{d+1} = \exp(\delta)(V(-V(b, \delta) + a, \delta')) + \mathfrak{a}_{d+1}.$$

Hence

$$\begin{aligned}
V(\text{BCH}((b, \delta), (-V(b, \delta) + a, \delta'))) + \mathfrak{a}_{d+1} \\
&= \text{BCH}(V(b, \delta), \exp(\delta)V(-V(b, \delta) + a, \delta')) + \mathfrak{a}_{d+1} \\
&= \text{BCH}(V(b, \delta), -V(b, \delta) + a) + \mathfrak{a}_{d+1} \\
&= a + \mathfrak{a}_{d+1},
\end{aligned}$$

where the first equality follows from the fact that γ is a group homomorphism. We conclude that $V(\mathfrak{g}) \cap (a + \mathfrak{a}_{d+1}) \neq \emptyset$ which is a contradiction.

Conversely, assume that $V(\mathfrak{g}) = \mathfrak{a}$. Clearly $\text{pr}_{\mathfrak{a}}(\mathfrak{g})$ is closed in \mathfrak{a} because $\text{pr}_{\mathfrak{a}}$ is continuous. Assume that there exists some $a \in \mathfrak{a} \setminus \text{pr}_{\mathfrak{a}}(\mathfrak{g})$ and let d be such that $\text{pr}_{\mathfrak{a}}(\mathfrak{g}) \cap (a + \mathfrak{a}_d) \neq \emptyset$ but $\text{pr}_{\mathfrak{a}}(\mathfrak{g}) \cap (a + \mathfrak{a}_{d+1}) = \emptyset$. Choose $(b, \delta) \in \mathfrak{g}$ such that $b \in a + \mathfrak{a}_d$. By assumption there exists some $(b', \delta') \in \mathfrak{g}$ such that $V(b', \delta') = a - b$ and using (5.6) we find $a - b = V(b', \delta') = b'$. We thus obtain the contradiction

$$\text{pr}_{\mathfrak{a}}((b, \delta) + (b', \delta')) + \mathfrak{a}_{d+1} = a + \mathfrak{a}_{d+1}. \quad \square$$

Remark 5.6. Transitive subgroups of the holomorph of a finite group play an essential role in the theory of Hopf–Galois structures, see for example [14, 24, 38]. One can prove that for any finite p -group A there exists a filtration of A such that $\text{Hol}(A)^+$ is a Sylow p -subgroup of the usual holomorph of A (just considered as a group without a filtration). If $|A| = p^k$ for $k < p$, then such a filtration has length at most p , so A is a Lazard group and so is any subgroup of $\text{Hol}(A)^+$. Hence in such a case, it follows from Proposition 5.5 that we can classify all regular p -groups of the holomorph of A up to conjugation by classifying the t -surjective sub Lie rings of $\text{aff}(\mathbf{Laz}^{-1}(A))^+$.

Proposition 5.7. *Let $(\mathfrak{a}, \triangleright)$ be a Lazard post-Lie ring. Then*

$$W : \mathfrak{a} \rightarrow \mathfrak{a} : a \mapsto V(a, \mathcal{L}_a)$$

is a bijective map and $\mathbf{S}(\mathfrak{a}, \triangleright) = (\mathfrak{a}, \cdot, \circ)$ with $a \cdot b = \text{BCH}(a, b)$, $a \circ b = a \cdot \exp(\mathcal{L}_{\Omega(a)})(b)$ is a Lazard skew brace, where $\Omega := W^{-1}$. Moreover, $W : \mathbf{Laz}(\mathfrak{a}^\circ) \rightarrow (\mathfrak{a}, \circ)$ is an isomorphism of filtered groups.

If \mathfrak{a} is abelian, then

$$W(a) = \sum_{k=1}^{\infty} \frac{1}{k!} \mathcal{L}_a^{k-1}(a).$$

Proof. Let $(\mathfrak{a}, \triangleright)$ be a Lazard post-Lie ring and let \mathfrak{g} be the regular Lazard sub Lie ring of $\text{aff}(\mathfrak{a})^+$ associated to $(\mathfrak{a}, \triangleright)$ as in Proposition 3.4. Then W is a bijection since it is the composition of the bijections $\mathfrak{a} \rightarrow \mathfrak{g} : a \mapsto (a, \mathcal{L}_a)$ and $V : \mathfrak{g} \rightarrow \mathfrak{a}$, recall that the latter is a bijection by Proposition 5.5. In the case that \mathfrak{a} is abelian, the explicit formula of W follows from Lemma 5.4.

By Proposition 5.5 we find that $\gamma(\mathfrak{g})$ is a regular Lazard subgroup of $\text{Hol}(\mathbf{Laz}(\mathfrak{a}))^+$, thus we obtain a Lazard skew brace $(\mathfrak{a}, \cdot, \circ)$ where $(\mathfrak{a}, \cdot) = \mathbf{Laz}(\mathfrak{a})$. Also $a \circ b = a \cdot \lambda_a(b)$ where $\lambda_a \in \text{Aut}(\mathfrak{a})$ is uniquely determined by the fact that $(a, \lambda_a) \in \gamma(\mathfrak{g})$. Since $\gamma(\mathfrak{a}) = \{(W(a), \exp(\mathcal{L}_a)) \mid a \in \mathfrak{a}\}$ we find $\lambda_a = \exp(\mathcal{L}_{\Omega(a)})$.

Also, $W : \mathbf{Laz}(\mathfrak{a}^\circ) \rightarrow (\mathfrak{a}, \circ)$ is a group isomorphism because it is the composition of the group homomorphisms $\mathbf{Laz}(\mathfrak{a}^\circ) \rightarrow \mathbf{Laz}(\mathfrak{g}) : a \mapsto (a, \mathcal{L}_a)$, $\gamma : \mathbf{Laz}(\mathfrak{g}) \rightarrow \gamma(\mathfrak{g})$ and $\text{pr}_{\mathfrak{a}} : \gamma(\mathfrak{g}) \rightarrow (\mathfrak{a}, \circ)$. \square

Note that in the case that \mathfrak{a} is abelian, one obtains the group of flows as in [2]. It is less trivial to see how the above construction is related to the formal integration as described by Bai, Guo, Sheng and Tang in [5]. Let \mathbb{F} be a field of characteristic

0. Then Bai, Guo, Sheng and Tang proved that to any connected complete post-Lie \mathbb{F} -algebra $(\mathfrak{a}, \triangleright)$ one can associate a skew brace $(\mathfrak{a}, \cdot, \circ)$ where

$$a \cdot b = \text{BCH}(a, b), \quad a \circ b = a \cdot \exp(\mathcal{L}_{\Omega'(a)})(b).$$

We first note that in our terminology their notion of a connected complete post-Lie \mathbb{F} -algebra is a Lazard post-Lie \mathbb{F} -algebra $(\mathfrak{a}, \triangleright)$ such that moreover $\mathfrak{a}_i \triangleright \mathfrak{a}_j \subseteq \mathfrak{a}_{i+j}$, or equivalently, such that $\mathfrak{a}_i^\circ = \mathfrak{a}_i$ for all $i \geq 0$. Note that under this extra assumption, if $\mathfrak{a}_i = 0$ for some i , then \mathfrak{a}_{i-1} is contained in the socle hence $\text{Soc}(\mathfrak{a}, \triangleright) \neq 0$. In [17, Section 3] an example is given of a left nilpotent pre-Lie \mathbb{R} -algebra with trivial socle, so it can be given a filtration which makes it a Lazard Lie \mathbb{R} -algebra but not a connected complete \mathbb{R} -algebra in the sense of Bai, Guo, Sheng and Tang.

We quickly recall their construction: Let $(\mathfrak{a}, \triangleright)$ be a connected complete post-Lie \mathbb{F} -algebra $(\mathfrak{a}, \triangleright)$ and consider the universal enveloping algebra $\mathcal{U}(\mathfrak{a})$ together with the filtration where $\mathcal{U}(\mathfrak{a})_0 = \mathcal{U}(\mathfrak{a})$ and $\mathcal{U}(\mathfrak{a})_i$ is the subspace generated by $x_1 \cdots x_n$ where $x_j \in \mathfrak{a}_{d_j}$ such that $d_1 + \cdots + d_n = i$. This makes $\mathcal{U}(\mathfrak{a})$ into a filtered \mathbb{F} -algebra. On $\mathcal{U}(\mathfrak{a})$ we define a new multiplication $*$ which for $a \in \mathfrak{a}$, $b \in \mathcal{U}(\mathfrak{a})$ is given by $a * b = ab + \delta_a(b)$, where δ_a is the unique derivation of $\mathcal{U}(\mathfrak{a})$ with $\delta_a(z) = a \triangleright z$ for $z \in \mathfrak{a}$. In this way, $(\mathcal{U}(\mathfrak{a}), *)$ becomes a filtered \mathbb{F} -algebra (where we keep the same filtration as before). This operation is then extended to the completion $\hat{\mathcal{U}}(\mathfrak{a})$ with respect to the filtration in order to obtain a complete \mathbb{F} -algebra $(\hat{\mathcal{U}}(\mathfrak{a}), *)$. For $a \in \mathfrak{a}$, we define $\Omega'(a) = \log_* \exp(a)$, where the exponential map with respect to the usual multiplication in $\hat{\mathcal{U}}(\mathfrak{a})$ is first applied, followed by the logarithmic map in $(\hat{\mathcal{U}}(\mathfrak{a}), *)$.

Recall that \mathfrak{a} embeds into $\hat{\mathcal{U}}(\mathfrak{a})$. One sees that $\rho : \mathfrak{aff}(\mathfrak{a}) \rightarrow \text{End}(\hat{\mathcal{U}}(\mathfrak{a}))$, given by $\rho_{(a, \delta)}(b) = ab + \delta(b)$ is a homomorphism of filtered Lie \mathbb{F} -algebras. Here we do not distinguish between a derivation δ of \mathfrak{a} and its unique extension to a derivation of $\hat{\mathcal{U}}(\mathfrak{a})$. Clearly, $a * b = \rho_{(a, \mathcal{L}_a)}(b)$ for all $a \in \mathfrak{a}$ and $b \in \hat{\mathcal{U}}(\mathfrak{a})$, thus $\exp_*(a) = \exp(\rho_{(a, \mathcal{L}_a)})(1)$. Since derivations map 1 to 0, also $\exp_*(a) = \exp(\rho_{(a, \mathcal{L}_a)}) \exp(\rho_{(0, -\mathcal{L}_a)})(1)$ where \exp_* denotes the exponential in $(\hat{\mathcal{U}}(\mathfrak{a}), *)$. By Proposition 2.5 and the definition of W , we find

$$\exp(\rho_{(a, \mathcal{L}_a)}) \exp(\rho_{(0, -\mathcal{L}_a)}) = \exp(\rho_{(W(a), 0)}),$$

thus $\exp_*(a) = \exp(W(a))$. We conclude that $\Omega = \Omega'$.

Proposition 5.8. *Let (A, \cdot, \circ) be a Lazard skew brace. Then*

$$\Omega : A \mapsto A : a \mapsto U(a, \lambda_a),$$

is a bijection and $\mathbf{L}(A, \cdot, \circ) = (\mathbf{Laz}^{-1}(A, \cdot), \triangleright)$ with $a \triangleright b = \log(\lambda_{W(a)})(b)$, is a Lazard Lie ring, where $W := \Omega^{-1}$. Moreover, $\Omega : (A, \circ) \rightarrow \mathfrak{a}^\circ$ is an isomorphism of filtered Lie rings, where $(\mathfrak{a}, \triangleright) = \mathbf{L}(A, \cdot)$.

If (A, \cdot) is abelian, then

$$\Omega(a) = \sum_{k=2}^{\infty} \frac{1}{k} (-1)^{k+1} (\lambda_a - \text{id})^{k-1}(a).$$

Proof. Let (A, \cdot, \circ) be Lazard skew brace and let G be the associated regular Lazard subgroup of $\text{Hol}(A, \cdot)^+$. Then Ω is a bijection since it is the composition of the bijections $A \mapsto G : a \mapsto (a, \lambda_a)$ and $U : G \rightarrow A$ with U . The fact that U is bijective, follows from Proposition 5.5.

By Proposition 5.5 we find that $\gamma^{-1}(G)$ is a Lazard sub Lie ring of $\mathfrak{aff}(\mathbf{Laz}^{-1}(A, \cdot))^+$, thus we obtain a Lazard post-Lie ring $(\mathbf{Laz}^{-1}(A, \cdot), \triangleright)$. We find $a \triangleright b = \mathcal{L}_a$ where \mathcal{L}_a is uniquely determined by the fact that $(a, \mathcal{L}_a) \in \gamma^{-1}(G)$. Since $\gamma^{-1}(G) = \{(\Omega(a), \log(\lambda_a)) \mid a \in A\}$ we find that $\mathcal{L}_a = \log(\lambda_{\Omega^{-1}(a)})$.

Also, Ω is $(A, \circ) \rightarrow \mathbf{L}(A, \cdot, \circ)^\circ$ is an isomorphism of filtered Lie rings since it is the composition of the isomorphisms $(A, \circ) \rightarrow G : a \mapsto (a, \lambda_a)$, $\gamma^1 : G \rightarrow \gamma^{-1}(G)$ and $\text{pr}_A : \gamma^{-1}(G) \rightarrow \mathbf{Laz}^{-1}(A, \cdot)$. \square

Theorem 5.9. *The constructions in Proposition 5.7 and Proposition 5.8 are mutually inverse and functorial. The map Ω associated to a Lazard post-Lie ring $(\mathfrak{a}, \triangleright)$ coincides with the map Ω associated to $\mathbf{S}(\mathfrak{a}, \triangleright)$.*

Lazard sub post-Lie rings of $(\mathfrak{a}, \triangleright)$ and Lazard sub skew braces of $\mathbf{S}(\mathfrak{a}, \triangleright)$ coincide and moreover, Lazard left ideals, Lazard strong left ideals and Lazard ideals of $(\mathfrak{a}, \triangleright)$ and $\mathbf{S}(\mathfrak{a}, \triangleright)$ coincide.

Proof. It is straightforward to verify that \mathbf{S} and \mathbf{L} are mutually inverse constructions and that their associated maps Ω coincide. We prove the functoriality of \mathbf{S} , the proof for \mathbf{L} is analogous. Let $f : (\mathfrak{a}, \triangleright) \rightarrow (\mathfrak{b}, \triangleright)$ be a homomorphism of Lazard post-Lie rings and set $(\mathfrak{a}, \cdot, \circ) = \mathbf{S}(\mathfrak{a}, \triangleright)$ and $(\mathfrak{b}, \cdot, \circ) = \mathbf{S}(\mathfrak{b}, \triangleright)$. Let $W_{\mathfrak{a}}$ and $W_{\mathfrak{b}}$ denote their respective maps W . Then by Theorem 2.3 $f : (\mathfrak{a}, \cdot) \rightarrow (\mathfrak{b}, \cdot)$ is a homomorphism of Lazard groups. As W is an isomorphism between $\mathbf{Laz}(\mathfrak{a}^\circ)$ and (\mathfrak{a}, \circ) , it is sufficient to prove $fW_{\mathfrak{a}} = W_{\mathfrak{b}}f$ in order to conclude that $f : (\mathfrak{a}, \circ) \rightarrow (\mathfrak{b}, \circ)$ is a group homomorphism. This equality follows direct from the fact that $f\mathcal{L}_a = \mathcal{L}_{f(a)}f$ for all $a \in \mathfrak{a}$.

The equivalence of Lazard post-Lie rings and Lazard sub skew braces is a direct corollary of Theorem 5.9. In particular, note that for \mathfrak{b} a Lazard sub post-Lie ring of $(\mathfrak{a}, \triangleright)$ we have $W_{\mathfrak{b}}(b) = W_{\mathfrak{a}}(b)$. In particular, $W_{\mathfrak{a}}(\mathfrak{b}) = \mathfrak{b}$.

Clearly the Lazard sub Lie rings of \mathfrak{a} invariant under $\{\mathcal{L}_x \mid x \in \mathfrak{a}\}$ and the ones invariant under $\{\exp(\mathcal{L}_x) \mid x \in \mathfrak{a}\}$ coincide, which yields the equivalence of Lazard left ideals. Taking into account Theorem 2.3, the equivalence of Lazard strong left ideals follows.

Let \mathfrak{b} be a Lazard strong left ideal of $(\mathfrak{a}, \triangleright)$. As $W : \mathbf{Laz}(\mathfrak{a}^\circ) \rightarrow (A, \circ)$ is a group isomorphism, we find that \mathfrak{b} is an ideal of \mathfrak{a}° if and only if $W(\mathfrak{b})$ is a normal subgroup of (A, \circ) . Since $W(\mathfrak{b}) = \mathfrak{b}$, the equivalence of Lazard ideals of $(\mathfrak{a}, \triangleright)$ and $\mathbf{S}(\mathfrak{a}, \triangleright)$ follows. \square

Remark 5.10. Recall that a skew brace $(A, +, \circ)$ is *square-free* if $a * a = 0$ for all $a \in A$. Similarly, we say that a post-Lie ring $(\mathfrak{a}, \triangleright)$ is *square-free* if $a \triangleright a = 0$ for all $a \in \mathfrak{a}$. Note that the property of being square-free is preserved by \mathbf{S} and \mathbf{L} and $W = \Omega = \text{id}$ in this case. Therefore these classes behave particularly well under the correspondence and eliminate most of the computations involved.

The construction \mathbf{L} coincides (under the necessary assumptions) with the geometric construction in [5, Theorem 4.3] (or in [9, Section 3]). We first recall this construction. Let (G, \cdot, \circ) be a skew brace such that (G, \cdot) and (G, \circ) are Lie groups and the map $a \mapsto (a, \lambda_a)$ is smooth (or equivalently, a post-Lie group). One defines a post-Lie structure on \mathfrak{g} , the Lie algebra associated to (G, \cdot) , by starting with the group homomorphism $\lambda : G \rightarrow \text{Aut}(G, \cdot) : a \mapsto \lambda_a$, which naturally yields a group homomorphism $\alpha : G \rightarrow \text{Aut}(\mathfrak{g}) : a \mapsto d\lambda_a$. Subsequently, differentiation yields a map $d\alpha : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{g})$ and we define $a \triangleright' b = (d\alpha)_a(b)$.

Let $(\mathfrak{a}, \cdot, \circ) = \mathbf{S}(\mathfrak{a}, \triangleright)$ for a finite dimensional Lazard Lie \mathbb{R} -algebra $(\mathfrak{a}, \triangleright)$. Then (\mathfrak{a}, \cdot) is a Lie group and its associated Lie algebra associated is isomorphic to \mathfrak{a} such that under this identification, the exponential map $\mathfrak{a} \rightarrow (\mathfrak{a}, \cdot)$ is just the identity. Thus, under this identification of (\mathfrak{a}, \cdot) and \mathfrak{a} we have $\lambda_g = \alpha_g$. Hence, $b \mapsto a \triangleright' b$ is the differential of the map $\lambda : a \mapsto \exp(\mathcal{L}_{\Omega(a)})(b)$. We first determine $d\Omega$. Since $\Omega W = \text{id}$, it is sufficient to determine dW . By (5.2), we find $dW = \text{id}$ hence also

$d\Omega = \text{id}$. The differential of the map $\exp(\mathcal{L}) : \mathfrak{a} \rightarrow \text{Aut}(\mathfrak{a}) : a \mapsto \exp(\mathcal{L}_a)$ in $a \in \mathfrak{a}$ is

$$d(\exp(\mathcal{L}))(a) = \left. \frac{d}{dt} \right|_{t=0} \exp(\mathcal{L}_{ta}) = \mathcal{L}_a.$$

We conclude that $(d\lambda)_a = \mathcal{L}_a$ and thus $a \triangleright b = a \triangleright b$, which proves the statement.

Example 5.11. Let A be a Lazard ring and let \mathcal{L}_a denote left multiplication by a . Then A^+ is a Lazard pre-Lie algebra for $a \triangleright b = ab$ and it is well-known that the associated group of flows is given by $a \circ b = a + ab + b$. We can verify this directly: clearly $W(a) = \exp(a) - 1$, hence $\Omega(a) = \log(1 + a)$. We now find

$$a \circ b = a + \exp(\mathcal{L}_{\log(1+a)})(b) = a + ab + b.$$

Conversely, starting from the brace $(A, +, \circ)$ we find $\Omega(a) = \log(1 + a)$ and thus $W(a) = \exp(a) - 1$. we recover the original multiplication as

$$a \triangleright b = \log(\lambda_{W(a)})(b) = \log(\mathcal{L}_{1+W(a)})(b) = a + \log(\mathcal{L}_{\exp(a)})(b) = ab.$$

Recall that this is precisely the classical correspondence between two-sided braces and Jacobson radical rings [30].

We are now ready to prove Theorem 1.5.

Proof of Theorem 1.5. Let $(\mathfrak{a}, \triangleright)$ be a post-Lie ring such that $(\mathfrak{a}, +)$ is a p -group of L -nilpotency class $k < p$. With its canonical filtration $(\mathfrak{a}, \triangleright)$ is a Lazard post-Lie ring since both \mathfrak{a} and \mathfrak{a}° are p -groups with $\mathfrak{a}_p = \mathfrak{a}_p^\circ = \{0\}$. By Theorem 5.9 we obtain a Lazard skew p -brace of L -nilpotency class less than p .

Conversely, let (A, \cdot, \circ) be a skew p -brace of L -nilpotency class $k < p$. Then with its canonical filtration it becomes a Lazard skew brace since (A, \cdot) and (A, \circ) are p -groups and $A_p = (A, \circ)_p = \{1\}$. BY Theorem 5.9 we obtain a Lazard post-Lie ring of L -nilpotency class less than p whose additive group is a p -group.

At last, recall that any skew brace of order p^k is left nilpotent by [13, Proposition 4.4] hence it is L -nilpotent. The L -nilpotency class of a skew brace or post-Lie ring of order p^k clearly can not exceed k . \square

Let (A, \cdot, \circ) be a skew brace and $a \in A$. For an integer n we denote the n -th power of a in (A, \cdot) by a^n and its n -th power in (A, \circ) is denoted by $a^{\circ n}$. The following result extends [37, Proposition 15, Lemma 17].

Proposition 5.12. *Let (A, \cdot, \circ) be a Lazard skew brace and let n be an integer, then $\{a^n \mid a \in A\} = \{a^{\circ n} \mid a \in A\}$ and $\{a \in A \mid a^n = 1\} = \{a \in A \mid a^{\circ n} = 1\}$, and these sets are Lazard ideals of (A, \cdot, \circ) .*

Proof. We prove the statement for the first set, the second part is analogous. Let $(\mathfrak{a}, \triangleright) = \mathbf{L}(A, \cdot, \circ)$. Then it is straightforward to see that $n\mathfrak{a}$ is a Lazard ideal of $(\mathfrak{a}, \triangleright)$, hence it is also a Lazard ideal of (A, \cdot, \circ) by Theorem 5.9. Since $(A, \cdot) = \mathbf{Laz}(\mathfrak{a})$, we find $n\mathfrak{a} = \{a^n \mid a \in A\}$. Also, as $W : \mathbf{Laz}(\mathfrak{a}^\circ) \rightarrow (A, \circ)$ is a group isomorphism we find $W(n\mathfrak{a}) = \{a^{\circ n} \mid a \in A\}$, but since W maps any Lazard sub post-Lie ring to itself, we conclude that $\{a^n \mid a \in A\} = \{a^{\circ n} \mid a \in A\}$. \square

Proposition 5.13. *Let $(\mathfrak{a}, \triangleright)$ be a Lazard post-Lie ring. Then $\text{Fix}(\mathfrak{a}, \triangleright) = \text{Fix}(\mathbf{S}(\mathfrak{a}, \triangleright))$, $\text{Soc}(\mathfrak{a}, \triangleright) = \text{Soc}(\mathbf{S}(\mathfrak{a}, \triangleright))$ and $\text{Ann}(\mathfrak{a}, \triangleright) = \text{Ann}(\mathbf{S}(\mathfrak{a}, \triangleright))$. In particular, if \mathfrak{a} is nilpotent, then $(\mathfrak{a}, \triangleright)$ is right nilpotent if and only if $\mathbf{S}(\mathfrak{a}, \triangleright)$ is right nilpotent.*

Proof. The first part of the statement follows from the fact that $W(a) = a$ whenever $a \triangleright a = 0$, which in particular is the case when $a \in \text{Fix}(\mathfrak{a}, \triangleright)$ or $a \in \text{Soc}(\mathfrak{a}, \triangleright)$. The second part of the statement now follows from [1, Lemma 5.3 and Lemma 5.4] and the analogous statement for post-Lie rings. \square

6. FORMAL DIFFERENTIATION THROUGH ROOTS OF UNITY

Let $n \geq 1$. Recall that a *primitive n -th root of unity* of a ring R is an element $\xi \in R$ such that $\xi^n = 1$ and $\xi^k - 1$ not a zero divisor for $1 \leq k < n$.

Lemma 6.1. *Let R be a ring, let M, N be R -modules, $\xi \in R$ a primitive n -th root of unity where n is invertible in R , and $f : M \rightarrow N$ a map such that for every $m \in M$ there exist elements $\alpha_0, \dots, \alpha_d \in N$, with $d < n$, such that*

$$f(rm) = \alpha_0 + r\alpha_1 + r^2\alpha_2 + \dots + r^d\alpha_d.$$

Then $\frac{1}{n} \sum_{i=0}^{n-1} \xi^{ik} f(\xi^{-i}m) = \alpha_k$ for all $0 \leq k \leq d$.

Proof. Evaluating the formula yields

$$\frac{1}{n} \sum_{i=0}^{n-1} \xi^{ik} f(\xi^{-i}m) = \frac{1}{n} \sum_{i=0}^{n-1} \sum_{j=0}^d \xi^{i(k-j)} \alpha_j = f_k$$

since $\sum_{i=0}^{n-1} \xi^{i(k-j)} f_j$ equals 0 for $j \neq k$ and it equals $n f_k$ for $j = k$. \square

For $k = 1$, it makes sense to see the above result as a discrete version of taking the directional derivative through m and then evaluating at 0. Therefore, it seems justified to introduce the notation

$$df : M \rightarrow N : m \mapsto \frac{1}{n} \sum_{i=0}^{n-1} \xi^i f(\xi^{-i}m).$$

Although the analogy is not explicitly mentioned, this technique plays a prominent role in [34, 36]. We can extend the formula of [36, Theorem 12] to the skew brace setting.

Proposition 6.2. *Let p^k be a prime power, $\xi \in \mathbb{Z}$ a primitive $(p-1)$ -th root of unity modulo p^k . Let (A, \cdot, \circ) be a filtered skew brace of order p^k such that $A_p = \{1\}$ and such that $A_i * A_j \subseteq A_{i+j}$ for all $i, j \geq 1$. Then*

$$a \triangleright b = \sum_{i=0}^{p-2} \xi^i \lambda_{\xi^{-i}a}(b),$$

for all $a, b \in A$ where the sum is taken in $\mathbf{Laz}^{-1}(A, \cdot)$.

Proof. It is clear that (A, \cdot, \circ) with the given filtration is a Lazard skew brace. By Theorem 5.9 we know that $(A, \cdot, \circ) = \mathbf{S}(\mathfrak{a}, \triangleright)$ for some Lazard post-Lie ring $(\mathfrak{a}, \triangleright)$. In particular, \mathfrak{a} has a natural structure of a $\mathbb{Z}/p^k\mathbb{Z}$ -module since $|\mathfrak{a}| = p^k$. Consider the map $\lambda : \mathfrak{a} \rightarrow \text{End}(\mathfrak{a}) : a \mapsto \lambda_a$. From Proposition 5.7 we find

$$\lambda_{ta} = \exp(\mathcal{L}_{W(ta)}) = \sum_{k=0}^{\infty} \sum_{i=1}^{\infty} \frac{1}{k!i!} \mathcal{L}_{\mathcal{L}_{ta}^{i-1}(ta)}^k = \sum_{k=0}^{\infty} \sum_{i=1}^{\infty} \frac{1}{k!i!} t^{ki} \mathcal{L}_{\mathcal{L}_a^{i-1}(a)}^k,$$

but by the assumption on the filtration we find that all powers t^j disappear for $j \geq p$ and the degree 1 coefficient is \mathcal{L}_a , hence the assumptions for Lemma 6.1 are satisfied and we find $d\lambda(a) = \mathcal{L}_a$ which concludes the proof. \square

Note that a filtration as in Proposition 6.2 exists precisely if $A^{\{p\}} = \{1\}$ where $A^{\{1\}} = A$ and $A^{\{k+1\}}$ is the subgroup of (A, \cdot) generated by all $a * b$ and $aba^{-1}b^{-1}$ where $a \in A^{\{i\}}$, $b \in A^{\{k+1-i\}}$, $1 \leq i \leq k$. For braces this is precisely the chain of ideals $A^{[k]}$ used to define strong nilpotency as in [33].

We remark a potential advantage of the above way of differentiating when working over characteristic p^k ; namely that in Lemma 6.1 we can have terms of degree more than p , whereas in the formulas for exp and log these terms are not allowed since the coefficients of degree p or more do not exist. As for example the field

\mathbb{F}_{p^n} has a primitive $(p^n - 1)$ -th root of unity. In particular, this means that there are possible uses in the case $p = 2$. It would be very interesting to see whether this idea of differentiation indeed has some meaningful application in such cases. In particular, the following question is natural to ask:

Question 6.3. Let $(A, +, \circ)$ be an \mathbb{F}_{p^n} -brace (as in [11, Definition 2]) such that $A^{[p^n-1]} = 1$ and let $\xi \in \mathbb{F}_{p^n}$ be a primitive $(p^n - 1)$ -th root of unity. Does the operation

$$a \triangleright b = - \sum_{i=0}^{p^n-2} \xi^i \lambda_a(\xi^{-i} b),$$

yield a pre-Lie \mathbb{F}_{p^n} -algebra structure on the abelian Lie \mathbb{F}_{p^n} -algebra $(A, +)$?

ACKNOWLEDGEMENTS

The author was supported by Fonds Wetenschappelijk Onderzoek – Vlaanderen, via a PhD Fellowship fundamental research (1160524N) and a Senior Research Project (G004124N), and by Vrije Universiteit Brussel via the project OZR3762.

REFERENCES

- [1] E. Acri, R. Lutowski, and L. Vendramin. Retractability of solutions to the Yang–Baxter equation and p -nilpotency of skew braces. *Internat. J. Algebra Comput.*, 30(01):91–115, 2020.
- [2] A. A. Agrachev and R. V. Gamkrelidze. Chronological Algebras and Nonstationary Vector Fields. *J. Sov. Math.*, 17:1650–1675, 1981.
- [3] L. Auslander. Simply transitive groups of affine motions. *Amer. J. Math.*, 99(4):809–826, 1977.
- [4] D. Bachiller. Counterexample to a conjecture about braces. *Journal of Algebra*, 453:160–176, 2016.
- [5] C. Bai, L. Guo, Y. Sheng, and R. Tang. Post-groups, (Lie-)Butcher groups and the Yang–Baxter equation. *Mathematische Annalen*, 2023.
- [6] Y. Benoist. Une nilvariété non affine. *J. Differential Geom.*, 41(1):21–52, 1995.
- [7] D. Burde. Affine structures on nilmanifolds. *Internat. J. Math.*, 7(5):599–616, 1996.
- [8] D. Burde. Left-symmetric algebras, or pre-lie algebras in geometry and physics. *Open Mathematics*, 4(3):323–357, 2006.
- [9] D. Burde, K. Dekimpe, and S. Deschamps. Affine actions on nilpotent Lie groups. *Forum Mathematicum*, 21(5):921–934, 2009.
- [10] D. Burde, K. Dekimpe, and K. Vercammen. Affine actions on Lie groups and post-Lie algebra structures. *Linear Algebra and Its Applications*, 437(5):1250–1263, 2012.
- [11] F. Catino, I. Colazzo, and P. Stefanelli. Skew left braces with non-trivial annihilator. *Journal of Algebra and Its Applications*, 18(02):1950033, 2019.
- [12] Cayley. On the Analytical Forms Called Trees. *Amer. J. Math.*, 4(1-4):266–268, 1881.
- [13] F. Cedó, A. Smoktunowicz, and L. Vendramin. Skew left braces of nilpotent type. *Proceedings of the London Mathematical Society*, 118(6):1367–1392, 2019.
- [14] L. N. Childs. On the Hopf Galois theory for separable field extensions. *Communications in Algebra*, 17(4):809–825, 1989.
- [15] S. Cicalò, W. A. De Graaf, and M. Vaughan-Lee. An effective version of the Lazard correspondence. *Journal of Algebra*, 352(1):430–450, 2012.
- [16] C. Curry, K. Ebrahimi-Fard, and H. Munthe-Kaas. What is a post-Lie algebra and why is it useful in geometric integration. In F. A. Radu, K. Kumar, I. Berre, J. M. Nordbotten, and I. S. Pop, editors, *Numerical Mathematics and Advanced Applications ENUMATH 2017*, pages 429–437, Cham, 2019. Springer International Publishing.
- [17] D. Fried. Distality, completeness, and affine structures. *Journal of Differential Geometry*, 24(3):265–273, 1986.
- [18] L. Guarnieri and L. Vendramin. Skew braces and the Yang–Baxter equation. *Math. Comp.*, 86(307):2519–2534, 2017.
- [19] E. I. Khukhro. *p-Automorphisms of Finite p-Groups*. London Mathematical Society Lecture Note Series. Cambridge University Press, 1998.
- [20] H. Kim. Complete left-invariant affine structures on nilpotent Lie groups. *Journal of Differential Geometry*, 24(3):373–394, 1986.
- [21] J.-L. Koszul. Domaines bornés homogènes et orbites de groupes de transformations affines. *Bull. Soc. Math. France*, 89:515–533, 1961.

- [22] M. Lazard. Sur les groupes nilpotents et les anneaux de Lie. *Annales scientifiques de l'École normale supérieure*, 71(2):101–190, 1954.
- [23] A. I. Malcev. Nilpotent torsion-free groups. *Izv. Akad. Nauk SSSR Ser. Mat.*, 13:201–212, 1949.
- [24] I. Martin-Lyons and P. J. Truman. Skew braceoids. *Journal of Algebra*, 638:751–787, 2024.
- [25] J. Milnor. On fundamental groups of complete affinely flat manifolds. *Advances in Math.*, 25(2):178–187, 1977.
- [26] M. F. Newman, E. A. O'Brien, and M. R. Vaughan-Lee. Groups and nilpotent Lie rings whose order is the sixth power of a prime. *Journal of Algebra*, 278(1):383–401, 2004.
- [27] D. Puljić. Classification of braces of cardinality p^4 . pages 1–18, 2022.
- [28] D. Puljić, A. Smoktunowicz, and K. Nejabati Zenouz. Some braces of cardinality p^4 and related Hopf–Galois extensions. *New York Journal of Mathematics*, 28:494–522, 2022.
- [29] W. Rump. Modules over braces. *Algebra Discrete Math.*, -(2):127–137, 2006.
- [30] W. Rump. Braces, radical rings, and the quantum Yang–Baxter equation. *J. Algebra*, 307(1):153–170, 2007.
- [31] W. Rump. The brace of a classical group. *Note di Matematica*, 34(1):115–144, 2014.
- [32] D. Segal. The structure of complete left-symmetric algebras. *Mathematische Annalen*, 293(1):569–578, 1992.
- [33] A. Smoktunowicz. On Engel groups, nilpotent groups, rings, braces and the Yang–Baxter equation. *Trans. Amer. Math.*, 370(9):6535–6564, 2018.
- [34] A. Smoktunowicz. A new formula for Lazard's correspondence for finite braces and pre-Lie algebras. *Journal of Algebra*, 594:202–229, 2022.
- [35] A. Smoktunowicz. From pre-Lie rings back to braces. pages 1–24, 2022.
- [36] A. Smoktunowicz. On the passage from finite braces to pre-Lie rings. *Advances in Mathematics*, 409:108683, 2022.
- [37] A. Smoktunowicz. On the passage from finite braces to pre-Lie rings. *arXiv:2202.00085v3*, 2022.
- [38] L. Stefanello and S. Trappeniers. On the connection between Hopf–Galois structures and skew braces. *Bulletin of the London Mathematical Society*, 2023.
- [39] B. Vallette. Homology of generalized partition posets. *Journal of Pure and Applied Algebra*, 208(2):699–725, 2007.
- [40] E. B. Vinberg. The theory of homogeneous convex cones. *Trudy Moskov. Mat. Obsč.*, 12:303–358, 1963.

DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSELS, BELGIUM

Email address: senne.trappeniers@vub.be

URL: <https://sites.google.com/view/sennetrappeniers/>