

Around the codifferential of products of p -forms and generalized interior products

E. Huguet,¹ J. Queva,² and J. Renaud³

¹*Université Paris Cité, APC-Astroparticule et Cosmologie (UMR-CNRS 7164),
Batiment Condorcet, 10 rue Alice Domon et Léonie Duquet, F-75205 Paris Cedex 13,
France.*

²*Université de Corse – CNRS UMR 6134 SPE, Campus Grimaldi BP 52, 20250 Corte,
France.*

³*Université Gustave Eiffel, APC-Astroparticule et Cosmologie (UMR-CNRS 7164),
Batiment Condorcet, 10 rue Alice Domon et Léonie Duquet, F-75205 Paris Cedex 13,
France.*

(*Electronic mail: jacques.renaud@u-pem.fr)

(*Electronic mail: queva@univ-corse.fr)

(*Electronic mail: huguet@apc.univ-paris7.fr)

(Dated: 18 July 2024)

Identities pertaining to the de Rham codifferential δ in differential geometry are scattered in the literature. This article gathers such formulas involving usual differential operators (Lie derivative, Schouten-Nijenhuis bracket, ...), while adding a few new ones using a natural extension of the interior product, to provide a compact handy summary.

I. INTRODUCTION

Computations with p -forms involve differential operators such as the exterior derivatives d and the Lie derivatives \mathcal{L}_v , both of which fulfilling a Leibniz identity with respect to the exterior product \wedge . In stark contrast is the codifferential δ which is built out of d and the Hodge operator $*$ and for which there is no such identity. However, some other identities involving δ can be found scattered in the literature, often in a wider setting. This short paper aims to gather such formulas in one compact space making it a useful toolbox for the practitioners of the field. Those formulas are extended further using a generalized interior product, producing new (up to our knowledge) formulas such as Eqs. (60)–(64). Our interest in such formulas came as a by-product of our studies on restrictions of differential operators from flat ambient space to cosmological spaces, more precisely computations needed for the generalization to p -form of earlier work¹.

Section II deals with the relationship between the codifferential δ and the Schouten-Nijenhuis bracket brought back to forms, facts known for a long time once cast in the framework of Gerstenhaber algebras. Some explicit and little known formulas are gleaned along the way. In Sec. III the interior product i_v is generalized in a somewhat natural manner, it provides a set of new identities of their own and in peculiar it involves naturally the Schouten-Nijenhuis bracket. A set of noteworthy formulas involving the differential operators d , \mathcal{L}_v , δ and \square are recorded. Our conventions regarding differential geometry follow those of Ref. 2 and needed formulas are briefly recalled in appendix A. The outline of some longer calculations are gathered in appendix B in order not to clutter the main body of the paper.

Notation. The usual interior and exterior product i_v and j_v , see App. A, are extended to 1-forms in a natural way. For $\lambda \in \Omega^1$, $\beta \in \Omega^b$ we note

$$i^\lambda \beta = i_{\sharp \lambda} \beta, \quad (1)$$

$$j^\lambda \beta = j_{\sharp \lambda} \beta = \lambda \wedge \beta. \quad (2)$$

They correspond to each other through the Hodge operator

$$i^\lambda \beta = (-1)^{b+1} *^{-1} j^\lambda * \beta. \quad (3)$$

These products will be further generalized in Sec. III with definition 2.

Notation. For ease of use, given an integer n , the (anti-)commutator between two operators X and Y is written as

$$[X, Y]_n = XY - (-1)^n YX, \quad (4)$$

recovering the commutator for even values of n and anticommutator for odd values. When there is no ambiguity we set, as usual, $[X, Y] = [X, Y]_0 = XY - YX$ and $[X, Y]_+ = [X, Y]_1 = XY + YX$.

II. THE SCHOUTEN-NIJENHUIS BRACKET ON FORMS AND ITS PROPERTIES

Schouten^{3,4}, and then Schouten-Nijenhuis^{5,6}, bracket came from the natural endeavor in differential geometry of generalizing the Lie derivative not just along vector but also along tensors. In particular it is the unique extension generalizing the Lie derivative on multivectors^{7,8} (i.e. the sections of $\wedge^p TM$). In addition to this, it fulfills a wealth of algebraic relations making it one of the simplest incarnation of a Gerstenhaber algebra⁹⁻¹¹. In this section we expose how, once brought back to forms, it arises out of the codifferential of the product of two forms. The bracket can be viewed from many angles^{12,13}, in particular it plays a key role in Poisson geometry^{14,15}. For additional material regarding Schouten brackets and its generalizations we refer the reader to Ref. 16 for a beautiful overview.

Definition 1. The Schouten-Nijenhuis bracket¹⁷, brought back to forms, is the unique bilinear application fulfilling:

$$\llbracket \alpha, \beta \rrbracket = -(-1)^{(a-1)(b-1)} \llbracket \beta, \alpha \rrbracket \in \Omega^{a+b-1}, \quad \alpha \in \Omega^a, \beta \in \Omega^b, \quad (5)$$

$$\llbracket \alpha, \beta \wedge \gamma \rrbracket = \llbracket \alpha, \beta \rrbracket \wedge \gamma + (-1)^{(a-1)b} \beta \wedge \llbracket \alpha, \gamma \rrbracket, \quad \alpha \in \Omega^a, \beta \in \Omega^b, \gamma \in \Omega^c, \quad (6)$$

$$\llbracket f, g \rrbracket = 0, \quad f, g \in \Omega^0, \quad (7)$$

$$\llbracket \lambda, \beta \rrbracket = \flat(\mathcal{L}_{\sharp\lambda} \sharp\beta), \quad \lambda \in \Omega^1, \beta \in \Omega^b. \quad (8)$$

The ‘‘usual’’ Schouten-Nijenhuis bracket is naturally defined on multivectors, it is lowered to forms using the musical applications \flat and \sharp . Nevertheless, this bracket has some particularities compared to the usual one due to the Lie derivative acting on the \sharp operator. In fact Eq. (8) leads to

$$\llbracket \lambda, \beta \rrbracket = (\mathcal{L}_{\sharp\lambda} \tilde{g})^{mn} j_m i_n \beta + \mathcal{L}_{\sharp\lambda} \beta, \quad (9)$$

where β is a b -form, \tilde{g} is the corresponding metrics on forms and j_m, i_n stand for j_{e_m} and i_{e_n} . When $\sharp\lambda$ is a Killing vector it simplifies to $\llbracket \lambda, \beta \rrbracket = \mathcal{L}_{\sharp\lambda} \beta$.

Property 1. The Schouten-Nijenhuis bracket fulfills a graded Jacobi identity^{5,17,18}

$$(-1)^{(a-1)(c-1)} \llbracket \alpha, \llbracket \beta, \gamma \rrbracket \rrbracket + (-1)^{(b-1)(a-1)} \llbracket \beta, \llbracket \gamma, \alpha \rrbracket \rrbracket + (-1)^{(c-1)(b-1)} \llbracket \gamma, \llbracket \alpha, \beta \rrbracket \rrbracket = 0. \quad (10)$$

Equations (5), (6) and (10), that is graded antisymmetry, graded Leibniz identity and graded Jacobi identity, makes the bracket a Gerstenhaber algebra. As usual, the graded Jacobi identity can be written in the suggestive form

$$\llbracket \alpha, \llbracket \beta, \gamma \rrbracket \rrbracket = \llbracket \llbracket \alpha, \beta \rrbracket, \gamma \rrbracket + (-1)^{(a-1)(b-1)} \llbracket \beta, \llbracket \alpha, \gamma \rrbracket \rrbracket, \quad (11)$$

making $\llbracket \alpha, \cdot \rrbracket$ a derivative of degree $a - 1$ with respect to Ω equipped with the Schouten-Nijenhuis bracket (the ‘‘Schouten degree’’ of a form being its exterior degree shifted by minus one).

Remark. For $\alpha \in \Omega^a$, $\beta \in \Omega^b$ and $\gamma \in \Omega^c$ through elementary algebra one gets, in order to expand the bracket with respect to the left argument, the following useful identity

$$\llbracket \alpha \wedge \beta, \gamma \rrbracket = \alpha \wedge \llbracket \beta, \gamma \rrbracket + (-1)^{ab} \beta \wedge \llbracket \alpha, \gamma \rrbracket. \quad (12)$$

For $\phi \in \Omega^0$ a function and $\lambda \in \Omega^1$ a 1-form notice that:

$$\llbracket \phi, \lambda \rrbracket = -\llbracket \lambda, \phi \rrbracket = -\mathcal{L}_{\sharp\lambda} \phi = -i^\lambda d\phi = -(\lambda, d\phi) = -i^{d\phi} \lambda, \quad (13)$$

which extends by induction on the degree to

$$\llbracket \phi, \alpha \rrbracket = -i^{d\phi} \alpha = (-1)^a \llbracket \alpha, \phi \rrbracket, \quad \phi \in \Omega^0, \alpha \in \Omega^a. \quad (14)$$

Property 2. Let $\alpha \in \Omega^a$ and $\beta \in \Omega^b$ be two forms, and e^k a co-frame, setting $c = a + b - 1$, then:

$$\begin{aligned} \llbracket \alpha, \beta \rrbracket &= \frac{1}{a!b!} \left\{ a\alpha_{k_1 \dots k_{a-1}r}(d\beta_{k_a \dots k_c}, e^r) - b(d\alpha_{k_1 \dots k_a}, e^r)\beta_{k_{a+1} \dots k_{c+r}} \right. \\ &\quad \left. + ab\alpha_{k_1 \dots k_{a-1}r}\beta_{k_a \dots k_{c-1}s}\tilde{c}_{k_c}^{rs} \right\} e^{k_1} \wedge \dots \wedge e^{k_c}, \end{aligned} \quad (15)$$

where $\tilde{c}_k^{rs} = g^{ri}g^{sj}g_{kl}c_{ij}^l$ and c are the anholonomy coefficients of the frame.

Remark. The last term in Eq. (15) vanishes as soon as the frame is an holonomic one.

Proof. Straightforward expanding first α , using Eq. (12), bringing:

$$\llbracket \alpha, \beta \rrbracket = \frac{1}{a!} \left\{ e^{m_1} \wedge \dots \wedge e^{m_a} \wedge \llbracket \alpha_{m_1 \dots m_a}, \beta \rrbracket + a\alpha_{m_1 \dots m_a} e^{m_1} \wedge \dots \wedge e^{m_{a-1}} \wedge \llbracket e^{m_a}, \beta \rrbracket \right\}, \quad (16)$$

using the skew-symmetry of the coefficient. Then, permuting the arguments in the brackets and using again this formula, gives rise to:

$$\begin{aligned} \llbracket \alpha, \beta \rrbracket &= \frac{1}{a!b!} \left\{ -(-1)^b b \llbracket \alpha_{m_1 \dots m_a}, e^{n_b} \rrbracket \beta_{n_1 \dots n_b} e^{m_1} \wedge \dots \wedge e^{m_a} \wedge e^{n_1} \wedge \dots \wedge e^{n_{b-1}} \right. \\ &\quad - a\alpha_{m_1 \dots m_a} \llbracket \beta_{n_1 \dots n_b}, e^{m_a} \rrbracket e^{m_1} \wedge \dots \wedge e^{m_{a-1}} \wedge e^{n_1} \wedge \dots \wedge e^{n_b} \\ &\quad \left. + ab\alpha_{m_1 \dots m_a} \beta_{n_1 \dots n_b} e^{m_1} \wedge \dots \wedge e^{m_{a-1}} \wedge e^{n_1} \wedge \dots \wedge e^{n_{b-1}} \wedge \llbracket e^{m_a}, e^{n_b} \rrbracket \right\}. \end{aligned} \quad (17)$$

Shuffling the indices and relabeling them in a straightforward manner, with $c = a + b - 1$, simplifies to

$$\begin{aligned} \llbracket \alpha, \beta \rrbracket &= \frac{1}{a!b!} \left\{ b \llbracket \alpha_{k_1 \dots k_a}, e^r \rrbracket \beta_{k_{a+1} \dots k_{c+r}} - a\alpha_{k_1 \dots k_{a-1}r} \llbracket \beta_{k_a \dots k_c}, e^r \rrbracket \right. \\ &\quad \left. + ab\alpha_{k_1 \dots k_{a-1}r}\beta_{k_a \dots k_{c-1}s} (\llbracket e^r, e^s \rrbracket)_{k_c} \right\} e^{k_1} \wedge \dots \wedge e^{k_c}. \end{aligned} \quad (18)$$

Using, in addition, Eq. (13) to express $\llbracket \alpha_{k_1 \dots k_a}, e^r \rrbracket$ and $\llbracket \beta_{k_a \dots k_c}, e^r \rrbracket$ and that $\llbracket e^r, e^s \rrbracket = b\mathcal{L}_{\#e^r}\#e^s = \tilde{c}_k^{rs}e^k$ gives the final formula. \square

The Schouten-Nijenhuis bracket is the measure of the failure of δ to be a derivation with respect to the exterior product \wedge , which is embodied in the following (already known) theorem for which we provide a pedestrian proof.

Theorem 1. Let $\alpha \in \Omega^a$ and $\beta \in \Omega^b$, then:

$$\delta(\alpha \wedge \beta) = (\delta\alpha) \wedge \beta + (-1)^a \alpha \wedge (\delta\beta) + (-1)^a \llbracket \alpha, \beta \rrbracket. \quad (19)$$

Proof. First let $\alpha = \phi \in \Omega^0$ be a function ($a = 0$) and $\beta \in \Omega^b$ a b -form, then:

$$\delta(\phi\beta) = \delta(\phi \wedge \beta) = \phi(\delta\beta) + \llbracket \phi, \beta \rrbracket, \quad (20)$$

which follows from the definition of the codifferential $\delta\beta = (-1)^b *^{-1} d * \beta$, and that, on the one hand, one has $\llbracket \phi, \beta \rrbracket = -i^{d\phi}\beta$ (cf. Eq. (14)) and, on the other hand, one has $i^{d\phi}\beta = \phi\delta\beta - \delta(\phi\beta)$. Then the rest of the proof relies on an induction on a . \square

Remark. The result of Theorem 1 can be viewed as the commutator between δ and j^α and is registered as such in Eq. (55). Also, it entails a nice identity for the Laplace-de Rham operator, $\square = -(d\delta + \delta d)$, acting on a product of two forms:

$$\square(\alpha \wedge \beta) = (\square\alpha) \wedge \beta + \alpha \wedge (\square\beta) - \llbracket \alpha, d\beta \rrbracket + (-1)^a \llbracket d\alpha, \beta \rrbracket - (-1)^a d \llbracket \alpha, \beta \rrbracket, \quad (21)$$

which can also be viewed as a commutator, see Eq. (57). In the case that one of the two forms in the product is a zero-form or a 1-form Eq. (21) simplifies to the cases below

$$\square(\phi\psi) = \psi(\square\phi) + \phi(\square\psi) + 2(d\phi, d\psi), \quad \phi, \psi \in \Omega^0, \quad (22)$$

$$\square(\phi\lambda) = \phi(\square\lambda) + \mathcal{L}_{\sharp d\phi}\lambda + *^{-1}\mathcal{L}_{\sharp d\phi}*\lambda \quad (23)$$

$$= (\square\phi)\lambda + \mathcal{L}_{\sharp d\phi}\lambda - (\delta\lambda)d\phi + \delta(\lambda \wedge d\phi), \quad \phi \in \Omega^0, \lambda \in \Omega^1, \quad (24)$$

$$\square(\phi\beta) = (\square\phi)\beta + \phi(\square\beta) + \mathcal{L}_{\sharp d\phi}\beta + \flat\mathcal{L}_{\sharp d\phi}\sharp\beta \quad (25)$$

$$= \phi(\square\beta) + \mathcal{L}_{\sharp d\phi}\beta + *^{-1}\mathcal{L}_{\sharp d\phi}*\beta, \quad \phi \in \Omega^0, \beta \in \Omega^b, \quad (26)$$

$$\square(\lambda \wedge \tau) = (\square\lambda) \wedge \tau + \lambda \wedge (\square\tau) - \flat\mathcal{L}_{\sharp\lambda}\sharp d\tau + \flat\mathcal{L}_{\sharp\tau}\sharp d\lambda + d\flat\mathcal{L}_{\sharp\lambda}\sharp\tau \quad (27)$$

$$= -(\delta d\lambda) \wedge \tau + \lambda \wedge (\square\tau) + (\delta\tau)\lambda \\ - *^{-1}\mathcal{L}_{\sharp\lambda}*\tau + *^{-1}\mathcal{L}_{\sharp\tau}*\lambda + d*^{-1}\mathcal{L}_{\sharp\lambda}*\tau, \quad \lambda, \tau \in \Omega^1. \quad (28)$$

Remark. Equation (19) provides an alternative representation of the bracket as it generates it:

$$\llbracket \alpha, \beta \rrbracket = (-1)^a [\delta(\alpha \wedge \beta) - (\delta\alpha) \wedge \beta - (-1)^a \alpha \wedge (\delta\beta)], \quad (29)$$

and makes the Gerstenhaber algebra an *exact* Gerstenhaber algebra, that is a Batalin-Vilkovisky algebra^{19,20}. Equation (29), or Eq. (19), has been known for a long time^{21,22} (see also Ref. 15) and can be recovered from Gerstenhaber original work⁹ through Hochschild-Kostant-Rosenberg theorem²³, or be found by other means²⁴. As the bracket is generated by δ it behaves nicely with respect to it, namely:

$$\delta \llbracket \alpha, \beta \rrbracket = \llbracket \delta\alpha, \beta \rrbracket - (-1)^a \llbracket \alpha, \delta\beta \rrbracket. \quad (30)$$

In addition making use of Eqs. (29) and (30), of the graded antisymmetry and Leibniz identity allows for a straightforward proof of the graded Jacobi identity Eq. (10). Moreover Eq. (29) gives rise to another representation of the bracket⁶:

$$\llbracket \alpha, \beta \rrbracket = -(\nabla_m \alpha) \wedge i^m \beta + (-1)^{a+1} (i^m \alpha) \wedge \nabla_m \beta, \quad (31)$$

where ∇ stands for the Levi-Civita connection.

Remark. The peculiar case of $a = 1$, that is $\alpha = \lambda$ being a 1-form, produces

$$\llbracket \lambda, \beta \rrbracket = \flat\mathcal{L}_{\sharp\lambda}\sharp\beta = *^{-1}\mathcal{L}_{\sharp\lambda}*\beta + (\delta\lambda)\beta, \quad (32)$$

and setting $v = \sharp\lambda$ provides a nice formula, namely:

$$*^{-1}\mathcal{L}_v* = \flat\mathcal{L}_v\sharp + \text{div}(v)\text{Id}, \quad (33)$$

by using Theorem 1 and $\delta(\alpha \wedge \beta) = -\alpha \wedge \delta\beta - *^{-1}\mathcal{L}_{\sharp\alpha}*\beta$, which holds thanks the Hodge operator properties and Cartan formula.

III. A GENERALIZED INTERIOR PRODUCT AND ITS INTERPLAY WITH THE SCHOUTEN-NIJENHUIS BRACKET

We want to extend the interior and exterior products i^λ and j^λ from $\lambda \in \Omega^1$ to any form $\alpha \in \Omega^a$. The extension of j is very natural: we note $j^\alpha \beta = \alpha \wedge \beta$, and we ask for i^α to correspond to j^α as in Eq. (3) through

$$i^\alpha \beta = \pm *^{-1} j^\alpha * \beta, \quad (34)$$

tweaking its sign such that it coincides with the $a = 1$ case and that $i^\phi \alpha = \phi \alpha$ for $\phi \in \Omega^0$. This leads to the definition below. Various generalized products have been introduced^{17,21,24} and can be introduced, here we advocate for its “naturalness” with respect to the Hodge operator.

Definition 2. Let $\alpha \in \Omega^a$ and $\beta \in \Omega^b$ be two forms, we define the generalized interior product as

$$i^\alpha \beta = (-1)^{a(b+1)} *^{-1} j^\alpha * \beta \in \Omega^{b-a}. \quad (35)$$

As a consequence we have $i^\alpha \beta = 0$ whenever $a > b$.

For $\alpha = \flat v \in \Omega^1$ a 1-form it agrees with the usual interior product (Eq. (3)) and, through direct computations, it also fulfills:

$$i^\phi \alpha = \phi \alpha, \quad (36)$$

$$i^\alpha \omega = * \alpha, \quad (37)$$

$$* i^\alpha \beta = (-1)^{a(b+1)} j^\alpha * \beta, \quad (38)$$

$$i^\alpha * \beta = (-1)^{ab} * j^\alpha \beta, \quad (39)$$

$$i^\alpha i^\beta \gamma = i^\beta \wedge^\alpha \gamma = (-1)^{ab} i^{\alpha \wedge \beta} \gamma = (-1)^{ab} i^\beta i^\alpha \gamma, \quad (40)$$

$$i^\alpha * \beta = (-1)^{ab} i^\beta * \alpha, \quad (41)$$

$$i^{*\alpha} \beta = (-1)^{a+b+ab+n} i^{*\beta} \alpha, \quad (42)$$

$$i^{*\alpha} * \beta = \text{sgn}(g) (-1)^{n(a+b)+a(b+1)} i^\beta \alpha, \quad (43)$$

for $\alpha \in \Omega^a$, $\beta \in \Omega^b$ and $\gamma \in \Omega^c$ forms, ϕ a function and ω the volume-form. For $\alpha, \beta \in \Omega^p$ two forms of the same degree one recovers the product between p -forms:

$$i^\alpha \beta = (\alpha, \beta). \quad (44)$$

Property 3. Let $\alpha \in \Omega^a$ and $\beta \in \Omega^b$ be two forms such that $a \leq b$, then the interior product is explicitly given by

$$i^\alpha \beta = \frac{1}{a!(b-a)!} \alpha_{m_1 \dots m_a} \beta_{n_1 \dots n_b} \tilde{g}(e^{m_1}, e^{n_1}) \dots \tilde{g}(e^{m_a}, e^{n_a}) e^{n_{a+1}} \wedge \dots \wedge e^{n_b}, \quad (45)$$

with $\tilde{g}(e^a, e^b) = g(\sharp e^a, \sharp e^b) = \langle e^a, \sharp e^b \rangle$. In an orthonormal basis it reads

$$i^\alpha \beta = \frac{1}{a!(b-a)!} \alpha_{m_1 \dots m_a} \beta_{n_1 \dots n_b} \eta^{m_1 n_1} \dots \eta^{m_a n_a} e^{n_{a+1}} \wedge \dots \wedge e^{n_b}. \quad (46)$$

Proof. Direct computation using the fact that $e^{m_1} \wedge \dots \wedge e^{m_a} \wedge \cdot = j^{e^{m_1}} \dots j^{e^{m_a}} \cdot \equiv j^{m_1} \dots j^{m_a} \cdot$ and commuting $*$ and j using Eq. (A4), in order to get

$$i^\alpha \beta = \frac{1}{a!} \alpha_{m_1 \dots m_a} i^{m_a} \dots i^{m_1} \beta. \quad (47)$$

Then, by writing $\beta = \frac{1}{b!} \beta_{n_1 \dots n_b} j^{n_1} \dots j^{n_b} 1$ and using Eq. (A1) to anti-commute the i s and the j s, thus generating the $g(\sharp e^{m_r}, \sharp e^{n_s}) = \tilde{g}(e^{m_r}, e^{n_s})$ terms, one obtains the end result. \square

Unfortunately, for generic forms $\alpha \in \Omega^a$ and $\beta \in \Omega^b$ there is no remarkable identity for $i^\alpha j^\beta$, because of the loss of a Leibniz identity. However, for $\lambda \in \Omega^1$ a 1-form, there remains the two special cases below

$$[i^\alpha, j^\lambda]_a \beta = i^\lambda \alpha \beta, \quad (48)$$

$$[i^\lambda, j^\alpha]_a \beta = j^\lambda i^\alpha \beta. \quad (49)$$

The interior product can also either act on a bracket or take a bracket as its argument:

$$i^{[\alpha, \beta]} \gamma = (-1)^a i^{\alpha \wedge \beta} d\gamma + (-1)^b d i^{\alpha \wedge \beta} \gamma + (-1)^{(a-1)(b-1)} i^\alpha d i^\beta \gamma - i^\beta d i^\alpha \gamma, \quad (50)$$

$$i^\lambda [\alpha, \beta] = [i^\lambda \alpha, \beta] - (-1)^a [\alpha, i^\lambda \beta] + (-1)^a [i^{d\lambda} \alpha \wedge \beta - (i^{d\lambda} \alpha) \wedge \beta - \alpha \wedge (i^{d\lambda} \beta)], \quad (51)$$

with $\alpha \in \Omega^a$, $\beta \in \Omega^b$ and $\gamma \in \Omega^c$ and the latter equation holds only for $\lambda \in \Omega^1$ a 1-form. For $\lambda = d\phi \in \Omega^1$ an exact 1-form it simplifies strikingly to

$$i^{d\phi} \llbracket \alpha, \beta \rrbracket = \llbracket i^{d\phi} \alpha, \beta \rrbracket - (-1)^a \llbracket \alpha, i^{d\phi} \beta \rrbracket, \quad (52)$$

and can be recovered using the graded Jacobi identity Eq. (10) and Eq. (14). In a similar vein setting $\alpha = \phi \in \Omega^0$ a function in Eq. (50) simplifies it to

$$i^{\llbracket \phi, \beta \rrbracket} \gamma = -i^{d\phi} \beta \gamma = -i^\beta (d\phi) \wedge \gamma + (-1)^b (d\phi) \wedge i^\beta \gamma = -[i^\beta, j^{d\phi}]_b \gamma, \quad (53)$$

and can be recovered, then, directly from Eq. (48) and Eq. (14).

Of interest for practical computations are the various commutators involving differential operators with j^α and i^α :

$$[d, j^\alpha]_a \beta = j^{d\alpha} \beta, \quad (54)$$

$$[\delta, j^\alpha]_a \beta = j^{\delta\alpha} \beta + (-1)^a \llbracket \alpha, \beta \rrbracket, \quad (55)$$

$$[\mathcal{L}_v, j^\alpha] \beta = j^{\mathcal{L}_v \alpha} \beta, \quad (56)$$

$$[\square, j^\alpha] \beta = j^{\square\alpha} \beta - \llbracket \alpha, d\beta \rrbracket + (-1)^a \llbracket d\alpha, \beta \rrbracket - (-1)^a d \llbracket \alpha, \beta \rrbracket, \quad (57)$$

$$[i^\alpha, d]_a \beta = -i^{\delta\alpha} \beta + (-1)^{b(a+1)} *^{-1} \llbracket \alpha, * \beta \rrbracket, \quad (58)$$

$$[i^\alpha, \delta]_a \beta = i^{d\alpha} \beta, \quad (59)$$

$$[i^\alpha, \mathcal{L}_v] \beta = -i^{\llbracket \flat v, \alpha \rrbracket} \beta = -i^{\flat \mathcal{L}_v \# \alpha} \beta = -i^{*-1 \mathcal{L}_v * \alpha} \beta - \text{div}(v) i^\alpha \beta, \quad (60)$$

$$[i^\alpha, \square] \beta = -i^{\square\alpha} \beta - (-1)^{ab} *^{-1} \llbracket d\alpha, * \beta \rrbracket + (-1)^{ab} *^{-1} d \llbracket \alpha, * \beta \rrbracket + (-1)^{ab+a} *^{-1} \llbracket \alpha, d * \beta \rrbracket, \quad (61)$$

in which the Schouten-Nijenhuis bracket appears naturally in the right-hand side. Equation (60) is obtained from Eq. (50) by specializing $\lambda = \flat v \in \Omega^1$ a 1-form thus leading to

$$i^{\llbracket \lambda, \alpha \rrbracket} \beta = -i^{\lambda \wedge \alpha} d\beta + (-1)^a d i^{\lambda \wedge \alpha} \beta + i^\lambda d i^\alpha \beta - i^\alpha d i^\lambda \beta, \quad (62)$$

then using Cartan formula in conjunction with Eq. (40) leads directly to the commutator of the Lie derivative with i^α . Similarly, setting $\alpha = \flat v \in \Omega^1$ a 1-form in Eq. (61) produces another notable identity between the Lie derivative and the codifferential:

$$\begin{aligned} [\mathcal{L}_v, \delta] \beta &= [\square, i_v] \beta + [d, i^{d\flat v}] \beta \\ &= [\square, i_v] \beta + i_{\# \delta \flat v} \beta + (-1)^b *^{-1} \llbracket d\flat v, * \beta \rrbracket, \end{aligned} \quad (63)$$

with the very special case for $\lambda = \flat v$ with v a Killing vector

$$[i^\lambda, \square] \beta = (d i^{d\lambda} - i^{d\lambda} d) \beta. \quad (64)$$

Remark. Some miscellaneous identities can be obtained, such as

$$\llbracket \phi_a, \dots \llbracket \phi_1, \alpha \rrbracket \dots \rrbracket = (-1)^a i^{d\phi_a} \dots i^{d\phi_1} \alpha = (-1)^a i^{d\phi_1 \wedge \dots \wedge d\phi_a} \alpha = (-1)^a (d\phi_1 \wedge \dots \wedge d\phi_a, \alpha), \quad (65)$$

with ϕ_1, \dots, ϕ_a functions and $\alpha \in \Omega^a$ a form of degree a . Also, for λ_1 and λ_2 two 1-forms, one easily gets from Eq. (10) that

$$\llbracket \lambda_1, \llbracket \lambda_2, \alpha \rrbracket \rrbracket - \llbracket \lambda_2, \llbracket \lambda_1, \alpha \rrbracket \rrbracket = \llbracket \lambda_3, \alpha \rrbracket, \quad (66)$$

with $\lambda_3 = \flat[\# \lambda_1, \# \lambda_2]$.

Finally, it can be interesting to use Tulczyjew⁷, or Michor⁸, differential operator $\Theta^\alpha = [i^\alpha, d]_a$, cf. Eq. (58), fulfilling

$$\Theta^\phi \alpha = -(d\phi) \wedge \alpha, \quad \phi \in \Omega^0, \alpha \in \Omega^a, \quad (67)$$

$$\Theta^\lambda \alpha = \mathcal{L}_{\# \lambda} \alpha, \quad \lambda \in \Omega^1, \alpha \in \Omega^a \quad (68)$$

$$\Theta^{\alpha \wedge \beta} = i^\beta \Theta^\alpha + (-1)^a \Theta^\beta i^\alpha = (-1)^{ab} [i^\alpha \Theta^\beta + (-1)^b \Theta^\alpha i^\beta], \quad \alpha \in \Omega^a, \beta \in \Omega^b, \quad (69)$$

$$[\Theta^\alpha, d]_{a+1} = 0, \quad \alpha \in \Omega^a, \quad (70)$$

$$i^{\llbracket \alpha, \beta \rrbracket} \gamma = (-1)^a i^\beta \Theta^\alpha \gamma + (-1)^{(a-1)(b-1)} \Theta^\alpha i^\beta \gamma, \quad \alpha \in \Omega^a, \beta \in \Omega^b, \gamma \in \Omega^c, \quad (71)$$

$$\Theta \llbracket \alpha, \beta \rrbracket = (-1)^{(a-1)(b-1)} \Theta^\alpha \Theta^\beta - \Theta^\beta \Theta^\alpha, \quad \alpha \in \Omega^a, \beta \in \Omega^b. \quad (72)$$

Appendix A: Overview of needed formulas of differential geometry

Let (M, g) be a smooth pseudo-riemannian manifold of dimension n , TM and T^*M being its tangent and cotangent fiber bundles, respectively. For p an integer such that $1 \leq p \leq n$ then $\Omega^p(M) \equiv \Omega^p = \wedge^p T^*M$ is the space of p -forms, smooth functions on M are identified with zero-forms: $\Omega^0(M) = C^\infty(M, \mathbb{R})$.

The natural pairing between a 1-form λ and a vector v is written $\langle \lambda, v \rangle$, the musical applications \flat and \sharp relate this pairing to the metric as $\langle \lambda, v \rangle = g(\sharp \lambda, v)$ and $g(u, v) = \langle \flat u, v \rangle$. Of prime importance here are the ‘creator operator’ j_v , such that $j_v \alpha = (\flat v) \wedge \alpha$ with $v \in TM$ and $\alpha \in \Omega^a$, and the interior product i_v (or insertion operator i_v) fulfilling an antisymmetry relation $i_u i_v = -i_v i_u$, a Leibniz identity $i_v \alpha \wedge \beta = (i_v \alpha) \wedge \beta + (-1)^a \alpha \wedge (i_v \beta)$ and interacting with j_v as

$$i_u j_v + j_v i_u = g(u, v), \quad u, v \in TM. \quad (\text{A1})$$

We recall that the interior product appears in the Lie derivative through Cartan formula: $\mathcal{L}_v = i_v d + di_v$.

With the Hodge operator $*$, fulfilling $*1 = \omega$ and $*\omega = \text{sgn}(g)$ with ω the volume form, the product between p -forms is defined as

$$(\alpha, \beta) = *^{-1}(\alpha \wedge * \beta), \quad \alpha, \beta \in \Omega^p, \quad (\text{A2})$$

and the codifferential as $\delta \alpha = (-1)^a *^{-1} d * \alpha$. For $\alpha \in \Omega^a$ a a -form, $*$ interacts with i_v and j_v as

$$* i_v \alpha = -(-1)^a j_v * \alpha, \quad (\text{A3})$$

$$* j_v \alpha = (-1)^a i_v * \alpha, \quad (\text{A4})$$

see say App. A of Ref. 1 for additional identities. The generalized interior product stems from Eq. (A3).

Appendix B: Sketches of a few long computations

1. Eq. (50)

In order to establish Eq. (50) one relies on the representation of the bracket as it appears in Eq. (29) and one is led to consider the term $i^{\delta(\alpha \wedge \beta)} \gamma$. Using Eq. (12) and explicit formulas for δ , it reads

$$\begin{aligned} i^{\delta(\alpha \wedge \beta)} \gamma &= (-1)^{(a+b-1)(c+1)} *^{-1} (\delta(\alpha \wedge \beta)) \wedge * \gamma \\ &= (-1)^{(a+b-1)c} *^{-1} \{ (-1)^{a+b+1} \delta(\alpha \wedge \beta \wedge * \gamma) + \alpha \wedge \beta \wedge (\delta * \gamma) + \llbracket \alpha \wedge \beta, * \gamma \rrbracket \} \\ &= (-1)^{(a+b)c} d *^{-1} \alpha \wedge \beta \wedge * \gamma + (-1)^{(a+b)c+1} *^{-1} \alpha \wedge \beta \wedge d \gamma \\ &\quad + (-1)^{(a+b-1)c} *^{-1} \alpha \wedge * (*^{-1} \llbracket \beta, * \gamma \rrbracket) + (-1)^{(a+b-1)c+ab} *^{-1} \beta \wedge * (*^{-1} \llbracket \alpha, * \gamma \rrbracket). \end{aligned} \quad (\text{B1})$$

Owing to its definition, Eq. (35), we recognize in the above expression the generalized interior products $i^{\alpha \wedge \beta}$, i^α and i^β . The remaining terms: $*^{-1} \llbracket \beta, * \gamma \rrbracket$ and $*^{-1} \llbracket \alpha, * \gamma \rrbracket$, are reexpressed using Eq. (58) backward. Bringing all together with the remaining terms of Eq. (29): $i^{(\delta \alpha) \wedge \beta}$ and $i^{\alpha \wedge (\delta \beta)}$, produces Eq. (50).

2. Eqs. (54)–(61)

First notice that among the set of Eqs. (54)–(61) a few are known formulas rewritten as commutators, namely: Eq. (54) is Leibniz’ identity for the exterior derivative, Eq. (55) is Eq. (19), Eq. (56) is Leibniz’ identity for the Lie derivative, Eq. (57) is Eq. (21) which is obtained from Eq. (54) and (55). The remaining formulas, Eqs. (58)–(61), are found from the previous by conjugation with the Hodge operator: $x\beta \rightarrow *^{-1} x * \beta$.

- ¹E. Hugueta, J. Queva, and J. Renaud, *J. Math. Phys.* **63**, 072301 (2022), arXiv:2201.03253 [math-ph].
- ²M. Fecko, *Differential geometry and Lie groups for physicists*, paperback edition ed. (Cambridge University Press, 2011).
- ³J. A. Schouten, *Nederl. Akad. Wetensch. Proc.* **43**, 449 (1940).
- ⁴J. A. Schouten, in *Convegno Int. Geom. Diff.* (Cremonese, 1953) pp. 1–7.
- ⁵A. Nijenhuis, *Indagationes Mathematicae (Proceedings)* **58**, 390 (1955).
- ⁶A. Nijenhuis, *Indagationes Mathematicae (Proceedings)* **58**, 398 (1955).
- ⁷W. Tulczyjew, *Bull. Acad. Polon. Sci.* **22**, 937 (1974).
- ⁸P. W. Michor, in *Proc. of the Winter School "Geometry and Physics"* (Circ. Matematico di Palermo, 1987) pp. 207–215.
- ⁹M. Gerstenhaber, *Annals of Mathematics* **78**, 267 (1963).
- ¹⁰Y. Kosmann-Schwarzbach, Graded poisson brackets and field theory, in *Modern Group Theoretical Methods in Physics: Proceedings of the Conference in Hono* edited by J. Bertrand, M. Flato, J.-P. Gazeau, M. Irac-Astaud, and D. Sternheimer (Springer, 1995) pp. 189–196.
- ¹¹C. Roger, *Archivum Mathematicum* **045**, 301 (2009).
- ¹²L. K. Norris, *J. Math. Phys.* **38**, 2694 (1997).
- ¹³E. Gozzi and D. Mauro, *J. Math. Phys.* **41**, 1916 (2000), arXiv:hep-th/9907065.
- ¹⁴K. Bhaskara and K. Viswanath, *Poisson Algebras and Poisson Manifolds*, Pitman research notes in mathematics series (Longman Scientific & Technical, 1988).
- ¹⁵I. Vaisman, *Lectures on the geometry of Poisson manifolds*, Vol. 118 (Birkhäuser, 1994).
- ¹⁶Y. Kosmann-Schwarzbach, *Journal of Geometric Mechanics* **13**, 459 (2021), arXiv:2105.14828 [math.HO].
- ¹⁷C.-M. Marle, *Journal of Geometry and Physics* **23**, 350 (1997).
- ¹⁸Y. Mikami, in *Noncommutative Differential Geometry and Its Applications to Physics: Proceedings of the Workshop at Shonan, Japan, June 1999*, edited by Y. Maeda, H. Moriyoshi, H. Omori, D. Sternheimer, T. Tate, and S. Watamura (Springer, 2001) pp. 131–143.
- ¹⁹I. A. Batalin and G. A. Vilkovisky, *Physics Letters B* **102**, 27 (1981).
- ²⁰E. Witten, *Mod. Phys. Lett. A* **05**, 487 (1990).
- ²¹J.-L. Koszul, in *Élie Cartan et les mathématiques d'aujourd'hui - Lyon, 25-29 juin 1984*, Astérisque No. S131 (Société mathématique de France, 1985) pp. 257–271.
- ²²I. Vaisman, *Annales de l'Institut Fourier* **40**, 951 (1990).
- ²³G. Hochschild, B. Kostant, and A. Rosenberg, *Trans. Amer. Math. Soc.* **102**, 383 (1962).
- ²⁴B. Coll and J.-J. Ferrando, *J. Math. Phys.* **45**, 2405 (2004), arXiv:gr-qc/0306102.