

Learning Coefficients in Semi-Regular Models

Yuki Kurumadani*

1 Abstract

Statistical learning models such as multilayer neural networks and mixed distributions are widely used, and understanding the accuracy of these models is crucial for their use. Recent advances have clarified theoretical learning accuracy in Bayesian inference, where metrics such as generalization loss and free energy are used to measure the accuracy of predictive distributions. It has become clear that the asymptotic behavior of these metrics is determined by a rational number specific to each statistical model, known as the learning coefficient (real log canonical threshold) [2]. It is known that for learning models that satisfy the regularity conditions for the posterior distribution to converge to a normal distribution, the learning coefficient becomes $d/2$ where d is the number of the model's parameters [2]. However, for singular models that do not satisfy the regularity conditions, except for a few models such as reduced-rank regression [3], the exact method for calculating the learning coefficient has not yet been discovered.

The problem of determining the learning coefficient is known to be reducible to the problem of finding the normal crossing of Kullback-Leibler divergence in relation to algebraic geometry [2]. In this context, it is crucial to perform appropriate coordinate transformations and blow-ups. This paper attempts to derive appropriate variable transformations and blow-ups from the properties of the log-likelihood ratio function. That is, instead of dealing with the Kullback-Leibler information itself, it uses the properties of the log-likelihood ratio function before taking the expectation to calculate the real log canonical threshold. This approach has not been considered in previous research.

Using these variable transformations and blow-ups, this paper provides the exact values of the learning coefficients and their calculation methods for statistical models that meet simple conditions next to the regular conditions (referred to as **semi-regular models**), and as specific examples, provides the learning coefficients for semi-regular models with two parameters and for those models where the random variables take a finite number of values.

2 Introduction

2.1 Definitions and Assumptions

Throughout this paper, we consider statistical models $p(x|\theta)$ with continuous parameters $\theta = (\theta_1, \dots, \theta_d) \in \Theta(\subset \mathbb{R}^d)(d \geq 1)$, and we denote the true distribution by $q(x)$. The set of possible

*Graduate School of Engineering Science, Osaka University (kurumadani@sigmath.es.osaka-u.ac.jp)

values for data x is denoted by χ . The statistical model is defined as **realizable**, meaning there exists a parameter θ_* such that $q(x) = p(x|\theta_*)$ a.s.. Such parameters θ_* are referred to as realization parameters, and the entire set of these parameters is denoted by Θ_* . **The prior distribution $\varphi(\theta)$ is assumed to satisfy $\varphi(\theta_*) > 0$ for any realization parameter θ_* .** The random variable that follows the true distribution q is denoted by X , and $\mathbb{E}_X[\cdot]$ denotes the operation of taking the average concerning the random variable X . In this paper, we assume that the operations of taking expectations and partial derivatives with respect to θ are interchangeable.

The Kullback-Leibler divergence is defined as:

$$K(\theta) := \mathbb{E}_X \left[\log \frac{p(X|\theta_*)}{p(X|\theta)} \right]$$

and is assumed to be analytic around $\theta = \theta_*$. The log-likelihood ratio function is given by:

$$f(x|\theta) := \log \frac{p(x|\theta_*)}{p(x|\theta)}$$

and is assumed to be L^2 integrable and analytic around $\theta = \theta_*$.

For fixed data x , the m -th order terms of the Taylor expansion of the log-likelihood ratio function $f(x|\theta)$ at $\theta = \theta_*$ for $\theta_1, \dots, \theta_s$ ($s \leq d$) are defined as:

$$F_m(x|\theta_1, \dots, \theta_s) := \sum_{\substack{i_1 + \dots + i_s = m \\ i_1, \dots, i_s \in \mathbb{Z}_{\geq 0}}} \frac{1}{i_1! \dots i_s!} \times \left. \frac{\partial^m f(x|\theta)}{\partial \theta_1^{i_1} \dots \partial \theta_s^{i_s}} \right|_{\theta = \theta_*} \times (\theta_1 - \theta_{1*})^{i_1} \dots (\theta_s - \theta_{s*})^{i_s}$$

The Fisher information matrix at $\theta = \theta_*$ is denoted by I , i.e.,

$$I = \text{Cov} \left(\left. \frac{\partial \log p(X|\theta)}{\partial \theta_i} \right|_{\theta = \theta_*}, \left. \frac{\partial \log p(X|\theta)}{\partial \theta_j} \right|_{\theta = \theta_*} \right)_{i,j=1, \dots, d}$$

and its rank is denoted by r . The Hessian matrix of $K(\theta)$ at $\theta = \theta_*$ is denoted by J , i.e.,

$$J = \left(\left. \frac{\partial^2 K(\theta)}{\partial \theta_i \partial \theta_j} \right|_{\theta = \theta_*} \right)_{i,j=1, \dots, d}$$

Generally, I and J do not coincide, but under the conditions of this paper, as will be seen later in Remark 3.5, they do coincide.

2.2 What is a Learning Coefficient?

First, let us describe the framework of Bayesian theory. Consider a statistical model $p(x|\theta)$ with parameters $\theta \in \mathbb{R}^d$ ($d \geq 1$), where $x \in \chi$, and the true distribution is denoted by $q(x)$. The posterior distribution of the parameter θ given data X_1, \dots, X_n independently drawn from $q(x)$ is expressed as:

$$p(\theta|X_1, \dots, X_n) = \frac{\varphi(\theta) \prod_{i=1}^n p(X_i|\theta)}{Z_n}$$

where $Z_n := \int \varphi(\theta) \prod_{i=1}^n p(X_i|\theta) d\theta$ represents the marginal likelihood function. When a new observation x is given, a statistical model marginalized over its posterior distribution is called a predictive distribution, represented by:

$$p^*(x) := \int p(x|\theta) p(\theta|X_1, \dots, X_n) d\theta$$

This predictive distribution is used to estimate the true distribution $q(x)$ under Bayesian theory.

The accuracy of the predictive distribution is measured by metrics such as the generalization loss G_n and the free energy F_n :

$$G_n := - \int_{\mathcal{X}} q(x) \log p^*(x) dx$$

$$F_n := - \log Z_n = - \log \int \varphi(\theta) \prod_{i=1}^n p(X_i|\theta) d\theta$$

These metrics are known to take smaller values when the predictive distribution closely approximates the true distribution.

When the true distribution $q(x)$ is realizable by the statistical model $p(x|\theta)$, the generalization loss G_n and the free energy F_n exhibit the following asymptotic behavior (i.e., the behavior as the sample size n becomes large) using a positive rational number λ and an integer m greater than or equal to 1 [2].

$$\mathbb{E}[G_n] = - \int_{\mathcal{X}} q(x) \log q(x) dx + \frac{\lambda}{n} - \frac{m-1}{n \log n} + o\left(\frac{1}{n \log n}\right)$$

$$\mathbb{E}[F_n] = -n \int_{\mathcal{X}} q(x) \log q(x) dx + \lambda \log n - (m-1) \log \log n + O(1)$$

Since the first term on the right-hand side of both expressions does not depend on the statistical model $p(x|\theta)$, the asymptotic behaviors are determined by λ in the second term. Thus, when comparing two learning models using these metrics, the value of λ becomes crucial to determine which model better approximates the true distribution. Generally, this λ is referred to as the **learning coefficient**. The learning coefficient λ is defined for the trio of the statistical model $p(x|\theta)$, the true distribution $q(x)$, and the prior distribution $\varphi(\theta)$.

2.3 Methods for Calculating Learning Coefficients Using Algebraic Geometry

It is known that the concept of learning coefficients coincides with the algebraic geometry concept of the **real log canonical threshold**. The real log canonical threshold is defined using a technique known as resolution of singularities. Here, resolution of singularities refers to transforming an analytic function F into normal crossing as specified by the following theorem:

Theorem 2.1 (Resolution of Singularities). [1][2, Theorem 2.3] Let $F(x)$ be a real analytic function defined near the origin in \mathbb{R}^d and assume $F(0) = 0$. Then, there exists an open set $W \subset \mathbb{R}^d$ containing the origin, a real analytic manifold U , and a proper analytic map $g : U \rightarrow W$ satisfying the following conditions:

- (1) Define $W_0 := F^{-1}(0)$ and $U_0 := g^{-1}(W_0)$. The map $g : U - U_0 \rightarrow W - W_0$ is an analytic isomorphism.
- (2) At any point Q in U_0 , by taking local coordinates $u = (u_1, \dots, u_d)$ with Q as the origin, we can express:

$$\begin{aligned} F(g(u)) &= a(u)u_1^{k_1}u_2^{k_2} \cdots u_d^{k_d} \\ |g'(u)| &= \left| b(u)u_1^{h_1}u_2^{h_2} \cdots u_d^{h_d} \right| \end{aligned} \tag{2.1}$$

where $k_i, h_i (i = 1, \dots, d)$ are non-negative integers, and $a(u), b(u)$ are real analytic functions defined near the origin in \mathbb{R}^d with $a(0) \neq 0, b(0) \neq 0$.

The expression as in (2.1) is referred to as **normal crossing**.

It should be noted that Theorem 2.1 is a local statement concerning the neighborhood of the origin in \mathbb{R}^d . In other words, if there are multiple points where singularities need to be resolved, it is necessary to find (W, U, g) as guaranteed by this Theorem 2.1 at each point.

Definition 2.1 (Real Log Canonical Threshold). Let F be a real analytic function defined on an open set O in \mathbb{R}^d , and let C be a compact set containing O . For each point P in C satisfying $F(P) = 0$, by applying a coordinate transformation such that point P is moved to the origin of \mathbb{R}^d , Theorem 2.1 can be applied. Fix (W, U, g) as guaranteed by Theorem 2.1(2). Additionally, denote the non-negative integers h_i, k_i given by Theorem 2.1(2) in the neighborhood of any point $Q \in U_0$ as $h_i^{(Q)}, k_i^{(Q)}$.

- (1) Define the **real log canonical threshold** λ_P at point P of the function F as:

$$\lambda_P = \inf_{Q \in U_0} \left\{ \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} \right\}$$

where $(h_i + 1)/k_i = \infty$ if $k_i = 0$. It is known that this is well-defined, i.e., it does not depend on the choice of (W, U, g) . [2, Theorem 2.4]

- (2) Define the **real log canonical threshold** λ for the compact set C of the function F as [2, Definition 2.7]:

$$\lambda = \inf_{P \in C} \lambda_P$$

- (3) In (2), for the point $P (\in C)$ that gives the minimum value, the maximum number of i such that $\lambda_P = (h_i^{(Q)} + 1)/k_i^{(Q)}$ is satisfied is called the **multiplicity**. (If there are multiple points $P (\in C)$ that give the minimum value, the maximum number of i among each is called the multiplicity.)

In this paper, we apply Theorem 2.1 to the analytic function F defined as the Kullback-Leibler divergence $K(\theta)$. As previously seen, we assumed a prior distribution $\varphi(\theta_*) > 0$ in this paper. Under this condition, the **learning coefficient** λ is equal to the **real log canonical**

threshold for the compact set $\Theta_* = \{\theta \in \Theta | K(\theta) = 0\}$ [2, Theorem 6.6, Definition 6.4]. That is, by performing the resolution of singularities guaranteed by Theorem 2.1 at each point P of Θ_* , we obtain the real log canonical threshold λ_P . The minimum value of λ_P as point P moves over the entire Θ_* coincides with the learning coefficient λ . Moreover, if the real log canonical threshold λ_P at some points P on Θ_* can be calculated, it is clear from the definition that this provides an upper bound for the learning coefficient. That is, $\lambda \leq \lambda_P$ holds.

It should be noted that, although $K(P_1) = K(P_2) = 0$ holds for any elements $P_1, P_2 \in \Theta_*$, the differential structure of the function K in the neighborhoods of these two points may not be identical, and thus the real log canonical thresholds $\lambda_{P_1}, \lambda_{P_2}$ may not coincide. Therefore, to obtain the learning coefficient, it is necessary to calculate the real log canonical threshold λ_P for all points P in Θ_* .

2.4 Learning Coefficients for Models Satisfying Regularity Conditions

It is known that for learning models satisfying the regularity conditions, which allow the posterior distribution to converge to a normal distribution, the learning coefficient λ is given by $\lambda = d/2$, where d is the dimension of the parameter space [2]. Here, the regularity conditions are defined as follows:

Definition 2.2 (Regularity). A true model $q(x)$ is said to be **regular** with respect to the statistical model $p(x|\theta)$ if it satisfies the following three conditions:

- (1) There is only one element in the set of realization parameters Θ_* .
- (2) θ_* is an interior point of the parameter set Θ , meaning there exists an open neighborhood $\tilde{\Theta} \subset \Theta$ around θ_* .
- (3) The Fisher information matrix $I \in \mathbb{R}^{d \times d}$ is a positive definite matrix.

This paper aims to generalize this formula. More specifically, it considers the case where the rank r of the Fisher information matrix I is $0 < r < d$ to provide the learning coefficient λ .

2.5 Upper Bound of Learning Coefficients for General Models

Assuming a prior distribution $\varphi(\theta_*) > 0$ throughout this paper, it is known that the real log canonical threshold λ of $K(\theta)$ at $\theta = \theta_*$ satisfies[2, Theorem 7.2]:

$$\lambda \leq \frac{d}{2} \tag{2.2}$$

Furthermore, let $\theta = (u, v) \in \Theta$ ($u \in \mathbb{R}^{d_1}, v \in \mathbb{R}^{d_2}$), and if $K(u_*, v) = 0$ holds for any v in the neighborhood of $\theta = \theta_* = (u_*, v_*)$, it is known that

$$\lambda \leq \frac{d_1}{2} \tag{2.3}$$

holds true. [2, Theorem 7.3]

3 Main Theorem

3.1 Overview of the Main Theorem

Definition 3.1 (Semi-Regularity). A statistical model $p(x|\theta)$ is said to be **semi-regular**¹ at the parameter θ_* that realizes the true model $q(x)$ if the rank r of the Fisher information matrix $I \in \mathbb{R}^{d \times d}$ at $\theta = \theta_*$ is greater than zero.

Clearly, a regular model is a semi-regular model. In the following, we arbitrarily fix an element θ_* in Θ_* and consider the real log canonical threshold at this point. We assume (by translating if necessary) that $\theta_* = 0$. In the subsequent discussion, we will denote it as θ_* for statements that hold regardless of whether $\theta_* = 0$, but it can be read as $\theta_* = 0$ without loss of generality. As we have already confirmed, if the real log canonical threshold at a certain realization parameter can be calculated, it provides an upper bound for the learning coefficient.

Assumption 1. In the Main Theorem of this paper, we assume the following (1)-(3) for semi-regular models.

- (1) For the r parameters $\theta_1, \dots, \theta_r$, the r random variables

$$\left. \frac{\partial f(X|\theta)}{\partial \theta_1} \right|_{(\theta_1, \dots, \theta_d)=0}, \dots, \left. \frac{\partial f(X|\theta)}{\partial \theta_r} \right|_{(\theta_1, \dots, \theta_d)=0}$$

are linearly independent².

- (2) For the remaining $d - r$ parameters $\theta_{r+1}, \dots, \theta_d$, let m be an integer greater than or equal to 1. The derivatives of the log-likelihood ratio function $f(X|\theta)$ with respect to $\theta_{r+1}, \dots, \theta_d$ up to order $m - 1$ at $(\theta_1, \dots, \theta_d) = 0$ are zero with probability 1. That is,

$$F_1(X|\theta_{r+1}, \dots, \theta_d) = \dots = F_{m-1}(X|\theta_{r+1}, \dots, \theta_d) = 0 \quad a.s.$$

holds.

The maximum value among such m is redefined as m . For convenience, if $m = 1$, it is treated as the case where $r = d$ in (1), and if (2) holds for any $m \geq 2$, it is denoted as $m = \infty$.

- (3) If $2 \leq m < \infty$, then for each of the $d - r$ parameters $(\theta_{r+1}, \dots, \theta_d) \neq 0$, one of the following holds:

- (i) $F_m(X|\theta_{r+1}, \dots, \theta_d) = 0$ (a.s.)
- (ii) $F_m(X|\theta_{r+1}, \dots, \theta_d)$ and the r random variables in (1) are linearly independent.

For $m = 1$, it is always treated as satisfying (ii) for convenience.

¹This terminology is not a general term.

²In this paper, random variables are said to be linearly independent if they are so with probability 1.

Remark 3.1. In Theorem 4.1 described later, we will see that any semi-regular model can satisfy this Assumption through appropriate coordinate transformations.

For semi-regular models that satisfy Assumption 1(1)(2), the Taylor expansion of the log-likelihood ratio function $f(X|\theta)$ in terms of the parameters θ does not contain terms of order less than $m - 1$ for $(\theta_{r+1}, \dots, \theta_d)$, implying:

$$f(X|\theta) = F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d) + (\text{higher order terms}) \text{ a.s.}$$

Taking the expected value with respect to X yields the Taylor expansion of $K(\theta)$ at $\theta = 0$:

$$K(\theta) = \mathbb{E}_X [F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d)] + (\text{higher order terms})$$

However, it turns out that many of the lower order terms vanish upon taking the expectation. This formulation is presented in Main Theorem 1.

Main Theorem 1. For semi-regular models that satisfy Assumption 1(1)(2), the Taylor expansion of $K(\theta)$ at $\theta = 0$ can be expressed as:

$$K(\theta) = \frac{1}{2} \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] + (\text{higher order terms})$$

The (higher order terms) do not include:

- Terms up to the $2m$ -th order that consist only of $\theta_{r+1}, \dots, \theta_d$
- Terms that are first order in $\theta_1, \dots, \theta_r$ and up to m -th order in $\theta_{r+1}, \dots, \theta_d$
- Second-order terms that consist only of $\theta_1, \dots, \theta_r$

In the case where $m = \infty$, $F_m(X|\theta_{r+1}, \dots, \theta_d)$ is considered to be zero a.s.

Main Theorem 2. Consider a semi-regular model that satisfies Assumption 1(1)(2)(3) and where $m < \infty$. Consider the following blow-up g at the origin O :

- (a) Perform one blow-up centered at the origin of \mathbb{R}^d .
- (b) If the exceptional surface from (a) is $\{\theta_i = 0\}$ (where $i = r + 1, \dots, d$), then further perform a blow-up centered at the subvariety $\{(\theta_1, \dots, \theta_d) \mid \theta_1 = \dots = \theta_r = \theta_i = 0\}$.
- (c) If the exceptional surface from (b) is $\{\theta_i = 0\}$, repeat (b) until the total number of blow-ups reaches m .

That is, at the m -th blow-up, when the exceptional surface is $\{\theta_i = 0\}$, the map $g = g_i$ can be expressed as follows for $(i = r + 1, \dots, d)$:

$$g_i: (\theta'_1, \dots, \theta'_{i-1}, \theta_i, \theta'_{i+1}, \dots, \theta'_d) \mapsto (\theta_1, \dots, \theta_{i-1}, \theta_i, \theta_{i+1}, \dots, \theta_d);$$




Fig 1: Main Theorem 1 (case $m < \infty$). Assuming that the derivatives of the log-likelihood ratio function are zero (a.s.) in the blue area (including the endpoints), the derivatives of the log-likelihood ratio function in the red area (including the endpoints) become random variables with expected value zero. This implies that the coefficients of $K(\theta)$'s Taylor expansion in these regions are all zero.

$$\begin{aligned}\theta_1 &= \theta_i^m \theta'_1, \dots, \theta_r = \theta_i^m \theta'_r, \\ \theta_{r+1} &= \theta_i \theta'_{r+1}, \dots, \theta_{i-1} = \theta_i \theta'_{i-1}, \theta_{i+1} = \theta_i \theta'_{i+1}, \dots, \theta_d = \theta_i \theta'_d\end{aligned}$$

Defining the subset S of $U_0 := g^{-1}(O)$ by local coordinates $(\theta'_1, \dots, \theta'_{i-1}, \theta_i, \theta'_{i+1}, \dots, \theta'_d)$ as

$$S := \bigcup_{i=r+1}^d \left\{ (\theta'_1, \dots, \theta'_d) \mid \begin{array}{l} (\theta'_1, \dots, \theta'_r, \theta_i) = 0 \\ F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d) = 0(a.s.) \end{array} \right\} \subset U_0$$

then, on $U_0 \setminus S$, normal crossing of $K(\theta)$ is obtained, and

$$\inf_{Q \in U_0 \setminus S} \left\{ \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} \right\} = \frac{d - r + rm}{2m}$$

is satisfied (multiplicity is 1). See Definition 2.1 for the symbols $k_i^{(Q)}, h_i^{(Q)}$.

Corollary 3.1. [Formula for the Real Log Canonical Threshold] Consider a semi-regular model that satisfies Assumption 1(1)(2).

- (1) If $m < \infty$ and all parameters $(\theta_{r+1}, \dots, \theta_d) \neq 0$ satisfy Assumption 1(3)(ii), then the real log canonical threshold λ_O at the origin O of $K(\theta)$ is given by the following formula (with multiplicity 1):

$$\lambda_O = \frac{d - r + rm}{2m}$$

In this case, the only point in the parameter space Θ near the origin where $K(\theta) = 0$ is the origin itself.

- (2) If $m = \infty$, a blow-up centered at the subvariety $W_0 := \{(\theta_1, \dots, \theta_d) \mid \theta_1 = \dots = \theta_r = 0\}$ obtains a normal crossing for $K(\theta)$, and the real log canonical threshold λ_O at the origin O is given by the following formula (with multiplicity 1):

$$\lambda_O = \frac{r}{2}$$

In this case, the set of points in the parameter space Θ near the origin where $K(\theta) = 0$ is W_0 .

Remark 3.2. In the case of regular models ($r = d$), substituting $r = d$ into the result of Corollary 3.1(1) gives the real log canonical threshold:

$$\lambda_O = \frac{d - d + d \times m}{2 \times m} = \frac{d}{2}$$

which indeed matches the results of prior research. Thus, this result generalizes the case for regular models. Additionally, the results of Corollary 3.1(1) are consistent with the findings of previous research (2.2), and the results of Corollary 3.1(2) align with the findings of previous research (2.3). Furthermore, the result of (2) matches the result of (1) when taking $m \rightarrow \infty$.

The real log canonical threshold that can be directly calculated using Corollary 3.1 is for cases where the multiplicity is 1, which geometrically means that the origin is a non-singular point in the set of realization parameters Θ_* . However, even if it is a singular point, it is possible to calculate the real log canonical threshold by further progressing the blow-up at the points contained in the set S using the Main Theorem 2.

Using Corollary 3.1 suggests that one can discuss the real log canonical threshold from the linear independence (or dependence) of the random variables that appear as coefficients in the lower-order terms of the Taylor expansion of the log-likelihood ratio function, without specifically considering the Taylor expansion of $K(\theta)$. In other words, there is **no need to take expectations with respect to the random variable X** .

Remark 3.3. When considering 2λ as an indicator of the complexity of a statistical model, the result for a regular model implies that 2λ coincides with the number of parameters d . In the case of a singular model, from (2.2), it is understood that 2λ is less than or equal to d .

From this perspective, the result of Corollary 3.1(1) suggests that for the linearly independent parameters $\theta_1, \dots, \theta_r$, each counts as 1, and for the parameters $\theta_{r+1}, \dots, \theta_d$ whose derivatives up to $m - 1$ are all zero, each counts as $1/m$.

Furthermore, the result of Corollary 3.1(2) suggests that parameters $\theta_{r+1}, \dots, \theta_d$, for which any number of derivatives are zero, do not affect the real log canonical threshold, and each linearly independent parameter $\theta_1, \dots, \theta_r$ can be counted as 1.

Remark 3.4. Consider the real log canonical threshold of Corollary 3.1(1) from the perspective of ideals. For the parameter $(\theta_1, \dots, \theta_d)$, consider the ideal

$$I := (\theta_1, \dots, \theta_r, \theta_{r+1}^m + \dots + \theta_d^m)$$

The real log canonical threshold of Corollary 3.1(1) is the same as the real log canonical threshold of the ideal I , i.e., the polynomial

$$\theta_1^2 + \dots + \theta_r^2 + (\theta_{r+1}^m + \dots + \theta_d^m)^2$$

can resolve singularities by the same blow-ups.

Before proving Main Theorem 1, 2, some practical examples are introduced.

Example 3.1 (Case of $m = 2$). Let X be a random variable following a binomial distribution $\text{Bin}(2, \theta)$ with parameter θ where $0 < \theta < 1$.

$$\tilde{p}(X = x|\theta) = \binom{2}{x} \theta^x (1 - \theta)^{2-x} = \begin{cases} (1 - \theta)^2, & (x = 0) \\ 2\theta(1 - \theta), & (x = 1) \\ \theta^2, & (x = 2) \end{cases}$$

Consider a mixed distribution model with parameters (θ_1, θ_2) given by:

$$p(X = x|\theta_1, \theta_2) := \frac{1}{2} \cdot \tilde{p}\left(X = x \left| \theta_1 - \theta_2 + \frac{1}{2} \right.\right) + \frac{1}{2} \cdot \tilde{p}\left(X = x \left| \theta_2 + \frac{1}{2} \right.\right) \quad (x = 0, 1, 2) \quad (3.1)$$

Assuming the true distribution is $\tilde{p}(X|1/2)$, note that this model realizes the true distribution at $(\theta_1, \theta_2) = 0$.

Let us verify that the Main Theorem 1 holds at $(\theta_1, \theta_2) = 0$. By setting $d = 2, r = 1$ and demonstrating that $F_1(X|\theta_1)$ and $F_2(X|\theta_2)$ are non-zero, it follows that Assumption 1(1)(2) is satisfied in the case of $m = 2$.

Using a computer program mathematica, we obtain³:

$$K(\theta) = \theta_1^2 + 8\theta_2^4 + 8\theta_1^2\theta_2^2 - 16\theta_1\theta_2^3 + \dots \quad (3.2)$$

$$F_1(X|\theta_1) = \left. \frac{\partial f(X|\theta)}{\partial \theta_1} \right|_{(\theta_1, \theta_2)=0} \theta_1 = \begin{cases} 2\theta_1, & x = 0 \\ 0, & x = 1 \\ -2\theta_1, & x = 2 \end{cases} \quad (3.3)$$

$$F_1(X|\theta_2) = \left. \frac{\partial f(X|\theta)}{\partial \theta_2} \right|_{(\theta_1, \theta_2)=0} \theta_2 = 0 \quad (3.4)$$

$$F_2(X|\theta_2) = \left. \frac{1}{2} \frac{\partial^2 f}{\partial \theta_2^2} \right|_{(\theta_1, \theta_2)=0} \theta_2^2 = \begin{cases} -4\theta_2^2, & x = 0 \\ 4\theta_2^2, & x = 1 \\ -4\theta_2^2, & x = 2 \end{cases} \quad (3.5)$$

Terms of order five and higher are omitted in (3.2). It is verified with probability 1 that (3.3) and (3.5) are non-zero.

The applicability of Main Theorem 1 is verified using (3.3) and (3.5), yielding:

$$\begin{aligned} \frac{1}{2} \mathbb{E}_X \left[\{F_1(X|\theta_1) + F_2(X|\theta_2)\}^2 \right] &= \frac{1}{2} \left\{ (2\theta_1 - 4\theta_2^2)^2 \cdot \frac{1}{4} + (0 + 4\theta_2^2) \cdot \frac{1}{2} + (-2\theta_1 - 4\theta_2^2)^2 \cdot \frac{1}{4} \right\} \\ &= \theta_1^2 + 8\theta_2^4 \end{aligned}$$

which shows that θ_1^2 and $8\theta_2^4$ are not part of the (higher order terms) in Main Theorem 1, and other terms are included in (higher order terms). Thus, as claimed by the Main Theorem 1, the

³See Appendix for calculations.

(higher order terms) do not include: terms of degree four or less consisting only of θ_2 , terms of first degree in θ_1 and second degree or less in θ_2 , terms of second degree consisting only of θ_1 .

Next, verify that Main Theorem 2 holds. First, note that (3.3) and (3.5) are linearly independent for any $\theta_2 \neq 0$ when $\theta_1 = 1$, so this statistical model satisfies Assumption 1(3)(ii).

Thus, applying Corollary 3.1(1) for $(d, r, m) = (2, 1, 2)$, the real log canonical threshold at the origin should be $3/4$ (multiplicity is 1). This is verified by performing the blow-up g_1 centered at $(\theta_1, \theta_2) = 0$ and seeking the normal crossing of $K(\theta)$.

(a) First, transform (3.2) with $\theta_2 = \theta_1 \theta'_2$ using power series h_1, a_1 to get,

$$K(\theta) = \theta_1^2 \{1 + 8\theta_1^2 \theta_2'^4 + \theta_1 h_1(\theta_1, \theta_2')\} = \theta_1^2 a_1(\theta_1, \theta_2')$$

Since on any point of $g_1^{-1}(0) = \{(\theta_1, \theta_2') | \theta_1 = 0\}$,

$$\forall \theta_2', a_1(0, \theta_2') = 1 \neq 0$$

normal crossings are obtained in this local coordinate (θ_1, θ_2') .

(b) Next, transform (3.2) with $\theta_1 = \theta_2 \theta'_1$ using power series h_2, a_2 to get,

$$K(\theta) = \theta_2^2 \{\theta_1'^2 + 8\theta_2^2 + \theta_2 h_2(\theta_1', \theta_2)\} = \theta_2^2 a_2(\theta_1', \theta_2)$$

where $g_1^{-1}(0) = \{(\theta_1', \theta_2) | \theta_2 = 0\}$ on any point except at $(\theta_1', \theta_2) = 0$,

$$\forall \theta_1' \neq 0, a_2(\theta_1', 0) = \theta_1'^2 \neq 0$$

shows a normal crossing is achieved.

Therefore, it is sufficient to find the normal crossings at the point $(\theta_1', \theta_2) = 0$. Further blow-up g_2 is performed centered at this point.

(b1) First, transform using $\theta_2 = \theta_1' \theta_2''$ with power series h_3, a_3 to get,

$$K(\theta) = \theta_1'^4 \theta_2''^2 \{1 + 8\theta_2''^2 + \theta_1' h_3(\theta_1', \theta_2'')\} = \theta_1'^4 \theta_2''^2 a_3(\theta_1', \theta_2'')$$

where $g_2^{-1}(0) = \{(\theta_1', \theta_2'') | \theta_1' = 0\}$ on any point,

$$\forall \theta_2'', a_3(0, \theta_2'') = 1 + 8\theta_2''^2 \neq 0$$

shows a normal crossing is achieved.

(b2) Next, transform using $\theta_1' = \theta_2 \theta_1''$ with power series h_4, a_4 to get,

$$K(\theta) = \theta_2^4 \{\theta_1''^2 + 8 + \theta_2 h_4(\theta_1'', \theta_2)\} = \theta_2^4 a_4(\theta_1'', \theta_2)$$

where $g_2^{-1}(0) = \{(\theta_1'', \theta_2) | \theta_2 = 0\}$ on any point,

$$\forall \theta_1'', a_4(\theta_1'', 0) = \theta_1''^2 + 8 \neq 0$$

shows a normal crossing is achieved.

Summarizing the normal crossings for each local coordinate (see Definition 2.1 for notation $k_i^{(Q)}, h_i^{(Q)}$):

No.	loc.coord.	$K(\theta)$	Jacobian	$(k_1^{(Q)}, k_2^{(Q)})$	$(h_1^{(Q)}, h_2^{(Q)})$	λ
(a)	(θ_1, θ_2)	$\theta_1^2 a_1(\theta_1, \theta_2)$	θ_1	$(2, 0)$	$(1, 0)$	1
(b)	$(\theta_1', \theta_2) \neq 0$	$\theta_2^2 a_2(\theta_1', \theta_2)$	θ_2	$(0, 2)$	$(0, 1)$	1
(b1)	(θ_1', θ_2'')	$\theta_1^4 \theta_2''^2 a_3(\theta_1', \theta_2'')$	$\theta_1^2 \theta_2''$	$(4, 2)$	$(2, 1)$	3/4
(b2)	(θ_1'', θ_2)	$\theta_2^4 a_4(\theta_1'', \theta_2)$	θ_2^2	$(0, 4)$	$(0, 2)$	3/4

Thus, the real log canonical threshold is confirmed to be 3/4 (multiplicity is 1).

Example 3.2 (Case of $m = \infty$). Let X be a random variable following a Bernoulli distribution $\text{Bin}(1, \theta)$, where θ is the parameter ($0 < \theta < 1$).

$$\tilde{p}(X = x|\theta) = \theta^x (1 - \theta)^{1-x} = \begin{cases} 1 - \theta, & (x = 0) \\ \theta, & (x = 1) \end{cases}$$

Similar to Example 3.1, consider a mixed distribution model with parameters (θ_1, θ_2) .

$$p(X = x|\theta_1, \theta_2) := \frac{1}{2} \cdot \tilde{p}\left(X = x \middle| \theta_1 - \theta_2 + \frac{1}{2}\right) + \frac{1}{2} \cdot \tilde{p}\left(X = x \middle| \theta_2 + \frac{1}{2}\right) \quad (x = 0, 1) \quad (3.6)$$

The true distribution is $\tilde{p}(X|1/2)$. Simple calculations show that

$$p(X = x|\theta_1, \theta_2) = \begin{cases} \frac{1-\theta_1}{2}, & (x = 0) \\ \frac{1+\theta_1}{2}, & (x = 1) \end{cases}$$

indicating that this model follows a Bernoulli distribution with parameter $(1 + \theta_1)/2$ and does not depend on θ_2 .

The Taylor expansion of $f(X|\theta)$ and $K(\theta)$ at $(\theta_1, \theta_2) = 0$ is computed as follows:

$$\begin{aligned} f(X|\theta) &= \begin{cases} \sum_{n=1}^{\infty} \frac{1}{n} \theta_1^n, & (x = 0) \\ \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \theta_1^n, & (x = 1) \end{cases} \\ K(\theta) &= \sum_{n \geq 2, n: \text{even}}^{\infty} \frac{1}{n} \theta_1^n \end{aligned} \quad (3.7)$$

Verify that Main Theorem 1 holds at $(\theta_1, \theta_2) = 0$.

$$\left. \frac{\partial^n f}{\partial \theta_2^n} \right|_{(\theta_1, \theta_2)=0} = 0 (\forall n \geq 1), \quad \left. \frac{\partial f}{\partial \theta_1} \right|_{(\theta_1, \theta_2)=0} = \begin{cases} 1, & (x = 0) \\ -1, & (x = 1) \end{cases} \neq 0$$

thus satisfying Assumption 1(1)(2) for $m = \infty$,

$$\frac{1}{2} \mathbb{E}_X [F_1(\theta_1)^2] = \frac{1}{2} \mathbb{E}_X \left[\left(\left. \frac{\partial f}{\partial \theta_1} \right|_{(\theta_1, \theta_2)=0} \right)^2 \theta_1^2 \right] = \frac{1}{2} \theta_1^2$$

matching the second order term in (3.7). Furthermore, as asserted in Main Theorem 1, terms consisting only of θ_2 , first-degree terms in θ_1 , and second-degree terms consisting only of θ_1 are not included in the (higher order terms).

Verify that Corollary 3.1 holds. Since Assumption 1(1)(2) is met for $m = \infty$, Corollary 3.1(2) can be applied, giving the real log canonical threshold at the origin as $1/2$ (multiplicity is 1). Indeed, from (3.7),

$$K(\theta) = \theta_1^2 a(\theta_1, \theta_2), \quad a(0, 0) = \frac{1}{2} \neq 0$$

indicates that a normal crossing is already achieved at $(\theta_1, \theta_2) = 0$, confirming that the real log canonical threshold is indeed $1/2$ (multiplicity is 1).

3.2 Proof of Main Theorem 1

The proof of Main Theorem 1 is carried out in several steps. Initially, we present formulas for higher derivatives of the log-likelihood ratio function

$$f(x|\theta) = \log \frac{p(x|\theta_*)}{p(x|\theta)}$$

and the Kullback-Leibler divergence $K(\theta)$

$$K(\theta) = \mathbb{E}_X [f(X|\theta)].$$

For this purpose, we define the quantity $G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)$ for $1 \leq i_1, \dots, i_n \leq d$ and $x \in \chi, \theta \in \Theta$ as given by Equation (3.8).

Consider partitions of the set $\{1, \dots, n\}$, where $n \geq 1$. For example, for $n = 2$, there is one possible partition: $\{\{1\}, \{2\}\}$. For $n = 3$, there are four possible partitions:

$$\{\{\{1\}, \{2\}, \{3\}\}, \{\{1\}, \{2, 3\}\}, \{\{2\}, \{1, 3\}\}, \{\{3\}, \{1, 2\}\}\}$$

The set partitions are structured as:

$$\bigcup_i U_i = \{1, \dots, n\}, \quad U_i \cap U_j = \{\}, \quad U_i \neq \{\}, \{1, \dots, n\}$$

Each level of the set is arbitrarily ordered and presented as a sequence. For instance, for $n = 2$, it can be represented as $(T_1 = ((1), (2)))$. For $n = 3$, it can be represented as $(T_1 = ((1), (2), (3)), T_2 = ((1), (2, 3)), T_3 = ((2), (1, 3)), T_4 = ((3), (1, 2)))$. These sequences are then relabeled with $\{i_1, \dots, i_n\}$ replacing $\{1, \dots, n\}$, denoted as S_{i_1, \dots, i_n} . For example:

$$\begin{aligned} S_{1,3} &= (T_1 = ((1), (3))) \\ S_{1,1,4} &= (T_1 = ((1), (1), (4)), T_2 = ((1), (1, 4)), T_3 = ((1), (1, 4)), T_4 = ((4), (1, 1))) \\ S_{1,1,1} &= (T_1 = ((1), (1), (1)), T_2 = ((1), (1, 1)), T_3 = ((1), (1, 1)), T_4 = ((1), (1, 1))) \end{aligned}$$

Here, we write:

$$k \in U, U \in T, T \in S_{i_1, \dots, i_n}$$

to denote that the sequence S_{i_1, \dots, i_n} includes T , and T includes U , which contains the integer k . The length of U is denoted by $|U|$, and we define:

$$G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta) := \sum_{T \in S_{i_1, \dots, i_n}} \prod_{U \in T} \frac{\partial^{|U|} \log p(x|\theta)}{\prod_{k \in U} \partial \theta_k} \quad (3.8)$$

For instance,

$$\begin{aligned} G_{\theta_1 \theta_3}(x, \theta) &= \frac{\partial \log p(x|\theta)}{\partial \theta_1} \frac{\partial \log p(x|\theta)}{\partial \theta_3} \\ G_{\theta_1 \theta_1 \theta_4}(x, \theta) &= \left\{ \frac{\partial \log p(x|\theta)}{\partial \theta_1} \right\}^2 \frac{\partial \log p(x|\theta)}{\partial \theta_4} + 2 \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_1} \frac{\partial^2 \log p(x|\theta)}{\partial \theta_1 \partial \theta_4} \\ &\quad + \frac{\partial \log p(x|\theta)}{\partial \theta_4} \frac{\partial^2 \log p(x|\theta)}{\partial \theta_1^2} \\ G_{\theta_1 \theta_1 \theta_1}(x, \theta) &= \left\{ \frac{\partial \log p(x|\theta)}{\partial \theta_1} \right\}^3 + 3 \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_1} \frac{\partial^2 \log p(x|\theta)}{\partial \theta_1^2} \end{aligned}$$

Clearly, the definition of $G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)$ does not depend on the definition of S_{i_1, \dots, i_n} (the method of ordering sets into sequences). In the following, $G_{\theta_1 \theta_1 \theta_4}(x, \theta)$ and $G_{\theta_1 \theta_1 \theta_1}(x, \theta)$ are also denoted as $G_{\theta_1^2 \theta_4}(x, \theta)$ and $G_{\theta_1^3}(x, \theta)$, respectively.

Lemma 3.1. The function $G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)$ satisfies the recurrence relation

$$G_{\theta_{i_1} \dots \theta_{i_{n+1}}}(x, \theta) = G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta) \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_{i_{n+1}}} + \frac{\partial^n \log p(x|\theta)}{\partial \theta_{i_1} \dots \partial \theta_{i_n}} \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_{i_{n+1}}} + \frac{\partial G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)}{\partial \theta_{i_{n+1}}}$$

Proof. Consider elements of $S_{i_1, \dots, i_{n+1}}$ that either include or exclude the sequence $U = (i_{n+1})$. For sequences that include $U = (i_{n+1})$, the differential of $\log p(x|\theta)$ corresponding to the sequence $((i_1, \dots, i_n), (i_{n+1}))$ is

$$\frac{\partial^n \log p(x|\theta)}{\partial \theta_{i_1} \dots \partial \theta_{i_n}} \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_{i_{n+1}}}$$

and for others, it corresponds to

$$G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta) \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_{i_{n+1}}}$$

For example, the differential of $\log p(x|\theta)$ corresponding to the sequence $((i_1), (i_2, \dots, i_n), (i_{n+1}))$ is,

$$\left(\frac{\partial \log p(x|\theta)}{\partial \theta_{i_1}} \frac{\partial^{n-1} \log p(x|\theta)}{\partial \theta_{i_2} \dots \partial \theta_{i_n}} \right) \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_{i_{n+1}}}$$

where

$$\frac{\partial \log p(x|\theta)}{\partial \theta_{i_1}} \frac{\partial^{n-1} \log p(x|\theta)}{\partial \theta_{i_2} \cdots \partial \theta_{i_n}}$$

is one of the components constructing $G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)$.

On the other hand, sequences that do not include $U = (i_{n+1})$ correspond to

$$\frac{\partial G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)}{\partial \theta_{i_{n+1}}}$$

by the product rule. For instance, for the elements of $S_{i_1, \dots, i_{n+1}}$ corresponding to the sequences $((i_1, i_{n+1}), (i_2, \dots, i_n)), ((i_1), (i_2, \dots, i_{n+1}))$, the differential of $\log p(x|\theta)$ is,

$$\begin{aligned} & \frac{\partial^2 \log p(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_{n+1}}} \frac{\partial^{n-1} \log p(x|\theta)}{\partial \theta_{i_2} \cdots \partial \theta_{i_n}} + \frac{\partial \log p(x|\theta)}{\partial \theta_{i_1}} \frac{\partial^n \log p(x|\theta)}{\partial \theta_{i_2} \cdots \partial \theta_{i_{n+1}}} \\ &= \frac{\partial}{\partial \theta_{i_{n+1}}} \left(\frac{\partial \log p(x|\theta)}{\partial \theta_{i_1}} \frac{\partial^{n-1} \log p(x|\theta)}{\partial \theta_{i_2} \cdots \partial \theta_{i_n}} \right) \end{aligned}$$

Hence, the recurrence relation as stated in the lemma is verified. \square

Proposition 3.1.

(1)

$$\frac{\partial^n f(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} = - \frac{\frac{\partial^n p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}}}{p(x|\theta)} + G_{\theta_{i_1} \dots \theta_{i_n}}(x, \theta)$$

(2) When the order of partial differentiation and integration can be exchanged,

$$\frac{\partial^n K}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \Big|_{\theta=\theta_*} = \mathbb{E}_X \left[G_{\theta_{i_1} \dots \theta_{i_n}}(X, \theta_*) \right]$$

which means that the first term on the right side of (1) becomes a random variable with expected value zero at $\theta = \theta_*$.

Proof.

(1) Demonstrate using induction on n . For $n = 1$, the case is evident from

$$\frac{\partial f(x|\theta)}{\partial \theta_{i_1}} = - \frac{\frac{\partial p(x|\theta)}{\partial \theta_{i_1}}}{p(x|\theta)}$$

For a general n , using the induction hypothesis,

$$\begin{aligned}
\frac{\partial^{n+1} f(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_{n+1}}} &= \frac{\partial}{\partial \theta_{i_{n+1}}} \left\{ -\frac{\frac{\partial^n p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}}}{p(x|\theta)} + G_{\theta_{i_1} \cdots \theta_{i_n}}(x, \theta) \right\} \\
&= -\frac{\frac{\partial^{n+1} p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_{n+1}}}}{p(x|\theta)} + \frac{\frac{\partial^n p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}}}{p(x|\theta)} \cdot \frac{\partial p(x|\theta)}{\partial \theta_{i_{n+1}}} + \frac{\partial G_{\theta_{i_1} \cdots \theta_{i_n}}(x, \theta)}{\partial \theta_{i_{n+1}}} \\
&= -\frac{\frac{\partial^{n+1} p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_{n+1}}}}{p(x|\theta)} + \left\{ G_{\theta_{i_1} \cdots \theta_{i_n}}(x, \theta) + \frac{\partial^n \log p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \right\} \cdot \frac{\partial \log p(x|\theta)}{\partial \theta_{i_{n+1}}} + \frac{\partial G_{\theta_{i_1} \cdots \theta_{i_n}}(x, \theta)}{\partial \theta_{i_{n+1}}} \\
&= -\frac{\frac{\partial^{n+1} p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_{n+1}}}}{p(x|\theta)} + G_{\theta_{i_1} \cdots \theta_{i_{n+1}}}(x, \theta)
\end{aligned}$$

The last equality uses Lemma 3.1.

- (2) Given the assumption that differentiation and integration can be interchanged, using the result from (1), it suffices to show

$$\mathbb{E}_X \left[\frac{\frac{\partial^n p(X|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}}}{p(X|\theta)} \right] \Bigg|_{\theta=\theta_*} = 0$$

as

$$\begin{aligned}
\mathbb{E}_X \left[\frac{\frac{\partial^n p(X|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}}}{p(X|\theta)} \right] \Bigg|_{\theta=\theta_*} &= \int_{\mathcal{X}} \frac{\frac{\partial^n p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \Big|_{\theta=\theta_*}}{p(x|\theta_*)} \cdot q(x) dx = \int_{\mathcal{X}} \frac{\frac{\partial^n p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \Big|_{\theta=\theta_*}}{p(x|\theta_*)} \cdot p(x|\theta_*) dx \\
&= \int_{\mathcal{X}} \frac{\partial^n p(x|\theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \Big|_{\theta=\theta_*} dx = \frac{\partial^n}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \int_{\mathcal{X}} p(x|\theta) dx \Big|_{\theta=\theta_*} = \frac{\partial^n 1}{\partial \theta_{i_1} \cdots \partial \theta_{i_n}} \Big|_{\theta=\theta_*} = 0
\end{aligned}$$

□

Remark 3.5. Using Proposition 3.1, if we express the derivatives of the log-likelihood ratio

function $f(x|\theta)$ up to the fourth order, defining $Y_{i_1 \dots i_n} := \frac{\partial^n \log p(x|\theta)}{\partial \theta_{i_1} \dots \partial \theta_{i_n}}$, we have:

$$\begin{aligned} \frac{\partial f(x|\theta)}{\partial \theta_{i_1}} &= -\frac{\frac{\partial p(x|\theta)}{\partial \theta_{i_1}}}{p(x|\theta)} \\ \frac{\partial^2 f(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_2}} &= -\frac{\frac{\partial^2 p(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_2}}}{p(x|\theta)} + Y_{i_1} Y_{i_2} \end{aligned} \quad (3.9)$$

$$\frac{\partial^3 f(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_2} \partial \theta_{i_3}} = -\frac{\frac{\partial^3 p(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_2} \partial \theta_{i_3}}}{p(x|\theta)} + Y_{i_1} Y_{i_2} Y_{i_3} + Y_{i_1, i_2} Y_{i_3} + Y_{i_2, i_3} Y_{i_1} + Y_{i_3, i_1} Y_{i_2} \quad (3.10)$$

$$\begin{aligned} \frac{\partial^4 f(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_2} \partial \theta_{i_3} \partial \theta_{i_4}} &= -\frac{\frac{\partial^4 p(x|\theta)}{\partial \theta_{i_1} \partial \theta_{i_2} \partial \theta_{i_3} \partial \theta_{i_4}}}{p(x|\theta)} + Y_{i_1} Y_{i_2} Y_{i_3} Y_{i_4} \\ &\quad + Y_{i_1, i_2} Y_{i_3} Y_{i_4} + Y_{i_1, i_3} Y_{i_2} Y_{i_4} + Y_{i_1, i_4} Y_{i_2} Y_{i_3} \\ &\quad + Y_{i_2, i_3} Y_{i_1} Y_{i_4} + Y_{i_2, i_4} Y_{i_1} Y_{i_3} + Y_{i_3, i_4} Y_{i_1} Y_{i_2} \\ &\quad + Y_{i_1, i_2} Y_{i_3, i_4} + Y_{i_1, i_3} Y_{i_2, i_4} + Y_{i_1, i_4} Y_{i_2, i_3} \\ &\quad + Y_{i_1, i_2, i_3} Y_{i_4} + Y_{i_1, i_2, i_4} Y_{i_3} + Y_{i_1, i_3, i_4} Y_{i_2} + Y_{i_2, i_3, i_4} Y_{i_1} \end{aligned} \quad (3.11)$$

In particular, using (3.9), we find:

$$\begin{aligned} \frac{\partial^2 K(\theta)}{\partial \theta_{i_1} \partial \theta_{i_2}} \Big|_{\theta=\theta_*} &= \mathbb{E}_X \left[\frac{\partial \log p(X|\theta)}{\partial \theta_{i_1}} \Big|_{\theta=\theta_*} \cdot \frac{\partial \log p(X|\theta)}{\partial \theta_{i_2}} \Big|_{\theta=\theta_*} \right] \\ &= \text{Cov} \left(\frac{\partial \log p(X|\theta)}{\partial \theta_{i_1}} \Big|_{\theta=\theta_*}, \frac{\partial \log p(X|\theta)}{\partial \theta_{i_2}} \Big|_{\theta=\theta_*} \right) \end{aligned}$$

Thus, under the conditions of this paper, the Fisher information matrix I and J coincide.

Based on Proposition 3.1(2), it is necessary to determine the expectation of the random variable $G(X, 0)$ when considering the derivatives of $K(\theta)$. Below, we provide specific calculations for $G(X, 0)$ under Assumption 1(1)(2).

Proposition 3.2. Assume that Assumption 1(1)(2) is satisfied. That is, for any non-negative integers (i_{r+1}, \dots, i_d) that satisfy $i_{r+1} + \dots + i_d \leq m - 1$, the log-likelihood ratio function $f(X|\theta)$ satisfies:

$$\frac{\partial^{i_{r+1} + \dots + i_d} f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0} = 0 \quad a.s.$$

Under this condition, the following holds:

- (1) For $i_{r+1} + \dots + i_d \leq 2m - 1$,

$$G_{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) = 0 \quad (a.s.)$$

(2) For $1 \leq j \leq r$, $i_{r+1} + \dots + i_d \leq m - 1$,

$$G_{\theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) = 0 \text{ (a.s.)}$$

(3) For $1 \leq j \leq r$, $i_{r+1} + \dots + i_d = m$,

$$G_{\theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) = \frac{\partial f(X|\theta)}{\partial \theta_j} \Big|_{(\theta_1, \dots, \theta_d)=0} \cdot \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0} \text{ a.s.}$$

(4) For $i_{r+1} + \dots + i_d = 2m$,

$$\begin{aligned} & G_{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) \\ &= \frac{1}{2} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \Big|_{\theta=0} \cdot \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}-j_{r+1}} \dots \partial \theta_d^{i_d-j_d}} \Big|_{\theta=0} \text{ a.s.} \end{aligned}$$

Proof.

(1) For any tuple of non-negative integers (i_{r+1}, \dots, i_d) that satisfies $i_{r+1} + \dots + i_d \leq 2m - 1$, each term of $G_{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}$ takes the form:

$$\prod_{U \in T} \frac{\partial^{|U|} \log p(x|\theta)}{\prod_{k \in U} \partial \theta_k} \tag{3.12}$$

where T consists of multiple non-empty proper subsets of the set

$$\underbrace{\{r+1, \dots, r+1\}}_{\# = i_{r+1}}, \dots, \underbrace{\{d, \dots, d\}}_{\# = i_d}$$

Thus, if $\sum_{k=r+1}^d i_k \leq 2m - 1$, it includes some U_1 where $|U_1| \leq m - 1$. By assumption,

$$\frac{\partial^{|U_1|} \log p(x|\theta)}{\prod_{k \in U_1} \partial \theta_k} \Big|_{(\theta_1, \dots, \theta_d)=0} = 0 \text{ (a.s.)} \tag{3.13}$$

and hence from (3.12) it follows that $G_{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) = 0$ (a.s.).

(2) Similarly to (1), each term of $G_{\theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}$ takes the form of (3.12), where T consists of multiple non-empty proper subsets of the set

$$\underbrace{\{j, r+1, \dots, r+1\}}_{\# = i_{r+1}}, \dots, \underbrace{\{d, \dots, d\}}_{\# = i_d}$$

Thus including some U_2 that does not contain element $\{j\}$. By assumption,

$$\left. \frac{\partial^{|U_2|} \log p(x|\theta)}{\prod_{k \in U_2} \partial \theta_k} \right|_{(\theta_1, \dots, \theta_d)=0} = 0 \text{ (a.s.)} \quad (3.14)$$

and hence from (3.12) it follows that $G_{\theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) = 0$ (a.s.).

(3) As with (2), consider that all T other than

$$T_0 = ((j), \underbrace{(r+1, \dots, r+1)}_{\# = i_{r+1}}, \dots, \underbrace{(d, \dots, d)}_{\# = i_d})$$

contain terms in the form of (3.14) as factors and therefore do not need to be considered when computing G . Given that such T_0 is unique,

$$\begin{aligned} G_{\theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) &= \prod_{U \in T_0} \left. \frac{\partial^{|U|} \log p(X|\theta)}{\prod_{k \in U} \partial \theta_k} \right|_{(\theta_1, \dots, \theta_d)=0} \\ &= \left. \frac{\partial \log p(X|\theta)}{\partial \theta_j} \right|_{(\theta_1, \dots, \theta_d)=0} \cdot \left. \frac{\partial^m \log p(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} \\ &= \left. \frac{\partial f(X|\theta)}{\partial \theta_j} \right|_{(\theta_1, \dots, \theta_d)=0} \cdot \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} \quad \text{a.s.} \end{aligned}$$

is shown.

(4) Similar to (1),

$$j_{r+1} + \dots + j_d = m, \quad i_h \geq j_h \geq 0$$

for the integer tuples (j_{r+1}, \dots, j_d) , define

$$T_0 = (\underbrace{(r+1, \dots, r+1)}_{\# = j_{r+1}}, \dots, \underbrace{(d, \dots, d)}_{\# = j_d}, \underbrace{(r+1, \dots, r+1)}_{\# = i_{r+1} - j_{r+1}}, \dots, \underbrace{(d, \dots, d)}_{\# = i_d - j_d})$$

Since T_0 is the only tuple considered in calculating G because other T s contain factors in the form of (3.13), note that the number of such T_0 depending on (j_{r+1}, \dots, j_d) is

$$\binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \times \begin{cases} \frac{1}{2} & \text{if } (2j_{r+1}, \dots, 2j_d) = (i_{r+1}, \dots, i_d) \\ 1 & \text{else} \end{cases}$$

Given this, it has been demonstrated:

$$\begin{aligned}
G_{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) &= \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \prod_{U \in T_0} \frac{\partial^{|U|} \log p(X|\theta)}{\prod_{k \in U} \partial \theta_k} \Big|_{(\theta_1, \dots, \theta_d) = 0} \\
&= \frac{1}{2} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \frac{\partial^m \log p(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \Big|_{\theta=0} \cdot \frac{\partial^m \log p(X|\theta)}{\partial \theta_{r+1}^{i_{r+1} - j_{r+1}} \dots \partial \theta_d^{i_d - j_d}} \Big|_{\theta=0} \\
&= \frac{1}{2} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \Big|_{\theta=0} \cdot \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1} - j_{r+1}} \dots \partial \theta_d^{i_d - j_d}} \Big|_{\theta=0} \quad \text{a.s.}
\end{aligned}$$

Note that in the transformation, we used the fact that

$$\binom{i_{r+1}}{i_{r+1} - j_{r+1}} \dots \binom{i_d}{i_d - j_d} = \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d}$$

□

Example 3.3. Consider the case of $d = 2, m = 2$ in Proposition 3.2(1)(4). Suppose for any non-negative integer pair (i_1, i_2) satisfying $i_1 + i_2 \leq 1$, the log-likelihood function $f(X|\theta)$ satisfies

$$\frac{\partial^{i_1+i_2} f(X|\theta)}{\partial \theta_1^{i_1} \partial \theta_2^{i_2}} \Big|_{(\theta_1, \theta_2) = 0} = 0 \quad \text{a.s.}$$

Specifically, assume

$$\frac{\partial f(X|\theta)}{\partial \theta_1} \Big|_{(\theta_1, \theta_2) = 0} = \frac{\partial f(X|\theta)}{\partial \theta_2} \Big|_{(\theta_1, \theta_2) = 0} = 0 \quad \text{a.s.} \quad (3.15)$$

For $(i_1, i_2) = (1, 2)$ in Proposition 3.2(1), we expect

$$G_{\theta_1 \theta_2^2}(X, 0) = 0 \quad \text{a.s.}$$

Indeed, from Remark 3.5(3.10),

$$\begin{aligned}
&G_{\theta_1 \theta_2^2}(X, \theta) \\
&= \frac{\partial \log p(X|\theta)}{\partial \theta_1} \left(\frac{\partial \log p(X|\theta)}{\partial \theta_2} \right)^2 + 2 \frac{\partial^2 \log p(X|\theta)}{\partial \theta_1 \partial \theta_2} \cdot \frac{\partial \log p(X|\theta)}{\partial \theta_2} + \frac{\partial^2 \log p(X|\theta)}{\partial \theta_2^2} \cdot \frac{\partial \log p(X|\theta)}{\partial \theta_1}
\end{aligned}$$

where each term includes a first derivative of f , which are all zero at $\theta = 0$ by (3.15), confirming that indeed $G_{\theta_1 \theta_2^2}(X, 0) = 0$ a.s.

Next, for $(i_1, i_2) = (1, 3)$ in Proposition 3.2(4),

$$\begin{aligned}
G_{\theta_1\theta_2^3}(X, 0) &= \frac{1}{2} \sum_{\substack{j_1+j_2=2 \\ i_h \geq j_h \geq 0}} \binom{1}{j_1} \binom{3}{j_2} \frac{\partial^2 f(X|\theta)}{\partial \theta_1^{j_1} \partial \theta_2^{j_2}} \Big|_{\theta=0} \cdot \frac{\partial^2 f(X|\theta)}{\partial \theta_1^{1-j_1} \partial \theta_2^{3-j_2}} \Big|_{\theta=0} \quad \text{a.s.} \\
&= \frac{3}{2} \frac{\partial^2 f(X|\theta)}{\partial \theta_2^2} \Big|_{\theta=0} \cdot \frac{\partial^2 f(X|\theta)}{\partial \theta_1 \partial \theta_2} \Big|_{\theta=0} + \frac{3}{2} \frac{\partial^2 f(X|\theta)}{\partial \theta_1 \partial \theta_2} \Big|_{\theta=0} \cdot \frac{\partial^2 f(X|\theta)}{\partial \theta_2^2} \Big|_{\theta=0} \quad \text{a.s.} \\
&= 3 \frac{\partial^2 f(X|\theta)}{\partial \theta_2^2} \Big|_{\theta=0} \cdot \frac{\partial^2 f(X|\theta)}{\partial \theta_1 \partial \theta_2} \Big|_{\theta=0} \quad \text{a.s.}
\end{aligned}$$

which should hold true. Indeed, considering only terms involving second derivatives as per Remark 3.5(3.11),

$$G_{\theta_1\theta_2^3}(X, 0) = 3 \frac{\partial^2 f(X|\theta)}{\partial \theta_2^2} \Big|_{\theta=0} \cdot \frac{\partial^2 f(X|\theta)}{\partial \theta_1 \partial \theta_2} \Big|_{\theta=0} \times 3 \quad \text{a.s.}$$

confirming that indeed Proposition 3.2(4) holds.

Proof of Main Theorem 1.

There are five points to be proven regarding the terms appearing in the Taylor expansion of $K(\theta)$ at $\theta = 0$:

- (1) The coefficients of the terms of order 0 in $(\theta_1, \dots, \theta_r)$ and order up to $2m-1$ in $(\theta_{r+1}, \dots, \theta_d)$ are zero.
- (2) The coefficients of the terms of order 1 in $(\theta_1, \dots, \theta_r)$ and order up to $m-1$ in $(\theta_{r+1}, \dots, \theta_d)$ are zero.
- (3) The terms of order 1 in $(\theta_1, \dots, \theta_r)$ and order m in $(\theta_{r+1}, \dots, \theta_d)$ are represented by:

$$\mathbb{E}_X [F_1(X|\theta_1, \dots, \theta_r) F_m(X|\theta_{r+1}, \dots, \theta_d)]$$

- (4) The terms of order 0 in $(\theta_1, \dots, \theta_r)$ and order $2m$ in $(\theta_{r+1}, \dots, \theta_d)$ are represented by:

$$\frac{1}{2} \mathbb{E}_X [F_m^2(X|\theta_{r+1}, \dots, \theta_d)]$$

- (5) The terms of order 2 in $(\theta_1, \dots, \theta_r)$ and order 0 in $(\theta_{r+1}, \dots, \theta_d)$ are represented by:

$$\frac{1}{2} \mathbb{E}_X [F_1^2(X|\theta_1, \dots, \theta_r)]$$

Assuming the interchange of differentiation and integration, using Proposition 3.1(2), for any tuple of non-negative integers (i_1, \dots, i_d) , the coefficient of the term $\theta_1^{i_1} \dots \theta_d^{i_d}$ in the Taylor expansion of $K(\theta)$ at $\theta = 0$ can be expressed as

$$\frac{1}{i_1! \dots i_d!} \left. \frac{\partial^{i_1 + \dots + i_d} K(\theta)}{\partial \theta_1^{i_1} \dots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} = \frac{1}{i_1! \dots i_d!} \mathbb{E}_X \left[G_{\theta_1^{i_1} \dots \theta_d^{i_d}}(X, 0) \right] \quad (3.16)$$

First, concerning points (1) and (2), according to Proposition 3.2(1)(2), the right-hand side G is zero as a random variable almost surely. Therefore, (3.16) = 0. Thus, the theorem is demonstrated.

Next, for point (3), as per Proposition 3.2(3) for $1 \leq j \leq r$ and $i_{r+1} + \dots + i_d = m$,

$$G_{\theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) = \left. \frac{\partial f(X|\theta)}{\partial \theta_j} \right|_{(\theta_1, \dots, \theta_d)=0} \cdot \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} \quad \text{a.s.}$$

From (3.16), the term in question is given by,

$$\begin{aligned} & \sum_{j=1}^r \sum_{\substack{i_{r+1} + \dots + i_d = m \\ i_h \geq 0}} \frac{1}{i_{r+1}! \dots i_d!} \left. \frac{\partial^{m+1} K(\theta)}{\partial \theta_j \partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} \times \theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d} \\ &= \sum_{j=1}^r \sum_{\substack{i_{r+1} + \dots + i_d = m \\ i_h \geq 0}} \frac{1}{i_{r+1}! \dots i_d!} \mathbb{E}_X \left[\left. \frac{\partial f(X|\theta)}{\partial \theta_j} \right|_{\theta=0} \cdot \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{\theta=0} \right] \times \theta_j \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d} \\ &= \mathbb{E}_X [F_1(X|\theta_1, \dots, \theta_r) F_m(X|\theta_{r+1}, \dots, \theta_d)] \end{aligned}$$

Thus, the expression can be represented as shown.

(5) is a special case of (4) ($m = 1$), so the proof is completed by demonstrating (4). By Proposition 3.2(4), for $i_{r+1} + \dots + i_d = 2m$,

$$\begin{aligned} & G_{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}(X, 0) \\ &= \frac{1}{2} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \right|_{\theta=0} \cdot \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}-j_{r+1}} \dots \partial \theta_d^{i_d-j_d}} \right|_{\theta=0} \quad \text{a.s.} \end{aligned}$$

Thus, from (3.16),

$$\begin{aligned} & \left. \frac{\partial^{2m} K(\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{\theta=0} \\ &= \frac{1}{2} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \mathbb{E}_X \left[\left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \right|_{\theta=0} \cdot \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}-j_{r+1}} \dots \partial \theta_d^{i_d-j_d}} \right|_{\theta=0} \right] \end{aligned}$$

Using this, we derive from the target formula:

$$\begin{aligned}
\frac{1}{2} \mathbb{E}_X [F_m^2(X|\theta_{r+1}, \dots, \theta_d)] &= \frac{1}{2} \mathbb{E}_X \left[\left(\sum_{\substack{k_{r+1} + \dots + k_d = m \\ k_h \geq 0}} \frac{\theta_{r+1}^{k_{r+1}} \dots \theta_d^{k_d}}{k_{r+1}! \dots k_d!} \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{k_{r+1}} \dots \partial \theta_d^{k_d}} \Big|_{\theta=0} \right)^2 \right] \\
&= \frac{1}{2} \sum_{\substack{k_{r+1} + \dots + k_d = m \\ k_h \geq 0}} \sum_{\substack{l_{r+1} + \dots + l_d = m \\ l_h \geq 0}} \frac{\theta_{r+1}^{k_{r+1} + l_{r+1}} \dots \theta_d^{k_d + l_d}}{k_{r+1}! \dots k_d! l_{r+1}! \dots l_d!} \\
&\quad \times \mathbb{E}_X \left[\frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{k_{r+1}} \dots \partial \theta_d^{k_d}} \Big|_{\theta=0} \cdot \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{l_{r+1}} \dots \partial \theta_d^{l_d}} \Big|_{\theta=0} \right] \\
&= \frac{1}{2} \sum_{\substack{i_{r+1} + \dots + i_d = 2m \\ i_h \geq 0}} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \frac{\theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}}{j_{r+1}! \dots j_d! (i_{r+1} - j_{r+1})! \dots (i_d - j_d)!} \\
&\quad \times \mathbb{E}_X \left[\frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \Big|_{\theta=0} \cdot \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1} - j_{r+1}} \dots \partial \theta_d^{i_d - j_d}} \Big|_{\theta=0} \right] \\
&= \sum_{\substack{i_{r+1} + \dots + i_d = 2m \\ i_h \geq 0}} \frac{1}{i_{r+1}! \dots i_d!} \cdot \frac{1}{2} \sum_{\substack{j_{r+1} + \dots + j_d = m \\ i_h \geq j_h \geq 0}} \binom{i_{r+1}}{j_{r+1}} \dots \binom{i_d}{j_d} \\
&\quad \times \mathbb{E}_X \left[\frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{j_{r+1}} \dots \partial \theta_d^{j_d}} \Big|_{\theta=0} \cdot \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1} - j_{r+1}} \dots \partial \theta_d^{i_d - j_d}} \Big|_{\theta=0} \right] \times \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d} \\
&= \sum_{\substack{i_{r+1} + \dots + i_d = 2m \\ i_h \geq 0}} \frac{1}{i_{r+1}! \dots i_d!} \frac{\partial^{2m} K(\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \Big|_{\theta=0} \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}
\end{aligned}$$

In the algebraic transformation, the variable change $k_h + l_h = i_h$, $k_h = j_h$ (for $h = r + 1, \dots, d$) was performed. Notice the domains before and after the variable transformation are as follows:

$$\begin{aligned}
&\{k_{r+1} + \dots + k_d = m, l_{r+1} + \dots + l_d = m, k_h \geq 0, l_h \geq 0\} \\
&= \{j_{r+1} + \dots + j_d = m, i_{r+1} + \dots + i_d - (j_{r+1} + \dots + j_d) = m, j_h \geq 0, i_h - j_h \geq 0\} \\
&= \{j_{r+1} + \dots + j_d = m, i_{r+1} + \dots + i_d = 2m, i_h \geq j_h \geq 0\}
\end{aligned}$$

□

Remark 3.6. Although it is outside the scope of this paper, semi-regularity in Main Theorem 1 is not essential, and it is possible to generalize to arbitrary orders (n, m) instead of $(1, m)$. Furthermore, it is also possible to generalize to tuples of three or more natural numbers.

3.3 Proof of Main Theorem 2

In this section, we provide the proof of Main Theorem 2. By utilizing the close relationship between the linear independence of random variables and the positive definiteness of quadratic forms (Lemma 3.2), we can rephrase the linear independence assumption of Assumption 1 as the positive definiteness of a certain quadratic form. By using the Taylor expansion of $K(\theta)$ obtained in Main Theorem 1 and repeating specific blow-ups, we can achieve normal crossings and evaluate the real log canonical threshold.

First, we show the close relationship between the linear independence of random variables and the positive definiteness of a certain matrix.

Lemma 3.2. [Characterization of the Linear Independence of Random Variables] For n random variables X_1, \dots, X_n , let $\Sigma := (\mathbb{E}[X_i X_j])_{1 \leq i, j \leq n} \in \mathbb{R}^{n \times n}$.

- (1) Σ is non-negative definite, and the following two conditions are equivalent:
 - (a) Σ is positive definite.
 - (b) X_1, \dots, X_n are linearly independent as random variables.
- (2) Let V be the vector space over \mathbb{R} spanned by the random variables X_1, \dots, X_n . Then, $\text{rank}(\Sigma) = \dim(V)$.

Proof. Let $A(X) := (X_1, \dots, X_n)^\top \in \mathbb{R}^n$, then $\Sigma = \mathbb{E}_X [A(X)A(X)^\top]$. In the following, let $V = \langle A(X) \rangle$ denote the vector space over \mathbb{R} spanned by the random variables X_1, \dots, X_n .

- (1) For $\mathbf{u} := (u_1, \dots, u_n)^\top \in \mathbb{R}^n$,

$$\mathbf{u}^\top \Sigma \mathbf{u} = \mathbf{u}^\top \mathbb{E}_X [A(X)A(X)^\top] \mathbf{u} = \mathbb{E}_X \left[\left\| A(X)^\top \mathbf{u} \right\|^2 \right] \geq 0$$

$$\mathbf{u}^\top \Sigma \mathbf{u} = 0 \Leftrightarrow A(X)^\top \mathbf{u} = 0 \text{ a.s.}$$

Therefore, the two conditions (a) and (b) are equivalent.

- (2) Let $r := \text{rank}(\Sigma)$. There exists a real symmetric matrix $P \in \mathbb{R}^{d \times d}$ such that, using a regular diagonal matrix $D \in \mathbb{R}^{r \times r}$,

$$P \mathbb{E}[A(X)A(X)^\top] P^\top = \left(\begin{array}{c|c} D & 0 \\ \hline 0 & 0 \end{array} \right)$$

Here, if we set $PA(X) = (B(X), C(X))^\top$, where $B(X) \in \mathbb{R}^r$ and $C(X) \in \mathbb{R}^{d-r}$, then

$$\mathbb{E} [B(X)B(X)^\top] = D, \quad \mathbb{E} [C(X)C(X)^\top] = 0$$

From (1), $B(X)$ is linearly independent, and $C(X) = 0$ (a.s.) follows. Consequently,

$$\dim V = \dim \langle A(X) \rangle = \dim \langle PA(X) \rangle = \dim \langle B(X), C(X) \rangle = r$$

follows.

□

Next, we show a key lemma for the proof of Main Theorem 2.

Lemma 3.3.

(1) When Assumption 1(1) is satisfied, the following equivalence holds:

$$\mathbb{E}_X [F_1^2(X|\theta_1, \dots, \theta_r)] = 0 \Leftrightarrow (\theta_1, \dots, \theta_r) = 0$$

(2) Let a be a non-zero constant, and assume $m < \infty$ and that Assumption 1(1)(2)(3) is satisfied.

(i) When $(\theta_{r+1}, \dots, \theta_d)$ satisfies Assumption 1(3)(i), the following equivalence holds:

$$\begin{aligned} \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] &= 0 \\ \Leftrightarrow (\theta_1, \dots, \theta_r) = 0, F_m(X|\theta_{r+1}, \dots, \theta_d) &= 0 \text{ a.s.} \end{aligned}$$

(ii) When $(\theta_{r+1}, \dots, \theta_d) \neq 0$ satisfies Assumption 1(3)(ii), the following holds:

$$\mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] > 0$$

In particular, in either case (i) or (ii), if $(\theta_1, \dots, \theta_r) \neq 0$, then

$$\mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] > 0$$

holds.

Proof.

(1) Let

$$A(X) := \left[\frac{\partial f(X|\theta)}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_d)=0}, \dots, \frac{\partial f(X|\theta)}{\partial \theta_r} \Big|_{(\theta_1, \dots, \theta_d)=0} \right]^\top \in \mathbb{R}^r$$

then

$$\mathbb{E}_X [F_1^2(X|\theta_1, \dots, \theta_r)] = [\theta_1, \dots, \theta_r] \mathbb{E}_X [A(X)A(X)^\top] \begin{bmatrix} \theta_1 \\ \vdots \\ \theta_r \end{bmatrix}$$

is represented as a non-negative quadratic form. Since the components of $A(X)$ are linearly independent by Assumption 1(1), Lemma 3.2(1) implies that $\mathbb{E}_X [A(X)A(X)^\top]$ is positive definite, establishing the proposition.

(2) If Assumption 1(3)(i) holds, then $F_m(X|\theta_{r+1}, \dots, \theta_d) = 0$ a.s., and the proposition follows from (1). Therefore, we only need to consider when Assumption 1(3)(ii) holds. Define

$$R(X|\theta_{r+1}, \dots, \theta_d) := \left[A(X)^\top, a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d) \right]^\top \in \mathbb{R}^{r+1}$$

then,

$$\begin{aligned} & \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] \\ &= [\theta_1, \dots, \theta_r, 1] \mathbb{E}_X \left[R(X|\theta_{r+1}, \dots, \theta_d) R(X|\theta_{r+1}, \dots, \theta_d)^\top \right] \begin{bmatrix} \theta_1 \\ \vdots \\ \theta_r \\ 1 \end{bmatrix} \end{aligned}$$

is expressed as a non-negative definite quadratic form. When Assumption 1(3)(ii) is satisfied, the random variable $a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d)$ and each component of $A(X)$ are linearly independent, and by Lemma 3.2(1),

$$\mathbb{E}_X \left[R(X|\theta_{r+1}, \dots, \theta_d) R(X|\theta_{r+1}, \dots, \theta_d)^\top \right]$$

is positive definite, thus,

$$\mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + a \cdot F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] > 0$$

is obtained. □

Proof of Main Theorem 2.

When Assumption 1(1)(2) is satisfied for $m < \infty$, we can use Main Theorem 1 to express the Taylor expansion of $K(\theta)$ at $(\theta_1, \dots, \theta_d) = 0$ as

$$K(\theta) = \frac{1}{2} \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] + (\text{higher order terms})$$

Here, the (higher order terms) are specifically expressed as the sum of the following four terms:

- $f_0(\theta_{r+1}, \dots, \theta_d)$: terms of order $2m + 1$ or higher
- $f_1(\theta_1, \dots, \theta_d)$: first-degree homogeneous in $\theta_1, \dots, \theta_r$ and of order $m + 1$ or higher in $\theta_{r+1}, \dots, \theta_d$
- $f_2(\theta_1, \dots, \theta_d)$: of order at least second-degree in $\theta_1, \dots, \theta_r$ and at least first-degree in $\theta_{r+1}, \dots, \theta_d$
- $f_3(\theta_1, \dots, \theta_r)$: third-degree or higher

In the following, we demonstrate these in the case of the lowest degree terms, namely:

- $f_0(\theta_{r+1}, \dots, \theta_d)$: homogeneous of degree $2m + 1$
- $f_1(\theta_1, \dots, \theta_d)$: first-degree homogeneous in $\theta_1, \dots, \theta_r$ and homogeneous of degree $m + 1$ in $\theta_{r+1}, \dots, \theta_d$
- $f_2(\theta_1, \dots, \theta_d)$: second-degree homogeneous in $\theta_1, \dots, \theta_r$ and first-degree homogeneous in $\theta_{r+1}, \dots, \theta_d$
- $f_3(\theta_1, \dots, \theta_r)$: homogeneous of third-degree

The general case is proven similarly.

We will consider the real log canonical threshold by performing the following blow-ups:

- (a) Perform a blow-up centered at the origin of \mathbb{R}^d once.
- (b) If the exceptional surface in (a) is $\{\theta_i = 0\}$ (where $i = r + 1, \dots, d$), perform another blow-up centered at the subvariety $\{(\theta_1, \dots, \theta_d) \mid \theta_1 = \dots = \theta_r = \theta_i = 0\}$.
- (c) If the exceptional surface in (b) is $\{\theta_i = 0\}$, repeat (b) until the total number of blow-ups reaches m .

Let's first consider (a).

(a-1) Consider the case where the exceptional surface is $\{\theta_i = 0\} (i = 1, \dots, r)$. For example, in the case $i = 1$, that is, when we perform a blow-up with $\{\theta_2 = \theta_1 \theta'_2, \dots, \theta_d = \theta_1 \theta'_d\}$, the exceptional surface is $\{\theta_1 = 0\}$, and

$$\begin{aligned}
K(\theta) &= \frac{1}{2} \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] \\
&\quad + f_0(\theta_{r+1}, \dots, \theta_d) + f_1(\theta_1, \dots, \theta_d) + f_2(\theta_1, \dots, \theta_d) + f_3(\theta_1, \dots, \theta_r) \\
&= \frac{1}{2} \theta_1^2 \left\{ \mathbb{E}_X \left[\{F_1(X|1, \theta'_2, \dots, \theta'_r) + \theta_1^{m-1} F_m(X|\theta'_{r+1}, \dots, \theta'_d)\}^2 \right] \right. \\
&\quad + \theta_1^{2m-1} f_0(\theta'_{r+1}, \dots, \theta'_d) + \theta_1^m f_1(1, \theta'_2, \dots, \theta'_d) \\
&\quad \left. + \theta_1 f_2(1, \theta'_2, \dots, \theta'_d) + \theta_1 f_3(1, \theta'_2, \dots, \theta'_r) \right\} \\
&= \frac{1}{2} \theta_1^2 a(\theta_1, \theta'_2, \dots, \theta'_d)
\end{aligned}$$

(where a is an analytic function). Considering the point on $U_0 = g^{-1}(O)$ in this local coordinate system $(\theta_1, \theta'_2, \dots, \theta'_d)$, which satisfies $\theta_1 = 0$,

$$a(0, \theta'_2, \dots, \theta'_d) = \begin{cases} \mathbb{E}_X [F_1^2(X|1, \theta'_2, \dots, \theta'_r)] & (m \geq 2) \\ \mathbb{E}_X \left[\{F_1(X|1, \theta'_2, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_d)\}^2 \right] & (m = 1) \end{cases}$$

and by Lemma 3.3(1)(2), we obtain $a(0, \theta'_2, \dots, \theta'_d) > 0$. Therefore, in this local coordinate system, the normal crossing of $K(\theta)$ is obtained at any point Q on U_0 , and

$$\inf_{Q \in U_0} \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \frac{d-1+1}{2} = \frac{d}{2}$$

(multiplicity is 1).

(a-2) Next, consider the case where the exceptional surface is $\{\theta_i = 0\} (i = r+1, \dots, d)$. For example, in the case $i = d$, that is, when we perform a blow-up with $\{\theta_1 = \theta_d \theta'_1, \dots, \theta_{d-1} = \theta_d \theta'_{d-1}\}$, the exceptional surface is $\{\theta_d = 0\}$, and

$$\begin{aligned} K(\theta) &= \frac{1}{2} \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d)\}^2 \right] \\ &\quad + f_0(\theta_{r+1}, \dots, \theta_d) + f_1(\theta_1, \dots, \theta_d) + f_2(\theta_1, \dots, \theta_d) + f_3(\theta_1, \dots, \theta_r) \\ &= \frac{1}{2} \theta_d^2 \left\{ \mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + \theta_d^{m-1} F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1)\}^2 \right] \right. \\ &\quad + \theta_d^{2m-1} f_0(\theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_d^m f_1(\theta'_1, \dots, \theta'_{d-1}, 1) \\ &\quad \left. + \theta_d f_2(\theta'_1, \dots, \theta'_{d-1}, 1) + \theta_d f_3(\theta'_1, \dots, \theta'_r) \right\} \\ &= \frac{1}{2} \theta_d^2 a(\theta'_1, \dots, \theta'_{d-1}, \theta_d) \end{aligned}$$

(where a is an analytic function). Considering the point on $U_0 = g^{-1}(O)$ in this local coordinate system $(\theta'_1, \dots, \theta'_{d-1}, \theta_d)$, which satisfies $\theta_d = 0$,

$$a(\theta'_1, \dots, \theta'_{d-1}, 0) = \begin{cases} \mathbb{E}_X [F_1^2(X|\theta'_1, \dots, \theta'_r)] & (m \geq 2) \\ \mathbb{E}_X [\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1)\}^2] & (m = 1) \end{cases}$$

holds. For any point Q on U_0 that satisfies $(\theta'_1, \dots, \theta'_r) \neq 0$, by Lemma 3.3(1)(2), the normal crossing of $K(\theta)$ is obtained, and

$$\min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \frac{d-1+1}{2} = \frac{d}{2}$$

(multiplicity is 1).

Next, consider the points on U_0 that satisfy $(\theta'_1, \dots, \theta'_r) = 0$. First, for the case of $m = 1$,

$$S = \left\{ (\theta'_1, \dots, \theta'_d) \mid \begin{array}{l} (\theta'_1, \dots, \theta'_r, \theta_d) = 0 \\ F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1) = 0 \quad (\text{a.s.}) \end{array} \right\} \subset U_0$$

For points Q on U_0 that are not included in S , by Lemma 3.3(2), $a(\theta'_1, \dots, \theta'_{d-1}, 0) > 0$, so the normal crossing of $K(\theta)$ is obtained, and

$$\inf_{Q \in U_0 \setminus S} \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \frac{d}{2}$$

(multiplicity is 1). Therefore, for $m = 1$, the theorem is shown.

On the other hand, for $m \geq 2$, $a(\theta'_1, \dots, \theta'_{d-1}, 0) = 0$, and the normal crossing of $K(\theta)$ is not obtained. Therefore, to obtain the normal crossing of $K(\theta)$, it is necessary to further blow-up centered at the subvariety $\{(\theta'_1, \dots, \theta'_{d-1}, \theta_d) \mid \theta'_1 = \dots = \theta'_r = \theta_d = 0\}$.

Hereafter, we assume $m \geq 2$ and use the transformed coordinates without the notation $'$. In this case, note that $r < d$.

For (b), in the local coordinates where the exceptional surface in (a) is $\{\theta_d = 0\}$, further blow-up is performed centered at the subvariety $\{(\theta_1, \dots, \theta_d) \mid \theta_1 = \dots = \theta_r = \theta_d = 0\}$. The same argument applies to other local coordinates.

(b-1) First, consider the case where the exceptional surface is $\{\theta_i = 0\} (i = 1, \dots, r)$. For example, in the case $i = 1$, that is, when we perform a blow-up with $\{\theta_2 = \theta_1\theta'_2, \dots, \theta_r = \theta_1\theta'_r, \theta_d = \theta_1\theta'_d\}$, the exceptional surface is $\{\theta_1 = 0\}$, and

$$\begin{aligned}
K(\theta) &= \frac{1}{2}\theta_d^2 \left\{ \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + \theta_d^{m-1}F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1)\}^2 \right] \right. \\
&\quad \left. + \theta_d^{2m-1}f_0(\theta_{r+1}, \dots, \theta_{d-1}, 1) + \theta_d^m f_1(\theta_1, \dots, \theta_{d-1}, 1) \right. \\
&\quad \left. + \theta_d f_2(\theta_1, \dots, \theta_{d-1}, 1) + \theta_d f_3(\theta_1, \dots, \theta_r) \right\} \\
&= \frac{1}{2}\theta_1^4 \theta_d'^2 \left\{ \mathbb{E}_X \left[\{F_1(X|1, \theta'_2, \dots, \theta'_r) + \theta_1^{m-2} \theta_d'^{m-1} F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1)\}^2 \right] \right. \\
&\quad \left. + \theta_1^{2m-3} \theta_d'^{2m-1} f_0(\theta_{r+1}, \dots, \theta_{d-1}, 1) \right. \\
&\quad \left. + \theta_1^{m-1} \theta_d'^m f_1(1, \theta'_2, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 1) \right. \\
&\quad \left. + \theta_1 \theta_d' f_2(1, \theta'_2, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 1) + \theta_1^2 \theta_d' f_3(1, \theta'_2, \dots, \theta'_r) \right\} \\
&= \frac{1}{2}\theta_1^4 \theta_d'^2 a(\theta_1, \theta'_2, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, \theta'_d)
\end{aligned}$$

(where a is an analytic function). Considering the point on $U_0 = g^{-1}(O)$ in this local coordinate system $(\theta_1, \theta'_2, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, \theta'_d)$, which satisfies $\theta_1 = 0$,

$$\begin{aligned}
&a(0, \theta'_2, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, \theta'_d) \\
&= \begin{cases} \mathbb{E}_X [F_1^2(X|1, \theta'_2, \dots, \theta'_r)] & (m \geq 3) \\ \mathbb{E}_X [\{F_1(X|1, \theta'_2, \dots, \theta'_r) + \theta_d' F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1)\}^2] & (m = 2) \end{cases}
\end{aligned}$$

and by Lemma 3.3(1)(2),

$$a(0, \theta'_2, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, \theta'_d) > 0$$

is obtained. Therefore, in this local coordinate system, the normal crossing of $K(\theta)$ is obtained at any point Q on U_0 , and

$$\inf_{Q \in U_0} \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \min \left(\frac{d+r-1+1}{4}, \frac{d}{2} \right) = \frac{d+r}{4}$$

(multiplicity is 1).

(b-2) Next, consider the case where the exceptional surface is $\{\theta_d = 0\}$. That is, when we perform a blow-up with $\{\theta_1 = \theta_d \theta'_1, \dots, \theta_r = \theta_d \theta'_r\}$,

$$\begin{aligned} K(\theta) &= \frac{1}{2} \theta_d^2 \left\{ \mathbb{E}_X \left[\{F_1(X|\theta_1, \dots, \theta_r) + \theta_d^{m-1} F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1)\}^2 \right] \right. \\ &\quad + \theta_d^{2m-1} f_0(\theta_{r+1}, \dots, \theta_{d-1}, 1) + \theta_d^m f_1(\theta_1, \dots, \theta_{d-1}, 1) \\ &\quad \left. + \theta_d f_2(\theta_1, \dots, \theta_{d-1}, 1) + \theta_d f_3(\theta_1, \dots, \theta_r) \right\} \\ &= \frac{1}{2} \theta_d^4 \left\{ \mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + \theta_d^{m-2} F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1)\}^2 \right] \right. \\ &\quad + \theta_d^{2m-3} f_0(\theta_{r+1}, \dots, \theta_{d-1}, 1) \\ &\quad + \theta_d^{m-1} f_1(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 1) \\ &\quad \left. + \theta_d f_2(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 1) + \theta_d^2 f_3(\theta'_1, \dots, \theta'_r) \right\} \\ &= \frac{1}{2} \theta_d^4 a(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_d) \end{aligned}$$

(where a is an analytic function). Considering the point on $U_0 = g^{-1}(O)$ in this local coordinate system $(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_d)$, which satisfies $\theta_d = 0$,

$$\begin{aligned} &a(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 0) \\ &= \begin{cases} \mathbb{E}_X [F_1^2(X|\theta'_1, \dots, \theta'_r)] & (m \geq 3) \\ \mathbb{E}_X [\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1)\}^2] & (m = 2) \end{cases} \end{aligned}$$

and for any point Q on $\{\theta_d = 0\}$ that satisfies $(\theta'_1, \dots, \theta'_r) \neq 0$, by Lemma 3.3(1)(2), the normal crossing of $K(\theta)$ is obtained, and

$$\min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \frac{d+r}{4}$$

(multiplicity is 1).

Next, consider any point on $\{\theta_d = 0\}$ that satisfies $(\theta'_1, \dots, \theta'_r) = 0$. First, for the case of $m = 2$,

$$S = \left\{ (\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_d) \mid \begin{array}{l} (\theta'_1, \dots, \theta'_r, \theta_d) = 0 \\ F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1) = 0 \quad (\text{a.s.}) \end{array} \right\} \subset U_0$$

For points Q on U_0 that are not included in S , by Lemma 3.3(2), $a(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 0) > 0$ so the normal crossing of $K(\theta)$ is obtained, and

$$\inf_{Q \in U_0 \setminus S} \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \inf \left\{ \frac{d+r}{4}, \frac{d}{2} \right\} = \frac{d+r}{4}$$

(multiplicity is 1). Therefore, for $m = 2$, the theorem is shown.

On the other hand, for $m \geq 3$, $a(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_{d-1}, 0) = 0$, and the normal crossing is not obtained. Therefore, to obtain the normal crossing of $K(\theta)$, it is necessary to further blow-up centered at the subvariety $\{(\theta'_1, \dots, \theta'_r, \theta_{r+1}, \dots, \theta_d) \mid \theta'_1 = \dots = \theta'_r = \theta_d = 0\}$.

Assuming $m \geq 3$ and using the transformed coordinates without the notation $'$, we have:

$$\begin{aligned} K(\theta) = & \frac{1}{2} \theta_d^4 \left\{ \mathbb{E}_X \left[\left\{ F_1(X|\theta_1, \dots, \theta_r) + \theta_d^{m-2} F_m(X|\theta_{r+1}, \dots, \theta_{d-1}, 1) \right\}^2 \right] \right. \\ & + \theta_d^{2m-3} f_0(\theta_{r+1}, \dots, \theta_{d-1}, 1) + \theta_d^{m-1} f_1(\theta_1, \dots, \theta_{d-1}, 1) \\ & \left. + \theta_d f_2(\theta_1, \dots, \theta_{d-1}, 1) + \theta_d^2 f_3(\theta_1, \dots, \theta_r) \right\} \end{aligned}$$

In this coordinate system, we need to find the normal crossing of $K(\theta)$ at any point on the subvariety $\{(\theta_1, \dots, \theta_d) \mid \theta_1 = \dots = \theta_r = \theta_d = 0\}$.

(c) Repeating the above discussion, after performing $m - 1$ blow-ups, that is, for the initial parameters $(\theta_1, \dots, \theta_d)$, consider the transformation

$$\theta_1 = \theta_d^{m-1} \theta'_1, \dots, \theta_r = \theta_d^{m-1} \theta'_r, \theta_{r+1} = \theta_d \theta'_{r+1}, \dots, \theta_{d-1} = \theta_d \theta'_{d-1}$$

The Jacobian of this transformation is $\theta_d^{d-r-1} \times \theta_d^{r(m-1)} = \theta_d^{d+r(m-2)-1}$, and $K(\theta)$ can be expressed as:

$$\begin{aligned} K(\theta) = & \frac{1}{2} \mathbb{E}_X \left[\left\{ F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d) \right\}^2 \right] \\ & + f_0(\theta_{r+1}, \dots, \theta_d) + f_1(\theta_1, \dots, \theta_d) + f_2(\theta_1, \dots, \theta_d) + f_3(\theta_1, \dots, \theta_r) \\ = & \frac{1}{2} \theta_d^{2(m-1)} \left\{ \mathbb{E}_X \left[\left\{ F_1(X|\theta'_1, \dots, \theta'_r) + \theta_d F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1) \right\}^2 \right] \right. \\ & + \theta_d^3 f_0(\theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_d^2 f_1(\theta'_1, \dots, \theta'_{d-1}, 1) \\ & \left. + \theta_d f_2(\theta'_1, \dots, \theta'_{d-1}, 1) + \theta_d^{m-1} f_3(\theta'_1, \dots, \theta'_r) \right\} \end{aligned}$$

In this coordinate system, perform one blow-up centered at $\{(\theta'_1, \dots, \theta'_{d-1}, \theta_d) \mid \theta'_1 = \dots = \theta'_r = \theta_d = 0\}$.

(c-1) First, consider the case where the exceptional surface is $\{\theta'_i = 0\} (i = 1, \dots, r)$. For example, in the case $i = 1$, that is, when

$$\{\theta'_2 = \theta'_1 \theta''_2, \dots, \theta'_r = \theta'_1 \theta''_r, \theta_d = \theta'_1 \theta'_d\}$$

we perform a blow-up, the exceptional surface is $\{\theta'_1 = 0\}$, and the Jacobian is

$$\theta_1^r \cdot \theta_d^{d+r(m-2)-1} = \theta_1^{d+r(m-1)-1} \cdot \theta_d^{d+r(m-2)-1}$$

and

$$\begin{aligned} K(\theta) &= \frac{1}{2} \theta_d^{2(m-1)} \left\{ \mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + \theta_d F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1)\}^2 \right] \right. \\ &\quad + \theta_d^3 f_0(\theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_d^2 f_1(\theta'_1, \dots, \theta'_{d-1}, 1) \\ &\quad \left. + \theta_d f_2(\theta'_1, \dots, \theta'_{d-1}, 1) + \theta_d^{m-1} f_3(\theta'_1, \dots, \theta'_r) \right\} \\ &= \frac{1}{2} \theta_1^{2m} \theta_d^{2(m-1)} \left\{ \mathbb{E}_X \left[\{F_1(X|1, \theta''_2, \dots, \theta''_r) + \theta'_d F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1)\}^2 \right] \right. \\ &\quad + \theta'_1 \theta_d^3 f_0(\theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta'_1 \theta_d^2 f_1(1, \theta''_2, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 1) \\ &\quad \left. + \theta'_1 \theta'_d f_2(1, \theta''_2, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_1^m \theta_d^{m-1} f_3(1, \theta''_2, \dots, \theta''_r) \right\} \\ &= \frac{1}{2} \theta_1^{2m} \theta_d^{2(m-1)} a(\theta'_1, \theta''_2, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_d) \end{aligned}$$

(where a is an analytic function). Considering the point on $U_0 = g^{-1}(O)$ in this local coordinate system $(\theta'_1, \theta''_2, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_d)$, which satisfies $\theta'_1 = 0$,

$$a(0, \theta''_2, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_d) = \mathbb{E}_X \left[\{F_1(X|1, \theta''_2, \dots, \theta''_r) + \theta'_d F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1)\}^2 \right]$$

and by Lemma 3.3(1)(2),

$$a(0, \theta''_2, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_d) > 0$$

Thus, in this local coordinate system, the normal crossing of $K(\theta)$ is obtained at any point Q on U_0 , and

$$\min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \min \left\{ \frac{d+r(m-1)-1+1}{2m}, \frac{d+r(m-2)-1+1}{2(m-1)} \right\} = \frac{d-r+rm}{2m}$$

Therefore, considering the local coordinates obtained from the first $m-1$ blow-ups,

$$\inf_{Q \in U_0} \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \inf_{1 \leq m' \leq m} \left\{ \frac{d-r+rm'}{2m'} \right\} = \frac{d-r+rm}{2m}$$

(multiplicity is 1).

(c-2) Next, consider the case where the exceptional surface is $\{\theta_d = 0\}$. That is, when we perform a blow-up with $\{\theta'_1 = \theta_d \theta''_1, \dots, \theta'_r = \theta_d \theta''_r\}$, the Jacobian is $\theta_d^r \cdot \theta_d^{d+r(m-2)-1} = \theta_d^{d+r(m-1)-1}$, and

$$\begin{aligned}
K(\theta) &= \frac{1}{2} \theta_d^{2(m-1)} \left\{ \mathbb{E}_X \left[\left\{ F_1(X|\theta'_1, \dots, \theta'_r) + \theta_d F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1) \right\}^2 \right] \right. \\
&\quad \left. + \theta_d^3 f_0(\theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_d^2 f_1(\theta'_1, \dots, \theta'_{d-1}, 1) \right. \\
&\quad \left. + \theta_d f_2(\theta'_1, \dots, \theta'_{d-1}, 1) + \theta_d^{m-1} f_3(\theta'_1, \dots, \theta'_r) \right\} \\
&= \frac{1}{2} \theta_d^{2m} \left\{ \mathbb{E}_X \left[\left\{ F_1(X|\theta''_1, \dots, \theta''_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1) \right\}^2 \right] \right. \\
&\quad \left. + \theta_d f_0(\theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_d f_1(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 1) \right. \\
&\quad \left. + \theta_d f_2(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 1) + \theta_d^m f_3(\theta''_1, \dots, \theta''_r) \right\} \\
&= \frac{1}{2} \theta_d^{2m} a(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, \theta_d)
\end{aligned}$$

(where a is an analytic function). Considering the point on $U_0 = g^{-1}(O)$ in this local coordinate system $(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, \theta_d)$, which satisfies $\theta_d = 0$,

$$a(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 0) = \mathbb{E}_X \left[\left\{ F_1(X|\theta''_1, \dots, \theta''_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1) \right\}^2 \right]$$

and

$$S = \left\{ (\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, \theta_d) \mid \begin{array}{l} (\theta''_1, \dots, \theta''_r, \theta_d) = 0 \\ F_m(X|\theta'_{r+1}, \dots, \theta'_{d-1}, 1) = 0 \text{ (a.s.)} \end{array} \right\} \subset U_0$$

At points Q on U_0 not included in S , by Lemma 3.3(2), $a(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 0) > 0$, so the normal crossing of $K(\theta)$ is obtained, and considering the local coordinates obtained from the first $m-1$ blow-ups,

$$\inf_{Q \in U_0 \setminus S} \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} = \inf_{1 \leq m' \leq m} \left\{ \frac{d-r+rm'}{2m'} \right\} = \frac{d-r+rm}{2m}$$

(multiplicity is 1). Note that at points included in S , $a(\theta''_1, \dots, \theta''_r, \theta'_{r+1}, \dots, \theta'_{d-1}, 0) = 0$, and the normal crossing of $K(\theta)$ is not obtained.

Thus, the theorem is proved. □

Remark 3.7.

From the proof of Main Theorem 2, the following can be understood. Considering the change in the log-likelihood ratio function f before and after the blow-up,

$$\begin{aligned} f(X|\theta) &= F_1(X|\theta_1, \dots, \theta_r) + F_m(X|\theta_{r+1}, \dots, \theta_d) + (\text{higher order terms}) \\ &= \theta_i^m \{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d) + g(\theta_i)\} \end{aligned}$$

can be expressed (where $g(0) = 0$). On the other hand, $K(\theta)$ can be expressed as

$$K(\theta) = \frac{1}{2} \theta_i^{2m} \left\{ \mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d)\}^2 \right] + h(\theta_i) \right\}$$

(where $h(0) = 0$). From this, it follows that

$$\begin{aligned} & \lim_{\theta \rightarrow 0} \frac{\mathbb{E}_X [f(X|\theta)^2]}{K(\theta)} \\ &= 2 \lim_{\theta_i \rightarrow 0} \frac{\mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d) + g(\theta_i)\}^2 \right]}{\mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d)\}^2 \right] + h(\theta_i)} \\ &= 2 \lim_{\theta_i \rightarrow 0} \frac{\mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d)\}^2 \right]}{\mathbb{E}_X \left[\{F_1(X|\theta'_1, \dots, \theta'_r) + F_m(X|\theta'_{r+1}, \dots, \theta'_{i-1}, 1, \theta'_{i+1}, \dots, \theta'_d)\}^2 \right]} \\ &= 2 \end{aligned}$$

follows. This is consistent with [2, Theorem 6.3].

3.4 Proof of Corollary 3.1

Lemma 3.4.

When Assumption 1(1)(2) is satisfied for $m = \infty$, the following holds for any $(\theta_1, \dots, \theta_d) \in \mathbb{R}^d$ sufficiently close to 0.

(1)

$$f(X|\theta_1 = 0, \dots, \theta_r = 0, \forall \theta_{r+1}, \dots, \forall \theta_d) = 0 \quad \text{a.s.}$$

(2) The following r random variables are linearly independent.

$$\frac{\partial f(X|\theta)}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_r)=0}, \dots, \frac{\partial f(X|\theta)}{\partial \theta_r} \Big|_{(\theta_1, \dots, \theta_r)=0} \quad (3.17)$$

Therefore, for $m = \infty$, Assumption 1(1) (in the case $d = r$) is satisfied for the parameters $(\theta_1, \dots, \theta_r)$ (here, $(\theta_{r+1}, \dots, \theta_d)$ are treated as constants).

Proof.

- (1) Since Assumption 1(2) is satisfied for $m = \infty$, in the Taylor expansion of f at $(\theta_1, \dots, \theta_d) = 0$, there are no terms consisting only of $\theta_{r+1}, \dots, \theta_d$. In other words,

$$f(X|\theta_1 = 0, \dots, \theta_r = 0, \forall \theta_{r+1}, \dots, \forall \theta_d) = 0 \quad \text{a.s.}$$

holds.

- (2) From Lemma 3.2, the linear independence in (3.17) is equivalent to the positive definiteness of the matrix formed by the product of these two elements. Denote this determinant by $D(\theta_{r+1}, \dots, \theta_d)$. It suffices to show that $D > 0$ for any $(\theta_{r+1}, \dots, \theta_d)$ sufficiently close to 0. When $(\theta_{r+1}, \dots, \theta_d) = 0$, by Assumption 1(1),

$$\left. \frac{\partial f(X|\theta)}{\partial \theta_1} \right|_{(\theta_1, \dots, \theta_d)=0}, \dots, \left. \frac{\partial f(X|\theta)}{\partial \theta_r} \right|_{(\theta_1, \dots, \theta_d)=0}$$

are linearly independent, so $D(0, \dots, 0) > 0$ holds. Therefore, for $(\theta_{r+1}, \dots, \theta_d)$ sufficiently close to 0, $D(\theta_{r+1}, \dots, \theta_d) > 0$ follows. □

Proof of Corollary 3.1.

- (1) Based on the assumption that $m < \infty$ and for all parameters $(\theta_{r+1}, \dots, \theta_d) \neq 0$, $F_m(X|\theta_{r+1}, \dots, \theta_d)$ is linearly independent of the following r random variables:

$$\left. \frac{\partial f(X|\theta)}{\partial \theta_1} \right|_{(\theta_1, \dots, \theta_d) \neq 0}, \dots, \left. \frac{\partial f(X|\theta)}{\partial \theta_r} \right|_{(\theta_1, \dots, \theta_d) \neq 0}$$

Specifically, for all parameters $(\theta_{r+1}, \dots, \theta_d) \neq 0$,

$$F_m(X|\theta_{r+1}, \dots, \theta_d) \neq 0 \quad \text{a.s.},$$

thus, $S = \{\}$ in Main Theorem 2, and the blow-up g in Main Theorem 2 yields a normal crossing of $K(\theta)$ at every point Q on $U_0 := g^{-1}(0)$. The real log canonical threshold λ_O at $\theta = 0$ is given by the following expression (with multiplicity 1):

$$\lambda_O = \inf_{Q \in U_0} \left\{ \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} \right\} = \inf_{Q \in U_0 \setminus S} \left\{ \min_{i=1, \dots, d} \frac{h_i^{(Q)} + 1}{k_i^{(Q)}} \right\} = \frac{d - r + rm}{2m}$$

Moreover, since normal crossings of $K(\theta)$ are obtained by the blow-up centered at the origin O , near the origin O using the notation from Theorem 2.1, we have:

$$\begin{aligned} K^{-1}(0) &= \{\theta \in \Theta \mid K(\theta) = 0\} = \left\{ g(u) \mid a(u)u_1^{k_1} \cdots u_d^{k_d} = 0 \right\} = \{g(u) \mid u_1 \cdots u_d = 0\} \\ &= \{O\} \end{aligned}$$

Particularly, the blow-up g satisfies the conditions of the resolution theorem (Theorem 2.1).

- (2) From Lemma 3.4(2), if we treat only $(\theta_1, \dots, \theta_r)$ as parameters and consider $(\theta_{r+1}, \dots, \theta_d)$ as arbitrary constants sufficiently close to zero, Assumption 1(1) is satisfied in the case where $d = r$. Furthermore, it is clear that Assumption 1(2),(3)(ii) is satisfied when $m = 1$. Therefore, Main Theorem 2 and Corollary 3.1(1) can be applied in the case of $(d, r, m) = (r, r, 1)$, yielding a normal crossing of $K(\theta)$ through a blow-up centered on the subvariety W_0 , and obtaining $\lambda_O = r/2$ (with multiplicity 1). Additionally, by the same argument as in (1), it can be seen that the set of points satisfying $K(\theta) = 0$ in the neighborhood of the origin in \mathbb{R}^d is W_0 .

□

4 Conditions for Applying Main Theorem 1

To use Main Theorem 2, it is necessary to satisfy Assumption 1 concerning the log-likelihood ratio function f . In the case of a semi-regular model, it is confirmed that this assumption can always be satisfied by making appropriate variable transformations.

Theorem 4.1. For any statistical model that is semi-regular at $(\theta_1, \dots, \theta_d) = 0$, there exists a variable transformation such that the transformed statistical model satisfies Assumption 1.

To satisfy Assumption 1, the variable transformation can be specifically constructed in the following order: (i)-(v). Here, we first describe the concrete method of construction. In this method of construction, general theories about variable transformations such as Lemma 4.1 and Corollary 4.1 will be used, but their proofs will be deferred. References to such propositions are made by citing them as [Corollary 4.1] at the relevant points in the discussion. Unless otherwise confusing, the same notation θ will be used before and after the variable transformation.

Proof.

- (i) Let V_1 be the vector space over \mathbb{R} generated by the first derivatives of f with respect to $\theta_1, \dots, \theta_d$. From [Lemma 4.1], the dimension of V_1 is r . Therefore, by suitably permuting coordinates, we can take as a basis:

$$\left\{ \frac{\partial f}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_d)=0}, \dots, \frac{\partial f}{\partial \theta_r} \Big|_{(\theta_1, \dots, \theta_d)=0} \right\} \quad (4.1)$$

This satisfies Assumption 1(1). In the following, consider the set of n -th derivatives of f with respect to variables $\theta_{r+1}, \dots, \theta_d$ ($n \geq 1$):

$$D_n := \left\{ \frac{\partial^n f}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0} \mid i_{r+1} + \cdots + i_d = n \right\}$$

We denote the vector space over \mathbb{R} generated by D_n as W_n .

- (ii) For the first derivatives of f with respect to $\theta_{r+1}, \dots, \theta_d$ (e.g., the derivative with respect to θ_{r+1}), consider the linear relationship

$$\frac{\partial f}{\partial \theta_{r+1}} \Big|_{(\theta_1, \dots, \theta_d)=0} = \sum_{k=1}^r a_k \cdot \frac{\partial f}{\partial \theta_k} \Big|_{(\theta_1, \dots, \theta_d)=0}, \quad a_k \in \mathbb{R}$$

which defines a coordinate transformation

$$\theta'_1 := \theta_1 + a_1 \cdot \theta_{r+1}, \dots, \theta'_r := \theta_r + a_r \cdot \theta_{r+1} \quad (4.2)$$

Performing this transformation affects no other first derivatives, and in the new coordinates,

$$\left. \frac{\partial f}{\partial \theta_{r+1}} \right|_{(\theta_1, \dots, \theta_d)=0} = 0 \quad \text{a.s.}$$

as stated in [Corollary 4.1]. By repeating this process, the transformed space W_1 becomes $\{0\}$.

- (iii) Starting with $n = 2$, if $\dim(V_1 + W_n) = r$, then $V_1 + W_n = V_1$, so any element of D_n can be represented as a linear combination from (4.1). Specifically,

$$\left. \frac{\partial^n f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} = \sum_{k=1}^r a_k \cdot \left. \frac{\partial f}{\partial \theta_k} \right|_{(\theta_1, \dots, \theta_d)=0}, \quad a_k \in \mathbb{R}$$

Given this representation, the coordinate transformation

$$\begin{cases} \theta'_1 := \theta_1 + \frac{a_1}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \\ \vdots \\ \theta'_r := \theta_r + \frac{a_r}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \end{cases} \quad (4.3)$$

can be performed without changing other derivatives of order n or lower, ensuring that

$$\left. \frac{\partial^n f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} = 0 \quad (\text{a.s.})$$

[Corollary 4.1]. This coordinate transformation can be applied to any element of D_n , resulting in $W_n = \{0\}$. Next, by incrementing n by 1 and repeating as long as $\dim(V_1 + W_n) = r$, W_n can be kept at $\{0\}$ for this n .

- (iv) If $W_n = \{0\}$ for all n , this satisfies Assumption 1(1)(2) in the case $m = \infty$.
- (v) Consider the case where for some n , $\dim(V_1 + W_n) > r$. Let us denote this particular n as $m (< \infty)$. Since $W_1 = \cdots = W_{m-1} = \{0\}$, Assumption 1(2) is satisfied. Therefore, we only need to perform a coordinate transformation that satisfies Assumption 1(3).

Let $r + s := \dim(V_1 + W_m)$ ($s \geq 1$) and use elements B_1, \dots, B_s of D_m to form a basis of the vector space $V_1 + W_m$:

$$\left\{ \left. \frac{\partial f}{\partial \theta_1} \right|_{(\theta_1, \dots, \theta_d)=0}, \dots, \left. \frac{\partial f}{\partial \theta_r} \right|_{(\theta_1, \dots, \theta_d)=0}, B_1, \dots, B_s \right\}.$$

Then, any element $B \in D_m \setminus \{B_1, \dots, B_s\}$, also being an element of $V_1 + W_m$, can be expressed as

$$B := \left. \frac{\partial^m f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} = \sum_{k=1}^r a_k \cdot \left. \frac{\partial f}{\partial \theta_k} \right|_{(\theta_1, \dots, \theta_d)=0} + \sum_{j=1}^s b_j \cdot B_j$$

Performing the coordinate transformation

$$\begin{cases} \theta'_1 := \theta_1 + \frac{a_1}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \\ \vdots \\ \theta'_r := \theta_r + \frac{a_r}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \end{cases} \quad (4.4)$$

allows for

$$B = \sum_{j=1}^s b_j B_j$$

without changing other derivatives up to order m [Corollary 4.1]. By repeating this for all elements of $D_m \setminus \{B_1, \dots, B_s\}$, any element of D_m can be expressed as a linear combination of $\{B_1, \dots, B_s\}$. Therefore, W_m is generated over \mathbb{R} by $\{B_1, \dots, B_s\}$. Since these generators are linearly independent from the generators of V_1 (4.1), it follows that $V_1 \cap W_m = \{0\}$.

This configuration satisfies Assumption 1(3). Regarding the m -th term F_m of the Taylor expansion of f around $\theta = 0$, it suffices to show that for parameters $(\theta_{r+1}, \dots, \theta_d)$ for which $F_m \neq 0$ (a.s.), F_m is linearly independent from (4.1), which follows since $F_m \in W_m$ and thus $V_1 \cap W_m = \{0\}$.

□

Remark 4.1. Ultimately, to satisfy Assumption 1, it is necessary to perform the variable transformations (4.2), (4.3), and (4.4) defined by the linear dependency of the random variables. Furthermore, when $m < \infty$, performing these coordinate transformations corresponds to establishing the conditions:

$$V_1 = V_1 + W_1 = V_1 + W_2 = \cdots = V_1 + W_{m-1} \subsetneq V_1 + W_m,$$

where $W_1 = \cdots = W_{m-1} = \{0\}$, $V_1 \cap W_m = \{0\}$.

To complete the proof of Theorem 4.1, it is necessary to prove Lemma 4.1 and Corollary 4.1 used in the proof. These proofs will be provided in the following sections, but here we will first describe specific examples of constructing the variable transformations.

Example 4.1 (Case of $m = 2$). In the statistical model (3.1) of Example 3.1, the variable transformation

$$\theta'_1 = \theta_1 + \theta_2 \quad (4.5)$$

was performed from the beginning. This is actually the variable transformation (4.2). In fact, it is more natural to set the statistical model as

$$p(X = x | \theta_1, \theta_2) := \frac{1}{2} \cdot \tilde{p} \left(X = x \middle| \theta_1 + \frac{1}{2} \right) + \frac{1}{2} \cdot \tilde{p} \left(X = x \middle| \theta_2 + \frac{1}{2} \right) \quad (x = 0, 1, 2),$$

but

$$\frac{\partial f}{\partial \theta_1} \Big|_{(\theta_1, \theta_2)=0} = \begin{cases} 2, & x = 0 \\ 0, & x = 1 \\ -2, & x = 2 \end{cases}, \quad \frac{\partial f}{\partial \theta_2} \Big|_{(\theta_1, \theta_2)=0} = \begin{cases} 2, & x = 0 \\ 0, & x = 1 \\ -2, & x = 2 \end{cases}$$

and there is a linear dependency between them:

$$\frac{\partial f}{\partial \theta_2} \Big|_{(\theta_1, \theta_2)=0} = \frac{\partial f}{\partial \theta_1} \Big|_{(\theta_1, \theta_2)=0}.$$

The variable transformation (4.2) determined from this is exactly (4.5). As calculated in Example 3.1, after the variable transformation,

$$\frac{\partial f}{\partial \theta_2} \Big|_{(\theta'_1, \theta_2)=0} = 0.$$

It has already been confirmed in Example 3.1 that the statistical model after this variable transformation satisfies Assumption 1 in the case of $m = 2$.

Example 4.2 (Case of $m = \infty$). Consider the statistical model with parameters (a, b)

$$p(x|a, b) = \frac{1}{1 + e^{-(a+1)(b+1)x}}, \quad q(x) = \frac{1}{1 + e^{-x}} \quad (x = 1, -1),$$

which realizes the true distribution at the origin $(a, b) = (0, 0)$. We consider variable transformations that satisfy Assumption 1 at the origin.

First, regarding the first derivatives of the log-likelihood ratio function,

$$\frac{\partial f(x|a, b)}{\partial a} \Big|_{(a, b)=0} = -\frac{x}{1 + e^x}, \quad \frac{\partial f(x|a, b)}{\partial b} \Big|_{(a, b)=0} = -\frac{x}{1 + e^x},$$

since these are non-zero, it is a semi-regular model. These satisfy the linear dependency

$$\frac{\partial f(x|a, b)}{\partial b} \Big|_{(a, b)=0} = \frac{\partial f(x|a, b)}{\partial a} \Big|_{(a, b)=0}.$$

Performing the corresponding coordinate transformation (4.2): $a' = a + b$, we indeed get

$$\frac{\partial f(x|a, b)}{\partial a'} \Big|_{(a', b)=0} = -\frac{x}{1 + e^x}, \quad \frac{\partial f(x|a, b)}{\partial b} \Big|_{(a', b)=0} = 0.$$

Next, for the second derivatives,

$$\frac{\partial^2 f(x|a, b)}{\partial b^2} \Big|_{(a', b)=0} = 2 \cdot \frac{x}{1 + e^x},$$

which again satisfies the linear dependency

$$\left. \frac{\partial^2 f(x|a, b)}{\partial b^2} \right|_{(a', b)=0} = -2 \cdot \left. \frac{\partial f(x|a, b)}{\partial a'} \right|_{(a', b)=0}.$$

Performing the corresponding coordinate transformation (4.3): $a'' = a' - b^2$, we indeed get

$$\begin{aligned} \left. \frac{\partial f(x|a, b)}{\partial a''} \right|_{(a'', b)=0} &= -\frac{x}{1 + e^x}, \\ \left. \frac{\partial f(x|a, b)}{\partial b} \right|_{(a'', b)=0} &= \left. \frac{\partial^2 f(x|a, b)}{\partial b^2} \right|_{(a'', b)=0} = 0. \end{aligned}$$

Next, for the third derivatives,

$$\left. \frac{\partial^3 f(x|a, b)}{\partial b^3} \right|_{(a'', b)=0} = -6 \cdot \frac{x}{1 + e^x},$$

which again satisfies the linear dependency

$$\left. \frac{\partial^3 f(x|a, b)}{\partial b^3} \right|_{(a'', b)=0} = 6 \cdot \left. \frac{\partial f(x|a, b)}{\partial a''} \right|_{(a'', b)=0}.$$

Performing the corresponding coordinate transformation (4.3): $a''' = a'' + b^3$, we indeed get

$$\begin{aligned} \left. \frac{\partial f(x|a, b)}{\partial a'''} \right|_{(a''', b)=0} &= -\frac{x}{1 + e^x}, \\ \left. \frac{\partial f(x|a, b)}{\partial b} \right|_{(a''', b)=0} &= \left. \frac{\partial^2 f(x|a, b)}{\partial b^2} \right|_{(a''', b)=0} = \left. \frac{\partial^3 f(x|a, b)}{\partial b^3} \right|_{(a''', b)=0} = 0. \end{aligned}$$

By iterating this infinitely many times, the coordinate transformation $(a, b) \mapsto (\tilde{a}, b)$

$$\tilde{a} := a + b - b^2 + b^3 + \dots = a + \frac{b}{b + 1} \tag{4.6}$$

satisfies, for any $m \geq 1$,

$$\left. \frac{\partial^m f(x|a, b)}{\partial b^m} \right|_{(\tilde{a}, b)=0} = 0.$$

Thus, by performing the variable transformation (4.6), it is confirmed that Assumption 1 is satisfied in the case of $m = \infty$.

4.1 On Assumption 1(1)

Lemma 4.1. If the rank of the Fisher information matrix I is $r (> 0)$, then the dimension of the \mathbb{R} -vector space V_1 generated by the first derivatives of f with respect to $\theta_1, \dots, \theta_d$ is r .

Proof. Let

$$A(X) := (A_1(X), \dots, A_d(X))^\top \in \mathbb{R}^d,$$

$$A_i(X) := \left. \frac{\partial f(X|\theta)}{\partial \theta_i} \right|_{\theta=\theta_*} \quad (i = 1, \dots, d).$$

As noted in Remark 3.5, since $\mathbb{E}_X [A(X)A(X)^\top] \in \mathbb{R}^{d \times d}$ coincides with the Fisher information matrix I , and the rank of this matrix is r . Therefore, it follows from Lemma 3.2(2) that $\dim V_1 = r$. □

4.2 On Assumption 1(2) and (3)

Proposition 4.1. [Properties of Coordinate Transformation] Let r be an integer greater than or equal to 1. Fix some non-negative integers i_{r+1}, \dots, i_d such that $(i_{r+1} + \dots + i_d \geq 1)$. For $k = 1, \dots, r$, define

$$a_k(\theta_{r+1}, \dots, \theta_d) := \frac{c_k}{i_{r+1}! \dots i_d!} \cdot \theta_{r+1}^{i_{r+1}} \dots \theta_d^{i_d}, \quad c_k \in \mathbb{R},$$

and consider the coordinate transformation $\varphi : (\theta_1, \dots, \theta_d) \mapsto (\theta'_1, \dots, \theta'_d)$ given by

$$\begin{aligned} \theta'_1 &= \theta_1 + a_1(\theta_{r+1}, \dots, \theta_d), \dots, \theta'_r = \theta_r + a_r(\theta_{r+1}, \dots, \theta_d), \\ \theta'_{r+1} &= \theta_{r+1}, \dots, \theta'_d = \theta_d. \end{aligned}$$

This transformation satisfies the following:

- (1) $\varphi(0) = 0$ and $|\det \varphi'(0)| = 1$, and it is bijective and analytic.
- (2) For $j = 1, \dots, r$, $\frac{\partial}{\partial \theta'_j} = \frac{\partial}{\partial \theta_j}$ holds.
- (3) For any $(h_{r+1}, \dots, h_d) \in \mathbb{Z}_{\geq 0}^{d-r}$ such that $h_{r+1} + \dots + h_d \leq i_{r+1} + \dots + i_d$,

$$\begin{aligned} & \left. \frac{\partial^{h_{r+1} + \dots + h_d}}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} \right|_{(\theta'_1, \dots, \theta'_d)=0} \\ &= \left. \frac{\partial^{h_{r+1} + \dots + h_d}}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} \right|_{\theta=0} - \begin{cases} 0 & \text{if } (h_{r+1}, \dots, h_d) \neq (i_{r+1}, \dots, i_d), \\ \sum_{k=1}^r c_k \cdot \left. \frac{\partial}{\partial \theta_k} \right|_{\theta=0} & \text{if } (h_{r+1}, \dots, h_d) = (i_{r+1}, \dots, i_d), \end{cases} \end{aligned}$$

holds (where $\theta := (\theta_1, \dots, \theta_d)$).

Proof. (1) is evident. (2) follows from the relationship between partial derivatives before and

after the variable transformation:

$$\begin{aligned}\frac{\partial}{\partial \theta'_j} &= \sum_{k=1}^d \frac{\partial}{\partial \theta_k} \cdot \frac{\partial \theta_k}{\partial \theta'_j} = \sum_{k=1}^r \frac{\partial}{\partial \theta_k} \cdot \frac{\partial \theta_k}{\partial \theta'_j} + \sum_{k=r+1}^d \frac{\partial}{\partial \theta_k} \cdot \delta_{k,j} \\ &= \frac{\partial}{\partial \theta_j} - \sum_{k=1}^r \frac{\partial a_k}{\partial \theta_j} \cdot \frac{\partial}{\partial \theta_k} \quad (j = r+1, \dots, d) \\ \frac{\partial}{\partial \theta'_j} &= \frac{\partial}{\partial \theta_j} \quad (j = 1, \dots, r)\end{aligned}$$

(3) can be expressed as follows: Let $m := i_{r+1} + \dots + i_d (\geq 1)$, $l := h_{r+1} + \dots + h_d (\leq m)$. The derivatives are expressed as follows:

$$\begin{aligned}\frac{\partial^l}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} &= \prod_{j=r+1}^d \left(\frac{\partial}{\partial \theta_j} - \sum_{k=1}^r \frac{\partial a_k}{\partial \theta_j} \cdot \frac{\partial}{\partial \theta_k} \right)^{h_j} \\ &= \prod_{j=r+1}^d \left(\frac{\partial^{h_j}}{\partial \theta_j^{h_j}} - \sum_{k=1}^r \frac{\partial^{h_j} a_k}{\partial \theta_j^{h_j}} \cdot \frac{\partial}{\partial \theta_k} + \dots \right) \quad (4.7)\end{aligned}$$

$$= \frac{\partial^l}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} - \sum_{k=1}^r \frac{\partial^l a_k}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} \cdot \frac{\partial}{\partial \theta_k} + \dots \quad (4.8)$$

and the “...” in (4.7) and (4.8) represent terms derived from derivatives of $a_k(\theta_{r+1}, \dots, \theta_d)$ up to order $l-1$ with respect to $\theta_{r+1}, \dots, \theta_d$. Since $a_k(\theta_{r+1}, \dots, \theta_d)$ is a homogeneous polynomial of degree m with respect to $\theta_{r+1}, \dots, \theta_d$, at $\theta = (\theta_1, \dots, \theta_d) = 0$ the “...” in (4.7) and (4.8) becomes zero, and

$$\left. \frac{\partial^l a_k}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} \right|_{\theta=0} = \begin{cases} 0 & (h_{r+1}, \dots, h_d) \neq (i_{r+1}, \dots, i_d) \\ c_k & (h_{r+1}, \dots, h_d) = (i_{r+1}, \dots, i_d) \end{cases}$$

Therefore, the following holds:

$$\begin{aligned}& \left. \frac{\partial^l}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} \right|_{(\theta'_1, \dots, \theta'_d)=0} \\ &= \left. \frac{\partial^l}{\partial \theta_{r+1}^{h_{r+1}} \dots \partial \theta_d^{h_d}} \right|_{\theta=0} - \begin{cases} 0 & (h_{r+1}, \dots, h_d) \neq (i_{r+1}, \dots, i_d) \\ \sum_{k=1}^r c_k \cdot \frac{\partial}{\partial \theta_k} \Big|_{\theta=0} & (h_{r+1}, \dots, h_d) = (i_{r+1}, \dots, i_d) \end{cases}\end{aligned}$$

□

Corollary 4.1. [Coordinate transformation to satisfy Assumption 1(2),(3)] Let r be an integer satisfying $1 \leq r \leq d-1$, and let n be an integer greater than or equal to 1. Define

$$D_n := \left\{ \left. \frac{\partial^n f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} \mid i_{r+1} + \dots + i_d = n \right\} \quad (4.9)$$

Let B be an element of D_n defined as

$$B := \frac{\partial^n f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

Assume that B can be expressed using $B_1, \dots, B_s \in D_n$ and real numbers c_k, b_j ($k = 1, \dots, r, j = 1, \dots, s$) as follows:

$$B = \sum_{k=1}^r c_k \cdot \frac{\partial f(X|\theta)}{\partial \theta_k} \Big|_{(\theta_1, \dots, \theta_d)=0} + \sum_{j=1}^s b_j \cdot B_j \quad \text{a.s.}$$

Then, applying the coordinate transformation $\varphi : (\theta_1, \dots, \theta_d) \mapsto (\theta'_1, \dots, \theta'_d)$ given by Proposition 4.1;

$$\theta'_k := \theta_k + \frac{c_k}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \quad (k = 1, \dots, r), \quad c_k \in \mathbb{R}$$

without affecting the other lower-order derivatives, the transformed B (denoted as B') becomes

$$B' = \sum_{j=1}^s b_j B_j \quad \text{a.s.}$$

Proof. We need to demonstrate the following two points. (Below, the inverse transformation of φ is denoted as $\phi : (\theta'_1, \dots, \theta'_d) \mapsto (\theta_1, \dots, \theta_d)$.)

(1) For $k = 1, \dots, r$,

$$\frac{\partial f(X|\phi(\theta'))}{\partial \theta'_k} \Big|_{(\theta'_1, \dots, \theta'_d)=0} = \frac{\partial f(X|\theta)}{\partial \theta_k} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

(2) For any $(h_{r+1}, \dots, h_d) \in \mathbb{Z}_{\geq 0}^{d-r}$ such that $h_{r+1} + \dots + h_d \leq i_{r+1} + \dots + i_d$,

$$\begin{aligned} & \frac{\partial^{h_{r+1} + \dots + h_d} f(X|\phi(\theta'))}{\partial \theta_{r+1}^{h_{r+1}} \cdots \partial \theta_d^{h_d}} \Big|_{(\theta'_1, \dots, \theta'_d)=0} \\ &= \begin{cases} \frac{\partial^{h_{r+1} + \dots + h_d} f(X|\theta)}{\partial \theta_{r+1}^{h_{r+1}} \cdots \partial \theta_d^{h_d}} \Big|_{(\theta_1, \dots, \theta_d)=0} & \text{if } (h_{r+1}, \dots, h_d) \neq (i_{r+1}, \dots, i_d) \\ \sum_{j=1}^s b_j B_j \quad \text{a.s.} & \text{if } (h_{r+1}, \dots, h_d) = (i_{r+1}, \dots, i_d) \end{cases} \end{aligned}$$

(1) can be directly applied from Proposition 4.1(2).

For (2), when $(h_{r+1}, \dots, h_d) = (i_{r+1}, \dots, i_d)$, it follows from Proposition 4.1(3) that

$$\begin{aligned} B' &= \left. \frac{\partial^n f(X|\phi(\theta'))}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta'_1, \dots, \theta'_d)=0} \\ &= \left. \frac{\partial^n f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} - \sum_{k=1}^r c_k \cdot \left. \frac{\partial f(X|\theta)}{\partial \theta_k} \right|_{(\theta_1, \dots, \theta_d)=0} \\ &= \sum_{j=1}^s b_j \cdot B_j \quad \text{a.s.} \end{aligned}$$

Additionally, when $(h_{r+1}, \dots, h_d) \neq (i_{r+1}, \dots, i_d)$ and $h_{r+1} + \dots + h_d \leq n$, it follows from Proposition 4.1(3) that

$$\left. \frac{\partial^{h_{r+1} + \dots + h_d} f(X|\phi(\theta'))}{\partial \theta_{r+1}^{h_{r+1}} \cdots \partial \theta_d^{h_d}} \right|_{(\theta'_1, \dots, \theta'_d)=0} = \left. \frac{\partial^{h_{r+1} + \dots + h_d} f(X|\theta)}{\partial \theta_{r+1}^{h_{r+1}} \cdots \partial \theta_d^{h_d}} \right|_{(\theta_1, \dots, \theta_d)=0}$$

holds true. □

Remark 4.2. Let us consider the meaning of Corollary 4.1. For simplicity, assume $s = 0$ and denote the random variables $A_k(X|\theta)$ as A_k . According to the assumption of Corollary 4.1, using the real numbers c_k ($k = 1, \dots, r$), we can express

$$\left. \frac{\partial^{i_{r+1} + \dots + i_d} f(X|\theta)}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta_1, \dots, \theta_d)=0} = \sum_{k=1}^r c_k A_k \quad \text{a.s.}$$

Then, according to Corollary 4.1(2), after the variable transformation, we have

$$\left. \frac{\partial^{i_{r+1} + \dots + i_d} f(X|\phi(\theta'))}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\theta'_1, \dots, \theta'_d)=0} = 0 \quad \text{a.s.}$$

Let's interpret this from a different perspective.

When we perform a Taylor expansion of the log-likelihood ratio function f with respect to the parameter θ (noting that the coefficients are random variables), and group the terms by linearly independent random variables, we get:

$$\begin{aligned} f &= A_1 \theta_1 + \dots + A_r \theta_r + \frac{\sum_{k=1}^r c_k A_k}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} + (\text{higher order terms}) \quad \text{a.s.} \\ &= \sum_{k=1}^r A_k \left(\theta_k + \frac{c_k}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \right) + (\text{higher order terms}) \quad \text{a.s.} \end{aligned}$$

Here, if we apply the variable transformation given in Corollary 4.1

$$\theta'_k := \theta_k + \frac{c_k}{i_{r+1}! \cdots i_d!} \theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d} \quad (k = 1, \dots, r)$$

then we can rewrite it as

$$f = A_1\theta'_1 + \cdots + A_r\theta'_r + (\text{higher order terms}) \text{ a.s.}$$

This shows that the coefficient (random variable) of the term involving $\theta_{r+1}^{i_{r+1}} \cdots \theta_d^{i_d}$ can indeed be made zero (a.s.).

4.3 The Case of $m = \infty$

Finally, let us consider the case where $m = \infty$. In this scenario, it is necessary to iteratively apply the coordinate transformation from Corollary 4.1 an infinite number of times. Conceptually, it is understood that a coordinate transformation for $m = \infty$ exists, but it must be constructed inductively, and it is not possible to explicitly represent the overall picture of the transformation. Moreover, while this transformation is clearly a bijection near the origin and maps the origin to the origin, the transformation is a formal power series, and it is not clear whether it is analytic.

For example, in Example 4.2, the variable transformation (4.6) was provided because it could be inductively understood as being in the form of an infinite geometric series. Generally, it is difficult to infer the overall picture of transformations in this manner, and obtaining this overall view may require a different approach.

Using the following Proposition 4.2, by solving the **local equation** for the set of realization parameters Θ_* near the origin with respect to $(\theta_1, \dots, \theta_r)$, the variable transformation can be explicitly constructed. Moreover, if Θ_* is an analytic set, then this solution will be an analytic function. Furthermore, as will be discussed later, this is unique, indicating that it is identical to the coordinate transformation obtained by Corollary 4.1.

Proposition 4.2. [Coordinate transformation for the case $m = \infty$] Assume that the statistical model $p(x|\theta)$ satisfies Assumption 1(1). If there exist r analytic functions $\varphi_i(\theta_{r+1}, \dots, \theta_d)$ ($i = 1, \dots, r$) such that the set of realizable parameters Θ_* can be expressed near the origin as

$$\Theta_* = \{(\theta_1, \dots, \theta_r, \theta_{r+1}, \dots, \theta_d) \mid \theta_i = -\varphi_i(\theta_{r+1}, \dots, \theta_d) \quad i = 1, \dots, r\} \quad (4.10)$$

then, by the variable transformation

$$\tilde{\theta}_i := \theta_i + \varphi_i(\theta_{r+1}, \dots, \theta_d) \quad (i = 1, \dots, r) \quad (4.11)$$

for any non-negative integers i_{r+1}, \dots, i_d (with $i_{r+1} + \cdots + i_d \geq 1$), we have

$$\left. \frac{\partial^{i_{r+1} + \cdots + i_d} f}{\partial \theta_{r+1}^{i_{r+1}} \cdots \partial \theta_d^{i_d}} \right|_{(\tilde{\theta}_1, \dots, \tilde{\theta}_r, \theta_{r+1}, \dots, \theta_d) = 0} = 0 \text{ a.s.}$$

In other words, for the case $m = \infty$, Assumption 1(1)(2) is satisfied. Specifically, in the transformed coordinates, we can express

$$\Theta_* = \{(\tilde{\theta}_1, \dots, \tilde{\theta}_r, \theta_{r+1}, \dots, \theta_d) \mid \tilde{\theta}_1 = \cdots = \tilde{\theta}_r = 0\}.$$

Proof. By performing the variable transformation (4.11), the set of realizable parameters Θ_* can be expressed near the origin as:

$$\begin{aligned}\Theta_* &= \{(\theta_1, \dots, \theta_r, \theta_{r+1}, \dots, \theta_d) \mid \theta_i = -\varphi_i(\theta_{r+1}, \dots, \theta_d) \quad i = 1, \dots, r\} \\ &= \{(\tilde{\theta}_1, \dots, \tilde{\theta}_r, \theta_{r+1}, \dots, \theta_d) \mid \tilde{\theta}_1 = \dots = \tilde{\theta}_r = 0\}\end{aligned}$$

Noting that near the origin, regardless of the values of $(\theta_{r+1}, \dots, \theta_d)$, setting $(\tilde{\theta}_1, \dots, \tilde{\theta}_r) = 0$, we have

$$p(x \mid \tilde{\theta}_1 = 0, \dots, \tilde{\theta}_r = 0, \forall \theta_{r+1}, \dots, \forall \theta_d) = q(x) \quad \text{a.s.}$$

This implies that, considering the Taylor expansion of the log-likelihood ratio function at the origin, there are no terms that depend solely on $(\theta_{r+1}, \dots, \theta_d)$. Therefore, for any non-negative integers i_{r+1}, \dots, i_d (with $i_{r+1} + \dots + i_d \geq 1$), we have

$$\frac{\partial^{i_{r+1} + \dots + i_d} f}{\partial \theta_{r+1}^{i_{r+1}} \dots \partial \theta_d^{i_d}} \Big|_{(\tilde{\theta}_1, \dots, \tilde{\theta}_r, \theta_{r+1}, \dots, \theta_d) = 0} = 0 \quad \text{a.s.}$$

□

Using Proposition 4.2, if in the neighborhood of the origin, the local equations for the set of realization parameters Θ_* can be solved with respect to the linearly independent parameters $(\theta_1, \dots, \theta_r)$, and if one can find analytic functions $\varphi_1, \dots, \varphi_r$ that satisfy (4.10), then a variable transformation satisfying Assumption 1 for the case $m = \infty$ can be obtained.

Such analytic functions can be guaranteed to exist by using the implicit function theorem.

Corollary 4.2. Assume Assumption 1(1) is satisfied. If the rank of the Jacobian of the r local equations of the set of realization parameters Θ_* at the origin is r , then functions $\varphi_1, \dots, \varphi_r$ uniquely exist and satisfy (4.10). In particular, when the origin is a non-singular point of the $d - r$ dimensional analytic manifold Θ_* , analytic functions φ_i that satisfy (4.10) exist, and Assumption 1(2) is satisfied for $m = \infty$.

Proof. This is a direct consequence of the implicit function theorem. Note that the implicit function theorem holds both in the class of analytic functions and the class of formal power series.

□

Example 4.3 (Case of $m = \infty$). Consider the statistical model with parameters (a, b) treated in Example 4.2:

$$p(x \mid a, b) = \frac{1}{1 + e^{-(a+1)(b+1)x}}, \quad \Theta_* = \{(a, b) \mid (a+1)(b+1) = 1\} \subset \mathbb{R}^2$$

The local equation of Θ_* can be solved for a near the origin:

$$a = -\frac{b}{b+1}$$

Thus, by Proposition 4.2, the coordinate transformation $(a, b) \mapsto (\tilde{a}, b)$ that satisfies $m = \infty$ is given by:

$$\tilde{a} := a + \frac{b}{b+1} \tag{4.12}$$

This matches the coordinate transformation (4.6) obtained in Example 4.2.

Let us visualize this coordinate transformation. As shown in Figure 2, the set of realizable parameters Θ_* is locally transformed to be represented by $\{\tilde{a} = 0\}$ before and after the coordinate transformation. This means that near the origin $(\tilde{a}, b) = (0, 0)$,

$$\Theta_* = \{(\tilde{a}, b) \mid \tilde{a} = 0, \forall b\}$$

and it can be seen that Θ_* is defined by a single parameter \tilde{a} near the origin.




Fig 2: Changes in the set of realizable parameters Θ_* before and after the coordinate transformation

Remark 4.3. Changing the statistical model $p(x|a, b)$ in Example 4.2 to:

$$p(x|a, b) = \frac{1}{1 + e^{-abx}}, \quad q(x) = \frac{1}{2} \quad (x = 1, -1)$$

In this case, the true distribution is realized at $(a, b) = (0, 0)$, but

$$\left. \frac{\partial f(x|a, b)}{\partial a} \right|_{(a,b)=0} = \left. \frac{\partial f(x|a, b)}{\partial b} \right|_{(a,b)=0} = 0$$

Therefore, this is not a semi-regular model. In this case, the set of realizable parameters Θ_* is given by:

$$\Theta_* = \{(a, b) \mid ab = 0\} \subset \mathbb{R}^2$$

which forms a one-dimensional variety with a singularity at the origin.

5 Example on the Derivation of the Real Log Canonical Threshold

Using the Main Theorem obtained so far for semi-regular models, we calculate the learning coefficient for a specific model.

5.1 The Case of Two Parameters

Here, we fix an arbitrary realizable parameter θ_* at a single point and translate it to the origin O , and consider the real log canonical threshold at the origin. We assume that the statistical model $p(x|\theta)$ at the origin satisfies the semi-regular condition (i.e., the rank $r > 0$ of the Fisher information matrix), and denote the real log canonical threshold at the origin by λ_O . Note that the learning coefficient is the minimum value of the real log canonical thresholds calculated for each realizable parameter.

By performing the variable transformation in Theorem 4.1, a semi-regular model with two parameters can be classified into one of the following three types.

No.	Property of f at $\theta = 0$	r	λ_O (Multiplicity)	Ideal	Geometry of Θ_* near $\theta = 0$
1	$\frac{\partial f}{\partial \theta_1}, \frac{\partial f}{\partial \theta_2}$: lin. ind.	2	1 (1)	(θ_1, θ_2)	pt $\{(\theta_1, \theta_2) = (0, 0)\}$
2	$\frac{\partial f}{\partial \theta_1}, \frac{\partial^m f}{\partial \theta_2^m}$: lin. ind. $\frac{\partial f}{\partial \theta_2} = \dots = \frac{\partial^{m-1} f}{\partial \theta_2^{m-1}} = 0$	1	$\frac{m+1}{2m}$ (1)	(θ_1, θ_2^m)	pt $\{(\theta_1, \theta_2) = (0, 0)\}$
3	$\frac{\partial f}{\partial \theta_1}$: lin. ind. $\forall m, \frac{\partial^m f}{\partial \theta_2^m} = 0$	1	$\frac{1}{2}$ (1)	(θ_1)	line $\{\theta_1 = 0\}$

The above results for the real log canonical threshold λ_O are all consequences of Corollary 3.1 (No.3 corresponds to the case $m = \infty$). From this, we find that for a statistical model with two parameters, where the Fisher information matrix at all realizable parameters has a non-zero rank, the learning coefficient is given by:

$$\lambda = \frac{m+1}{2m}, \quad m = 1, 2, \dots, \infty$$

(multiplicity is 1). Notably, the minimum value of the learning coefficient is $1/2$ and the maximum value is 1, parameterized by $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$. Moreover, Θ_* does not contain singularities.

Remark 5.1. The real log canonical threshold at the origin for the statistical model considered in Example 3.1 (3.1) was $3/4$. Applying the general theory to this statistical model with $(d, r, m) = (2, 1, 2)$, we immediately obtain:

$$\lambda_O = \frac{m+1}{2m} = \frac{3}{4}$$

Additionally, the real log canonical threshold at the origin for the statistical model considered in Example 3.2 (3.6) was $1/2$. Applying the general theory to this statistical model with

$(d, r, m) = (2, 1, \infty)$, we immediately obtain:

$$\lambda_O = \frac{1}{2}$$

The same discussion can be applied to the cases where $r = d$ or $r = d - 1$. We summarize this in the following proposition.

Proposition 5.1 (Case of $r \geq d - 1$). For a statistical model with d parameters, where the rank r of the Fisher information matrix at all realizable parameters is $r = d - 1$ or $r = d$, the learning coefficient λ can be expressed using a positive integer m as follows (multiplicity is 1):

$$\lambda = \frac{1 + (d - 1)m}{2m}, \quad m = 1, 2, \dots, \infty$$

Here, $m = \infty$ represents $\lambda = (d - 1)/2$. Notably, the minimum value of the learning coefficient is $(d - 1)/2$ and the maximum value is $d/2$. Furthermore, Θ_* does not contain singularities.

Remark 5.2. In the case of non-semi-regular models, there are exceptions to the above. For example, at $\theta = 0$, if the random variables

$$\left. \frac{\partial f}{\partial \theta_1^2} \right|_{(\theta_1, \theta_2)=0}, \quad \left. \frac{\partial f}{\partial \theta_2^3} \right|_{(\theta_1, \theta_2)=0}$$

are linearly independent and all derivatives of f up to those orders are zero (e.g., $K(\theta) = \theta_1^4 + \theta_2^6$), then this case applies. (Although methods for resolving such singularities similar to Euclid's algorithm are known, they are beyond the scope of this paper.)

However, if the model is semi-regular at other realizable parameters, the above results can be applied at those points to obtain an upper bound on the learning coefficient.

5.2 Formula for the Case When the Random Variable X Takes a Finite Number of Values

In the case of a semi-regular model where the random variable takes a finite number of values, the real log canonical threshold can sometimes be easily calculated.

Assume that the random variable X takes a finite number of values, and for the true model $q(X)$, assume without loss of generality that $q(X = x) \neq 0$ for all $x \in \chi$.

Theorem 5.1. Let X be a discrete random variable taking $N (< \infty)$ values. For $N - 1 (\leq d)$ variables $\theta_1, \dots, \theta_{N-1}$,

$$\left. \frac{\partial f(X|\theta)}{\partial \theta_1} \right|_{(\theta_1, \dots, \theta_d)=0}, \dots, \left. \frac{\partial f(X|\theta)}{\partial \theta_{N-1}} \right|_{(\theta_1, \dots, \theta_d)=0}$$

are linearly independent as random variables, the rank of the Fisher information matrix is $r = N - 1 (> 0)$, and the real log canonical threshold λ_O at the origin $\theta = 0$ for $K(\theta)$ is given by the following formula (multiplicity is 1):

$$\lambda_O = \frac{N - 1}{2}$$

Lemma 5.1. Let X be a random variable taking distinct values $\{x_1, x_2, \dots, x_N\}$, and consider $N - 1 (\leq d)$ variables $\theta_1, \dots, \theta_{N-1}$ such that

$$\frac{\partial p(X|\theta)}{\partial \theta_1} \Big|_{\theta=\theta_*}, \dots, \frac{\partial p(X|\theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*} \quad (5.1)$$

are linearly independent as random variables.

(1) The $(N - 1) \times (N - 1)$ matrix

$$\begin{pmatrix} \frac{\partial p(X=x_1|\theta)}{\partial \theta_1} \Big|_{\theta=\theta_*} & \dots & \frac{\partial p(X=x_1|\theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*} \\ \vdots & \ddots & \vdots \\ \frac{\partial p(X=x_{N-1}|\theta)}{\partial \theta_1} \Big|_{\theta=\theta_*} & \dots & \frac{\partial p(X=x_{N-1}|\theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*} \end{pmatrix}$$

is a regular matrix.

(2) If $B(X)$ is a function of X such that

$$\sum_{n=1}^N B(X = x_n) = 0, \quad (5.2)$$

then $B(X)$ is linearly dependent on the random variables in (5.1). That is, there exist real numbers a_1, \dots, a_{N-1} such that the following equation as a function of X holds:

$$B(X) = a_1 \cdot \frac{\partial p(X|\theta)}{\partial \theta_1} \Big|_{\theta=\theta_*} + \dots + a_{N-1} \cdot \frac{\partial p(X|\theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*}.$$

Proof. Since the random variable X takes values only in $\{x_1, x_2, \dots, x_N\}$, for all parameters θ ,

$$p(X = x_1|\theta) + \dots + p(X = x_N|\theta) = 1$$

holds. Differentiating this equation with respect to θ_i ($i = 1, \dots, d$), we obtain:

$$\frac{\partial p(X = x_N|\theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} = - \sum_{n=1}^{N-1} \frac{\partial p(X = x_n|\theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} \quad (5.3)$$

Also, in general, let $B(X)$ be a function of X satisfying (5.2), and assume

$$B(X = x_n) = \sum_{i=1}^{N-1} a_i \cdot \frac{\partial p(X = x_n|\theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} \quad (n = 1, \dots, N - 1) \quad (5.4)$$

for some real numbers a_1, \dots, a_{N-1} . From (5.2), (5.3), and (5.4), we get

$$\begin{aligned} B(X = x_N) &= - \sum_{n=1}^{N-1} B(X = x_n) = - \sum_{n=1}^{N-1} \sum_{i=1}^{N-1} a_i \cdot \frac{\partial p(X = x_n | \theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} \\ &= - \sum_{i=1}^{N-1} a_i \cdot \sum_{n=1}^{N-1} \frac{\partial p(X = x_n | \theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} = \sum_{i=1}^{N-1} a_i \cdot \frac{\partial p(X = x_N | \theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} \end{aligned}$$

Thus, equation (5.4) can be extended to the equality as a function of X :

$$B(X) = \sum_{i=1}^{N-1} a_i \cdot \frac{\partial p(X | \theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} \quad (5.5)$$

(1) To show the linear independence of columns, assume

$$\sum_{i=1}^{N-1} a_i \cdot \frac{\partial p(X = x_n | \theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} = 0 \quad (n = 1, \dots, N-1)$$

Let $B(X) \equiv 0$ obviously satisfies (5.2), and since (5.4) holds, this equation can be extended to the equality for the random variable (5.5)

$$\sum_{i=1}^{N-1} a_i \cdot \frac{\partial p(X | \theta)}{\partial \theta_i} \Big|_{\theta=\theta_*} = 0$$

From the linear independence of the random variables in (5.1), we get $a_i = 0$.

(2) From (1), the real solution (a_1, \dots, a_{N-1}) of the following equation exists uniquely.

$$\begin{pmatrix} B(X = x_1) \\ \vdots \\ B(X = x_{N-1}) \end{pmatrix} = \begin{pmatrix} \frac{\partial p(X = x_1 | \theta)}{\partial \theta_1} \Big|_{\theta=\theta_*} & \cdots & \frac{\partial p(X = x_1 | \theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*} \\ \vdots & \ddots & \vdots \\ \frac{\partial p(X = x_{N-1} | \theta)}{\partial \theta_1} \Big|_{\theta=\theta_*} & \cdots & \frac{\partial p(X = x_{N-1} | \theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*} \end{pmatrix} \begin{pmatrix} a_1 \\ \vdots \\ a_{N-1} \end{pmatrix}$$

Using (5.5), this can be extended to the equality as a random variable:

$$B(X) = a_1 \cdot \frac{\partial p(X | \theta)}{\partial \theta_1} \Big|_{\theta=\theta_*} + \cdots + a_{N-1} \cdot \frac{\partial p(X | \theta)}{\partial \theta_{N-1}} \Big|_{\theta=\theta_*}$$

□

Proof of Theorem 5.1.

$$\frac{\partial f(X | \theta)}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_d)=0}, \dots, \frac{\partial f(X | \theta)}{\partial \theta_{N-1}} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

are linearly independent as random variables. By multiplying both sides by $p(X|\theta = 0) = q(X) \neq 0$, it is equivalent to

$$\frac{\partial p(X|\theta)}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_d)=0}, \dots, \frac{\partial p(X|\theta)}{\partial \theta_{N-1}} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

being linearly independent as random variables.

From the assumption, it is clear that $r \geq N - 1$. For $i = N, \dots, d$,

$$B(X) := \frac{\partial p(X|\theta)}{\partial \theta_i} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

and from (5.3),

$$\sum_{n=1}^N B(X = x_n) = 0$$

is satisfied. Thus, Lemma 5.1(2) can be applied, and

$$\frac{\partial f(X|\theta)}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_d)=0}, \dots, \frac{\partial f(X|\theta)}{\partial \theta_{N-1}} \Big|_{(\theta_1, \dots, \theta_d)=0}, \frac{\partial f(X|\theta)}{\partial \theta_i} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

are linearly dependent. Therefore, we get $r = N - 1$. Then, performing the coordinate transformation guaranteed by Theorem 4.1, after the transformation, for $i = N, \dots, d$,

$$\frac{\partial f(X|\theta)}{\partial \theta_i} \Big|_{(\theta_1, \dots, \theta_d)=0} = 0$$

can be assumed.

We show that this statistical model corresponds to the case $m = \infty$ in Assumption 1. That is, taking $m(\geq 2)$ such that all derivatives of the log-likelihood ratio function with respect to $\{\theta_N, \dots, \theta_d\}$ up to $m - 1$ times are zero (a.s.), and performing the coordinate transformation guaranteed by Theorem 4.1, for any non-negative integers (i_N, \dots, i_d) such that $i_N + \dots + i_d = m$, we need to show that

$$\frac{\partial^m f(X|\theta)}{\partial \theta_N^{i_N} \dots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0} = 0 \tag{5.6}$$

is satisfied. This can be shown by applying Proposition 3.2(1), where

$$\frac{\partial^m f(X|\theta)}{\partial \theta_N^{i_N} \dots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0} = - \frac{\frac{\partial^m p(X|\theta)}{\partial \theta_N^{i_N} \dots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0}}{q(X)}$$

and, similar to the discussion in (5.3),

$$B(X) := \frac{\partial^m p(X|\theta)}{\partial \theta_N^{i_N} \dots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

satisfies (5.2). Therefore, applying Lemma 5.1(2) shows that

$$\frac{\partial^m f(X|\theta)}{\partial \theta_N^{i_N} \cdots \partial \theta_d^{i_d}} \Big|_{(\theta_1, \dots, \theta_d)=0}, \frac{\partial f(X|\theta)}{\partial \theta_1} \Big|_{(\theta_1, \dots, \theta_d)=0}, \dots, \frac{\partial f(X|\theta)}{\partial \theta_{N-1}} \Big|_{(\theta_1, \dots, \theta_d)=0}$$

are linearly dependent as random variables. Thus, performing the coordinate transformation ensured by Theorem 4.1 will satisfy (5.6). Consequently, this statistical model corresponds to the case $m = \infty$ in Assumption 1(2), and from Corollary 3.1(2),

$$\lambda = \frac{r}{2} = \frac{N-1}{2}$$

is obtained (multiplicity is 1). □

As a simple application, we calculate the learning coefficient for a mixture of Bernoulli distributions.

Corollary 5.1 (Learning Coefficient for a Mixture of Bernoulli Distributions with Two Mixture Components). Consider a mixture distribution with two components of a probability distribution $\tilde{p}(x|\theta)$ with a single parameter $\theta \in \mathbb{R}$.

$$p(x|\theta_1, \theta_2, \tau) := \tau \cdot \tilde{p}(x|\theta_1) + (1 - \tau) \cdot \tilde{p}(x|\theta_2)$$

where the parameter τ represents the mixing proportion ($0 \leq \tau \leq 1$). Let \tilde{p} be the Bernoulli distribution, i.e.,

$$\tilde{p}(x|\theta) = \begin{cases} 1 - \theta & x = 0 \\ \theta & x = 1 \end{cases}$$

(where $0 < \theta < 1$). When the true distribution $q(x)$ consists of a single mixture component

$$q(x) = \tilde{p}(x|\theta_*) \quad (0 < \theta_* < 1)$$

the learning coefficient is $1/2$.

Proof. Consider the set of realizable parameters $(\theta_1, \theta_2, \tau)$, Θ_* , by considering the case $x = 1$,

$$\Theta_* = \{(\theta_1, \theta_2, \tau) \mid \tau\theta_1 + (1 - \tau)\theta_2 = \theta_*\}$$

Fix an arbitrary realizable parameter $(Q_1, Q_2, T) \in \Theta_*$ and translate it so that it moves to the origin, redefining this statistical model as $p(x|\theta_1, \theta_2, \tau)$.

$$p(x|\theta_1, \theta_2, \tau) := (\tau + T) \cdot \tilde{p}(x|\theta_1 + Q_1) + (1 - \tau - T) \cdot \tilde{p}(x|\theta_2 + Q_2)$$

By symmetry, we can assume $T \neq 1$. This statistical model is a random variable taking two values, and

$$\frac{\partial f}{\partial \theta_2} \Big|_{(\theta_1, \theta_2, \tau)=0} = \begin{cases} \frac{1-T}{1-\theta_*}, & (x=0) \\ -\frac{1-T}{\theta_*}, & (x=1) \end{cases}$$

is a non-zero random variable, making this statistical model semi-regular. Since (Q_1, Q_2, T) is arbitrary, this statistical model is semi-regular for all realizable parameters, from Theorem 5.1, the learning coefficient is $1/2$ (multiplicity is 1). □

Remark 5.3. Instead of using Theorem 5.1, one can also determine the real log canonical threshold by explicitly performing a variable transformation using Proposition 4.2. The local equation for Θ_* is given by

$$(\tau + T)(\theta_1 + Q_1) + (1 - \tau - T)(\theta_2 + Q_2) - \theta_* = 0$$

Solving this equation for θ_2 yields the solution:

$$\begin{aligned} \theta_2 &= \frac{\theta_* - (\tau + T)(\theta_1 + Q_1)}{1 - \tau - T} - Q_2 \\ &= \frac{TQ_1 + (1 - T)Q_2 - (\tau + T)(\theta_1 + Q_1)}{1 - \tau - T} - Q_2 \\ &= -\frac{\tau + T}{1 - \tau - T}\theta_1 - \frac{Q_1 - Q_2}{1 - \tau - T}\tau \end{aligned}$$

The variable transformation defined by

$$\tilde{\theta}_2 = \theta_2 + \frac{\tau + T}{1 - \tau - T}\theta_1 + \frac{Q_1 - Q_2}{1 - \tau - T}\tau$$

transforms the local equation into:

$$\begin{aligned} &(\tau + T)(\theta_1 + Q_1) + (1 - \tau - T)(\theta_2 + Q_2) - \theta_* \\ &= (\tau + T)(\theta_1 + Q_1) + (1 - \tau - T) \left(\tilde{\theta}_2 - \frac{\tau + T}{1 - \tau - T}\theta_1 - \frac{Q_1 - Q_2}{1 - \tau - T}\tau + Q_2 \right) - \theta_* \\ &= (1 - \tau - T)\tilde{\theta}_2 \end{aligned}$$

which can be realized only if $\tilde{\theta}_2 = 0$. Thus, treating (θ_1, τ) as arbitrary constants in the neighborhood of zero,

$$\frac{\partial f}{\partial \tilde{\theta}_2} \Big|_{\tilde{\theta}_2=0} = -\frac{1 - \tau - T}{\tilde{p}(x|\theta_*)} = \begin{cases} -\frac{1-\tau-T}{1-\theta_*} & x=0 \\ -\frac{1-\tau-T}{\theta_*} & x=1 \end{cases}$$

which is a non-zero random variable. Therefore, applying Corollary 3.1 in the case $(d, r, m) = (3, 1, \infty)$, the real log canonical threshold of the realizable parameter (Q_1, Q_2, T) is $1/2$ (multiplicity is 1).

6 Conclusion

In this paper, we first clarified the relationship between the Taylor expansion of Kullback-Leibler divergence and the log-likelihood ratio function (Proposition 3.1). We then applied these findings to semi-regular models, where the rank of the Fisher information matrix is non-zero, and provided a formula related to the Taylor expansion of Kullback-Leibler divergence in Main Theorem 1. In Main Theorem 2, using the Taylor expansion derived in Main Theorem 1, we performed specific blow-ups and obtained an evaluation concerning the real log canonical thresholds. Specifically, we derived a formula for the exact value of the real log canonical threshold in cases where certain conditions of linear independence are met (Corollary 3.1).

The discussions above require Assumption 1, but Theorem 4.1 shows that, by appropriate variable transformations, Assumption 1 can be satisfied for any semi-regular model.

As practical examples of using the Main Theorem, we provided formulas for the real log canonical threshold for models where the number of parameters d and the rank of the Fisher information matrix is d or $d - 1$ (Proposition 5.1), as well as for discrete random variables that take a finite number of values (Theorem 5.1). We also provided an application example that gives the exact learning coefficient for a mixture of Bernoulli distributions with two components (Corollary 5.1).

Corollary 3.1 can directly compute the real log canonical threshold only for non-singular points in the set of realization parameters Θ_* . As future research, we aim to generalize the methods used in this study further and derive the real log canonical threshold at singular points.

Acknowledgments

I am grateful to Professor Joe Suzuki of Osaka University for teaching me the basics of Bayesian theory and for providing a research theme that bridges algebraic geometry and statistics, which has been my specialty. I also thank him for his valuable comments, advice, and for checking this manuscript. I also express my gratitude to Professor Sumio Watanabe, who proposed the concept of the learning coefficient.

References

- [1] Heisuke Hironaka. Resolution of Singularities of an Algebraic Variety Over a Field of Characteristic Zero. *Annals of Mathematics*, 79:109-326, 1964.
- [2] S. Watanabe. *Algebraic Geometry and Statistical Learning Theory*, volume 25. Cambridge University Press, New York, USA, 2009.
- [3] M. Aoyagi and S. Watanabe. Stochastic complexities of reduced rank regression in Bayesian estimation. *Neural Networks*, 18(7):924-933, 2005.

Appendix A Mathematica Output Results

Example 3.1

```
In[*]:=  $\tilde{p}[x_, \theta_] := \text{Binomial}[2, x] \theta^x (1 - \theta)^{2-x}$ 
In[*]:=  $p[x_, \theta1_, \theta2_] := \frac{1}{2} \tilde{p}\left[x, \theta1 - \theta2 + \frac{1}{2}\right] + \frac{1}{2} \tilde{p}\left[x, \theta2 + \frac{1}{2}\right]$ 
In[*]:=  $f[x_, \theta1_, \theta2_] := \text{Log}\left[\frac{\tilde{p}\left[x, \frac{1}{2}\right]}{p[x, \theta1, \theta2]}\right]$  (*Log-likelihood ratio function*)
In[*]:=  $K[\theta1_, \theta2_] := \text{Sum}\left[f[x, \theta1, \theta2] \tilde{p}\left[x, \frac{1}{2}\right], \{x, \theta, 2\}\right]$  (*Kullback-Leibler divergence*)
In[*]:=  $a[m_, n_] := \text{Simplify}\left[\frac{1}{m! n!} D[K[\theta1, \theta2], \{\theta1, m\}, \{\theta2, n\}] /. \{\theta1 \rightarrow \theta, \theta2 \rightarrow \theta\}\right]$ 
(*coefficient of the term  $\theta1^m \theta2^n$  in the Taylor expansion of  $K(\theta)$ *)
In[*]:=  $c[m_] := \text{Sum}\left[a[n, m - n] \theta1^n \theta2^{m-n}, \{n, \theta, m\}\right]$  (*mth order terms of  $K(\theta)$ *)
```

Taylor expansion of $K(\theta)$

```
In[*]:= c[2] (*2nd order terms*)
```

```
Out[*]=
 $\theta1^2$ 
```

```
In[*]:= c[3] (*3rd order terms*)
```

```
Out[*]=
 $0$ 
```

```
In[*]:= c[4] (*4th order terms*)
```

```
Out[*]=
 $8 \theta1^2 \theta2^2 - 16 \theta1 \theta2^3 + 8 \theta2^4$ 
```

derivative of the log - likelihood ratio function

```
In[*]:=  $df[x_, m_, n_] := \text{Simplify}\left[\frac{1}{m! n!} D[f[x, \theta1, \theta2], \{\theta1, m\}, \{\theta2, n\}] /. \{\theta1 \rightarrow \theta, \theta2 \rightarrow \theta\}\right]$ 
(*coefficient of the term  $\theta1^m \theta2^n$  in the Taylor expansion of  $f(x|\theta)$ *)
```

```
In[*]:= Print[df[0, 1, 0], "(x=0)"];
Print[df[1, 1, 0], "(x=1)"];
Print[df[2, 1, 0], "(x=2)"];
(*coefficient of the term  $\theta1$  in the Taylor expansion of  $f(x|\theta)$ *)
```

```
2 (x=0)
```

```
 $0$  (x=1)
```

```
-2 (x=2)
```

```
In[*]:= Print[df[0, 0, 1], "(x=0)"];
Print[df[1, 0, 1], "(x=1)"];
Print[df[2, 0, 1], "(x=2)"];
(*coefficient of the term  $\theta2$  in the Taylor expansion of  $f(x|\theta)$ *)
```

$\theta(x=0)$

$\theta(x=1)$

$\theta(x=2)$

```
In[*]:= Print[df[0, 0, 2], "(x=0)"];  
Print[df[1, 0, 2], "(x=1)"];  
Print[df[2, 0, 2], "(x=2)"];  
(*coefficient of the term  $\theta^2$  in the Taylor expansion of  $f(x|\theta)$ *)  
-4 (x=0)  
4 (x=1)  
-4 (x=2)
```