

# EXISTENCE AND NONEXISTENCE OF MINIMIZERS FOR CLASSICAL CAPILLARITY PROBLEMS IN PRESENCE OF NONLOCAL REPULSION AND GRAVITY

GIULIO PASCALE

**ABSTRACT.** We investigate, under a volume constraint and among sets contained in a Euclidean half-space, the minimization problem of an energy functional given by the sum of a capillarity perimeter, a nonlocal repulsive term and a gravitational potential energy. The capillarity perimeter assigns a constant weight to the portion of the boundary touching the boundary of the half-space. The nonlocal term is represented by a double integral of a positive kernel  $g$ , while the gravitational term is represented by the integral of a positive potential  $G$ .

We first establish existence of volume-constrained minimizers in the small mass regime, together with several qualitative properties of minimizers. The existence result holds even for rather general choices of kernels in the nonlocal term, including attractive-repulsive ones.

When the nonlocal kernel  $g(x) = 1/|x|^\beta$  with  $\beta \in (0, 2]$ , we also obtain nonexistence of volume constrained minimizers in the large mass regime.

Finally, we prove a generalized existence result of minimizers holding for all masses, meaning that the infimum of the problem is realized by a finite disjoint union of sets thought located at “infinite distance” one from the other.

These results stem from an application of quantitative isoperimetric inequalities for the capillarity problem in a half-space.

## CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Existence of minimizers for small masses	6
4. Nonexistence of minimizers for large masses	14
5. Generalized minimizers	21
References	28

## 1. INTRODUCTION

The classical liquid drop model for the atomic nucleus in the Euclidean space  $\mathbb{R}^n$ , for  $n \geq 2$ , aims to characterize minimizers of the functional

$$P(E) + \int_E \int_E \frac{1}{|y-x|^\beta} dy dx$$

among sets with a given volume, where  $0 < \beta < n$  is a given parameter and  $P(E)$  denotes the perimeter of  $E \subset \mathbb{R}^n$ . There is a clear competition between the two terms in the energy, since the ball at the same time minimizes the perimeter, by the isoperimetric inequality [De 58], [Mag12, Theorem 14.1], and maximizes the second term, by the Riesz rearrangement inequality [Rie30], [LL01, Theorem 3.7]. The physically relevant case is when  $\beta = 1$  and  $n = 3$ , that is when the second term is the Coulombic energy. This case goes back to Gamow’s liquid drop model for atomic nuclei [Gam30], subsequently developed by von Weizsäcker [Wei35], Bohr [Boh36; BW39], and many other researchers. This model is used to explain various properties of nuclear matter [CPS74; CS62; MS96; PTM90], but it also arises in the Ohta-Kawasaki model for diblock copolymers [OK86] and in many other physical situations, see [CM75; CK93; Gen79; EK93; GDM95; KN86; Mam94; Nag95; NKD94]. For a more specific account on the physical background of this kind of problems, we refer to [Mur02].

In the last decades, the model for general  $\beta$  and  $n$  has gained renewed interest in mathematics literature, in order to investigate existence and non-existence of minimizers and the minimality of the ball. In [KM13; KM14], Knüpfer and Muratov proved that balls are the only minimizers in the small mass regime when  $n = 2$  and when  $3 \leq n \leq 7$  with  $0 < \beta < n - 1$ . At the same time they obtained nonexistence results when  $n \geq 2$  and  $\beta \in (0, 2)$ . See

*Date:* June 18, 2024.

*2020 Mathematics Subject Classification.* Primary: 49J40, 49Q10. Secondary: 49Q20, 28A75, 26B30.

*Key words and phrases.* Isoperimetric problem, capillarity, Riesz energy, gravitational potential, existence, nonexistence, generalized existence.

also the alternative proofs [FKN16; LO14; Jul14] in the case  $\beta = 1$  and  $n = 3$  and [MZ14] in the case  $n = 2$  with  $\beta$  sufficiently small. Later on, Bonacini and Cristoferi [BC14] proved existence and uniqueness results for every  $n$  and  $0 < \beta < n - 1$ . Finally, Figalli, Fusco, Maggi, Millot and Morini [Fig+15] studied the case  $0 < \beta < n$  for every  $n$ , even replacing the perimeter  $P(E)$  by the fractional perimeter  $P_s(E)$ ,  $0 < s \leq 1$ . We refer to [CMT17; NO23a] for a review on the topic and to [CNT22; Fra19; FL15; FN21; Jul17; NO23b] and references therein for further results on the nonlocal liquid drop model. A variant of the problem with a constant background has been studied by [ACO09; CS13; CP10; CP11; EFK20; FL19; KMN16]; see also [AFM13; Ala+19; CN17; FNV18; GMS13; GMS14; Mur10; Nam20; Ono22; ST11] for further results on related problems.

In this paper we prove existence and nonexistence results of minimizers in a capillarity context with nonlocal and gravitational terms. If  $E$  is a measurable set in the half-space  $\{x_n > 0\} \subset \mathbb{R}^n$ ,  $n \geq 2$ , and  $\lambda \in (-1, 1)$ , we define the weighted perimeter functional

$$P_\lambda(E) := P(E, \mathbb{R}^n \setminus H) - \lambda \mathcal{H}^{n-1}(\partial^* E \cap \partial H),$$

where  $\mathcal{H}^k$ , with  $k \geq 0$ , denotes the  $k$ -dimensional Hausdorff measure in  $\mathbb{R}^n$ , and  $\partial^* E$  denotes the reduced boundary of  $E$ , see Section 2. Interpreting the perimeter as a measure of the surface tension of a liquid drop, the constant  $\lambda$  basically represent the relative adhesion coefficient between a liquid drop and the solid walls of the container given by  $\{x_n > 0\}$ . If  $m > 0$ , minimizers for the isoperimetric capillarity problem

$$(1.1) \quad \inf \{P_\lambda(E) : E \subset \{x_n > 0\}, |E| = m\}$$

are given by suitably truncated balls lying on the boundary of the half-space, see [Mag12, Theorem 19.21]. More precisely, if  $B^\lambda := \{x \in B_1(0) \subset \mathbb{R}^n : \langle x, e_n \rangle > \lambda\}$ ,  $m > 0$  and

$$B^\lambda(m) := \frac{m^{\frac{1}{n}}}{|B^\lambda|^{\frac{1}{n}}}(B^\lambda - \lambda e_n),$$

minimizers for (1.1) are sets of the form

$$B^\lambda(m, x) := B^\lambda(m) + x,$$

with  $x \in \{x_n = 0\}$ , see also [PP24, Fig. 1]

If  $g : \mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty)$ , we define the Riesz-type potential energy

$$\mathcal{R}(E) := \int_E \int_E g(y - x) dy dx.$$

Finally, given a function  $G : (0, \infty) \rightarrow (0, \infty)$ , we define the gravity-type potential energy

$$\mathcal{G}(E) := \int_E G(x_n) dx.$$

If  $m > 0$  and we denote

$$\mathcal{F}^\lambda(E) := P_\lambda(E) + \mathcal{R}(E) + \mathcal{G}(E),$$

we consider the minimization problem

$$(1.2) \quad \inf \{\mathcal{F}^\lambda(E) : E \subset \{x_n > 0\}, |E| = m\}.$$

In the context of minimization of energies with general Riesz-type potential in the Euclidean space  $\mathbb{R}^n$ , Novaga and Pratelli in [NP21] showed the existence of minimizers in a generalized sense. Later on, Carazzato, Fusco and Pratelli in [CFP23] showed that the ball is the unique minimizer in the small mass regime when the nonlocal kernel  $g$  is radial and decreasing. Pegon in [Peg21] proved that, if the kernel  $g$  decays sufficiently fast at infinity and if the volume is sufficiently large, then minimizers exist and converge to a ball as the volume goes to infinity. Then, Merlet and Pegon [MP22] proved that in the planar case minimizers are actually balls in the large mass regime. In [NO22], Novaga and Onoue obtained existence of minimizers for any volume and convergence to a ball as volume goes to infinity, if the Riesz potential decays sufficiently fast and even if the perimeter  $P(E)$  is replaced by the fractional perimeter  $P_s(E)$ ,  $0 < s < 1$ . We refer to [BNO23; CN18; MW21; MS19; Rig00] and references therein for further results on variational problems involving nonlocal energies.

The first result in this paper is an existence result in the capillarity context and in the small mass regime, together with a qualitative properties of volume constrained minimizers.

**Theorem 1.1.** *Let  $g$  be a  $\mathcal{R}$ -admissible  $q$ -decreasing function,  $q \geq 0$ , and let  $G$  be a  $\mathcal{G}$ -admissible function. There exists a mass  $\tilde{m} = \tilde{m}(n, \lambda, g, G, q) > 0$  such that, for every  $m \in (0, \tilde{m})$ , there exists a minimizer of  $\mathcal{F}^\lambda$  in the class*

$$\mathcal{A}_m := \{\Omega \subset \mathbb{R}^n \setminus H \text{ measurable} : |\Omega| = m\}.$$

*Moreover, if  $g$  is also infinitesimal, minimizers are indecomposable and, if in addition  $g$  is symmetric, minimizers are essentially bounded.*

*Finally, if  $g$  is also 0-decreasing, infinitesimal and symmetric and  $G$  is coercive, minimizers have no holes, i.e., if  $E$  is a minimizer of  $\mathcal{F}^\lambda$  in  $\mathcal{A}_m$ , there is no set  $F \subset \mathbb{R}^n \setminus (H \cup E)$  with  $|F| > 0$  such that*

$$P_\lambda(E) = P_\lambda(E \cup F) + P(F, \mathbb{R}^n \setminus H) + \lambda \mathcal{H}^{n-1}(\partial^* F \cap \partial H).$$

Let us make some comments on the definitions present in Theorem 1.1, while referring to Section 2 for their precise enunciation. The ‘‘admissibility’’ requirements on the kernels just refer to some necessary integrability conditions. The infinitesimality of  $g$  and the coercivity of  $G$  concern the behavior of these functions as the variable diverge, while the symmetry of  $g$  is referred to the symmetry with respect to the origin. The  $q$ -decreasing property is a way more general condition than the classical radial decreasing condition, denoted in this setting as 0-decreasing property. We point out that attractive-repulsive kernels of the type

$$(1.3) \quad g(x) = |x|^{\beta_1} + \frac{1}{|x|^{\beta_2}}, \quad \beta_1 > 0, \quad \beta_2 \in (0, n),$$

are  $q$ -decreasing for any  $q \geq \beta_1$ , even if they diverge positively as  $|x| \rightarrow +\infty$ , see Definition 2.2 and Remark 2.4. In particular attractive-repulsive kernels as in (1.3) represent a possible choice in the definition of  $\mathcal{F}^\lambda$  in Theorem 1.1. Minimization problems for attractive-repulsive functionals have been widely studied in the last years. Existence and nonexistence results are addressed in [BCT18; FL18; FL21], while stability and uniqueness of minimizers have been respectively studied in [BCT24; Lop19]. We refer to [Car23; CP22; CPT23; CDM16] for further results about analogous problems.

We remark that, by a symmetry argument, analyzing the Euler-Lagrange equation of problem (1.2), it is possible to verify that the sets  $B^\lambda(m, x)$  are not volume constrained minimizers of  $\mathcal{F}^\lambda$ ; actually, the isoperimetric bubbles  $B^\lambda(m, x)$  are not even volume constrained critical points of  $\mathcal{F}^\lambda$ . It is left as a future project to study quantitative properties of minimizers to (1.2), such as the proximity of minimizers from bubbles  $B^\lambda(m, x)$  in terms of the smallness of the mass.

For large masses and for suitable choices of  $g$ , the repulsive interaction dominates and the variational problem in Theorem 1.1 does not admit a minimizer.

**Theorem 1.2.** *Let*

$$g(x) = \frac{1}{|x|^\beta}, \quad 0 < \beta < n, \quad x \in \mathbb{R}^n \setminus \{0\}$$

*and let  $G$  be  $\mathcal{G}$ -admissible. For every  $\beta \in (0, 2]$ , there exists  $\tilde{m} > 0$ , depending on  $n, \lambda, \beta, G$ , such that for all  $m \geq \tilde{m}$  the minimization problem*

$$\inf \{ \mathcal{F}^\lambda(E) : E \subset \mathbb{R}^n \setminus H, |E| = m \}$$

*has no minimizers.*

Therefore, for a general  $g$ , existence may fail for masses large enough, since minimizers tend to split in two or more components which then move apart one from the other in order to decrease the nonlocal energy. To capture this phenomenon, it is convenient to introduce a generalized energy defined as

$$\tilde{\mathcal{F}}^\lambda(E) := \inf_{h \in \mathbb{N}} \tilde{\mathcal{F}}_h^\lambda(E),$$

where

$$\tilde{\mathcal{F}}_h^\lambda(E) := \inf \left\{ \sum_{i=1}^h \mathcal{F}^\lambda(E^i) : E = \bigcup_{i=1}^h E^i, E^i \cap E^j = \emptyset \text{ for } 1 \leq i \neq j \leq h \right\}.$$

Note that in this functional the interaction between different components is not evaluated, which corresponds to consider them ‘‘at infinite distance’’ one from the other. By considering  $\tilde{\mathcal{F}}^\lambda$  instead of  $\mathcal{F}^\lambda$ , we can prove the following generalized existence result.

**Theorem 1.3.** *Let  $g$  be a  $\mathcal{R}$ -admissible  $q$ -decreasing function,  $q \geq 0$ , and let  $G$  be a  $\mathcal{G}$ -admissible function. For every  $m > 0$  there exists a minimizer of  $\tilde{\mathcal{F}}^\lambda$  in the class*

$$\mathcal{A}_m = \{\Omega \subset \mathbb{R}^n \setminus H \text{ measurable} : |\Omega| = m\}.$$

More precisely, there exist a set  $E \in \mathcal{A}$  and a subdivision  $E = \cup_{j=1}^h E^j$ , with pairwise disjoint sets  $E^j$ , such that

$$\tilde{\mathcal{F}}^\lambda(E) = \sum_{j=1}^h \mathcal{F}^\lambda(E^j) = \inf \{ \tilde{\mathcal{F}}^\lambda(\Omega) : \Omega \in \mathcal{A} \}.$$

Moreover, for every  $1 \leq j \leq h$ , the set  $E^j$  is a minimizer of both the standard and the generalized energy for its volume, i.e.

$$\tilde{\mathcal{F}}^\lambda(E^j) = \mathcal{F}^\lambda(E^j) = \min \{ \tilde{\mathcal{F}}^\lambda(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = |E^j| \}.$$

**Strategy of the proof and comments.** The proof of Theorem 1.1 is divided into several steps. In the spirit of [KM14], the existence of minimizers in the small mass regime follows by the direct method of the calculus of variations, see Theorem 3.1, once we show that for sufficiently small mass every minimizing sequence of the energy may be replaced by another minimizing sequence where all sets have uniformly bounded diameter, see Lemma 3.5. We remark that we heavily use the quantitative isoperimetric inequality for the capillarity problem proved in [PP24], which estimates the Fraenkel asymmetry of a competitor with respect to the optimal sets in terms of the energy deficit. Note that it is unclear at the moment how to apply stronger isoperimetric inequalities of Fuglede-type [CL12; Fug89] for nearly spherical sets in the present capillarity framework; instead, stronger isoperimetric inequalities have been used as fundamental tools, for example, in [AFM13; BC14; CFP23]. In fact, the classical Fuglede's method relies on the precise knowledge of the eigenvalues of the Laplace-Beltrami operator, which is not available for  $P_\lambda$  on optimal sets for generic  $\lambda \in (-1, 1)$ . Moreover, in our case it is in general not possible to globally parametrize  $C^1$ -close boundaries one on the other as normal graphs.

The boundedness result in Theorem 1.1 follows once we show that minimizers enjoy uniform density estimates at boundary points. In order to do so, we prove that, under suitable conditions on the Riesz potential, minimizers are  $(K, r_0)$ -quasiminimal sets for all masses, see Definition 3.8 and Lemma 3.11. Indeed quasiminimal sets have well-known topological regularity properties (Theorem 3.9), which easily guarantee boundedness, see Theorem 3.6. Note that the lack of symmetry of the problem, due to the presence of gravitational potential and the fact that ambient space is a half-space, forces us to deal with the vertical direction in a separate way, see Lemma 3.10.

The absence of holes is based on the combination of some techniques from [KM14] and [NP21]. We firstly prove some density estimates which improve, under suitable hypotheses on  $g$ , the analogous estimates for quasiminimal sets, by providing bounds independent of the minimizer, see Lemma 4.2. In fact, this allows to prove the boundedness in the vertical direction with a bound independent of the minimizer, see Lemma 4.8, and to obtain absence of holes arguing by contradiction, see Theorem 4.7.

The proof of Theorem 1.2 is based on the combination of some techniques from [FN21] and [KM14] and exploits some estimates on the diameter and the nonlocal potential energy of minimizers. We remark that the range of the exponent  $\beta$  in Theorem 1.2 is the same as the analogous nonexistence results in the classical setting [CNT22; FKN16; FN21; KM14; LO14].

The proof of Theorem 1.3 is inspired by [NP21] and exploits the isoperimetric inequality for the capillarity functional  $P_\lambda$ . In our case the argument must be modified to take into account the presence of the gravitational energy and, as before, estimates in the vertical direction must be treated separately. We remark that also the possible choices for the kernels  $g$  in our Theorem 1.3 allow for more freedom than those considered in [NP21].

**Organization.** In Section 2 we collect definitions and facts on sets of finite perimeter and capillarity functional. In Section 3 we begin to prove Theorem 1.1, in particular we prove existence, boundedness and indecomposability of minimizers. In Section 4 we prove Theorem 1.2, together with absence of holes in minimizers in the small mass regime, thus completing the proof of Theorem 1.1. In Section 5 we prove Theorem 1.3.

**Acknowledgments.** The author is member of INdAM - GNAMPA. The author is grateful to Marco Pozzetta for many suggestions and for stimulating discussions.

## 2. PRELIMINARIES

From now on and for the rest of the paper we assume that  $\lambda \in (-1, 1)$  and  $n \in \mathbb{N}$  with  $n \geq 2$  are fixed.

### List of symbols.

- $|\cdot|$  denotes Lebesgue measure in  $\mathbb{R}^n$ .
- $B^\lambda = \{x \in B : \langle x, e_n \rangle > \lambda\}$ .

- $B^\lambda(m) := \frac{m^{\frac{1}{n}}}{|B^\lambda|^{\frac{1}{n}}}(B^\lambda - \lambda e_n)$ , for any  $m > 0$ .
- $B^\lambda(m, x) := B^\lambda(m) + x$ , for any  $x \in \{x_n = 0\}$ . In particular  $B^\lambda(m) = B^\lambda(m, 0)$ .
- $c(\cdot), C(\cdot)$  denote strictly positive constants, that may change from line to line.
- $H := \{x_n \leq 0\}$ .
- $\mathcal{H}^d$  denotes  $d$ -dimensional Hausdorff measure in  $\mathbb{R}^n$ , for  $d \geq 0$ .
- $Q_{\frac{r}{2}}$  denotes a generic cube in  $\mathbb{R}^n$  of side  $r > 0$ .
- $R_\lambda := \max \left\{ \sqrt{1 - \lambda^2}, 1 - \lambda \right\}$ .

**2.1. Sets of finite perimeter.** We recall basic definitions and properties regarding sets of finite perimeter, referring to [AFP00; Mag12] for a complete treatment on the subject. The perimeter of a measurable set  $E \subset \mathbb{R}^n$  in an open set  $A \subset \mathbb{R}^n$  is defined by

$$(2.1) \quad P(E, A) := \sup \left\{ \int_E \operatorname{div} T(x) \, dx : T \in C_c^1(A; \mathbb{R}^n), \|T\|_\infty \leq 1 \right\}.$$

Denoting  $P(E) := P(E, \mathbb{R}^n)$ , we say that  $E$  is a set of finite perimeter if  $P(E) < +\infty$ . In such a case, the characteristic function  $\chi_E$  has a distributional gradient  $D\chi_E$  that is a vector-valued Radon measure on  $\mathbb{R}^n$  such that

$$\int_E \operatorname{div} T(x) \, dx = - \int_{\mathbb{R}^n} T \, dD\chi_E, \quad \forall T \in C_c^1(\mathbb{R}^n; \mathbb{R}^n).$$

It can be proved that the set function  $P(E, \cdot)$  defined in (2.1) is the restriction of a nonnegative Borel measure to open sets. The measure  $P(E, \cdot)$  coincides with the total variation  $|D\chi_E|$  of the distributional gradient, and it is concentrated on the reduced boundary

$$\partial^* E := \left\{ x \in \operatorname{spt} |D\chi_E| : \exists v^E(x) := - \lim_{r \rightarrow 0} \frac{D\chi_E(B_r(x))}{|D\chi_E(B_r(x))|} \text{ with } |v^E(x)| = 1 \right\}.$$

Introducing the sets of density  $t \in [0, 1]$  points for  $E$  defined by

$$E^{(t)} := \left\{ x \in \mathbb{R}^n : \lim_{r \rightarrow 0} \frac{|E \cap B_r(x)|}{|B_r(x)|} = t \right\},$$

we have that the reduced boundary coincides both with  $\mathbb{R}^n \setminus (E^{(1)} \cup E^{(0)})$  and with the set  $E^{(1/2)}$  up to  $\mathcal{H}^{n-1}$ -negligible sets. The vector  $v^E$  is called the generalized outer normal of  $E$ . Moreover  $P(E, \cdot) = \mathcal{H}^{n-1} \llcorner \partial^* E$ , and the distributional gradient can be written as  $D\chi_E = -v^E \mathcal{H}^{n-1} \llcorner \partial^* E$ .

**2.2. Preliminary results on the capillarity functional.** We recall basic properties on the functional  $P_\lambda$ . Let  $E \subset \mathbb{R}^n \setminus H$  with measure  $|E| = m \in (0, +\infty)$ . Note that

$$P_\lambda(E) = \int_{\partial^* E \setminus H} 1 - \lambda \langle e_n, v^E \rangle \, d\mathcal{H}^{n-1};$$

in particular  $P_\lambda(E) > 0$  [PP24, Remark 2.1, Corollary 2.4].

The sets  $B^\lambda(m, x)$  uniquely minimize the problem

$$\inf \{ P_\lambda(E) : E \subset \mathbb{R}^n, |E| = m \},$$

and this is encoded in the isoperimetric inequality [Mag12, Theorem 19.21], [PP24, Theorem 3.5]

$$(2.2) \quad P_\lambda(E) \geq n |B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}}.$$

The previous isoperimetric inequality can be strengthened in a quantitative version. We define the Fraenkel asymmetry

$$\alpha_\lambda(E) := \inf \left\{ \frac{|E \Delta B^\lambda(m, x)|}{m} : x \in \{x_n = 0\} \right\},$$

and the isoperimetric deficit

$$D_\lambda(E) := \frac{P_\lambda(E) - P_\lambda(B^\lambda(m))}{P_\lambda(B^\lambda(m))}.$$

Then the following sharp quantitative isoperimetric inequality holds

**Theorem 2.1** ([PP24]). *There exists a constant  $c(n, \lambda) > 0$  such that for any measurable set  $E \subset \mathbb{R}^n \cap \{x_n > 0\}$  with finite measure there holds*

$$(2.3) \quad \alpha_\lambda(E)^2 \leq c(n, \lambda) D_\lambda(E).$$

2.3. **Definitions.** We give some definitions for the Riesz-type potential  $\mathcal{R}$  and the gravity-type potential  $\mathcal{G}$ .

**Definition 2.2.** A function  $g : \mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty)$  is  $\mathcal{R}$ -admissible if  $\mathcal{R}(B_1) < \infty$ .

A  $\mathcal{R}$ -admissible function  $g : \mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty)$  is  $q$ -decreasing, for some  $q \in [0, \infty)$ , if for every  $x \in \mathbb{R}^n \setminus \{0\}$  and every  $\alpha > 1$  it holds

$$g(\alpha x) \leq \alpha^q g(x).$$

A  $\mathcal{R}$ -admissible function  $g : \mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty)$  is infinitesimal if

$$\lim_{|x| \rightarrow +\infty} g(x) = 0.$$

A  $\mathcal{R}$ -admissible function  $g : \mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty)$  is symmetric if

$$g(-x) = g(x) \quad \forall x \in \mathbb{R}^n \setminus \{0\}.$$

Given two measurable sets  $L, M \subset \mathbb{R}^n \setminus H$  we let

$$\mathcal{R}(L, M) := \int_L \int_M g(y-x) dy dx.$$

*Remark 2.3.* The functions  $\frac{1}{|x|^\beta}$ , for  $\beta \in (0, n)$ , are  $\mathcal{R}$ -admissible, 0-decreasing, infinitesimal and symmetric.

*Remark 2.4.* The attractive-repulsive kernels  $|x|^{\beta_1} + \frac{1}{|x|^{\beta_2}}$ , for  $\beta_1 > 0$  and  $\beta_2 \in (0, n)$ , are  $\mathcal{R}$ -admissible  $\beta_1$ -decreasing symmetric functions. At the same time they diverge positively as  $|x| \rightarrow +\infty$ .

**Definition 2.5.** A function  $G : (0, \infty) \rightarrow (0, \infty)$  is  $\mathcal{G}$ -admissible if

$$(2.4) \quad \sup_{t \in (0, 2)} G(t) < \infty,$$

and

$$(2.5) \quad G(\alpha t) \leq \alpha^n G(t), \quad \forall \alpha > 1, t > 0.$$

A  $\mathcal{G}$ -admissible function  $G : (0, \infty) \rightarrow (0, \infty)$  is coercive if

$$G(t) \rightarrow +\infty \quad \text{as } t \rightarrow +\infty.$$

*Remark 2.6.* The identity function  $G(t) = t$  on  $(0, +\infty)$  is a  $\mathcal{G}$ -admissible function.

*Remark 2.7.* Conditions (2.4) and (2.5) easily imply

$$G(t) = G(t \cdot 1) \leq t^n G(1) \leq c(G) t^n, \quad \forall t > 1.$$

### 3. EXISTENCE OF MINIMIZERS FOR SMALL MASSES

3.1. **Existence.** The goal of this Section is to prove the following

**Theorem 3.1.** *Let  $g$  be a  $\mathcal{R}$ -admissible  $q$ -decreasing function and let  $G$  be a  $\mathcal{G}$ -admissible function. There exists a mass  $\bar{m} = \bar{m}(n, \lambda, g, G, q) > 0$  such that, for all  $m \in (0, \bar{m})$ , there exists a minimizer of  $\mathcal{F}^\lambda$  in the class*

$$\mathcal{A}_m := \{\Omega \subset \mathbb{R}^n \setminus H \text{ measurable} : |\Omega| = m\}.$$

We begin by proving some preparatory lemmas, which estimate the energy of some competitors.

**Lemma 3.2.** *Let  $g$  be  $\mathcal{R}$ -admissible and  $G$  be  $\mathcal{G}$ -admissible. There exists a constant  $c = c(n, \lambda, g, G)$  such that*

$$\mathcal{R}(B^\lambda(m)) \leq c m, \quad \mathcal{G}(B^\lambda(m)) \leq c m$$

for every  $0 < m \leq |B^\lambda|$ .

*Proof.* Let us denote by  $\bar{Q}_l \subset \mathbb{R}^n \setminus H$ , with  $l > 0$ , the cube  $[-l, l] \times \dots \times [-l, l] \times [0, 2l]$ . For any  $N \in \mathbb{N}$  the cube  $\bar{Q}_1$  is the essential union of  $(2N)^n$  disjoint isometric cubes  $Q_{\frac{1}{2N}}^i$  of side  $1/2N$ . If  $\bar{Q}_{\frac{1}{2N}} \subset \bar{Q}_1$  is the cube

$\left[-\frac{1}{2N}, \frac{1}{2N}\right] \times \dots \times \left[-\frac{1}{2N}, \frac{1}{2N}\right] \times \left[0, \frac{1}{2N}\right]$ , evidently

$$\mathcal{R}(\bar{Q}_1) \geq \sum_{i=0}^{(2N)^n} \mathcal{R}(Q_{\frac{1}{2N}}^i) = (2N)^n \mathcal{R}(\bar{Q}_{\frac{1}{2N}}).$$

Moreover

$$2^n \sup_{(0,2)} G = |\bar{Q}_1| \sup_{(0,2)} G = (2N)^n \frac{|\bar{Q}_1|}{(2N)^n} \sup_{(0,2)} G = (2N)^n \int_{\bar{Q}_{\frac{1}{2N}}} \sup_{(0,2)} G dx \geq (2N)^n \mathcal{G}(\bar{Q}_{\frac{1}{2N}}).$$

For any  $0 < r \leq 1$  we denote by  $N$  the integer part of  $\frac{1}{2r}$ , so that  $(2r)^{-1} \leq 2N \leq r^{-1}$ . The above estimates, together with  $4rN \geq 1$ , imply that

$$\mathcal{R}(\bar{Q}_r) \leq (4r)^n N^n \mathcal{R}(\bar{Q}_{\frac{1}{2N}}) \leq 2^n \mathcal{R}(\bar{Q}_1) r^n$$

and

$$\mathcal{G}(\bar{Q}_r) \leq (4r)^n N^n \mathcal{G}(\bar{Q}_{\frac{1}{2N}}) \leq 4^n \left( \sup_{(0,2)} G \right) r^n.$$

If  $r = \frac{\frac{1}{m^n}}{|B^\lambda|^{\frac{1}{n}}} \leq 1$ , since  $B^\lambda(|B^\lambda|) \subset \bar{Q}_1$ , we get  $B^\lambda(m) \subset \bar{Q}_r$  and we conclude that

$$\mathcal{R}(B^\lambda(m)) \leq \mathcal{R}(\bar{Q}_r) \leq cr^n \leq cm$$

and

$$\mathcal{G}(B^\lambda(m)) \leq \mathcal{G}(\bar{Q}_r) \leq cr^n \leq cm.$$

□

**Corollary 3.3.** *Let  $g$  be  $\mathcal{R}$ -admissible and infinitesimal and let  $G$  be  $\mathcal{G}$ -admissible. For every  $m \geq 1$  there exists  $E \subset \mathbb{R}^n \setminus H$  with  $|E| = m$  such that  $\mathcal{F}^\lambda(E) \leq cm$  for some  $c$  depending on  $n, \lambda, g$  and  $G$ .*

*Proof.* Let us consider the set  $E$  given by a collection of  $N \geq 1$  spherical caps  $\{B^\lambda(v, x_i)\}_{1 \leq i \leq N}$  of equal volume  $v$  and with centers located at  $x_i = iRe_1, i = 1, \dots, N$ , with  $R$  large enough so that  $B^\lambda(v, x_i)$  are pairwise disjoint. We choose the number  $N$  as the smallest integer for which the volume of each spherical cap does not exceed  $\min\{1, |B^\lambda|\}$ . In particular  $Nv = m$  and  $N = \left\lceil \frac{m}{\min\{1, |B^\lambda|\}} \right\rceil$ . Note that, by [PP24, Lemma 3.3], since  $v \leq 1 \leq m = |E|$ ,

$$\begin{aligned} P_\lambda(E) &= P_\lambda\left(\bigcup_{i=1}^N B^\lambda(v, x_i)\right) = \sum_{i=1}^N P_\lambda(B^\lambda(v, x_i)) = c(n, \lambda) N v^{\frac{n-1}{n}} \leq c(n, \lambda) \left( \frac{m}{\min\{1, |B^\lambda|\}} + 1 \right) v^{\frac{n-1}{n}} \\ &\leq c(n, \lambda) (m v^{\frac{n-1}{n}} + v^{\frac{n-1}{n}}) \leq c(n, \lambda) (m 1^{\frac{n-1}{n}} + m 1^{\frac{n-1}{n}}) = c(n, \lambda) m. \end{aligned}$$

Moreover, let  $R$  be so large that  $g(x - y) < \frac{1}{N}$  for every  $x \in B^\lambda(v, x_j), y \in B^\lambda(v, x_k)$  with  $j \neq k$ . Then, by Lemma 3.2, since  $v \leq 1 \leq m = |E|$ ,

$$\begin{aligned} \mathcal{R}(E) &= \int_{\bigcup_{i=1}^N B^\lambda(v, x_i)} \int_{\bigcup_{i=1}^N B^\lambda(v, x_i)} g(y - x) dy dx \\ &= \sum_{i=1}^N \mathcal{R}(B^\lambda(v, x_i)) + \sum_{i=1}^N \int_{B^\lambda(v, x_i)} \int_{\widehat{B^\lambda(v, x_1) \cup \dots \cup B^\lambda(v, x_i) \cup \dots \cup B^\lambda(v, x_N)}} g(y - x) dy dx \\ &\leq c(n, \lambda, g, G) N v + N \frac{1}{N} v (m - v) \leq c(n, \lambda, g, G) m + (m - v) \leq c(n, \lambda, g, G) m, \end{aligned}$$

where the  $B^\lambda(v, x_1) \cup \dots \cup \widehat{B^\lambda(v, x_i)} \cup \dots \cup B^\lambda(v, x_N)$  denotes union over all the bubbles except for  $B^\lambda(v, x_i)$ . Finally, by Lemma 3.2

$$\begin{aligned} \mathcal{F}^\lambda(E) &= P_\lambda(E) + \mathcal{R}(E) + \mathcal{G}(E) \\ &\leq c(n, \lambda) |E| + c(n, \lambda, g, G) |E| + \sum_{i=1}^N \mathcal{G}(B^\lambda(v, x_i)) \\ &\leq c(n, \lambda) |E| + c(n, \lambda, g, G) |E| + c(n, \lambda, g, G) N v = c(n, \lambda, g, G) m. \end{aligned}$$

□

**Lemma 3.4.** *Let  $E \subset \mathbb{R}^n \setminus H$  be a set of finite perimeter. Let  $g$  be  $\mathcal{R}$ -admissible and  $q$ -decreasing and let  $G$  be  $\mathcal{G}$ -admissible. If  $\alpha > 1$ , then*

$$\mathcal{F}^\lambda(\alpha E) \leq \alpha^{2n+q} \mathcal{F}^\lambda(E).$$

*Proof.* Note that, if  $E \subset \mathbb{R}^n \setminus H$ , then  $\alpha E \subset \mathbb{R}^n \setminus H$ . Since  $\alpha > 1$ , by the positivity of  $P_\lambda$  we get

$$P_\lambda(\alpha E) = \alpha^{n-1} P_\lambda(E) \leq \alpha^{2n+q} P_\lambda(E).$$

Since  $g$  is  $q$ -decreasing, we have

$$\begin{aligned}\mathcal{R}(\alpha E) &= \int \int_{(\alpha E)^2} g(y-x) \, dy \, dx \\ &= \alpha^{2n} \int \int_{E \times E} g(\alpha(y-x)) \, dy \, dx \\ &\leq \alpha^{2n+q} \int \int_{E \times E} g(y-x) \, dy \, dx = \alpha^{2n+q} \mathcal{R}(E).\end{aligned}$$

Finally, by (2.5) we get

$$\int_{\alpha E} G(x_n) \, dx = \alpha^n \int_E G(\alpha x_n) \, dx \leq \alpha^{2n+q} \int_E G(x_n) \, dx.$$

□

The following lemma allows to suitably localize minimizing sequences with sufficiently small volume.

**Lemma 3.5.** *Let  $g$  be  $\mathcal{R}$ -admissible and  $q$ -decreasing and let  $G$  be  $\mathcal{G}$ -admissible. There exists  $\bar{m} > 0$ , depending on  $n, \lambda, g, G$  and  $q$ , such that, for every  $m \in (0, \bar{m})$  and every set of finite perimeter  $F \subset \mathbb{R}^n \setminus H$  with  $|F| = m$ , there exists a set of finite perimeter  $L$  with*

$$(3.1) \quad \mathcal{F}^\lambda(L) \leq \mathcal{F}^\lambda(F) \quad \text{and} \quad L \subset \bar{Q}_1 := [-1, 1] \times \cdots \times [-1, 1] \times [0, 2].$$

*Proof.* Throughout the proof we will assume that  $\bar{m} < \frac{|B^\lambda|}{4^n}$ . If  $\mathcal{F}^\lambda(B^\lambda(m)) \leq \mathcal{F}^\lambda(F)$ , then the assertion of the lemma is proved by choosing  $L = B^\lambda(m)$ . Then we can assume, by Lemma 3.2, that

$$(3.2) \quad \mathcal{F}^\lambda(F) < \mathcal{F}^\lambda(B^\lambda(m)) \leq c(n, \lambda, g, G) \max \left\{ m, m^{\frac{n-1}{n}} \right\}.$$

By Lemma 3.2

$$\begin{aligned}D_\lambda(F) &= \frac{c(n, \lambda)}{m^{\frac{n-1}{n}}} (P_\lambda(F) - P_\lambda(B^\lambda(m))) \\ &\leq \frac{c(n, \lambda)}{m^{\frac{n-1}{n}}} ([\mathcal{R}(B^\lambda(m)) - \mathcal{R}(F)] + [\mathcal{G}(B^\lambda(m)) - \mathcal{G}(F)]) \\ &\leq \frac{c(n, \lambda)}{m^{\frac{n-1}{n}}} (\mathcal{R}(B^\lambda(m)) + \mathcal{G}(B^\lambda(m))) \\ &\leq c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{\frac{1}{n}}.\end{aligned}$$

By the quantitative isoperimetric inequality (2.3)

$$\alpha_\lambda(F) \leq c(n, \lambda) \sqrt{D_\lambda(F)} \leq c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{\frac{1}{2n}}$$

and, after a suitable translation,

$$|B^\lambda(m) \Delta F| \leq c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{1 + \frac{1}{2n}}.$$

Since  $|F| = m = |B^\lambda(m)|$  we also have

$$|B^\lambda(m) \Delta F| = 2|F \setminus B^\lambda(m)|$$

and

$$(3.3) \quad |F \setminus B^\lambda(m)| \leq c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{1 + \frac{1}{2n}}.$$

For any  $\varrho > 0$  let  $F_1 = F \cap B_\varrho(0)$  and  $F_2 = F \setminus B_\varrho(0)$ . Note that for every  $\varepsilon > 0$  there exists  $\bar{m}$  sufficiently small such that, if  $\varrho \geq \frac{m^{\frac{1}{n}}}{|B^\lambda|^{\frac{1}{n}}} R_\lambda =: \varrho_m$ , with  $m < \bar{m}$ , then

$$(3.4) \quad |F_2| \leq \varepsilon |F_1|.$$



Indeed, since  $B^\lambda(|B^\lambda|) \subset B_{R_\lambda}(0)$ , we get  $B^\lambda(m) \subset B_\rho(0)$ . Moreover, by (3.3) and for sufficiently small  $\bar{m}$  we estimate

$$\begin{aligned} |F_1| &= |F \cap B_\rho(0)| \geq |F \cap B^\lambda(m)| \\ &= |F| - |F \setminus B^\lambda(m)| \\ &\geq |B^\lambda| \left( \frac{m}{|B^\lambda|} \right) - c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{1+\frac{1}{2n}} \\ &\geq \frac{|B^\lambda|}{2} \left( \frac{m}{|B^\lambda|} \right). \end{aligned}$$

and

$$|F \setminus B_\rho(0)| \leq |F \setminus B^\lambda(m)| \leq c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{1+\frac{1}{2n}} \leq c(n, \lambda, g, G) \left( \frac{m}{|B^\lambda|} \right)^{\frac{1}{2n}} |F_1| \leq \varepsilon |F_1|.$$

Let us define the monotonically decreasing function  $U(\rho) = |F \setminus B_\rho(0)|$ .

We now distinguish two cases. Let us firstly prove (3.1) when we assume that

$$(3.5) \quad \Sigma := P_\lambda(F_1) + P_\lambda(F_2) - P_\lambda(F) > \frac{1}{2} \mathcal{F}^\lambda(F_2) \quad \forall \rho \in \left( \rho_m, \frac{R_\lambda}{2} \right)$$

By (3.3) we have

$$U(\rho_m) = |F \setminus B^\lambda(m)| \leq c(n, \lambda, g, G) m^{1+\frac{1}{2n}} \leq c(n, \lambda, g, G) \rho_m^{n+\frac{1}{2}}$$

Furthermore, by (2.2) and (3.5) we have

$$-2 \frac{dU(\rho)}{d\rho} = \Sigma > \frac{1}{2} \mathcal{F}^\lambda(F_2) \geq \frac{1}{2} P_\lambda(F \setminus B_\rho(0)) \geq c(n, \lambda) U^{\frac{n-1}{n}}(\rho).$$

In particular we have

$$\begin{cases} \frac{dU(\rho)}{d\rho} \leq -c(n, \lambda) U^{\frac{n-1}{n}}(\rho) & \text{for a.e. } \rho \in \left( \rho_m, \frac{R_\lambda}{2} \right) \\ U(\rho_m) \leq c(n, \lambda, g, G) \rho_m^{n+\frac{1}{2}}. \end{cases}$$

By ODE comparison we deduce that, if  $\bar{m} < 1$ ,

$$\begin{aligned} U(\rho)^{\frac{1}{n}} &\leq U(\rho_m)^{\frac{1}{n}} - c(n, \lambda) (\rho - \rho_m) \\ &\leq c(n, \lambda, g, G) \rho_m^{1+\frac{1}{2n}} - c(n, \lambda) \rho + c(n, \lambda) \rho_m \\ &= c(n, \lambda, g, G) m^{\frac{1}{n}+\frac{1}{2n^2}} - c(n, \lambda) \rho + c(n, \lambda) m^{\frac{1}{n}} \\ &\leq c(n, \lambda, g, G) m^{\frac{1}{n}} - c(n, \lambda) \rho + c(n, \lambda) m^{\frac{1}{n}} \\ &= c(n, \lambda, g, G) m^{\frac{1}{n}} - c(n, \lambda) \rho. \end{aligned}$$

For  $\bar{m}$  sufficiently small, it follows that  $U(\rho) = 0$  for  $\rho \geq \frac{R_\lambda}{2}$ , and we obtain (3.1) with  $L = F$ .

Let us prove (3.1) assuming that

$$(3.6) \quad \Sigma \leq \frac{1}{2} \mathcal{F}^\lambda(F_2)$$

holds for some  $\rho_0 \in \left( \rho_m, \frac{R_\lambda}{2} \right)$ . Let  $m_1 := |F_1|$ ,  $m_2 := |F_2|$  and  $\gamma := \frac{m_2}{m_1} \leq \varepsilon$ , with  $\varepsilon$  that will be chosen suitably small later. Let us also denote  $\tilde{F} = l F_1$ , with  $l := (1 + \gamma)^{\frac{1}{n}}$ . In particular  $|\tilde{F}| = m$  and, if  $\varepsilon$  is sufficiently small,

$$\tilde{F} = (1 + \gamma)^{\frac{1}{n}} F_1 = (1 + \gamma)^{\frac{1}{n}} (F \cap B_{\rho_0}(0)) \subset B_{\rho_0 \sqrt[1+\gamma]{1+\gamma}}(0) \subset B_{R_\lambda}(0) \subset \bar{Q}_1.$$

By Lemma 3.4

$$(3.7) \quad \begin{aligned} \mathcal{F}^\lambda(\tilde{F}) &= \mathcal{F}^\lambda(l F_1) \leq l^{2n+q} \mathcal{F}^\lambda(F_1) \\ &= \mathcal{F}^\lambda(F_1) + (l^{2n+q} - 1) \mathcal{F}^\lambda(F_1). \end{aligned}$$

Choosing  $\varepsilon \leq 1$ , we have  $1 \leq l \leq 2^{\frac{1}{n}}$ , and by Taylor's formula we obtain  $l^{2n+q} - 1 = (1 + \gamma)^{2+q/n} - 1 \leq \gamma K$  for some  $K > 0$  independent of  $\gamma$  and for  $\varepsilon$  sufficiently small. By (3.7) we arrive at

$$\mathcal{F}^\lambda(\tilde{F}) - \mathcal{F}^\lambda(F_1) \leq \gamma K \mathcal{F}^\lambda(F_1).$$

By the definition of  $\Sigma$  and since  $\mathcal{R}(F_1) + \mathcal{R}(F_2) \leq \mathcal{R}(F)$

$$(3.8) \quad \begin{aligned} \mathcal{F}^\lambda(\tilde{F}) - \mathcal{F}^\lambda(F) &\leq \mathcal{R}(F_1) + \mathcal{G}(F_1) + \mathcal{R}(F_2) + \mathcal{G}(F_2) - \mathcal{R}(F) - \mathcal{G}(F) + \Sigma - \mathcal{F}^\lambda(F_2) + \gamma K \mathcal{F}^\lambda(F_1) \\ &\leq -\frac{1}{2} \mathcal{F}^\lambda(F_2) + \gamma K \mathcal{F}^\lambda(F_1). \end{aligned}$$

By positivity of  $\mathcal{R}$  and  $\mathcal{G}$  and the isoperimetric inequality (2.2), we have  $\mathcal{F}^\lambda(F_2) > P_\lambda(F_2) \geq c(n, \lambda) m_2^{\frac{n-1}{n}}$ . By (3.6) we obtain

$$(3.9) \quad \begin{aligned} \mathcal{F}^\lambda(F) - \mathcal{F}^\lambda(F_1) &= P_\lambda(F) + \mathcal{R}(F) + \mathcal{G}(F) - P_\lambda(F_1) - \mathcal{R}(F_1) - \mathcal{G}(F_1) - P_\lambda(F_2) + P_\lambda(F_2) \\ &\geq -\frac{1}{2} \mathcal{F}^\lambda(F_2) + \mathcal{R}(F) + \mathcal{G}(F) - \mathcal{R}(F_1) - \mathcal{G}(F_1) + P_\lambda(F_2) \\ &= -\frac{1}{2} P_\lambda(F_2) - \frac{1}{2} \mathcal{R}(F_2) - \frac{1}{2} \mathcal{G}(F_2) + \mathcal{R}(F) + \mathcal{G}(F) - \mathcal{R}(F_1) - \mathcal{G}(F_1) + P_\lambda(F_2) \geq 0, \end{aligned}$$

that is  $\mathcal{F}^\lambda(F_1) \leq \mathcal{F}^\lambda(F)$ . By (3.2), since  $\gamma m \leq 2m_2$  and  $\gamma \leq \varepsilon$ , (3.8) turns into

$$\begin{aligned} \mathcal{F}^\lambda(\tilde{F}) - \mathcal{F}^\lambda(F) &\leq -c(n, \lambda) m_2^{\frac{n-1}{n}} + \gamma K \mathcal{F}^\lambda(F) \\ &\leq -c(n, \lambda) m_2^{\frac{n-1}{n}} + C(n, \lambda, g, G, q) \max \left\{ m_2, \varepsilon^{\frac{1}{n}} m_2^{\frac{n-1}{n}} \right\}. \end{aligned}$$

Since  $m_2 \leq c(n, \lambda) \varepsilon$  by (3.4), for  $\varepsilon$  sufficiently small (3.1) follows with  $L = \tilde{F}$ .  $\square$

Now we are ready to prove Theorem 3.1.

*Proof of Theorem 3.1.* By Lemma 3.5 there exists a minimizing sequence with uniformly bounded sets. The lower semicontinuity of  $P_\lambda$  [PP24, Lemma 3.9] and the continuity of  $\mathcal{R}$  and  $\mathcal{G}$  under strong  $L^1$  convergence allow to conclude the proof.  $\square$

**3.2. Boundedness and indecomposability of minimizers.** In this section we will prove two qualitative properties of volume constrained minimizers of  $\mathcal{F}^\lambda$ , namely boundedness and indecomposability. We begin with the following

**Theorem 3.6.** *Let  $g$  be  $\mathcal{R}$ -admissible, infinitesimal and symmetric and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer of  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > 0$ . Then  $E$  is essentially bounded.*

*Remark 3.7.* We remark that Theorem 3.6 proves boundedness of minimizers without requiring decreasing properties of the Riesz-type kernel, but only infinitesimality and symmetry.

Before giving the proof, we recall the definition and some properties of the so-called  $(K, r_0)$ -quasiminimal sets.

**Definition 3.8.** Let  $E \subset \mathbb{R}^n \setminus H$  be a set of finite perimeter with finite measure, and let  $K \geq 1$ ,  $r_0 > 0$ . We say that  $E$  is a  $(K, r_0)$ -quasiminimal set (relatively in  $\mathbb{R}^n \setminus H$ ) if

$$P(E, \mathbb{R}^n \setminus H) \leq K P(F, \mathbb{R}^n \setminus H),$$

for any  $F \subset \mathbb{R}^n \setminus H$  such that  $E \Delta F \subset \subset B_r(x)$ , for some ball  $B_r(x) \subset \mathbb{R}^n$  with  $r \leq r_0$  and  $x \in \{x_n \geq 0\}$ .

**Theorem 3.9.** *Let  $E \subset \mathbb{R}^n \setminus H$  be a  $(K, r_0)$ -quasiminimal set, for some  $K \geq 1$ ,  $r_0 > 0$ . Then there exist  $c = c(n, K, r_0) \in \left(0, \frac{1}{2}\right]$  and  $r'_0 = r'_0(n, K, r_0) \in (0, r_0]$  such that*

$$c \leq \frac{|E \cap B_r(x)|}{|B_r(x) \setminus H|} \leq 1 - c \quad \forall x \in \overline{\partial E \setminus H}, \forall r \in (0, r'_0].$$

*In particular the set  $E^{(1)}$  of points of density 1 for  $E$  is an open representative for  $E$ .*

The proof of Theorem 3.9 follows, for instance, by repeatedly applying [Kin+13, Theorem 4.2] with  $X = \{x_n \geq 0\}$  in domains  $\Omega = X \cap B_{r_0}(x)$  for  $x \in X$ , in the notation of [Kin+13, Theorem 4.2]. Observe also that in [Kin+13], the perimeter functional coincides with the relative perimeter in  $\mathbb{R}^n \setminus H$ , hence the definition of quasiminimal set in [Kin+13, Definition 3.1] coincides with Definition 3.8. Alternatively, Theorem 3.9 follows also by adapting

the classical argument in the proof of [Mag12, Theorem 21.11] working with  $(K, r_0)$ -quasiminimal sets instead of  $(\Lambda, r_0)$ -minimizers.

The aim of the following lemmas is to prove that minimizers of  $\mathcal{F}^\lambda$  are  $(K, r_0)$ -quasiminimal sets, in order to apply Theorem 3.9.

**Lemma 3.10.** *Let  $g$  be  $\mathcal{R}$ -admissible, infinitesimal and symmetric and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer of  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > 0$ . Then there exists  $\bar{x}_n > 0$ , depending on  $n$ ,  $g$ ,  $G$ ,  $E$ , such that*

$$(3.10) \quad |E \cap \{x_n > \bar{x}_n\}| = 0.$$

*Proof.* Let us define, for every  $t > 0$ ,

$$E_t := E \cap \{x_n \leq t\}, \quad V(t) := |E \cap \{x_n > t\}|.$$

Fix  $x_0 \in \partial^* E$  such that  $x_0 \in \partial^* E \cap \{0 < x_n < t\}$  and  $r_0 > 0$  such that  $B_{r_0}(x_0) \subset \subset \{0 < x_n < t\}$  for any  $t$  large enough. By [Mag12, Lemma 17.21] there exist  $\sigma_0, c_0 \in (0, \infty)$ , depending on  $E$ ,  $x_0$  and  $r_0$ , such that for every  $\sigma \in (-\sigma_0, \sigma_0)$  we can find a set of finite perimeter  $F$ , given by a suitable local variation of  $E$ , such that

$$(3.11) \quad F \Delta E \subset \subset B_{r_0}(x_0) \quad |F| = |E| + \sigma, \quad |P(F, B_{r_0}(x_0)) - P(E, B_{r_0}(x_0))| \leq c_0 |\sigma|.$$

Now consider  $t_0 = t_0(E) > 0$  large enough such that  $V(t_0) < \sigma_0$ , and set  $\sigma = V(t)$  for  $t > t_0$ . Then there exists  $\tilde{F}$  such that (3.11) holds. Define also  $\tilde{E}_t := \tilde{F} \cap \{0 < x_n \leq t\}$ , so that

$$|\tilde{E}_t| = |\tilde{F}| - |\tilde{F} \cap \{x_n > t\}| = |\tilde{F}| - V(t) = |E| + \sigma - \sigma = |E|.$$

Moreover by [Mag12, Lemma 17.9, Lemma 17.21] and properties of local variations, we get

$$\begin{aligned} |\tilde{E}_t \Delta E_t| &= |\tilde{F} \Delta E| \leq c(E) \|\tilde{F} - E\| = c(E)V(t), \\ |P(\tilde{E}_t, B_{r_0}(x_0)) - P(E, B_{r_0}(x_0))| &\leq c(E)V(t). \end{aligned}$$

By the minimality of  $E$

$$P_\lambda(E) + \mathcal{R}(E) + \mathcal{G}(E) \leq P_\lambda(\tilde{E}_t) + \mathcal{R}(\tilde{E}_t) + \mathcal{G}(\tilde{E}_t).$$

Since  $\mathcal{H}^{n-1}(\partial^* E \cap \partial H) = \mathcal{H}^{n-1}(\partial^* \tilde{E}_t \cap \partial H)$ , we get

$$\begin{aligned} P(E, \mathbb{R}^n \setminus H) + \mathcal{R}(E) + \mathcal{G}(E) &\leq P(\tilde{E}_t, \mathbb{R}^n \setminus H) + \mathcal{R}(\tilde{E}_t) + \mathcal{G}(\tilde{E}_t) \\ &\leq P(E_t, \mathbb{R}^n \setminus (H \cup B_{r_0}(x_0))) + P(E, B_{r_0}(x_0)) + c(E)V(t) + \mathcal{R}(\tilde{E}_t) + \mathcal{G}(\tilde{E}_t) \\ &= P(E, \{x_n < t\}) + |V'(t)| + \mathcal{R}(\tilde{E}_t) + \mathcal{G}(\tilde{E}_t) + c(E)V(t) \\ &= P(E, \mathbb{R}^n \setminus H) - P(E, \{x_n > t\}) + |V'(t)| + \mathcal{R}(\tilde{E}_t) + \mathcal{G}(\tilde{E}_t) + c(E)V(t). \end{aligned}$$

Then

$$P(E, \{x_n > t\}) \leq |V'(t)| + \mathcal{R}(\tilde{E}_t) - \mathcal{R}(E) + \mathcal{G}(\tilde{E}_t) - \mathcal{G}(E) + c(E)V(t)$$

By Fubini theorem and symmetry of  $g$

$$\begin{aligned} \mathcal{R}(\tilde{E}_t) - \mathcal{R}(E) &= \int_{\tilde{E}_t \setminus E} \int_{\tilde{E}_t} g(y-x) dy dx + \int_{\tilde{E}_t \cap E} \int_{\tilde{E}_t \setminus E} g(y-x) dy dx \\ &\quad - \int_{E \setminus \tilde{E}_t} \int_E g(y-x) dy dx - \int_{E \cap \tilde{E}_t} \int_{E \setminus \tilde{E}_t} g(y-x) dy dx \\ &= \int_{\tilde{E}_t \setminus E} \int_{\tilde{E}_t} g(y-x) dy dx + \int_{\tilde{E}_t \setminus E} \int_{\tilde{E}_t \cap E} g(x-y) dy dx \\ &\quad - \int_{E \setminus \tilde{E}_t} \int_E g(y-x) dy dx - \int_{E \setminus \tilde{E}_t} \int_{E \cap \tilde{E}_t} g(x-y) dy dx \\ &= \int_{\tilde{E}_t \setminus E} \int_{\tilde{E}_t} g(y-x) dy dx + \int_{\tilde{E}_t \setminus E} \int_{\tilde{E}_t \cap E} g(y-x) dy dx \\ &\quad - \int_{E \setminus \tilde{E}_t} \int_E g(y-x) dy dx - \int_{E \setminus \tilde{E}_t} \int_{E \cap \tilde{E}_t} g(y-x) dy dx \\ &\leq \int_{\tilde{E}_t \Delta E} \left( \int_{\tilde{E}_t} g(y-x) dy + \int_E g(y-x) dy \right) dx \end{aligned}$$

Since  $g$  is infinitesimal there exists  $R_g > 0$  such that

$$g(x) < 1 \quad \forall x : |x| > R_g.$$

Then

$$\begin{aligned} \mathcal{R}(\tilde{E}_t) - \mathcal{R}(E) &\leq \int_{\tilde{E}_t \Delta E} \left( 2 \int_{B_{R_g}(0)} g(z) dz + |\tilde{E}_t \setminus B_{R_g}(0)| + |E \setminus B_{R_g}(0)| \right) dx \\ &\leq 2 \int_{\tilde{E}_t \Delta E} \left( \int_{B_{R_g}(0)} g(z) dz + |E| \right) dx \\ &\leq c(g, m)(|\tilde{E}_t \Delta E_t| + |E_t \Delta E|) \leq c(g, E)V(t). \end{aligned}$$

By Remark 2.7  $(\sup_{B_{r_0}(x_0)} G) < \infty$  and

$$\mathcal{G}(\tilde{E}_t) - \mathcal{G}(E) = \int_{\tilde{E}_t \setminus E} G dx - \int_{E \setminus \tilde{E}_t} G dx \leq \int_{\tilde{E}_t \setminus E} G dx \leq \left( \sup_{B_{r_0}(x_0)} G \right) |\tilde{E}_t \setminus E|.$$

Therefore, for almost every  $t$  sufficiently large,

$$(3.12) \quad P(E, \{x_n > t\}) \leq |V'(t)| + c(n, g, G, E)V(t).$$

Finally, if  $c_{iso} = c_{iso}(n)$  is the constant in the classical isoperimetric inequality and  $t$  is large enough, (3.12) yields

$$\begin{aligned} c_{iso} V(t)^{\frac{n-1}{n}} = c_{iso} |E \setminus E_t|^{\frac{n-1}{n}} &\leq P(E \setminus E_t) = \mathcal{H}^{n-1}(\partial^* E_t \cap \{x_n = t\}) + P(E, \{x_n > t\}) \\ &\leq 2|V'(t)| + c(n, g, G, E)V(t) < 2|V'(t)| + \frac{c_{iso}}{2} V(t)^{\frac{n-1}{n}} \end{aligned}$$

and

$$-V'(t) \geq cV(t)^{\frac{n-1}{n}}.$$

Therefore ODE comparison implies that  $V(t)$  vanishes at some  $t = \bar{x}_n < +\infty$ .  $\square$

**Lemma 3.11.** *Let  $g$  be  $\mathcal{R}$ -admissible, infinitesimal and symmetric and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer of  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > 0$ . Then  $E$  is a  $(K, r_0)$ -quasiminimal set, for suitable  $K \geq 1$  and  $r_0 > 0$ , depending on  $n, \lambda, g, G, E$ .*

*Proof.* Let us consider  $x_1, x_2 \in \partial^* E \setminus H$  and  $t_0 > 0$  such that we have  $B_{t_0}(x_1) \cap B_{t_0}(x_2) = \emptyset$  and  $B_{t_0}(x_1) \cup B_{t_0}(x_2) \subset \subset \mathbb{R}^n \setminus H$ . By applying [Mag12, Lemma 17.21] we find two positive constants  $\sigma_0$  and  $c_0$ , depending on  $E$ , such that, given  $|\sigma| < \sigma_0$ , there exist two sets of finite perimeter  $F_1$  and  $F_2$  with

$$(3.13) \quad E \Delta F_k \subset \subset B_{t_0}(x_k), \quad |F_k| = |E| + \sigma, \quad \left| P(E, B_{t_0}(x_k)) - P(F_k, B_{t_0}(x_k)) \right| \leq c_0 |\sigma|, \quad k \in \{1, 2\}.$$

Let  $r_0 = r_0(n, \lambda, g, G, E) > 0$  to be determined later. At the moment assume that

$$r_0 < \min \left\{ \frac{t_0}{2}, \frac{\sigma_0^{\frac{1}{n}}}{\omega_n}, \frac{|x_1 - x_2| - 2t_0}{2} \right\}.$$

In particular, if a ball of radius  $r_0$  intersects  $B_{t_0}(x_1)$  (resp.  $B_{t_0}(x_2)$ ), then it is disjoint from  $B_{t_0}(x_2)$  (resp. from  $B_{t_0}(x_1)$ ). Let  $F$  be such that  $E \Delta F \subset \subset B_r(x) \cap (\mathbb{R}^n \setminus H)$ , where  $r < r_0$ . Then, by the definition of  $r_0$ ,

$$\||E| - |F|\| \leq |E \Delta F| \leq \omega_n r^n < \omega_n r_0^n \leq \sigma_0$$

and we can compensate for the volume deficit  $\||E| - |F|\|$  between  $E$  and  $F$  by modifying  $F$  inside either  $B_{t_0}(x_1)$  or  $B_{t_0}(x_2)$ . Precisely, by the definition of  $r_0$ , we may assume without loss of generality that  $B_r(x)$  does not intersect  $B_{t_0}(x_1)$ , set  $\sigma = |E| - |F|$ , and consider  $F_1$  verifying (3.13), so that

$$(3.14) \quad E \Delta F_1 \subset \subset B_{t_0}(x_1), \quad E \Delta F \subset \subset B_r(x) \cap (\mathbb{R}^n \setminus H) \subset \subset \mathbb{R}^n \setminus \left( H \cup \overline{B_{t_0}(x_1)} \right).$$

By (3.13)  $\sigma = |F_1| - |E|$  and, if we define

$$\tilde{F} = (F \cap B_r(x)) \cup (F_1 \cap B_{t_0}(x_1)) \cup (E \setminus (B_r(x) \cup B_{t_0}(x_1))),$$

then  $|\tilde{F}| = |E|$  and  $\tilde{F} \Delta E \subset \subset \{x_n > 0\}$ . By the minimality of  $E$

$$(3.15) \quad (1 - |\lambda|)P(E, \mathbb{R}^n \setminus H) \leq P_\lambda(E) \leq P_\lambda(\tilde{F}) + \mathcal{R}(\tilde{F}) - \mathcal{R}(E) + \mathcal{G}(\tilde{F}) - \mathcal{G}(E).$$

By (3.13) and (3.14) we get

$$\begin{aligned}
(3.16) \quad P_\lambda(\tilde{F}) &\leq (1 + |\lambda|)P(\tilde{F}, \mathbb{R}^n \setminus H) \\
&\leq (1 + |\lambda|) \left[ P(\tilde{F}, \mathbb{R}^n \setminus (H \cup \overline{B_{t_0}(x_1)})) + P(\tilde{F}, B_{t_0}(x_1)) + P(\tilde{F}, \partial B_{t_0}(x_1)) \right] \\
&= (1 + |\lambda|) \left[ P(F, \mathbb{R}^n \setminus (H \cup \overline{B_{t_0}(x_1)})) + P(F_1, B_{t_0}(x_1)) + P(F, \partial B_{t_0}(x_1)) \right] \\
&\leq (1 + |\lambda|) \left[ P(F, \mathbb{R}^n \setminus (H \cup B_{t_0}(x_1))) + P(E, B_{t_0}(x_1)) + c_0(E)|\sigma| \right] \\
&\leq (1 + |\lambda|) P(F, \mathbb{R}^n \setminus H) + c_0(\lambda, E)|F\Delta E|.
\end{aligned}$$

As in the proof of Lemma 3.10 one estimates

$$(3.17) \quad \mathcal{R}(\tilde{F}) - \mathcal{R}(E) \leq c(g, E)|F\Delta E|,$$

and by Remark 2.7, (3.10) and if  $t_0 < 1$

$$\begin{aligned}
(3.18) \quad \mathcal{G}(\tilde{F}) - \mathcal{G}(E) &\leq \left( \sup_{\tilde{F} \setminus E} G \right) |\tilde{F} \setminus E| \leq \left( \sup_{\tilde{F} \setminus E} G \right) (|\tilde{F}\Delta F| + |F\Delta E|) \\
&\leq c(G, E)(\bar{x}_n + 1)^n |F\Delta E| = c(n, g, G, E)|F\Delta E|.
\end{aligned}$$

However, by the relative isoperimetric inequality [CGR07; FM23] and (3.14)

$$\begin{aligned}
(3.19) \quad |F\Delta E| &= |F\Delta E|^{\frac{1}{n}} |F\Delta E|^{\frac{n-1}{n}} \leq c(n) |F\Delta E|^{\frac{1}{n}} P(F\Delta E, \mathbb{R}^n \setminus H) \\
&\leq c(n) |F\Delta E|^{\frac{1}{n}} (P(F, \mathbb{R}^n \setminus H) + P(E, \mathbb{R}^n \setminus H)) \\
&\leq c(n) r_0 (P(F, \mathbb{R}^n \setminus H) + P(E, \mathbb{R}^n \setminus H)).
\end{aligned}$$

Putting together (3.15)-(3.19) we obtain

$$((1 - |\lambda|) - c(n, \lambda, g, G, E) r_0) P(E, \mathbb{R}^n \setminus H) \leq ((1 + |\lambda|) + c(n, \lambda, g, G, E) r_0) P(F, \mathbb{R}^n \setminus H)$$

If  $r_0$  is sufficiently small, we conclude the proof.  $\square$

By Theorem 3.9 and Lemma 3.11, from now on we can identify any minimizer  $E$  of  $\mathcal{F}^\lambda$ , with  $|E| = m$ ,  $m > 0$ ,  $g$   $\mathcal{R}$ -admissible infinitesimal symmetric function and  $G$   $\mathcal{G}$ -admissible function, with the open set  $E^{(1)}$  of points of density 1 for  $E$ .

Now we are ready to prove Theorem 3.6.

*Proof of Theorem 3.6.* By Lemma 3.11 and Theorem 3.9 there exist  $r > 0$  and  $c > 0$  such that for every  $x \in \partial E \setminus H$  we have  $|E \cap B_r(x)| \geq cr^n$ . If  $E$  were not bounded, one would easily get  $|E| = \infty$ .  $\square$

Now we prove indecomposability of minimizers.

**Theorem 3.12.** *Let  $g$  be  $\mathcal{R}$ -admissible and infinitesimal and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer of  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > 0$ . Then  $E$  is indecomposable.*

*Proof.* We argue by contradiction. Assume that there exist two sets of finite perimeter  $E_1$  and  $E_2$  such that  $|E_1 \cap E_2| = 0$ ,  $E = E_1 \cup E_2$  and  $P(E) = P(E_1) + P(E_2)$ . If  $R > 0$  is sufficiently large, letting  $e_1 = (1, 0, \dots, 0)$  and defining  $E_R := E_1 \cup (E_2 + e_1 R)$ , we have  $|E_R| = m$ ,  $P_\lambda(E_R) = P_\lambda(E)$  and  $\mathcal{G}(E_R) = \mathcal{G}(E)$ . At the same time, the nonlocal energy decreases, precisely

$$\liminf_{R \rightarrow \infty} \left( \int_{E_1} \int_{E_2 + e_1 R} g(y-x) dy dx + \int_{E_2 + e_1 R} \int_{E_1} g(y-x) dy dx \right) = 0,$$

and

$$\begin{aligned}
\liminf_{R \rightarrow \infty} \mathcal{F}^\lambda(E_R) &= P_\lambda(E) + \mathcal{R}(E_1) + \mathcal{R}(E_2) + \mathcal{G}(E) \\
&< P_\lambda(E) + \mathcal{R}(E_1) + \mathcal{R}(E_2) + \int_{E_1} \int_{E_2} g(y-x) dy dx + \int_{E_2} \int_{E_1} g(y-x) dy dx + \mathcal{G}(E) \\
&= \mathcal{F}^\lambda(E).
\end{aligned}$$

Therefore, if  $R$  is sufficiently large, we obtain  $\mathcal{F}^\lambda(E_R) < \mathcal{F}^\lambda(E)$ , in contradiction with the minimizing property of  $E$ .  $\square$

## 4. NONEXISTENCE OF MINIMIZERS FOR LARGE MASSES

The goal of this Section is to prove Theorem 1.2. We begin by proving some preparatory lemmas. Let us start with a non-optimality criterion.

**Lemma 4.1.** *Let  $g$  be  $\mathcal{R}$ -admissible,  $q$ -decreasing and infinitesimal and let  $G$  be  $\mathcal{G}$ -admissible. There exists  $\varepsilon > 0$ , depending on  $n, \lambda, g, G$  and  $q$ , such that the following holds. Let  $F \subset \mathbb{R}^n \setminus H$  be a set of finite perimeter and assume there exist two sets of finite perimeter  $F_1, F_2 \subset F$  such that  $|F_1|, |F_2| > 0, |F_1 \cap F_2| = 0, |F \setminus (F_1 \cup F_2)| = 0$  and*

$$(4.1) \quad \Sigma := P_\lambda(F_1) + P_\lambda(F_2) - P_\lambda(F) \leq \frac{1}{2} \mathcal{F}^\lambda(F_2).$$

Then, if

$$(4.2) \quad |F_2| \leq \varepsilon \min\{1, |F_1|\},$$

there exists a set  $G \subset \mathbb{R}^n \setminus H$  with  $|G| = |F|$  and  $\mathcal{F}^\lambda(G) < \mathcal{F}^\lambda(F)$ .

*Proof.* Let us denote  $m := |F|, m_1 := |F_1|, m_2 := |F_2|$  and  $\gamma := \frac{m_2}{m_1} \leq \varepsilon$ . Let us define the sets  $\tilde{F}$  and  $\hat{F}$  in the following way:  $\tilde{F}$  is given by  $\tilde{F} = lF_1$ , with  $l := \sqrt[n]{1 + \gamma}$ , so that  $|\tilde{F}| = |F|$ , and  $\hat{F}$  is given by a collection of  $N \geq 1$  spherical caps  $\{B^\lambda(v, x_i)\}_{1 \leq i \leq N}$  of equal volume  $v$  and with centers located at  $x_i = iRe_1, i = 1, \dots, N$ , with  $R$  large enough so that the  $B^\lambda(v, x_i), 1 \leq i \leq N$  are pairwise disjoint. The number  $N$  is the smallest integer for which the volume of each spherical cap does not exceed  $\min\{1, |B^\lambda|\}$ . Hence  $Nv = m$  and  $N = \left\lceil \frac{m}{\min\{1, |B^\lambda|\}} \right\rceil$ . If there exists  $R > 0$  such that, for the corresponding  $\hat{F}$ , one has  $\mathcal{F}^\lambda(\hat{F}) < \mathcal{F}^\lambda(F)$ , then the proof is concluded with  $G = \hat{F}$ . So we can assume that for any  $R > 0$  there holds  $\mathcal{F}^\lambda(\hat{F}) \geq \mathcal{F}^\lambda(F)$ . Hence for  $R$  large enough we claim that

$$(4.3) \quad \mathcal{F}^\lambda(F) \leq \mathcal{F}^\lambda(\hat{F}) \leq c(n, \lambda, g, G) \max\left\{m, m^{\frac{n-1}{n}}\right\}.$$

Indeed, if  $m \geq 1$ , estimate (4.3) follows by the same computations done in the proof of Corollary 3.3. If instead  $m < 1$ , by [PP24, Lemma 3.3] we find

$$\begin{aligned} P_\lambda(E) &= P_\lambda\left(\bigcup_{i=1}^N B^\lambda(v, x_i)\right) = \sum_{i=1}^N P_\lambda(B^\lambda(v, x_i)) = c(n, \lambda) N v^{\frac{n-1}{n}} \leq c(n, \lambda) \left(\frac{m}{\min\{1, |B^\lambda|\}} + 1\right) v^{\frac{n-1}{n}} \\ &\leq c(n, \lambda) (m v^{\frac{n-1}{n}} + v^{\frac{n-1}{n}}) \leq c(n, \lambda) (m + m^{\frac{n-1}{n}}) = c(n, \lambda) m^{\frac{n-1}{n}}. \end{aligned}$$

Arguing as in Corollary 3.3, if  $R$  is so large that  $g(x - y) < \frac{1}{N}$  for every  $x \in B^\lambda(v, x_j), y \in B^\lambda(v, x_k)$  with  $j \neq k$ , then

$$\mathcal{R}(E) \leq c(n, \lambda, g, G) m \leq c(n, \lambda, g, G) m^{\frac{n-1}{n}}$$

and

$$\mathcal{F}^\lambda(E) = P_\lambda(E) + \mathcal{R}(E) + \mathcal{G}(E) \leq c(n, \lambda) |E|^{\frac{n-1}{n}} + c(n, \lambda, g, G) |E|^{\frac{n-1}{n}} + \sum_{i=1}^N \mathcal{G}(B^\lambda(v, x_i)) \leq c(n, \lambda, g, G) m^{\frac{n-1}{n}},$$

therefore (4.3) holds.

We want to show that if  $\varepsilon$  sufficiently small, then  $\mathcal{F}^\lambda(\tilde{F}) < \mathcal{F}^\lambda(F)$ , implying the claim with  $G = \tilde{F}$ . By Lemma 3.4

$$(4.4) \quad \mathcal{F}^\lambda(\tilde{F}) = \mathcal{F}^\lambda(lF_1) \leq l^{2n+q} \mathcal{F}^\lambda(F_1) = \mathcal{F}^\lambda(F_1) + (l^{2n+q} - 1) \mathcal{F}^\lambda(F_1).$$

Choosing  $\varepsilon \leq 1$ , we have  $1 \leq l \leq 2^{\frac{1}{n}}$ , and by Taylor's formula we obtain  $l^{2n+q} - 1 = (1 + \gamma)^{2+q/n} - 1 \leq \gamma K$  for some  $K > 0$  independent of  $\gamma$ , for  $\varepsilon$  sufficiently small. By (4.4) we arrive at

$$\mathcal{F}^\lambda(\tilde{F}) - \mathcal{F}^\lambda(F_1) \leq \gamma K \mathcal{F}^\lambda(F_1).$$

By the definition of  $\Sigma$  and since  $\mathcal{R}(F_1) + \mathcal{R}(F_2) \leq \mathcal{R}(F)$

$$(4.5) \quad \begin{aligned} \mathcal{F}^\lambda(\tilde{F}) - \mathcal{F}^\lambda(F) &\leq \mathcal{R}(F_1) + \mathcal{G}(F_1) + \mathcal{R}(F_2) + \mathcal{G}(F_2) - \mathcal{R}(F) - \mathcal{G}(F) + \Sigma - \mathcal{F}^\lambda(F_2) + \gamma K \mathcal{F}^\lambda(F_1) \\ &\leq -\frac{1}{2} \mathcal{F}^\lambda(F_2) + \gamma K \mathcal{F}^\lambda(F_1). \end{aligned}$$

By positivity of  $\mathcal{R}$  and  $\mathcal{G}$  and the isoperimetric inequality, we have  $\mathcal{F}^\lambda(F_2) > P_\lambda(F_2) \geq c(n, \lambda)m_2^{\frac{n-1}{n}}$ . As in (3.9) we obtain  $\mathcal{F}^\lambda(F_1) \leq \mathcal{F}^\lambda(F)$ . By (4.3), since  $\gamma m \leq 2m_2$  and  $\gamma \leq \varepsilon$ , (4.5) turns into

$$\mathcal{F}^\lambda(\tilde{F}) - \mathcal{F}^\lambda(F) \leq -c(n, \lambda)m_2^{\frac{n-1}{n}} + \gamma K \mathcal{F}^\lambda(F) \leq -c(n, \lambda)m_2^{\frac{n-1}{n}} + C(n, \lambda, g, G, q) \max \left\{ m_2, \varepsilon^{\frac{1}{n}} m_2^{\frac{n-1}{n}} \right\}.$$

Since  $m_2 \leq \varepsilon$  by (4.2), for  $\varepsilon$  sufficiently small the assertion of the lemma holds with  $G = \tilde{F}$ .  $\square$

Next lemma is an improvement of the standard density estimate for quasiminimizers.

**Lemma 4.2.** *Let  $g$  be  $\mathcal{R}$ -admissible,  $q$ -decreasing and infinitesimal, and let  $G$  be  $\mathcal{G}$ -admissible. Then there exists  $c = c(n, \lambda, g, G, q) > 0$  such that the following holds. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer of  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > 0$ . Then for almost every  $x \in E$  there holds*

$$|E \cap B_1(x)| \geq c \min\{1, m\}.$$

*Proof.* For  $r > 0$  and  $x \in E$ , let  $F_1^r := E \setminus B_r(x)$  and  $F_2^r := E \cap B_r(x)$ . Note that  $|F_1^r| + |F_2^r| = m$  and  $|F_2^r| \leq \omega_n r^n$ . Then there exists  $C > 0$ , depending on  $n, \lambda, g, G$  and  $q$ , such that (4.2) holds for all  $r \leq r_1 := C \min \left\{ 1, \sqrt[n]{m} \right\}$ . Note that we can choose  $C \leq 1$ . Since  $E$  is a minimizer, Lemma 4.1 implies that (4.1) cannot be satisfied for any  $r \leq r_1$ . Equivalently, recalling also [PP24, Corollary 2.4], for all  $r \leq r_1$  we have

$$(4.6) \quad \Sigma^r := P_\lambda(F_1^r) + P_\lambda(F_2^r) - P_\lambda(E) > \frac{1}{2} \mathcal{F}^\lambda(F_2^r) > \frac{1}{2} P_\lambda(F_2^r) \geq \frac{1-\lambda}{4} P(F_2^r).$$

At the same time, for almost every  $r$  we have

$$\Sigma^r = 2\mathcal{H}^{n-1}(E^{(1)} \cap \partial B_r(x)).$$

By (4.6), for a constant  $c(n, \lambda) \in \left(0, \frac{1}{2}\right)$  there holds

$$(4.7) \quad 2\mathcal{H}^{n-1}(E^{(1)} \cap \partial B_r(x)) > c(n, \lambda) (\mathcal{H}^{n-1}(\partial^* E \cap B_r(x)) + \mathcal{H}^{n-1}(E^{(1)} \cap \partial B_r(x))),$$

for almost every  $r$ . Let us now distinguish two cases. If there exists  $r_2 \in \left(\frac{r_1}{2}, r_1\right)$  such that  $|E \cap B_{r_2}| \geq \frac{1}{2} \omega_n r_2^n$ , then by the choice of  $r_1$  we get

$$|E \cap B_1| \geq |E \cap B_{r_2}| \geq c(n) \left(\frac{r_1}{2}\right)^n = c(n, \lambda, g, G, q) \min\{1, m\},$$

and the proof is concluded.

Let us assume that  $|E \cap B_r| < \frac{1}{2} \omega_n r^n$  for all  $r \in \left(\frac{r_1}{2}, r_1\right)$ . Then we rearrange terms in (4.7) and apply the relative isoperimetric inequality [Mag12, Proposition 12.37] to the right-hand side to obtain

$$\mathcal{H}^{n-1}(E^{(1)} \cap \partial B_r(x)) \geq c(n, \lambda) |E \cap B_r(x)|^{\frac{n-1}{n}}.$$

Let us denote  $U(r) := |E \cap B_r(x)|$ . Then  $\frac{dU(r)}{dr} = \mathcal{H}^{n-1}(E^{(1)} \cap \partial B_r(x))$  for all  $r \in \left(\frac{r_1}{2}, r_1\right)$  and

$$\frac{dU(r)}{dr} \geq c(n, \lambda) U^{\frac{n-1}{n}}(r) \quad \forall r \in \left(\frac{r_1}{2}, r_1\right).$$

For  $\mathcal{H}^n$ -a.e.  $x \in E$ , we have  $U(r) > 0$  for all  $r > 0$  and ODE comparison in  $r \in \left(\frac{r_1}{2}, r_1\right)$  implies that

$$U^{1/n}(r) \geq U^{1/n}\left(\frac{r_1}{2}\right) + c(n, \lambda) \left(r - \frac{r_1}{2}\right) \geq c(n, \lambda) \left(r - \frac{r_1}{2}\right) \quad \forall r \in \left(\frac{r_1}{2}, r_1\right).$$

Then the lemma follows as

$$c(n, \lambda, g, G) \min\{1, m\} = c(n, \lambda) r_1^n \leq U(r_1) = |E \cap B_{r_1}(x)| = \left| E \cap B_{C \min\{1, \sqrt[n]{m}\}}(x) \right| \leq |E \cap B_1(x)|.$$

$\square$

*Remark 4.3.* We remark that the density estimate in Lemma 4.2 is more precise than the one provided in Section 3.2. Indeed, in Lemma 3.11  $K$  and  $r_0$  depend on the minimizer, and consequently  $c$  in Theorem 3.9 also inherits this dependence. At the same time, Lemma 4.2 requires  $g$  to be  $q$ -decreasing, which is not required in Lemma 3.11.

The following lemma will imply Theorem 1.2 for  $\beta \in (0, 1)$ .

**Lemma 4.4.** *Let*

$$g(x) = \frac{1}{|x|^\beta}, \quad 0 < \beta < n, x \in \mathbb{R}^n \setminus \{0\},$$

and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E$  be a minimizer of  $\mathcal{F}^\lambda$  with  $|E| = m$  and  $m \geq 1$ . Then

$$(4.8) \quad cm^{\frac{1}{\beta}} \leq \text{diam } E \leq Cm,$$

for some  $C, c > 0$  depending only on  $n, \beta, \lambda, G$ .

*Proof.* By Theorem 3.6 and Theorem 3.12 we know that  $E$  is essentially bounded and indecomposable. In particular  $d := \text{diam } E < \infty$ . By Corollary 3.3 we get the existence of  $c(n, \lambda, \beta, G) > 0$  such that

$$\frac{m^2}{d^\beta} \leq \int_E \int_E \frac{1}{|x-y|^\beta} dy dx = \mathcal{R}(E) \leq \mathcal{F}^\lambda(E) \leq cm,$$

which implies the first bound in (4.8).

In order to prove the upper bound in (4.8), we may clearly assume that  $\frac{d}{\sqrt{2}} > 3$ . Recalling that we identify  $E$  with the bounded open set  $E^{(1)}$ , we let  $x^{(1)}, x^{(2)} \in \bar{E}$  such that

$$|x^{(1)} - x^{(2)}| = d.$$

Up to a rotation with respect to an axis orthogonal to  $\{x_n = 0\}$ , we can write

$$x^{(2)} - x^{(1)} = \langle x^{(2)} - x^{(1)}, e_1 \rangle e_1 + \langle x^{(2)} - x^{(1)}, e_n \rangle e_n.$$

In particular,

$$\max \left\{ \left| \langle x^{(2)} - x^{(1)}, e_1 \rangle \right|, \left| \langle x^{(2)} - x^{(1)}, e_n \rangle \right| \right\} \geq \frac{d}{\sqrt{2}}.$$

Assume for simplicity that

$$\left| \langle x^{(2)} - x^{(1)}, e_n \rangle \right| \geq \frac{d}{\sqrt{2}},$$

the remaining case being analogous. Up to relabeling, assume also that

$$\langle x^{(2)}, e_n \rangle > \langle x^{(1)}, e_n \rangle$$

Let  $N$  be the largest integer smaller than  $\frac{d}{3\sqrt{2}}$ , i.e.  $N := \left\lfloor \frac{d}{3\sqrt{2}} \right\rfloor$ . Since  $E$  is indecomposable, for every  $j = 1, \dots, N$  there holds

$$\left| E \cap \{3j-1 + \langle x^{(1)}, e_n \rangle < x_n < 3j + \langle x^{(1)}, e_n \rangle\} \right| > 0.$$

For every  $j = 1, \dots, N$ , let

$$x_j \in E \cap \{3j-1 + \langle x^{(1)}, e_n \rangle < x_n < 3j + \langle x^{(1)}, e_n \rangle\}.$$

The balls  $B_1(x_j)$ ,  $j = 1, \dots, N$ , are pairwise disjoint and, for a suitable choice of  $x_j$ , we can apply Lemma 4.2 to get

$$m = |E| \geq \sum_{j=1}^N |B_1(x_j) \cap E| \geq c(n, \lambda, \beta, G) N \geq c(n, \lambda, \beta, G) d.$$

□

The following lemma will imply Theorem 1.2 for  $\beta = 1$ .

**Lemma 4.5.** *Let*

$$g(x) = \frac{1}{|x|^\beta}, \quad \beta \in (0, n), x \in \mathbb{R}^n \setminus \{0\}$$

and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer for  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > 0$ . Then

$$\int_E \int_E \frac{1}{|x-y|^{\beta-1}} dy dx \leq c(n) m.$$



*Proof.* Let  $v \in \mathbb{S}^{n-1} \setminus \{\pm e_n\}$  and  $t \in \mathbb{R}$ . Denote

$$E_{v,t}^+ := E \cap \{\langle v, x \rangle > t\}$$

$$E_{v,t}^- := E \cap \{\langle v, x \rangle < t\}.$$

Let  $v_1 := \frac{v_{hor}}{|v_{hor}|}$ , where  $v_{hor}$  is the orthogonal projection of  $v$  on  $\{x_n = 0\}$ . For any  $\rho \geq 0$ , the set

$$E_{v,t}^+ \cup (E_{v,t}^- - \rho v_1)$$

has measure  $m$  and, by minimality of  $E$ ,

$$(4.9) \quad \mathcal{F}^\lambda(E_{v,t}^+ \cup (E_{v,t}^- - \rho v_1)) \geq \mathcal{F}^\lambda(E).$$

For any  $\rho > 0$  and for a.e.  $t \in \mathbb{R}$

$$P_\lambda(E_{v,t}^+ \cup (E_{v,t}^- - \rho v_1)) = P_\lambda(E_{v,t}^+) + P_\lambda(E_{v,t}^-) \leq P_\lambda(E) + 2\mathcal{H}^{n-1}(E \cap \{\langle v, x \rangle = t\}).$$

For any  $\rho \geq 0$  we have

$$\begin{aligned} \int_{E_{v,t}^+ \cup (E_{v,t}^- - \rho v_1)} \int_{E_{v,t}^+ \cup (E_{v,t}^- - \rho v_1)} \frac{1}{|x-y|^\beta} dy dx &= \int_{E_{v,t}^+} \int_{E_{v,t}^+} \frac{1}{|x-y|^\beta} dy dx + \int_{E_{v,t}^-} \int_{E_{v,t}^-} \frac{1}{|x-y|^\beta} dy dx \\ &\quad + 2 \int_{E_{v,t}^+} \int_{E_{v,t}^-} \frac{1}{|x-y+\rho v_1|^\beta} dy dx. \end{aligned}$$

Moreover

$$\int_{E_{v,t}^+} \int_{E_{v,t}^-} \frac{1}{|x-y+\rho v_1|^\beta} dy dx \rightarrow 0$$

as  $\rho \rightarrow \infty$ . Hence, by (4.9), letting  $\rho \rightarrow \infty$ , we get

$$\begin{aligned} &P_\lambda(E) + 2\mathcal{H}^{n-1}(E \cap \{\langle v, x \rangle = t\}) + \int_{E_{v,t}^+} \int_{E_{v,t}^+} \frac{1}{|x-y|^\beta} dy dx + \int_{E_{v,t}^-} \int_{E_{v,t}^-} \frac{1}{|x-y|^\beta} dy dx + \mathcal{G}(E_{v,t}^+) + \mathcal{G}(E_{v,t}^-) \\ &\geq P_\lambda(E) + \int_{E_{v,t}^+} \int_{E_{v,t}^+} \frac{1}{|x-y|^\beta} dy dx + \int_{E_{v,t}^-} \int_{E_{v,t}^-} \frac{1}{|x-y|^\beta} dy dx + 2 \int_{E_{v,t}^+} \int_{E_{v,t}^-} \frac{1}{|x-y|^\beta} dy dx + \mathcal{G}(E). \end{aligned}$$

Then

$$\begin{aligned} \mathcal{H}^{n-1}(E \cap \{\langle v, x \rangle = t\}) &\geq \int_{E_{v,t}^+} \int_{E_{v,t}^-} \frac{1}{|x-y|^\beta} dy dx \\ &= \int_E \int_E \chi_{\{\langle v, \cdot \rangle < t\}}(y) \chi_{\{\langle v, \cdot \rangle > t\}}(x) \frac{1}{|x-y|^\beta} dy dx. \end{aligned}$$

Integrating the last inequality with respect to  $t \in \mathbb{R}$ , by Fubini's theorem we get

$$\begin{aligned} m &\geq \int_E \int_E \int_{-\infty}^{+\infty} \chi_{\{\langle v, \cdot \rangle < t\}}(y) \chi_{\{\langle v, \cdot \rangle > t\}}(x) dt \frac{1}{|x-y|^\beta} dy dx = \int_E \int_E \int_{-\infty}^{+\infty} \chi_{\langle \langle v, y \rangle, \langle v, x \rangle \rangle}(t) dt \frac{1}{|x-y|^\beta} dy dx \\ &= \int_E \int_E \frac{\langle v, x-y \rangle_+}{|x-y|^\beta} dy dx. \end{aligned}$$

Further integrating over  $\mathbb{S}^{n-1} \setminus \{\pm e_n\}$ , since

$$\int_{\mathbb{S}^{n-1}} \langle v, x-y \rangle_+ dv = c(n)|x-y|,$$

by symmetry of  $\mathbb{S}^{n-1}$ , we conclude that

$$|\mathbb{S}^{n-1}|m \geq \int_E \int_E \int_{\mathbb{S}^{n-1}} \langle v, x-y \rangle_+ dv \frac{1}{|x-y|^\beta} dy dx = c(n) \int_E \int_E \frac{1}{|x-y|^{\beta-1}} dy dx.$$

□

The following lemma will imply Theorem 1.2 for  $\beta \in (1, 2]$ .

**Lemma 4.6.** *Let*

$$g(x) = \frac{1}{|x|^\beta}, \quad 0 < \beta < n, \quad x \in \mathbb{R}^n \setminus \{0\}$$

and let  $G$  be  $\mathcal{G}$ -admissible. Let  $E \subset \mathbb{R}^n \setminus H$  be a minimizer for  $\mathcal{F}^\lambda$  with  $|E| = m$ ,  $m > \omega_n$ . Then, for  $1 \leq r \leq \frac{\text{diam} E}{2}$ ,

$$|E \cap B_r(x)| \geq c(n, \lambda, \beta, G) r \quad \text{for a.e. } x \in E.$$

*Proof.* Let  $N := \left\lfloor \frac{r-1}{3} \right\rfloor$  and  $x \in E$ . If  $r < 4$ , by Lemma 4.2

$$|E \cap B_r(x)| \geq |E \cap B_1(x)| \geq c(n, \lambda, \beta, G) = \frac{c(n, \lambda, \beta, G)}{4} 4 \geq c(n, \lambda, \beta, G) r.$$

So we can assume that  $r \geq 4$ , in particular  $N \geq 1$ . Since  $E$  is indecomposable by Theorem 3.12, for every  $i = 0, \dots, N-1$  there holds

$$|E \cap (B_{3i+3}(x) \setminus B_{3i+2}(x))| > 0.$$

For every  $i = 0, \dots, N-1$ , let

$$y_i \in E \cap (B_{3i+3}(x) \setminus B_{3i+2}(x)).$$

The balls  $B_1(y_i)$ ,  $i = 0, \dots, N-1$ , are pairwise disjoint and, for a suitable choice of  $y_i$ , by Lemma 4.2 there exists  $c(n, \lambda, \beta, G)$  such that  $|E \cap B_1(y_i)| \geq c$  for  $i = 0, \dots, N-1$ . Finally

$$|E \cap B_r(x)| \geq \sum_{i=0}^{N-1} |E \cap B_1(y_i)| + |E \cap B_1(x)| \geq (N+1)c \geq c \frac{r-1}{3} \geq c r.$$

□

Now we are ready to prove Theorem 1.2.

*Proof of Theorem 1.2.* Lemma 4.4 and Lemma 4.5 easily imply Theorem 1.2 for  $\beta \in (0, 1]$  and mass  $m$  sufficiently large. Then it remains to consider  $\beta \in (1, 2]$ . Let  $E \subset \mathbb{R}^n \setminus H$  be a volume constrained minimizer for  $\mathcal{F}^\lambda$  with  $|E| = m$ . By Lemma 4.4, for  $m$  large enough, we can assume that  $\text{diam } E > 4$ . We observe first that

$$(4.10) \quad \frac{1}{r^{\beta-1}} |\{(x, y) \in E \times E : |x - y| < r\}| = \frac{1}{r^{\beta-1}} \int_E |E \cap B_r(x)| dx \leq \frac{\omega_n r^n}{r^{\beta-1}} |E| \xrightarrow{r \rightarrow 0} 0.$$

Applying the coarea formula on  $\mathbb{R}^n \times \mathbb{R}^n$  for the Lipschitz function  $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  given by  $f(x, y) := |x - y|$ , observing that  $|\nabla f| = \sqrt{2}$  and that

$$\frac{d}{dr} |\{(x, y) \in E \times E : |x - y| < r\}| = \frac{1}{\sqrt{2}} \int_{\{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n : |x - y| = r\}} \chi_{E \times E}(x, y) d\mathcal{H}^{2n-1}(x, y) \quad \text{for a.e. } r > 0,$$

and integrating by parts, we estimate

$$\begin{aligned} \int_E \int_E \frac{1}{|x - y|^{\beta-1}} dy dx &= \frac{1}{\sqrt{2}} \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{\chi_{E \times E}(x, y)}{|x - y|^{\beta-1}} |\nabla f| dx dy \\ &= \frac{1}{\sqrt{2}} \int_0^{+\infty} \int_{\{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n : |x - y| = r\}} \frac{\chi_{E \times E}(x, y)}{r^{\beta-1}} d\mathcal{H}^{2n-1}(x, y) dr \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^{+\infty} \frac{1}{r^{\beta-1}} \frac{d}{dr} |\{(x, y) \in E \times E : |x - y| < r\}| dr \\ &= \lim_{\varepsilon \rightarrow 0^+} -\frac{1}{\varepsilon^{\beta-1}} |\{(x, y) \in E \times E : |x - y| < \varepsilon\}| + \\ &\quad - \int_\varepsilon^{+\infty} (1 - \beta) \frac{1}{r^\beta} |\{(x, y) \in E \times E : |x - y| < r\}| dr. \end{aligned}$$

Exploiting (4.10) and Lemma 4.6 we deduce

$$\begin{aligned} \int_E \int_E \frac{1}{|x - y|^{\beta-1}} dy dx &= (\beta - 1) \int_0^{+\infty} |\{(x, y) \in E \times E : |x - y| < r\}| \frac{dr}{r^\beta} \\ &= (\beta - 1) \int_0^{+\infty} \int_E \frac{|E \cap B_r(x)|}{r^\beta} dx dr \\ &\geq (\beta - 1) \int_1^{\frac{\text{diam } E}{2}} \int_E \frac{|E \cap B_r(x)|}{r^\beta} dx dr \\ &\geq c \int_1^{\frac{\text{diam } E}{2}} \frac{|E|}{r^{\beta-1}} dr. \end{aligned}$$

The final right-hand side in the previous chain of inequalities is bounded from below by  $c |E| (\text{diam } E)^{2-\beta}$  if  $\beta < 2$ , and by  $c |E| \log(\text{diam } E)$  if  $\beta = 2$ . Combining this bounds with Lemma 4.5, we get a contradiction for  $|E|$  large enough. □

**4.1. Absence of holes in minimizers.** As a corollary of the estimates proved in the last section, we prove here a further qualitative property of volume constrained minimizers of  $\mathcal{F}^\lambda$ . The next theorem essentially tells that volume constrained minimizers of  $\mathcal{F}^\lambda$  do not have ‘‘interior holes’’.

**Theorem 4.7.** *Let  $g$  be  $\mathcal{R}$ -admissible, 0-decreasing, infinitesimal and symmetric and let  $G$  be  $\mathcal{G}$ -admissible and coercive. There exists  $\bar{m} > 0$ , depending on  $n, \lambda, g$  and  $G$ , such that, for all  $m \in (0, \bar{m})$ , every minimizer  $E$  of  $\mathcal{F}^\lambda$  with  $|E| = m$  has the following property. There is no set  $F \subset \mathbb{R}^n \setminus (H \cup E)$  with  $|F| > 0$  such that*

$$(4.11) \quad P_\lambda(E) = P_\lambda(E \cup F) + P(F, \mathbb{R}^n \setminus H) + \lambda \mathcal{H}^{n-1}(\partial^* F \cap \partial H).$$

We begin with a preparatory lemma.

**Lemma 4.8.** *Let  $g$  be  $\mathcal{R}$ -admissible,  $q$ -decreasing, infinitesimal and symmetric and let  $G$  be  $\mathcal{G}$ -admissible and coercive. There exist  $\bar{m} > 0$  and  $\bar{T} > 0$ , depending on  $n, \lambda, g, G, q$  such that, for all  $m \in (0, \bar{m})$ , every volume constrained minimizer  $E$  of  $\mathcal{F}^\lambda$  with  $|E| = m$  satisfies*

$$|E \cap \{x_n > \bar{T}\}| = 0.$$

*Proof.* By Lemma 3.10 there exists  $\bar{T}_E < \infty$ , depending on  $n, \lambda, g, G$  and  $E$  with

$$\bar{T}_E := \sup\{t : |E \cap \{x_n > t\}| > 0\}.$$

Let  $x_E \in E$  such that

$$(x_E)_n \geq \frac{1}{2} \bar{T}_E.$$

By Lemma 4.2 there exists  $c = c(n, \lambda, g, G, q) > 0$  such that, if  $\bar{m} < 1$ , then

$$|E \cap B_1(x_E)| \geq c m.$$

Therefore

$$(4.12) \quad \mathcal{F}^\lambda(E) \geq P_\lambda(E) + \int_{E \cap B_1(x_E)} G \geq P_\lambda(B^\lambda(m)) + c m \inf_{((x_E)_{n-1}, (x_E)_{n+1})} G.$$

On the other hand, by Lemma 3.2, if  $\bar{m} \leq |B^\lambda|$  we have

$$(4.13) \quad \mathcal{F}^\lambda(E) \leq \mathcal{F}^\lambda(B^\lambda(m)) \leq P_\lambda(B^\lambda(m)) + c(n, \lambda, g, G)m.$$

Putting together (4.12) and (4.13) we obtain

$$\inf_{((x_E)_{n-1}, (x_E)_{n+1})} G \leq c(n, \lambda, g, G, q).$$

Since  $G$  is coercive, then  $(x_E)_n$ , and in particular also  $\bar{T}_E$ , is bounded by a constant independent of  $E$ , and we conclude the proof.  $\square$

*Remark 4.9.* We remark that Lemma 4.8 is a stronger result than Lemma 3.10. Indeed, the bound in Lemma 4.8 does not depend on the minimizer. At the same time, Lemma 3.10 does not require that  $g$  is  $q$ -decreasing and  $G$  is coercive.

Now we are ready to prove Theorem 4.7.

*Proof of Theorem 4.7.* If  $\bar{m}$  is sufficiently small, Lemma 4.8 guarantees that there exists  $\bar{T} > 0$ , depending on  $n, \lambda, g, G$ , such that

$$|E \cap \{x_n > \bar{T}\}| = 0.$$

Assume that there exists a set  $F \subset \mathbb{R}^n \setminus (H \cup E)$  with  $v := |F| > 0$  and such that (4.11) holds. We aim to find a contradiction if  $\bar{m}$  is chosen suitably small. Let  $\bar{m} \leq |B^\lambda|$ . By the minimality of  $E$ , the isoperimetric inequality (2.2), the relative isoperimetric inequality outside convex sets [CGR07; FM23] and since  $P_\lambda(B^\lambda(m)) = n|B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}}$  [PP24, Lemma 3.3], we find

$$\begin{aligned} n|B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}} + \mathcal{R}(B^\lambda(m)) + \mathcal{G}(B^\lambda(m)) &= \mathcal{F}^\lambda(B^\lambda(m)) \geq \mathcal{F}^\lambda(E) \geq P_\lambda(E) \\ &= P_\lambda(E \cup F) + P(F, \mathbb{R}^n \setminus H) + \lambda \mathcal{H}^{n-1}(\partial^* F \cap \partial H) \\ &\geq P_\lambda(E \cup F) + (1 - |\lambda|)P(F, \mathbb{R}^n \setminus H) \\ &\geq n|B^\lambda|^{\frac{1}{n}} (m + v)^{\frac{n-1}{n}} + (1 - |\lambda|) n \left(\frac{\omega_n}{2}\right)^{\frac{1}{n}} v^{\frac{n-1}{n}} \\ &\geq n|B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}} + (1 - |\lambda|) n \left(\frac{\omega_n}{2}\right)^{\frac{1}{n}} v^{\frac{n-1}{n}} \end{aligned}$$

which gives, by Lemma 3.2,

$$(4.14) \quad v \leq \left( \frac{\mathcal{R}(B^\lambda(m)) + \mathcal{G}(B^\lambda(m))}{(1 - |\lambda|) n^n \sqrt{\frac{\omega_n}{2}}} \right)^{\frac{n}{n-1}} \leq c(n, \lambda, g, G) \left( \frac{m}{(1 - |\lambda|) n^n \sqrt{\frac{\omega_n}{2}}} \right)^{\frac{n}{n-1}}.$$

Since  $\bar{m} < c(n, \lambda, g, G)$ , also  $v$  is bounded by a suitable  $\bar{v}(n, \lambda, g, G)$ . By [NP21, Lemma 3.5] there exists a continuous and increasing function  $\varphi : (0, \infty) \rightarrow (0, \infty)$ , with  $\varphi(0) = 0$ , such that for every two sets  $F_1, F_2 \subset \mathbb{R}^n \setminus H$  one has

$$\mathcal{R}(F_1, F_2) \leq |F_1| \varphi(|F_2|).$$

Then

$$\mathcal{R}(E \cup F) - \mathcal{R}(E) = \mathcal{R}(F, F) + 2\mathcal{R}(F, E) \leq v \varphi(\bar{v}) + 2v \varphi(\bar{m}) \leq c(n, \lambda, g, G)v.$$

Let us prove that also  $F$  is essentially contained in  $\{0 < x_n \leq \bar{T}\}$ . To this end, assume by contradiction that

$$(4.15) \quad \mathcal{H}^{n-1}(\partial^* F \setminus (H \cup \partial^* E)) > 0.$$

Let us denote

$$\begin{aligned} \Sigma_E &:= \mathcal{H}^{n-1}(\partial^* E \setminus (H \cup \partial^* F)) \\ \Sigma &:= \mathcal{H}^{n-1}((\partial^* E \cap \partial^* F) \setminus H) \\ \Sigma_F &:= \mathcal{H}^{n-1}(\partial^* F \setminus (H \cup \partial^* E)) \\ \Theta_E &:= \mathcal{H}^{n-1}(\partial^* E \cap \partial H) \\ \Theta_F &:= \mathcal{H}^{n-1}(\partial^* F \cap \partial H). \end{aligned}$$

By (4.11) we obtain

$$\begin{aligned} \Sigma_E + \Sigma - \lambda \Theta_E &= P_\lambda(E) \\ &= P_\lambda(E \cup F) + P(F, \mathbb{R}^n \setminus H) + \lambda \Theta_F \\ &= \Sigma_E + \Sigma_F - \lambda \Theta_E - \lambda \Theta_F + \Sigma + \Sigma_F + \lambda \Theta_F \\ &= \Sigma_E + 2\Sigma_F + \Sigma - \lambda \Theta_E. \end{aligned}$$

In particular, we get  $\Sigma_F = 0$ , contradicting (4.15). Therefore  $F \subset \{0 < x_n \leq \bar{T}\}$  and we also deduce

$$(4.16) \quad \mathcal{G}(E \cup F) - \mathcal{G}(E) = \mathcal{G}(F) \leq \int_{F(0, \bar{T})} \sup G \, dx = c(n, \lambda, g, G)v.$$

By (4.11), we obtain

$$\begin{aligned} \mathcal{F}^\lambda(E \cup F) &= P_\lambda(E \cup F) + \mathcal{R}(E \cup F) + \mathcal{G}(E \cup F) \\ &= P_\lambda(E) - P(F, \mathbb{R}^n \setminus H) - \lambda \mathcal{H}^{n-1}(\partial^* F \cap \partial H) + \mathcal{R}(E \cup F) + \mathcal{G}(E \cup F) \\ &\leq P_\lambda(E) - (1 - |\lambda|)P(F, \mathbb{R}^n \setminus H) + \mathcal{R}(E \cup F) + \mathcal{G}(E \cup F) \\ (4.17) \quad &\leq P_\lambda(E) - (1 - |\lambda|)n \left( \frac{\omega_n}{2} \right)^{\frac{1}{n}} v^{\frac{n-1}{n}} + \mathcal{R}(E) + cv + \mathcal{G}(E) + cv \\ &= \mathcal{F}^\lambda(E) - (1 - |\lambda|)n \left( \frac{\omega_n}{2} \right)^{\frac{1}{n}} v^{\frac{n-1}{n}} + cv < \mathcal{F}^\lambda(E), \end{aligned}$$

where the last inequality holds if  $v$  is sufficiently small, hence by (4.14) as soon as  $\bar{m}$  is sufficiently small.

Let  $t \in (0, \infty)$  be such that, if  $D = \{x \in E \cup F, x_n < t\}$ , then  $|D| = m$ . Clearly  $\mathcal{F}^\lambda(D) \leq \mathcal{F}^\lambda(E \cup F)$ , hence (4.17) implies  $\mathcal{F}^\lambda(D) < \mathcal{F}^\lambda(E)$ , contradicting the minimality of  $E$ .  $\square$

*Remark 4.10.* Note that if the function  $G$  were globally bounded, Theorem 4.7 could be easily extended to minimizers of  $\mathcal{F}^\lambda$  in the class

$$\mathcal{A}_m = \{\Omega \subset \mathbb{R}^n \setminus H \text{ measurable} : |\Omega| = m\}$$

for every mass  $m > 0$ .

Indeed, in the proof of Theorem 4.7 we exploited Lemma 4.8 just to get the estimate (4.16), which is trivial in case  $G$  were assumed to be globally bounded.

Now we are ready to complete the proof of Theorem 1.1.

*Proof of Theorem 1.1.* Theorem 1.1 follows by Theorem 3.1, Theorem 3.6, Theorem 3.12 and Theorem 4.7.  $\square$

## 5. GENERALIZED MINIMIZERS

Let us give the following definition.

**Definition 5.1.** If  $E \subset \mathbb{R}^n \setminus H$  is a measurable set,  $g$  is a  $\mathcal{R}$ -admissible function,  $G$  is a  $\mathcal{G}$ -admissible function and  $\varepsilon_1, \varepsilon_2 > 0$ , we define the functional

$$\begin{aligned} \mathcal{F}_\varepsilon^\lambda(E) &:= P_\lambda(E) + \varepsilon_1 \mathcal{R}(E) + \varepsilon_2 \mathcal{G}(E) \\ &= P_\lambda(E) + \varepsilon_1 \int_E \int_E g(y-x) dy dx + \varepsilon_2 \int_E G(x_n) dx. \end{aligned}$$

*Remark 5.2.* We remark that minimizing the functional  $\mathcal{F}^\lambda$  in the small mass regime is equivalent to minimizing the functional  $\mathcal{F}_\varepsilon^\lambda$  for  $\varepsilon_1, \varepsilon_2$  small and among sets of a fixed volume. Indeed, let for instance  $|E| = |B^\lambda|$ , if  $m > 0$  and  $\bar{\varepsilon} := \frac{m^{\frac{1}{n}}}{|B^\lambda|^{\frac{1}{n}}}$ , then  $\tilde{E} := \bar{\varepsilon}E$  has volume  $m$  and by scaling we have

$$\mathcal{F}^\lambda(\tilde{E}) = \bar{\varepsilon}^{n-1} \left( P_\lambda(E) + \bar{\varepsilon}^{n+1} \int_E \int_E g(\bar{\varepsilon}(y-x)) dy dx + \bar{\varepsilon} \int_E G(\bar{\varepsilon}x_n) dx \right).$$

In particular we deduce that

- for every  $\mathcal{R}$ -admissible function  $g_1$ ,  $\mathcal{G}$ -admissible function  $G_1$  and  $m > 0$ , there exist  $\bar{\varepsilon} > 0$ , a  $\mathcal{R}$ -admissible function  $g_2$  and a  $\mathcal{G}$ -admissible function  $G_2$  such that, if

$$\mathcal{F}^\lambda(E) = P_\lambda(E) + \int_E \int_E g_1(y-x) dy dx + \int_E G_1(x_n) dx$$

and

$$\mathcal{F}_{\bar{\varepsilon}}^\lambda(E) = P_\lambda(E) + \bar{\varepsilon}^{n+1} \int_E \int_E g_2(y-x) dy dx + \bar{\varepsilon} \int_E G_2(x_n) dx,$$

then  $\inf_{|E|=m} \mathcal{F}^\lambda(E)$  is proportional to  $\inf_{|E|=|B^\lambda|} \mathcal{F}_{\bar{\varepsilon}}^\lambda(E)$  and the variational problems are equivalent.

- for every  $\mathcal{R}$ -admissible function  $g_2$ ,  $\mathcal{G}$ -admissible function  $G_2$ ,  $\varepsilon_1, \varepsilon_2 > 0$  and  $m > 0$ , there exist a  $\mathcal{R}$ -admissible function  $g_1$  and a  $\mathcal{G}$ -admissible function  $G_1$  such that, if

$$\mathcal{F}^\lambda(E) = P_\lambda(E) + \int_E \int_E g_1(y-x) dy dx + \int_E G_1(x_n) dx$$

and

$$\mathcal{F}_\varepsilon^\lambda(E) = P_\lambda(E) + \varepsilon_1 \int_E \int_E g_2(y-x) dy dx + \varepsilon_2 \int_E G_2(x_n) dx,$$

then  $\inf_{|E|=|B^\lambda|} \mathcal{F}_\varepsilon^\lambda(E)$  is proportional to  $\inf_{|E|=m} \mathcal{F}^\lambda(E)$  and the variational problems are equivalent.

From now on for the rest of the section, we assume that  $\varepsilon_1, \varepsilon_2 > 0$  and  $g, G$  as in Definition 5.1 are given. Hence we also define the generalized energy corresponding to  $\mathcal{F}_\varepsilon^\lambda$  as

$$\tilde{\mathcal{F}}_\varepsilon^\lambda(E) := \inf_{h \in \mathbb{N}} \tilde{\mathcal{F}}_{\varepsilon,h}^\lambda(E),$$

where

$$\tilde{\mathcal{F}}_{\varepsilon,h}^\lambda(E) := \inf \left\{ \sum_{i=1}^h \mathcal{F}_\varepsilon^\lambda(E^i) : E = \bigcup_{i=1}^h E^i, E^i \cap E^j = \emptyset \text{ for } 1 \leq i \neq j \leq h \right\}.$$

The goal of this Section is to prove the following version of Theorem 1.3, suitably modified for the functional  $\mathcal{F}_\varepsilon^\lambda$ .

**Theorem 5.3.** *Let  $g$  be  $\mathcal{R}$ -admissible and  $q$ -decreasing and let  $G$  be  $\mathcal{G}$ -admissible. For every  $\varepsilon_1, \varepsilon_2 > 0$  there exists a minimizer of  $\tilde{\mathcal{F}}_\varepsilon^\lambda$  in the class*

$$\mathcal{A} := \left\{ \Omega \subset \mathbb{R}^n \setminus H \text{ measurable} : |\Omega| = |B^\lambda| \right\}.$$

More precisely, there exist a set  $E \in \mathcal{A}$  and a subdivision  $E = \bigcup_{j=1}^h E^j$ , with pairwise disjoint sets  $E^j$ , such that

$$\tilde{\mathcal{F}}_\varepsilon^\lambda(E) = \sum_{j=1}^h \mathcal{F}_\varepsilon^\lambda(E^j) = \inf \left\{ \tilde{\mathcal{F}}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \right\}.$$

Moreover, for every  $1 \leq j \leq h$ , the set  $E^j$  is a minimizer of both the standard and the generalized energy for its volume, i.e.

$$(5.1) \quad \tilde{\mathcal{F}}_\varepsilon^\lambda(E^j) = \mathcal{F}_\varepsilon^\lambda(E^j) = \min \left\{ \tilde{\mathcal{F}}_\varepsilon^\lambda(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = |E^j| \right\}.$$

Note that an analogous version of Lemma 3.4 holds.

**Lemma 5.4.** *Let  $E \subset \mathbb{R}^n \setminus H$  be a set of finite perimeter. Let  $g$  be  $\mathcal{R}$ -admissible and  $q$ -decreasing and let  $G$  be  $\mathcal{G}$ -admissible. If  $\alpha > 1$ , then*

$$\mathcal{F}_\varepsilon^\lambda(\alpha E) \leq \alpha^{2n+q} \mathcal{F}_\varepsilon^\lambda(E).$$

We begin by proving some preparatory lemmas. The next geometric lemma allows to modify an excessively long and thin set decreasing its energy.

**Lemma 5.5.** *Let  $g$  be  $\mathcal{R}$ -admissible and  $G$  be  $\mathcal{G}$ -admissible. For every  $\bar{m} > 0$  there exists  $L(n, \lambda, \bar{m}) > 0$  such that the following holds. Let  $E \subset \mathbb{R}^n \setminus H$ , and let  $a < b$  be two numbers with  $b > a + 2L$  and such that*

$$\left| \{x \in E : a \leq x_1 \leq b\} \right| < \bar{m}.$$

*Then there exist two numbers  $a^+ \in [a, a + L]$  and  $b^- \in [b - L, b]$  such that, denoting  $E^- = E \setminus ([a^+, b^-] \times \mathbb{R}^{n-2} \times (0, \infty))$  and  $m = |E| - |E^-| < \bar{m}$ , one has*

$$(5.2) \quad \mathcal{F}_\varepsilon^\lambda(E^-) \leq \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^\frac{1}{n}m^\frac{n-1}{n}.$$

*Proof.* It is sufficient to prove the claim for bounded sets  $E$  such that  $\partial E \setminus \partial H$  is a smooth hypersurface with  $\mathcal{H}^{n-1}(\{x \in \partial E \setminus H : \nu^E(x) = \pm e_j\}) = 0$  for any  $j = 1, \dots, n$ . Indeed, if  $E$  is a generic set of finite perimeter satisfying the hypotheses of Lemma 5.5, let  $E_i \xrightarrow{L^1} E$  be the sequence of sets given by [PP24, Lemma 2.3]. For  $i$  sufficiently large,

$$\left| \{x \in E_i : a \leq x_1 \leq b\} \right| < \bar{m}$$

holds. Then there exist  $a_i^+ \in [a, a + L]$  and  $b_i^- \in [b - L, b]$  such that, if we set  $E_i^- = E_i \setminus ([a_i^+, b_i^-] \times (0, \infty))$  and  $m_i = |E_i| - |E_i^-|$ , we obtain

$$\mathcal{F}_\varepsilon^\lambda(E_i^-) \leq \mathcal{F}_\varepsilon^\lambda(E_i) - \frac{1}{2}n|B^\lambda|^\frac{1}{n}m_i^\frac{n-1}{n}.$$

Up to subsequence,  $a_i^+$  and  $b_i^-$  converge to certain  $a^+ \in [a, a + L]$  and  $b^- \in [b - L, b]$  respectively. By [PP24, Lemma 2.3]  $P_\lambda(E_i) \rightarrow P_\lambda(E)$ . By the lower semicontinuity of  $P_\lambda$  ([PP24, Lemma 3.9]), the continuity of  $\mathcal{R}$  and  $\mathcal{G}$  under strong  $L^1$  convergence and the properties of  $\{E_i\}$ , if  $E^- = E \setminus ([a^+, b^-] \times (0, \infty))$  then

$$\begin{aligned} \mathcal{F}_\varepsilon^\lambda(E^-) &= P_\lambda(E^-) + \varepsilon_1 \mathcal{R}(E^-) + \varepsilon_2 \mathcal{G}(E^-) \\ &\leq \liminf_i (P_\lambda(E_i^-) + \varepsilon_1 \mathcal{R}(E_i^-) + \varepsilon_2 \mathcal{G}(E_i^-)) \\ &\leq \liminf_i \left( P_\lambda(E_i) + \varepsilon_1 \mathcal{R}(E_i) + \varepsilon_2 \mathcal{G}(E_i) - \frac{1}{2}n|B^\lambda|^\frac{1}{n}m_i^\frac{n-1}{n} \right) \\ &= \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^\frac{1}{n}m^\frac{n-1}{n}. \end{aligned}$$

So let us fix  $\bar{m}$  and consider  $a$  and  $b$  as in the claim, with  $L$  to be determined later. For  $t \in \mathbb{R}$  let

$$\sigma(t) := \mathcal{H}^{n-1}(E \cap \{x \in \mathbb{R}^n : x_1 = t\}).$$

If  $c := \frac{a+b}{2}$ , let

$$\varphi(t) := \int_t^c \sigma(s) ds.$$

We note that there exists  $a^+ \in [a, a + L] \subset [a, c)$  such that, if

$$m_1 := \left| \{x \in E : a^+ < x_1 < c\} \right|,$$

then

$$(5.3) \quad \sigma(a^+) \leq \frac{1}{8}n|B^\lambda|^\frac{1}{n}m_1^\frac{n-1}{n}.$$

Indeed, assume by contradiction that for every  $t \in (a, a + L)$  it holds

$$-\varphi'(t) = \sigma(t) > \frac{1}{8}n|B^\lambda|^\frac{1}{n}\varphi(t)^\frac{n-1}{n}.$$

Then  $\varphi|_{(a, a+L)}$  is a positive decreasing function satisfying

$$\begin{cases} \varphi(a) \leq \bar{m}, \\ |\varphi'(t)| > \frac{1}{8}n|B^\lambda|^\frac{1}{n}\varphi(t)^\frac{n-1}{n}. \end{cases}$$

By standard ODE comparison there exists a constant  $d > 0$  depending only on  $n$ ,  $\lambda$  and  $\bar{m}$  such that, if  $a+d < a+L$ , then  $\varphi(t) \rightarrow 0$  as  $t \rightarrow (a+d)^-$ . Hence  $L$  could be chosen so big that  $a+d < a+L < c$ , and then  $\varphi(t) = 0$  for any  $t \in (a+d, c)$ . It follows that there exists  $a^+$  such that (5.3) holds. Similarly, up to choosing a larger  $L$ , we have the existence of  $b^- \in [b-L, b] \subset (c, b]$  such that, if

$$m_2 := \left| \{x \in E : c < x_1 < b^-\} \right|,$$

then

$$\sigma(b^-) \leq \frac{1}{8} n |B^\lambda|^{\frac{1}{n}} m_2^{\frac{n-1}{n}}.$$

Let  $E^- := E \setminus ([a^+, b^-] \times (0, \infty))$  and  $F := E \setminus E^-$ . Then

$$|F| = m = m_1 + m_2,$$

and, by isoperimetric inequality (2.2), there holds

$$P_\lambda(F) \geq n |B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}}.$$

Hence

$$\begin{aligned} P_\lambda(E^-) &\leq P_\lambda(E) - P_\lambda(F) + 2(\sigma(a^+) + \sigma(b^-)) \\ &\leq P_\lambda(E) - P_\lambda(F) + \frac{1}{4} n |B^\lambda|^{\frac{1}{n}} \left( m_1^{\frac{n-1}{n}} + m_2^{\frac{n-1}{n}} \right) \\ &\leq P_\lambda(E) - P_\lambda(F) + \frac{1}{2} n |B^\lambda|^{\frac{1}{n}} (m_1 + m_2)^{\frac{n-1}{n}} \\ &\leq P_\lambda(E) - \frac{1}{2} n |B^\lambda|^{\frac{1}{n}} (m_1 + m_2)^{\frac{n-1}{n}}. \end{aligned}$$

Since  $E^- \subset E$ , then  $\mathcal{R}(E^-) \leq \mathcal{R}(E)$  and  $\mathcal{G}(E^-) \leq \mathcal{G}(E)$ , and we deduce (5.2).  $\square$

The following variant of Lemma 5.5 concerns the case of the vertical direction when we modify a part lying on the hyperplane  $\{x_n = 0\}$ .

**Lemma 5.6.** *Let  $g$  be  $\mathcal{R}$ -admissible and  $G$  be  $\mathcal{G}$ -admissible. For every  $\bar{m} \in \mathbb{R}$  there exists  $L(n, \lambda, \bar{m}) > 0$  such that the following holds. Let  $E \subset \mathbb{R}^n \setminus H$ , and let  $b$  be a number with  $b > L$  and such that*

$$\left| \{x \in E : 0 < x_n \leq b\} \right| < \bar{m}.$$

*There exists then  $b^- \in [b-L, b]$  such that, denoting  $E^- = E \setminus (\mathbb{R}^{n-1} \times [0, b^-])$  and  $m = |E| - |E^-| \leq \bar{m}$ , one has*

$$(5.4) \quad \mathcal{F}_\varepsilon^\lambda(E^-) \leq \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2} n |B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}}.$$

*Proof.* As in the proof of Lemma 5.5, we may assume that  $\partial E \setminus \partial H$  is a smooth hypersurface with  $\mathcal{H}^{n-1}(\{x \in \partial E \setminus H : \nu^{E_i}(x) = \pm e_j\}) = 0$  for any  $j = 1, \dots, n$ . Let us fix  $\bar{m}$  and consider  $b$  as in the claim, with  $L$  to be determined later. For almost every  $t \in \mathbb{R}$ , let

$$\sigma(t) := \mathcal{H}^{n-1}(E \cap \{x \in \mathbb{R}^n : x_1 = t\}).$$

Let

$$\varphi(t) := \int_0^t \sigma(s) ds.$$

As in the proof of Lemma 5.5, we note that there exists  $b^- \in [b-L, b] \subset (0, b]$  such that, if

$$m := \left| \{x \in E : 0 < x_n < b^-\} \right|,$$

then

$$(5.5) \quad \sigma(b^-) \leq \frac{1}{4} n |B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}}.$$

If  $E^- := E \setminus (\mathbb{R}^{n-1} \times [0, b^-])$  and  $F := E \setminus E^-$ , then

$$|F| = m$$

and, by isoperimetric inequality (2.2),

$$P_\lambda(F) \geq n |B^\lambda|^{\frac{1}{n}} m^{\frac{n-1}{n}}.$$

We establish that

$$\begin{aligned} P_\lambda(E^-) &\leq P_\lambda(E) - P_\lambda(F) + 2\sigma(b^-) \\ &\leq P_\lambda(E) - P_\lambda(F) + \frac{1}{2}n|B^\lambda|^{\frac{1}{n}}m^{\frac{n-1}{n}} \\ &\leq P_\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}}m^{\frac{n-1}{n}}. \end{aligned}$$

Since  $E^- \subset E$ , then  $\mathcal{R}(E^-) \leq \mathcal{R}(E)$ ,  $\mathcal{G}(E^-) \leq \mathcal{G}(E)$  and we deduce (5.4).  $\square$

We now prove a uniform boundedness result.

**Lemma 5.7.** *Let  $g$  be  $\mathcal{R}$ -admissible and  $q$ -decreasing and let  $G$  be  $\mathcal{G}$ -admissible. Let  $\varepsilon_1, \varepsilon_2 > 0$ . For every  $m \in (0, \infty)$  there exist  $R > 0$  and  $\bar{h} \in \mathbb{N}$ , depending on  $n, \lambda, m, \varepsilon_1, \varepsilon_2, g, G$  and  $q$ , such that*

$$\inf \left\{ \mathcal{F}_\varepsilon^\lambda(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = m \right\} \geq \inf \left\{ \tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = m \right\},$$

where

$$\tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}(\Omega) := \inf \left\{ \sum_{i=1}^{\bar{h}} \mathcal{F}_\varepsilon^\lambda(\Omega^i) : \Omega = \bigcup_{i=1}^{\bar{h}} \Omega^i, \Omega^i \cap \Omega^j = \emptyset, \text{diam } \Omega^i \leq R \quad \forall 1 \leq i \neq j \leq \bar{h} \right\}.$$

*Proof.* Let  $M(n, \lambda, m, \varepsilon_1, \varepsilon_2, g, G, q) \in \mathbb{N}$  be a natural number to be determined later and let us denote  $\bar{m} = m/M$ . Let  $E \subset \mathbb{R}^n \setminus H$  be a bounded set with  $|E| = m$  and

$$(5.6) \quad \mathcal{F}_\varepsilon^\lambda(E) \leq \inf \left\{ \mathcal{F}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \right\} + \frac{n|B^\lambda|^{\frac{1}{n}}}{3} \left( \frac{m}{M^2} \right)^{\frac{n-1}{n}}.$$

This is possible since the infimum is reached by a sequence of bounded sets. Let  $t_0 < t_1 < \dots < t_{M-1} < t_M$  be real numbers such that

$$\left| E \cap \left( (t_i, t_{i+1}) \times \mathbb{R}^{n-2} \times (0, \infty) \right) \right| = \bar{m},$$

for every  $0 \leq i \leq M-1$  and let  $L(n, \lambda, \bar{m})$  be given by Lemma 5.5. For every  $0 \leq i \leq M-1$  let us define the interval  $I_i$  in the following way. If  $t_{i+1} - t_i \leq 2L$  we set  $I_i = \emptyset$ , otherwise we apply Lemma 5.5 with  $a = t_i$  and  $b = t_{i+1}$  and we set  $I_i = [a^+, b^-]$ . If  $m_i = |E \cap (I_i \times \mathbb{R}^{n-2} \times (0, \infty))|$ , then

$$(5.7) \quad m_i \leq \frac{m}{M^2}.$$

Indeed, if  $I_i = \emptyset$ , then (5.7) is clearly true. If  $I_i \neq \emptyset$ , we set

$$E' = \alpha \left( E \setminus (I_i \times \mathbb{R}^{n-2} \times (0, \infty)) \right),$$

with  $\alpha = \left( \frac{m}{m-m_i} \right)^{\frac{1}{n}}$ . Note that  $\frac{m_i}{m} \leq \frac{1}{M}$  by construction. By Lemma 5.4 and (5.2) we have

$$\begin{aligned} \mathcal{F}_\varepsilon^\lambda(E') &\leq \left( \frac{m}{m-m_i} \right)^{2+\frac{q}{n}} \mathcal{F}_\varepsilon^\lambda \left( E \setminus (I_i \times \mathbb{R}^{n-2} \times (0, \infty)) \right) \\ &\leq \left( \frac{1}{1-\frac{m_i}{m}} \right)^{2+\frac{q}{n}} \left( \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}}m_i^{\frac{n-1}{n}} \right). \end{aligned}$$

Moreover, if  $M$  is large enough,

$$(5.8) \quad \begin{aligned} \mathcal{F}_\varepsilon^\lambda(E') &\leq \left( 1 + \left( 3 + \frac{q}{n} \right) \frac{m_i}{m} \right) \left( \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}}m_i^{\frac{n-1}{n}} \right) \\ &\leq \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{3}n|B^\lambda|^{\frac{1}{n}}m_i^{\frac{n-1}{n}}. \end{aligned}$$

Estimates (5.6) and (5.8) imply that

$$\mathcal{F}_\varepsilon^\lambda(E') \leq \inf \left\{ \mathcal{F}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \right\} + \frac{n|B^\lambda|^{\frac{1}{n}}}{3} \left( \frac{m}{M^2} \right)^{\frac{n-1}{n}} - \frac{n|B^\lambda|^{\frac{1}{n}}}{3} m_i^{\frac{n-1}{n}},$$

and, since  $|E'| = m$ , (5.7) holds.



Let

$$\tilde{E} = E \setminus \left( \bigcup_{i=0}^{M-1} I_i \times \mathbb{R}^{n-2} \times (0, \infty) \right)$$

and  $\mu = \sum_{i=0}^{M-1} m_i$ , so that  $|\tilde{E}| = m - \mu$ . By Lemma 5.5 and the subadditivity of power function with exponent less than 1 we get

$$\mathcal{F}_\varepsilon^\lambda(\tilde{E}) \leq \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}} \sum_{i=0}^{M-1} m_i^{\frac{n-1}{n}} \leq \mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}} \mu^{\frac{n-1}{n}}.$$

The set  $F := \left(\frac{m}{m-\mu}\right)^{\frac{1}{n}} \tilde{E}$  has volume  $m$ . We can use Lemma 5.4 to obtain

$$\mathcal{F}_\varepsilon^\lambda(F) \leq \left(\frac{m}{m-\mu}\right)^{2+\frac{q}{n}} \mathcal{F}_\varepsilon^\lambda(\tilde{E}) \leq \left(\frac{m}{m-\mu}\right)^{2+\frac{q}{n}} \left(\mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}} \mu^{\frac{n-1}{n}}\right).$$

If  $\mu$  is small enough, which happens as soon as  $M$  is large enough thanks to (5.7), we deduce

$$\begin{aligned} \mathcal{F}_\varepsilon^\lambda(F) &\leq \left(\frac{m}{m-\mu}\right)^{2+\frac{q}{n}} \left(\mathcal{F}_\varepsilon^\lambda(E) - \frac{1}{2}n|B^\lambda|^{\frac{1}{n}} \mu^{\frac{n-1}{n}}\right) \\ &\leq \mathcal{F}_\varepsilon^\lambda(E) + C\mu \left( \inf \{ \mathcal{F}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \} + c \left(\frac{m}{M^2}\right)^{\frac{n-1}{n}} \right) + (1 + C\mu) \left(-\frac{1}{2}n|B^\lambda|^{\frac{1}{n}} \mu^{\frac{n-1}{n}}\right) \\ &\leq \mathcal{F}_\varepsilon^\lambda(E). \end{aligned}$$

Note that  $\tilde{E}$  is the union of at most  $M + 1$  sets, each contained in a slab having width at most equal to  $2L$  by Lemma 5.5. In particular,  $F$  is the union of at most  $M + 1$  parts and each of them has horizontal width at most equal  $3L$ .

If we repeat the arguments in the remaining directions, with care to apply also Lemma 5.6 in the  $n$ -th direction, we get the boundedness of the pieces in all the  $n$  directions. Then there exist  $R \in (0, \infty)$ ,  $\bar{h} \in \mathbb{N}$  and  $G \subset \mathbb{R}^n$  such that  $|G| = m$ ,  $G = \bigcup_{i=1}^{\bar{h}} G_i$ ,  $G_i \cap G_j = \emptyset$ ,  $\text{diam } G_i \leq R$  and  $\mathcal{F}_\varepsilon^\lambda(G) \leq \mathcal{F}_\varepsilon^\lambda(E)$ . Finally

$$\mathcal{F}_\varepsilon^\lambda(G) \geq \sum_{i=1}^{\bar{h}} \mathcal{F}_\varepsilon^\lambda(G_i) \geq \tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}(G).$$

□

Now we are ready to prove Theorem 5.3.

*Proof of Theorem 5.3.* We begin by proving the existence of  $h' \in \mathbb{N}$  and of a sequence  $\{G_i\}_{i \in \mathbb{N}} \subset \mathcal{A}$  such that

$$(5.9) \quad \inf \{ \tilde{\mathcal{F}}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \} = \lim_{i \rightarrow \infty} \tilde{\mathcal{F}}_{\varepsilon, h'}^\lambda(G_i).$$

Let  $h'(n, \lambda, \varepsilon_1, \varepsilon_2, g, G, q)$  be an integer to be determined later and consider a sequence  $\{E_i\}_{i \in \mathbb{N}} \subset \mathcal{A}$  such that

$$(5.10) \quad K := \inf \{ \tilde{\mathcal{F}}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \} = \lim_{i \rightarrow \infty} \tilde{\mathcal{F}}_\varepsilon^\lambda(E_i).$$

For every  $i \in \mathbb{N}$  let  $h(i) \in \mathbb{N}$  such that there exists a subdivision  $E_i = E_i^1 \cup E_i^2 \cup \dots \cup E_i^{h(i)}$  with

$$(5.11) \quad \tilde{\mathcal{F}}_\varepsilon^\lambda(E_i) > \left(1 - \frac{1}{i+1}\right) \sum_{j=1}^{h(i)} \mathcal{F}_\varepsilon^\lambda(E_i^j).$$

Without loss of generality, we can assume  $h(i) \rightarrow \infty$ , so that  $h(i) > h'$  for  $i$  large enough. Let us fix a generic  $i \in \mathbb{N}$ . For simplicity of notation, let us denote  $h = h(i)$  and  $m_j = |E_i^j|$  for every  $1 \leq j \leq h$ . Let us also assume, without loss of generality, that  $m_j$  is decreasing with respect to  $j$ . By (5.11) we get

$$\tilde{\mathcal{F}}_\varepsilon^\lambda(E_i) \geq \frac{1}{2} \sum_{j=1}^h P_\lambda(E_i^j) \geq \frac{1}{2} \sum_{j=1}^h n|B^\lambda|^{\frac{1}{n}} m_j^{\frac{n-1}{n}} \geq \frac{1}{2\sqrt[n]{m_1}} \sum_{j=1}^h n|B^\lambda|^{\frac{1}{n}} m_j = \frac{1}{2\sqrt[n]{m_1}} n|B^\lambda|^{\frac{n+1}{n}}.$$

If  $i$  is large enough, by (5.10) we deduce that

$$(5.12) \quad m_1 \geq \left(\frac{n|B^\lambda|^{\frac{n+1}{n}}}{4K}\right)^n.$$

For every such  $i$ , we define

$$G_i = \alpha \bigcup_{j=1}^{h'} E_i^j,$$

with

$$(5.13) \quad \alpha = \left( \frac{|B^\lambda|}{|B^\lambda| - \sum_{j>h'} m_j} \right)^{\frac{1}{n}} \leq 1 + c_1 \sum_{j=h'+1}^h m_j,$$

where  $c_1$  is a constant depending on  $n$ ,  $\lambda$  and  $K$  (that is on  $n$ ,  $\lambda$ ,  $\varepsilon_1$ ,  $\varepsilon_2$ ,  $g$  and  $G$ ). Note that also  $G_i$  belongs to  $\mathcal{A}$ . By Lemma 5.4 we deduce

$$(5.14) \quad \begin{aligned} \tilde{\mathcal{F}}_{\varepsilon, h'}^\lambda(G_i) &\leq \sum_{j=1}^{h'} \mathcal{F}_\varepsilon^\lambda(\alpha E_i^j) \leq \alpha^{2n+q} \sum_{j=1}^{h'} \mathcal{F}_\varepsilon^\lambda(E_i^j) \\ &\leq \left( 1 + c_2(n, \lambda, \varepsilon_1, \varepsilon_2, g, G, q) \sum_{j=h'+1}^h m_j \right) \sum_{j=1}^{h'} \mathcal{F}_\varepsilon^\lambda(E_i^j). \end{aligned}$$

By (5.11) we get

$$\begin{aligned} \tilde{\mathcal{F}}_\varepsilon^\lambda(E_i) &> \left( 1 - \frac{1}{i+1} \right) \sum_{j=1}^h \mathcal{F}_\varepsilon^\lambda(E_i^j) \\ &\geq \left( 1 - \frac{1}{i+1} \right) \sum_{j=1}^{h'} \mathcal{F}_\varepsilon^\lambda(E_i^j) + \frac{1}{2} \sum_{j=h'+1}^h P_\lambda(E_i^j) \\ &\geq \left( 1 - \frac{1}{i+1} \right) \sum_{j=1}^{h'} \mathcal{F}_\varepsilon^\lambda(E_i^j) + \frac{1}{2} n |B^\lambda|^{\frac{1}{n}} \sum_{j=h'+1}^h m_j^{\frac{n-1}{n}}. \end{aligned}$$

If  $i$  is large enough, by (5.10) and (5.11)

$$\sum_{j=1}^{h'} \mathcal{F}_\varepsilon^\lambda(E_i^j) \leq 2K.$$

By (5.14)

$$(5.15) \quad \tilde{\mathcal{F}}_{\varepsilon, h'}^\lambda(G_i) - \tilde{\mathcal{F}}_{\varepsilon, h}^\lambda(E_i) \leq 2K \left( c_2(n, \lambda, \varepsilon_1, \varepsilon_2, g, G, q) \sum_{j=h'+1}^h m_j + \frac{1}{i+1} \right) - \frac{1}{2} n |B^\lambda|^{\frac{1}{n}} \sum_{j=h'+1}^h m_j^{\frac{n-1}{n}}.$$

Now we can define  $h' \in \mathbb{N}$  so that

$$h' \geq \left( \frac{4Kc_2(n, \lambda, \varepsilon_1, \varepsilon_2, g, G, q)}{n} \right)^n.$$

Since  $m_j \leq \frac{|B^\lambda|}{h'}$  for  $j > h'$ , if  $i$  is large enough we get from (5.15)

$$\tilde{\mathcal{F}}_{\varepsilon, h'}^\lambda(G_i) \leq \tilde{\mathcal{F}}_{\varepsilon, h}^\lambda(E_i) + \frac{2K}{i+1}.$$

By (5.10) we finally deduce that  $\{G_i\}_{i \in \mathbb{N}} \subset \mathcal{A}$  satisfies (5.9).

Let us now show that for every  $m > 0$  there exist  $\bar{h} \in \mathbb{N}$ , a bounded set  $E$  with  $|E| = m$  and a subdivision  $E = \bigcup_{k=1}^{\bar{h}} E^k$  such that

$$(5.16) \quad \tilde{\mathcal{F}}_\varepsilon^\lambda(E) \leq \sum_{k=1}^{\bar{h}} \mathcal{F}_\varepsilon^\lambda(E^k) \leq \inf \{ \mathcal{F}_\varepsilon^\lambda(\Omega) : \Omega \in \mathbb{R}^n \setminus H, |\Omega| = m \}.$$

Let  $R$  and  $\bar{h}$  be as in Lemma 5.7. By Lemma 5.7 there is a sequence of sets  $\{\Omega_i\}_{i \in \mathbb{N}}$  of volume  $m$  such that

$$(5.17) \quad \inf \{ \mathcal{F}_\varepsilon^\lambda(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = m \} \geq \lim_{i \rightarrow +\infty} \tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}(\Omega_i),$$

where  $\tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}$  is defined in Lemma 5.7. For every  $i \in \mathbb{N}$  there exists a partition  $\Omega_i = \Omega_i^1 \cup \Omega_i^2 \cup \dots \cup \Omega_i^{\bar{h}}$  with  $\text{diam}(\Omega_i^j) \leq R$  and

$$(5.18) \quad \sum_{j=1}^{\bar{h}} \mathcal{F}_{\varepsilon}^{\lambda}(\Omega_i^j) \leq \tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}(\Omega_i) + \frac{1}{i}.$$

Up to a subsequence there exist  $m_j \in (0, \infty)$ , with  $1 \leq j \leq \bar{h}$ , such that

$$m_j = \lim_{i \rightarrow \infty} |\Omega_i^j| \quad \forall 1 \leq j \leq \bar{h}, \quad m = \sum_{j=1}^{\bar{h}} m_j.$$

Let us fix  $1 \leq k \leq \bar{h}$  and consider the sets  $\{\Omega_i^k\}_{i \in \mathbb{N}}$ . Since their diameters are uniformly bounded by  $R$ , up to translations we can assume that all the  $\Omega_i^k$  are pairwise disjoint and contained in a fixed ball with radius  $R$ . Therefore the characteristic functions  $f_i = \chi_{\Omega_i^k}$  have uniformly bounded supports and are bounded in  $BV$ . Up to a subsequence, we can assume that  $f_i$  is weakly\* convergent in  $BV$ , and in particular strongly convergent in  $L^1$ , to a certain function  $f$ . Then  $f$  is the characteristic function of a bounded set  $E^k$  with volume  $m_k$ . By the lower-semicontinuity of the perimeter under weak\*  $BV$ -convergence and the continuity of  $\mathcal{R}$  and  $\mathcal{G}$  under strong  $L^1$  convergence, we obtain that

$$(5.19) \quad \mathcal{F}_{\varepsilon}^{\lambda}(E^k) \leq \liminf_{i \rightarrow \infty} \mathcal{F}_{\varepsilon}^{\lambda}(\Omega_i^k).$$

Up to a translation we can assume that the sets  $E^k$  are pairwise disjoint. In particular the set  $E = \bigcup_{k=1}^{\bar{h}} E^k$  is bounded with  $|E| = m$ . By (5.17), (5.18) and (5.19) we get

$$\begin{aligned} \tilde{\mathcal{F}}_{\varepsilon}^{\lambda}(E) &\leq \sum_{k=1}^{\bar{h}} \mathcal{F}_{\varepsilon}^{\lambda}(E^k) \leq \sum_{k=1}^{\bar{h}} \liminf_{i \rightarrow \infty} \mathcal{F}_{\varepsilon}^{\lambda}(\Omega_i^k) \leq \liminf_{i \rightarrow \infty} \sum_{k=1}^{\bar{h}} \mathcal{F}_{\varepsilon}^{\lambda}(\Omega_i^k) \\ &\leq \liminf_{i \rightarrow \infty} \tilde{\mathcal{F}}_{\varepsilon, \bar{h}}^{\lambda, R}(\Omega_i) \leq \inf \{ \mathcal{F}_{\varepsilon}^{\lambda}(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = m \}, \end{aligned}$$

so (5.16) is proved.

We can now conclude the proof of the theorem. Let  $\{G_i\}_{i \in \mathbb{N}}$  as in (5.9) and let us consider a subdivision  $G_i = G_i^1 \cup G_i^2 \cup \dots \cup G_i^{h'}$  such that

$$(5.20) \quad \inf \{ \tilde{\mathcal{F}}_{\varepsilon}^{\lambda}(\Omega) : \Omega \in \mathcal{A} \} = \lim_{i \rightarrow \infty} \sum_{j=1}^{h'} \mathcal{F}_{\varepsilon}^{\lambda}(G_i^j).$$

Up to a subsequence there exist  $\mu_j > 0$ ,  $1 \leq j \leq h'$ , such that

$$\mu_j = \lim_{i \rightarrow \infty} |G_i^j| \quad \forall 1 \leq j \leq h', \quad |B^{\lambda}| = \sum_{j=1}^{h'} \mu_j.$$

Let

$$K_j := \inf \{ \mathcal{F}_{\varepsilon}^{\lambda}(\Omega) : \Omega \subset \mathbb{R}^n \setminus H, |\Omega| = \mu_j \}.$$

By (5.20)

$$(5.21) \quad \inf \{ \tilde{\mathcal{F}}_{\varepsilon}^{\lambda}(\Omega) : \Omega \in \mathcal{A} \} = \sum_{j=1}^{h'} K_j.$$

By (5.16) for every  $1 \leq j \leq h'$  there exist  $\bar{h}(j) \in \mathbb{N}$ , a bounded set  $E_j \subset \mathbb{R}^n \setminus H$  with  $|E_j| = \mu_j$  and a subdivision in pairwise disjoint sets  $E_j = \bigcup_{k=1}^{\bar{h}(j)} E_{j,k}$  such that

$$(5.22) \quad \tilde{\mathcal{F}}_{\varepsilon}^{\lambda}(E_j) \leq \sum_{k=1}^{\bar{h}(j)} \mathcal{F}_{\varepsilon}^{\lambda}(E_{j,k}) \leq K_j.$$

Since the sets  $E_j$  are bounded, up to translations we can assume that the set  $E = \bigcup_{j=1}^{h'} E_j$  has volume  $|B^{\lambda}|$ . Therefore  $E$  is the disjoint union of all the sets  $E_{j,k}$  with  $1 \leq j \leq h'$  and  $1 \leq k \leq \bar{h}(j)$ . Let us denote these sets as  $E^l$  with

$1 \leq l \leq h$  and  $h = \sum_{j=1}^{h'} \bar{h}(j)$ . By (5.21) and (5.22) we deduce that

$$\tilde{\mathcal{F}}_\varepsilon^\lambda(E) \leq \sum_{l=1}^h \mathcal{F}_\varepsilon^\lambda(E^l) \leq \sum_{j=1}^{h'} K_j = \inf \left\{ \tilde{\mathcal{F}}_\varepsilon^\lambda(\Omega) : \Omega \in \mathcal{A} \right\},$$

that is  $E$  is a minimizer of  $\tilde{\mathcal{F}}_\varepsilon^\lambda$  and the subdivision  $E = \bigcup_{l=1}^h E^l$  is optimal.

The proof of (5.1) for a given  $1 \leq \bar{j} \leq h$  easily follows as in the proof of [NP21, Proposition 1.2].  $\square$

Now we are ready to prove Theorem 1.3.

*Proof of Theorem 1.3.* With the notation of Theorem 1.3, if  $\Omega \subset \mathbb{R}^n \setminus H$  is a measurable set with  $|\Omega| = m$ , let

$$\mathcal{F}^\lambda(\Omega) = P_\lambda(\Omega) + \int_\Omega \int_\Omega g(y-x) dy dx + \int_\Omega G(x_n) dx$$

and

$$\tilde{\mathcal{F}}^\lambda(\Omega) := \inf_{h \in \mathbb{N}} \tilde{\mathcal{F}}_h^\lambda(\Omega),$$

where

$$\tilde{\mathcal{F}}_h^\lambda(\Omega) := \inf \left\{ \sum_{i=1}^h \mathcal{F}^\lambda(\Omega^i) : \Omega = \bigcup_{i=1}^h \Omega^i, \Omega^i \cap \Omega^j = \emptyset \text{ for } 1 \leq i \neq j \leq h \right\}.$$

If  $F \subset \mathbb{R}^n \setminus H$  has measure  $|B^\lambda|$  and  $\bar{\varepsilon} = \frac{m^{\frac{1}{n}}}{|B^\lambda|^{\frac{1}{n}}}$ , the set  $\tilde{F} := \bar{\varepsilon}F$  has volume  $m$  and by Remark 5.2 there exist  $\tilde{g}$   $\mathcal{R}$ -admissible and  $\tilde{G}$   $\mathcal{G}$ -admissible such that

$$\mathcal{F}^\lambda(\tilde{F}) = \bar{\varepsilon}^{n-1} \left( P_\lambda(F) + \bar{\varepsilon}^{n+1} \int_F \int_F \tilde{g}(y-x) dy dx + \bar{\varepsilon} \int_F \tilde{G}(x_n) dx \right) =: \bar{\varepsilon}^{n-1} \tilde{\mathcal{F}}_\varepsilon^\lambda(F).$$

Note that  $\tilde{\mathcal{F}}^\lambda(\tilde{F}) = \bar{\varepsilon}^{n-1} \tilde{\mathcal{F}}_\varepsilon^\lambda(F)$  and that, if  $g$  is  $q$ -decreasing, also  $\tilde{g}$  is  $q$ -decreasing (see Remark 5.2). By Theorem 5.3 there exists  $E \subset \mathbb{R}^n \setminus H$  with  $|E| = |B^\lambda|$  which minimizes  $\tilde{\mathcal{F}}_\varepsilon^\lambda$ . Then the set  $\tilde{E} := \bar{\varepsilon}E$  minimizes  $\tilde{\mathcal{F}}^\lambda$  among sets with volume  $m$  and Theorem 1.3 easily follows.  $\square$

## REFERENCES

- [AFM13] E. Acerbi, N. Fusco, and M. Morini. “Minimality via second variation for a nonlocal isoperimetric problem”. In: *Commun. Math. Phys.* 322.2 (2013), pp. 515–557.
- [Ala+19] S. Alama, L. Bronsard, R. Choksi, and I. Topaloglu. “Droplet breakup in the liquid drop model with background potential”. In: *Commun. Contemp. Math.* 21.3 (2019), p. 23.
- [ACO09] G. Alberti, R. Choksi, and F. Otto. “Uniform energy distribution for an isoperimetric problem with long-range interactions”. In: *J. Am. Math. Soc.* 22 (2009), pp. 596–605.
- [AFP00] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000, pp. xviii+434.
- [BNO23] K. Bessas, M. Novaga, and F. Onoue. “On the shape of small liquid drops minimizing nonlocal energies”. In: *ESAIM: COCV* 29.86 (2023), p. 26.
- [Boh36] N. Bohr. “Neutron capture and nuclear constitution”. In: *Nature* 137 (1936), pp. 344–348.
- [BW39] N. Bohr and J. Wheeler. “The mechanism of nuclear fission”. In: *Phys. Rev.* 56.5 (1939), pp. 426–450.
- [BC14] M. Bonacini and R. Cristoferi. “Local and global minimality results for a nonlocal isoperimetric problem on  $\mathbb{R}^N$ ”. In: *SIAM J. Math. Anal.* 46.4 (2014), pp. 2310–2349.
- [BCT24] M. Bonacini, R. Cristoferi, and I. Topaloglu. “Stability of the ball for attractive-repulsive energies”. In: *SIAM J. Math. Anal.* 56 (2024), pp. 588–615.
- [BCT18] A. Burchard, R. Choksi, and I. Topaloglu. “Nonlocal shape optimization via interactions of attractive and repulsive potentials”. In: *Indiana Univ. Math. J.* 67.1 (2018), pp. 375–395.
- [Car23] D. Carazzato. “A note on some non-local variational problems”. In: *Rend. Lincei Mat. Appl.* 34 (2023), pp. 265–293.
- [CFP23] D. Carazzato, N. Fusco, and A. Pratelli. “Minimality of balls in the small volume regime for a general Gamow-type functional”. In: *Advances in Calculus of Variations* 16.2 (2023), pp. 503–515.
- [CP22] D. Carazzato and A. Pratelli. *On the existence and boundedness of minimizing measures for a general form of non-local energies*. 2022. arXiv: 2205.09412.
- [CPT23] D. Carazzato, A. Pratelli, and I. Topaloglu. *On the existence of minimizing sets for a weakly-repulsive non-local energy*. 2023. arXiv: 2307.01830.
- [CM75] C. M. Care and N. H. March. “Electron crystallization”. In: *Adv. Phys.* 24.1 (1975), pp. 101–116.

- [CDM16] J. A. Carrillo, M. Delgadino, and A. Mellet. “Regularity of local minimizers of the interaction energy via obstacle problems”. In: *Commun. Math. Phys.* 343.3 (2016), pp. 747–781.
- [CN17] A. Cesaroni and M. Novaga. “Isoperimetric problems for a nonlocal perimeter of Minkowski type”. In: *Geometric Flows 2* (2017), pp. 86–93.
- [CN18] A. Cesaroni and M. Novaga. “The isoperimetric problem for nonlocal perimeters”. In: *Discrete Contin. Dyn. Syst.* 11.3 (2018), pp. 425–440.
- [CK93] L. Q. Chen and A. G. Khachaturyan. “Dynamics of simultaneous ordering and phase separation and effect of long-range Coulomb interactions”. In: *Phys. Rev. Lett.* 70.10 (1993), pp. 1477–1480.
- [CGR07] J. Choe, M. Ghomi, and M. Ritoré. “The relative isoperimetric inequality outside convex domains in  $\mathbf{R}^n$ ”. In: *Calc. Var. Partial Differential Equations* 29.4 (2007), pp. 421–429.
- [CMT17] R. Choksi, C. B. Muratov, and I. Topaloglu. “An Old Problem Resurfaces Nonlocally: Gamow’s Liquid Drops Inspire Today’s Research and Applications”. In: *Notices of the AMS* (2017).
- [CNT22] R. Choksi, R. Neumayer, and I. Topaloglu. “Anisotropic liquid drop models”. In: *Adv. Calc. Var.* 15.1 (2022), pp. 109–131.
- [CP10] R. Choksi and M. A. Peletier. “Small volume fraction limit of the diblock copolymer problem: I. Sharp interface functional”. In: *SIAM J. Math. Anal.* 42 (2010), pp. 1334–1370.
- [CP11] R. Choksi and M. A. Peletier. “Small volume-fraction limit of the diblock copolymer problem: II. Diffuse interface functional”. In: *SIAM J. Math. Anal.* 43.2 (2011), pp. 739–763.
- [CL12] M. Cicalese and G. P. Leonardi. “A selection principle for the sharp quantitative isoperimetric inequality”. In: *Arch. Ration. Mech. Anal.* 206.2 (2012), pp. 617–643.
- [CS13] M. Cicalese and E. Spadaro. “Droplet minimizers of an isoperimetric problem with long-range interactions”. In: *Commun. Pure Appl. Math.* 66 (2013), pp. 1298–1333.
- [CPS74] S. Cohen, F. Plasil, and W. J. Swiatecki. “Equilibrium configurations of rotating charged or gravitating liquid masses with surface tension. II”. In: *Annals of Physics* 82.2 (1974), pp. 557–596.
- [CS62] S. Cohen and W. J. Swiatecki. “The deformation energy of a charged drop: IV. Evidence for a discontinuity in the conventional family of saddle point shapes”. In: *Annals of Physics* 19.1 (1962), pp. 67–164.
- [De 58] E. De Giorgi. “Sulla proprietà isoperimetrica dell’ipersfera, nella classe degli insiemi aventi frontiera orientata di misura finita”. In: *Atti Accad. Naz. Lincei Mem. Cl. Sci. Fis. Mat. Nat., Sez. I* 8 (1958), pp. 33–44.
- [EK93] V. J. Emery and S. A. Kivelson. “Frustrated electronic phase-separation and high-temperature superconductors”. In: *Physica C* 209.4 (1993), pp. 597–621.
- [EFK20] L. Emmert, R. L. Frank, and T. König. “Liquid drop model for nuclear matter in the dilute limit”. In: *SIAM J. Math. Anal.* 52.2 (2020), pp. 1980–1999.
- [Fig+15] A. Figalli, N. Fusco, F. Maggi, V. Millot, and M. Morini. “Isoperimetry and stability properties of balls with respect to nonlocal energies”. In: *Commun. Math. Phys.* 336 (2015), pp. 441–507.
- [Fra19] R. L. Frank. “Non-spherical equilibrium shapes in the liquid drop model”. In: *J. Math. Phys.* 60.071506 (2019).
- [FKN16] R. L. Frank, R. Killip, and P. T. Nam. “Nonexistence of large nuclei in the liquid drop model”. In: *Lett. Math. Phys.* 106 (2016), pp. 1033–1036.
- [FL15] R. L. Frank and E. H. Lieb. “A compactness lemma and its application to the existence of minimizers for the liquid drop model”. In: *SIAM J. Math. Anal.* 47.6 (2015), pp. 4436–4450.
- [FL18] R. L. Frank and E. H. Lieb. “A “liquid-solid” phase transition in a simple model for swarming, based on the “no flat-spots” theorem for subharmonic functions”. In: *Indiana Univ. Math. J.* 67.4 (2018), pp. 1547–1569.
- [FL19] R. L. Frank and E. H. Lieb. “Periodic energy minimizers for a one-dimensional liquid drop model”. In: *Lett. Math. Phys.* 109.9 (2019), pp. 2069–2081.
- [FL21] R. L. Frank and E. H. Lieb. “Proof of spherical flocking based on quantitative rearrangement inequalities”. In: *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 22.3 (2021), pp. 1241–1263.
- [FN21] R. L. Frank and P. T. Nam. “Existence and nonexistence in the liquid drop model”. In: *Calc. Var. Partial Differential Equations* 60.223 (2021).
- [FNV18] R. L. Frank, P. T. Nam, and H. Van Den Bosch. “The ionization conjecture in Thomas–Fermi–Dirac–von Weizsäcker theory”. In: *Commun. Pure Appl. Math.* 71.3 (2018), pp. 577–614.
- [Fug89] B. Fuglede. “Stability in the isoperimetric problem for convex or nearly spherical domains in  $\mathbf{R}^n$ ”. In: *Trans. Amer. Math. Soc.* 314.2 (1989), pp. 619–638.

- [FM23] N. Fusco and M. Morini. “Total positive curvature and the equality case in the relative isoperimetric inequality outside convex domains”. In: *Calc. Var. Partial Differential Equations* 62.3 (2023), Paper No. 102, 32.
- [Gam30] G. Gamow. “Mass defect curve and nuclear constitution”. In: *Proc. R. Soc. Lond. A* 126 (1930), pp. 632–644.
- [Gen79] P. G. de Gennes. “Effect of cross-links on a mixture of polymers”. In: *J. Physique Lett.* 40.4 (1979), pp. 69–72.
- [GDM95] S. C. Glotzer, E. A. Di Marzio, and M. Muthukumar. “Reaction-controlled morphology of phase-separating mixtures”. In: *Phys. Rev. Lett.* 74.11 (1995), pp. 2034–2037.
- [GMS13] D. Goldman, C. B. Muratov, and S. Serfaty. “The  $\Gamma$ -limit of the two-dimensional Ohta-Kawasaki energy. I. Droplet density”. In: *Arch. Rational Mech. Anal.* 210 (2013), pp. 581–613.
- [GMS14] D. Goldman, C. B. Muratov, and S. Serfaty. “The  $\Gamma$ -limit of the two-dimensional Ohta-Kawasaki energy. II. Droplet arrangement via the renormalized energy”. In: *Arch. Rational Mech. Anal.* 212 (2014), pp. 445–501.
- [Jul14] V. Julin. “Isoperimetric problem with a Coulombic repulsive term”. In: *Indiana Univ. Math. J.* 63 (2014), pp. 77–89.
- [Jul17] V. Julin. “Remark on a nonlocal isoperimetric problem”. In: *Nonlinear Analysis* 154 (2017), pp. 174–188.
- [Kin+13] J. Kinnunen, R. Korte, A. Lorent, and N. Shanmugalingam. “Regularity of sets with quasiminimal boundary surfaces in metric spaces”. In: *J. Geom. Anal.* 23.4 (2013), pp. 1607–1640.
- [KM13] H. Knüpfer and C. B. Muratov. “On an isoperimetric problem with a competing nonlocal term I. The planar case”. In: *Commun. Pure Appl. Math.* 66 (2013), pp. 1129–1162.
- [KM14] H. Knüpfer and C. B. Muratov. “On an isoperimetric problem with a competing nonlocal term II. The general case”. In: *Commun. Pure Appl. Math.* 67 (2014), pp. 1974–1994.
- [KMN16] H. Knüpfer, C. B. Muratov, and M. Novaga. “Low density phases in a uniformly charged liquid”. In: *Commun. Math. Phys.* 345 (2016), pp. 141–183.
- [KN86] V. F. Kovalenko and E. L. Nagaev. “Photoinduced magnetism”. In: *Sov. Phys. Uspekhi* 29.4 (1986), pp. 297–321.
- [LL01] E. H. Lieb and M. Loss. *Analysis*. 2nd ed. Vol. 14. Graduate Studies in Mathematics. American Mathematical Society, 2001.
- [Lop19] O. Lopes. “Uniqueness and radial symmetry of minimizers for a nonlocal variational problem”. In: *Commun. Pure Appl. Math.* 18.5 (2019), pp. 2265–2282.
- [LO14] J. Lu and F. Otto. “Nonexistence of minimizers for Thomas–Fermi–Dirac–vonWeizsäcker model”. In: *Commun. Pure Appl. Math.* 67.10 (2014), pp. 1605–1617.
- [Mag12] F. Maggi. *Sets of finite perimeter and geometric variational problems*. Vol. 135. Cambridge Studies in Advanced Mathematics. An introduction to geometric measure theory. Cambridge University Press, Cambridge, 2012, pp. xx+454.
- [Mam94] R. F. Mamin. “Domain structure of a new type near a photostimulated phase transition; autosolitons”. In: *JETP Lett.* 60 (1994), pp. 52–56.
- [MW21] A. Mellet and Y. Wu. “An isoperimetric problem with a competing nonlocal singular term”. In: *Calc. Var. Partial Differential Equations* 60.106 (2021).
- [MP22] B. Merlet and M. Pegon. “Large mass rigidity for a liquid drop model in 2D with kernels of finite moments”. In: *J. Éc. polytech. Math.* 9 (2022), pp. 63–100.
- [Mur02] C. B. Muratov. “Theory of domain patterns in systems with long-range interactions of Coulomb type”. In: *Phys. Rev. E* 66.3 (2002), pp. 1–25.
- [Mur10] C. B. Muratov. “Droplet phases in non-local Ginzburg-Landau models with Coulomb repulsion in two dimensions”. In: *Commun. Math. Phys.* 299 (2010), pp. 45–87.
- [MS19] C. B. Muratov and T. M. Simon. “A Nonlocal Isoperimetric Problem with Dipolar Repulsion”. In: *Commun. Math. Phys.* 372 (2019), pp. 1059–1115.
- [MZ14] C. B. Muratov and A. Zaleski. “On an isoperimetric problem with a competing non-local term: quantitative results”. In: *Ann. Global Anal. Geom.* 47 (2014), pp. 63–80.
- [MS96] W. D. Myers and W. J. Swiatecki. “Nuclear properties according to the Thomas-Fermi model”. In: *Nuclear Physics A* 601.2 (1996), pp. 141–167.
- [Nag95] E. L. Nagaev. “Phase separation in high-temperature superconductors and related magnetic systems”. In: *Phys. Uspekhi* 38.5 (1995), pp. 497–521.
- [Nam20] P. T. Nam. “The Ionization Problem”. In: *EMS Newsl.* 12 (2020), pp. 22–27.

- [NO22] M. Novaga and F. Onoue. “Existence of minimizers for a generalized liquid drop model with fractional perimeter”. In: *Nonlinear Analysis* 224.113078 (2022).
- [NO23a] M. Novaga and F. Onoue. *Local and Nonlocal Liquid Drop Models*. 2023. arXiv: [2303.03189](https://arxiv.org/abs/2303.03189).
- [NO23b] M. Novaga and F. Onoue. “Local Hölder regularity of minimizers for nonlocal variational problems”. In: *Commun. Contemp. Math.* 25.10 (2023), p. 2250058.
- [NP21] M. Novaga and A. Pratelli. “Minimisers of a general Riesz-type problem”. In: *Nonlinear Analysis* 209.112346 (Aug. 2021).
- [NKD94] I. A. Nyrkova, A. R. Khokhlov, and M. Doi. “Microdomain structures in polyelectrolyte systems: calculation of the phase diagrams by direct minimization of the free energy”. In: *Macromolecules* 27.15 (1994), pp. 4220–4230.
- [OK86] T. Ohta and K. Kawasaki. “Equilibrium morphology of block copolymer melts”. In: *Macromolecules* 19.10 (1986), pp. 2621–2632.
- [Ono22] F. Onoue. “Nonexistence of minimizers for a nonlocal perimeter with a Riesz and a background potential”. In: *Rend. Sem. Mat. Univ. Padova* 147 (2022), pp. 111–137.
- [PP24] G. Pascale and M. Pozzetta. *Quantitative isoperimetric inequalities for classical capillarity problems*. 2024. arXiv: [2402.04675](https://arxiv.org/abs/2402.04675).
- [Peg21] M. Pegon. “Large mass minimizers for isoperimetric problems with integrable nonlocal potentials”. In: *Nonlinear Analysis* 211.112395 (2021).
- [PTM90] N. A. Pelakasis, J. A. Tsamopoulos, and G. D. Manolis. “Equilibrium shapes and stability of charged and conducting drops”. In: *Phys. Fluids A* 2.8 (1990), pp. 1328–1340.
- [Rie30] F. Riesz. “Sur une inégalité intégrale”. In: *J. Lond. Math. Soc.* 5 (1930), pp. 162–168.
- [Rig00] S. Rigot. “Ensembles quasi-minimaux avec contrainte de volume et rectifiabilité uniforme”. In: *Mém. Soc. Math. Fr.* 2.82 (2000).
- [ST11] P. Sternberg and I. Topaloglu. “On the global minimizers of a nonlocal isoperimetric problem in two dimensions”. In: *Interfaces Free Bound.* 13 (2011), pp. 155–169.
- [Wei35] C. F. von Weizsäcker. “Zur theorie der Kernmassen”. In: *Zeitschrift für Physik* 96.7 - 8 (1935), pp. 431–458.

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI "RENATO CACCIOPPOLI", UNIVERSITÀ DEGLI STUDI DI NAPOLI "FEDERICO II",  
 VIA CINTIA - MONTE SANT'ANGELO, 80126 NAPOLI, ITALY  
 Email address: [giulio.pascale@unina.it](mailto:giulio.pascale@unina.it) 