

Identities and Generating Functions of Products of Generalized Fibonacci numbers, Catalan and Harmonic Numbers

Vladimir V. Kruchinin

Tomsk State University of Control Systems and Radioelectronics
Tomsk
Russian Federation
kru@ie.tusur.ru

Maria Y. Perminova

Tomsk State University of Control Systems and Radioelectronics
Tomsk
Russian Federation
pmy@fdo.tusur.ru

Abstract

We considered the properties of generalized Fibonacci and Lucas numbers class. The analogues of well-known Fibonacci identities for generalized numbers are obtained. We gained a new identity of product convolution of generalized Fibonacci and Lucas numbers. We wrote down generating functions of generalized Fibonacci and Lucas numbers products, their multisections, harmonic numbers and Catalan numbers

1 Introduction

The Fibonacci, Lucas, and Catalan numbers are among the most studied objects of combinatorics. There is a huge variety of publications and monographs [1, 2, 3, 4]. Studies on the products of such numbers, e.g., Fibonacci and Catalan numbers product have appeared relatively recently [16]. There are many different generalizations of Fibonacci and Lucas numbers [5, 6, 7, 8, 9, 10]. Kalman and R. Mena proposed a generalization of Fibonacci numbers of the form

$$F(n) = \begin{cases} 0, & n \leq 0, \\ 1, & n = 1, \\ aF(n-1) + bF(n-2), & n > 1. \end{cases} \quad (1)$$

where $a, b \in \mathbb{R}$. The generating function for generalized Fibonacci numbers is

$$F(x) = \frac{x}{1 - ax - bx^2}.$$

a	b	$F_{a,b}(n)$	$L_{a,b}(n)$
1	1	<u>A000045</u>	<u>A000032</u>
2	1	<u>A000129</u>	<u>A002203</u>
3	1	<u>A006190</u>	<u>A006497</u>
4	1	<u>A001076</u>	<u>A014448</u>
5	1	<u>A052918</u>	<u>A087130</u>
6	1	<u>A005668</u>	<u>A085447</u>
7	1	<u>A054413</u>	<u>A086902</u>
8	1	<u>A041025</u>	<u>A086902</u>
9	1	<u>A099371</u>	<u>A087798</u>
1	2	<u>A001045</u>	<u>A014551</u>
1	3	<u>A006130</u>	<u>A075118</u>
2	2	<u>A002605</u>	<u>A080040</u>
2	3	<u>A015518</u>	<u>A102345</u>
3	2	<u>A007482</u>	<u>A206776</u>
3	3	<u>A030195</u>	<u>A172012</u>

Table 1: Generalized Fibonacci and Lucas numbers written in OEIS

They also introduced generalized Lucas numbers

$$L(n) = F(n+1) + bF(n-1).$$

The generating function for generalized Lucas numbers is

$$L(x) = \frac{2 - ax}{1 - ax - bx^2}.$$

They presented the identity accomplished for generalized Fibonacci numbers

$$F(n+m) = bF(n-1)F(m) + F(n)F(m+1). \quad (2)$$

We note if $a, b \in \mathbb{Z}$ are integers. Then $F(n)$ and $L(n)$ are Lucas sequences [19]. When setting the parameters values a and b , we obtain a particular sequence. Then fixing the parameters a and b , we explicitly indicate their value, e.g., $a = 1, b = 1, F_{1,1}(n)$ are Fibonacci numbers, $L_{1,1}(n)$ are Lucas numbers, $a = 2, b = 1, F_{2,1}(n)$ are Pell numbers and $L_{2,1}(n)$ are Pell-Lucas numbers. Table 1 presents the sequences $F(n)$ and $L(n)$ for different values a and b , written in OEIS [13].

We write down a number of identities to obtain generating functions in the following sections.

Proposition 1. *We accomplish the identity for generalized Fibonacci and Lucas numbers*

$$2F(n+m) = F(m)L(n) + L(m)F(n). \quad (3)$$

Proof. We use the identity (2)

$$F(n+m) = bF(n-1)F(m) + F(n)F(m+1).$$

and

$$F(n+m) = bF(n)F(m-1) + F(n+1)F(m).$$

We add

$$\begin{aligned} 2F(n+m) &= bF(n)F(m-1) + F(n+1)F(m) + bF(n-1)F(m) + F(n)F(m+1), \\ 2F(n+m) &= [bF(n)F(m-1) + F(n)F(m+1)] + [F(n+1)F(m) + bF(n-1)F(m)]. \end{aligned}$$

We group

$$2F(n+m) = F(n)[bF(m-1) + F(m+1)] + F(m)[F(n+1) + bF(n-1)].$$

We make a replacement

$$L(n) = F(n+1) + bF(n-1).$$

We obtain the sought-for formula

$$2F(n+m) = F(m)L(n) + L(m)F(n).$$

□

The consequence of this statement at $n = m$ is the identity

$$F(2n) = F(n)L(n). \tag{4}$$

We accomplish Johnson's identity for generalized Fibonacci numbers (2003) [12].

Proposition 2. *We accomplish the identity for all $p, q, c, d \in \mathbb{N}$, $r \in \mathbb{Z}$, at $p + q = c + d$*

$$F(p)F(q) - F(c)F(d) = (-b)^r (F(p-r)F(q-r) - F(c-r)F(d-r)). \tag{5}$$

Proof. We use the identity (2)

$$\begin{aligned} F(p+q-1) &= bF(p-1)F(q-1) + F(p)F(q), \\ F(c+d-1) &= bF(c-1)F(d-1) + F(c)F(d). \end{aligned}$$

Then

$$bF(p-1)F(q-1) + F(p)F(q) = bF(c-1)F(d-1) + F(c)F(d).$$

We transform to

$$F(p)F(q) - F(c)F(d) = -b(F(p-1)F(q-1) - F(c-1)F(d-1)).$$

Applying the identity r times for the right side, we obtain the sought-for identity

$$F(p)F(q) - F(c)F(d) = (-b)^r (F(p-r)F(q-r) - F(c-r)F(d-r)).$$

□

We accomplish the multiple-angle recurrence identity for generalized Fibonacci numbers.

Proposition 3. *We accomplish the identity for all $a, b \in \mathbb{R}$*

$$F(mn + j) = L(m)F(m(n - 1) + j) - (-b)^m F(m(n - 2) + j), \quad (6)$$

where $L(m)$ are generalized Fibonacci numbers.

Proof. We write down generalized Johnson's identity (5) $p = m(n - 1) + j$, $q = m + 1$, $c = m(n - 1) + j + 1$, $d = m$, $r = m$. We obtain

$$F(m(n - 2) + m + j)F(m + 1) - F(m(n - 2) + m + j + 1)F(m) = (-b)^m F(m(n - 2) + j),$$

$$F(m(n - 2) + m + j + 1)F(m) = F(m(n - 2) + m + j)F(m + 1) - (-b)^m F(m(n - 2) + j).$$

We add to the left and to the right parts

$$bF(m(n - 1) + j)F(m - 1).$$

We obtain

$$\begin{aligned} & bF(m(n - 1) + j)F(m - 1) + F(m(n - 1) + j + 1)F(m) = \\ & F(m(n - 1) + j)F(m + 1) + bF(m(n - 1) + j)F(m - 1) - (-b)^m F(m(n - 2) + j). \end{aligned}$$

Whence the left part of the identity is equal to $F(mn + j)$. In the right part the first two summands are equal to $L(m)F(m(n - 1) + j)$. Then we obtain the sought-for recurrence formula

$$F(mn + j) = L(m)F(m(n - 1) + j) - (-b)^m F(m(n - 2) + j).$$

□

Proposition 4. *We accomplish the identity for generalized Fibonacci numbers*

$$F(n + j)F(m) - F(j)F(n + m) = (-1)^j b^j F(n)F(m - j). \quad (7)$$

Proof. Using Johnson identity (5) at $p = n + j$, $q = m$, $c = j$, $d = n + m$, $r = j$ we obtain the sought-for identity

$$F(n + j)F(m) - F(j)F(n + m) = (-1)^j b^j F(n)F(m - j).$$

□

2 The identity for generalized Fibonacci and Lucas numbers

We consider the property of numbers sequences $F(n)$ and $L(n)$. Let we have some convolution of the form

$$p(n) = \sum_{i=0}^n r(i) r(n - i),$$

where $r(n) \in R$.

We prove the following theorem.

Theorem 5. We accomplish the identity for numbers sequences $F(n)$, $L(n)$ and $p(n)$, $k, m \in \mathbb{N}$

$$p(n) F(kn + 2m) = \sum_{i=0}^n r(i)L(ki + m)r(n - i)F(k(n - i) + m). \quad (8)$$

Proof. We obtain the identity (8) at $n = 0$

$$p(0) F(0 + 2m) = p(0) L(m)F(m).$$

Whence we obtain the equality based on the identity (4). We now consider the set of pairs of terms in the right part (8) such that

$$r(i)r(n - i) = r(n - i)r(i).$$

Then for the sum

$$r(i)L(ki + m)r(n - i)F(k(n - i) + m) + r(n - i)L(k(n - i) + m)r(i)F(ki + m)$$

we accomplish

$$r(i)r(n - i) [L(ki + m)F(k(n - i) + m) + L(k(n - i) + m)F(ki + m)].$$

We write down on the basis of the identity (3)

$$L(ki + m)F(k(n - i) + m) + L(k(n - i) + m)F(ki + m) = 2F(ki + m + k(n - i) + m) = 2F(kn + 2m).$$

The set of pairs of terms in the right part (8) at odd n is even and the obtained identity is accomplished for all pairs. The set of pairs of terms in the right part (8) at even n is odd. We consider $n = j + j$. Then

$$r(j)L(kj + m)r(j)F(kj + m) = r(j)r(j)L(kj + m)F(kj + m).$$

We obtain on the basis of the identity (4)

$$L(kj + m)F(kj + m) = F(2kj + 2m) = F(kn + 2m).$$

Then we can write down the right part of the identity (8) as

$$\sum_{i=0}^n r(i)L(ki + m)r(n - i)F(k(n - i) + m) = F(kn + 2m) \sum_{i=0}^n r(i)r(n - i).$$

Knowing that

$$p(n) = \sum_{i=0}^n r(i)r(n - i)$$

whence we obtain

$$p(n) F(kn + 2m) = \sum_{i=0}^n r(i)L(ki + m)r(n - i)F(k(n - i) + m).$$

We complete the proof. □

We consider the examples of using the identity. Since $r(n) = 1$, and then the identity (8) is

$$(n+1)F(kn+2m) = \sum_{i=0}^n L(ki+m)F(k(n-i)+m).$$

We obtain at $m = n$

$$(n+1)F((k+1)n) = \sum_{i=0}^n L(ki+n)F(k(n-i)+n).$$

We now consider the case $r(n) = F(n)$. We substitute into the identity (8)

$$p(n)F(n+2m) = \sum_{i=0}^n F(i)L(i+m)F(n-i)F(n-i+m).$$

where

$$p(n) = \sum_{i=0}^n F(i)F(n-i).$$

Hoggart obtained the following explicit expression for the convolution of Fibonacci numbers [14]

$$\sum_{i=0}^n F_{1,1}(i)F_{1,1}(n-i) = \frac{1}{5}((n-1)F_{1,1}(n) + (n+1)F_{1,1}(n-1)).$$

Using his method we can obtain the convolution formula for generalized Fibonacci numbers

$$\sum_{i=0}^n F(i)F(n-i) = \frac{(n-1)F(n+1) + b(n+1)F(n-1)}{a^2 + 4b}. \quad (9)$$

Whence at $m = 0$, $k = 1$, $a = 1$, $b = 1$ we obtain the identity

$$\sum_{i=0}^n F_{1,1}(i)^2 L_{1,1}(n-i)F_{1,1}(n-i) = F_{1,1}(n) \frac{(n-1)F_{1,1}(n+1) + (n+1)F_{1,1}(n-1)}{5}.$$

For Pell numbers we have $m = 0$, $k = 1$, $a = 2$, $b = 1$

$$\sum_{i=0}^n F_{2,1}(i)^2 L_{2,1}(n-i)F_{2,1}(n-i) = F_{2,1}(n) \frac{(n-1)F_{2,1}(n+1) + (n+1)F_{2,1}(n-1)}{8}.$$

For Jacobsthal numbers we have $m = 0$, $k = 1$, $a = 1$, $b = 2$

$$\sum_{i=0}^n F_{1,2}(i)^2 L_{1,2}(n-i)F_{1,2}(n-i) = F_{1,2}(n) \frac{(n-1)F_{1,2}(n+1) + (n+1)F_{1,2}(n-1)}{9}.$$

We can replace the property $r(i)r(n-i) = r(n-i)r(i)$ by the property $T(n, k) = T(n, n-k)$ for a triangle $T(n, k)$, the triangle is symmetric in relation to main diagonal. Then the identity (8) is

$$F(kn+2m) \sum_{k=0}^n T(n, k) = \sum_{k=0}^n T(n, k)F(ki+m)L(k(n-i)+m).$$

We write down the identity for Pascal's triangle

$$2^n F(kn + 2m) = \sum_{k=0}^n \binom{n}{k} F(ki + m)L(k(n - i) + m).$$

Similarly,

$$\binom{2n}{n} F(kn + 2m) = \sum_{k=0}^n \binom{n}{k}^2 F(ki + m)L(k(n - i) + m).$$

The identity for Euler numbers of the first kind $E_1(n, k)$

$$(n + 1)! F(kn + 2m) = \sum_{k=0}^n E_1(n, k) F(ki + m)L(k(n - i) + m).$$

Leibniz's harmonic triangle, Losanitsch's triangle, A027907 triangle of trinomial coefficients and others have such properties [13]. The obtained identity (8) allows us to write down the functional equation of the form

$$A(x) = B(x)C(x),$$

where

$$A(x) = \sum_{n=0}^{\infty} p(n) F(kn + 2m)x^n,$$

$$B(x) = \sum_{n=0}^{\infty} r(n) F(kn + m)x^n,$$

$$C(x) = \sum_{n=0}^{\infty} r(n) L(kn + m)x^n.$$

Now knowing two out of the three functions we can find the third function. We consider obtaining generating functions based on this equation.

3 Generating functions for the product of generalized Fibonacci and harmonic numbers

We consider the case $r(n) = \frac{1}{n}$, $m = 0$, $k = 1$. Then the identity (8) is

$$p(n)F_1(n) = \sum_{i=0}^{n-1} \frac{L(i)}{i} \frac{F(n-i)}{(n-i)},$$

where $F(n)$ are generalized Fibonacci numbers, $L(n)$ are adjoined numbers

$$p(n) = \sum_{i=1}^{n-1} \frac{1}{i(n-i)} = \frac{2}{n} H(n-1),$$

where $H(n)$ are harmonic numbers. We find the generating function for $p(n)F(n)$. We note that

$$p(n)F(n) = [x^n]U_1(x) = [x^n]U_2(x)U_3(x),$$

where

$$U_2(x) = \sum_{n=1}^{\infty} \frac{F(n)}{n} x^n,$$

$$U_3(x) = \sum_{n=1}^{\infty} \frac{L(n)}{n} x^n.$$

Then

$$U_2(x, a, b) = \int \frac{1}{(1 - ax - bx^2)} dx = \frac{1}{\sqrt{4b + a^2}} \log \left(\frac{-\sqrt{4b + a^2}x + ax - 2}{\sqrt{4b + a^2}x + ax - 2} \right),$$

$$U_3(x, a, b) = \int \frac{2 - ax}{(1 - ax - bx^2)} dx = -\log(1 - ax - bx^2).$$

Then the sought-for generating function is

$$U_1(x, a, b) = \frac{1}{\sqrt{a^2 + 4b}} \log \left(\frac{2 - \sqrt{a^2 + 4b}x - ax}{2 + \sqrt{a^2 + 4b}x - ax} \right) \log(1 - ax - bx^2).$$

And

$$\frac{2}{n} H(n-1) F(n) = [x^n]U_1(x, a, b).$$

Hence we can write down

$$H(n-1)F(n) = [x^n] \frac{x dU_1(x, a, b)}{2 dx}.$$

Then the generating function for the product $H(n)F(n+1)$ is

$$U_4(x, a, b) = \frac{\left((a + 2bx) \log \left(\frac{2 - \sqrt{a^2 + 4b}x - ax}{2 + \sqrt{a^2 + 4b}x - ax} \right) + \sqrt{a^2 + 4b} \log(1 - ax - bx^2) \right)}{2\sqrt{a^2 + 4b}(1 - ax - bx^2)}.$$

We now find the generating function $U_5(x, a, b)$ for $H(n)F(n)$. We write down

$$H(n)F(n) = H(n-1)F(n) + \frac{1}{n}F(n).$$

Whence

$$H(n)F(n) = [x^n] \frac{x dU_1(x, a, b)}{2 dx} + U_2(x, a, b).$$

Then the generating function for the product $H(n)F(n)$ is

$$U_5(x, a, b) = -\frac{\frac{(2-ax)}{\sqrt{4b+a^2}} \log \left(\frac{2-\sqrt{a^2+4b}x-ax}{2+\sqrt{a^2+4b}x-ax} \right) + x \log(1 - ax - bx^2)}{2(1 - ax - bx^2)}.$$

For example, we obtain at $a = 1$ and $b = 1$

$$H(n)F_{1,1}(n) = [x^n] \frac{5x \log(1-x-x^2) + \sqrt{5}(2-x) \log\left(-\frac{(\sqrt{5}+1)x-2}{(\sqrt{5}-1)x+2}\right)}{10(x^2+x-1)}.$$

The given sequence is written in A372199. The presented series converges at $x = \frac{1}{2}$. Hence it follows that

$$\sum_{n=1}^{\infty} \frac{H(n)F_1(n)}{2^n} = \frac{3}{\sqrt{5}} \log\left(\frac{3+\sqrt{5}}{3-\sqrt{5}}\right) + \log 4.$$

We obtain the generating function for the product of harmonic numbers and Pell numbers at $a = 2$ and $b = 1$

$$H(n)F_{2,1}(n) = [x^n] \frac{2x \log(1-2x-x^2) + \sqrt{2}(1-x) \log\left(\frac{1-(\sqrt{2}+1)x}{1+(\sqrt{2}-1)x}\right)}{4(x^2+2x-1)}.$$

The given sequence is written in A372210. The presented series converges at $x = \frac{1}{3}$. Hence it follows that

$$\sum_{n=1}^{\infty} \frac{1}{3^n} H(n)F_{2,1}(n) = -\frac{3}{4} \left(\sqrt{2} \log\left(\frac{2-\sqrt{2}}{2+\sqrt{2}}\right) + \log\left(\frac{2}{9}\right) \right).$$

The obtained generating functions $U_4(x, a, b)$ and $U_5(x, a, b)$ allow us to write down the generating function for the product $H(n)F(n+j+1)$, $j \geq 0$. For this purpose we use the identity (2)

$$H(n)F(n+j+1) = F(j+1)H(n)F(n+1) + bF(j)H(n)F(n).$$

Whence the generating function has the expression

$$U_6(x, a, b) = F(j+1)U_5(x, a, b) + bF(j)U_4(x, a, b). \quad (10)$$

4 Generating functions for the product of generalized Fibonacci and Catalan numbers

We consider the application of the obtained identity (8) at $r(n) = C(n)$. Where $C(n)$ are Catalan numbers

$$C(n) = \frac{1}{n+1} \binom{2n}{n}.$$

Then the identity is

$$C(n+1)F(n+2m) = \sum_{i=0}^n C(i)L(i+m)C(n-i)F(n-i+m). \quad (11)$$

We write down the known recurrence relation

$$C(n+1) = \sum_{i=0}^n C(i)C(n-i), C(0) = 1.$$

We write down the result obtained by Barry and Mwafise [16]

$$C(n)p(n+1) = [x^n]V_1(x, a, b) = [x^n]\frac{1}{x}\sqrt{\frac{1-2ax-\sqrt{1-4ax-16bx^2}}{(2(a^2+4b))}}, \quad (12)$$

where

$$p(n) = [x^n]\frac{1}{1-ax-bx^2}.$$

We write down the generating function

$$V_2(x, a, b) = \sum_{n=0}^{\infty} C(n) L(n+1)x^n.$$

Corollary 6. *The generating function for the product of generalized Lucas and Catalan numbers of the form*

$$C(n) L(n+1)$$

has the following expression

$$V_2(x, a, b) = \frac{1}{x} \left(1 - \sqrt{\frac{1-2ax+\sqrt{1-4ax-16bx^2}}{2}} \right). \quad (13)$$

Proof. We obtain the recurrence equation based on the identity (8) at $m = 1$

$$C(n+1)F(n+2) = \sum_{i=0}^n C(i)L(i+1)C(n-i)F(n-i+1).$$

Then we can write down the functional equation linking the generating functions $V_1(x, a, b)$ and $V_2(x, a, b)$

$$V_1(x, a, b) = 1 + x V_1(x, a, b) V_2(x, a, b).$$

Whence we find the explicit expression for the sought-for generating function

$$V_2(x, a, b) = \frac{1}{x} \left(1 - \frac{1}{V_1(x, a, b)} \right).$$

□

Corollary 7. *The generating function for the product of generalized Fibonacci and Catalan numbers $C(n)F(n)$ has the following expression*

$$V_3(x, a, b) = \frac{1}{2bx} \left(1 - \sqrt{\frac{2b\sqrt{-16bx^2-4ax+1}+4abx+2b+a^2}{4b+a^2}} \right).$$

Proof. By definition we have

$$L(n+1) = F(n+2) + bF(n) = aF(n+1) + 2bF(n).$$

Then

$$C(n)L(n+1) = aC(n)F(n+1) + 2bF(n)C(n)$$

$$V_2(x, a, b) = aV_1(x, a, b) + 2bV_3(x, a, b)$$

$$V_3(x, a, b) = \frac{1}{2b} (V_2(x, a, b) - aV_1(x, a, b)).$$

Hence knowing the generating functions for the right part terms of the expression we obtain the sought-for expression. \square

Then for the product of Fibonacci and Catalan numbers the generating function is

$$F_1(n)C(n) = [x^n] \frac{1}{2x} \left(1 - \frac{\sqrt{2\sqrt{1-4x-16x^2}+4x+3}}{\sqrt{5}} \right).$$

We write the generating function in A119694. At $x = \frac{1}{8}$, the series converges. Then

$$\sum_{n=1}^{\infty} \frac{1}{2^{3n}} C(n)F_1(n) = 4 \left(1 - \frac{3}{\sqrt{10}} \right).$$

For the product of Pell and Catalan numbers the generating function is

$$F_2(n)C(n) = [x^n] \frac{1}{2x} \left(1 - \frac{\sqrt{2\sqrt{1-8x-16x^2}+8x+6}}{2\sqrt{2}} \right).$$

We write the sequence in A372216. At $x = \frac{1}{16}$ the series converges. Then

$$\sum_{n=1}^{\infty} \frac{1}{2^{4n}} C(n)F_2(n) = 8 - 2\sqrt{13 + \sqrt{7}}.$$

The generating functions $V_1(x, a, b)$ and $V_3(x, a, b)$ allow us to write down the generating function for the product $C(n)F(n+j+1)$, $j \geq 0$. For this purpose we use the identity (2). Whence the generating function has the expression

$$V_4(x, a, b) = F(j+1)V_1(x, a, b) + bF(j)V_3(x, a, b). \quad (14)$$

4.1 Generating functions for multisections product of generalized Fibonacci and Lucas numbers, Catalan and harmonic numbers

The generating functions for multisections of Fibonacci and Lucas numbers are known [17, 18]. We find the expression of the generating function for multisections of generalized Fibonacci numbers of the form $F(mn+j)$. For this purpose we prove the following proposition

Proposition 8. For multisections of generalized Fibonacci numbers $F(mn+j)$, $m, n, j \in N$, $a, b \in R$ the generating function is

$$Q(x) = \frac{F(j) + (-b)^j F(m-j)x}{1 - L(m)x + (-b)^m x^2}, \quad (15)$$

where $L(m)$ is generalized Lucas number with parameters a, b .

Proof. We consider the obtained multiple-angle recurrence (6) at $j = 0$

$$F(mn) = L(m)F(m(n-1)) - (b)^m F(m(n-2)).$$

We easily see the generating function is

$$F(mn) = [x^n] \frac{F(m)x}{1 - L(m)x + (-b)^m x^2}.$$

We now use the identity (6). Then we select $F(n+j)$ in the left part of $F(n+j)$ and replace the argument n by mn

$$F(mn+j) = \frac{F(j)}{F(m)} F(m(n+1)) + \frac{(-b)^j F(m-j)}{F(m)} F(mn).$$

Whence

$$F(mn+j) = [x^n] \frac{F(j)}{F(m)} \frac{F(m)x}{x(1 - L(m)x + (-b)^m x^2)} + \frac{(-b)^j F(m-j)}{F(m)} \frac{F(m)x}{(1 - L(m)x + (-b)^m x^2)}.$$

Then we obtain the sought-for formula after simplification. \square

We use the relation between generalized Fibonacci and Lucas numbers. We obtain the generating function for generalized Lucas numbers

$$Q_L(x) = \frac{L(j) - (-b)^j L(m-j)x}{1 - L(m)x + (-b)^m x^2}. \quad (16)$$

We note that at $j = 0$ multisection of generalized Fibonacci numbers of the form $F(mn)$ belongs to generalized Fibonacci numbers with parameters $a = L(m)$, $b = -(-b')^m$, where a' and b' are parameters of the sequence $F(n)$ generating the multisection, $L(m)$ are generalized Lucas numbers with parameters a' and b' . Then we can use all identities valid for generalized Fibonacci numbers for multisections of the form $F(mn)$. Then we can use the function $U_5(x, a, b)$ to determine the generating functions of multisections product of generalized Fibonacci numbers of the form $F(mn)$ and harmonic numbers $H(n)$. We can use the function $V_3(x, a, b)$ to determine the generating functions of multisections product of generalized Fibonacci numbers of the form $F(mn)$ and Catalan numbers $H(n)$. We cannot use the proposed approach for multisections of generalized Lucas numbers since $L(0) \neq 0$. We consider some examples.

1. Let $a = 2$, $b = 1$, $F_{2,1}(n)$ are Pell numbers, $j = 0$ and $m = 2$. Then $F_{2,1}(2) = 2$, $L_{2,1}(2) = 6$, the generating function for bisection of Pell numbers is

$$F_{2,1}(2n) = [x^n] \frac{2x}{1 - 6x - x^2}.$$

Whence

$$H(n)F_{2,1}(2n) = [x^n] 2U_5(x, 6, -1),$$

$$C(n)F_{2,1}(2n) = [x^n] 2V_3(x, 6, -1).$$

2. Let $a = 1$, $b = 2$, $F_{1,2}(n)$ are Jacobsthal numbers, $j = 0$ and $m = 3$. Then $F_{1,2}(3) = 3$, $L_{2,1}(3) = 7$ the generating function for trisection of Jacobsthal numbers is

$$F_{2,1}(3n) = [x^n] \frac{3x}{1 - 7x - 8x^2}.$$

Whence

$$H(n)F_{1,2}(3n) = [x^n] 3U_5(x, 7, 8)$$

$$C(n)F_{1,2}(3n) = [x^n] 3V_3(x, 7, 8)$$

We consider obtaining generating functions for the product $H(n)F(mn+j)$ and $C(n)F(mn+j)$ at $0 \leq j < m$. For this purpose we use the identity (7) and replace the variable n by nm . We obtain

$$F(mn+j)F(m) - F(j)F(mn+m) = (-1)^j b^j F(mn)F(m-j).$$

Then

$$F(mn+j) = \frac{F(j)}{F(m)} F(m(n+1)) + \frac{(-1)^j b^j F(m-j)}{F(m)} F(mn).$$

We multiply the left and right parts by $H(n)$

$$H(n)F(mn+j) = \frac{F(j)}{F(m)} H(n)F(m(n+1)) + \frac{(-1)^j b^j F(m-j)}{F(m)} H(n)F(mn).$$

We now proceed to the generating functions

$$H(n)F(m(n+1)) = [x^n] F(m)U_4(x, L(m), -(-b)^m),$$

and

$$H(n)F(mn) = [x^n] F(m)U_5(x, L(m), -(-b)^m).$$

After simplification we obtain

$$H(n)F(mn+j) = [x^n] F(j)U_4(x, L(m), -(-b)^m) + (-b)^j F(m-j)U_5(x, L(m), -(-b)^m).$$

We perform similar actions for the product $C(n)F(mn+j)$ and write down the generating function

$$C(n)F(mn+j) = [x^n] F(j)V_1(x, L(m), -(-b)^m) + (-b)^j F(m-j)V_3(x, L(m), -(-b)^m).$$

We use the formula for generalized Lucas numbers

$$L(mn + j) = F(mn + j + 1) + bF(mn + j - 1)$$

After simplification we obtain

$$H(n)L(mn + j) = [x^n]L(j)U_4(x, L(m), -(-b)^m) - (-b)^j L(m - j)U_5(x, L(m), -(-b)^m)$$

and

$$C(n)L(mn + j) = [x^n]L(j)V_1(x, L(m), -(-b)^m) - (-b)^j L(m - j)V_3(x, L(m), -(-b)^m).$$

We consider the examples.

1. We specify Jacobsthal numbers $a = 1, b = 2, j = 3, m = 4$. Then the generating function for the product $H(n)F_{1,2}(4n + 1)$ is

$$H(n)F_{1,2}(4n + 3) = [x^n] \frac{(8x + 7) \log\left(\frac{x-1}{16x-1}\right) + (24x - 9) \log(16x^2 - 17x + 1)}{96x^2 - 102x + 6}.$$

2. We specify the sequence of numbers A010892, $m = 4, j = 1, a = 1, b = -1$. Then the product $(n)F_{1,-1}(4n + 1)$ has the generating function

$$C(n)F_{1,-1}(4n + 1) = [x^n] \frac{\sqrt{\sqrt{16x^2 + 4x + 1} - 2x - 1}}{\sqrt{6}x}.$$

3. We specify the sequence of Jacobsthal-Lucas numbers A014551, $m = 4, j = 1, a = 1, b = -1$. Then the product $H(n)L_{1,2}(2n)$ has the generating function

$$H(n)L_{1,2}(2n) = [x^n] \frac{3x \log\left(\frac{x-1}{4x-1}\right) + (5x - 2) \log(4x^2 - 5x + 1)}{8x^2 - 10x + 2}.$$

4. We specify the sequence A002249, $a = 1, b = -2, j = 1, m = 3$. Then the product $C(n)L_{1,-2}(2n)$ has the generating function

$$C(n)L_{1,-2}(3n + 1) = [x^n] \frac{\sqrt{\sqrt{128x^2 + 16x + 1} - 8x - 1}}{\sqrt{8}x}.$$

5 Conclusion

1. We wrote down the identities for generalized Fibonacci and Lucas numbers.
2. We obtained the new identity for generalized Fibonacci and Lucas numbers

$$p(n)F(kn + 2m) = \sum_{i=0}^n r(i)L(ki + m)r(n - i)F(k(n - i) + m),$$

where

$$p(n) = \sum_{i=0}^n r(i)r(n - i).$$

3. We obtained the generating functions for the products of the form

$$\begin{aligned} H(n)F(n+1) &= [x^n]U_4(x, a, b), \\ H(n)F(n) &= [x^n]U_5(x, a, b), \\ C(n)F(n) &= [x^n]V_3(x, a, b). \end{aligned}$$

4. We obtained the generating functions for the products of the form

$$\begin{aligned} H(n)F(n+j+1) &= [x^n]F(j+1)U_5(x, a, b) + bF(j)U_4(x, a, b), \\ C(n)F(n+j+1) &= [x^n]F(j+1)V_1(x, a, b) + bF(j)V_3(x, a, b). \end{aligned}$$

5. We obtained the generating functions for multisections of generalized Fibonacci and Lucas numbers

$$\begin{aligned} Q_F(x) &= \frac{F(j) + (-b)^j F(m-j)x}{1 - L(m)x + (-b)^m x^2}, \\ Q_L(x) &= \frac{L(j) - (-b)^j L(m-j)x}{1 - L(m)x + (-b)^m x^2}. \end{aligned}$$

6. We obtained the generating functions for multisections product of generalized Fibonacci and Lucas numbers, Catalan and harmonic numbers

$$\begin{aligned} H(n)F(mn+j) &= [x^n]F(j)U_4(x, L(m), -(-b)^m) + (-b)^j F(m-j)U_5(x, L(m), -(-b)^m), \\ C(n)F(mn+j) &= [x^n]F(j)V_1(x, L(m), -(-b)^m) + (-b)^j F(m-j)V_3(x, L(m), -(-b)^m), \\ H(n)L(mn+j) &= [x^n]L(j)U_4(x, L(m), -(-b)^m) - (-b)^j L(m-j)U_5(x, L(m), -(-b)^m), \\ C(n)L(mn+j) &= [x^n]L(j)V_1(x, L(m), -(-b)^m) - (-b)^j L(m-j)V_3(x, L(m), -(-b)^m). \end{aligned}$$

References

- [1] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, Wiley, New York, 2001.
- [2] D. B. Andrica and O. Bagdasar, *Recurrent Sequences: Key Results, Applications, and Problems*, Springer Nature, 2020.
- [3] R. Honsberger, *A Second Look at the Fibonacci and Lucas Numbers*. Ch. 8 in *Mathematical Gems III*, Math. Assoc. America, Washington, DC, 1985.
- [4] R. Stanley, *Catalan Numbers*, Cambridge Univ. Press, 2015.
- [5] M. Bicknell-Johnson and C. P. Spears, *Classes of Identities for the Generalized Fibonacci Numbers $G_n = G_{(n-1)} + G_{(n-c)}$ for Matrices with Constant Valued Determinants*, *Fib. Quart.* 34 (1996), 121–128.
- [6] A. Dujella, *Generalized Fibonacci Numbers and the Problem of Diophantus*, *Fib. Quart.* 34, 1996, 164–175.

- [7] A. F. Horadam, Generating Functions for Powers of a Certain Generalized Sequence of Numbers, *Duke Math. J.* 32, 1965, 437–446.
- [8] S. Falcon and A. Plaza, On the Fibonacci k-numbers, *Chaos, Solitons Fractals*, 32 (5), 2007, 1615–1624.
- [9] M. Edson and O. Yayenie, A New Generalization of Fibonacci Sequences and Extended Binet’s Formula, *Integers* 9, 2009, 639–654.
- [10] Y. Simsek, Construction of General Forms of Ordinary Generating Functions for more Families of Numbers and Multiple Variables Polynomials. *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* Vol. 117, art. num. 130, 2023.
- [11] D. Kalman and R. Mena, The Fibonacci Numbers-Exposed, *The Mathematical Magazine*, 2 2002.
- [12] P. Chandra and E. W. Weisstein, Fibonacci Number. From MathWorld—A Wolfram Web Resource. Available at <https://mathworld.wolfram.com/FibonacciNumber.html>
- [13] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, Available at <https://oeis.org/>.
- [14] V.E. Hoggatt Jr. and M. Bicknell-Johnson, Fibonacci convolution sequences, *Fibonacci Quart.* 15, 1977, 117–122.
- [15] M. Thongmoon, New Identities for the Even and Odd Fibonacci and Lucas Numbers, *Int. J. Contemp. Math. Sciences*, Vol. 4, 2009, 671–676.
- [16] P. Barry and A. M. Mwafise, Classical and Semi-Classical Orthogonal Polynomials Defined by Riordan Arrays and Their Moment Sequences, *J. Integer Seq.*, Vol. 21, 2018, article 18.1.5.
- [17] J. Shallit, An interesting continued fraction, *Math. Mag.*, Vol. 48, No. 4, 1975, 207–211.
- [18] I. M. Gessel and J. Li, Compositions and Fibonacci identities, *J. Int. Seq.* 16, 2013, 13.4.5.
- [19] D. H. Lehmer, An Extended Theory of Lucas’ Functions. *Annals of Mathematics*, Vol. 31, No. 3, 1930, 419–448. Available at <https://doi.org/10.2307/1968235>.