# THE METRIC DIMENSION OF THE TOTAL GRAPH OF A SEMIRING

DAVID DOLŽAN

ABSTRACT. We calculate the metric dimension of the total graph of a direct product of finite commutative antinegative semirings with their sets of zero-divisors closed under addition.

#### 1. INTRODUCTION

Of the many ways one can prescribe a graph to an algebraic structure, one of the most important ones is the total graph. The total graph of a ring or a semiring S is a graph with all elements of S as vertices, and for distinct  $x, y \in S$ , the vertices x and y are adjacent if and only if x + y is a zero-divisor in S. Total graph has been firstly studied in [2] for commutative rings and in [9] for non-commutative rings. Later, it has been studied in [10, 11] in the setting of commutative semirings and in [15] in the case of matrices over the Boolean semiring.

In a simple undirected graph, there exist a couple of natural ways to distinguish the vertices from one another - such ideas appear already in Sumner [26] and Entringer and Gassman [12]. Among them, one of the most important ones is the metric dimension of a graph. Let us explain the definition. Choose an ordered subset  $W = \{w_1, w_2, \ldots, w_k\}$  of the vertex set of graph G. Then, for a vertex v in G, the vector  $(d(v, w_1), d(v, w_2), \ldots, d(v, w_k))$  is called the *representation* of v with respect to W. A set W is called a *resolving set* for G if distinct vertices of G have distinct representations with respect to W. A resolving set of minimal cardinality for G is called a *basis* of G and the cardinality of the basis is defined as the *metric dimension* of G, and denoted by dim(G).

The metric dimension was studied by Slater [24], while trying to determine the location of an intruder in a network. The notion was then also studied by Harary and Melter [13]. The metric dimension has later been used in many applications, for example in pharmaceutical chemistry [4], robot navigation in space [16], combinatorial optimization [23] and sonar and coast guard long range navigation [24, 25].

In view of studying the metric dimensions of graphs corresponding to different algebraic structures, the metric dimension of a zero-divisor graph of a commutative ring was studied in [18, 19, 20, 21, 22], a zero-divisor graph of a matrix semiring in [8], a total graph of a finite commutative ring in [6], an annihilating-ideal graph of a finite ring in [7], and a commuting graph of a dihedral group in [1].

Date: June 6, 2024.

<sup>2010</sup> Mathematics Subject Classification. 16Y60, 05C25.

Key words and phrases. semiring, zero-divisor, graph, metric dimension.

The author acknowledges the financial support from the Slovenian Research Agency (research core funding No. P1-0222).

In this paper, we study the metric dimension of the total graph of a direct product of finite commutative antinegative semirings. A semiring is a set S equipped with binary operations + and  $\cdot$  such that (S, +) is a commutative monoid with identity element 0 and  $(S, \cdot)$  is a monoid with identity element 1, while both operations in S are connected by distributivity. We also assume that 0 annihilates S. Semiring S is called *commutative* if we have ab = ba for all  $a, b \in S$  and *antinegative* or *zero-sum-free* (or simply, an *antiring*), if a + b = 0 for  $a, b \in S$  implies a = b = 0. Commutative semirings often arise naturally, since the set of nonnegative integers (or reals) with the usual operations of addition and multiplication forms a commutative semiring. But there are many more examples, as distributive lattices, tropical semirings, dioids, fuzzy algebras, inclines and bottleneck algebras are all different examples of commutative semirings. The simplest semiring is the Boolean semiring  $\mathcal{B} = \{0, 1\}$ , where 1 + 1 = 1. The theory of semirings also provides many applications, for example in optimization theory, automatic control, models of discrete event networks and graph theory ([3, 5, 17, 27]).

This paper is organized as follows. In the next section, we deal with the preliminary concepts and lemmas that come useful later, while we also make an argument as to why it makes more sense to study the total graph of the semiring, where the vertex set is limited to the set of all zero-divisors. In the main section, we determine the metric dimension of a direct product of antinegative finite commutative semirings, with their set of zero-divisors closed under addition. Such semirings for example contain the Boolean semiring, all finite distributive lattices with zero and exactly one atom, all finite semifields, all polynomial semirings over finite antirings modulo a monome, etc. It turns out that the metric dimension behaves somewhat differently in different settings, with the exceptions consting of semifields and semirings such that all their elements except the identity are zero divisors. The main results of this paper are Proposition 3.2, Theorem 3.3 and Theorem 3.6, which together cover all possible cases. It is perhaps interesting to compare these results with [6, Theorem 3.5 and Theorem 3.7], where even for the direct product of two finite fields, the metric dimension of its total graph is not completely determined.

### 2. Preliminaries

Throughout this paper, we shall suppose that S is a semiring, and that Z(S) denotes the set of all zero-divisors in S,  $Z(S) = \{x \in S; \text{ there exists } 0 \neq y \in S \text{ such that } xy = 0 \text{ or } yx = 0\}$ . We denote by  $\Gamma(S)$  the *total graph* of S. The vertex set  $V(\Gamma(S))$  of  $\Gamma(S)$  is the set of all elements in S and an unordered pair of vertices  $x, y \in V(\Gamma(S)), x \neq y$ , is an edge x - y in  $\Gamma(S)$  if  $x + y \in Z(S)$ . The sequence of edges  $x_0 - x_1, x_1 - x_2, ..., x_{k-1} - x_k$ in  $\Gamma$  is called a path of length k. The distance between vertices x and y is the length of the shortest path between them, denoted by d(x, y), while the diameter diam $(\Gamma)$  of the graph  $\Gamma$  is the longest distance between any two vertices of the graph.

We shall need the following definition.

**Definition 2.1.** Let v be a vertex of a graph G. Then the open neighbourhood of v is  $N(v) = \{u \in V(G); \text{ there exists an edge } uv \text{ in } G\}$  and the closed neighbourhood of v us  $N[v] = N(v) \cup \{v\}$ . Two distinct vertices u and v of G are twins if N(u) = N(v) or N[u] = N[v].

Twins turn out to be a very important notion when studying the metric dimension of a graph due to the following lemma.

**Lemma 2.2** ([14]). Suppose u and v are twins in a connected graph G and the set W is a resolving set for G. Then u or v is in W.

The next lemma shows that in order to study the metric dimension, it makes sense to limit ourselves to the subgraph with the vertex set equal to the set of all zero-divisors.

**Lemma 2.3.** Let S be an antinegative semiring. Then  $\Gamma(S)$  is a disconnected graph (where the set of all non zero-divisors are isolated vertices). The subgraph  $\Gamma_1(S)$  with the vertex set equal to Z(S) is connected with diam $(\Gamma_1(S)) \leq 2$ .

*Proof.* Choose and  $s \in S$  such that  $s \notin Z(S)$  and suppose that s - w is an edge in  $\Gamma(S)$  for some  $w \in S$ . This implies that  $s + w \in Z(S)$ , so there exists  $z \neq 0$  such that (s + w)z = 0. Since S is antinegative, this yields sz = 0, a contradiction. On the other hand, if  $s \in Z(S)$ , then 0 - s is an edge in  $\Gamma_1(S)$ .

Therefore, we shall limit ourselves to studying the subgraph  $\Gamma_1(S)$  from here onwards. We shall now investigate twins in the graph  $\Gamma_1(S)$ .

**Lemma 2.4.** Let  $n \ge 1$  and  $S = S_1 \times S_2 \times \ldots \times S_n$ , where  $S_i$  is a finite antinegative commutative semiring with  $Z(S_i)$  closed under addition for every  $i = 1, 2, \ldots, n$ . Choose  $a = (a_1, a_2, \ldots, a_n), b = (b_1, b_2, \ldots, b_n) \in Z(S)$ . Then a and b are twins in  $\Gamma_1(S)$  if and only if either  $a_i, b_i \in Z(S_i)$  or  $a_i, b_i \notin Z(S_i)$  for every  $i = 1, 2, \ldots, n$ .

Proof. Choose any  $1 \leq i \leq n$ . Notice that the fact that  $S_i$  is antinegative implies that  $a_i + b_i \in Z(S_i)$  yields  $a_i, b_i \in Z(S_i)$ : if there exists  $x \in S_i$  such that  $0 = (a_i + b_i)x = a_ix + b_ix$ , we have  $a_ix = b_ix = 0$ . Since  $Z(S_i)$  closed under addition for every  $i = 1, 2, \ldots, n$ , we also have that  $a_i, b_i \in Z(S_i)$  implies  $a_i + b_i \in Z(S_i)$ . So, there is an edge between  $a = (a_1, a_2, \ldots, a_n)$  and  $c = (c_1, c_2, \ldots, c_n)$  in  $\Gamma_1(S)$  if and only if there exists  $1 \leq i \leq n$  such that  $a_i$  and  $c_i$  are both in  $Z(S_i)$ . Thus, the neighbourhood of any vertex is completely determined by which of its components belong to the set of zero-divisors. Therefore, two elements  $a = (a_1, a_2, \ldots, a_n)$  and  $b = (b_1, b_2, \ldots, b_n)$  in S are twins if and only if either  $a_i, b_i \in Z(S_i)$  or  $a_i, b_i \notin Z(S_i)$  for every  $i = 1, 2, \ldots, n$ .

### 3. Metric dimension

In this section, we examine the metric dimension of a total graph of a direct product of antinegative commutative semirings. Firstly, let us establish the lower bound for the metric dimension.

**Lemma 3.1.** Let  $n \ge 1$  and  $S = S_1 \times S_2 \times \ldots \times S_n$ , where  $S_i$  is a finite antinegative commutative semiring with  $Z(S_i)$  closed under addition for every  $i = 1, 2, \ldots, n$ . Then  $\dim(\Gamma_1(S)) \ge |Z(S)| - 2^n + 1$ .

Proof. Suppose that W is a resolving set for  $\Gamma_1(S)$  and choose a subset  $\emptyset \neq X \subseteq \{1, 2, \ldots, n\}$ . Define  $M_X = M_1 \times M_2 \times \ldots \times M_n \subset Z(S)$ , where  $M_i = Z(S_i)$  if  $i \in X$  and  $M_i = S_i \setminus Z(S_i)$  if  $i \notin X$ . By Lemma 2.4, all elements of  $M_x$  are twins, so by Lemma 2.2, W contains all but perhaps one element of the set  $M_X$ . Observe that  $a = (a_1, a_2, \ldots, a_n) \in S$  is a vertex in  $\Gamma_1(S)$  if and only if there exists  $i \in \{1, 2, \ldots, n\}$  such that  $a_i \in Z(S_i)$ .

Therefore, every vertex in  $\Gamma_1(S)$  lies in exactly one of the sets  $M_X$  for a chosen set  $X \neq \emptyset$ . So, W contains all elements of Z(S) apart from perhaps as many as there are nontrivial subsets of the set  $\{1, 2, \ldots, n\}$ . Since there are exactly  $2^n - 1$  such nontrivial subsets, we have  $|W| \geq |Z(S)| - 2^n + 1$ . Because this is true for any resolving set W, we have  $\dim(\Gamma_1(S)) \geq |Z(S)| - 2^n + 1$  as stated.

Let us examine the interesting special case of the metric dimension of the total graph of the direct product of a number of Boolean semirings.

**Remark.** If S is an antinegative finite commutative semiring with  $|S| \ge 3$  and |Z(S)| = 1, then  $\Gamma_1(S)$  is a trivial graph (since it has only one vertex) - obviously, dim $(\Gamma_1(S)) = 0$  in this case. We shall therefore from here onwards always limit ourselves to the case where  $\Gamma_1(S)$  is non-trivial.

**Proposition 3.2.** Let  $n \ge 2$  and  $S = S_1 \times S_2 \times \ldots \times S_n$ , where  $S_i = \mathcal{B}$  for every  $i = 1, 2, \ldots, n$ . Then  $\dim(\Gamma_1(S)) = n$  for  $n \ne 2$  and  $\dim(\Gamma_1(S)) = n - 1$  for n = 2.

*Proof.* Assume first that  $n \geq 3$ . For every i = 1, 2, ..., n, let us denote by  $\overline{e_i}$  the element with 1 at every component, except the *i*-th component, which is equal to 0. Let W = $\{\overline{e_1}, \overline{e_2}, \ldots, \overline{e_n}\}$ . Now, choose distinct  $x = (x_1, x_2, \ldots, x_n)$  and  $y = (y_1, y_2, \ldots, y_n)$  in S. There exists  $i \in \{1, 2, ..., n\}$  such that  $x_i \neq y_i$ . Without loss of generality, we can assume  $x_i = 0$  and  $y_i = 1$ . Then  $x - \overline{e_i}$  is an edge in  $\Gamma(S_1)$  and  $y - \overline{e_i}$  is not, therefore x and y have different representations with respect to W. Thus, W is a resolving set, which proves that  $\dim(\Gamma_1(S)) \leq n$ . On the other hand, Lemma 2.3 tells us that  $\dim(\Gamma_1(S)) \leq 2$ , so any representation of a vertex that is not a member of a resolving set, has to be a vector with every component in  $\{1, 2\}$ . Assume that W is a resolving set and denote w = |W|. Since there are at most  $2^w$  different representations of vertices that are not in W and there are  $2^n - 1$  vertices in  $\Gamma_1(S)$ , we have  $2^w + w \ge 2^n - 1$ . Obviously,  $w \mapsto 2^w + w$ is an increasing function, and  $2^{n-1} + n - 1 \ge 2^n - 1$  implies  $n \ge 2^{n-1}$ , which further implies  $n \leq 2$ . Therefore for  $n \geq 3$ , we can conclude that  $w \geq n$ . So, we have proved that  $\dim(\Gamma_1(S)) \ge n$  for all  $n \ge 3$  and thus  $\dim(\Gamma_1(S)) = n$ . Finally, let us examine the case n=2. Then  $S=\mathcal{B}\times\mathcal{B}$  and one can easily check that  $w=\{(1,0)\}$  is a resolving set, so  $\dim(\Gamma_1(S)) = 1.$ 

We can now prove the following theorem.

**Theorem 3.3.** Let  $n \ge 1$  and  $S = S_1 \times S_2 \times \ldots \times S_n$ , where  $S_i$  is a finite antinegative commutative semiring with  $Z(S_i)$  closed under addition for every  $i = 1, 2, \ldots, n$ . Furthermore, assume that for every  $i = 1, 2, \ldots, n$  the following condition holds:  $|Z(S_i)| \ge 2$  or there exists  $j \ne i$  such that  $|S_i \setminus Z(S_i)| \ge 2$ . Then dim $(\Gamma_1(S)) = |Z(S)| - 2^n + 1$ .

Proof. By Lemma 3.1, we already know that  $\dim(\Gamma_1(S)) \geq |Z(S)| - 2^n + 1$ . Choose any subset  $\emptyset \neq X \subseteq \{1, 2, \ldots, n\}$  and define  $M_X = M_1 \times M_2 \times \ldots \times M_n \subset Z(S)$ , where  $M_i = Z(S_i)$  if  $i \in X$  and  $M_i = S_i \setminus Z(S_i)$  if  $i \notin X$ . Now, choose  $t_X \in M_X$  and denote  $M'_X = M_X \setminus \{t_X\}$ . If  $M'_X \neq \emptyset$ , choose  $s_X \in M'_X$ . Define  $W = \bigcup_{\emptyset \neq X \subseteq \{1, 2, \ldots, n\}} M'_X$ and let us prove that W is a resolving set for  $\Gamma_1(S)$ . Choose any distinct elements  $u = (u_1, u_2, \ldots, u_n), v = (v_1, v_2, \ldots, v_n) \in Z(S)$  such that  $u, v \notin W$ . By the construction of W and Lemma 2.4, there exists  $i \in \{1, 2, \ldots, n\}$  such that  $u_i \in Z(S_i)$  and  $v_i \notin Z(S_i)$  or vice versa. Without any loss of generality, we can assume that  $u_i \in Z(S_i)$  and  $v_i \notin Z(S_i)$ . Define  $Y = \{i\}$  and observe  $M'_Y$  is nonempty by our assumption. Therefore,  $u - s_Y$  is an edge in  $\Gamma(S_1)$  since  $Z(S_i)$  closed under addition, while  $v - s_Y$  is not an edge in  $\Gamma(S_1)$  since  $S_i$  is an antiring for every i = 1, 2, ..., n. Thus, u and v have different representations with respect to W and therefore W is a resolving set. We can use the same proof as in the proof of Lemma 3.1, to see that  $|W| = |Z(S)| - 2^n + 1$ , which now implies that  $\dim(\Gamma_1(S)) = |Z(S)| - 2^n + 1$ .

Let us illustrate the above with an example.

**Example 3.4.** Suppose  $S = S_1 \times S_2 \times \ldots \times S_n$ , where  $n \ge 2$  and  $S_i$  is a finite antinegative semifield for every  $i = 1, 2, \ldots, n$ . Suppose that at least two of the semifields  $S_i$  are not isomorphic to  $\mathcal{B}$ . Then all the assumptions of Theorem 3.3 are satisfied, so dim $(\Gamma_1(S)) = |Z(S)| - 2^n + 1 = |S_1||S_2| \ldots |S_n| - (|S_1| - 1)(|S_2| - 1) \ldots (|S_n| - 1) - 2^n + 1$ .

On the other hand, if  $S = S_1 \times S_2 \times \ldots \times S_n$ , where  $S_i = \mathcal{B}$  for every  $i = 1, 2, \ldots, n$ . Then by Proposition 3.2, dim $(\Gamma_1(S)) = n$  for  $n \neq 2$  and dim $(\Gamma_1(S)) = n - 1$  for n = 2.

Compare these results with results in [6, Theorem 3.5 and Theorem 3.7], where even for the direct product of two finite fields, the metric dimension of the total graph is not known in all cases. Also, it is interesting to compare these results with results in [21, Theorem 6.1], where similar results have been obtained, but in a different setting of the zero-divisor graphs of finite commutative rings, where again the field with two elements is an exception to the rule.

Let us further investigate the other possible cases that are not covered by Proposition 3.2 and Theorem 3.3. Before we do that, we need the following definition.

**Definition 3.5.** Let  $S = S_1 \times S_2 \times \ldots \times S_n$  a semiring. Then for every  $i = 1, 2, \ldots, n$ , define  $W_i = \{0, 1\} \subseteq S_i$ . Then  $\operatorname{supp}(S) = W_1 \times W_2 \times \ldots \times W_n \subseteq S$  is a support of S.

- **Theorem 3.6.** (1) Let  $S = B_1 \times B_2 \times \ldots \times B_m \times Z_1 \times Z_2 \times \ldots \times Z_n$ , where  $m, n \ge 1$ ,  $B_i = \mathcal{B}$  for  $i = 1, 2, \ldots, m$ , and  $Z_i$  is an antinegative finite commutative semiring with  $Z(Z_i)$  closed under addition,  $|Z_i| \ge 3$ ,  $|Z_i \setminus Z(Z_i)| = 1$  for every  $i = 1, 2, \ldots, n$ . Then  $\dim(\Gamma_1(S)) = |S| + m - 2^{m+n} = |Z(S)| - 2^{m+n} + m + 1$ .
  - (2) Let  $S = R \times B_1 \times B_2 \times \ldots \times B_m \times Z_1 \times Z_2 \times \ldots \times Z_n$ , where  $m, n \ge 0$  and  $mn \ne 0$ , R is an antinegative finite commutative semiring with  $|R| \ge 3$  and |Z(R)| = 1,  $B_i = \mathcal{B}$  for  $i = 1, 2, \ldots, m$ , and  $Z_i$  is an antinegative finite commutative semiring with  $Z(Z_i)$  closed under addition,  $|Z_i| \ge 3$ ,  $|Z_i \setminus Z(Z_i)| = 1$  for every  $i = 1, 2, \ldots, n$ . Then dim $(\Gamma_1(S)) = |S| - 2^{m+n+1} - |R| + 3 = |Z(S)| - 2^{m+n+1} + 2$ .

*Proof.* Define  $T = \{v \in Z(S); \text{ there exists } v \neq u \in Z(S) \text{ such that } u \text{ and } v \text{ are twins in } \Gamma_1(S)\}, N = Z(S) \setminus T \text{ and } T' = T \setminus \text{supp}(S).$  Now, let us look at both cases separately.

(1) Obviously, N contains exactly all elements from S that have no twins. It follows from Lemma 2.4 that N contains exactly all elements from Z(S) of the form  $(b_1, b_2, \ldots, b_m, 1, 1, \ldots, 1)$  for every  $i = 1, 2, \ldots, m$ . Since T is a complement of N in Z(S), observe that then T' contains exactly all elements of the form  $(b_1, b_2, \ldots, b_m, z_1, z_2, \ldots, z_2)$  for all  $b_i \in B_i$   $(1 \le i \le m)$  and all  $z_j \in Z_j$   $(1 \le j \le n)$ , except for those that have  $z_j \in \{0, 1\}$  for every j. Therefore  $|T'| = |S| - 2^{n+m}$ .

By Lemma 2.2, every resolving set W has to contain at least |T'| elements (and we can limit ourselves to the case where W contains T' without any loss of generality, since in every  $Z_i$  any zero-divisor is interchangeable with 0 in the sense that it has the same neighbourhood).

Now, define  $X = \{(b_1, b_2, \dots, b_m, 0, 0, \dots, 0); b_i \in B_i \text{ for every } i = 1, 2, \dots, m\}$ and observe that  $|X| = 2^m$ . Note that by definition,  $X \cap T' = \emptyset$  and also that d(x,t) = d(x',t) = 1 for every  $x, x' \in X$  and every  $t \in T'$ . If we want to have different representations for different elements from X, we therefore have to add some elements of the form  $(b_1, b_2, \ldots, b_m, 1, 1, \ldots, 1)$  to T', since only these kind of elements can be at distance more than one from elements in X. We now reason similarly, as in the proof of Proposition 3.2. Suppose we add w elements to T'. By Lemma 2.3, this means that we have at most  $2^w$  possible different new representations of vertices from X. There are  $2^m$  vertices in X, which implies  $2^w + w \ge 2^m$ . Obviously,  $w \mapsto 2^w + w$  is an increasing function, and  $2^{m-1} + m - 1 \ge 2^m$  implies  $m - 1 \ge 2^m$ , which is a contradiction, so  $w \ge m$ . We have therefore proved that  $\dim(\Gamma_1(S)) \ge |S| + m - 2^{n+m}$ .

On the other hand, for every i = 1, 2, ..., n + m denote by  $\overline{e_i}(x)$  the element of S with 1 at every component, except the *i*-th component, which is equal to x. Define  $T'' = \{\overline{e_1}(0), \overline{e_2}(0), ..., \overline{e_m}(0)\}$  and  $W = T' \cup T''$  for T' as above. Observe that  $|W| = |S| + m - 2^{n+m}$ . Let us prove that W is a resolving set. Choose  $b = (b_1, b_2, ..., b_m, z_1, z_2, ..., z_n) \neq b' = (b'_1, b'_2, ..., b'_m, z'_1, z'_2, ..., z'_n) \in Z(S) \setminus W$ for some  $b_i, b'_i \in B_i$  and some  $z_j, z, j' \in Z_j$  for every  $1 \leq i \leq m, 1 \leq j \leq n$ . Suppose firstly that there exists  $1 \leq j \leq n$  such that  $z_j \neq z'_j$ . Since  $b, b' \notin W$ , we know that  $z_j = 0$  and  $z'_j = 1$  or vice versa. Without loss of generality, we can assume the former. Now, choose  $x \in Z(S_j) \setminus \{0\}$  and note that  $\overline{e_j}(x) \in T'$ . This now implies that  $d(b, \overline{e_j}(x)) = 1$ , while  $d(b', \overline{e_j}(x)) > 1$ . Suppose now that  $z_j = z'_j$  for all  $1 \leq j \leq n$ . Since  $b \neq b'$ , there exists  $1 \leq i \leq m$  such that  $b_i \neq b'_i$ . Then we have  $d(b, \overline{e_i}(0)) \neq d(b', \overline{e_i}(0))$ . This implies that W is indeed a resolving set, so  $\dim(\Gamma_1(S)) \leq |S| + m - 2^{n+m}$ , which now proves our assertion. Note also that every element of S except for (1, 1, ..., 1) is a zero divisor, so  $|S| + m - 2^{n+m} = |Z(S)| - 2^{m+n} + m + 1$ .

(2) Since N consists of exactly all those elements from Z(S) that have no twins, Lemma 2.4 this time shows that N contains exactly all the elements from Z(S) of the form  $(0, b_1, b_2, \ldots, b_m, 1, 1, \ldots, 1)$ , where  $b_i \in B_i$  for every  $i = 1, 2, \ldots, m$ . Again, T is the complement of N in Z(S), so we can reason similarly as above that  $|T'| = |S| - 2^{m+n+1} - (|R| - 2)$ , since in this case we have to exclude from S all the elements of the form  $(r, b_1, b_2, \ldots, b_m, z_1, z_2, \ldots, z_2)$  for all  $r \in \{0, 1\}$ , all  $b_i \in B_i$  and all  $z_j \in \{0, 1\}$ , as well as |R| - 2 elements of the form  $(r, 1, 1, \ldots, 1)$  for  $r \notin \{0, 1\}$  since they are not zero-divisors at all.

Suppose now that W is a resolving set. Using Lemma 2.2, we can reason that W has to contain at least |T'| elements. We can also limit ourselves to the case where W contains T' without any loss of generality, since in every  $Z_i$  any zero-divisor is interchangeable with 0 and in R every non-zero element is interchangeable with 1 in the sense that they have the same neighbourhood.

Let us assume that W = T' and choose b = (1, 0, 0, ..., 0) and b' = (0, 0, ..., 0)in S. Since m and n are not both equal to zero, b and b' are zero-divisors. By the construction of T', we know that  $b, b' \notin W$ . However, for every  $w = (w_1, w_2, ..., w_{n+m+1}) \in W$ , we have one of the following two possible cases.

- (a) We have  $w_1 = 0$  and thus (since  $w \notin \text{supp}(S)$ ) there exists  $1 \leq j \leq n$  such that  $w_{1+m+j} \notin \{0,1\}$ . But since all elements of  $S_{1+m+j}$  except 1 are zero-divisors, we have  $w_{1+m+j} \in Z(S_{1+m+j})$ , so in this case d(b,w) = d(b',w) = 1.
- (b) We have  $w_1 \neq 0$ . Since w is a zero divisor there exists  $2 \leq j \leq n+m+1$  such that  $w_j$  is a zero-divisor. Again, we have d(b, w) = d(b', w) = 1.

So, we have proved that b and b' have the same representations with respect to W, which is a contradiction with the fact that W is a resolving set. Therefore  $T' \subsetneq W$ , so  $|W| \ge |T'| + 1 = |S| - 2^{m+n+1} - (|R| - 2) + 1 = |S| - 2^{m+n+1} - |R| + 3$ . This implies that we have  $\dim(\Gamma_1(S)) \ge |S| - 2^{n+m+1} - |R| + 3$ .

Let us now prove that we also have  $\dim(\Gamma_1(S)) \leq |S| - 2^{n+m+1} - |R| + 3$ . For every  $i = 1, 2, \ldots, n + m + 1$  denote by  $\overline{e_i}(x)$  the element of S with 1 at every component, except the *i*-th component, which is equal to x. Let us define  $W = T' \cup \{e_1(0)\}$  for the set T' defined as above. Observe that  $|W| = |S| - 2^{m+n+1} - (|R| - 2) + 1 = |S| - 2^{m+n+1} - |R| + 3$ . Also, Let us prove that W is a resolving set. Choose  $b = (r_1, b_1, b_2, \dots, b_m, z_1, z_2, \dots, z_n) \neq b' =$  $(r'_1, b'_1, b'_2, \ldots, b'_m, z'_1, z'_2, \ldots, z'_n) \in Z(S) \setminus W$  for some  $r_1, r'_1 \in R$ , some  $b_i, b'_i \in B_i$ and some  $z_j, z'_j \in Z_j$  for every  $1 \le i \le m, 1 \le j \le n$ . Suppose firstly that there exists  $1 \leq j \leq n$  such that  $z_j \neq z'_j$ . Since  $b, b' \notin W$ , we know that  $z_j = 0$  and  $z'_i = 1$  or vice versa. Without loss of generality, we can assume the former. Now, choose  $x \in Z_j \setminus \{0,1\}$  and note that  $\overline{e_j}(x) \in T'$  by the construction of the set T'. This now implies that  $d(b, \overline{e_j}(x)) = 1$ , while  $d(b', \overline{e_j}(x)) > 1$ . Suppose now that  $z_j = z'_j$  for all  $1 \le j \le n$ . Since  $b \ne b'$ , we have  $r_1 \ne r'_1$  or there exists  $1 \leq i \leq m$  such that  $b_i \neq b'_i$ . Suppose firstly that  $b_i \neq b'_i$  for some  $1 \leq i \leq m$ . Without any loss of generality, we can assume that  $b_i = 0$  and  $b'_i = 1$ . Choose  $r \in R \setminus \{0\}$  and define  $s = e_1(r)e_i(0)$ . Again, note that  $s \in T'$  and that we now have d(b,s) = 1 and d(b',s) > 1. Finally, suppose  $b_i = b'_i$  and  $z_j = z'_j$  for all  $1 \leq i \leq m$  and  $1 \leq j \leq n$ . Then we have  $r_1 \neq r'_1$  and since  $b, b' \notin W$  we either have  $r_1 = 0$  or  $r'_1 = 1$  or vice versa. Again, without loss of generality, assume the former. Observe that  $d(b, e_1(0)) = 1$  and  $d(b', e_1(0)) > 1$ . This implies that W is indeed a resolving set, so  $\dim(\Gamma_1(S)) \leq |S| - 2^{n+m+1} - |R| + 3$ . Note also that all elements of S except for elements of the form (r, 1, ..., 1)for some  $r \in R \setminus \{0\}$  are zero-divisors, so |S| = |Z(S)| + |R| - 1 and thus  $\dim(\Gamma_1(S)) = |S| - 2^{n+m+1} - |R| + 3 = |Z(S)| - 2^{m+n+1} + 2$ , which now proves our assertion.

Observe that Proposition 3.2, Theorem 3.3 and Theorem 3.6 now cover all possible cases of direct product of antinegative finite commutative semirings with their sets of zerodivisors closed under addition, so we have completely determined the metric dimension in this setting.

## **Statements and Declarations**

The author states that there is no conflict of interest.

#### References

- 1. Faisal Ali, Muhammad Salman, and Shuliang Huang, On the commuting graph of dihedral group, Comm. Algebra 44 (2016), no. 6, 2389–2401. MR 3492163
- David F. Anderson and Ayman Badawi, The total graph of a commutative ring, J. Algebra 320 (2008), no. 7, 2706–2719. MR 2441996

- François Baccelli and Jean Mairesse, Ergodic theorems for stochastic operators and discrete event networks, Idempotency (Bristol, 1994), Publ. Newton Inst., vol. 11, Cambridge Univ. Press, Cambridge, 1998, pp. 171–208. MR 1608402
- P. J. Cameron and J. H. van Lint, *Designs, graphs, codes and their links*, London Mathematical Society Student Texts, vol. 22, Cambridge University Press, Cambridge, 1991. MR 1148891
- Raymond Cuninghame-Green, Minimax algebra, Lecture Notes in Economics and Mathematical Systems, vol. 166, Springer-Verlag, Berlin-New York, 1979. MR 580321
- David Dolžan, The metric dimension of the total graph of a finite commutative ring, Canad. Math. Bull. 59 (2016), no. 4, 748–759, [Paging previously given as 1–12]. MR 3563754
- 7. \_\_\_\_\_, The metric dimension of the annihilating-ideal graph of a finite commutative ring, Bull. Aust. Math. Soc. **103** (2021), no. 3, 362–368. MR 4255540
- The Metric Dimension of the Zero-Divisor Graph of a Matrix Semiring, Bull. Malays. Math. Sci. Soc. 46 (2023), no. 6, 201. MR 4656836
- David Dolžan and Polona Oblak, The total graphs of finite rings, Comm. Algebra 43 (2015), no. 7, 2903–2911. MR 3354069
- 10. \_\_\_\_\_, The total graphs of finite commutative semirings, Results Math. **72** (2017), no. 1-2, 193–204. MR 3684424
- 11. Shahabaddin Ebrahimi Atani and Fatemeh Esmaeili Khalil Saraei, *The total graph of a commutative semiring*, An. Stiint. Univ. "Ovidius" Constanța Ser. Mat. **21** (2013), no. 2, 21–33. MR 3096881
- R. C. Entringer and Larry D. Gassman, Line-critical point determining and point distinguishing graphs, Discrete Math. 10 (1974), 43–55. MR 347680
- Frank Harary and Robert A. Melter, On the metric dimension of a graph, Ars Combin. 2 (1976), 191–195. MR 457289
- Carmen Hernando, Mercè Mora, Ignacio M. Pelayo, Carlos Seara, and David R. Wood, *Extremal graph theory for metric dimension and diameter*, Electron. J. Combin. 17 (2010), no. 1, Research Paper 30, 28. MR 2595490
- L. John and Harita Vijay, On the total graph of the semiring of matrices over the Boolean semiring, Bull. Kerala Math. Assoc. 13 (2016), no. 2, 199–207. MR 3585523
- Samir Khuller, Balaji Raghavachari, and Azriel Rosenfeld, Landmarks in graphs, Discrete Appl. Math. 70 (1996), no. 3, 217–229. MR 1410574
- P. Li, A heuristic method to compute the approximate postinverses of a fuzzy matrix, IEEE Transactions on Fuzzy Systems 22 (2014), 1347–1351.
- Shikun Ou, Dein Wong, Fenglei Tian, and Qi Zhou, Fixing number and metric dimension of a zero-divisor graph associated with a ring, Linear Multilinear Algebra 69 (2021), no. 10, 1789–1802. MR 4279157
- S. Pirzada, M. Aijaz, and S. P. Redmond, Upper dimension and bases of zero-divisor graphs of commutative rings, AKCE Int. J. Graphs Comb. 17 (2020), no. 1, 168–173. MR 4145403
- S. Pirzada and Rameez Raja, On the metric dimension of a zero-divisor graph, Comm. Algebra 45 (2017), no. 4, 1399–1408. MR 3576665
- 21. Rameez Raja, S. Pirzada, and Shane Redmond, On locating numbers and codes of zero divisor graphs associated with commutative rings, J. Algebra Appl. 15 (2016), no. 1, 1650014, 22. MR 3393943
- 22. Shane Redmond and Steve Szabo, When metric and upper dimensions differ in zero divisor graphs of commutative rings, Discrete Math. Lett. 5 (2021), 34–40. MR 4227399
- András Sebő and Eric Tannier, On metric generators of graphs, Math. Oper. Res. 29 (2004), no. 2, 383–393. MR 2065985
- Peter J. Slater, *Leaves of trees*, Proceedings of the Sixth Southeastern Conference on Combinatorics, Graph Theory, and Computing (Florida Atlantic Univ., Boca Raton, Fla., 1975), Congressus Numerantium, No. XIV, Utilitas Math., Winnipeg, Man., 1975, pp. 549–559. MR 0422062
- Fault-tolerant locating-dominating sets, vol. 249, 2002, Combinatorics, graph theory and computing (Louisville, KY, 1999), pp. 179–189. MR 1898670
- 26. David P. Sumner, Point determination in graphs, Discrete Math. 5 (1973), 179-187. MR 316319
- Shan Zhao and Xue-ping Wang, Invertible matrices and semilinear spaces over commutative semirings, Inform. Sci. 180 (2010), no. 24, 5115–5124. MR 2726681

D. DOLŽAN: DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICS AND PHYSICS, UNIVER-SITY OF LJUBLJANA, JADRANSKA 19, SI-1000 LJUBLJANA, SLOVENIA, AND INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, JADRANSKA 19, SI-1000 LJUBLJANA, SLOVENIA; E-MAIL: DAVID.DOLZAN@FMF.UNI-LJ.SI