SKEW KNÖRRER'S PERIODICITY THEOREM

YANG LIU, YUAN SHEN, AND XIN WANG

ABSTRACT. In this paper, we introduce a class of twisted matrix algebras of $M_2(E)$ and twisted direct products of $E \times E$ for an algebra E. Let A be a noetherian Koszul Artin-Schelter regular algebra, $z \in A_2$ be a regular central element of A and $B = A_P[y_1, y_2; \sigma]$ be a graded double Ore extension of A. We use the Clifford deformation $C_{A^{!}}(z)$ of Koszul dual $A^{!}$ to study the noncommutative quadric hypersurface $B/(z + y_1^2 + y_2^2)$. We prove that the stable category of graded maximal Cohen-Macaulay modules over $B/(z + y_1^2 + y_2^2)$ is equivalent to certain bounded derived categories, which involve a twisted matrix algebra of $M_2(C_{A^{!}}(z))$ or a twisted direct product of $C_{A^{!}}(z) \times C_{A^{!}}(z)$ depending on the values of P. These results are presented as skew versions of Knörrer's periodicity theorem. Moreover, we show $B/(z + y_1^2 + y_2^2)$ may not be a noncommutative graded isolated singularity even if A/(z) is.

INTRODUCTION

Throughout the paper, k is an algebraically closed field of characteristic 0. All vector spaces and algebras are over k. Unless otherwise stated, the tensor product \otimes means \otimes_k .

Knörrer's periodicity theorem is a powerful tool in the study of the Cohen-Macaulay representation theory and also the singularities of (commutative) quadric hypersurfaces (see [7]). It implies that the stable category $\underline{\mathrm{mcm}} S$ of maximal Cohen-Macaulay modules over a quadric hypersurface $S = \mathbb{k}[x_1, \dots, x_n]/(f)$ is equivalent to $\underline{\mathrm{mcm}} S^{\#\#}$ where $f \in (x_1, \dots, x_2)^2$ and $S^{\#\#} = \mathbb{k}[x_1, \dots, x_n][u, v]/(f + u^2 + v^2)$ is the second double branched cover of S, which helps us to reduce the computation of the Cohen-Macaulay representation of a quadric hypersurface.

A noetherian Koszul Artin-Schelter regular algebra A is regarded as a noncommutative polynomial and a coordinate ring of a class of noncommutative projective spaces. Let $z \in A_2$ be a nonzero regular central element of A. The quotient algebra A/(z) is an analogue of (commutative) quadric hypersurface, usually called a *noncommutative quadric hypersurface*. There have been many studies in noncommutative quadric hypersurfaces for the past few years (for example, [1–6,9,10,16,18,19]). To study graded Cohen-Macaulay modules of a noncommutative quadric hypersurface A/(z), there are two useful tools. One is a finite dimensional algebra C(A/(z)) introduced by Smith and Van den Bergh in [16], and the other one is a Clifford deformation $C_{A'}(z)$ of the Koszul dual $A^{!}$ of Aintroduced by He and Ye in [2]. The Clifford deformation $C_{A'}(z)$ is a strongly \mathbb{Z}_2 -graded algebra and $C_{A'}(z)_0 \cong C(A/(z))$ (see [2]), and so $C_{A'}(z)$ may provide more information about A/(z).

²⁰¹⁰ Mathematics Subject Classification. 16S37, 16E65, 16G50, 16W50, 18E30.

Key words and phrases. noncommutative quadric hypersurface, skew Knörrer's periodicity theorem, double Ore extension, twisted matrix algebra.

Similar to the commutative case, the noncommutative Knörrer's periodicity theorem, that is, $\underline{\mathrm{mcm}} A/(z)$ and $\underline{\mathrm{mcm}}(A/(z))^{\#\#}$ are equivalent, was proved for a noncommutative quadric hypersurface A/(z) in [1, 2, 10]. He, Ma and Ye proved a generalized version of Knörrer's periodicity theorem for tensor products of two noncommutative quadric hypersurfaces with the help of Clifford deformations (see [3, Theorem 2.15]).

In noncommutative setting, it is reasonable to consider noncommutative versions of double branched covers and twisted tensor products of noncommutative quadric hypersurfaces. For example, $T = A[u; \sigma_1][v; \sigma_2]/(z + u^2 + v^2)$ defined by two-step iterated Ore extensions for some graded automorphisms σ_1 and σ_2 seems to make more sense as a class of second noncommutative double branched covers, and a natural question is how we can describe $\underline{\mathrm{mcm}} T$ by $\underline{\mathrm{mcm}} A/(z)$. If $\sigma_1 = \sigma_{2|A}$ and $\sigma_2(u) = u$, it is a special case of the noncommutative Knörrer's periodicity theorem given by Mori and Ueyama (see [10, Theorem 1.3]), and this case can also reduce to the second double branched cover $(A/(z))^{\#\#}$ by a Zhang-twist (see [20]). Nevertheless, we are curious abut how to describe $\underline{\mathrm{mcm}} T$ in general.

On the other hand, Hu, Matsuno and Mori completely classified noncommutative conics up to isomorphism of noncommutative projective schemes in [6], that is a class of noncommutative quadric hypersurfaces defined by 3-dimensional noetherian Koszul Artin-Schelter regular algebras. It inspires us to discuss the classification of noncommutative quadric hypersurfaces obtained by 4dimensional noetherian Koszul Artin-Schelter regular algebras. To achieve this goal, it is necessary to compute the Cohen-Macaulay representation of such noncommutative quadric hypersurfaces. Note that Zhang and Zhang introduced an approach called double Ore extension to obtain a classification of 4-dimensional noetherian Koszul Artin-Schelter regular algebras (see [21,22]). Also, double Ore extensions include two-step iterated Ore extensions. Therefore, it is reasonable to study relations between the graded maximal Cohen-Macaulay modules over noncommutative quadric hypersurfaces A/(z) and $A_P[y_1, y_2; \sigma]/(z + y_1^2 + y_2^2)$ where $A_P[y_1, y_2; \sigma]$ is a double Ore extension of A.

To our end, we introduce twisted algebra structures on a matrix algebra $M_2(E)$ and a direct product $E \times E$ for an algebra E. The twisted matrix algebra of $M_2(E)$ (see Theorem 2.7) comes from a twisting system Θ (see Definition 2.6) consisting of two linear maps $\theta^{(i)} : E \to M_2(E)$ for $i \in \mathbb{Z}_2$ and an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$, where $M_2(\mathbb{k})_0$ consists of diagonal matrices and $M_2(\mathbb{k})_1$ consists of anti-diagonal matrices. Denote the new twisted matrix algebra by $\Theta M_2(E)$, and it is a \mathbb{Z}_2 -graded algebra. This twisted algebra can be viewed as a generalization of Zhangtwists (see [20]) for graded algebras, but they have different properties. Furthermore, taking the subalgebra $\Theta M_2(E)_0$ of $\Theta M_2(E)$ provides an approach to obtaining a twisted algebra structure on $E \times E$ (see Corollary 2.14) and denote the new twisted algebra by $\theta_{\times}(E \times E)$ where the twisting system θ_{\times} consists of a linear map θ from E to $M_2(E)$ and an invertible basis of $\mathbb{k} \times \mathbb{k}$. If E is a \mathbb{Z}_2 -graded algebra, then $\Theta M_2(E)$ is $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded and also \mathbb{Z}_2 -graded by taking the total degrees, and $\theta_{\times}(E \times E)$ is \mathbb{Z}_2 -graded.

Let A be a noetherian Koszul Artin-Schelter regular algebra, $z \in A_2$ be a nonzero regular central element and $B = A_P[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of A where $P = \{p_{11}, p_{12}\} \subseteq \mathbb{k}$. The main object we study is the noncommutative quadric hypersurface $B/(z + y_1^2 + y_2^2)$. By the deformation method introduced by He-Ye, $\underline{\mathrm{mcm}} B/(z+y_1^2+y_2^2)$ is totally determined by the Clifford deformation $C_{B'}(z+y_1^2+y_2^2)$ of B' ([2, Theorem 0.2]). We aim to describe $C_{B'}(z+y_1^2+y_2^2)$ by the

Clifford deformation $C_{A^{\dagger}}(z)$ of A^{\dagger} . It should be noted that the semi-trivial extension (see Theorem 1.6) is a useful construction for our goal. To make the noncommutative quadric hypersurface $B/(z + y_1^2 + y_2^2)$ well-defined, it forces $p_{12} = \pm 1$. We divide it into two cases.

If $p_{12} = 1$, p_{11} should be 0. In this case, the Clifford deformation $C_{B'}(z + y_1^2 + y_2^2)$ is a twisted matrix algebra of $M_2(C_{A'}(z))$, and $\underline{\mathrm{mcm}}(B/(z + y_1^2 + y_2^2))$ can be described by a semi-trivial extension.

Theorem 0.1. (Theorem 4.4 and Theorem 4.8) Let $B = A_{\{1,0\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A with deg $y_1 = \text{deg } y_2 = 1$ and $z \in A_2$ be a nonzero regular central element of A. Suppose B is noetherian and $z + y_1^2 + y_2^2$ is a central element of B. Then

(1) there exists a twisting system Θ of $M_2(C_{A^1}(z))$ such that

$$C_{B'}(z+y_1^2+y_2^2) \cong {}^{\Theta}M_2(C_{A'}(z)),$$

as \mathbb{Z}_2 -graded algebras;

(2) there exists a semi-trivial extension $\Lambda = S \ltimes_{\psi} M(1)$ of a \mathbb{Z}_2 -graded subalgebra S of $C_{A^1}(z)$ and a \mathbb{Z}_2 -graded S-bimodule M (a subspace of $C_{A^1}(z)$) through a map $\psi : M(1) \otimes_S M(1) \to S$ such that

 $\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2} \,^{\Theta}M_2(C_{A^!}(z))) \cong D^b(\mathrm{mod}\,^{\Theta}M_2(C_{A^!}(z))_0) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2}\Lambda),$

as triangulated categories.

If $p_{12} = -1$, we can reduce p_{11} to be 0 in case $p_{11} \neq \pm 2\sqrt{-1}$. Then one obtains a twisted direct product $\Gamma = {}^{\theta_{\times}}(C_{A'}(z) \times C_{A'}(z))$ for some twisting system θ_{\times} and the Clifford deformation $C_{B'}(z + y_1^2 + y_2^2)$ is a semi-trivial extension constructed by Γ . In particular, the 0-th component of $C_{B'}(z + y_1^2 + y_2^2)$ is exactly a Zhang-twisted algebra (see [20]) of Γ .

Theorem 0.2. (Theorem 5.9) Let $B = A_{\{-1,0\}}[y_1, y_2; \sigma]$ be a Z-graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A with deg $y_1 = \text{deg } y_2 = 1$ and $z \in A_2$ be a nonzero regular central element of A. Suppose B is noetherian and $z + y_1^2 + y_2^2$ is a central element of B. Then

(1) there exist a twisted direct product Γ of $C_{A^{!}}(z) \times C_{A^{!}}(z)$ and a semi-trivial extension $\Gamma \ltimes_{\psi} (_{\mu}\Gamma(1))$ of \mathbb{Z}_{2} -graded algebra Γ and \mathbb{Z}_{2} -graded Γ -bimodule $_{\mu}\Gamma(1)$ through some map ψ where μ is a \mathbb{Z}_{2} -graded automorphism of Γ such that

$$C_{B^{!}}(z+y_{1}^{2}+y_{2}^{2}) \cong \Gamma \ltimes_{\psi} (_{\mu}\Gamma(1))$$

as \mathbb{Z}_2 -graded algebras.

(2) there are equivalences of triangulated categories

 $\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b\left(\mathrm{gr}_{\mathbb{Z}_2}\,\Gamma\ltimes_\psi\left({}_{\mu}\Gamma(1)\right)\right) \cong D^b\left(\mathrm{mod}\,{}^{\nu}\Gamma\right),$

where ${}^{\nu}\Gamma$ is a Zhang-twisted algebra of Γ for the left Zhang-twisting system $\nu = \{\nu_0 = id, \nu_1 = \mu\}$ of \mathbb{Z}_2 -graded algebra Γ .

In particular, Theorem 0.1 reduces to the noncommutative Knörrer's periodicity theorem if σ is a diagonal map. Theorem 0.1 and Theorem 0.2 can be viewed as skew versions of Knörrer's periodicity theorem.

For the exceptional case $p_{12} = -1$ and $p_{11} = \pm 2\sqrt{-1}$, we have any noncommutative quadric hypersurface $A_{\{-1,p_{11}\}}[y_1, y_2; \sigma]/(z+y_1^2+y_2^2)$ is not a noncommutative graded isolated singularity in the sense of [17] (see Proposition 5.1), or whose noncommutative projective scheme is not smooth. Hence, we are also interested in whether $A_{\{\pm 1,0\}}[y_1, y_2; \sigma]/(z+y_1^2+y_2^2)$ is a noncommutative graded isolated singularity in case A/(z) is. We give a negative answer to this question (see Proposition 5.12). It should be emphasized that this result is a totally different phenomena to the usual second double branched covers and tensor products of noncommutative quadric hypersurfaces, which preserve the property of noncommutative graded isolated singularity (see [2, Theorem 0.4] and [3, Theorem 2.5]).

This paper is organized as follows. In Section 1, we recall some definitions and results, including noncommutative quadric hypersurfaces, Clifford deformations and semi-trivial extensions. In Section 2, we introduce a class of twisted matrix algebras and a class of twisted direct products. In Section 3, we recall the definition of double Ore extensions and show some basic facts about the Clifford deformations for noncommutative quadric hypersurfaces obtaining by double Ore extensions. We devote Section 4 and Section 5 to proving Theorem 0.1 and Theorem 0.2 respectively.

1. Preliminaries

Let $C = \bigoplus_{g \in G} C_g$ be a *G*-graded algebra where *G* is an abelian group. Write $\operatorname{Gr}_G C$ for the category of *G*-graded right *C*-modules with morphisms of *C*-module homomorphisms preserving grading and $\operatorname{gr}_G C$ for the full subcategory of $\operatorname{Gr}_G C$ consisting of finitely generated *G*-graded right *C*-modules. Let *M* be a *G*-graded right *C*-module, then M(h) stands for the shifting *G*-graded right *C*-module where $M(h)_g = M_{g+h}$ for any $g \in G$ and some $h \in G$. For a *G*-graded automorphism μ of *C*, the twisted *G*-graded right *C*-module M_{μ} is the abelian group *M* with the right *C*-action $m \cdot c = m\mu(c)$ for any $m \in M, c \in C$. Similarly, we can define shifting and twisted *G*-graded left *C*-modules (or, *C*-bimodules).

For $M, M' \in \operatorname{Gr}_G C$, write

$$\operatorname{Hom}_{C}(M, M') = \bigoplus_{g \in G} \operatorname{Hom}_{\operatorname{Gr}_{G}}(M, M'(g))$$

and $\underline{\operatorname{Ext}}_{C}^{i}$ for the *i*-th derived functor of $\underline{\operatorname{Hom}}_{C}$. In particular, \Bbbk is always a *G*-graded algebra concentrated in degree 0, and denote the *G*-graded linear dual functor $\underline{\operatorname{Hom}}_{\Bbbk}(-, \Bbbk)$ by $(-)^{*}$. For any nonzero $k \in \Bbbk$, define a *G*-graded algebra automorphism ξ_{k} of *C* by $\xi_{k}(c) = k^{\deg c}c$ for any homogeneous element *c* of *C*.

A set of G-graded linear automorphisms $\nu = \{\nu_g \mid g \in G\}$ of C is called a *left Zhang-twisting* system of C, if $\nu_l(\nu_h(x)y) = \nu_{hl}(x)\nu_l(y)$ for all $g, h, l \in G$ and $x \in C_g, y \in C_h$. If $\nu = \{\nu_g \mid g \in G\}$ is left Zhang-twisting system of C, the graded vector space $C = \bigoplus_{g \in G} C_g$ with the multiplication $x * y = \nu_h(x)y$, for all $x \in C_g, y \in C_h$, is a G-graded algebra, called a *Zhang-twisted algebra* of A by ν , denoted by νA (see [20, Section 4]).

Definition 1.1. Let C be a G-graded algebra, and $\sigma = (\sigma_{ij})$ and $\varphi = (\varphi_{ij})$ be two algebra homomorphisms from C to $M_2(C)$ where σ_{ij} and φ_{ij} are G-graded linear transformations of C for any i, j = 1, 2. If

$$\sum_{k=1}^{2} \sigma_{ki} \varphi_{kj} = \delta_{ij} \operatorname{id}_{C} \quad \text{and} \quad \sum_{k=1}^{2} \varphi_{jk} \sigma_{ik} = \delta_{ij} \operatorname{id}_{C},$$

where δ_{ij} is the Kronecker symbol for any i, j = 1, 2, we say σ is *invertible*, φ is *t-invertible*, φ is an *inverse* of σ and σ is a *t-inverse* of φ .

In this case, it is not hard to obtain that the inverse of σ and the *t*-inverse of φ are both unique.

Remark 1.2. Definition 1.1 is a graded version of [21, Definition 1.8]. Note that the invertible form and *t*-invertible form are not the same. The letter "*t*" of *t*-invertible stands for the word "transpose" since the transpose of φ has the the same invertible property of σ .

A non-negative \mathbb{Z} -graded algebra C with $C_0 = \mathbb{k}$ and dim $C_i < \infty$ for any $i \ge 1$ is called a *locally finite connected* \mathbb{Z} -graded algebra. A locally finite connected \mathbb{Z} -graded algebra A is called a *Koszul algebra* (see [13]), if the trivial module \mathbb{k}_A has a free resolution

$$\cdots \to P_n \to \cdots \to P_1 \to P_0 \to \mathbb{k} \to 0,$$

where P_i is a \mathbb{Z} -graded free right A-module generated in degree i for any $i \geq 0$. Then $A \cong T(V)/(R)$ for some finite dimensional vector space V with degree 1 and some vector space $R \subseteq V \otimes V$. The Koszul dual of A is the quadratic algebra $A^! = T(V^*)/(R^{\perp})$, where V^* is the dual vector space of V and $R^{\perp} \subseteq V^* \otimes V^*$ is the orthogonal complement of R. The Koszul dual $A^!$ is also Koszul.

Definition 1.3. A locally finite connected \mathbb{Z} -graded algebra D is called an *Artin-Schelter Gorenstein algebra* of injective dimension d if

- (1) injdim $_DD = \text{injdim } D_D = d < \infty$,
- (2) $\underline{\operatorname{Ext}}_{D}^{i}(\Bbbk_{D}, D_{D}) = \underline{\operatorname{Ext}}_{D}^{i}({}_{D}\Bbbk_{,D} D) = 0$ if $i \neq d$ and $\underline{\operatorname{Ext}}_{D}^{d}(\Bbbk_{D}, D_{D}), \underline{\operatorname{Ext}}_{D}^{d}({}_{D}\Bbbk_{,D} D)$ are both 1-dimensional.

In addition, D has finite global dimension, then D is called an *Artin-Schelter regular algebra* of dimension d.

Let A = T(V)/(R) be a noetherian Koszul Artin-Schelter regular algebra of dimension d and $z \in A_2$ be a nonzero regular central element of A. The quotient algebra S = A/(z) is a noncommutative quadric hypersurface, and it is a noetherian Koszul Artin-Schelter Gorenstein algebra of injective dimension d - 1 by [16, Lemma 5.1(a)].

The noncommutative projective scheme qgr $S = \operatorname{gr}_{\mathbb{Z}} S/\operatorname{tors}_{\mathbb{Z}} S$ where $\operatorname{tors}_{\mathbb{Z}} S$ is the full subcategory of $\operatorname{gr}_{\mathbb{Z}} S$ consisting of finite dimensional \mathbb{Z} -graded right S-modules. We say S is a noncommutative graded isolated singularity if qgr S has finite global dimension (see [17]).

A finitely generated \mathbb{Z} -graded right S-module N is called a maximal Cohen-Macaulay module if the local cohomology

$$\mathrm{H}^{i}_{\mathfrak{m}}(N) = \lim_{m \to \infty} \underline{\mathrm{Ext}}^{i}_{S}(S/S_{\geq n}, N) = 0, \quad \text{if } i \neq d-1$$

where $\mathfrak{m} = S_{\geq 1}$. Denote the full subcategory of $\operatorname{gr}_{\mathbb{Z}} S$ consisting of all the maximal Cohen-Macaulay modules by mcm S and the stable category of mcm S by mcm S which is a triangulated category.

In this paper, we use the Clifford deformation introduced by He-Ye in [2] to discuss $\underline{\mathrm{mcm}}(S)$. Recall some results in [2] here. By [15, Proposition 5.1], $A^!$ is a Koszul Frobenius algebra. Choose $\hat{z} \in V^{\otimes 2}$ as a lift of z. There is a Clifford map

satisfying $(\theta_z \otimes 1 - 1 \otimes \theta_z)(V^* \otimes R^{\perp} \cap R^{\perp} \otimes V^*) = 0$. Note that the map θ_z is independent of the choice of \hat{z} . Then we have an algebra

$$C_{A^{!}}(z) = T(V^{*})/(f - \theta_{z}(f) : f \in R^{\perp}),$$

which is called a *Clifford deformation* of $A^!$. The \mathbb{Z} -graded algebra $T(V^*)$ is also a \mathbb{Z}_2 -graded algebra if we set the subspace $\bigoplus_{n\geq 0} (V^*)^{\otimes 2n}$ with degree 0 and $\bigoplus_{n\geq 0} (V^*)^{\otimes 2n+1}$ with degree 1. Then $C_{A^!}(z)$ is also a \mathbb{Z}_2 -graded algebra induced by the \mathbb{Z}_2 -grading of $T(V^*)$.

Theorem 1.4. [2, Theorem 0.1 and Theorem 0.2] Retain the notations above. Then

- (1) $C_{A^{!}}(z)$ is a strongly \mathbb{Z}_2 -graded Frobenius algebra;
- (2) S is a noncommutative graded isolated singularity if and only if $C_{A^{!}}(z)$ is a \mathbb{Z}_{2} -graded semisimple algebra;
- (3) there are equivalences of triangulated categories

$$\underline{\mathrm{mcm}} S \cong D^{b}(\mathrm{gr}_{\mathbb{Z}_{2}} C_{A^{!}}(z)) \cong D^{b}(\mathrm{mod} C_{A^{!}}(z)_{0}).$$

Let $\sigma = (\sigma_{ij}) : A \to M_2(A)$ be an algebra homomorphism such that each σ_{ij} is a \mathbb{Z} -graded linear transformation of A for any i, j = 1, 2. Write $\sigma_{|_V} = (\sigma_{ij}) : V \to M_2(V)$ for the restriction of σ on $A_1 = V$. One can extend $\sigma_{|_V}$ to an algebra homomorphism

(1.1)
$$\widetilde{\sigma} = (\widetilde{\sigma}_{ij}) : T(V) \to M_2(T(V))$$

such that

$$\widetilde{\sigma}(R) \subseteq M_2(R),$$

where each $\tilde{\sigma}_{ij}$ is a Z-graded linear transformation of T(V) for any i, j = 1, 2. Applying the Z-graded linear dual functor $(-)^*$ to each component of $\tilde{\sigma}$, one obtains an algebra homomorphism

(1.2)
$$\widetilde{\sigma}^* = (\widetilde{\sigma}^*_{ij}) : T(V^*) \to M_2(T(V^*)).$$

For any $f \in \mathbb{R}^{\perp}$, $r \in \mathbb{R}$ and i, j = 1, 2, we have $\tilde{\sigma}_{ij}^*(f)(r) = f(\tilde{\sigma}_{ij}(r)) = 0$. It implies

(1.3)
$$\widetilde{\sigma}^*(R^{\perp}) \subset M_2(R^{\perp}).$$

Suppose $\sigma_{ij}(z) = \delta_{ij}z$ for any i, j = 1, 2. Then the lift map $\tilde{\sigma} = (\tilde{\sigma}_{ij})$ defined in (1.1) satisfies

$$\widetilde{\sigma}_{ij}(\hat{z}) - \delta_{ij}\hat{z} \in R,$$

for any i, j = 1, 2. So the dual map $\tilde{\sigma}^* = (\tilde{\sigma}^*_{ij}) : T(V^*) \to M_2(T(V^*))$ defined in (1.2) satisfies $\tilde{\sigma}^*_{ij}(f - f(\hat{z})) = \tilde{\sigma}^*_{ij}(f) - \delta_{ij}f(\hat{z}) = \tilde{\sigma}^*_{ij}(f) - f(\tilde{\sigma}_{ij}(\hat{z})) = \tilde{\sigma}^*_{ij}(f) - \tilde{\sigma}^*_{ij}(f)(\hat{z}) \in (g - g(\hat{z}) \mid g \in R^{\perp}),$ for any $f \in R^{\perp}$ and i, j = 1, 2, where the first equality holds since $\tilde{\sigma}^*$ is an algebra homomorphism and $f(\hat{z}) \in k$, and the final belonging holds by (1.3). Hence, $\tilde{\sigma}^*$ induces an algebra homomorphism

(1.4)
$$\sigma^{!} = (\sigma^{!}_{ij}) : C_{A^{!}}(z) \to M_{2}(C_{A^{!}}(z)),$$

where each $\sigma_{ij}^!$ is a \mathbb{Z}_2 -graded linear transformation of $C_{A'}(z)$ for any i, j = 1, 2.

Lemma 1.5. If σ is invertible, then $\sigma^!$ is t-invertible.

Proof. Write φ for the inverse of σ . We have $\delta_{ij}z = \sum_{k=1}^{2} \varphi_{ik}\sigma_{jk}(z) = \varphi_{ij}(z)$ for i, j = 1, 2. Hence, φ' is well defined by (1.4) and is the *t*-inverse of σ' .

We close this section by introducing a class of extensions of algebras, which is called *semi-trivial* extension in [12].

Theorem 1.6. Let E be an algebra and M be an E-bimodule. Let $\psi : M \otimes_E M \to E$ be an E-bimodule homomorphism such that $m\psi(m' \otimes m'') = \psi(m \otimes m')m''$ for any $m, m', m'' \in M$. Then the vector space $E \oplus M$ with the multiplication

$$(x,m) \odot (x',m') = (xx' + \psi(m \otimes m'), xm' + mx'), \qquad \forall x, x' \in E, m, m' \in M,$$

is an algebra, called a semi-trivial extension of E by M through ψ , denoted by $E \ltimes_{\psi} M$.

Proof. It is clear (1,0) is the identity. Since ψ is an *E*-bimodule homomorphism and $m\psi(m'\otimes m'') = \psi(m \otimes m')m''$ for any $m, m', m'' \in M$, the associativity of this multiplication holds.

Remark 1.7. (1) The semi-trivial extension $E \ltimes_{\psi} M$ is the trivial extension $E \ltimes M$ if $\psi = 0$.

- (2) A semi-trivial extension $E \ltimes_{\psi} M$ can be viewed as a \mathbb{Z}_2 -graded algebra (or, a superalgebra) if $\deg(x, 0) = 0$, $\deg(0, m) = 1$ for any $x \in E, m \in M$.
- (3) Suppose E is a G-graded algebra, M is a G-graded E-bimodule and ψ is a G-graded map where G is an abelian group. Then $E \ltimes_{\psi} M$ is a $\mathbb{Z}_2 \times G$ -graded algebra whose homogeneous space $(E \ltimes_{\psi} M)_{(0,g)} = E_g$ and $(E \ltimes_{\psi} M)_{(1,g)} = M_g$ for any $g \in G$, and also a G-graded algebra by forgetting the first \mathbb{Z}_2 -degree.

In particular, if $G = \mathbb{Z}_2$, we always view $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded algebra $E \ltimes_{\psi} M$ as a \mathbb{Z}_2 -graded algebra by forgetting the first \mathbb{Z}_2 -degree in the sequel, that is,

$$(E \ltimes_{\psi} M)_0 = E_0 \oplus M_0, \qquad (E \ltimes_{\psi} M)_1 = E_1 \oplus M_1.$$

Example 1.8. (1) Let *E* be an algebra and $\psi : E \otimes_E E \to E$ be the multiplication of *E*. Then one obtains a semi-trivial extension $E \ltimes_{\psi} E$.

(2) Let E be a \mathbb{Z}_2 -graded algebra and μ be a \mathbb{Z}_2 -graded algebra automorphism of E such that $\mu^2 = \text{id.}$ Then

$$\psi: \ _{\mu}E(1) \otimes_{E} _{\mu}E(1) \rightarrow E$$
$$a \otimes b \qquad \mapsto \quad \mu(a)b,$$

is a \mathbb{Z}_2 -graded *E*-bimodule homomorphism and satisfies $\psi(a \otimes b) \cdot c = a\psi(b \otimes c)$ for any $a, b, c \in {}_{\mu}E(1)$. Hence, we have a semi-trivial extension $E \ltimes_{\psi}({}_{\mu}E(1))$ which is a \mathbb{Z}_2 -graded algebra (see Remark 1.7(3) when $G = \mathbb{Z}_2$). More precisely, $(E \ltimes_{\psi}({}_{\mu}E(1)))_0 = E_0 \oplus E_1$ and $(E \ltimes_{\psi}({}_{\mu}E(1)))_1 = E_1 \oplus E_0$ with multiplication

$$(a,b) \odot (a',b') = (aa' + \mu(b)b', \mu(a)b' + ba'),$$

for any homogeneous elements $(a, b), (a', b') \in E \ltimes_{\psi} (\mu E(1))$.

2. Two Classes of Twisted Structures

In this section, we introduce a class of twisting systems for matrix algebra $M_2(E)$ over some algebra E and a class of twisted matrix algebras. As an application, there is a twisted structure on the direct product $E \times E$ a by choosing a subalgebra of a twisted matrix algebra.

2.1. Twisted matrix algebras. Let $M_2(\Bbbk)$ be a \mathbb{Z}_2 -graded algebra whose \mathbb{Z}_2 -grading is given by

$$M_2(\mathbb{k})_0 = \begin{pmatrix} \mathbb{k} & 0 \\ 0 & \mathbb{k} \end{pmatrix}, \qquad M_2(\mathbb{k})_1 = \begin{pmatrix} 0 & \mathbb{k} \\ \mathbb{k} & 0 \end{pmatrix}.$$

Let $\mathbb{I}_1^{(0)}, \mathbb{I}_2^{(0)}$ be invertible and a basis of $M_2(k)_0$, and $\mathbb{I}_1^{(1)}, \mathbb{I}_2^{(1)}$ be invertible and a basis of $M_2(k)_1$. We say $\mathbb{I} = \{\mathbb{I}_j^{(i)} \mid i \in \mathbb{Z}_2, j = 1, 2\}$ is an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$. Let $\gamma_1, \gamma_2 \in \mathbb{k}$ such that

$$\gamma_1 \mathbb{I}_1^{(0)} + \gamma_2 \mathbb{I}_2^{(0)} = I_2$$

where I_2 is the 2 × 2 identity matrix. For any $i, i' \in \mathbb{Z}_2, j = 1, 2$, write

(2.1)
$$L_{j}^{(ii')} = \begin{pmatrix} l_{1j1}^{(ii')} & l_{1j2}^{(ii')} \\ l_{2j1}^{(ii')} & l_{2j2}^{(ii')} \end{pmatrix} \in M_{2}(\mathbb{k}),$$

such that

$$\mathbb{I}_{j}^{(i)}(\mathbb{I}_{1}^{(i')},\mathbb{I}_{2}^{(i')}) = (\mathbb{I}_{1}^{(i+i')},\mathbb{I}_{2}^{(i+i')})L_{j}^{(ii')}.$$

More precisely, for any $i, i' \in \mathbb{Z}_2, j, j' = 1, 2$,

(2.2)
$$\mathbb{I}_{j}^{(i)}\mathbb{I}_{j'}^{(i')} = \sum_{s=1}^{2} \mathbb{I}_{s}^{(i+i')}l_{sjj'}^{(ii')}$$

It is not hard to check that each $L_j^{(ii')}$ is invertible for any $i, i' \in \mathbb{Z}_2$ and j = 1, 2.

(1) For any $i, i', i'' \in \mathbb{Z}_2$ and j, j', j'', t = 1, 2, Lemma 2.1.

$$\sum_{s=1}^{2} l_{tjs}^{(i,i'+i'')} l_{sj'j''}^{(i'i'')} = \sum_{s=1}^{2} l_{tsj''}^{(i+i',i'')} l_{sjj'}^{(ii')}.$$

- (2) For any $i \in \mathbb{Z}_2$ and $s, j = 1, 2, \sum_{t=1}^2 l_{sjt}^{(i0)} \gamma_t = \delta_{sj}$. (3) For any $i \in \mathbb{Z}_2$ and $s, t = 1, 2, \sum_{j=1}^2 l_{sjt}^{(0i)} \gamma_j = \delta_{st}$.

Proof. (1) It follows by the associativity of $\mathbb{I}_{i}^{(i)}\mathbb{I}_{i'}^{(i)}\mathbb{I}_{i''}^{(i'')}$.

(2) By (2.2) with i' = 0, $\sum_{s=1}^{2} \left(\sum_{t=1}^{2} l_{sjt}^{(i0)} \gamma_t \right) \mathbb{I}_s^{(i)} = \sum_{t=1}^{2} \mathbb{I}_j^{(i)} \mathbb{I}_t^{(0)} \gamma_t = \mathbb{I}_j^{(i)}$.

(3) By (2.2) with
$$i = 0$$
, $\sum_{s=1}^{2} \left(\sum_{j=1}^{2} \gamma_{j} l_{sjt}^{(0i')} \right) \mathbb{I}_{s}^{(i')} = \sum_{j=1}^{2} \gamma_{j} \mathbb{I}_{j}^{(0)} \mathbb{I}_{t}^{(i')} = \mathbb{I}_{t}^{(i')}$.

Let E be an algebra. The basis I of $M_2(\mathbb{k})$ provides a decomposition for \mathbb{Z}_2 -graded vector space $M_2(E)$ as follows, for any $i \in \mathbb{Z}_2$

$$M_2(E)_i = \left\{ \mathbb{I}_1^{(i)} x \mid x \in E \right\} \oplus \left\{ \mathbb{I}_2^{(i)} x \mid x \in E \right\}$$

To introduce a twisted system of the matrix algebra $M_2(E)$, we consider a class of linear maps from E to the matrix algebra $M_2(E)$.

Proposition 2.2. Let $\theta^{(i)} = (\theta^{(i)}_{jj'}) : E \to M_2(E)$ be a linear map for $i \in \mathbb{Z}_2$. Then the following are equivalent:

Proof. $(1) \Leftrightarrow (2)$ follows by Lemma 2.1(1).

 $(2) \Rightarrow (3)$ Write $(L_j^{(ii')})^{-1} = (g_{j'jj''}^{(ii')})$ for any $i, i' \in \mathbb{Z}_2$ and j, j', j'' = 1, 2. Multiplying the equation in (2) by $g_{pjv}^{(i,i'+i'')}$ on both sides and taking the sum over v from 1 to 2 for any p = 1, 2, one obtains (3).

(3) \Rightarrow (2) Multiply the equation in (3) by $l_{vjp}^{(i,i'+i'')}$ for any j, v = 1, 2 on both sides and take the sum over p from 1 to 2.

Lemma 2.3. Let $\theta^{(i)} = (\theta^{(i)}_{jj'}) : E \to M_2(E)$ be a linear map for $i \in \mathbb{Z}_2$. Suppose $\theta^{(0)}$ and $\theta^{(1)}$ are *t*-invertible with the *t*-inverses $\varphi^{(0)}$ and $\varphi^{(1)}$ respectively, and satisfy the equivalent conditions of Proposition 2.2. Then

$$\sum_{u,j=1}^{2} l_{pqu}^{(ii')} \theta_{uj}^{(i')} \left(x \varphi_{rj}^{(i')}(x') \right) = \sum_{t,j=1}^{2} l_{tjr}^{(ii')} \theta_{pt}^{(i+i')} (\varphi_{qj}^{(i)}(x)) x'$$

for any $x, x' \in E$, $i, i' \in \mathbb{Z}_2$ and p, q, r = 1, 2.

Proof. Rewrite the indexes of Proposition 2.2(3):

$$\sum_{s,u=1}^{2} l_{psu}^{(ii')} \theta_{uj'}^{(i')} \left(\theta_{sj}^{(i)}(x)x' \right) = \sum_{t,u=1}^{2} l_{tju}^{(ii')} \theta_{pt}^{(i+i')}(x) \theta_{uj'}^{(i')}(x'),$$

for any $x, x' \in E$, $i, i' \in \mathbb{Z}_2$ and j, j', p = 1, 2.

Replacing x and x' by $\varphi_{qj}^{(i)}(x)$ and $\varphi_{rj'}^{(i')}(x')$ respectively and taking the sum over j, j' both from 1 to 2, the result follows.

Lemma 2.4. Retain the assumptions of Lemma 2.3. Then $\theta^{(0)}(1) \in GL_2(\mathbb{k})$ if and only if $\theta^{(1)}(1) \in GL_2(\mathbb{k})$. In this case, $\varphi^{(i)}(1) \in GL_2(\mathbb{k})$ for $i \in \mathbb{Z}_2$.

Proof. Suppose
$$\theta^{(0)}(1) \in GL_2(\mathbb{k}), \ (\varphi^{(0)}(1))^T \theta^{(0)}(1) = ((\varphi^{(0)})^T \theta^{(0)})(1) = I_2.$$
 So $\varphi^{(0)}(1) \in GL_2(\mathbb{k}).$

Taking x = x' = 1 and i = 0, i' = 1 in Lemma 2.3, we have

$$l_{pqr}^{(01)} = \sum_{t,j=1}^{2} l_{tjr}^{(01)} \theta_{pt}^{(1)}(\varphi_{qj}^{(0)}(1)) = \sum_{t,j=1}^{2} l_{tjr}^{(01)} \theta_{pt}^{(1)}(1) \varphi_{qj}^{(0)}(1)$$

for any p, q, r = 1, 2. Write those equations in the form of matrices:

$$\overline{L}_{1,2;r}^{(01)} = \theta^{(1)}(1)\overline{L}_{1,2;r}^{(01)}(\varphi^{(0)}(1))^T,$$

where $\overline{L}_{1,2;r}^{(01)} = \begin{pmatrix} l_{11r}^{(01)} & l_{12r}^{(01)} \\ l_{21r}^{(01)} & l_{22r}^{(01)} \end{pmatrix}$ for any r = 1, 2. Note that for any $r = 1, 2, \{\mathbb{I}_1^{(0)}\mathbb{I}_r^{(1)}, \mathbb{I}_2^{(0)}\mathbb{I}_r^{(1)}\}$ is also a basis of $M_2(k)_1$ and

$$(\mathbb{I}_1^{(0)}\mathbb{I}_r^{(1)},\mathbb{I}_2^{(0)}\mathbb{I}_r^{(1)}) = (\mathbb{I}_1^{(1)},\mathbb{I}_2^{(1)})\overline{L}_{1,2;r}^{(01)}.$$

So $\overline{L}_{1,2;r}^{(01)}$ is invertible and

$$\theta^{(1)}(1) = \overline{L}_{1,2;r}^{(01)} \theta^{(0)}(1) \left(\overline{L}_{1,2;r}^{(01)}\right)^{-1} \in GL_2(\mathbb{k}).$$

Similar to the invertibility of $\varphi^{(0)}(1)$, one obtains that $\varphi^{(1)}(1) \in GL_2(\mathbb{k})$.

Similarly, if $\theta^{(1)}(1) \in GL_2(\mathbb{k})$, then $\varphi^{(1)}(1) \in GL_2(\mathbb{k})$, and

$$\theta^{(0)}(1) = \overline{L}_{1,2;r}^{(11)} \theta^{(1)}(1) \left(\overline{L}_{1,2;r}^{(11)}\right)^{-1} \in GL_2(\mathbb{k}),$$

where $\overline{L}_{1,2;r}^{(11)} = \begin{pmatrix} l_{11r}^{(11)} & l_{12r}^{(11)} \\ l_{21r}^{(11)} & l_{22r}^{(11)} \end{pmatrix}$ is invertible for r = 1, 2.

Lemma 2.5. Retain the assumptions of Lemma 2.3. Suppose $\theta^{(0)}(1) \in GL_2(k)$ (or $\theta^{(1)}(1) \in$ $GL_2(\mathbb{k})).$

- (1) For any $i, i' \in \mathbb{Z}_2$ and q, r, s = 1, 2, then $\sum_{p=1}^2 l_{pqr}^{(ii')} \varphi_{ps}^{(i+i')}(1) = \sum_{j=1}^2 l_{sjr}^{(ii')} \varphi_{qj}^{(i)}(1)$. (2) For any $x \in E$ and v = 1, 2, then $\sum_{r,j=1}^2 \theta_{vj}^{(0)} \left(x \varphi_{rj}^{(0)}(1) \right) \gamma_r = x \gamma_v$. (3) For any $i \in \mathbb{Z}_2$ and q, s = 1, 2, then $\sum_{t,j,u=1}^2 \gamma_u l_{sjt}^{(0i)} \varphi_{uj}^{(0)}(1) \theta_{tq}^{(i)}(1) = \delta_{sq}$.

Proof. (1) Taking $x = 1, x' = \varphi_{ps}^{(i+i')}(1)$ in Lemma 2.3 and by Lemma 2.4, the result follows.

(2) Taking i' = 0 and x' = 1 in Lemma 2.3, one obtains that

$$\sum_{u,j=1}^{2} l_{pqu}^{(i0)} \theta_{uj}^{(0)} \left(x \varphi_{rj}^{(0)}(1) \right) = \sum_{t,j=1}^{2} l_{tjr}^{(i0)} \theta_{pt}^{(i)} (\varphi_{qj}^{(i)}(x)),$$

for any p, q, r = 1, 2. By Lemma 2.1(2), we have

$$\sum_{r,u,j=1}^{2} l_{pqu}^{(i0)} \theta_{uj}^{(0)} \left(x \varphi_{rj}^{(0)}(1) \right) \gamma_r = \sum_{r,t,j=1}^{2} l_{tjr}^{(i0)} \theta_{pt}^{(i)} (\varphi_{qj}^{(i)}(x)) \gamma_r = \sum_{j=1}^{2} \theta_{pj}^{(i)} (\varphi_{qj}^{(i)}(x)) = \delta_{pq} x,$$

and then multiply the equation above by $g_{vqp}^{(i0)}$ which is the (v, p)-th element of $(L_q^{(i0)})^{-1}$ for any v, p, q = 1, 2. The result follows.

(3) By Lemma 2.1(3), Lemma 2.4 and taking i = 0 in (1), one obtains

$$\sum_{t,j,u=1}^{2} \gamma_{u} l_{sjt}^{(0i)} \varphi_{uj}^{(0)}(1) \theta_{tq}^{(i)}(1) = \sum_{p,t,u=1}^{2} \gamma_{u} l_{put}^{(0i)} \varphi_{ps}^{(i)}(1) \theta_{tq}^{(i)}(1) = \sum_{t=1}^{2} \varphi_{ts}^{(i)}(1) \theta_{tq}^{(i)}(1) = \delta_{sq}.$$

Definition 2.6. Let $\theta^{(i)} = (\theta^{(i)}_{jj'}) : E \to M_2(E)$ be a *t*-invertible linear map for $i \in \mathbb{Z}_2$ and $\mathbb{I} = \{\mathbb{I}_j^{(i)} \mid i \in \mathbb{Z}_2, j = 1, 2\} \text{ be an invertible } \mathbb{Z}_2 \text{-graded basis of } M_2(\mathbb{k}). \text{ The pair } \Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ is called a *twisting system* of $M_2(E)$ if

(1) $\theta^{(0)}, \theta^{(1)}$ satisfy equivalent conditions of Proposition 2.2,

(2)
$$\theta^{(1)}(1) \in GL_2(\mathbb{k})$$
 (or equivalently, $\theta^{(0)}(1) \in GL_2(\mathbb{k})$).

Theorem 2.7. Let $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ be a twisting system of $M_2(E)$. Then the \mathbb{Z}_2 -graded vector space $M_2(E)$ with the multiplication as follows, for any $x, x' \in E, i, i' \in \mathbb{Z}_2$ and j, j' = 1, 2,

$$\mathbb{I}_{j}^{(i)}x \star \mathbb{I}_{j'}^{(i')}x' = \sum_{s=1}^{2} \mathbb{I}_{j}^{(i)}\mathbb{I}_{s}^{(i')}\theta_{sj'}^{(i')}(x)x' = \sum_{s,t=1}^{2} \mathbb{I}_{t}^{(i+i')}l_{tjs}^{(ii')}\theta_{sj'}^{(i')}(x)x',$$

is a \mathbb{Z}_2 -graded algebra with the identity $\sum_{j,s=1}^2 \gamma_s(\theta^{(0)})_{sj}^{-1}(1)\mathbb{I}_j^{(0)}$, called this new algebra the twisted matrix algebra of $M_2(E)$ by Θ , denoted by $\Theta M_2(E)$.

Proof. It is straightforward to check the multiplication \star satisfies the associative law by Proposition 2.2(1).

Write $\varphi^{(0)}$ for the *t*-inverse of $\theta^{(0)}$ and $\mathbb{I}^{(0)} = \sum_{j,s=1}^{2} \gamma_s \varphi^{(0)}_{sj}(1) \mathbb{I}^{(0)}_j$. By Lemma 2.5(2) and Lemma 2.1(2), for any $x \in E$, $i \in \mathbb{Z}_2$ and j = 1, 2,

$$\mathbb{I}_{j}^{(i)}x \star \mathbb{I}^{(0)} = \sum_{s,t,j',u=1}^{2} \mathbb{I}_{t}^{(i)} l_{tjs}^{(i0)} \theta_{sj'}^{(0)}(x) \gamma_{u} \varphi_{uj'}^{(0)}(1) = \sum_{s,t=1}^{2} \mathbb{I}_{t}^{(i)} l_{tjs}^{(i0)} \gamma_{s} x = \mathbb{I}_{j}^{(i)} x.$$

On the other hand, by Lemma 2.5(3), for any $x \in E$, $i \in \mathbb{Z}_2$ and j = 1, 2,

$$\mathbb{I}^{(0)} \star \mathbb{I}_{j}^{(i)} x = \sum_{s,t,j',u=1}^{2} \mathbb{I}_{t}^{(i)} \gamma_{u} \varphi_{uj'}^{(0)}(1) l_{tj's}^{(0i)} \theta_{sj}^{(i)}(1) x = \mathbb{I}_{j}^{(i)} x.$$

Hence, $\mathbb{I}^{(0)}$ is the identity.

The last theorem provides a twisted algebra structure on the matrix algebra $M_2(E)$, but the identity seems complicated. It is natural to find a twisting system such that the identity matrix is the identity of the associated twisted matrix algebra. We have an analogous result of [20, Proposition 2.4].

Theorem 2.8. Let $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ be a twisting system of $M_2(E)$. Let $\Upsilon = (v^{(0)}, v^{(1)})$ be a pair of linear maps from E to $M_2(E)$ satisfying

$$v^{(i)}(x) = \theta^{(i)}(x)(\varphi^{(i)}(1))^T,$$

where $\varphi^{(i)}$ is the t-inverse of $\theta^{(i)}$ for any $i \in \mathbb{Z}_2$ and $x \in E$. Then

- (1) $v^{(i)}$ is t-invertible with the t-inverse map E from $M_2(E)$ mapping x to $\varphi^{(i)}(x)(\theta^{(i)}(1))^T$ for any $i \in \mathbb{Z}_2$;
- (2) $\Upsilon = (\{\upsilon^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ is a twisting system of $M_2(E)$;
- (3) the identity of the twisted matrix algebra ${}^{\Upsilon}M_2(E)$ is the identity matrix I_2 ;
- (4) $\Theta M_2(E) \cong \Upsilon M_2(E)$ as \mathbb{Z}_2 -graded algebras.

Proof. (1) It is straightforward.

(2) It is clear that $v^{(i)}(1) = \theta^{(i)}(1)(\varphi^{(i)}(1))^T = I_2 \in GL_2(\mathbb{k})$ for any $i \in \mathbb{Z}_2$ by Lemma 2.4. For any $x, x' \in E, i, i', i'' \in \mathbb{Z}_2$, and j, j', j'' = 1, 2,

$$\sum_{s,t,u=1}^{2} l_{vtu}^{(i+i',i'')} l_{tjs}^{(ii')} v_{uj''}^{(i'')} \left(v_{sj'}^{(i')}(x)x' \right) = \sum_{s,t,u,p,q=1}^{2} l_{vtu}^{(i+i',i'')} l_{tjs}^{(ii')} \theta_{up}^{(i'')} \left(\theta_{sq}^{(i')}(x)x' \right) \varphi_{j'q}^{(i')}(1) \varphi_{j''p}^{(i'')}(1)$$

$$= \sum_{s,t,u,p,q=1}^{2} l_{vjs}^{(i,i'+i'')} l_{tqu}^{(i'i'')} \theta_{st}^{(i'+i'')}(x) \theta_{up}^{(i'')}(x') \varphi_{j'q}^{(i')}(1) \varphi_{j''p}^{(i'')}(1)$$

$$= \sum_{s,t,u,p,q=1}^{2} l_{vjs}^{(i,i'+i'')} l_{qj'u}^{(i'i'')} \theta_{st}^{(i'+i'')}(x) \theta_{up}^{(i'')}(x') \varphi_{qt}^{(i'+i'')}(1) \varphi_{j''p}^{(i'')}(1)$$

$$= \sum_{s,u,q=1}^{2} l_{vjs}^{(i,i'+i'')} l_{qj'u}^{(i'i'')} v_{sq}^{(i'+i'')}(x) v_{uj''}^{(i'')}(x'),$$

where the second equality follows by the Proposition 2.2(1) and the third equality follows by Lemma 2.5(1).

(3) By Theorem 2.7, the identity of ${}^{\Upsilon}M_2(E)$ is

$$\sum_{j,s,t=1}^{2} \gamma_{s} \varphi_{st}^{(0)}(1) \theta_{jt}^{(0)}(1) \mathbb{I}_{j}^{(0)} = \sum_{j,s,t=1}^{2} \gamma_{s} \theta_{jt}^{(0)}(\varphi_{st}^{(0)}(1)) \mathbb{I}_{j}^{(0)} = \sum_{j=1}^{2} \gamma_{j} \mathbb{I}_{j}^{(0)} = I_{2}$$

(4) Define two \mathbb{Z}_2 -graded linear maps:

$$\begin{array}{ccccc} F: & {}^{\Theta}M_2(E) & \to & {}^{\Upsilon}M_2(E) & & G: & {}^{\Upsilon}M_2(E) & \to & {}^{\Theta}M_2(E) \\ & \mathbb{I}_j^{(i)}x & \mapsto & \sum_{s=1}^2 \mathbb{I}_s^{(i)}x\theta_{sj}^{(i)}(1), & & & \mathbb{I}_j^{(i)}x & \mapsto & \sum_{s=1}^2 \mathbb{I}_s^{(i)}x\varphi_{js}^{(i)}(1). \end{array}$$

It is straightforward to verify that F and G are the inverse to each other. Moreover, F is an algebra homomorphism, since for any $i, i' \in \mathbb{Z}_2$ and j = 1, 2

$$\begin{split} F\left(\mathbb{I}_{j}^{(i)}x\right) \star F\left(\mathbb{I}_{j'}^{(i')}x'\right) &= \sum_{p=1}^{2} \mathbb{I}_{p}^{(i)}x\theta_{pj}^{(i)}(1) \star \sum_{q=1}^{2} \mathbb{I}_{q}^{(i')}x'\theta_{qj'}^{(i')}(1) \\ &= \sum_{p,q,s,t=1}^{2} \mathbb{I}_{t}^{(i+i')}l_{tps}^{(ii')}v_{sq}^{(i)}(x\theta_{pj}^{(i)}(1))\theta_{qj'}^{(i')}(1)x' \\ &= \sum_{p,q,s,t,u=1}^{2} \mathbb{I}_{t}^{(i+i')}\theta_{pj}^{(i)}(1)\theta_{qj'}^{(i')}(1)l_{tps}^{(ii')}\theta_{su}^{(i)}(x\varphi_{qu}^{(i)}(1))x' \\ &= \sum_{p,t=1}^{2} \mathbb{I}_{t}^{(i+i')}\theta_{pj}^{(i)}(1)\left(\sum_{q,u,j''=1}^{2} l_{tpu}^{(ii')}\theta_{uj''}^{(i')}(x)(\theta_{qj'}^{(i')}(1))\right)x' \\ &= \sum_{p,t=1}^{2} \mathbb{I}_{t}^{(i+i')}\theta_{pj}^{(i)}(1)\left(\sum_{q,u,j''=1}^{2} l_{tpu}^{(ii')}\theta_{uj''}^{(i')}\varphi_{qj''}^{(i')}(x)(\theta_{qj'}^{(i')}(1))\right)x' \\ &= \sum_{p,u,t=1}^{2} \mathbb{I}_{t}^{(i+i')}\theta_{pj}^{(i)}(1)l_{tpu}^{(ii')}\theta_{uj'}^{(i')}(x)x' \\ &= \sum_{u,s,t=1}^{2} \mathbb{I}_{t}^{(i+i')}l_{sju}^{(ii')}\theta_{ts}^{(i+i')}(1)\theta_{uj'}^{(i')}(x)x' \\ &= \sum_{u,s,t=1}^{2} \mathbb{I}_{t}^{(i+i')}l_{sju}^{(ii')}\theta_{ts}^{(i+i')}(1)\theta_{uj'}^{(i')}(x)x' \\ &= F\left(\mathbb{I}_{j}^{(i)}x \star \mathbb{I}_{j'}^{(i')}x'\right), \end{split}$$

where the fourth and the fifth equalities are by Lemma 2.3, and the seventh equality is by Proposition 2.2(c).

Before that, the twisting systems on matrix algebras depend on an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$. In fact, we can fix an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$ to obtain twisted matrix algebras.

Theorem 2.9. Let $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ be a twisting system of $M_2(E)$. Suppose $\mathbb{J} = \{\mathbb{J}_j^{(i)} \mid i \in \mathbb{Z}_2, j = 1, 2\}$ be an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$. Then there exists a twisting system $\Omega = (\{\omega^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{J})$ of $M_2(E)$ such that $\Theta M_2(E) \cong \Omega M_2(E)$ as \mathbb{Z}_2 -graded algebras.

Proof. There exist two invertible matrices $U^{(0)}$ and $U^{(1)}$ of $M_2(\Bbbk)$ such that

$$(\mathbb{I}_1^{(i)}, \mathbb{I}_2^{(i)}) = (\mathbb{J}_1^{(i)}, \mathbb{J}_2^{(i)})U^{(i)},$$

for any $i \in \mathbb{Z}_2$. Write $\Omega = (\omega^{(0)}, \omega^{(1)})$, where

$$y^{(i)}: E \rightarrow M_2(E)$$

 $x \mapsto U^{(i)}\theta^{(i)}(x)(U^{(i)})^{-1}$

for any $i \in \mathbb{Z}_2$. It is straightforward to check that $\Omega = (\{\omega^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{J})$ is a twisting system of $M_2(E)$ by Definition 2.6.

It is also not hard to obtain that the isomorphism between ${}^{\Theta}M_2(E)$ and ${}^{\Omega}M_2(E)$ as \mathbb{Z}_2 -graded vector spaces defined by $U^{(0)}$ and $U^{(1)}$ is an algebra isomorphism.

The following result follows by Theorem 2.8 and Theorem 2.9 immediately.

Corollary 2.10. Let $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ be a twisting system of $M_2(E)$. Suppose $\mathbb{J} = \{\mathbb{J}_j^{(i)} | i \in \mathbb{Z}_2, j = 1, 2\}$ be an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$. Then there exists a twisting system $\Omega = (\{\omega^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{J})$ of $M_2(E)$ such that $\Theta M_2(E) \cong \Omega M_2(E)$ as \mathbb{Z}_2 -graded algebras and the identity of $\Omega M_2(E)$ is I_2 .

This corollary implies that we can always fix an invertible \mathbb{Z}_2 -graded basis of $M_2(\mathbb{k})$ and assume the identity of a twisted matrix algebra is I_2 .

We turn to graded cases. Let E be a G-graded algebra where G is an abelian group. Suppose $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ is a twisting system of $M_2(E)$ where each $\theta_{jj'}^{(i)}$ is a G-graded linear map for any $i \in \mathbb{Z}_2$ and j, j' = 1, 2. Then the twisted matrix algebra $\Theta M_2(E)$ is a $\mathbb{Z}_2 \times G$ -graded algebra, where each component is

$${}^{\Theta}M_2(E)_{(i,g)} = \left\{ \mathbb{I}_1^{(i)} x \mid x \in E_g \right\} \oplus \left\{ \mathbb{I}_2^{(i)} x \mid x \in E_g \right\},$$

for any $i \in \mathbb{Z}_2$ and $g \in G$.

Remark 2.11. In particular, if E is a \mathbb{Z}_2 -graded algebra, then $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded algebra ${}^{\Theta}M_2(E)$ has the following two approaches to becoming a \mathbb{Z}_2 -graded algebra.

(1) By taking the total degrees, ${}^{\Theta}M_2(E)$ becomes a \mathbb{Z}_2 -graded algebra where

$${}^{\Theta}M_2(E)_0 = \begin{pmatrix} E_0 & E_1 \\ E_1 & E_0 \end{pmatrix}, \qquad {}^{\Theta}M_2(E)_1 = \begin{pmatrix} E_1 & E_0 \\ E_0 & E_1 \end{pmatrix}$$

(2) By forgetting the first degree, ${}^{\Theta}M_2(E)$ becomes a \mathbb{Z}_2 -graded algebra where

$${}^{\Theta}M_2(E)_0 = \begin{pmatrix} E_0 & E_0 \\ E_0 & E_0 \end{pmatrix}, \qquad {}^{\Theta}M_2(E)_1 = \begin{pmatrix} E_1 & E_1 \\ E_1 & E_1 \end{pmatrix}.$$

These are the only two good \mathbb{Z}_2 -grading on $M_2(E)$ (see [11]).

Remark 2.12. Let E be a \mathbb{Z}_2 -graded algebra. By taking the total degrees, ${}^{\Theta}M_2(E)$ and ${}^{\Theta'}M_2(E)$ become \mathbb{Z}_2 -graded for some twisting systems Θ and Θ' of $M_2(E)$. In general, $\operatorname{gr}_{\mathbb{Z}_2} {}^{\Theta}M_2(E)$ and $\operatorname{gr}_{\mathbb{Z}_2} {}^{\Theta'}M_2(E)$ are not equivalent and even not derived equivalent (see Example 4.9). It is different from the one of Zhang-twists (see [20, Theorem 3.1]).

2.2. Twisted direct products. Let E be an algebra and $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ be a twisting system of $M_2(E)$. The 0-th homogeneous space

$$\begin{pmatrix} E & 0 \\ 0 & E \end{pmatrix}$$

of \mathbb{Z}_2 -graded algebra ${}^{\Theta}M_2(E)$ provides a subalgebra of ${}^{\Theta}M_2(E)$, which is isomorphic to $E \times E$ as vector spaces. It implies that there is a kind of twisted algebra structure on $E \times E$ induced by the one of $M_2(E)$. It is analogous to Definition 2.6 and Theorem 2.7, we just list them.

Let $\varepsilon_1 = (a_1, a_2), \varepsilon_2 = (b_1, b_2)$ be an invertible k-basis of $\mathbb{k} \times \mathbb{k}$ where $a_1 a_2 b_1 b_2 \neq 0$, and $\{l_{p;jj'} \mid p, j, j' = 1, 2\} \subseteq \mathbb{k}$ such that $\varepsilon_j \varepsilon'_j = \sum_{p=1}^2 \varepsilon_p l_{p;jj'}$ for any j, j' = 1, 2.

Definition 2.13. Let $\theta = (\theta_{jj'}) : E \to M_2(E)$ be a *t*-invertible linear map and $\varepsilon = \{\varepsilon_1, \varepsilon_2\}$ be an invertible basis of $\mathbb{k} \times \mathbb{k}$. We say $\theta_{\times} = (\theta, \varepsilon)$ is a *twisting system* of $E \times E$, if $\theta(1) \in GL_2(\mathbb{k})$ and

$$\sum_{s,u=1}^{2} l_{p;su} \theta_{uj'} \left(\theta_{sj}(x) x' \right) = \sum_{t,u=1}^{2} l_{t;ju} \theta_{pt}(x) \theta_{uj'}(x'),$$

for any $x, x' \in E$ and j, j', p = 1, 2.

Corollary 2.14. Let $\theta_{\times} = (\theta, \varepsilon)$ be a twisting system of $E \times E$. Then the vector space $E \times E$ with the multiplication

$$\varepsilon_j x \star \varepsilon_{j'} x' = \sum_{s,t=1}^2 \varepsilon_t l_{t;js} \theta_{sj'}(x) x', \qquad \forall x, x' \in E, j, j' = 1, 2,$$

is an algebra, called the twisted direct product of $E \times E$ by θ , denoted by $\theta_{\times}(E \times E)$.

In particular, if $\theta : E \to M_2(E)$ is the diagonal map, then $\theta_{\times} = (\theta, \varepsilon)$ is always a twisting system of $E \times E$ for any invertible k-basis ε and $\theta_{\times}(E \times E)$ is just the usual direct product $E \times E$.

Remark 2.15. To distinguish from Zhang-twisting system, we add a subscript " \times " for θ to represent a twisting system of a direct product.

Let *E* be a *G*-graded algebra where *G* is an abelian group. Suppose $\theta_{\times} = (\theta, \varepsilon)$ is a twisting system of $E \times E$, where each $\theta_{jj'}$ is a *G*-graded linear map for any j, j' = 1, 2. Then the twisted direct product $\theta_{\times}(E \times E)$ is a *G*-graded algebra, where for any $g \in G$,

$${}^{\theta_{\times}}(E \times E)_a = E_g \oplus E_g.$$

3. Double Ore extensions

In this section, we recall the definition of double Ore extensions in [21] and show some basic properties of noncommutative quadric hypersurfaces defined by double Ore extensions.

Definition 3.1. We say B is a \mathbb{Z} -graded right double Ore extension of a \mathbb{Z} -graded algebra A if B generated by A and two homogeneous variables y_1, y_2 satisfying

- (1) a homogeneous relation $y_2y_1 = p_{12}y_1y_2 + p_{11}y_1^2 + \tau_1y_1 + \tau_2y_2 + \tau_0$ for some $p_{12}, p_{11} \in \mathbb{k}$ and homogeneous elements $\tau_0, \tau_1, \tau_2 \in A$,
- (2) B is a \mathbb{Z} -graded free left A-modules with basis $\{y_1^i y_2^j \mid i, j \ge 0\},\$
- (3) $y_1A + y_2A \subseteq Ay_1 + Ay_2 + A$.

Similarly, one can define a \mathbb{Z} -graded left double Ore extension. An algebra B is called a \mathbb{Z} -graded double Ore extension of A if it is a \mathbb{Z} -graded left and a \mathbb{Z} -graded right double Ore extension of A with the same generating set $\{y_1, y_2\}$.

The condition (3) of the definition of \mathbb{Z} -graded right double Ore extension is equivalent to the existence of an algebra homomorphism $\sigma = (\sigma_{ij}) : A \to M_2(A)$ and a σ -derivation $\delta = (\delta_i) : A \to A^{\oplus 2}$ where all σ_{ij} and δ_i are \mathbb{Z} -graded linear transformations of A for i, j = 1, 2, such that

$$y_i a = \sigma_{i1}(a)y_1 + \sigma_{i2}(a)y_2 + \delta_i(a),$$

for any $a \in A$ and i = 1, 2. We write $A_P[y_1, y_2; \sigma, \delta, \tau]$ for a \mathbb{Z} -graded right double Ore extension where $P = \{p_{12}, p_{11}\}$ and $\tau = \{\tau_0, \tau_1, \tau_2\}$, and write $A_P[y_1, y_2; \sigma]$ for convenience if $\tau = \{0, 0, 0\}$ and $\delta = 0$.

We will focus on the trimmed case $A_P[y_1, y_2; \sigma]$ in the sequel. By [21, Lemma 1.10], σ in the trimmed case $A_P[y_1, y_2; \sigma]$ satisfies following conditions:

(3.1)

$$\sigma_{21}\sigma_{11} = p_{12}\sigma_{11}\sigma_{21},$$

$$\sigma_{22}\sigma_{12} = p_{12}\sigma_{12}\sigma_{22},$$

$$\sigma_{21}\sigma_{12} + p_{12}\sigma_{22}\sigma_{11} = p_{12}\sigma_{11}\sigma_{22} + p_{12}^{2}\sigma_{12}\sigma_{21}.$$

There is a conjecture about the relation between the invertibility of σ and the condition a right double Ore extension being a double Ore extension (see [21]). It is true in the connected graded case by [21, Lemma 1.9 and Proposition 1.13].

Lemma 3.2. Let A be a locally finite connected \mathbb{Z} -graded algebra and $B = A_P[y_1, y_2; \sigma]$ be a connected \mathbb{Z} -graded right double Ore extension of A where $p_{12} \neq 0$. Then B is a \mathbb{Z} -graded double Ore extension of A if and only if σ is invertible.

Let A = T(V)/(R) be a noetherian Koszul Artin-Schelter regular algebra and $B = A_P[y_1, y_2; \sigma]$ be a Z-graded double Ore extension of A with $p_{12} \neq 0$ and deg $y_1 = \deg y_2 = 1$. By [23, Theorem 2.1] (or, [14, Proposition 2.10]) and [21, Theorem 0.2], B is a Koszul Artin-Schelter regular algebra. Clearly, $J = k \langle y_1, y_2 \rangle / (y_2 y_1 - p_{12} y_1 y_2 - p_{11} y_1^2)$ is a Z-graded subalgebra of B and a noetherian Koszul Artin-Schelter regular algebra. We identify elements of J as the elements of B and write R_J for the vector space spanned by $y_2 y_1 - p_{12} y_1 y_2 - p_{11} y_1^2$ in the sequel. The following result is clear. Lemma 3.3.

(1)
$$B^{!} = T(V^{*} \oplus \Bbbk y_{1}^{*} \oplus \Bbbk y_{2}^{*})/(R^{\perp} \oplus R_{J}^{\perp} \oplus R_{\tau}), where$$

 $R_{J}^{\perp} = Span_{\Bbbk}\{(y_{2}^{*})^{2}, y_{1}^{*}y_{2}^{*} + p_{12}y_{2}^{*}y_{1}^{*}, (y_{1}^{*})^{2} + p_{11}y_{2}^{*}y_{1}^{*}\},$
 $R_{\tau} = Span_{\Bbbk}\{v^{*}y_{i}^{*} + y_{1}^{*}\sigma_{1i}^{*}(v^{*}) + y_{2}^{*}\sigma_{2i}^{*}(v^{*}) \mid v^{*} \in V^{*}, i = 1, 2\};$

- (2) $A^!$ and $J^!$ are \mathbb{Z} -graded subalgebras of $B^!$;
- (3) $B^!$ is a \mathbb{Z} -graded free $A^!$ -module and a \mathbb{Z} -graded free $J^!$ -module on both sides;
- (4) $B^!$ has a k-basis $\{e, y_1^*e, y_2^*e, y_1^*y_2^*e \mid e \in \mathcal{B}_A\}$ where \mathcal{B}_A is a k-basis of $A^!$.

Lemma 3.4. Retain the notations above. Suppose B is noetherian and z + h is a nonzero regular central element of B for some $z \in A_2$ and $h \in J_2$. Write $C_{A^1}(z) = T(V^*)/(\widetilde{R^{\perp}})$ and $C_{J^1}(h) = T(\mathbb{k}y_1^* \oplus \mathbb{k}y_2^*)/(\widetilde{R_J^{\perp}})$, where

$$\widetilde{R^{\perp}} = \{ f - f(\hat{z}) \mid f \in R^{\perp} \}, \qquad \widetilde{R^{\perp}_J} = \{ g - g(\hat{h}) \mid g \in R^{\perp}_J \},$$

and $\hat{z} \in V^{\otimes 2}$ and $\hat{h} \in (\Bbbk y_1 \oplus \Bbbk y_2)^{\otimes 2}$ are lifts of z and h respectively. Then

- (1) $C_{B^{!}}(z+h) \cong T(V^{*} \oplus \Bbbk y_{1}^{*} \oplus \Bbbk y_{2}^{*})/(\widetilde{R^{\perp}} \oplus \widetilde{R_{J}^{\perp}} \oplus R_{\tau})$ as \mathbb{Z}_{2} -graded algebras;
- (2) $C_{A^{!}}(z)$ and $C_{J^{!}}(h)$ are \mathbb{Z}_{2} -graded subalgebras of $C_{B^{!}}(z+h)$;
- (3) $C_{B^{!}}(z+h)$ is a \mathbb{Z}_{2} -graded free $C_{A^{!}}(z)$ -module and a \mathbb{Z}_{2} -graded free $C_{J^{!}}(h)$ -module on both sides;
- (4) $\operatorname{gldim}_{\mathbb{Z}_2} C_{B^!}(z+h) \ge \max\{\operatorname{gldim}_{\mathbb{Z}_2} C_{A^!}(z), \operatorname{gldim}_{\mathbb{Z}_2} C_{J^!}(h)\}.$

Proof. Clearly, $z \in A_2$ and $h \in J_2$ are both central elements of A and J respectively.

(1,2) follow by the definition of Clifford deformations and Lemma 3.3.

(3) Since $C_{A^{!}}(z)$, $C_{J^{!}}(h)$ and $C_{B^{!}}(z+h)$ are Clifford deformations, a class of PBW deformations, they have the same forms of k-bases of $A^{!}$, $J^{!}$ and $B^{!}$ respectively. By Lemma 3.3(4), the map of multiplication

$$C_{J^{!}}(h) \otimes C_{A^{!}}(z) \to C_{B^{!}}(z+h)$$

is an isomorphism as left \mathbb{Z}_2 -graded $C_{J^1}(h)$ -modules and right \mathbb{Z}_2 -graded $C_{A^1}(z)$ -modules.

Similarly, one can check that the natural map $C_{A^{!}}(z) \otimes C_{J^{!}}(h) \to C_{B^{!}}(z+h)$ is an isomorphism as right \mathbb{Z}_{2} -graded $C_{J^{!}}(h)$ -modules and left \mathbb{Z}_{2} -graded $C_{A^{!}}(z)$ -modules.

(4) It follows by a graded version of [8, Theorem 7.2.6].

Our main goal is to present a skew version of the classical Knörrer's periodicity theorem. To this end, we consider noncommutative quadric hypersurface having the form $B/(z + y_1^2 + y_2^2)$ in the rest of this paper where $z \in A_2$ is a nonzero regular central element. To make such class of noncommutative quadric hypersurfaces well defined, $y_1^2 + y_2^2$ should be a central element of B and also of J. By an easy calculation, we have the following result.

Lemma 3.5. Suppose the element $y_1^2 + y_2^2$ is central in *B*, then $P = \{1, 0\}$ or $P = \{-1, p_{11}\}$.

With respect to the values of p_{12} , we discuss in two cases in the following.

16

4. A Skew version of Knörrer's Periodicity Theorem with $p_{12} = 1$

Let $B = A_{\{1,0\}}[y_1, y_2; \sigma]$ be a Z-graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A = T(V)/(R) with deg $y_1 = \deg y_2 = 1$, and $z \in A_2$ be a nonzero regular central element of A. It is not hard to obtain the following conditions for σ .

Lemma 4.1. The element $z + y_1^2 + y_2^2$ is a central element of B if and only if

(1) $\sigma_{11}^2 + \sigma_{21}^2 = \sigma_{12}^2 + \sigma_{22}^2 = \mathrm{id}_A;$ (2) $\sigma_{11}\sigma_{12} + \sigma_{21}\sigma_{22} + \sigma_{12}\sigma_{11} + \sigma_{22}\sigma_{21} = 0;$ (3) $\sigma(z) = \begin{pmatrix} z & 0 \\ 0 & z \end{pmatrix}$.

By Lemma 4.1(3) and the construction (1.4), we have the following algebra homomorphism

$$\sigma^{!} = (\sigma^{!}_{ij}) : C_{A^{!}}(z) \to M_2(C_{A^{!}}(z)),$$

where any $\sigma_{ij}^{!}$ is a \mathbb{Z}_2 -graded map for i, j = 1, 2. Then

Corollary 4.2. (1) $(\sigma_{11}^!)^2 + (\sigma_{21}^!)^2 = (\sigma_{12}^!)^2 + (\sigma_{22}^!)^2 = \operatorname{id}_{C_{\mathfrak{a}!}(z)};$ (2) $\sigma_{12}^! \sigma_{11}^! + \sigma_{22}^! \sigma_{21}^! + \sigma_{11}^! \sigma_{12}^! + \sigma_{21}^! \sigma_{22}^! = 0;$ (3) $\sigma_{11}^! \sigma_{21}^! = \sigma_{21}^! \sigma_{11}^!;$ (4) $\sigma_{12}^! \sigma_{22}^! = \sigma_{22}^! \sigma_{12}^!;$ (5) $\sigma_{22}^! \sigma_{11}^! - \sigma_{12}^! \sigma_{21}^! = \sigma_{11}^! \sigma_{22}^! - \sigma_{21}^! \sigma_{12}^!;$ (6) $\sigma^!$ is t-invertible.

Proof. (1)–(5) follow by Lemma 4.1 and (3.1). (6) holds by Lemma 3.2 and Lemma 1.5,

For now on, we construct a twisting system on $M_2(C_{A^{\dagger}}(z))$. We choose an invertible \mathbb{Z}_2 -graded basis $\mathbb{I} = \{\mathbb{I}_1^{(0)}, \mathbb{I}_2^{(0)}, \mathbb{I}_1^{(1)}, \mathbb{I}_2^{(1)}\}$ of $M_2(\mathbb{k})$ as follows,

$$(4.1) \quad \mathbb{I}_{1}^{(0)} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbb{I}_{2}^{(0)} = \begin{pmatrix} -\sqrt{-1} & 0 \\ 0 & \sqrt{-1} \end{pmatrix}, \quad \mathbb{I}_{1}^{(1)} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbb{I}_{2}^{(1)} = \begin{pmatrix} 0 & \sqrt{-1} \\ -\sqrt{-1} & 0 \end{pmatrix}.$$

Define linear maps $\theta^{(0)}$ and $\theta^{(1)}$ from $C_{A^{\dagger}}(z)$ to $M_2(C_{A^{\dagger}}(z))$ by

(4.2)
$$\theta^{(0)} = (\theta_{jj'}^{(0)}) = \begin{pmatrix} \mathrm{id} & \sigma_{12}^! \sigma_{11}^! + \sigma_{22}^! \sigma_{21}^! \\ 0 & \sigma_{22}^! \sigma_{11}^! - \sigma_{12}^! \sigma_{21}^! \end{pmatrix}, \quad \theta^{(1)} = (\theta_{jj'}^{(1)}) = \sigma^! \circ \xi_{-1} = (\sigma_{ij}^! \circ \xi_{-1}).$$

Clearly, $\theta_{jj'}^{(i)}$ is a \mathbb{Z}_2 -graded linear map for any $i \in \mathbb{Z}_2, j, j' = 1, 2$. Those two linear maps satisfy the following properties.

Lemma 4.3. Retain the notations above. Then

- (1) $\theta_{12}^{(0)}(aa') = a\theta_{12}^{(0)}(a') + \theta_{12}^{(0)}(a)\theta_{22}^{(0)}(a')$ for any $a, a' \in C_{A^{!}}(z)$; (2) $\theta^{(0)}$ and $\theta^{(1)}$ are algebra homomorphisms;
- $\begin{array}{l} (3) \ \ \theta_{12}^{(0)}\theta_{22}^{(0)} + \theta_{22}^{(0)}\theta_{12}^{(0)} = 0; \\ (4) \ \ (\theta_{22}^{(0)})^2 (\theta_{12}^{(0)})^2 = \mathrm{id}_{C_{4^{\dagger}}(z)}; \end{array}$

Proof. (1) holds since $\sigma^!$ is an algebra homomorphism. (2) follows by (1). (3,4) follow by Corollary 4.2. **Theorem 4.4.** Let $\theta^{(0)}$ and $\theta^{(1)}$ be linear maps from $C_{A^{!}}(z)$ to $M_2(C_{A^{!}}(z))$ defined in (4.2) and \mathbb{I} be the basis of $M_2(\mathbb{k})$ defined in (4.1). Then

- (1) $\Theta = (\{\theta^{(i)}\}_{i \in \mathbb{Z}_2}, \mathbb{I})$ is a twisting system of $M_2(C_{A^!}(z));$
- (2) $C_{B^{!}}(z+y_{1}^{2}+y_{2}^{2}) \cong {}^{\Theta}M_{2}(C_{A^{!}}(z))$ as \mathbb{Z}_{2} -graded algebras where the \mathbb{Z}_{2} -grading of ${}^{\Theta}M_{2}(C_{A^{!}}(z))$ comes from the total degrees (see Remark 2.11(1)).

Proof. (1) Write φ for the inverse of σ . By Lemma 1.5 and its proof, $\varphi^! = (\varphi_{ij}^!)$ exits and is the *t*-inverse of $\sigma^!$. Then one obtains that $\theta^{(0)}$ and $\theta^{(1)}$ are *t*-invertible with the *t*-inverses

$$\begin{pmatrix} \mathrm{id} & 0 \\ \varphi_{21}^{!}\varphi_{11}^{!} + \varphi_{22}^{!}\varphi_{12}^{!} & \varphi_{22}^{!}\varphi_{11}^{!} - \varphi_{21}^{!}\varphi_{12}^{!} \end{pmatrix}, \qquad \varphi^{!} \circ \xi_{-1}$$

respectively. Clearly, $\theta^{(1)}(1) = I_2 \in GL_2(\mathbb{k})$, and the invertible matrices defined in (2.1) are $L_1^{(ii')} = I_2$ for any $i, i' \in \mathbb{Z}_2$, and

$$L_2^{(00)} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \ L_2^{(01)} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \ L_2^{(10)} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \ L_2^{(11)} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Then by Corollary 4.2 and Lemma 4.3, it is straightforward but tedious to check Θ satisfies the following simplified equations of Proposition 2.2(3)

$$\begin{aligned} \theta_{pj'}^{(i')} \left(\theta_{11}^{(i)}(x)x' \right) &+ l_{p2u}^{(ii')} \theta_{uj'}^{(i')} \left(\theta_{21}^{(i)}(x)x' \right) = & \theta_{p1}^{(i+i')}(x)\theta_{1j'}^{(i')}(x') + \theta_{p2}^{(i+i')}(x)\theta_{2j'}^{(i')}(x'), \\ \theta_{pj'}^{(i')} \left(\theta_{12}^{(i)}(x)x' \right) &+ l_{p2u}^{(ii')} \theta_{uj'}^{(i')} \left(\theta_{22}^{(i)}(x)x' \right) = & l_{122}^{(ii')} \theta_{p1}^{(i+i')}(x)\theta_{2j'}^{(i')}(x') + & l_{221}^{(ii')} \theta_{p2}^{(i+i')}(x)\theta_{1j'}^{(i')}(x'), \end{aligned}$$

where $u = \{1, 2\} - \{p\}$ for any $x, x' \in E, i, i' \in \mathbb{Z}_2$ and p = 1, 2.

(b) By Lemma 3.4(1), $C_{B^{!}}(z + y_1^2 + y_2^2)$ is generated by V^*, y_1^*, y_2^* subject to the relations consisting of the relations of $C_{A^{!}}(z)$, and

$$\begin{split} &(y_1^*)^2-1, (y_2^*)^2-1, y_1^*y_2^*+y_2^*y_1^*,\\ &v^*y_i^*+y_1^*\sigma_{1i}^*(v^*)+y_2^*\sigma_{2i}^*(v^*), \qquad \forall v^*\in V^*, i=1,2. \end{split}$$

We list the precise multiplication of ${}^{\Theta}M_2(C_{A^{!}}(z))$ below: for any $x, x' \in C_{A^{!}}(z)$,

TABLE 1. Multiplication Table of $\Theta M_2(C_{A'}(z))$.

Then one obtains that there exists a \mathbb{Z}_2 -graded algebra isomorphism from $C_{B'}(z+y_1^2+y_2^2)$ to $\Theta M_2(C_{A^!}(z))$ mapping

$$y_1^* \mapsto \mathbb{I}_1^{(1)}, \qquad y_2^* \mapsto \mathbb{I}_2^{(1)}, \qquad v^* \mapsto \mathbb{I}_1^{(0)}v^*, \ \forall v^* \in V^*.$$

Proposition 4.5. Retain the assumptions of Theorem 4.4. The homogeneous element $e = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ is a full idempotent of ${}^{\Theta}M_2(C_{A^!}(z))$.

As a consequence, $\operatorname{gr}_{\mathbb{Z}_2} {}^{\Theta} M_2(C_{A^{!}}(z))$ and $\operatorname{gr}_{\mathbb{Z}_2} e\left({}^{\Theta} M_2(C_{A^{!}}(z))\right) e$ is equivalent.

Proof. Note that $e = 2^{-1} \left(\mathbb{I}_1^{(0)} + \sqrt{-1} \mathbb{I}_2^{(0)} \right)$. By Table 1, one obtains

$$2^{-1} \left(\mathbb{I}_1^{(i)} + \sqrt{-1} \mathbb{I}_2^{(i)} \right) \star e = 2^{-1} \left(\mathbb{I}_1^{(i)} + \sqrt{-1} \mathbb{I}_2^{(i)} \right),$$

and

$$\begin{split} \mathbb{I}_{1}^{(i)}x &= 2^{-1} \left(\mathbb{I}_{1}^{(i)} + \sqrt{-1}\mathbb{I}_{2}^{(i)} \right) \star e \star \mathbb{I}_{1}^{(0)}x + 2^{-1} \left(\mathbb{I}_{1}^{(i+1)} + \sqrt{-1}\mathbb{I}_{2}^{(i+1)} \right) \star e \star \mathbb{I}_{1}^{(1)}x, \\ \mathbb{I}_{2}^{(i)}x &= 2^{-1} \left(\mathbb{I}_{1}^{(i+1)} + \sqrt{-1}\mathbb{I}_{2}^{(i+1)} \right) \star e \star \sqrt{-1}\mathbb{I}_{1}^{(1)}x - 2^{-1} \left(\mathbb{I}_{1}^{(i)} + \sqrt{-1}\mathbb{I}_{2}^{(i)} \right) \star e \star \sqrt{-1}\mathbb{I}_{1}^{(0)}x, \\ \text{ny } i \in \mathbb{Z}_{2} \text{ and } x \in C_{A^{1}}(z). \text{ Hence, } e \text{ is a full idempotent.} \end{split}$$

for any $i \in \mathbb{Z}_2$ and $x \in C_{A^{!}}(z)$. Hence, e is a full idempotent.

To understand the multiplication of ${}^{\Theta}M_2(C_{A'}(z))$ further more, we introduce four linear transformations of $C_{A^{!}}(z)$:

We show some parts of the multiplication of ${}^{\Theta}M_2(C_{A^!}(z))$ in the usual matrix forms involving the linear transforms listed above:

(4.3)
$$\begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} * \begin{pmatrix} a' & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} \Xi_1(a)a' & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} * \begin{pmatrix} 0 & a' \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \Phi_1(a)a' \\ 0 & 0 \end{pmatrix},$$
$$\begin{pmatrix} (4.3) \\ 0 & 0 \end{pmatrix} * \begin{pmatrix} a' & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \Xi_2(a)a' \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & a \\ 0 & 0 \end{pmatrix} * \begin{pmatrix} 0 & a' \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} \Phi_2(a)a' & 0 \\ 0 & 0 \end{pmatrix}.$$

Here are some useful properties of those four liner transformations.

Lemma 4.6. (1) $\Xi_1^2 = \Xi_1, \Xi_2^2 = \Xi_2.$ (2) $\Xi_1(ab) = a\Xi_2(b) + \Xi_1(a)\theta_{22}^{(0)}(b) = a\Xi_1(b) + \Xi_1(a)\theta_{22}^{(0)}(b) - a\theta_{22}^{(0)}(b)$ for any $a, b \in C_{A^1}(z)$.

(3)
$$\Xi_2(ab) = a\Xi_2(b) + \Xi_2(a)\theta_{22}^{(0)}(b) = a\Xi_1(b) + \Xi_2(a)\theta_{22}^{(0)}(b) - a\theta_{22}^{(0)}(b)$$
 for any $a, b \in C_{A^1}(z)$.

- (b) $\Xi_2(0) = \Xi_2(0) + \Xi_2(0) + \Xi_2(0) = \Xi_1(0) + \Xi_2(0) = \Xi_2(0)$
- (6) $\Phi_1(\Xi_1(a)b) = \Phi_1(a)\Phi_1(b), \ \Phi_1(a\Xi_1(b)) = \Phi_1(a)\Phi_1((\Xi_1(b)), \ \Phi_1(\Phi_2(a)\Xi_2(b)) = \Xi_2(a)\Phi_2(b)$ for any $a, b \in C_{A^{!}}(z)$.
- (7) $\Phi_2(\Xi_2(a)b) = \Phi_2(a)\Phi_1(b), \Phi_2(\Phi_1(a)\Xi_2(b)) = \Xi_1(a)\Phi_2(b)$ for any $a, b \in C_{A^!}(z)$. (8) $(\theta_{11}^{(1)} \sqrt{-1}\theta_{21}^{(1)})\Phi_1 = (\theta_{22}^{(1)} + \sqrt{-1}\theta_{12}^{(1)})\Phi_1 = \Xi_1$. (9) $(\theta_{11}^{(1)} + \sqrt{-1}\theta_{21}^{(1)})\Phi_2 = (\sqrt{-1}\theta_{12}^{(1)} \theta_{22}^{(1)})\Phi_2 = \Xi_2$.

Proof. All follow by Corollary 4.2 and Lemma 4.3.

By Lemma 4.6(1), we write S for the \mathbb{Z}_2 -graded eigenspace for the eigenvalue 1 of Ξ_1 and M for the \mathbb{Z}_2 -graded eigenspace for the eigenvalue 1 of Ξ_2 .

- **Lemma 4.7.** (1) The \mathbb{Z}_2 -graded eigenspace S is a \mathbb{Z}_2 -graded subalgebra of $C_{A^1}(z)$ and $S \cong \Phi_1(S)$ as \mathbb{Z}_2 -graded algebras.
 - (2) The \mathbb{Z}_2 -graded eigenspace M is a \mathbb{Z}_2 -graded S-bimodule with the S-action defined by

$$s \cdot m \cdot s' = \Phi_1(s)ms',$$

for any $s, s' \in S$ and $m \in M$.

- (3) For any $m, m' \in M$, $\Phi_2(m)m' \in S$.
- (4) The map

$$\psi: \quad M(1) \otimes_S M(1) \quad \to \quad S$$
$$m \otimes m' \quad \mapsto \quad \Phi_2(m)m'$$

is a Z₂-graded S-bimodule homomorphism satisfying $m\psi(m' \otimes m'') = \psi(m \otimes m') \cdot m''$ for any homogeneous elements $m, m', m'' \in M$.

Proof. (1) Clearly $\Xi_1(1) = 1 \in S$. By Lemma 4.6(2), for any $s, s' \in S$,

$$\Xi_1(ss') = s\Xi_1(s') + \Xi_1(s)\theta_{22}^{(0)}(s') - s\theta_{22}^{(0)}(s') = ss'.$$

Hence, S is a \mathbb{Z}_2 -graded subalgebra of $C_{A^!}(z)$.

By Lemma 4.6(6), the restriction map $\Phi_{1|S} : S \to \Phi_1(S)$ is a surjective \mathbb{Z}_2 -graded algebra homomorphism. If $s \in \operatorname{Ker} \Phi_{1|S}$, then $s = \Xi_1(s) = (\theta_{11}^{(1)} - \sqrt{-1}\theta_{21}^{(1)})\Phi_1(s) = 0$ by Lemma 4.6(8). So $\Phi_{1|S}$ is an isomorphism.

(2) By Lemma 4.6(3,4),

$$\begin{aligned} \Xi_2(\Phi_1(s)ms') = &\Phi_1(s)\Xi_2(ms') + \Xi_2(\Phi_1(s))\theta_{22}^{(0)}(ms') \\ = &\Phi_1(s)\left(m\Xi_2(s') + \Xi_2(m)\theta_{22}^{(0)}(s') - m\theta_{22}^{(0)}(s')\right) \\ = &\Phi_1(s)ms', \end{aligned}$$

for any $s, s' \in S$ and $m \in M$. It implies that M is a $(\Phi_1(S), S)$ -bimodule. By (1), the result follows.

(3) By Lemma 4.6(2,5), $\Xi_1(\Phi_2(m)m') = \Phi_2(m)\Xi_2(m') + \Xi_1(\Phi_2(m))\theta_{22}^{(0)}(m') = \Phi_2(m)m'$, for any $m, m' \in M$.

(4) For any $s \in S$ and $m, m' \in M(1)$,

$$\psi(ms\otimes m') = \Phi_2(\Xi_2(m)s)m' = \Phi_2(m)\Phi_1(s)m' = \Phi_2(m)(s\cdot m') = \psi(m\otimes s\cdot m'),$$

by Lemma 4.6(7). Combining with (3), ψ is well defined.

For any $s, s' \in S$ and $m, m' \in M(1)$,

$$\psi(s \cdot m \otimes m' \cdot s') = \Phi_2(\Phi_1(s)\Xi_2(m))m's' = \Xi_1(s)\Phi_2(m)m's' = s\psi(m \otimes m')s',$$

by Lemma 4.6(7).

The last lemma allows us to construct a semi-trivial extension

$$\Lambda = S \ltimes_{\psi} M(1)$$

which is a $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded algebra and also a \mathbb{Z}_2 -graded algebra by forgetting the first \mathbb{Z}_2 -grading (see Remark 1.7 (3)). In general, Λ is not strongly \mathbb{Z}_2 -graded (see Example 4.9 below).

Theorem 4.8. Let $B = A_{\{1,0\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A with deg $y_1 = \deg y_2 = 1$ and $z \in A_2$ be a nonzero regular central element of A. Suppose B is noetherian and $z + y_1^2 + y_2^2$ is a central element of B. Then there are equivalences of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2} \,^{\Theta}M_2(C_{A^!}(z))) \cong D^b(\mathrm{mod}\,^{\Theta}M_2(C_{A^!}(z))_0) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2}\,\Lambda).$$

Proof. By Theorem 4.4 and Theorem 1.4, one obtains that there are equivalences of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2} \,^{\Theta} M_2(C_{A^!}(z))) \cong D^b(\mathrm{mod}\,^{\Theta} M_2(C_{A^!}(z))_0).$$

By Proposition 4.5, $\operatorname{gr}_{\mathbb{Z}_2}^{\Theta} M_2(C_{A^!}(z))$ is equivalent to $\operatorname{gr}_{\mathbb{Z}_2} e\left({}^{\Theta}M_2(C_{A^!}(z))\right)e$. By Table 1, one obtains

$$\begin{aligned} &e \star \mathbb{I}_{1}^{(0)} x \star e = 2^{-1} \left(\mathbb{I}_{1}^{(0)} \Xi_{1}(x) + \sqrt{-1} \mathbb{I}_{2}^{(0)} \Xi_{1}(x) \right), \\ &e \star \mathbb{I}_{2}^{(0)} x \star e = -2^{-1} \sqrt{-1} \left(\mathbb{I}_{1}^{(0)} \Xi_{1}(x) + \sqrt{-1} \mathbb{I}_{2}^{(0)} \Xi_{1}(x) \right), \\ &e \star \mathbb{I}_{1}^{(1)} x \star e = 2^{-1} \left(\mathbb{I}_{1}^{(1)} \Xi_{2}(x) - \sqrt{-1} \mathbb{I}_{2}^{(0)} \Xi_{2}(x) \right), \\ &e \star \mathbb{I}_{2}^{(1)} x \star e = 2^{-1} \sqrt{-1} \left(\mathbb{I}_{1}^{(1)} \Xi_{2}(x) - \sqrt{-1} \mathbb{I}_{2}^{(0)} \Xi_{2}(x) \right), \end{aligned}$$

for any $x \in C_{A^{!}}(z)$. It implies

Note th

$$e\left({}^{\Theta}M_{2}(C_{A^{!}}(z))\right)e = \left\{ \begin{pmatrix} \Xi_{1}(x) & \Xi_{2}(x') \\ 0 & 0 \end{pmatrix} \mid x, x' \in C_{A^{!}}(z) \right\} = \begin{pmatrix} S & M \\ 0 & 0 \end{pmatrix}.$$

that $e\left({}^{\Theta}M_{2}(C_{A^{!}}(z))\right)_{0}e = \begin{pmatrix} S_{0} & M_{1} \\ 0 & 0 \end{pmatrix}$ and $e\left({}^{\Theta}M_{2}(C_{A^{!}}(z))\right)_{1}e = \begin{pmatrix} S_{1} & M_{0} \\ 0 & 0 \end{pmatrix}$. By (4.3), we have

have $e\left({}^{\Theta}M_2(C_{A^!}(z))\right)e \cong \Lambda$ as \mathbb{Z}_2 -graded algebras.

Hence, there is an equivalence of triangulated categories

$$D^{b}(\operatorname{gr}_{\mathbb{Z}_{2}}^{\Theta}M_{2}(C_{A^{!}}(z))) \cong D^{b}(\operatorname{gr}_{\mathbb{Z}_{2}}^{\Lambda}\Lambda).$$

Example 4.9. Let A be a noetherian Koszul Artin-Schelter regular algebra. Let σ_1 and σ_2 be graded automorphisms of A satisfying

$$\sigma_1^2 = \sigma_2^2 = \mathrm{id}_A, \qquad \sigma_1 \sigma_2 = \sigma_2 \sigma_1.$$

Naturally, σ_2 can be extended to be a graded automorphism of $A[y_1; \sigma_1]$ which preserves y_1 , still denoted by σ_2 . Then we have an Ore extension $A[y_1; \sigma_1][y_2; \sigma_2]$ is isomorphic to the double Ore extension $B = A_{\{1,0\}}[y_1, y_2; \sigma]$, where

$$\sigma = \begin{pmatrix} \sigma_1 & 0\\ 0 & \sigma_2 \end{pmatrix} : A \to M_2(A).$$

Let $z \in A_2$ be a nonzero regular central element of A such that $\sigma_1(z) = \sigma_2(z) = z$. By Lemma 4.1, $z + y_1^2 + y_2^2$ is a central element of B. In this case, the linear maps in the twisting system are

$$\theta^{(0)} = \begin{pmatrix} \mathrm{id} & 0\\ 0 & \sigma_{22}^! \sigma_{11}^! \end{pmatrix}, \quad \theta^{(1)} = \begin{pmatrix} \sigma_1^! & 0\\ 0 & \sigma_2^! \end{pmatrix} \circ \xi_{-1}$$

and the liner transformations of $C_{A^{!}}(z)$ are

$$\Xi_1 = 2^{-1} (\mathrm{id} + \sigma_2^! \sigma_1^!), \quad \Xi_2 = 2^{-1} (\mathrm{id} - \sigma_2^! \sigma_1^!), \quad \Phi_1 = 2^{-1} (\sigma_1^! + \sigma_2^!) \xi_{-1}, \quad \Phi_2 = 2^{-1} (\sigma_1^! - \sigma_2^!) \xi_{-1}.$$

 So

$$S = \left\{ x \in C_{A^{!}}(z) \mid \sigma_{1}^{!}(x) = \sigma_{2}^{!}(x) \right\}, \qquad M = \left\{ x \in C_{A^{!}}(z) \mid \sigma_{1}^{!}(x) = -\sigma_{2}^{!}(x) \right\},$$

and $\Lambda = S \ltimes_{\psi} M(1)$ with multiplication

$$(s,m) \odot (s',m') = (ss' + \sigma_1^! \xi_{-1}(m)m', \sigma_1^! \xi_{-1}(s)m' + ms'),$$

for any homogeneous elements $s, s' \in S$ and $m, m' \in M$. By Theorem 4.8, there is an equivalence of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2}\Lambda)$$

There are some special cases.

(1) Suppose $\sigma_1 = \sigma_2 = \sigma$. Then $S = C_{A'}(z), M = 0$, and $\Lambda = C_{A'}(z)$. Hence, there are equivalences of triangulated categories

 $\underline{\operatorname{mcm}}(A[y_1;\sigma][y_2;\sigma]/(z+y_1^2+y_2^2)) \cong D^b(\operatorname{gr}_{\mathbb{Z}_2} C_{A^!}(z)) \cong \underline{\operatorname{mcm}}(A/(z)).$

It is exactly the noncommutative Knörrer's periodicity theorem for noncommutative quadric hypersurfaces (see [10, Theorem 1.3], or [2, Theorem 0.4]).

(2) Suppose $\sigma_1 = \xi_{-1}$ and σ_2 = id. Then $S_0 = C_{A^{!}}(z)_0, S_1 = 0, M_0 = 0, M_1 = C_{A^{!}}(z)_1$. So \mathbb{Z}_2 -graded algebra Λ is concentrated in degree 0 with $\Lambda_0 = C_{A^{!}}(z)$. Hence, there is an equivalence of triangulated categories

$$\underline{\mathrm{mcm}}(A[y_1;\xi_{-1}][y_2]/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{mod}\,C_{A^!}(z))^{\times 2}.$$

It is equivalent to [10, Theorem 1.4] (up to Zhang-twist) which provides a tool called two point reduction for computing a classes of noncommutative quadric hypersurfaces defined by (± 1) -skew polynomial algebras (see [10, Lemma 6.18]). We will also give another approach to [10, Theorem 1.4] in Example 5.10.

Example 4.10. Let $A = k_{-1}[x_1, x_2]$ and the nonzero regular central element $z = x_1^2 + x_2^2$. Let $B = A_{\{1,0\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of A where $\sigma = (\sigma_{ij}) : A \to M_2(A)$ satisfies

$$\sigma_{11}(x_1) = hx_1, \quad \sigma_{12}(x_1) = hx_2, \quad \sigma_{21}(x_1) = hx_2, \quad \sigma_{22}(x_1) = -hx_1, \\ \sigma_{11}(x_2) = hx_2, \quad \sigma_{12}(x_2) = hx_1, \quad \sigma_{21}(x_2) = hx_1, \quad \sigma_{22}(x_1) = -hx_2,$$

where $2h^2 = 1$. It is a special case of the class \mathbb{Z} with f = 1 in [22]. By [22, Proposition 5.16], B is noetherian. It is not hard to obtain that σ satisfies Lemma 4.1, so $x_1^2 + x_2^2 + y_1^2 + y_2^2$ is a central element of B. In this case, the linear maps in the twisting system of $M_2(C_{A^1}(z))$ are

$$\theta^{(0)} = \begin{pmatrix} \mathrm{id} & 0\\ 0 & \xi_{-1} \end{pmatrix}, \quad \theta^{(1)} = \begin{pmatrix} \sigma_{11}^{!} & \sigma_{12}^{!}\\ \sigma_{21}^{!} & \sigma_{22}^{!} \end{pmatrix} \circ \xi_{-1},$$

and two liner transformations of $C_{A^{!}}(z)$ are

$$\Xi_1 = 2^{-1} (\mathrm{id} + \xi_{-1}), \qquad \Xi_2 = 2^{-1} (\mathrm{id} - \xi_{-1}).$$

So the \mathbb{Z}_2 -graded algebra S equals $C_{A^{!}}(z)_0$ concentrated in degree 0 and the \mathbb{Z}_2 -graded S-module M equals $C_{A^{!}}(z)_1$ concentrated in degree 1. The other linear transformations are $\Phi_{1|S} = \mathrm{id}_S$ and Φ_2 mapping

$$x_1^* \mapsto -2h(x_1^* - \sqrt{-1}x_2^*), \qquad x_2^* \mapsto 2\sqrt{-1}h(x_1^* + \sqrt{-1}x_2^*).$$

Then the semi-trivial extension $\Lambda = S \ltimes_{\psi} M(1)$ is a $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded algebra where $\Lambda_{(0,0)} = C_{A^{!}}(z)_0$, $\Lambda_{(1,0)} = C_{A^{!}}(z)_1$ and $\Lambda_{(0,1)} = \Lambda_{(1,1)} = 0$ and with multiplication

$$(s,m) \odot (s',m') = (ss' + \Phi_2(m)m', sm' + ms'),$$

for any $s, s' \in C_{A^{!}}(z)_{0}, m, m' \in C_{A^{!}}(z)_{1}$. Then Λ is a \mathbb{Z}_{2} -graded algebra concentrated in degree 0 by forgetting the first grading and Λ_{0} is also a \mathbb{Z}_{2} -graded algebra. In fact, one can find a \mathbb{Z}_{2} -graded algebra isomorphism between Λ_{0} and $C_{A^{!}}(z)$. But we give another way to discuss here.

We view Λ_0 as a \mathbb{Z}_2 -graded algebra. It is not hard to know that $\Phi_{2|M_1}$ is a linear automorphism of M_1 by Lemma 4.6(5). We define a left Zhang-twisting system $\nu = \{\nu_0, \nu_1\}$ of \mathbb{Z}_2 -graded algebra $C_{A^{!}}(z)$ by

$$\nu_0 = \mathrm{id}, \qquad \nu_{1|C_{A!}(z)_0} = \mathrm{id}, \qquad \nu_{1|C_{A!}(z)_1} = \Phi_{2|M_1}.$$

So $\Lambda_0 \cong {}^{\nu}C_{A^{\dagger}}(z)$ as \mathbb{Z}_2 -graded algebras. By Theorem 4.8, there are equivalences of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{gr}_{\mathbb{Z}_2}\Lambda) \cong D^b(\mathrm{mod}\,\Lambda_0)^{\times 2} \cong D^b(\mathrm{mod}\,^{\nu}C_{A^!}(z))^{\times 2}.$$

Moreover, $C_{A'}(z) \cong \mathbb{k}\mathbb{Z}_2 \times \mathbb{k}\mathbb{Z}_2$ as \mathbb{Z}_2 -graded algebras, so the algebra ${}^{\nu}C_{A'}(z)$ is \mathbb{Z}_2 -graded semisimple by [20, Corollary 4.4] and ${}^{\nu}C_{A'}(z) \cong \mathbb{k}\mathbb{Z}_2 \times \mathbb{k}\mathbb{Z}_2 \cong C_{A'}(z)$ since ${}^{\nu}C_{A'}(z)$ is commutative by an easy computation. It implies that $B/(z+y_1^2+y_2^2)$ is a noncommutative graded isolated singularity and there is an equivalence of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathrm{mod}\,\mathbb{k})^{\times 8}.$$

5. A Skew version of Knörrer's Periodicity Theorem with $p_{12} = -1$

Let $B = A_{\{-1,p_{11}\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A = T(V)/(R) with deg $y_1 = \deg y_2 = 1$. In this case, p_{11} can be nonzero. We will always assume $p_{11} = 0$ except the following special values of p_{11} .

Proposition 5.1. Let $B = A_{\{-1,p_{11}\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A and $z \in A_2$ be a nonzero regular central element of A. Suppose B is noetherian and $z + y_1^2 + y_2^2$ is a regular central element of B. If $p_{11} = \pm 2\sqrt{-1}$, then $B/(z + y_1^2 + y_2^2)$ is not a noncommutative graded isolated singularity.

Proof. Write

$$\sigma' = \begin{pmatrix} \sigma_{11} + \frac{p_{11}}{2} \sigma_{12} & \sigma_{12} \\ \sigma_{21} + \frac{p_{11}}{2} \sigma_{22} - \frac{p_{11}}{2} \sigma_{11} - \frac{p_{11}^2}{4} \sigma_{12} & \sigma_{22} - \frac{p_{11}}{2} \sigma_{12} \end{pmatrix} : A \to M_2(A),$$

and

$$\varphi' = \begin{pmatrix} \varphi_{11} + \frac{p_{11}}{2}\varphi_{21} & \varphi_{12} + \frac{p_{11}}{2}\varphi_{22} - \frac{p_{11}}{2}\varphi_{11} - \frac{p_{11}^2}{4}\varphi_{21} \\ \varphi_{21} & \varphi_{22} - \frac{p_{11}}{2}\varphi_{21} \end{pmatrix} : A \to M_2(A),$$

where $\varphi = (\varphi_{ij})$ is the inverse of σ .

It is routine to check that σ' is an invertible algebra homomorphism with the inverse φ' and $A_{\{-1,0\}}[y_1, y_2; \sigma']$ is a \mathbb{Z} -graded double Ore extension by Lemma 3.2. Then the algebra homomorphism from B to $A_{\{-1,0\}}[y_1, y_2; \sigma']$ mapping

$$y_1 \mapsto y_1, \quad y_2 \mapsto y_2 + \frac{p_{11}}{2}y_1, \quad a \mapsto a, \ \forall a \in A,$$

is an isomorphism, and the image of $z + y_1^2 + y_2^2$ is $z + y_2^2$ since $p_{11} = \pm 2\sqrt{-1}$. So

$$B/(z+y_1^2+y_2^2) \cong A_{\{-1,0\}}[y_1,y_2;\sigma']/(z+y_2^2).$$

It is well known $k_{-1}[y_1, y_2]/(y_2^2)$ is not a noncommutative graded isolated singularity, and so

$$\operatorname{gldim}_{\mathbb{Z}_2} C_{k-1[y_1,y_2]!}(y_2^2) > 0$$

by Theorem 1.4(2). Write $D = A_{\{-1,0\}}[y_1, y_2; \sigma']$. By Lemma 3.4(4),

$$\operatorname{gldim}_{\mathbb{Z}_2} C_{D^!}(z+y_2^2) \ge \operatorname{gldim}_{\mathbb{Z}_2} C_{k_{-1}[y_1,y_2]^!}(y_2^2) > 0.$$

The result holds by Theorem 1.4(2).

g

Lemma 5.2. Let $B = A_{\{-1,p_{11}\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of A where $p_{11} \neq \pm 2\sqrt{-1}$. Suppose the element $y_1^2 + y_2^2$ is a central element in B, then there exists a \mathbb{Z} -graded double Ore extension $A_{\{-1,0\}}[y_1, y_2; \sigma']$ such that

$$B/(a+y_1^2+y_2^2) \cong A_{\{-1,0\}}[y_1,y_2;\sigma']/(a+y_1^2+y_2^2),$$

for any $a \in A$.

Proof. Write

Since p_1

$$\sigma' = \begin{pmatrix} \sigma_{11} + \frac{p_{11}}{2} \sigma_{12} & c\sigma_{12} \\ c^{-1} \left(\sigma_{21} + \frac{p_{11}}{2} \sigma_{22} - \frac{p_{11}}{2} \sigma_{11} - \frac{p_{11}^2}{4} \sigma_{12} \right) & \sigma_{22} - \frac{p_{11}}{2} \sigma_{12} \end{pmatrix} : A \to M_2(A),$$

for some $c \in \mathbb{k}$ satisfying $c^2 = 1 + \frac{p_{11}^2}{4}$. It is similar to the proof of Proposition 5.1 to obtain that $A_{\{-1,0\}}[y_1, y_2; \sigma']$ is a \mathbb{Z} -graded double Ore extension. There is a \mathbb{Z} -graded algebra isomorphism ρ from B to $A_{\{-1,0\}}[y_1, y_2; \sigma']$ mapping

$$y_1 \mapsto c^{-1}y_1, \quad y_2 \mapsto y_2 + \frac{p_{11}}{2}c^{-1}y_1, \quad x \mapsto x, \ \forall x \in A.$$

$$1_1 \neq \pm 2\sqrt{-1}, \text{ then } \rho(a+y_1^2+y_2^2) = a+y_1^2+y_2^2.$$

In the rest of this section, $B = A_{\{-1,0\}}[y_1, y_2; \sigma]$ be a Z-graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A = T(V)/(R) with deg $y_1 = \text{deg } y_2 = 1$, and $z \in A_2$ be a nonzero regular central element of A. There are results similar to Lemma 4.1 and Corollary 4.2.

24

Lemma 5.3. The element $z + y_1^2 + y_2^2$ is a regular central element of B if and only if

(1) $\sigma_{11}^2 + \sigma_{21}^2 = \sigma_{12}^2 + \sigma_{22}^2 = \mathrm{id}_A;$ (2) $\sigma_{11}\sigma_{12} + \sigma_{21}\sigma_{22} = \sigma_{12}\sigma_{11} + \sigma_{22}\sigma_{21};$ (3) $\sigma(z) = \begin{pmatrix} z & 0 \\ 0 & z \end{pmatrix}.$

By Lemma 5.3(3), the construction (1.4), 3.2 and Lemma 1.5, we have the following *t*-invertible algebra homomorphism

$$\sigma^! = (\sigma_{ij}^!) : C_{A^!}(z) \to M_2(C_{A^!}(z)).$$

Corollary 5.4. If $z + y_1^2 + y_2^2 \in B$ is a regular central element, then

 $\begin{array}{ll} (1) & (\sigma_{11}^{!})^{2} + (\sigma_{21}^{!})^{2} = (\sigma_{12}^{!})^{2} + (\sigma_{22}^{!})^{2} = \mathrm{id}_{C_{A^{!}}(z)};\\ (2) & \sigma_{12}^{!}\sigma_{11}^{!} + \sigma_{22}^{!}\sigma_{21}^{!} = \sigma_{11}^{!}\sigma_{12}^{!} + \sigma_{21}^{!}\sigma_{22}^{!};\\ (3) & \sigma_{11}^{!}\sigma_{21}^{!} + \sigma_{21}^{!}\sigma_{11}^{!} = 0;\\ (4) & \sigma_{12}^{!}\sigma_{22}^{!} + \sigma_{22}^{!}\sigma_{12}^{!} = 0;\\ (5) & \sigma_{22}^{!}\sigma_{11}^{!} + \sigma_{12}^{!}\sigma_{21}^{!} = \sigma_{11}^{!}\sigma_{22}^{!} + \sigma_{21}^{!}\sigma_{12}^{!}. \end{array}$

Now we introduce a linear map

(5.1)
$$\theta = (\theta_{ij}) = \begin{pmatrix} \mathrm{id} & \sigma_{12}^{!} \sigma_{11}^{!} + \sigma_{22}^{!} \sigma_{21}^{!} \\ 0 & \sigma_{22}^{!} \sigma_{11}^{!} + \sigma_{12}^{!} \sigma_{21}^{!} \end{pmatrix} : C_{A^{!}}(z) \to M_{2}(C_{A^{!}}(z)),$$

where each θ_{ij} is a \mathbb{Z}_2 -graded map for any i, j = 1, 2. Using the fact σ' is an algebra homomorphism, one obtains the following properites of θ .

Lemma 5.5. Retain the notations above. Then

- (1) θ is an algebra homomorphism;
- (2) $\theta_{12}\theta_{22} + \theta_{22}\theta_{12} = 0;$
- (3) $\theta_{22}^2 + \theta_{12}^2 = \mathrm{id}_{C_4!(z)}.$

For the \mathbb{Z}_2 -graded algebra $\mathbb{k} \times \mathbb{k}$, we choose an invertible \mathbb{k} -basis $\varepsilon = \{\varepsilon_1 = (1, 1), \varepsilon_2 = (1, -1)\}$.

Theorem 5.6. Let θ be the algebra homomorphisms from $C_{A^{!}}(z)$ to $M_{2}(C_{A^{!}}(z))$ defined in (5.1). Then $\theta_{\times} = (\theta, \varepsilon)$ is a twisting system of $C_{A^{!}}(z) \times C_{A^{!}}(z)$.

Proof. The linear map

$$\begin{pmatrix} \mathrm{id} & 0\\ -\varphi_{21}^{!}\varphi_{11}^{!} - \varphi_{22}^{!}\varphi_{12}^{!} & \varphi_{22}^{!}\varphi_{11}^{!} + \varphi_{21}^{!}\varphi_{12}^{!} \end{pmatrix} : C_{A^{!}}(z) \to M_{2}(C_{A^{!}}(z))$$

is the *t*-inverse of θ . Clearly, $\theta(1) = I_2 \in GL_2(\mathbb{k})$.

In this case, $l_{1;jj'} = \delta_{jj'}$ and $l_{2;jj'} = \delta_{jj'} + 1$ for any j, j' = 1, 2. It is straightforward but tedious to check the following simplified conditions of Definition 2.13 hold

$$\theta_{1j'} (\theta_{1j}(x)x') + \theta_{2j'} (\theta_{2j}(x)x') = x\theta_{jj'}(x') + \theta_{12}(x)\theta_{uj'}(x'), \theta_{1j'} (\theta_{2j}(x)x') + \theta_{2j'} (\theta_{1j}(x)x') = \theta_{22}(x)\theta_{uj'}(x'),$$

where $u \in \{1, 2\} - j$ for any j, j' = 1, 2 and $x, x' \in C_{A'}(z)$.

25

Hence, we have a \mathbb{Z}_2 -graded algebra $\Gamma = {}^{\theta_{\times}} (C_{A^!}(z) \times C_{A^!}(z))$ by Corollary 2.14, and the multiplication is as follows

(5.2)
$$\begin{aligned} \varepsilon_1 x \star \varepsilon_1 x' &= \varepsilon_1 x x', \quad \varepsilon_1 x \star \varepsilon_2 x' &= \varepsilon_1 \theta_{12}(x) x' + \varepsilon_2 \theta_{22}(x) x', \\ \varepsilon_2 x \star \varepsilon_1 x' &= \varepsilon_2 x x', \quad \varepsilon_2 x \star \varepsilon_2 x' &= \varepsilon_1 \theta_{22}(x) x' + \varepsilon_2 \theta_{12}(x) x', \end{aligned}$$

for any $x, x' \in C_{A^{!}}(z)$.

Lemma 5.7. The following map

$$\begin{split} \mu : & \Gamma & \to & \Gamma \\ \varepsilon_1 a & \mapsto & \varepsilon_1 \sigma_{11}^! \xi_{-1}(a) + \varepsilon_2 \sigma_{21}^! \xi_{-1}(a), \\ \varepsilon_2 a & \mapsto & \varepsilon_1 \sigma_{21}^! \xi_{-1}(a) + \varepsilon_2 \sigma_{11}^! \xi_{-1}(a), \end{split}$$

is a \mathbb{Z}_2 -graded algebra isomorphism satisfying $\mu^2 = \mathrm{id}$.

Proof. It is straightforward to check that μ is an algebra homomorphism by (5.2). For any $a \in C_{A^{!}}(z)$, one obtains that

$$\mu^{2}(\varepsilon_{1}a) = \varepsilon_{1}\left((\sigma_{11}^{!})^{2} + (\sigma_{21}^{!})^{2}\right)(a) + \varepsilon_{2}\left(\sigma_{21}^{!}\sigma_{11}^{!} + \sigma_{11}^{!}\sigma_{21}^{!}\right)(a) = \varepsilon_{1}a,$$

$$\mu^{2}(\varepsilon_{2}a) = \varepsilon_{1}\left(\sigma_{11}^{!}\sigma_{21}^{!} + \sigma_{21}^{!}\sigma_{11}^{!}\right)(a) + \varepsilon_{2}\left((\sigma_{21}^{!})^{2} + (\sigma_{11}^{!})^{2}\right)(a) = \varepsilon_{2}a.$$

Hence, $\mu^2 = \text{id}$ and μ is an isomorphism.

By Example 1.8(2), the following map

$$\psi: \ _{\mu}\Gamma(1) \otimes_{\Gamma} _{\mu}\Gamma(1) \rightarrow \Gamma$$
$$\varepsilon_{i}a \otimes \varepsilon_{i'}a' \quad \mapsto \quad \mu(\varepsilon_{i}a)\varepsilon_{i'}a'$$

is a \mathbb{Z}_2 -graded Γ -bimodule homomorphism, and we have a semi-trivial extension

$$\Gamma \ltimes_{\psi} (_{\mu} \Gamma(1)).$$

By Remark 1.7, it is $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded and also \mathbb{Z}_2 -graded by forgetting the first degree of $\mathbb{Z}_2 \times \mathbb{Z}_2$ degrees. So viewing this semi-trivial extension as a \mathbb{Z}_2 -graded algebra, the 0-th homogeneous space is also a \mathbb{Z}_2 -graded algebra.

Lemma 5.8. Retain the notations above. Then

ν = {ν₀ = id, ν₁ = μ} is a left Zhang-twisting system of Z₂-graded algebra Γ;
 ν Γ ≅ (Γ κ_ψ (μΓ(1)))₀ as Z₂-graded algebras.

Proof. Straightforwardly.

Theorem 5.9. Let $B = A_{\{-1,0\}}[y_1, y_2; \sigma]$ be a Z-graded double Ore extension of a noetherian Koszul Artin-Schelter regular algebra A and $z \in A_2$ be a nonzero regular central element of A. Suppose B is noetherian and $z + y_1^2 + y_2^2$ is a central element of B. Then

- (1) $C_{B^{!}}(z+y_{1}^{2}+y_{2}^{2}) \cong \Gamma \ltimes_{\psi} (_{\mu}\Gamma(1))$ as \mathbb{Z}_{2} -graded algebras;
- (2) there are equivalences of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b\left(\mathrm{gr}_{\mathbb{Z}_2}\,\Gamma\ltimes_{\psi}(_{\mu}\Gamma(1))\right) \cong D^b\left(\mathrm{mod}\,^{\nu}\Gamma\right),$$

where $\nu = \{\nu_0 = \mathrm{id}, \nu_1 = \mu\}$ is a left Zhang-twisting system of \mathbb{Z}_2 -graded algebra Γ .

26

Proof. (1) By Lemma 3.4(1), $C_{B^{!}}(z + y_1^2 + y_2^2)$ is generated by V^*, y_1^*, y_2^* subject to the relations consisting of the relations of $C_{A^{!}}(z)$ and

$$\begin{split} &(y_1^*)^2-1,(y_2^*)^2-1,y_1^*y_2^*-y_2^*y_1^*,\\ &v^*y_i^*+y_1^*\sigma_{1i}^*(v^*)+y_2^*\sigma_{2i}^*(v^*),\qquad \forall v^*\in V^*,i=1,2. \end{split}$$

Comparing with the multiplication list of $\Gamma \ltimes_{\psi} (\mu \Gamma(1))$ (see (5.2)), one obtains that the \mathbb{Z}_2 -graded algebra homomorphism from $C_{B^1}(z + y_1^2 + y_2^2)$ to $\Gamma \ltimes_{\psi} (\mu \Gamma(1))$ mapping

$$y_1^* \mapsto (0, \varepsilon_1), \qquad y_2^* \mapsto (0, \varepsilon_2), \qquad v^* \mapsto (\varepsilon_1 v^*, 0), \ \forall v^* \in V^*,$$

is an isomorphism. So $\Gamma \ltimes_{\psi} ({}_{\mu}\Gamma(1))$ is strongly \mathbb{Z}_2 -graded by Theorem 1.4(1).

(2) follows by Theorem 1.4 and Lemma 5.8.

Example 5.10. Let A be a noetherian Koszul Artin-Schelter regular algebra. Let σ_1 and σ_2 be graded automorphisms of A satisfying

$$\sigma_1^2 = \sigma_2^2 = \mathrm{id}_A, \qquad \sigma_1 \sigma_2 = \sigma_2 \sigma_1.$$

We extend σ_2 to be a graded automorphism $\widetilde{\sigma_2}$ of $A[y_1; \sigma_1]$ such that $\widetilde{\sigma_2}|_A = \sigma_2$ and $\widetilde{\sigma_2}(y_1) = -y_1$. Then we have an Ore extension $A[y_1; \sigma_1][y_2; \widetilde{\sigma_2}]$ is isomorphic to the double Ore extension $B = A_{\{-1,0\}}[y_1, y_2; \sigma]$, where

$$\sigma = \begin{pmatrix} \sigma_1 & 0\\ 0 & \sigma_2 \end{pmatrix} : A \to M_2(A).$$

Let $z \in A_2$ be a nonzero regular central element of A such that $\sigma_1(z) = \sigma_2(z) = z$. By Lemma 5.3, $z + y_1^2 + y_2^2$ is a central element of B. In this case, the linear map θ in the twisting system θ_{\times} is

$$\theta = \begin{pmatrix} \mathrm{id} & 0\\ 0 & \sigma_2^! \sigma_1^! \end{pmatrix} : C_{A^!}(z) \to M_2(C_{A^!}(z)).$$

Suppose $\sigma_1 = \sigma_2 = \sigma$, θ is just the diagonal map. So $\Gamma = {}^{\theta_{\times}}(C_{A^!}(z) \times C_{A^!}(z)) = C_{A^!}(z) \times C_{A^!}(z)$, and one obtains that $\bar{\nu} = \{\bar{\nu}_0 = \mathrm{id}, \bar{\nu}_1 = \sigma^! \xi_{-1}\}$ is a left Zhang-twisting system of \mathbb{Z}_2 -graded algebra $C_{A^!}(z)$ such that

$${}^{\nu}\Gamma \cong {}^{\bar{\nu}}C_{A^{!}}(z) \times {}^{\bar{\nu}}C_{A^{!}}(z),$$

as \mathbb{Z}_2 -graded algebras, where $\nu = \{\nu_0 = id, \nu_1 = \mu\}$ is a left Zhang-twisting system of \mathbb{Z}_2 -graded algebra Γ . By Theorem 5.9, there is an equivalence of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b \,(\mathrm{mod}\,{}^{\nu}C_{A^!}(z))^{\times 2}$$

In particular, if $\sigma = \xi_{-1}$, $\bar{\nu}C_{A^{\dagger}}(z) = C_{A^{\dagger}}(z)$ and so $\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b (\mathrm{mod} C_{A^{\dagger}}(z))^{\times 2}$. It is exactly [10, Theorem 1.4] (also see Example 4.9).

Example 5.11. Let $A = k_{-1}[x_1, x_2]$ and the central element $z = x_1^2 + x_2^2$. Let $B = A_{\{-1,0\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of A where $\sigma = (\sigma_{ij}) : A \to M_2(A)$ satisfies

$$\sigma_{11}(x_1) = -hx_1 + hx_2, \quad \sigma_{12}(x_1) = hx_1 + hx_2, \quad \sigma_{21}(x_1) = hx_1 + hx_2, \quad \sigma_{22}(x_1) = hx_1 - hx_2, \\ \sigma_{11}(x_2) = hx_1 - hx_2, \quad \sigma_{12}(x_2) = hx_1 + hx_2, \quad \sigma_{21}(x_2) = hx_1 + hx_2, \quad \sigma_{22}(x_2) = -hx_1 + hx_2,$$

where $4h^2 = 1$. It is a special case of T in [22]. One obtains that B is notherian by [22, Proposition 5.16] and σ satisfies Lemma 5.3. So $x_1^2 + x_2^2 + y_1^2 + y_2^2$ is a central element of B. In this case, the linear map θ in the twisting system θ_{\times} is

$$\theta = (\theta_{ij}) = \begin{pmatrix} \mathrm{id} & 0\\ 0 & \theta_{22} \end{pmatrix} : C_{A^{!}}(z) \to M_2(C_{A^{!}}(z)),$$

where $\theta_{22}(x_1^*) = x_2^*$, $\theta_{22}(x_2^*) = x_1^*$ and $\theta_{22}(x_1^*x_2^*) = x_1^*x_2^*$. Let $\Gamma = {}^{\theta_{\times}}(C_{A^!}(z) \times C_{A^!}(z))$ be the twisted direct product, μ be the \mathbb{Z}_2 -graded algebra isomorphism defined in Lemma 5.7 and ${}^{\nu}\Gamma$ be a Zhang-twisted algebra where $\nu = \{\nu_0 = \mathrm{id}, \nu_1 = \mu\}$ is a left Zhang-twisting system of \mathbb{Z}_2 -graded algebra Γ .

Write T_1, T_2, T_3, T_4 and T_5 for right ${}^{\nu}\Gamma$ -modules generated by $\{(1 - x_1^* x_2^*, 0), (x_1^* - x_2^*, 0)\}, \{\sqrt{-2h}(1 + x_1^* x_2^*, 0) - (x_1^* + x_2^*, 0)\}, \{\sqrt{2h}(0, 1 + x_1^* x_2^*) + (0, x_1^* + x_2^*)\}$ and $\{\sqrt{2h}(0, 1 + x_1^* x_2^*) - (0, x_1^* + x_2^*)\}$ respectively. It is routine to obtain that T_1, T_2, T_3, T_4, T_5 are all simple and non-isomorphic to each other, and ${}^{\nu}\Gamma \cong T_1^{\oplus 2} \oplus T_2 \oplus T_3 \oplus T_4 \oplus T_5$ as right ${}^{\nu}\Gamma$ -modules. So ${}^{\nu}\Gamma$ is semisimple and

$${}^{\nu}\Gamma \cong M_2(\Bbbk) \times \Bbbk^{\times 4}$$

By Theorem 5.9, there is an equivalence of triangulated categories

$$\underline{\mathrm{mcm}}(B/(z+y_1^2+y_2^2)) \cong D^b(\mathbb{k})^{\times 5}$$

Finally, we use a special case of the class \mathbb{R} constructed in [22] to show that noncommutative quadric hypersurfaces obtained by double Ore extension does not preserve the property of noncommutative graded isolated singularity. This behavior does not occur in the usual (second) double branched covers and tensor products of noncommutative quadric hypersurfaces.

Proposition 5.12. Let $A = k_{-1}[x_1, x_2]$ and $B = A_{\{-1,0\}}[y_1, y_2; \sigma]$ be a \mathbb{Z} -graded double Ore extension of A where $\sigma = (\sigma_{ij}) : A \to M_2(A)$ satisfies

$$\begin{aligned} \sigma_{11}(x_1) &= x_1 + x_2, \quad \sigma_{12}(x_1) = x_1, \quad \sigma_{21}(x_1) = x_2, \quad \sigma_{22}(x_1) = 0, \\ \sigma_{11}(x_2) &= 0, \quad \sigma_{12}(x_2) = x_1, \quad \sigma_{21}(x_2) = -x_2, \quad \sigma_{22}(x_2) = -x_1 + x_2. \end{aligned}$$

Then $B/(x_1^2 + x_2^2 + y_1^2 + y_2^2)$ is not a noncommutative graded isolated singularity.

Proof. By [21, Proposition 0.5], B is noetherian. It is straightforward to check that σ satisfies Lemma 5.3, and so $x_1^2 + x_2^2 + y_1^2 + y_2^2$ is a central element of B. So the noncommutative quadric hypersurface $B/(x_1^2 + x_2^2 + y_1^2 + y_2^2)$ is well defined. Write $z = x_1^2 + x_2^2$.

Note that $C_{A^{!}}(z)$ is generated by x_{1}^{*}, x_{2}^{*} subject to relations $x_{1}^{*}x_{2}^{*} - x_{2}^{*}x_{1}^{*}, (x_{1}^{*})^{2} - 1, (x_{2}^{*})^{2} - 1$. Write $\varepsilon = \{\varepsilon_{1} = (1, 1), \varepsilon_{2} = (1, -1)\}$. The linear map θ in the twisting system $\theta_{\times} = (\theta, \varepsilon)$ of $C_{A^{!}}(z) \times C_{A^{!}}(z)$ defined in (5.1) is

$$\theta = (\theta_{ij}) = \begin{pmatrix} \mathrm{id} & \theta_{12} \\ 0 & \theta_{22} \end{pmatrix} : C_{A^{!}}(z) \to M_2(C_{A^{!}}(z)),$$

where

$$\begin{aligned} \theta_{12}(x_1^*) &= x_1^* + x_2^*, \quad \theta_{12}(x_2^*) = x_1^* - x_2^*, \quad \theta_{12}(x_1^* x_2^*) = 2, \\ \theta_{22}(x_1^*) &= -x_2^*, \qquad \theta_{12}(x_2^*) = x_1^*, \qquad \theta_{22}(x_1^* x_2^*) = -x_1^* x_2^*. \end{aligned}$$

Let $\Gamma = {}^{\theta_{\times}}(C_{A^{!}}(z) \times C_{A^{!}}(z))$ be the twisted direct product, μ be the \mathbb{Z}_{2} -graded algebra isomorphism defined in Lemma 5.7 and ${}^{\nu}\Gamma$ be a Zhang-twisted algebra of Γ where $\nu = \{\nu_{0} = \mathrm{id}, \nu_{1} = \mu\}$ is a left Zhang-twisting system of \mathbb{Z}_{2} -graded algebra Γ . We list partial multiplication of ${}^{\nu}\Gamma$:

$$\begin{array}{ll} (1,0)*(a,b)=(a,0), & (x_1^*,0)*(0,b)=(0,0), \\ (x_1^*x_2^*,0)*(1,0)=(1,0), & (x_1^*x_2^*,0)*(x_1^*x_2^*,0)=(x_1^*x_2^*,0), \\ (x_1^*x_2^*,0)*(x_1^*,0)=(x_1^*,0), & (x_1^*x_2^*,0)*(x_2^*,0)=(x_2^*,0), \\ (x_1^*x_2^*,0)*(0,1)=(x_1^*x_2^*-1,0), & (x_1^*x_2^*,0)*(0,x_1^*x_2^*)=(1-x_1^*x_2^*,0), \\ (x_1^*x_2^*,0)*(0,x_1^*)=(x_1^*-x_2^*,0), & (x_1^*x_2^*,0)*(0,x_2^*)=(x_2^*-x_1^*,0), \\ (x_1^*,0)*(1,0)=(x_1^*,0), & (x_1^*,0)*(x_1^*x_2^*,0)=(x_2^*,0), \\ (x_1^*,0)*(x_1^*,0)=(x_1^*,0), & (x_1^*,0)*(x_1^*x_2^*,0)=(x_2^*,0), \\ (x_2^*,0)*(1,0)=(x_1^*,0), & (x_2^*,0)*(x_1^*x_2^*,0)=(x_2^*,0), \\ (x_2^*,0)*(x_1^*,0)=(-1,0), & (x_2^*,0)*(x_1^*x_2^*)=(x_1^*-x_2^*,0), \\ (x_2^*,0)*(0,1)=(x_2^*-x_1^*,0), & (x_2^*,0)*(x_1^*x_2^*)=(x_1^*-x_2^*,0), \\ (x_2^*,0)*(0,x_1^*)=(x_1^*x_2^*-1,0), & (x_2^*,0)*(0,x_2^*)=(1-x_1^*x_2^*,0), \end{array}$$

for any $a, b \in C_{A^{!}}(z)$.

One obtains that the right ${}^{\nu}\Gamma$ -module generated by $(1 - x_1^* x_2^*, 0)$ is nilpotent. So ${}^{\nu}\Gamma$ is not semisimple and $B/(x_1^2 + x_2^2 + y_1^2 + y_2^2)$ is not a noncommutative graded isolated singularity by Lemma 5.8, Theorem 5.9 and Theorem 1.4.

Acknowledgments. Yuan Shen is supported by Zhejiang Provincial Natural Science Foundation of China under Grant No. LY24A010006 and National Natural Science Foundation of China under Grant Nos. 11701515 and 12371101. Xin Wang is supported by Doctoral Research Fund of Shandong Jianzhu University under Grant No. XNBS1943.

References

- A. Conner, E. Kirkman, W. F. Moore and C. Walton, Noncommutative Knörrer periodicity and noncommutative Kleinian singularities, J. Algebra, 540 (2019), 234–273.
- [2] Ji-Wei He and Yu Ye, Clifford deformation of Koszul Frobenius algebras and noncommutative qudrics, Algebra Colloquium 31, (2024), 63–82.
- [3] Ji-Wei He, Xin-Chao Ma and Yu Ye, Generalized Knörrer's Periodicity Theorem, arXiv:2107.06438.
- [4] A. Higashitani and K. Ueyama, Combinatorial study of stable categories of graded Cohen-Macaulay modules over skew quadric hypersurfaces, Collectanea Math., 73 (2022), 43–54
- [5] H. Hu, Classification of noncommutative conics associated to symmetric regular superpotentials, J. Algebra Appl., 22(6) (2023), 2350136.
- [6] H. Hu, M. Matsuno and I. Mori, Noncommutative conics in Calabi-Yau quantum planes, J. Algebra, 620 (2023), 194–224.
- [7] H. Knörrer, Cohen-Macaulay modules on hypersurface singularities I, Invent. Math. 88 (1987), 153-164.
- [8] J. C. McConnell and J. C. Robson, Noncommutative Noetherian rings, vol. 30 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, revised ed., 2001.
- [9] I. Mori and K. Ueyama, Noncommutative matrix factorizitons with an application to skew exterior algebras, J. Algebra, 586 (2021), 1053–1087.
- [10] I. Mori, K. Ueyama, Noncommutative Knörrer's Periodicity theorem and noncommutative quadric surfaces, Algebra Number Theory, 16(2) (2022), 467—504.
- [11] C. Năstăsescu, F. Van Oystaeyen, Methods of Graded Rings, Lect. Notes Math. 1836, Springer, 2004.

- [12] I. Palmér, The global homological dimension of semi-trivial extensions of rings, Math. Scand., 37 (1975), 223-256.
- [13] S. B. Priddy, Koszul resolutions, Trans. Amer. Math. Soc. 152 (1970), 39-60.
- [14] Y. Shen, G.-S. Zhou and D.-M. Lu, Nakayama automorphisms of twisted tensor products, J. Algebra, 504 (2018), 445–478.
- [15] S. P. Smith, Some finite dimensional algebras related to elliptic curves, CMS Conf. Proc., 19 (1996), 315–348.
- [16] S. P. Smith and M. Van den Bergh, Noncommutative quadric surfaces, J. Noncommut. Geom. 7 (2013), 817– 856.
- [17] K. Ueyama, Graded maximal Cohen-Macaulay modules over noncommutative graded Gorenstein isolated singularities, J. Algebra, 383 (2013), 85–103.
- [18] K. Ueyama, On Knoörrer periodicity for quadric hypersurfaces in skew projectives spaces, Canda. Math. Bull., 62 (2019), 896–911.
- [19] K. Ueyama, Skew graded (A_∞) hypersurface singularities, C. R. Math. Acad. Sci. Paris, **361** (2023), 521–534.
- [20] J. J. Zhang, Twisted graded algebras and equivalences of graded categories, Proc. London Math. Soc., 72(2) (1996), 281–311.
- [21] J. J. Zhang and J. Zhang, Double Ore extension, J. Pure Appl. Algebra, 212(12) (2008), 2668–2690.
- [22] J. J. Zhang and J. Zhang, Double extension regular algebras of type (14641), J. Algebra, 322 (2009), 373–409.
- [23] C. Zhu, F. Van Oystaeyen and Y.-H. Zhang, Nakayama automorphisms of double Ore extensions of Koszul regular algebras, Mauscripta Math., 152 (2017), 555–584.

LIU: DEPARTMENT OF MATHEMATICS, ZHEJIANG SCI-TECH UNIVERSITY, HANGZHOU 310018, CHINA

Email address: wxly120@gmail.com

SHEN: DEPARTMENT OF MATHEMATICS, ZHEJIANG SCI-TECH UNIVERSITY, HANGZHOU 310018, CHINA

 $Email \ address: \ {\tt yuanshen} {\tt @zstu.edu.cn}$

WANG: SCHOOL OF SCIENCE, SHANDONG JIANZHU UNIVERSITY, JINAN 250101, CHINA

Email address: wangxin19@sdjzu.edu.cn