

A PRIORI ESTIMATION OF THE APPROXIMATION, OPTIMIZATION AND GENERALIZATION ERROR OF RANDOM NEURAL NETWORKS FOR SOLVING PARTIAL DIFFERENTIAL EQUATIONS

XIANLIANG XU AND ZHONGYI HUANG[†]

Department of Mathematics, Tsinghua University, Beijing 100084, China.

ABSTRACT. In recent years, there are numerous methods involving neural networks for solving partial differential equations (PDEs), such as Physics informed neural networks (PINNs), Deep Ritz method (DRM) and others. However, the optimization problems are typically non-convex, which makes these methods lead to unsatisfactory solutions. With weights sampled from some distribution, applying random neural networks to solve PDEs yields least squares problems that are easily solvable. In this paper, we focus on Barron type functions and demonstrate the approximation, optimization and generalization of random neural networks for solving PDEs.

1. INTRODUCTION

In many disciplines of science and engineering, PDEs have been utilized in modeling phenomena and characterizing physical laws. However, classical numerical methods such as, e.g., finite difference or finite elements methods are typically impractical in solving high-dimensional nonlinear PDEs. In recent years, inspired by the breakthrough of deep learning and rapid development of hardware, numerical methods involving neural networks have been proposed for solving numerous types of PDEs. Some methods, like PINNs [1] and DRM [2], train the neural networks by minimizing the designed loss functionals for the specific forms of PDEs. Although, compared to classical numerical methods, these methods involving neural networks may not suffer from the curse of dimensionality, they may lead to poor solutions due to the complex loss functionals [3]. Optimizing neural network parameters according to the loss functionals is typically a non-convex optimization problem, which are hard to solve.

Random neural networks [4] are neural networks whose weights of hidden layers are randomly initialized and only the parameters in the output layer are trained. Therefore, compared to the deterministic neural networks, random neural networks provide extremely faster learning speed as the non-convex optimization problem reduces to a convex optimization problem. Despite this, it is still unclear how to choose the priori distribution for the hidden parameters. In this paper, we focus on the approximation of two-layer random neural networks for Barron type functions due to their special integral representations. The form of the integral representation provides

E-mail address: xuxl19@mails.tsinghua.edu.cn; zhongyih@mail.tsinghua.edu.cn.

[†] Corresponding author.

* This work was partially supported by the NSFC Projects No. 12025104, 11871298, 81930119.

the sampling distribution. When applying random neural network in the framework of PINNs, we add a penalty, which yields a form similar to the ridge regression. Then we investigate the optimization error and generalization error of this method.

1.1. Notations. For $x \in \mathbb{R}^d$, $|x|_p$ denotes its p -norm and we use $|x|$ as shorthand for $|x|_2$. For the activation functions, we write $\sigma_k(x)$ for the ReLU^k activation function, i.e., $\sigma_k(x) := [\max(0, x)]^k$. For given probability measure P and a sequence of random variables $\{X_i\}_{i=1}^n$ distributed according to P , we denote the empirical measure of P by P_n , i.e. $P_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$.

2. MAIN RESULTS

Before presenting the approximation results, we first introduce the definition of the Barron space.

$$\mathcal{B}^s(\Omega) := \{f : \Omega \rightarrow \mathbb{C} : \|f\|_{\mathcal{B}^s(\Omega)} := \inf_{f_e \in L^1(\mathbb{R}^d)} \int_{\mathbb{R}^d} (1 + |\omega|_1)^s |\hat{f}_e(\omega)| d\omega < \infty\},$$

where the infimum is over extensions $f_e \in L^1(\mathbb{R}^d)$ and \hat{f}_e is the Fourier transform of f_e . Note that we choose 1-norm in the definition just for simplicity. In the following, we assume that Ω is a subset of $[0, 1]^d$.

Our method for approximating Barron functions by the random two-layer neural networks are based on the integral representations of these functions, which are introduced in [5]. Specifically, for $f \in \mathcal{B}^s(\Omega)$, without loss of generality, assume that the infimum is attained at f_e . Then for activation function σ with $\hat{\sigma}(a) \neq 0$ and some $a \neq 0$, we have

$$\begin{aligned} f_e(x) &= \int_{\mathbb{R}^d} e^{i\omega \cdot x} \hat{f}_e(\omega) d\omega \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi \hat{\sigma}(a)} \sigma\left(\frac{\omega}{a} \cdot x + b\right) \hat{f}_e(\omega) e^{-iab} db d\omega \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi \hat{\sigma}(a)} \sigma\left(\frac{\omega}{a} \cdot x + b\right) |\hat{f}_e(\omega)| \cos(\theta(\omega) - ab) db d\omega, \end{aligned}$$

where $\hat{f}_e(\omega) = e^{i\theta(\omega)} |\hat{f}_e(\omega)|$.

In the paper, we just consider the activation function $\sigma(t) = \sum_{i=0}^4 (-1)^i C_4^i \sigma_3(t+2-i)$, which is compactly supported on $[-2, 2]$. The reason is that we require the polynomial decay condition of the activation function, i.e., $|\sigma^{(k)}(t)| \leq C_p (1 + |t|)^{-p}$ for $0 \leq k \leq 2$ and some $p > 1$.

Theorem 1. *Let P_1 be the uniform distribution of the domain $\{\omega \in \mathbb{R}^d : |\omega|_1 \leq M\}$ and P_2 be the uniform distribution of the domain $\{b \in \mathbb{R} : |b| \leq 2M\}$ with a constant $M \geq 2$. Let $W_1, \dots, W_m \sim P_1$ and $B_1, \dots, B_m \sim P_2$, then there exists a \mathbb{R}^m -valued, $\sigma(W, B)$ -measurable vector A such that*

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \leq C \begin{cases} \frac{M^{d+7-s}}{m} \|f\|_{\mathcal{B}^s(\Omega)} \|\hat{f}_e\|_{L^\infty(\mathbb{R}^d)} + \left(\frac{1}{M^{s-3}} \|f\|_{\mathcal{B}^s(\Omega)} \right)^2, & 3 < s < 6, \\ \frac{M^{d+1}}{m} \|f\|_{\mathcal{B}^s(\Omega)} \|\hat{f}_e\|_{L^\infty(\mathbb{R}^d)} + \left(\frac{1}{M^{s-3}} \|f\|_{\mathcal{B}^s(\Omega)} \right)^2, & s \geq 6. \end{cases} \quad (1)$$

where

$$f_m(x) = \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i)$$

and C is a universal constant. Moreover,

$$|A_i| \leq C \|\hat{f}_e\|_{L^\infty(\mathbb{R}^d)} M^{d+1}.$$

Remark 1. By appropriately selecting the value of M , we can obtain that

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \begin{cases} m^{-\frac{2s-6}{d+s+1}}, & 3 < s < 6, \\ m^{-\frac{2s-6}{d+2s-5}}, & s \geq 6. \end{cases} \quad (2)$$

Remark 2. For other activation functions, like Sigmoidal function ($\text{Sig}(x) := (1 + e^{-x})^{-1}$) and Hyperbolic tangent function ($\text{Tanh}(x) := \frac{e^x - e^{-x}}{e^x + e^{-x}}$), let $\sigma(x) = \text{Sig}(x+1) - \text{Sig}(x)$ or $\sigma(x) = \text{Tanh}(x+1) - \text{Tanh}(x)$. We can deduce that

$$|\sigma^{(k)}(t)| \leq C e^{-|t|}$$

for $0 \leq k \leq 2$ and some universal constant C .

Therefore, for $f_{22}(x)$ in the proof of Theorem 1, we have that

$$\|f_{22}(x)\|_{H^2(\Omega)} \lesssim e^{-M} \|f\|_{\mathcal{B}^s(\Omega)},$$

which implies that

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \leq C \begin{cases} \frac{M^{d+7-s}}{m} \|f\|_{\mathcal{B}^s(\Omega)} \|\hat{f}_e\|_{L^\infty(\mathbb{R}^d)} + \left(\left(\frac{1}{M^{s-3}} + e^{-M} \right) \|f\|_{\mathcal{B}^s(\Omega)} \right)^2, & 3 < s < 6, \\ \frac{M^{d+1}}{m} \|f\|_{\mathcal{B}^s(\Omega)} \|\hat{f}_e\|_{L^\infty(\mathbb{R}^d)} + \left(\left(\frac{1}{M^{s-3}} + e^{-M} \right) \|f\|_{\mathcal{B}^s(\Omega)} \right)^2, & s \geq 6. \end{cases}$$

Theorem 3 in [5] implies that, when $\sigma \in W^{m,\infty}(\mathbb{R})$ is a non-constant periodic function, f has a similar integral representation for σ . Moreover, in this case, we do not truncate b , which may yields a better approximation rate.

For brevity and readability, we only consider the case where s is sufficiently large, for example, $2(s-3) = d+1$, i.e., $s = \frac{d+7}{2}$. At this time,

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \frac{1}{\sqrt{m}}, \quad |A_i| \lesssim \sqrt{m}.$$

When applying the random neural networks in the framework of PINNs, we consider the following equation.

$$\begin{cases} -\Delta u + Vu = f, & \text{in } \Omega, \\ u = g, & \text{on } \partial\Omega, \end{cases} \quad (3)$$

where $f, g \in L^\infty(\Omega)$.

Assume that the solution of the equation belongs to $\mathcal{B}^s(\Omega)$ with $s = \frac{d+7}{2}$. From the approximation result in Theorem 1, we can construct a random neural network u of the form

$$u(x) = \sum_{i=1}^m a_i \sigma(\omega_i \cdot x + b_i),$$

where $\omega_1, \dots, \omega_m \sim P_1$, $b_1, \dots, b_m \sim P_2$ and $a = (a_1, \dots, a_m) \in \mathbb{R}^m$ is to be determined.

Then the loss function of PINNs can be written as

$$\begin{aligned} L(a) &:= \int_{\Omega} (-\Delta u + Vu - f)^2 dx + \int_{\partial\Omega} (u - g)^2 dy \\ &= \int_{\Omega} \left(\sum_{i=1}^m a_i \left(-\sum_{k=1}^d \omega_{i,k}^2 \sigma''(\omega_i \cdot x + b_i) + V\sigma(\omega_i \cdot x + b_i) \right) - f(x) \right)^2 dx \\ &\quad + \int_{\partial\Omega} \left(\sum_{i=1}^m a_i \sigma(\omega_i \cdot y + b_i) - g(y) \right)^2 dy \\ &= \int_{\Omega} (a \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a \cdot f_2(y) + g_2(y))^2 dy, \end{aligned}$$

where $g_1(x) = -f(x)$, $g_2(y) = -g(y)$ and the i -th component of \mathbb{R}^m -valued functions $f_1(x)$ and $f_2(y)$ are

$$f_1^i(x) = -\sum_{k=1}^d \omega_{i,k}^2 \sigma''(\omega_i \cdot x + b_i) + V\sigma(\omega_i \cdot x + b_i) = |\omega_i|^2 \sigma''(\omega_i \cdot x + b_i) + V\sigma(\omega_i \cdot x + b_i)$$

and $f_2^i(y) = \sigma(\omega_i \cdot y + b_i)$ respectively.

To solve a , we consider the following loss function.

$$F(a) := L(a) + \lambda|a|^2 = \int_{\Omega} (a \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a \cdot f_2(y) + g_2(y))^2 dy + \lambda|a|^2.$$

For simplicity, we only consider the case where the number of samples in the interior and on the boundary are the same. Therefore, the empirical loss function is

$$F_n(a) := \frac{1}{n} \sum_{i=1}^n l(a; x_i, y_i) + \lambda|a|^2,$$

where

$$l(a; x_i, y_i) = |\Omega|(a \cdot f_1(x_i) + g_1(x_i))^2 + |\partial\Omega|(a \cdot f_2(y_i) + g_2(y_i))^2.$$

As $u \in \mathcal{B}^s(\Omega)$ with $s = \frac{d+7}{2}$, then Theorem 1 implies there is a random neural network $u_m = \sum_{i=1}^m \bar{a}_i \sigma(\omega_i \cdot x + b_i)$ such that

$$\mathbb{E} \left[\|u(x) - u_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \frac{1}{\sqrt{m}},$$

where $\bar{a} = (\bar{a}_1, \dots, \bar{a}_m)$, $|\bar{a}_i| \lesssim 1/\sqrt{m}$ for $1 \leq i \leq m$.

As $|\bar{a}| = \sqrt{\sum_{i=1}^m \bar{a}_i^2} = \mathcal{O}(1)$, we can use projected gradient descent method for $a \in \mathcal{A} := \{a \in \mathbb{R}^m : |a| \leq C\}$, where C is a constant such that $\bar{a} \in \mathcal{A}$. Specifically, the method of projected gradient descent consists of iteration of the following update rule for $t = 1, \dots, T$,

$$\begin{aligned} y_t &= a_t - \nu_t g_t, \text{ where } g_t \in \partial F_n(a_t) \\ a_{t+1} &= P_{\mathcal{A}}(y_t), \end{aligned}$$

where T is the total number of iterations, a_1 is an initial approximation and $P_{\mathcal{A}}$ is the projection operator onto the convex closed set \mathcal{A} .

Theorem 2. Choosing $\nu_t = \mathcal{O}(m^{\frac{d+1}{d+3}})$, then

$$\begin{aligned} & \mathbb{E}_{W,B} \mathbb{E}_{X,Y} \left[\int_{\Omega} (a_T \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a_T \cdot f_2(y) + g_2(y))^2 dy \right] \\ & \lesssim \frac{1}{\sqrt{m}} + \lambda + \frac{m^{3+\frac{6}{d+1}}}{\lambda^2} e^{-\frac{\lambda T}{m^{1+2/(d+1)}}} + \frac{m^{2+\frac{4}{d+1}}}{\lambda n} \log \log(mn). \end{aligned}$$

Taking $\lambda = \frac{1}{\sqrt{m}}$, $m = n^{\frac{d+1}{3d+7}}$ and $T = \mathcal{O}(\sqrt{n} \log(n))$, the final bound is $\mathcal{O}(n^{-\frac{d+1}{2(3d+7)}} \log \log(n))$.

Remark 3. In Theorem 1, we have neglected the dependency on the dimension. In fact, the final bound in Theorem 1 depends quadratically on $|\Omega|$ and $|\partial\Omega|$. Moreover, it is linear with respect to the constant in the Sobolev trace theorem. And the generalization error can also be measured by some appropriate Sobolev norm. Specifically, as shown in [6], we have that

$$\|u - u^*\|_{H^{\frac{1}{2}}(\Omega)}^2 \lesssim \|-\Delta u + Vu - f\|_{L^2(\Omega)}^2 + \|u - g\|_{L^2(\partial\Omega)}^2,$$

where u^* is the solution of equation(3).

Remark 4. [7] has utilized algorithm stability to derive excess risk of projected gradient descent method for λ -strongly convex and L -Lipschitz continuous loss. It is shown that $T = \mathcal{O}(n^2)$ iterations are required to attain the excess risk $\tilde{\mathcal{O}}(\frac{L^2}{\lambda n})$. Our result implies that $T = \mathcal{O}(\sqrt{n} \log(n))$ is sufficient to reach the same excess risk (omitting the logarithm term). Note that our method for deriving the generalization bound only requires the strong convexity and Lipschitz continuity. By setting $\nu_t = \frac{2}{\lambda(t+1)}$ and $a_T = \frac{2}{T(T+1)} \sum_{t=1}^T ta_t$ in the projected gradient descent method for non-smooth loss, our method can yield the same excess risk after $T = \mathcal{O}(n)$ steps.

Note that when s is large enough, we can choose not to truncate ω and b . Specifically, assume that $s > \alpha + \beta + 5$ with $\alpha > d, \beta > 1$, then for $f \in \mathcal{B}^s(\Omega)$, we have

$$\begin{aligned} f(x) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) \frac{(1 + |\omega|)^\alpha (1 + |b|)^\beta}{C_\alpha C_\beta} \frac{C_\alpha C_\beta}{(1 + |\omega|)^\alpha (1 + |b|)^\beta} db d\omega \\ &= \mathbb{E}_{P_1(\omega), P_2(b)} \left[\frac{1}{2\pi\hat{\sigma}(1)} \frac{\sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega) p_2(b)} \right], \end{aligned}$$

where $P_1(\omega) = p_1(\omega) d\omega, P_2(b) = p_2(b) db$ and $p_1(\omega) := \frac{C_\alpha}{(1+|\omega|)^\alpha}, p_2(b) := \frac{C_\beta}{(1+|b|)^\beta}$ are density functions in \mathbb{R}^d and \mathbb{R} respectively.

At this time, we can consider a smaller function space. For the function f that we want to approximate, assume that there is a function $f_e \in L^1(\mathbb{R}^d)$ such that $f_e|_{\Omega} = f$ and

$$\int_{\mathbb{R}^d} (1 + |\omega|)^s |\hat{f}_e(\omega)|^2 d\omega < \infty \quad (4)$$

with $s > \alpha + \beta + 5$ and $\alpha > d, 1 < \beta < 5$.

Theorem 3. Let $W_1, \dots, W_m \sim P_1$ and $B_1, \dots, B_m \sim P_2$, then there exists a \mathbb{R}^m -valued, $\sigma(W, B)$ -measurable vector A such that

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \frac{1}{m} \frac{d^{\beta+1}}{C_\alpha},$$

where

$$f_m(x) = \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i)$$

and

$$A_i = \frac{1}{2\pi \hat{\sigma}(1)} \frac{|\hat{f}(W_i)| \cos(\theta(W_i) - B_i)}{p_1(W_i) p_2(B_i)} I_{\{|B_i| \leq Cd(1+|W_i|)\}}, 1 \leq i \leq m.$$

Remark 5. Note that $\frac{1}{C_\alpha} \sim \frac{1}{\alpha-1} \mathcal{H}^{d-1}(S^{d-1})$, i.e., the hypervolume of the $(d-1)$ -dimensional unit sphere. Therefore, when d is sufficiently large, this term becomes small, and $\frac{d^{\beta+1}}{C_\alpha}$ does too.

Assume that the solution of equation(3) satisfies the condition(4), then we can derive the generalization bound as follows.

Theorem 4. Assume that $a_n = \arg \min_{a \in \mathbb{R}^m} F_n(a)$, then

$$\mathbb{E}_{W,B} \mathbb{E}_{X,Y} \left[\int_{\Omega} (a_n \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a_n \cdot f_2(y) + g_2(y))^2 dy \right] \lesssim \frac{m}{\lambda \sqrt{n}} + \frac{1}{m} + \frac{\lambda}{m}.$$

By taking $\lambda = \mathcal{O}(1)$, $m = n^{1/4}$, the final bound is $\mathcal{O}(n^{-1/4})$.

Moreover, a_n can be solved by using the closed form of the ridge regression and the complexity is $\mathcal{O}(m^2 n) = \mathcal{O}(n^{3/2})$.

3. PROOF

3.1. Proof of Theorem 1. Note that the activation function σ satisfies the polynomial decay condition

$$|\sigma^{(k)}(t)| \leq C_p (1 + |t|)^{-p}$$

for $0 \leq k \leq 2$ and any $p > 1$, as σ is compactly supported in $[-2, 2]$. We can just take $p = 2$.

Thus for any $x \in \Omega$ and $s \in \mathbb{N}$, we have

$$\begin{aligned} & \int_{\mathbb{R}} |\sigma^{(k)}(\omega \cdot x + b)|^s db \\ & \leq C_2 \int_{\mathbb{R}} (1 + |\omega \cdot x + b|)^{-2s} db \\ & \leq C_2 \int_{\mathbb{R}} (1 + \max(0, |b| - |\omega|_1))^{-2s} db \\ & \leq C_2 \int_{\{|b| \leq |\omega|_1\}} db + 2C_2 \int_{\{b > |\omega|_1\}} (1 + b - |\omega|_1)^{-2s} db \\ & = C_2 \left(2|\omega|_1 + 2 \frac{1}{2s-1} \right) \\ & \leq 2C_2 (1 + |\omega|_1). \end{aligned}$$

In the integral representation of f , we simultaneously truncate both ω and b .

$$\begin{aligned}
f(x) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&= \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&\quad + \int_{\{|\omega|_1 \leq M, |b| \leq 2M\}^c} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&:= f_1(x) + f_2(x).
\end{aligned}$$

Then, we use a random neural network to approximate $f_1(x)$ and take $f_2(x)$ as the remainder term.

We first estimate the $H^2(\Omega)$ norm of the remainder term $f_2(x)$. Note that $f_2(x)$ can be written as follows.

$$\begin{aligned}
f_2(x) &= \int_{\{|\omega|_1 \leq M, |b| \leq 2M\}^c} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&= \int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&\quad + \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| > 2M\}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&:= f_{21}(x) + f_{22}(x).
\end{aligned}$$

For $f_{21}(x)$, we have

$$\begin{aligned}
\|f_{21}(x)\|_{H^2(\Omega)}^2 &= \int_{\Omega} \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 dx \\
&\quad + \int_{\Omega} \sum_{k=1}^d \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \omega_k \sigma'(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 dx \\
&\quad + \int_{\Omega} \sum_{i=1}^d \sum_{j=1}^d \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \omega_i \omega_j \sigma''(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 dx.
\end{aligned}$$

For the first term,

$$\begin{aligned}
&\left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 \\
&\leq \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} |\sigma(\omega \cdot x + b)| |\hat{f}(\omega)| db d\omega \right)^2 \\
&\lesssim \left(\int_{\{|\omega|_1 > M\}} (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right)^2 \\
&\leq \left(\frac{1}{M^{s-1}} \|f\|_{B^s(\Omega)} \right)^2,
\end{aligned}$$

where the second inequality follows from the polynomial decay condition of σ .

For the second term,

$$\begin{aligned}
& \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \omega_k \sigma'(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 \\
& \leq \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} |\omega_k| |\sigma'(\omega \cdot x + b)| |\hat{f}(\omega)| db d\omega \right)^2 \\
& \lesssim \left(\int_{\{|\omega|_1 > M\}} |\omega_k| (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right)^2 \\
& \leq \left(\int_{\{|\omega|_1 > M\}} |\omega_k|^2 |\hat{f}(\omega)| d\omega \right) \left(\int_{\{|\omega|_1 > M\}} (1 + |\omega|_1)^2 |\hat{f}(\omega)| d\omega \right),
\end{aligned}$$

where the last inequality follows from the Cauchy inequality.

Adding k from 1 to d yields that

$$\begin{aligned}
& \int_{\Omega} \sum_{k=1}^d \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \frac{1}{2\pi \hat{\sigma}(1)} \omega_k \sigma'(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 dx \\
& \lesssim \left(\int_{\{|\omega|_1 > M\}} (1 + |\omega|_1)^2 |\hat{f}(\omega)| d\omega \right) \left(\int_{\{|\omega|_1 > M\}} (1 + |\omega|_1)^2 |\hat{f}(\omega)| d\omega \right) \\
& \leq \left(\frac{1}{M^{s-2}} \int_{\{|\omega|_1 > M\}} (1 + |\omega|_1)^s |\hat{f}(\omega)| d\omega \right)^2 \\
& \leq \left(\frac{1}{M^{s-2}} \|f\|_{B^s(\Omega)} \right)^2.
\end{aligned}$$

Similarly, for the third term, we have

$$\begin{aligned}
& \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \omega_i \omega_j \sigma''(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 \\
& \lesssim \left(\int_{\{|\omega|_1 > M\}} |\omega_i \omega_j| (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right)^2 \\
& \leq \left(\int_{\{|\omega|_1 > M\}} |\omega_i|^2 (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right) \left(\int_{\{|\omega|_1 > M\}} |\omega_j|^2 (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right).
\end{aligned}$$

Therefore,

$$\begin{aligned}
& \sum_{i=1}^d \sum_{j=1}^d \left(\int_{\{|\omega|_1 > M\}} \int_{\mathbb{R}} \frac{1}{2\pi \hat{\sigma}(1)} \omega_i \omega_j \sigma''(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \right)^2 \\
& \lesssim \left(\sum_{i=1}^d \int_{\{|\omega|_1 > M\}} |\omega_i|^2 (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right) \left(\sum_{j=1}^d \int_{\{|\omega|_1 > M\}} |\omega_j|^2 (1 + |\omega|_1) |\hat{f}(\omega)| d\omega \right) \\
& \leq \left(\int_{\{|\omega|_1 > M\}} (1 + |\omega|_1)^3 |\hat{f}(\omega)| d\omega \right)^2 \\
& \leq \left(\frac{1}{M^{s-3}} \|f\|_{B^s(\Omega)} \right)^2.
\end{aligned}$$

Combining the estimation above for the $H^2(\Omega)$ norm of $f_{21}(x)$, we can deduce that

$$\|f_{21}(x)\|_{H^2(\Omega)} \lesssim \frac{\|f\|_{B^s(\Omega)}}{M^{s-3}}.$$

For $f_{22}(x)$, recall that

$$f_{22}(x) = \int_{\{|\omega| \leq M\}} \int_{\{|b| > 2M\}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega.$$

Note that when $|\omega| \leq M$, $|b| > 2M$, we have

$$|\omega \cdot x + b| \geq |b| - |\omega \cdot x| \geq |b| - |\omega|_1 \geq M \geq 2,$$

which implies that $\sigma(\omega \cdot x + b) = 0$, as σ is compactly supported in $[-2, 2]$. Thus, $f_{22}(x) = 0$ and

$$\|f_2(x)\|_{H^2(\Omega)} = \|f_{21}(x)\|_{H^2(\Omega)} \lesssim \frac{\|f\|_{B^s(\Omega)}}{M^{s-3}}.$$

It remains to consider the stochastic approximation of the main term $f_1(x)$, recall that

$$f_1(x) = \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega.$$

Thus, we can express $f_1(x)$ in the form of expectation.

$$\begin{aligned} f_1(x) &= \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} \frac{1}{2\pi\hat{\sigma}(1)} \frac{\sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega)p_2(b)} p_1(\omega)p_2(b) db d\omega \\ &= \mathbb{E}_{P_1(\omega), P_2(b)} \left[\frac{1}{2\pi\hat{\sigma}(1)} \frac{\sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega)p_2(b)} \right], \end{aligned}$$

where $P_1(\omega) := p_1(\omega)d\omega$, $P_2(b) := p_2(b)db$ are uniform distributions over $\{|\omega|_1 \leq M\}$ and $\{|b| \leq 2M\}$ respectively, i.e., $p_1(\omega) = \frac{1}{C_d M^d}$, $p_2(b) = \frac{1}{4M}$.

Thus, $f_1(x)$ can be written as

$$f_1(x) = \mathbb{E}_{P_1(W), P_2(B)} [A\sigma(W \cdot x + B)],$$

where

$$A = \frac{1}{2\pi\hat{\sigma}(1)} \frac{|\hat{f}(W)| \cos(\theta(W) - B)}{p_1(W)p_2(B)}.$$

Taking $W_1, \dots, W_m \sim P_1$ and $B_1, \dots, B_m \sim P_2$, let

$$A_i = \frac{1}{2\pi\hat{\sigma}(1)} \frac{|\hat{f}(W_i)| \cos(\theta(W_i) - B_i)}{p_1(W_i)p_2(B_i)}, 1 \leq i \leq m,$$

we can construct the random neural network $f_m(x)$ as follows.

$$f_m(x) = \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i).$$

Next, we aim to estimate $\mathbb{E} \left[\|f_1(x) - f_m(x)\|_{H^2(\Omega)}^2 \right]$.

$$\begin{aligned}
& \mathbb{E} \left[\|f_1(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \\
&= \mathbb{E} \left[\left\| f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right\|_{H^2(\Omega)}^2 \right] \\
&= \mathbb{E} \left[\int_{\Omega} \left(f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right)^2 dx \right] \\
&\quad + \mathbb{E} \left[\sum_{k=1}^d \int_{\Omega} \left(\partial_k f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,k} \sigma'(W_i \cdot x + B_i) \right)^2 dx \right] \\
&\quad + \mathbb{E} \left[\sum_{s=1}^d \sum_{t=1}^d \int_{\Omega} \left(\partial_{s,t} f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,s} W_{i,t} \sigma''(W_i \cdot x + B_i) \right)^2 dx \right].
\end{aligned}$$

For the first term, applying Fubini's theorem yields that

$$\begin{aligned}
& \mathbb{E} \left[\int_{\Omega} \left(f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right)^2 dx \right] \\
&= \int_{\Omega} \mathbb{E} \left[\left(f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right)^2 \right] dx \\
&= \int_{\Omega} \frac{1}{m} \text{Var}(A\sigma(W \cdot x + B)) dx \\
&\leq \frac{1}{m} \int_{\Omega} P(A\sigma(W \cdot x + B))^2 dx \\
&= \frac{1}{m} \int_{\Omega} \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} \left(\frac{1}{2\pi \hat{\sigma}(1)} \frac{\sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega) p_2(b)} \right)^2 p_1(\omega) p_2(b) db d\omega dx \\
&\lesssim \frac{1}{m} C_d M^{d+1} \int_{\{|\omega| \leq M\}} (1 + |\omega|_1) |\hat{f}(\omega)|^2 d\omega \\
&\lesssim \frac{1}{m} C_d M^{d+1} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^1(\Omega)},
\end{aligned}$$

where the second inequality follows from the polynomial decay condition of σ .

Similarly, for the second term, we have

$$\begin{aligned}
& \mathbb{E} \left[\int_{\Omega} \left(\partial_k f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,k} \sigma'(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \leq \frac{1}{m} \int_{\Omega} \text{Var}(A W_k \sigma'(W \cdot x + B)) dx \\
& \leq \frac{1}{m} \int_{\Omega} P(A W_k \sigma'(W \cdot x + B))^2 dx \\
& = \frac{1}{m} \int_{\Omega} \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} \left(\frac{1}{2\pi \hat{\sigma}(1)} \frac{\omega_k \sigma'(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega) p_2(b)} \right)^2 p_1(\omega) p_2(b) db d\omega dx \\
& \lesssim \frac{1}{m} \int_{\Omega} \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} C_d M^{d+1} \omega_k^2 |\sigma'(\omega \cdot x + b)|^2 |\hat{f}(\omega)|^2 db d\omega dx \\
& \lesssim \frac{1}{m} C_d M^{d+1} \int_{\{|\omega|_1 \leq M\}} \omega_k^2 (1 + |\omega|_1) |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

Summing over k yields that

$$\begin{aligned}
& \mathbb{E} \left[\sum_{k=1}^d \int_{\Omega} \left(\partial_k f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,k} \sigma'(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \lesssim \frac{1}{m} C_d M^{d+1} \int_{\{|\omega|_1 \leq M\}} (1 + |\omega|_1)^2 (1 + |\omega|_1) |\hat{f}(\omega)|^2 d\omega \\
& \lesssim \frac{1}{m} C_d M^{d+1} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^3(\Omega)}.
\end{aligned}$$

And for the third term, we have

$$\begin{aligned}
& \mathbb{E} \left[\int_{\Omega} \left(\partial_{s,t} f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,s} W_{i,t} \sigma''(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \lesssim \frac{1}{m} \int_{\Omega} P(A W_s W_t \sigma''(W \cdot x + B))^2 dx \\
& \lesssim \frac{1}{m} C_d M^{d+1} \int_{\Omega} \int_{\{|\omega|_1 \leq M\}} \int_{\{|b| \leq 2M\}} \omega_s^2 \omega_t^2 |\sigma''(\omega \cdot x + b)|^2 |\hat{f}(\omega)|^2 db d\omega dx \\
& \lesssim \frac{1}{m} C_d M^{d+1} \int_{\{|\omega|_1 \leq M\}} \omega_s^2 \omega_t^2 (1 + |\omega|_1)^2 |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

Summing over i, j yields that

$$\begin{aligned}
& \mathbb{E} \left[\sum_{s=1}^d \sum_{t=1}^d \int_{\Omega} \left(\partial_{s,t} f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,s} W_{i,t} \sigma''(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \lesssim \frac{1}{m} C_d M^{d+1} \int_{\{|\omega|_1 \leq M\}} |\omega|^4 (1 + |\omega|_1)^2 |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

For $3 < s \leq 6$,

$$\int_{\{|\omega|_1 \leq M\}} |\omega|^4 (1 + |\omega|_1)^2 |\hat{f}(\omega)|^2 d\omega \leq M^{6-s} \int_{\{|\omega|_1 \leq M\}} (1 + |\omega|_1)^s |\hat{f}(\omega)|^2 d\omega \leq M^{6-s} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^s(\Omega)};$$

for $s > 6$,

$$\int_{\{|\omega|_1 \leq M\}} |\omega|^4 (1 + |\omega|_1)^2 |\hat{f}(\omega)|^2 d\omega \leq \int_{\{|\omega|_1 \leq M\}} (1 + |\omega|_1)^6 |\hat{f}(\omega)|^2 d\omega \leq \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^s(\Omega)}.$$

Therefore,

$$\mathbb{E} \left[\|f_1(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \begin{cases} \frac{1}{m} M^{d+7-s} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^s(\Omega)}, & 3 < s < 6, \\ \frac{1}{m} M^{d+1} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^s(\Omega)}, & s \geq 6. \end{cases}$$

Combining the estimation of the $H^2(\Omega)$ norm of $f_2(x)$, we have

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \begin{cases} \frac{1}{m} M^{d+7-s} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^s(\Omega)} + \left(\frac{1}{M^{s-3}} \|f\|_{B^s(\Omega)} \right)^2, & 3 < s < 6, \\ \frac{1}{m} M^{d+1} \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} \|f\|_{B^s(\Omega)} + \left(\frac{1}{M^{s-3}} \|f\|_{B^s(\Omega)} \right)^2, & s \geq 6. \end{cases}$$

Moreover, as

$$A_i = \frac{1}{2\pi\hat{\sigma}(1)} \frac{|\hat{f}(W_i)| \cos(\theta(W_i) - B_i)}{p_1(W_i)p_2(B_i)}, \quad 1 \leq i \leq m,$$

we can deduce that

$$|A_i| \lesssim \|\hat{f}\|_{L^\infty(\mathbb{R}^d)} M^{d+1}.$$

3.2. Proof of Theorem 2. Recall that the loss function and its empirical part are

$$F(a) = L(a) + \lambda|a|^2 = \int_{\Omega} (a \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a \cdot f_2(y) + g_2(y))^2 dy + \lambda|a|^2$$

and

$$F_n(a) = \frac{1}{n} \sum_{i=1}^n l(a; x_i, y_i) + \lambda|a|^2$$

respectively, where

$$l(a; x_i, y_i) = |\Omega| (a \cdot f_1(x_i) + g_1(x_i))^2 + |\partial\Omega| (a \cdot f_2(y_i) + g_2(y_i))^2.$$

Let ρ_0 be a constant to be determined. Construct a sequence $\rho_k = 2\rho_{k-1}$ for $k \in \mathbb{N}$ and then decompose \mathcal{A} into a sequence of disjoint sets as follows.

$$\mathcal{A}_k := \{a \in \mathcal{A} : \rho_{k-1} < F(a) - F(a^*) \leq \rho_k\},$$

where $a^* = \arg \min_{a \in \mathcal{A}} F(a)$. For brevity, we let $\rho_{-1} = 0$.

Recall that the i -th component of \mathbb{R}^m -valued functions $f_1(x)$ and $f_2(y)$ are

$$f_1^i(x) = -|\omega_i|^2 \sigma''(\omega_i \cdot x + b_i) + V \sigma(\omega_i \cdot x + b_i)$$

and $f_2^i(y) = \sigma(\omega_i \cdot y + b_i)$ respectively.

Therefore,

$$|F(a) - F(a^*)| \lesssim (\lambda + mM^4) |a - a^*| \lesssim \lambda + mM^4,$$

which implies

$$K := \max k \leq C \log\left(\frac{\lambda + mM^4}{\rho_0}\right),$$

as $\rho_K = 2^K \rho_0 = \mathcal{O}(\lambda + mM^4)$. For any constant $\delta \in (0, 1)$, set $\delta_k = \frac{\delta}{K+1}$.

By the fact that F is 2λ -strongly convex, we have $\lambda|a - a^*|^2 \leq F(a) - F(a^*)$. Thus, for any $a \in \mathcal{A}_k$, we can deduce that

$$|a - a^*|^2 \leq \frac{1}{\lambda}(F(a) - F(a^*)) \leq \frac{1}{\lambda}\rho_k.$$

Let $s(a; x, y) = l(a; x, y) - l(a^*; x, y)$, so that $F(a) - F_n(a) = (P - P_n)s(a; x, y)$.

We first derive the upper bound for $|s(a; x, y)|$ in \mathcal{A}_k .

$$\begin{aligned} & |s(a; x, y)| \\ &= \left| |\Omega|(a \cdot f_1(x) + g_1(x))^2 + |\partial\Omega|(a \cdot f_2(y) + g_2(y))^2 - |\Omega|(a^* \cdot f_1(x) + g_1(x))^2 - |\partial\Omega|(a^* \cdot f_2(y) + g_2(y))^2 \right| \\ &\lesssim \left| ((a - a^*) \cdot f_1(x))(a \cdot f_1(x) + g_1(x) + a^* \cdot f_1(x) + g_1(x)) \right| \\ &\quad + \left| ((a - a^*) \cdot f_2(y))(a \cdot f_2(y) + g_2(y) + a^* \cdot f_2(y) + g_2(y)) \right| \\ &\leq |a - a^*| |f_1(x)| (|a| |f_1(x)| + |a^*| |f_1(x)| + 2|g_1(x)|) \\ &\quad + |a - a^*| |f_2(y)| (|a| |f_2(y)| + |a^*| |f_2(y)| + 2|g_2(y)|) \\ &\lesssim \left(\sqrt{\frac{\rho_k}{\lambda}} \sqrt{m} M^2 \right) (\sqrt{m} M^2) \\ &= m \sqrt{\frac{\rho_k}{\lambda}} M^4, \end{aligned}$$

where the second inequality follows from that $|a|, |a^*| \leq C$ and $|f_1(x)|, |f_2(y)| \lesssim \sqrt{m} M^2$.

From the Hoeffding's inequality, we can deduce that with probability at least $1 - \delta_k$

$$\sup_{a \in \mathcal{A}_k} ((P - P_n)s(a; x, y)) \lesssim \mathbb{E}_{x, y} \sup_{a \in \mathcal{A}_k} ((P - P_n)s(a; x, y)) + mM^4 \sqrt{\frac{\rho_k}{\lambda}} \sqrt{\frac{\log(\frac{1}{\delta_k})}{n}}.$$

By the technique of symmetrization, we have

$$\begin{aligned} \mathbb{E}_{x, y} \sup_{a \in \mathcal{A}_k} ((P - P_n)s(a; x, y)) &\leq 2 \mathbb{E}_{z, \epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} \sum_{i=1}^n \epsilon_i (l(a; x_i, y_i) - l(a^*; x_i, y_i)) \\ &\lesssim \mathbb{E}_{x, \epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} \sum_{i=1}^n \epsilon_i ((a \cdot f_1(x_i) + g_1(x_i))^2 - (a^* \cdot f_1(x_i) + g_1(x_i))^2) \\ &\quad + \mathbb{E}_{y, \epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} \sum_{i=1}^n \epsilon_i ((a \cdot f_2(y_i) + g_2(y_i))^2 - (a^* \cdot f_2(y_i) + g_2(y_i))^2). \end{aligned}$$

Then from the contraction property of Rademacher complexity, we can deduce that

$$\begin{aligned}
& \mathbb{E}_{x,\epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} \sum_{i=1}^n \epsilon_i ((a \cdot f_1(x_i) + g_1(x_i))^2 - (a^* \cdot f_1(x_i) + g_1(x_i))^2) \\
& \lesssim \sqrt{m} M^2 \mathbb{E}_{x,\epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} \sum_{i=1}^n \epsilon_i (a - a^*) \cdot f_1(x_i) \\
& = \sqrt{m} M^2 \mathbb{E}_{x,\epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} (a - a^*) \cdot \left(\sum_{i=1}^n \epsilon_i f_1(x_i) \right) \\
& \leq \sqrt{m} M^2 \mathbb{E}_{x,\epsilon} \sup_{a \in \mathcal{A}_k} \frac{1}{n} |a - a^*| \left| \left(\sum_{i=1}^n \epsilon_i f_1(x_i) \right) \right| \\
& \lesssim \sqrt{m} M^2 \frac{1}{n} \sqrt{\frac{\rho_k}{\lambda}} \sqrt{n} \sqrt{m} M^2 \\
& = \frac{m M^4}{\sqrt{n}} \sqrt{\frac{\rho_k}{\lambda}}.
\end{aligned}$$

Combining above results together, we derive the following inequality with probability at least $1 - \delta_k$

$$\sup_{a \in \mathcal{A}_k} ((P - P_n)s(a; x, y)) \lesssim \frac{m M^4}{\sqrt{n}} \sqrt{\frac{\rho_k}{\lambda}} + m M^4 \sqrt{\frac{\rho_k}{\lambda}} \sqrt{\frac{\log(\frac{1}{\delta_k})}{n}}.$$

Note that for any $a \in \mathcal{A}_k$, we have

$$\frac{\rho_k}{2} = \rho_{k-1} \leq F(a) - F(a^*),$$

thus

$$\rho_k \leq \max\{\rho_0, 2(F(a) - F(a^*))\}.$$

For δ_k , it's obvious that

$$\frac{1}{\delta_k} = \frac{K+1}{\delta} \leq C \frac{1}{\delta} \log\left(\frac{\lambda + m M^4}{\rho_0}\right).$$

With the fact that

$$(F(a) - F(a^*)) - (F_n(a) - F_n(a^*)) = (P - P_n)s(a; x, y),$$

we have that with probability at least $1 - \delta_k$ for all $a \in \mathcal{A}_k$

$$F(a) - F(a^*) \leq F_n(a) - F_n(a^*) + C m M^4 \sqrt{\frac{\max\{\rho_0, 2(F(a) - F(a^*))\}}{n \lambda}} \left(1 + \sqrt{\log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + m M^4}{\rho_0}\right)}\right).$$

Since $\sum_{k=1}^K \delta_k = \delta$, the above inequality holds with probability at least $1 - \delta$ uniformly for all $a \in \mathcal{A}$.

By setting $\rho_0 = 1/n$, we can deduce that with probability at least $1 - \delta$, $F(a_T) - F(a^*) \leq \frac{\rho_0}{2} = \frac{1}{2n}$ or

$$F(a_T) - F(a^*) \leq F_n(a_T) - F_n(a^*) + C m M^4 \sqrt{\frac{2(F(a_T) - F(a^*))}{n \lambda}} \left(1 + \sqrt{\log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + m M^4}{\rho_0}\right)}\right).$$

Note that F_n is 2λ -strongly convex, $C(\lambda + mM^4)$ -Lipschitz continuous and $C(\lambda + mM^4)$ -smooth, thus Theorem 3.10 in [8] implies that the optimization error of projected gradient descent method with $\nu = \frac{1}{C(\lambda + mM^4)}$ is

$$\Delta_T := F_n(a_T) - \min_{a \in \mathcal{A}} F_n(a) \lesssim \frac{L^3}{\lambda^2} e^{-\frac{\lambda T}{L}},$$

where $L = C(\lambda + mM^4)$.

Therefore,

$$\begin{aligned} & F(a_T) - F(a^*) \\ & \leq F_n(a_T) - \min_{a \in \mathcal{A}} F_n(a) + \min_{a \in \mathcal{A}} F_n(a) - F_n(a^*) \\ & \quad + CmM^4 \sqrt{\frac{2(F(a_T) - F(a^*))}{n\lambda}} \left(1 + \sqrt{\log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + mM^4}{\rho_0}\right)}\right) \\ & \leq \Delta_T + CmM^4 \sqrt{\frac{2(F(a_T) - F(a^*))}{n\lambda}} \left(1 + \sqrt{\log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + mM^4}{\rho_0}\right)}\right) \\ & \leq \Delta_T + \frac{F(a_T) - F(a^*)}{2} + \frac{m^2 M^8}{n\lambda} \left(1 + \log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + mM^4}{\rho_0}\right)\right), \end{aligned}$$

where the last inequality follows from the mean inequality.

Through a simple algebraic transformation, we have

$$\begin{aligned} F(a_T) - F(a^*) & \leq C \left(\frac{(\lambda + mM^4)^3}{\lambda^2} e^{-\frac{\lambda T}{C(\lambda + mM^4)}} + \frac{m^2 M^8}{n\lambda} \left(1 + \log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + mM^4}{\rho_0}\right)\right) \right) \\ & = C \left(\frac{L^3}{\lambda^2} e^{-\frac{\lambda T}{L}} + \frac{m^{2+\frac{4}{d+1}}}{n\lambda} \left(1 + \log\left(\frac{1}{\delta}\right) + \log \log\left(\frac{\lambda + m^{1+\frac{2}{d+1}}}{\rho_0}\right)\right) \right). \end{aligned}$$

Integrating the tail term with respect to X, Y yields that

$$\mathbb{E}_{X,Y}[F(a_T) - F(a^*)] \lesssim \frac{L^3}{\lambda^2} e^{-\frac{\lambda T}{L}} + \frac{m^{2+\frac{4}{d+1}}}{n\lambda} \left(1 + \log \log\left(\frac{\lambda + m^{1+\frac{2}{d+1}}}{\rho_0}\right)\right).$$

On the other hand, as $u \in \mathcal{B}^s(\Omega)$, Theorem 1 implies that there is a random neural network u_m of the form

$$u_m(x) = \sum_{i=1}^m \bar{a}_i \sigma(W_i \cdot x + B_i)$$

such that

$$\mathbb{E}_{W,B} \left[\|u(x) - u_m(x)\|_{H^2(\Omega)}^2 \right] \leq \frac{C}{\sqrt{m}}$$

with $\bar{a} = (\bar{a}_1, \dots, \bar{a}_m)$, $|\bar{a}| \leq C$, i.e., $\bar{a} \in \mathcal{A}$.

Therefore,

$$\begin{aligned}
F(a^*) &= \int_{\Omega} (a^* \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a^* \cdot f_2(y) + g_2(y))^2 dy + \lambda |a^*|^2 \\
&\leq \int_{\Omega} (\bar{a} \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (\bar{a} \cdot f_2(y) + g_2(y))^2 dy + \lambda |\bar{a}| \\
&\lesssim \|u(x) - u_m(x)\|_{H^2(\Omega)}^2 + \lambda.
\end{aligned}$$

Finally, we have

$$\begin{aligned}
&\mathbb{E}_{W,B} \mathbb{E}_{X,Y} \left[\int_{\Omega} (a_T \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a_T \cdot f_2(y) + g_2(y))^2 dy \right] \\
&\lesssim \frac{1}{\sqrt{m}} + \lambda + \frac{L^3}{\lambda^2} e^{-\frac{\lambda T}{L}} + \frac{m^{2+\frac{4}{d+1}}}{\lambda n} (1 + \log \log(\frac{\lambda + m^{1+\frac{2}{d+1}}}{\rho_0})) \\
&\lesssim \frac{1}{\sqrt{m}} + \lambda + \frac{m^{3+\frac{6}{d+1}}}{\lambda^2} e^{-\frac{\lambda T}{m^{2+4/(d+1)}}} + \frac{m^{1+\frac{2}{d+1}}}{\lambda n} \log \log(mn).
\end{aligned}$$

3.3. Proof of Theorem 3. As $f_e|_{\Omega} = f$, we write f for f_e for simplicity. Thus,

$$\begin{aligned}
f(x) &= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) db d\omega \\
&= \int_{\mathbb{R}^d} \int_{\mathbb{R}} \frac{1}{2\pi\hat{\sigma}(1)} \sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b) \frac{(1+|\omega|)^{\alpha}(1+|b|)^{\beta}}{C_{\alpha}C_{\beta}} \frac{C_{\alpha}C_{\beta}}{(1+|\omega|)^{\alpha}(1+|b|)^{\beta}} db d\omega \\
&= \mathbb{E}_{P_1(\omega), P_2(b)} \left[\frac{1}{2\pi\hat{\sigma}(1)} \frac{\sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega)p_2(b)} \right],
\end{aligned}$$

where $P_1(\omega) = p_1(\omega)d\omega$, $P_2(b) = p_2(b)db$ and $p_1(\omega) := \frac{C_{\alpha}}{(1+|\omega|)^{\alpha}}$, $p_2(b) := \frac{C_{\beta}}{(1+|b|)^{\beta}}$ are density functions in \mathbb{R}^d and \mathbb{R} respectively.

Taking $W \sim P_1$, $B \sim P_2$ and let

$$A = \frac{1}{2\pi\hat{\sigma}(1)} \frac{|\hat{f}(W)| \cos(\theta(W) - B)}{p_1(W)p_2(B)} I_{\{|B| \leq Cd(1+|W|)\}},$$

then

$$f(x) = \mathbb{E}_{P_1(W), P_2(B)} [A\sigma(W \cdot x + B)],$$

as σ is compactly supported in $[-2, 2]$.

Therefore, we can construct a random neural network f_m as follows.

$$f_m(x) = \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i),$$

where $W_1, \dots, W_m \sim P_1$ and $B_1, \dots, B_m \sim P_2$,

$$A_i = \frac{1}{2\pi\hat{\sigma}(1)} \frac{|\hat{f}(W_i)| \cos(\theta(W_i) - B_i)}{p_1(W_i)p_2(B_i)} I_{\{|B_i| \leq Cd(1+|W_i|)\}}, 1 \leq i \leq m.$$

Next, we will estimate $\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right]$.

$$\begin{aligned}
& \mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \\
&= \mathbb{E} \left[\|f(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i)\|_{H^2(\Omega)}^2 \right] \\
&= \mathbb{E} \left[\int_{\Omega} \left(f(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right)^2 dx \right] \\
&\quad + \mathbb{E} \left[\sum_{k=1}^d \int_{\Omega} \left(\partial_k f(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,k} \sigma'(W_i \cdot x + B_i) \right)^2 dx \right] \\
&\quad + \mathbb{E} \left[\sum_{s=1}^d \sum_{t=1}^d \int_{\Omega} \left(\partial_{s,t} f(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,s} W_{i,t} \sigma''(W_i \cdot x + B_i) \right)^2 dx \right].
\end{aligned}$$

For the first term, we have

$$\begin{aligned}
& \mathbb{E} \left[\int_{\Omega} \left(f(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right)^2 dx \right] \\
&= \int_{\Omega} \mathbb{E} \left[\left(f(x) - \frac{1}{m} \sum_{i=1}^m A_i \sigma(W_i \cdot x + B_i) \right)^2 \right] dx \\
&= \int_{\Omega} \frac{1}{m} \text{Var}(A\sigma(W \cdot x + B)) dx \\
&\leq \frac{1}{m} \int_{\Omega} P(A\sigma(W \cdot x + B))^2 dx \\
&= \frac{1}{m} \int_{\Omega} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \left(\frac{1}{2\pi\hat{\sigma}(1)} \frac{\sigma(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega) p_2(b)} \right)^2 p_1(\omega) p_2(b) db d\omega dx \\
&\lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} \int_{\mathbb{R}^d} \int_{\mathbb{R}} (1 + |\omega|)^{\alpha} (1 + |b|)^{\beta} |\sigma(\omega \cdot x + b)|^2 |\hat{f}(\omega)|^2 db d\omega \\
&\lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} d^{\beta} \int_{\mathbb{R}^d} \int_{\mathbb{R}} (1 + |\omega|)^{\alpha+\beta} |\sigma(\omega \cdot x + b)|^2 |\hat{f}(\omega)|^2 db d\omega \\
&\lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} \int_{\mathbb{R}^d} d^{\beta+1} (1 + |\omega|)^{\alpha+\beta+1} |\hat{f}(\omega)|^2 d\omega,
\end{aligned}$$

where the third inequality follows from the fact σ is compactly supported in $[-2, 2]$ thus $|b| \leq Cd(1 + |\omega|)$ and the last inequality follows from the polynomial decay condition of σ .

Similarly, for the second term, we have

$$\begin{aligned}
& \mathbb{E} \left[\int_{\Omega} \left(\partial_k f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,k} \sigma'(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \leq \frac{1}{m} \int_{\Omega} \text{Var}(A W_k \sigma'(W \cdot x + B)) dx \\
& \leq \frac{1}{m} \int_{\Omega} P(A W_k \sigma'(W \cdot x + B))^2 dx \\
& = \frac{1}{m} \int_{\Omega} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \left(\frac{1}{2\pi \hat{\sigma}(1)} \frac{\omega_k \sigma'(\omega \cdot x + b) |\hat{f}(\omega)| \cos(\theta(\omega) - b)}{p_1(\omega) p_2(b)} \right)^2 p_1(\omega) p_2(b) db d\omega dx \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \omega_k^2 (1 + |\omega|)^{\alpha} (1 + |b|)^{\beta} |\sigma(\omega \cdot x + b)|^2 |\hat{f}(\omega)|^2 db d\omega \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} d^{\beta+1} \int_{\mathbb{R}^d} \omega_k^2 (1 + |\omega|)^{\alpha+\beta+1} |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

Summing over k yields that

$$\begin{aligned}
& \mathbb{E} \left[\sum_{k=1}^d \int_{\Omega} \left(\partial_k f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,k} \sigma'(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} d^{\beta+1} \int_{\mathbb{R}^d} (1 + |\omega|)^2 (1 + |\omega|)^{\alpha+\beta+1} |\hat{f}(\omega)|^2 d\omega \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} d^{\beta+1} \int_{\mathbb{R}^d} (1 + |\omega|)^{\alpha+\beta+3} |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

For the third term, we have

$$\begin{aligned}
& \mathbb{E} \left[\int_{\Omega} \left(\partial_{s,t} f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,s} W_{i,t} \sigma''(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \lesssim \frac{1}{m} \int_{\Omega} P(A W_s W_t \sigma''(W \cdot x + B))^2 dx \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} \int_{\Omega} \int_{\mathbb{R}^d} \int_{\mathbb{R}} \omega_s^2 \omega_t^2 (1 + |\omega|)^{\alpha} (1 + |b|)^{\beta} |\sigma''(\omega \cdot x + b)|^2 |\hat{f}(\omega)|^2 db d\omega dx \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} d^{\beta+1} \int_{\mathbb{R}^d} \omega_s^2 \omega_t^2 (1 + |\omega|)^{\alpha+\beta+1} |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

Summing over s, t gives

$$\begin{aligned}
& \mathbb{E} \left[\sum_{s=1}^d \sum_{t=1}^d \int_{\Omega} \left(\partial_{s,t} f_1(x) - \frac{1}{m} \sum_{i=1}^m A_i W_{i,s} W_{i,t} \sigma''(W_i \cdot x + B_i) \right)^2 dx \right] \\
& \lesssim \frac{1}{m} \frac{1}{C_{\alpha} C_{\beta}} d^{\beta+1} \int_{\mathbb{R}^d} (1 + |\omega|)^{\alpha+\beta+5} |\hat{f}(\omega)|^2 d\omega.
\end{aligned}$$

Therefore, when

$$\int_{\mathbb{R}^d} (1 + |\omega|)^{\alpha+\beta+5} |\hat{f}(\omega)|^2 d\omega < \infty,$$

combining above results yields that

$$\mathbb{E} \left[\|f(x) - f_m(x)\|_{H^2(\Omega)}^2 \right] \lesssim \frac{1}{m} \frac{d^{\beta+1}}{C_\alpha}.$$

3.4. Proof of Theorem 4.

By the definition of a_n , we have $F_n(a_n) \leq F_n(0)$, thus $\lambda|a_n|^2 \leq C$, i.e., $|a_n| \leq \sqrt{\frac{C}{\lambda}}$. Recall that

$$Pl(a; x, y) = \int_{\Omega} (a \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (a \cdot f_2(y) + g_2(y))^2 dy.$$

Therefore,

$$\begin{aligned} \mathbb{E}_{X,Y}[Pl(a_n; x, y)] &= \mathbb{E}_{X,Y}[(P - P_n)l(a_n; x, y) + P_n l(a_n; x, y)] \\ &\leq \mathbb{E}_{X,Y} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} (P - P_n)l(a; x, y) + P_n l(a_n; x, y) \right] \\ &\leq \mathbb{E}_{X,Y} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} (P - P_n)l(a; x, y) + P_n l(a_n; x, y) + \lambda|a_n|^2 \right] \\ &\leq \mathbb{E}_{X,Y} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} (P - P_n)l(a; x, y) + P_n l(\bar{a}; x, y) + \lambda|\bar{a}|^2 \right] \\ &= \mathbb{E}_{X,Y} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} (P - P_n)l(a; x, y) \right] + Pl(\bar{a}; x, y) + \lambda|\bar{a}|^2 \\ &= \mathbb{E}_{X,Y} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} (P - P_n)l(a; x, y) \right] + F(\bar{a}), \end{aligned}$$

where \bar{a} is the coefficient of the random neural network that provides the best approximation in Theorem 3.

By the technique of symmetrization, we have

$$\begin{aligned} &\mathbb{E}_{X,Y} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} (P - P_n)l(a; x, y) \right] \\ &\lesssim \mathbb{E}_{X,\epsilon} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} \frac{1}{n} \sum_{i=1}^n \epsilon_i (a \cdot f_1(x_i) + g_1(x_i))^2 \right] + \mathbb{E}_{Y,\epsilon} \left[\sup_{|a| \leq \sqrt{\frac{C}{\lambda}}} \frac{1}{n} \sum_{i=1}^n \epsilon_i (a \cdot f_2(y_i) + g_2(y_i))^2 \right]. \end{aligned}$$

It suffices to bound the Rademacher complexity.

$$\begin{aligned}
& \mathbb{E}_X \mathbb{E}_\epsilon \left[\sup_{|a| \leq \sqrt{\frac{c}{\lambda}}} \frac{1}{n} \sum_{i=1}^n \epsilon_i (a \cdot f_1(x_i) + g_1(x_i))^2 \right] \\
& \lesssim \mathbb{E}_X \mathbb{E}_\epsilon \left[\sup_{|a| \leq \sqrt{\frac{c}{\lambda}}} \frac{1}{n} \sum_{i=1}^n \epsilon_i (a \cdot f_1(x_i) + g_1(x_i)) \frac{1}{\sqrt{\lambda}} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} \right] \\
& = \frac{1}{n} \frac{1}{\sqrt{\lambda}} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} \mathbb{E}_X \mathbb{E}_\epsilon \left[\sup_{|a| \leq \sqrt{\frac{c}{\lambda}}} \frac{1}{n} a \cdot \left(\sum_{i=1}^n \epsilon_i f_1(x_i) \right) \right] \\
& \leq \frac{1}{n} \frac{1}{\sqrt{\lambda}} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} \mathbb{E}_X \mathbb{E}_\epsilon \left[\frac{1}{\sqrt{\lambda}} \left| \sum_{i=1}^n \epsilon_i f_1(x_i) \right| \right] \\
& \leq \frac{1}{n} \frac{1}{\lambda} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} \mathbb{E}_X (\mathbb{E}_\epsilon \left| \sum_{i=1}^n \epsilon_i f_1(x_i) \right|^2)^{1/2} \\
& = \frac{1}{n} \frac{1}{\lambda} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} \mathbb{E}_X \left(\sum_{i=1}^n |f_1(x_i)|^2 \right)^{1/2} \\
& \lesssim \frac{1}{n} \frac{1}{\lambda} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} \left(n \sum_{j=1}^m (|W_j|^2 + 1) \right)^{1/2} \\
& = \frac{1}{\lambda \sqrt{n}} \sum_{j=1}^m (|W_j|^2 + 1),
\end{aligned}$$

where the first inequality follows from the contraction property of Rademacher complexity and the fact that

$$|a \cdot f_1(x_i) + g_1(x_i)| \lesssim \frac{1}{\sqrt{\lambda}} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)} + 1 \lesssim \frac{1}{\sqrt{\lambda}} \sqrt{\sum_{j=1}^m (|W_j|^2 + 1)}, 1 \leq i \leq n,$$

as we choose $\lambda \leq m$.

On the other hand,

$$\begin{aligned}
\mathbb{E}_{W,B}[F(\bar{a})] &= \mathbb{E}_{W,B} \left[\int_{\Omega} (\bar{a} \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega} (\bar{a} \cdot f_2(y) + g_2(y))^2 dy \right] + \lambda \mathbb{E}_{W,B}[|\bar{a}|^2] \\
&\lesssim \mathbb{E}_{W,B}[\|u - u_m\|_{H^2\Omega}^2] + \lambda \mathbb{E}_{W,B}[|\bar{a}|^2] \\
&\lesssim \frac{1}{m} \frac{d^{\beta+1}}{C_\alpha} + \lambda \mathbb{E}_{W,B}[|\bar{a}|^2].
\end{aligned}$$

Recall that

$$\bar{a}_i = \frac{1}{m} \frac{1}{2\pi\hat{\sigma}(1)} \frac{|\hat{f}(W_i)| \cos(\theta(W_i) - B_i)}{p_1(W_i)p_2(B_i)} I_{\{|B_i| \leq Cd(1+|W_i|)\}}, 1 \leq i \leq m.$$

Therefore,

$$\begin{aligned}
\mathbb{E}_{W,B}|\bar{a}_i|^2 &\lesssim \frac{1}{m^2}\mathbb{E}_{W,B}\frac{1}{C_\alpha^2C_\beta^2}|\hat{f}(W_i)|^2(1+|W_i|)^{2\alpha}(1+|B_i|)^{2\beta}I_{\{|B_i|\leq Cd(1+|W_i|)\}} \\
&\lesssim \frac{1}{m^2}\mathbb{E}_{W,B}\frac{d^{2\beta}}{C_\alpha^2C_\beta^2}|\hat{f}(W_i)|^2(1+|W_i|)^{2\alpha}(1+|W_i|)^{2\beta} \\
&= \frac{1}{m^2}\mathbb{E}_W\frac{d^{2\beta}}{C_\alpha^2C_\beta^2}|\hat{f}(W_i)|^2(1+|W_i|)^{2\alpha}(1+|W_i|)^{2\beta} \\
&= \frac{1}{m^2}\frac{d^{2\beta}}{C_\alpha C_\beta^2}\int_{\mathbb{R}^d}|\hat{f}(\omega)|^2(1+|\omega|)^{\alpha+2\beta}d\omega \\
&\leq \frac{1}{m^2}\frac{d^{2\beta}}{C_\alpha C_\beta^2}\int_{\mathbb{R}^d}|\hat{f}(\omega)|^2(1+|\omega|)^{\alpha+\beta+5}d\omega.
\end{aligned}$$

Thus,

$$\mathbb{E}_{W,B}|\bar{a}|^2 \lesssim \frac{1}{m} \frac{d^{2\beta}}{C_\alpha}.$$

For $\mathbb{E}_{W,B}[F(\bar{a})]$, we have

$$\mathbb{E}_{W,B}[F(\bar{a})] \lesssim \frac{1}{m} \frac{d^{\beta+1}}{C_\alpha} + \frac{\lambda}{m} \frac{d^{2\beta}}{C_\alpha}.$$

As $\alpha > d + 2$,

$$\mathbb{E}\left[\sum_{j=1}^m(|W_j|^2 + 1)\right] \lesssim m,$$

which implies

$$\mathbb{E}_{W,B}\mathbb{E}_{X,Y}\left[\sup_{|a|\leq\sqrt{\frac{C}{\lambda}}}(P - P_n)l(a; x, y)\right] \lesssim \frac{1}{\lambda\sqrt{n}}\mathbb{E}_{W,B}\sum_{j=1}^m(|W_j|^2 + 1) \lesssim \frac{m}{\lambda\sqrt{n}}.$$

Combining all results above yields that

$$\mathbb{E}_{W,B}\mathbb{E}_{X,Y}\left[\int_{\Omega}(a_n \cdot f_1(x) + g_1(x))^2 dx + \int_{\partial\Omega}(a_n \cdot f_2(y) + g_2(y))^2 dy\right] \lesssim \frac{m}{\lambda\sqrt{n}} + \frac{1}{m} \frac{d^{2\beta}}{C_\alpha} + \frac{\lambda}{m} \frac{d^{2\beta}}{C_\alpha}.$$

4. CONCLUSION

In this draft, we make the priori distribution of random neural networks clear for approximating Barron type functions. We have considered two types of sampling distributions, one with bounded support and the other with unbounded support. In both settings, the approximation, optimization and generalization errors have been derived.

REFERENCES

- [1] M. Raissi, P. Perdikaris, and G. E. Karniadakis, ‘‘Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations,’’ *Journal of Computational physics*, vol. 378, pp. 686–707, 2019.
- [2] B. Yu *et al.*, ‘‘The deep ritz method: a deep learning-based numerical algorithm for solving variational problems,’’ *Communications in Mathematics and Statistics*, vol. 6, no. 1, pp. 1–12, 2018.

- [3] A. Krishnapriyan, A. Gholami, S. Zhe, R. Kirby, and M. W. Mahoney, “Characterizing possible failure modes in physics-informed neural networks,” *Advances in Neural Information Processing Systems*, vol. 34, pp. 26 548–26 560, 2021.
- [4] G.-B. Huang, Q.-Y. Zhu, and C.-K. Siew, “Extreme learning machine: theory and applications,” *Neurocomputing*, vol. 70, no. 1-3, pp. 489–501, 2006.
- [5] J. W. Siegel and J. Xu, “Approximation rates for neural networks with general activation functions,” *Neural Networks*, vol. 128, pp. 313–321, 2020.
- [6] S. Agmon, A. Douglis, and L. Nirenberg, “Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. i,” *Communications on pure and applied mathematics*, vol. 12, no. 4, pp. 623–727, 1959.
- [7] Y. Klochkov and N. Zhivotovskiy, “Stability and deviation optimal risk bounds with convergence rate $o(1/n)$,” *Advances in Neural Information Processing Systems*, vol. 34, pp. 5065–5076, 2021.
- [8] S. Bubeck *et al.*, “Convex optimization: Algorithms and complexity,” *Foundations and Trends[®] in Machine Learning*, vol. 8, no. 3-4, pp. 231–357, 2015.