

OPTIMAL CONTROL OF SEMILINEAR ELLIPTIC PARTIAL DIFFERENTIAL EQUATIONS WITH NON-LIPSCHITZIAN NONLINEARITIES

CONSTANTIN CHRISTOF*

CIT, Department of Mathematics, Technische Universität München,
Boltzmannstr. 3, 85748 Garching b. München, Germany

(Communicated by the associate editor name)

ABSTRACT. We study optimal control problems that are governed by semilinear elliptic partial differential equations that involve non-Lipschitzian nonlinearities. It is shown that, for a certain class of such PDEs, the solution map is Fréchet differentiable even though the differential operator contains a nondifferentiable term. We exploit this effect to establish first-order necessary optimality conditions for minimizers of the considered control problems. The resulting KKT-conditions take the form of coupled PDE-systems that are posed in non-Muckenhoupt weighted Sobolev spaces and raise interesting questions regarding the regularity of optimal controls, the derivation of second-order optimality conditions, and the analysis of finite element discretizations.

1. Introduction. This paper is concerned with optimal control problems of the following type:

$$\left. \begin{aligned} &\text{Minimize } J(y, u) \\ &\text{w.r.t. } y \in H_0^1(\Omega), \quad u \in L^2(\Omega), \\ &\text{s.t. } -\Delta y + \text{sgn}(y)|y|^\alpha = u \text{ in } \Omega, \quad y = 0 \text{ on } \partial\Omega, \\ &\text{and } u \in U_{\text{ad}}. \end{aligned} \right\} \quad (\mathbf{P})$$

Here, $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$, is a nonempty open bounded set with boundary $\partial\Omega$; the spaces $H_0^1(\Omega)$ and $L^2(\Omega)$ are defined as usual (see [1]); $J: H_0^1(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$ is a Fréchet differentiable objective function; Δ denotes the Laplace operator; $\text{sgn}: \mathbb{R} \rightarrow \mathbb{R}$ denotes the signum function; $\alpha \in (0, 1)$ is a given number; U_{ad} is a nonempty convex subset of $L^2(\Omega)$; and the governing partial differential equation (PDE) is understood in the weak sense. Note that the salient feature of **(P)** is that the state equation contains a Nemytskii operator that is neither differentiable nor Lipschitz continuous. The main purpose of this paper is to point out that **(P)** nevertheless possesses a Fréchet differentiable control-to-state map $S: L^2(\Omega) \rightarrow H_0^1(\Omega)$, $u \mapsto y$, and, thus, allows for the derivation of first-order necessary optimality conditions in qualified form. For the main results of our analysis, we refer the reader to [Theorem 4.8](#), [Corollary 4.9](#), and [Theorem 5.3](#).

2010 *Mathematics Subject Classification.* 49J20, 49K20, 49K40.

Key words and phrases. optimal control, nonsmooth optimization, optimality condition, KKT-system, non-Lipschitzian nonlinearity, semilinear partial differential equation, porous media flow.

*Corresponding author: Constantin Christof.

Before we begin with our study of the problem (P), let us give some background: Semilinear PDEs involving non-Lipschitzian nonlinearities of the type $\operatorname{sgn}(y)|y|^\alpha$, $\alpha \in (0, 1)$, arise, for instance, when modeling chemical reactions in porous media and processes in desalination plants; see [2, 4, 11, 13] and the references therein. In these fields, the exponent α is also known as the order of the reaction process [11], and the semilinearity $\operatorname{sgn}(y)|y|^\alpha$ as the *Freundlich isotherm* [4]. A main feature of the nonlinearity $\operatorname{sgn}(y)|y|^\alpha$, $\alpha \in (0, 1)$, is that it promotes the formation of so-called *dead zones* within Ω where the PDE-solution y vanishes identically [6, 11]. The price that one pays for this effect is that the term $\operatorname{sgn}(y)|y|^\alpha$ is neither Lipschitz continuous nor differentiable and, thus, induces a form of nonsmoothness that is often hard to handle analytically. The latter is in particular true when first-order necessary optimality conditions for optimal control problems like (P) are considered as these classically require at least some form of (directional) differentiability for the PDE-constraint [14]. Because of these difficulties, optimal control and inverse problems governed by partial differential equations involving non-Lipschitzian nonlinearities are rarely addressed in the literature. One of the few contributions on this topic is [11] which studies the existence of optimal controls for minimization problems that are governed by PDEs similar to that in (P). Compare also with [3] in this context where general regularization techniques are considered.

The main goal of the present paper is to point out that problems of the type (P) are amenable to classical techniques from optimal control theory even in the presence of the nonsmooth term $\operatorname{sgn}(y)|y|^\alpha$. The key observation is that, although nonsmooth and non-Lipschitzian, the state equation of (P) possesses a Fréchet differentiable solution map $S: L^2(\Omega) \rightarrow H_0^1(\Omega)$, $u \mapsto y$; see Theorem 4.8 and Corollary 4.9. This causes the reduced objective function $L^2(\Omega) \ni u \mapsto J(S(u), u) \in \mathbb{R}$ of (P) to be Fréchet differentiable as well, makes it possible to formulate a standard Bouligand stationarity condition for minimizers of (P) (see Proposition 5.2), and allows for the derivation of a KKT-type optimality system (see Theorem 5.3). Note that these results also imply that PDEs of the form $-\Delta y + \operatorname{sgn}(y)|y|^\alpha = u$ with $\alpha \in (0, 1)$ are, in a certain sense, better behaved than elliptic partial differential equations that involve a Lipschitz continuous, directionally differentiable semilinearity; see [8]. However, the nonsmoothness of the PDE governing (P) is not without consequences either. As the analysis of Section 5 shows, it causes the obtained KKT-systems to involve Sobolev spaces with singular weights that depend on the state, may blow up arbitrarily fast, and typically do not satisfy a Muckenhoupt property. At least to the best of the author's knowledge, this effect and the resulting KKT-conditions have not been documented so far in the literature. Note that the appearance of the weighted Sobolev spaces also causes the derivation of further results, e.g., on finite element error estimates, quadratic growth conditions, and regularity properties of optimal controls, to be a very interesting topic.

We would like to mention at this point that we consider (P) as a model problem in this paper. Extensions to more complicated differential operators etc. are possible without major problems. It should further be noted that the techniques that we use in Section 4 for analyzing the differentiability properties of the control-to-state map S of (P) are, in fact, part of a larger theoretical machinery for the analysis of elliptic variational inequalities (VIs) of the second kind that also allows for generalizations in various other directions; see Remark 4.10 and [7, 9]. We focus on the model PDE $-\Delta y + \operatorname{sgn}(y)|y|^\alpha = u$ in this paper to avoid obscuring the basic ideas of our analysis and the underlying mechanisms with technicalities.

We conclude this introduction with an overview of the content and the structure of the remainder of the paper.

Sections 2 and **3** are concerned with preliminaries. Here, we clarify the notation and recall basic results on the state equation of **(P)**. In **Section 4**, we establish the Fréchet differentiability of the control-to-state operator $S: L^2(\Omega) \rightarrow H_0^1(\Omega)$, $u \mapsto y$. The proof that we use in this section relies on Lipschitz continuity properties of S , a reformulation of the PDE $-\Delta y + \text{sgn}(y)|y|^\alpha = u$ as an elliptic variational inequality of the second kind, and techniques that have been developed in [7, 9]. **Section 5** is concerned with the derivation of first-order necessary optimality conditions for local minimizers of **(P)**. Here, we also show that the obtained KKT-systems and Stampacchia truncation arguments allow to establish improved $L^q(\Omega)$ -regularity properties for optimal controls of **(P)**. The paper concludes with some remarks on open questions and topics for future research.

2. Basic notation. In what follows, we use the symbols $\|\cdot\|$ and (\cdot, \cdot) to denote norms and inner products on real vector spaces, respectively, equipped with a subscript that clarifies the space under consideration. For the Euclidean norm on \mathbb{R}^d , $d \in \mathbb{N}$, we write $|\cdot|$. The space of linear and continuous functions from a normed space $(X, \|\cdot\|_X)$ to a normed space $(Y, \|\cdot\|_Y)$ is denoted by $\mathcal{L}(X, Y)$. In the special case $Y = \mathbb{R}$, we write $X^* := \mathcal{L}(X, \mathbb{R})$ for the topological dual space of X and $\langle \cdot, \cdot \rangle_X$ for a dual pairing, i.e., $\langle x^*, x \rangle_X := x^*(x)$ for all $x^* \in X^*$, $x \in X$. The modes of weak and strong convergence in a normed space $(X, \|\cdot\|_X)$ are denoted by \rightharpoonup and \rightarrow , respectively. If $(X, \|\cdot\|_X)$ embeds continuously into $(Y, \|\cdot\|_Y)$, then we write $X \hookrightarrow Y$. Recall that a function $F: X \rightarrow Y$ between normed spaces $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ is called Gâteaux differentiable if the directional derivative

$$F'(x; h) := \lim_{0 < \tau \rightarrow 0} \frac{F(x + \tau h) - F(x)}{\tau}$$

exists for all $x, h \in X$ and is linear and continuous in h . In this case, the operator $F'(x; \cdot) \in \mathcal{L}(X, Y)$ is called the Gâteaux derivative of F at x and denoted by $F'(x)$. If, for every $x \in X$, there exists $F'(x) \in \mathcal{L}(X, Y)$ such that

$$\lim_{0 < \|h\|_X \rightarrow 0} \frac{\|F(x+h) - F(x) - F'(x)h\|_Y}{\|h\|_X} = 0$$

holds, then F is called Fréchet differentiable with derivative $F': X \rightarrow \mathcal{L}(X, Y)$. If the argument of F has several components, then a partial Fréchet derivative with respect to (w.r.t.) a component z is denoted by ∂_z .

Given a Lebesgue measurable set $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$, we denote by $L^q(\Omega)$, $1 \leq q \leq \infty$, the standard real Lebesgue spaces on Ω , equipped with their usual norms. For level sets and complements of level sets of elements v of the Lebesgue spaces (defined up to sets of measure zero), we use the shorthand notation $\{v = c\}$ and $\{v \neq c\}$, $c \in \mathbb{R}$, respectively. The $\{0, 1\}$ -indicator function of a measurable set $D \subset \Omega$ is denoted by $\mathbf{1}_D \in L^\infty(\Omega)$. If Ω is a nonempty, open, and bounded set, then we write $W^{k,q}(\Omega)$, $H^k(\Omega)$, $k \in \mathbb{N}$, $1 \leq q \leq \infty$, for the Sobolev spaces on Ω , defined as in [1, Chapter 5] and again equipped with their usual norms. With $H_0^1(\Omega)$, we denote the closure of the set $C_c^\infty(\Omega)$ of smooth functions with compact support on Ω w.r.t. the $H^1(\Omega)$ -norm. Recall that the space $H_0^1(\Omega)$ is Hilbert when endowed with the inner product

$$(v, w)_{H_0^1(\Omega)} := \langle -\Delta v, w \rangle_{H_0^1(\Omega)} = \int_{\Omega} \nabla v \cdot \nabla w \, dx \quad \forall v, w \in H_0^1(\Omega).$$

Here, Δ and ∇ are the distributional Laplacian and the weak gradient, respectively. As usual, we denote by $H^{-1}(\Omega)$ the dual space of $H_0^1(\Omega)$ with pivot space $L^2(\Omega)$, i.e., we use the identifications $H_0^1(\Omega) \hookrightarrow L^2(\Omega) \cong L^2(\Omega)^* \hookrightarrow H^{-1}(\Omega)$. Recall that the Sobolev embeddings imply that we have

$$H_0^1(\Omega) \hookrightarrow L^q(\Omega) \quad \forall q \in \mathcal{Q}_d := \begin{cases} [1, \infty] & \text{if } d = 1, \\ [1, \infty) & \text{if } d = 2, \\ \left[1, \frac{2d}{d-2}\right] & \text{if } d > 2, \end{cases} \quad (1)$$

and, by duality,

$$L^q(\Omega) \hookrightarrow H^{-1}(\Omega) \quad \forall q \in \mathcal{Q}_d^* := \begin{cases} [1, \infty] & \text{if } d = 1, \\ (1, \infty] & \text{if } d = 2, \\ \left[\frac{2d}{d+2}, \infty\right] & \text{if } d > 2; \end{cases} \quad (2)$$

see [1, Theorem 5.7.2]. Recall further that the embeddings in (1) are compact except for the limit case $q = 2d/(d-2)$, $d > 2$; see [1, Theorem 5.7.7]. (Note that no regularity of the boundary $\partial\Omega$ is needed here due to the zero boundary conditions.) By Schauder's theorem, this also implies that the embeddings in (2) are compact, except for the case $q = 2d/(d+2)$, $d > 2$. We remark that additional symbols etc. are introduced in the remainder of this paper wherever necessary. This notation is clarified on its first appearance.

3. Well-posedness and Lipschitz stability of the governing PDE. We begin our study of the optimal control problem (P) by collecting preliminary results on the well-posedness and Lipschitz stability of the state equation

$$-\Delta y + \operatorname{sgn}(y)|y|^\alpha = u \text{ in } \Omega, \quad y = 0 \text{ on } \partial\Omega. \quad (\text{D})$$

For the sake of clarity, we restate our assumptions on the quantities in (D) in:

Assumption 3.1 (standing assumptions on the governing PDE).

- i) $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$, is a nonempty open bounded set;
- ii) $\alpha \in (0, 1)$ is a given exponent;
- iii) $u \in H^{-1}(\Omega)$ is a given right-hand side.

As usual, we understand solutions of (D) in the weak sense.

Definition 3.2 (weak solution). Given $u \in H^{-1}(\Omega)$, we call y a weak solution of (D) with right-hand side u if $y \in H_0^1(\Omega)$ holds and

$$\int_{\Omega} \nabla y \cdot \nabla v + \operatorname{sgn}(y)|y|^\alpha v \, dx = \langle u, v \rangle_{H_0^1(\Omega)} \quad \forall v \in H_0^1(\Omega). \quad (3)$$

The existence of a unique weak solution of (D) for all $u \in H^{-1}(\Omega)$ is a straightforward consequence of standard results from monotone operator theory.

Lemma 3.3 (unique solvability of (D)). The PDE (D) possesses a unique weak solution $y \in H_0^1(\Omega)$ for all $u \in H^{-1}(\Omega)$.

Proof. This follows from [14, Theorem 4.1] and trivial calculations. \square

Due to Lemma 3.3, it makes sense to introduce the following definition.

Definition 3.4 (solution operator). We denote by $S: H^{-1}(\Omega) \rightarrow H_0^1(\Omega)$, $u \mapsto y$, the solution operator of the partial differential equation (D).

Note that, in the remainder of this paper, the operator S is sometimes also considered on domains of definition that are smaller than $H^{-1}(\Omega)$ (e.g., $L^2(\Omega)$ in the context of the problem (P)). We denote these restrictions of S with the same symbol for the sake of simplicity. The next lemma establishes a connection between the PDE (D) and a certain elliptic variational inequality of the second kind.

Lemma 3.5 (equivalence to an elliptic VI). *Let $u \in H^{-1}(\Omega)$ be given. A function $y \in H_0^1(\Omega)$ is the weak solution of (D) with right-hand side u if and only if it solves the elliptic variational inequality of the second kind*

$$\left. \begin{aligned} y &\in H_0^1(\Omega), \\ (y, v - y)_{H_0^1(\Omega)} + \int_{\Omega} \frac{|v|^{\alpha+1}}{\alpha+1} dx - \int_{\Omega} \frac{|y|^{\alpha+1}}{\alpha+1} dx &\geq \langle u, v - y \rangle_{H_0^1(\Omega)} \\ &\forall v \in H_0^1(\Omega). \end{aligned} \right\} \quad (\text{V})$$

Proof. If $y \in H_0^1(\Omega)$ is the weak solution of (D), then our definition of the $H_0^1(\Omega)$ -inner product, (3), and the properties of convex subgradients imply

$$\begin{aligned} -(y, v - y)_{H_0^1(\Omega)} + \langle u, v - y \rangle_{H_0^1(\Omega)} &= \int_{\Omega} \operatorname{sgn}(y) |y|^{\alpha} (v - y) dx \\ &\leq \int_{\Omega} \frac{|v|^{\alpha+1}}{\alpha+1} dx - \int_{\Omega} \frac{|y|^{\alpha+1}}{\alpha+1} dx \quad \forall v \in H_0^1(\Omega). \end{aligned}$$

Thus, y is a solution of (V) as claimed. Note that this also shows that (V) possesses at least one solution for all $u \in H^{-1}(\Omega)$; see Lemma 3.3. As (V) can have at most one solution (as one may easily check by means of a trivial contradiction argument; see the next corollary and also [12, Theorem 4.1]), it follows that every solution of (V) coincides with the unique weak solution of (D). This completes the proof. \square

The main advantage of the variational inequality (V) is that it does not involve the signum function or potentially singular fractions (as appearing in (D) when using the reformulation $\operatorname{sgn}(y)|y|^{\alpha} = y/|y|^{1-\alpha}$). This is why we choose (V) as the point of departure for the analysis of the differentiability properties of S in Section 4 instead of (D). From Lemma 3.5, we also easily obtain that S is globally Lipschitz.

Corollary 3.6 (Lipschitz continuity of the solution operator). *The solution map $S: H^{-1}(\Omega) \rightarrow H_0^1(\Omega)$ of (D) (respectively, (V)) is globally one-Lipschitz, i.e.,*

$$\|S(u_1) - S(u_2)\|_{H_0^1(\Omega)} \leq \|u_1 - u_2\|_{H^{-1}(\Omega)} \quad \forall u_1, u_2 \in H^{-1}(\Omega).$$

Proof. If $u_1, u_2 \in H^{-1}(\Omega)$ are given and $y_i := S(u_i)$ for $i = 1, 2$, then, by choosing y_2 as the test function in the VI (V) for y_1 , by choosing y_1 as the test function in the VI (V) for y_2 , and by adding the resulting inequalities, we obtain that

$$\|y_1 - y_2\|_{H_0^1(\Omega)}^2 \leq \langle u_1 - u_2, y_1 - y_2 \rangle_{H_0^1(\Omega)} \leq \|u_1 - u_2\|_{H^{-1}(\Omega)} \|y_1 - y_2\|_{H_0^1(\Omega)}.$$

This proves the assertion of the corollary. \square

4. Differentiability of the PDE solution map. With the preliminaries in place, we can turn our attention to the analysis of the differentiability properties of the solution map $S: u \mapsto y$ of (D). To prove that this map possesses classical derivatives even though (D) contains terms that are merely Hölder continuous, we will exploit the characterization of S by means of (V) and techniques that have been developed for the analysis of elliptic VIs of the first and the second kind in [7, 9]. We begin with a technical lemma that is a special version of [7, Lemma 4.2.2].

Lemma 4.1 (Taylor-like expansion for powers of absolute value functions).

Let $\beta \in (0, 1)$. Then, for all $x \in \mathbb{R} \setminus \{0\}$, $t > 0$, and $z \in \mathbb{R}$, we have

$$\begin{aligned} & \frac{1}{t} \left(\frac{|x + tz|^{\beta+1} - |x|^{\beta+1}}{t} - (\beta + 1)|x|^{\beta-1}xz \right) \\ &= (\beta + 1)|x|^\beta \frac{1}{t} \left(\frac{|x + tz| - |x|}{t} - |x|^{-1}xz \right) \\ &+ \frac{1}{t^2}(|x + tz| - |x|)^2(\beta^2 + \beta) \int_0^1 (1-s) \left((1-s)|x| + s|x + tz| \right)^{\beta-1} ds. \end{aligned} \quad (4)$$

Further, it holds

$$\frac{1}{t} \left(\frac{|x + tz| - |x|}{t} - |x|^{-1}xz \right) = \frac{|x|(|x + tz| - |x|) - txz}{|x|(|x + tz| + |x|)^2} z^2 \quad (5)$$

and

$$\frac{1}{t^2}(|x + tz| - |x|)^2 = \frac{4x^2}{(|x| + |x + tz|)^2} z^2 + \frac{4txz + t^2z^2}{(|x| + |x + tz|)^2} z^2. \quad (6)$$

Proof. As the function $(0, c) \ni s \mapsto s^{\beta+1} \in \mathbb{R}$ is in $W^{2,q}(0, c)$ for all $1 \leq q < (1-\beta)^{-1}$ and all $c > 0$, Taylor's theorem implies

$$b^{\beta+1} - a^{\beta+1} = (\beta + 1)a^\beta(b - a) + (b - a)^2(\beta^2 + \beta) \int_0^1 (1-s)((1-s)a + sb)^{\beta-1} ds$$

for all $a > 0$ and $b \geq 0$. If we choose $a := |x| > 0$ and $b := |x + tz| \geq 0$ in the above and plug into the left-hand side of (4), then (4) follows immediately. The equalities in (5) and (6) are obtained by straightforward calculation. Indeed, by distinguishing between the cases $\text{sgn}(x + tz) = \text{sgn}(x)$ and $\text{sgn}(x + tz) \neq \text{sgn}(x)$, one easily checks that

$$\frac{1}{t} \left(\frac{|x + tz| - |x|}{t} - |x|^{-1}xz \right) = (|x + tz| - |x| - t|x|^{-1}xz) \frac{z^2}{(|x + tz| + |x|)^2}$$

holds for all $x \in \mathbb{R} \setminus \{0\}$, $t > 0$, and $z \in \mathbb{R}$. This proves (5). Using the third binomial identity, we may further compute that

$$\begin{aligned} \frac{1}{t^2}(|x + tz| - |x|)^2 &= \frac{(|x + tz|^2 - |x|^2)^2}{t^2} \frac{1}{(|x + tz| + |x|)^2} \\ &= \frac{(2txz + t^2z^2)^2}{t^2} \frac{1}{(|x + tz| + |x|)^2} \\ &= \frac{4x^2z^2 + 4txz^3 + t^2z^4}{(|x + tz| + |x|)^2} \\ &= \frac{4x^2}{(|x| + |x + tz|)^2} z^2 + \frac{4txz + t^2z^2}{(|x| + |x + tz|)^2} z^2. \end{aligned}$$

This establishes (6) and completes the proof of the lemma. \square

To prepare our sensitivity analysis, we next introduce some notation.

Definition 4.2 (difference quotient). Given $u, h \in H^{-1}(\Omega)$ and $\tau > 0$, we denote by $\delta_\tau \in H_0^1(\Omega)$ the difference quotient

$$\delta_\tau := \frac{S(u + \tau h) - S(u)}{\tau}. \quad (7)$$

By exploiting [Lemma 3.5](#), it is easy to establish that the difference quotients of S are also characterized by an elliptic variational inequality of the second kind.

Lemma 4.3 (VI for the difference quotients). *For every $u \in H^{-1}(\Omega)$ with state $y := S(u)$, every $h \in H^{-1}(\Omega)$, and every $\tau > 0$, the difference quotient δ_τ is the (necessarily unique) solution of the variational inequality*

$$\begin{aligned} \delta_\tau &\in H_0^1(\Omega), \\ (\delta_\tau, z - \delta_\tau)_{H_0^1(\Omega)} &+ \frac{1}{(\alpha+1)} \int_\Omega \frac{1}{\tau} \left(\frac{|y + \tau z|^{\alpha+1} - |y|^{\alpha+1}}{\tau} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha z \right) dx \\ &- \frac{1}{(\alpha+1)} \int_\Omega \frac{1}{\tau} \left(\frac{|y + \tau \delta_\tau|^{\alpha+1} - |y|^{\alpha+1}}{\tau} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha \delta_\tau \right) dx \\ &\geq \langle h, z - \delta_\tau \rangle_{H_0^1(\Omega)} \quad \forall z \in H_0^1(\Omega). \end{aligned} \quad (8)$$

Proof. To show that δ_τ solves (8), one chooses test functions of the form $v = y + \tau z$ in the VI (V) satisfied by $S(u + \tau h)$, plugs in the identity $S(u + \tau h) = y + \tau \delta_\tau$ obtained from (7), exploits the variational identity (3) satisfied by y , divides the resulting inequality by τ^2 , and artificially introduces the $|y|^{\alpha+1}$ -terms (which add up to zero in (8)). That (8) can have at most one solution again follows from a trivial contradiction argument; see [12, Theorem 4.1]. \square

Note that the integrands in (8) have the same form as the left-hand side of (4). From [Corollary 3.6](#), we further obtain that the family of difference quotients $\{\delta_\tau\}$ associated with a right-hand side $u \in H^{-1}(\Omega)$ and a perturbation $h \in H^{-1}(\Omega)$ is bounded in $H_0^1(\Omega)$ and, thus, possesses weakly convergent subsequences for $\tau \rightarrow 0$. In what follows, the main idea is to pass to the limit $\tau \rightarrow 0$ in (8) along weakly convergent subsequences of difference quotients to arrive at a limit variational inequality and to exploit this limit VI to show that the difference quotients δ_τ can have only one weak accumulation point in $H_0^1(\Omega)$ for $\tau \rightarrow 0$. By means of a trivial contradiction argument, it then follows that the whole family of difference quotients δ_τ converges weakly in $H_0^1(\Omega)$ to a unique $\delta \in H_0^1(\Omega)$ for $\tau \rightarrow 0$, and by means of bootstrapping and compactness arguments, that the solution operator S of (D) is Fréchet differentiable as a function $S: L^2(\Omega) \rightarrow H_0^1(\Omega)$. The most delicate point when arguing along these lines is, of course, the limit transition with the nonsmooth terms in (8). We proceed in several steps, starting with the following lemma.

Lemma 4.4 (behavior of the α - δ_τ -terms). *Let $u, h \in H^{-1}(\Omega)$ be given. Define $y := S(u)$. Suppose that $\delta_k := \delta_{\tau_k}$ are difference quotients as in (7) associated with a sequence $\{\tau_k\} \subset (0, \infty)$. Assume that $\tau_k \rightarrow 0$ holds and that $\delta_k \rightharpoonup \delta$ in $H_0^1(\Omega)$ for some $\delta \in H_0^1(\Omega)$. Then it holds*

$$\delta = 0 \text{ a.e. in } \{y = 0\} \quad (9)$$

and

$$\begin{aligned} \infty &> \liminf_{k \rightarrow \infty} \left[\frac{1}{(\alpha+1)} \int_\Omega \frac{1}{\tau_k} \left(\frac{|y + \tau_k \delta_k|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha \delta_k \right) dx \right] \\ &\geq \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{\delta^2}{|y|^{1-\alpha}} dx. \end{aligned} \quad (10)$$

Proof. By choosing the test function $z = 0$ in the VI (8) for δ_k , we obtain that

$$\begin{aligned}
& \langle h, \delta_k \rangle_{H_0^1(\Omega)} \\
& \geq \langle h, \delta_k \rangle_{H_0^1(\Omega)} - \|\delta_k\|_{H_0^1(\Omega)}^2 \\
& \geq \frac{1}{(\alpha+1)} \int_{\Omega} \frac{1}{\tau_k} \left(\frac{|y + \tau_k \delta_k|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha \delta_k \right) dx \\
& = \frac{1}{(\alpha+1)} \int_{\{y \neq 0\}} \frac{1}{\tau_k} \left(\frac{|y + \tau_k \delta_k|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha \delta_k \right) dx \\
& \quad + \frac{\tau_k^{\alpha-1}}{(\alpha+1)} \int_{\{y=0\}} |\delta_k|^{\alpha+1} dx.
\end{aligned} \tag{11}$$

Note that the left-hand side of (11) is bounded. This establishes the first inequality in (10). As difference quotients of convex functions dominate the derivatives that they approximate, we further have that the integrands in the integrals in (11) are all nonnegative. In combination with the convergence $\delta_k \rightarrow \delta$ in $L^2(\Omega)$ obtained from the compactness of the embedding $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$, this yields

$$0 = \limsup_{k \rightarrow \infty} (\alpha+1) \tau_k^{1-\alpha} \langle h, \delta_k \rangle_{H_0^1(\Omega)} \geq \limsup_{k \rightarrow \infty} \int_{\{y=0\}} |\delta_k|^{\alpha+1} dx = \int_{\{y=0\}} |\delta|^{\alpha+1} dx.$$

Thus, δ satisfies (9) as claimed. By revisiting (11), by noting that the first term on the right-hand side of (4) is nonnegative (again due to the properties of difference quotients of convex functions), and by invoking Fatou's lemma [5, Corollary 2.8.4] (keeping in mind that all of the involved integrands are nonnegative and that $\delta_k \rightarrow \delta$ holds in $L^2(\Omega)$), we further obtain that

$$\begin{aligned}
& \liminf_{k \rightarrow \infty} \left[\int_{\Omega} \frac{1}{\tau_k} \left(\frac{|y + \tau_k \delta_k|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha \delta_k \right) dx \right] \\
& \geq \liminf_{k \rightarrow \infty} \left[\int_{\{y \neq 0\}} \frac{1}{\tau_k} \left(\frac{|y + \tau_k \delta_k|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha+1) \operatorname{sgn}(y) |y|^\alpha \delta_k \right) dx \right] \\
& \geq \liminf_{k \rightarrow \infty} \left[\int_{\{y \neq 0\}} \left(\frac{|y + \tau_k \delta_k| - |y|}{\tau_k} \right)^2 \right. \\
& \quad \left. \cdot (\alpha^2 + \alpha) \int_0^1 (1-s) \left((1-s)|y| + s|y + \tau_k \delta_k| \right)^{\alpha-1} ds dx \right] \\
& \geq \frac{\alpha^2 + \alpha}{2} \int_{\{y \neq 0\}} \frac{\delta^2}{|y|^{1-\alpha}} dx.
\end{aligned}$$

This completes the proof. \square

Lemma 4.4 motivates the following definition.

Definition 4.5 (weighted Sobolev space). Given a right-hand side $u \in H^{-1}(\Omega)$ with associated state $y := S(u)$, we denote by V_y the subspace of $H_0^1(\Omega)$ given by

$$V_y := \left\{ v \in H_0^1(\Omega) : \int_{\{y \neq 0\}} \frac{v^2}{|y|^{1-\alpha}} dx < \infty \text{ and } v = 0 \text{ a.e. in } \{y = 0\} \right\}.$$

We further define $(\cdot, \cdot)_{V_y}$ to be the bilinear form

$$(\cdot, \cdot)_{V_y} : V_y \times V_y \rightarrow \mathbb{R}, \quad (v, w)_{V_y} := (v, w)_{H_0^1(\Omega)} + \alpha \int_{\{y \neq 0\}} \frac{vw}{|y|^{1-\alpha}} dx.$$

Note that V_y is Hilbert when endowed with $(\cdot, \cdot)_{V_y}$ as the inner product and that the limit δ in [Lemma 4.4](#) satisfies $\delta \in V_y$ by [\(9\)](#) and [\(10\)](#). The next lemma shows how to handle the limit transition with the nonsmooth terms involving the test function z in the variational inequality [\(8\)](#).

Lemma 4.6 (behavior of the α - z -terms). *Let $u \in H^{-1}(\Omega)$ be given. Define $y := S(u)$. Suppose that $\{\tau_k\} \subset (0, \infty)$ is a sequence satisfying $\tau_k \rightarrow 0$. Then, for all $z \in V_y$, we have*

$$\begin{aligned} & \lim_{k \rightarrow \infty} \frac{1}{(\alpha + 1)} \int_{\Omega} \frac{1}{\tau_k} \left(\frac{|y + \tau_k z|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha + 1) \operatorname{sgn}(y) |y|^{\alpha} z \right) dx \\ &= \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{z^2}{|y|^{1-\alpha}} dx. \end{aligned}$$

Proof. Due to the identities in [Lemma 4.1](#), it holds

$$\begin{aligned} & \frac{1}{\tau_k} \left(\frac{|y + \tau_k z|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha + 1) \operatorname{sgn}(y) |y|^{\alpha} z \right) \\ &= (\alpha + 1) \frac{|y|(|y + \tau_k z| - |y|) - \tau_k y z}{(|y + \tau_k z| + |y|)^2} \frac{z^2}{|y|^{1-\alpha}} \\ &+ (\alpha^2 + \alpha) \left(\frac{4y^2}{(|y| + |y + \tau_k z|)^2} \right) \int_0^1 (1-s) \left(\frac{|y|}{(1-s)|y| + s|y + \tau_k z|} \right)^{1-\alpha} ds \frac{z^2}{|y|^{1-\alpha}} \\ &+ (\alpha^2 + \alpha) \left(\frac{4\tau_k y z + \tau_k^2 z^2}{(|y| + |y + \tau_k z|)^2} \right) \int_0^1 (1-s) \left(\frac{|y|}{(1-s)|y| + s|y + \tau_k z|} \right)^{1-\alpha} ds \frac{z^2}{|y|^{1-\alpha}} \end{aligned}$$

a.e. in $\{y \neq 0\}$. From the triangle inequality and elementary estimates, we further obtain that we have

$$\left| \frac{|y|(|y + \tau_k z| - |y|) - \tau_k y z}{(|y + \tau_k z| + |y|)^2} \right| \leq \frac{|y + \tau_k z| + |y| + \tau_k |z|}{|y + \tau_k z| + |y|} \leq 2$$

and

$$\begin{aligned} & \left(\frac{4y^2}{(|y| + |y + \tau_k z|)^2} \right) \int_0^1 (1-s) \left(\frac{|y|}{(1-s)|y| + s|y + \tau_k z|} \right)^{1-\alpha} ds \leq 4 \int_0^1 (1-s)^{\alpha} ds \\ & \leq \frac{4}{\alpha + 1} \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{4\tau_k y z + \tau_k^2 z^2}{(|y| + |y + \tau_k z|)^2} \right| \int_0^1 (1-s) \left(\frac{|y|}{(1-s)|y| + s|y + \tau_k z|} \right)^{1-\alpha} ds \\ & \leq \frac{4\tau_k |y| |z| + \tau_k^2 z^2}{(|y| + |y + \tau_k z|)^2} \int_0^1 (1-s)^{\alpha} ds \\ & \leq \frac{5}{\alpha + 1} \end{aligned}$$

a.e. in $\{y \neq 0\}$. As $z = 0$ holds a.e. in $\{y = 0\}$ and since $z^2 |y|^{\alpha-1}$ is an element of $L^1(\{y \neq 0\})$ by our assumption $z \in V_y$, the above allows us to invoke Lebesgue's

dominated convergence theorem [5, Theorem 2.8.1] to obtain

$$\begin{aligned}
& \lim_{k \rightarrow \infty} \frac{1}{(\alpha + 1)} \int_{\Omega} \frac{1}{\tau_k} \left(\frac{|y + \tau_k z|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha + 1) \operatorname{sgn}(y) |y|^{\alpha} z \right) dx \\
&= \lim_{k \rightarrow \infty} \frac{1}{(\alpha + 1)} \int_{\{y \neq 0\}} \frac{1}{\tau_k} \left(\frac{|y + \tau_k z|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha + 1) \operatorname{sgn}(y) |y|^{\alpha} z \right) dx \\
&= \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{z^2}{|y|^{1-\alpha}} dx.
\end{aligned}$$

This completes the proof. \square

By putting everything together, we can now pass to the limit in (8) to arrive at a VI for the weak accumulation points of the difference quotients $\{\delta_{\tau}\}$ for $\tau \rightarrow 0$.

Proposition 4.7 (VI for limits of difference quotients). *Let $u, h \in H^{-1}(\Omega)$ be given. Define $y := S(u)$. Suppose that $\delta_k := \delta_{\tau_k}$ are difference quotients as in (7) associated with a sequence $\{\tau_k\} \subset (0, \infty)$. Assume that $\tau_k \rightarrow 0$ holds and that $\delta_k \rightharpoonup \delta$ in $H_0^1(\Omega)$ for some $\delta \in H_0^1(\Omega)$. Then δ_k converges even strongly in $H_0^1(\Omega)$ to δ and δ is the solution of the variational inequality*

$$\begin{aligned}
& \delta \in V_y, \\
& (\delta, z - \delta)_{H_0^1(\Omega)} + \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{z^2}{|y|^{1-\alpha}} dx - \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{\delta^2}{|y|^{1-\alpha}} dx \geq (h, z - \delta)_{H_0^1(\Omega)} \quad (12) \\
& \forall z \in V_y.
\end{aligned}$$

Proof. Due to (8), we know that

$$\begin{aligned}
& \langle h, \delta_k - z \rangle_{H_0^1(\Omega)} + (\delta_k, z)_{H_0^1(\Omega)} \\
&+ \frac{1}{(\alpha + 1)} \int_{\Omega} \frac{1}{\tau_k} \left(\frac{|y + \tau_k z|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha + 1) \operatorname{sgn}(y) |y|^{\alpha} z \right) dx \\
&\geq \|\delta_k\|_{H_0^1(\Omega)}^2 + \frac{1}{(\alpha + 1)} \int_{\Omega} \frac{1}{\tau_k} \left(\frac{|y + \tau_k \delta_k|^{\alpha+1} - |y|^{\alpha+1}}{\tau_k} - (\alpha + 1) \operatorname{sgn}(y) |y|^{\alpha} \delta_k \right) dx
\end{aligned}$$

holds for all k and all $z \in V_y$. By taking the limes superior for $k \rightarrow \infty$ on the left and the right of this inequality, by exploiting the weak lower semicontinuity of the function $H_0^1(\Omega) \ni v \mapsto \|v\|_{H_0^1(\Omega)}^2 \in \mathbb{R}$, and by using Lemmas 4.4 and 4.6, we get

$$\begin{aligned}
& \langle h, \delta - z \rangle_{H_0^1(\Omega)} + (\delta, z)_{H_0^1(\Omega)} + \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{z^2}{|y|^{1-\alpha}} dx \\
&\geq \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{\delta^2}{|y|^{1-\alpha}} dx + \limsup_{k \rightarrow \infty} \|\delta_k\|_{H_0^1(\Omega)}^2 \\
&\geq \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{\delta^2}{|y|^{1-\alpha}} dx + \liminf_{k \rightarrow \infty} \|\delta_k\|_{H_0^1(\Omega)}^2 \\
&\geq \frac{\alpha}{2} \int_{\{y \neq 0\}} \frac{\delta^2}{|y|^{1-\alpha}} dx + \|\delta\|_{H_0^1(\Omega)}^2 \quad \forall z \in V_y.
\end{aligned} \tag{13}$$

This shows that δ satisfies (12) and, since the choice $z = \delta \in V_y$ is allowed in (13) by Lemma 4.4, that $\|\delta_k\|_{H_0^1(\Omega)} \rightarrow \|\delta\|_{H_0^1(\Omega)}$ holds. In combination with the weak convergence $\delta_k \rightharpoonup \delta$ in $H_0^1(\Omega)$ and the binomial identities, it now follows immediately that δ_k converges also strongly in $H_0^1(\Omega)$ to δ and the proof is complete. \square

As a straightforward consequence of [Proposition 4.7](#), we now obtain our first main result.

Theorem 4.8 (Gâteaux differentiability of the solution operator S). *The solution operator $S: H^{-1}(\Omega) \rightarrow H_0^1(\Omega)$ of the PDE [\(D\)](#) is Gâteaux differentiable at all points $u \in H^{-1}(\Omega)$. Further, the Gâteaux derivative $S'(u) \in \mathcal{L}(H^{-1}(\Omega), H_0^1(\Omega))$ of S at a point $u \in H^{-1}(\Omega)$ with state $y := S(u)$ is precisely the solution operator $H^{-1}(\Omega) \ni h \mapsto \delta \in V_y \subset H_0^1(\Omega)$ of the variational problem*

$$\delta \in V_y, \quad (\delta, z)_{V_y} = \langle h, z \rangle_{H_0^1(\Omega)} \quad \forall z \in V_y. \quad (14)$$

Proof. Let $u, h \in H^{-1}(\Omega)$ be fixed and define $y := S(u)$. From [Corollary 3.6](#), we obtain that the family of difference quotients $\{\delta_\tau\}$ associated with u and h is bounded in $H_0^1(\Omega)$ and, thus, possesses weak accumulation points in $H_0^1(\Omega)$ for $\tau \rightarrow 0$. From [Proposition 4.7](#), we obtain that all of these weak accumulation points are, in fact, strong accumulation points of the family $\{\delta_\tau\}$ for $\tau \rightarrow 0$ and solutions of the variational inequality [\(12\)](#). As [\(12\)](#) can have at most one solution (cf. the proof of [Corollary 3.6](#) or [[12](#), Theorem 4.1]), it follows by contradiction that there can only be one accumulation point δ , that $\delta_\tau \rightarrow \delta$ holds in $H_0^1(\Omega)$ for $\tau \rightarrow 0$, and that δ is uniquely characterized by [\(12\)](#). This shows that S is directionally differentiable at u in direction h with directional derivative δ . By choosing test functions of the form $\delta + sz$ in [\(12\)](#) for arbitrary $s \in (0, 1)$ and $z \in V_y$, by dividing by s in the resulting inequality, and by letting s go to zero, one further easily checks that [\(12\)](#) can be recast as [\(14\)](#). As [\(14\)](#) is precisely the variational problem that characterizes the Riesz representative of $h \in H^{-1}(\Omega) \subset V_y^*$ in the Hilbert space V_y and since $V_y \hookrightarrow H_0^1(\Omega)$ holds, it follows that the directional derivative $\delta \in H_0^1(\Omega)$ depends linearly and continuously on the direction $h \in H^{-1}(\Omega)$. Thus, S is Gâteaux differentiable as claimed, with the Gâteaux derivative being characterized by [\(14\)](#). This completes the proof. \square

Due to the Lipschitz estimate for S in [Corollary 3.6](#) and the compactness of the embedding $L^q(\Omega) \hookrightarrow H^{-1}(\Omega)$ for all $\max(1, 2d/(d+2)) < q \leq \infty$, [Theorem 4.8](#) also implies the following Fréchet differentiability result for S .

Corollary 4.9 (Fréchet differentiability of the solution operator S). *The solution map S of the PDE [\(D\)](#) is Fréchet differentiable as a function from $L^q(\Omega)$ to $H_0^1(\Omega)$ for all $\max(1, 2d/(d+2)) < q \leq \infty$. Its Fréchet derivatives $S'(u) \in \mathcal{L}(L^q(\Omega), H_0^1(\Omega))$ are characterized by the variational identity [\(14\)](#) in [Theorem 4.8](#).*

Proof. The assertion follows from a classical contradiction argument. We begin with the case $q \neq \infty$. Suppose that there are $\max(1, 2d/(d+2)) < q < \infty$ and $u \in L^q(\Omega)$ such that S is not Fréchet differentiable at u as a function $S: L^q(\Omega) \rightarrow H_0^1(\Omega)$. Then there exist a number $\varepsilon > 0$ and sequences $\{h_k\} \subset L^q(\Omega)$ and $\{\tau_k\} \subset (0, \infty)$ such that $\|h_k\|_{L^q(\Omega)} = 1$ for all k , $\tau_k \rightarrow 0$ for $k \rightarrow \infty$, and

$$\varepsilon \leq \liminf_{k \rightarrow \infty} \frac{\|S(u + \tau_k h_k) - S(u) - \tau_k S'(u) h_k\|_{H_0^1(\Omega)}}{\tau_k}.$$

As $\{h_k\} \subset L^q(\Omega)$ is bounded and $q \in (1, \infty)$, we may assume without loss of generality that $h_k \rightharpoonup h$ holds in $L^q(\Omega)$ for some $h \in L^q(\Omega)$. Due to the compactness of the embedding $L^q(\Omega) \hookrightarrow H^{-1}(\Omega)$, [Corollary 3.6](#), the Gâteaux differentiability of S at u in direction h obtained from [Theorem 4.8](#), and $S'(u) \in \mathcal{L}(H^{-1}(\Omega), H_0^1(\Omega))$,

it follows that

$$\begin{aligned}
\varepsilon &\leq \liminf_{k \rightarrow \infty} \frac{\|S(u + \tau_k h_k) - S(u) - \tau_k S'(u) h_k\|_{H_0^1(\Omega)}}{\tau_k} \\
&\leq \liminf_{k \rightarrow \infty} \left(\frac{\|S(u + \tau_k h) - S(u) - \tau_k S'(u) h\|_{H_0^1(\Omega)}}{\tau_k} \right. \\
&\quad \left. + \frac{\|S(u + \tau_k h_k) - S(u + \tau_k h)\|_{H_0^1(\Omega)}}{\tau_k} + \|S'(u) h - S'(u) h_k\|_{H_0^1(\Omega)} \right) \\
&\leq 0 + \liminf_{k \rightarrow \infty} \|h_k - h\|_{H^{-1}(\Omega)} + 0 = 0.
\end{aligned}$$

This produces the contradiction $0 < \varepsilon \leq 0$ and shows that S indeed has to be Fréchet differentiable as a function $S: L^q(\Omega) \rightarrow H_0^1(\Omega)$ for all $\max(1, 2d/(d+2)) < q < \infty$. To obtain the assertion for $q = \infty$, it now suffices to use Hölder's inequality. This completes the proof. \square

We conclude this section with some comments.

Remark 4.10.

- Note that neither *Theorem 4.8* nor *Corollary 4.9* makes a statement about the continuity of the derivative $u \mapsto S'(u)$. The question of whether/on which sets the operator S is continuously differentiable is inherently related to the questions of how the space V_y varies with y and of whether the Meyers-Serrin identity $H = W$ holds in V_y . Compare also with the remarks after *Theorem 5.3* in this context. We leave this topic for future research.
- It is possible to prove analogues of *Theorem 4.8* and *Corollary 4.9* for PDEs of the form

$$-\Delta y - \nabla \cdot \left(\frac{\nabla y}{|\nabla y|^{1-\alpha}} \right) = u \text{ in } \Omega, \quad y = 0 \text{ on } \partial\Omega, \quad (15)$$

with $\alpha \in (0, 1)$; see [7, Section 4.3.5]. As one cannot exploit the compactness of Sobolev embeddings when studying the PDE (15), doing so, however, requires more involved instruments from the field of sensitivity analysis of elliptic VIs of the second kind; cf. [7, Chapters 1, 4].

5. Necessary optimality conditions for the optimal control problem. With the differentiability results in *Theorem 4.8* and *Corollary 4.9* at hand, we can turn our attention to the derivation of first-order necessary optimality conditions for the optimal control problem

$$\left. \begin{aligned}
&\text{Minimize } J(y, u) \\
&\text{w.r.t. } y \in H_0^1(\Omega), \quad u \in L^2(\Omega), \\
&\text{s.t. } -\Delta y + \text{sgn}(y)|y|^\alpha = u \text{ in } \Omega, \quad y = 0 \text{ on } \partial\Omega, \\
&\text{and } u \in U_{\text{ad}}.
\end{aligned} \right\} \quad (\text{P})$$

We begin by recalling our assumptions on the quantities in (P).

Assumption 5.1 (standing assumptions on (P)).

- i) $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$, is a nonempty open bounded set;
- ii) $J: H_0^1(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$ is a Fréchet differentiable function;
- iii) $\alpha \in (0, 1)$ is a given exponent;
- iv) U_{ad} is a nonempty and convex subset of $L^2(\Omega)$.

Note that, to be able to ensure that (P) possesses an optimal control-state pair, one requires more assumptions on U_{ad} and J than stated in [Assumption 5.1](#). For the discussion of necessary optimality conditions, however, this is irrelevant. As a first consequence of [Corollary 4.9](#), we obtain the following result.

Proposition 5.2 (Bouligand stationarity condition). *Suppose that $\bar{u} \in L^2(\Omega)$ is a locally optimal control of (P) with associated state $\bar{y} := S(\bar{u})$, i.e., an $L^2(\Omega)$ -local minimizer of the reduced problem*

$$\text{Minimize } J(S(u), u) \quad \text{s.t. } u \in U_{\text{ad}}.$$

Then \bar{u} is a solution of the variational inequality

$$\bar{u} \in U_{\text{ad}}, \quad \langle \partial_y J(\bar{y}, \bar{u}), S'(\bar{u})(u - \bar{u}) \rangle_{H_0^1(\Omega)} + (\partial_u J(\bar{y}, \bar{u}), u - \bar{u})_{L^2(\Omega)} \geq 0 \quad \forall u \in U_{\text{ad}}. \quad (16)$$

Here, $\partial_y J(\bar{y}, \bar{u}) \in H^{-1}(\Omega)$ and $\partial_u J(\bar{y}, \bar{u}) \in L^2(\Omega) \cong L^2(\Omega)^*$ denote the partial Fréchet derivatives of J w.r.t. y and u at (\bar{y}, \bar{u}) , respectively.

Proof. The assertion of this proposition follows immediately from [Corollary 4.9](#), the Fréchet differentiability of J , the chain rule, and the convexity of U_{ad} . \square

By exploiting the characterization of $S'(\bar{u})$ by means of the variational identity [\(14\)](#), we can reformulate [\(16\)](#) to arrive at a more tangible optimality condition. The resulting stationarity system is the second main result of this work.

Theorem 5.3 (KKT-system). *Let $\bar{u} \in L^2(\Omega)$ be a control with associated state $\bar{y} := S(\bar{u})$. Then \bar{u} satisfies the Bouligand stationarity condition [\(16\)](#) if and only if there exists an adjoint state \bar{p} such that the following KKT-system holds:*

$$\begin{aligned} \bar{u} \in U_{\text{ad}}, \quad \bar{y} \in H_0^1(\Omega), \quad \bar{p} \in V_{\bar{y}}, \\ -\Delta \bar{y} + \text{sgn}(\bar{y})|\bar{y}|^\alpha = \bar{u} \text{ in } \Omega, \quad \bar{y} = 0 \text{ on } \partial\Omega, \\ (\bar{p}, z)_{H_0^1(\Omega)} + \alpha \int_{\{\bar{y} \neq 0\}} \frac{\bar{p}z}{|\bar{y}|^{1-\alpha}} dx = \langle \partial_y J(\bar{y}, \bar{u}), z \rangle_{H_0^1(\Omega)} \quad \forall z \in V_{\bar{y}}, \\ (\bar{p} + \partial_u J(\bar{y}, \bar{u}), u - \bar{u})_{L^2(\Omega)} \geq 0 \quad \forall u \in U_{\text{ad}}. \end{aligned} \quad (17)$$

Here, the state equation is again understood weakly and $V_{\bar{y}}$ is defined by

$$V_{\bar{y}} := \left\{ v \in H_0^1(\Omega) : \int_{\{\bar{y} \neq 0\}} \frac{v^2}{|\bar{y}|^{1-\alpha}} dx < \infty \text{ and } v = 0 \text{ a.e. in } \{\bar{y} = 0\} \right\}. \quad (18)$$

Proof. Suppose that \bar{u} and $\bar{y} := S(\bar{u})$ satisfy [\(16\)](#). Then the Riesz representation theorem implies that there exists a unique \bar{p} satisfying

$$\bar{p} \in V_{\bar{y}}, \quad (\bar{p}, z)_{V_{\bar{y}}} = \langle \partial_y J(\bar{y}, \bar{u}), z \rangle_{H_0^1(\Omega)} \quad \forall z \in V_{\bar{y}}. \quad (19)$$

Using [\(14\)](#), we may compute that

$$\begin{aligned} \langle \partial_y J(\bar{y}, \bar{u}), S'(\bar{u})(u - \bar{u}) \rangle_{H_0^1(\Omega)} &= (\bar{p}, S'(\bar{u})(u - \bar{u}))_{V_{\bar{y}}} \\ &= \langle u - \bar{u}, \bar{p} \rangle_{H_0^1(\Omega)} = (\bar{p}, u - \bar{u})_{L^2(\Omega)} \end{aligned}$$

holds for all $u \in U_{\text{ad}}$. If we plug this identity into [\(16\)](#) and collect everything, then we obtain that \bar{u} and \bar{y} satisfy [\(17\)](#) as claimed.

To prove that every tuple (\bar{u}, \bar{y}) that satisfies [\(17\)](#) for some $\bar{p} \in V_{\bar{y}}$ is a solution of the variational inequality [\(16\)](#), we can proceed along the exact same lines in reverse. This establishes the assertion. \square

Note that the nondifferentiability of the function $\mathbb{R} \ni s \mapsto \text{sgn}(s)|s|^\alpha \in \mathbb{R}$ present in (D) manifests itself in (17) in the form of the singular weight $|\bar{y}|^{\alpha-1} \mathbf{1}_{\{\bar{y} \neq 0\}}$ and the weighted Sobolev space $V_{\bar{y}}$. We remark that the appearance of these quantities significantly complicates the analysis of the regularity properties of the adjoint state \bar{p} . As the state \bar{y} may be arbitrarily smooth and, thus, may go to zero arbitrarily fast when approaching the zero level set $\{\bar{y} = 0\}$, one cannot expect that the function $|\bar{y}|^{\alpha-1}$ is integrable or satisfies a Muckenhoupt property. In particular, it is typically completely unclear whether results like the theorem of Meyers-Serrin (i.e., the statement $H = W$) hold in the space $V_{\bar{y}}$ or not and whether \bar{p} possesses, e.g., some form of $W^{2,q}$ -regularity; cf. [15] and the references therein. Note that the latter question is of particular interest as it is crucial for the derivation of a-priori error estimates for finite element discretizations of (P). The only thing that can be established rather easily for the adjoint state \bar{p} in (17) is improved $L^q(\Omega)$ -regularity as the following result shows.

Corollary 5.4 (higher Lebesgue regularity for the adjoint state). *Suppose that \bar{u} , \bar{y} , and \bar{p} satisfying (17) are given and that $\partial_y J(\bar{y}, \bar{u}) \in L^s(\Omega)$ holds for some $\max(1, 2d/(d+2)) < s \leq \infty$. Define*

$$\mathcal{R}_s := \begin{cases} [1, \infty] & \text{if } s > d/2, \\ [1, \infty) & \text{if } s = d/2, \\ \left[1, \left(\frac{1}{s} - \frac{2}{d}\right)^{-1}\right) & \text{if } s < d/2. \end{cases}$$

Then it holds $\bar{p} \in L^r(\Omega)$ for all $r \in \mathcal{R}_s$.

Proof. Define $\bar{p}_k := \bar{p} - \min(k, \max(-k, \bar{p}))$, $k \geq 0$. Then \bar{p}_k is an element of $V_{\bar{y}}$ for all $k \geq 0$ since $\bar{p}_k = 0 - \min(k, \max(-k, 0)) = 0$ holds a.e. in $\{\bar{y} = 0\}$ and

$$\int_{\{\bar{y} \neq 0\}} \frac{\bar{p}_k^2}{|\bar{y}|^{1-\alpha}} dx \leq \int_{\{\bar{y} \neq 0\}} \frac{\bar{p}^2}{|\bar{y}|^{1-\alpha}} dx < \infty \quad \forall k \geq 0.$$

In particular, \bar{p}_k is a valid test function in (19). In combination with (1) and the lemma of Stampacchia [1, Theorem 5.8.2], this allows us to conclude that

$$c_q \|\bar{p}_k\|_{L^q(\Omega)}^2 \leq \|\bar{p}_k\|_{H_0^1(\Omega)}^2 \leq (\bar{p}_k, \bar{p}_k)_{V_{\bar{y}}} \leq (\bar{p}, \bar{p}_k)_{V_{\bar{y}}} = \int_{\Omega} \partial_y J(\bar{y}, \bar{u}) \bar{p}_k dx$$

holds for all $q \in \mathcal{Q}_d$ with some constants $c_q > 0$ and \mathcal{Q}_d as in (1). To obtain the assertion, it now suffices to invoke [10, Lemma 3.4]. This completes the proof. \square

Prototypical examples of functions J that are covered by Corollary 5.4 are tracking-type objectives of the form

$$J: H_0^1(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}, \quad J(y, u) := \frac{1}{2} \|y - y_D\|_{L^2(\Omega)}^2 + \frac{\nu}{2} \|u\|_{L^2(\Omega)}^2,$$

involving a desired state $y_D \in L^2(\Omega)$ and a Tikhonov parameter $\nu > 0$. For this type of objective function, we obtain from Theorem 5.3 and Corollary 5.4 that the adjoint state \bar{p} associated with a Bouligand stationary control \bar{u} has to satisfy $\bar{p} \in V_{\bar{y}} \cap L^r(\Omega)$ for all $r \in \mathcal{R}_d$, where \mathcal{R}_d is given by

$$\mathcal{R}_d := \begin{cases} [1, \infty] & \text{if } d < 4, \\ [1, \infty) & \text{if } d = 4, \\ \left[1, \frac{2d}{d-4}\right) & \text{if } d > 4, \end{cases}$$

where $V_{\bar{y}}$ is defined as in (18), and where d is the dimension of Ω . If, additionally, we assume that U_{ad} has the form $U_{\text{ad}} = \{u \in L^2(\Omega) : u_a \leq u \leq u_b \text{ a.e. in } \Omega\}$ for some $u_a, u_b \in H^1(\Omega)$ satisfying $u_a \leq 0 \leq u_b$ a.e. in Ω , then the projection formula

$$\bar{u} = \max\left(u_a, \min\left(u_b, -\frac{\bar{p}}{\nu}\right)\right)$$

obtained from the last line in (17) and Stampacchia's lemma [1, Theorem 5.8.2] also imply that \bar{u} inherits all of these regularity properties from \bar{p} , i.e., we have

$$\bar{u} \in V_{\bar{y}} \cap L^r(\Omega) \quad \forall r \in \mathcal{R}_d.$$

As already mentioned, whether optimal controls \bar{u} and their associated adjoint states \bar{p} possess higher regularity properties than those above in the situation of (P) (e.g., Lipschitz regularity or a form of $W^{2,q}$ -regularity) is an open problem. The same is true for the derivation of (no-gap) second-order optimality and quadratic growth conditions for problems of the type (P). We leave these topics for future research.

References

- [1] H. Attouch, G. Buttazzo, and G. Michaille. *Variational Analysis in Sobolev and BV Spaces*. SIAM, Philadelphia, 2006.
- [2] H. W. Alt and D. Phillips. A free boundary problem for semilinear elliptic equations. *J. Reine Angew. Math*, 368:63–107, 1986.
- [3] V. Barbu. *Optimal Control of Variational Inequalities*. Research Notes in Mathematics. Pitman, 1984.
- [4] J. W. Barrett and P. Knabner. Finite element approximation of the transport of reactive solutes in porous media. Part I: Error estimates for nonequilibrium adsorption processes. *SIAM J. Numer. Anal*, 34(1):201–227, 1997.
- [5] V. I. Bogachev. *Measure Theory*. Springer, 2007.
- [6] J. W. Barrett and R. M. Shanahan. Finite element approximation of a model reaction-diffusion problem with a non-Lipschitz nonlinearity. *Numer. Math.*, 59(1):217–242, 1991.
- [7] C. Christof. *Sensitivity Analysis of Elliptic Variational Inequalities of the First and the Second Kind*. PhD thesis, Technische Universität Dortmund, 2018.
- [8] C. Christof, C. Meyer, S. Walther, and C. Clason. Optimal control of a non-smooth semilinear elliptic equation. *Math. Control Relat. Fields*, 8(1):247–276, 2018.
- [9] C. Christof and G. Wachsmuth. Differential sensitivity analysis of variational inequalities with locally Lipschitz continuous solution operators. *Appl. Math. Optim.*, 81:23–62, 2020.
- [10] C. Christof and G. Wachsmuth. Semismoothness for solution operators of obstacle-type variational inequalities with applications in optimal control. *SIAM J. Control Optim.*, 61(3):1162–1186, 2023.
- [11] J. I. Díaz, T. Mingazzini, and Á. M. Ramos. On the optimal control for a semilinear equation with cost depending on the free boundary. *Netw. Heterog. Media*, 7(4):605–615, 2012.
- [12] R. Glowinski. *Numerical Methods for Nonlinear Variational Problems*. Springer Berlin, Heidelberg, 2013.
- [13] P. Knabner and R. Rannacher. A priori error analysis for the Galerkin finite element semi-discretization of a parabolic system with non-Lipschitzian nonlinearity. *Vietnam J. Math.*, 45:179–198, 2017.
- [14] F. Tröltzsch. *Optimal Control of Partial Differential Equations*, volume 112 of *Graduate Studies in Mathematics*. AMS, Providence, 2010.
- [15] V. V. Zhikov. Weighted Sobolev spaces. *Sb. Math.*, 189(8):1139, 1998.

Received xxxx 20xx; revised xxxx 20xx.

E-mail address: christof@cit.tum.de