# THE WEYL TRANSFORM OF A SMOOTH MEASURE ON A REAL-ANALYTIC SUBMANIFOLD

MANSI MISHRA AND M. K. VEMURI

ABSTRACT. If  $\mu$  is a smooth measure supported on a real-analytic submanifold of  $\mathbb{R}^{2n}$  which is not contained in any affine hyperplane, then the Weyl transform of  $\mu$  is a compact operator.

### 1. INTRODUCTION

Let  $f \in L^1(\mathbb{R}^n)$ . The Fourier transform of f, denoted by  $\hat{f}$ , is given by

$$\hat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{-2\pi i x \cdot \xi} dx, \quad \xi \in \mathbb{R}^n.$$

The Riemann-Lebesgue lemma [9, Theorem 1.2] states that if  $f \in L^1(\mathbb{R}^n)$ , then

$$\lim_{|\xi|\to\infty}\hat{f}(\xi)=0$$

More generally, the Fourier transform of a finite Borel measure  $\lambda$  on  $\mathbb{R}^n$  is given by

$$\hat{\lambda}(\xi) = \int_{\mathbb{R}^n} e^{-2\pi i x \cdot \xi} d\lambda(x), \quad \xi \in \mathbb{R}^n.$$

If  $\lambda$  is absolutely continuous with respect to the Lebesgue measure m on  $\mathbb{R}^n$ , i.e.,  $\lambda = fm$  for some  $f \in L^1(\mathbb{R}^n)$ , then the formula above reduces to the usual definition of the Fourier transform of f, and

$$\lim_{|\xi| \to \infty} \hat{\lambda}(\xi) = 0.$$

In general, the Fourier transform of a measure need not vanish at infinity. For example, the Fourier transform of  $\delta_0$ , the Dirac measure, is identically equal to 1. However, the decay of the Fourier transform of a measure can be deduced from certain curvature properties of the support of the measure.

Suppose M is a smooth submanifold of  $\mathbb{R}^n$ . By a smooth measure on M we mean a measure of the form  $\mu = \psi \sigma$  where  $\sigma$  is the measure on M induced by the Lebesgue measure on  $\mathbb{R}^n$ , and  $\psi$  is a smooth function on  $\mathbb{R}^n$  whose support intersects M in a compact set.

<sup>2010</sup> Mathematics Subject Classification. 22D10, 22E30, 43A05, 43A80, 53D55.

Key words and phrases. Absolute continuity; Compact operator; Real-analytic submanifold; Twisted convolution.

It is well known that if  $\mu$  is a smooth measure on a hypersurface in  $\mathbb{R}^n$ ,  $n \geq 2$ , whose Gaussian curvature is nonzero everywhere, then

$$\left|\widehat{\mu}(\xi)\right| \le A \left|\xi\right|^{(1-n)/2},$$

where A is a constant independent of  $\xi$  (see e.g. [8, p348, Theorem 1] and [4, Theorem 7.7.14]).

In [5], an analogue of this result for the Weyl transform was proved. Let  $\mathcal{H} = L^2(\mathbb{R}^n)$ , and  $\mathcal{B}(\mathcal{H})$  the set of bounded operators on  $\mathcal{H}$ . If  $f \in L^1(\mathbb{R}^{2n})$ , the Weyl Transform of f is the operator  $W(f) \in \mathcal{B}(\mathcal{H})$  defined by

$$(W(f)\varphi)(t) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x,y) e^{\pi i (x \cdot y + 2y \cdot t)} \varphi(t+x) \, dx \, dy.$$

More generally, if  $\lambda$  is a finite Borel measure on  $\mathbb{R}^{2n}$ , the Weyl Transform of  $\lambda$  is the operator  $W(\lambda) \in \mathcal{B}(\mathcal{H})$  defined by

$$(W(\lambda)\varphi)(t) = \int_{\mathbb{R}^{2n}} e^{\pi i (x \cdot y + 2y \cdot t)} \varphi(t+x) \, d\lambda(x,y).$$

The Weyl transform can be expressed in terms of the canonical representation of the Heisenberg group. Recall that the reduced Heisenberg group G is the set of triples

$$\{(x, y, z) | x, y \in \mathbb{R}^n, z \in \mathbb{C}, |z| = 1\}$$

with multiplication defined by

$$(x, y, z)(x', y', z') = \left(x + x', y + y', zz'e^{\pi i(x \cdot y' - y \cdot x')}\right).$$

According to the Stone-von Neumann Theorem [2, Theorem 1.50], there is a unique irreducible unitary representation  $\rho$  of G such that

$$\rho(0,0,z) = zI.$$

The standard realization of this representation is on the Hilbert space  $\mathcal{H}$  by the action

(1) 
$$(\rho(x, y, z)\varphi)(t) = ze^{\pi i(x \cdot y + 2y \cdot t)}\varphi(t+x).$$

Thus the Weyl transform of  $\lambda$  may be expressed as

(2) 
$$W(\lambda) = \int_{\mathbb{R}^{2n}} \rho(x, y, 1) \, d\lambda(x, y),$$

where the integral is the weak integral as defined in [7, Definition 3.26].

The analogue of the Riemann-Lebesgue lemma for the Weyl transform is the fact that W(f) is a compact operator if  $f \in L^1(\mathbb{R}^{2n})$  (see e.g. [2, Theorem 1.30] and [11, Theorem 1.3.3]). If  $\lambda$  is absolutely continuous with respect to the Lebesgue measure m on  $\mathbb{R}^{2n}$ , i.e.,

 $\mathbf{2}$ 

 $\lambda = fm$  for some  $f \in L^1(\mathbb{R}^{2n})$ , then  $W(\lambda)$  reduces to the usual definition of the Weyl transform of f, and hence  $W(\lambda)$  is a compact operator.

The main result of [5] is the following theorem.

**Theorem 1.1.** Suppose S is a compact connected smooth hypersurface in  $\mathbb{R}^{2n}$ ,  $n \geq 2$ , whose Gaussian curvature is positive everywhere. Let  $\mu$  be a smooth measure on S. Then  $W(\mu)$  is a compact operator. Moreover, when  $n \geq 6$  and p > n,

$$W(\mu) \in S^p(\mathcal{H}),$$

where  $S^p(\mathcal{H})$  denotes the  $p^{\text{th}}$  Schatten class of  $\mathcal{H}$ .

Naturally, the question of what happens if we take a submanifold of arbitrary codimension into consideration arises.

Let M be a smooth *m*-dimensional submanifold of  $\mathbb{R}^n$ ,  $1 \leq m \leq n-1$ , and let  $\mu$  be a smooth measure on M. It is well known that if M is of finite type, i.e., at each point, M has at most a finite order of contact with any affine hyperplane, then

$$\left|\widehat{\mu}(\xi)\right| \le A \left|\xi\right|^{-1/k}$$

where k is the type of M inside the support of  $\psi$  (see [8, p351, Theorem 2]). In particular,

(3) 
$$\lim_{|\xi| \to \infty} \widehat{\mu}(\xi) = 0$$

If we consider M to be a real-analytic submanifold of  $\mathbb{R}^n$ , then the condition of being finite type is equivalent to M not lying in any affine hyperplane.

The main result of this paper is the following theorem, which is an analogue of Equation (3) for the Weyl transform of a smooth measure supported on a real-analytic submanifold of finite type, and is proved in Section 3.

**Theorem 1.2.** Suppose M is a connected real-analytic submanifold of  $\mathbb{R}^{2n}$  which is not contained in an affine hyperplane. Let  $\mu$  be a smooth measure on M. Then  $W(\mu)$  is a compact operator.

In Section 2, we define and study the twisted convolution of finite Borel measures, which is an essential tool required to prove the main result. In Section 4, we prove that if n = 1, then the conclusion of Theorem 1.2 holds for a submanifold of finite type without the additional assumption of real analyticity. This also proves Theorem 1.1 partially for n = 1.

In [5], the quantum translation of an operator was defined, and as an application of Theorem 1.1, it was shown that for  $n \ge 6$ , there exists a non-zero compact operator on  $L^2(\mathbb{R}^n)$  whose quantum translates are linearly dependent. By using a similar argument, as an application of Theorem 1.2, we get the following result.

**Theorem 1.3.** There exists a non-zero compact operator T on  $L^2(\mathbb{R}^n)$  and distinct elements  $(x_1, y_1), \ldots, (x_{4n+1}, y_{4n+1}) \in \mathbb{R}^{2n}$  such that  $\{(x_1, y_1) \cdot T, \ldots, (x_{4n+1}, y_{4n+1}) \cdot T\}$  is a linearly dependent set, where  $(x_i, y_i) \cdot T = \rho(x_i, y_i, 1)T\rho(x_i, y_i, 1)^{-1}$  is the quantum translation of T by  $(x_i, y_i)$ ,  $i = 1, \ldots, 4n + 1$ .

## 2. Twisted convolution

Recall that if  $f, g \in L^1(\mathbb{R}^{2n})$ , the twisted convolution of f and g, denoted by  $f \nmid g$  is

$$f \natural g(x,y) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x - x', y - y') g(x', y') e^{\pi i (x \cdot y' - y \cdot x')} \, dx' \, dy', \quad (x,y) \in \mathbb{R}^{2n}$$

Twisted convolution turns  $L^1(\mathbb{R}^{2n})$  into a non-commutative Banach algebra. It is well known that the Weyl transform is an algebra homomorphism from  $L^1(\mathbb{R}^{2n})$  to  $\mathcal{B}(\mathcal{H})$ , i.e.,  $W(f \natural g) = W(f)W(g)$  (see e.g. [2, p26] and [11, p16]).

**Definition 2.1.** Let  $\mu$  and  $\nu$  be finite Borel measures on  $\mathbb{R}^{2n}$ . The twisted convolution of  $\mu$  and  $\nu$  is the measure on  $\mathbb{R}^{2n}$ , denoted by  $\mu \natural \nu$ , given by

$$\mu \natural \nu(E) = \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \chi_E(x + x', y + y') e^{\pi i (x \cdot y' - y \cdot x')} d\mu(x, y) d\nu(x', y'),$$

where E is a Borel subset of  $\mathbb{R}^{2n}$ .

It follows from Definition 2.1 that if  $f \in C_0(\mathbb{R}^{2n})$ , then

(4) 
$$\int_{\mathbb{R}^{2n}} f(x,y) \, d(\mu \,\natural \, \nu)(x,y) = \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} f(x+x',y+y') e^{\pi i (xy'-yx')} \, d\mu(x,y) \, d\nu(x',y').$$

Let  $f, g \in L^1(\mathbb{R}^{2n})$ . Let *m* denote the Lebesgue measure on  $\mathbb{R}^{2n}$ . Let  $\mu_f = fm$  and  $\mu_g = gm$ . If *E* is a Borel subset of  $\mathbb{R}^{2n}$ , then

$$\begin{split} \mu_{f} \natural \mu_{g}(E) &= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \chi_{E}(x+x',y+y') e^{\pi i (x \cdot y'-y \cdot x')} \, d\mu(x,y) \, d\nu(x',y') \\ &= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \chi_{E}(x+x',y+y') e^{\pi i (x \cdot y'-y \cdot x')} f(x,y) g(x',y') \, dm(x,y) \, dm(x',y') \\ &= \int_{\mathbb{R}^{2n}} \chi_{E}(x,y) \int_{\mathbb{R}^{2n}} e^{\pi i (x \cdot y'-y \cdot x')} f(x-x',y-y') g(x',y') \, dm(x',y') \, dm(x,y) \\ &= \int_{\mathbb{R}^{2n}} \chi_{E}(x,y) (f \natural g)(x,y) \, dm(x,y) \\ &= \mu_{f \natural g}(E). \end{split}$$

Therefore the two definitions of twisted convolution coincide for  $L^1$  functions.

Twisted convolution turns  $M(\mathbb{R}^{2n})$ , the set of finite Borel measures on  $\mathbb{R}^{2n}$ , into a noncommutative Banach algebra. The following theorem shows that the Weyl transform is an algebra homomorphism from  $M(\mathbb{R}^{2n})$  to  $\mathcal{B}(\mathcal{H})$ .

**Theorem 2.2.** Let  $\mu$  and  $\nu$  be finite Borel measures on  $\mathbb{R}^{2n}$ . Then

$$W(\mu \natural \nu) = W(\mu)W(\nu).$$

*Proof.* By Equations (2) and (4),

$$\begin{split} W(\mu \,\natural \, \nu) &= \int_{\mathbb{R}^{2n}} \rho(x, y, 1) \, d(\mu \,\natural \, \nu)(x, y) \\ &= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \rho(x + x', y + y', 1) e^{\pi i (x \cdot y' - y \cdot x')} \, d\mu(x, y) \, d\nu(x', y') \\ &= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \rho(x + x', y + y', e^{\pi i (x \cdot y' - y \cdot x')}) \, d\mu(x, y) \, d\nu(x', y') \quad \text{(by Equation (1))} \\ &= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \rho((x, y, 1)(x', y', 1)) \, d\mu(x, y) \, d\nu(x', y') \\ &= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \rho(x, y, 1) \rho(x', y', 1) \, d\nu(x', y') \, d\mu(x, y) \\ &= \int_{\mathbb{R}^{2n}} \rho(x, y, 1) W(\nu) \, d\mu(x, y) \\ &= W(\mu) W(\nu). \end{split}$$

#### 3. The proof

To prove Theorem 1.2, we need a result analogues to a result of Ragozin about the absolute continuity of the convolution of measures supported on analytic submanifolds of  $\mathbb{R}^n$  (see [6, Theorem 5.1]). In particular, Ragozin proved the absolute continuity of the convolution square of the surface measure on a compact analytic hypersurface of  $\mathbb{R}^n$ . Later, Thangavelu proved that the twisted convolution of the surface measure on a unit sphere in  $\mathbb{R}^{2n}$  with itself is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}^{2n}$  (see [10, Proposition 4.3]).

The following theorem is an analogue of [6, Theorem 5.1] for twisted convolution.

**Theorem 3.1.** Let  $M_1, \ldots, M_k$  be connected real-analytic submanifolds of  $\mathbb{R}^{2n}$  such that for some choice of points  $p_i \in M_i$ ,  $i = 1, \ldots, k$ ,  $T_{p_1}M_1 + \cdots + T_{p_k}M_k = \mathbb{R}^{2n}$ , where  $T_{p_i}M_i$ denotes the tangent space at  $p_i$  of  $M_i$ . If  $\mu_i$  is a smooth measure on  $M_i$ ,  $i = 1, \ldots, k$ , then  $\mu_1 \natural \cdots \natural \mu_k$  is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}^{2n}$ . Let  $\Sigma_k : M_1 \times \cdots \times M_k \to \mathbb{R}^{2n}$  be the map given by  $\Sigma_k(p_1, \ldots, p_k) = p_1 + \cdots + p_k$ . We need the following lemma to prove Theorem 3.1.

**Lemma 3.2.** There exists a smooth function  $\varphi_k : M_1 \times \cdots \times M_k \to \mathbb{C}$  such that, for a Borel set  $E \subseteq \mathbb{R}^{2n}$ ,

$$\mu_1 \natural \cdots \natural \mu_k(E) = (\varphi_k \mu_1 \times \cdots \times \mu_k) (\Sigma_k^{-1}(E)),$$

*i.e.*,  $\mu_1 
arrow \dots 
arrow \mu_k$  is the push-forward of the measure  $\varphi_k \mu_1 \times \dots \times \mu_k$  by  $\Sigma_k$ .

*Proof.* We will prove the result by induction on k.

Let k = 2. Observe that by Definition 2.1,

(5) 
$$\mu_1 \natural \mu_2(E) = (\varphi_2 \, \mu_1 \times \mu_2) \, (\Sigma_2^{-1}(E)),$$

where  $\varphi_2 : \mathbb{R}^{2n} \times \mathbb{R}^{2n} \to \mathbb{C}$  is given by  $\varphi_2((x_1, y_1), (x_2, y_2)) = e^{\pi i (x_1 \cdot y_2 - y_1 \cdot x_2)}$ .

Assume that there exists a function  $\varphi_{k-1}: M_1 \times \cdots \times M_{k-1} \to \mathbb{C}$  such that

$$\mu_1 \natural \cdots \natural \mu_{k-1}(E) = (\varphi_{k-1} \mu_1 \times \cdots \times \mu_{k-1}) (\Sigma_{k-1}^{-1}(E))$$

Then

$$((\mu_{1} \natural \cdots \natural \mu_{k-1}) \natural \mu_{k})(E)$$

$$= \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} \chi_{E}(x + x_{k}, y + y_{k}) e^{\pi i (x \cdot y_{k} - y \cdot x_{k})} d(\mu_{1} \natural \cdots \natural \mu_{k-1})(x, y) d\mu_{k}(x_{k}, y_{k})$$

$$= \int_{\mathbb{R}^{2n}} \cdots \int_{\mathbb{R}^{2n}} \left[ \chi_{E}(x_{1} + \cdots + x_{k}, y_{1} + \cdots + y_{k}) e^{\pi i ((x_{1} + \cdots + x_{k-1}) \cdot y_{k} - (y_{1} + \cdots + y_{k-1}) \cdot x_{k})} \right]$$

$$\varphi_{k-1}((x_{1}, y_{1}), \dots, (x_{k-1}, y_{k-1})) d\mu_{1}(x_{1}, y_{1}) \dots d\mu_{k}(x_{k}, y_{k})$$

$$= (\varphi_{k} \mu_{1} \times \cdots \times \mu_{k}) (\Sigma_{k}^{-1}(E)),$$

where  $\varphi_k : M_1 \times \cdots \times M_k \to \mathbb{C}$  is given by

$$\varphi_k((x_1, y_1), \dots, (x_k, y_k)) = \varphi_{k-1}((x_1, y_1), \dots, (x_{k-1}, y_{k-1}))e^{\pi i((x_1 + \dots + x_{k-1}) \cdot y_k - (y_1 + \dots + y_{k-1}) \cdot x_k)}.$$

Theorem 3.1 now follows by the argument in [6, Theorem 5.1]. Here, we give a more streamlined proof using the coarea formula.

Let  $M_1, \ldots, M_k$  be connected real-analytic submanifolds of  $\mathbb{R}^{2n}$ . Then  $M_1 \times \cdots \times M_k$ is a connected real-analytic manifold. Let  $\tau_{(M_1 \times \cdots \times M_k)}$  denote the Riemannian measure on  $M_1 \times \cdots \times M_k$ .

Observe that  $\Sigma_k$  is an analytic map. Let  $p_i \in M_i$ ,  $i = 1, \ldots, k$  be such that  $T_{p_1}M_1 + \cdots + T_{p_k}M_k = \mathbb{R}^{2n}$ . Then the rank of  $\Sigma_k$  is 2n at the point  $(p_1, \ldots, p_k)$ . Therefore the critical set of  $\Sigma_k$  is a proper analytic subvariety of  $M_1 \times \cdots \times M_k$ , and hence has  $\tau_{(M_1 \times \cdots \times M_k)}$  measure zero.

 $\mathbf{6}$ 

If  $\mu_i$  is a smooth measure on  $M_i$ , i = 1, ..., k, then  $\mu_1 \times \cdots \times \mu_k$  is a smooth measure on  $M_1 \times \cdots \times M_k$ , and so  $\varphi_k \mu_1 \times \cdots \times \mu_k$  is a smooth measure on  $M_1 \times \cdots \times M_k$ .

The proof of Theorem 3.1 now follows from the following lemma.

**Lemma 3.3.** Let M and N be Riemannian manifolds. Let  $\tau$  and  $\nu$  denote the Riemannian measures on M and N respectively. Let  $\mu = \psi \tau$  be a smooth measure on M. Suppose  $f: M \to N$  is a differentiable map. If f is a submersion everywhere except on a set of  $\tau$ -measure zero, then the push-forward of  $\mu$  by f is absolutely continuous with respect to  $\nu$ .

Proof. Let  $Z = \{x \in M \mid f \text{ is not a submersion at } x\}$ . Then  $\tau(Z) = 0$ , and so  $\mu(Z) = 0$ . Let  $\mathcal{J}_f$  denote the normal Jacobian of f, i.e., the absolute value of the determinant of df restricted to the orthogonal complement of its kernel. Then  $\mathcal{J}_f$  is strictly positive on the set of regular points of f, i.e., on  $M \setminus Z$ .

Let  $f_*\mu$  denote the push-forward of  $\mu$  by f. For  $x \in N$ , let  $\sigma_x$  denote the Riemannian measure on the manifold  $f^{-1}(x)$ . Let  $U \subseteq N$  be a Borel set. By the coarea formula (see [1, p159]), we have

$$f_*\mu(U) = \mu(f^{-1}(U))$$
  
=  $\mu(f^{-1}(U) \setminus Z)$   
=  $\int_{f^{-1}(U)\setminus Z} d\mu$   
=  $\int_{f^{-1}(U)\setminus Z} \psi d\tau$   
=  $\int_{U\setminus f(Z)} \int_{f^{-1}(x)} \frac{\psi(y)}{\mathcal{J}_f(y)} d\sigma_x(y) d\nu(x)$ 

i.e.,  $\frac{df_*\mu}{d\nu}(x) = \int_{f^{-1}(x)} \frac{\psi(y)}{\mathcal{J}_f(y)} d\sigma_x(y)$ , and so  $f_*\mu$  is absolutely continuous with respect to  $\nu$ .  $\Box$ 

We need the following lemma to prove Theorem 1.2.

**Lemma 3.4.** Let M be a connected real-analytic submanifold of  $\mathbb{R}^{2n}$ . Assume that M is not contained in any affine hyperplane. Then there exist points  $p_1, \ldots, p_k \in M$  such that

$$T_{p_1}M + \dots + T_{p_k}M = \mathbb{R}^{2n}.$$

*Proof.* Suppose  $\sum_{p \in M} T_p M \neq \mathbb{R}^{2n}$ . Then there exists a hyperplane Y in  $\mathbb{R}^{2n}$  such that  $\sum_{p \in M} T_p M \subseteq Y$ .

Fix  $p \in M$ . Let  $q \in M$ . Since M is connected, there exists a differentiable map  $\xi : [0,1] \to M$  such that  $\xi(0) = p$  and  $\xi(1) = q$ . By the fundamental theorem of calculus,

$$q = p + \int_0^1 \dot{\xi}(t) \, dt.$$

Since  $\dot{\xi}(t) \in Y$  for  $t \in [0, 1]$  and Y is closed, it follows that  $q \in p + Y$ . This implies that M is contained in the affine hyperplane p + Y, which is a contradiction.

Hence  $\sum_{p \in M} T_p M = \mathbb{R}^{2n}$ . By the Steinitz exchange lemma, there exist finitely many points  $p_1, \ldots, p_k \in M$  such that  $T_{p_1}M + \cdots + T_{p_k}M = \mathbb{R}^{2n}$ .

By possibly adding one more point, we can assume that the integer k obtained in Lemma 3.4 is even.

We are now in a position to prove Theorem 1.2. Let  $\widetilde{M} = \{-p \mid p \in M\}$ . Then  $\widetilde{M}$  is a connected real-analytic submanifold of  $\mathbb{R}^{2n}$ . Observe that if  $p \in M$ ,  $T_pM = T_{(-p)}\widetilde{M}$ . Therefore, by Lemma 3.4,

(6) 
$$T_{p_1}M + T_{(-p_2)}\widetilde{M} + \dots + T_{p_{k-1}}M + T_{(-p_k)}\widetilde{M} = \mathbb{R}^{2n}$$

Let  $\mu$  be a smooth measure on M. Let  $\tilde{\mu}$  denote the push-forward of  $\mu$  by the map which sends  $p \in M$  to -p. Then, by Theorem 3.1 and Equation (6), it follows that  $(\mu \natural \tilde{\mu})^{k/2}$  is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}^{2n}$ . Therefore  $W((\mu \natural \tilde{\mu})^{k/2})$  is a compact operator.

Observe that  $W(\tilde{\mu}) = W(\mu)^*$ , where  $W(\mu)^*$  is the adjoint of the operator  $W(\mu)$ . By Theorem 2.2,

$$W((\mu \natural \widetilde{\mu})^{k/2}) = (W(\mu)W(\widetilde{\mu}))^{k/2} = (W(\mu)W(\mu)^*)^{k/2}$$

Since  $W(\mu)W(\mu)^*$  is self-adjoint and  $(W(\mu)W(\mu)^*)^{k/2}$  is compact, it follows that  $W(\mu)W(\mu)^*$  is compact. By the polar decomposition, it follows that  $W(\mu)$  is compact.

# 4. Curve in $\mathbb{R}^2$

In this section, we prove the following result, which states that if n = 1, the assumption of real-analyticity can be removed from Theorem 1.2.

**Theorem 4.1.** Suppose  $M \subseteq \mathbb{R}^2$  is a connected smooth hypersurface of finite type. Let  $\mu$  be a smooth measure on M. Then  $W(\mu)$  is a compact operator.

We need the following lemma to prove this theorem.

**Lemma 4.2.** Let  $\gamma : [a,b] \to \mathbb{R}^2$  be a finite type unit-speed simple smooth curve. Let  $\delta : [c,d] \to \mathbb{R}^2$  be a unit-speed simple smooth curve. Let  $\mu$  and  $\nu$  be smooth measures on  $\operatorname{Im}(\gamma)$  and  $\operatorname{Im}(\delta)$  respectively. Then  $\mu \natural \nu$  is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}^2$ .

*Proof.* Let  $E \subseteq \mathbb{R}^2$  be a Borel set. Then, by Equation (5),

$$\mu \natural \nu(E) = (\varphi_2 \mu \times \nu) (\Sigma_2^{-1}(E)),$$

### THE WEYL TRANSFORM OF A SMOOTH MEASURE ON A REAL-ANALYTIC SUBMANIFOLD 9

where  $\Sigma_2 : \operatorname{Im}(\gamma) \times \operatorname{Im}(\delta) \to \mathbb{R}^2$  is the map given by  $\Sigma_2(\gamma(s), \delta(t)) = \gamma(s) + \delta(t)$ . Define  $S : [a, b] \times [c, d] \to \mathbb{R}^2$  by

$$S(s,t) = \gamma(s) + \delta(t).$$

Let  $T = \{(s,t) \in [a,b] \times [c,d] \mid \dot{\gamma}(s) = \pm \dot{\delta}(t)\}$ . Observe that T is the critical set of S. We claim that T has area zero. Suppose T has a positive area. Then, by Fubini's theorem, there exists  $t \in [c,d]$  such that the set  $Z = \{s \in [a,b] \mid \dot{\gamma}(s) = \pm \dot{\delta}(t)\}$  has positive length. By the Lebesgue differentiation theorem [3, Theorem 3.21], there exists  $s' \in [a,b]$  such that s' is a point of density of Z. Then there exists distinct  $s_j \in Z$  such that the sequence  $\{s_j\}$  converges to s', and so the sequence  $\{\dot{\gamma}(s_j)\}$  converges to  $\dot{\gamma}(s')$ . Let  $\vec{v}$  be a unit vector in  $\mathbb{R}^2$  perpendicular to  $\dot{\delta}(t)$ . Then

$$\langle \dot{\gamma}(s_j), \vec{v} \rangle = \langle \pm \delta(t), \vec{v} \rangle = 0, \quad j = 1, 2, \dots$$

Hence, all the coefficients in the Taylor expansion of  $\langle \dot{\gamma}(s), \vec{v} \rangle$  about s = s' are zero. This contradicts the fact that  $\gamma$  is a curve of finite type. Therefore T has area zero.

Since  $\operatorname{Im}(\gamma)$  and  $\operatorname{Im}(\delta)$  are smooth submanifolds of  $\mathbb{R}^2$ , it follows that  $\operatorname{Im}(\gamma) \times \operatorname{Im}(\delta)$  is a smooth submanifold of  $\mathbb{R}^4$ . Let  $\tau$  denote the Riemannian measure on  $\operatorname{Im}(\gamma) \times \operatorname{Im}(\delta)$ . Since  $\gamma \times \delta$  is a smooth map and the set T has area zero, it follows that  $(\gamma \times \delta)(T)$  has  $\tau$ -measure zero.

Observe that  $(\gamma \times \delta)(T)$  is the critical set of  $\Sigma_2$ . The proof now follows from Lemma 3.3.

We now prove Theorem 4.1. Suppose  $M \subseteq \mathbb{R}^2$  is a hypersurface of finite type. Let  $\mu$  be a smooth measure on M. Let  $\tilde{\mu}$  denote the push-forward of  $\mu$  by the map which sends  $p \in M$ to -p. It follows from Lemma 4.2 that  $\mu \natural \tilde{\mu}$  is absolutely continuous with respect to the Lebesgue measure on  $\mathbb{R}^2$ . Therefore  $W(\mu \natural \tilde{\mu})$  is compact. Observe that  $W(\tilde{\mu}) = W(\mu)^*$ . By Theorem 2.2,

$$W(\mu \natural \widetilde{\mu}) = W(\mu)W(\widetilde{\mu}) = W(\mu)W(\mu)^*.$$

Since  $W(\mu)W(\mu)^*$  is compact, it follows by the polar decomposition that  $W(\mu)$  is compact.

## References

- Isaac Chavel, *Riemannian geometry*, second ed., Cambridge Studies in Advanced Mathematics, vol. 98, Cambridge University Press, Cambridge, 2006, A modern introduction. MR 2229062
- Gerald B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, vol. 122, Princeton University Press, Princeton, NJ, 1989. MR 983366
- \_\_\_\_\_, Real analysis, second ed., Pure and Applied Mathematics (New York), John Wiley & Sons, Inc., New York, 1999, Modern techniques and their applications, A Wiley-Interscience Publication. MR 1681462

#### MANSI MISHRA AND M. K. VEMURI

- Lars Hörmander, The analysis of linear partial differential operators. I, second ed., Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 256, Springer-Verlag, Berlin, 1990, Distribution theory and Fourier analysis. MR 1065993
- Mansi Mishra and M. K. Vemuri, *The Weyl transform of a measure*, Proc. Indian Acad. Sci. Math. Sci. 133 (2023), no. 2, Paper No. 29, 11. MR 4642690
- David L. Ragozin, Rotation invariant measure algebras on Euclidean space, Indiana Univ. Math. J. 23 (1973/74), 1139–1154. MR 338688
- Walter Rudin, *Functional analysis*, second ed., International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991. MR 1157815
- 8. Elias M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals,* Princeton Mathematical Series, vol. 43, Princeton University Press, Princeton, NJ, 1993, With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III. MR 1232192
- Elias M. Stein and Guido Weiss, Introduction to Fourier analysis on Euclidean spaces, Princeton Mathematical Series, vol. No. 32, Princeton University Press, Princeton, NJ, 1971. MR 304972
- 10. Sundaram Thangavelu, Spherical means on the Heisenberg group and a restriction theorem for the symplectic Fourier transform, Rev. Mat. Iberoamericana 7 (1991), no. 2, 135–155. MR 1133375
- 11. \_\_\_\_\_, Harmonic analysis on the Heisenberg group, Progress in Mathematics, vol. 159, Birkhäuser Boston, Inc., Boston, MA, 1998. MR 1633042

DEPARTMENT OF MATHEMATICAL SCIENCES, IIT(BHU), VARANASI 221005. Email address: mansimishra1312@gmail.com