

On crossed modules of pre-Lie algebras

Xuedan Luo

*Chern Institute of Mathematics, Nankai University,
Tianjin, 300071, China*

Email: 1120230015@mail.nankai.edu.cn

Abstract

Given a Lie algebra \mathfrak{g} and a \mathfrak{g} -module V , it is due to Gerstenhaber that there is an isomorphism between $H^3(\mathfrak{g}, V)$ and the group of equivalence classes of crossed modules with kernel V and cokernel \mathfrak{g} . The goal of this article is to obtain this result for pre-Lie algebras and their modules.

1 Introduction

The low degree cohomologies of Lie algebras have some well-known applications. The 1-cocycles with respect to the adjoint representation define the derivations. Let \mathfrak{g} be a Lie algebra and V be a \mathfrak{g} -module. Given a 2-cocycle, one can associate $\mathfrak{g} \oplus V$ with a Lie algebra structure to get an exact sequence:

$$0 \rightarrow V \rightarrow \mathfrak{g} \oplus V \rightarrow \mathfrak{g} \rightarrow 0,$$

which is called an abelian extension of \mathfrak{g} by V . This association induces an bijection between $H^2(\mathfrak{g}, V)$ and the set of equivalence classes of abelian extensions, so the second cohomology classifies the abelian extensions.

The notion of crossed module was proposed by Whitehead [21] in the context of homotopy theory, which consists of a homomorphism $\mu : \mathfrak{m} \rightarrow \mathfrak{n}$ of Lie algebras and a compatible action of \mathfrak{n} on \mathfrak{m} . We attribute to Gerstenhaber [13, 14] the theorem stating that the group of equivalence classes of crossed modules with kernel V and cokernel \mathfrak{g} , is isomorphic to $H^3(\mathfrak{g}, V)$. One may express a crossed module with kernel V and cokernel \mathfrak{g} by the following exact sequence:

$$0 \rightarrow V \rightarrow \mathfrak{m} \xrightarrow{\mu} \mathfrak{n} \rightarrow \mathfrak{g} \rightarrow 0,$$

which is called a crossed module extension of \mathfrak{g} by V . So roughly, we may restate the result of Gerstenhaber by: the third cohomology classifies the crossed module extensions. The original proof is quite abstract using the *Ext* functor, and we recommend [20] for a more constructive and algebraic proof.

The goal of this article is to get the version of this result in pre-Lie algebras. For versions of this result in Lie 2-algebras, Lie-Rinehart algebras, pre-Lie Rinehart algebras and zinbiel algebras, see [17, 8, 5, 22].

The notion of pre-Lie algebra was introduced by Gerstenhaber [14] in the study of deformations of associative algebras. A pre-Lie algebra is a vector space \mathfrak{g} equipped with a map $\circ : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ such that the associator is left-symmetry, i.e., $(x, y, z) = (y, x, z)$, where $(x, y, z) := (x \circ y) \circ z - x \circ (y \circ z)$ for $x, y, z \in \mathfrak{g}$. It arises naturally from many areas, such as the left-invariant affine structures on Lie groups, complex structures on Lie algebras and so on, see [1, 2]. Its name is due to its closed relation to the Lie algebras, i.e., the commutator defines a Lie algebra on the underlying space. Moreover, the left multiplication of pre-Lie algebra gives a representation of the commutator Lie algebra.

The paper is organized as follows. Section 2 constitutes sufficient preparation. Section 3 states and proves our main theorem (Theorem 3.2) and get a corollary (Corollary 3.7). Section 4 summarizes some relations between crossed modules of pre-Lie algebras and crossed modules of Lie, Rota-Baxter Lie, and dendriform algebras.

2 Preliminaries

A **pre-Lie algebra** (\mathfrak{g}, \circ) is a vector space \mathfrak{g} equipped with a bilinear map $\circ : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ such that

$$(x \circ y) \circ z - x \circ (y \circ z) = (y \circ x) \circ z - y \circ (x \circ z), \quad \forall x, y, z \in \mathfrak{g}. \quad (1)$$

Let (\mathfrak{g}, \circ) be a pre-Lie algebra. Then there is a Lie algebra structure $[\cdot, \cdot] : \mathfrak{g} \wedge \mathfrak{g} \rightarrow \mathfrak{g}$ on \mathfrak{g} defined by $[x, y] := x \circ y - y \circ x$. We call $(\mathfrak{g}, [\cdot, \cdot])$ the sub-adjacent Lie algebra of (\mathfrak{g}, \circ) and denote it by \mathfrak{g}^c .

Definition 2.1. A **representation** of a pre-Lie algebra (\mathfrak{g}, \circ) is a pair (V, \circ_l, \circ_r) , where V is a vector space, $\circ_l : \mathfrak{g} \otimes V \rightarrow V$ is a representation of the sub-adjacent Lie algebra \mathfrak{g}^c on V , and $\circ_r : V \otimes \mathfrak{g} \rightarrow V$ is a bilinear operation satisfying

$$(x \circ_l u) \circ_r y - x \circ_l (u \circ_r y) = (u \circ_r x) \circ_r y - u \circ_r (x \circ_l y), \quad \forall x, y \in \mathfrak{g}, u \in V.$$

In this case, we also say that V is a \mathfrak{g} -module. There is a natural representation $(\mathfrak{g}, \circ_L, \circ_R)$ of \mathfrak{g} on itself (or on its two-sided ideal R , i.e., a subspace such that $R \circ \mathfrak{g} \subset R$, $\mathfrak{g} \circ R \subset R$) defined by $x \circ_L y = x \circ y$, $y \circ_R x = y \circ x$.

Definition 2.2. Given two pre-Lie algebras (\mathfrak{m}, \circ_m) and (\mathfrak{n}, \circ_n) , an **action** of \mathfrak{n} on \mathfrak{m} is a representation (\circ_l, \circ_r) of \mathfrak{n} on \mathfrak{m} satisfying 2 more equalities:

$$\begin{aligned} (x \circ_l u) \circ_m v - x \circ_l (u \circ_m v) &= (u \circ_r x) \circ_m v - u \circ_m (x \circ_l v), \\ (u \circ_m v) \circ_r x - u \circ_m (v \circ_r x) &= (v \circ_m u) \circ_r x - v \circ_m (u \circ_r x), \end{aligned}$$

for all $x \in \mathfrak{n}, u, v \in \mathfrak{m}$.

We see that the action reduces to a representation when the pre-Lie structure on \mathfrak{m} is trivial.

The cohomology of pre-Lie algebra (\mathfrak{g}, \circ) with a representation (V, \circ_l, \circ_r) is given in [11]. The set of n -cochains is

$$C^n(\mathfrak{g}, V) = \text{Hom}(\wedge^{n-1} \mathfrak{g} \otimes \mathfrak{g}, V), \quad n \geq 1.$$

The coboundary operator $d : C^n(\mathfrak{g}, V) \rightarrow C^{n+1}(\mathfrak{g}, V)$ is defined by

$$\begin{aligned} (df)(x_1, \dots, x_{n+1}) &= \sum_{i=1}^n (-1)^{i+1} x_i \circ_l f(x_1, \dots, \hat{x}_i, \dots, x_{n+1}) \\ &\quad + \sum_{i=1}^n (-1)^{i+1} f(x_1, \dots, \hat{x}_i, \dots, x_n, x_i) \circ_r x_{n+1} \\ &\quad - \sum_{i=1}^n (-1)^{i+1} f(x_1, \dots, \hat{x}_i, \dots, x_n, x_i \circ x_{n+1}) \\ &\quad + \sum_{1 \leq i < j \leq n} (-1)^{i+j} f([x_i, x_j], x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_{n+1}), \end{aligned}$$

for $x_1, \dots, x_{n+1} \in \mathfrak{g}$, $f \in C^n(\mathfrak{g}, V)$. Here $[x_i, x_j] := x_i \circ x_j - x_j \circ x_i$. In particular, for $n = 1$, $(df)(x, y) = x \circ_l f(y) + f(x) \circ_r y - f(x \circ y)$ and for $n = 2$,

$$\begin{aligned} (df)(x, y, z) &= x \circ_l f(y, z) - y \circ_l f(x, z) + f(y, x) \circ_r z - f(x, y) \circ_r z \\ &\quad - f(y, x \circ z) + f(x, y \circ z) - f([x, y], z). \end{aligned}$$

Definition 2.3. Let (\mathfrak{m}, \circ_m) , (\mathfrak{n}, \circ_n) be pre-Lie algebras with an action (\circ_l, \circ_r) of \mathfrak{n} on \mathfrak{m} . A **crossed module** is a homomorphism $\mu : \mathfrak{m} \rightarrow \mathfrak{n}$ of pre-Lie algebras satisfying

$$\begin{aligned} \mu(u \circ_r x) &= \mu(u) \circ_n x, & \mu(x \circ_l u) &= x \circ_n \mu(u), \\ \mu(u) \circ_l v &= u \circ_m v = u \circ_r \mu(v), & \forall x \in \mathfrak{n}, u, v \in \mathfrak{m}, \end{aligned}$$

denoted by $(\mathfrak{m} \xrightarrow{\mu} \mathfrak{n}, \circ_l, \circ_r)$.

Example 2.4. 1. Let (\mathfrak{n}, \circ) be a pre-Lie algebra, R be its two-sided ideal and i be the inclusion. Then $(R \xrightarrow{i} \mathfrak{n}, \circ_L, \circ_R)$ is a crossed module. In particular, $(\mathfrak{n} \xrightarrow{id} \mathfrak{n}, \circ_L, \circ_R)$ is a crossed module.

2. Let $f : \mathfrak{m} \rightarrow \mathfrak{n}$ be a homomorphism of pre-Lie algebras. Then $\ker(f)$ is a two-sided ideal of (\mathfrak{m}, \circ_m) , thus by the example above, $(\ker(f) \xrightarrow{i} \mathfrak{m}, \circ_L, \circ_R)$ is a crossed module.

3. Let (V, \circ_l, \circ_r) be a representation over pre-Lie algebra \mathfrak{g} . Endow V with the trivial pre-Lie structure, then $(V \xrightarrow{0} \mathfrak{g}, \circ_l, \circ_r)$ is a crossed module.

Let **preLie** be the category whose objects are pre-Lie algebras, morphisms are homomorphisms of pre-Lie algebras.

Definition 2.5. Let (\mathfrak{g}, \circ) be a pre-Lie algebra, and let (V, \circ_l, \circ_r) be a representation of it. A **crossed module extension** of \mathfrak{g} by V is an exact sequence in category **preLie**:

$$0 \rightarrow V \xrightarrow{i} \mathfrak{m} \xrightarrow{\mu} \mathfrak{n} \xrightarrow{\pi} \mathfrak{g} \rightarrow 0 \quad (2)$$

such that $\mathfrak{m} \xrightarrow{\mu} \mathfrak{n}$ is a crossed module.

Remark 2.6. Given an exact sequence (2) such that $\mathfrak{m} \xrightarrow{\mu} \mathfrak{n}$ is a crossed module, the action of \mathfrak{n} on \mathfrak{m} uniquely induces a representation of \mathfrak{g} on V , and we denote them by the same notation (\circ_l, \circ_r) . Choose any linear section $\rho : \mathfrak{g} \rightarrow \mathfrak{n}$ of π such that $\pi\rho = id$, this representation is given by

$$i(x \circ_l u) := \rho(x) \circ_l i(u), \quad i(u \circ_r x) := i(u) \circ_r \rho(x), \quad \forall x \in \mathfrak{g}, u \in V.$$

It is independent of the choice of ρ because if ρ_1 and ρ_2 are two sections of π , then $(\rho_1 - \rho_2) \in \ker \pi = im\mu$, thus $i(u) \circ_r (\rho_1 - \rho_2)(x) = i(u) \circ_r \mu(m)$ for some $m \in \mathfrak{m}$. By the crossed module properties, this equals $i(u) \circ_m m$, which vanishes since $i(V)$ is the center of \mathfrak{m} (say $i(v) \circ_m m = \mu(i(v)) \circ m = 0, m \in \mathfrak{m}, v \in V$). Similar argument applies to $\rho(x) \circ_l i(u)$.

Example 2.7. Given a crossed module $\mathfrak{m} \xrightarrow{\mu} \mathfrak{n}$, we get a natural crossed module extension (2), where $\mathfrak{g} = coker(\mu) = \mathfrak{n}/im\mu$ and $V = ker(\mu)$ as vector spaces. By the crossed module properties, $im\mu$ is an ideal of \mathfrak{n} , therefore \mathfrak{g} has an induced pre-Lie structure from \mathfrak{n} . By Remark 2.6, V is indeed a representation of \mathfrak{g} .

Definition 2.8. Two crossed module extensions of \mathfrak{g} by V are **elementary equivalent**, denoted by $E \leftrightarrow E'$, if there exists homomorphisms r and s of pre-Lie algebras such that

- the following diagram commutes:

$$\begin{array}{ccccccccc} E : & 0 & \longrightarrow & V & \xrightarrow{i} & \mathfrak{m} & \xrightarrow{\mu} & \mathfrak{n} & \xrightarrow{\pi} & \mathfrak{g} & \longrightarrow & 0 \\ & & & \downarrow id & & \downarrow r & & \downarrow s & & \downarrow id & & \\ E' : & 0 & \longrightarrow & V & \xrightarrow{i'} & \mathfrak{m}' & \xrightarrow{\mu'} & \mathfrak{n}' & \xrightarrow{\pi'} & \mathfrak{g} & \longrightarrow & 0; \end{array}$$

- they respects the actions in the crossed modules. That is, $r(u \circ_r x) = r(u) \circ_r s(x)$ and $r(x \circ_l u) = s(x) \circ_l r(u)$ for $x \in \mathfrak{n}, u \in \mathfrak{m}$.

It is easy to see that the relation \hookrightarrow is not symmetric for $n \geq 2$. However, it generates an equivalence relation. We define crossed module extensions E and E' to be **equivalent**, denoted by $E \sim E'$, if there exists a chain E_1, \dots, E_k such that

$$E \hookrightarrow E_1 \leftrightarrow E_2 \hookrightarrow \dots \hookrightarrow E_k \leftrightarrow E'.$$

The set of equivalence classes of crossed module extensions of \mathfrak{g} by V , which induce the given representation of \mathfrak{g} on V , is denoted by $CExt(\mathfrak{g}, V)$.

Remark 2.9. We remark that if two extensions are elementary equivalent, then they are equivalent. Suppose that $E \hookrightarrow E'$, then there exists a crossed module extension

$$E_1 : 0 \rightarrow V \rightarrow \mathfrak{m}/ker(r) \xrightarrow{\tilde{\mu}} \mathfrak{n}/ker(s) \rightarrow \mathfrak{g} \rightarrow 0,$$

where $\mathfrak{m}/ker(r) \xrightarrow{\tilde{\mu}} \mathfrak{n}/ker(s)$ is a crossed module with the induced homomorphism $\tilde{\mu}$ and action. We have $E \sim E'$ because $E \hookrightarrow E_1$ through the projections and $E_1 \leftrightarrow E'$ through the isomorphisms \tilde{r}, \tilde{s} induced by r, s .

We check that $ker\tilde{\mu} = V$ and $coker\tilde{\mu} = \mathfrak{g}$, i.e., $ker\tilde{\mu} = \tilde{r}^{-1}ker(\mu') = \tilde{r}^{-1}V = V$ since \tilde{r} is an isomorphism with $\tilde{r}|_V = id_V$, $coker\tilde{\mu} = (\mathfrak{n}/ker(s))/\tilde{\mu}(\mathfrak{m}/ker(r)) = (\mathfrak{n}/ker(s))/(\mu\mathfrak{m}/ker(s)) \cong \mathfrak{n}/\mu\mathfrak{m} = \mathfrak{g}$.

3 Classification of crossed modules via H^3

In this section, we prove our main theorem (Theorem 3.2), which says crossed modules of pre-Lie algebras with kernel V and cokernel \mathfrak{g} can be classified by $H^3(\mathfrak{g}, V)$. As an immediate consequence, we obtain Corollary 3.7, which shows the relation between crossed module extension of pre-Lie algebra and abelian extension of its sub-adjacent Lie algebra.

Lemma 3.1. Let F be a free pre-Lie algebra, V be an F -module. Then $H^n(F, V) = 0$ for $n \geq 2$.

Proof. Denote by F^c the sub-adjacent Lie algebra of F . The F -module (V, \circ_l, \circ_r) can be viewed as an F^c -module via $[X, v] := X \circ_l v - v \circ_r X$ for $X \in F^c, v \in V$. Then $\text{Hom}(F^c, V)$ is also an F^c -module with the representation $\rho : F^c \times \text{Hom}(F^c, V) \rightarrow \text{Hom}(F^c, V)$ given by $\rho(X)f(Y) := [X, f(Y)] - f([X, Y]_{F^c})$, for $X, Y \in F^c, f \in \text{Hom}(F^c, V)$. By Theorem 3.4 of [11], we have

$$H_{preLie}^n(F, V) \cong H_{Lie}^{n-1}(F^c, \text{Hom}(F^c, V)), \quad n \geq 1.$$

By Corollary 6.3 of [7], the sub-adjacent Lie algebra of free pre-Lie algebra is also free, thus F^c is free. Now, as a standard result in homological algebras (see [16]),

$$H_{Lie}^n(F^c, \text{Hom}(F^c, V)) = 0, \quad n \geq 2.$$

Thus $H_{preLie}^n(F, V) = 0$ for $n \geq 3$. For $n = 2$, let $Ext(F, V)$ be the set of equivalence class of abelian extensions of F by V , which is zero since F is free. By the fact that $H_{preLie}^2(F, V) \cong Ext(F, V)$, we obtain $H_{preLie}^2(F, V) = 0$. The proof is complete. \square

Theorem 3.2. Let $(\mathfrak{g}, \circ_{\mathfrak{g}})$ be a pre-Lie algebra and let (V, \circ_l, \circ_r) be a representation of \mathfrak{g} . There is a bijection between $CExt(\mathfrak{g}, V)$ and $H^3(\mathfrak{g}, V)$.

Proof. Since the proof is long, we only make precise here how to construct the bijection and its inverse, and details will be left to Lemma 3.3 to 3.6.

- We construct $T : CExt(\mathfrak{g}, V) \rightarrow H^3(\mathfrak{g}, V)$ as follows. Given a crossed module extension which induces the given representation of \mathfrak{g} on V :

$$0 \rightarrow V \xrightarrow{i} \mathfrak{m} \xrightarrow{\mu} \mathfrak{n} \xrightarrow{\pi} \mathfrak{g} \rightarrow 0, \quad (3)$$

where the action of \mathfrak{n} on \mathfrak{m} is denoted by $(\bar{\circ}_l, \bar{\circ}_r)$. The first step is to take a linear section ρ of π such that $\pi\rho = id$, and to calculate the default of ρ to be a pre-Lie homomorphism. That is, $\alpha(x, y) := \rho(x) \circ_{\mathfrak{n}} \rho(y) - \rho(x \circ_{\mathfrak{g}} y)$ for $x, y \in \mathfrak{g}$. Since π is a pre-Lie homomorphism, we have $\pi\alpha(x, y) = 0$. By the exactness of sequence (3), there exists $\beta \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{m})$ such that $\mu\beta = \alpha$. Choosing a linear section σ of μ such that $\mu\sigma = id$, we can take β as $\sigma\alpha$.

Next, we define $\theta \in C^3(\mathfrak{g}, \mathfrak{m})$ by

$$\begin{aligned} \theta(x, y, z) &= \rho(x)\bar{\circ}_l\beta(y, z) - \rho(y)\bar{\circ}_l\beta(x, z) + \beta(y, x)\bar{\circ}_r\rho(z) - \beta(x, y)\bar{\circ}_r\rho(z) \\ &\quad - \beta(y, x \circ_{\mathfrak{g}} z) + \beta(x, y \circ_{\mathfrak{g}} z) - \beta([x, y], z), \end{aligned} \quad (4)$$

where $[x, y] = x \circ_{\mathfrak{g}} y - y \circ_{\mathfrak{g}} x$. Despite θ has the form of $d\beta$, we can not say $\theta = d\beta$ because the map $\mathfrak{g} \otimes \mathfrak{m} \rightarrow \mathfrak{m}, (x, m) \mapsto \rho(x) \circ m$ is not a representation in general. But in case $\beta \in \text{Hom}(\mathfrak{g}^{\otimes 2}, i(V))$, this is just the given representation of \mathfrak{g} on V .

Finally, by the crossed module properties and $\alpha = \mu\beta$, we compute $\mu\theta = 0$, thus $\theta \in C^3(\mathfrak{g}, V)$. It is straightforward to check that θ is a 3-cocycle, thus we can define $T : CExt(\mathfrak{g}, V) \rightarrow H^3(\mathfrak{g}, V)$ by mapping the equivalence class of extension (3) to $[\theta]$.

- Conversely, the inverse map $S : H^3(\mathfrak{g}, V) \rightarrow CExt(\mathfrak{g}, V)$ is constructed by the following procedures. First, let F be the free pre-Lie algebra generated by \mathfrak{g} (see [11]), then we get an exact sequence:

$$0 \rightarrow K \xrightarrow{i} F \xrightarrow{\pi} \mathfrak{g} \rightarrow 0, \quad (5)$$

where π is the projection, $K = \ker(\pi)$ and i is the inclusion.

Second, π induces a map $\pi^* : C^n(\mathfrak{g}, V) \rightarrow C^n(F, V) (n \geq 1)$ by $f \mapsto f\pi$. Regarding the \mathfrak{g} -module V as an F -module and also trivial K -module by $X \circ u := \pi(X) \circ u$ for $X \in F, u \in V$, then π^* is a cochain map of complexes. That is, the following diagram commutes:

$$\begin{array}{ccccccc} \dots & \longrightarrow & C^2(\mathfrak{g}, V) & \xrightarrow{d} & C^3(\mathfrak{g}, V) & \xrightarrow{d} & C^4(\mathfrak{g}, V) & \longrightarrow & \dots \\ & & \pi^* \downarrow & & \pi^* \downarrow & & \pi^* \downarrow & & \\ \dots & \longrightarrow & C^2(F, V) & \xrightarrow{d} & C^3(F, V) & \xrightarrow{d} & C^4(F, V) & \longrightarrow & \dots \end{array} \quad (6)$$

Likewise, the inclusion $i : K \rightarrow F$ also induces a cochain map $i^* : C^n(F, V) \rightarrow C^n(K, V)$. For any 3-cocycle $\theta \in C^3(\mathfrak{g}, V)$, $\pi^*\theta$ is a 3-cocycle because $d\pi^*\theta =$

$\pi^*d\theta = 0$. By Lemma 3.1, $H^n(F, V) = 0$ ($n > 1$) for free pre-Lie algebra F , there exists an $\tilde{\alpha} \in C^2(F, V)$ such that $d\tilde{\alpha} = \pi^*\theta$. We thus get a map $\alpha := i^*\tilde{\alpha} \in C^2(K, V)$, which is a 2-cocycle because $d\alpha = di^*\tilde{\alpha} = i^*d\tilde{\alpha} = i^*\pi^*\theta = 0$.

Now, let $\mathfrak{m} = K \oplus V$ as vector space and define an operation $\circ_{\mathfrak{m}} : \mathfrak{m} \otimes \mathfrak{m} \rightarrow \mathfrak{m}$ by

$$(X + u) \circ (Y + v) := X \circ_F Y + \alpha(X, Y), \quad (7)$$

for $X, Y \in K, u, v \in V$. Since α is a 2-cocycle, $(\mathfrak{m}, \circ_{\mathfrak{m}})$ is a pre-Lie algebra. We thus get an exact sequence of pre-Lie algebras:

$$0 \rightarrow V \xrightarrow{i} \mathfrak{m} \xrightarrow{p} K \rightarrow 0, \quad (8)$$

where i is the inclusion and p is the projection.

Finally, splicing (8) and (5) together, we get a crossed module extension:

$$0 \rightarrow V \xrightarrow{i} \mathfrak{m} \xrightarrow{\mu} F \xrightarrow{\pi} \mathfrak{g} \rightarrow 0, \quad (9)$$

where $\mu(X + u) = X, X + u \in K \oplus V = \mathfrak{m}$. The action (\bar{o}_l, \bar{o}_r) of F on \mathfrak{m} is given by

$$\begin{aligned} X \circ (Y + v) &:= X \circ_F Y + \tilde{\alpha}(X, Y) + X \circ_l v, \\ (Y + v) \circ X &:= Y \circ_F X + \tilde{\alpha}(Y, X) + v \circ_r X, \end{aligned} \quad (10)$$

for $X \in F, Y + v \in K \oplus V = \mathfrak{m}$. This is an action because $d\tilde{\alpha}(X, Y, Z) = \pi^*\theta(X, Y, Z) = \theta(\pi X, \pi Y, \pi Z) = 0$ when one variable is in $K = \ker \pi$, and K acts trivially on V . We define the map $S : H^3(\mathfrak{g}, V) \rightarrow CExt(\mathfrak{g}, V)$ by mapping $[\theta]$ to equivalence class of crossed module extension (9).

• Lemma 3.3 to 3.6 below will show that T and S are well-defined and inverse to each other. Therefore, T is a bijection and the proof is complete. \square

Lemma 3.3. The above-mentioned $T : CExt(\mathfrak{g}, V) \rightarrow H^3(\mathfrak{g}, V)$ is well-defined.

Proof. We follow the notations in the proof of Theorem 3.2. We shall check that $[\theta]$ is independent of the choice of sections σ and ρ , and that equivalent extensions lead to 3-cocycles in the same cohomology class.

• For two sections σ and σ' of μ , we have $\mu(\sigma - \sigma') = 0$, which implies $\sigma - \sigma' \in \text{Hom}(\mathfrak{n}, V)$ and thus $\beta - \beta' = (\sigma - \sigma')\alpha \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, V)$. So $\theta - \theta' = d^2(\beta - \beta')$ and we have $[\theta - \theta'] = 0$.

• To check that $[\theta]$ is independent of the choice of ρ , let ρ and ρ' be two sections of π , then they induce $\beta, \beta' \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{m})$, and thus $\theta, \theta' \in C^3(\mathfrak{g}, V)$. Since $\pi(\rho - \rho') = 0$ and (3) is exact, there exists $h \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{m})$ such that $\rho - \rho' = \mu h$. To use this relation, we would like to express $\theta - \theta'$ in terms of $\rho - \rho'$.

Explicitly, since

$$\begin{aligned}
\rho(x)\bar{\sigma}_l\beta(y, z) - \rho'(x)\bar{\sigma}_l\beta'(y, z) &= (\rho - \rho')(x)\bar{\sigma}_l\beta(y, z) + \rho'(x)\bar{\sigma}_l(\beta - \beta')(y, z) \\
&= \mu h(x)\bar{\sigma}_l\sigma\alpha(y, z) + \rho'(x)\bar{\sigma}_l(\beta - \beta')(y, z) \\
&= h(x)\bar{\sigma}_r\mu\sigma\alpha(y, z) + \rho'(x)\bar{\sigma}_l(\beta - \beta')(y, z) \\
&= h(x)\bar{\sigma}_r\alpha(y, z) + \rho'(x)\bar{\sigma}_l(\beta - \beta')(y, z) \\
&= h(x)\bar{\sigma}_r(\rho(y) \circ_{\mathfrak{n}} \rho(z) - \rho(y \circ_{\mathfrak{g}} z)) \\
&\quad + \rho'(x)\bar{\sigma}_l(\beta - \beta')(y, z),
\end{aligned}$$

and similarly

$$\begin{aligned}
\beta(y, x)\bar{\sigma}_r\rho(z) - \beta'(y, x)\bar{\sigma}_r\rho'(z) &= (\rho(y) \circ_{\mathfrak{n}} \rho(x) - \rho(y \circ_{\mathfrak{g}} x)) \circ h(z) \\
&\quad + (\beta - \beta')(y, x)\bar{\sigma}_r\rho'(z),
\end{aligned}$$

we obtain

$$\begin{aligned}
(\theta - \theta')(x, y, z) &= \rho(x)\bar{\sigma}_l\beta(y, z) - \rho'(x)\bar{\sigma}_l\beta'(y, z) \\
&\quad + \beta(y, x)\bar{\sigma}_r\rho(z) - \beta'(y, x)\bar{\sigma}_r\rho'(z) \\
&\quad - \rho(y)\bar{\sigma}_l\beta(x, z) + \rho'(y)\bar{\sigma}_l\beta'(x, z) \\
&\quad - \beta(x, y)\bar{\sigma}_r\rho(z) + \beta'(x, y)\bar{\sigma}_r\rho'(z) \\
&\quad - \beta(y, x \circ_{\mathfrak{g}} z) + \beta(x, y \circ_{\mathfrak{g}} z) - \beta([x, y], z) \\
&\quad + \beta'(y, x \circ_{\mathfrak{g}} z) - \beta'(x, y \circ_{\mathfrak{g}} z) + \beta'([x, y], z) \\
&= \{h(x)\bar{\sigma}_r(\rho(y) \circ_{\mathfrak{n}} \rho(z) - \rho(y \circ_{\mathfrak{g}} z)) + \rho'(x)\bar{\sigma}_l(\beta - \beta')(y, z) \\
&\quad + (\rho(y) \circ_{\mathfrak{n}} \rho(x) - \rho(y \circ_{\mathfrak{g}} x))\bar{\sigma}_l h(z) + (\beta - \beta')(y, x)\bar{\sigma}_r\rho'(z) \\
&\quad - (\beta - \beta')(y, x \circ_{\mathfrak{g}} z) - c.p.(x, y)\} - (\beta - \beta')([x, y], z), \quad (11)
\end{aligned}$$

for $x, y, z \in \mathfrak{g}$. Define $b \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{m})$ by $b(x, y) := \rho(x)\bar{\sigma}_l h(y) - h(x)\bar{\sigma}_r\mu h(y) + h(x)\bar{\sigma}_r\rho(y) - h(x \circ_{\mathfrak{g}} y)$ such that $\mu b = \mu(\beta - \beta')$, then $\beta - \beta' - b \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, V)$. So

$$\begin{aligned}
(11) &= \{h(x)\bar{\sigma}_r(\rho(y) \circ_{\mathfrak{n}} \rho(z) - \rho(y \circ_{\mathfrak{g}} z)) + \rho'(x)\bar{\sigma}_l b(y, z) \\
&\quad + (\rho(y) \circ_{\mathfrak{n}} \rho(x) - \rho(y \circ_{\mathfrak{g}} x))\bar{\sigma}_l h(z) + b(y, x)\bar{\sigma}_r\rho'(z) - b(y, x \circ_{\mathfrak{g}} z) - c.p.(x, y)\} \\
&\quad - b([x, y], z) + d(\beta - \beta' - b),
\end{aligned}$$

which means that replacing $\beta - \beta'$ by b would not change the cohomology class of $\theta - \theta'$. This replacement yields $[\theta - \theta'] = [0]$ (this process requires comprehensive application of the definitions of the pre-Lie algebra \mathfrak{g} , the action of \mathfrak{n} on \mathfrak{m} , and the crossed module properties).

• It remains to show that equivalent extensions lead to 3-cocycles in the same cohomology class, which is sufficient to show for elementary equivalent extensions. Given elementary equivalent crossed module extensions $E \hookrightarrow E'$ (see definition 2.8), take sections ρ, ρ' of π, π' and sections σ, σ' of μ, μ' , respectively. We would

like to prove $[\theta - \theta'] = 0$. Since $[\theta']$ is independent of the choice of section, we can take $\rho' = s\rho$, then for $x, y, z \in \mathfrak{g}$,

$$\begin{aligned}
(\theta - \theta')(x, y, z) &= (r\theta - \theta')(x, y, z) \\
&= \{s\rho(x)\bar{\sigma}_l(r\sigma - \sigma's)(\rho(y) \circ_n \rho(z) - \rho(y \circ_{\mathfrak{g}} z)) \\
&\quad + (r\sigma - \sigma's)(\rho(y) \circ_n \rho(x) - \rho(y \circ_{\mathfrak{g}} x))\bar{\sigma}_r s\rho(z) \\
&\quad - (r\sigma - \sigma's)(\rho(y) \circ_n \rho(x \circ_{\mathfrak{g}} z) - \rho(y \circ_{\mathfrak{g}} (x \circ_{\mathfrak{g}} z))) - c.p.(x, y)\} \\
&\quad - (r\sigma - \sigma's)(\rho([x, y]) \circ_n \rho(z) - \rho([x, y] \circ_{\mathfrak{g}} z)). \tag{12}
\end{aligned}$$

The first equation holds because $r|_V = id_V$; the second one is guaranteed by the fact that r, s respect actions of crossed modules. For example, the first term in the second equation is obtained as follows:

$$\begin{aligned}
r(\rho(x)\bar{\sigma}_l\beta(y, z)) - \rho'(x)\bar{\sigma}_l\beta(y, z) &= s\rho(x)\bar{\sigma}_l r(\sigma(\rho(y) \circ_n \rho(z) - \rho(y \circ_{\mathfrak{g}} z))) \\
&\quad - s\rho(x)\bar{\sigma}_l \sigma'(s\rho(y) \circ_n s\rho(z) - s\rho(y \circ_{\mathfrak{g}} z)) \\
&= s\rho(x)\bar{\sigma}_l (r\sigma - \sigma's)(\rho(y) \circ_n \rho(z) - \rho(y \circ_{\mathfrak{g}} z)).
\end{aligned}$$

Now, define $\phi \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{m})$ by $\phi(x, y) := (r\sigma - \sigma's)(\rho(x) \circ_n \rho(y) - \rho(x \circ_{\mathfrak{g}} y))$, then direct computation shows that $\mu'\phi = 0$, hence $\phi \in \text{Hom}(\mathfrak{g} \otimes \mathfrak{g}, V)$. It follows from (12) that $\theta - \theta' = d\phi$, hence $[\theta - \theta'] = 0$, which completes the proof. \square

Lemma 3.4. The map $S : H^3(\mathfrak{g}, V) \rightarrow CExt(\mathfrak{g}, V)$ constructed in the proof of Theorem 3.2 is well-defined.

Proof. We claim that S does not depend on the choice of $\tilde{\alpha}$ and that 3-cocycles in the same cohomology class lead to equivalent crossed module extensions.

Indeed, suppose that there is another $\tilde{\alpha}' \in C^2(F, V)$ such that $d\tilde{\alpha}' = \pi^*\theta$, then we get a crossed module extension through it, see (7) and (10). We will show that this crossed module extension is equivalent to the one defined by $\tilde{\alpha}$. By $d(\tilde{\alpha} - \tilde{\alpha}') = 0$ and $H^2(F, V) = 0$, there exists $k \in C^1(F, V)$ such that $\tilde{\alpha} - \tilde{\alpha}' = dk$. It follows that $\alpha - \alpha' = i^*(\tilde{\alpha} - \tilde{\alpha}') = i^*dk = di^*k$. Define $\phi : \mathfrak{m} \rightarrow \mathfrak{m}$ by $\phi(Y + v) = Y + v + i^*k(Y)$ for $Y \in K, v \in V$. Then it is an isomorphism of pre-Lie algebras inducing the identities on V and K , and so the (elementary) equivalence of crossed module extension (9). Therefore, S is independent of the choice of $\tilde{\alpha}$.

Suppose there are 3-cocycles ξ', ξ such that $[\xi'] = [\xi]$. Then $\xi' - \xi = dl$ for some $l \in C^2(\mathfrak{g}, V)$. Hence $\pi^*\xi' = \pi^*\xi + \pi^*dl = d\tilde{\alpha} + d\pi^*l = d(\tilde{\alpha} + \pi^*l)$. Since S (or say the equivalence class of the resulting crossed module extension) is independent of the choice of $\tilde{\alpha}'$ that satisfies $d\tilde{\alpha}' = \pi^*\xi'$, we can choose $\tilde{\alpha}' = \tilde{\alpha} + \pi^*l$. This forces $\alpha' = i^*\tilde{\alpha}' = i^*\tilde{\alpha} + i^*\pi^*l = \alpha$, thus they correspond to the same extension (8). Now it is obvious that identity gives the (elementary) equivalence of crossed module extensions (9) and we conclude that S is well-defined. \square

Lemma 3.5. $ST = 1_{CExt}$.

Proof. • We first recall briefly how to construct $T : CExt \rightarrow H^3$ and $S : H^3 \rightarrow CExt$. Given a crossed module extension:

$$0 \rightarrow V \xrightarrow{i} \mathfrak{m} \xrightarrow{\mu} \mathfrak{n} \xrightarrow{p} \mathfrak{g} \rightarrow 0, \quad (13)$$

where the action of \mathfrak{n} on \mathfrak{m} is denoted by $(\bar{\circ}_l, \bar{\circ}_r)$, we can choose sections ρ of p and σ of μ to define a 3-cocycle θ (see (4)). Then T maps the equivalence class of (13) to $[\theta]$. Let F be the free pre-Lie algebra generated by \mathfrak{g} and $\pi : F \rightarrow \mathfrak{g}$ be the projection. For this 3-cocycle θ , there exists $\tilde{\alpha}_0 \in C^2(F, V)$ such that $d\tilde{\alpha}_0 = \pi^*\theta$. By this $\tilde{\alpha}_0$ we can construct a crossed module extension and S maps $[\theta]$ to equivalence class of it. Our goal is to show that the last crossed module extension is equivalent to the original one (13).

Observing that $d\tilde{\alpha}_0 = \pi^*\theta$ only depends on its values on $\pi(F)$, and that $\pi s\pi = \pi$ for any section s of π , we have

$$d\tilde{\alpha}_0(X, Y, Z) = d\tilde{\alpha}_0(s\pi X, s\pi Y, s\pi Z), \quad X, Y, Z \in F. \quad (14)$$

Write $F = \mathfrak{g} \oplus \ker\pi$ and choose section s to be the inclusion, i.e., $sx = (x, 0)$ for $x \in \mathfrak{g}$. Then by (14), $\pi^*\theta(x + X, y + Y, z + Z) = d\tilde{\alpha}_0(sx, sy, sz)$ for $x + X, y + Y, z + Z \in \mathfrak{g} \oplus \ker\pi$, so $\pi^*\theta$ is completely depended on the restriction of $\tilde{\alpha}_0$ on $(\mathfrak{g}, 0) \otimes (\mathfrak{g}, 0)$. Since S is independent of the choice of $\tilde{\alpha}$ that satisfies $d\tilde{\alpha} = \pi^*\theta$, we can adjust the values of $\tilde{\alpha}_0$ in other parts, like $(0, \ker\pi) \otimes (0, \ker\pi)$, and define a new $\tilde{\alpha}$ which is easier to work with.

Indeed, since F is free on \mathfrak{g} , the section $\rho : \mathfrak{g} \rightarrow \mathfrak{n}$ uniquely induces a pre-Lie homomorphism $G : F \rightarrow \mathfrak{n}$ such that $G(x, 0) = \rho x$ for $x \in \mathfrak{g}$. We can define $\tilde{\alpha} \in C^2(F, V)$ by

$$\tilde{\alpha}(x + X, y + Y) := \tilde{\alpha}_0(x, y) + f(x, Y) + f(X, y) + f(X, Y),$$

where $if(X, Y) := \sigma(GX) \circ \sigma(GY) - \sigma(GX \circ GY)$ for $X, Y \in F$. Now, we get a crossed module extensions as in the proof of Theorem 3.2:

$$0 \rightarrow V \rightarrow \ker\pi \oplus_{\tilde{\alpha}} V \xrightarrow{\mu_{\tilde{\alpha}}} F \xrightarrow{\pi} \mathfrak{g} \rightarrow 0. \quad (15)$$

The pre-Lie structure on $\ker\pi \oplus V$ is given by (7), and the action of F on $\ker\pi \oplus V$, denoted also by $(\bar{\circ}_l, \bar{\circ}_r)$, is given by (10).

• To show that crossed module (15) is equivalent to (13), we need to define $H : \ker\pi \oplus_{\tilde{\alpha}} V \rightarrow \mathfrak{m}$ appropriately to give the (elementary) equivalence:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & V & \longrightarrow & \ker\pi \oplus_{\tilde{\alpha}} V & \xrightarrow{\mu_{\tilde{\alpha}}} & F & \xrightarrow{\pi} & \mathfrak{g} & \longrightarrow & 0 \\ & & \parallel & & H \downarrow & & G \downarrow & & \parallel & & \\ 0 & \longrightarrow & V & \xrightarrow{i} & \mathfrak{m} & \xrightarrow{\mu} & \mathfrak{n} & \xrightarrow{p} & \mathfrak{g} & \longrightarrow & 0. \end{array} \quad (16)$$

The commutativity of the diagram indicates that we shall define $H : \ker\pi \oplus_{\tilde{\alpha}} V \rightarrow \mathfrak{m}$ by $H(Y + v) = \sigma GY + iv$. H is a pre-Lie homomorphism because $i\tilde{\alpha}(X, Y) = \sigma(GX) \circ_{\mathfrak{m}} \sigma(GY) - \sigma(GX \circ_{\mathfrak{n}} GY)$ for $X, Y \in \ker\pi$.

We next show that H and G respect the actions of crossed modules. Take $H(X\bar{\circ}_l(Y+v)) = GX\bar{\circ}_lH(Y+v)$, $X \in F, Y+v \in \ker\pi \oplus V$ as an example. This is equivalent to:

$$\begin{aligned}\sigma G(X \circ_F Y) + i\tilde{\alpha}(X, Y) + i(X \circ_l v) &= GX\bar{\circ}_l\sigma GY + GX\bar{\circ}_l i v \\ &= \mu\sigma GX\bar{\circ}_l\sigma GY + GX\bar{\circ}_l i v \\ &= \sigma GX \circ_m \sigma GY + GX\bar{\circ}_l i v.\end{aligned}$$

Since

$$\begin{aligned}GX\bar{\circ}_l i v - i(X \circ_l v) &= GX\bar{\circ}_l i v - i(\pi X \circ_l v) \\ &= GX\bar{\circ}_l i v - \rho\pi X\bar{\circ}_l i v \\ &= (G - \rho\pi)X\bar{\circ}_l i v,\end{aligned}$$

we have $i\tilde{\alpha}(X, Y) = \sigma(GX) \circ_m \sigma(GY) - \sigma(GX \circ_n GY) + (G - \rho\pi)X\bar{\circ}_l v$. We claim that $(G - \rho\pi)X \circ v = 0$ for all $X \in F, v \in V$, hence H, G respect the actions by the definition of $\tilde{\alpha}$. For $X \in \ker\pi$, $(G - \rho\pi)X = GX \in \ker p = \text{im } \mu$. Thus $GX = \mu m$ for some $m \in \mathfrak{m}$, and $GX \circ v = \mu m \circ i(v) = m \circ i(v) = m \circ \mu i(v) = 0$. For $X = (x, 0) \in \mathfrak{g} \oplus 0$, we have $(G - \rho\pi)X = G(x, 0) - \rho x = 0$.

• To sum up, ST maps equivalence class of crossed module (13) to $[\theta]$, and then to equivalence class of crossed module (15), which equals to the original one, hence $ST = 1_{CExt}$. \square

Lemma 3.6. $TS = 1_{H^3(\mathfrak{g}, V)}$

Proof. We begin by recalling the construction of TS . Given a $[\theta] \in H^3(\mathfrak{g}, V)$, we can find $\tilde{\alpha} \in C^2(F, V)$ such that $\pi^*\theta = d\tilde{\alpha}$, then we define a crossed module (15) by this $\tilde{\alpha}$. We next choose sections s of π and $\sigma_{\tilde{\alpha}}$ of $\mu_{\tilde{\alpha}}$ to get a map $\beta \in C^2(\mathfrak{g}, \ker\pi \oplus_{\tilde{\alpha}} V)$, which defines a new 3-cocycle θ' . Our goal is to show $[\theta] = [\theta']$. In fact, $\theta' - \theta = d(s^*\tilde{\alpha})$. We only need to prove $\pi^*\theta' - \pi^*\theta = \pi^*d(s^*\tilde{\alpha})$ since π^* is injective.

In particular, we can take $\sigma_{\tilde{\alpha}} : \ker\pi \rightarrow \ker\pi \oplus_{\tilde{\alpha}} V$ to be the inclusion, i.e., $\sigma_{\tilde{\alpha}}(X) = (X, 0)$ for $X \in \ker\pi$. Then $\beta(x, y) = (sx \circ sy - s(x \circ y), 0)$ for $x, y \in \mathfrak{g}$. Denote the projection of $\ker\pi \oplus_{\tilde{\alpha}} V$ onto $\ker\pi$ by pr . Then by (4) and (10) we have

$$\begin{aligned}\pi^*\theta'(X, Y, Z) &= s\pi X \circ pr\beta(\pi Y, \pi Z) + pr\beta(\pi Y, \pi X) \circ s\pi Z \\ &\quad - \beta(\pi Y, \pi X \circ \pi Z) - \beta(\pi Y \circ \pi X, \pi Z) \\ &\quad + \tilde{\alpha}(s\pi X, pr\beta(\pi Y, \pi Z)) + \tilde{\alpha}(pr\beta(\pi Y, \pi X), s\pi Z) - c.p.(X, Y) \\ &= \tilde{\alpha}(s\pi X, pr\beta(\pi Y, \pi Z)) + \tilde{\alpha}(pr\beta(\pi Y, \pi X), s\pi Z) - c.p.(X, Y), \\ &= \tilde{\alpha}(s\pi X, s\pi Y \circ s\pi Z - s(\pi Y \circ \pi Z)) \\ &\quad + \tilde{\alpha}(s\pi Y \circ s\pi X - s(\pi Y \circ \pi X), s\pi Z) - c.p.(X, Y),\end{aligned}$$

the second equation holds since the image of $\pi^*\theta'$ is in V . By (14) we have

$$\begin{aligned}\pi^*\theta(X, Y, Z) &= d\tilde{\alpha}(s\pi X, s\pi Y, s\pi Z), \\ &= s\pi X \circ \tilde{\alpha}(s\pi Y, s\pi Z) + \tilde{\alpha}(s\pi Y, s\pi X) \circ s\pi Z \\ &\quad - \tilde{\alpha}(s\pi Y, s\pi X \circ s\pi Z) - \tilde{\alpha}(s\pi Y \circ s\pi X, s\pi Z) - c.p.(X, Y),\end{aligned}$$

for $X, Y, Z \in F$. So

$$\begin{aligned}\pi^*(\theta - \theta')(X, Y, Z) &= s\pi X \circ \tilde{\alpha}(s\pi Y, s\pi Z) + \tilde{\alpha}(s\pi Y, s\pi X) \circ s\pi Z \\ &\quad - \tilde{\alpha}(s\pi Y, s(\pi X \circ \pi Z)) - \tilde{\alpha}(s(\pi Y \circ \pi X), s\pi Z) - c.p.(X, Y) \\ &= \pi^*d(s^*\tilde{\alpha})(X, Y, Z)\end{aligned}$$

as required. \square

- Relation with abelian extensions of the Lie algebra.

Let A be a Lie algebra, W be a A -module. An abelian extension of A by W is an exact sequence of Lie algebras:

$$0 \rightarrow W \rightarrow \hat{A} \rightarrow A \rightarrow 0,$$

where all arrows are homomorphisms of Lie algebras. Two abelian extension of A by W are equivalent if there exists a homomorphism γ such that the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & W & \longrightarrow & \hat{A}_2 & \longrightarrow & A \longrightarrow 0 \\ & & \parallel & & \gamma \downarrow & & \parallel \\ 0 & \longrightarrow & W & \longrightarrow & \hat{A}_1 & \longrightarrow & A \longrightarrow 0. \end{array}$$

Note that in this case γ must be an isomorphism. Denote by $AExt(A, W)$ the set of equivalence classes of abelian extensions of A by W .

Let \mathfrak{g} be a pre-Lie algebra, and V be a \mathfrak{g} -module. Then by remark 2.11 of [15], there is a \mathfrak{g}^c -module structure on $\text{Hom}(\mathfrak{g}, V)$ given by

$$(x \triangleright f)(y) := x \circ f(y) + f(x) \circ y - f(x \circ y), \quad \forall f \in \text{Hom}(\mathfrak{g}, V), \quad x, y \in \mathfrak{g}.$$

Corollary 3.7. There is a one-to-one correspondence between $CExt(\mathfrak{g}, V)$ and $AExt(\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V))$.

Proof. By [15], there is an isomorphism $\phi : H_{preLie}^n(\mathfrak{g}, V) \rightarrow H_{Lie}^{n-1}(\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V))$ induced by the cochain map $\phi : \text{Hom}(\wedge^{n-1}\mathfrak{g} \otimes \mathfrak{g}, V) \rightarrow \text{Hom}(\wedge^{n-1}\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V))$, which is defined by

$$(\phi f)(x_1, \dots, x_{n-1})(x_n) := f(x_1, \dots, x_n).$$

Combine it with theorem 3.2 and the well-known bijection between $H_{Lie}^2(\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V))$ and $AExt(\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V))$ yields the desire correspondence:

$$\begin{array}{ccc} H_{preLie}^3(\mathfrak{g}, V) & \xrightarrow{1-1} & H_{Lie}^2(\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V)) \\ \uparrow 1-1 & & \downarrow 1-1 \\ CExt(\mathfrak{g}, V) & \xrightarrow{\text{Corollary 3.7}} & AExt(\mathfrak{g}^c, \text{Hom}(\mathfrak{g}, V)). \end{array}$$

\square

4 Some relations on crossed modules of pre-Lie algebras

In this section, we summarize some relations between crossed modules of pre-Lie algebras and crossed modules of other algebras, which are similar to the relations between these algebras.

Denote the commutator Lie algebra of any associative algebra \mathfrak{m} by \mathfrak{m}^c . Let $(\mathfrak{m} \xrightarrow{\mu} \mathfrak{n}, \cdot_l, \cdot_r)$ be a crossed module of associative algebras, then as mentioned in [10], $(\mathfrak{m}^c \xrightarrow{\mu} \mathfrak{n}^c, \triangleright)$ is a crossed module of Lie algebras with the action $\triangleright : \mathfrak{n}^c \wedge \mathfrak{m}^c \rightarrow \mathfrak{m}^c$ given by $x \triangleright u := x \cdot_l u - u \cdot_r x$ for $x \in \mathfrak{n}^c, u \in \mathfrak{m}^c$.

Proposition 4.1 ([19], proposition 4.9). Let $((\mathfrak{m}, \circ_m) \xrightarrow{\mu} (\mathfrak{n}, \circ_n), \circ_l, \circ_r)$ be a crossed module of pre-Lie algebras and $\mathfrak{m}^c, \mathfrak{n}^c$ be their commutator Lie algebras. Then $(\mathfrak{m}^c \xrightarrow{\mu} \mathfrak{n}^c, \triangleright)$ is a crossed module of Lie algebras, where $x \triangleright u := x \circ_l u - u \circ_r x$ for $x \in \mathfrak{n}, u \in \mathfrak{m}$.

Proposition 4.2 ([23], proposition 4.8). Let $((\mathfrak{m}, [\cdot, \cdot]_m, T_m) \xrightarrow{\mu} (\mathfrak{n}, [\cdot, \cdot]_n, T_n), \rho)$ be a crossed module of Rota-Baxter Lie algebras. Define $\circ_m : \mathfrak{m} \otimes \mathfrak{m} \rightarrow \mathfrak{m}$, $\circ_n : \mathfrak{n} \otimes \mathfrak{n} \rightarrow \mathfrak{n}$, and $\circ_l : \mathfrak{n} \otimes \mathfrak{m} \rightarrow \mathfrak{m}$, $\circ_r : \mathfrak{m} \otimes \mathfrak{n} \rightarrow \mathfrak{m}$ by

$$\begin{aligned} x \circ_n y &:= [T_n x, y]_n, & u \circ_m v &:= [T_m u, v]_m, \\ x \circ_l u &:= \rho(T_n x)u, & u \circ_r x &:= -\rho(x)(T_m u), \end{aligned}$$

for $x, y \in \mathfrak{n}, u, v \in \mathfrak{m}$. Then $((\mathfrak{m}, \circ_m) \xrightarrow{\mu} (\mathfrak{n}, \circ_n), \circ_l, \circ_r)$ is a crossed module of pre-Lie algebras.

The notion of crossed module of dendriform algebras is given in [9], see definition 6.6 and also remark 2.5. It gives rise to a crossed module of associative algebra, see subsection 6.6 of [9]. The following proposition shows that it also gives a crossed module of pre-Lie algebras.


Proposition 4.3. Let $((\mathfrak{m}, \succ_m, \prec_m) \xrightarrow{\mu} (\mathfrak{n}, \succ_n, \prec_n), \succ, \prec)$ be a crossed module of dendriform algebras. Define $\circ_m : \mathfrak{m} \otimes \mathfrak{m} \rightarrow \mathfrak{m}$, $\circ_n : \mathfrak{n} \otimes \mathfrak{n} \rightarrow \mathfrak{n}$, and $\circ_l : \mathfrak{n} \otimes \mathfrak{m} \rightarrow \mathfrak{m}$, $\circ_r : \mathfrak{m} \otimes \mathfrak{n} \rightarrow \mathfrak{m}$ by

$$\begin{aligned} x \circ_n y &:= x \succ_n y - y \prec_n x, & u \circ_m v &:= u \succ_m v - v \prec_m u, \\ x \circ_l u &:= x \succ u - u \prec x, & u \circ_r x &:= u \succ x - x \prec u, \end{aligned}$$

for $x, y \in \mathfrak{n}, u, v \in \mathfrak{m}$. Then $((\mathfrak{m}, \circ_m) \xrightarrow{\mu} (\mathfrak{n}, \circ_n), \circ_l, \circ_r)$ is a crossed module of pre-Lie algebras.

Proof. By proposition 5.2 of [1], (\mathfrak{m}, \circ_m) and (\mathfrak{n}, \circ_n) are pre-Lie algebras. Since μ is a dendriform homomorphism, it is also a pre-Lie homomorphism. The fact that (\succ, \prec) is an action of dendriform algebras implies that (\circ_l, \circ_r) is an action of pre-Lie algebras. Moreover, $\mu(x \circ_l u) = \mu(x \succ u - u \prec x) = x \succ_n \mu u - \mu u \prec_n x = x \circ_n \mu u$ for $x \in \mathfrak{n}, u \in \mathfrak{m}$. Similar considerations show that $((\mathfrak{m}, \circ_m), (\mathfrak{n}, \circ_n), \mu, \circ)$ is a crossed module of pre-Lie algebras. \square

We summarize these relations in the following diagram:



where $X_{asso}, X_{Lie}, X_{preLie}, X_{RBLie}, X_{dend}$ are abbreviations of crossed modules of associative, Lie, pre-Lie, Rota-Baxter Lie and dendriform algebras respectively, and the arrow means that we can obtain one crossed module from the other.

References

- [1] Bai, C. M. (2021). An introduction to pre-Lie algebras. In: Algebra and Applications 1: Non-associative Algebras and Categories, Wiley Online Library, pp. 245-273.
- [2] Burde, D. (2006). Left-symmetric algebras and pre-Lie algebras in geometry and physics. *Cent. Eur. J. Math.* 4:323-357.
- [3] Casas, J. M., Casado, R. F., Khmaladze, E., Ladra, M. (2014). Universal enveloping crossed module of a Lie crossed module. *Homology Homotopy Appl.* 16:143-158.
- [4] Casas, J. M., Casado, R. F., Khmaladze, E., Ladra, M. (2017). More on crossed modules in Lie, Leibniz, associative and diassociative algebras. *J. Algebra Appl.* 16:1750107.
- [5] Casas, J. M., Ladra, M., Pirashvili, T. (2004). Crossed modules for Lie-Rinehart algebras. *J. Algebra* 274:192-201.
- [6] Chapton, F., Livernet, M. (2001). Pre-Lie algebras and the rooted trees operad. *Int. Math. Res. Not.* 8:395-408.
- [7] Chapton, F. (2010). Free pre-Lie algebras are free as Lie algebras. *Canad. Math. Bull.* 53:425-437.
- [8] Chen, L. Y., Liu M. J., Liu, J. F. (2022). Cohomologies and crossed modules for pre-Lie Rinehart algebras. *J. Geom. Phys.* 176:104501.
- [9] Das, A. (2022). Cohomology and deformations of dendriform algebras, and $Dend_\infty$ -algebras. *Commun. Algebra.* 50:1544-1567.
- [10] Donadze, G., Inassaridze, N., Khmaladze, E., Ladra, M. (2012). Cyclic homologies of crossed modules of algebras. *J. Noncommut. Geom.* 6:749-771.

- [11] Dzhumadil'daev, A. (1999). Cohomologies and deformations of right-symmetric algebras. *J. Math. Sci.* 93:836-876.
- [12] Gerstenhaber, M. (1963). The cohomology structure of an associative ring. *Ann. Math.* 78:267-288.
- [13] Gerstenhaber, M. (1964). A uniform cohomology theory for algebras. *Pros. Nat. Acad. Sci. U.S.A.* 51:626-629.
- [14] Gerstenhaber, M. (1966). On the deformation of rings and algebras: II. *Ann. Math.* 84:1-19.
- [15] Guan, A., Lazarev, A., Sheng, Y. H., Tang, R. (2020). Review of deformation theory I: concrete formulas for deformations of algebraic structures, *Adv. math. (China)* 49:257-277.
- [16] Hilton, P. J., Stammach, U. (1971). A course in homological algebra. (Grad. Texts in Math., vol. 4) Berlin Heidelberg New York: Springer.
- [17] Lang, H. L., Liu, Z. J. (2016). Crossed modules for Lie 2-algebras. *Appl. Categor. Struct.* 24:53-78.
- [18] Ma, Q. F., Song, L. N., Wang, Y. (2023). Non-abelian extensions of pre-Lie algebras. *Commun. Algebra.* 51:1370-1382.
- [19] Sheng, Y. H. (2019). Categorification of pre-Lie algebras and solutions of 2-graded classical Yang-Baxter equations. *Theo. Appl. Cate.* 34:269-294.
- [20] Wagemann, F. (2006). On Lie algebra crossed modules. *Commun. Algebra* 34:1699-1722.
- [21] Whitehead, J. (1941). On adding relations to homotopy groups. *Ann. Math.* 42:409-428.
- [22] Zhang, T. Zinbiel 2-algebras. arXiv:2104.12551.
- [23] Zhang, S. L., Liu, J. F. (2023). On Rota-Baxter Lie 2-algebras. *Theo. Appl. Cate.* 39:545-566.