

Alternating paths in oriented graphs with large semidegree

Jozef Skokan ^{*} Mykhaylo Tyomkyn [†]

June 7, 2024

Abstract

In new progress on conjectures of Stein, and Addario-Berry, Havet, Linhares Sales, Reed and Thomassé, we prove that every oriented graph with all in- and out-degrees greater than $5k/8$ contains an alternating path of length k . This improves on previous results of Klimošová and Stein, and Chen, Hou and Zhou.

1 Introduction

A classical topic in graph theory is establishing minimum degree conditions for existence of long paths and cycles in graphs. This research direction has a rich history, going back to works of Dirac [3], Posa [14] and others. The present paper is a part of the ongoing effort to establish analogous results in oriented graphs.

An oriented graph is a tuple $G = (V, \vec{D})$ where V is a finite set of *vertices* and \vec{D} is a set of (*directed*) *edges*. Each edge is an ordered pair of distinct vertices. Moreover, for all u and v at most one of (u, v) and (v, u) is an edge. The *in-degree* and *out-degree* of a vertex $v \in V$ are defined as $d^-(v) = |\{(u, v) \in \vec{D} : u \in V\}|$ and $d^+(v) = |\{(v, u) \in \vec{D} : u \in V\}|$, respectively. The semidegree of v is $d_0(v) = \min\{d^-(v), d^+(v)\}$. The minimum semidegree of the oriented graph $G = (V, \vec{D})$ is $\delta^0(G) := \min\{d^0(v) : v \in V\}$.

Analogously to undirected graphs, it is natural to ask for oriented graphs, what minimum semidegree conditions would guarantee occurrence of long paths. The first case to consider are the directed paths, i.e. paths whose edges are directed the same way. Here Jackson [8] proved, as a corollary of a more general result, that $\delta^0(G) > k/2$ implies that G contains a directed path with k edges. The proof follows a path extension argument, very similar to

^{*}Department of Mathematics, London School of Economics, Houghton Street, London, WC2A 2AE.

[†]Department of Applied Mathematics, Charles University. Email: tyomkyn@kam.mff.cuni.cz. Supported in part by ERC Synergy Grant DYNASNET 810115 and GAČR Grant 22-19073S.

the textbook proof of Dirac's theorem [3]. Subsequent years saw a large body of research on Hamilton cycles in oriented and directed graphs [4, 5, 6, 7, 9, 10, 11, 17].

Motivated by the above works, as well as by results on degree conditions for occurrence of trees in undirected and directed graphs (see [15]), Stein [15] conjectured the following generalization of Jackson's theorem [8].

Conjecture 1.1 (Stein [15]). *Every oriented graph with $\delta^0(G) > k/2$ contains every orientation of a path with k edges.*

While it is easy to see that for $\delta^0(G) > k$ we can find greedily every orientation of a k -edge path, at present we do not know to replace k with any smaller value (let alone $k/2$), for the statement of Conjecture 1.1 to hold universally.

After the directed paths, the next most natural case to consider are the *alternating* paths, that is, paths in which every vertex sees either just incoming or just outgoing edges (this case also provides a construction for the bound of $k/2$ in Conjecture 1.1: a $k/2$ -blowup of a directed cycle). Klimošová and Stein [12] proved for that $\delta^0(G) > 3k/4$ implies existence of an alternating path with k edges, and Chen, Hou and Zhou [2] improved their bound to $2k/3$ (up to additive constants). For the special case when k is linear in the number of vertices of the oriented graph, Stein and Zárata-Guerén [16] proved an approximate upper bound $(1/2 + o(1))k$.

Our goal here is to improve the general upper bound further to $5k/8$. Similarly to [2] and [12], we shall prove a slightly more general result, using the *pseudo-semidegree* $\bar{\delta}^0(G)$, which is defined, for an oriented graph without isolated vertices, as the minimum over all *positive* vertex in- and out-degrees.

For better proof clarity we decided not to optimize additive constants. Consequently, we find it more convenient to deal with paths with k vertices rather than edges.

Theorem 1.2. *Every oriented graph G with $\bar{\delta}^0(G) > 5k/8$ contains an alternating path with k vertices.*

As a corollary we obtain the following improvement towards a conjecture of Addario-Berry, Havet, Linhares Sales, Reed and Thomassé [1] (for the reduction we refer to [12]).

Corollary 1.3. *Every oriented graph G with more than $(5k + 4)|V(G)|/4$ edges contains an alternating path with k vertices.*

2 The proof of Theorem 1.2.

We start with some notation. Let $G = (V, \vec{D})$ be an oriented graph. For a vertex $v \in V$ and a vertex set $X \subseteq V$ let $N_X^+(v)$ and $N_X^-(v)$ denote $\{x \in X : (v, x) \in \vec{D}\}$ and $\{x \in X :$

$(x, v) \in \vec{D}$ respectively. Let $d_X^+(v) := |N_X^+(v)|$ and $d_X^-(v) := |N_X^-(v)|$. Let $N^+(v) := N_V^+(v)$ and $N^-(v) := N_V^-(v)$ denote the out- and in-neighbourhoods of v .

Suppose now that G is an oriented graph with $\bar{\delta}^0(G) > 5k/8$. Let P a maximum length alternating path in G . Assume for contradiction that P has strictly fewer than k vertices. We proceed according to the parity of $|P|$.

2.1 The even case

We first consider the case when P has an even number of vertices $2m < k$. Let the vertices of P be labeled p_1, \dots, p_{2m} , in the order of the underlying undirected path. Without loss of generality assume that the edge p_1p_2 is oriented toward p_2 . Let O and E be the sets of the odd and even indexed vertices p_i , respectively; we shall call them simply *odd* and *even* vertices. Note that every edge of P is directed from an odd to an even vertex. Therefore, by definition of $\bar{\delta}^0(G)$, we have $d^+(v) > 5k/8$ for every $v \in O$ and $d^-(w) > 5k/8$ for every $w \in E$.

Lemma 2.1. *There is an alternating spanning cycle on $V(P) = E \cup O$ such that each of its edges is directed from odd to even.*

Proof. We call an alternating path $P' \subseteq G$ *respectable* if $V(P') = E \cup O$ and each edge of P' is oriented from an odd to an even vertex (in particular, one end vertex of P' is odd and one is even). The set of respectable paths is not empty, since P is respectable. Call an even vertex $v \in E$ *terminal* if it is an end vertex of a respectable path, and let T denote the set of terminal vertices. Call an odd vertex $v \in O$ *starting* if it is an initial vertex of a respectable path, and let S denote the set of starting vertices. So, for instance, $p_{2m} \in T$ $p_1 \in S$, as witnessed by a respectable path P .

Suppose $v \in S$ is a starting vertex of some respectable path P' . If there is an edge $(v, w) \in \vec{D}$ from some $w \notin E \cup O$, we would be able to extend P' via the edge (v, w) obtaining a longer alternating path. Thus we may assume that $N^+(v) \subseteq E \cup O$. So, since $d_{E \cup O}^+(v) = d^+(v) > 5k/8 > (5/8) \cdot 2m$ and $|E| = |O| = m$, we must have $d_E^+(v) > 0$ and also $d_O^+(v) > 0$. Similarly, for every terminal $w \in T$ we have $N^-(w) \subseteq E \cup O$, and in particular $d_O^-(w) > 0$ and $d_E^-(w) > 0$.

Let now $a \in S$ be a starting vertex with $d_E^+(a) = \max\{d_E^+(v) : v \in S\}$. Fix some respectable path R starting at a , and denote its vertices $a = r_1, r_2, \dots, r_{2m}$ in the order of the underlying undirected path. Note that for each $r_i \in N_E^+(a)$ we have $r_{i-1} \in S$, since we have a respectable path $r_{i-1}, \dots, r_1, r_i, \dots, r_{2m}$. So, letting $A := \{r_{i-1} : r_i \in N_E^+(a)\}$, we have $A \subseteq S$ and $|A| = d_E^+(a)$.

Now, by maximality of $d_E^+(a)$, for each vertex $u \in A$ we have

$$d_E^+(u) \leq d_E^+(a) = |A|,$$

and therefore

$$d_O^+(u) \geq \frac{5}{8}k - |A| > \frac{5}{8} \cdot 2m - |A|.$$

Let $s := \sum_{u \in A} d_O^+(u)$. By the above, we have

$$s > \frac{5}{8}|A| \cdot 2m - |A|^2.$$

On the other hand, counting the edges via out-degrees, using $|O| = m$, yields

$$s \leq \frac{|A|^2}{2} + |A|(m - |A|).$$

It follows that

$$\frac{|A|^2}{2} + |A|(m - |A|) > \frac{5}{8}|A| \cdot 2m - |A|^2,$$

resulting in

$$d_E^+(a) = |A| > \frac{m}{2}.$$

Now let B be the set of all vertices $v \in T$ such that there exists a respectable path starting at a and terminating at v ; note that $B \neq \emptyset$, as R is respectable and starts at a , so $r_{2m} \in B$. Let $b \in B$ be a vertex satisfying $d_O^-(b) = \max\{d_O^-(v) : v \in B\}$. Without loss of generality we may assume that $b = r_{2m}$. Note now that for each $r_i \in N_O^-(b)$ we have $r_{i+1} \in B$ due to the respectable path $a = r_1, \dots, r_i, r_{2m}, \dots, r_{i+1}$. So, letting $C := \{r_{i+1} : r_i \in N_O^-(b)\}$, we have $C \subseteq B$ and $|C| = d_O^-(b)$.

By maximality of $d_O^-(b)$, for each vertex $w \in C$ we have

$$d_O^-(w) \leq d_O^-(b) = |C|,$$

and so

$$d_E^-(w) \geq \frac{5}{8}k - |C| > \frac{5}{8} \cdot 2m - |C|.$$

Let $t := \sum_{w \in C} d_E^-(w)$. We obtain

$$t > |C| \cdot \frac{5}{8} \cdot 2m - |C|^2.$$

On the other hand, counting the edges via in-degrees gives

$$t \leq \frac{|C|^2}{2} + |C|(m - |C|).$$

Hence,

$$\frac{|C|^2}{2} + |C|(m - |C|) > |C| \cdot \frac{5}{8} \cdot 2m - |C|^2,$$

which results in

$$d_O^-(b) = |C| > \frac{m}{2}.$$

Notice now that $|O| = m$, and we have shown that $N_O^-(b) \subseteq O$ and $\{r_i : r_{i+1} \in N_E^+(a)\} \subseteq O$ are of size greater than $m/2$. Therefore, by pigeonhole, for some i we must have $r_i \in N_O^-(b) = N_O^-(r_{2m})$ and $r_{i+1} \in N_E^+(a) = N_E^+(r_1)$. This gives the desired cycle via the cyclic ordering $r_1, \dots, r_i, r_{2m}, r_{2m-1}, \dots, r_{i+1}, r_1$. \square

Corollary 2.2. *We have $S = O$ and $T = E$. Consequently, for every $a \in O$ and $b \in E$ we have $N^+(a) \subseteq E \cup O$ and $N^-(b) \subseteq E \cup O$.*

Let now H be the simple bipartite graph obtained by taking all edges directed from odd to even vertices, and ignoring their directions. We claim that not only is H hamiltonian (we already know this) but it satisfies, with room to spare, the Moon-Moser degree condition for hamiltonicity of bipartite graphs. To this end, let $d(v)$ denote the degree of a vertex $v \in E \cup O$ in the graph H .

Lemma 2.3. *For every $1 \leq \ell \leq m/2$ we have*

$$|\{v \in O : d(v) \leq \ell + 1\}| < \ell.$$

And, symmetrically,

$$|\{w \in E : d(w) \leq \ell + 1\}| < \ell.$$

Proof. Suppose for a contradiction that for some $1 \leq \ell \leq m/2$ there exists a set $L \subseteq O$ (the statement for E is proved in the same way) of ℓ vertices of H -degree at most $\ell + 1$. Then each vertex $v \in L$ satisfies (using $d^+(v) > 5k/8$ and $k \geq 2m + 1$)

$$\begin{aligned} d_O^+(v) &> \frac{5}{8}k - \ell - 1 \geq \frac{5}{8}(2m + 1) + \frac{1}{8} - \ell - 1 \\ &= \frac{5}{4}m - \ell - \frac{1}{4}. \end{aligned}$$

It follows that

$$|N^+(v) \cap L| > \frac{5}{4}m - \ell - \frac{1}{4} - (m - \ell) = \frac{m - 1}{4} > \frac{\ell - 1}{2},$$

a contradiction, counting out-degrees inside L . \square

Lemma 2.4. *G contains an alternating path Q on $2m$ vertices, of the form $q_1 \dots q_{2m}$, where $q_3, \dots, q_{2m-1} \in O$, $q_4, \dots, q_{2m} \in E$, $q_2 \in O$, $q_1 \notin E \cup O$, and the edge q_1q_2 is directed from q_1 to q_2 .*

Proof. Recall that $d_O^+(v) > 0$ for every $v \in O$. So, there must be three vertices $u, q_2, q_3 \in O$ forming a directed path uq_2q_3 , i.e., $(u, q_2), (q_2, q_3) \in \vec{D}$. Since $d^-(q_2) > 0$, by the pseudo-semidegree condition we have $d^-(q_2) > 5k/8 > (5/8) \cdot 2m$ in addition to $d^+(q_2) > (5/8) \cdot 2m$,

as $q_2 \in O$. Since G cannot have multiple edges between a pair of vertices, we conclude that q_2 has an in-neighbour $q_1 \notin E \cup O$.

Let $O' = O \setminus q_2$, and let E' be the set E with an arbitrary vertex removed. Then, by Lemma 2.4, the bipartite graph $H' = H[E' \cup O']$ satisfies the Moon-Moser condition (Corollary 1 in [13]): for any $1 \leq \ell \leq |O'|/2 = (n-1)/2$ the number of vertices of H' -degree at most ℓ is less than ℓ , and symmetrically in E' . Therefore, H' has a hamilton cycle, and, a fortiori, a hamilton path q_3, \dots, q_{2m} . This yields a desired alternating path $Q = q_1, q_2, q_3, \dots, q_{2m}$ in G . □

Observe that Lemma 2.1 applies to any alternating path on $2m$ vertices, and in particular, to Q , in place of P . Now consider the vertex q_2 . Since $q_2 \in O$, we have $N^+(q_2) \subset E \cup O$. On the other hand, q_2 plays the role of an ‘even’ vertex in Q . So, by Lemma 2.1 we must have $N^-(q_2) \subset E \cup O \cup \{q_1\}$. Put together, this would imply $d^-(q_2) + d^+(q_2) \leq 2m$, contradicting the fact that $d_0(q_2) > (5/8) \cdot 2m$. This concludes the case when the maximum length path has an even number of vertices.

2.2 The odd case

Now consider the case when the maximum length path P has an odd number of vertices $2m+1 < k$. Let p_1, \dots, p_{2m+1} be the vertices of P in the order of the underlying undirected path. Without loss of generality assume that $p_1 p_2$ is oriented toward p_2 . We again call the even and odd indexed vertices of P *even* and *odd* respectively, and denote the corresponding sets by E and O . By $\bar{\delta}^0(G) > 5k/8$, we have $d^+(v) > 5k/8$ for every $v \in O$ and $d^-(w) > 5k/8$ for every $w \in E$.

The edge $p_{2m} p_{2m+1}$ is oriented toward p_{2m} . If there is a directed edge from p_{2m+1} to some vertex $w \notin P$, this would give rise to a longer alternating path. Thus we may assume that $N^+(p_{2m+1}) \subseteq E \cup O$.

Lemma 2.5. *There exists an odd vertex $v \in O$ such that on $E \cup O \setminus \{v\}$ there is an alternating spanning cycle such that each of its edges is directed from odd to even.*

Proof. This is similar to the proof of Lemma 2.1. Call a path on $2m+1$ vertices *respectable* if its vertex set is $V(P) = E \cup O$, and every edge is directed from odd to even (so the end vertices are both odd). Call an odd vertex *starting* if it is an end vertex of a respectable path. So, for instance, both p_1 and p_{2m+1} are starting, as witnessed by the respectable path P . Let S be the set of starting vertices.

For any starting vertex $v \in S$ we must have $N^+(v) \subseteq E \cup O$, otherwise we could extend P , and so we must have $d_E^+(v) > 0$. Let $a \in S$ be a starting vertex with $d_E^+(a) = \max\{d_E^+(v) : v \in S\}$. Fix some respectable path $a = r_1, \dots, r_{2m+1}$, and let $A = \{r_{i-1} : r_i \in N_E^+(a)\}$;

we have $A \subseteq S$. By maximality of $d_E^+(a)$ each $u \in A$ satisfies $d_E^+(u) \leq d_E^+(a) = |A|$, and therefore $d_O^+(u) > 5k/8 - |A|$. Double counting the edges similarly to the even case, we obtain $d_E^+(a) \geq m/2 + 1$ (we skip the straightforward calculations).

Let B be the set of all vertices $v \in S$ such that there is a respectable path with a and b as its end vertices; note that $B \neq \emptyset$. Let $b \in B$ be a vertex satisfying $d_O^+(b) = \max\{d_O^+(v) : v \in B\}$. Without loss of generality we may assume that $b = r_{2m+1}$. Similarly to the above we obtain $d_E^+(b) \geq m/2 + 1$.

Since $|E| = m$, by pigeonhole, there must be some $r_{i+2} \in N_E^+(a)$ and $r_i \in N_E^+(b)$. This gives the desired cycle $a, r_{i+2}, r_{i+3}, \dots, r_{2m+1} = b, r_i, r_{i-1}, \dots, r_1 = a$, with $r_{i+1} \in O$ playing the role of v in the lemma statement. \square

Hence, we now may assume the existence of an alternating spanning cycle C on $E \cup O \setminus \{v\}$. Let us label its vertices along the cycle by c_1, \dots, c_{2m} such that each even indexed c_i belongs to E , each odd indexed c_i is in O , and each edge is directed from odd to even. Note also that $v \notin V(C)$ sends at least one edge to an even vertex on C , say, to c_{2m} : $(v, c_{2m}) \in \vec{D}$. If for some $w \notin C$ we have $(v, w) \in \vec{D}$, we obtain a longer alternating path via $w, v, c_{2m}, c_{2m-1}, \dots, c_1$. So, $N^+(v) \subseteq E \cup O$. Therefore, as $5k/8 > m$, by pigeonhole, $N^+(v)$ will contain two adjacent vertices on C . Under re-labelling we may assume that they are c_1 and c_{2m} .

Recall that c_1 , like any odd vertex, satisfies $d^+(c_1) > 5k/8$. Additionally, since $(v, c_1) \in \vec{D}$, by the pseudo-semidegree condition we have $d^-(c_1) > 5k/8$. This means that by pigeonhole c_1 an out-neighbour or an in-neighbour $u \notin V(C) \cup \{v\}$. In the former case we obtain a longer alternating path via $u, c_1, c_2, \dots, c_{2m}, v$. In the latter case we also obtain a longer alternating path, namely $u, c_1, v, c_{2m}, c_{2m-1}, \dots, c_2$. This concludes the odd case and the proof.

References

- [1] L. Addario-Berry, F. Havet, C. Linhares Sales, B. Reed, and S. Thomassé, Oriented trees in digraphs, *Discrete Mathematics* 313 (2013), 967–974.
- [2] B. Chen, X. Hou, and H. Zhou, Long antipaths and anticycles in oriented graphs, *preprint*, arXiv:2401.05205 (2024).
- [3] G. A. Dirac, Some theorems on abstract graphs, *Proc. Lond. Math. Soc.* 2(1) (1952), 69–81.
- [4] A. Ghouila-Houri, Une condition suffisante d’existence d’un circuit hamiltonien, *C.R. Acad. Sci. Paris* 25 (1960), 495–497.
- [5] R. Häggkvist, Hamilton cycles in oriented graphs, *Combin. Probab. Comput.* 2 (1993), 25–32.

- [6] R. Häggkvist and A. Thomason, Oriented Hamilton cycles in digraphs, *Journal of Graph Theory* 20 (1995), 471–479.
- [7] R. Häggkvist and A. Thomason, Oriented Hamilton cycles in oriented graphs, In *Combinatorics, Geometry and Probability*, Cambridge University Press (1997), 339–353.
- [8] B. Jackson, Long paths and cycles in oriented graphs, *J. Graph Theory* 5(2) (1981), 145–157.
- [9] P. Keevash, D. Kühn, and D. Osthus, An exact minimum degree condition for Hamilton cycles in oriented graphs, *Journal of the London Math. Soc.* 79 (2009), 144–166.
- [10] L. Kelly, Arbitrary Orientations of Hamilton Cycles in Oriented Graphs *Electronic Journal of Combinatorics* 18(1) (2011), P186.
- [11] L. Kelly, D. Kuhn, and D. Osthus, A Dirac type result on Hamilton cycles in oriented graphs, *Combin. Probab. Comput.* 17 (2008), 689–709.
- [12] T. Klimošová and M. Stein, Antipaths in oriented graphs, *Discrete Mathematics* 346 (2023), 113515.
- [13] J. Moon and L. Moser, On Hamiltonian bipartite graphs, *Israel Journal of Mathematics* 1 (1963), 163–165.
- [14] 4] L. Pósa, Hamiltonian circuits in random graphs, *Discrete Mathematics* 14 (1976), 359–364.
- [15] M. Stein, Tree containment and degree conditions, *Discrete Mathematics and Applications* (A. M. Raigorodskii and M. Th. Rassias, eds.), Springer International Publishing, Cham (2020) 459–486.
- [16] M. Stein, C. Zárate-Guerén, Antidirected subgraphs of oriented graphs, *Combinatorics, Probability and Computing* (2024), First View, pp. 1–21.
- [17] C. Thomassen, Long cycles in digraphs with constraints on the degrees, In *Surveys in Combinatorics, London Math. Soc. Lecture Notes*, Cambridge University Press), 38 (1979), 211–228.